Subdiffusive bound on Fredkin and Motzkin dynamics

Catherine McCarthy,¹ Hansveer Singh,¹ Sarang Gopalakrishnan,² and Romain Vasseur¹

¹Department of Physics, University of Massachusetts, Amherst, MA 01003, USA

²Department of Electrical and Computer Engineering,

Princeton University, Princeton, NJ 08544, USA

We identify a pseudolocal conserved charge in the Fredkin and Motzkin quantum spin chains and explore its consequences for the hydrodynamics of systems with Fredkin- or Motzkin-type kinetic constraints. We use this quantity to formulate an exact upper bound $\mathcal{O}(L^{-5/2})$ on the gap of the Fredkin and Motzkin spin chains. Our results establish that transport in kinetically constrained dynamical systems with Fredkin or Motzkin constraints is subdiffusive, with dynamical exponent $z \geq 5/2$.

Introduction — Transport in one-dimensional lattice models at nonzero temperature is generically diffusive [1]. Finding exceptions to diffusive behavior has been a longstanding quest: potential exceptions range from disordered quantum models exhibiting many-body localization [2, 3] to stochastic dynamical systems with kinetic constraints [4], which were first introduced as toy models of the glass transition. (Subsequently, kinetically constrained models have also been explored for their exotic low-energy quantum properties [5–8].) In many such examples, transport is entirely absent [9]; others, such as those involving fractons, exhibit subdiffusive transport [10–14]. The theory of fracton hydrodynamics was recently developed to explain transport with dynamical spacetime scaling $x \sim t^{1/z}$ with dynamical exponent z = 2n for integer n, using multipole conservation laws. However, there are many examples of anomalous diffusion that fall outside this framework, for example with non-integer z. Dynamics with Fredkin or Motzkin constraints, which we explore here, is an important such example.

Dynamics with Motzkin and Fredkin constraints were first introduced in Ref. [15] and [16], respectively. These works constructed frustration-free Hamiltonians with a critical ground state. Both spin chains are closely related, and have very similar physical properties. Throughout this letter we will focus on the Fredkin spin chain for concreteness, but all our results extend to the Motzkin case as well [17]. The energy gap of the Fredkin Hamiltonian is rigorously upper-bounded as L^{-2} for a system of size L [18] —therefore, despite being a one-dimensional critical point, it is not described by a conformal field theory [19, 20]. The Fredkin Hamiltonian can be reinterpreted as the transfer matrix of a Markov process with some kinetic constraints [20, 21]; on this interpretation, the energy gap is interpreted as the inverse of the mixing time for a system of size L, which scales as L^{-z} . Numerical evidence from simulations of this stochastic process gives values of z ranging 8/3 [21] to 2.69 [19, 20] (Ref. [22] also reported $z \approx 5/2$ for the Motzkin Markov process), suggesting that the rigorous bound z > 2 is not tight. The physical significance of the dynamical exponent z > 2 is transparent from the Markov-process interpretation, as it implies subdiffusive transport. This subdiffusive behavior defines a new hydrodynamic universality class, which applies to systems with Fredkin or Motzkin kinetic constraints. Moreover, the fractional value of z is incompatible with simple linear hydrodynamics involving strictly local charges, of the type considered in Ref. [10]. Possible mechanisms for this exponent include intrinsically nonlinear mechanisms [23], pseudolocal conserved charges with power-law tails [24], etc. Despite the numerical evidence, there is currently no proof that z > 2for Fredkin-constrained dynamics, and it remains controversial whether subdiffusion truly persists to arbitrarily late times in the thermodynamic limit.

In this letter, we identify a locally-conserved charge in Fredkin quantum spin chains and use this charge to prove a subdiffusive lower bound on the dynamical exponent, $z \geq 5/2$. After translating the system into the Dyck combinatorial alphabet, we show that irreducible Dyck words are locally-conserved charges by the Fredkin dynamics, and explore some of their properties. We demonstrate that the charge possesses algebraically-decaying tails that underpin all nontrivial static correlations relevant to Fredkin-constrained systems. Despite the pseudolocal nature of the charge, we argue that it is relevant to transport by proving its extensivity and additivity. We then construct a variational spin-wave ansatz in terms of the new charge, from which we produce an exact upper bound for the gap of the Fredkin Hamiltonian $\Delta \leq \mathcal{O}(L^{-5/2})$ and consequently a lower bound for the dynamical exponent, z > 5/2. Our result implies that transport in systems with a Fredkin kinetic constraint is subdiffusive. Our results also extend, with minor modifications, to the Motzkin spin chain [15, 25], see supplemental material [17].

The Fredkin spin chain — The Fredkin quantum spin chain was originally proposed in Ref. [16] as the spin- $\frac{1}{2}$ version of the Motzkin spin chain [15, 25]. The Hamiltonian is given by

$$H = \frac{1}{8} \sum_{i} (1 + \sigma_i^z) (1 - \vec{\sigma}_{i+1} \cdot \vec{\sigma}_{i+2}) + (1 - \sigma_{i+2}^z) (1 - \vec{\sigma}_i \cdot \vec{\sigma}_{i+1}),$$
(1)

where σ_i^{α} are the usual Pauli operators with $\alpha = x, y, z$. The Fredkin Hamiltonian conserves the total magnetization, $S_{\text{tot}}^z = \frac{1}{2} \sum_i \sigma_i^z$ and exhibits charge-parity (CP) symmetry. Additionally, this Hamiltonian is frustration free and the ground state is exactly known [16]. For the majority of this work, we consider periodic boundary conditions and restrict to the $S_{\text{tot}}^z = 0$ sector where the ground state is a uniform superposition of all $S_{\text{tot}}^z = 0$ states.

Dyck words — As initially outlined in [16], the Fredkin Hamiltonian naturally lends itself to a combinatorial description in terms of the Dyck alphabet. We briefly summarize important aspects of the Dyck alphabet here. The Dyck alphabet consists of two symbols: open and closed parentheses. Dyck words are defined as strings of parentheses in which every parenthesis is matched when read from left to right; for instance, (()) and ()() are both Dyck words, while)()(is not. A Dyck word necessarily contains an even number of parentheses and can be identified by its semilength, ℓ , which is half of the total number of parentheses. The number of Dvck words of semilength ℓ is given by the $\ell^{\rm th}$ Catalan number, C_{ℓ} . A class of Dyck words that will be important for the conservation law are known as *irreducible Dyck* words. Irreducible Dyck words are those in which the first parenthesis in the word is matched by the final parenthesis in the word; for example, ((())) and (()()) are both irreducible Dyck words of semilength $\ell = 3$, while (())(), ()(()), and ()()() are also Dyck words of semilength $\ell = 3$ but are not irreducible. Since irreducible Dyck words of semilength ℓ are of the form (...), where ... must itself be a Dyck word, there are $C_{\ell-1}$ irreducible words of semilength ℓ .

Matched parentheses charges — To map a spinsystem to the Dyck alphabet, up spins are replaced with an open parenthesis and down spins are replaced with a closed parenthesis. Following the usual quantum-to-classical correspondence combined with the frustration-free property, the Fredkin Hamiltonian maps onto a classical Markov process of constrained hard-core random walkers [21], where an up spin corresponds to a particle, and down spin to a hole. The allowed moves are summarized in Fig. 1, which we will refer as "Fredkin constraint" in the remainder of this letter. In this language, the Fredkin Hamiltonian and the corresponding Markov process) locally conserves irreducible Dyck words, and the number of matched parentheses. This constraint can be generalized to longer-range interactions, and defines a new hydrodynamic universality class distinct from ordinary diffusion [21]. We will subsequently refer to the corresponding conserved charge as "matched parentheses charge."

To illustrate the conservation law, consider the configuration (()(())). There are two $\ell = 1$ words in this configuration in addition to two other irreducible words of semilengths $\ell = 2$ and 4; each of these four irre-




FIG. 1. Fredkin conserved charges and their hydrodynamics. Top: Effective time-dependent dynamical exponent z extracted from the Fredkin structure factor $\langle S^z(x,t)S^z(0,0)\rangle$ in the Fredkin Markov process. Inset: moves allowed by the Fredkin Hamiltonian in the Dyck combinatorial language. Allowed moves locally conserve the number of matched parentheses, while forbidden moves locally create or destroy matched sets of parentheses. Bottom: Collapsed structure factors for the matched parentheses charges (left) and the original S_z charge (right) at times t = 200-400. Both structure factors collapse after rescaling by $t^{-1/z} f(x/t^{1/z})$, for $z \approx 8/3$ for numerically accessible time scales.

ducible words corresponds to a single matched parentheses charge. We take the midpoint of the underlying irreducible Dyck word, which lives on a link, to be the position of the matched parentheses charges– for example, the $\ell = 2$ word in the above configuration has position n = 5. We label the matched parentheses charges as q_n^{ℓ} and adopt the convention that the lower index denotes position and the upper index denotes semilength; if either index is irrelevant to the discussion at hand, it will be omitted.

In the configuration (()(())), the Fredkin Hamiltonian permits swapping the second and third parentheses to perform the move: $((|)(|())) \rightarrow (|) (((|)))$. The charges affected by this move (initially q_2^1 and q_4^4) are underlined and their positions are marked with a vertical bar. The illustrated move consists of two distinct parts: the movement of the initial q_2^1 charge to the left, and the response of the original q_4^4 charge, which changes semilength from $\ell = 4$ to $\ell = 3$ as it moves to the right.

To generalize the above discussion, moves permitted by the Fredkin Hamiltonian can be decomposed into two processes: the movement of a single q^1 charge and the response of some other q^{ℓ} charge. As illustrated in the inset of Fig. 1, allowed moves correspond to moving a q^1 charge to either the left or right by a single link. At $S_{\text{tot}}^z = 0$, a q^1 charge necessarily passes through the end parenthesis of some other q^{ℓ} charge, which in turn causes the q^{ℓ} charge to both move by one site and change in semilength to $\ell \pm 1$. These processes are the only relevant dynamics at play, which may be readily seen by rewriting the Fredkin Hamiltonian in terms of hard-core bosonic creation and annihilation operators. Denoting $(a_i^{\ell})^{\dagger}$ as the creation operator for a matched parentheses charge centered on link *i* with semilength ℓ , the Hamiltonian can be rewritten as:

$$H = \frac{1}{2} \sum_{i,\ell} n_{i-1}^{\ell+1} n_{i-\ell}^1 + n_i^{\ell} n_{i-\ell-1}^1 + n_{i+1}^{\ell+1} n_{i+\ell}^1 + n_i^{\ell} n_{i+\ell+1}^1 - (a_{i-1}^{\ell+1})^{\dagger} (a_{i-\ell}^1)^{\dagger} a_i^{\ell} a_{i-\ell-1}^1 - (a_{i+1}^{\ell+1})^{\dagger} (a_{i+\ell}^1)^{\dagger} a_i^{\ell} a_{i+\ell+1}^1 - \text{h.c.}$$
(2)

where $n_i^{\ell} = (a_i^{\ell})^{\dagger} a_i^{\ell}$. In this representation one can see that the total matched parentheses charge is $Q = \sum_{i,\ell} n_i^{\ell}$.

Pseudo-locality — In general, $Q_{\text{tot}} = \sum_{i=1}^{L} q_i$ is related to S_{tot}^z : for example, $Q_{\text{tot}} = L/2$ for a system of L spins with $S_{\text{tot}}^z = 0$ with periodic boundary conditions. As a result, any spin- $\frac{1}{2}$ Hamiltonian that preserves S_{tot}^z acting on a system with periodic boundary conditions will globally conserve the number of matched parentheses charges. In contrast, the Fredkin Hamiltonian has a stronger, *local* conservation law governing the matched parentheses charge and so one has a corresponding continuity equation [17]. The locality of the matched parentheses charge is not entirely obvious a priori- the total matched parentheses charge may be written as the sum of terms $Q = \sum_{n} q_{n}$, but the support of each q_{n} is not obviously bounded. Here, we first show that the matched parentheses charge has power-law tails that account for nontrivial static correlations between parentheses. We then prove that the charge is additive and extensive, thus demonstrating that even with its power-law tails, the charge is relevant to transport.

First, we consider the source of the algebraicallydecaying tails of the matched parentheses charge. All static correlations between matched parentheses charges arise from the fact that a parenthesis may only function as the endpoint of a single q^{ℓ} charge. To illustrate this, consider a single q_0^1 charge, corresponding to a configuration () at the origin. Since parentheses may not be shared between charges, and since the existence of a $q_{\pm 1}^1$ charge would require one of the endpoints of the q_0^1 to instead be one of the endpoints of the $q_{\pm 1}^1$ charge, the simultaneous existence of both charges is impossible. Extending this logic to an arbitrary link ℓ , the presence of q_0^1 precludes the existence of any $q_{\pm \ell}^{\ell}$ charge. The tails of the matched parentheses charge therefore depend on the probability of finding a q^{ℓ} charge ℓ away from a q^1 charge. Since the probability of finding a q^{ℓ} charge on a generic site is given by $C_{\ell-1}/2^{2\ell} \sim \ell^{-3/2}$, all nontrivial static correlations between the presence of q_0^1 and any other charges are given solely by the probability of finding $q_{\pm \ell}^{\ell}$. Therefore, we find $\langle q_m^1 q_n \rangle \sim |m-n|^{-3/2}$.

For a charge to be relevant to thermodynamics, it must be extensive and additive; despite having power-law tails, the matched parentheses charge q^{ℓ} indeed exhibits both of these properties. To see this, consider a region of $2L \to \infty$ total sites with $Q = \sum_{n=1}^{2L} q_n$. If the system is divided into two L-site subsystems A and B, the total number of matched parentheses charges in subsystem A, given by $Q_A = \sum_{n=1}^{L} q_n$, grows with L, and so $\langle Q_A \rangle \sim L$. One could worry that in spite of the extensivity of the matched parentheses charge, the power-law tails of the charge may lead to $Q \neq Q_A + Q_B$. From showing exactly that $\langle Q_A^2 \rangle_c \sim L$, we conclude that fluctuations of the total charge are sufficiently suppressed to guarantee additivity $Q \simeq Q_A + Q_B$, up to subleading corrections of $\mathcal{O}(\sqrt{L})$ – for further details, see [17]. Therefore, fluctuations in the value of Q_A are suppressed as $\sqrt{\langle Q_A^2 \rangle_c} / \langle Q_A \rangle \sim 1 / \sqrt{L}$, so the local density of matched parentheses is well-defined.

Since the matched parentheses charge is locally conserved, extensive, and additive, it is relevant to hydrodynamics. Using standard Markov chain simulation techniques, we numerically compute the structure factor of the matched parentheses charge $\langle q(x,t)q(0,0)\rangle_c$. We find that this structure factor exhibits the same collapse form $t^{1/z} f(x/t^{1/z})$ with $z \approx 8/3$ as the original S^z charge (Fig. 1). Transport of the matched parentheses charge is therefore subdiffusive, a fact which we will exploit to rigorously constrain the transport and low-energy properties of Fredkin constrained models.

Exact subdiffusive bound on z — The gap of the Fredkin Hamiltonian scales polynomially with system size L as $\Delta \sim L^{-z}$ [18]. To produce a bound for the gap, and by extension the dynamical exponent, we construct a variational spin-wave ansatz in terms of the matched parentheses charge – see Ref. [26] for a similar approach applied to other systems with fracton-like constraints. We find that $z \geq z_{\rm SW} = 5/2$ and derive an exact expression for this bound in [17]; below, we summarize the procedure to do so.

We construct a spin-wave trial wave function in terms of the matched parentheses charge q for the state $|k\rangle = \sum_{n} e^{ik \cdot n} q_n |GS\rangle$ with momentum $k = 2\pi j/L$, for j = 0, 1, ..., L - 1, where the $S_{\text{tot}}^z = 0$ ground state $|GS\rangle$ is given by the uniform superposition of all $S_{\text{tot}}^z = 0$ states. By standard variational considerations, the energy of the spin-wave state $|k\rangle$, given by $E_{\text{SW}}(k) = \langle k|H|k\rangle / \langle k|k\rangle$, serves as an upper bound on the ground state energy of the Fredkin Hamiltonian within the k^{th} crystal momentum sector. The gap of the Fredkin Hamiltonian may




FIG. 2. Exact bound on gap scaling. Exact diagonalization (ED) data shown for system sizes of L = 10 - 24 sites plotted with the exact ~ $L^{-5/2}$ analytical bound. Inset: exact expression for the dispersion $E_{\rm SW}(k)$ of the spin-wave ansatz, evaluated in the limit $L \to \infty$.

then be bounded by evaluating the energy of the spinwave state in the $k = 2\pi/L$ sector.

Using translation invariance, we have $\langle k|H|k \rangle = L \sum_{r} e^{ik \cdot n} \langle GS|q_0Hq_n|GS \rangle$, and using $\langle k|k \rangle \sim L$ [17], the task of bounding the gap reduces to evaluating the Fourier transform of the correlator $C(n) \equiv \langle GS|q_0Hq_n|GS \rangle$. From $H|GS \rangle = 0$, we may write $C(n) = \langle GS|q_0[H,q_n|GS \rangle$ with $i[H,q_n] = j_n - j_{n+1}$, where j_n is the current of the matched parentheses charge on site n. This yields

$$C(n) = i \langle GS | q_0 j_{n+1} | GS \rangle - i \langle GS | q_0 j_n | GS \rangle.$$
(3)

We obtain exact expressions for the charge-current correlator $\langle GS|q_0j_n|GS \rangle$ in the $S_{\text{tot}}^z = 0$ ensemble for any system size L; for further details, see [17]. We find that $\sum_n n^2 C(n) = 0$, ruling out diffusive $\sim k^2$ contributions to $E_{\text{SW}}(k)$. Instead, the low momentum scaling of $E_{\text{SW}}(k) \sim k^{5/2}$ is governed by the long distance decay of $C(n) \sim n^{-7/2}$, which follows from the charge-current correlator $\langle GS|q_0j_{n+1}|GS \rangle$ decaying as $\sim n^{-5/2}$. This $\sim n^{-5/2}$ decay may be readily obtained by computing the probability that processes such as $(\dots()) \rightarrow (\dots)()$ occur, since these processes make up all possible contributions to the charge-current correlator [17].

Our results provide the exact energy of the spin-wave state $E_{SW}(k)$ for any system size L, which we verify constitutes an upper bound of the gap of the Fredkin Hamiltonian (Fig. 2). In the thermodynamic limit, we find the dispersion relation

$$\lim_{L \to \infty} E_{\rm SW}(k) = \frac{4\sqrt{2}}{3} \cos\left(\frac{k-\pi}{4}\right) \sin\left(\frac{k}{2}\right)^{5/2}, \qquad (4)$$

with $k \in [0, 2\pi]$. The scaling of the gap follows from the energy of the first excited spin-wave state $E_{SW}(k =$ $2\pi/L$) = $\Delta_{\rm SW} \sim L^{-5/2}$. Since the gap of the Fredkin Hamiltonian is upper-bounded as $\Delta \leq \Delta_{\rm SW}$, we obtain the following lower bound on the dynamical exponent:

$$z \ge z_{\rm SW} = 5/2. \tag{5}$$

The lower bound on the Fredkin dynamical exponent is the main result of this letter. To the best of our knowledge, this is the first proof that Fredkin-constrained systems exhibit subdiffusive transport; this follows from our physically-motivated spin-wave ansatz. We emphasize that since we obtain the energy of the spin-wave state exactly, even for finite system size L (see [17]), this result is entirely exact.

Discussion — In this work, we identified a locallyconserved quantity in the Fredkin spin chain and used it to prove a subdiffusive lower bound for the Fredkin dynamical exponent. This conservation law corresponds to the number of irreducible Dyck words [16], which we name the "matched parentheses charge". The matched parentheses charges have algebraically-decaying tails leading to non-trivial static correlations. However, in addition to obeying a local conservation law, these pseudo-local matched parentheses charges are both extensive and additive and thus relevant to transport. We find numerically that the matched parentheses charge features emergent scaling behavior with the same dynamical exponent $z \simeq 8/3$ as the original S_{tot}^z charge. We construct a variational ansatz from the matched parentheses charge for the first excited of the Fredkin spin chain in a fixed crystal momentum sector. This physicallymotivated trial wavefunction provides an upper bound $\mathcal{O}(L^{-5/2})$ on the gap of the Fredkin spin chain, thus providing a lower bound on the dynamical exponent of Fredkin constrained systems, i.e. $z \ge 5/2$. Our results also extend to the Motzkin spin chain [17].

The bound obtained from the spin-wave ansatz represents the first proof that transport in Fredkin and Motzkin constrained systems is subdiffusive. While obtaining a subdiffusive bound constitutes significant progress in understanding Fredkin/Motzkin dynamics, the exact value of the dynamical exponent still remains an open question. A derivation of the true dynamical exponent and detailing the mechanism responsible for its value would represent a clear direction for future research. We also note that Fredkin and Motzkin constrained systems seem to be – to the best of our knowledge - one of the only examples of a system with local interactions but with a conserved charge having nonlocal densities as relevant hydrodynamic modes. While the consequences of pseudo-local charges have mostly been explored for exponentially decaying tails [27–32], further exploring the consequences of densities with algebraically-decaying correlations would also be an interesting area for future research.

Acknowledgements – H.S. and R.V. thank Aaron Friedman and Brayden Ware for earlier collaborations on this topic. C.M. acknowledges support from NSF GRFP-1938059. This work was supported by the US Department of Energy, Office of Science, Basic Energy Sciences, under award No. DE-SC0023999 (H.S. and R.V.). S.G. and R.V. acknowledge hospitality of KITP during the DYNISQ22 follow-on program "Phases of active quantum matter". KITP is supported by grant NSF PHY-2309135.

- B. Bertini, F. Heidrich-Meisner, C. Karrasch, T. Prosen, R. Steinigeweg, and M. Žnidarič, Rev. Mod. Phys. 93, 025003 (2021).
- [2] R. Nandkishore and D. A. Huse, Annual Reviews of Condensed Matter Physics 6, 15 (2015).
- [3] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Rev. Mod. Phys. **91**, 021001 (2019).
- [4] F. Ritort and P. Sollich, Advances in physics 52, 219 (2003).
- [5] E. Lake and T. Senthil, Phys. Rev. Lett. 131, 043403 (2023).
- [6] E. Lake, M. Hermele, and T. Senthil, Phys. Rev. B 106, 064511 (2022).
- [7] A. Keesling, A. Omran, H. Levine, H. Bernien, H. Pichler, S. Choi, R. Samajdar, S. Schwartz, P. Silvi, S. Sachdev, et al., Nature 568, 207 (2019).
- [8] R. Verresen, M. D. Lukin, and A. Vishwanath, Phys. Rev. X 11, 031005 (2021).
- [9] A. Morningstar, V. Khemani, and D. A. Huse, 101, 214205.
- [10] A. Gromov, A. Lucas, and R. M. Nandkishore, Phys. Rev. Research 2 (2020), 10.1103/physrevresearch.2.033124.
- [11] J. Feldmeier, P. Sala, G. De Tomasi, F. Pollmann, and M. Knap, Phys. Rev. Lett. **125**, 245303 (2020).
- [12] J. Iaconis, S. Vijay, and R. Nandkishore, Phys. Rev. B 100 (2019), 10.1103/physrevb.100.214301.

- [13] P. Glorioso, J. Guo, J. F. Rodriguez-Nieva, and A. Lucas, "Breakdown of hydrodynamics below four dimensions in a fracton fluid," (2021), arXiv:2105.13365 [condmat.str-el].
- [14] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, Phys. Rev. X 10, 011047 (2020).
- [15] S. Bravyi, L. Caha, R. Movassagh, D. Nagaj, and P. W. Shor, Phys. Rev. Lett. **109**, 207202 (2012).
- [16] O. Salberger and V. Korepin, "Fredkin spin chain," (2016), arXiv:1605.03842 [quant-ph].
- [17] See Supplemental Material for additional details.
- [18] R. Movassagh, arXiv preprint arXiv:1609.09160 (2016).
- [19] X. Chen, E. Fradkin, and W. Witczak-Krempa, Physical Review B 96 (2017), 10.1103/physrevb.96.180402.
- [20] X. Chen, E. Fradkin, and W. Witczak-Krempa, Journal of Physics A: Mathematical and Theoretical 50, 464002 (2017).
- [21] H. Singh, B. A. Ware, R. Vasseur, and A. J. Friedman, Phys. Rev. Lett. **127**, 230602 (2021).
- [22] J. Richter and A. Pal, Phys. Rev. Res. 4, L012003 (2022).
- [23] V. Popkov, A. Schadschneider, J. Schmidt, and G. M. Schütz, Proceedings of the National Academy of Sciences 112, 12645 (2015).
- [24] J.-P. Bouchaud and A. Georges, Physics Reports 195, 127 (1990).
- [25] R. Movassagh and P. W. Shor, Proceedings of the National Academy of Sciences 113, 13278?13282 (2016).
- [26] O. Ogunnaike, J. Feldmeier, and J. Y. Lee, Phys. Rev. Lett. 131, 220403 (2023).
- [27] T. Prosen, Journal of Physics A: Mathematical and General 31, L645 (1998).
- [28] T. Prosen, Phys. Rev. E **60**, 3949 (1999).
- [29] T. Prosen, Phys. Rev. Lett. 106, 217206 (2011).
- [30] T. Prosen and E. Ilievski, Phys. Rev. Lett. 111, 057203 (2013).
- [31] B. Doyon, Communications in Mathematical Physics 351, 155 (2017).
- [32] B. Doyon, Journal of Statistical Physics 186, 25 (2022).

Supplemental material for "Subdiffusive bound on the Fredkin dynamical exponent"

Catherine McCarthy, Hansveer Singh, and Romain Vasseur Department of Physics, University of Massachusetts, Amherst, MA 01003, USA

> Sarang Gopalakrishnan Department of Electrical and Computer Engineering, Princeton University, Princeton, NJ 08544, USA (Dated: July 15, 2024)

I. PROOF OF ADDITIVITY OF MATCHED PARENTHESES CHARGE

In the main text, we argue that despite the fact that the matched parentheses charge has algebraically-decaying tails, the charge is extensive and additive and thus sufficiently local to be relevant to thermodynamics and hydrodynamics. In this section, we derive this statement explicitly. Consider a system of size 2L with open boundary conditions in a random (infinite temperature) state, with total number of matched parentheses Q. Let us cut the system into two halves A and B of size L, and denote by Q_A and Q_B the number of matched parentheses in subsystems A and B, respectively.

In this appendix, we derive the full probability distribution of the charge Q_A (or Q_B), and provide exact expressions for both $\langle Q_A \rangle$ and $\langle Q_A^2 \rangle_c$, showing that they scale asymptotically as ~ L. As a result, typical values of Q_A and Q_B are overwhelmingly likely to be equal to L/2, up to $\mathcal{O}(\sqrt{L})$ corrections. This implies that the matched parentheses charge is additive: $Q \simeq Q_A + Q_B$, again up to subleading $\mathcal{O}(\sqrt{L})$ corrections.

To establish this result, first consider a subsystem of L sites in a fixed charge sector with L - n open and n closed parentheses. The number of states that contain Q matched parentheses charge is given by Catalan's triangle:

$$T(L,Q) = \binom{L}{Q} - \binom{L}{Q-1},\tag{S1}$$

for $Q \leq \min(n, L - n)$. Generalizing to an infinite temperature state requires summing over all possible charge sectors. Therefore, for a system of L sites at infinite temperature, the number of possible states that have Q_A matched parentheses is given by the following expression:

$$T_{\rm inf}(L, Q_A) = (L - 2Q_A + 1)T(L, Q_A), \tag{S2}$$

for $Q_A \leq L/2$. The probability $P_L(Q_A)$ of finding a state with Q_A matched parentheses charges is thus given by:

$$P_L(Q_A) = \frac{T_{\inf}(L, Q_A)}{2^L}.$$
 (S3)

Given the probability of finding a subsystem of L sites that contains Q_A matched parentheses charges, it is straightforward to compute all moments of total matched parentheses charge Q_A in that interval.

We can derive the asymptotic behavior of these quantities in the limit $L \gg 1$ using Stirling's formula. Introducing the matched parentheses density $q_A = Q_A/L$, we find that the probability distribution $P_L(q_A) = P_L(Q_A = q_A L)L$ is maximized by the maximal allowed value of q_A , $q_A = q_* = 1/2$. Expanding about this maximum $q_A = 1/2 - \delta$ with $\delta > 0$, we find

$$P_L(\delta) \underset{L \to \infty}{\sim} 8L^{3/2} \sqrt{\frac{2}{\pi}} \delta^2 \mathrm{e}^{-2L\delta^2}.$$
 (S4)

A similar distribution in terms of height variables was derived in Refs. 1 and 2. Using this distribution, we can derive the asymptotic behavior of all cumulants of $Q_A = L/2 - L\delta$. This yields

$$\langle Q_A \rangle = \frac{L}{2} - \sqrt{\frac{2L}{\pi}} + \mathcal{O}(1),$$
 (S5)

and

$$\langle Q_A^2 \rangle_c = \left(\frac{3}{4} - \frac{2}{\pi}\right) L + \dots$$
(S6)

In the above setup, the matched parentheses charge only receives contributions from matched parentheses residing within the region. One could also consider situations where the matched parentheses charge receive contributions outside the subregion. For instance, if one considers a subregion $[-\ell/2, \ell/2]$ of a larger system of size $L \gg \ell$ then any matched parentheses with ends located at -y and y (including $y > \ell/2$) contributes to the matched parentheses charge in that interval. Despite this difference, we have numerically confirmed that this does not change the scaling behavior of the matched parentheses charge with subregion size.

II. DERIVATION OF EXACT BOUND ON GAP SCALING

In the main text, we construct a parameter-free variational spin-wave ansatz for the Fredkin Hamiltonian in terms of the matched parentheses charge. Here, we derive the an exact expression for the energy $E_{\rm SW}(k)$ of these spin-wave states, and use them to upper bound the gap of the Fredkin Hamiltonian, $\Delta \sim L^{-z}$, with L the size of the system. We emphasize that all of the expressions shown below are obtained exactly for finite-size systems.

A. Spin wave ansatz and charge-current correlations

We construct a (unnormalized) spin-wave ansatz in terms of the the matched parentheses charge as $|k\rangle = \sum_{n} e^{ik \cdot n} q_n |GS\rangle$. Recall that $q_n = \sum_{\ell} q_n^{\ell}$ counts the total number of matched parentheses centered on link n, while $q_n^{\ell} = (a_n^{\ell})^{\dagger} a_n^{\ell}$ corresponds to a set of matched parentheses separated by 2ℓ centered on link n. As in the main text, we adopt the convention that lower indices denote position while upper indices denote separation; if either the upper or lower index is irrelevant to the discussion at hand, it will be omitted. Here, we work in the $S_{\text{tot}}^z = 0$ ensemble, where the ground state of the Fredkin chain $|GS\rangle$ is given by the uniform superposition of all $S_{\text{tot}}^z = 0$ states.

Since $|k\rangle$ is a state with definite crystal momentum, the energy expectation value

$$E_{\rm SW}(k) = \frac{\langle k|H|k\rangle}{\langle k|k\rangle} = \frac{\sum_{n,m} e^{ik \cdot (n-m)} \langle GS|q_m H q_n |GS\rangle}{\sum_{n,m} e^{ik \cdot (n-m)} \langle GS|q_m q_n |GS\rangle},\tag{S7}$$

provides an upper bound on the ground state energy of the Hamiltonian H in the sector with momentum k. Since the lowest excited-state of the Fredkin Hamiltonian is in the $k = 2\pi/L$ sector, and since $H|GS\rangle = 0$, the variational spin-wave energy provides an upper bound on the Fredkin gap $\Delta = E(k = 2\pi/L) \leq E_{SW}(k = 2\pi/L)$. Because the total number of matched parentheses $\sum_{n} q_n$ is conserved, we have $E_{SW}(k = 0) = 0$, and so the spin-wave ansatz is guaranteed to be gapless as $k \to 0$.

Next, we invoke translation invariance and consider the quantity $\langle GS | q_0 H q_n | GS \rangle$. Using $H | GS \rangle = 0$, this correlator can be expressed as $\langle GS | q_0[H, q_n] | GS \rangle$. We note that $\dot{q}_n = i[H, q_n] = j_n - j_{n+1}$, where j_n is the current of the matched parentheses charge on site n; therefore, this quantity can be expressed as a discrete derivative:

$$\langle GS| q_0 H q_n | GS \rangle = i(\langle q_0 j_{n+1} \rangle_0 - \langle q_0 j_n \rangle_0), \tag{S8}$$

where we adopt the notation that $\langle \bullet \rangle_0$ corresponds to $\langle GS | \dots | GS \rangle$. Therefore, obtaining an expression for the charge-current correlations $\langle q_0 j_n \rangle_0$ will directly lead to an exact bound for the scaling of $E(k = 2\pi/L)$.

B. Source of charge-current correlations

In the picture of the matched parentheses charge, it becomes clear that the source of all Fredkin dynamics is the movement of q^1 charges, which are matched parentheses of the form (). When a q^1 charge moves, it necessarily disrupts the end parenthesis of a larger q^{ℓ} charge. This can be pictured as the q^1 moving past one of the matched end parentheses of the q^{ℓ} charge, which we will call moving "inside" or "outside" of the q^{ℓ} charge. The movement of the q^1 charge triggers the process $q_n^{\ell} \to q_{n\pm 1}^{\ell\pm 1}$.

All of these considerations are captured by the current operator $j_n = \sum_{\ell} j_n^{\ell}$, where each j_n^{ℓ} may be written explicitly

in terms of ladder operators as:

$$j_{n}^{\ell} = -\frac{i}{2} \Big((a_{n+\ell-1}^{\ell+1})^{\dagger} (a_{n}^{1})^{\dagger} a_{n+\ell}^{\ell} a_{n-1}^{1} - (a_{n-1}^{\ell+1})^{\dagger} (a_{n-\ell}^{1})^{\dagger} a_{n}^{\ell} a_{n-\ell-1}^{1} + (a_{n}^{\ell+1})^{\dagger} (a_{n+\ell-1}^{1})^{\dagger} a_{n-1}^{\ell} a_{n+\ell}^{1} - (a_{n-\ell}^{\ell+1})^{\dagger} (a_{n-1}^{1})^{\dagger} a_{n-\ell-1}^{\ell} a_{n}^{1} - \text{h.c.} \Big)$$
(S9)

Since the only source of dynamics from the Fredkin Hamiltonian involve locally moving q^1 charges, all nontrivial correlations between a charge sitting on link 0 and a current on site n must come from the coincidence of two scenarios: first, that the charge sitting on link 0 corresponds to matched parentheses separated by $\ell \pm 1$, and second, that a charge $q_{\pm(\ell\mp 1)}^1$ disturbs the end parenthesis of the $q_0^{\ell\pm 1}$ charge. This statement has an intuitive interpretation– a current may only be correlated with a q^{ℓ} charge if a q^1 charge either moves inside or outside a q^{ℓ} charge, which consequently changes the separation of the charge $q^{\ell} \rightarrow q^{\ell\pm 1}$. In order to move inside or outside of the q^{ℓ} charge at a given timestep, since there are no non-local q^1 hops encoded in the Fredkin Hamiltonian, there must be a q^1 charge located directly adjacent to one of the matched end parentheses of the q^{ℓ} charge immediately before the hop. Therefore, the task of understanding the static charge-current correlations in Fredkin reduces to finding the probability of these configurations. Exact expressions for these probabilities are derived in the following section.

C. Probabilities

First, we focus on the process $()(\ldots)^{\ell} \to (() \ldots)^{\ell+1}$, in which $(\ldots)^{\ell}$ pictorially represents a q^{ℓ} charge. We denote the probability of finding a configuration of the form $()(\ldots)^{\ell}$ or $(\ldots)^{\ell}()$ conditional on having a $(\ldots)^{\ell}$ as $p_{\text{out}}(\ell)$. From direct counting, we find that $p_{\text{out}}(\ell) = 7/16$ in the thermodynamic limit since the system is locally at infinite temperature. For a system of L sites in the $S_{\text{tot}}^z = 0$ ensemble, however, the expression for $p_{\text{out}}(\ell, L)$ becomes dependent on both ℓ and L due to additional correlations that arise from finite-size effects.

Since the Dyck word corresponding to any q^{ℓ} charge contains equal numbers of up and down spins, the $L - 2\ell$ sites in the system that are not part of the q^{ℓ} charge must also contain the same number of up and down spins when the system is at $S_{\text{tot}}^z = 0$. There are $\binom{L-2\ell}{(L-2\ell)/2}$ possible states for these $L - 2\ell$ sites. Consider a Dyck word of semilength $\ell + 1$ of the form $()(\ldots)^{\ell}$; since $()(\ldots)^{\ell}$ also contains only matched sets of parentheses, the $L - (\ell + 2)$ sites in the system that are not part of q^{ℓ} must contain the same number of up and down spins. Therefore, there are $\binom{L-2\ell-2}{(L-2\ell-2)/2}$ possible configurations for these $L - 2\ell - 2$ sites. Finally, taking care to avoid double-counting the $\binom{L-2\ell-4}{(L-2\ell-4)/2}$ configurations of the form $()(\ldots)^{\ell}()$, we obtain the following expression for $p_{\text{out}}(\ell, L)$:

$$p_{\rm out}(\ell,L) = \frac{2\binom{L-2\ell-2}{(L-2\ell-2)/2} - \binom{L-2\ell-4}{(L-2\ell-4)/2}}{\binom{L-2\ell}{(L-2\ell-4)/2}}.$$
(S10)

Next, we focus on the process $(() \dots)^{\ell} \to ()(\dots)^{\ell-1}$. We denote the probability of finding a configuration of the form $(() \dots)^{\ell}$ conditional on having a $(\dots)^{\ell}$ as $p_{in}(\ell)$. Unlike when a q^1 charge sits directly outside of a q^{ℓ} charge, a q^1 sitting directly inside of a q^{ℓ} charge is subject to strong correlations from being embedded within a Dyck word. In total, there are $C_{\ell-1}$ possible Dyck words corresponding to a q^{ℓ} charge, where C_{ℓ} are Catalan numbers. Recall that this was obtained from there fact that of the C_{ℓ} Dyck word of semilength ℓ , only those of the form $(\dots)^{\ell}$ correspond to q^{ℓ} charges; therefore, the \dots in $(\dots)^{\ell}$ must also be a Dyck word of semilength $\ell - 1$, of which there are $C_{\ell-1}$ possibilities. By similar logic, there are $\ell - 2$ ways to obtain a q^{ℓ} charge of the form $(() \dots)^{\ell}$, where \dots is a path of length $\ell - 2$. To obtain the expression for $p_{in}(\ell)$, we consider both words of the form $(() \dots)^{\ell}$ and $(\dots ())^{\ell}$, of which there are $2 C_{\ell-2}$; however, care must be taken to not count configurations like $(() \dots ())^{\ell}$ twice. Since \dots in these configurations must be a Dyck path of semilength $\ell - 3$, the expression for $p_{in}(\ell)$ is given by the following:

$$p_{\rm in}(\ell) = \frac{2C_{\ell-2} - C_{\ell-3}}{C_{\ell-1}}.$$
(S11)

We note that the boundary cases for $\ell \leq 2$ must be handled separately. Since the only possible configuration for a q^2 charge is (()), $p_{in}(2) = 1$. Additionally, since there is no possible way for a q^1 charge to be embedded in another q^1 charge, $p_{in}(1) = 0$. We also note that in the limit $\ell \gg 1$, the correlations that come from being embedded in a Dyck path become weaker and $p_{in}(\ell) \rightarrow 7/16$.

The expressions in eqs. S10 and S11 for $p_{in}(\ell)$ and $p_{out}(\ell, L)$ are conditioned on finding q^{ℓ} on a particular link; however, in order to obtain the expression for $\langle q_0 j_n \rangle$, these conditional probabilities need to be multiplied by the probability to have matched parentheses q_0^{ℓ} in the first place. To produce the relevant expressions, we write down

4

the probability $p_q(\ell, L)$ of finding a q^ℓ charge on any particular link in a system of L sites. Overall, there are $\binom{L}{L/2}$ possible states that the system can be found in as a while. Since there are $C_{\ell-1}$ possible Dyck paths corresponding to a q^ℓ charge, and since there are $\binom{L-2\ell}{(L-2\ell)/2}$ possible states that the the $L-2\ell$ sites outside of q^ℓ may take, $p_q(\ell, L)$ is given by the following expression:

$$p_q(\ell, L) = \frac{C_{\ell-1} \left(\frac{L-2\ell}{(L-2\ell)/2} \right)}{\binom{L}{L/2}}.$$
(S12)

D. Exact expression for $E_{SW}(k)$

With the probabilities $p_{in}(\ell)$, $p_{out}(\ell, L)$, and $p_q(\ell, L)$ obtained in the previous section, the charge-current correlator is given exactly by:

$$i\langle q_0 j_n \rangle_0 = \frac{1}{2} \left(p_{\text{out}}(n-1,L) p_q(n-1,L) - p_{\text{in}}(n+1) p_q(n+1,L) \right).$$
(S13)

From this, we may exactly compute the correlator $\langle GS | q_0 H q_n | GS \rangle = i(\langle q_0 j_{n+1} \rangle_0 - \langle q_0 j_n \rangle_0)$ explicitly in terms of probabilities. The exact dispersion for the spin-wave state is given by:

$$E_{\rm SW}(k) = \frac{\sum_{n} i e^{ik \cdot n} (\langle q_0 j_{n+1} \rangle_0 - \langle q_0 j_n \rangle_0)}{(3L - 4)/4(L - 1)},\tag{S14}$$

where we used the normalization $\langle k|k \rangle = \frac{L(3L-4)}{4(L-1)}$ valid for $k = 2\pi j/L$ with j odd – for j > 0 even, this factor should be replaced by $\langle k|k \rangle = 3(L/2)^2/(L-1)$. This expression may used to evaluated the exact value of $E_{\rm SW}(k = 2\pi/L)$ for any L.

The asymptotics of this expression may be derived in a straightforward way. In the thermodynamic limit $L \to \infty$, the correlator $\langle GS | q_0 H q_x | GS \rangle$ decays as

$$\langle GS | q_0 H q_x | GS \rangle \underset{x \to \infty}{\sim} \frac{15}{64\sqrt{\pi}x^{7/2}},$$
(S15)

at long distances, giving rise to a contribution $\sim k^{5/2}/(3\sqrt{2})$ to $E_{\rm SW}(k)$ at small momenta. In general, one would also expect a $\sim k^2$ contribution in $E_{\rm SW}(k)$ dominating this anomalous behavior coming from short distance physics. However, this diffusive contribution vanishes exactly because of the sum rule $\sum_n n^2 \langle GS | q_0 H q_n | GS \rangle = 0$, which follows from our exact expressions. In the thermodynamic limit where k becomes continuous, we find the exact dispersion relation

$$\lim_{L \to \infty} E_{\rm SW}(k) = \frac{4\sqrt{2}}{3} \cos\left(\frac{k-\pi}{4}\right) \sin\left(\frac{k}{2}\right)^{5/2},\tag{S16}$$

valid when $k \in [0, 2\pi]$. We thus conclude that $E_{\rm SW}(k) \sim k^{5/2}/(3\sqrt{2})$ as $k \to 0$, leading to the scaling $E_{\rm SW}(k = 2\pi/L) = \Delta_{\rm SW}(L) \sim L^{-5/2}$ as $L \to \infty$. This expression provides a rigorous upper bound on the Fredkin gap $\Delta = E(k = 2\pi/L)$. In turn, this upper bound on the gap $\Delta \sim L^{-z}$ translates to a lower bound on the dynamical exponent

$$z \ge z_{\rm SW} = \frac{5}{2},\tag{S17}$$

which is the main result of this letter.

III. RESULTS FOR MOTZKIN

The main text and the previous two sections have dealt with Fredkin spin chains and Fredkin-constrained systems. The Fredkin spin chain was originally proposed as the half-integer spin version of a model known as the Motzkin spin chain³. Given the similarities between the two models, it should come as no surprise that the same procedure as the one used for Fredkin may be employed to derive an upper bound on the scaling of the gap of the Motzkin Hamiltonian, which also serves as a lower bound on the Motzkin dynamical exponent.

Here, mirroring the format of the main text, we outline the procedure to bound the gap of the Motzkin Hamiltonian. After reviewing the Motzkin spin chain, we provide an overview of the relevant combinatorics and identify the spin-1 version of the matched parentheses charge. As with Fredkin, the Motzkin matched parentheses charge can be shown to be pseudolocal in that, despite having power-law tails, it is extensive and additive. The gap of the Motzkin Hamiltonian $\Delta_{\rm M}$ may then be bounded through the construction of a variational spin-wave ansatz, which features the same scaling as the gap of the Fredkin Hamiltonian, $\Delta_{\rm M} \leq \mathcal{O}(L^{-5/2})$. We then arrive at our main result of a bound on the Motzkin dynamical exponent, $z \geq 5/2$.

A. Background

1. The Motzkin spin chain

The Motzkin spin chain was originally proposed as an interesting example of a frustration-free spin-1 chain with a highly-entangled ground state³. With periodic boundary conditions, Motzkin Hamiltonain may be expressed as the sum of nearest-neighbor projectors, $H = \sum_{n,n+1} \prod_{n,n+1}$. Adopting the notation used by Ref.², the onsite Hilbert space is spanned by the states $|u\rangle$, $|d\rangle$ and $|0\rangle$, and the 2-site projector $\prod_{n,n+1}$ may be expressed as:

$$\Pi_{n,n+1} = |D\rangle_{n,n+1} \langle D| + |U\rangle_{n,n+1} \langle U| + |V\rangle_{n,n+1} \langle V|, \qquad (S18)$$

where:

$$|D\rangle = \frac{1}{\sqrt{2}} (|0d\rangle - |d0\rangle),$$

$$|U\rangle = \frac{1}{\sqrt{2}} (|0u\rangle - |u0\rangle),$$

$$|V\rangle = \frac{1}{\sqrt{2}} (|00\rangle - |ud\rangle).$$

(S19)

With open boundary conditions, the ground state of the Motzkin Hamiltonian for a system of L sites is given by the uniform superposition of all Motzkin words of length L (see the following subsection)³. When considering periodic boundary conditions, the Motzkin ground state is instead given by the uniform superposition of all $S_{tot}^z = 0$ states consisting of L sites.

2. Motzkin words

In the context of Fredkin-constrained systems, spin- $\frac{1}{2}$ systems may be recast in terms of the Dyck combinatorial alphabet by replacing up-spins with an open parenthesis and down-spins with a closed parentheses. The description of spin-1 models requires generalizing the Dyck alphabet to include a symbol for zero-spins, which may be accomplished by replacing zero-spins with a "flat" symbol, -. Strings of parentheses and flats can be interpreted as a combinatorial object known as a Motzkin path; as suggested by the model's name, this a natural combinatorial language for Motzkin-constrained systems. We briefly review some important aspects of Motzkin combinatorics here.

Motzkin words are strings of parentheses and flats that, read from left to right, do not contain unmatched parentheses. For example, the Motzkin words of length $\ell = 3$ are - - -, -(), (-), and ()-. The number of Motzkin words of length ℓ is given by the ℓ^{th} Motzkin number, $M_{\ell} = \sum_{k=0}^{\ell/2} {\ell \choose 2k} C_k$. Note that in Fredkin, we identify words by their *semilength*, given that Dyck words necessarily contain an even number of parentheses; in contrast, we identify Motzkin words by their *length* since the presence of flats allows for words to be comprised of an odd number of symbols.

Just as irreducible Dyck words are crucial to the construction of the Fredkin matched parentheses charge, irreducible Motzkin words will play the same role for this model. Motzkin words of length $\ell \geq 2$ that are irreducible begin and end with a matched set of parentheses; for example, the only irreducible Motzkin word of $\ell = 3$ is (-). Since irreducible Motzkin words must be of the form (...), where ... itself is a Motzkin word, there are $M_{\ell-2}$ irreducible Motzkin words of length ℓ . For the case of $\ell = 1$, a single flat – is itself an irreducible word.

B. The matched parentheses charge in Motzkin

As outlined in the main text, each matched parentheses charge in Fredkin corresponds to the midpoint of a single irreducible word. The conservation of the number of irreducible words in Fredkin therefore follows trivially from the

fact that number of open (and closed) parentheses is conserved within a fixed charge sector. In contrast, the number of irreducible words is not conserved in Motzkin for a fixed charge sector due to the allowed move $-- \leftrightarrow$ (), in which two flats (irreducible $\ell = 1$ words) merge into a single irreducible word of length $\ell = 2$.

A matched parentheses charge given by the midpoints of irreducible words may still be constructed for Motzkinconstrained system; however, the midpoints of paths of length $\ell \geq 2$ must correspond to two charges, while flats $(\ell = 1)$ only carry a single charge. This modification ensures the conservation of the total matched parentheses charge during the allowed move $-- \Leftrightarrow ()$, since there are two charges on either side of the process. We will continue to refer to these charges at the matched parentheses charges and label them as q_n^{ℓ} , where n gives the location of the underlying irreducible word's midpoint and ℓ denotes the word's length. Note that a consequence of the existence of odd-length words is that charges are permitted to live on both sites and links; therefore, the index for the matched parentheses charge runs from n = 1 to 2L. In the $S_{tot}^z = 0$ charge sector for a system of L sites with periodic boundary conditions, the total Motzkin matched parentheses charge is $Q = \sum_n q_n = L$.

In each model, the smallest irreducible word plays the role of a freely-moving object that facilitates the motion of larger words when it disrupts one of their endpoints. The allowed Fredkin move $()_n(\ldots)_{n+\ell}^{\ell} \to (()_{n+1} \ldots)_{n+\ell-1}^{\ell+1}$ is equivalent to the allowed Motzkin move $-_n(\ldots)_{n+\ell+1}^{\ell} \to (-_{n+2} \ldots)_{n+\ell}^{\ell+1}$ in that it shows the process of the smallest irreducible word of the model passing the end parenthesis of the word (\ldots) , which leads to the displacement of the charge(s) corresponding to the larger word. The fact that Motzkin charges may live on both links and sites leads to some notable differences in labelling the midpoint; in particular, the flat involved in this process changes its position by two, while the charges corresponding to the larger word only move one space. Note that the process $-- \to ()$ behaves differently, as the flats each move one space when forming the $\ell = 2$ word; therefore, this process must be dealt with carefully and separately.

Taking into account all of the considerations outlined above, the Motzkin Hamiltonian may be rewritten in terms of the matched parentheses charge. Writing the hard-core bosonic ladder operators for matched parentheses charges of length ℓ whose midpoint position is i as $a_i^{\ell(\dagger)}$, the Hamiltonian is given by:

$$H = \frac{1}{4} \left\{ \left(-\frac{1}{2} \sum_{i=1}^{2L} \sum_{\ell=2}^{L-1} \left((a_{i-1}^{\ell+1\dagger})^2 a_{i-\ell+1}^{1\dagger} (a_i^{\ell})^2 a_{i-\ell-1}^1 + (a_{i+1}^{\ell+1\dagger})^2 a_{i+\ell-1}^{1\dagger} (a_i^{\ell})^2 a_{i+\ell+1}^1 + \text{h.c.} \right) + \frac{1}{2} \sum_{i=1}^{2L} \sum_{\ell=2}^{L-1} \left((a_i^{\ell\dagger})^2 (a_i^{\ell})^2 n_{i-\ell-1}^1 + (a_{i-1}^{\ell+1\dagger})^2 (a_{i-1}^{\ell+1})^2 n_{i-\ell+1}^1 + (a_i^{\ell\dagger})^2 (a_i^{\ell})^2 n_{i+\ell+1}^1 + (a_{i+1}^{\ell+1\dagger})^2 (a_{i+1}^{\ell+1})^2 n_{i+\ell-1}^1 \right)$$
(S20)
$$- \sum_{i=1}^{2L} \left(((a_{i+1}^{2\dagger})^2 a_i^1 a_{i+2}^1 + \text{h.c}) - \frac{1}{2} (a_i^{2\dagger})^2 (a_i^2)^2 \right) + 2 \sum_{i=1}^{2L} n_i^1 n_{i+2}^1 \right\}.$$

C. Pseudolocality of the Motzkin matched parentheses charge

Like the matched parentheses charge in Fredkin, the Motzkin matched parentheses charge is *pseudolocal* in that it obeys a local conservation law despite possessing power-law tails. Here, we first show that the Motzkin matched parentheses charge has algebraically-decaying tails before proving that the charge in Motzkin is both extensive and additive. Both properties together ensure that this charge is a relevant for transport.

1. Power-law tails

The main text outlines how all static correlations in spin- $\frac{1}{2}$ systems may be traced back to the fact that each endpoint of an irreducible word uniquely belongs to a single irreducible word. The exact same argument holds for spin-1 systems. In a spin-1 system, the existence of a single q_0^1 charge precludes the simultaneous existence of any $q_{\pm \ell}^{\ell}$ charge; the only differences between the integer and half-integer spin models is that here, q_0^1 is a flat and $q_{\pm \ell}^{\ell}$ may live on either a link or a site.

The Motzkin numbers are known to asymptotically scale as $M_{\ell} \sim 3^{\ell} (3/\ell)^{3/2}$ as $\ell \to \infty$. Since there are $M_{\ell-2}$ possible irreducible Motzkin words of length ℓ and 3^{ℓ} possible ℓ -site states, the probability of finding a matched parentheses charge on any given site (or link) is given asymptotically by $M_{\ell-2}/3^{\ell} \sim \ell^{-3/2}$. Therefore, the decay of the tails of the Motzkin matched parentheses charge, like the Fredkin case, is captured by $\langle q_m^1 q_n \rangle \sim |m-n|^{-3/2}$.

2. Extensivity and additivity

Given the myriad of similarities between Fredkin and Motzkin, it is unsurprising that the Motzkin matched parentheses charge is also extensive and additive. To see this, consider the same setup outlined in Section I but for a spin-1 system. By only performing allowed Motzkin moves within subsystem A, the subsystem may be reordered into a state where all of the n_- flats come first, followed by all of the $n_1 + n_1$ parentheses. Since this state may be reached by only performing allowed Motzkin moves, the total charge of the subsystem $Q_A = \sum_{n=1}^{2L} q_n = (\sum_{n=1}^{n_-} q_n) + (\sum_{n=n_-+1}^{2L} q_n)$ remains unchanged after this reordering. The number of charges contributed by the flats trivially scales with n_- , and the number of charges attached to parentheses scales with $n_1 + n_1$ by the arguments presented in section I. Therefore, $Q_A \sim n_- + n_1 + n_1 = L$, so the charge is naturally extensive.

To show that the charge is additive, we derive the probability distribution of the charge in a subsystem. By counting the population of flats and matched parentheses separately, it can be shown that the number of states in subsystem A that contain Q_A charges is given by:

$$T_{\rm M}(L,Q_A) = \sum_{i={\rm mod}(Q_A,2)}^{Q_A} {\binom{L}{i}} T_{\rm inf}\Big(L-i,\frac{Q_A-i}{2}\Big),$$
(S21)

where $T_{inf}(L,Q)$ is given by eq. S2. Therefore, the probability finding Q_A charges in subsystem A is given by:

$$P_L(Q_A) = \frac{T_{\rm M}(L, Q_A)}{3^L}.$$
(S22)

Computing cumulants from this distribution, we find that both the mean and variance scale linearly with L, as in the Fredkin case.

D. Bound for the Motzkin gap scaling

We constructed a spin-wave ansatz in order to bound the scaling of the gap of the Fredkin Hamiltonian, $\Delta \sim L^{-z}$, with $z \geq 5/2$. Here, we employ a nearly identical approach in order to obtain the same bound for the Motzkin Hamiltonian.

1. The Motzkin spin-wave ansatz

We construct a spin-wave ansatz in terms of the Motzkin matched parentheses charge as $|k\rangle = \sum_{n} e^{ik \cdot n/2} q_n |GS\rangle$; the factor of 1/2 in the exponent is due to the fact that the matched parentheses charges may sit on both sites and links, and with our conventions *n* runs from 1 to 2*L*. As before, we work in the S_{tot}^z ensemble, where the ground state $|GS\rangle$ is the uniform superposition of all S_{tot}^z states. The energy expectation value is therefore given by:

$$E_{\rm SW}(k) = \frac{\langle k | H | k \rangle}{\langle k | k \rangle} = \frac{\sum_{n,m} e^{ik \cdot (n-m)/2} \langle GS | q_m H q_n | GS \rangle}{\sum_{n,m} e^{ik \cdot (n-m)/2} \langle GS | q_m q_n | GS \rangle}.$$
(S23)

We will admit that as in the Fredkin case, the norm just scales linearly with system size $\langle k|k \rangle \sim L$, so we will focus on the numerator in the following. Using translation invariance, we thus have

$$E_{\rm SW}(k) \sim \underbrace{\sum_{n} e^{ik \cdot n/2} \langle GS | q_0 H q_n | GS \rangle}_{E_{\rm SW}^0(k)} + \underbrace{\sum_{n} e^{ik \cdot (n-1)/2} \langle GS | q_1 H q_n | GS \rangle}_{E_{\rm SW}^1(k)}.$$
(S24)

In our conventions, n even corresponds to links, while n odd corresponds to sites.

2. Charge-current correlations

For Motzkin, the lattice continuity equation for the new conserved charge reads $i[H, q_n] = j_n - j_{n+1} + j'_n - j'_{n+2}$ with $j_n = \sum_{n=1}^{L-1} j_n^\ell$ and $j'_n = \sum_{\ell=2}^{L-1} j_n^{'\ell}$. The current operator j_n^ℓ associated with the movement of charges that are associated with matched sets of parentheses is given by:

$$j_{n}^{\ell} = \begin{cases} -\frac{i}{4} \left(a_{n-1}^{1} a_{n+1}^{1} (a_{n}^{2\dagger})^{2} - a_{n-2}^{1} a_{n}^{1} (a_{n-1}^{2\dagger})^{2} - \text{h.c.} \right), \text{ if } \ell = 1 \\ \frac{i}{4} \left((a_{n-1}^{\ell+1\dagger})^{2} a_{n-\ell+1}^{1\dagger} (a_{n}^{\ell})^{2} a_{n-\ell-1}^{1} - (a_{n}^{\ell+1\dagger})^{2} a_{n+\ell-2}^{1\dagger} (a_{n-1}^{\ell})^{2} a_{n+\ell}^{1} - \text{h.c.} \right), \text{ otherwise,} \end{cases}$$
(S25)

while the current operator j_n^{ℓ} corresponding the movement of flats is given by:

$$j_{n}^{\prime\ell} = -\frac{i}{8} \bigg((a_{n+\ell-2}^{\ell+1\dagger})^2 a_n^{1\dagger} (a_{n+\ell-1}^{\ell})^2 a_{n-2}^{1} - (a_{n-\ell}^{\ell+1\dagger})^2 a_{n-2}^{1\dagger} (a_{n-\ell-1}^{\ell})^2 a_n^{1} - \text{h.c.} \bigg).$$
(S26)

The origin of charge-current correlations may be understood in direct analogy to the source of charge-current correlations in Fredkin, as outlined in II B. In Motzkin, the movement of all q^{ℓ} charges, for $\ell \geq 1$, is facilitated by a single q^1 charge passing through one of the endpoints of the q^{ℓ} charge. As the q^1 charge disrupts one of the endpoints of the q^{ℓ} charge, the larger charge changes both length and position to become a $q_{i\pm 1}^{\ell\pm 1}$ charge.

As already noted, the continuity equation contains contributions from two different types of currents of the matched parentheses charges (j and j') due to differences between the movement of flats $(q^1 \text{ charges})$ and charges associated with $\ell \geq 2$ Motzkin words. During the process of passing through the endpoint of a larger word, the q^1 charge moves *two*, since the q^1 charge is may only live on sites. In contrast, the other charge involved in this process will move by only *one* space as it increases or decreases in length by one. Therefore, since $\ell = 1$ charges move different amounts than $\ell \geq 2$ charges, the continuity equation associated with the movement of q_n cares whether the matched parentheses current at n is caused by a $\ell = 1$ or $\ell \geq 1$ charges.

We may follow the procedure used in II A and write $\langle GS|q_mHq_n|GS \rangle = -i \langle GS|q_m[H,q_n]|GS \rangle$. The energy of the spin-wave state may then be expressed as a sum of terms of the form $\langle q_m j'_n \rangle$ and $\langle q_m j_n \rangle$. As was the case for Fredkin, these charge-current correlators may be expressed in terms of the probability of finding specific configurations– we derive these probabilities in the following section.

3. Probabilities

In subsection IIB, we outline how all charge-current correlations in Fredkin may be linked to the probability of finding configurations of the form ()(...) or (()...), since the $\ell = 1$ Dyck word facilitates the motion of larger words by passing through their endpoints. In Motzkin, the same type of reasoning holds, except the $\ell = 1$ Motzkin word is a single flat, -, instead of the $\ell = 1$ Dyck word, (). Therefore, charge-current correlations may be expressed in terms of the probability that configurations of the form -(...) or (-...) occur.

As was the case in Fredkin, the probability of finding a configuration like (-...) is independent of system size. There are $M(\ell)$ Motzkin paths of length ℓ , $M(\ell-2)$ of which are irreducible. For Fredkin, eq. S11 gives the probability that there is an $\ell = 1$ charge sitting directly inside of *either* of the endpoints of the $\ell > 1$ charge; for reasons that will become clear in the following section, we are interested in looking at only *one* of the endpoints for Motzkin. Therefore, conditional on having a word of length ℓ on a particular link, we have:

$$p_{\rm in}(\ell) = \frac{M(\ell-3)}{M(\ell-2)}.$$
 (S27)

This expression approaches 5/9 in the $\ell \to \infty$ limit.

Directly counting all possible configurations gives $p_{\text{out}} = 5/9$ but does not account for finite-size effects. Since there are $\frac{L!}{(n!)^2(L-2n)!}$ possible $S_{\text{tot}}^z = 0$ states with L - 2n spins-0, there are:

$$\Omega(L) = \sum_{n=0}^{L/2} \frac{L!}{(n!)^2 (L-2n)!},$$
(S28)

total possible states. Therefore, the probability of finding a configuration of the form -(...) is given by:

$$p_{\text{out}}(\ell, L) = \frac{\Omega(L - \ell - 1)}{\Omega(L - \ell)}.$$
(S29)

Both $p_{in}(\ell)$ and $p_{out}(\ell, L)$ given above are conditioned on having a Motzkin word on a particular link; obtaining the full probability requires removing that condition. The probability that any given link hosts a Motzkin word of length

 ℓ is given by:

$$p_q(\ell, L) = \frac{M(\ell - 2)\Omega(L - \ell)}{\Omega(L)}.$$
(S30)

Having worked out all relevant probabilities, we can now use them to exactly find $E_{SW}(k)$ and thus bound the scaling of the gap of the Motzkin Hamiltonian.

4. Motzkin spin-wave energy $E_{SW}(k)$

The energy of the spin-wave state splits into two parts, $E_{SW} = E_{SW}^0(k) + E_{SW}^1(k)$, as shown in S24. Each $E_{SW}^m(k)$ has contributions from both j_n currents and j'_n currents:

$$E_{\rm SW}^{m=0,1}(k) \sim \sum_{n} i e^{ik \cdot n/2} \Big(\big(\langle q_m j_{n+1} \rangle_0 - \langle q_m j_n \rangle_0 \big) + \big(\langle q_m j'_{n+2} \rangle_0 - \langle q_m j'_n \rangle_0 \big) \Big), \tag{S31}$$

where again $\langle \bullet \rangle_0$ corresponds to $\langle GS | \dots | GS \rangle$. The form of the $\langle q_m j'_n \rangle$ is the same for either m = 0, 1 and $|n| \geq 2$:

$$i\langle q_m j'_n \rangle_0 = \begin{cases} p_{\text{out}}(n-1,L)p_q(n-1,L) - p_{\text{in}}(n+1)p_q(n+1,L), \text{ for odd } n; \\ 0 \text{ otherwise.} \end{cases}$$
(S32)

The interpretation is straightforward: since j'_n is the current on link *n* associated with a flat moving across that link, there can be no j'_n current associated with even values of *n*. The expression for odd *n* is similar to the charge-current correlations in Fredkin (eq. S13).

The form of the $\langle q_m j_n \rangle_0$ correlator depends on whether m is even or odd due to the fact that j_n is the current associated with non- $\ell = 1$ irreducible words at n. Therefore, all charge-current correlations are driven by the presence of an $\ell = 1$ charge sitting at site m. For link m = 0, the j_n correlations vanish for $|n| \ge 2$ since flats can not live on links; for |n| = 1 and m = 0, all correlations are driven by the presence of a single flat on site n, and so $i\langle q_1 j_1 \rangle_0 = p_q(1, L)$. In contrast, flats are allowed to sit on site m = 1, and so there are nontrivial charge correlations for all values of n when m = 1. Therefore, we have:

$$i\langle q_1 j_n \rangle_0 = p_{\text{out}}(n-1,L)p_q(n-1,L) - p_{\text{in}}(n+2)p_q(n+2,L),$$
(S33)

for $n \ge 3$. For small values of n, we have $i\langle q_1 j_2 \rangle_0 = (p_q(2,L) - p_q(4,L))/2$ and $i\langle q_1 j_1 \rangle_0 = p_q(1,L) - p_{in}(4)p_q(4,L) + p_{out}(4,L)p_q - i\langle q_1 j_2 \rangle_0$.

Note that both $E_{\rm SW}^0$ and $E_{\rm SW}^1$ contain a diffusive $\sim k^2$ term at low momenta, but those contributions come in with opposite signs and cancel out in $E_{\rm SW}$. We will admit that this cancellation is due to a sum rule $\sum_{m,n} (m - n)^2 \langle GS | q_m H q_n | GS \rangle = 0$, similar to the one we showed in the Fredkin case. This rules out a potentially diffusive $\sim k^2$ contribution in $E_{\rm SW}(k)$. Instead, the low-energy dispersion is governed by the long distance of the correlator (S32), in close analogy with the Fredkin case (S13). The probabilities $p_{\rm in}$ and $p_{\rm out}$ are $\mathcal{O}(1)$ at long distances, whereas $p_q(\ell) \sim \ell^{-3/2}$. The charge current correlator thus scales as $i \langle q_0 j'_n \rangle_0 \sim n^{-5/2}$, and $E_{\rm SW}(k) \sim k^{5/2}$, mirroring the Fredkin spin-wave dispersion relation. Setting $k = 2\pi/L$, this provides an upper bound $\mathcal{O}(L^{-5/2})$ on the gap of the Motzkin spin chain.

^[1] X. Chen, E. Fradkin, and W. Witczak-Krempa, Physical Review B 96 (2017), 10.1103/physrevb.96.180402.

^[2] X. Chen, E. Fradkin, and W. Witczak-Krempa, Journal of Physics A: Mathematical and Theoretical 50, 464002 (2017).

^[3] S. Bravyi, L. Caha, R. Movassagh, D. Nagaj, and P. W. Shor, Phys. Rev. Lett. 109, 207202 (2012).