# Seiberg-Witten curves of $\widehat{D}$ -type Little Strings

BAPTISTE FILOCHE\* , STEFAN HOHENEGGER $^{\dagger}$  , and TARO KIMURA  $^{\ddagger}$ 

July 17, 2024

<sup>\*†</sup> Univ Lyon, Univ Claude Bernard Lyon 1, CNRS/IN2P3, IP2I Lyon, UMR 5822, F-69622, Villeurbanne, France

Dept. of Physics E. Pancini, Università di Napoli Federico II, via Cintia, 80126 Napoli, Italy

<sup>‡</sup> Institut de Mathématiques de Bourgogne, Université de Bourgogne, CNRS, UMR 5584, France

#### Abstract

Little Strings are a type of non-gravitational quantum theories that contain extended degrees of freedom, but behave like ordinary Quantum Field Theories at low energies. A particular class of such theories in six dimensions is engineered as the world-volume theory of an M5-brane on a circle that probes a transverse orbifold geometry. Its low energy limit is a supersymmetric gauge theory that is described by a quiver in the shape of the Dynkin diagram of the affine extension of an ADE-group. While the so-called  $\hat{A}$ -type Little String Theories (LSTs) are very well studied, much less is known about the  $\hat{D}$ -type, where for example the Seiberg-Witten curve (SWC) is only known in the case of the  $\hat{D}_4$  theory. In this work, we provide a general construction of this curve for arbitrary  $\hat{D}_M$  that respects all symmetries and dualities of the LST and is compatible with lower-dimensional results in the literature. For M = 4 our construction reproduces the same curve as previously obtained by other methods. The form in which we cast the SWC for generic  $\hat{D}_M$  allows to study the behaviour of the LST under modular transformations and provides insights into a dual formulation as a circular quiver gauge theory with nodes of Sp(M-4) and SO(2M).

t

<sup>\*</sup>b.filoche@ip2i.in2p3.fr

<sup>&</sup>lt;sup>†</sup>s.hohenegger@ipnl.in2p3.fr

<sup>&</sup>lt;sup>‡</sup>taro.kimura@u-bourgogne.fr

## Contents

T	Introduction	2
2	Little String Theories and Seiberg-Witten curves2.1 $\widehat{ADE}$ -type Little String Theories2.2 $\widehat{A}$ -type Seiberg-Witten curves	<b>4</b> 4 5
3	$\widehat{D}$ -type Seiberg-Witten curves3.1 $\widehat{D}$ -type LSTs3.2General Ansatz for the Seiberg Witten Curve	<b>7</b> 7 8
4	Seiberg-Witten Curve of $(\widehat{D}_4, 1)$ 4.1 Construction4.2 Modular properties4.3 Lower Dimensional Limit	<b>10</b> 10 12 13
5	Seiberg-Witten Curve of $(\widehat{D}_5, 1)$ 5.1 Construction5.2 Modular properties5.3 Lower dimensional limit	<b>15</b> 15 16 17
6	Generalisation to the $(\widehat{D}_M, 1)$ LSTs	18
7	Conclusion	20
Α	Modular functions and propertiesA.1 Definitions and propertiesA.2 Modular (sub)group	<b>22</b> 22 23
в	$\mathbb{M}$ matrices examples	<b>24</b>

റ

# 1 Introduction

Little String Theories (LSTs) [1–5] are a class of non-gravitational quantum theories, which at low energies behave like quantum field theories with point-like degrees of freedom, but whose UV-completion requires new extended degrees of freedom. Such theories arise naturally within the framework of String Theory, through suitable limits in which the gravitational sector is decoupled [6, 7]. They constitute interesting generalisations of (conformal) field theories that are obtained through different point-particle limits and have therefore attracted a lot of attention over the years: for example, through geometric methods similar to the case of (higher dimensional) conformal field theories, a classification has been provided in [8–10].

A very rich class of supersymmetric LSTs (dubbed Little String Orbifolds in [11]) can be constructed as the world-volume theory of N parallel M5-branes on a circle, probing a transverse orbifold geometry [12] (see also [13–32]) of the form  $\mathbb{R}^4_{\perp}/\Gamma$ , where  $\Gamma \subset SU(2)$ . At low energies, the world-volume theory on the M5-branes is a six-dimensional supersymmetric gauge theory, described by a quiver in the shape of the Dynkin diagram of the affine group  $\widehat{\Upsilon}$ , which is related to  $\Gamma$  through the McKay correspondence. Without additional flavour groups, the rank of the gauge nodes is fixed through the absence of anomalies and matter appears in the bifundamental representation.

Among these theories the case with  $\widehat{\Upsilon} = \widehat{A}_{M-1}$  (called  $\widehat{A}$ -type and denoted  $(\widehat{A}_{M-1}, N)$ theory in the following) has been studied in quite some detail: indeed, this class of theories enjoys many different dual descriptions, which allow to calculate important quantities explicitly. For example, the BPS-(instanton) partition function can be obtained by counting M2-branes stretched between the M5 branes [13–16]. Using a dual description in terms of F-theory compactified on a toric Calabi-Yau manifold  $X_{N,M}$  [33], this counting is captured by the topological string partition function on this threefold [11, 17, 18, 20], which in turn can be computed in an algorithmic fashion using the (refined) topological vertex [34, 35]. Another quantity of the  $(A_{M-1}, N)$  theory, which can be constructed explicitly using various different approaches [36, 37], is the Seiberg-Witten curve (SWC). The latter encodes important information about the non-perturbative structure of the theory, notably its symmetries. Furthermore, it affords interesting insights into related algebraic integrable systems |38-40|, where it is identified with their spectral curve. The SWC depends on MN + 2 independent parameters and can be written as a linear combination of a basis of theta functions that are defined on a genus 2 surface. This surface reflects the doubly periodic nature of the M5-brane setup (which is for example also visible in the double elliptic fibration structure of the threefold  $X_{N,M}$  mentioned above). Furthermore, for a generic  $(A_{M-1}, N)$  theory, this surface has a full period matrix: the third parameter, besides the two modular parameters, has an interpretation as the mass of the bifundamental matter from the perspective of the quiver gauge theory.

In this work, however, we shall be concerned with the case  $\widehat{\Upsilon} = \widehat{D}_M$  (for  $M \ge 4$ ) and N = 1 (*i.e.* a single M5-brane), which we shall denote as  $(\widehat{D}_M, 1)$  theory in the following. Due to technical difficulties and subtleties in realising dual descriptions, these so-called  $\widehat{D}$ -type LSTs are much less studied in the literature. Explicit computations of the instanton partition function have been performed in the topological vertex formalism by using so-called trivalent gluing techniques [41–43], using the blow-up equations [44, 45] or using dual realisations in term of two-dimensional  $\mathcal{N} = (0, 4)$  gauge theories [46]. From the perspective of the SWC, to date only the curve in the case M = 4 is known [47], either from the thermodynamic limit [48] of the instanton partition function or by using the quantum states formalism [49].

For general M, the geometric structure of the SWC is somewhat less restrictive than in the  $(\widehat{A}_{M-1}, 1)$  case: due to the absence of a mass parameter [50], a suitable set of basis functions consists of products of genus 1 theta functions (with two different modular parameters), rather than genuine genus 2 theta functions as in the  $\widehat{A}$ -type case. In this paper, we present a general ansatz for the SWC of the  $(\widehat{D}_M, 1)$  theory that is manifestly compatible with all symmetries. To further restrict this ansatz, we use the fact that the  $(\widehat{D}_M, 1)$  theory is dual to a circular quiver gauge theory with one gauge node of type Sp(M-4) and one node of type SO(2M). This requires that the SWC can equivalently be re-written in a way that is compatible with the symmetries of this dual theory. We argue for low values of M (notably M = 4 and M = 5) that consistency of these dual presentations imposes further conditions on the parameters and ultimately leaves only a few distinct forms of the curve (two in the case of M = 4 and three in the case of M = 5). By further considering a dimensional reduction (by assuming a natural behaviour of the remaining parameters in the decompactification limit) and comparing with known results in five-dimensions [51], we find further conditions on the parameters leaving a

unique form of the SWC. In particular in the case of M = 4, the latter agrees with known results in [47, 49]. By studying further examples up to M = 12, we find distinctive patterns, which allow us to conjecture the general form of the  $(\hat{D}_M, 1)$  SWC for generic  $M \ge 4$ . The latter could be identified with the spectral curves of new  $\hat{D}$ -type double elliptic integrable systems. Some steps have been taken in this direction through the construction of the quantum integrable system of D-type conformal matter (*i.e.* six-dimensional Super Conformal Field Theories (SCFTs)) [52] and quantum deformations of SWCs [53–55].

This paper is organised as follows: In Section 2, we provide more details on LSTs engineered from M5-branes probing transverse orbifold geometries and their SWCs, recalling in particular the  $(\widehat{A}_{M-1}, N)$  theories. In Section 3 we focus on the  $(\widehat{D}_M, 1)$  LSTs and present a general ansatz for the SWC that is compatible with all symmetries of the theory. In Section 4 we discuss in detail the case M = 4 and show how this ansatz can be further restricted by imposing duality of the  $(\widehat{D}_4, 1)$  theory to a quiver gauge theory with one Sp(0) and one SO(8) node and by comparing to known five-dimensional results. We show that the unique SWC obtained in this way is equivalent to the result in [47, 49]. In Section 5 we repeat this construction for the  $(\widehat{D}_5, 1)$  theory and show that it again leads to a unique SWC, which constitutes a novel result in the literature. Based on further examples up to M = 12, we present in Section 6 a general form for the SWC of the  $(\widehat{D}_M, 1)$  theory. Finally, Section 7 contains our conclusions, as well as an outlook on applications of our results. This paper is accompanied by two Appendices: in Appendix A we compile definitions as well as useful identities for modular objects that are used throughout the main body of this text. Appendix B contains the results of the SWC for certain M > 5, which corroborate our general conjecture in Section 6.

## 2 Little String Theories and Seiberg-Witten curves

### 2.1 *ADE*-type Little String Theories

Little String Theories (LSTs) arise in string theory (or related dual descriptions) through different decoupling limits [1–5, 56] (see also [6, 7] for reviews). They follow an  $\widehat{ADE}$ -classification [8] and can be engineered from numerous setups. For concreteness, we begin with a construction in M-theory [11, 17], consisting of N M5-branes (along directions  $x^{0,1,\dots,5}$ ), separated along a circle  $\mathbb{S}_6^1$  probing a transverse  $\mathbb{R}_{\perp}^4/\Gamma$  orbifold with a finite subgroup of SU(2),  $\Gamma \subset SU(2)$ :

In the low energy limit, this construction corresponds to a supersymmetric quiver gauge theory on  $\mathbb{S}_0^1 \times \mathbb{S}_1^1 \times \mathbb{R}_{\parallel}^3 \times \mathbb{S}_5^1$ . The McKay correspondence [57] associates the orbifold group  $\Gamma$  to an affine quiver group  $\widehat{\Upsilon}$  as follows

where  $\mathbb{BD}_M$  is the binary dihedral group of order 4M and  $\mathbb{BT}, \mathbb{BO}$  and  $\mathbb{BI}$  are respectively the binary isomorphism groups of the tetrahedron, the cube and the dodecahedron. The quiver

takes the form of the Dynkin diagram of  $\widehat{\Upsilon}$  (see Fig. 1) and mathematically is described by  $(\Upsilon_0, \Upsilon_1)$ , where  $\Upsilon_0$  is the set of nodes and  $\Upsilon_1$  the set of edges connecting the latter. Fig. 1 also display an *n*-coloring of the quivers, *i.e.* a particular assignment of the rank of each gauge node. To explain this, we introduce the *Cartan matrix* of the quiver  $(\Upsilon_0, \Upsilon_1)$  by

$$c_{ij} = 2\delta_{ij} - \sum_{e \in \Upsilon_1: i \to j} 1 - \sum_{e \in \Upsilon_1: j \to i} 1, \qquad \forall i, j \in \Upsilon_0.$$

$$(2.1)$$

The rank of the gauge group at each node in  $\Upsilon_0$  is uniquely fixed<sup>1</sup> by demanding the vanishing of the beta-functions for the gauge couplings [58]:

$$\sum_{j \in \Upsilon_0} c_{ij} \cdot n_j = 0, \qquad \forall i \in \Upsilon_0,$$
(2.2)

with  $n_j$  the rank of gauge group associated with the *j*-th node. A solution of the set of equations (2.2) is called an *n*-coloring of the quiver. For quivers of simply laced Lie groups, the coloring is fixed by a single integer N which corresponds to the number of M5-branes probing the orbifold. Concretely, in Fig. 1, the nodes (N) of the quiver represent gauge groups  $\widehat{A}_{N-1}$ , while lines denote matter in the bifundamental representation.<sup>2</sup> We shall denote a LST in this framework as  $(\widehat{\Upsilon}, N)$ .



Figure 1: N-coloring of  $\widehat{A}$ ,  $\widehat{D}$  and  $\widehat{E}$ -type quivers.

### **2.2** $\widehat{A}$ -type Seiberg-Witten curves

Reviewing first the case of  $\Gamma = \mathbb{Z}_M$ , the so-called A-type LSTs engineered from the M-brane setup in Table 2.1 can be described by a doubly elliptic Seiberg-Witten curve [36, 37]. To understand the origin of the latter, we recall [11, 13–17] that this M-brane configuration can be described by MN + 2 independent parameters: we refer the reader for more details to [11, 17] and only mention that in the following we denote the (complexified) radii of  $\mathbb{S}_1^1$  and  $\mathbb{S}_6^1$ (measured in units of the radius of  $\mathbb{S}_0^1$ ) by  $\tau$  and  $\rho$  respectively. Furthermore, following [17], we introduce a  $U(1)_S$  deformation of  $\mathbb{R}_+^4/\mathbb{Z}_M$  with respect to  $\mathbb{S}_0^1$ 

$$U(1)_S: (w_+, w_-) \longrightarrow (e^{2\pi i S} w_+, e^{-2\pi i S} w_-), \quad \text{with} \quad w_\pm = x^7 \pm i x^8, \quad (2.3)$$

<sup>&</sup>lt;sup>1</sup>Here we do not assume any additional flavor groups.

<sup>&</sup>lt;sup>2</sup>Certain quiver gauge theories, such as the  $\hat{A}_0$  theory with M = 1 has matter in the adjoint- rather than the bifundamental representation [17]. However, we shall not consider this case in the current paper.

which is compatible with the  $\mathbb{Z}_M$  action. The parameters  $(\tau, \rho, S)$  can be arranged into the period matrix  $\Omega$  of a genus 2 Riemann surface and the SWC (parametrised by  $(z_1, z_2)$ ) can be written in terms of genus 2 theta functions  $\Theta$  (see (A.1) for the definition) [36, 37]:

$$\sum_{i=0}^{M-1} \sum_{j=0}^{N-1} a_{ij} \Theta \begin{bmatrix} i/M & j/N \\ 0 & 0 \end{bmatrix} (z \mid \Omega) = 0, \quad z = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}, \quad \Omega = \begin{bmatrix} \frac{\tau}{M} & S \\ S & \frac{\rho}{N} \end{bmatrix}.$$
(2.4)

The  $\{a_{ij}\}$  in (2.4) depend on the remaining parameters of the M-brane configuration mentioned above.<sup>3</sup> Using a dual description in terms of F-theory compactified on a Calabi-Yau threefold  $X_{N,M}$  [11, 17, 18] with double elliptic-fibration structure, the  $\{a_{ij}\}$  have been characterised in [33] as (the product of) generating functions of Gromov-Witten invariants.

From the perspective of the  $\hat{A}_M$  quiver gauge theory represented by the leftmost diagram in Fig. 1, the parameter  $\tau$  is interpreted as a complexified gauge coupling (which is common to all M nodes in the quiver),  $\rho$  as the (common) parameter which extends all gauge group nodes to their affine forms  $\hat{A}_{N-1}$  and S as the mass of the bifundamental hypermultiplets. The remaining parameters [33] (appearing in the form of  $a_{ij}$  in (2.4)) encode further structure of the gauge- and matter spectrum, but the details shall not be important in the following. At the classical level, the theory inherits modular properties from the genus 2 formulation (2.4). The functional basis (*i.e.* the particular set  $\Theta \begin{bmatrix} i/M & j/N \\ 0 & 0 \end{bmatrix} (z | \Omega)$  of genus 2 theta functions) over which the curve (2.4) is written, corresponds to theta functions of polarised Abelian varieties [59, 60]. The modular group acting on the period matrix  $\Omega$  is therefore the paramodular group defined as [60]:

$$\Sigma_{M,N} := Sp(4, \mathbb{Q}) \cap \begin{bmatrix} \mathbb{Z} & \mathbb{Z} & \mathbb{Z}/M & \mathbb{Z}/N \\ \mathbb{Z} & \mathbb{Z} & \mathbb{Z}/M & \mathbb{Z}/N \\ M\mathbb{Z} & M\mathbb{Z} & \mathbb{Z} & M\mathbb{Z}/N \\ N\mathbb{Z} & N\mathbb{Z} & N\mathbb{Z}/M & \mathbb{Z} \end{bmatrix}, \qquad (2.5)$$

with,

$$Sp(4,\mathbb{Q}) = \left\{ \Lambda = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{M}_4(\mathbb{Q}) \mid \Lambda \begin{bmatrix} 0 & \mathbb{1} \\ -\mathbb{1} & 0 \end{bmatrix} \Lambda^T = \begin{bmatrix} 0 & \mathbb{1} \\ -\mathbb{1} & 0 \end{bmatrix} \right\},$$
(2.6)

where  $\mathbb{M}_4(\mathbb{Q})$  is the space of  $4 \times 4$  matrices with rational entries and non-vanishing determinant. The group  $\Sigma_{M,N}$  acts in the following way on the period matrix  $\Omega$ :

$$\Omega \to (A\Omega + B)(C\Omega + D)^{-1}, \quad \forall \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \Sigma_{M,N}.$$
 (2.7)

It is interesting to note that the linearly acting subgroup of the paramodular group (*i.e.* the subgroup with B = C = 0 in (2.7)) generates the web of dualities in LSTs that was described at the level of the instanton partition function in [18, 20, 21]. In this respect, the group  $\Sigma_{M,N}$  is a generalisation of the paramodular group found in [26, 27] for generic M, N > 1.

The  $(\widehat{A}_{M-1}, N)$  theory was shown to be dual to  $(\widehat{A}_{M'-1}, N')$  for any  $N', M' \in \mathbb{N}$  such that N'M' = NM and gcd(N', M') = gcd(N, M) =: p. The corresponding duality map at the level of the period matrix  $\Omega$ , was formulated in [18] in terms of the following  $Sp(4, \mathbb{Z})$  transformation

$$A = \begin{bmatrix} \frac{M}{p} & -\frac{N}{p} \\ 1 & -1 \end{bmatrix}, \quad \Omega \to (A^T)^{-1} \Omega A^{-1} = \begin{bmatrix} \frac{\tau'}{M'} & S' \\ S' & \frac{\rho'}{N'} \end{bmatrix}.$$
 (2.8)

<sup>&</sup>lt;sup>3</sup>Indeed, due to an overall scaling symmetry of (2.4), the moduli space of the SWC is indeed (NM + 2)-dimensional.

This transformation acts naturally on the basis of genus 2 theta functions  $\Theta\begin{bmatrix} i/M & j/N \\ 0 & 0 \end{bmatrix}(z \mid \Omega)$  (for  $i \in \{0 \dots M-1\}$  and  $j \in \{0 \dots N-1\}$  over which (2.4) is formulated: using (A.2) and noticing the equivalence of the following sets

$$\left\{A \cdot \left(\frac{i}{M}, \frac{j}{N}\right), 0 \le i \le M - 1, 0 \le j \le N - 1\right\}$$
$$\cong \left\{\left(\frac{k}{p} \mod 1, \frac{lp}{MN} \mod 1\right), 0 \le k \le p - 1, 0 \le l \le \frac{MN}{p} - 1\right\}, \quad (2.9)$$

it indeed follows that (2.8) (along with a suitable duality map of the remaining parameters  $\{a_{ij}\}$ ) leaves the SWC (2.4) invariant.

# 3 $\widehat{D}$ -type Seiberg-Witten curves

In this Section, we review some of the key properties of  $(\widehat{D}_M, 1)$  LSTs, which allow us to formulate a general ansatz for their SWCs. In the following Sections 4 and 5 we shall use this ansatz to provide the unique form of the curves in the cases M = 4 and M = 5 respectively, before generalising our results in Section 6 to generic  $M \ge 4$ .

## 3.1 $\widehat{D}$ -type LSTs

First of all, contrary to  $\hat{A}$ -type,  $\hat{D}$ -type LSTs do not allow for a mass-deformation parameter [50], *i.e.* there is no equivalent to the parameter S in the period matrix in (2.4). Therefore, also the genus 2 theta functions  $\Theta$  are effectively reduced to two types of genus 1 (*i.e.* Jacobi) theta functions. Secondly, the transverse  $\hat{D}$ -type orbifold itself has two important effects, which are of importance to the SWC:

- (i) the orbifold gives rise to a  $\mathbb{Z}_2$  symmetry, under which the SWC is invariant [47]
- (ii) M5-brane fractionalisation: as explained in [61] due to the orbifold, a single M5-brane splits into two 'half-branes' on the  $\mathbb{S}_6^1$  circle of Table 2.1. At the level of the quantum states introduced in [49], each M5 state can be described by the product of two 1/2 M5-brane states.

Finally, there is another point of view, which provides crucial information for the SWCs, namely a dual description of the  $(\hat{D}_M, 1)$  LSTs in terms of an Sp(M-4)-SO(2M) quiver gauge theory, which is schematically shown in Figure 2. Indeed, for M = 4 this duality was first argued in [46] at the level of the partition function and in [47] at the level of the SWC using fiber-base duality, while for general  $M \ge 4$ , it was established at the level of the partition function in [46].

To understand the implications of such a duality at the level of the SWC, we remark that, in the absence of a mass-deformation, curves with  $\hat{A}_1$  base can generally be written in the following form [47, 49]:

$$0 = \theta_3^2(0;\rho)\theta_2^2(0;\rho)\theta_1^2(z_2;\rho)c_1(\tau)F_1(z_1;\tau) - \theta_4^2(z_2;\rho)F_2(z_1;\tau),$$
(3.1)



Figure 2: Quiver diagram of  $(\widehat{D}_M, 1)$  LST with M - 3 SU(2) nodes (which are indicated explicitly) and its dual  $\widehat{A}_1$  base Sp(M - 4) - SO(2M) quiver theory

where the Jacobi theta functions  $\theta_{2,3}$  are defined in (A.3),  $c_1(\tau)$  is a modular constant and  $F_{1,2}$  correspond to elliptic gauge polynomials<sup>4</sup> of each of the two nodes of the (right) quiver in Figure 2. For example, in [47], these polynomials have been found for SO(8) and Sp(0) nodes to be respectively

$$F_1(z_1;\tau) = \prod_{i=1}^4 \frac{\theta_1(z_1 - a_i^{SO};\tau)\theta_1(z_1 + a_i^{SO};\tau)}{\theta_1^2(z_1;\tau)}, \quad \text{and} \quad F_2(z_1;\tau) = \frac{\theta_1^2(2z_1;\tau)}{\theta_1^8(z_1;\tau)}. \quad (3.2)$$

The existence of the dual description of the  $(\hat{D}_M, 1)$  theory requires that their SWC need to admit a presentation of the form (3.1). As we shall see in the following, by starting from a generic ansatz (based on a finite dimensional basis of theta-function building blocks), this requirement leads to non-trivial constraints.

### 3.2 General Ansatz for the Seiberg Witten Curve

We are now ready to discuss the general form of the SWC of the  $(\widehat{D}_M, 1)$ -LST. This curve is doubly elliptic and we shall denote the corresponding parameters by  $\tau$  and  $\rho$ , while the coordinates shall be called  $z_1$  (the coordinate on the  $\tau$ -parameterised elliptic curve) and  $\mathbb{Z}_2$ even in  $z_2$  (the coordinate on the  $\rho$ -parameterised elliptic curve). Following the discussion of the previous Subsection, we shall now provide a general ansatz for the  $(\widehat{D}_M, 1)$ -LST, which is based on the following principles:

- 1. The SWC is independently  $\mathbb{Z}_2$  even in  $z_1$  and  $z_2$ . This symmetry is due to the  $\mathbb{Z}_2$  symmetry of the  $\widehat{D}$ -type orbifold, explained in point (*i*) of the previous Subsection. The fact that it (a priori) can act independently on  $z_1$  and  $z_2$  is a consequence of the absence of a mass deformation in the  $\widehat{D}$ -type LSTs.
- 2. Using (pairs of) genus 1 theta functions as building blocks, the general form of the D-type-LST SWC can be cast in the following form

$$\sum_{0 \le i,j \le M} \widetilde{T}_i(z_2,\rho,\tau) \,\mathbb{M}_{ij}(\tau) \,\Phi_j(z_1,\tau) = 0\,.$$
(3.3)

<sup>&</sup>lt;sup>4</sup>The gauge polynomial is the polynomial whose roots are the Coulomb branch moduli. Elliptic gauge polynomials are their natural elliptic uplift.

Indeed, here  $\{\Phi_j\}_{j\in\{0...M\}}$  is a basis of  $\mathbb{Z}_2$  even theta functions of degree 2M (defined in (A.7)) and  $\widetilde{T}_i$  are  $\mathbb{Z}_2$  even combinations of elliptic Weyl characters of SU(1) and SU(2)[50, 62], concretely

$$\widetilde{T}_{i}(z_{2},\rho) = \begin{cases} \alpha_{i}\theta_{1}^{2}(z_{2};\rho), & \text{for } i \in \{0,1,M-1,M\}, \\ \alpha_{i}\theta_{1}^{2}(z_{2};\rho) + \beta_{i}\theta_{4}^{2}(z_{2};\rho), & \text{for } i \in \{2\dots M-2\}, \end{cases}$$
(3.4)

with  $\theta_1$  and  $\theta_4$  defined eq. (A.3).  $\alpha_i, \beta_i$  are  $2M - 2 = h^{\vee}(D_M)$ , which are independent of  $z_{1,2}$ , but may depend on  $\tau$  and  $\rho$  (as well as other Coulomb moduli of the  $\widehat{D}_M$ -LST).

3. The form (3.3) of the SWC must permit an alternative decomposition either on sections of line bundles with polarisation (2, 2M) or on products of sections of line bundles with polarisation (1, M). As detailed in [49], this is interpreted in term of quantum states as a consequence of the M5-brane fractionalisation discussed in point *(ii)* of the previous Subsection. This can alternatively be motivated by analysing the 4d SWC of the nodes involved in the dual construction (corresponding to the quiver on the right of Fig. 2). The gauge polynomial of SO(2M) and Sp(M-4) gauge theories can be decomposed into the product of two degree M polynomials [63]. We assume that this structure stems from a decomposition in term of sections of line bundle as explained above in the SWC of the LST.

The most general form of the SWC that is compatible with points 1. and 3. and permits a presentation of the form (3.3) in point 2., with the correct structure for the (elliptic) Weyl characters  $\tilde{T}_i$  takes the following form

$$\theta_{1}^{2}(z_{2};\rho) \left[ \sum_{0 \le i \le j \le \lfloor \frac{M}{2} \rfloor} a_{ij}^{(+)} X_{i}(z_{1};\tau) X_{j}(z_{1};\tau) + \sum_{1 \le i \le j \le \lfloor \frac{M-1}{2} \rfloor} a_{ij}^{(-)} Y_{i}(z_{1};\tau) Y_{j}(z_{1};\tau) \right] + \theta_{4}^{2}(z_{2};\rho) \left[ \sum_{0 \le i \le j \le \lfloor \frac{M}{2} \rfloor} b_{ij}^{(+)} X_{i}(z_{1};\tau) X_{j}(z_{1};\tau) + \sum_{1 \le i \le j \le \lfloor \frac{M-1}{2} \rfloor} b_{ij}^{(-)} Y_{i}(z_{1};\tau) Y_{j}(z_{1};\tau) \right] = 0,$$

$$(3.5)$$

where  $X_i$  and  $Y_i$  (with  $i \in \{0, \ldots, \lfloor \frac{M-1}{2} \rfloor$ ) are defined in (A.6) in Appendix A.1. Here  $a_{ij}^{\pm}$  and  $b_{ij}^{\pm}$  are a total of  $(M+1)^2/2 + 3/2$  parameters for M even and  $(M+1)^2/2$  parameters for M odd, which exceeds the expected number of 2M - 2. This implies that these parameters are not all independent and the ansatz (3.5) is too general. We therefore require further conditions to uniquely determine the SWC: such conditions are provided by demanding an alternative presentation of the curve (3.5) in the form of (3.1), compatible with a dual description as a quiver theory such as in the right part of Figure 2. Finally, any remaining ambiguities can be fixed by comparison with known results of the 5-dimensional compactification of the LSTs. To explain this reasoning concretely, we shall explain in detail in the following Section 4 the case of the  $(\widehat{D}_4, 1)$  theory (for which the SWC was previously constructed in [47, 49]) and in Section 5 the case of the  $(\widehat{D}_5, 1)$  theory (for which the SWC is not known in the literature). Finally, in Section 6, we present the generalisation of our results to theories  $(\widehat{D}_M, 1)$ , with generic  $M \ge 4$ .

# 4 Seiberg-Witten Curve of $(\widehat{D}_4, 1)$

In this Section, we focus on the theory  $(\hat{D}_4, 1)$  and explain how the general ansatz (3.5) can be further restricted to obtain the unique Seiberg-Witten curve. As we shall explain, our results agree with [47, 49], which acts as a verification of the general ansatz presented in the previous Subsection 3.2. We shall furthermore discuss modular properties of our solution and connect the result to known 5d SCFTs.

#### 4.1 Construction

We start from the general ansatz (3.5) for M = 4

$$\theta_{1}^{2}(z_{2};\rho)\left[a_{00}^{(+)}X_{0}^{2}+a_{11}^{(+)}X_{1}^{2}+a_{02}^{(+)}X_{0}X_{2}+a_{11}^{(-)}Y_{1}^{2}+a_{22}^{(+)}X_{2}^{2}+a_{01}^{(+)}X_{0}X_{1}+a_{12}^{(+)}X_{1}X_{2}\right]$$
  
+
$$\theta_{4}^{2}(z_{2};\rho)\left[b_{00}^{(+)}X_{0}^{2}+b_{11}^{(+)}X_{1}^{2}+b_{02}^{(+)}X_{0}X_{2}+b_{11}^{(-)}Y_{1}^{2}+b_{22}^{(+)}X_{2}^{2}+b_{01}^{(+)}X_{0}X_{1}+b_{12}^{(+)}X_{1}X_{2}\right]=0.$$
  
(4.1)

In order to rearrange this expression into the form (3.3), we systematically use Riemann's addition formula (A.8) in Appendix A.1 for genus 1 theta functions [60]. This allows us to systematically replace the basis functions  $X_iX_j$  and  $Y_iY_j$  by the degree 8 theta functions  $\Phi_k$  (defined in (A.7)), as is explicitly shown in eq. (A.9) of Appendix A.1. Indeed, using the shorthand notation:

$$\phi_i := \Phi_i(0, \tau), \quad \Phi_i := \Phi_i(z_1, \tau), \quad \theta_1 := \theta_1(z_2; \rho), \quad \theta_4 := \theta_4(z_2; \rho), \quad (4.2)$$

we find that (4.1) can equivalently be written as

$$\left[ \left(a_{00}^{(+)}\theta_{1}^{2} + b_{00}^{(+)}\theta_{4}^{2}\right)\phi_{0} + \left(\left(a_{11}^{(+)} - a_{11}^{(-)}\right)\theta_{1}^{2} + \left(b_{11}^{(+)} - b_{11}^{(-)}\right)\theta_{4}^{2}\right)\frac{\phi_{2}}{2} + \left(a_{22}^{(+)}\theta_{1}^{2} + b_{22}^{(+)}\theta_{4}^{2}\right)\phi_{4} \right] \Phi_{0} \right]$$

$$+ \left[ \left(a_{01}^{(+)}\theta_{1}^{2} + b_{01}^{(+)}\theta_{4}^{2}\right)\phi_{1} + \left(a_{12}^{(+)}\theta_{1}^{2} + b_{12}^{(+)}\theta_{4}^{2}\right)\phi_{3}\right] \Phi_{1} \right]$$

$$+ \left[ \left(a_{11}^{(+)} + a_{11}^{(-)}\right)\theta_{1}^{2} + \left(b_{11}^{(+)} + b_{11}^{(-)}\right)\theta_{4}^{2}\right)\frac{\phi_{0} + \phi_{4}}{2} + \left(a_{02}^{(+)}\theta_{1}^{2} + b_{02}^{(+)}\theta_{4}^{2}\right)\phi_{2}\right] \Phi_{2}$$

$$+ \left[ \left(a_{01}^{(+)}\theta_{1}^{2} + b_{01}^{(+)}\theta_{4}^{2}\right)\phi_{3} + \left(a_{12}^{(+)}\theta_{1}^{2} + b_{12}^{(+)}\theta_{4}^{2}\right)\phi_{1}\right] \Phi_{3}$$

$$+ \left[ \left(a_{00}^{(+)}\theta_{1}^{2} + b_{00}^{(+)}\theta_{4}^{2}\right)\phi_{4} + \left(\left(a_{11}^{(+)} - a_{11}^{(-)}\right)\theta_{1}^{2} + \left(b_{11}^{(+)} - b_{11}^{(-)}\right)\theta_{4}^{2}\right)\frac{\phi_{2}}{2} + \left(a_{22}^{(+)}\theta_{1}^{2} + b_{22}^{(+)}\theta_{4}^{2}\right)\phi_{0} \right] \Phi_{4} = 0.$$

$$\tag{4.3}$$

In order to further write (4.3) in the form of (3.3), we need to identify the coefficients of  $\Phi_i$  (for  $i \in \{0, \ldots, 4\}$ ) with quantities  $\sum_{0 \le i, j \le M} \widetilde{T}_i(z_2, \rho, \tau) \mathbb{M}_{ij}(\tau)$ , where in particular the  $z_2$  dependent part needs to respect the correct assignment of SU(2) and SU(1) characters: by identifying (some of) the coefficients  $a_{ij}^{\pm}$  and  $b_{ij}^{(\pm)}$  we have to create combinations of the  $\theta_{1,4}^2$ , which can be identified with the characters (3.4). A priori, there are multiple possibilities to do this, however, not all of them lead to a consistent structure (3.3) with one SU(2) and 4 SU(1) characters. However, the number of possibilities can be further reduced by assuming additional

symmetries:<sup>5</sup> we assume that the modular matrix  $\mathbb{M}$  defined by (3.3) remains invariant under the exchanges

$$\widetilde{T}_0 \leftrightarrow \widetilde{T}_4, \ \Phi_0 \leftrightarrow \Phi_4,$$
 and  $\widetilde{T}_1 \leftrightarrow \widetilde{T}_3, \ \Phi_1 \leftrightarrow \Phi_3.$  (4.4)

This is equivalent to stating that none of the Jacobi theta functions multiplied by  $a_{ij}^{\pm}$  or  $b_{ij}^{\pm}$  with  $(ij) \in \{(00), (22), (01), (12)\}$  can be formed into SU(2) characters such that

$$b_{00}^{(+)} = b_{22}^{(+)} = b_{01}^{(+)} = b_{12}^{(+)} = 0, \qquad (4.5)$$

and (4.3) takes the form

$$\left[ a_{00}^{(+)} \theta_1^2 \phi_0 + \left( (a_{11}^{(+)} - a_{11}^{(-)}) \theta_1^2 + (b_{11}^{(+)} - b_{11}^{(-)}) \theta_4^2 \right) \frac{\phi_2}{2} + a_{22}^{(+)} \theta_1^2 \phi_4 \right] \Phi_0 + \left[ a_{01}^{(+)} \phi_1 + a_{12}^{(+)} \phi_3 \right] \theta_1^2 \Phi_1$$

$$+ \left[ \left( (a_{11}^{(+)} + a_{11}^{(-)}) \theta_1^2 + (b_{11}^{(+)} + b_{11}^{(-)}) \theta_4^2 \right) \frac{\phi_0 + \phi_4}{2} + \left( a_{02}^{(+)} \theta_1^2 + b_{02}^{(+)} \theta_4^2 \right) \phi_2 \right] \Phi_2 + \left[ a_{01}^{(+)} \phi_3 + a_{12}^{(+)} \phi_1 \right] \theta_1^2 \Phi_3$$

$$+ \left[ a_{00}^{(+)} \theta_1^2 \phi_4 + \left( (a_{11}^{(+)} - a_{11}^{(-)}) \theta_1^2 + (b_{11}^{(+)} - b_{11}^{(-)}) \theta_4^2 \right) \frac{\phi_2}{2} + a_{22}^{(+)} \theta_1^2 \phi_1 \right] \Phi_4 = 0.$$

$$(4.6)$$

Among the terms highlighted in colour, one (combination) needs to assume the role of the SU(2) character. This can be achieved in two inequivalent and consistent ways

I) Setting  $a_{11}^{(+)} = b_{11}^{(+)} = a_{02}^{(+)} = b_{02}^{(+)} = 0$  such that the red and blue terms in (4.6) are identified and play the role of the SU(2) character, while the green term is eliminated:

$$\begin{bmatrix} a_{00}^{(+)}\theta_1^2 \phi_0 - \left(a_{11}^{(-)}\theta_1^2 + b_{11}^{(-)}\theta_4^2\right) \frac{\phi_2}{2} + a_{22}^{(+)}\theta_1^2 \phi_4 \end{bmatrix} \Phi_0 + \begin{bmatrix} a_{01}^{(+)} \phi_1 + a_{12}^{(+)} \phi_3 \end{bmatrix} \theta_1^2 \Phi_1$$

$$+ \begin{bmatrix} \left(a_{11}^{(-)}\theta_1^2 + b_{11}^{(-)}\theta_4^2\right) \frac{\phi_0 + \phi_4}{2} \end{bmatrix} \Phi_2 + \begin{bmatrix} a_{01}^{(+)} \phi_3 + a_{12}^{(+)} \phi_1 \end{bmatrix} \theta_1^2 \Phi_3$$

$$+ \begin{bmatrix} a_{00}^{(+)}\theta_1^2 \phi_4 - \left(a_{11}^{(-)}\theta_1^2 + b_{11}^{(-)}\theta_4^2\right) \frac{\phi_2}{2} + a_{22}^{(+)}\theta_1^2 \phi_0 \end{bmatrix} \Phi_4 = 0.$$

$$(4.7)$$

The same form can also be achieved by identifying  $a_{02}^{(+)} = -a_{11}^{(+)} \frac{\phi_0 + \phi_4}{\phi_2}$  and  $b_{02}^{(+)} = -b_{11}^{(+)} \frac{\phi_0 + \phi_4}{\phi_2}$ ). The curve (4.7) can indeed be cast into the form (3.3) with

$$\widetilde{\mathbf{T}}^{T} = \begin{bmatrix} a_{00}^{(+)}\theta_{1}^{2} \\ a_{01}^{(-)}\theta_{1}^{2} \\ \frac{1}{2}(a_{11}^{(-)}\theta_{1}^{2} + b_{11}^{(-)}\theta_{4}^{2}) \\ a_{12}^{(+)}\theta_{1}^{2} \\ a_{22}^{(+)}\theta_{1}^{2} \end{bmatrix}, \quad \mathbb{M}_{\widehat{D}_{4},\mathbf{I}}(\tau) = \begin{bmatrix} \phi_{0} & 0 & 0 & \phi_{0} \\ 0 & \phi_{1} & 0 & \phi_{3} & 0 \\ -\phi_{2} & 0 & \phi_{0} + \phi_{4} & 0 & -\phi_{2} \\ 0 & \phi_{3} & 0 & \phi_{1} & 0 \\ \phi_{4} & 0 & 0 & 0 & \phi_{0} \end{bmatrix}.$$
(4.8)

In fact, this curve is the result obtained in [47, 49] since the terms proportional to  $\theta_4^2$  are  $\beta_2 (\phi_2 \Phi_0 - (\phi_0 + \phi_4) \Phi_2 + \phi_2 \Phi_4) = 2\beta_2 Y_1^2$  which is indeed compatible with [49, eq. (3.86)].

<sup>&</sup>lt;sup>5</sup>In fact, by fixing some of the  $a_{ij}^{\pm}$  and  $b_{ij}^{(\pm)}$  we have found 7 independent candidates of the form  $\alpha \theta_1^2 + \beta \theta_4^2$  for the 5  $\widetilde{T}_i$  entering (3.3). We have carefully analysed all of these possible solutions and found that the only consistent ones are compatible with this assumption. Furthermore, we shall motivate this assumption by modular considerations in the following Subsection 4.2.

II) Setting  $a_{02}^{(+)} = b_{02}^{(+)} = 0$  and identifying  $a_{11}^{(-)} = a_{11}^{(+)}$  and  $b_{11}^{(-)} = b_{11}^{(+)}$ , such that the red and green terms are eliminated and the curve becomes

$$\left[ a_{00}^{(+)} \phi_0 + a_{22}^{(+)} \phi_4 \right] \theta_1^2 \Phi_0 + \left[ a_{01}^{(+)} \phi_1 + a_{12}^{(+)} \phi_3 \right] \theta_1^2 \Phi_1 + \left[ a_{11}^{(+)} \theta_1^2 + b_{11}^{(+)} \theta_4^2 \right] (\phi_0 + \phi_4) \Phi_2$$

$$+ \left[ a_{01}^{(+)} \phi_3 + a_{12}^{(+)} \phi_1 \right] \theta_1^2 \Phi_3 + \left[ a_{00}^{(+)} \phi_4 + a_{22}^{(+)} \phi_0 \right] \theta_1^2 \Phi_4 = 0.$$

$$(4.9)$$

This curve can be cast into the form (3.3) with

$$\widetilde{\mathbf{T}}^{T} = \begin{bmatrix} a_{00}^{(+)}\theta_{1}^{2} \\ a_{01}^{(+)}\theta_{1}^{2} \\ a_{11}^{(+)}\theta_{1}^{2} + b_{11}^{(+)}\theta_{4}^{2} \\ a_{12}^{(+)}\theta_{1}^{2} \\ a_{22}^{(+)}\theta_{1}^{2} \end{bmatrix}, \quad \mathbb{M}_{\widehat{D}_{4},\mathrm{II}}(\tau) = \begin{bmatrix} \phi_{0} & 0 & 0 & 0 & \phi_{4} \\ 0 & \phi_{1} & 0 & \phi_{3} & 0 \\ 0 & 0 & \phi_{0} + \phi_{4} & 0 & 0 \\ 0 & \phi_{3} & 0 & \phi_{1} & 0 \\ \phi_{4} & 0 & 0 & 0 & \phi_{0} \end{bmatrix}.$$
(4.10)

So far, both solutions are compatible with all symmetries of the  $(\hat{D}_4, 1)$  theory and cannot be transformed into one another through any obvious reparametrisations or symmetries. We shall see, however, in Subsection 4.3 that case I) is compatible with known results of the 5 dimensional theory and thus is the correct SWC to describe the  $(\hat{D}_4, 1)$  LST. Concerning case II), it is not clear whether it describes a consistent LST (or any 6 dimensional gauge theory).

#### 4.2 Modular properties

Before discussing the 5 dimensional limit of the  $(\hat{D}_4, 1)$  LST, we first discuss modular properties of the two potential SWC characterised by (4.8) and (4.10). Since modular transformations acting on the parameter  $\rho$  (and coordinate  $z_2$ ) only act on the elliptic Weyl characters  $\tilde{T}_i$  (and are thus relatively straight-forward), we shall focus on modular transformations acting on  $\tau$ .

From the perspective of the dual theory (*i.e.* the right quiver in Figure 2), the gauge polynomials (3.2) transform covariantly under an  $SL(2,\mathbb{Z})$  symmetry acting on the modular parameter  $\tau$  (and the coordinate  $z_1$ ), owing to the transformations (A.14) of the Jacobi theta functions. However, the picture is more complicated from the perspective of the  $(\hat{D}_4, 1)$  LST form in eq. (3.3), where in fact only a subgroup of this  $SL(2,\mathbb{Z})$  is manifestly realised. In the following we shall therefore discuss only transformations under the congruence subgroup  $\Gamma_0(4)$ ,<sup>6</sup> which is the largest subgroup we have found that preserves the general structure (3.3) of the SWC.  $\Gamma_0(4)$  is generated by

$$T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \qquad C = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, \qquad \widetilde{S}_4 = C \cdot S \cdot T^4 \cdot S = \begin{bmatrix} 1 & 0 \\ -4 & 1 \end{bmatrix}, \qquad (4.11)$$

where the action of S and T on Jacobi theta functions is explained in Appendix A.2 and the set  $\mathbf{\Phi} := {\Phi_1, \ldots, \Phi_4}^T$  is invariant under C by construction (which effectively acts as  $z_1 \to -z_1$ ). We start by discussing the action of T on  $\mathbb{M}$  and  $\mathbf{\Phi}$ . We can treat (4.8) and (4.10)

<sup>&</sup>lt;sup>6</sup>This group also corresponds to the modular subgroup acting on the coefficients in the change of basis between Weierstrass functions and theta functions given in [62].

simultaneously by introducing

$$\epsilon = \begin{cases} -1 & \text{for I} \end{pmatrix}, \\ 0 & \text{for II} \end{pmatrix}, \quad (\alpha_2, \beta_2) = \begin{cases} \left(\frac{a_{11}^{(-)}}{2}, \frac{b_{11}^{(-)}}{2}\right) & \text{for I} \end{pmatrix}, \\ \left(a_{11}^{(+)}, b_{11}^{(+)}\right) & \text{for II} \end{pmatrix}, \qquad \mathbb{M}_{\widehat{D}_4}(\epsilon) = \begin{bmatrix} \phi_0 & 0 & 0 & 0 & \phi_4 \\ 0 & \phi_1 & 0 & \phi_3 & 0 \\ \epsilon \phi_2 & 0 & \phi_0 + \phi_4 & 0 & \epsilon \phi_2 \\ 0 & \phi_3 & 0 & \phi_1 & 0 \\ \phi_4 & 0 & 0 & 0 & \phi_0 \end{bmatrix},$$

$$(4.12)$$

and  $(\alpha_0, \alpha_1, \alpha_3, \alpha_4) = \left(a_{00}^{(+)}, a_{01}^{(+)}, a_{12}^{(+)}, a_{22}^{(+)}\right)$ , such that the SWC can be written in the form

 $\theta_1^2(z_2;\rho) \begin{bmatrix} \alpha_0 & \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \end{bmatrix} \cdot \mathbb{M}_{\widehat{D}_4}(\epsilon) \cdot \Phi + \theta_4^2(z_2;\rho) \beta_2(\epsilon \phi_2(\Phi_0 + \Phi_4) + (\phi_0 + \phi_4)\Phi_2) = 0, \quad (4.13)$ 

covering both cases. Indeed, under the action of T, the matrix  $\mathbb{M}_{\widehat{D}_4}(\epsilon)$  transform as

$$\mathbb{M}_{\widehat{D}_{4}}(\epsilon) \xrightarrow{T} \begin{bmatrix} \phi_{0} & 0 & 0 & 0 & \phi_{4} \\ 0 & e^{i\pi/4}\phi_{1} & 0 & e^{i\pi/4}\phi_{3} & 0 \\ \epsilon i\phi_{2} & 0 & i(\phi_{0} + \phi_{4}) & 0 & \epsilon i\phi_{2} \\ 0 & -e^{i\pi/4}\phi_{3} & 0 & -e^{i\pi/4}\phi_{1} & 0 \\ \phi_{4} & 0 & 0 & 0 & \phi_{0} \end{bmatrix},$$
(4.14)

which can be compensated by the transformations

$$\alpha_2 \to -i\alpha_2, \qquad \beta_2 \to -i\beta_2, \qquad \alpha_1 \to e^{-i\pi/4}\alpha_1, \qquad \alpha_3 \to -e^{-i\pi/4}\alpha_3,.$$
(4.15)

Therefore T leaves the overall form of the SWC invariant for both cases I) and II).

The generator  $\widetilde{S}_4$  acts in the following way on  $\mathbb{M}_{\widehat{D}_4}(\epsilon)$ 

$$\mathbb{M}_{\widehat{D}_{4}}(\epsilon) \xrightarrow{\widetilde{S}_{4}} \begin{bmatrix} \phi_{4} & 0 & 0 & 0 & \phi_{0} \\ 0 & \phi_{3} & 0 & \phi_{1} & 0 \\ \epsilon \phi_{2} & 0 & \phi_{0} + \phi_{4} & 0 & \epsilon \phi_{2} \\ 0 & \phi_{1} & 0 & \phi_{3} & 0 \\ \phi_{0} & 0 & 0 & 0 & \phi_{4} \end{bmatrix} ,$$
(4.16)

which can be compensated by the permutations

$$\alpha_0 \longleftrightarrow \alpha_4, \quad \text{and} \quad \alpha_1 \longleftrightarrow \alpha_3. \quad (4.17)$$

Therefore also  $\widetilde{S}_4$  leaves the overall form of the SWC invariant for both cases I) and II), which are therefore both manifestly invariant under  $\Gamma_0(4)$ .

We remark, that invariance under  $\tilde{S}_4$  (as explained above) only applies to matrices  $\mathbb{M}$  that remain invariant under the exchanges (4.4). This property therefore serves as additional motivation for imposing this condition when searching for viable forms of the SWC in the previous Subsection.

### 4.3 Lower Dimensional Limit

So far, both solutions I) and II) have been compatible with all symmetries of the  $(\widehat{D}_4, 1)$  LST and are therefore completely equivalent. However, we shall now demonstrate that only solution I) correctly reproduces known results in the literature upon decompactification to 5 dimensions.

To describe the 5 dimensional limit, we shall use the same notation as in the previous Subsection to treat the two cases I) and II) in parallel. First, we take the limit  $Q_{\rho} = e^{2\pi i \rho} \to 0$ (in which the  $(\hat{D}_4, 1)$  LST is described in terms of a conformal field theory): let

$$w = e^{2\pi i z_1}$$
, and  $t = e^{2\pi i z_2}$ , (4.18)

and assume that  $\beta_2 = -Q_{\rho}^{-1/4} \widetilde{\beta}_2 + \mathfrak{o}(Q_{\rho}^{-1/4})$  (while  $\alpha_i = \widetilde{\alpha}_i + \mathfrak{o}(Q_{\rho}^0)$ ), then the leading order contribution of (4.13) takes the form

$$(t-1)^2 \begin{bmatrix} \widetilde{\alpha}_0 & \widetilde{\alpha}_1 & \widetilde{\alpha}_2 & \widetilde{\alpha}_3 & \widetilde{\alpha}_4 \end{bmatrix} \cdot \mathbb{M}_{\widehat{D}_4} \cdot \mathbf{\Phi} + t \widetilde{\beta}_2 (\epsilon \phi_2(\Phi_0 + \Phi_4) + (\phi_0 + \phi_4)\Phi_2) = 0.$$
(4.19)

We next extract the leading order in the limit  $Q_{\tau} = e^{2\pi i \tau} \to 0$ , which we shall perform in two steps: first we expand the  $\{\Phi_i\}_{i \in \{0...M\}}$  in powers of  $Q_{\tau}$ 

$$0 = 4\widetilde{\alpha}_{0} \left[1 + \mathcal{O}(Q_{\tau})\right] + \widetilde{\alpha}_{1} \left[2(w + w^{-1})Q_{\tau}^{\frac{1}{8}} + \mathcal{O}(Q_{\tau}^{9/8})\right] + \widetilde{\alpha}_{2} \left[2(w^{2} + 2\epsilon + w^{-2})Q_{\tau}^{\frac{1}{4}} + \mathcal{O}(Q_{\tau}^{5/4})\right] + \widetilde{\alpha}_{3} \left[2(w^{3} + w^{-3})Q_{\tau}^{\frac{5}{8}} + \mathcal{O}(Q_{\tau}^{13/8})\right] + \widetilde{\alpha}_{4} \left[4(w^{4} + w^{-4})Q_{\tau} + \mathcal{O}(Q_{\tau}^{2})\right] + \widetilde{\beta}_{2} \left[2(w^{2} + 2\epsilon + w^{-2})Q_{\tau}^{\frac{1}{4}} + \mathcal{O}(Q_{\tau}^{5/4})\right].$$

$$(4.20)$$

In order to perform the full limit, we need to make certain assumptions regarding the  $Q_{\tau}$  dependence of the parameters  $\tilde{\alpha}_{0,\dots,4}$  and  $\tilde{\beta}_2$ . As in the case of the  $Q_{\rho} \to 0$  limit, we shall assume a behaviour that preserves as much of the 6 dimensional symmetries as possible. Concretely, we maintain the exchange symmetries

$$\widetilde{\alpha}_0 \longleftrightarrow \widetilde{\alpha}_4, \quad \text{and} \quad \widetilde{\alpha}_1 \longleftrightarrow \widetilde{\alpha}_3, \quad (4.21)$$

which enforces that  $\tilde{\alpha}_0$  and  $\tilde{\alpha}_4$  (as well as  $\tilde{\alpha}_1$  and  $\tilde{\alpha}_3$ ) have the same leading behaviour in  $Q_{\tau}$ . However, this implies that the contribution with  $\tilde{\alpha}_4$  in (4.20) (as well as the contribution with  $\tilde{\alpha}_3$ ) is subleading and therefore drops out in the limit  $Q_{\tau} \to 0$ . In order to ensure that none of the remaining terms drop out in the limit (which we would interpret as a signal of a modification of the symmetries in the lower dimensional theory), we assume the following scaling behaviour

$$\widetilde{\alpha}_{0} = \widetilde{\alpha}'_{0} Q_{\tau}^{s} + \mathfrak{o}(Q_{\tau}^{s}), \qquad \widetilde{\alpha}_{1} = \widetilde{\alpha}'_{1} Q_{\tau}^{s-\frac{1}{8}} + \mathfrak{o}(Q_{\tau}^{s-1/8}), 
\widetilde{\alpha}_{2} = \widetilde{\alpha}'_{2} Q_{\tau}^{s-\frac{1}{4}} + \mathfrak{o}(Q_{\tau}^{s-1/4}), \qquad \widetilde{\beta}_{2} = \widetilde{\beta}'_{2} Q_{\tau}^{s-\frac{1}{4}} + \mathfrak{o}(Q_{\tau}^{s-1/4}). \qquad (4.22)$$

where  $s \in \mathbb{R}$  is undetermined and shall not be important in the following. We thus obtain the following limit for the SWC

$$(t-1)^2 \left( w^2 + w^{-2} + \frac{\widetilde{\alpha}'_1}{\widetilde{\alpha}'_2} (w+w^{-1}) + 2\frac{\widetilde{\alpha}'_0 + \epsilon \widetilde{\alpha}'_2}{\widetilde{\alpha}'_2} \right) + t \frac{\widetilde{\beta}'_2}{\widetilde{\alpha}'_2} \left( w^2 + 2\epsilon + w^{-2} \right) = 0.$$
(4.23)

This curve can be directly compared with the SWC of the Sp(N) with  $N_f$  fundamentals discussed in [51]: let

$$M_1 + M_1^{-1} + M_2 + M_2^{-1} = -\frac{\widetilde{\alpha}'_1}{\widetilde{\alpha}'_2}, \qquad (M_1 + M_1^{-1})(M_2 + M_2^{-1}) + 2 = \frac{2\widetilde{\alpha}'_0}{\widetilde{\alpha}'_2} + 2\epsilon, \qquad (4.24)$$

<sup>&</sup>lt;sup>7</sup>Here we assume a scaling of all parameters of the theory, which preserves as much structure as possible of the 6 dimensional theory. Indeed, a weaker scaling of  $\beta_2$  would imply that the lower dimensional theory has a different gauge group structure.

such that we can rewrite (4.23) in the language of [51]:

$$t^{2}p_{2}(w) + \left(\frac{\widetilde{\beta}_{2}'}{\widetilde{\alpha}_{2}'}(w^{2} + 2\epsilon + w^{-2}) + 2\left(M_{1} + M_{1}^{-1} + M_{2} + M_{2}^{-1}\right)(w + w^{-1}) - 2(M_{1} + M_{1}^{-1})(M_{2} + M_{2}^{-1}) - 4 - 2(w^{2} + w^{-2})\right)t + p_{2}(w^{-1}) = 0, \quad (4.25)$$

where we have defined

$$p_2(w) := w^{-2} \prod_{i=1}^{2} (w - M_i)(w - M_i^{-1}) = p_2(w^{-1}).$$
(4.26)

In order to match the contribution of order t in (4.25) (with  $(1 - M_i^{\pm}) = (1 - M_i)(1 - M_i^{-1})$ )

$$p_1(w) = -\frac{(w+1)^2 \prod_{i=1}^2 (1-M_i^{\pm})}{2w} + \frac{(w-1)^2 \prod_{i=1}^2 (1+M_i^{\pm})}{2w} + \frac{\beta_2}{\alpha_2} (w^2 + 2\epsilon + w^{-2}) - 2(w^2 - 2 + w^{-2}),$$
(4.27)

to [51] requires  $\epsilon = -1$  and  $\beta_2/\alpha_2 = 2 + q^{-1}$ , where q is the exponentiated complex gauge coupling of Sp(0). In this way, we can indeed interpret the resulting SWC as that of a Sp(0) with SO(4) flavor symmetry 5d SCFT [64, 65] since  $p_2(w)$  has the structure of the corresponding SO(4) gauge polynomial. This is indeed a consistent lower-dimensional limit of the 6 dimensional  $(\hat{D}_4, 1)$  LST. Since this fixes  $\epsilon = -1$ , this also suggests that the correct SWC of the latter theory is in fact described by case I) of Subsection 4.1, *i.e.* eq. (4.8). This is also compatible with the result obtained in [47, 49].

# 5 Seiberg-Witten Curve of $(\widehat{D}_5, 1)$

After having discussed in detail the construction of the SWC of the  $(\hat{D}_4, 1)$  LST in the previous Section, we now focus on the  $(\hat{D}_5, 1)$  theory, for which the result is not known in the literature. Following the same steps as in the previous case, we show that we find again a unique solution for the SWC.

#### 5.1 Construction

We explain the construction of the  $(\widehat{D}_5, 1)$  LST SWC following the same general ansatz (3.5) outlined in Section 3.2. Using the change of basis (A.10), we can reformulate this curve in the basis  $\{\Phi_i\}_{i \in \{0...5\}}$  of degree 10 theta functions

$$\begin{bmatrix} (a_{00}^{(+)}\theta_1^2 + b_{00}^{(+)}\theta_4^2)\phi_0 + \Xi_{11}^-\phi_2 + \Xi_{22}^-\phi_4 \end{bmatrix} \Phi_0 + \begin{bmatrix} (a_{01}^{(+)}\theta_1^2 + b_{01}^{(+)}\theta_4^2)\phi_1 + \Xi_{12}^-\phi_3 + \Xi_{22}^+\phi_5 \end{bmatrix} \Phi_1 + \\ \begin{bmatrix} (a_{02}^{(+)}\theta_1^2 + b_{02}^{(+)}\theta_4^2)\phi_2 + \Xi_{12}^+\phi_4 + \Xi_{11}^+\phi_5 \end{bmatrix} \Phi_2 + \begin{bmatrix} (a_{02}^{(+)}\theta_1^2 + b_{02}^{(+)}\theta_4^2)\phi_3 + \Xi_{12}^+\phi_1 + \Xi_{11}^+\phi_0 \end{bmatrix} \Phi_3 + \\ \begin{bmatrix} (a_{01}^{(+)}\theta_1^2 + b_{01}^{(+)}\theta_4^2)\phi_4 + \Xi_{12}^-\phi_2 + \Xi_{22}^+\phi_0 \end{bmatrix} \Phi_4 + \begin{bmatrix} (a_{00}^{(+)}\theta_1^2 + b_{00}^{(+)}\theta_4^2)\phi_5 + \Xi_{11}^-\phi_3 + \Xi_{22}^-\phi_1 \end{bmatrix} \Phi_5 = 0, \\ \end{bmatrix}$$

$$(5.1)$$

where we have defined

$$\Xi_{ij}^{\pm} := \frac{a_{ij}^{(+)} \pm a_{ij}^{(-)}}{2} \theta_1^2 + \frac{b_{ij}^{(+)} \pm b_{ij}^{(-)}}{2} \theta_4^2.$$
(5.2)

To simplify the assignment of SU(2) and SU(1) Weyl characters, we shall assume the symmetries<sup>8</sup>

$$\widetilde{T}_i \leftrightarrow \widetilde{T}_j, \ \Phi_i \leftrightarrow \Phi_j, \qquad \text{for} \qquad (i,j) \in \{(0,5), (1,4)\}, \qquad (5.3)$$

analogous to (4.4) in the case of the  $(\hat{D}_4, 1)$  LST. These symmetries impose

$$b_{00}^{(+)} = b_{01}^{(+)} = a_{02}^{(+)} = b_{02}^{(+)} = 0.$$
(5.4)

and we are left with three candidates  $(\Xi_{11}^{\pm}, \Xi_{12}^{\pm} \text{ and } \Xi_{22}^{\pm})$  for two SU(2) characters and two SU(1) characters. The natural choice that provides a symmetry among the SU(1) nodes is to break one of the  $\Xi_{ij}^{\pm}$  (for  $(ij) \in \{(11), (12), (22)\}$ ) into two SU(1) characters. There is a priori three possibilities, schematically:

$$\begin{array}{lll} \Xi_{11}^{\pm} \to SU(1)^2 & \Xi_{11}^{\pm} \to SU(2) & \Xi_{11}^{\pm} \to SU(2) \\ 1) & \Xi_{12}^{\pm} \to SU(2), & 2) & \Xi_{12}^{\pm} \to SU(1)^2, & 3) & \Xi_{12}^{\pm} \to SU(2). \\ & \Xi_{22}^{\pm} \to SU(2) & \Xi_{22}^{\pm} \to SU(2) & \Xi_{22}^{\pm} \to SU(1)^2 & (5.5) \end{array}$$

All three choices lead to SWC that are compatible with all symmetries of the  $(\hat{D}_5, 1)$  LST and (as we shall discuss in the next Subsection), have interesting modular properties: we shall in particular see that there exist modular transformations that transform the solutions 2) and 3) into one another. Furthermore, we shall see in Subsection 5.3, that choice 3) provides a (natural)  $Q_{\rho}, Q_{\tau} \to 0$  limit that is compatible with known 5 dimensional results in the literature. Since this is a strong indication that 3) is the correct 6 dimensional SWC of the  $(\hat{D}_5, 1)$  LST, we provide here the explicit details of this solution:

$$a_{11}^{(+)} = b_{11}^{(+)} = a_{22}^{(+)} = b_{22}^{(+)} = b_{12}^{(+)} = b_{12}^{(-)} = 0,$$
(5.6)

such that

$$\widetilde{\mathbf{T}}^{T} = \begin{bmatrix} a_{00}^{(+)}\theta_{1}^{2} \\ (a_{22}^{(+)} + a_{22}^{(-)})\theta_{1}^{2} \\ a_{11}^{(-)}\theta_{1}^{2} + b_{11}^{(-)}\theta_{4}^{2} \\ a_{12}^{(-)}\theta_{1}^{2} + b_{12}^{(-)}\theta_{4}^{2} \\ a_{01}^{(+)}\theta_{1}^{2} \\ (a_{22}^{(+)} - a_{22}^{(-)})\theta_{1}^{2} \end{bmatrix}, \quad \mathbb{M}_{\widehat{D}_{5}}(\tau) = \begin{bmatrix} \phi_{0} & 0 & 0 & 0 & \phi_{5} \\ 0 & \phi_{5} & 0 & 0 & \phi_{5} \\ 0 & \phi_{5} & 0 & -\phi_{3} \\ 0 & -\phi_{3} & \phi_{4} & \phi_{1} & -\phi_{2} & 0 \\ 0 & \phi_{1} & 0 & 0 & \phi_{4} & 0 \\ \phi_{4} & 0 & 0 & 0 & 0 & \phi_{1} \end{bmatrix}.$$
(5.7)

### 5.2 Modular properties

In the same way as in the  $(\widehat{D}_4, 1)$  LST in Subsection 4.2, we indicate the action of the generators of a  $\Gamma_0(5)$  modular group (acting on the modular parameter  $\tau$  and coordinate  $z_1$ ) on the solutions (5.5). The congruence subgroup  $\Gamma_0(5) \subset SL(2,\mathbb{Z})$  is generated by:

$$T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \quad \widetilde{S}_{5,1} = ST^2ST^3ST^2 = \begin{bmatrix} -3 & -5 \\ 5 & 8 \end{bmatrix}, \quad \widetilde{S}_{5,2} = ST^5S = \begin{bmatrix} -1 & 0 \\ 5 & -1 \end{bmatrix}.$$
(5.8)

<sup>&</sup>lt;sup>8</sup>We motivate this symmetry by the natural  $\mathbb{Z}_2$  action of the SU(1) nodes of the quiver which are labelled by  $\widetilde{T}_{0,1,4,5}$ .

The overall form of  $\mathbb{M}$  is left untouched by the action of T and  $\widetilde{S}_{5,2}$ . However,  $\widetilde{S}_{5,2} \circ \widetilde{S}_{5,1}$  acts non-trivially on  $\Phi$  and  $\mathbb{M}_{\widehat{D}_5}$  given in (5.7) (see (A.13)), in particular

$$\begin{bmatrix} \phi_0 & 0 & 0 & 0 & 0 & \phi_5 \\ 0 & \phi_5 & 0 & 0 & \phi_0 & 0 \\ -\phi_2 & 0 & \phi_0 & \phi_5 & 0 & -\phi_3 \\ 0 & -\phi_3 & \phi_4 & \phi_1 & -\phi_2 & 0 \\ 0 & \phi_1 & 0 & 0 & \phi_4 & 0 \\ \phi_4 & 0 & 0 & 0 & 0 & \phi_1 \end{bmatrix} \xrightarrow{\xrightarrow{}}_{\tilde{S}_{5,2} \circ \tilde{S}_{5,1}} \begin{bmatrix} \phi_0 & 0 & 0 & 0 & 0 & \phi_5 \\ 0 & 0 & \phi_0 & \phi_5 & 0 & 0 \\ -\phi_4 & \phi_5 & 0 & 0 & \phi_0 & -\phi_1 \\ 0 & \phi_3 & -\phi_4 & -\phi_1 & \phi_2 & 0 \\ 0 & 0 & \phi_2 & \phi_3 & 0 & 0 \\ \phi_2 & 0 & 0 & 0 & \phi_3 \end{bmatrix}$$
(5.9)

This matrix, however, corresponds precisely to the solution called in 2) in (5.5), indicating that the two are not independent, but are related through a non-trivial modular transformation (and are thus simply equivalent presentations of the same curve.)

#### 5.3 Lower dimensional limit

In order to compute the 5 dimensional limit of the SWC characterised by (5.7), we follow the same double limit  $Q_{\rho} \to 0$  and  $Q_{\tau} \to 0$  as explained in Section 4.3 in the case of the  $(\hat{D}_4, 1)$  LST, which preserves as much of the higher dimensional symmetries as possible. In the interest of brevity, we shall only display the final result, which we can cast in the form

$$(t-1)^{2} \left[ (w^{3}+w^{-3}) + \frac{\alpha_{2}}{\alpha_{3}}(w^{2}+w^{-2}) + \left(\frac{\alpha_{1}}{\alpha_{3}} - 1\right)(w+w^{-1}) + 2\frac{\alpha_{0}}{\alpha_{3}} - 2\frac{\alpha_{2}}{\alpha_{3}} \right] \\ + \left[ \frac{\beta_{3}}{\alpha_{3}}(w^{3}-(w+w^{-1})+w^{-3}) + \frac{\beta_{2}}{\alpha_{3}}(w^{2}-2+w^{-2}) \right] t = 0,$$
(5.10)

In order to compare to the results of [51], we define

$$\frac{\alpha_2}{\alpha_3} = -(M_1 + M_1^{-1} + M_2 + M_2^{-1} + M_3 + M_3^{-1}),$$

$$\frac{\alpha_1}{\alpha_3} = (1 + M_1 M_2 + M_1 M_3 + M_2 M_3)(1 + (M_1 M_2)^{-1} + (M_1 M_3)^{-1} + (M_2 M_3)^{-1}),$$

$$\frac{\alpha_0}{\alpha_3} = -\frac{(1 + M_1^2)(1 + M_2^2)(1 + M_3^2)}{2M_1 M_2 M_3}.$$
(5.11)

such that the lower dimensional curve can be written in the form

$$t^{2}p_{2}(w) + \left[\frac{\beta_{3}}{\alpha_{3}}(w^{3} - (w + w^{-1}) + w^{-3}) + \frac{\beta_{2}}{\alpha_{3}}(w^{2} - 2 + w^{-2}) - 2p_{2}(w)\right]t + p_{2}(w) = 0, \quad (5.12)$$

with

$$p_2(w) = \prod_{i=1}^3 (w - (M_i + M_i^{-1}) + w^{-1}).$$
(5.13)

Following [51], we redefine  $t \to p_2(w)^{-1}t$ , the SWC becomes:

$$t^{2} + \left[ \left( \frac{\beta_{3}}{\alpha_{3}} - 2 \right) (w^{3} + w^{-3}) + \left( \frac{\beta_{2} - 2\alpha_{2}}{\alpha_{3}} \right) (w^{2} + w^{-2}) + \left( -\chi_{c} + 2 - \frac{\beta_{3}}{\alpha_{3}} \right) (w + w^{-1}) + 2 \left( \chi_{s} - \frac{\beta_{2} - 2\alpha_{2}}{\alpha_{3}} \right) \right] t + p_{2}(w)^{2} = 0,$$
(5.14)

where we used the definition of  $SO(2N_f)$  characters for spinor and conjugate spinor representation [65]:

$$\chi_s := \frac{1}{2} \left( \prod_{i=1}^{N_f} \left( M_i^{-\frac{1}{2}} + M_i^{\frac{1}{2}} \right) + \prod_{i=1}^{N_f} \left( M_i^{-\frac{1}{2}} - M_i^{\frac{1}{2}} \right) \right),$$
(5.15a)

$$\chi_c := \frac{1}{2} \left( \prod_{i=1}^{N_f} \left( M_i^{-\frac{1}{2}} + M_i^{\frac{1}{2}} \right) - \prod_{i=1}^{N_f} \left( M_i^{-\frac{1}{2}} - M_i^{\frac{1}{2}} \right) \right).$$
(5.15b)

We can match the result of [51, eq. (4.31)] by identifying  $U_2 = q(\beta_2 - 2\alpha_2)/\alpha_3$  and  $\beta_3/\alpha_3 = 2 + q^{-1}$ . Furthermore, we impose  $M_3 = M_3^{-1} = -1$ , which has the interpretation of an exponentiated complex mass parameter, which leads to  $\chi_c = -\chi_s$ . This is a natural choice since in the  $Q_{\tau} \to 0$  limit the original SO(10) group is broken down to SO(5) which involves two Coulomb branch moduli, there is therefore a spurious parameter in (5.11). Thus we obtain:

$$t^{2} + q^{-1} \left( (w^{3} + w^{-3}) + U_{2}(w^{2} + w^{-2}) + (q\chi_{s} - 1)(w + w^{-1}) - 2(q\chi_{c} + U_{2}) \right) t + p_{2}(w)^{2} = 0.$$
(5.16)

This result matches the SWC of the Sp(1) with SO(5) flavor symmetry 5d SCFT [64, 65] as a specialisation of Sp(1) with 6 flavours. This therefore serves as a strong consistency check, that (5.7) provides the correct SWC of the  $(\hat{D}_5, 1)$  LST.

# 6 Generalisation to the $(\widehat{D}_M, 1)$ LSTs

In the same manner as for the  $(\widehat{D}_4, 1)$  and  $(\widehat{D}_5, 1)$  theories, we can analyse  $(\widehat{D}_M, 1)$  LSTs for generic  $M \geq 4$ . Here we only indicate a general pattern we have found by studying explicit cases up to M = 12: In the following we simply state the form of the modular matrix  $\mathbb{M}$  and  $\widetilde{\mathbf{T}}$ .

For M odd, we distinguish the cases

• M = 4M' + 3:

• 
$$M = 4M' + 1$$
  

$$\widetilde{\mathbf{T}}^{T} = \begin{bmatrix} \alpha_{0}\theta_{1}^{2} & \alpha_{1}\theta_{1}^{2} & \alpha_{2}\theta_{1}^{2} + \beta_{2}\theta_{4}^{2} & \cdots & \alpha_{4M'-1}\theta_{1}^{2} + \beta_{4M'-1}\theta_{4}^{2} & \alpha_{4M'}\theta_{1}^{2} & \alpha_{4M'+1}\theta_{1}^{2} \end{bmatrix}, \quad (6.1)$$

$$\mathbb{M}_{\widehat{D}_{4M'+1}} = \begin{bmatrix} \phi_{0} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \phi_{4M'+1} \\ 0 & \phi_{4M'+1} & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_{0} & \phi_{1} \\ -\phi_{2} & 0 & \phi_{0} & \cdots & 0 & 0 & \cdots & \phi_{4M'+3} & 0 & -\phi_{4M'-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ -\phi_{2M'+1} & 0 & 0 & \cdots & \phi_{0} & \phi_{4M'+1} & \cdots & 0 & 0 & -\phi_{2M'} \\ 0 & -\phi_{2M'} & 0 & \cdots & \phi_{4M'} & \phi_{1} & \cdots & 0 & -\phi_{2M'+1} & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & -\phi_{4M'-1} & \phi_{4M'+2} & \cdots & 0 & 0 & \cdots & \phi_{1} & -\phi_{2} & 0 \\ 0 & \phi_{1} & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_{4M'} & 0 \\ \phi_{4M'} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \phi_{1} \end{bmatrix}. \quad (6.2)$$

$$\widetilde{\mathbf{T}}^{T} = \begin{bmatrix} \alpha_0 \theta_1^2 & \alpha_1 \theta_1^2 & \alpha_2 \theta_1^2 + \beta_2 \theta_4^2 & \cdots & \alpha_{4M'+1} \theta_1^2 + \beta_{4M'+1} \theta_4^2 & \alpha_{4M'+2} \theta_1^2 & \alpha_{4M'+3} \theta_1^2 \end{bmatrix},$$
(6.3)

$$\mathbb{M}_{\widehat{D}_{4M'+3}} = \begin{bmatrix} \phi_0 & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \phi_{4M'+3} \\ 0 & \phi_{4M'+3} & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_0 & 0 \\ -\phi_2 & 0 & \phi_0 & \cdots & 0 & 0 & \cdots & \phi_{4M'+3} & 0 & -\phi_{4M'+1} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ -\phi_{2M'+2} & 0 & 0 & \cdots & \phi_1 & \phi_{4M'+2} & \cdots & 0 & -\phi_{2M'+2} & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & -\phi_{4M'+1} & \phi_{4M'+2} & \cdots & 0 & 0 & \cdots & \phi_1 & -\phi_2 & 0 \\ 0 & \phi_1 & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_{4M'+2} & 0 \\ \phi_{4M'+2} & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_1 \end{bmatrix} .$$
(6.4)

while for M even we distinguish

• M = 4M' + 2

• 
$$M = 4M'$$
  
 $\widetilde{\mathbf{T}} = \begin{bmatrix} \alpha_0 \theta_1^2 & \alpha_1 \theta_1^2 & \alpha_2 \theta_1^2 + \beta_2 \theta_4^2 & \cdots & \alpha_{4M'-2} \theta_1^2 + \beta_{4M'-2} \theta_4^2 & \alpha_{4M'-1} \theta_1^2 & \alpha_{4M'} \theta_1^2 \end{bmatrix}, \quad (6.5)$   
 $\mathbb{M}_{\widehat{D}_{4M'}} = \begin{bmatrix} \phi_0 & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_{4M'} & 0 \\ 0 & \phi_1 & 0 & \cdots & 0 & \cdots & 0 & \phi_{4M'-1} & 0 \\ -\phi_2 & 0 & \phi_2 & \cdots & 0 & \cdots & \phi_{4M'} & 0 & -\phi_{4M'-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots & \vdots \\ -\phi_{2M'} & 0 & 0 & \cdots & \phi_0 + \phi_{4M'} & \cdots & 0 & 0 & -\phi_{2M'} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots & \vdots \\ -\phi_{4M'-2} & 0 & \phi_{4M'} & \cdots & 0 & \cdots & \phi_0 & 0 & -\phi_2 \\ 0 & \phi_{4M'-1} & 0 & \cdots & 0 & \cdots & 0 & \phi_1 & 0 \\ \phi_{4M'} & 0 & 0 & \cdots & 0 & \cdots & 0 & \phi_0 \end{bmatrix}, \quad (6.6)$ 

$$\widetilde{\mathbf{T}} = \begin{bmatrix} \alpha_0 \theta_1^2 & \alpha_1 \theta_1^2 & \alpha_2 \theta_1^2 + \beta_2 \theta_4^2 & \cdots & \alpha_{4M'} \theta_1^2 + \beta_{4M'} \theta_4^2 & \alpha_{4M'+1} \theta_1^2 & \alpha_{4M'+2} \theta_1^2 \end{bmatrix}, \quad (6.7)$$
$$\mathbb{M}_{\widehat{D}_{4M'+2}} = \begin{bmatrix} \phi_0 & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & \phi_{4M'} \\ 0 & \phi_1 & 0 & \cdots & 0 & \cdots & 0 & \phi_{4M'-1} & 0 \\ -\phi_2 & 0 & \phi_2 & \cdots & 0 & \cdots & \phi_{4M'} & 0 & -\phi_{4M'-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & -\phi_{2M'+1} & 0 & \cdots & \phi_1 + \phi_{4M'+1} & \cdots & 0 & -\phi_{2M'+1} & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots & \vdots & \vdots \\ -\phi_{4M'-2} & 0 & \phi_{4M'} & \cdots & 0 & \cdots & \phi_0 & 0 & -\phi_2 \\ 0 & \phi_{4M'-1} & 0 & \cdots & 0 & \cdots & 0 & \phi_1 & 0 \\ \phi_{4M'} & 0 & 0 & \cdots & 0 & \cdots & 0 & \phi_0 \end{bmatrix}. \quad (6.8)$$

Explicit examples for the matrices  $\mathbb{M}$  for  $M \in \{6...9\}$  can be found in Appendix B.

These four classes of result can be expressed is a unifying manner from the dual perspective of  $\widehat{A}_1$  base theory (*i.e.* the right quiver in Figure 2). The results translate to the elliptic gauge polynomials of SO(2M) and Sp(M') with M' = M - 4 and can be understood as a direct generalisation of the case M = 4 (3.1). Elliptic gauge polynomials are finite products of Jacobi theta functions and can be systematically understood as a Higgsing of a SU(N) elliptic gauge polynomial [37]:

$$s_{SU(N)}(z_1;\tau) = \prod_{i=1}^{N} \frac{\theta_1(z_1 - a_i;\tau)}{\theta_1(z_1;\tau)}, \text{ and } \langle \varphi \rangle = (a_1, \dots, a_N),$$
 (6.9)

 $\langle \varphi \rangle$  corresponds to the gauge holonomies of SU(N). For SO(2M), we obtain:

$$F_1(z_1;\tau) = s_{SO(2M)}(z_1,\tau) = \prod_{i=1}^M \frac{\theta_1(z_1 - a_i^{SO};\tau)\theta_1(z_1 + a_i^{SO};\tau)}{\theta_1^2(z_1;\tau)}.$$
(6.10)

which corresponds to a Higgsing of SU(2M) gauge polynomial given by:

$$\langle \varphi \rangle = \left( \pm a_1^{SO}, \dots, \pm a_M^{SO} \right). \tag{6.11}$$

For Sp(M'), we have:

$$F_2(z_1;\tau) = s_{Sp(M')}(z_1,\tau) = \frac{\theta_1^2(2z_1;\tau)}{\theta_1^8(z_1;\tau)} \prod_{i=1}^{M'} \frac{\theta_1(z_1 - a_i^{Sp};\tau)\theta_1(z_1 + a_i^{Sp};\tau)}{\theta_1^2(z_1;\tau)}, \quad (6.12)$$

which corresponds to a Higgsing of SU(2M'+8) gauge polynomial given by:

$$\langle \varphi \rangle = \left(0, 0, \pm \frac{1}{2}, \pm \frac{\tau}{2}, \pm \frac{1+\tau}{2}, \pm a_1^{Sp}, \dots, \pm a_{M'}^{Sp}\right).$$
 (6.13)

We remark that  $F_2$  can be seen as a specialisation of  $F_1$  by using (A.5) and identifying  $(a_1^{SO}, \ldots, a_4^{SO}) = (0, \frac{1}{2}, \frac{\tau}{2}, \frac{\tau+1}{2})$  (which correspond to the so-called discrete holonomies). This result is in line with observations in lower dimensions. In 5 dimensions, SO or Sp gauge theories with flavors can be engineered from Type IIB String Theory through 5-brane webs and  $O7^{\pm}$  orientifold planes [51]. Typically, 5d  $\mathcal{N} = 1$  SO theories are constructed using  $O7^+$ -planes while Sp theories arise from constructions involving  $O7^-$ . In [51], it was pointed out that an  $O7^+$  plane admits an equivalent description as an  $O7^-$  along with  $8D7|_{\text{frozen}}$  i.e. with no moduli attached to them. We interpret the relation between  $F_1$  and  $F_2$  as a 6d manifestation of this observation.

### 7 Conclusion

In this paper we have constructed the Seiberg-Witten Curves (SWCs) for a family of Little String Theories (LSTs) engineered by a single M5-brane probing a transverse (binary) dihedral orbifold geometry  $\mathbb{R}^4_{\perp}/\mathbb{BD}_{M-2}$  (with  $M \geq 4$ ). The six-dimensional world-volume theory on the M5-brane can be described by a quiver gauge theory, as shown in the left part of Figure 2: the quiver is in the form of the Dynkin diagram of  $\hat{D}_M$ , with nodes of type SU(2) and SU(1)respectively and matter in the bifundamental representations. This theory, denoted  $(\hat{D}_M, 1)$  in this work, was shown to be dual [46] (see also [47] for the case M = 4) to a circular quiver gauge theory (*i.e.* with quiver group  $\hat{A}_0$ ) with one node of type Sp(M - 4) and one node of type SO(2M) and bifundamental matter, as shown in the right part of Figure 2. While a form for the SWC of the theory  $(\hat{D}_M, 1)$  with M = 4 was written down in [47, 49], a construction for general M is not known in the literature.

In (3.5) we provide a general ansatz for the SWC for generic  $M \ge 4$  that is compatible with important symmetries and dualities, notably a decomposition in the form of equation (3.3) in terms of (elliptic) Weyl characters  $\tilde{\mathbf{T}}$  associated with the different gauge nodes, a basis  $\{\Phi_j\}_{j\in\{0,\ldots,M\}}$  of  $\mathbb{Z}_2$ -even theta functions of degree 2M (see (A.7)) and a modular matrix  $\mathbb{M}_{ij}$ . The ansatz (3.5) depends on more than the 2M - 2 expected parameters, such that further conditions need to be imposed to determine the SWC: indeed, further parameters are fixed by demanding that the curve can also be cast in the form (3.1), thus reflecting the dual description of the  $(\hat{D}_M, 1)$  theory in terms of the Sp(M-4)-SO(2M) quiver gauge theory explained above. Finally, by assuming a natural dimensional reduction to 5 dimensions, and comparison with known results of the lower dimensional SWC [51] further puts restrictions on the possible solutions. In this way, we have found in Sections 4 and 5 respectively, a unique solution for the SWC of the  $(\hat{D}_4, 1)$  and  $(\hat{D}_5, 1)$  respectively. In the former case, the solution precisely agrees with the curve presented in [47, 49], while the latter constitutes a novel result.

We have systematically analysed in a similar fashion all  $(\hat{D}_M, 1)$  LSTs up to M = 12, which present a number of clear structured patterns (see Appendix B for the modular matrices up to M = 10). These have lead us to conjecture the general form of the  $(\hat{D}_M, 1)$  SWC, as explained in Section 6. Furthermore, using the dual form of the SWC, we also find the form (6.10) and (6.12) respectively, of the gauge potentials  $F_{1,2}$  entering into (3.1).

By determining the form of the SWC of the  $(D_M, 1)$  theory for generic M, our work casts further light on the  $\hat{D}$ -type LSTs, which are far less studied than their  $\hat{A}$ -type counterparts [36, 37]. For example, the relation between the two (equivalent) presentations (3.3) and (3.1) encodes details on the duality between the  $(\hat{D}_M, 1)$  and the Sp(M-4)-SO(2M) theory.

Our results may also hint to further  $\widehat{D}$ -type orbifold theories in 6 dimensions. Indeed, in the case of M = 4 (Section 4) and M = 5 (Section 5), the final form of the SWC could only be determined after taking a (natural) decompactification limit and comparison with lower dimensional results. In both cases, we found alternative forms of the curve ((4.10) and (5.5)respectively) that are compatible with all symmetries of the problem, and only whose natural dimensional reduction leads to a different curve than expected in the lower-dimensional theory [51]. It is an interesting question, whether these curves nevertheless correspond to viable SWC of – possibly deformed – theories in 6 dimensions and whether they are dual to the  $(\widehat{D}_M, 1)$  LSTs. If this is the case, there may exist more intricate limits than the ones considered in Sections 4.3 and 5.3, respectively, that makes them compatible with the known lower dimensional results. We have seen one example of such a scenario being realised, namely the case entitled 2) in (5.5), which we showed to be related to the case 3) by a modular transformation (an element of the congruence subgroup  $\Gamma_0(5)$ ). Since the latter SWC in turn reduces by a natural limit to a known 5 dimensional SWC, this implies that there exists also a (transformed) limit, that reduces the case 2) to a viable 5 dimensional result. Similar mechanisms can a priori not be ruled out for all other 6 dimensional SWCs that we found to be compatible with all higher dimensional symmetries, thus leaving the question whether they indeed realise viable gauge theories. We leave this question for future research.

Let us remark another potential application of our results. It has been established that the Coulomb branch of the supersymmetric vacua of eight supercharge theory is identified with the phase space of the algebraic integrable system [38, 39]. In this context, the SWC is identified with the spectral curve of the corresponding integrable system. Since the integrable system associated with the  $\hat{A}$ -type LST is known to be the double elliptic system [66–68], the SWC discussed in this paper would be identified with a spectral curve of new  $\hat{D}$ -type double elliptic integrable systems.

A further interesting question concerns the generalisation of our work to more general LSTs. Indeed, here we have only considered theories engineered by a single M5 brane probing a dihedral orbifold geometry. It would be interesting to study the generalisation to multiple M5-branes, *i.e.* LSTs of the type  $(\hat{D}_M, N)$  for  $N \ge 1$ . For N > 1, the existence of a duality generalising Figure 2 is not evident and it will be interesting to analyse the structure of the SWC in this case. Finally, instead of considering a transverse dihedral orbifold, it would be interesting to consider other transverse group actions, which lead to the  $\hat{E}$ -type LSTs.

### Acknowledgements

We are very grateful to B. Haghighat for useful correspondence on the SWC of the  $(\hat{D}_4, 1)$ LST. BF and SH are grateful to the Quantum Theory Center ( $\hbar$ QTC, D-IAS, University of Southern Denmark) for kind hospitality when this work was being finalised. BF and TK also acknowledge the Galileo Galilei Institute for Theoretical Physics (GGI) for support and hospitality. The work of TK was supported by CNRS through MITI interdisciplinary programs, EIPHI Graduate School (No. ANR-17-EURE-0002) and Bourgogne-Franche-Comté region.

## A Modular functions and properties

In this Appendix, we define modular functions and give some of their properties that are useful in the computation of SWC.

### A.1 Definitions and properties

We use the notation  $\Theta$  for genus 2 theta functions and  $\theta$  for genus 1 theta functions. Genus 2 theta functions are defined as:

$$\Theta\begin{bmatrix}\delta\\\epsilon\end{bmatrix}(z|\Omega) = \sum_{m\in\mathbb{Z}^2} \exp\left(i\pi(m+\delta)^T \cdot \Omega \cdot (m+\delta) + 2i\pi(z+\epsilon)^T(m+\delta)\right), \quad (A.1)$$

and verify:

$$\Theta\begin{bmatrix}\delta\\\mathbf{0}\end{bmatrix}(z|A\cdot\Omega\cdot A^T) = \Theta\begin{bmatrix}A\cdot\delta\\\mathbf{0}\end{bmatrix}(z\cdot A^{-1}|\Omega).$$
(A.2)

We define the standard Jacobi theta function as:

that are related by shifts in  $\mathbb{Z}/2 \oplus \tau \mathbb{Z}/2$ :

$$\theta_2(z;\tau) = \theta_1\left(z+\frac{1}{2};\tau\right), \quad \theta_3(z;\tau) = \theta_4\left(z+\frac{1}{2};\tau\right), \quad \theta_1(z;\tau) = -ie^{i\pi z}e^{i\pi\frac{\tau}{4}}\theta_4\left(z+\frac{\tau}{2};\tau\right),$$
(A.4)

and verify:

$$\theta_1(z;\tau)\theta_2(z;\tau)\theta_3(z;\tau)\theta_4(z;\tau) = \frac{\theta_1(2z;\tau)}{2\theta_2(0;\tau)\theta_3(0;\tau)\theta_4(0;\tau)}.$$
 (A.5)

In this work, we are interested in basis of modular forms that arise as sections of a degree r line bundle. Elements of such basis  $\{\varphi_i(z;\tau)\}_{i\in I}$  will be denoted as degree r modular forms and have the property that each  $\varphi_i(z;\tau)/\theta_1(z;\tau)^r$  has a pole of maximal order r in z in the

 $\tau \to 0$  limit, for all  $i \in I$ . For example, we define  $X_i$  ( $Y_i$ ), the elements of  $\mathbb{Z}_2$  even (odd) basis of degree M modular forms:

$$X_{i}(z_{1},\tau) = \theta \begin{bmatrix} \frac{i}{M} \\ 0 \end{bmatrix} (Mz_{1}|M\tau) + \theta \begin{bmatrix} \frac{M-i}{M} \\ 0 \end{bmatrix} (Mz_{1}|M\tau), \quad i \in \{0 \dots \lfloor M/2 \rfloor\},$$

$$Y_{i}(z_{1},\tau) = \theta \begin{bmatrix} \frac{i}{M} \\ 0 \end{bmatrix} (Mz_{1}|M\tau) - \theta \begin{bmatrix} \frac{M-i}{M} \\ 0 \end{bmatrix} (Mz_{1}|M\tau), \quad i \in \{1 \dots \lfloor (M-1)/2 \rfloor\},$$
(A.6)

and  $\Phi_i$  the elements of  $\mathbb{Z}_2$  even basis of degree 2M modular forms:

$$\Phi_{j}(z_{1},\tau) = \theta \begin{bmatrix} \frac{j}{2M} \\ 0 \end{bmatrix} (2Mz_{1}|2M\tau) + \theta \begin{bmatrix} \frac{2M-j}{2M} \\ 0 \end{bmatrix} (2Mz_{1}|2M\tau), \quad j \in \{0\dots M\},$$
(A.7)

the two basis are related by Riemann's addition formula

$$\theta\begin{bmatrix}\frac{i}{M}\\0\end{bmatrix}(z_1|\tau)\,\theta\begin{bmatrix}\frac{j}{M}\\0\end{bmatrix}(z_1|\tau) = \sum_{k\in\{0,\frac{1}{2}\}}\theta\begin{bmatrix}k+\frac{i-j}{2M}\\0\end{bmatrix}(0|2\tau)\,\theta\begin{bmatrix}k+\frac{i+j}{2M}\\0\end{bmatrix}(2z_1|2\tau)\,.$$
(A.8)

In particular, we have the following product rules for  $\widehat{D}_4$ :

$$X_{0}^{2} = \phi_{0}\Phi_{0} + \phi_{4}\Phi_{4}, \qquad X_{2}^{2} = \phi_{0}\Phi_{4} + \phi_{4}\Phi_{0}, \\ X_{0}X_{2} = 2\phi_{2}\Phi_{2}, \qquad X_{1}X_{2} = \phi_{1}\Phi_{3} + \phi_{3}\Phi_{1}, \\ X_{1}^{2} = \frac{\phi_{0} + \phi_{4}}{2}\Phi_{2} + \frac{\phi_{2}}{2}(\Phi_{0} + \Phi_{4}), \qquad Y_{1}^{2} = \frac{\phi_{0} + \phi_{4}}{2}\Phi_{2} - \frac{\phi_{2}}{2}(\Phi_{0} + \Phi_{4}), \\ X_{0}X_{1} = \phi_{1}\Phi_{1} + \phi_{3}\Phi_{3}. \qquad (A.9)$$

and product rules for  $\widehat{D}_5$ :

$$\begin{aligned} X_0^2 &= \phi_0 \Phi_0 + \phi_5 \Phi_5, & X_0 X_1 = \phi_1 \Phi_1 + \phi_4 \Phi_4, \\ X_1^2 &= \frac{\phi_0}{2} \Phi_2 + \frac{\phi_5}{2} \Phi_3 + \frac{\phi_4}{2} \Phi_5 + \frac{\phi_1}{2} \Phi_0, & Y_1^2 = \frac{\phi_0}{2} \Phi_2 + \frac{\phi_5}{2} \Phi_3 - \frac{\phi_4}{2} \Phi_5 - \frac{\phi_1}{2} \Phi_0, \\ X_1 X_2 &= \frac{\phi_1}{2} \Phi_3 + \frac{\phi_4}{2} \Phi_2 + \frac{\phi_2}{2} \Phi_4 + \frac{\phi_3}{2} \Phi_1, & Y_1 Y_2 = \frac{\phi_1}{2} \Phi_3 + \frac{\phi_4}{2} \Phi_2 - \frac{\phi_2}{2} \Phi_4 - \frac{\phi_3}{2} \Phi_1, \\ X_2^2 &= \frac{\phi_0}{2} \Phi_4 + \frac{\phi_5}{2} \Phi_1 + \frac{\phi_1}{2} \Phi_5 + \frac{\phi_4}{2} \Phi_0, & Y_2^2 = \frac{\phi_0}{2} \Phi_4 + \frac{\phi_5}{2} \Phi_1 - \frac{\phi_1}{2} \Phi_5 - \frac{\phi_4}{2} \Phi_0, \\ X_0 X_2 &= \phi_2 \Phi_2 + \phi_3 \Phi_3. & (A.10) \end{aligned}$$

### A.2 Modular (sub)group

Defining the action of elements of a modular subgroup  $\Gamma \subseteq SL(2,\mathbb{Z})$  on  $(z,\tau)$ :

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma : (z,\tau) \longmapsto \left(\frac{z}{c\tau+d}, \frac{a\tau+b}{c\tau+d}\right), \tag{A.11}$$

the set of theta functions used to formulate the SWC (3.3) is vector-valued under such modular transformations [69]:

$$\theta \begin{bmatrix} \frac{j}{2M} \\ 0 \end{bmatrix} (2Mz_1 | 2M(\tau+1)) \xrightarrow[T:\tau \to \tau+1]{} e^{i\pi \frac{j^2}{2M}} \theta \begin{bmatrix} \frac{j}{2M} \\ 0 \end{bmatrix} (2Mz_1 | 2M\tau),$$
(A.12a)

$$\theta\begin{bmatrix}\frac{j}{2M}\\0\end{bmatrix}(2Mz_1|2M\tau) \xrightarrow[S:z_1,\tau\to\frac{z_1}{\tau},\frac{-1}{\tau}]{\omega(z_1|\tau)} \sum_{k=0}^{2M-1} e^{-i\pi\frac{kj}{M}}\theta\begin{bmatrix}\frac{k}{2M}\\0\end{bmatrix}(2Mz_1|2M\tau),$$
(A.12b)

with  $\omega(z_1|\tau) = \sqrt{\frac{\tau}{2iM}} \exp\left(\frac{2iMz_1^2}{\tau}\right)$ ,  $S = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$  and  $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ . This generates the following transformation on the base of even modular forms of degree 2M:

$$\boldsymbol{\Phi} := \begin{bmatrix} \Phi_0 \\ \vdots \\ \Phi_M \end{bmatrix} \xrightarrow{T:\tau \to \tau+1} \left[ \exp\left(i\pi \frac{j^2}{2M}\right) \right]_{jj} \cdot \boldsymbol{\Phi}, \quad \boldsymbol{\Phi} \xrightarrow{S:z_1, \tau \to \frac{z_1}{\tau}, \frac{-1}{\tau}} \frac{\omega(z_1|\tau)}{\sqrt{2M}} \left[ 2\cos\left(\frac{ij\pi}{M}\right) \right]_{ij} \cdot \boldsymbol{\Phi}.$$
(A.13)

Standard Jacobi theta functions are re-scaled by modular transformations, in particular  $\theta_1$  verifies:

$$\theta_1(z_1;\tau) \xrightarrow[T:\tau \to \tau+1]{} e^{i\frac{\pi}{4}} \theta_1(z_1;\tau), \quad \theta_1(z_1;\tau) \xrightarrow[S:z_1,\tau \to \frac{z_1}{\tau},\frac{-1}{\tau}] - i\sqrt{\frac{\tau}{i}} e^{i\frac{\pi z_1^2}{\tau}} \theta_1(z_1;\tau).$$
(A.14)

We define the modular congruence subgroup  $\Gamma_0(M)$  as:

$$\Gamma_0(M) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, \mathbb{Z}), \ c \equiv 0 \mod M \right\}.$$
 (A.15)

## **B** M matrices examples

In this appendix, we give explicit examples of  $\mathbb{M}$  matrices entering (3.3) of the SWC for  $\widehat{D}_{6...9}$ . For  $(\widehat{D}_6, 1)$  LST, we have:

$$\mathbb{M}_{\widehat{D}_{6}}(\tau) = \begin{bmatrix} \phi_{0} & 0 & 0 & 0 & 0 & 0 & \phi_{6} \\ 0 & \phi_{1} & 0 & 0 & 0 & \phi_{5} & 0 \\ -\phi_{2} & 0 & \phi_{0} & 0 & \phi_{6} & 0 & -\phi_{4} \\ 0 & -\phi_{3} & 0 & \phi_{1} + \phi_{5} & 0 & -\phi_{3} & 0 \\ -\phi_{4} & 0 & \phi_{6} & 0 & \phi_{0} & 0 & -\phi_{2} \\ 0 & \phi_{5} & 0 & 0 & 0 & \phi_{1} & 0 \\ \phi_{6} & 0 & 0 & 0 & 0 & \phi_{0} \end{bmatrix},$$
(B.1)

for  $(\widehat{D}_7, 1)$  LST:

$$\mathbb{M}_{\widehat{D}_{7}}(\tau) = \begin{bmatrix} \phi_{0} & 0 & 0 & 0 & 0 & 0 & 0 & \phi_{7} \\ 0 & \phi_{7} & 0 & 0 & 0 & \phi_{0} & 0 \\ -\phi_{2} & 0 & \phi_{0} & 0 & 0 & \phi_{7} & 0 & -\phi_{5} \\ -\phi_{4} & 0 & 0 & \phi_{7} & \phi_{0} & 0 & 0 & -\phi_{3} \\ 0 & -\phi_{3} & 0 & \phi_{1} & \phi_{6} & 0 & -\phi_{4} & 0 \\ 0 & -\phi_{5} & \phi_{6} & 0 & 0 & \phi_{1} & -\phi_{2} & 0 \\ 0 & \phi_{1} & 0 & 0 & 0 & \phi_{6} & 0 \\ \phi_{6} & 0 & 0 & 0 & 0 & 0 & \phi_{1} \end{bmatrix},$$
(B.2)

for  $(\widehat{D}_8, 1)$  LST:

$$\mathbb{M}_{\widehat{D}_8}(\tau) = \begin{bmatrix} \phi_0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \phi_8 \\ 0 & \phi_1 & 0 & 0 & 0 & 0 & \phi_7 & 0 \\ -\phi_2 & 0 & \phi_0 & 0 & 0 & \phi_8 & 0 & -\phi_6 \\ 0 & -\phi_3 & 0 & \phi_1 & 0 & \phi_7 & 0 & -\phi_5 & 0 \\ -\phi_4 & 0 & 0 & 0 & \phi_0 + \phi_8 & 0 & 0 & 0 & -\phi_4 \\ 0 & -\phi_5 & 0 & \phi_7 & 0 & \phi_1 & 0 & -\phi_3 & 0 \\ -\phi_6 & 0 & \phi_8 & 0 & 0 & 0 & \phi_0 & 0 & -\phi_2 \\ 0 & \phi_7 & 0 & 0 & 0 & 0 & 0 & \phi_0 \end{bmatrix},$$
(B.3)

for  $(\widehat{D}_9, 1)$  LST:

## References

- Edward Witten. "Some comments on string dynamics". In: STRINGS 95: Future Perspectives in String Theory. July 1995, pp. 501–523. arXiv: hep-th/9507121.
- Paul S. Aspinwall and David R. Morrison. "Point like instantons on K3 orbifolds". In: Nucl. Phys. B 503 (1997), pp. 533-564. DOI: 10.1016/S0550-3213(97)00516-6. arXiv: hep-th/9705104.
- Kenneth A. Intriligator. "New string theories in six-dimensions via branes at orbifold singularities". In: Adv. Theor. Math. Phys. 1 (1998), pp. 271-282. DOI: 10.4310/ATMP.1997.v1.n2.a5. arXiv: hep-th/9708117.
- [4] Amihay Hanany and Alberto Zaffaroni. "Branes and six-dimensional supersymmetric theories". In: Nucl. Phys. B 529 (1998), pp. 180–206. DOI: 10.1016/S0550-3213(98)00355-1. arXiv: hep-th/9712145.
- Ilka Brunner and Andreas Karch. "Branes at orbifolds versus Hanany Witten in sixdimensions". In: JHEP 03 (1998), p. 003. DOI: 10.1088/1126-6708/1998/03/003. arXiv: hep-th/9712143.
- [6] Ofer Aharony. "A Brief review of 'little string theories". In: Class. Quant. Grav. 17 (2000). Ed. by O. Lechtenfeld, J. Louis, Dieter Lust, and H. Nicolai, pp. 929–938. DOI: 10.1088/0264-9381/17/5/302. arXiv: hep-th/9911147.

- [7] David Kutasov. "Introduction to little string theory". In: *ICTP Lect. Notes Ser.* 7 (2002).
   Ed. by C. Bachas, Juan Martin Maldacena, K. S. Narain, and S. Randjbar-Daemi, pp. 165–209.
- [8] Lakshya Bhardwaj, Michele Del Zotto, Jonathan J. Heckman, David R. Morrison, Tom Rudelius, and Cumrun Vafa. "F-theory and the Classification of Little Strings". In: *Phys. Rev. D* 93.8 (2016). [Erratum: Phys.Rev.D 100, 029901 (2019)], p. 086002. DOI: 10.1103/PhysRevD.93.086002. arXiv: 1511.05565 [hep-th].
- [9] Lakshya Bhardwaj. "Revisiting the classifications of 6d SCFTs and LSTs". In: JHEP 03 (2020), p. 171. DOI: 10.1007/JHEP03(2020)171. arXiv: 1903.10503 [hep-th].
- [10] Lakshya Bhardwaj. "Discovering T-dualities of little string theories". In: JHEP 02 (2024),
   p. 046. DOI: 10.1007/JHEP02(2024)046. arXiv: 2209.10548 [hep-th].
- Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Self-Duality and Self-Similarity of Little String Orbifolds". In: *Phys. Rev. D* 94.4 (2016), p. 046006. DOI: 10.1103/PhysRevD.94.046006. arXiv: 1605.02591 [hep-th].
- Julie D. Blum and Kenneth A. Intriligator. "Consistency conditions for branes at orbifold singularities". In: Nucl. Phys. B 506 (1997), pp. 223-235. DOI: 10.1016/S0550-3213(97)00450-1. arXiv: hep-th/9705030.
- Babak Haghighat, Amer Iqbal, Can Kozçaz, Guglielmo Lockhart, and Cumrun Vafa. "M-Strings". In: Commun. Math. Phys. 334.2 (2015), pp. 779–842. DOI: 10.1007/s00220-014-2139-1. arXiv: 1305.6322 [hep-th].
- Babak Haghighat, Can Kozcaz, Guglielmo Lockhart, and Cumrun Vafa. "Orbifolds of M-strings". In: *Phys. Rev. D* 89.4 (2014), p. 046003. DOI: 10.1103/PhysRevD.89.046003. arXiv: 1310.1185 [hep-th].
- [15] Stefan Hohenegger and Amer Iqbal. "M-strings, elliptic genera and N = 4 string amplitudes". In: Fortsch. Phys. 62 (2014), pp. 155–206. DOI: 10.1002/prop.201300035. arXiv: 1310.1325 [hep-th].
- [16] Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "M-strings, monopole strings, and modular forms". In: *Phys. Rev. D* 92.6 (2015), p. 066005. DOI: 10.1103/PhysRevD.92.066005. arXiv: 1503.06983 [hep-th].
- Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Instanton-monopole correspondence from M-branes on S<sup>1</sup> and little string theory". In: *Phys. Rev. D* 93.6 (2016), p. 066016.
   DOI: 10.1103/PhysRevD.93.066016. arXiv: 1511.02787 [hep-th].
- [18] Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Dual Little Strings from F-Theory and Flop Transitions". In: JHEP 07 (2017), p. 112. DOI: 10.1007/JHEP07(2017)112. arXiv: 1610.07916 [hep-th].
- [19] Ambreen Ahmed, Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Bound states of little strings and symmetric orbifold conformal field theories". In: *Phys. Rev. D* 96.8 (2017), p. 081901. DOI: 10.1103/PhysRevD.96.081901. arXiv: 1706.04425 [hep-th].
- [20] Brice Bastian, Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Dual little strings and their partition functions". In: *Phys. Rev. D* 97.10 (2018), p. 106004. DOI: 10.1103/PhysRevD.97.106004. arXiv: 1710.02455 [hep-th].

- [21] Brice Bastian, Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Triality in Little String Theories". In: *Phys. Rev. D* 97.4 (2018), p. 046004. DOI: 10.1103/PhysRevD.97.046004. arXiv: 1711.07921 [hep-th].
- Brice Bastian and Stefan Hohenegger. "Five-Brane Webs and Highest Weight Representations". In: JHEP 12 (2017), p. 020. DOI: 10.1007/JHEP12(2017)020. arXiv: 1706.08750 [hep-th].
- Brice Bastian, Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Beyond Triality: Dual Quiver Gauge Theories and Little String Theories". In: *JHEP* 11 (2018), p. 016.
   DOI: 10.1007/JHEP11(2018)016. arXiv: 1807.00186 [hep-th].
- Brice Bastian, Stefan Hohenegger, Amer Iqbal, and Soo-Jong Rey. "Five-Dimensional Gauge Theories from Shifted Web Diagrams". In: *Phys. Rev. D* 99.4 (2019), p. 046012.
   DOI: 10.1103/PhysRevD.99.046012. arXiv: 1810.05109 [hep-th].
- Brice Bastian and Stefan Hohenegger. "Dihedral Symmetries of Gauge Theories from Dual Calabi-Yau Threefolds". In: *Phys. Rev. D* 99.6 (2019), p. 066013. DOI: 10.1103/PhysRevD.99.066013. arXiv: 1811.03387 [hep-th].
- Brice Bastian and Stefan Hohenegger. "Symmetries in A-type little string theories. Part I. Reduced free energy and paramodular groups". In: JHEP 03 (2020), p. 062. DOI: 10.1007/JHEP03(2020)062. arXiv: 1911.07276 [hep-th].
- [27] Brice Bastian and Stefan Hohenegger. "Symmetries in A-type little string theories. Part II. Eisenstein series and generating functions of multiple divisor sums". In: *JHEP* 03 (2020), p. 016. DOI: 10.1007/JHEP03(2020)016. arXiv: 1911.07280 [hep-th].
- [28] Stefan Hohenegger. "From Little String Free Energies Towards Modular Graph Functions". In: JHEP 03 (2020), p. 077. DOI: 10.1007/JHEP03(2020)077. arXiv: 1911.08172 [hep-th].
- [29] Stefan Hohenegger. "Diagrammatic Expansion of Non-Perturbative Little String Free Energies". In: JHEP 04 (2021), p. 275. DOI: 10.1007/JHEP04(2021)275. arXiv: 2011.06323 [hep-th].
- [30] Stefan Hohenegger and Amer Iqbal. "Symmetric orbifold theories from little string residues". In: *Phys. Rev. D* 103.6 (2021), p. 066004. DOI: 10.1103/PhysRevD.103.066004. arXiv: 2009.00797 [hep-th].
- [31] Baptiste Filoche and Stefan Hohenegger. "Little string instanton partition functions and scalar propagators". In: JHEP 08 (2023), p. 114. DOI: 10.1007/JHEP08(2023)114. arXiv: 2212.09602 [hep-th].
- [32] Baptiste Filoche, Stefan Hohenegger, and Taro Kimura. "Non-perturbative Symmetries of Little Strings and Affine Quiver Algebras". In: JHEP 02 (2024), p. 233. DOI: 10.1007/JHEP02(2024)233. arXiv: 2311.03858 [hep-th].
- [33] Atsushi Kanazawa and Siu-Cheong Lau. "Local Calabi-Yau manifolds of type A via SYZ mirror symmetry". In: J. Geom. Phys. 139 (2019), pp. 103–138. DOI: 10.1016/j.geomphys.2018.12.015. arXiv: 1605.00342 [math.AG].
- [34] Mina Aganagic, Albrecht Klemm, Marcos Marino, and Cumrun Vafa. "The Topological vertex". In: Commun. Math. Phys. 254 (2005), pp. 425–478. DOI: 10.1007/s00220-004-1162-z. arXiv: hep-th/0305132.

- [35] Amer Iqbal, Can Kozcaz, and Cumrun Vafa. "The Refined topological vertex". In: JHEP 10 (2009), p. 069. DOI: 10.1088/1126-6708/2009/10/069. arXiv: hep-th/0701156.
- [36] Harry W. Braden and Timothy J. Hollowood. "The Curve of compactified 6-D gauge theories and integrable systems". In: JHEP 12 (2003), p. 023. DOI: 10.1088/1126-6708/2003/12/023. arXiv: hep-th/0311024.
- [37] Timothy J. Hollowood, Amer Iqbal, and Cumrun Vafa. "Matrix models, geometric engineering and elliptic genera". In: JHEP 03 (2008), p. 069. DOI: 10.1088/1126-6708/2008/03/069. arXiv: hep-th/0310272.
- [38] Ron Donagi and Edward Witten. "Supersymmetric Yang-Mills theory and integrable systems". In: Nucl. Phys. B 460 (1996), pp. 299–334. DOI: 10.1016/0550-3213(95)00609-5. arXiv: hep-th/9510101.
- [39] Nathan Seiberg and Edward Witten. "Gauge dynamics and compactification to threedimensions". In: Conference on the Mathematical Beauty of Physics (In Memory of C. Itzykson). June 1996, pp. 333–366. arXiv: hep-th/9607163.
- [40] Peter Koroteev and Shamil Shakirov. "The quantum DELL system". In: Lett. Math. Phys. 110.5 (2020), pp. 969–999. DOI: 10.1007/s11005-019-01247-y. arXiv: 1906.10354 [hep-th].
- [41] Hirotaka Hayashi and Kantaro Ohmori. "5d/6d DE instantons from trivalent gluing of web diagrams". In: JHEP 06 (2017), p. 078. DOI: 10.1007/JHEP06(2017)078. arXiv: 1702.07263 [hep-th].
- [42] Hirotaka Hayashi, Hee-Cheol Kim, and Kantaro Ohmori. "6d/5d exceptional gauge theories from web diagrams". In: JHEP 07 (2021), p. 128. DOI: 10.1007/JHEP07(2021)128. arXiv: 2103.02799 [hep-th].
- [43] Xing-Yue Wei, Yuji Sugimoto, Futoshi Yagi, and Sung-Soo Kim. "DE-type little strings from glued brane webs". In: JHEP 05 (2023), p. 214. DOI: 10.1007/JHEP05(2023)214. arXiv: 2212.07344 [hep-th].
- [44] Joonho Kim, Sung-Soo Kim, Ki-Hong Lee, Kimyeong Lee, and Jaewon Song. "Instantons from Blow-up". In: *JHEP* 11 (2019). [Erratum: JHEP 06, 124 (2020)], p. 092. DOI: 10.1007/JHEP11(2019)092. arXiv: 1908.11276 [hep-th].
- [45] Hee-Cheol Kim, Minsung Kim, and Yuji Sugimoto. "Blowup equations for little strings". In: JHEP 05 (2023), p. 029. DOI: 10.1007/JHEP05(2023)029. arXiv: 2301.04151 [hep-th].
- [46] Joonho Kim and Kimyeong Lee. "Little strings on  $D_n$  orbifolds". In: *JHEP* 10 (2017), p. 045. DOI: 10.1007/JHEP10(2017)045. arXiv: 1702.03116 [hep-th].
- [47] Babak Haghighat, Joonho Kim, Wenbin Yan, and Shing-Tung Yau. "D-type fiber-base duality". In: *JHEP* 09 (2018), p. 060. DOI: 10.1007/JHEP09(2018)060. arXiv: 1806.10335 [hep-th].
- [48] Nikita Nekrasov and Andrei Okounkov. "Seiberg-Witten theory and random partitions".
   In: Prog. Math. 244 (2006), pp. 525–596. DOI: 10.1007/0-8176-4467-9\_15. arXiv: hep-th/0306238.
- [49] Babak Haghighat and Rui Sun. "M5 branes and Theta Functions". In: JHEP 10 (2019),
   p. 192. DOI: 10.1007/JHEP10(2019)192. arXiv: 1811.04938 [hep-th].

- [50] Babak Haghighat, Wenbin Yan, and Shing-Tung Yau. "ADE String Chains and Mirror Symmetry". In: JHEP 01 (2018), p. 043. DOI: 10.1007/JHEP01(2018)043. arXiv: 1705.05199 [hep-th].
- [51] Hirotaka Hayashi, Sung-Soo Kim, Kimyeong Lee, and Futoshi Yagi. "Seiberg-Witten curves with O7-planes". In: JHEP 11 (2023), p. 178. DOI: 10.1007/JHEP11(2023)178. arXiv: 2306.11631 [hep-th].
- [52] Jin Chen, Yongchao Lü, and Xin Wang. "D-type minimal conformal matter: quantum curves, elliptic Garnier systems, and the 5d descendants". In: JHEP 10 (2023), p. 045. DOI: 10.1007/JHEP10(2023)045. arXiv: 2304.04383 [hep-th].
- [53] Jin Chen, Babak Haghighat, Hee-Cheol Kim, and Marcus Sperling. "Elliptic quantum curves of class  $S_k$ ". In: *JHEP* 03 (2021), p. 028. DOI: 10.1007/JHEP03(2021)028. arXiv: 2008.05155 [hep-th].
- [54] Jin Chen, Babak Haghighat, Hee-Cheol Kim, Marcus Sperling, and Xin Wang.
   "E-string quantum curve". In: Nucl. Phys. B 973 (2021), p. 115602. DOI: 10.1016/j.nuclphysb.2021.115602. arXiv: 2103.16996 [hep-th].
- [55] Jin Chen, Babak Haghighat, Hee-Cheol Kim, Kimyeong Lee, Marcus Sperling, and Xin Wang. "Elliptic quantum curves of 6d SO(N) theories". In: JHEP 03 (2022), p. 154. DOI: 10.1007/JHEP03(2022)154. arXiv: 2110.13487 [hep-th].
- [56] Nathan Seiberg. "New theories in six-dimensions and matrix description of M theory on  $T^5$  and  $T^5/Z_2$ ". In: *Phys. Lett. B* 408 (1997), pp. 98–104. DOI: 10.1016/S0370-2693(97)00805-8. arXiv: hep-th/9705221.
- [57] Michael R. Douglas and Gregory W. Moore. "D-branes, quivers, and ALE instantons". In: (Mar. 1996). arXiv: hep-th/9603167.
- [58] Albion E. Lawrence, Nikita Nekrasov, and Cumrun Vafa. "On conformal field theories in four-dimensions". In: *Nucl. Phys. B* 533 (1998), pp. 199–209. DOI: 10.1016/S0550-3213(98)00495-7. arXiv: hep-th/9803015.
- Grundlehren [59]Jun-ichi Igusa. Theta Functions. der mathematischen Wis-Heidelberg, 1972. 9783642653155. senschaften. Springer Berlin ISBN: DOI: 10.1007/978-3-642-65315-5.
- [60] C. Birkenhake and H. Lange. Complex Abelian Varieties. Grundlehren der mathematischen Wissenschaften. Springer Berlin Heidelberg, 2004. ISBN: 9783540204886. DOI: 10.1007/978-3-662-06307-1.
- [61] Michele Del Zotto, Jonathan J. Heckman, Alessandro Tomasiello, and Cumrun Vafa. "6d Conformal Matter". In: JHEP 02 (2015), p. 054. DOI: 10.1007/JHEP02(2015)054. arXiv: 1407.6359 [hep-th].
- [62] Nikita Nekrasov and Vasily Pestun. "Seiberg-Witten Geometry of Four-Dimensional  $\mathcal{N} = 2$  Quiver Gauge Theories". In: *SIGMA* 19 (2023), p. 047. DOI: 10.3842/SIGMA.2023.047. arXiv: 1211.2240 [hep-th].
- [63] Eric D'Hoker, I. M. Krichever, and D. H. Phong. "The Effective prepotential of N=2 supersymmetric SO(N(c)) and Sp(N(c)) gauge theories". In: Nucl. Phys. B 489 (1997), pp. 211–222. DOI: 10.1016/S0550-3213(97)00033-3. arXiv: hep-th/9609145.

- [64] Hirotaka Hayashi, Sung-Soo Kim, Kimyeong Lee, Masato Taki, and Futoshi Yagi.
   "More on 5d descriptions of 6d SCFTs". In: JHEP 10 (2016), p. 126. DOI: 10.1007/JHEP10(2016)126. arXiv: 1512.08239 [hep-th].
- [65] Hirotaka Hayashi, Sung-Soo Kim, Kimyeong Lee, and Futoshi Yagi. "Discrete theta angle from an O5-plane". In: JHEP 11 (2017), p. 041. DOI: 10.1007/JHEP11(2017)041. arXiv: 1707.07181 [hep-th].
- [66] H. W. Braden, A. Marshakov, A. Mironov, and A. Morozov. "On double elliptic integrable systems. 1. A Duality argument for the case of SU(2)". In: *Nucl. Phys. B* 573 (2000), pp. 553–572. DOI: 10.1016/S0550-3213(99)00739-7. arXiv: hep-th/9906240.
- [67] A. Mironov and A. Morozov. "Double elliptic systems: Problems and perspectives". In: 3rd International Workshop on Supersymmetries and Quantum Symmetries. July 1999. arXiv: hep-th/0001168.
- [68] H. W. Braden, A. Gorsky, A. Odessky, and V. Rubtsov. "Double elliptic dynamical systems from generalized Mukai-Sklyanin algebras". In: *Nucl. Phys. B* 633 (2002), pp. 414– 442. DOI: 10.1016/S0550-3213(02)00248-1. arXiv: hep-th/0111066.
- [69] M. Eichler and D. Zagier. The Theory of Jacobi Forms. Progress in Mathematics. 1985. DOI: 10.1007/978-1-4684-9162-3.