Empirical Mean and Frequency Estimation Under Heterogeneous Privacy: A Worst-Case Analysis

Syomantak Chaudhuri University of California, Berkeley syomantak@berkeley.edu Thomas A. Courtade University of California, Berkeley courtade@berkeley.edu

Abstract

Differential Privacy (DP) is the current gold-standard for measuring privacy. Estimation problems under DP constraints appearing in the literature have largely focused on providing equal privacy to all users. We consider the problems of empirical mean estimation for univariate data and frequency estimation for categorical data, two pillars of data analysis in the industry, subject to heterogeneous privacy constraints. Each user, contributing a sample to the dataset, is allowed to have a different privacy demand. The dataset itself is assumed to be worst-case and we study both the problems in two different formulations – the correlated and the uncorrelated setting. In the former setting, the privacy demand and the user data can be arbitrarily correlated while in the latter setting, there is no correlation between the dataset and the privacy demand. We prove some optimality results, under both PAC error and mean-squared error, for our proposed algorithms and demonstrate superior performance over other baseline techniques experimentally.

1 Introduction

Mean and frequency estimation are pillars of data analysis. Empirical mean estimation plays a crucial role in fields like Modern Portfolio Theory [1], Machine Learning [2], and Healthcare [3] to name a few. For service providers, relative frequencies of categorical data is an import statistic such as distribution of traffic across websites [4], age and geographical distribution of social media users [5], and so on. Thus, it is crucial to study the estimation of these statistics under the lens of privacy due to the increased demand for privacy, as reflected in recent laws such as GDPR and CCPA. We adopt the popular framework of Differential Privacy [6, 7], which has found real-world applications [8–10].

The trade-off between estimation accuracy and afforded privacy is one of the most fundamental questions in this field. An aspect of estimation subject to privacy constraints that has not been studied well in literature is the case when different users demand different levels of privacy for their data. Existing literature mostly focus on a common privacy level for the users (see Wang et al. [11] for references), but this does not reflect the real-world where different users have different privacy requirements [12]. These heterogeneous privacy demands naturally arise in social media platforms, where users can get more features by opting-in to share more data. For example, in a discontinued feature, users on Facebook could share their location with Facebook in order to get notified if a friend is nearby [13]. Thus, heterogeneity in privacy requirements is an important consideration and there needs to be a greater understanding of the accuracy-privacy trade-off. The presented work focuses on understanding this trade-off for mean and frequency estimation.

1.1 Our Contribution and Problem Description

We consider the problem of mean estimation of univariate data and frequency estimation of categorical data under heterogeneous privacy demands in the Central-DP model. In Central-DP, the users trust

Preprint. Under review.

the server and send their true data to the server [7]. The users provide the server with their datapoint (such as their salary in mean estimation or salary bracket in frequency estimation) and a privacy level. The server is then tasked with publishing an estimate of the mean or relative frequencies of the categories in the dataset, while respecting the individual privacy constraints. We consider the adversarial minimax setting where the dataset is the worst-case for the algorithm. This is in contrast to the statistical minimax setting where the dataset is sampled from the worst-case distribution. Under the adversarial regime, we formulate two distinct settings: the so-called correlated and uncorrelated settings. These are explained below, and formally defined in Section 2. Each problem is studied in each setting under both PAC error and mean-squared error.

Since we assume that we are given the heterogeneous privacy demands, simply considering the worst-case dataset might lead to arbitrary correlations between user data values and their individual privacy requirements. Correlations may be reasonable for certain cases. For example, users sharing their salary may demand higher privacy if they are have an extremely high or low salary. Such correlations make it harder to estimate frequencies of categories in which users demand a lot more privacy; this is a problem for mean estimation as well. We emphasize that such correlations, although often accepted to be present in real-world, have never been modeled in the literature prior to this work that the authors are aware of.

It is equally well-motivated to consider the uncorrelated setting where users do not have a correlation between their data and privacy demand. For example, for frequency estimation of users' geographicalregion, there may be no meaningful correlation between a user's location and their privacy demand. We formulate this "uncorrelated" setting by modeling the realization of the dataset as a uniformly random permutation of a worst-case dataset.

We propose algorithms for both PAC error and mean-squared error, for both correlated and uncorrelated settings, for both mean and frequency estimation. Thus, in this work, we consider eight different minimax rates. To obtain tight upper bounds on the minimax rates, we consider slightly different algorithms tailored to each combination of problem, setting, and error metric. However, we also present a fast heuristic algorithm that is agnostic to the underlying modeling assumptions and performs well in practice for all the eight problems. We compare our algorithms to natural baseline algorithms and show superior performance of our algorithms. We also prove the minimax optimality of our algorithms in some problems and error metrics.

Organization: We define the problem in Section 2 and describe our algorithm in Section 3. Experiments are presented in Section 4, followed by theoretical analysis of the minimax rate in Section 5. Extended discussion can be found in Section 6, followed by some concluding remarks.

1.2 Related Work

The two most common models for DP are the Central-DP model and the Local-DP model. In Local-DP, users do not trust the server and send noisy data to the server, ensuring their privacy (see Kasiviswanathan et al. [14] for more details). In this work, we focus on the Central-DP model.

The problem of frequency estimation under privacy constraints is well-studied under homogeneous DP [7, 8, 15, 16]. In terms of heterogeneity in privacy demands, a special case considered in the literature is the existence of a dataset requiring homogeneous privacy in conjunction with a public dataset [15, 17–24]. Mean estimation under privacy is well-studied as well (see Biswas et al. [25] for references). Although most work on mean estimation focuses on the statistical setting, one can use the ideas from empirical frequency estimation of Dwork et al. [7] and corresponding lower bounds from Vadhan [26] to come up with good error bounds for empirical mean estimation.

Alaggan et al. [27] and Jorgensen et al. [28] have proposed task-agnostic methods to deal with Heterogeneous DP (HDP). While their methods are versatile since they are task agnostic, one can not expect competitive performance from their methods for specific tasks like mean or frequency estimation. Further, neither of the methods can handle the case if there is a single user having no privacy demand (public data).

Mean estimation under Heterogeneous DP (HDP) is considered by Ferrando et al. [29] in the Local-DP model, assuming the variance of the distribution is known. HDP mean estimation under the Central-DP model has been studied in Fallah et al. [30], Chaudhuri and Courtade [31], Chaudhuri et al. [32], Cummings et al. [33]; these works have focused on the statistical setting where the data is sampled i.i.d. from some distribution. The problem of creating confidence intervals for the mean under sequential observations in Local-DP was studied by Waudby-Smith et al. [34] for univariate random variables with heterogeneity in privacy demand. Canonne and Sun [35] present the problem of testing whether two datasets are from the same multinomial distribution under a different privacy constraint for each dataset. HDP has also been considered in the context of mechanism design, auctions, and data valuation [30, 36, 37]. Acharya et al. [38] study the problem of ridge regression under HDP and provide an algorithm similar to one of our baseline techniques we use in the experiments. ¹

In the context of modeling correlations between privacy demand and data, the authors are only aware of the work by Ghosh and Roth [39] - a negative result in the context of auction design. The question of how to deal with correlations between data and privacy is a crucial one in HDP. However, the literature lacks any proper formulation of the problem itself.

2 Problem Definition

2.1 Notation

We denote positive real numbers by $\mathbb{R}_{>0}$. The notation [n] refers to the set $\{1, 2, \ldots, n\}$. The probability simplex in *n*-dimensions is represented by Δ_n . Throughout this work the ℓ_1 norm, $\|\cdot\|_1$, is interchangeably used for sum of elements of vectors with positive components. The notation $a \wedge b$ is used to denote min $\{a, b\}$. We shall use $\lceil \cdot \rceil$ to refer to the *modified* ceiling function $\lceil x \rceil = \min\{m \in \mathbb{Z} : m > x\}$. The notation $c\mathbf{1}$ shall denote the vector, of appropriate length, with all entries being equal to c. In this work, the notation $M(\cdot)$, for a randomized algorithm M mapping \mathcal{X}^n to a probability distribution on \mathcal{Y} , will interchangeably be used to refer to the output distribution or a sample from it. For vectors x and y, x/y shall represent element-wise division. The notations \leq and \simeq denote inequality and equality that hold up to a universal multiplicative constant.

2.2 **Problem Definition**

Heterogeneous Differential Privacy (HDP) permits users to have different privacy requirements. The standard definition for HDP is presented in Definition 2.1 [27, 30].

Definition 2.1 (Heterogeneous Differential Privacy). A randomized algorithm $M : \mathcal{X}^n \to \mathcal{Y}$ is said to be ϵ -DP for $\epsilon \in \mathbb{R}^n_{>0}$ if

$$\mathbb{P}\{M(X) \in S\} \le e^{\epsilon_i} \mathbb{P}\{M(X'^i) \in S\} \quad \forall i \in [n],$$
(1)

for all measurable sets $S \subseteq \mathcal{Y}$, where $X, X'^i \in \mathcal{X}^n$ are any two 'neighboring' datasets that differ arbitrarily in only the *i*-th component.

The probability in Definition 2.1 is taken over the randomized algorithm conditioned on the given datasets X, X'^i , i.e., it is a conditional probability. In the DP framework, smaller ϵ means higher privacy. The privacy levels ϵ can range from zero to infinity.

Let *n* be the number of users and let the heterogeneous privacy demand of the users be the vector ϵ . Without loss of generality, we assume it is arranged in a non-decreasing order, i.e., user *i* has privacy requirement ϵ_i , and $\epsilon_i \leq \epsilon_j$ for $i \leq j$. We operate in the Central-DP regime and assume that the privacy demand is public as stated in Assumption 2.2; we justify it in Section 2.3.

Assumption 2.2. The privacy demand ϵ is public.

For normalized *frequency estimation* with k categories (or bins), we have $\mathcal{X} = [k]$ and $\mathcal{Y} = [0, 1]^k$; the parameter k is assumed to be known. For *mean estimation*, without loss of generality, we consider $\mathcal{X} = [0, 1]$ and $\mathcal{Y} = [0, 1]$; one can consider arbitrary length intervals as long as the length of the interval is known, the position of the interval does not need to be known.

We study the problem in the adversarial minimax regime – for an algorithm, the worst case performance over all datasets is analyzed. We formulate two settings to capture possible correlations.

In order to define the two settings, let the underlying dataset be denoted by $D \in \mathcal{X}^n$.

¹Their method can perform arbitrarily worse than the minimax optimal ridge regression algorithm (see Prop algorithm in Section 4 to understand why).

- Correlated setting: In this setting, the dataset is simply D. In other words, user i has privacy demand ϵ_i and datapoint D_i for $i \in [n]$.
- Uncorrelated setting: In this setting, to break correlations between the dataset and the privacy demand, we assume that the actual realization of the dataset is a random uniform permutation of D. Let $\sigma : [n] \to [n]$ denote a random permutation over [n] then we shall denote the realization of the dataset as D_{σ} , i.e., user *i* has the datapoint $D_{\sigma(i)}$ and privacy demand ϵ_i . The notation $\sigma(i)$ indicates which element of D gets mapped to the *i*-th user.

Let $\mu(D)$ refer to the true empirical statistic of the dataset. In other words,

- for mean estimation, $\mu(D) \in [0,1]$ and it is defined as $\mu(D) = \frac{1}{n} \sum_{i=1}^{n} D_i$.
- for frequency estimation, $\mu(D) \in \Delta_k$ and the *j*-th component of $\mu(D)$ is defined as $\mu(D)_j = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{D_i = j\}, j \in [k].$

Note that this function is permutation-invariant $\mu(D) = \mu(D_{\sigma})$. Let \mathcal{M}_{ϵ} refer to the set of all ϵ -DP mechanisms from \mathcal{X}^n to \mathcal{Y} . We define the minimax rates of frequency estimation in Definition 2.3.

Definition 2.3 (Minimax Rates of Frequnecy Estimation). The PAC-minimax rate for k bins, privacy demand ϵ , and error probability β is the minimum value of the $(1 - \beta)$ -th quantile of the ℓ_{∞} error under the worst-case dataset for any ϵ -DP algorithm. The MSE-minimax rate is similarly defined. (A) **Correlated Setting:** The PAC-minimax rate is given by

$$\mathcal{R}_{c}^{f}(k,\beta,\epsilon) = \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [k]^{n}} \min_{\mathbb{P}\{\|M(D) - \mu(D)\|_{\infty} > \alpha\} \le \beta} \alpha , \qquad (2)$$

where the probability is taken over the randomness in the map M. The MSE-minimax rate is given by

$$\mathcal{E}_{c}^{f}(k,\boldsymbol{\epsilon}) = \min_{\boldsymbol{M}\in\mathcal{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{D}\in[k]^{n}} \mathbb{E}[\|\boldsymbol{M}(\boldsymbol{D}) - \boldsymbol{\mu}(\boldsymbol{D})\|_{\infty}^{2}],$$
(3)

where the expectation is taken over the randomness in the map M.

(B) Uncorrelated Setting: The PAC-minimax rate is given by

$$\mathcal{R}^{f}_{u}(k,\beta,\epsilon) = \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [k]^{n}} \min_{\mathbb{P}\{\|M(D_{\sigma}) - \mu(D)\|_{\infty} > \alpha\} \le \beta} \alpha ,$$
(4)

where the probability is taken over the randomness in the map M and the uniform permutation σ of the dataset D. The MSE-minimax rate is given by

$$\mathcal{E}_{u}^{f}(k,\boldsymbol{\epsilon}) = \min_{\boldsymbol{M}\in\mathcal{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{D}\in[k]^{n}} \mathbb{E}[\|\boldsymbol{M}(\boldsymbol{D}_{\sigma}) - \boldsymbol{\mu}(\boldsymbol{D})\|_{\infty}^{2}],$$
(5)

where the expectation is taken over the randomness in the map M and the uniform permutation σ of the dataset D.

Similarly, we define the minimax rates for mean estimation in Definition A.1, deferred to Appendix A.

2.3 Modeling Choice

On Assumption 2.2: In the worst-case, where arbitrary correlation between the data and the privacy demands are allowed, knowing the privacy demand can leak all information about the data. Indeed, one example would be the case where the privacy demand is equal to the data! Protecting the privacy demand in addition to the data is a more complex problem that is not yet studied in literature and hence, we simplify the problem by assuming the privacy demand itself is public and does not need to be protected. With this assumption, our algorithm does not leak any extra information than claimed. This assumption is indeed encountered in certain practical scenarios like social media where anyone can see if some user has decided to keep their profile and data private. Since no prior work has ever tried to model these correlations, we view our work as a first step in this crucial direction. There do exist cases where privacy demands should also be protected but that is beyond the scope of this work. One possible way to decrease such privacy risks is if the service provider allows the users to choose their privacy from a set of pre-determined privacy levels, as is typical in many practical systems.t

On the minimax rates and correlations: Since we consider the error bounds for a given privacy demand and an adversarial dataset, there could be an arbitrary correlation between the dataset and the privacy demand in the correlated setting. Considering arbitrary worst-case correlations avoids the

Algorithm 1 Heterogeneously Private Frequency (HPF)	Table 1: Weight different settings.	s for Algorithm 1 for
Input: weights $\boldsymbol{w} \in \Delta_n, X \in [k]^n$	ALGORITHM	WEIGHTS w
Sample i.i.d. $N_1, \ldots, N_k \sim \text{Laplace}(2 \ \frac{\boldsymbol{w}}{\boldsymbol{\epsilon}} \ _{\infty})$ Calculate $\boldsymbol{y} \in \mathbb{R}_n^k$:	$HPF-C\mathbb{P}$	$\arg\min_{\boldsymbol{w}\in\Delta_n}r_C(\boldsymbol{w},k,\beta,\boldsymbol{\epsilon})$
	HPF-Cℤ	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_C(\boldsymbol{w},k,1,\boldsymbol{\epsilon})$
$y_j = N_j + \sum w_i \mathbb{I}\{X_i = j\}, \ \forall j \in [k].$	HPF-U₽	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},k,\beta,\boldsymbol{\epsilon})$
$i{=}1$	HPF-UE	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},k,1,\boldsymbol{\epsilon})$
Set $y_j \leftarrow \max\{\min\{y_j, 1\}, 0\} \ \forall j \in [k].$	HPF-A	$\frac{1 - e^{-\epsilon}}{\ 1 - e^{-\epsilon}\ _1}$
iciui ii y		

need for obscure assumptions on joint distribution between privacy and data. This may or may not be desired, depending on the application. For example, if the categorical data consists of salary range of users, then the users in the highest and lowest salary range might demand a higher privacy leading to correlations between data and privacy demands. A counterexample is if the categorical data consists of different types of cancers in patients; in this case such correlations need not exist in a meaningful way. Thus, to model such setting without meaningful correlations, we formulate the uncorrelated setting. Another possible way to break the correlation is to go to the statistical minimax setting where the samples are generated i.i.d. from some distribution, independent of the privacy demands, and the goal is to estimate the underlying multinomial distribution.

The above described statistical minimax rate is not related to the correlated adversarial minimax rate. To see this, consider the case where all the users have no privacy constraint; the correlated adversarial minimax error is zero but the statistical minimax error is of the order $\frac{1}{\sqrt{n}}$ (PAC or square-root of MSE). Conversely, consider the case where n/2 users demand no privacy and rest of the users demand $\epsilon \rightarrow 0$ – the statistical rate is of the order $\frac{1}{\sqrt{n}}$ but the correlated adversarial minimax rate is of the order 1. However, the uncorrelated adversarial minimax rates and the statistical minimax rates are closely related (see Theorem 5.3).

We do not consider the worst case of ϵ since the worst-case is trivial when every user demands full privacy ($\epsilon \rightarrow 0$) and no meaningful estimation can be done. Other possible modeling choices, not presented in this work, include a different norm for errors, the statistical setting where the underlying dataset and privacy demand is jointly generated according to some distribution, and so on.

3 Algorithm Description

Let

$$r_C(\boldsymbol{w}, k, \beta, \boldsymbol{\epsilon})^2 = \left\| \boldsymbol{w} - \frac{1}{n} \right\|_1^2 + \log\left(\frac{k}{\beta}\right)^2 \left\| \frac{\boldsymbol{w}}{\boldsymbol{\epsilon}} \right\|_\infty^2, \tag{6}$$

$$r_U(\boldsymbol{w}, k, \beta, \boldsymbol{\epsilon})^2 = \left(\left\| \boldsymbol{w} - \frac{1}{n} \right\|_1^2 \wedge \log\left(\frac{k}{\beta}\right) \|\boldsymbol{w}\|_2^2 \right) + \log\left(\frac{k}{\beta}\right)^2 \left\| \frac{\boldsymbol{w}}{\boldsymbol{\epsilon}} \right\|_\infty^2.$$
(7)

These functions help us track the errors for problems. For frequency estimation, we propose a family of algorithms HPF with the skeleton code presented in Algorithm 1. Table 1 outlines our suggested weights to be used in HPF for the relevant setting. Note that these different weight suggestions are mainly to obtain theoretically good performance in the respective setting. For example, to get good theoretical performance under PAC metric for error level β in the correlated setting, one would use the weights given by HPF-CP from the table. The second-last letter of the algorithm names specify if the algorithm is for correlated setting (C) or uncorrelated setting (U). The last letter specifies if the algorithm is geared for optimizing the PAC error (P) error or the MSE error (E).

While these methods are intended to facilitate characterization of fundamental limits, an alternate fast, heuristic-based algorithm HPF-A is also described which can be used in either setting and error metric, i.e., it is agnostic to the modeling assumptions and error metric. *We stress that while we propose a different algorithm based on the setting and error metric for proving theoretical guarantees,*

HPF-A can be used agnostic to the setting since it enjoys good performance (see Section 4) across the settings and metric, and can be employed in practice.

Similarly, for mean estimation, we propose the family of algorithms HPM presented in Algorithm 2 and the corresponding weights are presented in Table 3, both deferred to Appendix A.

The performance guarantees for the algorithms and lower bounds on the minimax rate can be found in Section 5. Lemma 3.1 guarantees that the algorithms described in Algorithm 1 satisfy the privacy constraint; the proof can be found in Appendix D.

Lemma 3.1. The proposed family of algorithms HPF and HPM satisfy the ϵ -DP constraint defined in Definition 2.1.

Broadly, our proposed algorithms assign different weights to different users instead of equally weighing the users for the estimation task. For example, in frequency estimation, for weights $w \in \Delta_n$, if user *i* has a datapoint in bin *j*, then HPF adds w_i to bin *j* instead of $\frac{1}{n}$. In particular, if the users with more stringent privacy are weighed less, then the magnitude of noise that is required to be added to the frequency estimate for ensuring ϵ -DP can be significantly less. On the flip side, this reweighing introduces bias in the output as the data from less privacy stringent users dominates. Thus, the weights need to be carefully chosen to optimize this trade-off.

The minimization in the algorithm minimizes an upper bound for error metric in the relevant setting. For example, HPF-CP minimizes an upper bound for the $(1 - \beta)$ -th quantile of the ℓ_{∞} -error over the set of weights $w \in \Delta_n$. All the optimization problems are quadratic programs with linear constraints and can be solved using a numerical optimizer like cvxpy [40]. Faster variants of the algorithms which can be computed in $O(n \log n)$ time are discussed in Section 6.

4 **Representative Experiments**



Figure 1: (A): The negative log MSE for six algorithms for three experiments (higher is better) in the correlated regime is plotted. (B): The negative log of the 95-th empirical error quantile for the six algorithms for three experiments (higher is better) in the uncorrelated regime is plotted.

We only present some representative experiments for frequency estimation in this section in Figure 1. More extensive experiments and exact implementation details can be found in Appendix B. In Figure 1(A), we consider three different datasets and construct a privacy demand that is correlated with the categorical data. We aim to optimize the MSE (l_{∞} error) so the variant of HPF that we employ is HPF-CE. The negative of the log MSE is plotted for the different algorithms. Similarly, in Figure 1(B), we consider three different datasets and construct a privacy demand that is uncorrelated with the categorical data. We aim to optimize the PAC error with $\beta = 0.05$, i.e., the 95-th quantile of the error so the variant of HPF that we employ is HPF-UP with $\beta = 0.05$. The negative of the log of the 95-th empirical error quantile is plotted for the different algorithms.

Overall, in both correlated and uncorrelated settings, under both PAC and MSE, HPF-A is a superior algorithm due to it good performance and being agnostic to the modeling choices and the error metric.

5 Performance Analysis

Recall the functions r_C and r_U from (6) and (7) respectively. We denote their corresponding minimum values by

$$R_C(k,\beta,\epsilon) = \left(\min_{\boldsymbol{w}\in\Delta_n} r_C(\boldsymbol{w},k,\beta,\epsilon)\right) \wedge 1,$$
(8)

and

$$R_U(k,\beta,\boldsymbol{\epsilon}) = \left(\min_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},k,\beta,\boldsymbol{\epsilon})\right) \wedge 1.$$
(9)

These functions are relevant in tracking the upper bound values in the various settings.

We summarize our theoretical results in this section. Table 2 summarizes the results presented in this work. The theorems can be found in Appendix C. For the rest of the section, we focus on the question of minimax optimality - are our upper and lower bounds of the same order?

Table 2: A summary of the theoretical results presented in this paper. The setting column specifies the problem – (F)requency estimation or (M)ean estimation, the setting – (C)orrelated or (U)ncorrelated, and the error metric – (P)AC or MS(E). The algorithm column specifies the algorithm used to obtain the upper bound in the next column, the functions R_C and R_U are defined in (8) and (9) respectively. The Lower Bound column references the location where the lower bounds can be found and the last column is on whether our upper and lower bounds are of the same order.

Setting	Algorithm	Upper Bound (UB)	Lower Bound (LB)	$UB \simeq LB?$	
F, C, ℙ	HPF-C₽	$R_C(k,eta,oldsymbol{\epsilon})$	Theorem C.3	Yes	
F, C, E	HPF-Cℤ	$R_C(k, 1, \epsilon)^2$	×	N/A	
F, U, ₽	HPF-Uℙ	$R_U(k,eta,oldsymbol{\epsilon})$	Theorem C.3	Under certain assumptions	
F, U, E	HPF-Uℝ	$R_U(k,1,oldsymbol{\epsilon})^2$	Theorem C.4	Near-optimal	
M, C, ℙ	HPM-C₽	$R_C(1,eta,oldsymbol{\epsilon})$	Theorem C.3	Yes	
М, С, Е	HPM-CE	$R_C(1, e, \epsilon)^2$	×	N/A	
M, U, ℙ	HPM-Uℙ	$R_U(1,eta,oldsymbol{\epsilon})$	Theorem C.3	Under certain assumptions	
М, С, Е	HPM-UE	$R_U(1,e,oldsymbol{\epsilon})^2$	Theorem C.4	Yes	

5.1 Minimax Optimality in Correlated Regime

In the correlated regime, we present Theorem 5.1 on the minimax optimality of HPF-CP and HPM-CP in a certain regime of interest of the PAC minimax rate. The proof is presented in Appendix I.

Theorem 5.1 (PAC Minimax Optimality: Correlated Setting). (A) For $\frac{1}{2n} \leq \mathcal{R}_c^f(k, \beta, \epsilon) \leq \frac{1}{4}$, the lower bound in Theorem C.3 implies that error incurred by HPF-C \mathbb{P} is of the same order as the minimax rate, i.e.,

$$\mathcal{R}_c^f(k,\beta,\epsilon) \simeq R_C(k,\beta,\epsilon)$$

(B) For $\frac{1}{2n} \leq \mathcal{R}_c^m(\beta, \epsilon) \leq \frac{1}{4}$, the lower bound in Theorem C.3 implies that error incurred by HPM-CP is of the same order as the minimax rate, i.e.,

$$\mathcal{R}_c^m(\beta, \epsilon) \simeq R_C(1, \beta, \epsilon)$$

In particular, when the PAC minimax rate is at least $\frac{1}{2n}$ and at most $\frac{1}{4}$, then our upper bound is of the same order as the minimax rate, for both mean and frequency estimation. We emphasize that this really is the regime of interest for the problem being considered; without any privacy constraint, normalized mean and frequency count is naturally of the precision $\frac{1}{n}$. Theorem C.3 also assumes that the PAC minimax rate is bounded away from the trivial upper bound of $\frac{1}{2}$. Thus, for the practically important regime of values of the PAC minimax rate, our proposed algorithms are order optimal.

5.2 Minimax Optimality in Uncorrelated Regime

In the uncorrelated regime, to compare the PAC upper and lower bounds, we consider two regimes. This is primarily because of the lower bounds presented in Theorem C.3 are difficult to analyze without further simplifications. Let $Var(e^{-\epsilon})$ denote the empirical variance of the vector $e^{-\epsilon}$. Theorem 5.2 shows how the upper and lower bounds behave in the high privacy regime; the proof can be found in Appendix I.

Theorem 5.2 (PAC Minimax Optimality: Uncorrelated Setting).

Assume high privacy demand where $\|\boldsymbol{\epsilon}\|_{\infty} \leq 1$, for the two problems. (A) Frequency Estimation: In the regime $\frac{1}{2n} \leq \mathcal{R}_u^f(k, \beta, \boldsymbol{\epsilon})$, we have the following upper and lower bound on \mathcal{R}^f_u ,

$$\frac{\log(k/\beta)}{\|\boldsymbol{\epsilon}\|_1} \lesssim \mathcal{R}_u^f(k,\beta,\boldsymbol{\epsilon}) \lesssim \frac{\log(k/\beta)}{\|\boldsymbol{\epsilon}\|_1} + \frac{\sqrt{n^2 \mathsf{Var}(e^{-\boldsymbol{\epsilon}}) \wedge n \log(k/\beta)}}{\|\boldsymbol{\epsilon}\|_1}.$$
 (10)

The upper bound is achieved by HPF-UP and HPF-A. Further, if $\operatorname{Var}(e^{-\epsilon}) \lesssim \frac{\log^2(k/\beta)}{n^2}$, then we have

$$\mathcal{R}_{u}^{f}(k,\beta,\epsilon) \simeq \frac{\log(k/\beta)}{\|\epsilon\|_{1}}.$$
(11)

(B) Mean Estimation: In the regime $\frac{1}{2n} \leq \mathcal{R}_u^m(\beta, \epsilon)$, we have the following upper and lower bound on \mathcal{R}_u^m ,

$$\frac{\log(1/\beta)}{\|\boldsymbol{\epsilon}\|_1} \lesssim \mathcal{R}_u^m(\beta, \boldsymbol{\epsilon}) \lesssim \frac{\log(1/\beta)}{\|\boldsymbol{\epsilon}\|_1} + \frac{\sqrt{n^2 \mathsf{Var}(e^{-\boldsymbol{\epsilon}}) \wedge n \log(1/\beta)}}{\|\boldsymbol{\epsilon}\|_1}.$$
 (12)

The upper bound is achieved by HPM-UP and HPM-A. Further, if $\operatorname{Var}(e^{-\epsilon}) \leq \frac{\log^2(1/\beta)}{n^2}$, then we have

$$\mathcal{R}_{u}^{m}(\beta, \boldsymbol{\epsilon}) \simeq \frac{\log(k/\beta)}{\|\boldsymbol{\epsilon}\|_{1}}.$$
(13)

The high privacy assumption, i.e., $\|\epsilon\|_{\infty} \leq 1$, is helpful for dealing with the implicit lower bounds presented in terms of expectation in Theorem C.3. Such assumptions on the privacy level are frequent in the literature, even for homogeneous privacy [30, 41]. The downside to this regime is that it does not explain what occurs if there are some users desiring no privacy $(\epsilon \to \infty)$, i.e., there is some public data. While HPF-UP and HPM-UP can handle such ϵ , the theorem can not. We remark that the assumptions made in Theorem 5.2 might be a shortcoming of the analysis and not the algorithm.

For the homogeneous privacy demand ϵ , one can recover the minimax rate $1/n\epsilon$ for homogeneous DP^2 from the theorem since the variance term is zero.

Before presenting Theorem 5.3, we define another minimax rate closely related to the problem. For some set S, let $\mathcal{P}(S)$ denote the set of probability distributions on S. Similar to the MSE-minimax rates, we define the minimax rate for mean estimation under ϵ -DP from i.i.d. data as

$$\mathcal{E}_{stat}(\boldsymbol{\epsilon}) = \min_{\boldsymbol{M} \in \mathcal{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{P} \in \mathcal{P}(\{0,1\})} \mathbb{E}_{X_1,\dots,X_n \sim \boldsymbol{P}}[|\boldsymbol{M}(X^n) - \boldsymbol{\mu}(\boldsymbol{P})|^2].$$
(14)

We take a moment to comment on $\mathcal{E}_{stat}(\epsilon)$. For *n*-users with no privacy demand, let $\infty = (\infty, \ldots, \infty)$, then note that $\mathcal{E}_{stat}(\infty) \simeq \frac{1}{n}$ as this is the standard minimax rate for mean estimation without privacy constraints (see Wainwright [42] for reference). If there is no privacy demand, then we also have $\mathcal{E}_{u}^{f}(k, \infty), \mathcal{E}_{u}^{m}(\infty) = 0$. In Theorem 5.3, we show MSE-minimax optimality of the proposed element dependence of privacy demand element dependence. of the proposed algorithms for the uncorrelated setting in the regime of privacy demands where ϵ causes a non-trivial increase in the statistical rate \mathcal{E}_{stat} to be greater than $\frac{1}{2n}$. In fact, in this regime, the MSE-minimax rates for empirical mean and frequency estimation are nearly the same as the MSE-minimax rate for statistical mean estimation.

Theorem 5.3 (MSE Bounds for Uncorrelated Setting). For values of ϵ satisfying $\mathcal{E}_{stat}(\epsilon) \geq \frac{1}{2n}$, HPF-U \mathbb{E} and HPM-U \mathbb{E} are minimax optimal. Further,

$$\mathcal{E}_{stat}(\boldsymbol{\epsilon}) \lesssim \mathcal{E}_{u}^{f}(k, \boldsymbol{\epsilon}) \lesssim \log(k)^{2} \mathcal{E}_{stat}(\boldsymbol{\epsilon})$$
(15)

²See Dwork et al. [7] and Vadhan [26] for upper and lower bounds respectively in homogeneous DP for frequency estimation.

for frequency estimation and

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) \simeq \mathcal{E}_{stat}(\boldsymbol{\epsilon}) \tag{16}$$

for mean estimation.

The proof can be found in Appendix J.

6 Extended Discussion

Efficient and approximate versions of HPF and HPM algorithms: The minimizations in the HPF and HPM algorithms (except HPF-A and HPM-A) can be the bottleneck in terms of computation and numerical stability. In order to address these two concerns, one can consider making the upper bound loose by replacing the ℓ_1 -norm with $n\ell_2$ -norm (the latter upper bounds the former). These variants of the algorithm can be efficiently implemented in $O(n \log n)$ time using the algorithm presented by Chaudhuri et al. [32]. Some experiments with this variant of the algorithm are presented in Appendix M.

Free Privacy: An interesting phenomenon in the context of estimation under heterogeneous privacy constraints is the emergence of extra privacy given by the algorithm to some of the users than what is required, contrary to the intuition that invading more privacy should lead to improved performance³. When an HPF or HPM algorithm uses weights w (in either setting), the privacy guarantee for user i is $w_i/||w/\epsilon||_{\infty} \le \epsilon_i$. The equality only holds for every user when the weights are chosen proportional to the privacy parameter, i.e., $w \propto \epsilon$. In the context of minimax optimality, we know that this proportional weighing can be strictly sub-optimal. Thus, there is a phenomenon of free and extra privacy afforded to the users by the algorithm.

7 Conclusion

The problem of mean estimation of univariate data and frequency estimation of categorical data under heterogeneous differential privacy is considered for an adversarial data setting. We formulate two distinct problem settings, one for scenarios where a correlation exists between the dataset and users' privacy demands, and another for scenarios where no such correlation exists. For each problem, in each setting, we consider the minimax rate under mean squared error and PAC error. For each of the eight combinations of problem, setting, and error metric, we demonstrate good upper bounds on the corresponding minimax rates by analysis of an algorithm tailor-made for this purpose. For practical use, we also suggest a heuristic algorithm which experimentally demonstrate superior performance relative to baseline.

A possible line of future work can be on making the upper and lower bounds tighter for some of the minimax rates. Other directions include proving lower bounds for frequency estimation under Heterogeneous Local-DP. Another exciting line of work is to possibly provide privacy to the privacy demand of the users, i.e., solving the problem without Assumption 2.2.

Broader Impacts

This work considers an important problem in the field of Machine Learning and Statistics regarding privacy. Understanding heterogeneous privacy demand can lead to more tailored privacy offerings by services, leading to better user satisfaction. No negative consequences of this work are apparent to the authors.

³This is true in homogeneous privacy in general.

References

- Harry Markowitz. Portfolio selection. The Journal of Finance, 7(1):77-91, 1952. ISSN 00221082, 15406261. URL http://www.jstor.org/stable/2975974.
- [2] Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-efficient learning of deep networks from decentralized data. In *Artificial intelligence and statistics*, pages 1273–1282. PMLR, 2017.
- [3] Kornelia Batko and Andrzej Ślęzak. The use of big data analytics in healthcare. *Journal of big Data*, 9(1):3, 2022.
- [4] Tong Yang, Jie Jiang, Peng Liu, Qun Huang, Junzhi Gong, Yang Zhou, Rui Miao, Xiaoming Li, and Steve Uhlig. Elastic sketch: Adaptive and fast network-wide measurements. In *Proceedings of the 2018 Conference of the ACM Special Interest Group on Data Communication*, pages 561–575, 2018.
- [5] Koustuv Saha, Yozen Liu, Nicholas Vincent, Farhan Asif Chowdhury, Leonardo Neves, Neil Shah, and Maarten W Bos. Advertiming matters: Examining user ad consumption for effective ad allocations on social media. In *Proceedings of the 2021 CHI Conference on Human Factors* in Computing Systems, pages 1–18, 2021.
- [6] Cynthia Dwork, Krishnaram Kenthapadi, Frank McSherry, Ilya Mironov, and Moni Naor. Our data, ourselves: Privacy via distributed noise generation. In *Annual international conference on* the theory and applications of cryptographic techniques. Springer, 2006.
- [7] Cynthia Dwork, Frank McSherry, Kobbi Nissim, and Adam Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of cryptography conference*. Springer, 2006.
- [8] Úlfar Erlingsson, Vasyl Pihur, and Aleksandra Korolova. Rappor: Randomized aggregatable privacy-preserving ordinal response. In *Proceedings of the 21st ACM Conference on Computer* and Communications Security, Scottsdale, Arizona, 2014.
- [9] Apple. Learning with privacy at scale differential. 2017. URL https://api. semanticscholar.org/CorpusID:43986173.
- [10] John M. Abowd. The u.s. census bureau adopts differential privacy. In Proceedings of the 24th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining. Association for Computing Machinery, 2018.
- [11] Teng Wang, Xuefeng Zhang, Jingyu Feng, and Xinyu Yang. A comprehensive survey on local differential privacy toward data statistics and analysis. *Sensors*, 2020.
- [12] Mark S. Ackerman, Lorrie Faith Cranor, and Joseph Reagle. Privacy in e-commerce: examining user scenarios and privacy preferences. In *Proceedings of the 1st ACM Conference on Electronic Commerce*, EC '99, page 1–8. Association for Computing Machinery, 1999.
- [13] Hong Ping Li, Haibo Hu, and Jianliang Xu. Nearby friend alert: Location anonymity in mobile geosocial networks. *IEEE Pervasive Computing*, 12(4):62–70, 2012.
- [14] Shiva Prasad Kasiviswanathan, Homin K Lee, Kobbi Nissim, Sofya Raskhodnikova, and Adam Smith. What can we learn privately? *SIAM Journal on Computing*, 2011.
- [15] Peter Kairouz, Monica Ribero Diaz, Keith Rush, and Abhradeep Thakurta. (Nearly) dimension independent private erm with adagrad rates via publicly estimated subspaces. In *Conference on Learning Theory*. PMLR, 2021.
- [16] Tianhao Wang, Jeremiah Blocki, Ninghui Li, and Somesh Jha. Locally differentially private protocols for frequency estimation. In *USENIX Security Symposium*, 2017.
- [17] Raef Bassily, Albert Cheu, Shay Moran, Aleksandar Nikolov, Jonathan Ullman, and Steven Wu. Private query release assisted by public data. In *International Conference on Machine Learning*. PMLR, 2020.

- [18] Raef Bassily, Shay Moran, and Anupama Nandi. Learning from mixtures of private and public populations. Advances in Neural Information Processing Systems, 2020.
- [19] Terrance Liu, Giuseppe Vietri, Thomas Steinke, Jonathan Ullman, and Steven Wu. Leveraging public data for practical private query release. In *International Conference on Machine Learning*. PMLR, 2021.
- [20] Noga Alon, Raef Bassily, and Shay Moran. Limits of private learning with access to public data. *Advances in neural information processing systems*, 2019.
- [21] Anupama Nandi and Raef Bassily. Privately answering classification queries in the agnostic pac model. In *Algorithmic Learning Theory*. PMLR, 2020.
- [22] Ehsan Amid, Arun Ganesh, Rajiv Mathews, Swaroop Ramaswamy, Shuang Song, Thomas Steinke, Vinith M Suriyakumar, Om Thakkar, and Abhradeep Thakurta. Public data-assisted mirror descent for private model training. In *International Conference on Machine Learning*. PMLR, 2022.
- [23] Di Wang, Huangyu Zhang, Marco Gaboardi, and Jinhui Xu. Estimating smooth glm in noninteractive local differential privacy model with public unlabeled data. In *Algorithmic Learning Theory*. PMLR, 2021.
- [24] Alex Bie, Gautam Kamath, and Vikrant Singhal. Private estimation with public data. In *Advances in Neural Information Processing Systems*, 2022.
- [25] Sourav Biswas, Yihe Dong, Gautam Kamath, and Jonathan Ullman. Coinpress: Practical private mean and covariance estimation. Advances in Neural Information Processing Systems, 33: 14475–14485, 2020.
- [26] Salil Vadhan. The complexity of differential privacy. Tutorials on the Foundations of Cryptography: Dedicated to Oded Goldreich, pages 347–450, 2017.
- [27] Mohammad Alaggan, Sébastien Gambs, and Anne-Marie Kermarrec. Heterogeneous differential privacy. *Journal of Privacy and Confidentiality*, 2017.
- [28] Zach Jorgensen, Ting Yu, and Graham Cormode. Conservative or liberal? personalized differential privacy. In 2015 IEEE 31St international conference on data engineering, 2015.
- [29] Cecilia Ferrando, Jennifer Gillenwater, and Alex Kulesza. Combining public and private data. In NeurIPS 2021 Workshop Privacy in Machine Learning, 2021.
- [30] Alireza Fallah, Ali Makhdoumi, Azarakhsh Malekian, and Asuman Ozdaglar. Optimal and differentially private data acquisition: Central and local mechanisms. *arXiv preprint arXiv:2201.03968*, 2022.
- [31] Syomantak Chaudhuri and Thomas Courtade. Mean estimation under heterogeneous privacy: Some privacy can be free. In 2023 IEEE International Symposium on Information Theory (ISIT), 2023.
- [32] Syomantak Chaudhuri, Konstantin Miagkov, and Thomas A Courtade. Mean estimation under heterogeneous privacy demands. arXiv preprint arXiv:2310.13137, 2023.
- [33] Rachel Cummings, Hadi Elzayn, Emmanouil Pountourakis, Vasilis Gkatzelis, and Juba Ziani. Optimal data acquisition with privacy-aware agents. In 2023 IEEE Conference on Secure and Trustworthy Machine Learning (SaTML), pages 210–224. IEEE, 2023.
- [34] Ian Waudby-Smith, Steven Wu, and Aaditya Ramdas. Nonparametric extensions of randomized response for private confidence sets. In Andreas Krause, Emma Brunskill, Kyunghyun Cho, Barbara Engelhardt, Sivan Sabato, and Jonathan Scarlett, editors, *Proceedings of the 40th International Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pages 36748–36789. PMLR, 23–29 Jul 2023.
- [35] Clément L Canonne and Yucheng Sun. Private distribution testing with heterogeneous constraints: Your epsilon might not be mine. *arXiv preprint arXiv:2309.06068*, 2023.

- [36] Ameya Anjarlekar, Rasoul Etesami, and R Srikant. Striking a balance: An optimal mechanism design for heterogenous differentially private data acquisition for logistic regression. arXiv preprint arXiv:2309.10340, 2023.
- [37] Justin Kang, Kannan Ramchandran, and Ramtin Pedarsani. The fair value of data under heterogeneous privacy constraints in federated learning. In *International Workshop on Federated Learning in the Age of Foundation Models in Conjunction with NeurIPS 2023*, 2023.
- [38] Krishna Acharya, Franziska Boenisch, Rakshit Naidu, and Juba Ziani. Personalized differential privacy for ridge regression. *arXiv preprint arXiv:2401.17127*, 2024.
- [39] Arpita Ghosh and Aaron Roth. Selling privacy at auction. In *Proceedings of the 12th ACM* conference on Electronic commerce, pages 199–208, 2011.
- [40] Steven Diamond and Stephen Boyd. CVXPY: A Python-embedded modeling language for convex optimization. *Journal of Machine Learning Research*, 2016.
- [41] John Duchi, Michael Jordan, and Martin Wainwright. Local privacy and minimax bounds: Sharp rates for probability estimation. *Adv. Neural Inform. Process. Syst*, 2013.
- [42] Martin J Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*, volume 48. Cambridge university press, 2019.
- [43] Peter Kairouz, Keith Bonawitz, and Daniel Ramage. Discrete distribution estimation under local privacy. In *International Conference on Machine Learning*, pages 2436–2444. PMLR, 2016.
- [44] CA State Controller. University of california salary data, 2022. URL https://publicpay. ca.gov/Reports/RawExport.aspx. Accessed: 2024-01-22.
- [45] U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, and National Cancer Institute. U.s. cancer statistics data visualizations tool, based on 2022 submission data (1999-2020), 2020. URL https://gis.cdc.gov/Cancer/USCS/#/ AtAGlance/. Accessed: 2024-01-22.
- [46] David Wajc. Negative association: definition, properties, and applications. 2017.
- [47] Wei-Ning Chen, Peter Kairouz, and Ayfer Ozgur. Breaking the communication-privacy-accuracy trilemma. *Advances in Neural Information Processing Systems*, 33:3312–3324, 2020.
- [48] Michael Kearns and Lawrence Saul. Large deviation methods for approximate probabilistic inference. arXiv preprint arXiv:1301.7392, 2013.
- [49] Daniel Berend and Aryeh Kontorovich. On the concentration of the missing mass. *Electronic Communications in Probability*, 18(none):1 7, 2013. doi: 10.1214/ECP.v18-2359.
- [50] T-H Hubert Chan, Elaine Shi, and Dawn Song. Private and continual release of statistics. ACM *Transactions on Information and System Security (TISSEC)*, 14(3):1–24, 2011.

A Mean Estimation

Definition A.1 (Minimax Rates of Mean Estimation). (A) **Correlated Setting:** The PAC-minimax rate is given by

$$\mathcal{R}_{c}^{m}(\beta, \epsilon) = \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [0,1]^{n}} \min_{\mathbb{P}\{|M(D) - \mu(D)| > \alpha\} \le \beta} \alpha ,$$
(17)

where the probability is taken over the randomness in the map M. The MSE-minimax rate is given by

$$\mathcal{E}_{c}^{m}(\boldsymbol{\epsilon}) = \min_{\boldsymbol{M} \in \mathcal{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{D} \in [0,1]^{n}} \mathbb{E}[|\boldsymbol{M}(\boldsymbol{D}) - \boldsymbol{\mu}(\boldsymbol{D})|^{2}],$$
(18)

where the expectation is taken over the randomness in the map M. (B) **Uncorrelated Setting:** The PAC-minimax rate is given by

$$\mathcal{R}_{u}^{m}(\beta, \epsilon) = \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [0,1]^{n}} \min_{\mathbb{P}\{|M(D_{\sigma}) - \mu(D)| > \alpha\} \le \beta} \alpha,$$
(19)

where the probability is taken over the randomness in the map M and the uniform permutation σ of the dataset D. The MSE-minimax rate is given by

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) = \min_{\boldsymbol{M} \in \mathcal{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{D} \in [0,1]^{n}} \mathbb{E}[|\boldsymbol{M}(\boldsymbol{D}_{\sigma}) - \boldsymbol{\mu}(\boldsymbol{D})|^{2}],$$
(20)

where the expectation is taken over the randomness in the map M and the uniform permutation σ of the dataset D.

The family of algorithms HPM is presented in Algorithm 2, along with the choices of weights in Table 3.

Table 3: Weights for Algorithm 2 for different settings.

Algorithm 2 Heterogeneously Private Mean	ALGORITHM	WEIGHTS w		
(HPM)	HPM-C₽	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_C(\boldsymbol{w},1,\beta,\boldsymbol{\epsilon})$		
Input: weights $\boldsymbol{w} \in \Delta_n, X \in [0,1]^n$	HPM-CE	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n}r_C(\boldsymbol{w},e,1,\boldsymbol{\epsilon})$		
Sample $N \sim \text{Laplace}(\ \frac{\boldsymbol{w}}{\boldsymbol{\epsilon}}\ _{\infty})$	HPM-U₽	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},1,\beta,\boldsymbol{\epsilon})$		
return $\max\{\min\{N + \sum_{i=1}^{n} w_i X_i, 1\}, 0\}$	HPM-UE	$\operatorname{argmin}_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},e,1,\boldsymbol{\epsilon})$		
	HPM-A	$\frac{1 - e^{-\epsilon}}{\ 1 - e^{-\epsilon}\ _1}$		

B Experiment Details

Uniformly enforce ϵ_1 -**DP** (**UNI**): This approach offers the highest privacy level ϵ_1 to all the datapoints and uses the known minimax estimator for homogeneous ϵ_1 -DP mean and frequency estimation, depending on the problem. UNI can be arbitrarily worse than the optimal algorithm – consider a dataset with a single high and several low privacy datapoints⁴.

Proportional Weighing (Prop): This approach uses the weights $w \propto \epsilon$. This weighing minimizes the amount of Laplace noise required over all possible weights, but it comes at the cost of a higher bias. In the case that ϵ have a high variance, this weighing strategy can perform rather poorly. For example, in the case that there is just one user requiring no privacy ($\epsilon \rightarrow \infty$), this strategy would disregard the data of all other users.

Sampling Mechanism (SM): This mechanism was proposed by Jorgensen et al. [28]. Let $t = ||\epsilon||_{\infty}$. Then sample *i*-th datapoint independently with probability $(e^{\epsilon_i} - 1)/(e^t - 1)$. On the sub-sampled dataset, we can use any homogeneous *t*-DP algorithm and this will be ϵ -DP. A shortcoming of this mechanism is of just one user requires no privacy, the SM algorithm disregards the rest of the dataset.

Local Differential Private Frequency (LDP): A simple Local-DP approach to heterogeneous DP is to send the user *i*'s data to the server via an ϵ_i -DP channel.

⁴High privacy corresponds to a low value of ϵ .

- *Mean estimation:* the LDP channels adds Laplace(¹/_{εi}) noise to the user *i*'s data. *Frequency estimation:* we use *k*-RAPPOR [8, 43] to send the datapoints to the central server,

which essentially does randomized response on one-got encoding of user data. The exact scheme used by us for combining these estimates from different users at the server in outlined in Appendix L. While this is not a completely fair comparison with other algorithms since Local-DP is typically more noisy than Central-DP, it is still insightful to see the performance difference. In fact, our implementation of this Local-DP mechanism under heterogeneous privacy is on par with some of the baseline Central-DP techniques for frequency estimation.

For PAC error evaluation, we set $\beta = 0.05$ for all experiments for obtaining the weights for the PAC variants of HPF and HPM algorithms. We run the experiments on both synthetic and real datasets. Keeping the dataset and privacy demand fixed, several trials of the algorithms are performed for each experiment. For the uncorrelated setting, in each trial, the dataset is randomly permuted.

For each experiment on the PAC variants, we record the empirical ℓ_{∞} error of the algorithms for each trial and report the 95-th quantile error. For each experiment on the MSE variant, we report the mean squared l_{∞} -error.

B.1 Frequency Estimation under Correlated Setting

For the correlated setting, we sub-sample the University of California (UC) salary data for 2022 [44] consisting of the salaries of 50K employees in the UC system. We partition the salaries into 12 bins and assign less privacy demand to the people in the more central bins. In particular, a user that belongs to bin i has a privacy demand randomly sampled as $\log \epsilon \sim -|i - 6.5| + \text{Uniform}[-3,3]$. We sample the privacy demands once and keep it fixed, while running several trials. Results of the experiments are outlined in Table 4. Our proposed algorithms HPF-CP and HPF-CE outperforms the other algorithms, while HPF-A also performs competitively. The empirical error distributions are plotted in Figure 1(a) to illustrate the performance of different algorithms. The Prop algorithm exhibits a steep curve since this method adds the minimum amount of Laplace noise, at the cost of higher bias.

We also consider two synthetic datasets with n = 10000; the number of bins k is set as 5 and 20. The privacy demand is randomly sampled, with different distributions for users in different bins. UNI performs poorly since it adds significantly more noise, but LDP's performance is impressive since Local-DP typically performs significantly worse than Central-DP. In Table 4, in the synthetic datasets, HPF-A outperforms HPF-C \mathbb{P} and HPF-C \mathbb{E} .

B.2 Frequency Estimation under Uncorrelated Setting

For uncorrelated setting, we consider the number of new cancers in the state of Colorado in 2020 [45] containing data of about 17K surgeries with ten different types of cancers. We sample log $\epsilon \sim \text{Uniform}[-5, 5]$ and keep ϵ fixed. For each trial, the dataset, mapping each user to a type of cancer surgery, is randomly permuted. We perform $O(n \log n)$ trials to get meaningful results due to the random permutations. The plot for empirical cdf of the errors is shown in Figure 1(b). Our proposed algorithms HPF-A, HPF-UE, and HPF-UE perform better than the other baseline algorithms.

In a similar manner as the correlated setting, we also test the algorithms under synthetic datasets. From Table 4, it can be seen that while HPF-A performs superior to the other algorithms, HPF-UP and HPF-UE also enjoy good performance.

B.3 Mean Estimation under Correlated Setting:

For the correlated setting, we consider the UC Dataset and use the the method described in frequency estimation to get a correlated data and privacy demand. Results of the experiments are outlined in Table 5. We also consider a synthetic datasets with n = 10000 and use a method similar to the frequency estimation experiments to obtain correlated data and privacy demand.

B.4 Mean Estimation under Uncorrelated Setting:

For uncorrelated setting, we test the algorithms under a synthetic dataset. Overall, HPM-A performs superior as comapred to the other algorithms.

Table 4: We present the performance of various algorithms under different datasets and settings. The first column specifies the dataset, whether it is the correlated or the uncorrelated setting, and whether the reported numbers are empirical 95-th quantile error (\mathbb{P}) or mean squared error (\mathbb{E}) under ℓ_{∞} norm. Datasets of form (n, k) are synthetic datasets with n users and k bins. The variant of the HPF algorithm used is given by the regime and error metric for the dataset listed in the first column.

DATASET, REGIME, ERROR METRIC	HPF	HPF-A	PROP	UNI	SM	LDP
UC SALARY, C, ℙ	0.118	0.174	0.273	1.000	0.459	0.420
UC SALARY, C, E	0.011	0.030	0.074	0.396	0.180	0.087
(10000, 5), C, ℙ	0.105	0.083	0.202	1.000	0.924	0.253
(10000, 5), C, E	0.012	0.007	0.041	0.834	0.779	0.027
(10000, 20), C, ℙ	0.020	0.003	0.090	0.233	1.000	0.125
(10000, 20), C, E	3×10^{-4}	$ 7 \times 10^{-6}$	0.008	0.022	0.779	0.008
Cancer Types, U, \mathbb{P}	0.009	0.005	0.014	0.091	0.382	0.012
Cancer Types, U, $\mathbb E$	5×10^{-5}	1×10^{-5}	8×10^{-5}	0.003	0.036	7×10^{-5}
(10000, 5), U, ℙ	0.012	0.008	0.023	0.370	0.798	0.0152
(10000, 5), U, E	1×10^{-4}	3×10^{-5}	2×10^{-4}	0.043	0.198	1×10^{-4}
(10000, 20), U, ℙ	0.013	0.007	0.020	0.482	0.348	0.015
(10000, 20), U, E	1×10^{-4}	2 ×10 ⁻⁵	2×10^{-4}	0.095	0.028	1×10^{-4}

Table 5: We present the performance of various algorithms under different datasets and settings. The first column specifies the dataset, whether it is the correlated or the uncorrelated setting, and whether the reported numbers are empirical 95-th quantile error (\mathbb{P}) or mean squared error (\mathbb{E}) under ℓ_{∞} norm. The variant of the HPM algorithm used is given by the regime and error metric for the dataset listed in the first column.

DATASET, REGIME, ERROR METRIC	HPM	HPM-A	PROP	UNI	SM	LDP
UC SALARY, C, \mathbb{P}	0.064	0.005	0.005	0.066	0.005	1.000
UC SALARY, C, $\mathbb E$	2×10^{-5}	2×10^{-5}	2 ×10 ⁻⁵	0.001	2 ×10 ⁻⁵	0.832
10000, C, P	0.008	0.009	0.009	0.027	0.052	0.085
10000, C, E	4×10^{-5}	7×10^{-5}	8×10^{-4}	2×10^{-4}	2 ×10 ⁻⁵	0.002
10000, U, P	0.008	0.001	0.004	0.027	0.110	0.086
10000, U, E	4×10^{-5}	3×10^{-7}	5×10^{-6}	1×10^{-4}	0.003	0.002

C Upper and Lower Bounds

C.1 Upper Bounds

Recall the functions r_C and r_U from (6) and (7) respectively. We denote their corresponding minimum values by

$$R_C(k,\beta,\epsilon) = \left(\min_{\boldsymbol{w}\in\Delta_n} r_C(\boldsymbol{w},k,\beta,\epsilon)\right) \wedge 1,$$
(21)

and

$$R_U(k,\beta,\epsilon) = \left(\min_{\boldsymbol{w}\in\Delta_n} r_U(\boldsymbol{w},k,\beta,\epsilon)\right) \wedge 1.$$
(22)

These functions are relevant for expressing the PAC and MSE errors for either problems in either settings.

For a choice of weights $w \in \Delta_n$ in Algorithm 1 or Algorithm 2, the term $||w - 1/n||_1$ in r_C and r_U are errors due to having weights different from 1/n. The $||\frac{w}{\epsilon}||_{\infty}$ term accounts for noise needed to satisfy the privacy demand. The algorithms minimizing their respective errors over the choices of weights w.

Theorem C.1 provides upper bounds for the PAC-minimax rates in the two settings and the two problems; its proof can be found in Appendix E.

Theorem C.1 (PAC Upper Bounds).

(A) Correlated Setting: For frequency estimation

$$\mathcal{R}_{c}^{f}(k,\beta,\epsilon) \lesssim R_{C}(k,\beta,\epsilon), \tag{23}$$

and for mean estimation

$$\mathcal{R}_{c}^{m}(\beta, \epsilon) \lesssim R_{C}(1, \beta, \epsilon).$$
(24)
(B) Uncorrelated Setting: For frequency estimation

$$\mathcal{R}_{u}^{f}(k,\beta,\epsilon) \lesssim R_{U}(k,\beta,\epsilon),$$
(25)

and for mean estimation

$$\mathcal{R}^m_u(\beta, \epsilon) \lesssim R_U(1, \beta, \epsilon). \tag{26}$$

Since we are considering PAC error under ℓ_{∞} -norm for frequency estimation, the $\log(k/\beta)$ captures the effect of union bound. Simply substituting k = 1 recovers the guarantee for mean estimation.

Theorem C.2 provides an upper bound on the MSE-minimax rates for the two settings for both frequency and mean estimation, its proof can be found in Appendix F.

Theorem C.2 (MSE Upper Bounds).

(A) Correlated Setting: For frequency estimation

$$\mathcal{E}_c^f(k,\beta,\epsilon) \lesssim R_C(k,1,\epsilon)^2,\tag{27}$$

and for mean estimation

$$\mathcal{E}_{c}^{m}(\beta, \boldsymbol{\epsilon}) \lesssim R_{C}(e, 1, \boldsymbol{\epsilon})^{2}.$$
(28)

$$\mathcal{E}_{u}^{f}(k,\epsilon) \lesssim R_{U}(k,1,\epsilon)^{2},$$
(29)

and for mean estimation

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) \lesssim R_{U}(e, 1, \boldsymbol{\epsilon})^{2}.$$
(30)

We mention that for frequency estimation in Theorem C.2, the MSE upper bound has $\log(k)$ term instead of $\log(k/\beta)$ so we just set $\beta = 1$ in R_C and R_U for brevity. Similarly, for mean estimation, there is no $\log(k/\beta)$ term in our upper bounds; hence, we substitute the constant $\beta \leftarrow 1, k \leftarrow e$ in the R_C and R_U functions for brevity. We also assume $k \ge 2$ for frequency estimation to make simplifications like $\log(2k) \le \log(k)$ in the proof. Theorem C.2 is obtained by some concentration inequalities and the tail-sum identity for expectation, using the PAC bounds in Theorem C.1.

As one would expect, the upper bounds in the correlated setting is always greater than the corresponding one in uncorrelated setting – in both mean and frequency estimation under both PAC and MSE bounds.

For comparison, in frequency estimation, consider the straw-man approach of assigning every user equal weight⁵ and adding i.i.d. Laplace $(\frac{2}{n\epsilon_1})$ noise to each bin., leading to the PAC-error of the order $\frac{\log(k/\beta)}{n\epsilon_1}$ and MSE-error of the order $\frac{\log(k)^2}{n^2\epsilon_1^2}$ (in both settings). This method has a clear drawback in case there is one user demanding an extremely high privacy ($\epsilon_1 \rightarrow 0$) and the rest of the users requiring no privacy. It is simply better to ignore the first user and do non-private frequency calculation for the rest of the users; in correlated setting, this leads to a PAC-error of at most $\frac{1}{n}$ from Theorem C.1, and MSE-error of $\frac{1}{n^2}$ from Theorem C.2. In the uncorrelated setting, it leads to PAC-error of at most $\sqrt{\frac{\log(k/\beta)}{n}} \wedge \frac{1}{n}$ and MSE-error of $\frac{1}{n}$. A similar case can be made for mean estimation.

⁵This is the UNI algorithm defined in Section 4.

C.2 Lower Bounds

For frequency estimation, the trivial output of 1/2 achieves an ℓ_{∞} error of $\frac{1}{2}$. Therefore, we have $\mathcal{R}_{c}^{f}(k,\beta,\epsilon), \mathcal{R}_{u}^{f}(k,\beta,\epsilon) \leq \frac{1}{2}$. Similarly, for mean estimation we have $\mathcal{R}_{c}^{m}(\beta,\epsilon), \mathcal{R}_{u}^{m}(\beta,\epsilon) \leq \frac{1}{2}$.

Our lower bounds for PAC rates follow from a packing argument and are presented in Theorem C.3. We remind the readers that ϵ is assumed to be in non-decreasing order.

When the minimax rates are strictly less than half, we show that the minimax rate must satisfy the conditions in Theorem C.3. The proof can be found in Appendix G.

Theorem C.3 (Implicit PAC Lower Bounds). We have $\mathcal{R}_c^f(k, \beta, \epsilon), \mathcal{R}_u^f(k, \beta, \epsilon), \mathcal{R}_c^m(\beta, \epsilon), \mathcal{R}_u^m(\beta, \epsilon) \leq \frac{1}{2}$. (A) Correlated Setting: if $\mathcal{R}_c^f(k, \beta, \epsilon) < \frac{1}{2}$, then it also satisfies

$$\sum_{i=1}^{2n\mathcal{R}_c^f(k,\beta,\epsilon)]} \epsilon_i \gtrsim \log\left(\frac{k(1-\beta)}{\beta}\right).$$
(31)

Similarly, if $\mathcal{R}_c^m(\beta, \epsilon) < \frac{1}{2}$, then it also satisfies

$$\sum_{i=1}^{\lceil 2n\mathcal{R}_c^m(\beta,\epsilon)\rceil} \epsilon_i \gtrsim \log\left(\frac{(1-\beta)}{\beta}\right).$$
(32)

(B) Uncorrelated Setting:

Let $\{Z_i\}_{i=1}^n$ be sampled without replacement from the entries of ϵ . If $\mathcal{R}_u^f(k,\beta,\epsilon) < \frac{1}{2}$, then it also satisfies

$$\mathbb{E}\left[\exp\left\{-\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{f}(k,\beta,\epsilon)\rceil}Z_{i}\right\}\right] \leq \frac{k\beta}{k-1},$$
(33)

$$\mathbb{E}\left[\exp\left\{\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{f}(k,\beta,\epsilon)\rceil}Z_{i}\right\}\right] \ge k(1-\beta).$$
(34)

Similarly, if $\mathcal{R}_{u}^{m}(\beta, \epsilon) < \frac{1}{2}$, then it also satisfies

$$\mathbb{E}\left[\exp\left\{-\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{m}(\beta,\epsilon)\rceil}Z_{i}\right\}\right] \leq \beta,$$
(35)

$$\mathbb{E}\left[\exp\left\{\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{m}(\beta,\epsilon)\rceil}Z_{i}\right\}\right] \ge (1-\beta).$$
(36)

While the conditions presented in Theorem C.3 are implicit, it should be noted that the conditions imply a lower limit to the minimax rates.

We also present MSE lower bounds in the uncorrelated setting in Theorem C.4. In order to present the lower bound, we define another minimax rate closely related to the problem. For some set S, let $\mathcal{P}(S)$ denote the set of probability distributions on S. Similar to the MSE-minimax rates, we define the minimax rate for mean estimation under ϵ -DP from i.i.d. data as

$$\mathcal{E}_{stat}(\boldsymbol{\epsilon}) = \min_{\boldsymbol{M} \in \boldsymbol{M}_{\boldsymbol{\epsilon}}} \max_{\boldsymbol{P} \in \mathcal{P}(\{0,1\})} \mathbb{E}_{X_1,\dots,X_n \sim \boldsymbol{P}}[|\boldsymbol{M}(\boldsymbol{X}^n) - \boldsymbol{\mu}(\boldsymbol{P})|^2].$$
(37)

Theorem C.4 (MSE Lower Bounds). We have $\mathcal{E}_u^f(k, \epsilon), \mathcal{E}_u^m(\epsilon) \leq \frac{1}{4}$. Both $\mathcal{E}_u^f(k, \epsilon), \mathcal{E}_u^m(\epsilon)$ satisfy

$$\mathcal{E}_{u}^{f}(k,\boldsymbol{\epsilon}), \mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) \geq \left(\sqrt{\mathcal{E}_{stat}(\boldsymbol{\epsilon})} - \sqrt{\frac{1}{4n}}\right)_{+}^{2}.$$
(38)

In Theorem C.4, we use the notation $(x)_+$ to refer to the function $\max\{x, 0\}$. We interpret this lower bound in Section 5.2 and compare it with the upper bound. We mention that we do not present any lower bounds for the MSE-minimax rates \mathcal{E}_c^f and \mathcal{E}_c^m .

D Proof of Lemma 3.1

Lemma (3.1). The proposed family of algorithms HPF and HPM satisfy the ϵ -DP constraint defined in Definition 2.1.

Proof. We present the proof for frequency estimation (HPF); similar steps can be used to show ϵ -DP property for HPM. In this proof, we use the notation $\theta_{w}(X) \in \Delta_{k}$ to denote the vector where $\theta_{w}(X)_{i} = \sum_{j} w_{j} \mathbb{I}\{X_{j} = i\}$. Note that the algorithms compute $\theta_{w}(X)$ and adds i.i.d. Laplace $(2\|w/\epsilon\|_{\infty})$ noise in each component. We have

$$\frac{p\{M(X) = a\}}{p\{M(X'^{i}) = a\}} = \prod_{j=1}^{k} \exp\left(\frac{|\theta_{w}(X)_{j} - a_{j}| - |\theta_{w}(X'^{i})_{j} - a_{j}|}{2\|w/\epsilon\|_{\infty}}\right)$$
(39)

$$\leq \exp\left(\sum_{j=1}^{k} \frac{|\theta_{\boldsymbol{w}}(X)_j - \theta_{\boldsymbol{w}}(X'^i)_j|}{2\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}}\right)$$
(40)

$$\leq \exp \frac{w_i}{\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}} \leq \exp \epsilon_i,\tag{41}$$

where we used $\sum_{j} |\theta_{w}(X)_{j} - \theta_{w}(X'^{i})_{j}| \leq 2w_{i}$ since X'^{i} differs from X only in the *i*-th coordinate.

E Proof of Theorem C.1

Theorem (C.1).

(A) Correlated Setting: For frequency estimation

$$\mathcal{R}_c^f(k,\beta,\epsilon) \lesssim R_C(k,\beta,\epsilon),\tag{42}$$

and for mean estimation

$$\mathcal{R}_c^m(\beta, \epsilon) \lesssim R_C(1, \beta, \epsilon). \tag{43}$$

(B) Uncorrelated Setting: For frequency estimation

$$\mathcal{R}_{u}^{f}(k,\beta,\epsilon) \lesssim R_{U}(k,\beta,\epsilon), \tag{44}$$

and for mean estimation

$$\mathcal{R}_{u}^{m}(\beta, \epsilon) \lesssim R_{U}(1, \beta, \epsilon).$$
(45)

Proof. (A) We consider frequency estimation. Substituting k = 1 recovers the guarantees for mean estimation.

Suppose that with probability greater than $1 - \beta$, the algorithm is within α of the true frequency. Let $Q \in \{0,1\}^{k \times n}$ such that $Q(i,j) = \mathbb{I}\{D_j = i\}$. Further, let $N = (N_1, \ldots, N_k)$ where the components are i.i.d. Laplace $(2 \| \frac{w^*}{\epsilon} \|_{\infty})$. Then we have,

 \leq

$$\mathbb{P}\{\|Q\boldsymbol{w} + \boldsymbol{N} - \boldsymbol{\mu}(D)\|_{\infty} > \alpha\} = \mathbb{P}\{\|Q(\boldsymbol{w} - \mathbf{1}/n) + \boldsymbol{N}\|_{\infty} > \alpha\}$$
(46)

$$\mathbb{P}\{\|Q(\boldsymbol{w}-\boldsymbol{1}/n)\|_{\infty} > \alpha/2\} + \mathbb{P}\{\|\boldsymbol{N}\|_{\infty} > \alpha/2\}$$
(47)

$$\leq \mathbb{I}\{\alpha/2 \leq \|\boldsymbol{w} - \boldsymbol{1}/n\|_{\mathsf{TV}}\} + k \exp(-\frac{\alpha}{4\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}}), \qquad (48)$$

where we used the fact that $\|Q(w - 1/n)\|_{\infty} \leq \|w - 1/n\|_{\text{TV}}$. To see this, note that the *a*-th component of Q(w - 1/n) is given by $\sum_{i} Q(a, i)(w_i - \frac{1}{n})$, which is less than $\|w - 1/n\|_{\text{TV}}$ almost surely by straightforward definition of TV-distance. The second term is obtained by union bound on laplace tail probabilities.

To get a condition on α , it is sufficient to upper bound the first term by 0 and second term by β in (48). Thus, since α satisfies both conditions, we have $\alpha^2 \gtrsim \max\{\|\boldsymbol{w} - \mathbf{1}/n\|_1^2, \log^2(k/\beta)\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2\} \simeq \|\boldsymbol{w} - \mathbf{1}/n\|_1^2 + \log^2(k/\beta)\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2$. Taking minimum over α satisfying the $(1-\beta)$ quantile condition in the definition of PAC minimax rate implies $\alpha^2 \simeq \|\boldsymbol{w} - \mathbf{1}/n\|_1^2 + \log^2(k/\beta)\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2$. Optimizing over the weights \boldsymbol{w} and due to the projection to [0, 1] in the algorithm, we have $\mathcal{R}_c^f \lesssim R_C(k, \beta, \boldsymbol{\epsilon})$.

(B) We consider frequency estimation. Substituting k = 1 recovers the guarantees for mean estimation. Let $Q_{\sigma} \in \{0,1\}^{k \times n}$ such that $Q_{\sigma}(i,j) = \mathbb{I}\{D_{\sigma(j)} = i\}$. Thus, we have

$$\mathbb{P}\{\|Q_{\sigma}\boldsymbol{w} + \boldsymbol{N} - \boldsymbol{\mu}(D)\|_{\infty} > \alpha\} = \mathbb{P}\{\|Q_{\sigma}(\boldsymbol{w} - \mathbf{1}/n) + \boldsymbol{N}\|_{\infty} > \alpha\}$$

$$\leq \mathbb{P}\{\|Q_{\sigma}(\boldsymbol{w} - \mathbf{1}/n)\|_{\infty} > \alpha/2\} + \mathbb{P}\{\|\boldsymbol{N}\|_{\infty} > \alpha/2\}$$
(49)
(50)

= 1 (1 + 1)

We focus on the $\mathbb{P}\{\|Q_{\sigma}(w-1/n)\|_{\infty} > \alpha/2\}$ term. By previous part, it can be at most $\|w-1/n\|_{\mathsf{TV}}$ (almost surely). However, we also have another bound using the fact that the data is uniformly permuted. We shall now prove

$$\mathbb{P}\{\left|\sum_{i} Q_{\sigma}(a,i)(w_{i}-\frac{1}{n})\right| > \alpha/2\} \le 2\exp\left(-\frac{\alpha^{2}}{2\|\boldsymbol{w}\|_{2}^{2}}\right).$$
(51)

Since we restrict our attention to the *a*-th component in (51), we shall denote the vector $Q_{\sigma}(a, \cdot)$ by $z_{\sigma} \in \{0, 1\}^n$. We start by noting

$$\sum_{i} (w_i - \frac{1}{n}) z_{\sigma(i)} = \sum_{i} (w_i - \frac{1}{n}) (z_{\sigma(i)} - c) \text{ for any constant } c$$
(52)

$$=\sum_{i}(w_{i}-\frac{1}{n})(z_{\sigma(i)}-\mu) \text{ by setting } c=\mu=\sum_{i}z_{i}/n$$
(53)

$$=\sum_{i}w_{i}(z_{\sigma(i)}-\mu).$$
(54)

Define the random variables Y_1, \ldots, Y_n sampled without replacement from $z - \mu \mathbf{1}$. Note that the Y(s) are 'permutation' random variables and they are zero-mean. Thus, they satisfy Negative Association and we have,

=

$$\mathbb{P}\{\sum_{i} w_{i} Y_{i} > t\} = \mathbb{P}\{\exp(\lambda \sum_{i} w_{i} Y_{i}) > \exp(\lambda t)\}$$
(55)

$$\leq \frac{\mathbb{E}[\prod_{i} \exp(\lambda w_{i} Y_{i})]}{e^{\lambda t}}$$
(56)

$$\leq \frac{\prod_{i} \mathbb{E}[\exp(\lambda w_{i} Y_{i})]}{e^{\lambda t}}$$
(57)

$$\leq \frac{\exp(\lambda^2 \|\boldsymbol{w}\|_2^2/8)}{e^{\lambda t}}$$
(58)

$$\leq \exp(-2t^2/\|\boldsymbol{w}\|_2^2). \tag{59}$$

In (57) we used Negative association and the fact that each function is increasing in Y since $w_i \ge 0$ (see Corollary 4 and Lemma 8 from Wajc [46]). Thus, we proved (51). Overall, we have

$$\mathbb{P}\{\|Q_{\sigma}\boldsymbol{w}+\boldsymbol{N}-\boldsymbol{\mu}(D)\|_{\infty} > \alpha\} \le 2k\mathbb{I}\{\alpha \le \|\boldsymbol{w}-\boldsymbol{1}/n\|_{1}\}\exp\left(-\frac{\alpha^{2}}{2\|\boldsymbol{w}\|_{2}^{2}}\right) + k\exp\left(-\frac{\alpha}{4\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}}\right)$$
(60)

where (60) results from the fact that maximum error is the TV-distance established in (48). We can upper bound each term by $\beta/2$ to get the condition that

$$\alpha^2 \gtrsim \max\{\min\{\|\boldsymbol{w} - \mathbf{1}/n\|_1^2, \log(k/\beta)\|\boldsymbol{w}\|_2^2\}, \log^2(k/\beta)\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_\infty^2\}$$
(61)

$$\simeq \min\{\|\boldsymbol{w} - \mathbf{1}/n\|_{1}^{2}, \log(k/\beta)\|\boldsymbol{w}\|_{2}^{2}\} + \log^{2}(k/\beta)\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^{2}.$$
(62)

Taking minimum over such α , optimizing over the weights \boldsymbol{w} , and due to the projection to [0, 1] in the algorithm, we have $\mathcal{R}_{\boldsymbol{u}}^f \leq R_U(k, \beta, \boldsymbol{\epsilon})$.

Interesting Note: In (60), we used union bound and (51), which finally results in (62). In particular, there is a $\log(k/\beta)$ coefficient to the $||w||_2^2$ term. This is might be intuitive as the $\log(k)$ term results from union bound over the k coordinates. However, this method completely disregards that the matrix Q has only one-hot vectors and one can leverage this structure to get tighter inequalities. In particular, using the Kearns-Saul inequality, one can get a coefficient of the order $1 + \frac{\log(1/\beta)}{\log k}$. Thus, the effect of k is now opposite – a higher k results in a smaller coefficient!

F Proof of Theorem C.2

Theorem (C.2).

(A) Correlated Setting: For frequency estimation

$$\mathcal{E}_c^f(k,\beta,\epsilon) \lesssim R_C(k,1,\epsilon)^2,\tag{63}$$

and for mean estimation

$$\mathcal{E}_c^m(\beta, \boldsymbol{\epsilon}) \lesssim R_C(e, 1, \boldsymbol{\epsilon})^2.$$
(64)

(B) Uncorrelated Setting: For frequency estimation

$$\mathcal{E}_{u}^{f}(k,\boldsymbol{\epsilon}) \lesssim R_{U}(k,1,\boldsymbol{\epsilon})^{2}, \tag{65}$$

and for mean estimation

$$\mathcal{E}_u^m(\boldsymbol{\epsilon}) \lesssim R_U(e, 1, \boldsymbol{\epsilon})^2.$$
 (66)

Proof. We shall reuse parts of the proof of Theorem C.1 in Appendix E. We present the proof for frequency estimation, similar steps can be used for mean estimation.

(A) Begin by noting that

$$\mathbb{E}[\|\boldsymbol{Q}\boldsymbol{w} + \boldsymbol{N} - \boldsymbol{\mu}(\boldsymbol{D})\|_{\infty}^{2}] = \mathbb{E}[\|\boldsymbol{Q}(\boldsymbol{w} - \boldsymbol{1}/n) + \boldsymbol{N}\|_{\infty}^{2}]$$
(67)

$$\leq \mathbb{E}[\|Q(\boldsymbol{w}-\mathbf{1}/n)\|_{\infty}^{2}] + \mathbb{E}[\|\boldsymbol{N}\|_{\infty}^{2}].$$
(68)

From (48), recall

$$\mathbb{P}\{\|Q(\boldsymbol{w}-\boldsymbol{1}/n)\|_{\infty} > \alpha/2\} \le \mathbb{I}\{\alpha/2 \le \|\boldsymbol{w}-\boldsymbol{1}/n\|_{\mathsf{TV}}\}.$$
(69)

Therefore, using the above and tail-sum form of expectation, we get

$$\mathbb{E}[\|Q(w-1/n)\|_{\infty}^{2}] \lesssim \|w-1/n\|_{1}^{2}.$$
(70)

Using Lemma F.1 (it can be applied since squared magnitudes of Laplace distribution is equivalent to squared of exponential distribution), we get

$$\mathbb{E}[\|\boldsymbol{N}\|_{\infty}^2] \lesssim \log^2(k) \|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2.$$
(71)

Combining (70) and (71), we get the result. Note that for mean estimation, k = 1, one needs to use (76) instead of the result in Lemma F.1 which assumes $k \ge 2$.

(B) In the uncorrelated setting, we shall use the results of part (a) along with another inequality that we shall prove. Begin by noting that

$$\mathbb{E}[\|Q_{\sigma}\boldsymbol{w} + \boldsymbol{N} - \boldsymbol{\mu}(D)\|_{\infty}^{2}] = \mathbb{E}[\|Q_{\sigma}(\boldsymbol{w} - \boldsymbol{1}/n) + \boldsymbol{N}\|_{\infty}^{2}]$$
(72)

$$\leq \mathbb{E}[\|Q_{\sigma}(\boldsymbol{w}-\boldsymbol{1}/n)\|_{\infty}^{2}] + \mathbb{E}[\|\boldsymbol{N}\|_{\infty}^{2}].$$
(73)

(70) already gives us a bound on $\mathbb{E}[\|Q_{\sigma}(\boldsymbol{w}-\mathbf{1}/n)\|_{\infty}^2]$ but we shall also derive an alternate bound. From (51), we know that each element of the vector $\|Q_{\sigma}(\boldsymbol{w}-\mathbf{1}/n)\|_{\infty}^2$ behaves like a sub-exponential random variable. In particular, let $Y_i = |\sum_j Q_{\sigma}(i,j)(w_j-1/n)|^2 \ge 0$. Then, from (51), we have

$$\mathbb{P}\{Y_i \ge t\} \le 2 \exp\left(-\frac{2t}{\|\boldsymbol{w}\|_2^2}\right)$$

Using Lemma F.2, we have

$$\mathbb{E}[\max_{i} Y_i] \lesssim \log(k) \|\boldsymbol{w}\|_2^2,$$

completing the proof. Note that for mean estimation, k = 1, one can not use the fact $\log(2k) \lesssim \log(k)$ used above.

Lemma F.1. Let $k \ge 2$ and N_1, \ldots, N_k be i.i.d. $Exp(\lambda)$ random variables. Then $E[\max_{i=1}^k N_i^2] \simeq \frac{\log^2(k)}{\sqrt{2}}$.

Proof. Note that $\max_{i=1}^{k} N_i$ can be written as $\sum_{i=1}^{k} Y_i$ where Y_i is $Exp((k-i+1)\lambda)$ and independent of other Y_s . Thus,

$$E[\max_{i=1}^{k} N_i^2] = \mathsf{Var}(\sum_i Y_i) + E[\sum_i Y_i]^2$$
(74)

$$= \frac{1}{\lambda^2} \left(\sum_{i} 1/i^2 + (\sum_{i} 1/i)^2 \right)$$
(75)

$$\simeq \frac{1}{\lambda^2} \left(1 + \log^2(k) \right) \tag{76}$$

$$\simeq \frac{\log^2 k}{\lambda^2}.\tag{77}$$

Lemma F.2. Let Y_1, \ldots, Y_k non-negative be sub-exponential random variables satisfying $\mathbb{P}\{Y_i \ge t\} \le c_1 \exp(-c_2 t) \ \forall t > 0$, not necessarily independent. Then, $E[\max_{i=1}^k Y_i] \le \frac{(c_1+1)\log(2k)}{c_2}$.

Proof. Similar to the standard technqiue for proving upper bound on maximum of sub-gaussians, for $\lambda > 0$, we have

$$e^{\lambda \mathbb{E}[\max_{i=1}^{k} Y_i]} \le \mathbb{E}[e^{\lambda \max_{i=1}^{k} Y_i}]$$
(78)

$$\leq \mathbb{E}[\sum_{i} e^{\lambda Y_{i}}] \tag{79}$$

$$=k\int_{0}^{\infty} \mathbb{P}\{e^{\lambda Y} > t\}dt$$
(80)

$$\leq k \left(1 + \int_{1}^{\infty} \mathbb{P}\{Y > \log(t)/\lambda\} dt \right)$$
(81)

$$\leq k \left(1 + c_1 \int_0^\infty \exp(-c_2 \log(t)/\lambda) dt \right)$$
(82)

$$= k + \frac{kc_1}{c_2/\lambda - 1} \text{ for } \lambda < c_2$$
(83)
he desired result.

Plugging in $\lambda = c_2/(c_1 + 1)$ gives the desired result.

G Proof of Theorem C.3

Theorem (C.3). We have $\mathcal{R}_{c}^{f}(k,\beta,\epsilon), \mathcal{R}_{u}^{f}(k,\beta,\epsilon), \mathcal{R}_{c}^{m}(\beta,\epsilon), \mathcal{R}_{u}^{m}(\beta,\epsilon) \leq \frac{1}{2}$. (A) Correlated Setting: if $\mathcal{R}_{c}^{f}(k,\beta,\epsilon) < \frac{1}{2}$, then it also satisfies $\begin{bmatrix} 2n\mathcal{R}_{c}^{f}(k,\beta,\epsilon) \end{bmatrix} \qquad (1 (1 - \epsilon))$

$$\sum_{i=1}^{\ln \mathcal{R}_{c}^{j}(k,\beta,\epsilon)\rceil} \epsilon_{i} \gtrsim \log\left(\frac{k(1-\beta)}{\beta}\right).$$
(84)

Similarly, if $\mathcal{R}_c^m(\beta, \epsilon) < \frac{1}{2}$, then it also satisfies

$$\sum_{i=1}^{\lceil 2n\mathcal{R}_c^m(\beta,\epsilon)\rceil} \epsilon_i \gtrsim \log\left(\frac{(1-\beta)}{\beta}\right).$$
(85)

(B) Uncorrelated Setting:

Let $\{Z_i\}_{i=1}^n$ be sampled without replacement from the entries of ϵ . If $\mathcal{R}_u^f(k,\beta,\epsilon) < \frac{1}{2}$, then it also satisfies

$$\mathbb{E}\left[\exp\left\{-\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{f}(k,\beta,\epsilon)\rceil}Z_{i}\right\}\right] \leq \frac{k\beta}{k-1},$$
(86)

$$\mathbb{E}\left[\exp\left\{\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{f}(k,\beta,\epsilon)\rceil} Z_{i}\right\}\right] \ge k(1-\beta).$$
(87)

Similarly, if $\mathcal{R}_u^m(\beta, \epsilon) < \frac{1}{2}$, then it also satisfies

$$\mathbb{E}\left[\exp\left\{-\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{m}(\beta,\epsilon)\rceil}Z_{i}\right\}\right] \leq \beta,$$
(88)

$$\mathbb{E}\left[\exp\left\{\sum_{i=1}^{\lceil 2n\mathcal{R}_{u}^{m}(\beta,\epsilon)\rceil}Z_{i}\right\}\right] \ge (1-\beta).$$
(89)

Proof. $\mathcal{R}_c^f, \mathcal{R}_c^m, \mathcal{R}_u^f$ and \mathcal{R}_u^m are less than equal to half is trivial since the estimate of 1/2 or 1/2 achieves error at most half. We only illustrate the proof technique for frequency estimation. Similar steps can be used for mean estimation.

The proof technique adapts the homogeneous DP packing lower bounds (for example, see Vadhan [26]) to our setting. Let $M(\cdot)$ be an ϵ -DP randomized algorithm that is within $\alpha < \frac{1}{2}$ of $\mu(D)$, in l_{∞} norm, with probability greater than $1 - \beta$ for any dataset D.

Define k different datasets D^i for $i \in [k]$ in the following way: $\mu(D^i) = (1-t)e_1 + te_i$, where e_j is the j-th standard basis vector in \mathbb{R}^k . Without loss of generality, we assume the first nt indexes in D^i are the ones with value in bin i. Let $B_\alpha(D^i)$ be the l_∞ -ball of radius α centered at $\mu(D^i)$. Setting $t = \frac{\lceil 2\alpha n \rceil}{n}$, observe that $B_\alpha(D^i)$ are disjoint⁶.

(A) In the correlated setting, we have

$$-\beta \le \mathbb{E}[\mathbb{I}\{M(D^i) \in B_\alpha(D^i)\}] \,\forall i,\tag{90}$$

where the expectation is over the randomness in the algorithm \mathcal{M} . By DP constraint, we also have,

$$\mathbb{E}[\mathbb{I}\{M(D^i) \in B_{\alpha}(D^i)\}] \le e^{\sum_{i=1}^{nt} \epsilon_i} \mathbb{E}[\mathbb{I}\{M(D^1) \in B_{\alpha}(D^i)\}].$$
(91)

By (90) and (91), summing $i \in [k]$, and noting that $B_{\alpha}(D^{i})$ are disjoint, we get

$$k(1-\beta) \le e^{\sum_{i=1}^{nt} \epsilon_i}.$$
(92)

Thus, $\sum_{i=1}^{nt} \epsilon_i \ge \log k(1-\beta)$. In (91), if we instead use

1

1

$$\mathbb{P}\{M(D^i) \in B_{\alpha}(D^i)\} \le 1 - e^{-\sum_{i=1}^{nt} \epsilon_i} + e^{-\sum_{i=1}^{nt} \epsilon_i} \mathbb{P}\{M(D^1) \in B_{\alpha}(D^i)\},\$$

which is also a consequence of the DP definition (see Lemma 13 in Chaudhuri et al. [32]), then we obtain $\sum_{i=1}^{nt} \epsilon_i \ge \log(\frac{k-1}{k\beta})$. Thus, we can get

$$\sum_{i=1}^{nt} \epsilon_i \gtrsim \log \frac{(k-1)(1-\beta)}{\beta}.$$

(B) In the uncorrelated setting we have:

$$-\beta \le \mathbb{E}[\mathbb{I}\{M(D^i_{\sigma}) \in B_{\alpha}(D^i)\}] \,\forall i, \tag{93}$$

where the expectation is over the randomness in the algorithm \mathcal{M} and the permutation σ . Let $\gamma : [n] \to [n]$ be such that $\gamma(i) = \sigma^{-1}(i)$. By DP constraint, we also have,

$$\mathbb{E}[\mathbb{I}\{M(D^{i}_{\sigma})\in B_{\alpha}(D^{i})\}|\sigma] \le e^{\sum_{i=1}^{nt}\epsilon_{\gamma(i)}}\mathbb{E}[\mathbb{I}\{M(D^{1}_{\sigma})\in B_{\alpha}(D^{i})\}|\sigma].$$
(94)

By taking expectation over σ (94) and (93), summing $i \in [k]$, noting that $B_{\alpha}(D^i)$ are disjoint, we get

$$k(1-\beta) \le \mathbb{E}[e^{\sum_{i=1}^{nt} \epsilon_{\gamma(i)}}].$$
(95)

Using the same argument as part (A), using the alternate expression implied by DP, the following is obtained

$$E[e^{-\sum_{i=1}^{nt} \epsilon_{\gamma(i)}}] \le \frac{k\beta}{k-1}.$$
(96)

It should be noted that the above steps are only valid if $\alpha < \frac{1}{2}$. $\alpha = \frac{1}{2}$ is not admissible in the above due to our modified definition of the ceiling function rendering t to be greater than one.

⁶Note the slightly modified definition of ceiling function that we use which ensures the sets are disjoint.

H Proof of Theorem C.4

Our proof is based on the idea of [47], augmented by a more methodical step-by-step analysis.

Theorem (C.4). We have $\mathcal{E}_{u}^{f}(k, \epsilon), \mathcal{E}_{u}^{m}(\epsilon) \leq \frac{1}{4}$. Both $\mathcal{E}_{u}^{f}(k, \epsilon), \mathcal{E}_{u}^{m}(\epsilon)$ satisfy $\mathcal{E}_{u}^{f}(k, \epsilon), \mathcal{E}_{u}^{m}(\epsilon) \geq \left(\sqrt{\mathcal{E}_{stat}(\epsilon)} - \sqrt{\frac{1}{4n}}\right)_{+}^{2}$. (97)

Proof. We start with mean estimation.

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) = \min_{M \in \mathcal{M}_{\boldsymbol{\epsilon}}} \max_{D \in [0,1]^{n}} \mathbb{E}\left[|\mu(D) - M(D_{\sigma})|^{2} \right]$$
(98)

$$\geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in \{0,1\}^n} \mathbb{E}\left[|\mu(D) - M(D_{\sigma})|^2 \right]$$
(99)

$$\geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \mathbb{E}_{Z_1,\dots,Z_n \sim P} \left[\left(\mu(Z) - M(Z) \right)^2 \right], \tag{100}$$

$$= \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \mathbb{E}_{Z_1,\dots,Z_n \sim P} \left[(\mu(Z) - \mu(P))^2 + (\mu(P) - M(Z))^2 - 2(\mu(Z) - \mu(P))(\mu(P) - M(Z)) \right]$$
(101)

$$= \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left\{ \mathbb{E}\left[\left(\mu(P) - M(Z) \right)^2 \right] + \frac{\mathsf{Var}(P)}{n} - 2\mathbb{E}\left[\left(\mu(Z) - \mu(P) \right) \left(\mu(P) - M(Z) \right) \right] \right\},$$

$$(102)$$

$$\geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left\{ \mathbb{E}\left[\left(\mu(P) - M(Z) \right)^2 \right] + \frac{\mathsf{Var}(P)}{n} - 2\sqrt{\mathbb{E}\left[\left(\mu(Z) - \mu(P) \right)^2 \right] \mathbb{E}\left[\left(\mu(P) - M(Z) \right)^2 \right]} \right\}$$

$$= \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left\{ \mathbb{E}\left[\left(\mu(P) - M(Z) \right)^2 \right] + \frac{\mathsf{Var}(P)}{n} - 2\sqrt{\frac{\mathsf{Var}(P)}{n}} \mathbb{E}\left[\left(\mu(P) - M(Z) \right)^2 \right]} \right\},$$
(103)
(104)

$$= \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left\{ \underbrace{\frac{f(M,P) + \frac{\mathsf{Var}(P)}{n} - 2\sqrt{\frac{\mathsf{Var}(P)}{n}f(M,P)}}_{\text{Quadratic in }\sqrt{\frac{\mathsf{Var}(P)}{n}}} \right\}.$$
(105)

(100) follows by relaxing worst-case dataset to worst-case distribution where the data is sampled i.i.d. – this also helpds get rid of the permutation. (103) follows by Cauchy-Schwarz and we use the notation $f(M, P) = \mathbb{E}\left[\left(\mu(P) - M(Z)\right)^2\right]$ in (105). Now, we minimize the quadratic in (105) wrt $\sqrt{\frac{\operatorname{Var}(P)}{n}}$, which has a domain of $\left[0, \sqrt{\frac{1}{4n}}\right]$ for $P \in \mathcal{P}(\{0, 1\})$.

$$\geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left\{ f(M,P) + (f(M,P) \wedge \frac{1}{4n}) - 2\sqrt{(f(M,P) \wedge \frac{1}{4n})f(M,P)} \right\}, \quad (106)$$

$$= \min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left(\sqrt{f(M,P)} - \sqrt{\frac{1}{4n}} \right)_{+}^{2}$$
(107)

Now let $P^*(M) = \arg \max_{P \in \mathcal{P}(\{0,1\})} f(M, P)$.

For a given M, (107) is maximized wrt P if f(M, P) is maximized. Thus,

$$\min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \left(\sqrt{f(M,P)} - \sqrt{\frac{1}{4n}} \right)_{+}^{2} = \min_{M \in \mathcal{M}_{\epsilon}} \left(\sqrt{f(M,P^{*}(M))} - \sqrt{\frac{1}{4n}} \right)_{+}^{2}.$$
 (108)

The minimum over M is achieved when $f(M, P^*(M))$ is minimized. Noting that $\min_{M \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} f(M, P) = \min_{M \in \mathcal{M}_{\epsilon}} f(M, P^*(M)) = \mathcal{E}_{stat}(\epsilon)$, we have the desired result.

For frequency estimation,

$$\mathcal{E}_{u}^{f}(k, \boldsymbol{\epsilon}) = \min_{M \in \mathcal{M}_{\boldsymbol{\epsilon}}} \max_{D \in [k]^{n}} \mathbb{E}\left[\|\mu(D) - M(D_{\sigma})\|_{\infty}^{2} \right]$$
(109)

$$\geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [k]^n} \mathbb{E}\left[|\mu(D)_1 - M(D_{\sigma})_1|^2 \right]$$
(110)

(110) follows since the error in the first coordinate will be lower than the infinity norm. Next, we describe how we can now reduce this problem and follow the same steps as that of mean estimation.

Let
$$Q(p) = (p, 1-p, 0, 0, \dots, 0)$$
 be a multinomial distribution over k values. Then, we have

$$\min_{M \in \mathcal{M}_{\epsilon}} \max_{D \in [k]^n} \mathbb{E}\left[|\mu(D)_1 - M(D_{\sigma})_1|^2 \right] \geq \min_{M \in \mathcal{M}_{\epsilon}} \max_{p \in [0,1]} \mathbb{E}_{Z_1,\dots,Z_n \sim Q(p)} \left[|\mu(Z_1^n)_1 - M(Z_1^n)_1|^2 \right],$$
(111)

$$= \min_{\tilde{M} \in \mathcal{M}_{\epsilon}} \max_{P \in \mathcal{P}(\{0,1\})} \mathbb{E}_{W_1,\dots,W_n \sim P} \left[|\mu(W) - \tilde{M}(W)|^2 \right].$$
(112)

The first inequality follows by replacing worst-case dataset by expectation over the dataset generated by the multinomial distribution. Since a sample from the multinomial distribution is fully determined by the first-coordinate, we can restrict to algorithms \tilde{M} which only depend on the first-coordinate; thus, we have a change of variables and represent the problem as a scalar and now the same steps as that of mean estimation can be applied. We believe our bounds for frequency estimation can be made tighter with a more delicate analysis.

I Proof of Theorem 5.1 and Theorem 5.2

Theorem (5.1). (A) For $\frac{1}{2n} \leq \mathcal{R}_c^f(k, \beta, \epsilon) \leq \frac{1}{4}$, the lower bound (31) implies that error incurred by *HPF-C* \mathbb{P} is of the same order as the minimax rate, i.e.,

$$\mathcal{R}_c^f(k,\beta,\epsilon) \simeq R_C(k,\beta,\epsilon)$$

(B) For $\frac{1}{2n} \leq \mathcal{R}_c^m(\beta, \epsilon) \leq \frac{1}{4}$, the lower bound (32) implies that error incurred by HPM-CP is of the same order as the minimax rate, i.e.,

$$\mathcal{R}_c^m(\beta, \epsilon) \simeq R_C(1, \beta, \epsilon).$$

Proof. We present the proof for frequency estimation. Similar steps can be used for mean estimation. Let $t = \lceil 2n\mathcal{R}_C \rceil$. Since $\frac{1}{2n} \leq \mathcal{R}_c \leq \frac{1}{4}$, we have $\frac{t}{n} \simeq \mathcal{R}_c$ and $n - t \simeq n$. The lower bound in Theorem C.3(A) implies that t satisfies $\sum_{i=1}^{t} \epsilon_i \gtrsim \log(k/\beta)$ (we assume $1 - \beta \geq 1/2$ for meaningful regimes where we want low probability of error). Thus, we have

$$\epsilon_{t+1} \ge \log(k/\beta)/t$$
,

since ϵ is non-decreasing. Consider the weights w such that the first t entries are zero and the rest n-t entries are 1/(n-t). Using these weights, we get

$$R_C(k,\beta,\epsilon)^2 \lesssim \frac{t^2}{n^2} \simeq \mathcal{R}_c(k,\beta,\epsilon)^2,$$

where we used $\frac{1}{(n-t)\epsilon_{t+1}} \lesssim \frac{t}{n \log(k/\beta)}$.

Theorem (5.2). Assume high privacy demand where $\|\boldsymbol{\epsilon}\|_{\infty} \leq 1$, for the two problems. (A) Frequency Estimation: In the regime $\frac{1}{2n} \leq \mathcal{R}_{u}^{f}(k,\beta,\epsilon)$, we have the following upper and lower bound on \mathcal{R}_{u}^{f} ,

$$\frac{\log(k/\beta)}{\|\boldsymbol{\epsilon}\|_{1}} \lesssim \mathcal{R}_{u}^{f}(k,\beta,\boldsymbol{\epsilon}) \lesssim \frac{\log(k/\beta)}{\|\boldsymbol{\epsilon}\|_{1}} + \frac{\sqrt{n^{2}\mathsf{Var}(e^{-\boldsymbol{\epsilon}}) \wedge n\log(k/\beta)}}{\|\boldsymbol{\epsilon}\|_{1}}.$$
(113)

The upper bound is achieved by HPF-UP and HPF-A. Further, if $Var(e^{-\epsilon}) \lesssim \frac{\log^2(k/\beta)}{n^2}$, then we have

$$\mathcal{R}_{u}^{f}(k,\beta,\epsilon) \simeq \frac{\log(k/\beta)}{\|\epsilon\|_{1}}.$$
(114)

(B) Mean Estimation: In the regime $\frac{1}{2n} \leq \mathcal{R}_u^m(\beta, \epsilon)$, we have the following upper and lower bound on \mathcal{R}_u^m ,

$$\frac{\log(1/\beta)}{\|\boldsymbol{\epsilon}\|_{1}} \lesssim \mathcal{R}_{u}^{m}(\beta, \boldsymbol{\epsilon}) \lesssim \frac{\log(1/\beta)}{\|\boldsymbol{\epsilon}\|_{1}} + \frac{\sqrt{n^{2} \operatorname{Var}(e^{-\boldsymbol{\epsilon}}) \wedge n \log(1/\beta)}}{\|\boldsymbol{\epsilon}\|_{1}}.$$
(115)

The upper bound is achieved by HPM-UP and HPM-A. Further, if $Var(e^{-\epsilon}) \lesssim \frac{\log^2(1/\beta)}{n^2}$, then we have

$$\mathcal{R}_{u}^{m}(\beta, \epsilon) \simeq \frac{\log(k/\beta)}{\|\epsilon\|_{1}}.$$
(116)

Proof. We present the proof for frequency estimation. Similar steps can be used for mean estimation. In the upper bound, use the HPF-A weights $\boldsymbol{w} = \frac{1 - \exp(-\epsilon)}{\|1 - \exp(-\epsilon)\|_1}$. Using $\|\boldsymbol{w} - \mathbf{1}/n\|_1^2 \le n \|\boldsymbol{w} - \mathbf{1}/n\|_2^2$, we have

$$R_C(k,\beta,\epsilon)^2 \le \frac{\min\{n\|1 - e^{-\epsilon} - \frac{\|1 - e^{-\epsilon}\|_1}{n}\|_2^2, \log(k/\beta)\|1 - e^{-\epsilon}\|_2^2\} + \log^2(k/\beta)\left(\frac{1 - e^{-\epsilon_1}}{\epsilon_1}\right)^2}{\|1 - e^{-\epsilon}\|_1^2}$$

$$\leq \frac{\min\{n^{2} \operatorname{Var}(e^{-\epsilon}), \log(k/\beta) \| 1 - e^{-\epsilon} \|_{2}^{2}\} + \log^{2}(k/\beta)}{\| 1 - e^{-\epsilon} \|_{1}^{2}}$$
(118)

Using
$$1 - e^{-\epsilon} \gtrsim \epsilon$$
 for $\epsilon \le 1$, (119)

$$\leq \frac{\min\{n^2 \operatorname{Var}(e^{-\epsilon}), \log(k/\beta) \| 1 - e^{-\epsilon} \|_2^2\} + \log^2(k/\beta)}{\|\epsilon\|_1^2}$$
(120)

Using
$$||1 - e^{-\epsilon}||_2^2 = n \operatorname{Var}(e^{-\epsilon}) + \frac{||1 - e^{-\epsilon}||_1^2}{n} \le n \operatorname{Var}(e^{-\epsilon}) + \frac{||\epsilon||_1^2}{n},$$
 (121)

$$\leq \frac{\min\{n^{2}\mathsf{Var}(e^{-\epsilon}), \log(k/\beta)n\mathsf{Var}(e^{-\epsilon}) + \log(k/\beta)\frac{\|\epsilon\|_{1}^{2}}{n}\} + \log^{2}(k/\beta)}{\|\epsilon\|_{1}^{2}}$$

$$= \frac{\log^{2}(k/\beta)}{\|\epsilon\|_{1}^{2}} + \min\left\{\frac{n^{2}\mathsf{Var}(e^{-\epsilon})}{\|\epsilon\|_{1}^{2}}, \log(k/\beta)\left(\frac{1}{n} + \frac{n\mathsf{Var}(e^{-\epsilon})}{\|\epsilon\|_{1}^{2}}\right)\right\}$$
(123)

If in (120), we use $||1 - e^{-\epsilon}||_2^2 \le n$, then we would get

$$R_C^2 \lesssim \frac{\log^2(k/\beta)}{\|\boldsymbol{\epsilon}\|_1^2} + \min\left\{\frac{n^2 \mathsf{Var}(e^{-\boldsymbol{\epsilon}})}{\|\boldsymbol{\epsilon}\|_1^2}, \frac{n \log(k/\beta)}{\|\boldsymbol{\epsilon}\|_1^2}\right\}.$$

Coming to the lower bound, use Jensen's inequality in (33) to obtain

$$\log \frac{k-1}{k\beta} \le \frac{\lceil 2n\mathcal{R}_u \rceil}{n} \|\boldsymbol{\epsilon}\|_1 \simeq \mathcal{R}_u \|\boldsymbol{\epsilon}\|_1.$$
(124)

Let Z be a sample from the vector $\sim \epsilon$. Using Negative Association in (34), we get

$$k(1-\beta) \le \mathbb{E}[e^{Z}]^{\lceil 2n\mathcal{R}_{u}\rceil}$$
(125)

$$= \left(\frac{1}{n}\sum_{i}e^{\epsilon_{i}}\right)^{+2m\epsilon_{u}}$$
(126)

$$\leq \left(\frac{1}{n}\sum_{i}(1+(e-1)\epsilon_i)\right)^{\lceil 2n\mathcal{R}_u\rceil}$$
(127)

$$= \left(1 + \frac{(e-1)}{n} \|\boldsymbol{\epsilon}\|_1\right)^{\lceil 2n\mathcal{R}_u\rceil}.$$
(128)

In the above, (127) follows from $\|\boldsymbol{\epsilon}\|_{\infty} \leq 1$. Taking logarithm and using $\log(1+x) \leq x$, we get

$$\log k(1-\beta) \lesssim \mathcal{R}_u \|\boldsymbol{\epsilon}\|_1. \tag{129}$$

By adding (124) and (129), we get

$$\log \frac{k}{\beta} \lesssim \mathcal{R}_u \|\boldsymbol{\epsilon}\|_1. \tag{130}$$

J Proof for Theorem 5.3

Theorem (5.3). *For values of* ϵ *satisfying* $\mathcal{E}_{stat}(\epsilon) \geq \frac{1}{2n}$ *, we have*

$$\mathcal{E}_{stat}(\boldsymbol{\epsilon}) \lesssim \mathcal{E}_{u}^{f}(k, \boldsymbol{\epsilon}) \lesssim \log(k)^{2} \mathcal{E}_{stat}(\boldsymbol{\epsilon})$$
 (131)

for frequency estimation and

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) \simeq \mathcal{E}_{stat}(\boldsymbol{\epsilon}) \tag{132}$$

for mean estimation.

Proof. From Theorem C.4, we directly have $\mathcal{E}_{stat}(\epsilon) \lesssim \mathcal{E}_{u}^{f}(k, \epsilon), \mathcal{E}_{u}^{m}(k, \epsilon)$ in the regime of interest. Now, observe that

$$\mathcal{E}_{u}^{f}(k,\boldsymbol{\epsilon}) \lesssim \min_{\boldsymbol{w}\in\Delta_{n}} \left(\left\| \boldsymbol{w} - \frac{1}{n} \right\|_{1}^{2} \wedge \log(k) \|\boldsymbol{w}\|_{2}^{2} \right) + \log(k)^{2} \left\| \frac{\boldsymbol{w}}{\boldsymbol{\epsilon}} \right\|_{\infty}^{2},$$
(133)

$$\lesssim \log(k)^2 \min_{\boldsymbol{w} \in \Delta_n} \|\boldsymbol{w}\|_2^2 + \left\|\frac{\boldsymbol{w}}{\boldsymbol{\epsilon}}\right\|_{\infty}^2.$$
(134)

Similarly,

$$\mathcal{E}_{u}^{m}(\boldsymbol{\epsilon}) \lesssim \min_{\boldsymbol{w} \in \Delta_{n}} \|\boldsymbol{w}\|_{2}^{2} + \left\|\frac{\boldsymbol{w}}{\boldsymbol{\epsilon}}\right\|_{\infty}^{2}.$$
(135)

Using the result of Chaudhuri et al. [32] that $\mathcal{E}_{stat}(\epsilon) \simeq \min_{\boldsymbol{w} \in \Delta_n} \|\boldsymbol{w}\|_2^2 + \left\|\frac{\boldsymbol{w}}{\epsilon}\right\|_{\infty}^2$, we get the desired.

It may be noted that the statistical minimax rate considered in Chaudhuri et al. [32] is over all distributions on [0, 1] instead of our definition of all distributions on $\{0, 1\}$. However, bound their upper and lower bound work for this case as well and leads to identical result.

K A Special Case for \mathcal{E}_c^f

We consider a special case in this Section and prove a tight lower bound using a different technique. This Section is added purely for fun.

Consider the scenario where there are only two bins and out of n users, n/2 of them have privacy demand $\epsilon \to 0$ and the rest have a privacy demand of $\epsilon \to \infty$. Let the dataset be presented as

 $Z = (Z_1, \ldots, Z_n) \in \{0, 1\}$. The frequency is given by $\mu(Z) = (1 - \frac{\sum_i Z_i}{n}, \frac{\sum_i Z_i}{n})$. For this section, we use slightly different notations, that we described below.

For this special setting, we denote the minimax rate by $\hat{\mathcal{R}}_u(n)$. Let the set of all algorithms that are 0-DP in the first n/2 entries and ∞ -DP in the next n/2 entries of the input be represented as $\hat{\mathcal{M}}(n)$. In particular, we have the definition for minimax expected squared ℓ_{∞} error as

$$\hat{\mathcal{R}}_{u}(n) = \min_{M \in \mathcal{M}(n)} \max_{Z \in \{0,1\}^{n}} \mathbb{E}\left[\|\mu(Z) - M(Z_{\sigma})\|_{\infty}^{2} \right].$$
(136)

The above expectation is over the permutations and randomness in M. By the result in the following two subsections (see (145) and (137)), we have $\hat{\mathcal{R}}_u(n) \simeq \frac{1}{n}$. This implies that our upper bound, given by HPF-U, is tight in this regime.

K.1 Upper Bound

HPF-U, for this setting, just uses the empirical frequency from public data as the estimator. HPF-U will have a squared ℓ_{∞} error greater than $\frac{\log(2/\beta)}{n}$ with probability less than β . In terms of expected squared ℓ_{∞} error, one can integrate the tail bound in (60) to get the upper bound

$$\hat{\mathcal{R}}_u(n) \lesssim \frac{1}{n}.\tag{137}$$

K.2 Lower Bound

Let the data of the n/2 users with public data be $X = (X_1, \ldots, X_{n/2})$ which are samples without replacement from Z and let the rest of the private data be $Y = (Y_1, \ldots, Y_{n/2})$. Based on X, the algorithm would need to estimate $\mu(Z)$.

Out of the two bins (0,1), we can just focus on bin 1. Thus we denote $M(Z_{\sigma})_1$ by f(X). We have,

$$\hat{\mathcal{R}}_u(n) = \min_{M \in \mathcal{M}(n)} \max_{Z \in \{0,1\}^n} \mathbb{E}\left[\|\mu(Z) - M(Z_\sigma)\|_\infty^2 \right]$$
(138)

$$\geq \min_{M \in \mathcal{M}(n)} \max_{Z \in \{0,1\}^n} \mathbb{E}\left[|\mu(Z)_1 - f(X)|^2 \right]$$
(139)

$$= \min_{M \in \mathcal{M}(n)} \max_{Z \in \{0,1\}^n} \mathbb{E}\left[\left| \frac{\sum_{i=1}^{n/2} X_i + Y_i}{n} - f(X) \right|^2 \right]$$
(140)

$$\geq \min_{M \in \mathcal{M}(n)} \mathbb{E}_{Z_1, \dots, Z_n \sim \mathsf{Bern}(1/2)} \left[\left| \frac{\sum_{i=1}^{n/2} X_i + Y_i}{n} - f(X) \right|^2 \right], \tag{141}$$

where the last inequality follows by replacing the worst-case dataset Z by taking expectation over i.i.d. samples $Z_1, \ldots, Z_n \sim \text{Bern}(1/2)$. Now, since Z_1, \ldots, Z_n are i.i.d. Bern(1/2) and X, Yare samples without replacement from Z, we have the fact that $X_1, \ldots, X_{n/2}, Y_1, \ldots, Y_{n/2}$ are i.i.d. Bern(1/2). Since X and Y are now independent, and by the MMSE property of conditional expectation, we have

$$\mathbb{E}\left[\left|\frac{\sum_{i=1}^{n/2} X_i + Y_i}{n} - f(X)\right|^2\right] \ge \mathbb{E}\left[\left|\frac{\sum_{i=1}^{n/2} X_i + Y_i}{n} - \mathbb{E}\left[\frac{\sum_{i=1}^{n/2} X_i + Y_i}{n}|X\right]\right|^2\right]$$
(142)

=

=

$$\mathbb{E}\left[\left|\frac{\sum_{i=1}^{n/2} Y_i}{n} - \frac{1}{4}\right|^2\right] \tag{143}$$

$$\frac{1}{8n}.$$
 (144)

Thus, we have

$$\hat{\mathcal{R}}_u(n) \gtrsim \frac{1}{n}.$$
(145)

L Weights in Local DP

L.1 Frequency Estimation

Suppose user *i* has a datapoint in bin *j*, then present user *i*'s datapoint D_i as $e_j \in \mathbb{R}^k$. In the following, we shall represent user *i*'s true data as $D_i \in \{0, 1\}^k$.

In the k-RAPPOR scheme, user *i* flips each bit of the vector independently with probability $\frac{1}{1+\exp(\epsilon_i/2)}$ and sends the new vector \boldsymbol{x}_i to the server. After receiving the vector $\boldsymbol{x}_i \in \{0,1\}^k$ from user *i*, the server can compute $\operatorname{coth}(\frac{\epsilon_i}{4})(\boldsymbol{x}_i - \mathbf{1}/(1 + \exp(\epsilon_i/2)))$ which is an unbiased estimator of the user's data. For weights $\boldsymbol{w} \in \Delta_n$, let the overall estimate for $\mu(D)$ be $\sum_{i=1}^n w_i \operatorname{coth}(\frac{\epsilon_i}{4})(\boldsymbol{x}_i - \mathbf{1}/(1 + \exp(\epsilon_i/2)))$.

Now, representing this mechanism as M(D), we have

$$\mathbb{P}\{\|M(D)-\mu(D)\|_{\infty} > \alpha\} \le \mathbb{P}\{\|M(D)-\sum_{i} w_{i}\boldsymbol{D}_{i}\|_{\infty} > \alpha/2\} + \mathbb{P}\{\|\sum_{i} w_{i}\boldsymbol{D}_{i}-\mu(D)\|_{\infty} > \alpha/2\}$$
(146)

Now note that $\mathbb{E}[\coth\left(\frac{\epsilon_i}{4}\right)(\boldsymbol{x}_i-\mathbf{1}/(1+\exp\left(\epsilon_i/2\right)))] = \boldsymbol{D}_i$. For $j \in [k]$, let Y_i^j be the *j*-th component of $\coth\left(\frac{\epsilon_i}{4}\right)(\boldsymbol{x}_i-\mathbf{1}/(1+\exp\left(\epsilon_i/2\right))) - \boldsymbol{D}_i$. Then Y_i^j is either a $\coth\left(\frac{\epsilon_i}{4}\right) \operatorname{Bern}\left(\frac{e^{\epsilon_i/2}}{1+e^{\epsilon_i/2}}\right) - \frac{e^{\epsilon_i/2}}{e^{\epsilon_i/2}-1}$ RV (if $\boldsymbol{D}_i = \boldsymbol{e}_j$) or a $\coth\left(\frac{\epsilon_i}{4}\right) \operatorname{Bern}\left(\frac{1}{1+e^{\epsilon_i/2}}\right) - \frac{1}{e^{\epsilon_i/2}-1}$ RV (otherwise). For a given j, Y_i^j are independent as well.

Therefore, using Kearns-Saul concentration inequality [48, 49], we get

$$\mathbb{P}\{\|M(D) - \sum_{i} w_i \boldsymbol{D}_i\|_{\infty} > \alpha/2\} \le 2k \exp\left[-\frac{\alpha^2}{8\sum_{i} w_i^2 \coth\left(\frac{\epsilon_i}{4}\right)/\epsilon_i}\right]$$
(147)

For the $\mathbb{P}\{\|\sum_i w_i D_i - \mu(D)\|_{\infty} > \alpha/2\}$ term, exact same analysis as that of Theorem C.1 works. Thus,

• In the correlated setting, we have

$$\alpha^2 \lesssim \left(\min_{\boldsymbol{w} \in \Delta_n} \|\boldsymbol{w} - \mathbf{1}/n\|_1^2 + \log(k/\beta) \sum_i w_i^2 \coth\left(\frac{\epsilon_i}{4}\right)/\epsilon_i \right) \wedge 1.$$
 (148)

• In the uncorrelated setting, we have

$$\alpha^2 \lesssim \left(\min_{\boldsymbol{w} \in \Delta_n} \left[\|\boldsymbol{w} - \mathbf{1}/n\|_1^2 \wedge \log(k/\beta) \|\boldsymbol{w}\|_2^2 \right] + \log(k/\beta) \sum_i w_i^2 \coth\left(\frac{\epsilon_i}{4}\right)/\epsilon_i \right) \wedge 1.$$
(149)

For the purpose of implementation, we upper bound the ℓ_1 norm in the above by $n\ell_2$ norm.

L.2 Mean Estimation

Let user *i*'s data be $X_i \in [0, 1]$ then it sends to the server $Y_i = X_i + N_i$ where $N_i \sim \text{Laplace}(\frac{1}{\epsilon_i})$. The server's estimate is $\sum_i w_i Y_i$ for $w \in \Delta_n$.

Now, representing this mechanism as M(Y), we have

$$\mathbb{P}\{|M(Y) - \mu(X)| > \alpha\} \le \mathbb{P}\{|\sum_{i} w_i N_i| > \alpha/2\} + \mathbb{P}\{|\sum_{i} w_i X_i - \mu(X)| > \alpha/2\}.$$
 (150)

For the second term, exact same analysis as that of Theorem C.1 works so we focus on bounding the first term. Using the tail bound of Chan et al. [50] (Lemma 2.8), we have

$$\mathbb{P}\{|\sum_{i} w_{i}N_{i}| > \alpha/2\} \le \begin{cases} \exp\{-c\frac{\alpha^{2}}{\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^{2}}\} & \text{if } \alpha/2 \le \frac{2\sqrt{2}\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{2}^{2}}{\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}}\\ \exp\{-c'\frac{\alpha}{\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}}\} & \text{else.} \end{cases}$$
(151)

To simplify the problem, we just consider the sub-exponential tail bound as a heuristic for the optimization. Thus,

• In the correlated setting, we have

$$\boldsymbol{w}_{LDP} = \underset{\boldsymbol{w} \in \Delta_n}{\arg\min} \|\boldsymbol{w} - \mathbf{1}/n\|_1^2 + \log(1/\beta)^2 \|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2.$$
(152)

• In the uncorrelated setting, we have

$$\boldsymbol{w}_{LDP} = \underset{\boldsymbol{w}\in\Delta_n}{\arg\min}\left[\|\boldsymbol{w} - \mathbf{1}/n\|_1^2 \wedge \log(k/\beta)\|\boldsymbol{w}\|_2^2\right] + \log(1/\beta)^2\|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2.$$
(153)

Recall that these are exact same weights obtained in HPM-CP and HPM-UP!

M Faster Variants of HPF-CP and HPF-UP

Algorithm 3 Heterogeneously Private Frequency *Turbo* for different cases. HPF-CT and HPF-UT are for the correlated and the uncorrelated setting respectively.

Input: $\epsilon \in \mathbb{R}_{>0}^{n}, X \in [k]^{n}, \beta > 0$ HPF-CP-T: Set $w^{*} = \underset{w \in \Delta_{n}}{\operatorname{argmin}} n \left\| w - \frac{1}{n} \right\|_{2}^{2} + \log^{2}(k/\beta) \left\| \frac{w}{\epsilon} \right\|_{\infty}^{2}$. HPF-UP-T: Set $w^{*} = \underset{w \in \Delta_{n}}{\operatorname{argmin}} \left(n \left\| w - \frac{1}{n} \right\|_{2}^{2} \wedge \log \left(\frac{k}{\beta} \right) \| w \|_{2}^{2} \right) + \log^{2} \left(\frac{k}{\beta} \right) \left\| \frac{w}{\epsilon} \right\|_{\infty}^{2}$ Sample i.i.d. $N_{1}, \ldots, N_{k} \sim \operatorname{Laplace}(2 \| \frac{w^{*}}{\epsilon} \|_{\infty})$ Calculate $y \in \mathbb{R}^{k}$: $y_{j} = N_{j} + \sum_{i=1}^{n} w_{i}^{*} \mathbb{I}\{X_{i} = j\}, \forall j \in [k].$

Set $y_j \leftarrow \max\{\min\{y_j, 1\}, 0\} \forall j \in [k].$ return y

In the algorithms HPF-CP and HPF-UP, the minimization, although convex, can be a bottleneck and we attempt to consider more efficient variants. If we upper bound the ℓ_1 -norm by $n\ell_2$ -norm, then the optimization problem has a closed form solution which we can implement efficiently in $O(n \log n)$ time. Further, the arithmetic operations involved in finding the solution do not suffer from the numerical instability issues that cvxpy runs into in HPF-CP and HPF-UP when the entries of the privacy demand vector can range vary between several orders of magnitude. Increasing the number of users n also brings numerical instability for solving the weights using cvxpy.

These variants of the algorithm can be efficiently implemented since minimizations of form $\min_{\boldsymbol{w}\in\Delta_n} \|\boldsymbol{w}\|_2^2 + c \|\boldsymbol{w}/\boldsymbol{\epsilon}\|_{\infty}^2$ can be solved in $O(n\log n)$ time using the algorithm presented by Chaudhuri et al. [32]. In brief, for non-decreasing $\boldsymbol{\epsilon}$, construct the sequence

$$r_1 = \epsilon_1, \tag{154}$$

$$r_{k+1} = \frac{\sum_{i=1}^{k} r_i^2 + c}{\sum_{i=1}^{k} r_i} \wedge \epsilon_{k+1} \quad \forall k \in [n-1].$$
(155)

The solution can be obtained as $w^* = r/||r||_1$. We refer to these two faster variantspresented in Algorithm 3 as HPF-CT and HPF-UT respectively. For HPF-UT, one can solve two minimization problems – one using the term $n||w - 1/n||_2^2$ and one using the term $\log(k/\beta)||w||_2^2$ – and finally choose the weights corresponding to the one minimizing the two objective values.

We perform experiments with synthetic dataset in a similar manner as our earlier experiments to compare HPF-CP with HPF-CT in the correlated setting, and HPF-UP with HPF-UT in the uncorrelated setting. The graphs showing the empricial 95-th quantile of the l_{∞} error is presented in

Figure 2. Our proposed heuristic algorithm HPF-A performs the best and is the most efficient. In the correlated regime, HPF-CP performs better than HPF-CT at the cost of being more computationally difficult. In the uncorrelated regime, HPF-UP and HPF-UT perform exactly the same in Figure 2, indicating that the optimal weights must have been identical, which is possible if $\log(k/\beta) ||\boldsymbol{w}||_2^2$ term is smaller among the two terms in the minimization in both HPF-UP and HPF-UT.

Similar steps can be performed to speed up HPF-C \mathbb{E} and HPF-U \mathbb{E} .





(b) Correlared, Synthetic (10000, 20) Dataset



(c) Uncorrelared, Synthetic (10000, 5) Dataset

(d) Uncorrelared, Synthetic (10000, 20) Dataset

Figure 2: Comparison of performance of the HPF algorithms in Algorithm 1 and 3.