

CHAIN PROJECTION ORDERED CATEGORIES AND DRC-RESTRICTION SEMIGROUPS

YIN DIE AND SHOUFENG WANG*

ABSTRACT. In this paper we provide a theory of chain projection ordered categories and generalize that of chain projection ordered groupoids developed by East and Azeef Muhammed recently. By using chain projection ordered categories, we obtain a structure theorem for DRC-restriction semigroups. More specifically, we prove that the category of DRC-restriction semigroups together with $(2,1,1)$ -homomorphisms is isomorphic to the category of chain projection ordered categories together with chain projection ordered functors. Moreover, some special cases are also considered. As applications of the main theorem, we demonstrate that the existence of free projection-generated DRC-restriction semigroups associated to any strong two-sided projection algebra and reobtain the structures of projection-fundamental DRC-restriction semigroups. Our work may be regarded as an answer for the fourth problem proposed by East and Azeef Muhammed in [Advances in Mathematics, 437 (2024) 109447].

1. INTRODUCTION

Let (S, \cdot) be a semigroup. As usual, we denote the set of all idempotents in S by $E(S)$ and the set of all inverses of $x \in S$ by $V(x)$. Recall that $V(x) = \{a \in S \mid xax = x, axa = a\}$ for all $x \in S$. A semigroup S is called *regular* if $V(x) \neq \emptyset$ for any $x \in S$, and a regular semigroup S is called *inverse* if $E(S)$ is a commutative subsemigroup of S , or equivalently, the cardinal of $V(x)$ is equal to 1 for all $x \in S$ (see Howie [22] for example). The theory of inverse semigroups is perhaps the best developed within semigroup theory (see the monographs [30, 42]).

It is well known that a semigroup S is an inverse semigroup if and only if there exists a unary operation $\circ : S \rightarrow S, x \mapsto x^\circ$ satisfying the following axioms:

$$(1) \quad xx^\circ x = x, x^{\circ\circ} = x, (xy)^\circ = y^\circ x^\circ, xx^\circ yy^\circ = yy^\circ xx^\circ.$$

Inspired by this fact, in 1978 Nordahl and Scheiblich [41] introduced regular \circ -semigroups as a generalization of inverse semigroups in the range of regular semigroups. Recall that a unary semigroup (S, \cdot, \circ) is called a *regular \circ -semigroup* if the following axioms hold:

$$(2) \quad xx^\circ x = x, x^{\circ\circ} = x, (xy)^\circ = y^\circ x^\circ.$$

Inverse semigroups are regular \circ -semigroups obviously. But the converse is not true. Regular \circ -semigroups have been investigated extensively in literatures (for example, see [1, 3, 8, 9, 10, 23, 24, 26, 41, 43, 53, 62], etc.)

On the other hand, in 1991 Lawson introduced Ehresmann semigroups as a generalization of inverse semigroups in the range of non-regular semigroups by using generalized Green

Key words and phrases. Chain projection ordered categories; Strong two-sided projection algebras; DRC-restriction semigroups; Projection-fundamental DRC-restriction semigroups; Free projection-generated DRC-restriction semigroups.

*Corresponding author.

relation developed by “York School”. Let S be a semigroup and let $E \subseteq E(S)$. The relation $\tilde{\mathcal{R}}_E$ is defined on S by the rule that for any $x, y \in S$, we have $x\tilde{\mathcal{R}}_E y$ if

$$ex = x \text{ if and only if } ey = y \text{ for all } e \in E.$$

Dually, we have the relation $\tilde{\mathcal{L}}_E$ on S . Observe that both $\tilde{\mathcal{R}}_E$ and $\tilde{\mathcal{L}}_E$ are equivalences on S but $\tilde{\mathcal{R}}_E$ (respectively, $\tilde{\mathcal{L}}_E$) may not be a left congruence (respectively, a right congruence). Recall that a *band* is a semigroup in which every element is idempotent and a *semilattice* is a commutative band. In view of Lawson [31], (S, E) is called *E -Ehresmann* if

- E is a subsemilattice of S ,
- every $\tilde{\mathcal{R}}_E$ -class contains an element of E and $\tilde{\mathcal{R}}_E$ is a left congruence,
- every $\tilde{\mathcal{L}}_E$ -class contains an element of E and $\tilde{\mathcal{L}}_E$ is a right congruence.

If this is the case, E is called *the distinguished semilattice* of S , and the $\tilde{\mathcal{R}}_E$ -class (respectively, $\tilde{\mathcal{L}}_E$ -class) containing $a \in S$ has a unique element in E which will be denoted by a^+ (respectively, a^*). An E -Ehresmann semigroup (S, E) is called *E -restriction*, if $ae = (ae)^+a$ and $ea = a(ea)^*$ for all $e \in E$ and $a \in S$. We say that a semigroup S is *restriction* (respectively, *Ehresmann*) in the sequel if (S, E) is an E -restriction (respectively, E -Ehresmann) semigroup for some $E \subseteq E(S)$. Let (S, \cdot) be an inverse semigroup. Define $x^+ = xx^{-1}$ and $x^* = x^{-1}x$ for all $x \in S$. Then it is easy to check that $(S, \cdot, +, *)$ forms a restriction semigroup. From Lemmas 2.2 and 2.4 and their dual in Gould [17], we have the following characterizations of Ehresmann semigroups and restriction semigroups from a varietal perspective.

Lemma 1.1 ([17]). *A semigroup (S, \cdot) is Ehresmann if and only if there are two unary operations “+” and “*” on S such that the following axioms hold:*

$$x^+x = x, x^+y^+ = y^+x^+ = (x^+y^+)^+, x^+(xy)^+ = (xy)^+ = (xy^+)^+, x^{++} = x^+;$$

$$xx^* = x, x^*y^* = y^*x^* = (x^*y^*)^*, (xy)^*y^* = (xy)^* = (x^*y)^*, x^{**} = x^*.$$

*In this case, we denote this Ehresmann semigroup by $(S, \cdot, +, *)$. An Ehresmann $(S, \cdot, +, *)$ is restriction if and only if it additionally satisfies the following axioms*

$$\textbf{ample conditions} \quad (xy)^+x = xy^+, x(yx)^* = y^*x.$$

We observe that adequate semigroups and ample semigroups defined firstly in Fountain [14] are special Ehresmann semigroups and restriction semigroups, respectively. We also observe that restriction semigroups have appeared also as the type $SL2$ γ -semigroups [4, 5] in the early 1980s. Restriction semigroups also have arisen in the work of Jackson and Stokes [25] in the guise of twisted C -semigroups and in that of Manes [33] as guarded semigroups, motivated by consideration of closure operators and categories, respectively. The work of Manes has a forerunner in the restriction categories of Cockett and Lack [6], who were influenced by considerations of theoretical computer science. In fact the term “restriction semigroup” is exactly taken from the above work of Cockett and Lack. More detailed information on Ehresmann and restriction semigroups can be found in the survey articles of Gould [17] and Hollings [19].

In 2012, Jones [26] introduced P -Ehresmann and P -restriction semigroups from the view of variety and gave a common framework for restriction semigroups and regular $*$ -semigroups. A bi-unary semigroup $(S, \cdot, +, *)$ is called a *P -Ehresmann semigroup* if the

following axioms hold:

$$\begin{aligned} x^+x &= x, (xy)^+ = (xy^+)^+, (x^+y^+)^+ = x^+y^+x^+, x^+x^+ = x^+, x^{+*} = x^+, \\ xx^* &= x, (xy)^* = (x^*y)^*, (x^*y^*)^* = y^*x^*y^*, x^*x^* = x^*, x^{*+} = x^*. \end{aligned}$$

A P -Ehresmann semigroup $(S, \cdot, +, *)$ is called a P -restriction semigroup if it additionally satisfies the following axioms :

$$\textbf{P-ample conditions} \quad (xy)^+x = xy^+x^*, \quad x(yx)^* = x^+y^*x.$$

Let (S, \cdot, \circ) be a regular \circ -semigroup. Define $x^+ = xx^\circ$ and $x^* = x^\circ x$ for all $x \in S$. Then it is easy to check that $(S, \cdot, +, *)$ forms a P -restriction semigroup. Some works on P -Ehresmann and P -restriction semigroups can be found in [26, 27, 46, 47, 54, 55, 56, 58] and the references therein.

In 2021, Jones [29] introduced DRC semigroups and further generalized P -Ehresmann semigroups. A bi-ary semigroup $(S, \cdot, +, *)$ is called a DRC semigroup if the following axioms hold:

$$\begin{array}{l|l} \text{(i)} & x^+x = x & \text{(i)'} & xx^* = x \\ \text{(ii)} & (xy)^+ = (xy^+)^+ & \text{(ii)'} & (xy)^* = (x^*y)^* \\ \text{(iii)} & (xy)^+ = x^+(xy^+)^+ & \text{(iii)'} & (xy)^* = y^*(xy^*)^*y^* \\ \text{(iv)} & x^{++} = x^+ & \text{(iv)'} & x^{**} = x^* \\ \text{(v)} & x^{+*} = x^+ & \text{(v)'} & x^{*+} = x^* \end{array}$$

DRC Conditions

In this case, we denote $P(S) = \{x^+ \mid x \in S\}$ and call it *the set of projections* of S . By (v) and (v)', we have

$$(3) \quad P(S) = \{x^+ \mid x \in S\} = \{x^* \mid x \in S\}.$$

Observe that DRC semigroups were termed *York semigroups* by Jones [28], *DR semigroups satisfying the congruence conditions* by Stokes [49] and *reduced E-Fountain semigroups* by Stein [46, 47], respectively. Recently, Wang [59] introduced DRC-ample conditions for DRC semigroups by which DRC-restriction semigroups are discriminated from DRC semigroups. A DRC semigroup $(S, \cdot, +, *)$ is called a DRC -restriction semigroup if it also satisfies the following axioms:

$$\text{(vi)} \quad x(yx)^* = (yx^+)^*x \quad \text{(vi)'} \quad (xy)^+x = x(x^*y)^+$$

DRC-ample conditions

In view of the identities (ii)' and (ii), (vi) (respectively, (vi')) is exactly the axiom $x(y^*x)^* = (y^*x^+)^*x$ (respectively, $(xy^+)^+x = x(x^*y^+)^+$). This implies that (vi) and (vi)' are equivalent to the conditions that

$$(4) \quad x(ex)^* = (ex^+)^*x \text{ and } (xe)^+x = x(x^*e)^+ \text{ for all } x \in S \text{ and } e \in P(S),$$

respectively.

Moreover, a regular version of DRC-restriction semigroups also provided in Wang [59]. A unary semigroup (S, \cdot, \circ) is called a *generalized regular \circ -semigroup* if the following axioms hold:

$$(5) \quad xx^\circ x = x, x^{\circ\circ} = x, (xy)^\circ = (xy)^\circ xx^\circ, (xy)^\circ = y^\circ y(xy)^\circ, (xy)^\circ x = (x^\circ xy)^\circ.$$

Obviously, regular \circ -semigroups are generalized regular \circ -semigroups. But the converse does not hold (see [59, Example 3.3]). Let (S, \cdot, \circ) be a generalized regular \circ -semigroup.

Define $x^+ = xx^\circ$ and $x^* = x^\circ x$ for all $x \in S$. By [59, Proposition 3.7], $(S, \cdot, +, *)$ forms a DRC-restriction semigroup. DRC semigroups and their generalizations have been studied by several authors, see [28, 29, 46, 47, 48, 49, 51, 52, 57, 59, 60] for example.

On the relationship among Ehresman (respectively, restriction), P -Ehresmann (respectively, P -restriction) and DRC (respectively, DRC-restriction) semigroups, we have the following result.

Proposition 1.2 ([17, 26, 59]). *The following statements hold:*

- (1) *A P -Ehresmann (respectively, P -restriction) semigroup is exactly a DRC (respectively, DRC-restriction) semigroup in which $(ef)^+ = efe = (fe)^*$ for all projections e and f .*
- (2) *An Ehresman (respectively, restriction) semigroup is exactly a DRC (respectively, DRC-restriction) semigroup in which $(ef)^+ = ef = fe = (fe)^*$ for all projections e and f .*

In view of the previous statements, we have the following hierarchy of the above mentioned classes of semigroups, where “ $A \rightarrow B$ ” denote A is a subclass of B .




Diagram 1: A hierarchy of several subclasses of DRC-semigroups

According to Gould [17] and Hollings [21], in the study of structure theory of inverse semigroups three major approaches have emerged. These are: *categorical approach* developed in the texts [11, 12, 36, 37, 40, 44, 45]), *fundamental approach* developed in the texts [38, 39] and *covering approach* developed in the texts [34, 35], respectively. The main result on inverse semigroups obtained by using categorical approach was formulated by Lawson in [30] as the following Ehresmann–Schein–Nambooripad Theorem, due to its varied authorship.

Theorem 1.3 ([30]). *The category of inverse semigroups together with homomorphisms is isomorphic to the category of inductive groupoids together with inductive functors.*

On the other hand, the main result on inverse semigroups obtained by using fundamental approach is the theorem below.

Theorem 1.4 ([22]). *Let S be an inverse semigroup and $T_{E(S)}$ be the Munn semigroup of $E(S)$. Then there exists a homomorphism $\phi : S \rightarrow T_{E(S)}$ whose kernel is the maximum idempotent-separating congruence on S . Moreover, S is fundamental if and only if S is isomorphic to a full inverse subsemigroup of $T_{E(S)}$.*

Since our present paper does not involve covering approach, we will no longer list the results on inverse semigroups obtained by using this approach. Theorems 1.3 and 1.4 have been generalized in various directions, see for example [2, 7, 8, 9, 13, 14, 15, 16, 17, 18, 20, 23, 24, 26, 29, 31, 32, 40, 50, 51, 55, 56, 57, 59, 60, 61]. More specifically, for regular case, Nambooripad [40] generalized this two theorems to general regular semigroups; Imaoka and Fujiwara [24] and East and Azeef Muhammed [9] generalized Theorem 1.3 to locally

inverse regular \circ -semigroups and general regular \circ -semigroups respectively; East and Azeef Muhammed [9], Imaoka [23] and Jones [26] generalized Theorem 1.4 to regular \circ -semigroups. For non-regular cases, Fountain [14], Fountain Gomes and Gould [15] and Gomes and Gould [16] generalized Theorem 1.4 to Ehresmann semigroups and restriction semigroups; Jones [26, 29] and Wang [59] generalized Theorem 1.4 to P -restriction, P -Ehresmann, DRC-restriction and DRC semigroups, respectively. On the other hand, Lawson [31, 32] generalized Theorem 1.3 to Ehresmann semigroups and restriction semigroups; Wang [55, 56] generalized Theorem 1.3 to a special class of P -restriction and P -Ehresmann semigroups, respectively. Recently, Wang [57] generalized Theorem 1.3 to the whole class of DRC semigroups and so generalized Theorem 1.3 to all subclasses of DRC semigroups mentioned in Diagram 1.

Although Wang [57] investigated all subclasses of DRC semigroups mentioned in Diagram 1 by so-called categorical approach, we observe that in this process he actually used some kinds of generalized small categories or generalized groupoids introduced there instead of the usual small categories or groupoids. More recently, among other things East and Azeef Muhammed [8, 9] generalized Theorem 1.3 to the whole class of regular \circ -semigroups by using the theory of chain projection ordered groupoids established there rather than the theory of generalized groupoids of any kind. In particular, they proved the following theorem.

Theorem 1.5 ([8]). *The category of regular \circ -semigroups together with \circ -homomorphisms is isomorphic to the category of chain projection ordered groupoids together with chain projection ordered functors.*

In the end of their paper [8], East and Azeef Muhammed proposed the following natural problem.

Problem 1.6 (The Fourth Problem in [8]). *Can the theory developed in [8] be applied to other categories of “projection-based” semigroups such as P -Ehresmann semigroups and DRC semigroups?*

In this paper, we solve the above Problem 1.6 for DRC-restriction semigroups. We generalize the theory of chain projection ordered groupoids established by East and Azeef Muhammed in [8, 9] to that of chain projection ordered categories by which DRC-restriction semigroups are characterized. More specifically, we prove the category of DRC-restriction semigroups together with $(2,1,1)$ -homomorphisms is isomorphic to the category of chain projection ordered categories together with chain projection ordered functors. Some special cases such as P -restriction semigroups and generalized regular \circ -semigroups are also considered. Finally, some applications of the main theorem are given. Our results generalize and enrich the corresponding results obtained by East and Azeef Muhammed in [8, 9].

The paper is organized as follows. Section 2 gives some fundamental notions and results on ordered categories and groupoids, and Section 3 states some items on two-sided projection algebras. In Section 4, based on the results given in Sections 2 and 3, we establish the theory of chain projection ordered categories and construct the category of chain projection ordered categories together with chain projection ordered functors. Section 5 is devoted to proving that a DRC-restriction semigroup give rises to a chain projection ordered category, and Section 6 proves our category isomorphism theorem. In the final two sections, we consider two applications of our main result. Section 7 uses so-called chain semigroups to construct the free DRC-restriction semigroup over any arbitrary strong two-sided projection

algebra and Section 8 uses projection categories (groupoids) to reobtain the fundamental representation theorem for DRC-restriction semigroups appeared firstly in Wang [59].

2. ORDERED CATEGORIES

In this setion, we give some basic results on ordered categories and ordered groupoids. Assume that C is a nonempty set, “ \circ ” is a partial binary operation on C and

$$\mathbf{d} : C \rightarrow C, x \mapsto \mathbf{d}(x), \quad \mathbf{r} : C \rightarrow C, x \mapsto \mathbf{r}(x)$$

are maps. According to [31], $(C, \circ, \mathbf{d}, \mathbf{r})$ is called a (*small*) *category*, if the following conditions hold:

- (C1) For all $x, y \in C$, $x \circ y$ is defined if and only if $\mathbf{r}(x) = \mathbf{d}(y)$, in this case, $\mathbf{d}(x \circ y) = \mathbf{d}(x)$ and $\mathbf{r}(x \circ y) = \mathbf{r}(y)$.
- (C2) For all $x, y, z \in C$, if $x \circ y$ and $y \circ z$ are defined, then $(x \circ y) \circ z = x \circ (y \circ z)$.
- (C3) For all $x \in C$, $\mathbf{d}(x) \circ x$ and $x \circ \mathbf{r}(x)$ are defined, and $\mathbf{d}(x) \circ x = x = x \circ \mathbf{r}(x)$.

Remark 2.1. Let $(C, \cdot, \mathbf{d}, \mathbf{r})$ be a category and $x \in C$. Then $\mathbf{d}(x) \circ x$ and $x \circ \mathbf{r}(x)$ are defined by (C3), and so $\mathbf{r}(\mathbf{d}(x)) = \mathbf{d}(x)$ and $\mathbf{r}(x) = \mathbf{d}(\mathbf{r}(x))$ by (C1). This implies that $\{\mathbf{d}(x) \mid x \in C\} = \{\mathbf{r}(x) \mid x \in C\}$. We call

$$P_C = \{\mathbf{d}(x) \mid x \in C\} = \{\mathbf{r}(x) \mid x \in C\}$$

the *set of objects* of C . Clearly, $\mathbf{d}(e) = e = \mathbf{r}(e)$ for all $e \in P_C$.

Let $(C, \circ, \mathbf{d}, \mathbf{r})$ be a category and “ \leq ” be a partial order on C . Then $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ is called an *ordered category* if for all $a, b, c, d \in C$ and $p, q \in P_C$, the following statements hold:

- (O1) If $a \leq b$, then $\mathbf{d}(a) \leq \mathbf{d}(b)$ and $\mathbf{r}(a) \leq \mathbf{r}(b)$.
- (O2) If $a \leq b$, $c \leq d$, $\mathbf{r}(a) = \mathbf{d}(c)$, $\mathbf{r}(b) = \mathbf{d}(d)$, then $a \circ c \leq b \circ d$.
- (O3) If $p \leq \mathbf{d}(a)$, then exists a unique element $u \in C$ such that $u \leq a$ and $\mathbf{d}(u) = p$.
- (O4) If $q \leq \mathbf{r}(a)$, then exists a unique element $v \in C$ such that $v \leq a$ and $\mathbf{r}(v) = q$.

In this case, the elements u, v in (O3) and (O4) are denoted by $u = {}_p \downarrow a$ and $v = a \downarrow_q$, and call them the *left restriction* and *right restriction* on p and q of a , respectively.

Lemma 2.2. *Let $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category, $a, b \in C$ and $p, q \in P_C$.*

- (1) *If $p \leq \mathbf{d}(a)$, then ${}_p \downarrow a \leq a$, $\mathbf{d}({}_p \downarrow a) = p$ and $\mathbf{r}({}_p \downarrow a) \leq \mathbf{r}(a)$.*
- (2) *If $q \leq \mathbf{r}(a)$, then $a \downarrow_q \leq a$, $\mathbf{r}(a \downarrow_q) = q$ and $\mathbf{d}(a \downarrow_q) \leq \mathbf{d}(a)$.*
- (3) *If $a \leq b$, then $a = {}_{\mathbf{d}(a)} \downarrow b$. In particular, ${}_{\mathbf{d}(a)} \downarrow a = a$.*
- (4) *If $a \leq b$, then $a = b \downarrow_{\mathbf{r}(a)}$. In particular, $a \downarrow_{\mathbf{r}(a)} = a$.*
- (5) *If $p \leq \mathbf{d}(a)$, then ${}_p \downarrow a = a \downarrow_{\mathbf{r}({}_p \downarrow a)}$.*
- (6) *If $q \leq \mathbf{r}(a)$, then $a \downarrow_q = {}_{\mathbf{d}(a \downarrow_q)} \downarrow a$.*
- (7) *If $p \leq \mathbf{d}(a)$, $\mathbf{r}(a) = \mathbf{d}(b)$, then ${}_p \downarrow (a \circ b) = {}_p \downarrow a \circ {}_{\mathbf{r}({}_p \downarrow a)} \downarrow b$.*
- (8) *If $q \leq \mathbf{r}(b)$, $\mathbf{r}(a) = \mathbf{d}(b)$, then $(a \circ b) \downarrow_q = a \downarrow_{\mathbf{d}(b \downarrow_q)} \circ b \downarrow_q$.*
- (9) *If $p \leq q \leq \mathbf{d}(a)$, then ${}_p \downarrow ({}_q \downarrow a) = {}_p \downarrow a$.*
- (10) *If $p \leq q \leq \mathbf{r}(a)$, then $(a \downarrow_q) \downarrow_p = a \downarrow_p$.*
- (11) *If $p, q \in P_C$ and $p \leq q$, then ${}_p \downarrow q = p$.*
- (12) *If $p, q \in P_C$ and $p \leq q$, then $q \downarrow_p = p$.*

Proof. We only need to show items (1), (3), (5), (7), (9) and (11) by symmetry.

(1) By the definition of ${}_p \downarrow a$ and (O3), we have ${}_p \downarrow a \leq a$ and $\mathbf{d}({}_p \downarrow a) = p$, and so $\mathbf{r}({}_p \downarrow a) \leq \mathbf{r}(a)$ by (O1).

(3) The fact $a \leq b$ and (O1) give $\mathbf{d}(a) \leq \mathbf{d}(b)$, and so b has a unique left restriction on $\mathbf{d}(a)$ by (O3). Since $a \leq b$ and $\mathbf{d}(a) = \mathbf{d}(a)$, we have $a = \mathbf{d}(a) \downarrow b$.

(5) (O3) gives $p \downarrow a \leq a$ and so we have $p \downarrow a = a \downarrow_{\mathbf{r}(p \downarrow a)}$ by (4) in the present lemma.

(7) By (O3) and (O1), we have $p \downarrow a \leq a$ and $\mathbf{r}(p \downarrow a) \leq \mathbf{r}(a) = \mathbf{d}(b)$. By the definitions of $p \downarrow (a \circ b)$ and $\mathbf{r}(p \downarrow a) \downarrow b$ and (O3),

$$p \downarrow (a \circ b) \leq a \circ b, \mathbf{r}(p \downarrow a) \downarrow b \leq b, \mathbf{r}(p \downarrow a) = \mathbf{d}(\mathbf{r}(p \downarrow a) \downarrow b), \mathbf{d}(p \downarrow (a \circ b)) = p,$$

this together with the fact $\mathbf{r}(a) = \mathbf{d}(b)$ and (O2) implies that $p \downarrow a \circ_{\mathbf{r}(p \downarrow a)} \downarrow b \leq a \circ b$. On the other hand, we have $\mathbf{d}(p \downarrow a \circ_{\mathbf{r}(p \downarrow a)} \downarrow b) = \mathbf{d}(p \downarrow a) = p$ by (C1) and (O3). By (3) in the present lemma, we obtain $p \downarrow a \circ_{\mathbf{r}(p \downarrow a)} \downarrow b = p \downarrow (a \circ b)$.

(9) Item (O3) gives that $p \downarrow (q \downarrow a) \leq q \downarrow a \leq a$ and $\mathbf{d}(p \downarrow (q \downarrow a)) = p$, and so $p \downarrow (q \downarrow a) = p \downarrow a$ by (3) in the present lemma.

(11) Since $p \leq q$ and $\mathbf{d}(p) = p$ by Remark 2.1, we have $p = p \downarrow q$ by (3) in the present lemma. \square

Let $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category and $p \in P_C$. Denote

$$p^\downarrow = \{q \in P_C \mid q \leq p\}.$$

Let $a \in C$. By (1) and (2) in Lemma 2.2, we have $\mathbf{r}(p \downarrow a) \leq \mathbf{r}(a)$ and $\mathbf{d}(a \downarrow_q) \leq \mathbf{d}(a)$. Thus we can define the maps ν_a and μ_a as follows:

$$(6) \quad \nu_a : \mathbf{d}(a)^\downarrow \rightarrow \mathbf{r}(a)^\downarrow, p \mapsto p\nu_a = \mathbf{r}(p \downarrow a); \quad \mu_a : \mathbf{r}(a)^\downarrow \rightarrow \mathbf{d}(a)^\downarrow, q \mapsto q\mu_a = \mathbf{d}(a \downarrow_q).$$

By using the maps ν_a and μ_a ($a \in S$), we can restate (7) and (8) in Lemma 2.2 as follows: If $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ is an ordered category, $a, b \in C$ and $p, q \in P_C$ with $\mathbf{r}(a) = \mathbf{d}(b)$, $p \leq \mathbf{d}(a)$ and $q \leq \mathbf{r}(b)$. Then

$$(7) \quad p \downarrow (a \circ b) = p \downarrow a \circ_{p\nu_a} \downarrow b, \quad (a \circ b) \downarrow_q = a \downarrow_{q\mu_b} \circ b \downarrow_q.$$

Lemma 2.3. *Let $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category and $a \in C$. Then ν_a and μ_a are mutually inverse bijections. In particular, if $e \in P_C$, then $\nu_e = \mu_e = \text{id}_{e^\downarrow}$.*

Proof. Let $p \leq \mathbf{d}(a)$ and denote $q = p\nu_a = \mathbf{r}(p \downarrow a)$. Then by (5) and (1) in Lemma 2.2,

$$p\nu_a\mu_a = q\mu_a = \mathbf{d}(a \downarrow_q) = \mathbf{d}(p \downarrow a) = p.$$

Thus $\nu_a\mu_a = \text{id}_{\mathbf{d}(a)^\downarrow}$. Dually, we can prove that $\mu_a\nu_a = \text{id}_{\mathbf{r}(a)^\downarrow}$. If $e \in P_C$, then by Remark 2.1, we have $\mathbf{d}(e) = \mathbf{r}(e) = e$. Let $p \in \mathbf{d}(e)^\downarrow = e^\downarrow$. Then $p \in P_C$ and $p \leq e$, and so $p\nu_e = \mathbf{r}(p \downarrow e) = \mathbf{r}(p) = p$ by Lemma 2.2 (11) and Remark 2.1 again. This shows that $\nu_e = \text{id}_{e^\downarrow}$. Moreover, $\mu_e = \nu_e^{-1} = \text{id}_{e^\downarrow}^{-1} = \text{id}_{e^\downarrow}$. \square

Lemma 2.4. *Let $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category, $a \in C$ and $e, f \in P_C$ with $e \leq \mathbf{d}(a)$, $f \leq \mathbf{r}(a)$. Then $\nu_{e \downarrow a} = \nu_a \downarrow_{e^\downarrow}$ and $\mu_{a \downarrow f} = \mu_a \downarrow_{f^\downarrow}$.*

Proof. It is easy to see that $\nu_{e \downarrow a}$ and $\nu_a \downarrow_{e^\downarrow}$ have the same domain e^\downarrow by Lemma 2.2 (1). Let $t \leq e$. By (6) and Lemma 2.2 (9), we have $t\nu_{e \downarrow a} = \mathbf{r}(t \downarrow (e \downarrow a)) = \mathbf{r}(t \downarrow a) = t\nu_a$. Dually, $\mu_{a \downarrow f} = \mu_a \downarrow_{f^\downarrow}$. \square

Lemma 2.5. *Let $(C, \cdot, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category, $a, b \in C$ and $\mathbf{r}(a) = \mathbf{d}(b)$. Then*

$$\nu_{a \circ b} = \nu_a \nu_b, \quad \mu_{a \circ b} = \mu_b \mu_a.$$

Proof. Let $p = \mathbf{d}(a), q = \mathbf{r}(a) = \mathbf{d}(b), r = \mathbf{r}(b)$. Then $\mathbf{d}(a \circ b) = \mathbf{d}(a) = p$ and $\mathbf{r}(a \circ b) = \mathbf{r}(b) = r$, and so

$$\nu_a : p^\downarrow \rightarrow q^\downarrow, \nu_b : q^\downarrow \rightarrow r^\downarrow, \nu_{a \circ b} : p^\downarrow \rightarrow r^\downarrow, \nu_a \nu_b : p^\downarrow \rightarrow r^\downarrow$$

Let $e \in p^\downarrow$ (i.e. $s \leq p$) and denote $t = s \nu_a = \mathbf{r}(s \downarrow a)$. By Lemma 2.2 (7), we have

$$s \nu_{a \circ b} = \mathbf{r}(s \downarrow (a \circ b)) = \mathbf{r}(s \downarrow a \circ t \downarrow b) = \mathbf{r}(t \downarrow b) = t \nu_b = s \nu_a \nu_b.$$

Thus $\nu_{a \circ b} = \nu_a \nu_b$. Dually, we can prove that $\mu_{a \circ b} = \mu_b \mu_a$. \square

A category $(C, \circ, \mathbf{d}, \mathbf{r})$ is called a *groupoid* if for all $a \in C$, there exists $a' \in C$ such that $a \circ a' = \mathbf{d}(a)$ and $a' \circ a = \mathbf{r}(a)$. In this case, for $a \in C$, we have $\mathbf{r}(a') = \mathbf{r}(a \circ a') = \mathbf{r}(\mathbf{d}(a)) = \mathbf{d}(a)$. Dual argument gives $\mathbf{d}(a') = \mathbf{r}(a)$. The above a' is unique. In fact, if $a'' \in C$ and $a \circ a'' = \mathbf{d}(a)$ and $a'' \circ a = \mathbf{r}(a)$, then $\mathbf{d}(a) = \mathbf{r}(a'')$, and so

$$a' = \mathbf{d}(a') \circ a' = \mathbf{r}(a) \circ a' = (a'' \circ a) \circ a' = a'' \circ \mathbf{d}(a) = a'' \circ \mathbf{r}(a'') = a''.$$

Denote the unique element a' by a^{-1} . An ordered category $(C, \circ, \mathbf{d}, \mathbf{r}, \leq)$ is called an *ordered groupoid* if $(C, \circ, \mathbf{d}, \mathbf{r})$ is a groupoid.

Lemma 2.6. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered groupoid and $a, b \in C$.*

- (1) $a \circ a^{-1} = \mathbf{d}(a), a^{-1} \circ a = \mathbf{r}(a), \mathbf{d}(a) = \mathbf{r}(a^{-1}), \mathbf{r}(a) = \mathbf{d}(a^{-1})$.
- (2) If $a \leq b$, then $a^{-1} \leq b^{-1}$.
- (3) If $p, q \in P_C$ and $p \leq \mathbf{d}(a), q \leq \mathbf{r}(a)$, then $a \downarrow_q = (q \downarrow a^{-1})^{-1}, p \downarrow a = (a^{-1} \downarrow_p)^{-1}$.
- (4) $\mu_a = \nu_{a^{-1}}$.

Proof. (1) This follows from the statements before Lemma 2.6.

(2) Let $a \leq b$. Then $a = \mathbf{d}(a) \downarrow b$ by Lemma 2.2 (3). Item (O1) gives $\mathbf{d}(a) \leq \mathbf{d}(b)$, and so $\mathbf{d}(a) = \mathbf{d}(a) \downarrow \mathbf{d}(b)$ by Lemma 2.2 (11). Thus

$$\mathbf{r}(a^{-1}) = \mathbf{d}(a) = \mathbf{d}(a) \downarrow \mathbf{d}(b) = \mathbf{d}(a) \downarrow (b \circ b^{-1}) = \mathbf{d}(a) \downarrow b \circ_{\mathbf{r}(a) \downarrow b} \downarrow b^{-1} = a \circ_{\mathbf{r}(a) \downarrow} b^{-1}$$

by item (1) in the present lemma and Lemma 2.2 (7). This implies that

$$\begin{aligned} a^{-1} &= a^{-1} \circ \mathbf{r}(a^{-1}) = a^{-1} \circ a \circ_{\mathbf{r}(a) \downarrow} b^{-1} = \mathbf{r}(a) \circ_{\mathbf{r}(a) \downarrow} b^{-1} \\ &= \mathbf{d}(\mathbf{r}(a) \downarrow b^{-1}) \circ_{\mathbf{r}(a) \downarrow} b^{-1} = \mathbf{r}(a) \downarrow b^{-1} \leq b^{-1} \end{aligned}$$

by Lemma 2.2 (1) and (C3).

(3) Since $q \leq \mathbf{r}(a) = \mathbf{d}(a^{-1})$, we have $q \downarrow a^{-1} \leq a^{-1}$, and so $(q \downarrow a^{-1})^{-1} \leq a$ by item (2) in the present lemma. By Lemma 2.2 (1), we get $\mathbf{r}((q \downarrow a^{-1})^{-1}) = \mathbf{d}(q \downarrow a^{-1}) = q$. By Lemma 2.2 (4), we have $(q \downarrow a^{-1})^{-1} = a \downarrow_q$. The other identity can be proved dually.

(4) By (6), we have

$$\nu_{a^{-1}} : \mathbf{d}(a^{-1})^\downarrow \rightarrow \mathbf{r}(a^{-1})^\downarrow, q \mapsto \mathbf{r}(q \downarrow a^{-1}).$$

By (1) and (3) in the present lemma,

$$\nu_{a^{-1}} : \mathbf{r}(a)^\downarrow \rightarrow \mathbf{d}(a)^\downarrow, q \mapsto \mathbf{d}((q \downarrow a^{-1})^{-1}) = \mathbf{d}(a \downarrow_q).$$

This implies that $\nu_{a^{-1}} = \mu_a$ by (6) again. \square

3. TWO-SIDED PROJECTION ALGEBRAS

In this section, we shall give some basic notions and facts on two-sided projection algebras introduced by Jones in [26]. Let P be a non-empty set and “ \times ” be a binary operation on P . Then (P, \times) is called a *left projection algebra* if the following axioms hold:

- (L1) $e \times e = e$.
- (L2) $(e \times f) \times e = e \times f$.
- (L3) $e \times (f \times g) = (e \times f) \times (f \times g)$.
- (L4) $(e \times f) \times g = (e \times f) \times (e \times g)$.

Dually, let P be a non-empty set and “ \star ” be a binary operation on P . Then (P, \star) is called a *right projection algebra* if the following axioms hold:

- (R1) $e \star e = e$.
- (R2) $(f \star e) = e \star (f \star e)$.
- (R3) $(g \star f) \star e = (g \star f) \star (f \star e)$.
- (R4) $g \star (f \star e) = (g \star e) \star (f \star e)$.

Let P be a non-empty set and “ \times ” and “ \star ” be binary operations on P . Then (P, \times, \star) is called a *two-sided projection algebra*, or *projection algebra* for simplicity, if the following conditions hold:

- (P1) (P, \times) forms a left projection algebra.
- (P2) (P, \star) forms a right projection algebra.
- (P3) For all $e, f, g \in P$, we have $(e \star (f \times g)) \star g = (e \star f) \star g$ and $g \times ((g \star f) \times e) = g \times (f \times e)$.
- (P4) For all $e, f \in P$, we have $(e \times f) \star e = e \times f$ and $f \times (e \star f) = e \star f$.

Let (P, \times, \star) be a projection algebra. Define a relation “ \leq_P ” on P as follows: For all $e, f \in P$,

$$e \leq_P f \iff e = f \times e.$$

Then “ \leq_P ” is a partial order by [26, Lemma 5.2]. Recall from [59] that a projection algebra (P, \times, \star) is *strong* if the following axioms hold:

- (SP1) $e \times (((e \times f) \times g) \star f) = (e \times f) \times g$.
- (SP2) $(f \times (g \star (f \star e))) \star e = g \star (f \star e)$.

It is easy to see that (SP1) (respectively, (SP2)) is equivalent to the following condition: For all $e, f, g \in P$,

$$(8) \quad g \leq_P e \times f \implies e \times (g \star f) = g \quad (\text{respectively, } g \leq_P f \star e \implies (f \times g) \star e = g).$$

A strong projection algebra (P, \times, \star) is called

- *symmetric* if $e \times f = f \star e$ for all $e, f \in P$, and
- *commutative* if $e \times f = f \times e$ for all $e, f \in P$.

Lemma 3.1. *Let (P, \times, \star) be a projection algebra and $e, f, g \in P$.*

- (1) $e \leq_P f$ if and only if $e = e \times f = f \times e$. Moreover, $e \times f \leq_P e$.
- (2) $e \leq_P f$ if and only if $e = e \star f = f \star e$. Moreover, $e \star f \leq_P f$.
- (3) $(e \times f) \star f = e \star f$ and $e \times (e \star f) = e \times f$.
- (4) $(e \times f) \times (e \star f) = e \times f$ and $(e \times f) \star (e \star f) = e \star f$.
- (5) If $e \leq_P f$, then $g \times e \leq_P g \times f$ and $e \star g \leq_P f \star g$.

Proof. Items (1)–(4) follow from [56, Lemma 3.2], and (5) follows from [59, Lemma 5.2], respectively. \square

Let (P, \times, \star) be a projection algebra. For each $e \in P$, define

$$e^\downarrow = \{x \in P \mid x \leq_P e\}.$$

By (1) and (2) in Lemma 3.1, we have

$$(9) \quad e^\downarrow = \{e \times p \mid p \in P\} = \{p \star e \mid p \in P\}.$$

Let $p \in P$. Define θ_p and δ_p as follows:

$$(10) \quad \theta_p : P \rightarrow P, q \mapsto q\theta_p = q \star p, \quad \delta_p : P \rightarrow P, q \mapsto q\delta_p = p \times q.$$

Again by (1) and (2) in Lemma 3.1, for all $p, q \in P$ we have

$$(11) \quad p \leq_P q \iff p\theta_q = q\theta_p = p \iff p\delta_q = q\delta_p = p.$$

Moreover, (9) gives that

$$(12) \quad \text{im}\theta_p = \{q \star p \mid q \in P\} = p^\downarrow = \{p \times q \mid q \in P\} = \text{im}\delta_p.$$

Lemma 3.2. *Let (P, \times, \star) be a projection algebra and $p \leq_P q$. Then*

$$\theta_p = \theta_p\theta_q = \theta_q\theta_p = \theta_p\delta_q, \quad \delta_p = \delta_p\delta_q = \delta_q\delta_p = \delta_p\theta_q.$$

Proof. Let $t \in P$. By Lemma 3.1 (2), we have $t \star p \leq_P p$ and $(t \star p) \star p = t \star p$. Since $p \leq_P q$, it follows that $p \star q = p$ and $q \times (t \star p) = t \star p$ by (2) and (1) in Lemma 3.1. Firstly, by (R3) we have

$$t\theta_p\theta_q = (t \star p) \star q \stackrel{(R3)}{=} (t \star p) \star (p \star q) = (t \star p) \star p = t \star p = t\theta_p.$$

Secondly, by (R4) it follows that

$$t\theta_q\theta_p = (t \star q) \star p = (t \star q) \star (p \star q) \stackrel{(R4)}{=} t \star (p \star q) = t \star p = t\theta_p.$$

Finally, $t\theta_p\delta_q = q \times (t \star p) = t \star p = t\theta_p$. Thus $\theta_p = \theta_q\theta_p = \theta_p\theta_q = \theta_p\delta_q$. Dually, we have $\delta_p = \delta_p\delta_q = \delta_q\delta_p = \delta_p\theta_q$. \square

Let (P, \times, \star) be a projection algebra. Define a relation \mathcal{F}_P on P as follows: For all $p, q \in P$,

$$(13) \quad p\mathcal{F}_P q \iff p = q\delta_p, q = p\theta_q.$$

Let $p, q \in P$. Then $q\delta_p = p \times q$ and $p\theta_q = p \star q$ by the definitions of δ_p and θ_q . Thus

$$(14) \quad p\mathcal{F}_P q \text{ if and only if } p \times q = p \text{ and } p \star q = q.$$

Lemma 3.3. *Let (P, \times, \star) be a projection algebra and $p, q \in P$. Then $q\delta_p \leq_P p, p\theta_q \leq_P q$ and $q\delta_p \mathcal{F}_P p\theta_q$.*

Proof. Firstly, we have $q\delta_p \leq_P p$ and $p\theta_q \leq_P q$ by the facts that $q\delta_p = p \times q, p\theta_q = p \star q$ and (1) and (2) in Lemma 3.1. On the other hand,

$$\begin{aligned} (q\delta_p) \times (p\theta_q) &= (p \times q) \times (p \star q) = p \times q = q\delta_p, \\ (q\delta_p) \star (p\theta_q) &= (p \times q) \star (p \star q) = p \star q = p\theta_q \end{aligned}$$

by Lemma 3.1 (4). Thus $q\delta_p \mathcal{F}_P p\theta_q$. \square

Lemma 3.4. *Let (P, \times, \star) be a projection algebra and $p, q, r \in P$.*

- (1) *If $p \leq_P q$ and $q = q \times r$, then $p = p \times r$.*
- (2) *If $p = p \times q$ and $q \leq_P r$, then $p = p \times r$.*

Proof. By the facts that $p \leq_P q$, $q = q \times r$ and Lemma 3.1 (1), we obtain $p = p \times q = q \times p$ and $q = q \times r$. This together with (L4) gives that

$$p = p \times q = (q \times p) \times (q \times r) \stackrel{(L4)}{=} (q \times p) \times r = p \times r.$$

This proves (1). On the other hand, by the facts $p = p \times q$, $q \leq_P r$ and Lemma 3.1 (1), we obtain $p = p \times q$, $q = q \times r = r \times q$. This together with (L3), (L4) and (L2) implies that

$$\begin{aligned} p &= p \times q = p \times (r \times q) \stackrel{(L3)}{=} (p \times r) \times (r \times q) \\ &= (p \times r) \times q \stackrel{(L4)}{=} (p \times r) \times (p \times q) = (p \times r) \times p \stackrel{(L2)}{=} p \times r. \end{aligned}$$

This shows (2). \square

Lemma 3.5. *Let (P, \times, \star) be a projection algebra, $p_1, p_2, \dots, p_k, q \in P$ and $p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k$. Then*

$$p_k \delta_{p_{k-1}} \delta_{p_{k-2}} \dots \delta_{p_1} = p_1, \quad p_1 \theta_{p_2} \theta_{p_3} \dots \theta_{p_k} = p_k.$$

Proof. In view of the fact that $p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k$, (13) and (14), we have

$$\begin{aligned} p_k \delta_{p_{k-1}} \delta_{p_{k-2}} \dots \delta_{p_1} &= (p_{k-1} \times p_k) \delta_{p_{k-2}} \dots \delta_{p_1} \\ &= p_{k-1} \delta_{p_{k-2}} \dots \delta_{p_1} = (p_{k-2} \times p_{k-1}) \delta_{p_{k-3}} \dots \delta_{p_1} \\ &= p_{k-2} \delta_{p_{k-3}} \dots \delta_{p_1} = \dots = p_2 \delta_{p_1} = p_1 \times p_2 = p_1. \end{aligned}$$

The other identity can be proved dually. \square

Let (P, \times, \star) and (P', \times, \star) be two projection algebras. A map $\psi : P \rightarrow P'$ is called a *homomorphism* if

$$(y \delta_x) \psi = (x \times y) \psi = (x \psi) \times (y \psi) = (y \psi) \delta_{x \psi}, \quad (x \theta_y) \psi = (x \star y) \psi = (x \psi) \star (y \psi) = (x \psi) \theta_{y \psi}$$

for all $x, y \in P$.

Lemma 3.6. *Let (P, \times, \star) and (P', \times, \star) be two projection algebras, $\psi : P \rightarrow P'$ be a homomorphism and $p, q \in P$.*

- (1) *If $p \leq_P q$, then $p \psi \leq_{P'} q \psi$.*
- (2) *If $p \mathcal{F}_P q$, then $p \psi \mathcal{F}_{P'} q \psi$.*

Proof. Let $p \leq_P q$. Then $q \times p = p$. Since ψ is homomorphism, we have $(q \psi) \times (p \psi) = p \psi$, and so $p \psi \leq_{P'} q \psi$. This gives (1), and (2) can be proved similarly. \square

Lemma 3.7. *Let (P, \times, \star) be a strong projection algebra and $p, s \in P$. Then $\delta_{p \theta_s} = \delta_s \delta_p \theta_s$ and $\theta_{p \delta_s} = \theta_s \theta_p \delta_s$.*

Proof. Let $t \in P$. Then $(p \star s) \times t \leq_P p \star s$ by Lemma 3.1 (1), and so $(p \star s) \times t = (p \times ((p \star s) \times t)) \star s$ by (8). Now, (P3) gives that

$$t \delta_{p \theta_s} = (p \star s) \times t = (p \times ((p \star s) \times t)) \star s \stackrel{(P3)}{=} (p \times (s \times t)) \star s = t \delta_s \delta_p \theta_s.$$

Dually, we have $\theta_{p \delta_s} = \theta_s \theta_p \delta_s$. \square

Using Lemma 3.7 and mathematical induction, we have the following corollary.

Corollary 3.8. *Let (P, \times, \star) be a strong projection algebra and $p_1, p_2, \dots, p_k, q \in P$. Then*

$$\begin{aligned} \delta_{q \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}} &= \delta_{p_k} \dots \delta_{p_1} \delta_q \theta_{p_1} \dots \theta_{p_k}, \\ \theta_{q \delta_{p_k} \delta_{p_{k-1}} \dots \delta_{p_1}} &= \theta_{p_1} \dots \theta_{p_k} \theta_q \delta_{p_k} \dots \delta_{p_1}. \end{aligned}$$

4. CHAIN PROJECTION ORDERED CATEGORIES

In this section, we develop a theory of chain projection ordered categories, which generalizes the theory of chain projection ordered groupoids provided by East and Azeef Muhammed in [8] recently.

Definition 4.1. Let (P, \times, \star) be a strong projection algebra and k be a positive integer. Assume that $p_1, p_2, \dots, p_k \in P$ and $p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k$. Then we call $\mathbf{p} = (p_1, p_2, \dots, p_k)$ a P -path in P with length k from p_1 to p_k . Denote the set of all P -paths in P by $\mathcal{P}(P)$. For all $p \in P$, we identify P -path (p) with length 1 to p . Under this assumption, P is a subset of $\mathcal{P}(P)$.

Let (P, \times, \star) be a strong projection algebra. For

$$\mathbf{p} = (p_1, p_2, \dots, p_k), \mathbf{q} = (q_1, q_2, \dots, q_l) \in \mathcal{P}(P),$$

we define $\mathbf{d}(\mathbf{p}) = p_1, \mathbf{r}(\mathbf{p}) = p_k$, and if $\mathbf{r}(\mathbf{p}) = p_k = q_1 = \mathbf{d}(\mathbf{q})$, we let

$$\mathbf{p} \circ \mathbf{q} = (p_1, p_2, \dots, p_k = q_1, q_2, \dots, q_l).$$

It is easy to see that $(\mathcal{P}(P), \circ, \mathbf{d}, \mathbf{r})$ is a small category, $P_{\mathcal{P}(P)} = P$ and $\mathbf{d}(\mathbf{p}) = p = \mathbf{r}(\mathbf{p})$ for all $p \in P$. We call this category the *path category* of (P, \times, \star) . Let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$, $q \in P$ and $q \leq_P \mathbf{d}(\mathbf{p}) = p_1$. Denote

$$(15) \quad q \downarrow \mathbf{p} = (q_1, q_2, \dots, q_k), \text{ where } q_1 = q, q_i = q_{i-1} \theta_{p_i} = q_{i-1} \star p_i, i = 2, 3, \dots, k,$$

and call it the *left restriction* of \mathbf{p} on q . In this case, observe that $q \leq_P p_1$, it follows that $q_1 = q = q \star p_1 = q \theta_{p_1}$ by Lemma 3.1 (2). Combining (15) and Lemma 3.1 (2), we have

$$(16) \quad q_i = q \theta_{p_1} \theta_{p_2} \dots \theta_{p_i}, \quad q_i \leq_P p_i, \quad i = 1, 2, \dots, k.$$

Dually, let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$, $q \in P$ and $q \leq_P \mathbf{r}(\mathbf{p}) = p_k$. Denote

$$(17) \quad \mathbf{p} \downarrow_q = (q_1, q_2, \dots, q_k), \text{ where } q_k = q, q_i = q_{i+1} \delta_{p_i} = p_i \times q_{i+1}, i = k-1, k-2, \dots, 1$$

and call it the *right restriction* of \mathbf{p} on q . In this case, observe that $q \leq_P p_k$, it follows that $q_k = q = p_k \times q = q \delta_{p_k}$ by Lemma 3.1 (1). Combining (17) and Lemma 3.1 (1), we have

$$(18) \quad q_i = q \delta_{p_k} \dots \delta_{p_i}, \quad q_i \leq_P p_i, \quad i = k, k-1, \dots, 1.$$

Lemma 4.2. Let (P, \times, \star) be a strong projection algebra, $q, r \in P$ and $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$.

- (1) If $q \leq_P \mathbf{d}(\mathbf{p})$, then $q \downarrow \mathbf{p} \in \mathcal{P}(P)$, $\mathbf{d}(q \downarrow \mathbf{p}) = q$ and $\mathbf{r}(q \downarrow \mathbf{p}) \leq_P \mathbf{r}(\mathbf{p})$.
- (2) If $q \leq_P \mathbf{r}(\mathbf{p})$, then $\mathbf{p} \downarrow_q \in \mathcal{P}(P)$, $\mathbf{r}(\mathbf{p} \downarrow_q) = q$ and $\mathbf{d}(\mathbf{p} \downarrow_q) \leq_P \mathbf{d}(\mathbf{p})$.
- (3) If $q = \mathbf{d}(\mathbf{p})$, then $q \downarrow \mathbf{p} = \mathbf{p}$.
- (4) If $q = \mathbf{r}(\mathbf{p})$, then $\mathbf{p} \downarrow_q = \mathbf{p}$.
- (5) If $r \leq_P q \leq_P \mathbf{d}(\mathbf{p})$, then $r \downarrow (q \downarrow \mathbf{p}) = r \downarrow \mathbf{p}$.
- (6) If $r \leq_P q \leq_P \mathbf{r}(\mathbf{p})$, then $(\mathbf{p} \downarrow_q) \downarrow_r = \mathbf{p} \downarrow_r$.
- (7) If $r \leq_P \mathbf{d}(\mathbf{p})$, then $r \downarrow \mathbf{p} = \mathbf{p} \downarrow_{\mathbf{r}(r \downarrow \mathbf{p})}$.
- (8) If $r \leq_P \mathbf{r}(\mathbf{p})$, then $\mathbf{p} \downarrow_r = \mathbf{d}(\mathbf{p} \downarrow_r) \downarrow \mathbf{p}$.

Proof. We only need to show items (1), (3), (5) and (7) by symmetry.

(1) Let $q \downarrow \mathbf{p} = (q_1, q_2, \dots, q_k)$, where

$$q_1 = q, q_i = q_{i-1} \theta_{p_i} = q_{i-1} \star p_i, i = 2, 3, \dots, k.$$

Since $\mathbf{p} \in \mathcal{P}(P)$, we have $p_i \mathcal{F}_P p_{i+1}$, that is,

$$p_i \times p_{i+1} = p_i, p_{i+1} = p_i \star p_{i+1}, i = 1, 2, \dots, k-1.$$

Let $i \in \{1, 2, \dots, k-1\}$. By (R4) and (R1), we have

$$q_i \theta_{q_{i+1}} = q_i \star q_{i+1} = q_i \star (q_i \star p_{i+1}) = (q_i \star p_{i+1}) \star (q_i \star p_{i+1}) = q_i \star p_{i+1} = q_{i+1}.$$

On the other hand, by the fact that $q_{i+1} = q_i \star p_{i+1}$ and Lemma 3.1 (3), we have

$$q_{i+1} \delta_{q_i} = q_i \times q_{i+1} = q_i \times (q_i \star p_{i+1}) = q_i \times p_{i+1}.$$

Observe (16) and the fact that $p_i \mathcal{F}_P p_{i+1}$, it follows that $q_i \leq_P p_i = p_i \times p_{i+1}$. This together with (8) gives that $q_i = p_i \times (q_i \star p_{i+1})$. Combing (L4) and (L2), we obtain

$$\begin{aligned} q_{i+1} \delta_{q_i} &= q_i \times p_{i+1} = (p_i \times (q_i \star p_{i+1})) \times p_{i+1} \\ &= (p_i \times (q_i \star p_{i+1})) \times (p_i \times p_{i+1}) = (p_i \times (q_i \star p_{i+1})) \times p_i = p_i \times (q_i \star p_{i+1}) = q_i. \end{aligned}$$

This implies that $q_i \mathcal{F}_P q_{i+1}$, and so $q \downarrow \mathbf{p} \in \mathcal{P}(P)$. By the definition of $q \downarrow \mathbf{p}$, we have $\mathbf{d}(q \downarrow \mathbf{p}) = q_1 = q$. Moreover, (16) gives that $\mathbf{r}(q \downarrow \mathbf{p}) = q_k \leq_P p_k = \mathbf{r}(\mathbf{p})$.

(3) Let $\mathbf{p} = (p_1, p_2, \dots, p_k)$. Then $q = \mathbf{d}(\mathbf{p}) = p_1$, and so $q \downarrow \mathbf{p} = (q_1, q_2, \dots, q_k)$, where

$$q_i = q \theta_{p_1} \dots \theta_{p_i}, i = 1, 2, \dots, k.$$

Obviously, $q_1 = q = p_1$. Assume that $q_{i-1} = p_{i-1}$, $i \geq 2$. Since $\mathbf{p} \in \mathcal{P}(P)$, it follows that $p_{i-1} \mathcal{F}_P p_i$, and so $p_{i-1} \star p_i = p_i$. By (15), $q_i = q_{i-1} \star p_i = p_{i-1} \star p_i = p_i$. By mathematical induction, the result follows.

(5) Let

$$\begin{aligned} \mathbf{p} &= (p_1, p_2, \dots, p_k), q \downarrow \mathbf{p} = (q_1, q_2, \dots, q_k), \\ r \downarrow \mathbf{p} &= (r_1, r_2, \dots, r_k), r \downarrow (q \downarrow \mathbf{p}) = (s_1, s_2, \dots, s_k). \end{aligned}$$

By (15) and (16),

$$\begin{aligned} q_i &= q \theta_{p_1} \dots \theta_{p_i} = q_{i-1} \star p_i, r_i = r \theta_{p_1} \dots \theta_{p_i} = r_{i-1} \star p_i, s_i = r \theta_{q_1} \dots \theta_{q_i} = s_{i-1} \star q_i, \\ (19) \quad q_i &\leq_P p_i, r_i \leq_P p_i, s_i \leq_P q_i, i = 1, 2, \dots, k. \end{aligned}$$

When $i = 1$, $r_1 = r = s_1$. Let $r_{i-1} = s_{i-1}$. Then the fact $s_{i-1} \leq_P q_{i-1}$ gives $s_{i-1} = s_{i-1} \star q_{i-1}$ by Lemma 3.1 (2). This together with the fact $q_i = q_{i-1} \star p_i$ and (R3) implies that

$$s_i = s_{i-1} \star q_i = (s_{i-1} \star q_{i-1}) \star (q_{i-1} \star p_i) = (s_{i-1} \star q_{i-1}) \star p_i = s_{i-1} \star p_i = r_{i-1} \star p_i = r_i.$$

By mathematical induction, the result follows.

(7) Let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$ and $r \leq_P \mathbf{d}(\mathbf{p})$. By (16),

$$r \downarrow \mathbf{p} = (r, r \theta_{p_2}, \dots, r \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}).$$

Let $t = \mathbf{r}(r \downarrow \mathbf{p}) = r \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}$. Then $t \leq_P \mathbf{r}(\mathbf{p}) = p_k$ by item (1). In view of (18),

$$\mathbf{p} \downarrow t = (t \delta_{p_k} \delta_{p_{k-1}} \dots \delta_{p_1}, \dots, t \delta_{p_k} \delta_{p_{k-1}}, t \delta_{p_k}).$$

In the following statements, we will show $r \downarrow \mathbf{p} = \mathbf{p} \downarrow t$. In fact, by the fact that $t \leq_P \mathbf{r}(\mathbf{p}) = p_k$ and (11), we have $t \delta_{p_k} = t = r \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}$. On the other hand, by (16) and the fact that $p_{k-1} \mathcal{F}_P p_k$, we have

$$r \theta_{p_1} \theta_{p_2} \dots \theta_{p_{k-1}} \leq_P p_{k-1} = p_{k-1} \times p_k.$$

This together with (8) gives that

$$r \theta_{p_1} \theta_{p_2} \dots \theta_{p_k} \delta_{p_{k-1}} = p_{k-1} \times ((r \theta_{p_1} \theta_{p_2} \dots \theta_{p_{k-1}}) \star p_k) = r \theta_{p_1} \theta_{p_2} \dots \theta_{p_{k-1}}.$$

By mathematical induction, we have

$$r\theta_{p_1}\theta_{p_2}\cdots\theta_{p_i} = r\theta_{p_1}\theta_{p_2}\cdots\theta_{p_k}\delta_{p_k}\cdots\delta_{p_i}, \quad i = k, k-1, \dots, 1.$$

Thus $r\downarrow\mathbf{p} = \mathbf{p}|_t = \mathbf{p}|_{r(\downarrow\mathbf{p})}$. □

Lemma 4.3. *Let (P, \times, \star) be a strong projection algebra and $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$, $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{q})$.*

- (1) *If $r \in P$, $r \leq_P \mathbf{d}(\mathbf{p})$, $s = \mathbf{r}(\downarrow\mathbf{p})$, then $r\downarrow(\mathbf{p} \circ \mathbf{q}) = r\downarrow\mathbf{p} \circ_s \downarrow\mathbf{q}$.*
- (2) *If $l \in P$, $l \leq_P \mathbf{r}(\mathbf{q})$, $t = \mathbf{d}(\mathbf{q}|_l)$, then $(\mathbf{p} \circ \mathbf{q})|_l = \mathbf{p}|_t \circ \mathbf{q}|_l$.*

Proof. Assume that $\mathbf{p} = (p_1, p_2, \dots, p_k)$ and $\mathbf{q} = (q_1, q_2, \dots, q_l)$. Since $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{q})$, we have $p_k = q_1$. By (15),

$$r\downarrow\mathbf{p} = (r, r\theta_{p_2}, r\theta_{p_2}\theta_{p_3}, \dots, r\theta_{p_2}\cdots\theta_{p_k}),$$

and so $s = \mathbf{r}(\downarrow\mathbf{p}) = r\theta_{p_2}\cdots\theta_{p_k}$. Again by (15), we have

$$\begin{aligned} r\downarrow(\mathbf{p} \circ \mathbf{q}) &= r\downarrow(p_1, p_2, \dots, p_k, q_2, \dots, q_l) \\ &= (r, r\theta_{p_2}, r\theta_{p_2}\theta_{p_3}, \dots, r\theta_{p_2}\cdots\theta_{p_k} = s, s\theta_{q_2}, s\theta_{q_2}\theta_{q_3}, \dots, s\theta_{q_2}\cdots\theta_{q_l}) \\ &= (r, r\theta_{p_2}, r\theta_{p_2}\theta_{p_3}, \dots, r\theta_{p_2}\cdots\theta_{p_k}) \circ (s, s\theta_{q_2}, s\theta_{q_2}\theta_{q_3}, \dots, s\theta_{q_2}\cdots\theta_{q_l}) = r\downarrow\mathbf{p} \circ_s \downarrow\mathbf{q}. \end{aligned}$$

This proves item (1). Dually, we can prove (2). □

Definition 4.4. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category such that (P_C, \times, \star) is a strong projection algebra. Then $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ is called a *weak projection ordered category* if $p \leq q$ if and only if $p \leq_{P_C} q$ for all $p, q \in P_C$.

Theorem 4.5. *Let (P, \times, \star) be a strong projection algebra. Define a relation “ \leq ” on $\mathcal{P}(P)$ as follows: For all $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$,*

$$\mathbf{p} \leq_{\mathcal{P}} \mathbf{q} \iff \text{there exists } r \in P \text{ such that } r \leq_P \mathbf{d}(\mathbf{q}) \text{ and } \mathbf{p} = r\downarrow\mathbf{q}.$$

Then “ $\leq_{\mathcal{P}}$ ” is a partial order on $\mathcal{P}(P)$, and $(\mathcal{P}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\mathcal{P}}, P, \times, \star)$ (Note that $P_{\mathcal{P}(P)} = P$) forms a weak projection ordered category.

Proof. Let $\mathbf{p}, \mathbf{q}, \mathbf{m} \in \mathcal{P}(P)$. By Lemma 4.2 (3), we have $\mathbf{d}(\mathbf{p})\downarrow\mathbf{p} = \mathbf{p}$, and so $\mathbf{p} \leq_{\mathcal{P}} \mathbf{p}$. Let $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$ and $\mathbf{q} \leq_{\mathcal{P}} \mathbf{p}$. Then there exist $r, l \in P$ such that

$$r \leq_P \mathbf{d}(\mathbf{q}), l \leq_P \mathbf{d}(\mathbf{p}), \mathbf{p} = r\downarrow\mathbf{q}, \mathbf{q} = l\downarrow\mathbf{p}$$

By Lemma 4.2 (1), $r \leq_P \mathbf{d}(\mathbf{q}) = \mathbf{d}(l\downarrow\mathbf{p}) = l$. Dually, $l \leq_P r = \mathbf{d}(\mathbf{p})$. Thus $r = l = \mathbf{d}(\mathbf{p}) = \mathbf{d}(\mathbf{q})$, and hence $\mathbf{p} = r\downarrow\mathbf{q} = \mathbf{d}(\mathbf{q})\downarrow\mathbf{q} = \mathbf{q}$ by Lemma 4.2 (3). Assume that $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$ and $\mathbf{q} \leq_{\mathcal{P}} \mathbf{m}$. Then there exist $r, l \in P$ such that

$$r \leq_P \mathbf{d}(\mathbf{q}), l \leq_P \mathbf{d}(\mathbf{m}), \mathbf{p} = r\downarrow\mathbf{q}, \mathbf{q} = l\downarrow\mathbf{m}.$$

This together with Lemma 4.2 (1) gives that $r \leq_P \mathbf{d}(\mathbf{q}) = \mathbf{d}(l\downarrow\mathbf{m}) = l \leq_P \mathbf{d}(\mathbf{m})$. By Lemma 4.2 (5), we have $r\downarrow\mathbf{m} = r\downarrow(l\downarrow\mathbf{m}) = r\downarrow\mathbf{q} = \mathbf{p}$, and so $\mathbf{p} \leq_{\mathcal{P}} \mathbf{m}$. Thus “ $\leq_{\mathcal{P}}$ ” is a partial order.

Let $p, q \in P$. If $p \leq_{\mathcal{P}} q$, then there exists $r \in P$ such that $r \leq_P \mathbf{d}(\mathbf{q}) = q, p = r\downarrow\mathbf{q}$. By (15), we have $p = r$, and so $p \leq_P q$. Conversely, if $p \leq_P q$, then $p \leq_P q = \mathbf{d}(\mathbf{q})$. By (15), we obtain $p\downarrow\mathbf{q} = p$. This gives that $p \leq_{\mathcal{P}} q$. Thus for all $p, q \in P$, $p \leq_{\mathcal{P}} q$ if and only if $p \leq_P q$.

(O1) Let $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$. Then there exists $r \in P$ such that $r \leq_P \mathbf{d}(\mathbf{q})$ and $\mathbf{p} = r\downarrow\mathbf{q}$. By Lemma 4.2 (1),

$$\mathbf{d}(\mathbf{p}) = \mathbf{d}(r\downarrow\mathbf{q}) = r \leq_P \mathbf{d}(\mathbf{q}), \mathbf{r}(\mathbf{p}) = \mathbf{r}(r\downarrow\mathbf{q}) \leq_P \mathbf{r}(\mathbf{q}).$$

This implies that $\mathbf{d}(\mathbf{p}) \leq_{\mathcal{P}} \mathbf{d}(\mathbf{q})$ and $\mathbf{r}(\mathbf{p}) \leq_{\mathcal{P}} \mathbf{r}(\mathbf{q})$.

(O2) Let $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$, $\mathbf{n} \leq_{\mathcal{P}} \mathbf{m}$ and $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{n})$, $\mathbf{r}(\mathbf{q}) = \mathbf{d}(\mathbf{m})$. Then there exist $r, l \in P$ such that

$$\mathbf{p} = {}_r\downarrow\mathbf{q}, \mathbf{n} = {}_l\downarrow\mathbf{m}, r \leq_P \mathbf{d}(\mathbf{q}), l \leq_P \mathbf{d}(\mathbf{m}).$$

By Lemma 4.2 (1), $l = \mathbf{d}({}_l\downarrow\mathbf{m}) = \mathbf{d}(\mathbf{n}) = \mathbf{r}(\mathbf{p}) = \mathbf{r}({}_r\downarrow\mathbf{q})$, and so $\mathbf{p} \circ \mathbf{n} = ({}_r\downarrow\mathbf{q}) \circ ({}_l\downarrow\mathbf{m}) = {}_r\downarrow(\mathbf{q} \circ \mathbf{m})$ by Lemma 4.3 (1). Observe that $r \leq_P \mathbf{d}(\mathbf{q}) = \mathbf{d}(\mathbf{q} \circ \mathbf{m})$, it follows that $\mathbf{p} \circ \mathbf{n} \leq_{\mathcal{P}} \mathbf{q} \circ \mathbf{m}$.

(O3) Let $r \leq_P \mathbf{d}(\mathbf{q})$. By Lemma 4.2 (1), we have ${}_r\downarrow\mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{d}({}_r\downarrow\mathbf{q}) = r$. Obviously, ${}_r\downarrow\mathbf{q} \leq_{\mathcal{P}} \mathbf{q}$. Let $\mathbf{p} \in \mathcal{P}(P)$, $\mathbf{d}(\mathbf{p}) = r$ and $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$. Then there exists $l \in P$ such that $l \leq_P \mathbf{d}(\mathbf{q})$ and $\mathbf{p} = {}_l\downarrow\mathbf{q}$. This implies that $l = \mathbf{d}({}_l\downarrow\mathbf{q}) = \mathbf{d}(\mathbf{p}) = r$ by Lemma 4.2 (1) again. Thus $\mathbf{p} = {}_l\downarrow\mathbf{q} = {}_r\downarrow\mathbf{q}$.

(O4) Let $l \leq_P \mathbf{r}(\mathbf{q})$. By Lemma 4.2 (2), we have $\mathbf{q}|_l \in \mathcal{P}(P)$ and $\mathbf{r}(\mathbf{q}|_l) = l$. By Lemma 4.2 (8), we have $\mathbf{q}|_l = \mathbf{d}(\mathbf{q}|_l)\downarrow\mathbf{q}$, and so $\mathbf{q}|_l \leq_{\mathcal{P}} \mathbf{q}$. Now let $\mathbf{p} \in \mathcal{P}(P)$, $\mathbf{r}(\mathbf{p}) = l$ and $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$. Then there exists $m \leq_P \mathbf{d}(\mathbf{q})$ such that $\mathbf{p} = {}_m\downarrow\mathbf{q} = \mathbf{q}|_{\mathbf{r}({}_m\downarrow\mathbf{q})}$ by Lemma 4.2 (7). So $\mathbf{r}({}_m\downarrow\mathbf{q}) = \mathbf{r}(\mathbf{q}|_{\mathbf{r}({}_m\downarrow\mathbf{q})}) = \mathbf{r}(\mathbf{p}) = l$ by Lemma 4.2 (2). Thus $\mathbf{p} = \mathbf{q}|_{\mathbf{r}({}_m\downarrow\mathbf{q})} = \mathbf{q}|_l$. \square

In the following statements, we shall give some properties of general weak projection ordered categories. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a weak projection ordered category and $a \in C$. Then $\mathbf{d}(a) \in P_C$. By (6) and (10), we can define

$$(20) \quad \Theta_a = \theta_{\mathbf{d}(a)}\nu_a : P_C \longmapsto \mathbf{r}(a)^\downarrow, x \longmapsto \mathbf{r}(x \star \mathbf{d}(a)\downarrow a).$$

Dually, we have

$$(21) \quad \Delta_a = \delta_{\mathbf{r}(a)}\mu_a : P_C \longmapsto \mathbf{d}(a)^\downarrow, y \longmapsto \mathbf{d}(a|_{\mathbf{r}(a) \times y}).$$

It is easy to see that

$$(22) \quad \Theta_p = \theta_p, \quad \Delta_p = \delta_p.$$

for all $p \in P_C$ by (1) and (2) in Lemma 3.1, Remark 2.1 and Lemma 2.2 (11). Moreover, by (1) and (2) in Lemma 3.1, we can obtain that

$$(23) \quad \Theta_a|_{\mathbf{d}(a)^\downarrow} = \nu_a, \quad \Delta_a|_{\mathbf{r}(a)^\downarrow} = \mu_a.$$

Lemma 4.6. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a weak projection ordered category and $a, b \in C$.*

- (1) $\nu_a\theta_{\mathbf{r}(a)} = \nu_a = \nu_a\delta_{\mathbf{r}(a)}$ and $\mu_a\delta_{\mathbf{d}(a)} = \mu_a = \mu_a\theta_{\mathbf{d}(a)}$.
- (2) $\Theta_a\theta_{\mathbf{r}(a)} = \Theta_a = \theta_{\mathbf{d}(a)}\Theta_a$ and $\Delta_a\delta_{\mathbf{d}(a)} = \Delta_a = \delta_{\mathbf{r}(a)}\Delta_a$.
- (3) $\Theta_a\mu_a = \theta_{\mathbf{d}(a)}$.
- (4) If $a \leq b$, then $\Theta_a = \theta_{\mathbf{d}(a)}\nu_b$.
- (5) For all $p, q \in P_C$, if $p \leq \mathbf{d}(a)$ and $q \leq \mathbf{r}(a)$, then

$$\Theta_{p|a} = \theta_p\Theta_a, \quad \Delta_{a|q} = \delta_q\Delta_a, \quad \Theta_{a|q} = \theta_{q\mu_a}\Theta_a, \quad \Delta_{p|a} = \delta_{p\nu_a}\Delta_a.$$

- (6) If $\mathbf{r}(a) = \mathbf{d}(b)$, then $\Theta_{a \circ b} = \Theta_a\Theta_b$ and $\Delta_{a \circ b} = \Delta_b\Delta_a$.
- (7) $\Theta_a\Delta_a = \theta_{\mathbf{d}(a)}$ and $\Delta_a\Theta_a = \delta_{\mathbf{r}(a)}$.

Proof. We first observe that $p \leq q$ if and only if $p \leq_{P_C} q$ for all $p, q \in P_C$ as C is a weak projection ordered category.

(1) It is easy to see that $\nu_a\theta_{\mathbf{r}(a)}$, ν_a and $\nu_a\delta_{\mathbf{r}(a)}$ have the same domain $\mathbf{d}(a)^\downarrow$. Let $x \in \mathbf{d}(a)^\downarrow$. Then $x\nu_a \in \mathbf{r}(a)^\downarrow$, i.e. $x\nu_a \leq \mathbf{r}(a)$. By (1) and (2) in Lemma 3.1, we have

$$x(\nu_a\theta_{\mathbf{r}(a)}) = (x\nu_a)\theta_{\mathbf{r}(a)} = (x\nu_a) \star \mathbf{r}(a) = x\nu_a = \mathbf{r}(a) \times (x\nu_a) = x\nu_a\delta_{\mathbf{r}(a)}.$$

This implies that $\nu_a\theta_{\mathbf{r}(a)} = \nu_a = \nu_a\delta_{\mathbf{r}(a)}$. Dually, $\mu_a\delta_{\mathbf{d}(a)} = \mu_a = \mu_a\theta_{\mathbf{d}(a)}$.

(2) The fact $\Theta_a = \theta_{\mathbf{d}(a)}\nu_a$ and (1) imply that $\Theta_a\theta_{\mathbf{r}(a)} = \theta_{\mathbf{d}(a)}\nu_a\theta_{\mathbf{r}(a)} = \Theta_a$. On the other hand, let $t \in P_C$. Then $t \star \mathbf{d}(a) \leq \mathbf{d}(a)$ and so $(t \star \mathbf{d}(a)) \star \mathbf{d}(a) = t \star \mathbf{d}(a)$ by Lemma 3.1 (2). This implies that

$$t\theta_{\mathbf{d}(a)}\Theta_a = \mathbf{r}((t \star \mathbf{d}(a)) \star \mathbf{d}(a) \downarrow a) = \mathbf{r}(t \star \mathbf{d}(a) \downarrow a) = t\Theta_a.$$

Since $\theta_{\mathbf{d}(a)}\Theta_a$ and Θ_a have the same domain P_C , we have $\theta_{\mathbf{d}(a)}\Theta_a = \Theta_a$. Dually, $\Delta_a\delta_{\mathbf{d}(a)} = \Delta_a = \delta_{\mathbf{r}(a)}\Delta_a$.

(3) Obviously, $\Theta_a\mu_a$ and $\theta_{\mathbf{d}(a)}$ have the same domain P_C . Let $t \in P_C$. By Lemma 3.1 (2), $t \star \mathbf{d}(a) \leq \mathbf{d}(a)$. By using (5) and (1) in Lemma 2.2, we have

$$t\Theta_a\mu_a = \mathbf{d}(a \downarrow_{\mathbf{r}(t \star \mathbf{d}(a) \downarrow a)}) = \mathbf{d}(t \star \mathbf{d}(a) \downarrow a) = t \star \mathbf{d}(a) = t\theta_{\mathbf{d}(a)}.$$

Thus $\Theta_a\mu_a = \theta_{\mathbf{d}(a)}$.

(4) Let $a \leq b$. Then $a = \mathbf{d}(a) \downarrow b$ by Lemma 2.2 (3). Let $t \in P_C$. By Lemma 3.1 (2) and (O1), we have $t \star \mathbf{d}(a) \leq \mathbf{d}(a) \leq \mathbf{d}(b)$. By Lemma 2.2 (9),

$$t\Theta_a = \mathbf{r}((t \star \mathbf{d}(a)) \downarrow a) = \mathbf{r}((t \star \mathbf{d}(a)) \downarrow (\mathbf{d}(a) \downarrow b)) = \mathbf{r}((t \star \mathbf{d}(a)) \downarrow b) = t\theta_{\mathbf{d}(a)}\nu_b.$$

(5) Let $p \leq \mathbf{d}(a)$ and $t \in P_C$. By Lemma 3.1 (2), we have $t \star p \leq p \leq \mathbf{d}(a)$ and $(t \star p) \star \mathbf{d}(a) = t \star p$. By the definition of Θ_a , (1) and (9) in Lemma 2.2,

$$t\Theta_{p \downarrow a} = \mathbf{r}((t \star \mathbf{d}(p \downarrow a)) \downarrow (p \downarrow a)) = \mathbf{r}(t \star p \downarrow (p \downarrow a)) = \mathbf{r}((t \star p) \downarrow a) = \mathbf{r}((t \star p) \star \mathbf{d}(a) \downarrow a) = t\theta_p\Theta_a.$$

This implies that $\Theta_{p \downarrow a} = \theta_p\Theta_a$. Dually, $\Delta_{a \downarrow q} = \delta_q\Delta_a$ for all $q \in P_C$ with $q \leq \mathbf{r}(a)$. On the other hand, let $q \in P_C$ and $q \leq \mathbf{r}(a)$. By Lemma 2.2 (6), we obtain $a \downarrow_q = \mathbf{d}(a \downarrow_q) \downarrow a = q\mu_a \downarrow a$, and so $\Theta_{a \downarrow q} = \Theta_{q\mu_a \downarrow a} = \theta_{q\mu_a}\Theta_a$. Dually, $\Delta_{p \downarrow a} = \delta_{p\nu_a}\Delta_a$ for all $p \in P_C$ with $p \leq \mathbf{d}(a)$.

(6) Let $p = \mathbf{d}(a)$ and $q = \mathbf{r}(a) = \mathbf{d}(b)$. Using (1) of the present lemma, Lemma 2.5 and the fact that $\mathbf{d}(a \circ b) = \mathbf{d}(a)$ in order,

$$\Theta_a\Theta_b = \theta_{\mathbf{d}(a)}\nu_a\theta_{\mathbf{d}(b)}\nu_b = \theta_{\mathbf{d}(a)}\nu_a\theta_{\mathbf{r}(a)}\nu_b = \theta_{\mathbf{d}(a)}\nu_a\nu_b = \theta_{\mathbf{d}(a)}\nu_{a \circ b} = \theta_{\mathbf{d}(a \circ b)}\nu_{a \circ b} = \Theta_{a \circ b}.$$

Dually, $\Delta_{a \circ b} = \Delta_b\Delta_a$.

(7) Let $t \in P$. Then $t \star \mathbf{d}(a) \leq \mathbf{d}(a)$ By Lemma 3.1 (2). By Lemma 2.2 (1), we have $\mathbf{r}((t \star \mathbf{d}(a)) \downarrow a) \leq \mathbf{r}(a)$, which together with Lemma 3.1 (1) gives that $\mathbf{r}(a) \times \mathbf{r}((t \star \mathbf{d}(a)) \downarrow a) = \mathbf{r}((t \star \mathbf{d}(a)) \downarrow a)$. By (5) and (1) in Lemma 2.2,

$$t\Theta_a\Delta_a = \mathbf{d}(a \downarrow_{\mathbf{r}(a) \times \mathbf{r}((t \star \mathbf{d}(a)) \downarrow a)}) = \mathbf{d}(a \downarrow_{\mathbf{r}((t \star \mathbf{d}(a)) \downarrow a)}) = \mathbf{d}((t \star \mathbf{d}(a)) \downarrow a) = t \star \mathbf{d}(a) = t\theta_{\mathbf{d}(a)}.$$

Since $\Theta_a\Delta_a$ and $\theta_{\mathbf{d}(a)}$ have the same domain P_C , we have $\Theta_a\Delta_a = \theta_{\mathbf{d}(a)}$. Dually, we can prove that $\Delta_a\Theta_a = \delta_{\mathbf{r}(a)}$. \square

Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a weak projection ordered category. Then $p^\downarrow = \{x \in P_C \mid x \leq p\}$ is a subalgebra of (P_C, \times, \star) for all $p \in P_C$. In fact, if $x, y \in p^\downarrow$, then $x \leq p$ and $y \leq p$. By (1) and (2) in Lemma 3.1, we have $x \times y \leq x \leq p$ and $x \star y \leq y \leq p$. This yields that $x \times y, x \star y \in p^\downarrow$.

Lemma 4.7. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a weak projection ordered category. Then the following statements are equivalent:*

- (G1a) *For all $a \in C$ and $p, q \in P_C$, if $p \leq \mathbf{d}(a)$ and $q \leq \mathbf{r}(a)$, then $\delta_{p\nu_a} = \Delta_a\delta_p\Theta_a$ and $\theta_{q\mu_a} = \Theta_a\theta_q\Delta_a$.*
- (G1b) *For all $a \in C$ and $p \in P_C$, we have $\delta_p\Theta_a = \Delta_a\delta_p\Theta_a$ and $\theta_{p\Delta_a} = \Theta_a\theta_p\Delta_a$.*
- (G1c) *For all $a \in C$ and $p, q \in P_C$, if $p \leq \mathbf{d}(a)$ and $q \leq \mathbf{r}(a)$, then $\Delta_{p \downarrow a} = \Delta_a\delta_p$ and $\Theta_{a \downarrow q} = \Theta_a\theta_q$.*

(G1d) For all $a \in C$, the map $\nu_a : \mathbf{d}(a)^\downarrow \rightarrow \mathbf{r}(a)^\downarrow$, $p \mapsto \mathbf{r}(p \downarrow a)$ is a homomorphism between projection algebras.

Proof. Let $a \in C$ and denote $s = \mathbf{d}(a)$ and $t = \mathbf{r}(a)$. Then $s, t \in P_C$.

(G1a) \implies (G1b). Let $p \in P_C$. By using the definition of Θ_a , (G1a), the fact that $p\theta_s = p \star s \leq s = \mathbf{d}(a)$, Lemma 3.7 and Lemma 4.6 (2) in order, we have

$$\delta_p \Theta_a = \delta_{(p\theta_s)\nu_a} = \Delta_a \delta_{(p\theta_s)} \Theta_a = \Delta_a \delta_s \delta_p \theta_s \Theta_a = \Delta_a \delta_p \Theta_a.$$

Dually, $\theta_p \Delta_a = \Theta_a \theta_p \Delta_a$.

(G1a) \implies (G1c). Let $p \in P_C$ with $p \leq \mathbf{d}(a)$. By (5) and (1) in Lemma 2.2, we have $p \downarrow a = a \downarrow_r$, where $r = \mathbf{r}(p \downarrow a) \leq \mathbf{r}(a)$. By using Lemma 4.6 (5), (G1a), Lemma 4.6 (7) and Lemma 3.2, we have

$$\Delta_{p \downarrow a} = \delta_{p\nu_a} \Delta_a = \Delta_a \delta_p \Theta_a \Delta_a = \Delta_a \delta_p \theta_{\mathbf{d}(a)} = \Delta_a \delta_p.$$

Dually, $\Theta_{a \downarrow q} = \Theta_a \theta_q$ for all $q \in P_C$ with $q \leq \mathbf{r}(a)$.

(G1b) \implies (G1a). Let $p \in P_C$ with $p \leq \mathbf{d}(a)$. By Lemma 3.1 (2), we have $p = p \star \mathbf{d}(a) = p\theta_{\mathbf{d}(a)}$, and so $p\nu_a = (p\theta_{\mathbf{d}(a)})\nu_a = p(\theta_{\mathbf{d}(a)}\nu_a) = p\Theta_a$. In view of (G1b), we obtain that $\delta_{p\nu_a} = \delta_{p\Theta_a} = \Delta_a \delta_p \Theta_a$. Dually, $\theta_{q\nu_a} = \Theta_a \theta_q \Delta_a$ for all $q \in P_C$ with $q \leq \mathbf{r}(a)$.

(G1c) \implies (G1a). Let $p \in P_C$ with $p \leq \mathbf{d}(a)$. By Lemma 2.2 (1), we have $p\nu_a = \mathbf{r}(p \downarrow a) \leq \mathbf{r}(a)$ and so $\delta_{p\nu_a} \delta_{\mathbf{r}(a)} = \delta_{p\nu_a}$ by Lemma 3.2. Using (G1c) and (5) and (7) in Lemma 4.6, we have

$$\Delta_a \delta_p \Theta_a = \Delta_{p \downarrow a} \Theta_a = \delta_{p\nu_a} \Delta_a \Theta_a = \delta_{p\nu_a} \delta_{\mathbf{r}(a)} = \delta_{p\nu_a}.$$

Dually, $\theta_{q\nu_a} = \Theta_a \theta_q \Delta_a$ for all $q \in P_C$ with $q \leq \mathbf{r}(a)$.

(G1c) \implies (G1d). Assume that (G1c) holds. Then (G1a) also holds by the previous paragraph. Let $p, q \in P_C$ with $p, q \in \mathbf{d}(a)^\downarrow$. Then $p, q \leq \mathbf{d}(a)$. On one hand,

$$\begin{aligned} (q\nu_a) \times (p\nu_a) &= p\nu_a \delta_{q\nu_a} \quad (\text{by (10)}) \\ &= p\nu_a \Delta_a \delta_q \Theta_a \quad (\text{by (G1a)}) \\ &= p\nu_a \delta_{\mathbf{r}(a)} \mu_a \delta_q \theta_{\mathbf{d}(a)} \nu_a \quad (\text{by (20) and (21)}) \\ &= p\nu_a \mu_a \delta_q \nu_a \quad (\text{Lemma 4.6 (1) and Lemma 3.2}) \\ &= p \delta_q \nu_a \quad (\text{by Lemma 2.3}) \\ &= (q \times p) \nu_a. \quad (\text{by (10)}) \end{aligned}$$

This shows that ν_a preserves \times . On the other hand,

$$\begin{aligned} p\Theta_a \theta_{\mathbf{r}(q \downarrow a)} &= p\Theta_{a \downarrow_{\mathbf{r}(q \downarrow a)}} \quad (\text{by (G1c)}) \\ &= p\Theta_{q \downarrow a} \quad (\text{by Lemma 2.2 (5)}) \\ &= \mathbf{r}(p \star \mathbf{d}(q \downarrow a) \downarrow (q \downarrow a)) \quad (\text{by (20)}) \\ &= \mathbf{r}(p \star q \downarrow (q \downarrow a)) \quad (\text{by Lemma 2.2 (1)}) \\ &= \mathbf{r}(p \star q \downarrow a) \quad (\text{by Lemma 3.1 (2) and Lemma 2.2 (9)}) \\ &= (p \star q) \nu_a. \quad (\text{by (6)}) \end{aligned}$$

Moreover,

$$\begin{aligned}
p\Theta_a\theta_{\mathbf{r}(q\downarrow a)} &= \mathbf{r}(p\star\mathbf{d}(a)\downarrow a)\theta_{\mathbf{r}(q\downarrow a)} \quad (\text{by (20)}) \\
&= \mathbf{r}(p\downarrow a)\theta_{\mathbf{r}(q\downarrow a)} \quad (\text{by Lemma 3.1 (2)}) \\
&= \mathbf{r}(p\downarrow a) \star \mathbf{r}(q\downarrow a) \quad (\text{by (10)}) \\
&= (p\nu_a) \star (q\nu_a). \quad (\text{by (6)})
\end{aligned}$$

Thus $(p\star q)\nu_a = (p\nu_a) \star (q\nu_a)$. Therefore ν_a is a homomorphism.

(G1d) \implies (G1a). By Lemma 2.3 and (G1d), $\nu_a : \mathbf{d}(a)^\downarrow \rightarrow \mathbf{r}(a)^\downarrow$, $p \mapsto \mathbf{r}(p\downarrow a)$ is an isomorphism. Let $p \leq \mathbf{d}(a) = s$ and $e \in P_C$. We shall prove that $e\delta_{p\nu_a} = e\Delta_a\delta_p\Theta_a$ in the sequel. Denote $f = t \times e = e\delta_t$, where $t = \mathbf{r}(a)$. Then by Lemma 3.1 (1), we have $f \leq t$, i.e. $f \in \mathbf{r}(a)^\downarrow$, and so we can let $g = f\mu_a$. By Lemma 2.3, we have $f = f\mu_a\nu_a = g\nu_a$. Obviously, we obtain $g \in \mathbf{d}(a)^\downarrow$, i.e. $g \leq \mathbf{d}(a) = s$. By the facts that $p\nu_a \leq \mathbf{r}(a) = t$, $p \leq \mathbf{d}(a) = s$ and Lemma 3.2, we have $\delta_{p\nu_a} = \delta_t\delta_{p\nu_a}$ and $\delta_p\theta_s = \delta_p$. Using Lemma 3.2, the fact that ν_a is an isomorphism and the definitions of Δ_a and Θ_a in order, we have

$$\begin{aligned}
e\delta_{p\nu_a} &= e\delta_t\delta_{p\nu_a} = f\delta_{p\nu_a} = (g\nu_a)\delta_{p\nu_a} = (g\delta_p)\nu_a = (f\mu_a)\delta_p\nu_a = e\delta_t\mu_a\delta_p\nu_a \\
&= e\delta_t\mu_a\delta_p\theta_s\nu_a = e\delta_{\mathbf{r}(a)}\mu_a\delta_p\theta_{\mathbf{d}(a)}\nu_a = e\Delta_a\delta_p\Theta_a.
\end{aligned}$$

This shows $\delta_{p\nu_a} = \Delta_a\delta_p\Theta_a$. On the other hand, by Lemma 2.3 and (G1d),

$$\mu_a : \mathbf{r}(a)^\downarrow \rightarrow \mathbf{d}(a)^\downarrow, q \mapsto q\mu_a = \mathbf{d}(a\downarrow q)$$

is also an isomorphism. Thus $\theta_{q\mu_a} = \Theta_a\theta_q\Delta_a$ by dual arguments. \square

Definition 4.8. A weak projection ordered category $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ is called a *projection ordered category* if the conditions (G1a)–(G1d) in Lemma 4.7 hold.

Lemma 4.9. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category and $a \in C, p, q \in P_C$. Then

$$p\Delta_a\delta_q\Theta_a\theta_p = q\Theta_a\theta_p, \quad p\Theta_a\theta_q\Delta_a\delta_p = q\Delta_a\delta_p.$$

Proof. Denote $t = q\Theta_a$. By (G1b) and Lemma 3.1 (3),

$$p\Delta_a\delta_q\Theta_a\theta_p = p\delta_{q\Theta_a}\theta_p = p\delta_t\theta_p = (t \times p) \star p = t \star p = t\theta_p = q\Theta_a\theta_p.$$

The other identity can be proved dually. \square

Definition 4.10. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category, $\mathcal{P}(P_C)$ be the path category of P_C and $\varepsilon : \mathcal{P}(P_C) \rightarrow C$ be a map. Then ε is called an *evaluation map* if the following conditions hold: For all $p \in P_C$ and $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P_C)$,

- (E1) $\varepsilon(p) = \varepsilon(p, p) = p$.
- (E2) $\mathbf{d}(\varepsilon(\mathbf{p})) = \mathbf{d}(\mathbf{p})$ and $\mathbf{r}(\varepsilon(\mathbf{p})) = \mathbf{r}(\mathbf{p})$.
- (E3) If $\mathbf{p} \circ \mathbf{q}$ is defined, then $\varepsilon(\mathbf{p} \circ \mathbf{q}) = \varepsilon(\mathbf{p}) \circ \varepsilon(\mathbf{q})$.
- (E4) If $\mathbf{p} \leq \mathbf{q}$, then $\varepsilon(\mathbf{p}) \leq \varepsilon(\mathbf{q})$.

Proposition 4.11. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category, $\mathcal{P}(P_C)$ be the path category of P_C and $\varepsilon : \mathcal{P}(P_C) \rightarrow C$ be a map satisfying (E1)–(E3). Then the following statements are equivalent:

- (E4) If $\mathbf{p} \leq \mathbf{q}$, then $\varepsilon(\mathbf{p}) \leq \varepsilon(\mathbf{q})$.
- (E5) For all $\mathbf{p} \in \mathcal{P}(P_C)$ and $t \in P_C$, the fact that $t \leq \mathbf{d}(\mathbf{p})$ implies that $\varepsilon(t\downarrow\mathbf{p}) = t\downarrow\varepsilon(\mathbf{p})$.
- (E6) For all $\mathbf{p} \in \mathcal{P}(P_C)$ and $t \in P_C$, the fact that $t \leq \mathbf{r}(\mathbf{p})$ implies that $\varepsilon(\mathbf{p}\downarrow t) = \varepsilon(\mathbf{p})\downarrow t$.

Proof. Assume that (E5) holds and $\mathbf{p} \leq_{\mathcal{P}} \mathbf{q}$. Then $\mathbf{p} = r \downarrow \mathbf{q}$ for some $r \in P_C$ with $r \leq \mathbf{d}(\mathbf{q})$. This implies that $\varepsilon(\mathbf{p}) = \varepsilon(r \downarrow \mathbf{q}) = r \downarrow \varepsilon(\mathbf{q}) \leq \varepsilon(\mathbf{q})$. Conversely, assume that (E4) holds and $q \in P_C$ with $q \leq \mathbf{d}(\mathbf{p})$. By Theorem 4.5, we have $q \downarrow \mathbf{p} \leq_{\mathcal{P}} \mathbf{p}$ and $q \leq_{\mathcal{P}} \mathbf{d}(\mathbf{p})$. Using (E1), (E4) and (E2), we have $q = \varepsilon(q) \leq \varepsilon(\mathbf{d}(\mathbf{p})) = \mathbf{d}(\varepsilon(\mathbf{p}))$. On the other hand, (E4) gives $\varepsilon(q \downarrow \mathbf{p}) \leq \varepsilon(\mathbf{p})$, and (E2), (E1) and Lemma 4.2 (1) together imply that $\mathbf{d}(\varepsilon(q \downarrow \mathbf{p})) = \varepsilon(\mathbf{d}(q \downarrow \mathbf{p})) = \mathbf{d}(q \downarrow \mathbf{p}) = q$. By Lemma 2.2 (3), $\varepsilon(q \downarrow \mathbf{p}) =_q \downarrow \varepsilon(\mathbf{p})$. Thus (E4) is equivalent to (E5). Dually, we can prove that (E4) is equivalent to (E6). \square

Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category, $\mathcal{P}(P_C)$ be the path category of P_C and $\varepsilon : \mathcal{P}(P_C) \rightarrow C$ be an evaluation map, $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P_C)$. By (E3) and the fact that $\mathbf{p} = (p_1, p_2) \circ (p_2, p_3) \circ \dots \circ (p_{k-1}, p_k)$ in $\mathcal{P}(P)$, we have

$$(24) \quad \varepsilon(\mathbf{p}) = \varepsilon(p_1, p_2) \circ \varepsilon(p_2, p_3) \circ \dots \circ \varepsilon(p_{k-1}, p_k).$$

Lemma 4.12. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category with an evaluation map $\varepsilon : \mathcal{P}(P_C) \rightarrow C$ and*

$$a \in C, p, q, r, s \in P_C, p \mathcal{F}_{P_C} q, r \leq p, s \leq q, a \leq \varepsilon(p, q).$$

$$(1) \quad r \downarrow \varepsilon(p, q) = \varepsilon(r, r\theta_q) \text{ and } \varepsilon(p, q) \downarrow_s = \varepsilon(s\delta_p, s).$$

(2)

$$\begin{aligned} \mathbf{d}(\varepsilon(p, q)) &= p, & \nu_{\varepsilon(p, q)} &= \theta_q \downarrow_{p^\downarrow}, & \Theta_{\varepsilon(p, q)} &= \theta_p \theta_q, \\ \mathbf{r}(\varepsilon(p, q)) &= q, & \mu_{\varepsilon(p, q)} &= \delta_p \downarrow_{q^\downarrow}, & \Delta_{\varepsilon(p, q)} &= \delta_q \delta_p, \end{aligned}$$

$$(3) \quad a = \varepsilon(\mathbf{d}(a), \mathbf{r}(a)).$$

Proof. (1) By the fact that $r \leq p = \mathbf{d}((p, q))$ and (15), we have $r \downarrow (p, q) = (r, r\theta_q)$ and so $\varepsilon(r, r\theta_q) = \varepsilon(r \downarrow (p, q)) = r \downarrow \varepsilon(p, q)$ by (E5). Dually, $\varepsilon(p, q) \downarrow_s = \varepsilon(s\delta_p, s)$.

(2) By (E2), we obtain $\mathbf{d}(\varepsilon(p, q)) = \mathbf{d}((p, q)) = p$. This implies that $\nu_{\varepsilon(p, q)}$ and $\theta_q \downarrow_{p^\downarrow}$ have the same domain p^\downarrow . Let $r \in p^\downarrow$. Then $r \leq p$ and so

$$r\nu_{\varepsilon(p, q)} = \mathbf{r}(r \downarrow \varepsilon(p, q)) \stackrel{(E5)}{=} \mathbf{r}(\varepsilon(r \downarrow (p, q))) \stackrel{(4.1)}{=} \mathbf{r}(\varepsilon(r, r\theta_q)) \stackrel{(E2)}{=} \mathbf{r}(r, r\theta_q) = r\theta_q.$$

Observe that the range of θ_p is p^\downarrow , it follows that

$$\Theta_{\varepsilon(p, q)} = \theta_{\mathbf{d}(\varepsilon(p, q))} \nu_{\varepsilon(p, q)} = \theta_p \nu_{\varepsilon(p, q)} = \theta_p (\theta_q \downarrow_{p^\downarrow}) = \theta_p \theta_q.$$

The other three equalities can be proved dually.

(3) By the fact $a \leq \varepsilon(p, q)$, (O1) and (E2), we have $\mathbf{d}(a) \leq \mathbf{d}(\varepsilon(p, q)) = \mathbf{d}(p, q) = p$. Item (1) in the present lemma gives that $\mathbf{d}(a) \downarrow \varepsilon(p, q) = \varepsilon(\mathbf{d}(a), \mathbf{d}(a)\theta_q)$. On the other hand, by the fact that $a \leq \varepsilon(p, q)$ and Lemma 2.2 (3), we obtain $a = \mathbf{d}(a) \downarrow \varepsilon(p, q)$. So $a = \varepsilon(\mathbf{d}(a), \mathbf{d}(a)\theta_q)$. This implies that

$$\mathbf{r}(a) = \mathbf{r}(\varepsilon(\mathbf{d}(a), \mathbf{d}(a)\theta_q)) = \mathbf{r}(\mathbf{d}(a), \mathbf{d}(a)\theta_q) = \mathbf{d}(a)\theta_q$$

by (E2). Thus $a = \varepsilon(\mathbf{d}(a), \mathbf{r}(a))$. \square

Definition 4.13. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category, $b \in C$ and $(e, f) \in P_C \times P_C$. Then (e, f) is called a *b-linked pair* if

$$(25) \quad f = e\Theta_b\theta_f, \quad e = f\Delta_b\delta_e.$$

Lemma 4.14. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category, $b \in C$ and (e, f) be a b-linked pair. Denote*

$$(26) \quad e_1 = e\theta_{\mathbf{d}(b)}, e_2 = f\Delta_b, f_1 = e\Theta_b, f_2 = f\delta_{\mathbf{r}(b)}$$

- (1) $e_i \leq \mathbf{d}(b), f_i \leq \mathbf{r}(b), i = 1, 2.$
(2) $e = e \times \mathbf{d}(b), \mathbf{r}(b) \star f = f, e \mathcal{F}_{P_C} e_i, f_i \mathcal{F}_{P_C} f, i = 1, 2.$
(3) $e_i \downarrow b = b \downarrow_{f_i}, i = 1, 2.$

Proof. (1) Since $\text{im}\Delta_b = \text{im}\theta_{\mathbf{d}(b)} \subseteq \mathbf{d}(b)^\downarrow$ and $\text{im}\Theta_b = \text{im}\delta_{\mathbf{r}(b)} = \mathbf{r}(b)^\downarrow$, we have $e_i \leq \mathbf{d}(b)$ and $f_i \leq \mathbf{r}(b), i = 1, 2.$

(2) By (25) and (26), we have $e = f\Delta_b\delta_e = e_2\delta_e = e \times e_2.$ Using (25), (26) and (G1b), we can obtain

$$e\theta_{e_2} = e\theta_{f\Delta_b} = e\Theta_b\theta_f\Delta_b = f\Delta_b = e_2.$$

This implies that $e\mathcal{F}_{P_C}e_2.$ By the facts $e = e \times e_2, e_2 \leq \mathbf{d}(b)$ and Lemma 3.4 (2), we get $e = e \times \mathbf{d}(b).$ This together with (3) and (4) in Lemma 3.1 gives that

$$e \times e_1 = e \times e\theta_{\mathbf{d}(a)} = e \times (e \star \mathbf{d}(b)) = e \times \mathbf{d}(b) = e,$$

$$e \star e_1 = e \star e\theta_{\mathbf{d}(b)} = e \star (e \star \mathbf{d}(b)) = (e \times \mathbf{d}(b)) \star (e \star \mathbf{d}(b)) = e \star \mathbf{d}(b) = e\theta_{\mathbf{d}(b)} = e_1.$$

This shows that $e\mathcal{F}_{P_C}e_1.$ By (25) and (26), $f_1 \star f = f_1\theta_f = e\Theta_b\theta_f = f.$ Again by (25), (26) and (G1b),

$$f_1 \times f = f\delta_{f_1} = f\delta_{e\Theta_b} = f\Delta_b\delta_e\Theta_b = e\Theta_b = f_1.$$

Thus $f_1\mathcal{F}_{P_C}f.$ On the other hand, (26), (L4) and (L1) imply

$$f_2 \times f = f\delta_{\mathbf{r}(b)} \times f = (\mathbf{r}(b) \times f) \times f = (\mathbf{r}(b) \times f) \times (\mathbf{r}(b) \times f) = \mathbf{r}(b) \times f = f\delta_{\mathbf{r}(b)} = f_2.$$

By Lemma 3.1 (3), $f_2 \star f = (\mathbf{r}(b) \times f) \star f = \mathbf{r}(b) \star f.$ Since $f_1 \leq \mathbf{r}(b),$ we have $f = f_1 \star f \leq \mathbf{r}(b) \star f \leq f$ by (5) and (2) in Lemma 3.1, and so $\mathbf{r}(b) \star f = f = f_2 \star f.$ Thus $f_2\mathcal{F}_{P_C}f.$

(3) By (25), (26) and Lemma 2.3, we have $e_1\nu_b = e\theta_{\mathbf{d}(b)}\nu_b = e\Theta_b = f_1$ and

$$e_2\nu_b = f\Delta_b\nu_b = f\delta_{\mathbf{r}(b)}\nu_b = f\delta_{\mathbf{r}(b)} = f_2.$$

By Lemma 2.2 (5), $e_i \downarrow b = b \downarrow_{\mathbf{r}(e_i \downarrow b)} = b \downarrow_{e_i\nu_b} = b \downarrow_{f_i}, i = 1, 2.$ □

Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category with an evaluation map $\varepsilon : \mathcal{P}(P_C) \rightarrow C, b \in C, (e, f)$ be a b -linked pair and $e_1, e_2, f_1, f_2 \in P_C$ be defined as in (26). By (1) and (2) in Lemma 2.2, (E2), Lemmas 4.12 and 4.14, the following two elements are well defined:

$$(27) \quad \lambda(e, b, f) = \varepsilon(e, e_1) \circ e_1 \downarrow b \circ \varepsilon(f_1, f) = \varepsilon(e, e_1) \circ b \downarrow_{f_1} \circ \varepsilon(f_1, f),$$

$$(28) \quad \rho(e, b, f) = \varepsilon(e, e_2) \circ e_2 \downarrow b \circ \varepsilon(f_2, f) = \varepsilon(e, e_2) \circ b \downarrow_{f_2} \circ \varepsilon(f_2, f).$$

Definition 4.15. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star)$ be a projection ordered category with an evaluation map $\varepsilon : \mathcal{P}(P_C) \rightarrow C.$ Then $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ is called a *chain projection ordered category* if the following condition holds:

- (G2) For all $b \in C$ and b -linked pair $(e, f),$ we have $\lambda(e, b, f) = \rho(e, b, f),$ where $\lambda(e, b, f)$ and $\rho(e, b, f)$ are defined by (27) and (28) respectively.

Moreover, a chain projection ordered category $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ is called a

- *chain projection ordered groupoid* if $(C, \circ, \mathbf{d}, \mathbf{r})$ is a groupoid,
- *symmetric chain projection ordered category* if (P_C, \times, \star) is a symmetric strong projection algebra, and for all $p, q \in P_C, p\mathcal{F}_{P_C}q$ implies that $\varepsilon(p, q, p) = p,$
- *symmetric chain projection ordered groupoid* if it is symmetric and $(C, \circ, \mathbf{d}, \mathbf{r})$ is also a groupoid,

- *commutative chain projection ordered category* if (P_C, \times, \star) is a commutative strong projection algebra,
- *commutative chain projection ordered groupoid* if is commutative and $(C, \circ, \mathbf{d}, \mathbf{r})$ is also a groupoid.

Definition 4.16. Assume that $(C_1, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C \times, \star, \varepsilon_1)$ and $(C_2, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon_2)$ are chain projection ordered categories and $\phi : C_1 \rightarrow C_2, a \rightarrow a\phi$ be a map. Then ϕ is called a *chain projection ordered functor* if for all $a, b \in C_1$ and $e, f \in P_{C_1}$, the following conditions hold:

- (F1) $\mathbf{d}(a\phi) = (\mathbf{d}(a))\phi, \mathbf{r}(a\phi) = (\mathbf{r}(a))\phi.$
- (F2) If $a \circ b$ is defined, then $(a \circ b)\phi = (a\phi) \circ (b\phi).$
- (F3) If $a \leq b$, then $a\phi \leq b\phi.$
- (F4) $(e \times f)\phi = e\phi \times f\phi, (e \star f)\phi = e\phi \star f\phi.$
- (F5) For all $(p_1, p_2, \dots, p_k) \in \mathcal{P}(P_{C_1}),$

$$(\varepsilon_1((p_1, p_2, \dots, p_k)))\phi = \varepsilon_2((p_1\phi, p_2\phi, \dots, p_k\phi)).$$

Lemma 4.17. Let ϕ be a chain projection ordered functor between chain projection ordered groupoids $(C_1, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_1}, \times, \star, \varepsilon_1)$ and $(C_2, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_2}, \times, \star, \varepsilon_2).$ Then $(x^{-1})\psi = (x\psi)^{-1}$ for all $x \in C_1.$

Proof. By the facts $x \circ x^{-1} = \mathbf{d}(x)$ and $x^{-1} \circ x = \mathbf{r}(x)$ and (F1), (F2),

$$(x\psi) \circ (x^{-1}\psi) = (x \circ x^{-1})\psi = (\mathbf{d}(x))\psi = \mathbf{d}(x\psi), (x^{-1}\psi) \circ (x\psi) = \mathbf{r}(x\psi).$$

This implies that $(x\psi)^{-1} = x^{-1}\psi$ by the statements before Lemma 2.6. \square

The following proposition is obvious.

Proposition 4.18. The class of (respectively, symmetric, commutative) chain projection ordered categories (respectively, groupoids) together with chain projection ordered functors forms a category.

Lemma 4.19. Let ϕ be a chain projection ordered functor between chain projection ordered categories $(C_1, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_1}, \times, \star, \varepsilon_1)$ and $(C_2, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_2}, \times, \star, \varepsilon_2),$ and $x \in C_1, e, f \in P_{C_1}, e \leq \mathbf{d}(x), f \leq \mathbf{r}(x).$ Then $(e \downarrow x)\phi = e\phi \downarrow (x\phi), (x \downarrow f)\phi = (x\phi) \downarrow f\phi.$

Proof. (F1) gives $e\phi = (\mathbf{d}(e))\phi = \mathbf{d}(e\phi) \in P_{C_2},$ and (F3) gives $e\phi \leq (\mathbf{d}(x))\phi = \mathbf{d}(x\phi).$ This implies that $e\phi \downarrow (x\phi)$ is defined. By the fact that $e \downarrow x \leq x$ and (F3), we have $(e \downarrow x)\phi \leq x\phi,$ and by the fact $\mathbf{d}(e \downarrow x) = e$ and (F1), we obtain $\mathbf{d}((e \downarrow x)\phi) = (\mathbf{d}(e \downarrow x))\phi = e\phi.$ By Lemma 2.2 (3), we get $(e \downarrow x)\phi = e\phi \downarrow (x\phi).$ Dually, $(x \downarrow f)\phi = (x\phi) \downarrow f\phi. \square$

5. FROM DRC-RESTRICTION SEMIGROUPS TO CHAIN PROJECTION ORDERED CATEGORIES

In this section, we present some results on DRC-restriction semigroups and prove that each DRC-restriction semigroup give rises to a chain projection ordered category. We begin by giving the following lemma.

Lemma 5.1. Let (S, \cdot, \circ) be a generalized regular \circ -semigroup (respectively, a regular \circ -semigroup, an inverse semigroup) and denote $x^+ = xx^\circ$ and $x^* = x^\circ x$ for all $x \in S.$ Then

$(S, \cdot, +, *)$ is a DRC-restriction semigroup (respectively, P -restriction semigroup, restriction semigroup) and for all $x \in S$,

$$(29) \quad x^+ = xx^\circ, x^* = x^\circ x, x^* = x^{\circ+}, x^+ = x^{\circ*}.$$

Conversely, let $(S, \cdot, +, *)$ be a DRC-restriction semigroup (respectively, P -restriction semigroup, restriction semigroup) and for each $x \in S$, there exists $x^\circ \in S$ such that

$$(30) \quad x^+ = xx^\circ, x^* = x^\circ x, x^* = x^{\circ+}, x^+ = x^{\circ*}.$$

Then the above x° is unique and (S, \cdot, \circ) forms a generalized regular \circ -semigroup (respectively, a regular \circ -semigroup, an inverse semigroup).

Proof. By [59, Proposition 3.7], under the given assumption, $(S, \cdot, +, *)$ is a DRC-restriction semigroup, and (29) follows from (5). This gives the direct part. Conversely, let $(S, \cdot, +, *)$ be DRC-restriction semigroup satisfying (30). We first assert that the element x° in (30) is unique. In fact, let $y \in S$ and $x^+ = xy$, $x^* = yx$, $x^* = y^+$, $x^+ = y^*$. Then by (i) and (i)',

$$y = y^+y = x^*y = x^\circ xy = x^\circ x^+ = x^\circ x^{\circ*} = x^\circ.$$

Moreover, $xx^\circ x = x$, $x^\circ x x^\circ = x^* x^\circ = x^{\circ+} x^\circ = x^\circ$. This implies that $x^* = x^\circ x \mathcal{L} x \mathcal{R} x x^\circ = x^+$, and so x° is the unique inverse element of x satisfying $xx^\circ = x^+$ and $x^* = x^\circ x$. By [59, Proposition 3.6], (S, \cdot, \circ) is a generalized regular \circ -semigroup.

The other two special cases follow from [55, Proposition 6.1], (1) and Proposition 1.2, we omit the details. \square

Remark 5.2. By Lemma 5.1, a generalized regular \circ -semigroup (respectively, a regular \circ -semigroup, an inverse semigroup) is exactly an algebraic system $(S, \cdot, +, *, \circ)$ where $(S, \cdot, +, *)$ is a DRC-restriction semigroup (respectively, P -restriction semigroup, restriction semigroup) and “ \circ ” is a unary operation on S satisfying the axioms:

$$x^+ = xx^\circ, x^\circ x = x^*, x^* = x^{\circ+}, x^{\circ*} = x^+.$$

The following lemmas provide some properties of projections in DRC semigroups.

Lemma 5.3 ([57]). *Let $(S, \cdot, +, *)$ be a DRC semigroup and $e, f \in P(S)$.*

- (1) $e^+ = e = e^* \in E(S)$.
- (2) $(ef)^+ = e(ef)^+e$, $(ef)^* = f(ef)^*f$.
- (3) $(ef)^+f = ef = e(ef)^*$.
- (4) $(ef)^+(ef)^* = ef$.

Corollary 5.4 ([57]). *In DRC Conditions, the axiom (iii) (respectively, (iii)') can be replaced by the axiom $(x^+y^+)^+ = x^+(x^+y^+)^+x^+$ (respectively, $(x^*y^*)^* = y^*(x^*y^*)^*y^*$).*

Lemma 5.5 ([26, 59]). *Let $(S, \cdot, +, *)$ be a DRC semigroup. Define two binary operations “ \times_S ” and “ \star_S ” on $P(S)$ as follows: For all $e, f \in P(S)$,*

$$(31) \quad e \times_S f = (ef)^+, e \star_S f = (ef)^*.$$

Then $(P(S), \times_S, \star_S)$ forms a projection algebra and is called the projection algebra of S . In particular, if S is DRC-restriction, then $P(S)$ is strong. Moreover, if S is P -restriction, then $P(S)$ is symmetric; If S is restriction, then $P(S)$ is commutative.

Let $(S, \cdot, +, *)$ be a DRC semigroup. Define a partial order “ ω_S ” on $P(S)$ as follows: For all $e, f \in P(S)$,

$$e\omega_S f \iff fef = e.$$

By DRC Conditions and Lemma 5.3, for all $x, y \in S$ and $e, f \in P(S)$, we have

$$(32) \quad (xy)^+ \omega_S x^+, (xy)^* \omega_S y^*, (ef)^+ \omega_S e, (ef)^* \omega_S f.$$

Define binary relations “ \leq_S^r ” and “ \leq_S^l ” on S as follows: For all $a, b \in S$,

$$(33) \quad a \leq_S^r b \iff a = a^+b, a^+ \omega_S b^+,$$

$$(34) \quad a \leq_S^l b \iff a = ba^*, a^* \omega_S b^*.$$

By (i), (i)' and Lemma 5.3, we can obtain the following result easily.

Lemma 5.6. *Let $(S, \cdot, +, *)$ be a DRC semigroup. Then “ \leq_S^r ” and “ \leq_S^l ” are partial orders on S , and for all $e, f \in P(S)$,*

$$e \leq_S^r f \iff e\omega_S f \iff e \leq_S^l f \iff e \leq_{P(S)} f,$$

where $\leq_{P(S)}$ is the natural partial order of the projection algebra $(P(S), \times_S, \star_S)$.

Lemma 5.7 ([57]). *Let $(S, \cdot, +, *)$ be a DRC semigroup and $a, b \in S$.*

- (1) *If $a \leq_S^r b$ or $a \leq_S^l b$, then $a^+ \omega_S b^+$, $a^* \omega_S b^*$.*
- (2) *$a \leq_S^r b$ if and only if there exists $e \in P(S)$ such that $a = eb$ and $e\omega_S b^+$.*
- (3) *$a \leq_S^l b$ if and only if there exists $e \in P(S)$ such that $a = be$ and $e\omega_S b^*$.*
- (4) *For all $x \in S$ and $e \in P(S)$, if $e\omega_S x^+$, then there exists a unique element $z \in S$ such that $z^+ = e$ and $z \leq_S^l x$, this element is exactly ex .*
- (5) *For all $x \in S$ and $e \in P(S)$, if $e\omega_S x^*$, then there exists a unique element $z \in S$ such that $z^* = e$, $z \leq_S^r x$, this element is exactly xe .*

Lemma 5.8. *Let $(S, \cdot, +, *)$ be a DRC semigroup. Then $\leq_S^r = \leq_S^l$ if and only if S is DRC-restriction.*

Proof. Let $\leq_S^l \subseteq \leq_S^r$ and $x, y \in S$. By (32), $(x^*y)^+ \omega_S x^{**} = x^*$, and so $x(x^*y)^+ \leq_S^l x$ by Lemma 5.7 (3). By hypothesis, $x(x^*y)^+ \leq_S^r x$, which implies that

$$x(x^*y)^+ = (x(x^*y)^+)^+ x = (xx^*y)^+ x = (xy)^+ x$$

by (ii) and (i)'. This gives (vi)'. Conversely, assume that (vi)' is satisfied, $a, b \in S$ and $a \leq_S^l b$. Then $a = ba^*$ and $a^* \omega_S b^*$. Thus we have $b^*a^* = a^*$ and

$$a^+b = (ba^*)^+b = b(b^*a^*)^+ = b(a^*)^+ = ba^* = a, \quad b^+a^+b^+ = b^+(ba^*)^+b^+ = (ba^*)^+ = a^+$$

by DRC-conditions. This implies that $a \leq_S^r b$. Therefore $\leq_S^l \subseteq \leq_S^r$. The above statement together with its dual now gives the desired result. \square

Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup. By lemma 5.8, $\leq_S^r = \leq_S^l$. In this case, denote $\leq_S = \leq_S^r = \leq_S^l$ and call it the *natural partial order* of S .

Corollary 5.9. *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup and $a, b, c, d \in S$.*

- (1) *If $a \leq_S b$, then $a^+ \leq_S b^+$ and $a^* \leq_S b^*$.*
- (2) *$a \leq_S b$ if and only if there exists $e \in P(S)$ such that $a = eb$ and $e\omega_S b^+$.*
- (3) *$a \leq_S b$ if and only if there exists $e \in P(S)$ such that $a = be$ and $e\omega_S b^*$.*
- (4) *$a \leq_S b$ if and only if $a = a^+b = ba^*$.*
- (5) *If $a \leq_S b, c \leq_S d$ and $a^* = c^+, b^* = d^+$, then $ac \leq_S bd$.*

- (6) For all $x \in S$ and $e \in P(S)$, if $e \leq_S x^+$, then there exists a unique element $z \in S$ such that $z^+ = e$ and $z \leq_S x$, this element is exactly ex . In this case, we denote this unique element by ${}_e \downarrow x$, that is, ${}_e \downarrow x = ex$.
- (7) For all $x \in S$ and $e \in P(S)$, if $e \leq_S x^*$, then there exists a unique element $z \in S$ such that $z^* = e$ and $z \leq_S x$, this element is exactly xe . In this case, we denote this unique element by $x \downarrow_e$, that is, $x \downarrow_e = xe$.

Proof. By Lemma 5.7, we only need to prove (4) and (5).

(4) If $a \leq_S b$, then we have $a = a^+b = ba^*$ by (33), (34) and Lemma 5.8. Conversely, assume that $a = a^+b = ba^*$. Then $a^+ = (ba^*)^+ = b^+(ba^*)^+b^+ = b^+a^+b^+$ by (iii), and so $a^+\omega_S b^+$. By (33) and Lemma 5.8, we have $a \leq_S b$.

(5) Let $a \leq_S b$ and $c \leq_S d$. Then $a = a^+b, a^+\omega_S b^+, c = c^+d, c^+\omega_S d^+$. By DRC Conditions and the hypothesis,

$$(ac)^+ = (ac^+)^+ = (aa^*)^+ = a^+, (bd)^+ = (bd^+)^+ = (bb^*)^+ = b^+,$$

and so $(ac)^+\omega_S (bd)^+$. Moreover, $(ac)^+bd = a^+bd = ad = aa^*d = ac^+d = ac$. Thus $ac \leq_S bd$. \square

Let $(S, \cdot, +, *)$ be a DRC semigroup. In $(P(S), \times_S, \star_S)$, denote

$$\langle e \rangle = \{x \in P(S) \mid x \leq_{P(S)} e\} = \{x \in P(S) \mid x\omega_S e\}$$

for any $e \in P(S)$ (see Lemma 5.6). By (32), we can define

$$(35) \quad \rho_a : \langle a^+ \rangle \rightarrow \langle a^* \rangle, x \mapsto (xa)^*; \quad \sigma_a : \langle a^* \rangle \rightarrow \langle a^+ \rangle, y \mapsto (ay)^+$$

for all $a \in S$. In particular, by Lemma 5.3 we have

$$(36) \quad \rho_e = \sigma_e = \text{the identity map on } \langle e \rangle$$

for all $e \in P(S)$.

Lemma 5.10 ([59]). *Let $(S, \cdot, +, *)$ be a DRC semigroup. Then S is DRC-restriction if and only if for all $a \in S$, ρ_a and σ_a are mutually inverse bijections. In this case, for all $a \in S$, ρ_a and σ_a are mutually inverse isomorphisms between projection algebras.*

Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup. Define the *restricted product* “ \circ_S ” of S as follows:

$$a \circ_S b = \begin{cases} ab & \text{if } a^* = b^+, \\ \text{undefined} & \text{otherwise,} \end{cases}$$

where ab is the multiplication of a and b in S . Moreover, define the following maps:

$$(37) \quad \mathbf{d}_S : S \rightarrow S, x \rightarrow x^+, \quad \mathbf{r}_S : S \rightarrow S, x \rightarrow x^*.$$

$$(38) \quad \varepsilon_S : \mathcal{P}(P(S)) \rightarrow S, (p_1, p_2, \dots, p_k) \mapsto p_1 p_2 \cdots p_k.$$

where $\mathcal{P}(P(S))$ is the path category of $(P(S), \times_S, \star_S)$. Now we can state our main result in this section.

Theorem 5.11. *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup. Then*

$$\mathbf{C}(S) = (S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_S, \times_S, \star_S, \varepsilon_S)$$

is a chain projection ordered category, where $P_S = P(S)$. Moreover, we have the following results:

- If S is a generalized regular \circ -semigroup, then $\mathbf{C}(S)$ is a chain projection ordered groupoid.
- If S is P -restriction, then $\mathbf{C}(S)$ is symmetric chain projection ordered category.
- If S is a regular \circ -semigroup, then $\mathbf{C}(S)$ is a symmetric chain projection ordered groupoid.
- If S is restriction, Then $\mathbf{C}(S)$ is a commutative chain projection ordered category.
- If S is an inverse semigroup, then $\mathbf{C}(S)$ is a commutative chain projection ordered groupoid.

Proof. Let $x, y, z \in S$. By the definitions of \circ_S , \mathbf{d}_S and \mathbf{r}_S , $x \circ_S y$ is defined if and only if $x^* = y^+$, if and only if $\mathbf{r}_S(x) = \mathbf{d}_S(y)$. In this case,

$$\mathbf{d}_S(x \circ_S y) = (xy)^+ = (xy^+)^+ = (xx^*)^+ = x^+ = \mathbf{d}_S(x).$$

Dually, $\mathbf{r}_S(x \circ_S y) = \mathbf{r}_S(y)$. If $x \circ_S y$ and $y \circ_S z$ are defined, by the above discussions $(x \circ_S y) \circ_S z$ and $x \circ_S (y \circ_S z)$ are defined and

$$(x \circ_S y) \circ_S z = (xy)z = x(yz) = x \circ_S (y \circ_S z).$$

Since $\mathbf{r}_S(\mathbf{d}_S(x)) = \mathbf{r}_S(x^+) = (x^+)^* = x^+ = \mathbf{d}_S(x)$, it follows that $\mathbf{d}(x) \circ_S x$ is defined and $\mathbf{d}_S(x) \circ_S x = \mathbf{d}_S(x)x = x^+x = x$. Dually, $x \circ_S \mathbf{r}_S(x)$ is defined and $x \circ_S \mathbf{r}_S(x) = x$. Thus $C = (S, \circ_S, \mathbf{d}_S, \mathbf{r}_S)$ is a category whose set of objects is

$$P_S = \{\mathbf{d}_S(x) \mid x \in S\} = \{x^+ \mid x \in S\} = P(S).$$

By Lemma 5.9, $(S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S)$ is an ordered category, and for all $x \in S$ and $e, f \in P_S = P(S)$, if $e \leq_S \mathbf{d}_S(x) = x^+$ and $f \leq_S \mathbf{r}_S(x) = x^*$, then $e \downarrow x = ex$ and $x \downarrow f = xf$. By Lemma 5.5, (P_S, \times_S, \star_S) is a strong projection algebra. Moreover, $e \leq_S f$ if and only if $e \leq_{P_S} f$ for all $e, f \in P_S$ by Lemma 5.6. This shows that $(S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_S, \times_S, \star_S)$ forms a weak projection ordered category. Let $x \in S$. Then in $(S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_S, \times_S, \star_S)$,

$$\mathbf{d}_S(x)^\downarrow = \{e \in P_S \mid e \leq_S \mathbf{d}_S(x)\}, \quad \mathbf{r}_S(x)^\downarrow = \{f \in P_S \mid f \leq_S \mathbf{r}_S(x)\},$$

$$(39) \quad \nu_x : (x^+)^\downarrow = \mathbf{d}_S(x)^\downarrow \rightarrow \mathbf{r}_S(x)^\downarrow = (x^*)^\downarrow, \quad p \rightarrow \mathbf{r}_S(p \downarrow x) = (px)^*,$$

$$(40) \quad \mu_x : (x^*)^\downarrow = \mathbf{r}_S(x)^\downarrow \rightarrow \mathbf{d}_S(x)^\downarrow = (x^+)^\downarrow, \quad q \rightarrow \mathbf{d}_S(x \downarrow q) = (xq)^+.$$

By (35), we have

$$(41) \quad \nu_x = \rho_x \text{ and } \mu_x = \sigma_x$$

Lemma 5.10 gives that μ_x and ν_x are mutually inverse isomorphisms between projection algebras. Thus, $(S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_S, \times_S, \star_S)$ is a projection ordered category. Define

$$\varepsilon_S : \mathcal{P}(P_S) \rightarrow S, \quad (p_1, p_2, \dots, p_k) \rightarrow p_1 p_2 \dots p_k.$$

In the followings, we shall prove that ε_S is an evaluation map. Let $\mathbf{p} = (p_1, p_2, \dots, p_k), \mathbf{q} = (q_1, q_2, \dots, q_l) \in \mathcal{P}(P_S)$. Then

$$p_1, p_2, \dots, p_k, q_1, q_2, \dots, q_l \in P_S, \quad p_1 \mathcal{F}_{P_S} p_2 \mathcal{F}_{P_S} \dots \mathcal{F}_{P_S} p_k, \quad q_1 \mathcal{F}_{P_S} q_2 \mathcal{F}_{P_S} \dots \mathcal{F}_{P_S} q_l.$$

Firstly, let $p \in P_S$. Then $\varepsilon_S(p) = p$ and $\varepsilon_S(p, p) = pp = p$, and so (E1) holds. Secondly,

$$\begin{aligned} \mathbf{d}_S(\varepsilon_S(\mathbf{p})) &= \mathbf{d}_S(p_1 p_2 \dots p_k) = (p_1 p_2 \dots p_k)^+ = (p_1 (p_2 \dots p_k)^+)^+ \\ &= p_1 \times_S (p_2 \dots p_k)^+ = (p_2 \dots p_k)^+ \delta_{p_1} = \dots = p_k \delta_{p_{k-1}} \dots \delta_{p_1} = p_1 = \mathbf{d}((p_1, p_2, \dots, p_k)) \end{aligned}$$

by (ii) and Lemma 3.5. Dually, $\mathbf{r}_S(\varepsilon_S(\mathbf{p})) = p_k = \mathbf{r}((p_1, p_2, \dots, p_k))$. This gives (E2). If $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{q})$, i.e. $p_k = q_1$, then $\mathbf{p} \circ \mathbf{q} = (p_1, p_2, \dots, p_{k-1}, p_k = q_1, q_2, \dots, q_l)$. Since $p_k q_1 = p_k p_k = p_k$, we have

$$\begin{aligned} \varepsilon_S(\mathbf{p} \circ \mathbf{q}) &= p_1 p_2 \dots p_{k-1} p_k q_2 \dots q_l = (p_1 p_2 \dots p_{k-1} p_k)(q_1 q_2 \dots q_l) \\ &= \varepsilon_S((p_1, p_2, \dots, p_{k-1}, p_k)) \circ_S \varepsilon_S((q_1, q_2, \dots, q_l)). \end{aligned}$$

This implies (E3) is true. Let $q \in P_S$, $q \leq_S \mathbf{d}(\mathbf{p}) = p_1$, and $q \downarrow \mathbf{p} = (s_1, s_2, \dots, s_k)$, where

$$\begin{aligned} s_i &= q \theta_{p_1} \theta_{p_2} \dots \theta_{p_i} = (qp_1)^* \theta_{p_2} \dots \theta_{p_i} = ((qp_1)^* p_2)^* \theta_{p_3} \dots \theta_{p_i} = (qp_1 p_2)^* \theta_{p_3} \dots \theta_{p_i} \\ &= \dots = (qp_1 p_2 \dots p_i)^* = p_i^* (qp_1 p_2 \dots p_i)^* = p_i (qp_1 p_2 \dots p_i)^*, i = 1, 2, \dots, k. \end{aligned}$$

If $k = 1$, then $q \downarrow \mathbf{p} = q$. Since $q \leq_S \mathbf{d}(\mathbf{p}) = p_1$, we have $q = qp_1$, and so

$$\varepsilon_S(q \downarrow \mathbf{p}) = \varepsilon_S(q) = q = qp_1 = q \downarrow p_1 = q \downarrow \varepsilon_S(p_1) = q \downarrow \varepsilon_S(\mathbf{p}).$$

Denote $\mathbf{t} = (p_1, p_2, \dots, p_{k-1})$. Then $q \downarrow \mathbf{t} = (s_1, s_2, \dots, s_{k-1})$. Assume that $\varepsilon_S(q \downarrow \mathbf{t}) = q \downarrow \varepsilon_S(\mathbf{t})$. Then $s_1 s_2 \dots s_{k-1} = qp_1 p_2 \dots p_{k-1}$. Thus

$$\begin{aligned} \varepsilon_S(q \downarrow \mathbf{p}) &= s_1 s_2 \dots s_{k-1} s_k \\ &= qp_1 p_2 \dots p_{k-1} p_k (qp_1 p_2 \dots p_{k-1} p_k)^* = qp_1 p_2 \dots p_{k-1} p_k = q \downarrow \varepsilon_S(\mathbf{p}). \end{aligned}$$

By mathematical induction, for all $q \in P_S$ and $\mathbf{p} \in \mathcal{P}(P_S)$, $q \leq_S \mathbf{d}_S(\mathbf{p})$ implies that $\varepsilon_S(q \downarrow \mathbf{p}) = q \downarrow \varepsilon_S(\mathbf{p})$. This shows that (E5) is true, and so (E4) is also true by Proposition 4.11. Thus ε_S is an evaluation map.

Now let $a \in S$. Then by (20), (41) and (35),

$$(42) \quad p \Theta_a = p \theta_{\mathbf{d}_S(a)} \nu_a = p(\theta_{a^+} \nu_a) = (p \star_S a^+) \nu_a = ((pa^+)^* a)^* = (pa^+ a)^* = (pa)^*$$

for all $p \in P_S$. Dually,

$$(43) \quad p \Delta_a = (ap)^+ \text{ for all } p \in P_S.$$

Let $b \in S, e, f \in P_S$ and (e, f) be a b -linked pair. Then $\mathbf{d}_S(b) = b^+$ and $\mathbf{r}_S(b) = b^*$. By (26),

$$\begin{aligned} e_1 &= e \theta_{\mathbf{d}_S(b)} = e \star_S b^+ = (eb^+)^*, \quad e_2 = f \Delta_b = (bf)^+, \\ f_1 &= e \Theta_b = (eb)^*, \quad f_2 = f \delta_{\mathbf{r}_S(b)} = b^* \times_S f = (b^* f)^+. \end{aligned}$$

According to (27), Lemma 5.3 and DRC Conditions,

$$\begin{aligned} \lambda(e, b, f) &= \varepsilon_S(e, e_1) \circ_S e_1 \downarrow b \circ_S \varepsilon_S(f_1, f) = ee_1 e_1 b f_1 f \\ &= ee_1 b f_1 f = e(eb^+)^* b (eb)^* f = eb^+ (eb^+)^* b^+ b (eb)^* f = eb^+ b (eb)^* f = (eb)(eb)^* f = eb f. \end{aligned}$$

Dually, we can prove that $\rho(e, b, f) = eb f$. Thus $\lambda(e, b, f) = \rho(e, b, f)$. We have shown that $\mathbf{C}(S)$ is a chain projection ordered category.

Finally, if $(S, \cdot, +, *, \circ)$ is a generalized regular \circ -semigroup, then $\mathbf{d}_S(x) = x^+ = x \circ_S x^\circ$ and $\mathbf{r}(x) = x^* = x^\circ \circ_S x$ by (29). This shows that $(S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \circ)$ is a groupoid, and so $\mathbf{C}(S)$ is a chain projection ordered groupoid. The other cases can be proved by using Lemmas 5.1 and 5.5, we omit the details. \square

Lemma 5.12. *Let $(S_1, \cdot, +, *)$ and $(S_2, \cdot, +, *)$ be two DRC-restriction semigroups and θ be a $(2,1,1)$ -homomorphism from S_1 to S_2 . Then the rule $\mathbf{C}(\theta) : S_1 \rightarrow S_2$, $x \rightarrow x\theta$ provides a chain projection ordered functor from $\mathbf{C}(S_1)$ to $\mathbf{C}(S_2)$.*

Proof. Firstly, let $x, y \in S_1$. Then

$$\mathbf{d}(x\theta) = (x\theta)^+ = x^+\theta = (\mathbf{d}(x))\theta, \quad \mathbf{r}(x\theta) = (x\theta)^* = x^*\theta = (\mathbf{r}(x))\theta.$$

This gives (F1). If $x \circ y$ is defined, then $x^* = \mathbf{r}(x) = \mathbf{d}(y) = y^+$, and so $(x\theta)^* = x^*\theta = y^+\theta = (y\theta)^+$. This implies $(x\theta) \circ (y\theta)$ is defined and

$$(x \circ y)\theta = (xy)\theta = (x\theta)(y\theta) = (x\theta) \circ (y\theta).$$

Thus (F2) holds. Let $x \leq_{S_1} y$. Then $x = x^+y = yx^*$ by Corollary 5.9 (4), and so $x\theta = (x\theta)^+(y\theta) = (y\theta)(x\theta)^*$. This shows that $x\theta \leq_{S_2} y\theta$. So (F3) holds. Let $e, f \in P_{S_1}$. Then

$$(e \times_{S_1} f)\theta = (ef)^+\theta = ((e\theta)(f\theta))^+ = (e\theta) \times_{S_2} (f\theta).$$

Dually, $(e \star_{S_1} f)\theta = (e\theta) \star_{S_2} (f\theta)$. Thus (F4) is true. Finally, let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P_{S_1})$. Then

$$(\varepsilon_1(\mathbf{p}))\theta = (p_1 p_2 \dots p_k)\theta = (p_1\theta)(p_2\theta) \dots (p_k\theta) = \varepsilon_2((p_1\theta, p_2\theta, \dots, p_k\theta)).$$

Thus (F5) holds. The above discussion tells us that $\mathbf{C}(\theta)$ is a chain projection functor. \square

6. THE CATEGORY ISOMORPHISM

In this section, we first show how to construct a DRC-restriction semigroup from a chain projection ordered category, and then prove that the category of DRC-restriction semigroups together with (2,1,1)-homomorphisms is isomorphic to the category of chain projection ordered categories together with chain projection ordered functors.

Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category and $a, b \in C$. By Lemma 3.3,

$$(\mathbf{r}(a) \times \mathbf{d}(b), \mathbf{r}(a) \star \mathbf{d}(b)) \in \mathcal{F}_{P_C}, \quad \mathbf{r}(a) \times \mathbf{d}(b) \leq \mathbf{r}(a), \mathbf{r}(a) \star \mathbf{d}(b) \leq \mathbf{d}(b).$$

Thus $a \downarrow_{\mathbf{r}(a) \times \mathbf{d}(b)}, \varepsilon(\mathbf{r}(a) \times \mathbf{d}(b), \mathbf{r}(a) \star \mathbf{d}(b))$ and $\mathbf{r}(a) \star \mathbf{d}(b) \downarrow b$ are all defined. By (1) and (2) in Lemma 2.2 and Lemma 4.12, we can define

$$(44) \quad a \bullet b = a \downarrow_{\mathbf{r}(a) \times \mathbf{d}(b)} \circ \varepsilon(\mathbf{r}(a) \times \mathbf{d}(b), \mathbf{r}(a) \star \mathbf{d}(b)) \circ \mathbf{r}(a) \star \mathbf{d}(b) \downarrow b.$$

Lemma 6.1. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category and $a, b, c \in C$.*

- (1) *If $\mathbf{r}(a) = \mathbf{d}(b)$, then $a \bullet b = a \circ b$.*
- (2) *$\mathbf{r}(a \bullet b) = \mathbf{r}(a)\Theta_b$ and $\mathbf{d}(a \bullet b) = \mathbf{d}(b)\Delta_a$.*
- (3) *$\Theta_{a \bullet b} = \Theta_a\Theta_b$ and $\Delta_{a \bullet b} = \Delta_b\Delta_a$.*
- (4) *$\mathbf{r}((a \bullet b) \bullet c) = \mathbf{r}(a)\Theta_b\Theta_c = \mathbf{r}(a \bullet (b \bullet c))$ and $\mathbf{d}(a \bullet (b \bullet c)) = \mathbf{d}(c)\Delta_b\Delta_a = \mathbf{d}((a \bullet b) \bullet c)$.*

Proof. Denote $p = \mathbf{r}(a), q = \mathbf{d}(b), p' = q\delta_p = p \times q, q' = p\theta_q = p \star q$.

- (1) If $p = q$, then by (L1) and (R1), we have $p' = q' = p$, and so

$$a \bullet b = a \downarrow_p \circ \varepsilon(p, p) \circ p \downarrow b = a \circ p \circ b = (a \circ \mathbf{r}(a)) \circ b = a \circ b.$$

by (3) and (4) in Lemma 2.2, (E1) and (C3).

- (2) By (44), we have

$$\mathbf{r}(a \bullet b) = \mathbf{r}(q' \downarrow b) = q'\nu_b = p\theta_q\nu_b = \mathbf{r}(a)\theta_{\mathbf{d}(b)}\nu_b = \mathbf{r}(a)\Theta_b.$$

Dually, $\mathbf{d}(a \bullet b) = \mathbf{d}(b)\Delta_a$.

- (3) Let $t \in P_C$. By (R4), (P3), (R4) and (R3),

$$t\theta_{p'}\theta_{q'} = (t \star p') \star q' = t\theta_{q\delta_p}\theta_{p\theta_q} = (t \star (p \times q)) \star (p \star q) = ((t \star (p \times q)) \star q) \star (p \star q)$$

$$= ((t \star p) \star q) \star (p \star q) = (t \star p) \star (p \star q) = (t \star p) \star q = t\theta_p\theta_q.$$

This implies that $\theta_{p'}\theta_{q'} = \theta_p\theta_q$. Using (44), Lemma 4.6 (6), (G1c), Lemma 4.6 (5), Lemma 4.12, Lemma 3.2, Lemma 4.6 (5) and the fact $\theta_{p'}\theta_{q'} = \theta_p\theta_q$ in order, we have

$$\begin{aligned} \Theta_{a \bullet b} &= \Theta_{a|_{p'}} \Theta_{\varepsilon(p', q')} \Theta_{q'|_b} = \Theta_a \theta_{p'} \theta_{p'} \theta_{q'} \theta_{q'} \Theta_b \\ &= \Theta_a \theta_{p'} \theta_{q'} \Theta_b = \Theta_a \theta_p \theta_q \Theta_b = \Theta_a \theta_{\mathbf{r}(a)} \theta_{\mathbf{d}(b)} \Theta_b = \Theta_a \Theta_b. \end{aligned}$$

Dually, $\Delta_{a \bullet b} = \Delta_b \Delta_a$.

(4) By (2) and (3) of the present lemma, we have

$$\mathbf{r}((a \bullet b) \bullet c) = \mathbf{r}(a \bullet b) \Theta_c = \mathbf{r}(a) \Theta_b \Theta_c = \mathbf{r}(a) \Theta_{b \bullet c} = \mathbf{r}(a \bullet (b \bullet c)).$$

Dually, $\mathbf{d}(a \bullet (b \bullet c)) = \mathbf{d}(c) \Delta_b \Delta_a = \mathbf{d}((a \bullet b) \bullet c)$. \square

Lemma 6.2. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category and $a, b, c \in C$, and denote*

$$p = \mathbf{r}(a), q = \mathbf{d}(b), r = \mathbf{r}(b), s = \mathbf{d}(c).$$

(1) $(a \bullet b) \bullet c = a|_e \circ \varepsilon(e, e_1) \circ_{e_1} b \circ \varepsilon(f_1, f) \circ_f c$, where

$$e = s\Delta_b\delta_p, e_1 = e\theta_q, f_1 = e\Theta_b, f = p\Theta_b\theta_s.$$

(2) $a \bullet (b \bullet c) = a|_e \circ \varepsilon(e, e_2) \circ_b b|_{f_2} \circ \varepsilon(f_2, f) \circ_f c$, where

$$e = s\Delta_b\delta_p, e_2 = f\Delta_b, f_2 = f\delta_r, f = p\Theta_b\theta_s.$$

(3) $(a \bullet b) \bullet c = a \bullet (b \bullet c)$. This implies that (C, \bullet) forms a semigroup.

Proof. (1) Denote $p' = q\delta_p = p \times q, q' = p\theta_q = p \star q$. Then

$$a \bullet b = a' \circ \varepsilon' \circ b', a' = a|_{p'}, \varepsilon' = \varepsilon(p', q'), b' = q'|_b.$$

By Lemma 6.1 (2), we can let $t = \mathbf{r}(a \bullet b) = \mathbf{r}(b') = \mathbf{r}(q'|_b) = p\Theta_b$. Then

$$(a \bullet b) \bullet c = (a \bullet b)|_{t'} \circ \varepsilon(t', s') \circ_{s'} c, t' = s\delta_t = t \times s, s' = t\theta_s = t \star s.$$

Denote $\nu = \mathbf{d}(b'|_{t'})$ and $\mu = \mathbf{d}(\varepsilon'|_{\nu})$. By Lemma 2.2 (8),

$$(a \bullet b)|_{t'} = (a' \circ \varepsilon' \circ b')|_{t'} = a'|_{\mu} \circ \varepsilon'|_{\nu} \circ b'|_{t'}.$$

By Lemma 2.2 (2), (E2) and Lemma 3.1 (1),

$$\mu = \mathbf{d}(\varepsilon'|_{\nu}) \leq \mathbf{d}(\varepsilon') = \mathbf{d}(\varepsilon(p', q')) = \mathbf{d}(p', q') = p' = p \times q \leq p,$$

and so we have $a'|_{\mu} = (a|_{p'})|_{\mu} = a|_{\mu}$ by Lemma 2.2 (10). In view of Lemma 2.2 (2), we obtain

$$\varepsilon'|_{\nu} \leq \varepsilon' = \varepsilon(p', q'), \mathbf{d}(\varepsilon'|_{\nu}) = \mathbf{r}(a'|_{\mu}) = \mu, \mathbf{r}(\varepsilon'|_{\nu}) = \nu.$$

By Lemma 4.12 (3), $\varepsilon'|_{\nu} = \varepsilon(\mu, \nu)$. Moreover, since $b'|_{t'} \leq b' = q'|_b \leq b$ and $\mathbf{r}(b'|_{t'}) = t'$, we have $b'|_{t'} = b|_{t'}$ by Lemma 2.2 (3). This gives that

$$b'|_{t'} = b|_{t'} = \mathbf{d}(b|_{t'}) \downarrow b = \mathbf{d}(b'|_{t'}) \downarrow b = \nu \downarrow b$$

by Lemma 2.2 (6). Thus

$$\begin{aligned} (a \bullet b) \bullet c &= (a \bullet b)|_{t'} \circ \varepsilon(t', s') \circ_{s'} c \\ &= a'|_{\mu} \circ \varepsilon'|_{\nu} \circ b'|_{t'} \circ \varepsilon(t', s') \circ_{s'} c = a|_{\mu} \circ \varepsilon(\mu, \nu) \circ_{\nu} b \circ \varepsilon(t', s') \circ_{s'} c. \end{aligned}$$

In the sequel, we shall prove the followings:

$$(a) \mu = e, \quad (b) \nu = e_1, \quad (c) t' = f_1, \quad (d) s' = f.$$

(a) By Lemma 6.1 (4), we have $\mathbf{d}(a|_\mu) = \mathbf{d}((a \bullet b) \bullet c) = s\Delta_b\Delta_a$. Since $a|_\mu \leq a$, it follows that $a|_\mu = s\Delta_b\Delta_a \downarrow a$ by Lemma 2.2 (3), and so

$$\mu = \mathbf{r}(a|_\mu) = \mathbf{r}(s\Delta_b\Delta_a \downarrow a) = s\Delta_b\Delta_a\nu_a = s\Delta_b\delta_p\mu_a\nu_a = s\Delta_b\delta_p = e$$

by Lemma 2.2 (2) and Lemma 2.3.

(c) By (G1b), $t' = s\delta_t = s\delta_{p\Theta_b} = s\Delta_b\delta_p\Theta_b = e\Theta_b = f_1$.

(b) Since $\nu \downarrow b = b|_{t'}$, we have

$$\nu = \mathbf{d}(\nu \downarrow b) = \mathbf{d}(b|_{t'}) = t'\mu_b = f_1\mu_b = (e\Theta_b)\mu_b = e\Theta_{\mathbf{d}(b)} = e\theta_q = e_1$$

by Lemma 2.2 (1), (c) and Lemma 4.6 (3).

(d) $s' = t\theta_s = p\Theta_b\theta_s = f$.

(2) This is the dual of (1).

(3) By (1) and (2), we only need to prove

$$(45) \quad \varepsilon(e, e_1) \circ e_1 \downarrow b \circ \varepsilon(f_1, f) = \varepsilon(e, e_2) \circ b|_{f_2} \circ \varepsilon(f_2, f).$$

By the fact $f \leq s$ and Lemma 3.2, we have $\theta_f = \theta_s\theta_f$. By Lemma 4.9,

$$e\Theta_b\theta_f = s\Delta_b\delta_p\Theta_b\theta_f = s\Delta_b\delta_p\Theta_b\theta_s\theta_f = p\Theta_b\theta_s\theta_f = p\Theta_b\theta_f = f.$$

Dually, $e = f\Delta_b\delta_e$. This implies that (e, f) is a b -lined pair. By (G2), (45) follows. \square

Lemma 6.3. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category, $a \in C$ and $s, t \in P_C$.*

- (1) $a \bullet t = a|_{\mathbf{r}(a) \times t} \circ \varepsilon(\mathbf{r}(a) \times t, \mathbf{r}(a) \star t)$. In particular, if $\mathbf{r}(a) \mathcal{F}_{P_C} t$, then $a \bullet t = a \circ \varepsilon(\mathbf{r}(a), t)$, if $t \leq \mathbf{r}(a)$, then $a \bullet t = a|_t$.
- (2) $t \bullet a = \varepsilon(t \times \mathbf{d}(a), t \star \mathbf{d}(a)) \circ t \star \mathbf{d}(a) \downarrow a$. In particular, if $t \mathcal{F}_{P_C} \mathbf{d}(a)$, then $t \bullet a = \varepsilon(t, \mathbf{d}(a)) \circ a$, if $t \leq \mathbf{d}(a)$, then $t \bullet a = t \downarrow a$.
- (3) $s \bullet t = \varepsilon(s \times t, s \star t)$, $\mathbf{d}(s \bullet t) = s \times t$, $\mathbf{r}(s \bullet t) = s \star t$. In particular, if $s \mathcal{F}_{P_C} t$, then $s \bullet t = \varepsilon(s, t)$, $\mathbf{d}(s \bullet t) = s$, $\mathbf{r}(s \bullet t) = t$.
- (4) $s \bullet (s \times t) \bullet s = s \times t$, $t \bullet (s \star t) \bullet t = s \star t$.
- (5) $s \leq t$ (i.e. $s \leq_{P_C} t$) if and only if $s = t \bullet s \bullet t$.

Proof. (1) Since $t \in P_C$, we have $\mathbf{r}(t) = t = \mathbf{d}(t)$ and $\mathbf{r}(s) = s = \mathbf{d}(s)$ by Remark 2.1. Denote $\mathbf{d}(a) = p$ and $\mathbf{r}(a) = q$. Then $q \star t \leq t$ by Lemma 3.1 (2), and so $q \star t|t = q \star t$ by Lemma 2.2 (12). By (E2), $\mathbf{r}(\varepsilon(q \times t, q \star t)) = \mathbf{r}(q \times t, q \star t) = q \star t$. This together with (C3) gives that $\varepsilon(q \times t, q \star t) \circ q \star t = \varepsilon(q \times t, q \star t)$. Thus

$$\begin{aligned} a \bullet t &= a|_{q \times t} \circ \varepsilon(q \times t, q \star t) \circ q \star t|t = a|_{q \times t} \circ \varepsilon(q \times t, q \star t) \circ (q \star t) \\ &= a|_{q \times t} \circ \varepsilon(q \times t, q \star t) = a|_{\mathbf{r}(a) \times t} \circ \varepsilon(\mathbf{r}(a) \times t, \mathbf{r}(a) \star t). \end{aligned}$$

If $\mathbf{r}(a) = q \mathcal{F}_{P_C} t$, then $q \times t = q$ and $q \star t = t$, and so

$$a \bullet t = a|_q \circ \varepsilon(q, t) = a|_{\mathbf{r}(a)} \circ \varepsilon(\mathbf{r}(a), t) = a \circ \varepsilon(\mathbf{r}(a), t)$$

by Lemma 2.2 (3). If $t \leq \mathbf{r}(a) = q$, then $q \times t = q \star t = t$ by (1) and (2) in Lemma 3.1. In view of (E1), Lemma 2.2 (2) and (C3), we obtain that

$$a \bullet t = a|_{q \times t} \circ \varepsilon(q \times t, q \star t) = a|_t \circ \varepsilon(t, t) = a|_t \circ t = a|_t \circ \mathbf{r}(a|_t) = a|_t.$$

(2) This is the dual of item (1).

(3) By Lemma 3.1 (1), we get $s \times t \leq s$, and so $s|_{s \times t} = s \times t$ by Lemma 2.2 (11). Now (E2) gives that $\mathbf{d}(\varepsilon(s \times t, s \star t)) = \mathbf{d}(s \times t, s \star t) = s \times t$. Dually, we have $\mathbf{r}(\varepsilon(s \times t, s \star t)) = s \star t$. This together with (C3) implies that

$$(s \times t) \circ \varepsilon(s \times t, s \star t) = \varepsilon(s \times t, s \star t).$$

In view of item (1) in this lemma, we obtain

$$s \bullet t = s|_{s \times t} \circ \varepsilon(s \times t, s \star t) = (s \times t) \circ \varepsilon(s \times t, s \star t) = \varepsilon(s \times t, s \star t).$$

If $s \mathcal{F}_{P_C} t$, then $s \times t = s$, $s \star t = t$, and so $s \bullet t = \varepsilon(s, t)$.

(4) By item (3) of the present lemma, (1) and (2) in Lemma 3.1 and (E1),

$$s \bullet (s \times t) = \varepsilon(s \times (s \times t), s \star (s \times t)) = \varepsilon(s \times t, s \times t) = s \times t,$$

$$(s \times t) \bullet s = \varepsilon(((s \times t) \times s, (s \times t) \star s)) = \varepsilon(s \times t, s \times t) = s \times t.$$

This gives that $s \bullet (s \times t) \bullet s = s \times t$. Dually, $t \bullet (s \star t) \bullet t = s \star t$.

(5) If $s \leq t$, then $s \times t = s \star t = s = t \times s = t \star s$ by (1) and (2) in Lemma 3.1. By item (3) in the present lemma, we have $s \bullet t = \varepsilon(s \times t, s \star t) = \varepsilon(s, s) = s$ by (E1), and so

$$t \bullet s \bullet t = t \bullet s = \varepsilon(t \times s, t \star s) = \varepsilon(s, s) = s.$$

Conversely, assume that $s = t \bullet s \bullet t$. Since $\mathbf{d}(\varepsilon(s \times t, s \star t)) = s \times t$ by Lemma 4.12 (2) and $\mathbf{r}(t) = t$, it follows that

$$s = t \bullet s \bullet t = t \bullet \varepsilon(s \times t, s \star t) = t|_{t \times (s \times t)} \circ U \circ V = (t \times (s \times t)) \circ U \circ V$$

for some $U, V \in C$ by item (3) in the present lemma, (44), Lemma 3.1 (1) and Lemma 2.2 (12). This implies that $s = \mathbf{d}(s) = \mathbf{d}(t \times (s \times t)) = t \times (s \times t) \leq t$ by the fact that $s, t \times (s \times t) \in P_C$, Remark 2.1 and Lemma 3.1 (1). \square

Theorem 6.4. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category. Define*

$$a \bullet b = a|_{\mathbf{r}(a) \times \mathbf{d}(b)} \circ \varepsilon(\mathbf{r}(a) \times \mathbf{d}(b), \mathbf{r}(a) \star \mathbf{d}(b)) \circ \mathbf{r}(a) \star \mathbf{d}(b)|_b, \quad a^\clubsuit = \mathbf{d}(a), \quad a^\spadesuit = \mathbf{r}(a)$$

for all $a, b \in C$. Then $\mathbf{S}(C) = (C, \bullet, \clubsuit, \spadesuit)$ forms a DRC-restriction semigroup with set of projections $P(\mathbf{S}(C)) = P_C$. Moreover, we have the following results:

- If C is a chain projection ordered groupoid, then $\mathbf{S}(C)$ is a generalized regular \circ -semigroup.
- If C is symmetric chain projection ordered category, then $\mathbf{S}(C)$ is P -restriction.
- If C is a symmetric chain projection ordered groupoid, then $\mathbf{S}(C)$ is a regular \circ -semigroup.
- If C is commutative chain projection ordered category, then $\mathbf{S}(C)$ is restriction.
- If C is commutative chain projection ordered groupoid, then $\mathbf{S}(C)$ is an inverse semigroup.

Proof. By Lemma 6.2 (3), (C, \bullet) forms a semigroup. To show $\mathbf{S}(C)$ is a DRC-restriction semigroup, we need to prove (i)–(vi) by symmetry. Let $x, y \in C$.

(i) By (C3), $\mathbf{d}(x) \circ x$ is defined and $\mathbf{d}(x) \circ x = x$. In view of Lemma 6.1, we have $x^\clubsuit \bullet x = \mathbf{d}(x) \bullet x = \mathbf{d}(x) \circ x = x$. Thus (i) holds.

(ii) By the definition of the operation \bullet , we have $\mathbf{d}(x \bullet y) = \mathbf{d}(x|_{\mathbf{r}(x) \times \mathbf{d}(y)})$ and $\mathbf{d}(x \bullet y^\clubsuit) = \mathbf{d}(x|_{\mathbf{r}(x) \times \mathbf{d}(y^\clubsuit)})$. Since $\mathbf{d}(y^\clubsuit) = \mathbf{d}(\mathbf{d}(y)) = \mathbf{d}(y)$ by Remark 2.1, it follows that

$$(x \bullet y)^\clubsuit = \mathbf{d}(x \bullet y) = \mathbf{d}(x \bullet y^\clubsuit) = (x \bullet y^\clubsuit)^\clubsuit.$$

(iii) By Lemma 6.3 (3),

$$(46) \quad (x^\clubsuit \bullet y^\clubsuit)^\clubsuit = (\mathbf{d}(x) \bullet \mathbf{d}(y))^\clubsuit = \mathbf{d}(\mathbf{d}(x) \bullet \mathbf{d}(y)) = \mathbf{d}(x) \times \mathbf{d}(y).$$

This together with Lemma 6.3 (4) gives that

$$x^\clubsuit \bullet (x^\clubsuit \bullet y^\clubsuit)^\clubsuit \bullet x^\clubsuit = \mathbf{d}(x) \bullet (\mathbf{d}(x) \times \mathbf{d}(y)) \bullet \mathbf{d}(x) = \mathbf{d}(x) \times \mathbf{d}(y) = (x^\clubsuit \bullet y^\clubsuit)^\clubsuit.$$

By Corollary 5.4, (iii) is true.

(iv)–(v) In view of Remark 2.1, we have $(x^\clubsuit)^\clubsuit = \mathbf{d}(\mathbf{d}(x)) = \mathbf{d}(x) = x^\clubsuit$ and

$$(x^\clubsuit)^\spadesuit = (\mathbf{d}(x))^\spadesuit = \mathbf{r}(\mathbf{d}(x)) = \mathbf{d}(x) = x^\clubsuit.$$

By the above discussion and its dual, $\mathbf{S}(C) = (C, \bullet, \clubsuit, \spadesuit)$ forms a DRC semigroup with set of projections

$$(4\mathcal{P})(\mathbf{S}(C)) = \{x^\clubsuit \mid x \in C\} = \{\mathbf{d}(x) \mid x \in C\} = \{\mathbf{r}(x) \mid x \in C\} = \{x^\spadesuit \mid x \in C\} = P_C.$$

Let $e \in P(\mathbf{S}(C)) = P_C$ and $x \in C$. Then $\mathbf{r}(e) = e$ by Remark 2.1. Denote $t = e \star \mathbf{d}(x)$. By (1) and (2) in Lemma 3.1, we have $t \leq \mathbf{d}(x)$ and $t \times \mathbf{d}(x) = t \star \mathbf{d}(x) = t$. Moreover, Lemma 2.2 (11) gives that ${}_t\downarrow \mathbf{d}(x) = t$. Since $x^\clubsuit = \mathbf{d}(x)$, we have $(e \bullet x^\clubsuit)^\spadesuit = \mathbf{r}(e \bullet \mathbf{d}(x)) = \mathbf{r}(e \star \mathbf{d}(x)) = \mathbf{r}(t) = t$ by Lemma 6.3 (3). In view of Lemma 6.3 (2), (E1), Lemma 2.2 (1) and (C3), we have

$$\begin{aligned} (e \bullet x^\clubsuit)^\spadesuit \bullet x &= t \bullet x = \varepsilon(t \times \mathbf{d}(x), t \star \mathbf{d}(x)) \circ {}_{t \star \mathbf{d}(x)}\downarrow x = \\ &= \varepsilon(t, t) \circ {}_t\downarrow x = t \circ {}_t\downarrow x = \mathbf{d}({}_t\downarrow x) \circ {}_t\downarrow x = {}_t\downarrow x. \end{aligned}$$

Denote $u = \mathbf{r}({}_t\downarrow x)$. Then $u \in P_C$, and so $\mathbf{r}(u) = u$ by Remark 2.1. By Lemma 2.2 (1), we have $u \leq \mathbf{r}(x)$, and so

$$\mathbf{r}(x) \star u = \mathbf{r}(x) \star \mathbf{r}(u) = \mathbf{r}(x) \times u = \mathbf{r}(x) \times \mathbf{r}(u) = u$$

by (1) and (2) in Lemma 3.1. In view of Lemma 6.3 (2),

$$(e \bullet x)^\spadesuit = \mathbf{r}(e \bullet x) = \mathbf{r}({}_{e \star \mathbf{d}(x)}\downarrow x) = \mathbf{r}({}_t\downarrow x) = u.$$

This together with Lemma 6.3 (1), (E1), Lemma 2.2 (2) and (C3) yields that

$$\begin{aligned} x \bullet (e \bullet x)^\spadesuit &= x \bullet u = a|_{\mathbf{r}(x) \times \mathbf{r}(u)} \circ \varepsilon(\mathbf{r}(x) \times \mathbf{r}(u), \mathbf{r}(x) \star \mathbf{r}(u)) \\ &= x|_u \circ \varepsilon(u, u) = x|_u \circ u = x|_u \circ \mathbf{r}(x|_u) = x|_u. \end{aligned}$$

By the fact that $u = \mathbf{r}({}_t\downarrow x)$ and Lemma 2.2 (5), we have ${}_t\downarrow x = x|_u$. Thus $x \bullet (e \bullet x)^\spadesuit = (e \bullet x^\clubsuit)^\spadesuit \bullet x$. This proves (vi). Thus $\mathbf{S}(C)$ is a DRC-restriction semigroup by (4).

Finally, let $(C, \circ, \mathbf{d}, \mathbf{r}, ^{-1})$ be a groupoid and $x \in C$. By Lemmas 6.1 and 2.6,

$$x \bullet x^{-1} = x \circ x^{-1} = \mathbf{d}(x) = x^\clubsuit = \mathbf{d}(x) = \mathbf{r}(x^{-1}) = (x^{-1})^\spadesuit.$$

Dually, $x^{-1} \bullet x = x^\spadesuit = (x^{-1})^\clubsuit$. This implies that $\mathbf{S}(C) = (C, \bullet, \clubsuit, \spadesuit, -1)$ is a generalized regular \circ -semigroup by Remark 5.2. The other cases can be proved by Proposition 1.2, Lemma 5.1 and Remark 5.2, we omit the details. \square

Let $(S, \cdot, ^+, \star)$ be a DRC-restriction semigroup. We use $\mathbf{E}(S)$ to denote the subsemigroup of S generated by $P(S)$. We say that S is *projection-generated* if $\mathbf{E}(S) = S$.

Proposition 6.5. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category and $S = \mathbf{S}(C)$. Then*

$$P(S) = P_C, \quad \mathbf{E}(S) = \text{im}(\varepsilon) = \{\varepsilon(\mathbf{c}) \mid \mathbf{c} \in \mathcal{P}(P)\}.$$

Thus S is projection-generated if and only if ε is surjective.

Proof. By (47), $P(S) = P_C$. Let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P_C)$. Then by (24), Lemma 6.1 (1) and Lemma 6.3 (3),

$$\begin{aligned} \varepsilon(\mathbf{p}) &= \varepsilon(p_1, p_2) \circ \varepsilon(p_2, p_3) \circ \dots \circ \varepsilon(p_{k-1}, p_k) = \varepsilon(p_1, p_2) \bullet \varepsilon(p_2, p_3) \bullet \dots \bullet \varepsilon(p_{k-1}, p_k) \\ &= (p_1 \bullet p_2) \bullet (p_2 \bullet p_3) \bullet \dots \bullet (p_{k-1} \bullet p_k) = p_1 \bullet p_2 \bullet \dots \bullet p_k \in \mathbf{E}(S). \end{aligned}$$

Conversely, let $a \in \mathbf{E}(S)$. Then there exists a positive integer k and $p_1, p_2, \dots, p_k \in P$ such that $a = p_1 \bullet p_2 \bullet \dots \bullet p_k$. In the sequel, we shall prove that $a \in \text{im}\varepsilon$ by mathematical induction. When $k = 1$, we have $\varepsilon(p_1) = p_1$ by (E1), and so $p_1 \in \text{im}\varepsilon$. Let $p_1 \bullet p_2 \bullet \dots \bullet p_{k-1} = \varepsilon(\mathbf{p})$, where $\mathbf{p} \in \mathcal{P}(P)$. Denote $s = \mathbf{r}(\varepsilon(\mathbf{c})) \times p_k$ and $t = \mathbf{r}(\varepsilon(\mathbf{c})) \star p_k$. By Lemma 6.3 (1), (E6) and (E3),

$$a = p_1 \bullet p_2 \bullet \dots \bullet p_{k-1} \bullet p_k = \varepsilon(\mathbf{c}) \bullet p_k = \varepsilon(\mathbf{c})|_s \circ \varepsilon(s, t) = \varepsilon(\mathbf{c}|_s) \circ \varepsilon(s, t) = \varepsilon(\mathbf{c}|_s \circ (s, t)) \in \text{im}\varepsilon.$$

By mathematical induction, we have $a \in \text{im}\varepsilon$ for all $a \in \mathbf{E}(S)$. Thus $\mathbf{E}(S) \subseteq \text{im}\varepsilon$. This implies that $\mathbf{E}(S) = \text{im}\varepsilon$, and so S is projection-generated if and only if ε is surjective. \square

Lemma 6.6. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category and $S = \mathbf{S}(C)$. Then $\leq_S = \leq$.*

Proof. Let $x, y \in C$. Then

$$\begin{aligned} x \leq_S y &\iff x^\clubsuit = y^\clubsuit \bullet x^\clubsuit \bullet y^\clubsuit, x = x^\clubsuit \bullet y \quad (\text{by (33)}) \\ &\iff \mathbf{d}(x) = \mathbf{d}(y) \bullet \mathbf{d}(x) \bullet \mathbf{d}(y), x = \mathbf{d}(x) \bullet y \quad (\text{by (6.4)}) \\ &\iff \mathbf{d}(x) \leq \mathbf{d}(y), x = \mathbf{d}(x)|_y. \quad (\text{by (2) and (5) in Lemma 6.3}) \\ &\iff x \leq y. \quad (\text{by (O1), (O3) and Lemma 2.2 (3)}) \end{aligned}$$

Thus $\leq_S = \leq$. \square

Lemma 6.7. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category. Then the restricted product of $S = \mathbf{S}(C)$ is exactly “ \circ ” and $(P(S), \times_S, \star_S) = (P_C, \times, \star)$.*

Proof. Consider the restricted product of $\mathbf{S}(C)$: For all $x, y \in C$,

$$x \odot y = \begin{cases} x \bullet y & \text{if } x^\spadesuit = y^\spadesuit, \\ \text{undefined} & \text{otherwise,} \end{cases}$$

Let $x, y \in C$. Then $x^\spadesuit = \mathbf{r}(x), y^\spadesuit = \mathbf{d}(y)$, and so

$$x \odot y \text{ is defined} \iff \mathbf{r}(x) = \mathbf{d}(y) \iff x \circ y \text{ is defined,}$$

In this case, we have $x \odot y = x \bullet y = x \circ y$ by Lemma 6.1 (1). This shows that \odot is equal to \circ . On the other hand, we obtain $P(S) = P_C$ by (47). Let $e, f \in P_C = P(S)$. By Lemma 6.3 (3),

$$e \times_S f = (e \bullet f)^\clubsuit = \mathbf{d}(e \bullet f) = e \times f, \quad e \star_S f = (e \bullet f)^\spadesuit = \mathbf{r}(e \bullet f) = e \star f.$$

Thus $(P(S), \times_S, \star_S) = (P_C, \times, \star)$. \square

Lemma 6.8. *Let φ be a chain projection ordered functor from $(C_1, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_1}, \times, \star, \varepsilon_1)$ to $(C_2, \circ, \mathbf{d}, \mathbf{r}, \leq, P_{C_2}, \times, \star, \varepsilon_2)$. Then $\mathbf{S}\varphi : C_1 \rightarrow C_2$, $x \mapsto x\varphi$ provides a $(2, 1, 1)$ -homomorphism from $\mathbf{S}(C_1)$ to $\mathbf{S}(C_2)$.*

Proof. By (F1), $x^{\clubsuit}\varphi = (\mathbf{d}(x))\varphi = \mathbf{d}(x\varphi) = (x\varphi)^{\clubsuit}$. This together with its dual shows that $\mathbf{S}\varphi$ preserves “ \clubsuit ” and “ \spadesuit ”. Using (F2), Lemma 4.19, (F5), (F4) and (F1) in order,

$$\begin{aligned} (x \bullet y)\varphi &= (x|_{\mathbf{r}(x) \times \mathbf{d}(y)} \circ \varepsilon_1(\mathbf{r}(x) \times \mathbf{d}(y), \mathbf{r}(x) \star \mathbf{d}(y)) \circ_{\mathbf{r}(x) \star \mathbf{d}(y)} \downarrow y)\varphi \\ &= (x|_{\mathbf{r}(x) \times \mathbf{d}(y)})\varphi \circ (\varepsilon_1(\mathbf{r}(x) \times \mathbf{d}(y), \mathbf{r}(x) \star \mathbf{d}(y)))\varphi \circ_{(\mathbf{r}(x) \star \mathbf{d}(y))\varphi} \downarrow y\varphi \\ &= x\varphi|_{(\mathbf{r}(x) \times \mathbf{d}(y))\varphi} \circ \varepsilon_2((\mathbf{r}(x) \times \mathbf{d}(y))\varphi, (\mathbf{r}(x) \star \mathbf{d}(y))\varphi) \circ_{(\mathbf{r}(x) \star \mathbf{d}(y))\varphi} \downarrow y\varphi \\ &= x\varphi|_{\mathbf{r}(x)\varphi \times \mathbf{d}(y)\varphi} \circ \varepsilon_2(\mathbf{r}(x)\varphi \times \mathbf{d}(y)\varphi, \mathbf{r}(x)\varphi \star \mathbf{d}(y)\varphi) \circ_{\mathbf{r}(x)\varphi \star \mathbf{d}(y)\varphi} \downarrow y\varphi \\ &= x\varphi|_{\mathbf{r}(x\varphi) \times \mathbf{d}(y\varphi)} \circ \varepsilon_2(\mathbf{r}(x\varphi) \times \mathbf{d}(y\varphi), \mathbf{r}(x\varphi) \star \mathbf{d}(y\varphi)) \circ_{\mathbf{r}(x\varphi) \star \mathbf{d}(y\varphi)} \downarrow y\varphi = x\varphi \bullet y\varphi. \end{aligned}$$

Thus $\mathbf{S}\varphi$ is a (2,1,1)-homomorphism. \square

Lemma 6.9. *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup. Then we have $\mathbf{S}(\mathbf{C}(S)) = S$.*

Proof. By Theorem 5.11, $\mathbf{C}(S)$ is a chain projection ordered category and

$$(48) \quad \mathbf{d}_S(x) = x^+, \mathbf{r}_S(x) = x^*, e \times_S f = (ef)^+, e \star_S f = (ef)^*$$

for all $x \in S$ and $e, f \in P_S = P(S)$. Let $x, y \in S$. Then $x \circ_S y = xy$ if and only if

$$(49) \quad x^* = \mathbf{r}_S(x) = \mathbf{d}_S(y) = y^+$$

where xy is the product of x and y in (S, \cdot) . Moreover, for $x \in S, e, f \in P_S$ with $e \leq_S \mathbf{d}_S(x)$ and $f \leq_S \mathbf{r}_S(x)$, have $e|_x = ex$ and $x|_f = xf$. Define three operations on S as follows:

$$x \bullet y = x|_{\mathbf{r}_S(x) \times_S \mathbf{d}_S(y)} \circ \varepsilon_S(\mathbf{r}_S(x) \times_S \mathbf{d}_S(y), \mathbf{r}_S(x) \star_S \mathbf{d}_S(y))|_{\mathbf{r}_S(x) \star_S \mathbf{d}_S(y)} \downarrow y,$$

and $x^{\clubsuit} = \mathbf{d}_S(x), x^{\spadesuit} = \mathbf{r}_S(x)$, where ε_S is defined as in (38). By Theorem 6.4, $\mathbf{S}(\mathbf{C}(S)) = (S, \bullet, \clubsuit, \spadesuit)$ is a DRC-restriction semigroup, and

$$x^{\clubsuit} = \mathbf{d}_S(x) = x^+, x^{\spadesuit} = \mathbf{r}_S(x) = x^*$$

by (48). Furthermore,

$$\begin{aligned} (50) \quad x \bullet y &= x|_{\mathbf{r}_S(x) \times_S \mathbf{d}_S(y)} \circ \varepsilon_S(\mathbf{r}_S(x) \times_S \mathbf{d}_S(y), \mathbf{r}_S(x) \star_S \mathbf{d}_S(y)) \circ_{\mathbf{r}_S(x) \star_S \mathbf{d}_S(y)} \downarrow y \\ &= x|_{(x^*y^+)^+} \circ_S \varepsilon_S((x^*y^+)^+, (x^*y^+)^*) \circ_{(x^*y^+)^*} \downarrow y \\ &= x(x^*y^+)^+ \circ_S (x^*y^+)^+ (x^*y^+)^* \circ_S (x^*y^+)^* y \\ &= x(x^*y^+)^+ (x^*y^+)^* y = xx^*y^+y = xy. \quad (\text{by Lemma 5.3 (4), (i), (i)'}) \end{aligned}$$

Thus $\mathbf{S}(\mathbf{C}(S)) = S$. \square

Lemma 6.10. *Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category. Then we have $\mathbf{C}(\mathbf{S}(C)) = C$.*

Proof. Define three operations on C as follows:

$$x \bullet y = x|_{\mathbf{r}(x) \times \mathbf{d}(y)} \circ \varepsilon(\mathbf{r}(x) \times \mathbf{d}(y), \mathbf{r}(x) \star \mathbf{d}(y)) \circ_{\mathbf{r}(x) \star \mathbf{d}(y)} \downarrow y.$$

$$(51) \quad x^{\clubsuit} = \mathbf{d}(x), x^{\spadesuit} = \mathbf{r}(x).$$

Then by Theorem 6.4, $S = \mathbf{S}(C) = (C, \bullet, \clubsuit, \spadesuit)$ is a DRC-restriction semigroup and $P_C = P(S)$. By Theorem 5.11, we have $\mathbf{C}(S) = (C, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_{\mathbf{C}(S)}, \times_S, \star_S, \varepsilon_S)$ and $P_{\mathbf{C}(S)} = P(S)$. Let $x, y \in C$. Then $\mathbf{d}_S(x) = x^{\clubsuit} = \mathbf{d}(x)$ and $\mathbf{r}_S(x) = x^{\spadesuit} = \mathbf{r}(x)$. Thus $\mathbf{d}_S = \mathbf{d}, \mathbf{r}_S = \mathbf{r}$. Moreover, $x \circ_S y$ is defined if and only if $\mathbf{r}_S(x) = \mathbf{d}_S(y)$, if and only if $\mathbf{r}(x) = \mathbf{d}(y)$. In this case, we obtain $x \circ_S y = x \bullet y = x \circ y$ by Lemma 6.1. This shows that $\circ_S = \circ$. By Lemma 6.7, we get $(P_C, \times, \star) = (P_{\mathbf{C}(S)}, \times_S, \star_S)$. Lemma 6.6 gives that $\leq_S = \leq$, and the fact $P_C = P_{\mathbf{C}(S)}$ gives that $\mathcal{P}(P_C) = \mathcal{P}(P_{\mathbf{C}(S)})$.

Let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P_C)$. Then $p_1 \mathcal{F}_{P_C} p_2 \mathcal{F}_{P_C} \dots \mathcal{F}_{P_C} p_k$. If $k = 1$, we have $\varepsilon(\mathbf{p}) = \varepsilon(p_1) = p_1 = \varepsilon_S(p_1) = \varepsilon_S(\mathbf{p})$ by (E1). Denote $\mathbf{q} = (p_1, p_2, \dots, p_{k-1})$. Then $\mathbf{p} = \mathbf{q} \circ (p_{k-1}, p_k)$. Assume that $\varepsilon(\mathbf{q}) = \varepsilon_S(\mathbf{q}) = p_1 \bullet p_2 \bullet \dots \bullet p_{k-1}$. Then

$$\varepsilon_S(\mathbf{p}) = p_1 \bullet p_2 \bullet \dots \bullet p_{k-1} \bullet p_k = \varepsilon(\mathbf{q}) \bullet p_k.$$

By (E2), we have $\mathbf{r}(\varepsilon(\mathbf{q})) = \mathbf{r}(\mathbf{q}) = p_{k-1}$. Since $p_{k-1} \mathcal{F}_{P_C} p_k$, by Lemma 6.3 (1) and (E3) we obtain

$$\varepsilon_S(\mathbf{p}) = \varepsilon(\mathbf{q}) \bullet p_k = \varepsilon(\mathbf{q}) \circ \varepsilon(p_{k-1}, p_k) = \varepsilon(\mathbf{q} \circ (p_{k-1}, p_k)) = \varepsilon(\mathbf{p}).$$

Thus $\varepsilon_S = \varepsilon$. By the above discussions, we have $C = \mathbf{C}(S) = \mathbf{C}(\mathbf{S}(C))$. \square

Now we can give our main theorem in this paper.

Theorem 6.11. *The category \mathbf{DRS} of DRC-restriction semigroups together with $(2,1,1)$ -homomorphisms is isomorphic to the category \mathbf{CPOC} of chain projection ordered categories together with chain projection ordered functors.*

Proof. By Theorems 6.4, 5.11 and Lemmas 5.12, 6.8, 6.9 and 6.10,

$$\mathbf{S} : \mathbf{CPOC} \rightarrow \mathbf{DRS} : (C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon) \rightarrow \mathbf{S}(C),$$

$$\mathbf{C} : \mathbf{DRS} \rightarrow \mathbf{CPOC} : (S, \cdot, +, *) \rightarrow \mathbf{C}(S)$$

are mutually inverse functors. Thus the result holds. \square

Finally, we also have the following special cases.

Theorem 6.12. *The category of generalized regular \circ -semigroups (respectively, P -restriction semigroups, regular \circ -semigroups, restriction semigroups, inverse semigroups) together with $(2,1,1)$ -homomorphisms is isomorphic to the category of chain projection ordered groupoids (respectively, symmetric chain projection ordered categories, symmetric chain projection ordered groupoids, commutative chain projection ordered categories, commutative chain projection ordered groupoids) together with chain projection ordered functors.*

7. CHAIN SEMIGROUPS

In this section, we consider the chain semigroups determined by strong two-sided projection algebras and give a presentation of these semigroups by using Theorem 6.11. We first recall some basic results on categories given in [8]. Let $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r})$ be a category and \approx be an equivalence on C . Then \approx is called a *congruence* on \mathcal{C} if for all $a, b, u, v \in C$, the following conditions hold:

(V1) If $a \approx b$, then $\mathbf{d}(a) = \mathbf{d}(b)$ and $\mathbf{r}(a) = \mathbf{r}(b)$.

(V2) If $\mathbf{r}(u) = \mathbf{d}(a) = \mathbf{d}(b)$ and $a \approx b$, then $u \circ a = u \circ b$.

(V3) If $\mathbf{d}(v) = \mathbf{r}(a) = \mathbf{r}(b)$ and $a \approx b$, then $a \circ v = b \circ v$.

Let $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r})$ be a category and \approx be a congruence on \mathcal{C} . Let $a \in C$. Denote the \approx -class of C containing a by $[a]$. Moreover, denote $C/\approx = \{[a] \mid a \in C\}$. Define a partial binary operation “ \circ ” on C/\approx and two maps \mathbf{d}, \mathbf{r} from C/\approx to C/\approx as follows: For all $[a], [b] \in C/\approx$,

$$[a] \circ [b] = \begin{cases} [a \circ b] & \text{if } \mathbf{r}([a]) = \mathbf{d}([b]), \\ \text{undefined} & \text{otherwise,} \end{cases}$$

$$\mathbf{d}([a]) = [\mathbf{d}(a)], \mathbf{r}([a]) = [\mathbf{r}(a)].$$

Then it is easy to see that $\mathcal{C}/\approx = (C/\approx, \circ, \mathbf{d}, \mathbf{r})$ is a category, and is called *the quotient category* of \mathcal{C} with respect to \approx . In this case, the set of objects of \mathcal{C}/\approx is $P_{C/\approx} = \{[p] \mid p \in P_C\}$. Let $p, q \in P_C$. By (V1), it follows that $p = q$ if and only if $[p] = [q]$.

A congruence \approx on an ordered category $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r}, \leq)$ is called an *ordered congruence* if for all $a, b \in C$ and $p \in P_C$, the following condition holds:

$$(V4) \text{ If } a \approx b \text{ and } p \leq \mathbf{d}(a), \text{ then } p \downarrow a \approx p \downarrow b.$$

Lemma 7.1 ([8]). *Let \approx be an ordered congruence on an ordered category $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r}, \leq)$. Define the relation " \leq_{\approx} " on C/\approx as follows: For all $[a], [b] \in C/\approx$,*

$$(52) \quad [a] \leq_{\approx} [b] \iff \text{there exist } x \in [a], y \in [b] \text{ such that } x \leq y.$$

Let $[a], [b] \in C/\approx$. Then $[a] \leq_{\approx} [b]$ if and only if for all $y \in [b]$, there exists $x \in [a]$ such that $x \leq y$. Furthermore, $\mathcal{C}/\approx = (C/\approx, \cdot, \mathbf{d}, \mathbf{r}, \leq_{\approx})$ is an ordered category.

Lemma 7.2 ([8]). *Let $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r})$ be a category and Ω be a subset of $C \times C$. Assume that the following condition holds:*

$$(53) \quad (u, v) \in \Omega \implies \mathbf{d}(u) = \mathbf{d}(v) \text{ and } \mathbf{r}(u) = \mathbf{r}(v).$$

Denote the least congruence of \mathcal{C} containing Ω by Ω^{\sharp} . Then for all $a, b \in C$, $(a, b) \in \Omega^{\sharp}$ if and only if there exists a sequence

$$a = a_1 \rightarrow \cdots \rightarrow a_k = b,$$

where

$$a_i = b_i \circ u_i \circ c_i, a_{i+1} = b_i \circ v_i \circ c_i, b_i, c_i \in C, (u_i, v_i) \in \Omega \cup \Omega^{-1}, \\ \Omega^{-1} = \{(v, u) \mid (u, v) \in \Omega\}, i = 1, 2, \dots, k.$$

Lemma 7.3 ([8]). *Let $\mathcal{C} = (C, \circ, \mathbf{d}, \mathbf{r}, \leq)$ be an ordered category and Ω be a subset of $C \times C$ satisfying (53). Assume that the following condition holds:*

$$(54) \quad (a, b) \in \Omega \implies_p \downarrow a \approx_p \downarrow b \text{ for all } p \in P_C \text{ with } p \leq \mathbf{d}(a).$$

Then Ω^{\sharp} is an ordered congruence on \mathcal{C} .

Let (P, \times, \star) be a strong projection algebra and $p, e, f \in P$. Then (e, p, f) is called an *admissible triple* of P if

$$(55) \quad f = e\theta_p\theta_f = (e \star p) \star f, \quad e = f\delta_p\delta_e = e \times (p \times f),$$

that is, $f = (e \star p) \star f$, $e = e \times (p \times f)$.

Lemma 7.4. *Let (P, \times, \star) be a strong projection algebra and (e, p, f) be an admissible triple of P . Then*

$$(e, e\theta_p, f) = (e, e \star p, f), (e, f\delta_p, f) = (e, p \times f, f) \in \mathcal{P}(P),$$

where $\mathcal{P}(P)$ is the path category of P .

Proof. By Lemma 3.1 (1), we have $p \times f \leq_P p$. This together with (55) and (5) and (1) in Lemma 3.1 gives that $e = e \times (p \times f) \leq_P e \times p \leq_P e$, and so $e \times p = e$. By Lemma 3.1 (3), $e \times e\theta_p = e \times (e \star p) = e \times p = e$. In view of (R4) and (R1), we have

$$e \star (e\theta_p) = e \star (e \star p) = (e \star p) \star (e \star p) = e \star p = e\theta_p.$$

This implies that $e \mathcal{F}_P e\theta_p$. On the other hand, denote $t = e\theta_p$. By (55), (R4) and (R1),

$$e\theta_p \star f = t \star (t \star f) = (t \star f) \star (t \star f) = t \star f = e\theta_p\theta_f = f,$$

and Lemma 3.7 and (55) imply that

$$e\theta_p \times f = f\delta_{e\theta_p} = f\delta_p\delta_e\theta_p = e\theta_p.$$

This shows that $e\theta_p \mathcal{F}_P f$. Thus $(e, e\theta_p, f) \in \mathcal{P}(P)$. Dually, we have $(e, f\delta_p, f) \in \mathcal{P}(P)$. \square

Lemma 7.5. *Let (P, \times, \star) be a strong projection algebra, (e, p, f) be an admissible triple of P and $e' \in P$, $e' \leq_P e$. Denote $f' = e'\theta_p\theta_f$. Then (e', p, f') is also an admissible triple of P and*

$$e' \downarrow (e, e\theta_p, f) = (e', e'\theta_p, f'), \quad e' \downarrow (e, f\delta_p, f) = (e', f'\delta_p, f').$$

Proof. Since (e, p, f) is an admissible triple of P , we have $f = e\theta_p\theta_f$ and $e = f\delta_p\delta_e$. Denote $t = e'\theta_p$. By (R4) and (R1),

$$e'\theta_p\theta_{f'} = e'\theta_p\theta_{e'\theta_p\theta_f} = t\theta_{t\theta_f} = t \star (t \star f) = (t \star f) \star (t \star f) = t \star f = e'\theta_p\theta_f = f'.$$

On the other hand, using Lemma 3.7, Lemma 3.1 (3), Lemma 3.2, (55), (11) and the fact that $e' \leq_P e$ in order, we have

$$\begin{aligned} f'\delta_p\delta_{e'} &= e'\theta_p\theta_f\delta_p\delta_{e'} = e'\theta_{f\delta_p}\delta_{e'} = e' \times (e' \star f\delta_p) \\ &= e' \times f\delta_p = f\delta_p\delta_{e'} = f\delta_p\delta_e\delta_{e'} = e\delta_{e'} = e' \times e = e'. \end{aligned}$$

Thus (e', p, f') is an admissible triple of P . By (16),

$$e' \downarrow (e, e\theta_p, f) = (e', e'\theta_{e\theta_p}, e'\theta_{e\theta_p}\theta_f), \quad e' \downarrow (e, f\delta_p, f) = (e', e'\theta_{f\delta_p}, e'\theta_{f\delta_p}\theta_f).$$

According to Lemma 3.1 (5) and the fact that $e' \leq_P e$, we have $e' \star p \leq e \star p$, and so

$$e'\theta_{e\theta_p} = e' \star (e \star p) = (e' \star p) \star (e \star p) = e' \star p = e'\theta_p$$

and $e'\theta_{e\theta_p}\theta_f = e'\theta_p\theta_f = f'$ by (R4) and Lemma 3.1 (2). Moreover, the definition of f' , (P3) and Lemma 3.7 give

$$e'\theta_{f\delta_p}\theta_f = (e' \star (p \times f)) \star f = (e' \star p) \star f = e'\theta_p\theta_f = f'$$

and $e'\theta_{f\delta_p} = e'\theta_p\theta_f\delta_p = f'\delta_p$. This implies that

$$e' \downarrow (e, e\theta_p, f) = (e', e'\theta_p, f'), \quad e' \downarrow (e, f\delta_p, f) = (e', f'\delta_p, f').$$

Thus the result follows. \square

Let (P, \times, \star) be a strong projection algebra. Let $\Omega_1(P) = \{(p, p) \mid p \in P\}$,

$$\Omega_2(P) = \{((e, e \star p, f), (e, p \times f, f)) \mid (e, p, f) \text{ is an admissible triple of } P\}.$$

Write $\Omega(P) = \Omega_1(P) \cup \Omega_2(P)$. Obviously, $\Omega(P)$ satisfies (53). We use \approx_P to denote the congruence $\Omega^\sharp(P)$ on $\mathcal{P}(P)$ generated by $\Omega(P)$.

Lemma 7.6. *Let (P, \times, \star) be a strong projection algebra. Then \approx_P is an ordered congruence on $\mathcal{P}(P)$.*

Proof. Let $(\mathfrak{s}, \mathfrak{t}) \in \Omega(P)$. If

$$p, r \in P, \mathfrak{s} = (p, p), \mathfrak{t} = p, r \leq_P p = \mathbf{d}(\mathfrak{s}) = \mathbf{d}(\mathfrak{t}),$$

then $r\theta_p = r \star p = r$ by Lemma 3.1 (2), and so $r \downarrow \mathfrak{s} = (r, r\theta_p) = (r, r)$, $r \downarrow \mathfrak{t} = r$ by (16). This implies that $(r \downarrow \mathfrak{s}, r \downarrow \mathfrak{t}) \in \Omega(P)$, and hence $r \downarrow \mathfrak{s} \approx_P r \downarrow \mathfrak{t}$. Assume that (e, p, f) is an admissible triple of P and

$$\mathfrak{s} = (e, e\theta_p, f), \mathfrak{t} = (e, f\delta_p, f), e' \in P, e' \leq_P e = \mathbf{d}(\mathfrak{s}) = \mathbf{d}(\mathfrak{t}), f' = e'\theta_p\theta_f.$$

By Lemma 7.5, (e', p, f') is also an admissible triple of P and

$$e' \downarrow \mathfrak{s} = (e', e' \theta_p, f'), \quad e' \downarrow \mathfrak{t} = (e', f' \delta_p, f').$$

This implies that $(e' \downarrow \mathfrak{s}, e' \downarrow \mathfrak{t}) \in \Omega(P)$, and so $e' \downarrow \mathfrak{s} \approx_P e' \downarrow \mathfrak{t}$. The above statement shows that $\Omega(P)$ satisfies (54). Since $\Omega(P)$ satisfies (53), \approx_P is an ordered congruence on $\mathcal{P}(P)$ by Lemma 7.3. \square

Let (P, \times, \star) be a strong projection algebra. Denote $\mathcal{C}(P) = \mathcal{P}(P)/\approx_P$. By Lemmas 7.1 and 7.6, $(\mathcal{C}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\approx_P}, P_{\mathcal{C}(P)}, \times, \star)$ is an ordered category, and is called the *chain category* of (P, \times, \star) . In fact, $\mathcal{C}(P)$ is a weak projection ordered category. Several key points of this ordered category are listed as follows:

- The elements in $\mathcal{C}(P)$: For any $\mathfrak{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$, denote the \approx_P -class containing \mathfrak{p} by $[\mathfrak{p}] = [p_1, p_2, \dots, p_k]$. Then $\mathcal{C}(P) = \{[\mathfrak{p}] \mid \mathfrak{p} \in \mathcal{P}(P)\}$.
- For any $[\mathfrak{p}] = [p_1, p_2, \dots, p_k] \in \mathcal{C}(P)$, we have $\mathbf{d}([\mathfrak{p}]) = [\mathbf{d}(\mathfrak{p})] = [p_1]$ and $\mathbf{r}([\mathfrak{p}]) = [\mathbf{r}(\mathfrak{p})] = [p_k]$, and so

$$P_{\mathcal{C}(P)} = \{[p] \mid p \in P\}.$$

- For all $p, q \in P$, $[p] = [q]$ if and only if $p = q$. This implies that $(P_{\mathcal{C}(P)}, \times, \star)$ forms a strong projection algebra, where $[p] \times [q] = [p \times q]$ and $[p] \star [q] = [p \star q]$ for all $p, q \in P$. So we can identify $[p]$ with p for any $p \in P$, and identify $(P_{\mathcal{C}(P)}, \times, \star)$ with (P, \times, \star) .
- If $\mathfrak{p} = (p_1, p_2, \dots, p_k), \mathfrak{q} = (q_1, q_2, \dots, q_l) \in \mathcal{P}(P)$ and $\mathbf{r}[\mathfrak{p}] = [p_k] = [q_1] = \mathbf{d}[\mathfrak{q}]$ (i.e. $p_k = q_1$), then

$$[\mathfrak{p}] \circ [\mathfrak{q}] = [p \circ q] = [p_1, \dots, p_{k-1}, p_k = q_1, q_2, \dots, q_l].$$

- If $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}(P)$, then

$$\begin{aligned} [\mathfrak{p}] \leq_{\approx_P} [\mathfrak{q}] &\iff \text{there exist } \mathfrak{m}, \mathfrak{n} \in \mathcal{P}(P) \text{ such that } \mathfrak{m} \in [\mathfrak{p}], \mathfrak{n} \in [\mathfrak{q}] \text{ and } \mathfrak{m} \leq_{\mathcal{P}} \mathfrak{n} \\ &\iff \text{there exists } \mathfrak{m} \in \mathcal{P}(P) \text{ such that } \mathfrak{m} \in [\mathfrak{p}] \text{ and } \mathfrak{m} \leq_{\mathcal{P}} \mathfrak{q}. \end{aligned}$$

- For any $[\mathfrak{p}] \in \mathcal{C}(P)$ and $r, s \in P$, if $[r] \leq_{\approx_P} \mathbf{d}([\mathfrak{p}])$, $[s] \leq_{\approx_P} \mathbf{r}([\mathfrak{p}])$, then

$$[r] \downarrow [\mathfrak{p}] = [r \downarrow \mathfrak{p}], \quad [\mathfrak{p}] \downarrow_s = [\mathfrak{p} \downarrow_s].$$

- For all $p, q \in P$,

$$p \leq_{\mathcal{P}} q \iff p \leq_P q \iff [p] \leq_{P_{\mathcal{C}(P)}} [q] \iff [p] \leq_{\approx_P} [q].$$

To prove that $(\mathcal{C}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\approx_P}, P_{\mathcal{C}(P)}, \times, \star)$ is a projection ordered category, we need some calculations. Let $\mathfrak{c} = [p_1, p_2, \dots, p_k] \in \mathcal{C}(P)$. (Notice that q denotes $[q]$ and p_i denotes $[p_i]$ in the sequel). Then $\mathbf{d}(\mathfrak{c}) = p_1$ and $\mathbf{r}(\mathfrak{c}) = p_k$. By (16) and Lemma 3.2,

$$(56) \quad \nu_{\mathfrak{c}} : p_1 \downarrow \rightarrow p_k \downarrow, \quad q \mapsto q \nu_{\mathfrak{c}} = \mathbf{r}(q \downarrow \mathfrak{c}) = q \theta_{p_1} \theta_{p_2} \dots \theta_{p_k},$$

$$\Theta_{\mathfrak{c}} : P_{\mathcal{C}(P)} \rightarrow P_{\mathcal{C}(P)}, \quad q \mapsto q \Theta_{\mathfrak{c}} = q \theta_{\mathbf{d}(\mathfrak{c})} \nu_{\mathfrak{c}} = q \theta_{p_1} \theta_{p_1} \theta_{p_2} \dots \theta_{p_k} = q \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}.$$

This implies that

$$(57) \quad \Theta_{\mathfrak{c}} = \theta_{p_1} \theta_{p_2} \dots \theta_{p_k}.$$

Dually,

$$(58) \quad \mu_{\mathfrak{c}} : p_k \downarrow \rightarrow p_1 \downarrow, \quad q \mu_{\mathfrak{c}} = \mathbf{d}(\mathfrak{c} \downarrow q) = q \delta_{p_k} \delta_{p_{k-1}} \dots \delta_{p_1},$$

$$(59) \quad \Delta_{\mathfrak{c}} = \delta_{p_k} \delta_{p_{k-1}} \dots \delta_{p_1}.$$

Lemma 7.7. *Let (P, \times, \star) be a strong projection algebra. Then*

$$(\mathcal{C}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\approx_P}, P_{\mathcal{C}(P)}, \times, \star)$$

is a projection ordered category.

Proof. Let $\mathbf{c} = [p_1, p_2, \dots, p_k] \in \mathcal{C}(P)$ (Notice that q denotes $[q]$ and p_i denotes $[p_i]$ in the sequel) and $q \leq_{\approx_P} \mathbf{d}(\mathbf{c}) = p_1$. By (56), Corollary 3.8, (59) and (57),

$$\delta_{q\nu_{\mathbf{c}}} = \delta_{q\theta_{p_1}\theta_{p_2}\dots\theta_{p_k}} = \delta_{p_k}\delta_{p_{k-1}}\dots\delta_{p_1}\delta_q\theta_{p_1}\theta_{p_2}\dots\theta_{p_k} = \Delta_{\mathbf{c}}\delta_q\Theta_{\mathbf{c}}.$$

Dually, we have $\theta_{q\mu_{\mathbf{c}}} = \Theta_{\mathbf{c}}\theta_q\Delta_{\mathbf{c}}$ for all $q \in P$ with $q \leq_{\approx_P} \mathbf{r}(\mathbf{c}) = p_k$. Thus (G1a) is true, and so the result follows. \square

The following proposition is obvious.

Proposition 7.8. *Let (P, \times, \star) be a strong projection algebra. Then the natural map*

$$\varepsilon_P \approx_{\mathcal{P}}^{\natural}: \mathcal{P}(P_{\mathcal{C}(P)}) \rightarrow \mathcal{C}(P), ([p_1], [p_2], \dots, [p_k]) \mapsto [p_1, p_2, \dots, p_k]$$

is an evaluation map on $(\mathcal{C}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\approx_P}, P_{\mathcal{C}(P)}, \times, \star)$.

Proposition 7.9. *Let (P, \times, \star) be a strong projection algebra. Then $(\mathcal{C}(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{\approx_P}, P_{\mathcal{C}(P)}, \times, \star, \varepsilon_P)$ is a chain projection ordered category.*

Proof. Let $[\mathbf{p}] \in \mathcal{C}(P)$ and $[e], [f] \in P_{\mathcal{C}(P)}$, where $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$ and $e, f \in P$. Assume that $([e], [f])$ is a $[\mathbf{p}]$ -linked pair. Then

$$(60) \quad [f] = [e]\Theta_{[\mathbf{p}]} \theta_{[f]}, \quad [e] = [f]\Delta_{[\mathbf{p}]} \delta_{[e]}.$$

By the definition of ε_P and (26)–(28),

$$(61) \quad \lambda([e], [\mathbf{p}], [f]) = [e, e_1] \circ_{[e_1] \downarrow} [\mathbf{p}] \circ [f_1, f], \quad \rho([e], [\mathbf{p}], [f]) = [e, e_2] \circ [\mathbf{p}] \upharpoonright_{[f_2]} \circ [f_2, f],$$

where

$$[e_1] = [e]\theta_{[p_1]}, \quad [e_2] = [f]\Delta_{[\mathbf{p}]}, \quad [f_1] = [e]\Theta_{[\mathbf{p}]}, \quad [f_2] = [f]\delta_{[p_k]}, \quad e_1, e_2, f_1, f_2 \in P.$$

Denote

$$[e_1] \downarrow [\mathbf{p}] = [e_1] \downarrow \mathbf{p} = [u_1, \dots, u_k], \quad [\mathbf{p}] \upharpoonright_{[f_2]} = [\mathbf{p}] \upharpoonright_{f_2} = [v_1, \dots, v_k].$$

By (15), (16) and the facts $e_1 = e\theta_{p_1}$ and $f_2 = f\delta_{p_k}$, for all $1 \leq i \leq k$, we have

$$u_i = e\theta_{p_1} \dots \theta_{p_i}, \quad v_i = f\delta_{p_k} \dots \delta_{p_i}.$$

This together with (61) gives that

$$\lambda([e], [\mathbf{p}], [f]) = (e, u_1, \dots, u_k, f), \quad \rho([e], [\mathbf{p}], [f]) = (e, v_1, \dots, v_k, f).$$

Denote $u_0 = e$ and $v_{k+1} = f$. We shall show that (u_{i-1}, p_i, v_{i+1}) is an admissible triple of P for all $1 \leq i \leq k$, that is,

$$v_{i+1} = u_{i-1}\theta_{p_i}\theta_{v_{i+1}}, \quad u_{i-1} = v_{i+1}\delta_{p_i}\delta_{u_{i-1}}.$$

In view of (57), (59) and (60), we have

$$f = e\theta_{p_1} \dots \theta_{p_k}\theta_f, \quad e = f\delta_{p_k} \dots \delta_{p_1}\delta_e.$$

By Lemma 3.8,

$$\begin{aligned} u_{i-1}\theta_{p_i}\theta_{v_{i+1}} &= e\theta_{p_1} \dots \theta_{p_{i-1}}\theta_{p_i}\theta_f\delta_{p_k} \dots \delta_{p_{i+1}} \\ &= e\theta_{p_1} \dots \theta_{p_{i-1}}\theta_{p_i}\theta_{p_{i+1}} \dots \theta_{p_k}\theta_f\delta_{p_k} \dots \delta_{p_{i+1}} = f\delta_{p_k} \dots \delta_{p_{i+1}} = v_{i+1}. \end{aligned}$$

Dually,

$$\begin{aligned} v_{i+1}\delta_{p_i}\delta_{u_{i-1}} &= f\delta_{p_k}\cdots\delta_{p_{i+1}}\delta_{p_i}\delta_{e\theta_{p_1}\cdots\theta_{p_{i-1}}} \\ &= f\delta_{p_k}\cdots\delta_{p_{i+1}}\delta_{p_i}\delta_{p_{i-1}}\cdots\delta_{p_i}\delta_e\theta_{p_1}\cdots\theta_{p_{i-1}} = e\theta_{p_1}\cdots\theta_{p_{i-1}} = u_{i-1}. \end{aligned}$$

Thus (u_{i-1}, p_i, v_{i+1}) is an admissible triple for all $1 \leq i \leq k$, and so $(u_{i-1}, u_{i-1}\theta_{p_i}, v_{i+1}) = (u_{i-1}, u_i, v_{i+1})$ is $\approx_P^{\mathfrak{h}}$ -related to $(u_{i-1}, v_{i+1}\delta_{p_i}, v_{i+1}) = (u_{i-1}, v_i, v_{i+1})$. Therefore

$$(62) \quad [u_{i-1}, u_i, v_{i+1}] = [u_{i-1}, v_i, v_{i+1}] \text{ for all } 1 \leq i \leq k.$$

This gives that

$$\begin{aligned} \rho([e], [\mathfrak{p}], [f]) &= [e, v_1, \dots, v_k, f] = [u_0, v_1, v_2, v_3, v_4, \dots, v_k, v_{k+1}] \\ &= [u_0, u_1, v_2, v_3, v_4, \dots, v_k, v_{k+1}] = [u_0, u_1, u_2, v_3, v_4, \dots, v_k, v_{k+1}] \\ &= \cdots [u_0, u_1, u_2, u_3, u_4, \dots, u_k, v_{k+1}] = [e, u_1, \dots, u_k, f] = \lambda([e], [\mathfrak{p}], [f]) \end{aligned}$$

Hence (G2) holds, and the desired result now follows. \square

Definition 7.10. Let $(P, \times \star)$ be a strong projection algebra. By Proposition 7.9 and Theorem 6.4, we have the DRC-restriction semigroup $\mathbf{S}(\mathcal{C}(P))$ and call it the chain semigroup determined by (P, \times, \star) .

Let

$$[\mathfrak{p}] = [p_1, p_2, \dots, p_k], [\mathfrak{q}] = [q_1, q_2, \dots, q_l] \in \mathcal{C}(P).$$

Then $\mathbf{r}([\mathfrak{p}]) = [p_k]$, $\mathbf{d}([\mathfrak{q}]) = [q_1]$. Denote $[p'] = [p_k \times q_1]$, $[q'] = [p_k \star q_1]$. By the definition of ε_P , we have

$$[\mathfrak{p}] \bullet [\mathfrak{q}] = [\mathfrak{p}] \downarrow_{[p']} \circ [p', q'] \circ [q'] \downarrow_{[\mathfrak{q}]}.$$

By (16) and (18) we let $[\mathfrak{p}] \downarrow_{[p']} = [p'_1, \dots, p'_k]$, $[q'] \downarrow_{[\mathfrak{q}]} = [q'_1, \dots, q'_l]$, where $p' = p'_k$, $q' = q'_1$ and

$$p'_i = p' \delta_{p_k} \cdots \delta_{p_i}, q'_j = q' \theta_{q_1} \cdots \theta_{q_j}, i = 1, 2, \dots, k, j = 1, 2, \dots, l.$$

This implies that

$$\begin{aligned} [\mathfrak{p}] \bullet [\mathfrak{q}] &= [p'_1, \dots, p'_k] \circ [p'_k, q'_1] \circ [q'_1, \dots, q'_l] = [p'_1, \dots, p'_k, q'_1, \dots, q'_l], \\ [\mathfrak{p}]^+ &= \mathbf{d}([\mathfrak{p}]) = [p_1], [\mathfrak{p}]^* = \mathbf{r}([\mathfrak{p}]) = [p_k]. \end{aligned}$$

Moreover, $P(\mathbf{S}(\mathcal{C}(P))) = P_{\mathcal{C}(P)} = \{[p] \mid p \in P\}$.

Proposition 7.11. Let $(C, \circ, \mathbf{d}, \mathbf{r}, \leq, P_C, \times, \star, \varepsilon)$ be a chain projection ordered category. Then

$$\bar{\varepsilon} : \mathcal{C}(P_C) \rightarrow C, [\mathfrak{p}] \mapsto \varepsilon(\mathfrak{p})$$

is a chain projection ordered functor from $\mathcal{C}(P_C)$ to C , and so is a $(2,1,1)$ -homomorphism from $\mathbf{S}(\mathcal{C}(P_C))$ to $\mathbf{S}(C)$ such that $[p]\bar{\varepsilon} = p$.

Proof. We first show that $\bar{\varepsilon}$ is well-defined. Let $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}(P_C)$ and $\mathfrak{p} = \mathfrak{b} \circ \mathfrak{s} \circ \mathfrak{c}$, $\mathfrak{q} = \mathfrak{b} \circ \mathfrak{t} \circ \mathfrak{c}$, where $\mathfrak{b}, \mathfrak{c} \in \mathcal{P}(P_C)$, $(\mathfrak{s}, \mathfrak{t}) \in \Omega(P_C) \cup \Omega^{-1}(P_C)$. If $\mathfrak{s} = (p, p)$, $\mathfrak{t} = p$ or $\mathfrak{s} = p$, $\mathfrak{t} = (p, p)$, where $p \in P_C$, then by (E1) we have $\varepsilon(p, p) = p = \varepsilon(p)$, and so $\varepsilon(\mathfrak{s}) = \varepsilon(\mathfrak{t})$. Assume that (e, p, f) is an admissible triple in P_C . Then

$$(63) \quad f = e\theta_p\theta_f, e = f\delta_p\delta_e$$

Let $\mathfrak{s} = (e, e\theta_p, f)$, $\mathfrak{t} = (e, f\delta_p, f)$ or $\mathfrak{s} = (e, f\delta_p, f)$, $\mathfrak{t} = (e, e\theta_p, f)$. We assume the former case holds without loss of generality. By (24),

$$(64) \quad \varepsilon(\mathfrak{s}) = \varepsilon(e, e\theta_p) \circ \varepsilon(e\theta_p, f), \varepsilon(\mathfrak{t}) = \varepsilon(e, f\delta_p) \circ \varepsilon(f\delta_p, f).$$

By using the fact that $p \in P_C$ and (22), we have $\Theta_p = \theta_p, \Delta_p = \delta_p$, and hence $f = e\Theta_p\theta_f$ and $e = f\Delta_p\delta_e$ by (63). This implies that (e, f) is (in the sense of Definition 4.13) p -linked pair. By (G2), (27), (28), (26) and the fact that $\mathbf{d}(p) = p$, we have

$$\varepsilon(e, e_1) \circ_{e_1} \downarrow p \circ \varepsilon(f_1, f) = \varepsilon(e, e_2) \circ_{e_2} \downarrow p \circ \varepsilon(f_2, f),$$

where

$$e_1 = e\theta_{\mathbf{d}(p)} = e\theta_p, e_2 = f\Delta_p = f\delta_p, f_1 = e\Theta_p = e\theta_p, f_2 = f\delta_{\mathbf{r}(p)} = f\delta_p.$$

This gives that

$$(65) \quad \varepsilon(e, e\theta_p) \circ_{e\theta_p} \downarrow p \circ \varepsilon(e\theta_p, f) = \varepsilon(e, f\delta_p) \circ_{f\delta_p} \downarrow p \circ \varepsilon(f\delta_p, f).$$

Using the fact that $e\theta_p \leq p, f\delta_p \leq p$ and Lemma 2.2 (11), we get $e\theta_p \downarrow p = e\theta_p$ and $f\delta_p \downarrow p = f\delta_p$. Observe that $\mathbf{r}(\varepsilon(e, e\theta_p)) = e\theta_p$ and $\mathbf{r}(\varepsilon(e, f\delta_p)) = f\delta_p$ by Lemma 4.12. This together with (C3) provides

$$(66) \quad \varepsilon(e, e\theta_p) \circ_{e\theta_p} \downarrow p = \varepsilon(e, e\theta_p), \quad \varepsilon(e, f\delta_p) \circ_{f\delta_p} \downarrow p = \varepsilon(e, f\delta_p).$$

By (64), (65) and (66), we have $\varepsilon(\mathbf{s}) = \varepsilon(\mathbf{t})$. The above discussion shows that $\varepsilon(\mathbf{s}) = \varepsilon(\mathbf{t})$ in either case. Thus by (E3),

$$\varepsilon(\mathbf{p}) = \varepsilon(\mathbf{b} \circ \mathbf{s} \circ \mathbf{c}) = \varepsilon(\mathbf{b}) \circ \varepsilon(\mathbf{s}) \circ \varepsilon(\mathbf{c}) = \varepsilon(\mathbf{b}) \circ \varepsilon(\mathbf{t}) \circ \varepsilon(\mathbf{c}) = \varepsilon(\mathbf{b} \circ \mathbf{t} \circ \mathbf{c}) = \varepsilon(\mathbf{q}).$$

Let $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P_C)$ and $[\mathbf{p}] = [\mathbf{q}]$. Then $\mathbf{p} \approx_{P_C} \mathbf{q}$. By Lemma 7.2 and the above statements, we have $\varepsilon(\mathbf{p}) = \varepsilon(\mathbf{q})$. This shows that $\bar{\varepsilon}$ is well-defined.

In the sequel, we show that $\bar{\varepsilon}$ is a chain projection ordered functor. Let

$$[\mathbf{p}], [\mathbf{q}] \in \mathcal{C}(P_C), e, f \in P_C, \mathbf{p} = (p_1, p_2, \dots, p_k), \mathbf{q} = (q_1, q_2, \dots, q_k) \in \mathcal{P}(P_C).$$

(1) Firstly,

$$\bar{\varepsilon}(\mathbf{d}([\mathbf{p}])) = \bar{\varepsilon}([\mathbf{d}(\mathbf{p})]) = \bar{\varepsilon}([p_1]) = \varepsilon(p_1) = p_1 = \mathbf{d}(\mathbf{p}) = \mathbf{d}(\varepsilon(\mathbf{p})) = \mathbf{d}(\bar{\varepsilon}([\mathbf{p}])).$$

Dually, $\bar{\varepsilon}(\mathbf{r}([\mathbf{p}])) = p_k = \mathbf{r}(\bar{\varepsilon}([\mathbf{p}]))$.

(2) If $\mathbf{r}([\mathbf{p}]) = \mathbf{d}([\mathbf{q}])$, then $p_k = q_1$ and

$$\bar{\varepsilon}([\mathbf{p}] \circ [\mathbf{q}]) = \bar{\varepsilon}([\mathbf{p} \circ \mathbf{q}]) = \varepsilon(\mathbf{p} \circ \mathbf{q}) = \varepsilon(\mathbf{p}) \circ \varepsilon(\mathbf{q}) = \bar{\varepsilon}([\mathbf{p}]) \circ \bar{\varepsilon}([\mathbf{q}]).$$

(3) Let $[\mathbf{p}] \leq_{\mathcal{P}} [\mathbf{q}]$. Then there exist $\mathbf{m}, \mathbf{n} \in \mathcal{P}(P_C)$ such that $[\mathbf{p}] = [\mathbf{m}], [\mathbf{q}] = [\mathbf{n}]$ and $\mathbf{m} \leq_{\mathcal{P}} \mathbf{n}$ in $\mathcal{P}(P_C)$. Since ε is order-preserving, we have

$$\bar{\varepsilon}([\mathbf{p}]) = \bar{\varepsilon}([\mathbf{m}]) = \varepsilon(\mathbf{m}) \leq \varepsilon(\mathbf{n}) = \bar{\varepsilon}([\mathbf{n}]) = \bar{\varepsilon}([\mathbf{q}]).$$

(4) Let $[p], [q] \in P_{\mathcal{C}(P_C)}$, where $p, q \in P_C$. Then

$$\bar{\varepsilon}([p] \times [q]) = \bar{\varepsilon}([p \times q]) = \varepsilon(p \times q) = \varepsilon(p) \times \varepsilon(q) = \bar{\varepsilon}([p]) \times \bar{\varepsilon}([q]).$$

Dually, $\bar{\varepsilon}([p] \star [q]) = \bar{\varepsilon}([p]) \star \bar{\varepsilon}([q])$.

(5) Let $([p_1], [p_2], \dots, [p_k]) \in \mathcal{P}(P_{\mathcal{C}(P_C)})$. Then

$$\begin{aligned} (\varepsilon_{P_C}((([p_1], [p_2], \dots, [p_k])))\bar{\varepsilon} &= ([p_1, p_2, \dots, p_k])\bar{\varepsilon} \\ &= \varepsilon((p_1, p_2, \dots, p_k)) = \varepsilon(\varepsilon(p_1), \varepsilon(p_2), \dots, \varepsilon(p_k)) = \varepsilon([\varepsilon(p_1)]\bar{\varepsilon}, [\varepsilon(p_2)]\bar{\varepsilon}, \dots, [\varepsilon(p_k)]\bar{\varepsilon}). \end{aligned}$$

By Lemma 6.8, $\bar{\varepsilon}$ is a (2,1,1)-homomorphism from $\mathbf{S}(\mathcal{C}(P))$ to $\mathbf{S}(C)$. Moreover, since ε is an evaluation map, it follows that $([p])\bar{\varepsilon} = \varepsilon(p) = p$ for all $p \in P_C$ by the definition of $\bar{\varepsilon}$. \square

Theorem 7.12. *If (P, \times, \star) be a strong projection algebra, then $\mathbf{S}(\mathcal{C}(P))$ is a projection-generated DRC-restriction semigroup with projection algebra $P(\mathbf{S}(\mathcal{C}(P))) = \{[p] \mid p \in P\}$. Conversely, if $(S, \cdot, +, *)$ is a projection-generated DRC-restriction semigroup whose projection algebra is (P, \times, \star) , then there exists a surjective $(2,1,1)$ -homomorphism ϕ from $\mathbf{S}(\mathcal{C}(P))$ onto S such that $[p]\phi = p$ for all $p \in P$.*

Proof. Since ε_P (see Proposition 7.8) is an evaluation map on $\mathcal{C}(P)$ and is also surjective, by Proposition 7.9, Theorem 6.4 and Proposition 6.5, $\mathbf{S}(\mathcal{C}(P))$ is a projection-generated DRC-restriction semigroup with projection algebra $P(\mathbf{S}(\mathcal{C}(P))) = P_{\mathcal{C}(P)} = \{[p] \mid p \in P\}$.

Conversely, by Proposition 7.11 and Theorem 6.11, $\overline{\varepsilon_S}$ is a $(2,1,1)$ -homomorphism from $\mathbf{S}(\mathcal{C}(P_{\mathbf{C}(S)})) = \mathbf{S}(\mathcal{C}(P))$ to $\mathbf{S}(\mathbf{C}(S)) = S$ satisfying $[p]\overline{\varepsilon_S} = p$ for all $p \in P$. Since S is projection-generated, it follows that ε_S is surjective by Proposition 6.5, and hence $\overline{\varepsilon_S}$ is also surjective. \square

Theorem 7.13. *Let ψ be a homomorphism from the strong projection algebra (P, \times, \star) to the strong projection algebra (P', \times, \star) . Then*

$$\Phi : \mathcal{C}(P) \rightarrow \mathcal{C}(P'), \quad [p_1, p_2, \dots, p_k] \rightarrow [p_1\psi, p_2\psi, \dots, p_k\psi]$$

is a chain projection ordered functor, and so is a $(2,1,1)$ -homomorphism from $\mathbf{S}(\mathcal{C}(P))$ to $\mathbf{S}(\mathcal{C}(P'))$.

Proof. Let

$$\xi : \mathcal{P}(P) \rightarrow \mathcal{P}(P'), \quad (p_1, p_2, \dots, p_k) \mapsto (p_1\psi, p_2\psi, \dots, p_k\psi).$$

Then ξ is well-defined by Lemma 3.6. It is easy to prove that

$$(67) \quad (\mathbf{d}(\mathbf{p}))\xi = \mathbf{d}(\mathbf{p}\xi), (\mathbf{r}(\mathbf{p}))\xi = \mathbf{r}(\mathbf{p}\xi), (\mathbf{p} \circ \mathbf{q})\xi = (\mathbf{p}\xi) \circ (\mathbf{q}\xi)$$

for all $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$. Let $(\mathbf{s}, \mathbf{t}) \in \Omega(P) \cup \Omega^{-1}(P)$. If $\mathbf{s} = (p, p), \mathbf{t} = p, p \in P$, then $\mathbf{s}\xi = (p\psi, p\psi)$ and $\mathbf{t}\xi = p\psi$, and so $(\mathbf{s}\xi, \mathbf{t}\xi), (\mathbf{t}\xi, \mathbf{s}\xi) \in \Omega(P') \cup \Omega^{-1}(P')$. Now let (e, p, f) be an admissible triple of P and $\mathbf{s} = (e, e \star p, f), \mathbf{t} = (e, p \times f, f)$. Then $(e \star p) \star f = f$ and $e = e \times (p \times f)$. Since ψ is a homomorphism, it follows that $(e\psi, p\psi, f\psi)$ is an admissible triple of P' and

$$\mathbf{s}\xi = (e\psi, (e\psi) \star (p\psi), f\psi), \quad \mathbf{t}\xi = (e\psi, (p\psi) \times (e\psi), f\psi).$$

This implies that $(\mathbf{s}\xi, \mathbf{t}\xi), (\mathbf{t}\xi, \mathbf{s}\xi) \in \Omega(P') \cup \Omega^{-1}(P')$.

Let $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{p} = \mathbf{b} \circ \mathbf{s} \circ \mathbf{c}, \mathbf{q} = \mathbf{b} \circ \mathbf{t} \circ \mathbf{c}$, where $\mathbf{b}, \mathbf{c} \in \mathcal{P}(P), (\mathbf{s}, \mathbf{t}) \in \Omega(P) \cup \Omega^{-1}(P)$. Then by the above discussions, $(\mathbf{s}\xi, \mathbf{t}\xi) \in \Omega(P') \cup \Omega^{-1}(P')$. By (67),

$$\mathbf{p}\xi = (\mathbf{b} \circ \mathbf{s} \circ \mathbf{c})\xi = (\mathbf{b}\xi) \circ (\mathbf{s}\xi) \circ (\mathbf{c}\xi), \quad \mathbf{q}\xi = (\mathbf{b} \circ \mathbf{t} \circ \mathbf{c})\xi = (\mathbf{b}\xi) \circ (\mathbf{t}\xi) \circ (\mathbf{c}\xi).$$

This implies that $\mathbf{p}\xi \approx_{P'} \mathbf{q}\xi$, i.e. $[\mathbf{p}\xi] = [\mathbf{q}\xi]$.

Let $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $[\mathbf{p}] = [\mathbf{q}]$. Then $\mathbf{p} \approx_P \mathbf{q}$. By Lemma 7.2 and the statements in previous paragraph, we have $[\mathbf{p}]\Phi = [\mathbf{p}\xi] = [\mathbf{q}\xi] = [\mathbf{q}]\Phi$. Thus Φ is well-defined.

Let $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{p} = (p_1, p_2, \dots, p_k), \mathbf{q} = (q_1, q_2, \dots, q_l)$. Then

$$(\mathbf{d}([\mathbf{p}]))\Phi = [p_1]\Phi = [p_1\psi] = \mathbf{d}([\mathbf{p}]\Phi).$$

Dually, $(\mathbf{r}([\mathbf{p}]))\Phi = \mathbf{r}([\mathbf{p}]\Phi)$. If $\mathbf{r}([\mathbf{p}]) = \mathbf{d}([\mathbf{p}])$, i.e. $p_k = q_1$, by (67) we have

$$([\mathbf{p}] \circ [\mathbf{q}])\Phi = [\mathbf{p} \circ \mathbf{q}]\Phi = [(\mathbf{p} \circ \mathbf{q})\xi] = [(\mathbf{p}\xi) \circ (\mathbf{q}\xi)] = [\mathbf{p}\xi] \circ [\mathbf{q}\xi] = ([\mathbf{p}]\Phi) \circ ([\mathbf{q}]\Phi).$$

This shows that (F1) and (F2) hold.

Assume that $[\mathbf{p}] \leq_{\approx_P} [\mathbf{q}]$. Then there exist $\mathbf{m} \in [\mathbf{p}]$ and $\mathbf{n} \in [\mathbf{q}]$ such that $\mathbf{m} \leq_{\mathcal{P}} \mathbf{n}$, and so there exists $r \in P$ such that $r \leq_P \mathbf{d}(\mathbf{n})$ and $\mathbf{m} = r \downarrow \mathbf{n}$. Denote $\mathbf{m} = (m_1, m_2, \dots, m_u)$ and $\mathbf{n} = (n_1, n_2, \dots, n_v)$. Then by (16),

$$\mathbf{m} = (r, r\theta_{n_2}, \dots, r\theta_{n_2} \dots \theta_{n_v}), \quad \mathbf{n}\xi = (n_1\psi, n_2\psi, \dots, n_v\psi).$$

Since ψ is a homomorphism between projection algebras, we have $r\psi \leq_{P'} (\mathbf{d}(\mathbf{n}))\psi = n_1\psi = \mathbf{d}(\mathbf{n}\xi)$ and

$$\begin{aligned} \mathbf{m}\xi &= (r\psi, (r\theta_{n_2})\psi, \dots, (r\theta_{n_2}\theta_{n_3} \dots \theta_{n_v})\psi) = (r\psi, r\psi\theta_{n_2\psi}, \dots, r\psi\theta_{n_2\psi} \dots \theta_{n_v\psi}) \\ &= {}_{r\psi}\downarrow(n_1\psi, n_1\psi, \dots, n_v\psi) = {}_{r\psi}\downarrow\mathbf{n}\xi. \end{aligned}$$

This shows that $\mathbf{m}\xi \leq_{\mathcal{P}} \mathbf{n}\xi$, and so $[\mathbf{m}]\Phi = [\mathbf{m}\xi] \leq_{\approx_{P'}} [\mathbf{n}\xi] = [\mathbf{n}]\Phi$. As $\mathbf{m} \in [\mathbf{p}]$ and $\mathbf{n} \in [\mathbf{q}]$, we have $[\mathbf{m}] = [\mathbf{p}]$ and $[\mathbf{n}] = [\mathbf{q}]$. Hence $[\mathbf{p}]\Phi = [\mathbf{m}]\Phi \leq_{\approx_{P'}} [\mathbf{n}]\Phi = [\mathbf{q}]\Phi$. This shows that Φ satisfies (F3).

Take $[p], [q] \in P_{\mathcal{C}(P)}$, $p, q \in P$ arbitrarily. Then

$$([p] \times [q])\Phi = ([p \times q])\Phi = [(p \times q)\psi] = [(p\psi) \times (q\psi)] = [p\psi] \times [q\psi].$$

Dually, $([p] \star [q])\Phi = [p\psi] \star [q\psi]$. This gives (F4). Let $([p_1], [p_2], \dots, [p_k]) \in \mathcal{P}(P_{\mathcal{C}(P)})$. Then

$$\begin{aligned} (\varepsilon_P([p_1], [p_2], \dots, [p_k]))\Phi &= [p_1, p_2, \dots, p_k]\Phi = [p_1\psi, p_2\psi, \dots, p_k\psi] \\ &= \varepsilon_{P'}([p_1\psi], [p_2\psi], \dots, [p_k\psi]) = \varepsilon_{P'}([p_1]\Phi, [p_2]\Phi, \dots, [p_k]\Phi). \end{aligned}$$

Thus (F5) holds. This implies that Φ is a chain projection ordered functor. By Lemma 6.8, Φ is a (2,1,1)-homomorphism from $\mathbf{S}(\mathcal{C}(P))$ to $\mathbf{S}(\mathcal{C}(P'))$. \square

In the remainder of this section, we consider the presentations of chain semigroups. We first recall the notion of presentations of semigroups. Let X be a nonempty set and X^+ be the free semigroup on X . Let $R \subseteq X^+ \times X^+$ and R^\sharp be the congruence on X^+ generated by R . Let S be a semigroup. If $S \cong X^+/R^\sharp$, then we say that S has presentation $\langle X : R \rangle$. In this case, we shall identify S with X^+/R^\sharp , and call the elements in X and R the *generators and generations* of S , respectively. Obviously, S has presentation $\langle X : R \rangle$ if and only if there exists a surjective semigroup homomorphism $\pi : X^+ \rightarrow S$ such that the kernel of π is exactly R^\sharp .

Let (P, \times, \star) be a strong projection algebra. The following theorem gives a presentation of the chain semigroup $\mathbf{S}(\mathcal{C}(P))$.

Theorem 7.14. *Let (P, \times, \star) be a strong projection algebra. Then the chain semigroup $\mathbf{S}(\mathcal{C}(P))$ has a presentation $\mathbf{S}(\mathcal{C}(P)) \cong \langle X_P : R_P \rangle$, where $X_P = \{x_p \mid p \in P\}$ is an alphabet with a correspondence to P and R_P is the set of the following relations: For all $p, q \in P$,*

- (R1) $x_p^2 = x_p$.
- (R2) $x_{p \times q} x_q = x_p x_q$.
- (R3) $x_{p \times q} x_{p \star q} = x_p x_q$.
- (R4) $x_p x_{p \star q} = x_p x_q$.

To prove Theorem 7.14, we need a series of lemmas. For convenience, we denote $\sim = R_P^\sharp$, the congruence of X_P^+ generated by R_P .

Lemma 7.15. *Let (P, \times, \star) be a strong projection algebra and k be a positive integer. Let*

$$p_1, p_2, \dots, p_k, q \in P, p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k, q \leq_P p_k.$$

Then there exist $q_1, q_2, \dots, q_{k-1} \in P$ such that

$$q_1 \mathcal{F}_P q_2 \mathcal{F}_P \dots \mathcal{F}_P q_{k-1} \mathcal{F}_P q, x_{p_1} x_{p_2} \dots x_{p_{k-1}} x_q \sim x_{q_1} x_{q_2} \dots x_{q_{k-1}} x_q.$$

Proof. If $k = 1$, then $x_q \sim x_q$ obviously. Let $k \geq 2$. Denote $q_{k-1} = p_{k-1} \times q$. Then $q_{k-1} \leq_P p_{k-1}$ by Lemma 3.1 (1). By mathematical induction, there exist $q_1, q_2, \dots, q_{k-2} \in P$ such that

$$(68) \quad q_1 \mathcal{F}_P q_2 \mathcal{F}_P \dots \mathcal{F}_P q_{k-2} \mathcal{F}_P q_{k-1}, x_{p_1} x_{p_2} \dots x_{p_{k-2}} x_{q_{k-1}} \sim x_{q_1} x_{q_2} \dots x_{q_{k-2}} x_{q_{k-1}}.$$

By (R2),

$$(69) \quad x_{p_{k-1}} x_q \sim x_{p_{k-1} \times q} x_q = x_{q_{k-1}} x_q.$$

By (68) and (69), we have

$$x_{p_1} x_{p_2} \dots x_{p_{k-2}} x_{p_{k-1}} x_q \sim x_{p_1} x_{p_2} \dots x_{p_{k-2}} x_{q_{k-1}} x_q \sim x_{q_1} x_{q_2} \dots x_{q_{k-2}} x_{q_{k-1}} x_q.$$

We shall prove $q_{k-1} \mathcal{F}_P q$ in the sequel. On one hand, by the fact that $q_{k-1} = p_{k-1} \times q$, (L4) and (L1), we obtain

$$q_{k-1} \times q = (p_{k-1} \times q) \times q = (p_{k-1} \times q) \times (p_{k-1} \times q) = p_{k-1} \times q = q_{k-1}.$$

On the other hand, the fact $p_{k-1} \mathcal{F}_P p_k$ gives that $p_k = p_{k-1} \star p_k$, and the fact $q \leq p_k = p_{k-1} \star p_k$ and (8) imply $q = (p_{k-1} \times q) \star p_k$. By (R4), (R1) and the above statements,

$$\begin{aligned} q_{k-1} \star q &= (p_{k-1} \times q) \star ((p_{k-1} \times q) \star p_k) = \\ &= ((p_{k-1} \times q) \star p_k) \star ((p_{k-1} \times q) \star p_k) = ((p_{k-1} \times q) \star p_k) = q. \end{aligned}$$

By (14), $q_{k-1} \mathcal{F}_P q$. □

Lemma 7.16. *Let (P, \times, \star) be a strong projection algebra and $w = x_{q_1} x_{q_2} \dots x_{q_k} \in X_P^+$. Then there exist $p_1, p_2, \dots, p_k \in P$ such that $p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k$ and $w \sim x_{p_1} x_{p_2} \dots x_{p_k}$.*

Proof. When $k = 1$, it is obvious that $x_{q_1} \sim x_{q_1}$. Let $k \geq 2$ and $w = x_{q_1} x_{q_2} \dots x_{q_k}$. By mathematical induction, there exist $r_1, r_2, \dots, r_{k-1} \in P$ such that

$$r_1 \mathcal{F}_P r_2 \mathcal{F}_P \dots \mathcal{F}_P r_{k-1}, x_{q_1} x_{q_2} \dots x_{q_{k-1}} \sim x_{r_1} x_{r_2} \dots x_{r_{k-1}}.$$

Denote $p_{k-1} = r_{k-1} \times q_k$, $p_k = r_{k-1} \star q_k$. By Lemma 3.3, we have $p_{k-1} \mathcal{F}_P p_k$, and so

$$x_{r_{k-1}} x_{q_k} \stackrel{(R3)}{\sim} x_{r_{k-1} \times q_k} x_{r_{k-1} \star q_k} = x_{p_{k-1}} x_{p_k}.$$

Moreover, the fact $p_{k-1} = r_{k-1} \times q_k$ gives that $p_{k-1} \leq_P r_{k-1}$ by Lemma 3.1 (1). By Lemma 7.15, there exist $p_1, p_2, \dots, p_{k-2} \in P$ such that

$$p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_{k-2} \mathcal{F}_P p_{k-1}, x_{r_1} x_{r_2} \dots x_{r_{k-2}} x_{p_{k-1}} \sim x_{p_1} x_{p_2} \dots x_{p_{k-2}} x_{p_{k-1}}.$$

By the above statements,

$$\begin{aligned} p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_{k-2} \mathcal{F}_P p_{k-1} \mathcal{F}_P p_k, \\ w = x_{q_1} x_{q_2} \dots x_{q_{k-1}} x_{q_k} \sim x_{r_1} x_{r_2} \dots x_{r_{k-2}} x_{r_{k-1}} x_{q_k} \\ \sim x_{r_1} x_{r_2} \dots x_{r_{k-2}} x_{p_{k-1}} x_{p_k} \sim x_{p_1} x_{p_2} \dots x_{p_{k-1}} x_{p_k}. \end{aligned}$$

By mathematical induction, the result follows. □

Let (P, \times, \star) be a strong projection algebra and $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$. Define

$$w_{\mathbf{p}} = x_{p_1} x_{p_2} \cdots x_{p_k} \in X_P^+.$$

Let $w \in X_P^+$. By Lemma 7.16, there exists $\mathbf{p} \in \mathcal{P}(P)$ such that $w \sim w_{\mathbf{p}}$. Let $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{q})$. By (R1),

$$(70) \quad w_{\mathbf{p}} w_{\mathbf{q}} \sim w_{\mathbf{p} \circ \mathbf{q}}.$$

Lemma 7.17. *Let (P, \times, \star) be a strong projection algebra, $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{p} \approx_P \mathbf{q}$. Then $w_{\mathbf{p}} \sim w_{\mathbf{q}}$.*

Proof. By Lemma 7.2, we assume that $\mathbf{p}, \mathbf{q} \in \mathcal{P}(P)$ and $\mathbf{p} = \mathbf{b} \circ \mathbf{s} \circ \mathbf{c}, \mathbf{q} = \mathbf{b} \circ \mathbf{t} \circ \mathbf{c}$, where

$$\mathbf{b}, \mathbf{c} \in \mathcal{P}(P), (\mathbf{s}, \mathbf{t}) \in \Omega(P) \cup \Omega^{-1}(P).$$

By (70), we obtain that $w_{\mathbf{p}} \sim w_{\mathbf{b}} w_{\mathbf{s}} w_{\mathbf{c}}$ and $w_{\mathbf{q}} \sim w_{\mathbf{b}} w_{\mathbf{t}} w_{\mathbf{c}}$. We shall prove $w_{\mathbf{s}} \sim w_{\mathbf{t}}$ in the sequel. Let $\mathbf{s} = (p, p), \mathbf{t} = (p, p) \in P$. Then $w_{\mathbf{s}} = x_p x_p \stackrel{(R1)}{\sim} x_p = w_{\mathbf{t}}$. Let (e, p, f) be an admissible triple in P and $\mathbf{s} = (e, e \star p, f), \mathbf{t} = (e, p \times f, f)$. Then

$$w_{\mathbf{s}} = x_e x_{e \star p} x_f \stackrel{(R4)}{\sim} x_e x_p x_f \stackrel{(R2)}{\sim} x_e x_{p \times f} x_f = w_{\mathbf{t}}.$$

This shows that $w_{\mathbf{s}} \sim w_{\mathbf{t}}$, and so $w_{\mathbf{p}} \sim w_{\mathbf{q}}$. \square

Now we can give a proof of Theorem 7.14.

A proof of Theorem 7.14 By the freeness of X_P^+ we can assume that the unique semigroup homomorphism determined by the map $X_P \rightarrow \mathbf{S}(\mathcal{C}(P)), x_p \mapsto [p]$ is

$$\Psi : X_P^+ \rightarrow \mathbf{S}(\mathcal{C}(P)).$$

We first show that Ψ is surjective. Let $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P), \mathbf{c} = [\mathbf{p}] \in \mathbf{S}(\mathcal{C}(P))$. By the proof of Proposition 6.5, we have

$$(71) \quad \begin{aligned} \mathbf{c} &= [p_1, p_2, \dots, p_k] = \varepsilon_P([p_1], [p_2], \dots, [p_k]) \\ &= [p_1] \bullet [p_2] \bullet \dots \bullet [p_k] = (x_{p_1} \Psi) \bullet (x_{p_2} \Psi) \bullet \dots \bullet (x_{p_k} \Psi) = (x_{p_1} x_{p_2} \dots x_{p_k}) \Psi = w_{\mathbf{p}} \Psi. \end{aligned}$$

We next prove that $R_P^\# = \ker \Psi$. Let $(u, v) \in R_P$.

(1) In the case that $u = x_p^2, v = x_p$, we have

$$u \Psi = (x_p x_p) \Psi = (x_p \Psi) \bullet (x_p \Psi) = [p] \bullet [p] = [p] = x_p \Psi = v \Psi.$$

(2) Let $u = x_{p \times q} x_q$ and $v = x_p x_q$ for some $p, q \in P$. By (L4) and (L1),

$$(p \times q) \times q = (p \times q) \times (p \times q) = p \times q,$$

which together with Lemma 3.1 (3) gives that $(p \times q) \star q = p \star q$. By Lemma 6.3 (3),

$$\begin{aligned} u \Psi &= (x_{p \times q} x_q) \Psi = (x_{p \times q} \Psi) \bullet (x_q \Psi) = [p \times q] \bullet [q] = \varepsilon_P([p \times q] \times [q], [p \times q] \star [q]) \\ &= \varepsilon_P([(p \times q) \times q], [(p \times q) \star q]) = \varepsilon_P([p \times q], [p \star q]) = \varepsilon_P([p] \times [q], [p] \star [q]) \\ &= [p] \bullet [q] = x_p \Psi \bullet x_q \Psi = (x_p x_q) \Psi = v \Psi. \end{aligned}$$

(3) Let $u = x_{p \times q} x_{p \star q}$ and $v = x_p x_q$ for some $p, q \in P$. By Lemma 3.1 (4),

$$(p \times q) \times (p \star q) = p \times q, (p \times q) \star (p \star q) = p \star q.$$

Using similar calculations as in (2), we have

$$u \Psi = [p \times q] \bullet [p \star q] = \varepsilon_P([p \times q] \times [p \star q], [p \times q] \star [p \star q]) = \varepsilon_P([p \times q], [p \star q]) = v \Psi.$$

(4) This is the dual of (2).

From items (1)–(4) above, we can obtain $R_P^\sharp \subseteq \ker \Psi$. On the other hand, let $(u, v) \in \ker \Psi$. Then $u, v \in X_P^+$, $u\Psi = v\Psi$. By Lemma 7.16 we have $u \sim w_p$ and $v \sim w_q$ for some $p, q \in \mathcal{P}(P)$. Moreover, (71) and the fact that $R_P^\sharp \subseteq \ker \Psi$ give that

$$[p] = w_p\Psi = u\Psi = v\Psi = w_q\Psi = [q].$$

Thus $p \approx_P q$. By Lemma 7.17, $w_p \sim w_q$. So $u \sim w_p \sim w_q \sim v$. This implies that $\ker(\Psi) \subseteq R_P^\sharp$. Therefore, $R_P^\sharp = \ker \Psi$.

Denote the R_P^\sharp -class containing $w \in X_P^+$ by \bar{w} . Then

$$(72) \quad \bar{\Psi} : X_P^+/R_P^\sharp = \{\bar{w} \mid w \in X_P^+\} = \{\bar{w}_p \mid p \in \mathcal{P}(P)\} \rightarrow \mathbf{S}(\mathcal{C}(P)), \bar{w} \mapsto w\Psi$$

is a semigroup isomorphism. We observe that the multiplication on X_P^+/R_P^\sharp is as follows: For all $w_p, w_q \in X_P^+/R_P^\sharp$, $\bar{w}_p \bar{w}_q = \overline{w_p w_q}$. Moreover, let $p = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$. Define

$$(73) \quad \bar{w}_p^+ = (\bar{w}_p \bar{\Psi})^+ \bar{\Psi}^{-1} = (w_p \Psi)^+ \bar{\Psi}^{-1} = [p_1] \bar{\Psi}^{-1} = \overline{w_{p_1}} = \overline{x_{p_1}}, \quad \bar{w}_p^* = \overline{x_{p_k}}.$$

Then $(X_P^+/R_P^\sharp, \cdot, +, *)$ forms a DRC-restriction semigroup with respect to the above operations with the set of projections $\{\bar{x}_p \mid p \in P\}$, and $\bar{\Psi}$ is a (2,1,1)-isomorphism from X_P^+/R_P^\sharp onto $\mathbf{S}(\mathcal{C}(P))$. Thus the strong projection algebra of $\langle X_P : R_P \rangle$ is isomorphic to that of $\mathbf{S}(\mathcal{C}(P))$, and so is isomorphic to (P, \times, \star) .

Theorem 7.18. *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup and $(P(S), \times_S, \star_S)$ be the strong projection algebra of S . Assume that (P, \times, \star) is a strong projection algebra and $\phi : P \rightarrow P(S)$ is a homomorphism between strong projection algebras. Then there exists a unique (2,1,1)-homomorphism*

$$\Phi : \mathbf{S}(\mathcal{C}(P)) \rightarrow S, \quad [p_1, p_2, \dots, p_k] \mapsto (p_1\phi)(p_2\phi) \dots (p_k\phi).$$

such that the following diagram

$$(74) \quad \begin{array}{ccc} P & \xrightarrow{\phi} & P(S) \\ \iota_1 \downarrow & & \downarrow \iota_2 \\ \mathbf{S}(\mathcal{C}(P)) & \xrightarrow{\Phi} & S \end{array}$$

commutes, where

$$\iota_1 : P \rightarrow \mathbf{S}(\mathcal{C}(P)), \quad p \mapsto [p], \quad \iota_2 : P(S) \rightarrow S, \quad p \mapsto p.$$

Proof. Bt Theorem 7.14, it suffices to show that there exists a unique (2,1,1)-homomorphism $\Gamma : X_P^+/R_P^\sharp \rightarrow S$ such that the diagram

$$\begin{array}{ccc} P & \xrightarrow{\phi} & P(S) \\ \tau_1 \downarrow & & \downarrow \tau_2 \\ X_P^+/R_P^\sharp & \xrightarrow{\Gamma} & S \end{array}$$

commutes, where

$$\tau_1 : P \rightarrow X_P^+/R_P^\sharp, \quad p \mapsto \bar{x}_p, \quad \tau_2 : P(S) \rightarrow S, \quad x \mapsto x.$$

If this is the case, it is easy to prove that

$$(75) \quad \Phi = \overline{\Psi}^{-1} \Gamma : \mathbf{S}(\mathcal{C}(P)) \rightarrow S$$

is the unique (2,1,1)-homomorphism satisfying $\iota_1 \Phi = \phi \iota_2$, where $\overline{\Psi}$ is define in (72).

By the freeness of X_P^+ , we can assume that the unique semigroup homomorphism determined by the map $X_P \rightarrow S, x_p \mapsto p\phi$ is as follows:

$$\varphi : X_P^+ \rightarrow S, \quad x_p \rightarrow p\phi.$$

We shall show that $R_P \subseteq \ker \varphi$ in the sequel. Let $p, q \in P$ and $(u, v) \in R_P$. If $u = x_p^2$ and $v = x_p$, then $u\varphi = x_p^2\varphi = (p\phi)(p\phi) = p\phi = v\varphi$. If $u = x_{p \times q}$ and $v = x_p x_q$, then by using the fact that ϕ is homomorphism, Lemmas 5.5 and 5.3 (3), we have

$$\begin{aligned} u\varphi &= (x_{p \times q} x_q)\varphi = ((p \times q)\phi)(q\phi) = (p\phi \times_S q\phi)(q\phi) \\ &= ((p\phi)(q\phi))^+(q\phi) = (p\phi)(q\phi) = (x_p\varphi)(x_q\varphi) = (x_p x_q)\varphi = v\varphi. \end{aligned}$$

Dually, we also have $u\varphi = v\varphi$ if $u = x_p x_{p \star q}$ and $v = x_p x_q$. If $u = x_{p \times q} x_{p \star q}$, $v = x_p x_q$, then by using the fact that ϕ is homomorphism, Lemmas 5.5 and 5.3 (4), we obtain

$$\begin{aligned} u\varphi &= (x_{p \times q} x_{p \star q})\varphi = (p \times q)\phi(p \star q)\phi = (p\phi \times_S q\phi)(p\phi \star_S q\phi)\phi \\ &= ((p\phi)(q\phi))^+((p\phi)(q\phi))^* = (p\phi)(q\phi) = (x_p x_q)\varphi = v\varphi. \end{aligned}$$

This implies that $R_P \subseteq \ker \varphi$, and so $R_P^\# \subseteq \ker \varphi$. Thus we can define the semigroup homomorphism

$$\Gamma : X_P^+ / R_P^\# = \langle X_P : R_P \rangle \rightarrow S, \quad \overline{w} \rightarrow w\varphi.$$

By the statements after Lemma 7.16, we have $X_P^+ / R_P^\# = \{\overline{w_p} \mid \mathbf{p} \in \mathcal{P}(P)\}$. Let

$$\overline{w_p} \in X_P^+ / R_P^\#, \quad \mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P).$$

By (73), $\overline{w_p}^+ = \overline{x_{p_1}}$ and $\overline{w_p}^* = \overline{x_{p_k}}$. This shows that $\overline{w_p}^+ \Gamma = \overline{x_{p_1}} \Phi = x_{p_1} \varphi = p_1 \phi$. Observe $p_1 \mathcal{F}_P p_2 \mathcal{F}_P \dots \mathcal{F}_P p_k$ as $\mathbf{p} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P)$. Hence

$$p_1 \phi \mathcal{F}_{P(S)} p_2 \phi \mathcal{F}_{P(S)} \dots \mathcal{F}_{P(S)} p_k \phi.$$

In view of Lemma 3.5, in $(P(S), \times_S, \star_S)$ we have

$$\begin{aligned} p_1 \phi &= p_k \phi \delta_{p_{k-1} \phi} \delta_{p_{k-2} \phi} \dots \delta_{p_1 \phi} \\ &= (p_1 \phi \times_S (\dots (p_{k-2} \phi \times_S (p_{k-1} \phi \times_S p_k \phi)))) \\ &= (p_1 \phi (\dots (p_{k-2} \phi (p_{k-1} \phi p_k \phi)^+)^+)^+)^+ \\ &= ((p_1 \phi)(p_2 \phi) \dots (p_k \phi))^+ \quad (\text{by (ii) in DRC Conditions}) \\ &= ((x_{p_1} x_{p_2} \dots x_{p_k})\varphi)^+ = (w_p \varphi)^+ = (\overline{w_p} \Gamma)^+. \end{aligned}$$

Thus $(\overline{w_p} \Gamma)^+ = \overline{w_p}^+ \Gamma$. Dually, we have $(\overline{w_p} \Gamma)^* = \overline{w_p}^* \Gamma$. Let $p \in P$. Then

$$p(\tau_1 \Gamma) = \overline{x_p} \Gamma = x_p \varphi = p\phi = (p\phi)\tau_2 = p(\phi\tau_2).$$

Finally, since

$$\overline{w_p} = \overline{x_{p_1} x_{p_2} \dots x_{p_k}} = \overline{x_{p_1}} \overline{x_{p_2}} \dots \overline{x_{p_k}},$$

it follows that $X_P^+ / R_P^\#$ is generated by $\{\overline{x_p} \mid p \in P\}$. So Γ is necessarily unique. Moreover, combining (71), (72) and (75), we have

$$\Phi : \mathbf{S}(\mathcal{C}(P)) \rightarrow S, \quad [p_1, p_2, \dots, p_k] \mapsto (p_1 \phi)(p_2 \phi) \dots (p_k \phi),$$

which is the unique (2,1,1)-homomorphism such that the diagram (74) commutes. \square

Denote the category of DRC-restriction semigroups together with $(2,1,1)$ -homomorphisms by \mathbb{DRS} and the category of strong projection algebras together with projection algebra homomorphisms by \mathbb{SPA} , respectively. Define a functor $\mathcal{U} : \mathbb{DRS} \rightarrow \mathbb{SPA}$ as follows: For any two DRC-semigroups $(S, \cdot, +, *)$, $(T, \cdot, +, *)$ and a $(2,1,1)$ -homomorphism $\phi : S \rightarrow T$,

- (1) $\mathcal{U}(S) = (P(S), \times_S, \star_S)$.
- (2) $\mathcal{U}(\phi) : \mathcal{U}(S) = P(S) \rightarrow \mathcal{U}(T) = P(T), x \mapsto x\phi$.

Moreover, define a functor $\mathcal{V} : \mathbb{SPA} \rightarrow \mathbb{DRS}$ as follows: For any two strong projection algebras (P, \times, \star) and (Q, \times, \star) and a projection algebra homomorphism $\psi : P \rightarrow Q$,

- (1) $\mathcal{V}(P) = \mathbf{S}(\mathcal{C}(P))$.
- (2) $\mathcal{V}(\psi) : \mathcal{V}(P) = \mathbf{S}(\mathcal{C}(P)) \rightarrow \mathcal{V}(Q) = \mathbf{S}(\mathcal{C}(Q))$ is the $(2,1,1)$ homomorphism induced in Theorem 7.13.

To give further properties of the functors \mathcal{U} and \mathcal{V} above, we need some notions.

Definition 7.19. Let \mathbf{C} and \mathbf{D} be two (large) category and F and G be two functors from \mathbf{C} to \mathbf{D} . Assume that for each object C in \mathbf{C} , η_C is a morphism from $F(C)$ to $G(C)$ in \mathbf{D} . The family of morphisms $\eta = \{\eta_C \mid C \text{ is an object in } \mathbf{C}\}$ is called a *natural transformation* from F to G if for any two objects C, C' in \mathbf{C} and any morphism $\phi : C \rightarrow C'$ in \mathbf{C} , we have $\eta_{C'}G(\phi) = F(\phi)\eta_C$, that is, the following diagram

$$\begin{array}{ccc} F(C) & \xrightarrow{\eta_C} & G(C) \\ F(\phi) \downarrow & & G(\phi) \downarrow \\ F(C') & \xrightarrow{\eta_{C'}} & G(C') \end{array}$$

commutes.

Definition 7.20. Let \mathbf{C} and \mathbf{D} be two (large) category, $\mathcal{V} : \mathbf{C} \rightarrow \mathbf{D}$ and $\mathcal{U} : \mathbf{D} \rightarrow \mathbf{C}$ be two functors, and $\eta : \text{id}_{\mathbf{C}} \rightarrow \mathcal{U}\mathcal{V}$ be a natural transformation. Then $(\mathcal{V}, \mathcal{U}, \eta)$ is called an *adjunction* if for any object C in \mathbf{C} , any object D in \mathbf{D} and any morphism $\phi : C \rightarrow \mathcal{U}(D)$, there exists a unique morphism $\bar{\phi} : \mathcal{V}(C) \rightarrow D$ such that $\eta_C \mathcal{U}(\bar{\phi}) = \phi$, that is, the following diagram

$$\begin{array}{ccc} C & \xrightarrow{\phi} & \mathcal{U}(D) \\ \eta_C \downarrow & \nearrow \mathcal{U}(\bar{\phi}) & \\ \mathcal{U}(\mathcal{V}(C)) & & \end{array}$$

commutes. In this case, \mathcal{V} and \mathcal{U} are called *left and right adjoints*, respectively, and η is called the *unit* of the adjunction $(\mathcal{V}, \mathcal{U}, \eta)$. Moreover, elements in $\{\mathcal{V}(C) \mid C \text{ is an object in } \mathbf{C}\}$ are called the *\mathbf{C} -free objects in \mathbf{D}* .

Theorem 7.21. *Define*

$$\eta = \{\eta_P \mid P \text{ is a strong projection algebra}\} : \text{id}_{\mathbb{SPA}} \rightarrow \mathcal{U}\mathcal{V}$$

as follows: For any strong projection algebra P ,

$$\eta_P : P = \text{id}_{\mathbb{SPA}}(P) \rightarrow \mathcal{U}\mathcal{V}(P) = \{[p] \mid p \in P\}, p \mapsto [p].$$

Then $(\mathcal{V}, \mathcal{U}, \eta)$ is an adjunction, and so the \mathbb{SPA} -objects in \mathbb{DRS} are exactly chain semi-groups.

Proof. We first show that η is a natural transformation. Let P and Q be two strong projection algebras and $\phi : P \rightarrow Q$ be a homomorphism. Let $p \in P$. Then

$$p(\eta_P(\mathcal{UV}(\phi))) = [p](\mathcal{UV}(\phi)) = [p\phi] = (p\phi)\eta_Q = p(\phi\eta_Q) = p(\text{id}_{\mathbb{S}\mathbb{P}\mathbb{A}}(\phi)\eta_Q).$$

This shows that the following diagram

$$\begin{array}{ccc} P = \text{id}_{\mathbb{S}\mathbb{P}\mathbb{A}}(P) & \xrightarrow{\eta_P} & \mathcal{UV}(P) = P(\mathbf{S}(\mathcal{C}(P))) \\ \text{id}_{\mathbb{S}\mathbb{P}\mathbb{A}}(\phi) \downarrow & & \mathcal{UV}(\phi) \downarrow \\ Q = \text{id}_{\mathbb{S}\mathbb{P}\mathbb{A}}(Q) & \xrightarrow{\eta_Q} & \mathcal{UV}(Q) = P(\mathbf{S}(\mathcal{C}(Q))) \end{array}$$

commutes. Thus η is a natural transformation. Let (P, \times, \star) be a strong projection algebra, $(S, \cdot, +, *)$ be a DRC-restriction semigroup whose strong projection algebra is $(P(S), \times_S, \star_S)$ and $\phi : P \rightarrow \mathcal{U}(S) = P(S)$ be a projection algebra homomorphism. By Theorem 7.18, there exists a unique $(2,1,1)$ -homomorphism

$$\Phi : \mathcal{V}(P) = \mathbf{S}(\mathcal{C}(P)) \rightarrow S, [p_1, p_2, \dots, p_k] \mapsto (p_1\phi)(p_2\phi) \dots (p_k\phi).$$

such that $[p]\Phi = p\phi$. Thus

$$\mathcal{U}(\Phi) : P(\mathbf{S}(\mathcal{C}(P))) = \mathcal{UV}(P) \rightarrow \mathcal{U}(S) = P(S), [p] \mapsto p\phi.$$

Let $p \in P$. Then $p(\eta_P \mathcal{U}(\Phi)) = (p\eta_P) \mathcal{U}(\Phi) = [p] \mathcal{U}(\Phi) = p\phi$, that is, the following diagram

$$\begin{array}{ccc} P & \xrightarrow{\phi} & \mathcal{U}(S) = P(S) \\ \eta_P \downarrow & \nearrow \mathcal{U}(\Phi) & \\ P(\mathbf{S}(\mathcal{C}(P))) = \mathcal{UV}(P) & & \end{array}$$

commutes. Assume that $\sigma : \mathcal{V}(P) = \mathbf{S}(\mathcal{C}(P)) \rightarrow S$ is a $(2,1,1)$ -homomorphism such that the following diagram

$$\begin{array}{ccc} P & \xrightarrow{\phi} & \mathcal{U}(S) = P(S) \\ \eta_P \downarrow & \nearrow \mathcal{U}(\sigma) & \\ P(\mathbf{S}(\mathcal{C}(P))) = \mathcal{UV}(P) & & \end{array}$$

commutes. Then $[p]\sigma = [p]\mathcal{U}(\sigma) = p(\eta_P \mathcal{U}(\sigma)) = p\phi$ for any $p \in P$, and so the following diagram

$$\begin{array}{ccc} P & \xrightarrow{\phi} & P(S) \\ \iota_1 \downarrow & & \downarrow \iota_2 \\ \mathbf{S}(\mathcal{C}(P)) & \xrightarrow{\sigma} & S \end{array}$$

commutes. By Theorem 7.18, we have $\sigma = \Phi$. This gives the uniqueness of Φ . Thus $(\mathcal{V}, \mathcal{U}, \eta)$ is an adjunction. \square

8. PROJECTION-FUNDAMENTAL DRC-RESTRICTION SEMIGROUPS

In this section, by using Theorem 6.11 we reobtain the constructions of projection-fundamental DRC-restriction semigroups which originally given by Wang [59]. We first give some preliminary notions and results.

Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup and σ be an equivalence on S . Then σ is called a $(2,1,1)$ -congruence on S if for all $a, b, c \in S$,

$$a \sigma b \implies ac \sigma bc, ca \sigma cb, a^+ \sigma b^+, a^* \sigma b^*.$$

A $(2,1,1)$ -congruence σ on S is called *projection-separating* if for all $p, q \in P(S)$, the fact that $p \sigma q$ implies that $p = q$.

Lemma 8.1 ([26]). *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup. Then*

$$(76) \quad \mu_S = \{(a, b) \in S \times S \mid a^+ = b^+, a^* = b^* \text{ and } (ap)^+ = (bp)^+, (pa)^* = (pb)^* \text{ for all } p \in P(S)\}$$

is the maximum projection-separating $(2,1,1)$ -congruence on S .

Definition 8.2. A DRC-restriction semigroup $(S, \cdot, +, *)$ is called *projection-fundamental* if the unique $(2,1,1)$ -projection-separating congruence on S is the identity relation, that is, μ_S is the identity relation on S .

Proposition 8.3. *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup and $a, b \in S$. Assume that $\mathbf{d}_S, \mathbf{r}_S, \nu_a, \mu_a, \nu_b, \mu_b$ and $\Theta_a, \Delta_a, \Theta_b, \Delta_b$ are given in (37), (39), (40), (42) and (43), respectively. Then the following statements are equivalent:*

- (1) $(a, b) \in \mu_S$.
- (2) $\Theta_a = \Theta_b$ and $\Delta_a = \Delta_b$.
- (3) $\mathbf{d}_S(a) = \mathbf{d}_S(b)$ and $\Theta_a = \Theta_b$.
- (4) $\mathbf{r}_S(a) = \mathbf{r}_S(b)$ and $\Delta_a = \Delta_b$.
- (5) $\mathbf{d}_S(a) = \mathbf{d}_S(b)$ and $\Theta_a|_{\mathbf{d}_S(a)^\downarrow} = \Theta_b|_{\mathbf{d}_S(b)^\downarrow}$.
- (6) $\mathbf{r}_S(a) = \mathbf{r}_S(b)$ and $\Delta_a|_{\mathbf{r}_S(a)^\downarrow} = \Delta_b|_{\mathbf{r}_S(b)^\downarrow}$.
- (7) $\nu_a = \nu_b$.
- (8) $\mu_a = \mu_b$.

Proof. We first observe that $\mathbf{d}_S(a) = a^+$, $\mathbf{r}_S(a) = a^*$, and

$$p\Theta_a = p\theta_{\mathbf{d}_S(a)}\nu_a = (pa)^* \text{ and } p\Delta_a = p\delta_{\mathbf{r}_S(a)}\mu_a = (ap)^+ \text{ for all } p \in P(S).$$

By symmetry, we only need to show that (1), (2), (3), (5) and (7) are mutually equivalent. Lemma 8.1 gives that (1) implies that (2). That (3) implies (5) is trivial. The fact (23) provides that (5) implies (7). In the followings, we prove that (2) implies (3) and (7) implies (1). Assume that (2) holds. Then $\Theta_a = \Theta_b$ and $\Delta_a = \Delta_b$. This implies that

$$\mathbf{d}_S(a) = a^+ = (aa^*)^+ = a^*\Delta_a = a^*\Delta_b = (ba^*)^+ \leq_S b^+ = \mathbf{d}_S(b)$$

by (i)', (32) and Lemma 5.6. Dually, we have $\mathbf{d}_S(b) \leq_S \mathbf{d}_S(a)$. This implies that $\mathbf{d}_S(b) = \mathbf{d}_S(a)$. This together with the known fact that $\Theta_a = \Theta_b$ gives (3).

Now assume that (7) is true. Then $\nu_a = \nu_b$. This yields that $\mathbf{d}_S(a)^\downarrow = \text{dom}\nu_a = \text{dom}\nu_b = \mathbf{d}_S(b)^\downarrow$, and so $\mathbf{d}_S(a) = \mathbf{d}_S(b)$. Moreover, $\theta_{\mathbf{d}_S(a)} = \theta_{\mathbf{d}_S(b)}$. Thus $\Theta_a = \theta_{\mathbf{d}_S(a)}\nu_a = \theta_{\mathbf{d}_S(b)}\nu_b = \Theta_b$. This gives (3). On the other hand, by Lemma 2.3, we have $\mu_a = \nu_a^{-1} = \nu_b^{-1} = \mu_b$.

By the dual statements, we can obtain (4). Now Lemma 8.1 and (3), (4) together imply (1). \square

Let (P, \times, \star) be a strong projection algebra and $p \in P$. Then

$$p^\downarrow = \{x \in P \mid x \leq_P p\}$$

is a subalgebra of P . Let $p, q \in P$. Denote

$$M(p, q) = \{\alpha \mid \alpha \text{ is a projection algebra isomorphism from } p^\downarrow \text{ to } q^\downarrow\},$$

$$M(P) = \bigcup_{p, q \in P} M(p, q).$$

For all $\alpha, \beta \in M(P)$, define

$$\alpha \circ \beta = \begin{cases} \alpha\beta & \text{if } \text{ran}\alpha = \text{dom}\beta, \\ \text{undefined} & \text{otherwise,} \end{cases}$$

and $\mathbf{d}(\alpha) = \text{id}_{\text{dom}\alpha}$, $\mathbf{r}(\alpha) = \text{id}_{\text{ran}\alpha}$, where $\alpha\beta$ denotes the composition of α and β . Denote the inverse map of α by α^{-1} for all $\alpha \in M(P)$. Then it is easy to prove that $(M(P), \circ, \mathbf{d}, \mathbf{r}, ^{-1})$ is a groupoid with the set of objects $P_{M(P)} = \{\text{id}_{p^\downarrow} \mid p \in P\}$. We also observe the following fact:

$$(77) \quad p\alpha = q \text{ and } s\alpha \leq_P t\alpha \text{ for all } \alpha \in M(p, q) \text{ and } s, t \in p^\downarrow \text{ with } s \leq_P t,$$

Let $\alpha \in M(s, t)$ and X be a subalgebra of $\text{dom}\alpha = s^\downarrow$. Then $\alpha|_X : X \rightarrow X\alpha$, $x \mapsto x\alpha$ is a projection algebra isomorphism from X to $X\alpha$, and is called *the restriction of α on X* .

Lemma 8.4. *Let $\alpha \in M(s, t)$ and $p \in \text{dom}\alpha$. Then $\alpha|_{p^\downarrow} \in M(p, p\alpha)$.*

Proof. If $p \in \text{dom}\alpha$, then $p^\downarrow \subseteq \text{dom}\alpha = s^\downarrow$, and so $\alpha|_{p^\downarrow} : p^\downarrow \rightarrow p^\downarrow\alpha$, $x \mapsto x\alpha$. Let $x\alpha \in p^\downarrow\alpha$, where $x \leq_P p$. By (77), we have $x\alpha \leq_P p\alpha$ and so $x\alpha \in (p\alpha)^\downarrow$. This shows that $p^\downarrow\alpha \subseteq (p\alpha)^\downarrow$. Now let $y \in (p\alpha)^\downarrow$. Then $y \leq_P p\alpha \leq_P t$, and so $y \in t^\downarrow$. Denote $y = x\alpha$. Then $y = x\alpha \leq_P p\alpha$. Thus $x \leq_P p$ as α^{-1} is also order-preserving. This implies that $(p\alpha)^\downarrow \subseteq p^\downarrow\alpha$. Therefore $(p\alpha)^\downarrow = p^\downarrow\alpha$. We have shown that $\alpha|_{p^\downarrow} \in M(p, p\alpha)$. \square

Define a relation $\leq_{M(P)}$ on $M(P)$ as follows: For all $\alpha, \beta \in M(P)$,

$$\alpha \leq_{M(P)} \beta \iff \alpha = \beta|_{\text{dom}\alpha}.$$

That is, $\alpha \leq_{M(P)} \beta$ if and only if $\text{dom}\alpha \subseteq \text{dom}\beta$, and $x\alpha = x\beta$ for all $x \in \text{dom}\alpha$. It is easy to see that “ $\leq_{M(P)}$ ” is a partial order on $M(P)$.

Lemma 8.5. *Let (P, \times, \star) be a strong projection algebra. Then $(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{M(P)}, P_{M(P)}, \times, \star)$ is a projection ordered category and*

$$\text{id}_{p^\downarrow} \downarrow \alpha = \alpha|_{p^\downarrow}, \quad \alpha \downarrow \text{id}_{q^\downarrow} = \alpha|_{(q\alpha^{-1})^\downarrow}$$

for all $p \in P$ with $\text{id}_{p^\downarrow} \leq_{M(P)} \mathbf{d}(\alpha)$ and $q \in P$ with $\text{id}_{q^\downarrow} \leq_{M(P)} \mathbf{r}(\alpha)$, respectively.

Proof. Define \times and \star on $P_{M(P)}$ as follows: For all $p, q \in P$,

$$\text{id}_{p^\downarrow} \times \text{id}_{q^\downarrow} = \text{id}_{(p \times q)^\downarrow}, \quad \text{id}_{p^\downarrow} \star \text{id}_{q^\downarrow} = \text{id}_{(p \star q)^\downarrow}.$$

Then $(P_{M(P)}, \times, \star)$ forms a strong projection algebra isomorphic to (P, \times, \star) via the map $\text{id}_{p^\downarrow} \mapsto p$. Thus for all $p, q \in P$, $p \leq_P q$ if and only if $\text{id}_{p^\downarrow} \leq_{P_{M(P)}} \text{id}_{q^\downarrow}$.

Now let $p, q \in P$. Then

$$\text{id}_{p^\downarrow} \leq_{M(P)} \text{id}_{q^\downarrow} \iff p^\downarrow \subseteq q^\downarrow \iff p \leq_P q \iff \text{id}_{p^\downarrow} \leq_{P_{M(P)}} \text{id}_{q^\downarrow}.$$

This implies that $\leq_{M(P)}$ and $\leq_{P_{M(P)}}$ is the same on $P_{M(P)}$.

(O1) Let $\alpha \leq_{M(P)} \beta$. Then $\alpha = \beta|_{\text{dom}\alpha}$, which implies that $\text{dom}\alpha \subseteq \text{dom}\beta$ and $x\alpha = x\beta$ for all $x \in \text{dom}\alpha$. Hence $\mathbf{d}(\alpha) = \text{id}_{\text{dom}\alpha} \leq_{M(P)} \text{id}_{\text{dom}\beta} = \mathbf{d}(\beta)$. Dually, we have $\mathbf{r}(\alpha) \leq_{M(P)} \mathbf{r}(\beta)$.

(O2) Let $\alpha \leq_{M(P)} \beta, \gamma \leq_{M(P)} \delta$ and $\alpha \circ \gamma$ and $\beta \circ \delta$ be defined. Then

$$\begin{aligned} \alpha &= \beta|_{\text{dom}\alpha}, \gamma = \delta|_{\text{dom}\gamma}, \text{dom}\alpha \subseteq \text{dom}\beta, \text{dom}\gamma \subseteq \text{dom}\delta, \\ \text{ran}\alpha &= \text{dom}\gamma, \text{ran}\beta = \text{dom}\delta, \alpha \circ \gamma = \alpha\gamma, \beta \circ \delta = \beta\delta. \end{aligned}$$

Thus

$$\text{dom}(\alpha \circ \gamma) = \text{dom}(\alpha\gamma) = \text{dom}(\alpha) \subseteq \text{dom}(\beta) = \text{dom}(\beta \circ \delta).$$

Let $x \in \text{dom}(\alpha \circ \gamma) = \text{dom}\alpha$. Then

$$x(\alpha \circ \gamma) = x(\alpha\gamma) = (x\alpha)\gamma = (x\beta)\gamma = (x\beta)\delta = x\beta\delta = x(\beta \circ \delta).$$

This implies that $\alpha \circ \gamma \leq_{M(P)} \beta \circ \delta$.

(O3) Let $\text{id}_{p^\downarrow} \in P_{M(P)}, p \in P$ and $\text{id}_{p^\downarrow} \leq_{M(P)} \mathbf{d}(\alpha) = \text{id}_{\text{dom}\alpha}$. Then $p^\downarrow \subseteq \text{dom}\alpha$. By Lemma 8.4, we have $\alpha|_{p^\downarrow} \in M(p, p\alpha) \subseteq M(P)$. Obviously, $\alpha|_{p^\downarrow} \leq_{M(P)} \alpha$. Moreover, $\mathbf{d}(\alpha|_{p^\downarrow}) = \text{id}_{\text{dom}(\alpha|_{p^\downarrow})} = \text{id}_{p^\downarrow}$. Assume that $\beta \in M(P), \beta \leq_{M(P)} \alpha$ and $\mathbf{d}(\beta) = \text{id}_{p^\downarrow}$. Then $\text{id}_{\text{dom}\beta} = \mathbf{d}(\beta) = \text{id}_{p^\downarrow}$ and $\beta = \alpha|_{\text{dom}\beta}$, and so $\text{dom}\beta = p^\downarrow$ and $\beta = \alpha|_{\text{dom}\beta} = \alpha|_{p^\downarrow}$. Thus $\text{id}_{p^\downarrow} \downarrow \alpha = \alpha|_{p^\downarrow}$.

(O4) Let $\text{id}_{q^\downarrow} \in P_{M(P)}, q \in P$ and $\text{id}_{q^\downarrow} \leq_{M(P)} \mathbf{r}(\alpha) = \text{id}_{\text{ran}\alpha}$. Then

$$q^\downarrow \subseteq \text{ran}\alpha = \text{dom}\alpha^{-1}, q\alpha^{-1} \in \text{ran}\alpha^{-1} = \text{dom}\alpha, (q\alpha^{-1})^\downarrow \subseteq \text{dom}\alpha.$$

By Lemma 8.4, we have $\alpha|_{(q\alpha^{-1})^\downarrow} \in M(q\alpha^{-1}, q)$. Obviously, $\alpha|_{(q\alpha^{-1})^\downarrow} \leq_{M(P)} \alpha$. Moreover, $\mathbf{r}(\alpha|_{(q\alpha^{-1})^\downarrow}) = \text{id}_{\text{ran}(\alpha|_{(q\alpha^{-1})^\downarrow})} = \text{id}_{q^\downarrow}$. Assume that $\beta \in M(P), \beta \leq_{M(P)} \alpha$ and $\mathbf{r}(\beta) = \text{id}_{q^\downarrow}$. Then $\text{id}_{\text{ran}\beta} = \mathbf{r}(\beta) = \text{id}_{q^\downarrow}$ and $\beta = \alpha|_{\text{dom}\beta}$, and so $\text{ran}\beta = q^\downarrow$ and

$$\beta = \alpha|_{\text{dom}\beta} = \alpha|_{(q^\downarrow)\beta^{-1}} = \alpha|_{(q\beta^{-1})^\downarrow} = \alpha|_{(q\alpha^{-1})^\downarrow}.$$

Thus $\alpha|_{\text{id}_{q^\downarrow}} = \alpha|_{(q\alpha^{-1})^\downarrow}$. We have shown that $(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{M(P)}, P_{M(P)}, \times, \star)$ is a weak projection ordered category.

Now, let $p, q \in P$ and $\alpha \in M(p, q)$. Then

$$\begin{aligned} \mathbf{d}(\alpha)^\downarrow &= (\text{id}_{\text{dom}\alpha})^\downarrow = (\text{id}_{p^\downarrow})^\downarrow = \{\text{id}_{x^\downarrow} \mid x \in P, x \leq_P p\} = \{\text{id}_{x^\downarrow} \mid x \in P, x \in p^\downarrow\}, \\ \mathbf{r}(\alpha)^\downarrow &= (\text{id}_{\text{ran}\alpha})^\downarrow = (\text{id}_{q^\downarrow})^\downarrow = \{\text{id}_{y^\downarrow} \mid y \in P, y \leq_P q\} \\ &= \{\text{id}_{y^\downarrow} \mid y \in P, y \in q^\downarrow\} = \{\text{id}_{(x\alpha)^\downarrow} \mid x \in P, x \in p^\downarrow\}. \end{aligned}$$

Let $\text{id}_{x^\downarrow} \leq_{M(P)} \mathbf{d}(\alpha)$. By (6),

$$\text{id}_{x^\downarrow} \nu_\alpha = \mathbf{r}(\text{id}_{x^\downarrow} \downarrow \alpha) = \mathbf{r}(\alpha|_{x^\downarrow}) = \text{id}_{\text{ran}(\alpha|_{x^\downarrow})} = \text{id}_{(x\alpha)^\downarrow}.$$

Thus we have

$$(78) \quad \nu_\alpha : \mathbf{d}(\alpha)^\downarrow \rightarrow \mathbf{r}(\alpha)^\downarrow, \text{id}_{x^\downarrow} \mapsto \text{id}_{(x\alpha)^\downarrow}.$$

Lemma 2.3 gives that $\mu_\alpha = \nu_{\alpha^{-1}}$. Hence

$$(79) \quad \mu_\alpha : \mathbf{r}(\alpha)^\downarrow \rightarrow \mathbf{d}(\alpha)^\downarrow, \text{id}_{y^\downarrow} \mapsto \text{id}_{(y\alpha^{-1})^\downarrow}.$$

Since α is isomorphism, ν_α is an isomorphism. Thus $(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{M(P)}, P_{M(P)}, \times, \star)$ is a projection ordered category. \square

Let (P, \times, \star) be a strong projection algebra, $p, q \in P$ and $\alpha \in M(p, q)$. Then $\mathbf{d}(\alpha) = \text{id}_{p^\downarrow}$ and $\mathbf{r}(\alpha) = \text{id}_{q^\downarrow}$. Let $\text{id}_{x^\downarrow} \in P_{M(P)}$. By (20) and (78),

$$(\text{id}_{x^\downarrow})\Theta_\alpha = (\text{id}_{x^\downarrow})\theta_{\mathbf{d}(\alpha)}\nu_\alpha = (\text{id}_{x^\downarrow} \star \text{id}_{p^\downarrow})\nu_\alpha = (\text{id}_{(x \star p)^\downarrow})\nu_\alpha = \text{id}_{((x \star p)\alpha)^\downarrow}.$$

Thus we have

$$(80) \quad \Theta_\alpha : P_{M(P)} \rightarrow \mathbf{r}(\alpha)^\downarrow, \quad \text{id}_{x^\downarrow} \mapsto \text{id}_{((x \star p)\alpha)^\downarrow}.$$

Dually, we obtain

$$(81) \quad \Delta_\alpha : P_{M(P)} \rightarrow \mathbf{d}(\alpha)^\downarrow, \quad \text{id}_{x^\downarrow} \mapsto \text{id}_{((q \times x)\alpha^{-1})^\downarrow}.$$

In particular, for all $\text{id}_{x^\downarrow} \in P_{M(P)}$,

$$(82) \quad \Theta_{\text{id}_{x^\downarrow}} = \theta_{\text{id}_{x^\downarrow}}, \quad \Delta_{\text{id}_{x^\downarrow}} = \delta_{\text{id}_{x^\downarrow}}$$

Let (P, \times, \star) be a strong projection algebra, and $p, q \in P$ with $p\mathcal{F}_P q$. Then

$$\theta_q : P \rightarrow P, \quad x \mapsto x \star q, \quad \delta_p : P \rightarrow P, \quad y \mapsto p \times y.$$

By (12), we have $\text{ran}\theta_q = q^\downarrow$ and $\text{ran}\delta_p = p^\downarrow$. Denote

$$(83) \quad \gamma_{p,q} = \theta_q|_{p^\downarrow}, \quad \beta_{q,p} = \delta_p|_{q^\downarrow}$$

Then $\gamma_{p,q}$ and $\beta_{q,p}$ are maps from p^\downarrow to q^\downarrow and q^\downarrow to p^\downarrow , respectively. According to [59, (4.8)], [59, Lemma 4.5] and the fact that $p\mathcal{F}_P q$, we have the following lemma.

Lemma 8.6. *Let (P, \times, \star) be a strong projection algebra and $p, q \in P$ with $p\mathcal{F}_P q$. Then $\gamma_{p,q} \in M(p, q)$ and $\gamma_{p,q}^{-1} = \beta_{q,p}$.*

Let $f : A \rightarrow B$ and $g : B \rightarrow C$ be two bijections and $X \subseteq A$. Then

$$(84) \quad (fg)|_X = (f|_X)(g|_{Xf}).$$

Lemma 8.7. *Let (P, \times, \star) be a strong projection algebra and $(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{M(P)}, P_{M(P)}, \times, \star)$ be the projection ordered category determined by P and $\mathcal{P}(P_{M(P)})$ be the path category of $P_{M(P)}$. Define*

$$\varepsilon : \mathcal{P}(P_{M(P)}) \rightarrow M(P), \quad \text{id}_{p_1^\downarrow} \mapsto \text{id}_{p_1^\downarrow}, \quad (\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}, \dots, \text{id}_{p_k^\downarrow}) \mapsto \gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-1}, p_k}.$$

Then ε is an evaluation map.

Proof. Let

$$\mathbf{p} = (\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}, \dots, \text{id}_{p_k^\downarrow}), \quad \mathbf{s} = (\text{id}_{s_1^\downarrow}, \text{id}_{s_2^\downarrow}, \dots, \text{id}_{s_l^\downarrow}) \in \mathcal{P}(P_{M(P)})$$

and $p \in P$. Then $\varepsilon(\text{id}_{p^\downarrow}) = \text{id}_{p^\downarrow}$. By (83), we have $\varepsilon((\text{id}_{p^\downarrow}, \text{id}_{p^\downarrow})) = \gamma_{p,p} = \theta_p|_{p^\downarrow} = \text{id}_{p^\downarrow}$. This gives (E1). By (83) and (E1),

$$\begin{aligned} \mathbf{d}(\varepsilon(\mathbf{p})) &= \mathbf{d}(\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-1}, p_k}) \\ &= \mathbf{d}(\theta_{p_2}|_{p_1^\downarrow} \theta_{p_3}|_{p_2^\downarrow} \cdots \theta_{p_k}|_{p_{k-1}^\downarrow}) = \text{id}_{p_1^\downarrow} = \varepsilon(\text{id}_{p_1^\downarrow}) = \varepsilon(\mathbf{d}(\mathbf{p})). \end{aligned}$$

Dually, we have $\mathbf{r}(\varepsilon(\mathbf{p})) = \varepsilon(\mathbf{r}(\mathbf{p}))$. This proves (E2). If $\mathbf{r}(\mathbf{p}) = \mathbf{d}(\mathbf{s})$, then

$$\begin{aligned} \varepsilon(\mathbf{p} \circ \mathbf{s}) &= \varepsilon(\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}, \dots, \text{id}_{p_k^\downarrow} = \text{id}_{s_1^\downarrow}, \text{id}_{s_2^\downarrow}, \dots, \text{id}_{s_l^\downarrow}) \\ &= (\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-1}, p_k})(\gamma_{s_1, s_2} \gamma_{s_2, s_3} \cdots \gamma_{s_{l-1}, s_l}) = \varepsilon(\mathbf{p}) \circ \varepsilon(\mathbf{s}). \end{aligned}$$

Thus (E3) is true.

Let $\text{id}_{q^\downarrow} \in P_{M(P)}$ and $\text{id}_{q^\downarrow} \leq_{M(P)} \mathbf{d}(\mathbf{p})$. In the case that $k = 1$, by the fact that $\text{id}_{q^\downarrow} \leq_{M(P)} \mathbf{d}(\mathbf{p}) = \text{id}_{p_1^\downarrow}$, Lemma 2.2 (11) and (E1),

$$\varepsilon(\text{id}_{q^\downarrow} \downarrow \mathbf{p}) = \varepsilon(\text{id}_{q^\downarrow} \downarrow \text{id}_{p_1^\downarrow}) = \varepsilon(\text{id}_{q^\downarrow}) = \text{id}_{q^\downarrow} = \text{id}_{q^\downarrow} \downarrow \text{id}_{p_1^\downarrow} = \text{id}_{q^\downarrow} \downarrow \varepsilon(\text{id}_{p_1^\downarrow}) = \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p}).$$

Let $\text{id}_{q^\downarrow} \downarrow \mathbf{p} = (\text{id}_{q_1^\downarrow}, \text{id}_{q_2^\downarrow}, \dots, \text{id}_{q_k^\downarrow})$ and

$$\mathbf{p}' = (\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}, \dots, \text{id}_{p_{k-1}^\downarrow}), \quad \text{id}_{q^\downarrow} \downarrow \mathbf{p}' = (\text{id}_{q_1^\downarrow}, \text{id}_{q_2^\downarrow}, \dots, \text{id}_{q_{k-1}^\downarrow}).$$

Then by induction hypothesis,

$$\begin{aligned} \varepsilon(\text{id}_{q^\downarrow} \downarrow \mathbf{p}) &= \varepsilon(\text{id}_{q_1^\downarrow}, \text{id}_{q_2^\downarrow}, \dots, \text{id}_{q_k^\downarrow}) = \gamma_{q_1, q_2} \gamma_{q_2, q_3} \cdots \gamma_{q_{k-1}, q_k} \\ &= \varepsilon(\text{id}_{q^\downarrow} \downarrow \mathbf{p}') \circ \gamma_{q_{k-1}, q_k} = \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p}') \circ \gamma_{q_{k-1}, q_k}. \end{aligned}$$

Denote $Y = q^\downarrow(\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-2}, p_{k-1}})$. By (84),

$$\begin{aligned} \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p}) &= (\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-1}, p_k})|_{q^\downarrow} \\ &= (\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-2}, p_{k-1}})|_{q^\downarrow} \circ \gamma_{p_{k-1}, p_k}|_Y = \varepsilon(\mathbf{p}')|_{q^\downarrow} \circ \gamma_{p_{k-1}, p_k}|_Y = \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p}') \circ \gamma_{p_{k-1}, p_k}|_Y. \end{aligned}$$

We shall prove that $\gamma_{p_{k-1}, p_k}|_Y = \gamma_{q_{k-1}, q_k}$. By (16),

$$\begin{aligned} Y &= q^\downarrow(\gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-2}, p_{k-1}}) \\ &= (q \gamma_{p_1, p_2} \gamma_{p_2, p_3} \cdots \gamma_{p_{k-2}, p_{k-1}})^\downarrow = (q \theta_{p_2} \cdots \theta_{p_{k-1}})^\downarrow = q_{k-1}^\downarrow. \end{aligned}$$

This together with the fact that $q_{k-1} \leq_P p_{k-1}$ implies that

$$\gamma_{p_{k-1}, p_k}|_Y = (\theta_{p_k}|_{p_{k-1}^\downarrow})|_{q_{k-1}^\downarrow} = \theta_{p_k}|_{q_{k-1}^\downarrow}.$$

Let $t \in q_{k-1}^\downarrow$. By (11) and (16), we have $t = t \theta_{q_{k-1}} = t \star q_{k-1}$ and $q_k = q_{k-1} \theta_{p_k} = q_{k-1} \star p_k$. This together with (R3) implies that

$$t \theta_{q_k} = t \star q_k = (t \star q_{k-1}) \star (q_{k-1} \star p_k) = (t \star q_{k-1}) \star p_k = t \theta_{q_{k-1}} \theta_{p_k} = t \theta_{p_k}.$$

Thus $\gamma_{p_{k-1}, p_k}|_Y = \theta_{p_k}|_{q_{k-1}^\downarrow} = \theta_{q_k}|_{q_{k-1}^\downarrow} = \gamma_{q_{k-1}, q_k}$, and so $\varepsilon(\text{id}_{q^\downarrow} \downarrow \mathbf{p}) = \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p})$. By mathematical induction, for all $\text{id}_{q^\downarrow} \in P_{M(P)}$ and $\mathbf{p} \in \mathcal{P}(P_{M(P)})$, the fact that $\text{id}_{q^\downarrow} \leq_{M(P)} \mathbf{d}(\mathbf{p})$ implies that $\varepsilon(\text{id}_{q^\downarrow} \downarrow \mathbf{p}) = \text{id}_{q^\downarrow} \downarrow \varepsilon(\mathbf{p})$. This gives (E5). By Proposition 4.11, ε is an evaluation map. \square

Proposition 8.8. *Let (P, \times, \star) be a strong projection algebra. Then*

$$(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq_{M(P)}, P_{M(P)}, \times, \star, \varepsilon)$$

is a chain projection ordered category.

Proof. Let $\beta \in M(q, r)$, $\text{id}_{e^\downarrow}, \text{id}_{f^\downarrow} \in P_{M(P)}$, $e, f \in P$ and $(\text{id}_{e^\downarrow}, \text{id}_{f^\downarrow})$ be a β -linked pair. Then

$$(85) \quad \text{id}_{f^\downarrow} = \text{id}_{e^\downarrow} \Theta_\beta \theta_{\text{id}_{f^\downarrow}}, \quad \text{id}_{e^\downarrow} = \text{id}_{f^\downarrow} \Delta_\beta \delta_{\text{id}_{e^\downarrow}}.$$

By the definition of ε (see Lemma 8.7), (26)–(28), Lemma 8.5, (80) and (81),

$$\lambda(\text{id}_{e^\downarrow}, \beta, \text{id}_{f^\downarrow}) = \varepsilon(\text{id}_{e^\downarrow}, \text{id}_{e_1^\downarrow}) \circ \text{id}_{e_1^\downarrow} \downarrow \beta \circ \varepsilon(\text{id}_{f_1^\downarrow}, \text{id}_{f^\downarrow}) = \gamma_{ee_1} \beta|_{e_1^\downarrow} \gamma_{f_1 f} = \theta_{e_1}|_{e^\downarrow} \beta|_{e_1^\downarrow} \theta_f|_{f_1^\downarrow},$$

$$\rho(\text{id}_{e^\downarrow}, \beta, \text{id}_{f^\downarrow}) = \varepsilon(\text{id}_{e^\downarrow}, \text{id}_{e_2^\downarrow}) \circ \beta|_{\text{id}_{f_2^\downarrow}} \circ \varepsilon(\text{id}_{f_2^\downarrow}, \text{id}_{f^\downarrow}) = \gamma_{ee_2} \beta|_{e_2^\downarrow} \gamma_{f_2 f} = \theta_{e_2}|_{e^\downarrow} \beta|_{e_2^\downarrow} \theta_f|_{f_2^\downarrow},$$

where

$$\begin{aligned} \text{id}_{e_1^\downarrow} &= (\text{id}_{e^\downarrow}) \theta_{\text{id}_{q^\downarrow}} = \text{id}_{e^\downarrow} \star \text{id}_{q^\downarrow} = \text{id}_{(e \star q)^\downarrow}, \quad \text{id}_{e_2^\downarrow} = \text{id}_{f^\downarrow} \Delta_\beta = \text{id}_{((r \times f) \beta^{-1})^\downarrow}, \\ \text{id}_{f_1^\downarrow} &= \text{id}_{e^\downarrow} \Theta_\beta = \text{id}_{((e \star q) \beta)^\downarrow}, \quad \text{id}_{f_2^\downarrow} = \text{id}_{f_1^\downarrow} \delta_{\text{id}_{r^\downarrow}} = \text{id}_{r^\downarrow} \times \text{id}_{f^\downarrow} = \text{id}_{(r \times f)^\downarrow}. \end{aligned}$$

So

$$e_1 = e \star q, \quad e_2 = (r \times f)\beta^{-1}, \quad f_1 = (e \star q)\beta, \quad f_2 = r \times f.$$

We shall prove that $\lambda(\text{id}_{e\downarrow}, \beta, \text{id}_{f\downarrow}) = \rho(\text{id}_{e\downarrow}, \beta, \text{id}_{f\downarrow})$. This is equivalent to prove that $t\theta_{e_1}\beta\theta_f = t\theta_{e_2}\beta\theta_f$ for $t \in P$ with $t \in e\downarrow$. Since $t \leq_P e$, by (11) we have $t = t\theta_e = t \star e$. This together with the fact $e_1 = e\theta_q$ and (R3) implies that

$$t\theta_{e_1}\beta\theta_f = t\theta_e\theta_{e\theta_q}\beta\theta_f = ((t \star e) \star (e \star q))\beta\theta_f = ((t \star e) \star q)\beta\theta_f = t\theta_e\theta_q\beta\theta_f = t\theta_q\beta\theta_f.$$

On the other hand, let $p \in P$. By Lemma 4.14 (2),

$$\text{id}_{f\downarrow} = \mathbf{r}(\beta) \star \text{id}_{f\downarrow} = \text{id}_{r\downarrow} \star \text{id}_{f\downarrow} = \text{id}_{(r \star f)\downarrow},$$

and so $f = r \star f$. Since $p \star f \leq_P f = r \star f$ by Lemma 3.1 (2), we obtain $(r \times (p \star f)) \star f = p \star f$ by (8). This implies that

$$p\theta_f\delta_r\theta_f = (r \times (p \star f)) \star f = p \star f = p\theta_f.$$

Thus $\theta_f\delta_r\theta_f = \theta_f$. In view of (G1b), (80) and (81), we have

$$\text{id}_{(t \star e_2)\downarrow} = \text{id}_{t\downarrow} \star \text{id}_{e_2\downarrow} = \text{id}_{t\downarrow}\theta_{\text{id}_{e_2\downarrow}} = \text{id}_{t\downarrow}\theta_{\text{id}_{f\downarrow}\Delta_\beta} = \text{id}_{t\downarrow}\Theta_\beta\theta_{\text{id}_{f\downarrow}}\Delta_\beta = \text{id}_{((r \times ((t \star q)\beta \star f))\beta^{-1})\downarrow},$$

and so $t\theta_{e_2} = t \star e_2 = (r \times ((t \star q)\beta \star f))\beta^{-1}$. By the above statements,

$$t\theta_{e_2}\beta\theta_f = (r \times ((t \star q)\beta \star f))\beta^{-1}\beta\theta_f = (r \times ((t \star q)\beta \star f))\theta_f = t\theta_q\beta\theta_f\delta_r\theta_f = t\theta_q\beta\theta_f.$$

This gives that $t\theta_{e_1}\beta\theta_f = t\theta_{e_2}\beta\theta_f$. Therefore $\lambda(\text{id}_{e\downarrow}, \beta, \text{id}_{f\downarrow}) = \rho(\text{id}_{e\downarrow}, \beta, \text{id}_{f\downarrow})$. \square

Let (P, \times, \star) be a strong projection algebra. Define a binary operation “ \bullet ” and two unary operations “ $+$ ” and “ $*$ ” on $M(P)$ as follows: For all $\alpha \in M(p, q), \beta \in M(s, t)$,

$$\alpha \bullet \beta = \alpha|_{\mathbf{r}(\alpha) \times \mathbf{d}(\beta)} \circ \varepsilon(\mathbf{r}(\alpha) \times \mathbf{d}(\beta), \mathbf{r}(\alpha) \star \mathbf{d}(\beta)) \circ_{\mathbf{r}(\alpha) \star \mathbf{d}(\beta)} \downarrow \beta, \quad \alpha^+ = \mathbf{d}(\alpha), \quad \alpha^* = \mathbf{r}(\alpha).$$

By (44), Theorem 6.4 and Proposition 8.8, $\mathbf{S}(M(P)) = (M(P), \bullet, +, *)$ forms a DRC-restriction semigroup ([cf. Theorem 4.7 in [59]]). Since $\mathbf{d}(\alpha) = \text{id}_{p\downarrow}$ and $\mathbf{r}(\alpha) = \text{id}_{q\downarrow}$, we have $\alpha^+ = \text{id}_{p\downarrow}$ and $\alpha^* = \text{id}_{q\downarrow}$. Moreover,

$$\alpha|_{\mathbf{r}(\alpha) \times \mathbf{d}(\beta)} = \alpha|_{\text{id}_{q\downarrow} \times \text{id}_{s\downarrow}} = \alpha|_{\text{id}_{(q \times s)\downarrow}} = \alpha|_{((q \times s)\alpha^{-1})\downarrow}.$$

Similarly, we have

$$\mathbf{r}(\alpha) \star \mathbf{d}(\beta) \downarrow \beta = \beta|_{(q \star s)\downarrow}, \quad \varepsilon(\mathbf{r}(\alpha) \times \mathbf{d}(\beta), \mathbf{r}(\alpha) \star \mathbf{d}(\beta)) = \varepsilon(\text{id}_{(q \times s)\downarrow}, \text{id}_{(q \star s)\downarrow}) = \gamma_{q \times s, q \star s}.$$

Thus

$$(86) \quad \alpha \bullet \beta = \alpha|_{((q \times s)\alpha^{-1})\downarrow} \circ \gamma_{q \times s, q \star s} \circ \beta|_{(q \star s)\downarrow} = \alpha\theta_{q \star s}\beta|_{((q \times s)\alpha^{-1})\downarrow}.$$

If $\mathbf{r}(\alpha) = \mathbf{d}(\beta)$, i.e. $q = s$, we have $q \times s = q \star s = q = s$, and so $(q \times s)\alpha^{-1} = q\alpha^{-1} = p$. This yields that $\alpha \bullet \beta = \alpha\theta_q\beta|_{p\downarrow} = \alpha\beta$. In particular, for any $\alpha \in M(p, q)$, we have $\alpha^{-1} \in M(q, p)$, and so

$$\alpha \bullet \alpha^{-1} = \alpha\alpha^{-1} = \text{id}_{p\downarrow} = \alpha^+ = (\alpha^{-1})^*, \quad \alpha^{-1} \bullet \alpha = \alpha^{-1}\alpha = \text{id}_{q\downarrow} = \alpha^* = (\alpha^{-1})^+.$$

By Lemma 5.1, $\mathbf{S}(M(P))$ is a generalized regular \circ -semigroup with set of projections $P(\mathbf{S}(M(P))) = P_{M(P)} = \{\text{id}_{p\downarrow} \mid p \in P\}$. Let $p, q \in P$. By Lemma 6.3,

$$\text{id}_{p\downarrow} \bullet \text{id}_{q\downarrow} = \varepsilon(\text{id}_{p\downarrow} \times \text{id}_{q\downarrow}, \text{id}_{p\downarrow} \star \text{id}_{q\downarrow}) = \varepsilon(\text{id}_{(p \times q)\downarrow}, \text{id}_{(p \star q)\downarrow}) = \gamma_{p \times q, p \star q}.$$

In particular, if $p \mathcal{F}_P q$, then $\text{id}_{p\downarrow} \bullet \text{id}_{q\downarrow} = \gamma_{p, q}$.

Theorem 8.9 (cf. Theorem 5.3 in [59]). *Let $(S, \cdot, +, *)$ be a DRC-restriction semigroup and $(P(S), \times_S, \star_S)$ be the strong projection algebra of S . For all $a \in S$, denote*

$$\nu_a : (a^+)^\downarrow \rightarrow (a^*)^\downarrow, \quad x \mapsto (xa)^*,$$

where

$$(a^+)^\downarrow = \{x \in P(S) \mid x \leq_S a^+\}, \quad (a^*)^\downarrow = \{x \in P(S) \mid x \leq_S a^*\}.$$

Then $\phi_S : S \rightarrow \mathbf{S}(M(P(S)))$, $a \mapsto \nu_a$ is a $(2,1,1)$ -homomorphism with $\ker \phi_S = \mu_S$ and $p\phi_S = \text{id}_{p^\downarrow}$ for all $p \in P(S)$, and so S is fundamental if and only if ϕ is injective. Moreover, $\text{im}(\phi_S) \cong S/\mu_S$ is fundamental.

Proof. By Theorem 5.11, we can consider the chain projection ordered category

$$\mathbf{C}(S) = (S, \circ_S, \mathbf{d}_S, \mathbf{r}_S, \leq_S, P_S, \times_S, \star_S, \varepsilon_S).$$

Let $a \in S$. By (G1d) and (41),

$$\nu_a : (a^+)^\downarrow = \mathbf{d}_S(a)^\downarrow \rightarrow \mathbf{r}_S(a)^\downarrow = (a^*)^\downarrow, \quad x \mapsto \mathbf{r}_S(x \downarrow a) = (xa)^*$$

is an isomorphism, and so $\nu_a \in M(a^+, a^*)$. So we can define

$$\phi_S : \mathbf{C}(S) \rightarrow M(P(S)), \quad a \rightarrow \nu_a.$$

We shall prove that ϕ_S is a chain projection ordered functor. Let $a, b \in S$.

(F1) Firstly, we have $\mathbf{d}(a\phi_S) = \mathbf{d}(\nu_a) = \text{id}_{(a^+)^\downarrow}$. Since $a^+ \in P(S) = P_S$, it follows that $\nu_{a^+} = \text{id}_{(a^+)^\downarrow}$ by Lemma 2.3. This implies that $\mathbf{d}(a\phi_S) = \nu_{a^+} = a^+\phi_S = (\mathbf{d}_S(a))\phi_S$. Dually, $\mathbf{r}(a\phi_S) = (\mathbf{r}_S(a))\phi_S$.

(F2) Let $\mathbf{r}_S(a) = \mathbf{d}_S(b)$. Then $a^* = b^+$ and $\mathbf{r}(\nu_a) = \text{id}_{(a^*)^\downarrow} = \text{id}_{(b^+)^\downarrow} = \mathbf{d}(\nu_b)$. By Lemma 2.5,

$$(a \circ_S b)\phi_S = \nu_{a \circ_S b} = \nu_a \nu_b = \nu_a \circ \nu_b = (a\phi_S) \circ (b\phi_S).$$

(F3) Let $a \leq_S b$. By Lemma 2.2 (3), we obtain $a = \mathbf{d}_S(a) \downarrow b = a^+ \downarrow b$, this together with Lemma 2.4 gives that $\nu_a = \nu_{a^+ \downarrow b} = \nu_b \downarrow_{(a^+)^\downarrow} = \nu_b \downarrow_{\text{dom} \nu_a}$. This shows that $a\phi_S = \nu_a \leq_{M(P)} \nu_b = b\phi_S$ in $(M(P), \circ, \mathbf{d}, \mathbf{r}, \leq)$.

(F4) Let $e, f \in P_S = P(S)$. By Lemma 2.3,

$$(e \times_S f)\phi_S = \nu_{e \times_S f} = \text{id}_{(e \times_S f)^\downarrow} = \text{id}_{e^\downarrow} \times \text{id}_{f^\downarrow} = \nu_e \times \nu_f = (e\phi_S) \times (f\phi_S).$$

(F5) Let $\mathbf{c} = (p_1, p_2, \dots, p_k) \in \mathcal{P}(P(S))$. Using (24), (F2), Lemma 4.12 (2), (83) and Lemma 8.7, we have

$$\begin{aligned} (\varepsilon_S(\mathbf{c}))\phi_S &= (\varepsilon_S(p_1, p_2) \circ \dots \circ \varepsilon_S(p_{k-1}, p_k))\phi_S = (\varepsilon_S(p_1, p_2))\phi_S \circ \dots \circ (\varepsilon_S(p_{k-1}, p_k))\phi_S \\ &= \nu_{\varepsilon_S(p_1, p_2)} \circ \dots \circ \nu_{\varepsilon_S(p_{k-1}, p_k)} = \theta_{p_2 \downarrow_{p_1^\downarrow}} \circ \dots \circ \theta_{p_k \downarrow_{p_{k-1}^\downarrow}} \\ &= \gamma_{p_1, p_2} \circ \dots \circ \gamma_{p_{k-1}, p_k} = \varepsilon(\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}) \circ \dots \circ \varepsilon(\text{id}_{p_{k-1}^\downarrow}, \text{id}_{p_k^\downarrow}) = \varepsilon(\text{id}_{p_1^\downarrow}, \text{id}_{p_2^\downarrow}, \dots, \text{id}_{p_k^\downarrow}). \end{aligned}$$

Since $p_i \in P(S)$ for all i , it follows that $p_i\phi_S = \nu_{p_i} = \text{id}_{p_i^\downarrow}$ by Lemma 2.3. Thus $(\varepsilon_S(\mathbf{c}))\phi_S = \varepsilon((p_1\phi_S, p_2\phi_S, \dots, p_k\phi_S))$. By items (F1)–(F5), ϕ_S is a chain projection ordered functor. By Lemma 6.8 and Theorem 6.11, ϕ_S is a $(2,1,1)$ -homomorphism from $S = \mathbf{S}(\mathbf{C}(S))$ to $\mathbf{S}(M(P(S)))$, and $p\phi_S = \nu_p = \text{id}_{p^\downarrow}$ for all $p \in P(S)$. Finally,

$$\ker \phi_S = \{(a, b) \in S \times S \mid a\phi_S = b\phi_S\} = \{(a, b) \in S \times S \mid \nu_a = \nu_b\} = \mu_S$$

by Proposition 8.3. The remaining part of the theorem is obvious. \square

Theorem 8.10 (cf. Theorem 5.4 in [59]). *Let (P, \times, \star) be a strong projection algebra and $(S, \cdot, ^+, \star)$ be a DRC-restriction semigroup whose strong projection algebra is isomorphic to (P, \times, \star) . Then S is fundamental if and only if S is isomorphic to some full $(2,1,1)$ -subalgebra of $\mathbf{S}(M(P))$. In particular, $\mathbf{S}(M(P))$ is projection-fundamental.*

Proof. Assume that S is projection-fundamental. By Theorem 8.9,

$$\phi_S : S \rightarrow \mathbf{S}(M(P(S))) (\cong \mathbf{S}(M(P)))$$

is an injective $(2,1,1)$ -homomorphism and $p\phi_S = \text{id}_{p\downarrow}$ for all $p \in P(S)$. Thus

$$\text{im}\phi_S \supseteq P(S)\phi_S = \{p\phi_S \mid p \in P(S)\} = \{\text{id}_{p\downarrow} \mid p \in P(S)\} = P(\mathbf{S}(M(P(S)))).$$

This implies that $S \cong \text{im}\phi_S$ and $\text{im}\phi_S$ is a full $(2,1,1)$ -subalgebra of $\mathbf{S}(M(P))$.

Conversely, assume that T is a full $(2,1,1)$ -subalgebra of $\mathbf{S}(M(P))$. Let $\alpha, \beta \in T$ and $(\alpha, \beta) \in \mu_T$. By Lemma 8.1, we have $\text{id}_{\text{dom}\alpha} = \alpha^+ = \beta^+ = \text{id}_{\text{dom}\beta}$, and so we can denote $e^\downarrow = \text{dom}\alpha = \text{dom}\beta$ for some $e \in P$. Let $x \in e^\downarrow$. Then $x \in P$ and $x \leq_P e$, and so $\text{id}_{x\downarrow} \in P(\mathbf{S}(M(P))) \subseteq T$. This gives that $(\text{id}_{x\downarrow} \bullet \alpha, \text{id}_{x\downarrow} \bullet \beta) \in \mu_T$. Similar discussion gives that $\text{ran}(\text{id}_{x\downarrow} \bullet \alpha) = \text{ran}(\text{id}_{x\downarrow} \bullet \beta)$. By (86), we have $x^\downarrow\alpha = x^\downarrow\beta$, i.e. $(x\alpha)^\downarrow = (x\beta)^\downarrow$. Thus $x\alpha = x\beta$. We have shown that $\alpha = \beta$. Therefore μ_T is the identity relation and so T is projection-fundamental. \square

Theorem 8.11. *Let (P, \times, \star) be a strong projection algebra, $\mu = \mu_{\mathbf{S}(\mathcal{C}(P))}$ and $(S, \cdot, ^+, \star)$ be a projection-fundamental DRC-restriction semigroup whose strong projection algebra is isomorphic to (P, \times, \star) . Then $\mathbf{E}(S) \cong \mathbf{S}(\mathcal{C}(P))/\mu$. In particular if $(S, \cdot, ^+, \star)$ is projection-generated, then $S \cong \mathbf{S}(\mathcal{C}(P))/\mu$.*

Proof. Assume that $(S, \cdot, ^+, \star)$ is a projection-fundamental DRC-restriction semigroup whose strong projection algebra is (P, \times, \star) . By Theorem 5.11, $\mathbf{C}(S)$ is a chain projection ordered category. By Proposition 7.11, $\bar{\varepsilon} : \mathcal{C}(P_S) \rightarrow \mathbf{C}(S)$ is a chain projection ordered functor from $\mathcal{C}(P_S)$ to $\mathbf{C}(S)$, and so is a $(2,1,1)$ -homomorphism from $\mathbf{S}(\mathcal{C}(P_S))$ to $\mathbf{S}(\mathbf{C}(S)) = S$ (by Theorem 6.11) such that $[p]\bar{\varepsilon} = p$ for all $p \in P$. According to Theorem 8.9, there exists a $(2,1,1)$ -homomorphism

$$\phi_S : S \rightarrow \mathbf{S}(M(P)), \quad a \rightarrow \nu_a$$

such that $p \mapsto \text{id}_{p\downarrow}$ for all $p \in P$. This implies that

$$\bar{\varepsilon}\phi_S : \mathbf{S}(\mathcal{C}(P_S)) \rightarrow \mathbf{S}(M(P))$$

is also a $(2,1,1)$ -homomorphism such that $[p]\bar{\varepsilon}\phi_S = \text{id}_{p\downarrow}$ for all $p \in P$. In view of Theorem 8.9, we have a $(2,1,1)$ -homomorphism

$$\phi_{\mathbf{S}(\mathcal{C}(P_S))} : \mathbf{S}(\mathcal{C}(P_S)) \longrightarrow \mathbf{S}(M(P))$$

such that $[p] = \text{id}_{p\downarrow}$ for all $p \in P$. By Theorem 7.12, $\mathbf{S}(\mathcal{C}(P_S))$ is projection-generated, and so $\bar{\varepsilon}\phi_S = \phi_{\mathbf{S}(\mathcal{C}(P_S))}$. Since S is projection-fundamental, it follows that ϕ_S is injective by Theorem 8.9. This together with Theorem 8.9 gives that

$$\ker(\bar{\varepsilon}) = \ker(\bar{\varepsilon}\phi_S) = \ker(\phi_{\mathbf{S}(\mathcal{C}(P_S))}) = \mu_{\mathbf{S}(\mathcal{C}(P_S))}.$$

By Proposition 6.5,

$$\mathbf{E}(S) = \text{im}(\bar{\varepsilon}) \cong \mathbf{S}(\mathcal{C}(P_{\mathbf{C}(S)}))/\mu_{\mathbf{S}(\mathcal{C}(P_{\mathbf{C}(S)})})} \cong \mathbf{S}(\mathcal{C}(P))/\mu.$$

In particular, if $(S, \cdot, ^+, \star)$ is projection-generated, then $S = \mathbf{E}(S) \cong \mathbf{S}(\mathcal{C}(P))/\mu$ by Proposition 6.5 again. \square

Let (P, \times, \star) be a strong projection algebra. To end this section, we shall give some remarks on the semigroup $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$. By Theorem 8.11, up to isomorphism, there exists a unique projection-generated projection-fundamental DRC-restriction semigroup whose strong projection algebra is isomorphic to (P, \times, \star) . This semigroup is $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$. Moreover, if $(S, \cdot, +, *)$ is a projection-fundamental DRC-restriction semigroup with $P(S) \cong P$, then $\mathbf{E}(S) \cong \mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$. To obtain $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$, we only need to find a projection-fundamental DRC-restriction semigroup $(S, \cdot, +, *)$ with $P(S) \cong P$ and compute $\mathbf{E}(S)$. So we have the following three approaches to describe $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$.

Approach 1: The elements in $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$ are all $\mu_{\mathbf{S}(\mathcal{C}(P))}$ -classes in $\mathbf{S}(\mathcal{C}(P))$. Let

$$\mathbf{p} = [p_1, p_2, \dots, p_k], \mathbf{q} = [q_1, q_2, \dots, q_l] \in \mathbf{S}(\mathcal{C}(P)).$$

By Proposition 8.3 and (56), (57), (58), (59),

$$\begin{aligned} (\mathbf{p}, \mathbf{q}) \in \mu_{\mathbf{S}(\mathcal{C}(P))} &\iff \theta_{p_1}\theta_{p_2}\dots\theta_{p_k} = \theta_{q_1}\theta_{q_2}\dots\theta_{q_l}, \delta_{p_k}\delta_{p_{k-1}}\dots\delta_{p_1} = \delta_{q_l}\delta_{q_{l-1}}\dots\delta_{q_1} \\ &\iff \theta_{p_1}\theta_{p_2}\dots\theta_{p_k} = \theta_{q_1}\theta_{q_2}\dots\theta_{q_l}, p_1 = q_1 \\ &\iff p_1 = q_1, \theta_{p_2}\theta_{p_3}\dots\theta_{p_k}|_{p_1^\downarrow} = \theta_{q_2}\theta_{q_3}\dots\theta_{q_l}|_{q_1^\downarrow}. \end{aligned}$$

Approach 2: Since $\mathbf{S}(M(P))$ is projection-fundamental and its strong projection algebra is isomorphic to (P, \times, \star) . By Theorem 8.11, $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$ is isomorphic to $\mathbf{E}(\mathbf{S}(M(P)))$. The elements in $\mathbf{E}(\mathbf{S}(M(P)))$ are:

$$\text{id}_{p_1^\downarrow} \bullet \text{id}_{p_2^\downarrow} \bullet \dots \bullet \text{id}_{p_k^\downarrow}, p_1, p_2, \dots, p_k \in P, k \text{ is an integer.}$$

Approach 3: Let (P, \times, \star) be a strong projection algebra. Then we have the generalized regular \circ -semigroup $\mathbf{S}(M(P)) = (M(P), \bullet, +, *, ^{-1})$. Denote

$$\overline{\mathbf{S}(M(P))} = \{(\theta_p\alpha, \delta_q\alpha^{-1}) \mid \alpha \in M(p, q), p, q \in P\}$$

and define

$$\varphi : \mathbf{S}(M(P)) \rightarrow \overline{\mathbf{S}(M(P))}, \alpha \in M(p, q) \mapsto (\theta_p\alpha, \delta_q\alpha^{-1}).$$

We assert that φ is a bijection. Obviously, φ is surjective. Let $\alpha \in M(p, q), \beta \in M(s, t)$ and $(\theta_p\alpha, \delta_q\alpha^{-1}) = (\theta_s\beta, \delta_t\beta^{-1})$. Then $\theta_p\alpha = \theta_s\beta, \delta_q\alpha^{-1} = \delta_t\beta^{-1}$. Let $x \in P$. Then $x(\theta_p\alpha) = x(\theta_s\beta)$, i.e. $(x \star p)\alpha = (x \star s)\beta$. Take $x = p$. Then $q = p\alpha = (p \star p)\alpha = (p \star s)\beta \leq_P s\beta = t$. Similarly, take $x = s$. Then we have $t \leq_P q$. Thus $q = t$. Dually, we can obtain $p = s$ from the fact that $\delta_q\alpha^{-1} = \delta_t\beta^{-1}$. Thus $\alpha, \beta \in M(p, q)$. Take $x \in \text{dom}\alpha = p^\downarrow$ arbitrarily. Then $x \leq_P p$, and so $x \star p = x$ by Lemma 3.1 (1). Thus

$$x\alpha = (x \star p)\alpha = x\theta_p\alpha = x\theta_p\beta = (x \star p)\beta = x\beta.$$

This gives $\alpha = \beta$. Therefore φ is injective.

Since φ is bijective, we can transfer the operations on $\mathbf{S}(M(P))$ to $\overline{\mathbf{S}(M(P))}$ as follows. Let $\alpha \in M(p, q), \beta \in M(s, t)$. By (86), $\alpha \bullet \beta \in M(((q \times s)\alpha^{-1}, (q \star s)\beta))$. So we define

$$\begin{aligned} (\theta_p\alpha, \delta_q\alpha^{-1})(\theta_s\beta, \delta_t\beta^{-1}) &= (\alpha \bullet \beta)\varphi = (\alpha\gamma_{q \times s, q \star s}\beta)\varphi \\ &= (\theta_{(q \times s)\alpha^{-1}}\alpha\gamma_{q \times s, q \star s}\beta, \delta_{(q \star s)\beta}\beta^{-1}\gamma_{q \times s, q \star s}^{-1}\alpha^{-1}), \\ (\theta_p\alpha, \delta_q\alpha^{-1})^+ &= \alpha^+\varphi = \text{id}_{p^\downarrow}\varphi = (\theta_p\text{id}_{p^\downarrow}, \delta_p\text{id}_{p^\downarrow}^{-1}) = (\theta_p, \delta_p). \\ (\theta_p\alpha, \delta_q\alpha^{-1})^* &= \alpha^*\varphi = \text{id}_{q^\downarrow}\varphi = (\theta_q\text{id}_{q^\downarrow}, \delta_q\text{id}_{q^\downarrow}^{-1}) = (\theta_q, \delta_q). \end{aligned}$$

Let $\alpha \in M(p, q)$ and $\beta \in M(s, t)$. Then

$$\alpha \bullet \beta = \alpha\gamma_{q \times s, q \star s}\beta \in M(((q \times s)\alpha^{-1}, (q \star s)\beta)).$$

Assume that $x \in P$. Then we have

$$\begin{aligned}
x\theta_{(q \times s)\alpha^{-1}} &= x\theta_p\theta_{(q \times s)\alpha^{-1}} \quad (\text{by } (q \times s)\alpha^{-1} \leq_P p \text{ and Lemma 3.2}) \\
&= (x\theta_p)\alpha\alpha^{-1}\theta_{(q \times s)\alpha^{-1}} \\
&= (x\theta_p\alpha\theta_{q \times s})\alpha^{-1} \quad (\text{since } \alpha^{-1} \text{ is an isomorphism}) \\
&= x\theta_p\alpha\theta_{q \times s}\delta_q\alpha^{-1} \quad (\text{by } q \times s \leq_P q, \text{ and Lemma 3.2}) \\
&= x\theta_p\alpha\theta_{(q \star s)\delta_q}\delta_q\alpha^{-1} \quad (\text{by } (q \star s)\delta_q = q \times (q \star s) = q \times s \text{ and Lemma 3.1 (3)}) \\
&= x\theta_p\alpha\theta_q\theta_{q \star s}\delta_q\delta_q\alpha^{-1} \quad (\text{by Lemma 3.7}) \\
&= x\theta_p\alpha\theta_{q \star s}\delta_q\alpha^{-1} \quad (\text{by } (x\theta_p)\alpha \in q^\downarrow \text{ and Lemma 3.2})
\end{aligned}$$

(R4) gives that $q \star (q \star s) = (q \star s) \star (q \star s) = q \star s$ and Lemma 3.1 (2) gives that

$$(x\theta_p\alpha) \star (q \star s) \leq_P q \star s = q \star (q \star s).$$

This together with (8) implies that

$$(q \times ((x\theta_p\alpha) \star (q \star s))) \star (q \star s) = (x\theta_p\alpha) \star (q \star s).$$

Thus

$$\begin{aligned}
x\theta_{(q \times s)\alpha^{-1}}\alpha\gamma_{q \times s, q \star s}\beta &= x\theta_{(q \times s)\alpha^{-1}}\alpha\theta_{q \star s}\beta = x\theta_p\alpha\theta_{q \star s}\delta_q\alpha^{-1}\alpha\theta_{q \star s}\beta = x\theta_p\alpha\theta_{q \star s}\delta_q\theta_{q \star s}\beta \\
&= ((q \times ((x\theta_p\alpha) \star (q \star s))) \star (q \star s))\beta = ((x\theta_p\alpha) \star (q \star s))\beta \quad ((8)) \\
&= ((x\theta_p\alpha \star q) \star (q \star s))\beta \quad (x\theta_p\alpha \leq_P q, \text{ Lemma 3.1 (2)}) \\
&= ((x\theta_p\alpha \star q) \star s)\beta \quad ((R3)) \\
&= (x\theta_p\alpha \star s)\beta = x\theta_p\alpha\theta_s\beta.
\end{aligned}$$

Dually, $\delta_{(q \star s)\beta}\beta^{-1}\gamma_{q \times s, q \star s}^{-1}\alpha^{-1} = \delta_t\beta^{-1}\delta_q\alpha^{-1}$. Thus

$$(\theta_p\alpha, \delta_q\alpha^{-1})(\theta_s\beta, \delta_t\beta^{-1}) = (\theta_p\alpha\theta_s\beta, \delta_t\beta^{-1}\delta_q\alpha^{-1}).$$

So φ is a (2,1,1)-isomorphism from $\mathbf{S}(M(P))$ onto $\overline{\mathbf{S}(M(P))}$. Since $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$ is isomorphic to $\mathbf{E}(\mathbf{S}(M(P)))$, $\mathbf{S}(\mathcal{C}(P))/\mu_{\mathbf{S}(\mathcal{C}(P))}$ is isomorphic to $\mathbf{E}(\overline{\mathbf{S}(M(P))})$. Observe that the projections in $\overline{\mathbf{S}(M(P))}$ are: $(\theta_p, \delta_p), p \in P$, it follows that the elements in $\mathbf{E}(\overline{\mathbf{S}(M(P))})$ are:

$$(\theta_{p_1}\theta_{p_2}\cdots\theta_{p_k}, \delta_{p_k}\delta_{p_{k-1}}\cdots\delta_{p_1}), p_1, p_2, \dots, p_k \in P, k \text{ is an integer.}$$

Acknowledgements: The paper is supported jointly by the Nature Science Foundations of China (12271442, 11661082).

REFERENCES

- [1] C. L. Adair, Bands with an involution. *J. Algebra* 75 (1982), 297–314.
- [2] S. Armstrong, Structure of concordant semigroups. *J. Algebra* 118 (1988), 205–260.
- [3] K. Auinger, Strict regular \ast -semigroups, in: J.M. Howie, W.D. Munn, H.-J. Weinert (Eds.), *Proc. Conf. on Semigroups with Applications*, World Scientific, Singapore, 1992, pp. 190–204.
- [4] A. Batbedat, γ -demi-groupes, demi-modules, produit demi-direct, in *Semigroups Proceedings: Oberwolfach 1979*, Lecture Notes in Mathematics 855, Springer-Verlag, 1981, pp. 1–18.
- [5] A. Batbedat, J. B. Fountain, Connections between left adequate semi-groups and γ -semigroups. *Semigroup Forum* 22 (1981), 59–65.
- [6] J. R. Cockett, S. Lack, Restriction categories I: categories of partial maps. *Theoretical Computer Science* 270 (2002), 223–259.

- [7] D. DeWolf, D. Pronk, The Ehresmann-Schein-Nambooripad theorem for inverse categories. *Theory Appl. Categ.* 33 (2018), 27.
- [8] J. East, P. A. Azeef Muhammed, A groupoid approach to regular $*$ -semigroups. *Advances in Mathematics* 437 (2024), 109447.
- [9] J. East, P. A. Azeef Muhammed, A groupoid approach to regular $*$ -semigroups. [arXiv:2301.04845v1](https://arxiv.org/abs/2301.04845), 2023.
- [10] J. East, R. D. Gray, P. A. Azeef Muhammed, N. Ruskuc, Projection algebras and free projection- and idempotent-generated regular $*$ -semigroups. [arXiv:2406.09109v1](https://arxiv.org/abs/2406.09109), 2024.
- [11] C. Ehresmann, Gattungen von lokalen Strukturen. *Jahresber. Dtsch. Math.-Ver.* 60 (Abt.1) (1957), 49–77.
- [12] C. Ehresmann, Categories inductives et pseudo-groupes. *Ann. Inst. Fourier (Grenoble)* 10 (1960), 307–332.
- [13] D. G. FitzGerald, M. K. Kinyon, Trace and pseudo-products: restriction-like semigroups with a band of projections. *Semigroup Forum* 103 (2021), 848–866.
- [14] J. B. Fountain, Adequate semigroups. *Proc. Edinb. Math. Soc.* 22 (1979), 113–123.
- [15] J. B. Fountain, G. M. S. Gomes, V. Gould, A Munn type representation for a class of E-semiadequate semigroups. *J. Algebra* 218 (1999), 693–714.
- [16] G. M. S. Gomes, V. Gould, Fundamental Ehresmann semigroups. *Semigroup Forum* 63 (2001), 11–33.
- [17] V. Gould, Restriction and Ehresmann semigroups. *Proceedings of the International Conference on Algebra 2010*, (World Sci. Publ., Hackensack, NJ, 2012), 265–288.
- [18] V. Gould, Y. Wang, Beyond orthodox semigroups. *J. Algebra* 368 (2012), 209–230.
- [19] C. Hollings, From right PP monoids to restriction semigroups: a survey. *Eur. J. Pure Appl. Math.* 2 (2009), 21–57.
- [20] C. Hollings, The Ehresmann-Schein-Nambooripad theorem and its successors. *Eur. J. Pure Appl. Math.* 5 (2012), 414–450.
- [21] C. Hollings, Three approaches to inverse semigroups. *Eur. J. Pure Appl. Math.* 8 (2015), 294–323.
- [22] J. M. Howie, *Fundamentals of semigroup theory*, Clarendon Press, Oxford, 1995.
- [23] T. Imaoka, Some remarks on fundamental regular $*$ -semigroups, in: *Recent Developments in the Algebraic, Analytical, and Topological Theory of Semigroups*, Oberwolfach, 1981, in: *Lecture Notes in Math.*, vol.998, Springer, Berlin, 1983, pp.270–280.
- [24] T. Imaoka, K. Fujiwara, Characterization of locally inverse $*$ -semigroups. *Sci. Math. Jpn.* 57 (2003), 49–55.
- [25] M. Jackson and T. Stokes, An invitation to C -semigroups. *Semigroup Forum* 62 (2001), 279–310.
- [26] P. R. Jones, A common framework for restriction semigroups and regular $*$ -semigroups. *J. Pure Appl. Algebra* 216 (2012), 618–632.
- [27] P. R. Jones, Varieties of P -restriction semigroups. *Commun. Algebra* 42 (2014), 1811–1834.
- [28] P. R. Jones, Beyond Ehresmann semigroups, invited presentation. *International Conference on Semigroups*, University of Lisbon, Lisbon, 2018.
- [29] P. R. Jones, Generalized Munn representations of DRC semigroups. *Southeast Asian Bull. Math.* 45 (2021), 591–619.
- [30] M. V. Lawson, *Inverse semigroups, the theory of partial symmetries*, World Scientific, Singapore, 1998.
- [31] M. V. Lawson, Semigroups and ordered categories I. The reduced case. *J. Algebra* 141 (1991), 422–462.
- [32] M. V. Lawson, On Ehresmann semigroups. *Semigroup Forum* 103 (2021), 953–965.
- [33] E. Manes, Guarded and banded semigroups. *Semigroup Forum* 72 (2006), 94–120.
- [34] D. B. McAlister, groups, semilattices and inverse semigroups, *Trans. American Math. Soc.* 192 (1974), 227–244.
- [35] D. B. McAlister, groups, semilattices and inverse semigroups II. *Trans. American Math. Soc.* 196 (1974), 351–370.
- [36] J. Meakin, On the structure of inverse semigroups. *Semigroup Forum* 12 (1976), 6–14.
- [37] J. Meakin, Coextensions of inverse semigroups. *J. Algebra* 46 (1977), 315–333.

- [38] W. D. Munn, Uniform semilattices and bisimple inverse semigroups. *Quart. J. of Math.* 17 (1966), 151–159.
- [39] W. D. Munn, Fundamental inverse semigroups. *Quart. J. Math.* 21 (1970), 157–170.
- [40] K. S. S. Nambooripad, Structure of regular semigroups I. *Mem. Am. Math. Soc.* 22 (224) (1979), vii+119.
- [41] T. Nordahl, H. E. Scheiblich, Regular $*$ -semigroups. *Semigroup Forum* 16 (1978), 369–377.
- [42] M. Petrich, *Inverse semigroups*, John Wiley and Sons, New York, 1984.
- [43] L. Polak, A solution of the word problem for free $*$ -regular semigroups. *J. Pure Appl. Alg.* 157 (2001), 107–114 .
- [44] B. M. Schein, On the theory of generalized groups and generalized heaps, in *The theory of semigroups and its applications I*, University of Saratov, Saratov, 1965, 286–324 (Russian).
- [45] B. M. Schein, On the theory of inverse semigroups and generalised groups. *Amer. Math. Soc. Translations* 113 (1979), 89–122.
- [46] I. Stein, Algebras of reduced E -Fountain semigroups and the generalized ample identity. *Commun. Algebra* 50 (2022), 2557–2570.
- [47] I. Stein, Algebras of reduced E -Fountain semigroups and the generalized ample identity II. *Journal of Algebra and Its Applications*, DOI: 10.1142/S0219498825500914
- [48] I. Stein, The algebra of the monoid of order-preserving functions on an n -set and other reduced E -Fountain semigroups. <https://arxiv.org/pdf/2404.08075v1>
- [49] T. Stokes, Domain and range operations in semigroups and rings. *Commun. Algebra* 43 (2015), 3979–4007.
- [50] T. Stokes, Ordered Ehresmann semigroups and categories. *Commun. Algebra* 50 (2022), 4805–4821.
- [51] T. Stokes, Ehresmann-Schein-Nambooripad theorems for classes of binary semigroups. *Semigroup Forum* 107 (2023), 751–779.
- [52] T. Stokes, Laws for generalised interior operations in semigroups. *Semigroup Forum* 106 (2023), 692–719.
- [53] M. B. Szendrei, A new interpretation of free orthodox and generalized inverse $*$ -semigroups, *Semigroups. Theory and Applications (Proc. Conf. Oberwolfach, 1986)*, *Lecture Notes in Math.*, 1320, Springer, Berlin, 1988, 358–371.
- [54] S. F. Wang, On algebras of P -Ehresmann semigroups and their associate partial semigroups. *Semigroup Forum* 95 (2017), 569–588.
- [55] S. F. Wang, An Ehresmann-Schein-Nambooripad-type theorem for a class of P -restriction semigroups. *Bull. Malays. Math. Sci. Soc.* 42 (2019), 535–568.
- [56] S. F. Wang, An Ehresmann–Schein–Nambooripad theorem for locally Ehresmann P -Ehresmann semigroups. *Period. Math. Hung.* 80 (2020), 108–137.
- [57] S. F. Wang, An Ehresmann-Schein-Nambooripad type theorem for DRC-semigroups. *Semigroup Forum* 104 (2022), 731–757.
- [58] S. F. Wang, K. P. Shum, On orthodox P -restriction semigroups. *J. Algebra Appl.* 22 (2023), 2350018 (16 pages).
- [59] S. F. Wang, A Munn type representation for DRC-restriction semigroups. *Period Math Hung.* 88 (2024), 148–171.
- [60] S. F. Wang, On d -semigroups, r -semigroups, dr -semigroups and their subclasses. *Semigroup Forum* 106 (2023), 230–270.
- [61] Y. Wang, Beyond regular semigroups. *Semigroup Forum* 92 (2016), 414–448.
- [62] M. Yamada, P -systems in regular semigroups. *Semigroup Forum* 24 (1982), 173–178.

SCHOOL OF MATHEMATICS, YUNNAN NORMAL UNIVERSITY, KUNMING, YUNNAN, 650500, CHINA.
Email address: 204585080@qq.com

SCHOOL OF MATHEMATICS, YUNNAN NORMAL UNIVERSITY, KUNMING, YUNNAN, 650500, CHINA.
Email address: wsf1004@163.com