

# Fermionic Casimir Energy in Horava-Lifshitz Scenario

E. R. Bezerra de Mello<sup>1,\*</sup> and M. B. Cruz<sup>2,†</sup>

<sup>1</sup>*Departamento de Física, Universidade Federal da Paraíba,  
Caixa Postal 5008, 58051-970, João Pessoa, Paraíba, Brazil.*

<sup>2</sup>*Centro de Ciências Exatas e Sociais Aplicadas, Universidade Estadual da Paraíba,  
R. Alfredo Lustosa Cabral, s/n, Salgadinho, Patos - PB, 58706-550 - Brazil.*

(Dated: July 17, 2024)

## Abstract

In this work, we investigate the violation of Lorentz symmetry through the Casimir effect. The Casimir effect is one of the most intriguing aspects of modern physics, representing a macroscopic quantum-origin force between two neutral conducting surfaces, and it stands as a triumph of Quantum Field Theory. Here, we examine the Casimir effects associated with a massive fermionic quantum field confined in the region between two large and parallel plates within the Horava-Lifshitz framework of Lorentz symmetry violation. To calculate the Casimir energy and consequently the Casimir pressure, we impose a MIT bag boundary condition on two plates, compatible with the higher-order derivative term in the modified Dirac equation. Our results indicate a strong influence of Lorentz violation on the Casimir effect. We observe that the Casimir energy is affected, both in intensity and sign, potentially exhibiting repulsive or attractive force between the plates, depending on the critical exponent associated with the Horava-Lifshitz formalism.

Keywords: Massive fermionic fields, Lorentz symmetry violation, Horava-Lifshitz, Casimir effect, MIT bag model.

---

\*Electronic address: [emello@fisica.ufpb.br](mailto:emello@fisica.ufpb.br)

†Electronic address: [messiasdebritocruz@servidor.uepb.edu.br](mailto:messiasdebritocruz@servidor.uepb.edu.br)

## I. INTRODUCTION

The Casimir effect refers to the interaction between neutral objects due to the boundary conditions imposed on the quantum fluctuations of fields at zero point. In the context of Quantum Field Theory (QFT), the vacuum state is understood as the sum of all states of quantum oscillators in their ground states. Therefore, the energy associated with the quantum vacuum results in an infinite value. In QFT, various phenomena are associated with the quantum vacuum, with the Casimir effect being one of the principal. This phenomenon is not only significant from a fundamental standpoint but also holds great technical importance in micro and nano mechanical systems.

In 1948, H. B. Casimir theorized the Casimir effect [1], originally considering the vacuum state associated with the electromagnetic field confined between two large, parallel, conducting, and neutral plates. Due to the boundary conditions imposed on the quantum field at the plates, only vacuum excitations with specific wavelengths are allowed between them. In this way, the Casimir energy can be obtained by subtracting the energy associated with these quantum fluctuations from the energy of the vacuum state in the absence of boundaries. This procedure is called renormalization. Using this method, Casimir found an attractive force between the plates, given by:

$$F = -A \frac{\pi^2 \hbar c}{240 a^4}, \quad (1)$$

where  $A = L^2$  represents the area of the plates and  $a$  denotes the distance between them, with  $a \ll L$ . This phenomenon has been confirmed experimentally ten years later by M. J. Sparnaay [2]; however, due to the low precision, many others experiments were carried out in the 90s, confirming the Casimir effect with a high degree of precision [3–5].

As previously mentioned, the Casimir effect was initially associated with the quantum vacuum for electromagnetic fields. However, in principle, this phenomenon can also occur for other types of relativistic quantum fields, such as scalar and fermionic fields, when subjected to specific boundary conditions. Studies on the Casimir effect considering scalar quantum fields have been conducted in various contexts, including different boundary conditions, temperature, and Lorentz symmetry violation [6–13]. In the context of fermionic quantum fields, several studies have been developed for different scenarios [14–24]. Typically, these studies employ the MIT bag model [23, 25], originally used to confine quarks in hadrons, assuming a spherical shape for them, as a boundary condition.

The standard QFT represents a successful partnership between the theory of Special Relativity and Quantum Mechanics, keeping Lorentz symmetry invariance. However, the unification of General Relativity with Quantum Mechanics remains one of the major challenge in contemporary physics, with Loop Quantum Gravity [26, 27] and String Theory [28, 29] being primary candidates. In [30], P. Horava proposed a formalism, named Horava-Lifshitz (HL) model, as an attempt to construct a renormalizable quantum field theory for gravity, introducing an anisotropy between space and time coordinates. This results in the breaking of Lorentz symmetry, an effect also expected in quantum gravity theories [31]. Although this breaking takes place only at Planck energy scales, residual signatures of spacetime anisotropy may be observable in phenomena at lower energy scales, such as in the study of the Casimir effects.

Motivated by the relevance of studying a purported violation of Lorentz symmetry, due to the fundamental aspects of Quantum Gravity, our aim is to utilize the Casimir effect as a laboratory. With the success of highly precise experimental techniques, we believe in the feasibility of investigating aspects related to the violation of Lorentz symmetry. In this sense the analysis of Casimir energy associated with massless scalar quantum field confined in the region between two large plates, and in the context of HL formalism, has been developed in [32] and [33]. In these papers it was imposed on the field Dirichlet, Newman and mixed boundary conditions at the plates. Considering scalar massive quantum field, this investigation was carried out in [34], representing an important extension of previous analysis. The analysis of Casimir effects associated with massless fermionic quantum field, has been developed in [24]. For the later, the MIT bag boundary condition was imposed on the field on the two large and parallel plates. For both case, scalar and fermionic quantum fields, it was observed that a non-vanishing Casimir energy takes place, only when the critical exponent associated with the space and time anisotropy, is a odd number. The objective of this paper is to extend the analysis of fermionic Casimir effect in the context of HL scenario, considering at this time massive field. As will see, a non-vanishing mass represents a more technical extension than considered in [24].

This paper is structured as follows. In Section II, we discuss the modified Dirac Lagrangian within the framework of the Horava-Lifshitz approach, considering only the case where the parameter  $\xi$ , associated with Lorentz symmetry breaking, called the critical exponent, is an odd number. We explicitly present solutions for massive fields confined in

the region between two large and parallel plates. Also in this section, we discuss how it is possible to formulate a modified MIT bag model compatible with the higher-order spatial derivative in the Dirac equation. In Section III, we present the formal expression for the vacuum energy. By using the generalized Abel-Plana summation formula, we provide an integral expression for renormalized Casimir energy. Considering specific values of the parameter  $\xi$ , we present asymptotic results for this observable for large and small values of mass. In addition, we present plots that exhibit the behaviors of the Casimir energy with the mass of the field. The most relevant results are summarized and discussed in section IV. In the Appendices A and B, we provide some important calculations carried out during the development of the work. Here, we adopt natural units  $\hbar = c = 1$ , and the metric signature convention  $(+, -, -, -)$ .

## II. THE HORAVA-LIFSHITZ MODEL

In this section, we introduce the theoretical model in which the Dirac equation is modified in the context of the Horava-Lifshitz (HL) approach. Our goal is to obtain solutions for this equation that are compatible with the MIT bag model boundary condition, applied to the fields on two parallel plates with areas  $L^2$ , separated by a distance  $a$ , being  $a \ll L$ , as shown in Fig. 1.




Figure 1: Two large parallel plates with area  $L^2$  separated by a distance  $a$ .

In the HL context, the Lagrangian density adopted to describe the dynamics of a massive

fermionic field, as presented in [35], is given by:

$$\mathcal{L} = \bar{\psi} \left\{ i\gamma^0 \partial_t + l^{\xi-1} (i\gamma^i \partial_i)^\xi - m \right\} \psi. \quad (2)$$

Here, the parameter  $l$  has the dimension of length and is introduced to provide the correct dimension for the Lagrangian above. The exponent  $\xi$ , known as the critical exponent, is associated with the violation of Lorentz symmetry. Note that for  $\xi = 1$ , we recover the usual Dirac Lagrangian without Lorentz violation.

From this point on, we will assume that the critical exponent  $\xi$  takes only odd integer values. This choice is based on results obtained in [24], where it was demonstrated that for even values of  $\xi$ , the Casimir energy is zero. Thus, the modified Dirac equation takes the following form:

$$\hat{H}\psi = \left[ -l^{\xi-1} \gamma^0 (\nabla^2)^{\frac{\xi-1}{2}} (i\gamma^i \partial_i) + m\gamma^0 \right] \psi = i \frac{\partial \psi}{\partial t}, \quad (3)$$

where  $\gamma^j$  given below, represents the Dirac matrices:

$$\gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \quad \text{and} \quad \gamma^j = \begin{pmatrix} 0 & \sigma^j \\ -\sigma^j & 0 \end{pmatrix}. \quad (4)$$

In the definition above,  $I$  represents the  $2 \times 2$  identity matrix, and  $\sigma^j$ ,  $j = 1, 2, 3$  the Pauli matrices.

As mentioned previously, our focus is on calculating the Casimir energy and, consequently, the Casimir pressure associated with the massive fermionic quantum field in the Horava-Lifshitz framework. To achieve this objective, we consider that the fields are confined between two parallel plates as illustrated in Fig. 1, considering  $L \gg a$ . Therefore, assuming the time dependence of the fermionic wave function in the form  $e^{\mp\omega t}$ , we obtain the positive and negative energy solutions for Eq. (3), given by:

$$\psi_\beta^{(+)}(x) = A_\beta e^{-i\omega_\beta t} \begin{pmatrix} \varphi^{(+)}(\vec{x}) \\ \chi^{(+)}(\vec{x}) \end{pmatrix} \quad \text{with} \quad \chi^{(+)}(\vec{x}) = -\frac{il^{\xi-1}(\nabla^2)^{\frac{\xi-1}{2}}}{\omega + m} \sigma^j \partial_j \varphi^{(+)}(\vec{x}) \quad (5)$$

and

$$\psi_\beta^{(-)}(x) = A_\beta e^{i\omega_\beta t} \begin{pmatrix} \varphi^{(-)}(\vec{r}) \\ \chi^{(-)}(\vec{x}) \end{pmatrix} \quad \text{with} \quad \varphi^{(-)}(\vec{r}) = \frac{il^{\xi-1}(\nabla^2)^{\frac{\xi-1}{2}}}{\omega + m} \sigma^j \partial_j \chi^{(-)}(\vec{x}), \quad (6)$$

where  $\omega_\beta$  is the dispersion relation, given by

$$\omega_\beta = l^{\xi-1} \sqrt{(\vec{k}^2)^\xi + \mu^{2\xi}}. \quad (7)$$

Here, the parameter  $\beta$  represents the momentum components  $k_x, k_y, k_z$ . Furthermore, note that in Eq. (7) we have replaced the mass parameter,  $m$ , with  $m = \mu^\xi l^{\xi-1}$ .

The Eqs. (5) and (6) represent two-component spinors,  $\varphi$  and  $\chi$ , where  $\psi_\beta^{(+)}$  and  $\psi_\beta^{(-)}$  correspond to particles and antiparticles, respectively. The components of the spinors have the following explicit form:

$$\varphi^{(+)}(\vec{x}) = e^{i(k_x x + k_y y)} \left( \varphi_+ e^{ik_z z} + \varphi_- e^{-ik_z z} \right), \quad \chi^{(-)}(\vec{x}) = e^{-i(k_x x + k_y y)} \left( \chi_+ e^{ik_z z} + \chi_- e^{-ik_z z} \right). \quad (8)$$

As said before, the fermionic modes are characterized by the set of quantum numbers  $k_x, k_y, k_z$ , with  $k_x$  and  $k_y$  being continuous quantum numbers; however,  $k_z$  must be discrete to satisfy the boundary conditions on the plates. Therefore, assuming the standard approach to derive the MIT bag boundary condition, and considering the modified Lagrangian from Eq. (2), we obtain:

$$\left( 1 + i l^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^\mu n_\mu \right) \psi \Big|_{z=0,a} = 0, \quad (9)$$

with  $n_\mu$  being the unit vector normal directed outward from the boundaries. For the plate configuration we are considering, we have  $n_\mu = -\delta_\mu^z$  for the plate at  $z = 0$  and  $n_\mu = \delta_\mu^z$  at  $z = a$ .

Now, let's analyze the solutions on the plates to find a transcendental relationship. Considering the positive-energy solution from Eq. (5), on the surface  $z = a$  with  $\vec{n} = -\hat{z}$ , the Eq. (9) provides:

$$i l^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \begin{pmatrix} 0 & \sigma^z \\ -\sigma^z & 0 \end{pmatrix} \begin{pmatrix} \varphi \\ \chi \end{pmatrix} = - \begin{pmatrix} \varphi \\ \chi \end{pmatrix}. \quad (10)$$

Considering that  $\nabla^2 \varphi = -\vec{k}^2 \varphi$ , we have:

$$\begin{cases} i^\xi l^{\xi-1} (\vec{k}^2)^{\frac{\xi-1}{2}} \sigma_3 \chi = -\varphi \\ i^\xi l^{\xi-1} (\vec{k}^2)^{\frac{\xi-1}{2}} \sigma_3 \varphi = \chi. \end{cases} \quad (11)$$

Now, substituting  $\chi(\vec{x})$  given in (5) into (11), we notice that the second equation is not satisfied for arbitrary values of the critical exponent  $\xi \neq 1$ . However, if we are inclined to accept that the modified Dirac equation can confine electrons in a finite region, we must appropriately modify the MIT bag model so that for  $\xi = 1$ , we recover the standard MIT model. In Appendix A we propose a higher-order derivative Lagrangian associated with

the MIT bag model that solves this problem. Accepting this modified model, the equations relating the two-component spinors  $\varphi$  and  $\chi$  on the two flat boundaries, coincide with those corresponding to  $\xi = 1$ , as shown in Eqs. (A9) and (A10) on the plate  $z = a$ , for positive and negative energy solutions, respectively. They are given by:

$$\sigma^3 \varphi^{(+)} = \frac{\sigma^i \partial_i \varphi^{(+)}}{w + m} \quad \text{and} \quad \sigma^3 \chi^{(-)} = -\frac{\sigma^i \partial_i \chi^{(-)}}{w + m}. \quad (12)$$

From the boundary condition on the plate at  $z = 0$ , we obtain the following relations between the spinors given in (8):

$$\begin{aligned} \varphi_+ &= -\frac{m(\omega + m) + k_z^2 - k_z \sigma^3 (\sigma^1 k_x + \sigma^2 k_y)}{(m - ik_z)(\omega + m)} \varphi_-, \\ \chi_- &= -\frac{m(\omega + m) + k_z^2 - k_z \sigma^3 (\sigma^1 k_x + \sigma^2 k_y)}{(m + ik_z)(\omega + m)} \chi_+. \end{aligned} \quad (13)$$

Finally, considering the boundary condition for the fermionic functions at  $z = a$ , we arrive at the following transcendental equation:

$$\frac{am}{k_z a} \sin(k_z a) + \cos(k_z a) = 0. \quad (14)$$

We just want to emphasize that the above equation coincides with the standard equation for the Lorentz preserving Dirac equation.

### III. THE INFLUENCE OF LORENTZ SYMMETRY BREAKING IN CASIMIR EFFECT

In this section, we will focus on calculating the Casimir energy associated with the quantum fermionic vacuum fluctuations considering the Horava-Lifshitz Lorentz violation scenario. To this objective, we adopt the formalism described in [36] and present the explicit expansion of the field operator in terms of the normalized positive and negative energy fermionic modes:

$$\psi(x) = \sum_{r,n} \int dk_x \int dk_y \left[ a_{r,\beta} \psi_{r,\beta}^{(+)} + b_{r,\beta}^\dagger \psi_{r,\beta}^{(-)} \right], \quad (15)$$

where  $a_{r,\beta}$  and  $b_{r,\beta}^\dagger$  are the annihilation and creation operators for particles and antiparticles, respectively, with momentum  $\vec{k}$  and polarization  $r$ .

The standard Dirac Lagrangian provides the following Hamiltonian density operator, as demonstrated in [37]:

$$\mathcal{H} = i\psi^\dagger \dot{\psi}. \quad (16)$$

Following a similar procedure, we verify that the modified Dirac Lagrangian in Eq. (2) yields the same expression for the Hamiltonian density.

Therefore, the vacuum energy,  $E_0$ , is determined by calculating the expected value of the Hamiltonian operator from (16) in the vacuum state, as shown below:

$$E_0 = i \int_V d^3x \langle 0 | \psi^\dagger \dot{\psi} | 0 \rangle . \quad (17)$$

Substituting the field operator given in (15) into (17), we obtain:

$$E_0 = \sum_r \sum_n \int d^2\vec{k} \omega_\beta \langle 0 | a_{r,\beta}^\dagger a_{r,\beta} - b_{r,\beta} b_{r,\beta}^\dagger | 0 \rangle . \quad (18)$$

Furthermore, by using the standard anti-commutation relations for the operators  $a_{r,\beta}$  and  $b_{r,\beta}$ , the above expression provides

$$E_0 = -\frac{L^2 l^{\xi-1}}{4\pi^2} \sum_n \int d^2\vec{k} \sqrt{(\vec{k}^2)^\xi + \mu^{2\xi}} . \quad (19)$$

Here, we have  $\vec{k}^2 = k_x^2 + k_y^2 + k_z^2$ . Expressing  $k_x$  and  $k_y$  in polar coordinates, the above expression reads,

$$E_0 = -\frac{L^2 l^{\xi-1}}{\pi} \int_0^\infty k dk \sum_{n=1}^\infty \sqrt{[k^2 + (\alpha_n/a)^2]^\xi + \mu^{2\xi}} , \quad (20)$$

where we have defined  $\alpha_n = k_n a$ , with  $k_n = k_z$  being the  $n$ -th root of the transcendental equation given in (14). As we can see, the vacuum energy above is divergent. Therefore, in order to obtain a finite and well-defined result, namely the Casimir energy, we have to renormalize this expression. This can be done by employing the Abel-Plana summation formula [38]:

$$\sum_{n=1}^\infty \frac{\pi f(\alpha_n)}{1 - \frac{\sin(2\alpha_n)}{2\alpha_n}} = -\frac{\pi a m f(0)}{2(am+1)} + \int_0^\infty dz f(z) - i \int_0^\infty dt \frac{f(it) - f(-it)}{\frac{t+am}{t-am} e^{2t} + 1} . \quad (21)$$

To develop the summation over  $n$  in (20) we must rewrite the integrand in an appropriate manner. At this point we follow the steps adopted in [39]. So let us start with the denominator appearing in the left hand side of the summation (21). By using (14), it is possible to obtain the identity below:

$$1 - \frac{\sin(2\alpha_n)}{2\alpha_n} = 1 + \frac{am}{(am)^2 + \alpha_n^2} . \quad (22)$$

The function  $f(x)$  in Eq. (21) is defined as

$$f(x) = \sqrt{(x^2 + k^2 a^2)^\xi + \mu^{2\xi} a^{2\xi}} \left( 1 + \frac{am}{(am)^2 + x^2} \right) . \quad (23)$$



Thus, substituting (21) into (20), the vacuum energy becomes,

$$E_0 = -\frac{L^2 l^{\xi-1}}{\pi^2 a^\xi} \int_0^\infty k dk \left\{ -\frac{\pi a m}{2(am+1)} f(0) + \int_0^\infty f(z) dz - i \int_0^\infty \frac{f(it) - f(-it)}{\frac{t+am}{t-am} e^{2t} + 1} dt \right\}. \quad (24)$$

Note that the equation above involves the mass  $m$  and also the quantity  $\mu$ , which is related to the mass. Although being possible to use just one of them, here we will keep both notations. In the final expressions, we will express  $\mu$  in terms of the mass  $m$ .

The first term inside the brackets in (24) corresponds to the vacuum energy in the presence of only one plate, while the second term is related to the vacuum energy in the absence of plates. Therefore, these terms are divergent, but after renormalization, they are discarded. Consequently, the Casimir energy is associated with the third term, and is given by:

$$E_C = \frac{iL^2 l^{\xi-1} a}{\pi^2} \int_0^\infty k dk \int_0^\infty \frac{u-m}{(u+m)e^{2au} + u-m} \left(1 + \frac{m}{am^2 - au^2}\right) \times \left\{ \sqrt{[(iu)^2 + k^2]^\xi + \mu^{2\xi}} - \sqrt{[(-iu)^2 + k^2]^\xi + \mu^{2\xi}} \right\} du, \quad (25)$$

where we have performed the substitution  $t = au$ . At this point, we should divide the integral with respect to the variable  $u$  into two intervals: the first is  $[0, \sqrt{k^2 + \mu^2}]$  and the second is  $[\sqrt{k^2 + \mu^2}, \infty)$ . However, we find that the integral in the first interval vanishes, and, consequently, the resulting Casimir energy is given by,

$$E_C = -\frac{2L^2 l^{\xi-1}}{\pi^2} \sin\left(\frac{\pi\xi}{2}\right) \int_0^\infty k dk \int_{\sqrt{k^2 + \mu^2}}^\infty \frac{\left[a(u-m) - \frac{m}{u+m}\right] \sqrt{(u^2 - k^2)^\xi - \mu^{2\xi}}}{(u+m)e^{2au} + u-m} du. \quad (26)$$

In order to obtain a more suitable expression that allows us to analyze asymptotic limits for the dimensionless quantity  $am$ , we will adopt the following change of coordinates (see Appendix B),  $x = \sqrt{u^2 - k^2 - \mu^2}$ . Expressing  $x = r \cos \theta$  and  $k = r \sin \theta$ , the Casimir energy can be written as:

$$E_C = -\frac{mL^2}{\pi^2} \sin\left(\frac{\pi\xi}{2}\right) \int_0^\infty r^3 \frac{g(\sqrt{r^2 + \mu^2})}{\sqrt{r^2 + \mu^2}} dr \int_0^1 \sqrt{(r^2 v / \mu^2 + 1)^\xi - 1} dv, \quad (27)$$

where

$$g(y) = \frac{a(y-m) - \frac{m}{y+m}}{(y+m)e^{2ay} + y-m}. \quad (28)$$

Finally, adopting the variable change  $z = ar$ , we obtain

$$E_C = -\frac{L^2 m}{\pi^2 a^2} \sin\left(\frac{\pi\xi}{2}\right) \int_0^\infty \frac{\sqrt{\mu^2 a^2 + z^2} - ma - \frac{ma}{\sqrt{\mu^2 a^2 + z^2 + ma}}}{\left(\sqrt{\mu^2 a^2 + z^2} + ma\right) e^{2\sqrt{\mu^2 a^2 + z^2}} + \sqrt{\mu^2 a^2 + z^2} - ma} \times \frac{z^3 dz}{\sqrt{z^2 + \mu^2 a^2}} \int_0^1 \sqrt{(z^2 v / (\mu a)^2 + 1)^\xi - 1} dv. \quad (29)$$

As mentioned earlier, the final results will be expressed in terms of the mass  $m$ . For this, we consider  $a\mu = (am)^{1/\xi}\rho_\xi$ , where  $\rho_\xi = (a/l)^{1-1/\xi}$ . Because the parameter  $l$  is of the order of the inverse of Planck energy scale, the parameter  $\rho_\xi$  is much greater than unity.

Unfortunately, for  $m \neq 0$  it is not possible to perform the integral in (29), even for  $\xi = 1$ . Note that, for  $\xi = 1$ , we recover the Casimir energy without the influence of Lorentz symmetry violation. In this case, the result is consistent with the Casimir energy obtained in [21] when Lorentz violation is neglected. Moreover, for  $\xi > 1$ , Eq. (29) presents an additional difficulty, that is an extra integral in the variable  $v$ . However, for massless fermionic fields the integral in (29) can be exactly solved, and the result agrees with the result found in [24]. Therefore, to provide a better understanding about the behavior of Casimir energy for massive fields with Lorentz violation, we present in Fig. 2 three plots of the Casimir energy for  $\xi = 1, 3, 5$ . For these three graphs, we consider only as an illustrative example  $\frac{a}{l} = 10^2$ . From them we can observe a change in sign and intensity that is strongly dependent on  $\xi$ . For  $\xi = 1, 5$  the forces between the plates are attractive, being repulsive for  $\xi = 3$ , and the decay becomes more pronounced with increasing of  $\xi$ .

As mentioned earlier, we cannot obtain a closed expression for the integrals in Eq. (29), i.e., to express it in terms of spacial functions. However, by this result it is possible to explore some important behaviors. These limiting cases can be interpreted as corresponding to small mass or distances between plates, and large mass or distances between them, respectively. Let's analyze each case separately:

- However, before to develop these behaviors, let us obtain the Casimir energy for the case of massless fermions. After some intermediate steps, from (29), we get:


$$E_C = -\frac{L^2 l^{\xi-1} 2^{-2(\xi+2)} (2^{\xi+2} - 1) \Gamma(2 + \xi) \zeta(3 + \xi) \sin(\pi\xi/2)}{\pi^2 a^{\xi+2}}. \quad (30)$$

Consequently, the Casimir pressure is expressed as:


$$P_C = \frac{F_C}{L^2} = -\frac{l^{\xi-1} 2^{-2(\xi+2)} (2^{\xi+2} - 1) \Gamma(3 + \xi) \zeta(3 + \xi) \sin(\pi\xi/2)}{\pi^2 a^{\xi+3}}. \quad (31)$$

Both expressions above coincide with obtained previously in [24].


Now, let's consider the limiting cases when  $am \ll 1$  and  $am \gg 1$ . For  $am \ll 1$ , the obtained result for any values of  $\xi$ , is a long expression and not very enlightening. Therefore, we consider only the cases exhibited in Fig. 2, namely, the cases  $\xi = 1, 3, 5$ .



(a) For  $\xi = 1$ .



(b) For  $\xi = 3$ .



(c) For  $\xi = 5$ .

Figure 2: The behavior of the Casimir energy, Eq. (29), in units of  $L^2/a^3$ , as function of  $am$ , considering different values of  $\xi$ , as  $\xi = 1$  (a),  $\xi = 3$  (b), and  $\xi = 5$  (c). Thus, Lorentz symmetry violation affects the Casimir energy both in intensity and in sign.

- For the case where  $am \ll 1$ , we have:

- For  $\xi = 1$ , the integral in  $v$  in Eq. (29) gives us:

$$\int_0^1 \sqrt{(z^2 v / (\mu a)^2 + 1)^\xi - 1} dv = \frac{2z}{3am}. \quad (32)$$

Now, substituting this result into (29), the integral over  $z$  can be approximated evaluated providing for the Casimir energy the approximated result:

$$E_C \approx -\frac{L^2}{\pi^2 a^3} \left( \frac{7\pi^4}{2880} - \frac{\pi^2 am}{24} \right). \quad (33)$$

As to the the Casimir pressure is get:

$$P_C \approx -\frac{7\pi^2}{960a^4} \left( 1 - \frac{80am}{7\pi^2} \right). \quad (34)$$

As expected, this result agrees with that obtained in [21] for the case where Lorentz symmetry is not violated.

- For  $\xi = 3$ , the integral in  $v$  is

$$\begin{aligned} \int_0^1 \sqrt{(z^2 v / (\mu a)^2 + 1)^\xi - 1} dv &= \frac{2(a^2 \mu^2 + z^2) \sqrt{3a^4 \mu^4 + 3a^2 \mu^2 z^2 + z^4}}{5a^3 \mu^3 z} \\ &+ \frac{3^{3/4} a^2 \mu^2 F \left( \frac{2\sqrt[4]{3} a \mu z}{\sqrt{3a^2 \mu^2 + z^2}}, \frac{1}{\sqrt{2}(1+\sqrt{3})} \right)}{5z^2} - \frac{2 \cdot 3^{3/4} a^2 \mu^2 K \left[ \frac{1}{\sqrt{2}(1+\sqrt{3})} \right]}{5z^2}. \end{aligned} \quad (35)$$

Where the special functions  $F$  and  $K$  represent the elliptic integral of the first and second kinds, respectively [40]. Thus, substituting the above expression into (29), and considering  $a\mu = \sqrt[3]{ma}(a/l)^{2/3}$ , the Casimir energy reads,

$$E_C \approx \frac{L^2}{\pi^2 a^3} \left( \frac{l}{a} \right)^2 \left( \frac{31\pi^6}{40320} - \frac{7\pi^4 am}{960} \right). \quad (36)$$

Therefore, the Casimir pressure is

$$P_C \approx \left( \frac{l}{a} \right)^2 \left( \frac{31\pi^4}{8064a^4} - \frac{7\pi^2 am}{240a^4} \right). \quad (37)$$

- Finally, for  $\xi = 5$ , the integral in  $v$  is given by

$$\begin{aligned} \int_0^1 \sqrt{(z^2 v / (\mu a)^2 + 1)^\xi - 1} &= -\frac{(\sqrt{5} - 1)\pi^{3/2} a^2 \mu^2}{7z^2 \Gamma(7/10) \Gamma(4/5)} \\ &+ \frac{2(a^2 \mu^2 + z^2)^{7/2} {}_2F_1 \left( -\frac{7}{10}, -\frac{1}{2}; \frac{3}{10}; \frac{a^{10} \mu^{10}}{(a^2 \mu^2 + z^2)^5} \right)}{7a^5 \mu^5 z^2}. \end{aligned} \quad (38)$$

In the above expression,  ${}_2F_1$  represents the hypergeometric function [40].

For this case, an approximated expression can be obtained for the integral with respect to  $z$  in (29), consequently, in this approximation, the Casimir energy is expressed by

$$E_C \approx -\frac{L^2}{\pi^2 a^3} \left(\frac{l}{a}\right)^4 \left(\frac{127\pi^8}{215040} - \frac{31\pi^6 am}{8064}\right), \quad (39)$$

and the pressure by

$$P_C \approx -\left(\frac{l}{a}\right)^4 \left(\frac{127\pi^6}{30720a^4} - \frac{31\pi^4 am}{1344a^4}\right). \quad (40)$$

Thus, we can observe that the critical exponent significantly impacts the Casimir energy and, consequently, the Casimir pressure. In the case where  $\xi = 1$ , i.e., the scenario without Lorentz violation, the standard result is obtained, as expected. However, the new results for  $\xi > 1$  can provide a fresh perspective on the study of potential Lorentz violation.

- Now, for the case where  $am \gg 1$ , the integral with respect to  $v$  can be approximated by:

$$\int_0^1 \sqrt{(z^2 v / (\mu a)^2 + 1)^\xi - 1} dv \approx \frac{2\sqrt{\xi} z}{3a\mu}. \quad (41)$$

Thus, the integration with respect to  $z$  in (29) yields:

$$I_\xi \approx (am)^{\frac{1}{\xi}} (a/l)^{1-\frac{1}{\xi}} e^{-2(am)^{\frac{1}{\xi}} (a/l)^{1-\frac{1}{\xi}}}. \quad (42)$$

Substituting this result back into (29), we get

$$E_C \approx -\frac{L^2 (am)^{\frac{1}{\xi}} (a/l)^{1-\frac{1}{\xi}} \sin(\pi\xi/2)}{\pi^2 a^3} e^{-2(am)^{\frac{1}{\xi}} (a/l)^{1-\frac{1}{\xi}}}. \quad (43)$$

Here we can see that the Casimir energy decays exponentially with  $(am)^{\frac{1}{\xi}} (a/l)^{1-\frac{1}{\xi}}$ . Therefore, in this limit, the Casimir pressure also quickly approaches zero. This result is consistent with the behavior of the graphs in Fig. 2. Additionally, we can note that in this limit, calculations can proceed for any odd  $\xi$ .

#### IV. CONCLUDING REMARKS

In this paper we have analyzed the fermionic Casimir effects in the HL formalism of Lorentz violation, considering that the field is confined in the region between two large parallel plates. In order to implement this confinement, we had to modify the Lagrangian density associated with MIT bag model, compatible with the higher-order spatial derivative system. This proposed model is explicitly given in Eq. (A1) of the Appendix A. In this way, we found the transcendental equation (14), which discretizes the moment along the direction orthogonal to the plates. By using the generalized Abel-Plana summation formula, Eq. (21), we were able to provide an expression for the Casimir energy in (29), from the general vacuum energy, Eq. (20).

In fact in our analysis we have considered only the cases where the critical exponent,  $\xi$ , is an odd number [41]. As in the case of  $\xi = 1$ , our result to Casimir energy, Eq. (29), cannot be expressed in a closed expression, i.e., in terms of combinations of special functions. Furthermore, for  $\xi > 1$ , this expression presents an additional difficulty represented by an extra integral in the variable  $v$ . In principle it is possible to find a general result for this integral, but it is a long expression and not very enlightening. So, in view of this fact, we decided to present for this integral only results with  $\xi = 1, 3, 5$ , given in (32), (35) and (38), respectively. To provide a better understanding about the behavior of the fermionic Casimir energy, we presented in Fig. 2 three plots that exhibit the behavior of of the Casimir energy for  $\xi = 1, 3, 5$  as function of dimensionless parameter  $am$ . For these three graphs, we consider, only as an illustrative example,  $\frac{a}{l} = 10^2$ . As we can observe the corresponding energies present a strong dependence on the critical exponent. They change their signs and intensities.

Another analysis developed in this paper concerns the behavior of the Casimir energy and pressure in the limiting cases of the dimensionless parameter  $am \ll 1$  and  $am \gg 1$ . For  $am \ll 1$ , we present the asymptotic expressions for the Casimir energy and pressure, considering  $\xi = 1, 3, 5$ . These expressions are given by, (33) and (34), (36) and (37), and finally (39) and (40), respectively. In the limiting case of large  $am$ , we were able to present the behavior of the Casimir energy in a compact form, for any value of odd  $\xi$ . This expression was given in (43), where we can see a strong exponential decay, accentuated when  $\xi > 1$ .

Therefore, we hope that the results presented here can guide theoretical and experimental studies in exploring aspects related to Lorentz symmetry violation. This will enable us to further improve our theoretical models, including the one in this current work, as well as our experimental framework.

We would like to finish this paper by saying that in previous work [21], we have analyzed the Casimir energy in a CPT-even, aether-like Lorentz symmetry violation. In this context the Lorentz symmetry is broken in the Planck scale energy, and this violation is implemented by the emergence of a nonzero vacuum expectation value of some vector and tensor components, which implies in preferential directions, therefore, space-time anisotropy [28]. Because in this present analysis we have analyzed the fermionic Casimir effects considering HL Lorentz violation scenario, we hope to have filled the gap left.

### Acknowledgments

We would like to thank Aram A. Saharian for helpful discussions.

### Appendix A: MIT bag model in the Horava-Lifshitz scenario

In this appendix, we introduce a modified Lagrangian in the Horava-Lifshitz Lorentz violation scenario for the MIT bag model, which provides consistent equations for the two-component spinors  $\varphi$  and  $\chi$ . We impose that the positive and negative energy solutions, given in Eqs. (5) and (6), satisfy the MIT boundary condition on the plates located at  $z = 0$  and  $z = a$ , as shown in Fig. 1. The proposed Lagrangian density given below, is very similar with the standard MIT model:

$$\begin{aligned} \mathcal{L} = & \left\{ \frac{i}{2} \left[ \bar{\Psi} \gamma^0 (\partial_0 \Psi) - (\partial_0 \bar{\Psi}) \gamma^0 \Psi \right] + \frac{i l^{\xi-1}}{2} \left[ \bar{\Psi} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^i (\partial_i \Psi) - (\partial_i \bar{\Psi}) (\nabla^2)^{\frac{\xi-1}{2}} \gamma^i \Psi \right] \right. \\ & \left. - m \bar{\Psi} \Psi - B \right\} \theta_v - \frac{1}{2} \Delta_s \bar{\Psi} M \Psi, \end{aligned} \quad (\text{A1})$$

where  $M$  is a specific  $4 \times 4$  matrix operator, as indicated below,

$$M = \begin{pmatrix} \hat{\alpha} & 0 \\ 0 & \hat{\beta} \end{pmatrix}. \quad (\text{A2})$$

Here,  $\hat{\alpha}$  and  $\hat{\beta}$  are higher-order differential operators for  $\xi > 1$ . Both operators reduce to the  $2 \times 2$  identity matrix when  $\xi = 1$ . Furthermore, for positive-energy solutions, we impose

$\hat{\alpha} = l^{2(\xi-1)}(\nabla^2)^{\xi-1}$  and  $\hat{\beta} = 1$ , and for negative-energy solutions, we impose  $\hat{\alpha} = 1$  and  $\hat{\beta} = l^{2(\xi-1)}(\nabla^2)^{\xi-1}$ .

In Eq. (A1),  $\theta_v$  is the Heaviside function, which is equal to unity inside the bag and vanishes outside it. The factor  $B$  is related to the energy-momentum tensor on the bag, and  $\Delta_s$  is the Dirac delta function given by the spatial derivative of  $\theta_v$ , as shown below,

$$\frac{\partial \theta_v}{\partial x^i} = n_i \Delta_s, \quad (\text{A3})$$

where  $n_i$  is the unit vector normal to the surface.

Therefore, using the usual Euler-Lagrange formalism, the dynamics of the system are governed by the modified Dirac equation:

$$\frac{\partial \mathcal{L}}{\partial \bar{\Psi}} - \partial_0 \left[ \frac{\partial \mathcal{L}}{\partial (\partial_0 \bar{\Psi})} \right] - \partial_i \left[ \frac{\partial \mathcal{L}}{\partial (\partial_i \bar{\Psi})} \right] = 0. \quad (\text{A4})$$

Therefore, substituting Eq. (A1) into the equation above, we obtain

$$\left[ i\gamma^0 (\partial_0 \Psi) + il^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^i \partial_i \Psi - m\Psi \right] \theta_v - \frac{i}{2} \left[ l^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^\mu n_\mu \Psi - iM\Psi \right] \Delta_s = 0. \quad (\text{A5})$$

From Eq. (A5), we can derive two different equations: (i) In the region inside the bag, where  $\Delta_s = 0$  and  $\theta_v = 1$ , we have

$$i\gamma^0 (\partial_0 \Psi) + il^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^i \partial_i \Psi - m\Psi = 0. \quad (\text{A6})$$

(ii) On the surface of the bag, where  $\Delta_s = \infty$  and  $\theta_v = 0$ , we obtain

$$l^{\xi-1} (\nabla^2)^{\frac{\xi-1}{2}} \gamma^\mu n_\mu \Psi = iM\Psi, \quad (\text{A7})$$

$$\begin{cases} i^\xi l^{\xi-1} (\vec{k}^2)^{\frac{\xi-1}{2}} \sigma_3 \chi = -\hat{\alpha} \varphi \\ i^\xi l^{\xi-1} (\vec{k}^2)^{\frac{\xi-1}{2}} \sigma_3 \varphi = \hat{\beta} \chi. \end{cases} \quad (\text{A8})$$

Therefore, considering the positive and negative-energy solutions on the plate at  $z = a$ , both equations above are consistent, resulting:

$$\sigma^3 \varphi^{(+)} = -\frac{\sigma^i \partial_i \varphi^{(+)}}{w + m}, \quad (\text{A9})$$

and

$$\sigma^3 \chi^{(-)} = -\frac{\sigma^i \partial_i \chi^{(-)}}{w + m}. \quad (\text{A10})$$



## Appendix B: Some mathematical manipulation

In this appendix we present the procedure adopted to express the Casimir energy, given in (26), in the more workable expression, Eq. (29). Let's consider the integral in the following form:

$$\int_0^\infty k dk \int_{\sqrt{k^2+\mu^2}}^\infty f(u) \sqrt{(u^2 - k^2)^\xi - \mu^{2\xi}} du. \quad (\text{B1})$$

The first step will be to perform a change of coordinates,  $x = \sqrt{u^2 - k^2 - \mu^2}$ . Thus, the integral above can be rewritten as:

$$\int_0^\infty k dk \int_0^\infty x \frac{f(\sqrt{x^2 + k^2 + \mu^2})}{\sqrt{x^2 + k^2 + \mu^2}} \sqrt{(x^2 + \mu^2)^\xi - \mu^{2\xi}} dx. \quad (\text{B2})$$

Next, we should adopt a transformation to polar coordinates,  $x = r \cos \theta$  and  $k = r \sin \theta$ , which gives us:

$$\int_0^\infty r^3 \frac{f(\sqrt{r^2 + \mu^2})}{\sqrt{r^2 + \mu^2}} dr \int_0^{\pi/2} \sin \theta \cos \theta \sqrt{(r^2 \cos^2 \theta + \mu^2)^\xi - \mu^{2\xi}} d\theta. \quad (\text{B3})$$

Finally, considering  $w = \cos \theta$  and then  $t = w^2$ , we can represent the integral (B1) as:

$$\frac{\mu^\xi}{2} \int_0^\infty r^3 \frac{f(\sqrt{r^2 + \mu^2})}{\sqrt{r^2 + \mu^2}} dr \int_0^1 \sqrt{(r^2 t / \mu^2 + 1)^\xi - 1} dt. \quad (\text{B4})$$

- [1] H. B. G. Casimir, *Indagat. Math* **10**, 261 (1948); *Kon. Ned. Akad. Wetensch. Proc.* **51**, 793 (1948); *Front. Phys.* **65**, 342 (1987); *Kon. Ned. Akad. Wetensch. Proc.* **100**, N3-4, 61 (1997).
- [2] M. J. Sparnaay, *Physica* **24**, 751 (1958).
- [3] P. H. G. M. van Blokland and J. T. G. Overbeek, *J. Chem. Soc. Faraday Trans. I* **74**, 2637 (1978).
- [4] S. K. Lamoreaux, *Phys. Rev. Lett.* **78**, 5 (1997).
- [5] U. Mohideen and A. Roy, *Phys. Rev. Lett.* **81**, 4549 (1998); B. W. Harris, F. Chen, and U. Mohideen, *Phys. Rev. A* **62**, 052109 (2000).
- [6] F. Khelili, *Phys. Rev. D* **85** (2012), 125013.
- [7] A. Erdas and K. P. Seltzer, *Phys. Rev. D* **88** (2013), 105007.
- [8] A. L. C. Rego, B. W. Mintz, C. Farina and D. T. Alves, *Phys. Rev. D* **87** (2013) no.4, 045024.

- [9] A. Erdas and K. P. Seltzer, *Int. J. Mod. Phys. A* **29** (2014), 1450091.
- [10] E. R. B. de Mello and M. B. Cruz, *Int. J. Mod. Phys. A* **38** (2023) no.11, 2350062.
- [11] M. B. Cruz, E. R. Bezerra de Mello, and A. Yu. Petrov, *Phys. Rev. D* **96**, 045019 (2017).
- [12] Š. Vedl, D. J. George and G. K. Brennen, *Phys. Rev. D* **109** (2024) no.1, 016018.
- [13] C. Bórquez and B. Droguett, *Eur. Phys. J. C* **84** (2024) no.1, 62.
- [14] E. Elizalde, F. C. Santos and A. C. Tort, *Int. J. Mod. Phys. A* **18** (2003), 1761-1772. [
- [15] A. Bulgac, P. Magierski and A. Wirzba, *EPL* **72** (2005), 327-333.
- [16] *Annals Phys.* **317** (2005), 220-237 [erratum: *Annals Phys.* **321** (2006), 1274-1275].
- [17] A. A. Saharian, *Class. Quant. Grav.* **25** (2008), 165012.
- [18] L. P. Teo, *Phys. Rev. D* **91** (2015) no.12, 125030.
- [19] L. P. Teo, *Phys. Rev. D* **91** (2015) no.8, 085034.
- [20] A. K. Grigoryan, A. R. Mkrtchyan and A. A. Saharian, *Int. J. Mod. Phys. D* **26** (2017) no.07, 1750064.
- [21] M. B. Cruz, E. R. Bezerra de Mello and A. Y. Petrov, *Phys. Rev. D* **99** (2019), 085012.
- [22] S. Bellucci, I. Brevik, A. A. Saharian and H. G. Sargsyan, *Eur. Phys. J. C* **80** (2020), 281.
- [23] P. J. Steinhardt, *Phys. Rev. D* **21** (1980), 2390.
- [24] D. R. da Silva, M. B. Cruz and E. R. Bezerra de Mello, *Int. J. Mod. Phys. A* **34** (2019) no.20, 1950107.
- [25] M. V. Barnhill, III and A. Halprin, *Phys. Rev. D* **21** (1980), 1916.
- [26] A. Ashtekar, *Loop Quantum Gravity: Four Recent Advances and a Dozen Frequently Asked Questions*. 11th Marcel Grossmann Meeting on General Relativity, 126-147 (2007).
- [27] R. Gambini, S. Rastgoo and J. Pullin, *Class. Quant. Grav.* **28** (2011), 155005.
- [28] V. A. Kostelecky and S. Samuel, *Phys. Rev. D* **39** (1989), 683.
- [29] V. A. Kostelecky and R. Potting, *Phys. Lett. B* **381** (1996), 89-96.
- [30] P. Horava, *Phys. Rev. D* **79** (2009), 084008.
- [31] J. W. Moffat, *Class. Quant. Grav.* **27** (2010), 135016.
- [32] A. F. Ferrari, H. O. Girotti, M. Gomes, A. Yu. Petrov and A. J. da Silva, *Mod. Phys. Lett. A* **28**, 1350052 (2013).
- [33] I. J. Morales Ulion, E. R. Bezerra de Mello and A. Y. Petrov, *Int. J. Mod. Phys. A* **30** 36, 1550220 (2015).
- [34] R. V. Maluf, D. M. Dantas and C. A. S. Almeida, *Eur. Phys. J. C* **80** (2020), 442.

- [35] T. Mariz, J. R. Nascimento and A. Y. Petrov, Phys. Rev. D **101** (2020) no.10, 105008.
- [36] C. Itzykson and J. B. Zuber, *Quantum Field Theory*. International Series In Pure and Applied Physics, New York: McGraw-Hill, 1980.
- [37] F. Mandl, S. Shaw, Quantum Field Theory, second ed., John Wiley & Sons, London (2010).
- [38] A. A. Saharian, PoS IC **2006** (2006) 019 [[hep-th/0609093](#)].
- [39] S. Bellucci, A.A. Saharian, Phys. Rev. D **80**, 105003 (2009).
- [40] *Handbook of Mathematical Functions*, edited by M. Abramowitz and I.A. Stegun (Dover, New York, 1972).
- [41] Considering even values for  $\xi$ , the corresponding Hamiltonian operator becomes diagonal, and spin up and down components of the fermionic field do not interact themselves