Mean-Variance Optimization for Participating Life Insurance Contracts[∗]

Felix Fießinger^{†,‡} and Mitja Stadje[§]

December 20, 2024

Abstract

This paper studies the equity holders' mean-variance optimal portfolio choice problem for (non-)protected participating life insurance contracts. We derive explicit formulas for the optimal terminal wealth and the optimal strategy in the multi-dimensional Black-Scholes model, showing the existence of all necessary parameters. Moreover, we provide a numerical analysis of the Black-Scholes market. The equity holders on average increase their investment into the risky asset in bad economic states and decrease their investment over time.

Keywords: optimal portfolio, portfolio insurance, mean-variance optimization, participating life insurance, non-concave utility maximization

JEL: C61, G11, G22

1 Introduction

This paper investigates a mean-variance optimization from the perspective of the equity holders of an insurance company for the two standard designs of participating life insurance contracts, i.e., with a protected or non-protected guarantee. In a participating life insurance contract, the policyholder gets a (possibly protected) guarantee and participates proportionally at maturity from the portfolio value exceeding a pre-defined threshold higher than the guarantee. The policyholders receive at least the guarantee value at maturity if the guarantee is protected. If the guarantee is not protected, then the policyholders get, at most, the portfolio value where, initially, the insurance company gives additional equity to the premium for the investment, i.e., the equity has only limited liability.

The portfolio theory research goes back to the 1950s with the pioneering work of Markowitz [\[26,](#page-33-0) [27\]](#page-33-1), who used variance to measure the riskiness of stock returns in a one-period setting. Mean-variance was later also analyzed in dynamic settings, see for instance Hakansson [\[18\]](#page-32-0), Samuelson [\[36\]](#page-33-2) or Merton [\[29,](#page-33-3) [30\]](#page-33-4), and has been extended in various directions. For a more detailed overview of mean-variance portfolio optimizations, see the literature review from Zhang et al. [\[38\]](#page-33-5). In these works, the management is typically assumed to maximize the mean-variance of the entire portfolio, and no distinction between equity holders and debt holders or policyholders is made.

This paper aims to provide an explicit formula for the optimal terminal wealth (from the perspective of the equity holders of an insurance company), show that the optimal solution exists (which is non-trivial since the solution includes an additional parameter), and prove an analytical

[∗]Declarations of interest: none †University of Ulm, Institute of Insurance Science and Institute of Mathematical Finance, Faculty of Mathematics and Economics, Ulm, Germany. Email: felix.fiessinger@uni-ulm.de

[‡]Corresponding author §University of Ulm, Institute of Insurance Science and Institute of Mathematical Finance, Faculty of Mathematics and Economics, Ulm, Germany. Email: mitja.stadje@uni-ulm.de

formula for the optimal strategy. Moreover, we discuss the characteristics of the terminal wealth and the optimal strategy in a numerical analysis.

While the optimal investment problem for participating life insurance contracts was solved for expected utility (EU), an analysis for mean-variance is lacking, which is the contribution of our paper. This analysis is relevant as mean-variance is widely spread in the industry, and most finance papers and textbooks use it as the benchmark to measure risk; see, for instance, Cochrane [\[11\]](#page-32-1). The reason is that mean-variance provides a straightforward interpretation of risk vs. reward, admits tractable statistical properties, and also, due to historical contingencies, has become the leading standard taught in business schools, making it easier for a risk manager to justify its use compared to specifying a utility function. Finally, due to the nonlinearity of variance, the mathematical analysis differs from EU. In particular, one needs to specify an equivalent problem which, contrary to EU-maximization, adds an additional parameter λ to the model. The existence of such a λ is not apparent, and we are only able to derive its existence and compute it in a semi-explicit way in the case of the Black-Scholes model.

As discussed in the beginning, the insurance company offering a participating life insurance contract pays back a surplus to the policyholder in good economic states. Insurance policies with profit participation play an essential role in the life insurance sector. According to the European Insurance Overview 2023 [\[16\]](#page-32-2), issued by the European Insurance and Occupational Pensions Authority (EIOPA), policyholders spent 2022 around a quarter of their gross premiums on profit participation insurance policies in the life sector (includes life, health, and pension insurance). In Croatia, Italy, and Belgium, these policies have a market share of over 50 % in the life sector. Several publications are concerned with the valuation and hedging of such insurance policies, e.g., Bryis and de Varenne [\[6\]](#page-32-3), Bacinello and Persson [\[2\]](#page-32-4), Gatzert and Kling [\[17\]](#page-32-5), Schmeiser and Wagner [\[37\]](#page-33-6) or Mirza and Wagner [\[32\]](#page-33-7). In recent years, the study of optimal investments for participating life insurance contracts in continuous time started. Lin et al. [\[25\]](#page-33-8) analyzed 2017 the optimal investment for an EU-problem for a specific S-shaped utility function. Afterwards, Nguyen and Stadje [\[33\]](#page-33-9) examined a similar setting for more general S-shaped utility functions under a Value at Risk constraint and mortality risk. He et al. [\[19\]](#page-32-6) made another generalization by considering a weighted utility function between the insurer and the policyholders. Moreover, Dong et al. [\[15\]](#page-32-7) added to the EU-problem a Value at Risk and a portfolio insurance constraint. Chen et al. [\[8\]](#page-32-8) considered Value at Risk, expected shortfall, and Average Value at Risk constraints. To the best of our knowledge, we are the first to consider participating insurance contracts in continuous time under mean-variance optimization.

A significant factor of an optimal policy design is the payoff structure: For a participating life insurance contract, the payoff has at least one point where it is non-differentiable, e.g., at the threshold where the proportional surplus participation starts. (The payoff of the non-protected product also has a second point of non-differentiability at the guarantee value.) At these points, the payoff changes its slope, resulting in a payoff that is generally neither convex nor concave (see Figure [2](#page-4-0) for an illustration of the insurer's payoff). In such situations, one often uses concavification techniques; see, for instance, Larsen [\[22\]](#page-32-9) or Reichlin [\[35\]](#page-33-10). Liang et al. [\[24\]](#page-33-11) used a generalization of this method. For alternative approaches in non-concave portfolio optimization, see, for instance, Kraft and Steffensen [\[21\]](#page-32-10), Dai et al. [\[14\]](#page-32-11), or Qian and Yang [\[34\]](#page-33-12). In this work, we use the specific structure of a payoff in a complete market to carefully compare different solution candidates and show that the optimal terminal wealth actually has a closed and, up to the implicit parameters whose existence proof and computation in our setting is delicate, simple form. We also use a standard Lagrangian approach to get the optimal terminal wealth and use the price density process to give the analytic formula for the optimal strategy. This method was used to solve several other optimization problems in complete markets; see, for instance, Basak and Shapiro [\[4\]](#page-32-12), Cuoco et al. [\[13\]](#page-32-13), Chen et al. [\[8\]](#page-32-8), Chen et al. [\[7\]](#page-32-14), Nguyen and Stadje [\[33\]](#page-33-9), Mi et al. [\[31\]](#page-33-13), Chen et al. [\[10\]](#page-32-15), or Avanzi et al. [\[1\]](#page-32-16). To use this Lagrangian approach for mean-variance, we start to give an equivalent problem in the spirit of Zhou and Li [\[39\]](#page-33-14). The ansatz described above then leads to having two multipliers instead of only one Lagrangian multiplier. We obtain a system of non-linear equations with several variables yielding implicit functions whose properties, through various intermediate results, then entail the existence of a solution. Although mean-variance does not respect first-order stochastic dominance, we do get similar results compared to EUoptimization regarding the general form of the optimal solution, but with the terminal wealth having a simpler structure. In particular, in a complete market, the optimal terminal wealth is piecewise linear in the price density. Compared to the papers discussed above, the main difference of our ansatz is to consider an optimization functional, where we include an option-like payoff of the participation instead of adding further conditions to a non-option-like payoff. We show in Proposition [3.6](#page-7-0) that our optimization corresponds to a target problem, where one aims to be as close as possible to certain exogenously given constants. This alternative formulation is close to Avanzi et al. [\[1\]](#page-32-16), who minimize a variance-like function under solvency conditions. However, while they face a 1-target problem our optimization corresponds to a 2-target problem, where the exogenously given target depends on the region of the terminal wealth. An additional constraint ensures that mass can only be shifted from one region to another at a rate which depends on the amount of participation. Target-based optimization is common in the literature as an alternative to mean-variance, see for instance Avanzi et al. [\[1\]](#page-32-16), Menoncin and Vigna [\[28\]](#page-33-15), or Li and Forsyth [\[23\]](#page-32-17). Finally, our numerical results show a somewhat different investment behavior compared to EU-maximization, in particular for S-shaped utility functions. Specifically, we observe that the investment gets more conservative with shorter maturities, and the equity holders on average increase their investment into the risky asset in bad economic states. Moreover, the insurer invests more riskily when offering a non-protected participation life insurance product than when offering a protected one due to a reduced downside potential. Since variance is the most widely used risk measure in industry and the cornerstone of most modern portfolio theory and the finance literature, these results might be potentially relevant from a descriptive point of view.

Section [2](#page-2-0) describes participating life insurance contracts and briefly introduces the functional to optimize. In Section [3,](#page-5-0) we explicitly show the optimal terminal wealth, the optimal strategy, and the existence of the necessary parameters in the Black-Scholes model. Section [4](#page-12-0) analyzes some numerical results for the Black-Scholes model. Finally, Section [5](#page-18-0) concludes the paper.

2 Model Setup

2.1 Participating life insurance contracts

Participating insurance contracts or, in general, insurance contracts with some profit participation play a crucial role in the life sector. Figure [1](#page-3-0) shows that, on average, around a quarter of the gross premiums 2022 in the life sector are spent on policies with profit participation. In some countries, like Croatia, Belgium, or Italy, even more than 50 % of the gross premiums are invested in such policies.

This paper focuses on two standard designs of participating life insurance contracts. Both products offer a guarantee value G and a proportional surplus participation rate α_2 when the portfolio value exceeds a threshold, which we denote by k_2 . Note that k_2 is always higher or equal to G. The difference between the two designs is that the first product offers a non-protected guarantee, whereas the second provides a protected one. Protected means, in this case, that the policyholders get at least their guarantee value, independent of the economic situation at maturity. In contrast, in the non-protected case, the insurance company declares bankruptcy if the portfolio value is below G. In this case, the policyholders only get the portfolio value. For the portfolio, it is essential to note that the initial portfolio value x_0 is the sum of the premiums from the policyholders plus some initial capital from the insurer (i.e., the equity holders). There is no rule on how to set the guarantee value. One possible example of setting G is to take the sum of the premiums in addition to a guaranteed interest rate below the risk-free interest rate.

Figure 1: Market share in 2022 of the gross premium separated by the line of business in the life sector. Data source: European Insurance Overview from the EIOPA [\[16\]](#page-32-2)

Moreover, product designers often set the threshold k_2 as the sum of the premiums divided by the share of the policyholders in the portfolio.

Hence, we conclude the following payoffs V for the policyholders (pol) and the insurer (ins) for the contracts with non-protected (non) resp. protected (pro) guarantees and terminal portfolio value $X_T \geq 0$:

$$
V_{\text{pol}}^{\text{non}}(X_T) = \begin{cases} X_T & \text{if } X_T < G, \\ G & \text{if } G \le X_T < k_2, \\ G + \alpha_2(X_T - k_2) & \text{if } X_T \ge k_2, \end{cases}
$$
\n
$$
V_{\text{pol}}^{\text{pro}}(X_T) = \begin{cases} G & \text{if } X_T < k_2, \\ G + \alpha_2(X_T - k_2) & \text{if } X_T \ge k_2, \end{cases}
$$
\n
$$
V_{\text{ins}}^{\text{non}}(X_T) = \begin{cases} 0 & \text{if } X_T < G, \\ X_T - G & \text{if } G \le X_T < k_2, \\ X_T - G - \alpha_2(X_T - k_2) & \text{if } X_T \ge k_2, \end{cases}
$$
\n
$$
V_{\text{ins}}^{\text{pro}}(X_T) = \begin{cases} X_T - G & \text{if } X_T < k_2, \\ X_T - G - \alpha_2(X_T - k_2) & \text{if } X_T \ge k_2, \\ X_T - G - \alpha_2(X_T - k_2) & \text{if } X_T \ge k_2. \end{cases}
$$

Note that solely $V_{\text{ins}}^{\text{pro}}$ can attain negative values. For a more detailed explanation of participating insurance, we refer to Nguyen and Stadje [\[33\]](#page-33-9).

2.2 Optimization Functional

Before defining the optimization functional, let us introduce the basic financial market, which in Section [3](#page-5-0) and [4](#page-12-0) will become the Black-Scholes model.

Let $(\Omega, \mathcal{F}, (\mathcal{F})_{t\in[0,T]}, \mathbb{P})$ be a filtered probability space with time horizon $T > 0$. The filtration is generated by the d -dimensional Brownian Motion W satisfying the usual conditions. We consider an arbitrage-free market with a risk-free asset B and a deterministic interest rate $r_t \geq 0$ following the price process $dB_t = B_t r_t dt$, and d risky assets S^i , $i \in \{1, ..., d\}$ with adapted price processes. Let the dynamic strategy u represent the fraction of wealth invested in the corresponding risky asset, while the remaining money is invested in the risk-free asset with the corresponding wealth

 $\overline{1}$ For simplicity, we assume that the management or the regulator does not allow the total wealth to become negative. However, if either we extend the following definitions also for $X_T < 0$, or let the insurer cover in both cases all losses stemming from a negative terminal portfolio value, all our results hold (see also Remark [3.3](#page-6-0)[c\)](#page-6-1).

process X_t and initial value $X_0 = x_0$. We denote by U the set of all admissible strategies given by all u, which are progressively measurable and induce an integrable X_T .

Next, we introduce a general functional to optimize, the previously defined participating life insurance products being special cases. We look for the optimal strategy $\hat{u} \in \mathcal{U}$ such that

$$
J(0, T, \hat{u}, x_0) = \sup_{u \in \mathcal{U}} J(0, T, u, x_0), \tag{2.1}
$$

where the value functional J is defined as

$$
J(0,T,u,x_0) := \mathbb{E}[F(0,T,u,x_0)] - \gamma \text{Var}(F(0,T,u,x_0))
$$

with the risk aversion parameter $\gamma > 0$. Hence, we are looking for the mean-variance optimal strategy for the (continuous) function F , which we define for $s < t$ as

$$
F(s, t, u, x) := \alpha ((X_t - k_1)_+ - k_0) - \alpha_2 (X_t - k_2)_+
$$

=
$$
\begin{cases} -\alpha k_0 & X_t < k_1, \\ \alpha (X_t - k_1 - k_0) & k_1 \le X_t < k_2, \\ \tilde{\alpha} (X_t - k_2) + \alpha (k_2 - k_1 - k_0) & X_t \ge k_2, \end{cases}
$$

where $X_s = x, 0 \le k_0, k_1 \le k_2 < \infty$ with $k_0 + k_1 \le k_2, 0 \le \alpha_2 < \alpha < \infty$ with $\tilde{\alpha} := \alpha - \alpha_2$. Note that $0 < \tilde{\alpha} \leq \alpha$ holds. We also write $F(X_T)$ instead of $F(0, T, u, x)$ when the trading strategy is clear (for instance, for the optimal terminal wealth \hat{X}_T), suppressing the initial value.

Now, one can observe that if $\alpha = 1$, $k_0 = 0$, and $k_1 = G$, the function F reduces to the payoff of the insurer for the non-protected participating life insurance contract $V_{\text{ins}}^{\text{non}}$. If $\alpha = 1$, $k_0 = G$, and $k_1 = 0$, F reduces to the payoff of the insurer for the protected product $V_{\text{ins}}^{\text{pro}}$. In the following Figure [2,](#page-4-0) we show the payoff of the insurer for these two variants of participating life insurance contracts.

Figure 2: Payoffs for the insurer with $k_1 = 2.5$ (left) resp. $k_0 = 2.5$ (right), $k_2 = 7$, $\alpha = 1$, $\alpha_2 = 0.5$.

Remark 2.1. Since the problem in this paper is a variant of a mean-variance optimization, it inherits also the time-inconsistency of the classical mean-variance approach, see, for instance, Basak and Chabakauri [\[3\]](#page-32-18) or Björk and Murgoci [\[5\]](#page-32-19). There are several ways to approach this timeinconsistency. We take in this paper the perspective of a so-called pre-committed investor. Such an investor optimizes once in the beginning and sticks to this strategy, i.e., the chosen strategy of the investor is not dynamically reoptimized. Note that due to the function F , our problem is not covered by the general theory from Björk and Murgoci $[5]$ which assumes a sophisticated investor, i.e., she/he knows that her/his optimal strategy changes over time and only optimizes over all strategies she/he would actually follow.

3 Optimization in a Black-Scholes market

This section assumes that the underlying financial market follows the Black-Scholes model, i.e., the market is complete. Then, the price dynamics for the d risky assets S^i , $i \in \{1, ..., d\}$ are given by

$$
dS_t^i = S_t^i \mu_t^i dt + S_t^i \sigma_t^i dW_t,
$$

where W is the Brownian Motion generating the filtration \mathcal{F} , μ_i is the deterministic drift of the *i*'th asset and σ^{ij} is the deterministic volatility between the *i*'th and the *j*'th asset. We assume that σ_t is bounded, bounded away from zero and invertible. Then, there exists a unique price density process ξ with the wealth process X and the price density process ξ admitting the following dynamics

$$
dX_t = X_t \left[r_t + u_t^T (\mu_t - r) \right] dt + X_t u_t^T \sigma_t dW_t, \qquad (3.1)
$$

$$
d\xi_t = -\xi_t r_t dt - \xi_t \kappa_t^T dW_t, \qquad (3.2)
$$

with $X_0 = x_0$ and $\xi_0 = 1$, where $\kappa_t = (\sigma_t)^{-1}(\mu_t - r_t)$ is the Sharpe ratio process and \cdot^T denotes the transpose of a vector. The term $\xi_T(\omega)$, $\omega \in \Omega$, can be interpreted as the Arrow-Debreu value per probability unit in state ω at time T. Note that $\xi_T(\omega)$ can be written as a decreasing function of the stock price, and therefore attains high values in times of a bad economy and low values in times of a good economy. We assume that the processes r_t and μ_t are integrable and the processes σ_t and κ_t are square-integrable over [0, T] to ensure that the previous SDEs and all of the following integrals are well-defined.

3.1 Derivation of the optimal terminal wealth

The mean-variance optimization [\(2.1\)](#page-4-1) is challenging to solve directly due to the term $(\mathbb{E}[X])^2$ in the decomposition formula $Var(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$. Hence, we show in the following Lemma [3.1](#page-5-1) that if an optimal strategy exists, we can alternatively look for the optimal strategy considering the value functional \tilde{J} which we define as:

$$
\tilde{J}(0,T,u,x_0) := \mathbb{E}[\lambda F(0,T,u,x_0) - \gamma F(0,T,u,x_0)^2]
$$
\n(3.3)

with $\lambda = 1 + 2\gamma \mathbb{E} [F(0,T,\hat{u},x_0)]$ where \hat{u} is the optimal strategy.

Lemma 3.1. If \hat{u} is an optimal strategy for J, it is also an optimal strategy for \tilde{J} .

Consequently, the optimal terminal wealth also coincides for the two problems maximizing J resp. J. This result is a slight generalization of Theorem 3.1 in Zhou and Li $[39]$, who showed this lemma in the case of F being the identity. We will include the proof in [A](#page-18-1)ppendix Λ for the reader's convenience.

Since we have a complete market, we optimize with the following three steps. First, we use a Lagrangian approach to find the optimal terminal wealth \ddot{X} (using the alternative problem). Second, we derive the optimal strategy, and third, we determine the λ and the Lagrangian multiplier y since the optimal terminal wealth and strategy depend on these values initially. In the following, we suppress the dependence on λ and y for the sake of simplicity in the notation unless stated otherwise in some proofs, and we use the convention that $(a, a] = \emptyset$ and $[a, b] = \emptyset$ if $b < a$.

Theorem 3.2. The optimal terminal wealth \hat{X}_T is given by:

$$
\hat{X}_T := \begin{cases}\nk_2 + \frac{\lambda \tilde{\alpha} - y\xi_T}{2\gamma \tilde{\alpha}^2} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0) & \xi_T \in (0, \xi_1^*],\\
k_2 & \xi_T \in (\tilde{\alpha}\hat{\xi}, \xi_2^*],\\
k_0 + k_1 + \frac{\lambda \alpha - y\xi_T}{2\gamma \alpha^2} & \xi_T \in (\alpha \hat{\xi}, \xi_3^*],\\
0 & else,\n\end{cases} \tag{3.4}
$$

where y is the Lagrangian multiplier which solves $\mathbb{E}[\xi_T \hat{X}_T(y)] = \xi_0 x_0$, $\lambda = 1 + 2\gamma \mathbb{E}[F(0, T, \hat{u}, x_0)],$ and

$$
\hat{\xi} := \max \left\{ 0, \frac{\lambda - 2\gamma \alpha (k_2 - k_1 - k_0)}{y} \right\},
$$
\n
$$
\bar{\xi} := \frac{\lambda \alpha}{y} + \frac{2\gamma \alpha^2 k_0}{y},
$$
\n
$$
\tilde{\xi}_1^* := \tilde{\alpha} \hat{\xi} - \frac{2\gamma \tilde{\alpha}}{y} \left(\sqrt{\max \left\{ 0, (\alpha (k_0 + k_1) - \alpha_2 k_2)^2 - \alpha^2 k_0^2 + \frac{\lambda}{\gamma} (\alpha k_1 - \alpha_2 k_2) \right\}} - \tilde{\alpha} k_2 \right),
$$
\n
$$
\xi_1^* := \max \left\{ 0, \min \left\{ \tilde{\alpha} \hat{\xi}, \tilde{\xi}_1^* \right\} \right\},
$$
\n
$$
\tilde{\xi}_2^* := \frac{\alpha \lambda}{y} - \frac{\gamma \alpha^2 (k_2 - k_1)^2 - 2\gamma \alpha^2 k_0 (k_2 - k_1) + \lambda \alpha k_1}{y k_2},
$$
\n
$$
\xi_2^* := \max \left\{ \tilde{\alpha} \hat{\xi}, \min \left\{ \alpha \hat{\xi}, \tilde{\xi}_2^* \right\} \right\},
$$
\n
$$
\xi_3^* := \max \left\{ \alpha \hat{\xi}, \bar{\xi} - \frac{2\gamma \alpha^2}{y} \left(\sqrt{k_1^2 + k_1 \left(2k_0 + \frac{\lambda}{\gamma \alpha} \right)} - k_1 \right) \right\}.
$$

In particular, such \hat{u}, t, y exist. Moreover, let ξ^* be defined by:

$$
\xi^* := \begin{cases} \xi_3^* & ,if \xi_3^* > \alpha \hat{\xi}, \\ \xi_2^* & ,if \xi_3^* = \alpha \hat{\xi}, \xi_2^* > \tilde{\alpha} \hat{\xi}, \\ \xi_1^* & ,if \xi_3^* = \alpha \hat{\xi}, \xi_2^* = \tilde{\alpha} \hat{\xi}. \end{cases}
$$

Then, it holds that $\xi^* > 0$ and $\hat{X}_T > 0$ for $\xi \in (0, \xi^*)$ and $\hat{X}_T = 0$ for $\xi > \xi^*$.

- **Remark 3.3.** (a) The terminal wealth \hat{X}_T denotes the wealth before distributing the wealth to the insurer and the policyholders. The terminal wealth of the insurer is given by $F(\hat{X}_T) =$ $\alpha((\hat{X}_T-k_1)_+-k_0)-\alpha_2(\hat{X}_T-k_2)_+.$ In particular, in the case of a protected participating life insurance contract, i.e., $\alpha = 1$, $k_1 = 0$, the insurer makes a loss if $\hat{X}_T < k_0$. In the case of a non-protected participating life insurance contract, the insurer cannot make a loss by construction.
- (b) If $\alpha_2 = \alpha$ resp. $\tilde{\alpha} = 0$, the surplus over k_2 is fully distributed to the policyholders. In this case, it holds that $\tilde{\xi}_1^* = 0$ and $\tilde{\alpha}\hat{\xi} = 0$, i.e., the result is similar with the exception that the first case, $\hat{X}_T = k_2 + \frac{\lambda \tilde{\alpha} - y\xi_T}{2\gamma \tilde{\alpha}^2}$ $\frac{\tilde{\alpha}-y\xi_T}{2\gamma\tilde{\alpha}^2}-\frac{\alpha}{\tilde{\alpha}}$ $\frac{\alpha}{\tilde{\alpha}}(k_2-k_1-k_0)$, does not need to be considered. The proofs are similar, but we exclude this case for the ease of exposition.
- (c) We can generalize the result for $X_T \in \mathbb{R}$ when we extend the function F to $\mathbb{R}_{\leq 0}$ as $F(X_T) =$ $\alpha(X_T - k_0)$. Then, the optimal terminal wealth has the same structure as before, but with the additional case that $\hat{X}_T = k_0 + \frac{\lambda \alpha - y \xi_T}{2\gamma \alpha^2}$ if $\xi_T > \bar{\xi}$. Hence, the restriction to $X_T \ge 0$ is not crucial for the structure of the solution. Note that $F(\hat{X}_T(\xi_T = \bar{\xi})) = 0$, and that for $\xi_T > \bar{\xi}$ the function F is linearly decreasing in ξ_T . Again, the proofs are similar to before, but we nevertheless restrict to $X_T \geq 0$ to avoid too technical proofs.

(d) The representation of \hat{X}_T also holds in a general complete market (without necessarily using a Black-Scholes market model) if we assume the existence of the multipliers y and λ . However, that these exist is then not clear. Note that the price density process, ξ , in a different complete market would change, but still admit a geometric form. For example, if we replace the Brownian Motion W by a Poisson process N with parameter λ_2 (in the Black-Scholes model), the price density admits a dynamic of the form (see, e.g., Cont and Tankov [\[12,](#page-32-20) p.303]):

$$
d\xi_{t-} = \xi_{t-}(\lambda_2 - \lambda_1 - r_t)dt - \xi_{t-} \ln \frac{\lambda_2}{\lambda_1} dN_t,
$$

where $\lambda_1 > 0$ is an appropriately chosen constant.

(e) The way how to compute the parameters λ and y is given in the proof of Proposition [A.1.](#page-21-0)

From the definition formula of \hat{X}_T , we can derive the following two propositions, which are proven in Appendix [A:](#page-18-1)

Proposition 3.4. It holds that $(0, \xi_1^*] \cup (\tilde{\alpha}\hat{\xi}, \xi_2^*] \cup (\alpha\hat{\xi}, \xi_3^*] = (0, \xi^*]$ with these constants defined in Theorem [3.2,](#page-6-2) i.e., the three intervals of the terminal wealth are connected.

Proposition 3.5. It holds that \hat{X}_T as a function of ξ is continuous and non-increasing in $(0,\xi^*)\cup$ (ξ^*, ∞) with ξ^* as in Theorem [3.2.](#page-6-2) If $k_1 > 0$, then \hat{X}_T is always discontinuous at ξ^* .

Therefore, \hat{X}_T as a function of ξ_T is non-increasing, taking the value 0 for $\xi_T > \xi^*$ and one possible discontinuity point. See the black lines in Figure [3](#page-11-0) for a visualization.

In the following proposition, we give an alternative formulation for our optimization problem. Again, we give the proof in Appendix [A.](#page-18-1)

Proposition 3.6. Let $\tilde{L}(x) = \begin{cases} \gamma \alpha^2 k_0^2, & \text{if } x < k_1, \\ \gamma \alpha^2 (x - k_1 - k_0)^2, & \text{if } k_1 \leq x \end{cases}$ $\gamma \alpha^2 (x - k_1 - k_0)^2$, if $k_1 \le x < k_2$, $\gamma \tilde{\alpha}^2 (x - k_2 + \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0))^2$, if $x \ge k_2$. and consider the opti-

mization problem $\min_{u \in \mathcal{U}} \mathbb{E}[\tilde{L}(X_T)]$ under the constraint $\mathbb{E}[\alpha(X_T(\lambda) - k_1) \mathbb{1}_{k_1 \leq X_T(\lambda) < k_2}$ $\tilde{\alpha}(X_T(\lambda) - k_2 + \frac{\alpha}{\tilde{\alpha}})$ $\frac{\alpha}{\tilde{\alpha}}(k_2 - k_1) \hat{\mathbb{1}}_{X_T(\lambda) \ge k_2}] = \alpha k_0$ and the budget constraint $\mathbb{E}[\xi_T X_T(y)] = \xi_0 x_0$, where λ and y are the respective Lagrangian multipliers.

Then, the optimal terminal wealth is identical to the one of our main problem, i.e., given by $(3.4).$ $(3.4).$

Note that the loss function \tilde{L} is continuous and non-decreasing in x, and that the definition and therefore the interpretation of the parameter λ is different than for mean-variance. However, Proposition [3.6](#page-7-0) still gives an equivalent formulation. As already stated in the introduction, this formulation is a 2-target problem. Depending on the region of the terminal wealth, we want to minimize the distance to different exogenously given target values $(k_1 + k_0)$ if the terminal wealth $X \in [k_1, k_2)$ respectively to $k_2 - \frac{\alpha}{\tilde{\alpha}}$ $\frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)$ if $X \ge k_2$). Moreover, the first constraint entails that one can only reduce mass in one region when one simultaneously increases the mass in another region where the reduction and the enlargement are, in general, not of equal size. Especially, the weight of the region $[k_2, \infty)$ is heavily influenced by the participation rate α_2 through the pre-factor $\tilde{\alpha} (= \alpha - \alpha_2)$.

Proof of Theorem [3.2.](#page-6-2) We use the Lagrangian multiplier method to prove this theorem. Therefore, choose $\lambda > 0$ and $y > 0$ as in Proposition [A.1](#page-21-0) in the appendix and we define the Lagrangian function L for $X \geq 0$ as

$$
L(X, y) := \lambda F(0, T, u, x_0) - \gamma F(0, T, u, x_0)^2 - y\xi X
$$
\n
$$
= \lambda \left[\alpha ((X - k_1)_+ - k_0) - \alpha_2 (X - k_2)_+ \right] - \gamma \left[\alpha ((X - k_1)_+ - k_0) - \alpha_2 (X - k_2)_+ \right]^2 - y\xi X
$$
\n
$$
= \lambda \left[\alpha (X - k_1 - k_0) \mathbb{1}_{k_1 \le X < k_2} + \tilde{\alpha} (X - k_2) \mathbb{1}_{X \ge k_2} + \alpha (k_2 - k_1 - k_0) \mathbb{1}_{X \ge k_2} - \alpha k_0 \mathbb{1}_{X < k_1} \right]
$$
\n(3.5)

$$
-\gamma [\alpha(X - k_1 - k_0) \mathbb{1}_{k_1 \le X < k_2} + \tilde{\alpha}(X - k_2) \mathbb{1}_{X \ge k_2} + \alpha(k_2 - k_1 - k_0) \mathbb{1}_{X \ge k_2} - \alpha k_0 \mathbb{1}_{X < k_1}]^2
$$

\n
$$
- y\xi X,
$$

\n
$$
= \lambda \alpha(X - k_1 - k_0) \mathbb{1}_{k_1 \le X < k_2} + \lambda [\tilde{\alpha}(X - k_2) + \alpha(k_2 - k_1 - k_0)] \mathbb{1}_{X \ge k_2} - \lambda \alpha k_0 \mathbb{1}_{X < k_1}
$$

\n
$$
-\gamma \alpha^2 (X - k_1 - k_0)^2 \mathbb{1}_{k_1 \le X < k_2} - \gamma [\tilde{\alpha}(X - k_2) + \alpha(k_2 - k_1 - k_0)]^2 \mathbb{1}_{X \ge k_2} - \gamma \alpha^2 k_0^2 \mathbb{1}_{X < k_1}
$$

\n
$$
- y\xi X
$$

\n
$$
= \begin{cases}\n-\lambda \alpha k_0 - \gamma \alpha^2 k_0^2 - y\xi X & X \in [0, k_1), \\
\lambda \alpha(X - k_1 - k_0) - \gamma \alpha^2 (X - k_1 - k_0)^2 - y\xi X & X \in [k_1, k_2), \\
\lambda [\tilde{\alpha}(X - k_2) + \alpha(k_2 - k_1 - k_0)] - \gamma [\tilde{\alpha}(X - k_2) + \alpha(k_2 - k_1 - k_0)]^2 - y\xi X & X \in [k_2, \infty),\n\end{cases}
$$

where we suppress the ω , write X instead of X_T , ξ instead of ξ_T , and use y as the multiplier. The following paragraph considers L as a function of X. Obviously, L is not smooth in k_1 and k_2 , but it is continuous in $[0, \infty)$. Hence, we optimize L in the regions $[0, k_1]$, $[k_1, k_2]$, and $[k_2, \infty)$ separately and compare afterwards the minimal points. Now, for the individual optimization, we get:

$$
\frac{\partial L}{\partial X} = \begin{cases}\n-y\xi & X \in (0, k_1), \\
\lambda \alpha - 2\gamma \alpha^2 (X - k_1 - k_0) - y\xi & X \in (k_1, k_2), \\
\lambda \tilde{\alpha} - 2\gamma \tilde{\alpha} [\tilde{\alpha}(X - k_2) + \alpha(k_2 - k_1 - k_0)] - y\xi & X \in (k_2, \infty).\n\end{cases}
$$

Hence, it follows that we get the following possible maximal points (if they are in the respective interval): $X_1 = 0$, $X_2 = k_0 + k_1 + \frac{\lambda \alpha - y\xi}{2\gamma \alpha^2}$, and $X_3 = k_2 + \frac{\lambda \tilde{\alpha} - y\xi}{2\gamma \tilde{\alpha}^2}$ $\frac{\lambda \tilde{\alpha}-y\xi}{2\gamma \tilde{\alpha}^2}-\frac{\alpha}{\tilde{\alpha}}$ $\frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)$. To have X_2 and X_3 in the correct interval, we get the following conditions (since $\xi > 0$):

$$
X_2 \in [k_1, k_2] \qquad \Leftrightarrow \qquad \xi \in \left[\frac{\lambda \alpha - 2\gamma \alpha^2 (k_2 - k_1 - k_0)}{y}, \frac{\lambda \alpha}{y} + \frac{2\gamma \alpha^2 k_0}{y}\right] = \left[\alpha \hat{\xi}, \bar{\xi}\right],\tag{3.6}
$$

$$
X_3 \in [k_2, \infty) \qquad \Leftrightarrow \qquad \xi \in \left(0, \frac{\lambda \tilde{\alpha} - 2\gamma \tilde{\alpha} \alpha (k_2 - k_1 - k_0)}{y}\right] = \left(0, \tilde{\alpha} \hat{\xi}\right]. \tag{3.7}
$$

Moreover, for the maximum of L , we must consider the different boundary values of the corresponding intervals. The first boundary value k_1 is always dominated by X_1 due to L being non-increasing in [0, k₁]. The second boundary value k₂ is dominated by X_2 or by X_3 if ξ is in one of the intervals from [\(3.6\)](#page-8-0) or [\(3.7\)](#page-8-1). If $\xi > \xi$, then L is non-increasing in $[0, \infty)$ since then $\frac{\partial L}{\partial X} \leq 0$ for all $X \geq 0$ and hence k_2 is dominated by X_1 . If $\xi \in [\tilde{\alpha}\hat{\xi}, \alpha \hat{\xi}]$, then L has a local maximum in k_2 since $\frac{\partial L}{\partial X} \geq 0$ for $X \in (k_1, k_2)$ and $\frac{\partial L}{\partial X} \leq 0$ for $X > k_2$. Hence, we add

$$
X_4 = k_2 \quad \text{for} \quad \xi \in [\tilde{\alpha}\hat{\xi}, \alpha\hat{\xi}] \tag{3.8}
$$

to the list of potential maximum points. We note that the intervals of ξ for X_2 , X_3 , and X_4 are disjoint except the lower (resp. upper) boundary points of X_2 (resp. X_3) with the upper (resp. lower) boundary point of X_4 . However, in this case X_2 (resp. X_3) coincides with X_4 . It is remarkable that due to the disjointness of the related intervals for X_2 , X_3 , and X_4 , we do not have to compare these values themselves with each other (for the potential maximum of L). Thus, we replace the closed intervals for ξ by left-open, right-closed intervals and denote by X^* the combined solution of X_2 , X_3 , and X_4 with $X^* = 0$ for $\xi > \overline{\xi}$, i.e.,

$$
X^* := \begin{cases} k_2 + \frac{\lambda \tilde{\alpha} - y\xi}{2\gamma \tilde{\alpha}^2} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0) & \xi \in (0, \tilde{\alpha}\hat{\xi}], \\ k_2 & \xi \in (\tilde{\alpha}\hat{\xi}, \alpha\hat{\xi}], \\ k_0 + k_1 + \frac{\lambda \alpha - y\xi}{2\gamma \alpha^2} & \xi \in (\alpha\hat{\xi}, \bar{\xi}], \\ 0 & \xi \in (\bar{\xi}, \infty). \end{cases}
$$

Note that X^* takes the same values as \hat{X}_T in [\(3.4\)](#page-6-3), but the intervals are not truncated. Moreover, we observe that X^* is continuous in ξ on $(0,\overline{\xi}]$. Therefore, we only have to compare $L(X^*,y)$, y fixed, with $L(X_1, y) = -\gamma \alpha^2 k_0^2 - \lambda \alpha k_0$. Thus, we compare these values in the following separately for the three cases, i.e., depending on ξ , which value out of $\{X_2, X_3, X_4\}$ X^{*} attains. Note that we consider, from now on, L as a function of ξ . To emphasize this, we write $L(X, y, \xi)$.

Case 1: $\xi \in (\alpha \hat{\xi}, \bar{\xi}],$ i.e., $X^* = X_2 \in [k_1, k_2]$: Then, we get:

$$
L(X_2, y, \xi) = \lambda \alpha \frac{\lambda \alpha - y\xi}{2\gamma \alpha^2} - \gamma \alpha^2 \left(\frac{\lambda \alpha - y\xi}{2\gamma \alpha^2}\right)^2 - y\xi(k_0 + k_1) - y\xi \frac{\lambda \alpha - y\xi}{2\gamma \alpha^2}
$$

=
$$
\frac{y^2}{4\gamma \alpha^2} \xi^2 - y\left(k_0 + k_1 + \frac{\lambda}{2\gamma \alpha}\right)\xi + \frac{\lambda^2}{4\gamma}.
$$
 (3.9)

Then $L(X_2, y, \xi) \ge -\gamma \alpha^2 k_0^2 - \lambda \alpha k_0 = L(X_1, y, \xi)$ if $\xi \le \xi$ or $\xi \ge \xi$ with

$$
\xi_{\pm} = \frac{2\gamma\alpha^2}{y^2} \left(y \left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha} \right) \pm \sqrt{y^2 \left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha} \right)^2 - \frac{y^2}{\gamma\alpha^2} \left(\frac{\lambda^2}{4\gamma} + \gamma\alpha^2 k_0^2 + \lambda\alpha k_0 \right)} \right)
$$
\n
$$
= \frac{\lambda\alpha}{y} + \frac{2\gamma\alpha^2 (k_0 + k_1)}{y} \pm \frac{2\gamma\alpha^2}{y^2} \sqrt{y^2 k_1^2 + \frac{y^2 k_1 \lambda}{\gamma\alpha} + 2y^2 k_0 k_1}
$$
\n
$$
= \frac{\lambda\alpha}{y} + \frac{2\gamma\alpha^2 (k_0 + k_1)}{y} \pm \frac{2\gamma\alpha^2}{y} \sqrt{k_1^2 + k_1 \left(2k_0 + \frac{\lambda}{\gamma\alpha} \right)}
$$
\n
$$
= \bar{\xi} + \frac{2\gamma\alpha^2}{y} \left(\pm \sqrt{k_1^2 + k_1 \left(2k_0 + \frac{\lambda}{\gamma\alpha} \right)} + k_1 \right).
$$

We notice that $\xi_+ > \bar{\xi}$ and $\max \{\alpha \hat{\xi}, \xi_-\} = \xi_3^*$ where ξ_3^* is defined as in Theorem [3.2.](#page-6-2) If $\xi_- < \alpha \hat{\xi}$, X_2 is not in the necessary interval $[k_1, k_2)$ which entails that $(\alpha \hat{\xi}, \xi_3^*] = \emptyset$. Otherwise, it holds that $\xi = \xi_3^*$ by which we get [\(3.4\)](#page-6-3) for this case, i.e., the interval for ξ being in the interval $(\alpha \hat{\xi}, \xi_3^*]$ for the optimal wealth \hat{X}_T .

<u>Case 2:</u> $\xi \in (0, \tilde{\alpha}\hat{\xi}],$ i.e., $X^* = X_3 \in [k_2, \infty)$: Then it holds:

$$
L(X_3, y, \xi) = \lambda \left(\frac{\lambda \tilde{\alpha} - y\xi}{2\gamma \tilde{\alpha}} - \tilde{\alpha} \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) + \alpha (k_2 - k_1 - k_0) \right) - \gamma \left(\tilde{\alpha} \frac{\lambda \tilde{\alpha} - y\xi}{2\gamma \tilde{\alpha}^2} - \tilde{\alpha} \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) + \alpha (k_2 - k_1 - k_0) \right)^2 - y\xi k_2 - y\xi \frac{\lambda \tilde{\alpha} - y\xi}{2\gamma \tilde{\alpha}^2} + y\xi \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) = \frac{y^2}{4\gamma \tilde{\alpha}^2} \xi^2 - y \left(k_2 - \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) + \frac{\lambda}{2\gamma \tilde{\alpha}} \right) \xi + \frac{\lambda^2}{4\gamma}.
$$
(3.10)

Note that $k_2 - \frac{\alpha}{\tilde{\alpha}}$ $\frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0) = \frac{\alpha}{\tilde{\alpha}}(k_0 + k_1) - \frac{\alpha_2}{\tilde{\alpha}}k_2$. First we assume that the discriminant, i.e., the term under the square root in the following equation, is non-negative. Then, we have $L(X_3, y, \xi) \ge -\gamma \alpha^2 k_0^2 - \lambda \alpha k_0 = L(X_1, y, \xi)$ if $\xi \le \xi$ or $\xi \ge \xi$ with

$$
\xi_{\pm} = \frac{2\gamma \tilde{\alpha}^2}{y^2} \left(y \left(k_2 - \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) + \frac{\lambda}{2\gamma \tilde{\alpha}} \right) \right) \n\pm \frac{2\gamma \tilde{\alpha}^2}{y^2} \sqrt{y^2 \left(k_2 - \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) + \frac{\lambda}{2\gamma \tilde{\alpha}} \right)^2 - \frac{y^2}{\gamma \tilde{\alpha}^2} \left(\frac{\lambda^2}{4\gamma} + \gamma \alpha^2 k_0^2 + \lambda \alpha k_0 \right)}
$$

$$
\begin{split}\n&=\frac{2\gamma\tilde{\alpha}^{2}}{y^{2}}\left(y\left(k_{2}-\frac{\alpha}{\tilde{\alpha}}(k_{2}-k_{1}-k_{0})+\frac{\lambda}{2\gamma\tilde{\alpha}}\right)\right) \\
&+\frac{2\gamma\tilde{\alpha}^{2}}{y^{2}}\sqrt{y^{2}\left(\frac{\alpha^{2}}{\tilde{\alpha}^{2}}(2k_{0}k_{1}+k_{1}^{2})+\frac{\alpha_{2}^{2}}{\tilde{\alpha}^{2}}k_{2}^{2}-2\frac{\alpha\alpha_{2}}{\tilde{\alpha}^{2}}(k_{0}+k_{1})k_{2}+\frac{\lambda}{\gamma\tilde{\alpha}^{2}}(\alpha k_{1}-\alpha_{2}k_{2})\right)} \\
&=\frac{\lambda\tilde{\alpha}}{y}-\frac{2\gamma\alpha\tilde{\alpha}(k_{2}-k_{1}-k_{0})}{y}+\frac{2\gamma\tilde{\alpha}^{2}k_{2}}{y} \\
&+\frac{2\gamma\tilde{\alpha}}{y}\sqrt{\alpha^{2}(2k_{0}k_{1}+k_{1}^{2})+\alpha_{2}^{2}k_{2}^{2}-2\alpha\alpha_{2}(k_{0}+k_{1})k_{2}+\frac{\lambda}{\gamma}(\alpha k_{1}-\alpha_{2}k_{2})} \\
&=\tilde{\alpha}\hat{\xi}+\frac{2\gamma\tilde{\alpha}}{y}\left(\pm\sqrt{(\alpha(k_{0}+k_{1})-\alpha_{2}k_{2})^{2}-\alpha^{2}k_{0}^{2}+\frac{\lambda}{\gamma}(\alpha k_{1}-\alpha_{2}k_{2})}+\tilde{\alpha}k_{2}\right).\n\end{split}
$$

Again, we notice that $\xi_+ > \tilde{\alpha}\hat{\xi}$ and if the discriminant is non-negative, we get $\xi_- = \tilde{\xi}_1^*$ where we can ignore this case if $\xi_-\leq 0$ due to the assumption that ξ has to be in the interval $(0,\tilde{\alpha}\hat{\xi}]$. Now, if the discriminant is negative, then $L(X_3, y, \xi) \ge -\gamma \alpha^2 k_0^2 - \lambda \alpha k_0$ for all ξ and $\tilde{\xi}_1^* > \tilde{\alpha} \hat{\xi}$. Hence, X_3 is optimal for all $\xi \in (0, \tilde{\alpha}\hat{\xi}]$ and it holds that $\xi_1^* = \tilde{\alpha}\hat{\xi}$ which implies (3.4) for this case.

Case 3: $\xi \in (\tilde{\alpha}\hat{\xi}, \alpha\hat{\xi}],$ i.e., $X^* = X_4 = k_2$: Now, we get:

$$
L(X_4, y, \xi) = \lambda \alpha (k_2 - k_1 - k_0) - \gamma \alpha^2 (k_2 - k_1 - k_0)^2 - y \xi k_2.
$$
 (3.11)

Then $L(X_4, y, \xi) \ge -\gamma \alpha^2 k_0^2 - \lambda \alpha k_0 = L(X_1, y, \xi)$ if

$$
\xi \le \frac{\lambda \alpha}{y} - \frac{\gamma \alpha^2 (k_2 - k_1)^2 - 2\gamma \alpha^2 k_0 (k_2 - k_1) + \lambda \alpha k_1}{y k_2} = \tilde{\xi}_2^*.
$$

Hence, it follows that $\xi \in (\tilde{\alpha}\hat{\xi}, \tilde{\xi}_2^*]$ which implies (3.4) in this case as well. Hence, (3.4) holds in total.

Finally, we must have that $\xi^* > 0$ since otherwise $(0, \xi_1^*] \cup (\tilde{\alpha}\hat{\xi}, \xi_2^*] \cup (\alpha\hat{\xi}, \xi_3^*] = \emptyset$ which implies that $\hat{X}_T \equiv 0$. Hence, the budget constraint $\mathbb{E}[\xi_T \hat{X}_T(y)] = \xi_0 x_0$ cannot be fulfilled, which is a contradiction to Proposition [A.1](#page-21-0) whose proof is given below. The property of \hat{X}_T follows directly from Proposition [3.4](#page-7-1) and $y < \infty$. \Box

We deduce from Theorem [3.2:](#page-6-2)

Corollary 3.7. The optimal payoff of the insurer is given by:

$$
F(\hat{X}_T) = \begin{cases} \frac{\lambda \tilde{\alpha} - y\xi_T}{2\gamma \tilde{\alpha}} & \xi_T \in (0, \xi_1^*],\\ \alpha(k_2 - k_1 - k_0) & \xi_T \in (\tilde{\alpha}\hat{\xi}, \xi_2^*],\\ \frac{\lambda \alpha - y\xi_T}{2\gamma \alpha} & \xi_T \in (\alpha \hat{\xi}, \xi_3^*],\\ -\alpha k_0 & else, \end{cases}
$$

where $\hat{\xi}$, ξ_1^* , ξ_2^* , and ξ_3^* are defined as in Theorem [3.2.](#page-6-2)

Proof. This proof follows immediately by simple calculations when we plug (3.4) into the function F. \Box

In the following Figure [3,](#page-11-0) we derive the shape of the optimal value \hat{X}_T and the corresponding wealth of the insurance company as a function of ξ_T .

Non−protected participating life insurance

Protected participating life insurance

Figure 3: The optimal value \hat{X}_T and the optimal terminal wealth of the insurance company as a function of ξ_T with $k_1 = 2.5$ (left) resp. $k_0 = 2.5$ (right), $k_2 = 7$, $\alpha = 1$, $\alpha_2 = 0.5$, $\lambda = 3$, $y = 1$, $\gamma = 0.25$.

3.2 Derivation of the optimal strategy

Next, we derive the optimal strategy for our problem where the proof is somewhat technical and therefore given in Appendix [A.](#page-18-1)

Theorem 3.8. The optimal solution \hat{u} is non-negative and given by:

$$
\hat{u}_t = (\sigma_t^T)^{-1} \kappa_t \frac{v_t}{\hat{X}_t},
$$

where

$$
v_{t} = \left(k_{2} + \frac{\lambda}{2\gamma\tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}}(k_{2} - k_{1} - k_{0})\right) \frac{e^{-\int_{t}^{T} r_{s}ds}}{\sqrt{\int_{t}^{T} ||\kappa_{s}||^{2} ds}} \varphi\left(d_{1}\left(\xi_{1}^{*}, t\right)\right)
$$

+
$$
\frac{y}{2\gamma\tilde{\alpha}^{2}} \xi_{t} e^{\int_{t}^{T} -(2r_{s} - ||\kappa_{s}||^{2}) ds} \left[\Phi\left(d_{2}\left(\xi_{1}^{*}, t\right)\right) - \frac{1}{\sqrt{\int_{t}^{T} ||\kappa_{s}||^{2} ds}} \varphi\left(d_{2}\left(\xi_{1}^{*}, t\right)\right)\right]
$$

+
$$
k_{2} \frac{e^{-\int_{t}^{T} r_{s}ds}}{\sqrt{\int_{t}^{T} ||\kappa_{s}||^{2} ds}} \left(\varphi\left(d_{1}\left(\xi_{2}^{*}, t\right)\right) - \varphi\left(d_{1}\left(\tilde{\alpha}\xi, t\right)\right)\right)
$$

+
$$
\left(k_{0} + k_{1} + \frac{\lambda}{2\gamma\alpha}\right) \frac{e^{-\int_{t}^{T} r_{s}ds}}{\sqrt{\int_{t}^{T} ||\kappa_{s}||^{2} ds}} \left(\varphi\left(d_{1}\left(\xi_{3}^{*}, t\right)\right) - \varphi\left(d_{1}\left(\alpha\xi, t\right)\right)\right)
$$

+
$$
\frac{y}{2\gamma\alpha^{2}} \xi_{t} e^{\int_{t}^{T} -(2r_{s} - ||\kappa_{s}||^{2}) ds} \left[\left(\Phi\left(d_{2}\left(\xi_{3}^{*}, t\right)\right) - \Phi\left(d_{2}\left(\alpha\xi, t\right)\right)\right)\right]
$$

-
$$
\frac{1}{\sqrt{\int_{t}^{T} ||\kappa_{s}||^{2} ds}} \left(\varphi\left(d_{2}\left(\xi_{3}^{*}, t\right)\right) - \varphi\left(d_{2}\left(\alpha\xi, t\right)\right)\right)
$$

$$
\hat{X}_{t} = \left(k_{2} + \frac{\lambda}{2\gamma\tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}}(k_{2} - k_{1} - k_{0})\
$$

+
$$
k_2 e^{-\int_t^T r_s ds} \left(\Phi(d_1(\xi_2^*, t)) - \Phi(d_1(\tilde{\alpha}\hat{\xi}, t)) \right)
$$

+ $\left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha} \right) e^{-\int_t^T r_s ds} \left(\Phi(d_1(\xi_3^*, t)) - \Phi(d_1(\alpha\hat{\xi}, t)) \right)$
- $\frac{y}{2\gamma\alpha^2} \xi_t e^{\int_t^T -(2r_s - ||\kappa_s||^2) ds} \left(\Phi(d_2(\xi_3^*, t)) - \Phi(d_2(\alpha\hat{\xi}, t)) \right)$

with

$$
d_1(x,t) = \frac{\ln x - \ln \xi_t + \int_t^T (r_s - \frac{\|\kappa_s\|^2}{2}) ds}{\sqrt{\int_t^T \|\kappa_s\|^2 ds}},
$$

$$
d_2(x,t) = \frac{\ln x - \ln \xi_t + \int_t^T (r_s - \frac{3\|\kappa_s\|^2}{2}) ds}{\sqrt{\int_t^T \|\kappa_s\|^2 ds}} = d_1(x,t) - \sqrt{\int_t^T \|\kappa_s\|^2 ds},
$$

where Φ denotes the cdf and ϕ the density of a standard normal distributed random variable. The values $\hat{\xi}$, ξ_1^* , ξ_2^* , and ξ_3^* are defined as in Theorem [3.2.](#page-6-2)

4 Numerical Results

In the following section, we discuss the numerical results for the Black-Scholes model with a given calibration of the parameters with a setting as in Section [3.](#page-5-0) The first two figures show the impact of different values for the risk aversion resp. the participation rate and of different optimization strategies (EU instead of mean-variance) on the optimal terminal wealth as a function of ξ_T . The remaining four figures show simulation outcomes for the optimal terminal wealth and the optimal strategy. We analyze the results for the non-protected resp. the protected participation life insurance product, and compare those (a) to each other, (b) to a product with no surplus participation, and (c) to the results with an EU-optimization.

For the participating life insurance contract, we use the parameters $k_0 = 0$, $k_1 = 2.5$, $k_2 = 7$, $\alpha = 1$, and $\alpha_2 = 0.25$ for the non-protected insurance product. We change the values of k_0 and k_1 , i.e., $k_0 = 2.5$ and $k_1 = 0$, to get the protected product. The initial wealth, i.e., the sum of the premiums and the initial capital from the insurance company, is given by $x_0 = 4$, the time horizon by $T = 10$, and the risk aversion by $\gamma = 0.25$. We assume a constant risk-free interest rate of $r := r_t \equiv 0.02$ for the financial market and consider one risky asset with constant mean $\mu := \mu_t \equiv 0.08$ and volatility $\sigma_t \equiv 0.2$. In particular, the Sharpe ratio is also constant and given by $\kappa_t \equiv 0.3$. For the implementation, we used a time step of $\delta = 0.01$, i.e., we have 1000 intermediate points. Moreover, we simulate 10000 realizations of the Brownian Motion.

In Figure [4,](#page-13-0) we present the optimal terminal wealth as a function of ξ_T for different values of risk aversion parameter γ . (Note that the risk aversion parameters are not comparable for the different optimization strategies.) For both the non-protected and the protected case, we compare the optimal terminal wealth stemming from our mean-variance optimization with the EU-optimization for participating life insurance contracts from Lin et al. [\[25\]](#page-33-8). To measure the utility, we use, as Lin et al., the following S-shaped utility function: $U(x) = \begin{cases} x^{\tilde{\gamma}} & x \geq 0, \\ z & \tilde{\gamma} \end{cases}$ $-\tilde{\lambda}(-x)^{-\tilde{\gamma}} \quad x < 0$

with $\lambda = 2$ and different values for $\tilde{\gamma}$ (Lin et al. [\[25\]](#page-33-8) used $\tilde{\gamma} = 0.5$ in their numerical analysis). Moreover, we added the case that there is no additional surplus participation of the policyholders, i.e., $\alpha_2 = 0$, $k_0 = 0$, and $k_1 = 0$. Both participating optimizations share that the optimal wealth function has three points of significant behavioral change, i.e., at ξ_1^*, ξ_2^* and ξ_3^* as in Theorem [3.2.](#page-6-2) This theorem shows that the optimal wealth decreases before ξ_1^* and between ξ_2^* and ξ_3^* , and that the optimal wealth is constant elsewhere. The main difference in the shape of the functions is that we have a piecewise linearity in the mean-variance case but a total non-linearity in the EU case.

This effect is already apparent from the respective formulas. Noticeably, the distance between the second and third point of behavioral change is much higher in the EU-optimization than in the mean-variance one. Moreover, the figure shows a moral hazard of the insurance companies offering a non-protected participating life insurance contract since the insurance companies favor a portfolio value of 0 over a portfolio value at the guarantee value level. We can observe this effect for both the mean-variance and the EU-optimization in the figure due to the drop-down of the optimal terminal wealth to zero. Protected products have no such effect when optimizing mean-variance since the insurance company always profits from higher portfolio values. While this effect also occurs when optimizing EU, the probability is smaller since the drop-down is at a far higher value (compared to the non-protected product).

Figure 4: Comparison of the optimal terminal wealth as a function of ξ_T of mean-variance with EU optimized participating life insurance products and no participation in the case of a not protected guarantee (left) and a protected one (right). Note that the risk aversion parameters are not comparable for mean-variance and EU.

In Figure [5,](#page-14-0) we show the influence of the participation rate α_2 on the optimal terminal wealth as a function of ξ_T . We can see that a higher rate α_2 leads to a more extended plateau at $k_2 = 7$. This effect is not surprising when checking the formula (3.4) of the optimal terminal wealth \hat{X}_T since a higher α_2 leads to a smaller $\tilde{\alpha}$. Thus, the intermediate interval $(\tilde{\alpha}\hat{\xi},\xi_2^*)$ gets bigger since for our parametrization, it always holds that $\xi_2^* = \alpha \hat{\xi}$. Moreover, we can observe that ξ^* (defined in Theorem [3.2](#page-6-2) as the point where in the non-protected case the optimal terminal wealth drops down to 0) is increasing for an increasing participation rate α_2 . This effect stems from the different values of the variables λ and y when changing the participation rate. Additionally, the figure shows that for minimal values of ξ_T , the optimal terminal wealth is higher for large participation rates, which we can derive directly from the formula of \hat{X}_T in [\(3.4\)](#page-6-3). For the economic interpretation, we infer that a higher participation rate of the policyholders leads to a higher portfolio value in bad economic states, because for a higher α_2 , the upside potential of the insurer is reduced. Hence, the optimal strategy should be less risky, which implies a higher portfolio value. For good economic states, the opposite effect happens. A somewhat surprising result is that in very good economic states (happening with an extremely low probability), the optimal terminal wealth is higher for higher participation rates, which occurs due to changes in the Lagrange multiplier of the budget constraint.

In both, Figure [4](#page-13-0) and Figure [5,](#page-14-0) we can observe a jump of the optimal terminal wealth in the non-protected case, which is typical for problems with option-like constraints, see for

Figure 5: Influence of the participation rate α_2 on the optimal terminal wealth as a function of ξ_T of a not protected (left) and a protected (right) participation life insurance.

instance Basak and Shapiro $[4]$, Chen et al. $[8]$, Nguyen and Stadje $[33]$, Chen et al. $[9]$, or Kraft and Steffensen [\[21\]](#page-32-10). The reason is that for certain outcomes, the insurance company's incentive change and therefore it can gain additional utility by changing to a different regime.^{[2](#page-14-1)} Moreover, the piecewise linearity observed also occurs in, e.g., Avanzi et al. [\[1\]](#page-32-16) who considered a minimization of a quadratic function under solvency constraints including option-like constraints as the expected shortfall. This piecewise linearity is typical for problems with some kind of quadratic penalties, whereas the option-like constraints entail the existence of the plateaus and the jumps.

In Figure [6,](#page-15-0) we show the optimal wealth process and the optimal share into the risky asset over time for the non-protected participating life insurance product. In both figures, the red line shows the average of the 10000 realizations, and each of the ten black lines shows a single realization. Note that for the optimal strategy, we used a weighted average with the weight given by the absolute investment amount (which also holds for all other averages in this section). Furthermore, we get $\lambda \approx 3.423$ and $y \approx 0.860$. The average optimal wealth process develops from 4 to approximately 7.6, where the increase is slightly larger in the beginning than in the end. A terminal wealth of about 7.6 corresponds to a terminal wealth for the insurer of 4.95 and a wealth of 2.65 for the policyholder. Note that this is higher than the guarantee value, which is 2.5. When exercising the optimal strategy, we start with high investment into the risky asset (approximately 110% of our wealth), which decreases relatively constant over time. The final optimal investment share into the risky asset is around 45%. When analyzing the strategies in more detail, one observes that the most risky investments are taken when the economy has poorly evolved until then, i.e., when ξ_t is high, while the least risky investments are taken in cases where the economy has developed well. These results are reasonable since when the wealth is below the guarantee, the insurer has nothing to lose anymore, while the wealth is already over the second threshold k_2 (i.e., the policyholder also participates in the surplus over k_2), the upside potential of the insurer is reduced, whereas the downside potential is not (or only slightly).

In Figure [7,](#page-16-0) we show the optimal wealth process and the optimal share into the risky asset over time for the protected participating life insurance product. Again, the red line shows the average

 $\frac{2}{3}$ For instance, if the entire wealth for certain outcomes is transferred to the policyholders, the insurance company will target a terminal wealth of zero in these states in order to gain additional wealth in states where it benefits more. (Note that the overall expectation is fixed due to the budget constraint.)

Figure 6: Optimal wealth and optimal strategy for the non-protected participating life insurance product.

of the 10000 realizations, and each of the ten black lines shows a single realization. Moreover, the variables λ and y are given by $\lambda \approx 2.893$ and $y \approx 1.003$. In this case, the optimal value evolves only to around 6.7, corresponding to a terminal wealth of the insurance company of 4.2 since we are below k_2 , i.e., there is no surplus participation of the policyholder. The optimal strategy starts risky but not as risky as in the non-protected case, with around 75% of the wealth invested into the risky asset. It also decreases over time, but the reduction is smaller, and the final optimal investment percentage is at around 35% of the wealth. As in the non-protected case, the realizations in the best economic states correspond with the least risky investments and vice versa. One of the shown strategies is rather inconspicuous over most of the time but has a major change close to the final time point $T = 10$, i.e., a drop-down from $u_{9.93} = 0.2$ to $u_{9.99} = 0.00001$. Such an effect (also in the other way, i.e., an upward move) happens for several realizations when their wealth for time points t, which are close to the maturity T , is close to the second threshold k_2 since the optimal terminal wealth as a function of ξ has a small plateau at k_2 , i.e., if $\xi_t \in (\tilde{\alpha}\hat{\xi}, \xi_2^*]$. Hence, for $\xi_t \approx \tilde{\alpha}\hat{\xi}$ (resp. $\xi_t \approx \tilde{\alpha}\xi_2^*$) for t close to T, the upside (resp. downside) potential is reduced and the optimal strategy is to invest safely (resp. risky).

In Figure [8,](#page-16-1) we compare the optimal wealth before splitting it between the policyholders and the insurance company and the optimal strategy of our two participating life insurance products with mean-variance to a non-participating investment, i.e., we set $\alpha_2 = 0$, $k_0 = 0$, and $k_1 = 0$. Moreover, we set the initial wealth such that the mean of \hat{X}_T is approximately equal to (4 non-protected product,

7.59. Therefore, we get the following initial values: $x_0 =$ 4.692 protected product, 4.665 no participation. The lines

show again an average of 10000 realizations each. The figure shows that the investor makes the riskiest investment when offering a non-protected insurance product and approximately identical investments when offering one of the other two products. When offering the protected product compared to the non-participation product, it is remarkable that the investor invests slightly safer in the beginning and somewhat riskier close to maturity. This investment behavior leads to the highest wealth gain for the non-protected product since, on average, the risky asset performs better than the risk-less asset (due to $\mu > r$). For the other two products, the wealth gain is approximately equal. The protected life insurance investment strategy is relatively close to the non-participating strategy since a fixed-guarantee payment only minimally influences the optimal strategy. Then, the only remaining difference in the payoff shape is when the wealth is above k_2 ,

Figure 7: Optimal wealth and optimal strategy for the protected participating life insurance product.

which happens in our chosen parametrization either relatively late or not at all (see Figure [7\)](#page-16-0). The payoff structure for the non-protected insurance product differs highly in the values due to the reduced downside potential below k_1 .

Figure 8: Comparison of the optimal wealth (left) and strategy (right) between a (non-)protected participating life insurance product and a non-participating investment.

In Figure [9,](#page-17-0) we compare the optimal wealth and strategy for the non-protected and the protected life insurance contract when considering mean-variance resp. EU-optimization. We take the result for optimizing EU from Lin et al. [\[25\]](#page-33-8) as in Figure [4](#page-13-0) with parameter values $\tilde{\gamma} = 0.125$ and $\lambda = 2$. As in the other figures, the lines show the average of 10000 realizations, and we state the total wealth (before splitting it between the policyholders and the insurance company). As in the previous figure, we again choose the initial value such that the terminal wealth is approximately equal for all products, i.e., we take $x_0 = \begin{cases} 4 & \text{non-protected product,} \\ 4.692 & \text{protected product} \end{cases}$ for the mean-variance

optimization and $x_0 = \begin{cases} 3.31 & \text{non-protected product} \\ 3.59 & \text{protected product} \end{cases}$ for the EU-optimization. Both optimizations have in common that the insurance company offering a non-protected product invests riskier than the insurance company offering the protected product due to the reduced downside potential. When comparing the strategies, we observe that the strategies optimizing EU become riskier over time (except for the final time points). In contrast, the strategies optimizing mean-variance get less risky over time, as already discussed. Since we chose the two (not comparable) risk aversion factors such that the initial investments in the risky asset are similar, the wealth gain of the EU-optimization is higher. The decreasing portion invested into the risky asset is typical for a pre-commitment mean-variance optimization strategy (this can be seen by implementing the optimal mean-variance strategy by Zhou and Li [\[39\]](#page-33-14) when using typical ranges for the parameters), which is structurally different from EU-optimization strategies.

Figure 9: Comparison of the optimal wealth (left) and the optimal strategy (right) between a (non-)protected participating life insurance contract optimizing mean-variance and EU with Sshaped utility from Lin et al. [\[25\]](#page-33-8).

In Figure [10,](#page-18-2) we show another comparison for the optimal wealth and strategy for the nonprotected and the protected life insurance contract when considering mean-variance resp. EUoptimization. Here, we take for the EU-optimization an exponential utility function $U(x)$ $1 - e^{-\hat{\gamma}x}$, where $\hat{\gamma} = 0.5$ denotes the risk aversion of the investor. Note that of course this risk aversion parameter is not directly comparable to the risk aversion parameter from our meanvariance model. (For small risks though U is actually related to mean-variance by a Taylorexpansion.) The optimal wealth and strategy are calculated based on Korn and Trautmann [\[20\]](#page-32-22). Due to their assumptions, we change the risk-free interest rate to $r = 0$. Again, we show the average of 10000 realizations, state the total wealth (before splitting it between the policyholders and the insurance company), and choose the initial value such that the terminal wealth is approximately equal for all products, i.e., we take $x_0 = \begin{cases} 4 & \text{non-protected product,} \\ 4.77 & \text{protected product} \end{cases}$ for the mean-variance optimization and $x_0 = \begin{cases} 6.224 & \text{non-protected product} \\ 6.262 & \text{protected product} \end{cases}$ for the EU-optimization. We obtain again that the insurance company offering a non-protected product invests riskier than the insurance company offering the protected product due to the reduced downside potential, where the difference between the strategies is larger for the mean-variance optimization. In particular, we observe for exponential EU-optimization that the optimal strategy is relatively similar for

the non-protected and the protected life insurance products. Furthermore, these strategies are decreasing for longer maturities and slightly increasing for short maturities, and are closer to the mean-variance optimization than the ones discussed in Figure [9](#page-17-0) for S-shaped utility. However, the change in the strategies for exponential EU-maximization over time is less than for mean-variance optimization.

Figure 10: Comparison of the optimal wealth (left) and the optimal strategy (right) between a (non-)protected participating life insurance contract optimizing mean-variance, or EU with an exponential utility function.

5 Conclusion

In this paper, we derived explicit analytic formulas for general contracts, including participating life insurance contracts, when optimizing mean-variance in the multi-dimensional Black-Scholes market. Moreover, by showing the existence of all arising parameters, we showed the existence of the optimal solution. A numerical analysis shows that the mean-variance optimal strategy compared to the EU optimal strategy becomes more conservative the shorter the maturity is and is increased in particular in bad economic states. Future research directions include possibly generalizing this approach to other stock market models.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Appendix

A Proofs

Proof of Lemma [3.1.](#page-5-1) We show this lemma by contradiction. In particular, we assume that a \hat{u} exists, which is an optimal solution for J, but not for J. Consequently, there exists a strategy u which is better than \hat{u} for \hat{J} , i.e.,

$$
\tilde{J}(0,T,u,x_0) - \tilde{J}(0,T,\hat{u},x_0) > 0
$$

$$
\Leftrightarrow \quad \lambda \left(\mathbb{E} \left[F(0, T, u, x_0) - F(0, T, \hat{u}, x_0) \right] \right) - \gamma \left(\mathbb{E} \left[F(0, T, u, x_0)^2 - F(0, T, \hat{u}, x_0)^2 \right] \right) > 0. \tag{A.1}
$$

The next step is to define the function $G(x, y) := y - \gamma x + \gamma y^2$. We observe that G is a convex function, and it holds that

$$
G \left(\mathbb{E} \left[F(0, T, u, x_0)^2 \right], \mathbb{E} \left[F(0, T, u, x_0) \right] \right) = \mathbb{E} \left[F(0, T, u, x_0) \right] - \gamma \text{Var} \left(F(0, T, u, x_0) \right) = J(0, T, u, x_0).
$$

We note that $\frac{\partial}{\partial x} G(x, y) = -\gamma$ and $\frac{\partial}{\partial y} G(x, y) = 1 + 2\gamma y$. Then, the convexity of G implies

$$
G \left(\mathbb{E} \left[F(0, T, u, x_0)^2 \right], \mathbb{E} \left[F(0, T, u, x_0) \right] \right)
$$

$$
\geq G \left(\mathbb{E} \left[F(0, T, \hat{u}, x_0)^2 \right], \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \right)
$$

$$
+ \begin{pmatrix} -\gamma \\ 1 + 2\gamma \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \end{pmatrix} \cdot \begin{pmatrix} \mathbb{E} \left[F(0, T, u, x_0)^2 \right] - \mathbb{E} \left[F(0, T, \hat{u}, x_0)^2 \right] \\ \mathbb{E} \left[F(0, T, u, x_0) \right] - \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \end{pmatrix}
$$

$$
= G \left(\mathbb{E} \left[F(0, T, \hat{u}, x_0)^2 \right], \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \right)
$$

$$
- \gamma \left(\mathbb{E} \left[F(0, T, u, x_0)^2 \right] - \mathbb{E} \left[F(0, T, \hat{u}, x_0)^2 \right] \right) + (1 + 2\gamma \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \left(\mathbb{E} \left[F(0, T, u, x_0) \right] - \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \right) > G \left(\mathbb{E} \left[F(0, T, \hat{u}, x_0)^2 \right], \mathbb{E} \left[F(0, T, \hat{u}, x_0) \right] \right), \tag{A.2}
$$

where we used $(A.1)$ and $\lambda = 1 + 2\gamma \mathbb{E}[F(0,T,\hat{u},x_0)]$. Hence, $(A.2)$ implies that $J(0,T,u,x_0)$ $J(0, T, \hat{u}, x_0)$, i.e., \hat{u} is not optimal, which is a contradiction and implies the lemma. \Box

Proof of Proposition [3.4.](#page-7-1) We prove this proposition by showing the following two statements from which the claim follows directly: (i) If $\xi_3^* > \alpha \hat{\xi}$, then it holds that $\xi_2^* = \alpha \hat{\xi}$, and (ii) if $\xi_2^* > \tilde{\alpha}\hat{\xi}$, then it holds that $\xi_1^* = \tilde{\alpha}\hat{\xi}$.

We start with the proof of (i). Therefore, we reformulate the condition $\xi_3^* > \alpha \hat{\xi}$ to get a condition for λ . When plugging in the formulas as defined in Theorem [3.2,](#page-6-2) it holds that $\xi_3^* > \alpha \hat{\xi}$ if and only if

$$
\frac{2\gamma\alpha^2(k_0+k_1) - 2\gamma\alpha^2\sqrt{k_1^2 + k_1\left(2k_0 + \frac{\lambda}{\gamma\alpha}\right)}}{y} > -\frac{2\gamma\alpha^2(k_2 - k_1 - k_0)}{y}.
$$
 (A.3)

Hence, it follows that $y < \infty$ and $k_2 > \sqrt{k_1^2 + 2k_0k_1 + \frac{\lambda k_1}{\gamma \alpha}}$ which is again equivalent to $\frac{\lambda k_1}{\gamma \alpha}$ $k_2^2 - k_1^2 - 2k_0k_1$. Thus, we get that $\xi_3^* > \alpha \hat{\xi}$ holds if and only if $k_1 = 0$ or $\lambda < \gamma \alpha \left(\frac{k_2^2}{k_1} - k_1 - 2k_0\right)$ since $k_2 > 0$. Before showing the implication (i), let us also reformulate the equality $\xi_2^* = \alpha \hat{\xi}$. First, $\xi_2^* = \alpha \hat{\xi}$ if and only if $\tilde{\xi}_2^* \geq \alpha \hat{\xi}$ which is equivalent to:

$$
-\frac{\gamma\alpha^2(k_2-k_1)^2 - 2\gamma\alpha^2k_0(k_2-k_1) + \lambda\alpha k_1}{yk_2} \ge -\frac{2\gamma\alpha^2(k_2-k_1-k_0)}{y}.
$$

Since $y < \infty$ (otherwise [\(A.3\)](#page-19-2) would be wrong), this is equivalent to:

$$
2\gamma\alpha^2(k_0(k_2 - k_1) + k_2(k_2 - k_1 - k_0)) - \gamma\alpha^2(k_2^2 - 2k_1k_2 + k_1^2) \ge \lambda\alpha k_1.
$$

This again is equivalent to:

$$
\lambda k_1 \leq \gamma \alpha \left(2k_0k_2 - 2k_0k_1 + 2k_2^2 - 2k_1k_2 - 2k_0k_2 - k_2^2 + 2k_1k_2 - k_1^2 \right) = \gamma \alpha \left(k_2^2 - k_1^2 - 2k_0k_1 \right).
$$

Now, if $k_1 = 0$ and if $\lambda < \gamma \alpha \left(\frac{k_2^2}{k_1} - k_1 - 2k_0\right)$ the inequality is fulfilled since $k_2 > 0$. Thus, statement (i) follows.

Second, we prove (ii). As in the proof of (i), we start by reformulating the first condition. We note that $\xi_2^* > \tilde{\alpha}\hat{\xi}$ if and only if $\tilde{\xi}_2^* > \tilde{\alpha}\hat{\xi}$ which is equivalent to:

$$
\frac{\alpha\lambda}{y} - \frac{\gamma\alpha^2(k_2 - k_1)^2 - 2\gamma\alpha^2k_0(k_2 - k_1) + \lambda\alpha k_1}{yk_2} > \frac{\tilde{\alpha}\lambda}{y} - \frac{2\gamma\alpha\tilde{\alpha}(k_2 - k_1 - k_0)}{y}.
$$

From this inequality, it follows that $y < \infty$ and

$$
\lambda(\alpha - \tilde{\alpha})k_2 - \lambda \alpha k_1 + \gamma \alpha^2 \left(-k_2^2 + 2k_1k_2 - k_1^2 + 2k_0k_2 - 2k_0k_1 \right) > 2\gamma \alpha \tilde{\alpha} \left(-k_2^2 + k_1k_2 + k_0k_2 \right).
$$

Since $\alpha - \tilde{\alpha} = \alpha_2$, this is equivalent to:

$$
\lambda(\alpha_2 k_2 - \alpha k_1) > -\gamma \alpha \left(k_2^2 (\tilde{\alpha} - \alpha_2) + 2k_1 k_2 \alpha_2 + 2k_0 k_2 \alpha_2 - \alpha k_1^2 - 2\alpha k_0 k_1 \right).
$$

As before, we also reformulate the equality $\xi_1^* = \tilde{\alpha}\hat{\xi}$. This is indeed true if $\tilde{\xi}_1^* \geq \tilde{\alpha}\hat{\xi}$. If the term under the square root in the definition of $\tilde{\xi}_1^*$ (see Theorem [3.2\)](#page-6-2) is 0, then $\tilde{\xi}_1^* \geq \tilde{\alpha}\hat{\xi}$ is fulfilled since $\frac{2\gamma\tilde{\alpha}^2 k_2}{y} \geq 0$. Now, if this term is positive, $\tilde{\xi}_1^* \geq \tilde{\alpha}\hat{\xi}$ is equivalent to:

$$
\frac{2\gamma\tilde{\alpha}^2k_2}{y} \ge \frac{2\gamma\tilde{\alpha}}{y} \sqrt{(\alpha(k_0+k_1)-\alpha_2k_2)^2 - \alpha^2k_0^2 + \frac{\lambda}{\gamma}(\alpha k_1 - \alpha_2k_2)}.
$$

Then, since $y < \infty$, this is equivalent to:

$$
\tilde{\alpha}^2 k_2^2 \ge \alpha^2 k_0^2 + 2\alpha^2 k_0 k_1 + \alpha^2 k_1^2 - 2\alpha \alpha_2 k_0 k_2 - 2\alpha \alpha_2 k_1 k_2 + \alpha_2^2 k_2^2 - \alpha^2 k_0^2 + \frac{\lambda}{\gamma} (\alpha k_1 - \alpha_2 k_2),
$$

which is again equivalent to:

$$
\lambda(\alpha_2k_2-\alpha k_1)\geq -\gamma\alpha\left(k_2^2\left(\frac{\tilde{\alpha}^2-\alpha_2^2}{\alpha}\right)+2k_1k_2\alpha_2+2k_0k_2\alpha_2-\alpha k_1^2-2\alpha k_0k_1\right).
$$

Now, we get that $\frac{\tilde{\alpha}^2 - \alpha_2^2}{\alpha} = \frac{(\tilde{\alpha} - \alpha_2)(\tilde{\alpha} + \alpha_2)}{\alpha} = \tilde{\alpha} - \alpha_2$ since $\tilde{\alpha} + \alpha_2 = \alpha$. Hence, the inequality is fulfilled if $\lambda(\alpha_2 k_2 - \alpha k_1) > -\gamma \alpha (k_2^2(\tilde{\alpha} - \alpha_2) + 2k_1 k_2 \alpha_2 + 2k_0 k_2 \alpha_2 - \alpha k_1^2 - 2\alpha k_0 k_1)$ which implies statement (ii) and finishes the proof. \Box

Proof of Proposition [3.5.](#page-7-2) Using the two properties proven in the proof of Proposition [3.4,](#page-7-1) we only have to check the boundary values $\xi_1^* = \tilde{\alpha}\hat{\xi}$ and $\xi_2^* = \alpha\hat{\xi}$ for continuity. However, this follows immediately by plugging in the values. Now, if $k_1 > 0$, we see that $\hat{X}_T > 0$ if $\xi^* \neq \xi_3^*$. If $\xi^* = \xi_3^*$, then we note that

$$
\hat{X}_T(\xi_3^*) = k_0 + k_1 + \frac{\lambda \alpha - \lambda \alpha - 2\gamma \alpha^2 k_0 + 2\gamma \alpha^2 \left(\sqrt{k_1^2 + k_1 \left(2k_0 + \frac{\lambda}{\gamma \alpha}\right)} - k_1\right)}{2\gamma \alpha^2}
$$

$$
= k_0 + k_1 - k_0 - k_1 + \sqrt{k_1^2 + k_1 \left(2k_0 + \frac{\lambda}{\gamma \alpha}\right)} = \sqrt{k_1 \left(k_1 + 2k_0 + \frac{\lambda}{\gamma \alpha}\right)}.
$$

Hence, $\hat{X}_T(\xi_3^*) > 0$ if $k_1 > 0$ and $\lambda > -2\gamma \alpha k_0 - \gamma \alpha k_1$. From the proof of Proposition [A.3,](#page-23-0) we see that $\lambda > C$ with $C \ge -2\gamma \alpha k_0$. Thus, $\hat{X}_T(\xi_3^*) > 0$ if $k_1 > 0$. \Box

Proof of Proposition [3.6.](#page-7-0) For this proof, we show that the alternative formulation of this proposition admits the same Lagrange function as our main problem given in [\(3.5\)](#page-7-3). Therefore, we first reformulate the problem as a maximization problem by replacing $\min_{u \in \mathcal{U}} \mathbb{E}[\tilde{L}(X_T)]$ with $\max_{u \in \mathcal{U}} \mathbb{E}[-\tilde{L}(X_T)].$ Then, we get for the Lagrange function with $X \geq 0$:

$$
L(X, y, \lambda) = -\tilde{L}(X) - \lambda[\alpha k_0 - \alpha(X - k_1)1_{k_1 \le X < k_2} - \tilde{\alpha}(X - k_2 + \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1))1_{X \ge k_2}] - y\xi X
$$

$$
= \begin{cases}\n-\gamma \alpha^{2} k_{0}^{2} - \lambda \alpha k_{0} - y\xi X, & \text{if } X \in [0, k_{1}), \\
-\gamma \alpha^{2} (X - k_{1} - k_{0})^{2} - \lambda \alpha k_{0} + \lambda \alpha (X - k_{1}) - y\xi X, & \text{if } X \in [k_{1}, k_{2}), \\
-\gamma \tilde{\alpha}^{2} (X - k_{2} + \frac{\alpha}{\tilde{\alpha}} (k_{2} - k_{1} - k_{0}))^{2} - \lambda \alpha k_{0} \\
+ \lambda \tilde{\alpha} (X - k_{2} + \frac{\alpha}{\tilde{\alpha}} (k_{2} - k_{1})) - y\xi X, & \text{if } X \in [k_{2}, \infty), \\
-\gamma \alpha^{2} k_{0}^{2} - \lambda \alpha k_{0} - y\xi X, & \text{if } X \in [0, k_{1}), \\
-\gamma \alpha^{2} (X - k_{1} - k_{0})^{2} + \lambda \alpha (X - k_{1} - k_{0}) - y\xi X, & \text{if } X \in [k_{1}, k_{2}), \\
-\gamma (\tilde{\alpha} (X - k_{2}) + \alpha (k_{2} - k_{1} - k_{0}))^{2} \\
+ \lambda (\tilde{\alpha} (X - k_{2}) + \alpha (k_{2} - k_{1} - k_{0})) - y\xi X, & \text{if } X \in [k_{2}, \infty).\n\end{cases}
$$

By comparing the terms with (3.5) , we observe that the Lagrange functions are identical which gives us the proof. \Box

Proof of Theorem [3.8.](#page-11-1) From [\(3.1\)](#page-5-2) and [\(3.2\)](#page-5-3), we conclude using Itô's formula that $\xi_t \hat{X}_t$ is a martingale. Hence, we get using Theorem [3.2:](#page-6-2)

$$
\hat{X}_t = \mathbb{E}\left[\frac{\xi_T}{\xi_t}\hat{X}_T\middle|\mathcal{F}_t\right]
$$
\n
$$
= \left(k_2 + \frac{\lambda}{2\gamma\tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)\right) \mathbb{E}\left[\frac{\xi_T}{\xi_t}\mathbb{1}_{\xi_T \leq \xi_1^*}\middle|\mathcal{F}_t\right] - \frac{y}{2\gamma\tilde{\alpha}^2}\mathbb{E}\left[\frac{\xi_T^2}{\xi_t}\mathbb{1}_{\xi_T \leq \xi_1^*}\middle|\mathcal{F}_t\right]
$$
\n
$$
+ k_2 \mathbb{E}\left[\frac{\xi_T}{\xi_t}\mathbb{1}_{\tilde{\alpha}\hat{\xi} < \xi_T \leq \xi_2^*}\middle|\mathcal{F}_t\right] + \left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha}\right) \mathbb{E}\left[\frac{\xi_T}{\xi_t}\mathbb{1}_{\alpha\hat{\xi} < \xi_T \leq \xi_2^*}\middle|\mathcal{F}_t\right]
$$
\n
$$
- \frac{y}{2\gamma\alpha^2}\mathbb{E}\left[\frac{\xi_T^2}{\xi_t}\mathbb{1}_{\alpha\hat{\xi} < \xi_T \leq \xi_2^*}\middle|\mathcal{F}_t\right].
$$
\n(A.4)

Now, the claim for \hat{X}_t follows from the formula for the conditional expectation of log-normal distributions (see [\(B.1\)](#page-31-0)) using that it holds conditionally on \mathcal{F}_t :

$$
\frac{\xi_T}{\xi_t} \sim \mathcal{LN}\left(-\int_t^T r_s + \frac{\|\kappa_s\|^2}{2} \mathrm{d}s, \int_t^T \|\kappa_s\|^2 \mathrm{d}s\right),
$$
\n
$$
\frac{\xi_T^2}{\xi_t} \sim \mathcal{LN}\left(\ln \xi_t - 2\int_t^T r_s + \frac{\|\kappa_s\|^2}{2} \mathrm{d}s, 4\int_t^T \|\kappa_s\|^2 \mathrm{d}s\right),
$$

where LN denotes a log-normal distribution.

The next step is to calculate the volatility process of \hat{X}_t , i.e., the term before the dW_t in its SDE, denoted by $\sigma_{\hat{X}_t}$ using [\(3.2\)](#page-5-3). There, we get $\sigma_{\hat{X}_t} = (-\kappa_t^T \xi_t)$ $\partial \hat{X}_t$ $\frac{\partial \Lambda_t}{\partial \xi_t} = \kappa_t^T v_t$, where we used that $\partial d_1(x)$ $\frac{d_1(x)}{\partial \xi_t} = \frac{\partial d_2(x)}{\partial \xi_t}$ $rac{d_2(x)}{\partial \xi_t} = \frac{1}{-\varepsilon \sqrt{\int_0^T} }$ $-\xi_t\sqrt{\int_t^T\left\|\kappa_s\right\|^2\mathrm{d}s}$. Now, we get the optimal strategy \hat{u}_t by comparing the

volatility processes from \hat{X}_t which is $\sigma_{\hat{X}_t} = \hat{X}_t \hat{u}^T \sigma_t$. Hence, the claim follows. \Box

A.1 Proposition [A.1](#page-21-0)

Since the equation $\mathbb{E}[\xi_T \hat{X}_T(y)] = \xi_0 x_0$ depends on λ and $\lambda = 1 + 2\gamma \mathbb{E}[F(0,T,\hat{u},x_0)]$ depends on y , we have to solve these two equations together. We show that these variables always exist in the following Proposition [A.1.](#page-21-0)

Proposition A.1. There always exists a solution for λ and y and an equation system that can be numerically solved to determine them.

We split this proof into two lemmas. In the first Lemma [A.2,](#page-22-0) we give the equation system to solve for the two parameters y and λ , where we added y and λ as a superscript to help the reader follow the interdependence. In the second Lemma $A.3$, we show that these parameters exist:

Lemma A.2. The variables y and λ are the solution of the following equation system:

$$
x_0 = \left(k_2 + \frac{\lambda}{2\gamma\tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)\right)e^{-\int_0^T r_s ds} \Phi\left(d_1\left(\xi_1^{*,y,\lambda}, 0\right)\right) - \frac{y}{2\gamma\tilde{\alpha}^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \Phi\left(d_2\left(\xi_1^{*,y,\lambda}, 0\right)\right) + k_2 e^{-\int_0^T r_s ds} \left(\Phi\left(d_1\left(\xi_2^{*,y,\lambda}, 0\right)\right) - \Phi\left(d_1\left(\tilde{\alpha}\xi^{y,\lambda}, 0\right)\right)\right) + \left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha}\right)e^{-\int_0^T r_s ds} \left(\Phi\left(d_1\left(\xi_3^{*,y,\lambda}, 0\right)\right) - \Phi\left(d_1\left(\alpha\xi^{y,\lambda}, 0\right)\right)\right) - \frac{y}{2\gamma\alpha^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \left(\Phi\left(d_2\left(\xi_3^{*,y,\lambda}, 0\right)\right) - \Phi\left(d_2\left(\alpha\xi^{y,\lambda}, 0\right)\right)\right), \lambda = 1 - 2\gamma\alpha k_0 + 2\gamma\alpha k_0 \left(\Phi\left(d_0\left(\xi_3^{*,y,\lambda}, 0\right)\right) - \Phi\left(d_0\left(\alpha\xi^{y,\lambda}, 0\right)\right) + \Phi\left(d_0\left(\xi_2^{*,y,\lambda}, 0\right)\right)\right) - \Phi\left(d_0\left(\tilde{\alpha}\xi^{y,\lambda}, 0\right)\right) + \Phi\left(d_0\left(\xi_1^{*,y,\lambda}, 0\right)\right) + \lambda\left(\Phi\left(d_0\left(\xi_3^{*,y,\lambda}, 0\right)\right) - \Phi\left(d_0\left(\alpha\xi^{y,\lambda}, 0\right)\right) + \Phi\left(d_0\left(\xi_1^{*,y,\lambda}, 0\right)\right)\right) - y\left(\frac{1}{\alpha}\Phi\left(d_1\left(\xi_3^{*,y,\lambda}, 0\right)\right) - \frac{1}{\alpha}\Phi\left(d_1\left(\alpha\xi^{y,\lambda}, 0\right)\right) + \frac{1}{\tilde
$$

with

$$
d_0(x,t) = \frac{\ln x - \ln \xi_t + \int_t^T r_s + \frac{\|\kappa_s\|^2}{2} ds}{\sqrt{\int_t^T \|\kappa_s\|^2 ds}}.
$$

The values $\hat{\xi}^{y,\lambda}, \xi_1^{*,y,\lambda}$ $\zeta_1^{*,y,\lambda}, \zeta_2^{*,y,\lambda}$ $x_2^{*,y,\lambda}$, and $\xi_3^{*,y,\lambda}$ $3^{*,y,\lambda}$ are defined as in Theorem [3.2](#page-6-2) and the functions d_1 and d_2 are defined as in Theorem [3.8.](#page-11-1)

Proof. The first formula follows from the definition $\mathbb{E}[\xi_T \hat{X}_T(y)] = \xi_0 x_0 = x_0$ and [\(A.4\)](#page-21-1) for $t = 0$. The second formula follows from the definition $\lambda = 1 + 2\gamma \mathbb{E} [F(0, T, \hat{u}, x_0)]$ and the following properties for arbitrary $0 \le a < b < \infty$:

$$
\hat{X}_T = \left(k_2 + \frac{\lambda \tilde{\alpha} - y\xi_T}{2\gamma \tilde{\alpha}^2} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)\right) \mathbb{1}_{\xi_T \in (0,\xi_1^*]} + k_2 \mathbb{1}_{\xi_T \in [\tilde{\alpha}\hat{\xi},\xi_2^*]} \n+ \left(k_0 + k_1 + \frac{\lambda \alpha - y\xi_T}{2\gamma \alpha^2}\right) \mathbb{1}_{\xi_T \in [\alpha\hat{\xi},\xi_3^*]}, \n\ln \xi_T \sim \mathcal{N}\left(-\int_0^T r_s + \frac{\|\kappa_s\|^2}{2} \mathrm{d}s, \int_0^T \|\kappa_s\|^2 \mathrm{d}s\right), \n\mathbb{E}[\xi_T \mathbb{1}_{\xi_T \in [a,b]}] = \Phi\left(d_1(b,0)\right) - \Phi\left(d_1(a,0)\right), \n\mathbb{E}[\mathbb{1}_{\xi_T \in [a,b]}] = \mathbb{P}(\xi_T \in [a,b]) - \Phi\left(d_2(b,0)\right) - \Phi\left(d_2(a,0)\right)
$$
\n(A.6)

$$
\mathbb{E}[\mathbb{1}_{\xi_T \in [a,b]}] = \mathbb{P}(\xi_T \in [a,b]) = \Phi(d_0(b,0)) - \Phi(d_0(a,0)).
$$
\n(A.6)

Moreover, we conclude from [\(3.4\)](#page-6-3) (combined with Proposition [3.5\)](#page-7-2) that $\mathbb{1}_{\hat{X}_T \geq k_2} = \mathbb{1}_{\xi_T \in (0,\xi_1^*]}$ + $\mathbb{1}_{\xi_T \in (\tilde{\alpha}\hat{\xi},\xi_2^*]}$, $\mathbb{1}_{\hat{X}_T \in [k_1,k_2)} = \mathbb{1}_{\xi_T \in (\alpha\hat{\xi},\xi_3^*]}$, and $\mathbb{1}_{\hat{X}_T < k_1} = \mathbb{1}_{\hat{X}_T = 0} = \mathbb{1}_{\xi_T \notin (0,\xi_1^*] \cup (\tilde{\alpha}\hat{\xi},\xi_2^*] \cup (\alpha\hat{\xi},\xi_3^*]}$ Indeed, we get:

$$
\mathbb{E}\left[F(0,T,\hat{u},x_0)\right] = \mathbb{E}\left[-\alpha k_0 1\!\!1_{\hat{X}_T < k_1} + \alpha(\hat{X}_T - k_1 - k_0)1\!\!1_{\hat{X}_T \in [k_1,k_2)} + \tilde{\alpha}(\hat{X}_T - k_2)1\!\!1_{\hat{X}_T \ge k_2}\right]
$$

+
$$
\mathbb{E}\left[\alpha(k_2 - k_1 - k_0)\mathbb{1}_{\hat{X}_T \ge k_2}\right]
$$

\n= $-\alpha k_0 \mathbb{P}(\hat{X}_T < k_1) + \alpha \mathbb{E}[(\hat{X}_T - k_1 - k_0)\mathbb{1}_{\xi_T \in (\alpha \hat{\xi}, \xi_3^*]}]$
\n+ $\mathbb{E}\left[\left(\tilde{\alpha}(\hat{X}_T - k_2) + \alpha(k_2 - k_1 - k_0)\right)\left(\mathbb{1}_{\xi_T \in (0, \xi_1^*]} + \mathbb{1}_{\xi_T \in (\tilde{\alpha} \hat{\xi}, \xi_2^*)}\right)\right]$
\n= $-\alpha k_0 + \alpha k_0 \left(\mathbb{P}(\xi_T \in (0, \xi_1^*]) + \mathbb{P}(\xi_T \in (\tilde{\alpha} \hat{\xi}, \xi_2^*]) + \mathbb{P}(\xi_T \in (\alpha \hat{\xi}, \xi_3^*])\right)$
\n+ $\frac{\lambda}{2\gamma} \mathbb{P}(\xi_T \in (\alpha \hat{\xi}, \xi_3^*]) - \frac{y}{2\gamma \alpha} \mathbb{E}\left[\xi_T \mathbb{1}_{\xi_T \in (\alpha \hat{\xi}, \xi_3^*]}\right] + \frac{\lambda}{2\gamma} \mathbb{P}(\xi_T \in (0, \xi_1^*))$
\n- $\frac{y}{2\gamma \tilde{\alpha}} \mathbb{E}\left[\xi_T \mathbb{1}_{\xi_T \in (0, \xi_1^*]} + \alpha(k_2 - k_1 - k_0)\mathbb{P}(\xi_T \in (\tilde{\alpha} \hat{\xi}, \xi_2^*]),$

where we used the above-discussed results in the second and (3.4) in the third equation. The claim follows with $(A.5)$ and $(A.6)$. \Box

Lemma A.3. The equation system from Lemma $A.2$ admits a solution.

Proof. For this existence proof, we make the following two definitions by writing the two equations as functions depending on y and λ :

$$
f_1(y,\lambda) := -x_0 + \left(k_2 + \frac{\lambda}{2\gamma\tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)\right)e^{-\int_0^T r_s ds} \Phi(d_1(\xi_1^*, 0))
$$

\n
$$
- \frac{y}{2\gamma\tilde{\alpha}^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_1^*, 0)) + k_2 e^{-\int_0^T r_s ds} \left(\Phi(d_1(\xi_2^*, 0)) - \Phi\left(d_1\left(\tilde{\alpha}\tilde{\xi}, 0\right)\right)\right)
$$

\n
$$
+ \left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha}\right)e^{-\int_0^T r_s ds} \left(\Phi(d_1(\xi_3^*, 0)) - \Phi\left(d_1\left(\alpha\tilde{\xi}, 0\right)\right)\right)
$$

\n
$$
- \frac{y}{2\gamma\alpha^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \left(\Phi(d_2(\xi_3^*, 0)) - \Phi\left(d_2\left(\alpha\tilde{\xi}, 0\right)\right)\right),
$$

\n
$$
f_2(y,\lambda) := 1 - 2\gamma\alpha k_0 + 2\gamma\alpha k_0 \left(\Phi(d_0(\xi_3^*, 0)) - \Phi\left(d_0\left(\alpha\tilde{\xi}, 0\right)\right) + \Phi(d_0(\xi_2^*, 0))\right)
$$

\n
$$
- \Phi\left(d_0\left(\tilde{\alpha}\tilde{\xi}, 0\right)\right) + \Phi(d_0(\xi_3^*, 0))\right)
$$

\n
$$
+ \lambda \left(-1 + \Phi(d_0(\xi_3^*, 0)) - \Phi\left(d_0\left(\alpha\tilde{\xi}, 0\right)\right) + \Phi(d_0(\xi_1^*, 0))\right)
$$

\n
$$
- y \left(\frac{1}{\alpha}\Phi(d_1(\xi_3^*, 0)) - \frac{1}{\alpha}\Phi\left(d_1\left(\alpha\tilde{\xi}, 0\right)\right) + \frac{1}{\alpha}\Phi(d_1(\xi_1^*, 0))\right)
$$

\n
$$
+ 2\gamma\alpha(k_2 - k_1 - k_0) \left(\Phi(d_0(\
$$

Note that we suppress again the dependence of $\hat{\xi}$ and $\xi_i^*, i \in \{1, 2, 3\}$, on y and λ . By definition, the equation system from Lemma [A.2](#page-22-0) is solved if there exist λ^*, y^* such that $f_1(y^*, \lambda^*) = 0$ $f_2(y^*,\lambda^*).$

To show this, we define

$$
C := \begin{cases} -2\gamma\alpha k_0 & x_0 e^{\int_0^T r_s ds} \le k_1, \\ 2\gamma\alpha (x_0 e^{\int_0^T r_s ds} - k_1 - k_0) & x_0 e^{\int_0^T r_s ds} \in (k_1, k_2], \\ 2\gamma \left(\tilde{\alpha}x_0 e^{\int_0^T r_s ds} + \alpha_2 k_2 - \alpha (k_0 + k_1)\right) & x_0 e^{\int_0^T r_s ds} > k_2, \end{cases}
$$

and $h:(C,\infty) \to \mathbb{R}_{\geq 0}$ an arbitrary continuous function with $\lim_{x\to C} h(x) = 0$ if $x_0e^{\int_0^T r_s ds} > k_1$ resp. $\liminf_{x\to C} h(x) \ge 0$ if $x_0e^{\int_0^T r_s ds} \le k_1$, and $\lim_{x\to\infty} h(x) = \infty$. Then, we make the following four statements, which are shown at the end of this proof:

$$
\lim_{y \to 0} f_1(y, \lambda) \ge 0 \text{ for all } \lambda > C,
$$
\n(A.8)

$$
\lim_{y \to \infty} f_1(y, \lambda) \le -x_0 \text{ for all } \lambda > C,
$$
\n(A.9)

$$
\liminf_{\lambda \searrow C} f_2(h(\lambda), \lambda) \ge 1,\tag{A.10}
$$

$$
\limsup_{\lambda \to \infty} f_2(h(\lambda), \lambda) \le 0. \tag{A.11}
$$

Due to the continuity of $y \mapsto f_1(y, \cdot)$ and $\lambda \mapsto f_2(h(\lambda), \lambda)$, there exists for each $\lambda > C$ a $y^*_{\lambda} \in [0, \infty)$ and for all functions h a $\lambda_h^* \in (C, \infty)$ such that $f_1(y_\lambda^*, \lambda) = 0$ and $f_2(h(\lambda_h^*), \lambda_h^*) = 0$. If there exists more than one solution, then we take y^*_{λ} as the smallest one of those (well-defined due to y being bounded from below and the continuity of f_1 and f_2). Moreover, we have the following four statements, which are also shown at the end of this proof:

$$
\lambda \mapsto y_{\lambda}^* \text{ is continuous in } \lambda \text{ on } (C, \infty), \tag{A.12}
$$

$$
\lim_{\lambda \searrow C} y_{\lambda}^* = 0 \text{ if } x_0 e^{\int_0^T r_s ds} > k_1,
$$
\n(A.13)

$$
\liminf_{\lambda \searrow C} y_{\lambda}^* \ge 0 \text{ if } x_0 e^{\int_0^T r_s ds} \le k_1,\tag{A.14}
$$

$$
\lim_{\lambda \to \infty} y_{\lambda}^* = \infty. \tag{A.15}
$$

Now, if we consider y^*_{λ} as a function of λ , it fulfills the assumptions of h. Hence, there exists a λ^* such that $f_2(y^*_{\lambda^*}, \lambda^*) = 0$. On the other hand, by the definition of $\lambda \mapsto y_\lambda$, we have that $f(y_{\lambda^*}^*, \lambda^*) = 0$, which implies the claim.

To finalize the proof, we show the remaining eight statements afterwards. But first, we note that it holds by definition (see Theorem [3.2\)](#page-6-2):

$$
0 \le \xi_1^* \le \tilde{\alpha}\hat{\xi} \le \xi_2^* \le \alpha\hat{\xi} \le \xi_3^* \le \bar{\xi}.\tag{A.16}
$$

Proof of $(A.8)$. First of all, we claim that

$$
\xi_3^* > 0 \text{ if } \lambda > C,\tag{A.17}
$$

$$
\xi_1^* > 0 \text{ if and only if } \lambda > 2\gamma\alpha(k_2 - k_1 - k_0),\tag{A.18}
$$

which we will show directly after the proof of $(A.8)$. So for now assume that $(A.17)$ and $(A.18)$ hold, and let $\lambda > C$.

By $(A.17), \xi_3^*$ $(A.17), \xi_3^*$ $y \to 0$ + ∞ if $\lambda > C$. Moreover, it follows directly from the definition that

$$
\hat{\xi} \xrightarrow{y \to 0} \begin{cases} 0, & \text{if } \lambda \le 2\gamma\alpha(k_2 - k_1 - k_0), \\ +\infty, & \text{if } \lambda > 2\gamma\alpha(k_2 - k_1 - k_0). \end{cases} \tag{A.19}
$$

Thus, if $\lambda \leq 2\gamma\alpha(k_2-k_1-k_0)$ then it holds that (i) $\Phi(d_1(\xi_3^*,0)) - \Phi(d_1(\alpha \hat{\xi},0)) \xrightarrow{y\to 0} 1$. On the other hand, if $\lambda > 2\gamma\alpha (k_2 - k_1 - k_0)$, then it holds that (ii) $\Phi(d_1(\xi_1^*, 0)) \xrightarrow{y \to 0} 1$ as $(A.18)$ entails that ξ_1^* $\frac{y\rightarrow0}{x}$ ∞ . Now, if $x_0e^{\int_0^T r_s ds} \leq k_1$, it holds that (iii) $\left(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha}\right)e^{-\int_0^T r_s ds} > x_0$ for $\lambda > C =$ $-2\gamma\alpha k_0$. Furthermore, if $x_0e^{\int_0^T r_s ds} \in (k_1, k_2]$, we deduce that (iv) $(k_0 + k_1 + \frac{\lambda}{2\gamma\alpha})e^{-\int_0^T r_s ds} > x_0$ for $\lambda > C = 2\gamma\alpha(x_0e^{\int_0^T r_s ds} - k_1 - k_0)$ and, if $x_0e^{\int_0^T r_s ds} \leq k_2$, it follows directly that (v) $\left(k_2+\frac{\lambda}{2\gamma\tilde{\alpha}}-\frac{\alpha}{\tilde{\alpha}}\right)$ $\frac{\alpha}{\tilde{\alpha}}(k_2-k_1-k_0)\Bigl)\,e^{-\int_0^Tr_s\text{d}s} \,\,>\,\, x_0 \,\,\text{ for }\,\, \lambda \,\,>\,\, 2\gamma\alpha (k_2-k_1-k_0) (\geq \,\,C). \,\,\,\, \text{ Finally, if}\,\,$ $x_0e^{\int_0^T r_s ds} > k_2$, then it holds that $C > 2\gamma\alpha(k_2-k_1-k_0)$ and (vi) $\left(k_2+\frac{\lambda}{2\gamma\tilde{\alpha}}-\frac{\alpha}{\tilde{\alpha}}\right)$ $\frac{\alpha}{\tilde{\alpha}}(k_2 - k_1 - k_0)$ $e^{-\int_0^T r_s ds} >$ x_0 for $\lambda > C = 2\gamma \left(\tilde{\alpha} x_0 e^{\int_0^T r_s ds} + \alpha_2 k_2 - \alpha (k_0 + k_1) \right)$. Note that $\tilde{\alpha} \hat{\xi} \leq \xi_2^* \leq \alpha \hat{\xi}$ (see [\(A.16\)](#page-24-2)) and hence (vii) ξ_2^* shows the same limiting behavior as $\hat{\xi}$ for $y \to 0$. Summarizing these results, we conclude for $\lambda > C$:

(a) Properties (ii), (v), and (vi) imply:

$$
\lim_{y \to 0} \left(k_2 + \frac{\lambda}{2\gamma \tilde{\alpha}} - \frac{\alpha}{\tilde{\alpha}} (k_2 - k_1 - k_0) \right) e^{-\int_0^T r_s ds} \Phi(d_1(\xi_1^*, 0))
$$
\n
$$
\begin{cases}\n> x_0 & \text{if } \lambda > 2\gamma \alpha (k_2 - k_1 - k_0) \text{ and } x_0 e^{\int_0^T r_s ds} \le k_2, \\
> x_0 & \text{if } x_0 e^{\int_0^T r_s ds} > k_2, \\
\ge 0 & \text{else.} \n\end{cases}
$$

(b) One can see directly that $\lim_{y\to 0} \frac{y}{2}$ $\frac{y}{2\gamma\tilde{\alpha}^2}e^{\int_0^T -2r_s +\left\|\kappa_s\right\|^2\mathrm{d}s}\Phi\left(d_2\left(\xi_1^*,0\right)\right)=0.$

(c) It holds that $\lim_{y\to 0} k_2 e^{-\int_0^T r_s ds} \left(\Phi(d_1(\xi_2^*,0)) - \Phi(d_1(\tilde{\alpha}_2^*,0)) \right) = 0$ due to (vii).

(d) Properties (i), (iii), and (iv) imply:

$$
\lim_{y \to 0} \left(k_0 + k_1 + \frac{\lambda}{2\gamma \alpha} \right) e^{-\int_0^T r_s ds} \left(\Phi \left(d_1 \left(\xi_3^*, 0 \right) \right) - \Phi \left(d_1 \left(\alpha \hat{\xi}, 0 \right) \right) \right)
$$
\n
$$
\begin{cases}\n> x_0 & \text{if } \lambda \in (C, 2\gamma \alpha (k_2 - k_1 - k_0)] \text{ and } x_0 e^{\int_0^T r_s ds} \le k_2, \\
\ge 0 & \text{else.} \n\end{cases}
$$

(e) One can see directly that $\lim_{y\to 0} \frac{y}{2x}$ $\frac{y}{2\gamma\alpha^2}e^{\int_0^T -2r_s +\|\kappa_s\|^2\mathrm{d}s} \left(\Phi\left(d_2\left(\xi_3^*,0\right)\right)-\Phi\left(d_2\left(\alpha\hat{\xi},0\right)\right)\right)=0.$ Thus, the claim follows. \Box

Proof of $(A.17)$. We show this equation by considering the three different cases of C : <u>Case 1:</u> $x_0e^{\int_0^T r_s ds} \le k_1$, i.e., $C = -2\gamma \alpha k_0$:

Let $\lambda > C$, i.e., there exists an $\varepsilon > 0$ such that $\lambda = C + 2\gamma\alpha\varepsilon$. First, we notice that $\xi_3^* > 0$ if $\xi_3^* := \frac{\lambda \alpha}{u}$ $\frac{\Delta \alpha}{y} + \frac{2 \gamma \alpha^2 k_0}{y}$ $\frac{\alpha^2 k_0}{y} - \frac{2\gamma \alpha^2}{y}$ \hat{y} $\left(\begin{array}{c} \end{array} \right)$ $k_1^2+k_1\left(2k_0+\frac{\lambda}{\gamma\alpha}\right)-k_1$ \setminus > 0 . Now, it holds if we plug in λ : $\tilde{\xi}_3^* =$ $2\gamma\alpha^2$ \hat{y} $\left(-k_0 + \varepsilon + k_0 + k_1 - \sqrt{k_1^2 + k_1(2k_0 - 2k_0 + 2\varepsilon)}\right) = \frac{2\gamma\alpha^2}{k_0^2}$ \hat{y} $\left(k_1+\varepsilon-\sqrt{(k_1+\varepsilon)^2-\varepsilon^2}\right).$

Since $y, \gamma, \alpha, \varepsilon > 0$ and $k_1 \geq 0$, we get that $\tilde{\xi}_3^* > 0$ if and only if $(k_1 + \varepsilon)^2 > (\sqrt{(k_1 + \varepsilon)^2 - \varepsilon^2})^2$, which is equivalent to $\varepsilon^2 > 0$. Thus, the claim, i.e., $\xi_3^* > 0$ for $\lambda > C$. $\underline{\text{Case 2: }} x_0 e^{\int_0^T r_s \text{d}s} \in (k_1, k_2], \text{ i.e., } C = 2 \gamma \alpha (x_0 e^{\int_0^T r_s \text{d}s} - k_1 - k_0)$

Let $\lambda > C$, i.e., there exists an $\varepsilon > 0$ such that $\lambda = C + 2\gamma\alpha\varepsilon$. As in the first case, we get that $\xi_3^* > 0$ if $\tilde{\xi}_3^* > 0$ (defined as in Case 1). Then, we get after plugging in λ :

$$
\tilde{\xi}_{3}^{*} = \frac{2\gamma\alpha^{2}}{y} \left(x_{0} e^{\int_{0}^{T} r_{s} ds} - k_{1} - k_{0} + \varepsilon + k_{0} + k_{1} - \sqrt{k_{1}^{2} + k_{1} \left(2k_{0} + 2x_{0} e^{\int_{0}^{T} r_{s} ds} - 2k_{1} - 2k_{0} + 2\varepsilon \right)} \right)
$$
\n
$$
= \frac{2\gamma\alpha^{2}}{y} \left(x_{0} e^{\int_{0}^{T} r_{s} ds} + \varepsilon - \sqrt{-k_{1}^{2} + 2k_{1} x_{0} e^{\int_{0}^{T} r_{s} ds} - \left(x_{0} e^{\int_{0}^{T} r_{s} ds} \right)^{2} + \left(x_{0} e^{\int_{0}^{T} r_{s} ds} \right)^{2} + 2\varepsilon k_{1}} \right)
$$
\n
$$
= \frac{2\gamma\alpha^{2}}{y} \left(x_{0} e^{\int_{0}^{T} r_{s} ds} + \varepsilon - \sqrt{-\left(x_{0} e^{\int_{0}^{T} r_{s} ds} - k_{1} \right)^{2} + \left(x_{0} e^{\int_{0}^{T} r_{s} ds} \right)^{2} + 2\varepsilon k_{1}} \right).
$$

Since $y, \gamma, \alpha, \varepsilon > 0$ and $x_0 e^{\int_0^T r_s ds} \ge 0$, we get that $\tilde{\xi}_3^* > 0$ if and only if

$$
\left(x_0 e^{\int_0^T r_s ds} + \varepsilon\right)^2 > -\left(x_0 e^{\int_0^T r_s ds} - k_1\right)^2 + \left(x_0 e^{\int_0^T r_s ds}\right)^2 + 2\varepsilon k_1.
$$

Since $\left(x_0e^{\int_0^T r_s ds} + \varepsilon\right)^2 = \left(x_0e^{\int_0^T r_s ds}\right)^2 + 2\varepsilon x_0e^{\int_0^T r_s ds} + \varepsilon^2$, this is equivalent to:

$$
0 < \left(x_0 e^{\int_0^T r_s ds} - k_1\right)^2 - 2\varepsilon k_1 + 2\varepsilon x_0 e^{\int_0^T r_s ds} + \varepsilon^2 = \left(\left(x_0 e^{\int_0^T r_s ds} - k_1\right) + \varepsilon\right)^2.
$$

Now, the claim, i.e., $\xi_3^* > 0$ for $\lambda > C$, follows.

 $\underline{\text{Case 3: }} x_0 e^{\int_0^T r_s \text{d}s} > k_2 \text{, i.e., } C = 2\gamma \left(\tilde{\alpha} x_0 e^{\int_0^T r_s \text{d}s} + \alpha_2 k_2 - \alpha (k_0 + k_1) \right)$

Then, we get immediately that $\hat{\xi} > 0$ for $\lambda > C(> 2\gamma\alpha(k_2 - k_1 - k_0))$ since $\alpha_2 + \tilde{\alpha} = \alpha$). Hence, it holds that $\xi_3^* > 0$ due to $\xi_3^* \geq \alpha \hat{\xi}$ (see [\(A.16\)](#page-24-2)). \Box

Proof of [\(A.18\)](#page-24-1). First, let $\lambda > 2\gamma\alpha (k_2 - k_1 - k_0)$, i.e., there exists an $\varepsilon > 0$ such that $\lambda =$ $2\gamma\alpha(k_2 - k_1 - k_0) + 2\gamma\tilde{\alpha}\varepsilon$. We note that $\hat{\xi} > 0$ for this λ which implies that $\xi_1^* > 0$ if and only if $\tilde{\xi}_1^*>0$. Now, if the term under the square root in the maximum in the formula of $\tilde{\xi}_1^*$ (for the definition, see Theorem [3.2\)](#page-6-2) is negative, the claim follows immediately. Hence, we assume that the term is non-negative. If we plug in λ into $\tilde{\xi}_1^*$, we get:

$$
\tilde{\xi}_1^* = \frac{2\gamma\tilde{\alpha}}{y} \left(\frac{\lambda}{2\gamma} - \alpha(k_2 - k_1 - k_0) + \tilde{\alpha}k_2 - \sqrt{(\alpha(k_0 + k_1) - \alpha_2k_2)^2 - \alpha^2k_0^2 + \frac{\lambda}{\gamma}(\alpha k_1 - \alpha_2 k_2)} \right)
$$
\n
$$
= \frac{2\gamma\tilde{\alpha}}{y} \left(\tilde{\alpha}\varepsilon + \tilde{\alpha}k_2 - \sqrt{\alpha_2^2k_2^2 + 2\alpha^2k_1k_2 - 2\alpha\alpha_2k_2^2 - \alpha^2k_1^2 + 2\alpha\tilde{\alpha}k_1\varepsilon - 2\tilde{\alpha}\alpha_2k_2\varepsilon} \right)
$$
\n
$$
= \frac{2\gamma\tilde{\alpha}^2}{y} \left(k_2 + \varepsilon - \frac{1}{\tilde{\alpha}}\sqrt{k_2^2\alpha_2(\alpha_2 - 2\alpha) + \alpha^2k_1(2k_2 - k_1) + 2\tilde{\alpha}\varepsilon(\alpha k_1 - \alpha_2k_2)} \right).
$$

Since $y, \gamma, \tilde{\alpha}, \varepsilon > 0$ and $k_2 \ge 0$, it holds that $\tilde{\xi}_1^* > 0$ if and only if

$$
k_2^2 + 2\varepsilon k_2 + \varepsilon^2 > \frac{1}{\tilde{\alpha}^2} \left(k_2^2 \alpha_2(-\tilde{\alpha} - \alpha) + \alpha^2 k_1 (2k_2 - k_1) + 2\tilde{\alpha}\varepsilon(\alpha k_1 - \alpha_2 k_2) \right).
$$

This is equivalent to:

$$
L := k_2^2 \left(1 + \frac{\alpha_2}{\tilde{\alpha}} + \frac{\alpha \alpha_2}{\tilde{\alpha}^2} \right) + 2\varepsilon \left(k_2 \left(1 + \frac{\alpha_2}{\tilde{\alpha}} \right) - k_1 \frac{\alpha}{\tilde{\alpha}} \right) + \varepsilon^2 - \frac{\alpha^2}{\tilde{\alpha}^2} k_1 (2k_2 - k_1) > 0.
$$

Now, we notice with $\alpha = \tilde{\alpha} + \alpha_2$ that

$$
1 + \frac{\alpha_2}{\tilde{\alpha}} + \frac{\alpha \alpha_2}{\tilde{\alpha}^2} = \frac{\tilde{\alpha}^2 + \alpha_2 \tilde{\alpha} + \tilde{\alpha} \alpha_2 + \alpha_2^2}{\tilde{\alpha}^2} = \frac{(\tilde{\alpha} + \alpha_2)^2}{\tilde{\alpha}^2} = \frac{\alpha^2}{\tilde{\alpha}^2}.
$$

Then, it follows with L as above since $1 + \frac{\alpha_2}{\tilde{\alpha}} = \frac{\alpha}{\tilde{\alpha}}$ $\frac{\alpha}{\tilde{\alpha}}$:

$$
\frac{\tilde{\alpha}^2}{\alpha^2}L = k_2^2 + 2\frac{\tilde{\alpha}}{\alpha}\varepsilon(k_2 - k_1) + \frac{\tilde{\alpha}^2}{\alpha^2}\varepsilon^2 - 2k_1k_2 + k_1^2
$$

= $(k_2 - k_1)^2 + 2\frac{\tilde{\alpha}}{\alpha}\varepsilon(k_2 - k_1) + \left(\frac{\tilde{\alpha}}{\alpha}\varepsilon\right)^2 = \left((k_2 - k_1) + \frac{\tilde{\alpha}}{\alpha}\varepsilon\right)^2.$

Hence, $L > 0$. Second, we note that $\hat{\xi} = 0$ (and thus also ξ_1^*) for $\lambda \leq 2\gamma\alpha(k_2 - k_1 - k_0)$ by definition. Hence, the claim $\xi_1^* > 0$ if and only if $\lambda > 2\gamma\alpha (k_2 - k_1 - k_0)$ follows. \Box

Proof of [\(A.9\)](#page-24-3). By definition, it holds that $\bar{\xi} \xrightarrow{y \to \infty} 0$. Then, we get the claim due to [\(A.16\)](#page-24-2) and $\lim_{x\to 0} \Phi(d_i(x, 0)) = 0$ for $i \in \{1, 2\}.$ \Box

Proof of [\(A.10\)](#page-24-4). Recall that h is a continuous function from $(C, \infty) \to \mathbb{R}_{\geq 0}$ with some limiting properties at C and ∞ , and we plug this function into f_2 with $y = h(\lambda)$. To start, we give a small overview of the following proof: To show $(A.10)$, we must find the limiting behavior of $\xi_1^*, \tilde{\alpha}\hat{\xi}, \xi_2^*,$

 $\alpha \hat{\xi}$, and ξ_3^* when $\lambda \searrow C$. However, this behavior heavily depends on $x_0e^{\int_0^T r_s ds}$, i.e., the different values of C and the limiting properties of h . Hence, we have to distinguish several cases. First, we differentiate between the different limiting properties for $\lambda \searrow C$ depending if $x_0 e^{\int_0^T r_s ds} > k_1$ is true. If this is true, we first analyze the limit of ξ_3^* . After that, we have to separate the cases depending if C is bigger, smaller, or equal to $2\gamma\alpha(k_2 - k_1 - k_0)$. For the equality case, we even have to separate along the limiting behavior of $\hat{\xi}$ to derive the limiting behavior for ξ_1^* and ξ_2^* . If $x_0e^{\int_0^T r_s ds} > k_1$ is not true, we only have to separate along the possible limiting values of h.

<u>Case 1:</u> $x_0e^{\int_0^T r_s ds} > k_1$, i.e., $\lim_{x \to C} h(x) = 0$: Since $\lim_{\lambda \searrow C} h(\lambda) = 0$ and $C > -2\gamma \alpha k_0$, it holds that (i) $\xi_3^* \to +\infty$ for $\lambda \searrow C$, i.e., $y =$ $h(\lambda) \xrightarrow{\lambda \searrow C} 0$. Indeed, it holds: If $x_0 e^{\int_0^T r_s ds} > k_2$, we get that $C > 2\gamma \alpha (k_2 - k_1 - k_0)$ and hence $\hat{\xi} \xrightarrow{\lambda \searrow C} +\infty$ since $y = h(\lambda) \xrightarrow{\lambda \searrow C} 0$. Now, [\(A.16\)](#page-24-2) implies that ξ_3^* $\longrightarrow^{\lambda\searrow C} +\infty$. If $x_0 e^{\int_0^T r_s ds} \leq k_2$, it holds that $C = 2\gamma\alpha \left(e^{\int_0^T r_s ds} - k_1 - k_0\right)$ since $x_0 e^{\int_0^T r_s ds} > k_1$ by assumption and we get that:

$$
\xi_3^* \ge \frac{\alpha \lambda + 2\gamma \alpha^2 \left(k_0 + k_1 - \sqrt{k_1^2 + k_1(2k_0 + \frac{\lambda}{\alpha \gamma})}\right)}{h(\lambda)} =: \frac{D}{h(\lambda)}.
$$

When calculating the limit for $\lambda \searrow C$ for D, it holds that

$$
\lim_{\lambda \to C} D = 2\gamma \alpha^2 \left(x_0 e^{\int_0^T r_s ds} - k_1 - k_0 + k_0 + k_1 - \sqrt{k_1^2 + k_1(2k_0 + 2x_0 e^{\int_0^T r_s ds} - 2k_1 - 2k_0)} \right)
$$

= $2\gamma \alpha^2 \left(x_0 e^{\int_0^T r_s ds} - \sqrt{-k_1^2 + 2k_1 x_0 e^{\int_0^T r_s ds}} \right).$

Now, we see that $\lim_{\lambda\searrow C} D>0$ if and only if $(x_0e^{\int_0^T r_s ds}-k_1)^2>0$ which is true by assumption. Thus, ξ_3^* $\longrightarrow^{\lambda\setminus C}$ + ∞ since $h(\lambda) \longrightarrow^{\lambda\setminus C} 0$, i.e., (i) is proven.

Next, if $C > 2\gamma \alpha (k_2 - k_1 - k_0)$, we get that ((ii).1) ξ_1^* $\longrightarrow^{\lambda\setminus C}$ + ∞ due to [\(A.18\)](#page-24-1) and $h(\lambda) \longrightarrow^{\lambda\setminus C} 0$ and thus also ((ii).2) $\alpha \hat{\xi}, \xi_2^*, \tilde{\alpha} \hat{\xi} \xrightarrow{\lambda \searrow C} +\infty$ due to [\(A.16\)](#page-24-2). If $C < 2\gamma \alpha (k_2 - k_1 - k_0)$, we get that $((iii).1) \alpha \hat{\xi} \stackrel{\lambda \searrow C}{\longrightarrow} 0$ due to $(A.19)$ and $y = h(\lambda) \stackrel{\lambda \searrow C}{\longrightarrow} 0$ which can be factored out. Hence, also ((iii).2) $\xi_1^*, \tilde{\alpha}\hat{\xi}, \xi_2^*$ $\longrightarrow^{\lambda\setminus C} 0$ due to [\(A.16\)](#page-24-2). The remaining case that $C = 2\gamma\alpha (k_2 - k_1 - k_0)$ needs a closer look. First, we note that then $k_2 = x_0 e^{\int_0^T r_s ds}$ and hence $k_2 > k_1$ by the assumption of Case 1. There, we have to distinguish three more cases for the limiting behavior of $\xi_1^*, \tilde{\alpha}\hat{\xi},$ ξ_2^* , and $\alpha \hat{\xi}$ depending on the behavior of $\hat{\xi}$. Note that $\hat{\xi}$ can have multiple accumulation points for $\lambda \searrow C$ (all in $[0,\infty]$) despite $\hat{\xi}$ being continuous in λ . Now, let λ_n be a sequence converging to the $\liminf_{n \to \infty}$, $\lim_{n \to \infty} f_2(h(\lambda_n), \lambda_n) = \liminf_{\lambda \searrow C} f_2(h(\lambda), \lambda)$. Note that $\lambda_n \xrightarrow{n \to \infty} C$ from above. We show [\(A.10\)](#page-24-4) then for each accumulation point separately by possibly switching to subsequences, i.e., we can assume without loss of generality that $\lim_{n\to\infty} \hat{\xi}^{\lambda_n}$ exists (in $[0,\infty]$) and have to consider the following cases:

Case 1.1: $\hat{\xi}^{\lambda_n} \xrightarrow{n \to \infty} 0$:

Then, it follows immediately that (iv) $\xi_1^{*,\lambda_n}, \xi_2^{*,\lambda_n} \xrightarrow{n \to \infty} 0$ due to [\(A.16\)](#page-24-2). Case 1.2: $\hat{\xi}^{\lambda_n} \xrightarrow{n \to \infty} +\infty$:

Here, we first note that when applying the same argument as in the proof of $(A.18)$ (with $\varepsilon = 0$) to the second term in the definition of $\tilde{\xi}_1^{*,\lambda_n}$, this second term is non-negative for $\lambda = C$, i.e., $\tilde{\xi}_1^{*,\lambda_n} \geq \tilde{\alpha} \hat{\xi}^{\lambda_n}$ when taking the limit $n \to \infty$. Then, we obtain $((v).1) \xi_1^{*,\lambda_n} \xrightarrow{n \to \infty} +\infty$. In particular, we get $((v).2) \xi_1^{*,\lambda_n}, \tilde{\alpha}\hat{\xi}^{\lambda_n}, \xi_2^{*,\lambda_n}, \alpha \hat{\xi}^{\lambda_n} \xrightarrow{n \to \infty} +\infty$ due to $(A.16)$. <u>Case 1.3:</u> $\hat{\xi}^{\lambda_n} \xrightarrow{n \to \infty} c \in (0, \infty)$:

First, since $k_2 > k_1$, we get by plugging in $\lambda = 2\gamma\alpha (k_2 - k_1 - k_0)$ into $\tilde{\xi}_2^*$ that $\tilde{\xi}_2^{*,\lambda_n} \xrightarrow{n \to \infty} +\infty$. The reason is that $y_n = h(\lambda_n) \xrightarrow{n \to \infty} 0$ and the nominator in $\tilde{\xi}_2^{*,\lambda_n}$ converges to a positive number, i.e.:

$$
\alpha \lambda k_2 - \gamma \alpha^2 (k_2 - k_1)^2 + 2\gamma \alpha^2 k_0 (k_2 - k_1) - \lambda_n \alpha k_1
$$

\n
$$
\xrightarrow{n \to \infty} 2\gamma \alpha^2 k_2 (k_2 - k_1 - k_0) - \gamma \alpha^2 (k_2 - k_1)^2 + 2\gamma \alpha^2 k_0 (k_2 - k_1) - 2\gamma \alpha^2 k_1 (k_2 - k_1 - k_0)
$$

\n
$$
= 2\gamma \alpha^2 (k_2 - k_1 - k_0) (k_2 - k_1) - \gamma \alpha^2 (k_2 - k_1) (k_2 - k_1 - 2k_0)
$$

\n
$$
= \gamma \alpha^2 (k_2 - k_1) (2k_2 - 2k_1 - 2k_0 - k_2 + k_1 + 2k_0) = \gamma \alpha^2 (k_2 - k_1)^2 > 0.
$$

Hence, ((vi).1) $\xi_2^{*,\lambda_n} \xrightarrow{n \to \infty} \alpha c$. Second, we note that $\tilde{\xi}_1^{*,\lambda_n} \geq \tilde{\alpha} \hat{\xi}^{\lambda_n}$ for $n \to \infty$ by the same argument as in Case 1.2. Thus, $((vi).2) \xi_1^{*,\lambda_n} \xrightarrow{n \to \infty} \tilde{\alpha}c$.

Summarizing, we get in all subcases of Case 1, that $\Phi(d_0(\xi_3^*,0)) \xrightarrow{\lambda \searrow C} 1$ since $\lim_{x\to\infty} \Phi(d_0(x,0))$ = 1 due to (i) and $-\Phi(d_0(\alpha \hat{\xi}^{\lambda_n},0)) + \Phi(d_0(\xi_2^{*,\lambda_n},0)) - \Phi(d_0(\tilde{\alpha} \hat{\xi}^{\lambda_n},0)) + \Phi(d_0(\xi_1^{*,\lambda_n},0)) \xrightarrow{n \to \infty} 0$ due to either (ii), (iii), (iv), (v), or (vi) depending on C. Hence, it holds that $\Phi(d_0(\xi_3^{*,\lambda_n},0))$ – $\Phi(d_0(\alpha\hat{\xi}^{\lambda_n},0)) + \Phi(d_0(\xi_2^{*,\lambda_n},0)) - \Phi(d_0(\alpha\hat{\xi}^{\lambda_n},0)) + \Phi(d_0(\xi_1^{*,\lambda_n},0)) \xrightarrow{n\to\infty} 1$ since $\lim_{x\to\infty} \Phi(d_0(x,0)) =$ 1 and $\lambda_n \xrightarrow{n \to \infty} C$ from above. In particular, we get that $\lambda(-1+\Phi(d_0(\xi_3^{*,\lambda_n},0))-\Phi(d_0(\alpha \hat{\xi}^{\lambda_n},0))+\Phi(d_0(\alpha \hat{\xi}^{\lambda_n},0))$ $\Phi(d_0(\xi_1^{*,\lambda_n},0))) + 2\gamma\alpha(k_2 - k_1 - k_0)(\Phi(d_0(\xi_2^{*,\lambda_n},0)) - \Phi(d_0(\tilde{\alpha}\hat{\xi}^{\lambda_n},0))) \xrightarrow{n\to\infty} 0$ since $C = 2\gamma\alpha(k_2 - k_1 - k_0)$ $k_1 - k_0$) and $\lambda_n \xrightarrow{n \to \infty} C$ from above. Thus, the claim follows.

<u>Case 2:</u> $x_0e^{\int_0^T r_s ds} \le k_1$, i.e., $\liminf_{x \searrow C} h(x) \ge 0$ and $C = -2\gamma \alpha k_0$:

For this case, we have to distinguish two more subcases depending on the possible limit of h. However, h can have multiple accumulation points. As before, we consider each accumulation point separately, denote by λ_n a sequence converging to the lim inf, and assume, without loss of generality by possibly switching to a subsequence, that $\lambda_n \xrightarrow{n \to \infty} C$ from above and $\lim_{n \to \infty} h(\lambda_n)$ exists (in $[0, \infty]$). This gives us the following cases:

Case 2.1: $\lim_{n\to\infty} h(\lambda_n) > 0$:

Then, we get that $\bar{\xi}^{\lambda_n} \xrightarrow{n \to \infty} 0$ since $\bar{\xi}^{\lambda_n} = \frac{\alpha(\lambda_n - C)}{h(\lambda_n)}$ $\frac{(\lambda_n - C)}{h(\lambda_n)}$ for $C = -2\gamma \alpha k_0$ which implies that $\hat{\xi}^{\lambda_n}, \xi_1^{*,\lambda_n}, \xi_2^{*,\lambda_n}, \xi_3^{*,\lambda_n} \xrightarrow{n \to \infty} 0$ using [\(A.16\)](#page-24-2). Thus, it holds that $\lim_{n \to \infty} f_2(h(\lambda_n), \lambda_n) = 1$ $2\gamma\alpha k_0 - C = 1$ using that $\lim_{x\to 0} \Phi(d_i(x, 0)) = 0$ for $i \in \{0, 1\}.$

<u>Case 2.2:</u> $\lim_{n\to\infty} h(\lambda_n) = 0$:

Here, we first rewrite
$$
f_2
$$
 into:

$$
f_2(h(\lambda_n), \lambda_n) := 1 + 2\gamma \alpha k_0 \left(-1 + \Phi \left(d_0 \left(\xi_3^{*, \lambda_n}, 0 \right) \right) - \Phi \left(d_0 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) + \Phi \left(d_0 \left(\xi_1^{*, \lambda_n}, 0 \right) \right) \right) + \lambda_n \left(-1 + \Phi \left(d_0 \left(\xi_3^{*, \lambda_n}, 0 \right) \right) - \Phi \left(d_0 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) + \Phi \left(d_0 \left(\xi_1^{*, \lambda_n}, 0 \right) \right) \right) - h(\lambda_n) \left(\frac{1}{\alpha} \Phi \left(d_1 \left(\xi_3^{*, \lambda_n}, 0 \right) \right) - \frac{1}{\alpha} \Phi \left(d_1 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) + \frac{1}{\tilde{\alpha}} \Phi \left(d_1 \left(\xi_1^{*, \lambda_n}, 0 \right) \right) \right) + 2\gamma \alpha (k_2 - k_1) \left(\Phi \left(d_0 \left(\xi_2^{*, \lambda_n}, 0 \right) \right) - \Phi \left(d_0 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) \right) = 1 + (\lambda_n - C) \left(-1 + \Phi \left(d_0 \left(\xi_3^{*, \lambda_n}, 0 \right) \right) - \Phi \left(d_0 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) + \Phi \left(d_0 \left(\xi_1^{*, \lambda_n}, 0 \right) \right) \right) - h(\lambda_n) \left(\frac{1}{\alpha} \Phi \left(d_1 \left(\xi_3^{*, \lambda_n}, 0 \right) \right) - \frac{1}{\alpha} \Phi \left(d_1 \left(\alpha \hat{\xi}^{\lambda_n}, 0 \right) \right) + \frac{1}{\tilde{\alpha}} \Phi \left(d_1 \left(\xi_1^{*, \lambda_n}, 0 \right) \right) \right) + 2\gamma \alpha (k_2 - k_1) \left(\Phi \left(d_0 \left(\xi_2^{*, \lambda_n}, 0 \right) \right) - \Phi \left(d_0 \left(\tilde{\alpha
$$

where we used that $C = -2\gamma\alpha k_0$. Now, the claim follows, i.e., $\liminf_{\lambda \searrow C} f_2(h(\lambda), \lambda) \geq 1$ since $\lim_{n\to\infty} h(\lambda_n) = 0$, $\lambda_n \xrightarrow{n\to\infty} C$ from above, $\xi_2^{*,\lambda_n} \geq \tilde{\alpha}\hat{\xi}^{\lambda_n}$, and $\Phi(\tilde{d}_0(\cdot,0))$ being nondecreasing. \Box

Proof of $(A.11)$. For this proof, we have to consider three cases depending on the limiting behavior of $\frac{\lambda}{h(\lambda)}$ for $\lambda \to \infty$. As in the proof of $(A.10)$, we have here possibly multiple accumulation points that we treat separately and denote by λ_n the sequence converging to the lim sup. Note that $\lambda_n \xrightarrow{n \to \infty} \infty$ in this case. Hence, we can assume, without loss of generality by possibly switching to a subsequence, that $\lim_{n\to\infty}\frac{\lambda_n}{h(\lambda_n)}$ $\frac{\lambda_n}{h(\lambda_n)}$ exists (in $[0,\infty]$) and consider the three cases that $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = 0$, $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = c \in (0, \infty)$, and $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = \infty$:

<u>Case 1:</u> $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = 0$:

It holds that $\bar{\xi}^{\lambda_n} \xrightarrow{n \to \infty} 0$ using $y_n = h(\lambda_n)$ and the assumption. Therefore, we also get that $\hat{\xi}^{\lambda_n}, \xi_1^{*,\lambda_n}, \xi_2^{*,\lambda_n}, \xi_3^{*,\lambda_n} \xrightarrow{n \to \infty} 0$ due to [\(A.16\)](#page-24-2). Thus, the claim follows using $\lim_{x \to 0} \Phi(d_i(x, 0)) = 0$ for $i \in \{0, 1\}$.

<u>Case 2:</u> $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = c \in (0,\infty)$:

In this case, it holds by assumption that $\hat{\xi}^{\lambda_n} \xrightarrow{n \to \infty} c$, and $\bar{\xi}^{\lambda_n}$, $\xi_3^{*,\lambda_n} \xrightarrow{n \to \infty} \alpha c$. Moreover, it holds that $\lim_{n\to\infty} \xi_2^{*,\lambda_n} \in [\tilde{\alpha}c,\alpha c]$ by $(A.16)$ and $\xi_1^{*,\lambda_n} \xrightarrow{n\to\infty} \tilde{\alpha}c$ since $\frac{\lambda_n - \sqrt{l(\lambda_n)}}{h(\lambda_n)}$ $\frac{-\sqrt{l(\lambda_n)}}{h(\lambda_n)} \xrightarrow{n \to \infty} c$ for any affine function l. Next, we note that $\Phi(d_i(ac, 0)) \in (0, 1)$ for all $a > 0$ and $i \in \{0, 1\}$. In total, we get that $\lim_{n\to\infty} -1 + \Phi(d_0(\xi_3^{*,\lambda_n},0)) - \Phi(d_0(\alpha \hat{\xi}^{\lambda_n},0)) + \Phi(d_0(\xi_1^{*,\lambda_n},0)) = -1 + \Phi(d_0(\tilde{\alpha}c,0)) < 0$ and $\lim_{n\to\infty}\frac{1}{\alpha}\Phi(d_1(\xi_3^{*,\lambda_n},0))-\frac{1}{\alpha}\Phi(d_1(\alpha\xi^{\lambda_n},0))+\frac{1}{\tilde{\alpha}}\Phi(d_1(\xi_1^{*,\lambda_n},0))=\frac{1}{\tilde{\alpha}}\Phi(d_1(\tilde{\alpha}c,0))>0.$ Thus, the claim follows since $\lambda_n, h(\lambda_n) \xrightarrow{n \to \infty} +\infty$.

<u>Case 3:</u> $\lim_{n\to\infty} \frac{\lambda_n}{h(\lambda_n)} = \infty$:

In this case, we get that $\xi_1^{*,\lambda_n} \xrightarrow{n \to \infty} \infty$ as $\lim_{n \to \infty} \frac{\lambda_n - \sqrt{l(\lambda_n)}}{h(\lambda_n)} = \infty$ for any affine function l. Hence, it holds that $\hat{\xi}^{\lambda_n}, \bar{\xi}^{\lambda_n}, \xi_2^{*,\lambda_n}, \xi_3^{*,\lambda_n} \xrightarrow{n \to \infty} \infty$ due to [\(A.16\)](#page-24-2). Thus, we get the claim using $\lim_{n\to\infty} h(\lambda_n) = \infty$, $\lim_{n\to\infty} \lambda_n = \infty$, and $\lim_{x\to\infty} \Phi(d_i(x, 0)) = 1$ for $i \in \{0, 1\}$. \Box

Proof of [\(A.12\)](#page-24-7). To show this, it is sufficient to show that (i) $\lambda \mapsto f_1(\cdot, \lambda)$ is strictly increasing, (ii) $y \mapsto f_1(y, \cdot)$ is non-increasing and (iii) f_1 is jointly continuous on $\mathbb{R}_{>0} \times (C, \infty)$. Indeed, let $\lambda, \lambda_n > C$ with $\lambda_n \xrightarrow{n \to \infty} \lambda$. Due to the existence of the zero root (see the main part of the proof) and the uniqueness of the zero root (since $\lambda \mapsto f_1(\cdot, \lambda)$ is strictly increasing), there exist then unique $y_{\lambda}^*, y_{\lambda_n}^* \in (0, \infty)$ such that $f_1(y_{\lambda}^*, \lambda) = 0 = f_1(y_{\lambda_n}^*, \lambda_n)$. Then, we have to show that $y_{\lambda_n}^* \xrightarrow{n \to \infty} y_{\lambda}^*$. Due to $y \mapsto f_1(y, \cdot)$ being non-increasing and $\lambda_n \xrightarrow{n \to \infty} \lambda$, it holds that $y_{\lambda_n}^* \in [y_{\max_{k \in \mathbb{N}} {\{\lambda,\lambda_k\}}}, y_{\min_{k \in \mathbb{N}} {\{\lambda,\lambda_k\}}}]$ (maximum and minimum exist due to the convergence of λ_n to λ). Then, there exist a subsequence $y^*_{\lambda_{n_l}}$ and a \tilde{y} such that $y^*_{\lambda_{n_l}} \xrightarrow{l \to \infty} \tilde{y}$. Moreover, we know that $f_1(y^*_{\lambda}, \lambda) = 0 = f_1(y^*_{\lambda_{n_l}}, \lambda_{n_l}) \stackrel{l \to \infty}{\longrightarrow} f_1(\tilde{y}, \lambda)$ due to the joint continuity of f_1 . Now, the uniqueness of the zero root implies that $\tilde{y} = y_{\lambda}^*$. Hence, all subsequences converge to y_{λ}^* and thus also the sequence itself. Therefore, $(A.12)$ would follow provided we can show (i) , (ii) , and (iii) from the beginning:

First, the joint continuity (i.e., property (iii)) follows directly from the definition of f_1 .

Second, we derive property (i), i.e., the strict monotonicity in λ , from the formula of \hat{X}_T (see Theorem [3.2\)](#page-6-2), which gives us the claim. Note that we add in the following paragraph a superscript to \hat{X}_T , ξ_1^* , ξ_2^* , and ξ_3^* when we take these values for a certain fixed λ .

Indeed, if for all $\omega \in \Omega$, we can show that $\hat{X}_T^{\lambda_1}(\xi_T(\omega)) \geq \hat{X}_T^{\lambda_2}(\xi_T(\omega))$ with a strict inequality for a set with positive probability for all $\lambda_1 > \lambda_2 (> C)$, then also $f_1(\cdot, \lambda_1) > f_1(\cdot, \lambda_2)$ since $f_1 = \mathbb{E}[\xi_T \hat{X}_T] - x_0$ and $\xi_T > 0$. Therefore, let $\lambda_1 > \lambda_2 (> C)$: Due to $(A.17)$ and Proposition [3.4,](#page-7-1) we conclude that for all $\lambda > C$ (and hence for λ_1 and λ_2) $\hat{X}_T \neq 0$, i.e., $\hat{X}_T(\xi_T) > 0$ for ξ_T small enough. The formula of \hat{X}_T (see Theorem [3.2\)](#page-6-2) implies that for fixed y the slope remains unchanged in each interval $(0,\xi_1^*]$, $(\tilde{\alpha}\hat{\xi},\xi_2^*]$, resp. $(\alpha\hat{\xi},\xi_3^*]$ when changing λ , but the interval boundaries change. Note that the slope (as a function of ξ_T) is strictly negative in $(0,\xi_1^*]$ and $(\alpha \hat{\xi}, \xi_3^*]$, and constant otherwise. Therefore, due to \hat{X}_T being non-increasing and a non-increasing function getting larger when being shifted to the right, it is sufficient to show that all interval boundaries do not decrease when λ increases and at least one boundary value strictly increases when λ increases:

It follows directly from the definition of \hat{X}_T that $\lim_{\xi_T \to 0} \hat{X}_T^{\lambda_1}(\xi_T) > \lim_{\xi_T \to 0} \hat{X}_T^{\lambda_2}(\xi_T)$ (> 0).

Hence, due to having the same slope and the continuity of $\hat{X}^{\lambda_1}_{T}$ (resp. $\hat{X}^{\lambda_2}_{T}$) in ξ_T on $(0,\xi_1^{*,\lambda_1}]$ (resp. $(0,\xi_1^{*,\lambda_2})$, we conclude that $\xi_1^{*,\lambda_1} > \xi_1^{*,\lambda_2}$ since $\hat{X}_T^{\lambda_1}(\xi_T = \xi_1^{*,\lambda_1}) = k_2 = \hat{X}_T^{\lambda_2}(\xi_T = \xi_1^{*,\lambda_2})$, i.e., the interval is strictly increasing in λ . Next, we observe immediately from its definition that $\hat{\xi}^{\lambda_1} \geq \hat{\xi}^{\lambda_2}$. For ξ_2^* and ξ_3^* , we show this property by proving that their derivatives with respect to λ are nonnegative. Let $\tilde{\xi}_2^*$ be defined as in Theorem [3.2](#page-6-2) and $\tilde{\xi}_3^* := \bar{\xi} - \frac{2\gamma\alpha^2}{y}$ \overline{y} $\left(\sqrt{k_1^2 + k_1\left(2k_0 + \frac{\lambda}{\gamma \alpha}\right)} - k_1\right)$ with $\bar{\xi}$ as in Theorem [3.2.](#page-6-2) Then, we get:

$$
\frac{\partial}{\partial \lambda} \tilde{\xi}_{2}^{*} = \frac{\alpha}{y} - \frac{\alpha k_{1}}{yk_{2}} = \frac{\alpha}{y} \left(1 - \frac{k_{1}}{k_{2}} \right) \ge 0
$$
\n
$$
\Rightarrow \frac{\partial}{\partial \lambda} \xi_{2}^{*} \ge 0,
$$
\n
$$
\frac{\partial}{\partial \lambda} \tilde{\xi}_{3}^{*} = \begin{cases} \frac{\alpha}{y} & \text{if } k_{1} = 0, \\ \frac{\alpha}{y} - \frac{2\gamma \alpha^{2}}{y} & \frac{\frac{1}{2} \cdot \frac{k_{1}}{\gamma \alpha}}{\sqrt{k_{1}^{2} + k_{1} (2k_{0} + \frac{\lambda}{\gamma \alpha})}} = \frac{\alpha}{y} \left(1 - \frac{k_{1}}{\sqrt{k_{1}^{2} + k_{1} (2k_{0} + \frac{\lambda}{\gamma \alpha})}} \right) & \text{if } k_{1} > 0, \\ \ge 0, & \ge \frac{\alpha}{\alpha \lambda} \xi_{3}^{*} \ge 0, \end{cases}
$$

where $\frac{\partial^-}{\partial\lambda}$ denotes the left side derivative with respect to λ . Summarizing, ξ_1^* strictly increases and ξ_2^* and ξ_3^* non-decrease in λ . Thus, property (i) follows.

Third, we prove property (ii), i.e., that f_1 is non-increasing in y. Let $\hat{X}_T(y)$ be a function of y as in Theorem [3.2.](#page-6-2) It follows directly from the definitions of $\hat{\xi}$, $\xi_1^*, \xi_2^*,$ and ξ_3^* that they are non-increasing in y for fixed λ . Now, let $y_1 < y_2$ and $\xi_T > 0$ arbitrary. Then, the claim follows if $\hat{X}_T(y_1,\xi_T) \geq \hat{X}_T(y_2,\xi_T)$. If ξ_T is in the same interval for both values y_1 and y_2 , the claim is obvious due to [\(A.7\)](#page-23-2). Since $\hat{\xi}$, ξ_1^*, ξ_2^* , and ξ_3^* are non-increasing in y and \hat{X}_T is non-increasing in ξ_T (check Remark [3.3\(](#page-6-0)c)), the claim also follows if ξ_T is in different intervals for y_1 and y_2 . \Box

Proof of [\(A.13\)](#page-24-8). For the proof of this equation, we have to consider the two cases $x_0e^{\int_0^T r_s ds} < k_2$ and $x_0e^{\int_0^T r_s ds} \ge k_2$. Note that if $x_0e^{\int_0^T r_s ds} = k_2$, we get that $C = 2\gamma\alpha(x_0e^{\int_0^T r_s ds} - k_1 - k_0)$ $2\gamma \left(\tilde{\alpha} x_0 e^{\int_0^T r_s ds} + \alpha_2 k_2 - \alpha (k_0 + k_1) \right)$. Moreover, notice that $x_0 e^{\int_0^T r_s ds} > k_1$ by assumption.

 $\frac{\text{Case 1: }}{x_0 e^{\int_0^T r_s ds}} < k_2$, i.e., $C = 2\gamma \alpha (x_0 e^{\int_0^T r_s ds} - k_1 - k_0)$:

For $\lambda_{\varepsilon} = C + 2\gamma \alpha e^{\int_0^T r_s ds} \varepsilon$, it holds that $\hat{\xi} = 0$ for ε small enough and hence also $\xi_1^* = \xi_2^* = 0$ due to [\(A.16\)](#page-24-2). However, it holds with [\(A.17\)](#page-24-0) that $\xi_3^* > 0$. Note that $\lim_{x\to 0} \Phi(d_j(x,0)) = 0$ for $j \in \{0, 1, 2\}$. Thus, f_1 reduces for all $\varepsilon > 0$ small enough to

$$
f_1(y, C + 2\gamma \alpha e^{\int_0^T r_s ds} \varepsilon) = -x_0 + \left(k_0 + k_1 + \frac{C + 2\gamma \alpha e^{\int_0^T r_s ds} \varepsilon}{2\gamma \alpha} \right) e^{-\int_0^T r_s ds} \Phi(d_1(\xi_3^*, 0))
$$

$$
- \frac{y}{2\gamma \alpha^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_3^*, 0))
$$

$$
= -x_0 + (x_0 + \varepsilon) \Phi(d_1(\xi_3^*, 0)) - \frac{y}{2\gamma \alpha^2} e^{\int_0^T -2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_3^*, 0)).
$$
 (A.20)

Now, if $\limsup_{\varepsilon\to 0} y^*_{\lambda_{\varepsilon}} > 0$, it follows that $\liminf_{\varepsilon\to 0} f_1(y^*_{\lambda_{\varepsilon}}, \lambda_{\varepsilon}) < 0$ (since $\Phi(d_1(\xi_3^*, 0)) < 1$ for all $y > 0$) which is a contradiction to [\(A.8\)](#page-23-1). Thus, the claim follows, i.e., $\lim_{\lambda \to C} y^*_{\lambda} = 0$, when taking the limit of $\varepsilon \to 0$ (and hence $\lambda \to C$) on both sides, since $y^*_{\lambda} \geq 0$. Note that for every y, the existence of a solution λ_y of $(A.20)$ was already ensured in the main part of the proof.

Case 2:
$$
x_0e^{\int_0^T r_s ds} \ge k_2
$$
, i.e., $C = 2\gamma \left(\tilde{\alpha}x_0e^{\int_0^T r_s ds} + \alpha_2k_2 - \alpha(k_0 + k_1) \right)$:
\nFor $\lambda_{\varepsilon} = C + 2\gamma \tilde{\alpha}e^{\int_0^T r_s ds} \varepsilon$, it holds that:
\n $f_1(y, \lambda_{\varepsilon}) = -x_0 + (x_0 + \varepsilon)\Phi(d_1(\xi_1^*, 0)) - \frac{y}{2\gamma \tilde{\alpha}^2} e^{\int_0^T - 2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_1^*, 0))$
\n $+ k_2e^{-\int_0^T r_s ds} \Phi(d_1(\xi_2^*, 0)) - \Phi(d_1(\tilde{\alpha}\xi, 0))\Big)$
\n $+ \left(\frac{\tilde{\alpha}}{\alpha}(x_0 + \varepsilon) + \frac{\alpha_2}{\alpha}k_2e^{-\int_0^T r_s ds}\right) \Phi(d_1(\xi_3^*, 0)) - \Phi(d_1(\alpha \hat{\xi}, 0))\right)$
\n $- \frac{y}{2\gamma \alpha^2} e^{\int_0^T - 2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_3^*, 0)) - \Phi(d_2(\alpha \hat{\xi}, 0))$
\n $\leq -x_0 + x_0 \Phi(d_1(\xi_1^*, 0)) + \Phi(d_1(\xi_2^*, 0)) - \Phi(d_1(\tilde{\alpha}\hat{\xi}, 0)) + \Phi(d_1(\xi_3^*, 0)) - \Phi(d_1(\alpha \hat{\xi}, 0))$
\n $+ \varepsilon \Phi(d_1(\xi_1^*, 0)) + \Phi(d_1(\xi_3^*, 0)) - \Phi(d_1(\alpha \hat{\xi}, 0))$
\n $- \frac{y}{2\gamma} e^{\int_0^T - 2r_s + ||\kappa_s||^2 ds} \Phi(d_2(\xi_3^*, 0)) - \frac{1}{\alpha^2} \Phi(d_2(\alpha \hat{\xi}, 0)) + \frac{1}{\tilde{\alpha}^2} \Phi(d_2(\xi_1^*, 0))$
\n $\leq -x_0 + x_0 + \varepsilon - \frac{y}{2\$

where we used that $k_2e^{-\int_0^T r_s ds} \leq x_0$, $\tilde{\alpha} + \alpha_2 = \alpha$, and $\frac{\tilde{\alpha}}{\alpha} \leq 1$ in the first inequality. Moreover, we used $(A.16)$ and $\Phi(d_1(\xi_3^*,0)) \leq 1$ for all $y > 0$ in the second inequality. Hence, the claim follows, i.e., $\lim_{\varepsilon\to 0} y_{\lambda_{\varepsilon}}^* = 0$ by the same argument as in Case 1. Note that the existence of a solution was already ensured in the main part of the proof. \Box

Proof of [\(A.14\)](#page-24-9). There is nothing to show since we already have that $y^*_{\lambda} \in [0, \infty)$. \Box

Proof of $(A.15)$. We prove this by contradiction. Therefore, we assume that there exists an $L > 0$ such that $\limsup_{\lambda \to \infty} y^*_{\lambda} \leq L$. Under this assumption, it holds that $\xi_1^* \xrightarrow{\lambda \to \infty} \infty$ since $\lambda - \sqrt{l(\lambda)} \xrightarrow{\lambda \to \infty} \infty$ for all affine functions l and y^*_{λ} being bounded. Then $(A.16)$ implies that $\hat{\xi}, \bar{\xi}, \xi_2^*, \xi_3^* \xrightarrow{\lambda \to \infty} \infty$ and we know that $\Phi(d_1(\xi_3^*, 0)) \ge \Phi(d_1(\alpha \hat{\xi}, 0))$. However, then $\lim_{\lambda \to \infty} f_1(y, \lambda) = \infty$ for all $y \leq L$ which is a contradiction to y_{λ}^* being the zero root of f_1 . \Box

Since all statements are proved, we have shown the lemma. \Box

B Additional lemma

The following lemma restates a well-known result for log-normal distributions, which is used, e.g., in deriving the pricing formula of a put or call in a Black-Scholes model. The proof is just a straightforward calculation. However, we will give it for the sake of completeness.

Lemma B.1. Let $0 \le a \le b \le +\infty$ and $X \sim \mathcal{LN}(\mu, \sigma^2)$, where \mathcal{LN} denotes a log-normal distribution. Then, it holds that:

$$
\mathbb{E}\left[X\mathbb{1}_{X\in[a,b]}\right] = e^{\mu + \frac{\sigma^2}{2}} \left(\Phi\left(\frac{\ln b - \mu - \sigma^2}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu - \sigma^2}{\sigma}\right)\right),\tag{B.1}
$$

where Φ denotes the cdf of a standard normal distribution with $\Phi(+\infty) = 1$ and $\Phi(-\infty) = 0$. The formula remains unchanged when we replace the interval $[a, b]$ by $(a, b]$, $[a, b)$, or (a, b) .

Proof. It holds with $y := \ln x$ and $z := \frac{y - \mu - \sigma^2}{\sigma^2}$ $rac{1-\sigma^2}{\sigma}$:

$$
\mathbb{E}\left[X\mathbb{1}_{X\in[a,b]}\right] = \int_{\ln a}^{\ln b} \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(y - \frac{(y-\mu)^2}{2\sigma^2}\right) dy
$$

$$
= \exp\left(-\frac{1}{2\sigma^2}\left(\mu^2 - (\mu + \sigma^2)^2\right)\right)\int_{\frac{\ln b - \mu - \sigma^2}{\sigma}}^{\frac{\ln b - \mu - \sigma^2}{\sigma}} \frac{1}{\sqrt{2\pi}}\exp\left(\frac{1}{2}z^2\right)dz
$$

$$
= \exp\left(\mu + \frac{\sigma^2}{2}\right)\left(\Phi\left(\frac{\ln b - \mu - \sigma^2}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu - \sigma^2}{\sigma}\right)\right).
$$

 \Box

References

- [1] Benjamin Avanzi, Hayden Lau, and Mogens Steffensen. Optimal reinsurance design under solvency constraints. Scandinavian Actuarial Journal, 2024(4):383–416, 2024.
- [2] Anna Rita Bacinello and Svein-Arne Persson. Design and pricing of equity-linked life insurance under stochastic interest rates. The Journal of Risk Finance, 3(2):6–21, 2002.
- [3] Suleyman Basak and Georgy Chabakauri. Dynamic mean-variance asset allocation. The Review of Financial Studies, 23(8):2970–3016, 2010.
- [4] Suleyman Basak and Alexander Shapiro. Value-at-risk-based risk management: optimal policies and asset prices. The review of financial studies, 14(2):371–405, 2001.
- [5] Tomas Bjork and Agatha Murgoci. A general theory of markovian time inconsistent stochastic control problems. Available at SSRN 1694759, 2010.
- [6] Eric Briys and François De Varenne. On the risk of insurance liabilities: debunking some common pitfalls. Journal of Risk and Insurance, pages 673–694, 1997.
- [7] An Chen, Peter Hieber, and Thai Nguyen. Constrained non-concave utility maximization: An application to life insurance contracts with guarantees. European Journal of Operational Research, 273(3):1119–1135, 2019.
- [8] An Chen, Thai Nguyen, and Mitja Stadje. Optimal investment under var-regulation and minimum insurance. Insurance: Mathematics and Economics, 79:194–209, 2018.
- [9] An Chen, Mitja Stadje, and Fangyuan Zhang. On the equivalence between value-at-risk-and expected shortfallbased risk measures in non-concave optimization. arXiv preprint arXiv:2002.02229, 2020.
- [10] An Chen, Mitja Stadje, and Fangyuan Zhang. On the equivalence between value-at-risk-and expected shortfallbased risk measures in non-concave optimization. Insurance: Mathematics and Economics, 117:114–129, 2024.
- [11] John Cochrane. Asset pricing: Revised edition. Princeton university press, 2009.
- [12] Rama Cont and Peter Tankov. Financial modelling with jump processes. Chapman and Hall/CRC, 2004.
- [13] Domenico Cuoco, Hua He, and Sergei Isaenko. Optimal dynamic trading strategies with risk limits. Operations Research, 56(2):358–368, 2008.
- [14] Min Dai, Steven Kou, Shuaijie Qian, and Xiangwei Wan. Non-concave utility maximization without the concavification principle. Available at SSRN 3422276, 2019.
- [15] Yinghui Dong, Sang Wu, Wenxin Lv, and Guojing Wang. Optimal asset allocation for participating contracts under the var and pi constraint. Scandinavian Actuarial Journal, 2020(2):84–109, 2020.
- [16] EIOPA. European insurance overview 2023. Technical report, European Insurance and Occupational Pensions Authority (EIOPA), 2023.
- [17] Nadine Gatzert and Alexander Kling. Analysis of participating life insurance contracts: A unification approach. Journal of Risk and Insurance, 74(3):547–570, 2007.
- [18] Nils H Hakansson. Capital growth and the mean-variance approach to portfolio selection. Journal of Financial and Quantitative Analysis, 6(1):517–557, 1971.
- [19] Lin He, Zongxia Liang, Yang Liu, and Ming Ma. Weighted utility optimization of the participating endowment contract. Scandinavian Actuarial Journal, 2020(7):577–613, 2020.
- [20] Ralf Korn and Siegfried Trautmann. Optimal control of option portfolios and applications: Optimale steuerung von optionsportefeuilles und anwendungen. Or-Spektrum, 21:123–146, 1999.
- [21] Holger Kraft and Mogens Steffensen. A dynamic programming approach to constrained portfolios. European Journal of Operational Research, 229(2):453–461, 2013.
- [22] Kasper Larsen. Optimal portfolio delegation when parties have different coefficients of risk aversion. Quantitative Finance, 5(5):503–512, 2005.
- [23] Yuying Li and Peter A Forsyth. A data-driven neural network approach to optimal asset allocation for target based defined contribution pension plans. Insurance: Mathematics and Economics, 86:189–204, 2019.
- [24] Zongxia Liang, Yang Liu, Ming Ma, et al. A unified formula of the optimal portfolio for piecewise hara utilities. arXiv preprint arXiv:2107.06460, 2021.
- [25] Hongcan Lin, David Saunders, and Chengguo Weng. Optimal investment strategies for participating contracts. Insurance: Mathematics and Economics, 73:137–155, 2017.
- [26] Harry Markowitz. Portfolio Selection. Journal of Finance, 7(1):77–91, March 1952.
- [27] Harry M. Markowitz. Portfolio Selection: Efficient Diversification of Investments. Yale University Press, 1959.
- [28] Francesco Menoncin and Elena Vigna. Mean–variance target-based optimisation for defined contribution pension schemes in a stochastic framework. Insurance: Mathematics and Economics, 76:172–184, 2017.
- [29] Robert C Merton. An analytic derivation of the efficient portfolio frontier. Journal of financial and quantitative analysis, 7(4):1851–1872, 1972.
- [30] Robert C Merton. Optimum consumption and portfolio rules in a continuous-time model. In Stochastic optimization models in finance, pages 621–661. Elsevier, 1975.
- [31] Hui Mi, Zuo Quan Xu, and Dongfang Yang. Optimal management of dc pension plan with inflation risk and tail var constraint. arXiv preprint arXiv:2309.01936, 2023.
- [32] Charbel Mirza and Joël Wagner. Policy characteristics and stakeholder returns in participating life insurance: which contracts can lead to a win-win? European Actuarial Journal, 8:291–320, 2018.
- [33] Thai Nguyen and Mitja Stadje. Nonconcave optimal investment with value-at-risk constraint: An application to life insurance contracts. SIAM journal on control and optimization, 58(2):895–936, 2020.
- [34] Shuaijie Qian and Chen Yang. Non-concave utility maximization with transaction costs. arXiv preprint arXiv:2307.02178, 2023.
- [35] Christian Reichlin. Utility maximization with a given pricing measure when the utility is not necessarily concave. Mathematics and Financial Economics, 7(4):531–556, 2013.
- [36] Paul A Samuelson. Lifetime portfolio selection by dynamic stochastic programming. Stochastic optimization models in finance, pages 517–524, 1975.
- [37] Hato Schmeiser and Joël Wagner. A proposal on how the regulator should set minimum interest rate guarantees in participating life insurance contracts. Journal of Risk and Insurance, 82(3):659–686, 2015.
- Yuanyuan Zhang, Xiang Li, and Sini Guo. Portfolio selection problems with markowitz's mean-variance framework: a review of literature. Fuzzy Optimization and Decision Making, 17:125–158, 2018.
- [39] Xun Yu Zhou and Duan Li. Continuous-time mean-variance portfolio selection: A stochastic lq framework. Applied Mathematics and Optimization, 42:19–33, 2000.