Variance Norms for Kernelized Anomaly Detection

Thomas Cass^{1,2}, Lukas Gonon¹, Nikita Zozoulenko^{1,*}

¹ Department of Mathematics, Imperial College London, London, UK ² Institute for Advanced Study, Princeton, USA

* Corresponding author, email: n.zozoulenko23@imperial.ac.uk

Abstract

We present a unified theory for Mahalanobis-type anomaly detection on Banach spaces, using ideas from Cameron-Martin theory applied to non-Gaussian measures. This approach leads to a basis-free, data-driven notion of anomaly distance through the so-called variance norm of a probability measure, which can be consistently estimated using empirical measures. Our framework generalizes the classical \mathbb{R}^d , functional $(L^2[0,1])^d$, and kernelized settings, including the general case of non-injective covariance operator. We prove that the variance norm depends solely on the inner product in a given Hilbert space, and hence that the kernelized Mahalanobis distance can naturally be recovered by working on reproducing kernel Hilbert spaces.

Using the variance norm, we introduce the notion of a kernelized nearest-neighbour Mahalanobis distance for semi-supervised anomaly detection. In an empirical study on 12 real-world datasets, we demonstrate that the kernelized nearest-neighbour Mahalanobis distance outperforms the traditional kernelized Mahalanobis distance for multivariate time series anomaly detection, using state-of-the-art time series kernels such as the signature, global alignment, and Volterra reservoir kernels. Moreover, we provide an initial theoretical justification of nearest-neighbour Mahalanobis distances by developing concentration inequalities in the finite-dimensional Gaussian case.

Keywords: Mahalanobis distance; covariance operator; kernel methods; semi-supervised anomaly detection; multivariate time series

Contents

1	Intr	roduction	2
	1.1	Mahalanobis distance on \mathbb{R}^d and $(L^2[0,1])^d$	3
	1.2	Overview of the unified framework	5
	1.3	Organization of the paper	6
2	The	eoretical Foundations of Variance Norms	7
	2.1	Covariance operators	7
	2.2	Extended Cameron-Martin spaces	1
	2.3	Hilbert space setting	4
	2.4	Mahalanobis distance and conformance score 1	5
	2.5	Finite-dimensional concentration inequalities 1	8
	2.6	Regularized variance norms	1
	2.7	Kernelized variance norms	4
3	Cor	nputing Variance Norms 2	5
	3.1	Hilbert Case	5
	3.2	Algorithms	7
	3.3	Banach case	0
4	Apr	olication to Multivariate Time Series 3	1
	4.1	Time series kernels 3	
		4.1.1 Static and integral class kernels	
		4.1.2 Global alignment kernel	
		4.1.3 Volterra reservoir kernel	
		4.1.4 Signature kernels	
	4.2	Experiments	
		4.2.1 Experimental setup	
		4.2.2 Hyper-parameter selection	
		4.2.3 Pre-processing	
		4.2.4 Data and results	
		4.2.5 Discussion	

1 Introduction

The field of anomaly detection encompasses various techniques to identify data that deviate significantly from expected patterns in a given dataset. Some widely used methods include the Mahalanobis distance [Mah36; DJM00], density-based local outlier factors [Bre+00], clustering methods [HXD03], nearest-neighbour distances [HKF04; VF11], angle-based outlier detection [KSZ08], isolation forests [LTZ08], support vector machines [AGA13], and neural networks [Ruf+20; Pan+21]. For a review of both classical and modern approaches to anomaly detection, we refer to [Ruf+21]. This paper concerns the semi-supervised setting, where data is only available for the non-anomalous class.

In this article we propose a novel framework for Mahalanobis-type anomaly detection on any Banach or Hilbert space, based on a generalized notion of variance norm [Sha+23] and ideas from Cameron-Martin spaces (see e.g. [Bog15; Lif12]). Our extended framework includes the classical Mahalanobis distance on \mathbb{R}^d [Mah36; DJM00], the functional Mahalanobis distance on $(L^2[0,1])^d$ [PL15; BBC20], and the kernelized Mahalanobis distance [RL01] as special cases. Notably, our formulation includes the general case of non-injective covariance operators, which is not addressed in the current literature. Using the variance norm, we derive theory and computational formulas for the Mahalanobis distance, as well as the nearest-neighbour Mahalanobis distance, in the infinite-dimensional setting. One key contribution of this paper is the introduction of a kernelized nearest-neighbour Mahalanobis distance, which we demonstrate can be efficiently computed with the same time complexity as the kernelized Mahalanobis distance.

We begin by giving a brief overview of the current literature on Mahalanobis distance, to motivate the need for a general theory.

1.1 Mahalanobis distance on \mathbb{R}^d and $(L^2[0,1])^d$

A classical approach to semi-supervised anomaly detection on \mathbb{R}^d is via the Mahalanobis distance [Mah36; DJM00]. Suppose we have a random vector X in \mathbb{R}^d with non-singular covariance matrix $\Sigma \in \mathbb{R}^{d \times d}$ and mean $\mathfrak{m} \in \mathbb{R}^d$. The Mahalanobis distance $d_M(\cdot; X)$ is defined as the covariance-adjusted distance to the mean, and can be expressed in any of the three following ways for a new sample $y \in \mathbb{R}^d$

$$d_M(y;X) := \sqrt{(y-\mathfrak{m})^T \Sigma^{-1}(y-\mathfrak{m})}$$

$$= \left\| \Sigma^{-\frac{1}{2}} (y-\mathfrak{m}) \right\|_{\mathbb{R}^d}$$

$$= \sqrt{\sum_{i=1}^d \frac{1}{\lambda_i} \langle y-\mathfrak{m}, e_i \rangle^2},$$
(1)

where $(e_n, \lambda_n)_{n=1}^N$ are the eigenvector-eigenvalue pairs of the covariance matrix Σ . Anomaly detection is performed by classifying new samples with a Mahalanobis distance greater than some fixed threshold as anomalous. This method was originally used by Mahalanobis [Mah36], but has since also been employed in a variety of fields such as chemometrics [DJM00], unmanned vehicle detection [LKK10], medicine [Wan+11], cybersecurity [Dan+20], supervised classification [XNZ08], data clustering [Bro+22], and financial market anomaly detection [Aky+22].

In the case of Gaussian data, the distribution of $d_M(y; X)$ is explicitly known, which allows for the construction of a test statistic (see e.g. [DJM00]). A popular method to extend the use of Mahalanobis distances to non-Gaussian data is through the use of *kernel learning* [CS01; Sch09], where samples are embedded into RKHSs using non-linear feature maps. The kernelized Mahalanobis distance was first studied in [RL01] by replacing the implicit dot products in (1) with inner products of the feature map, or equivalently, by positive definite kernel evaluations. The method has since been used in supervised classification [WYT07; PH09; Cha+20] and anomaly detection [Lah+17; SCZ18; Dan+20]. Recently, the Mahalanobis distance was generalized to the Hilbert space $L^2[0,1]$ in the context of functional data analysis [PL15; BBC20]. This extension uses Hilbert-Schmidt covariance operators to define functional analogues of the Mahalanobis distance. In this setting, we consider a stochastic process $(X(t))_{t\in[0,1]}$ in $L^2[0,1]$ with continuous covariance function $a(s,t) := \operatorname{Cov}[X(s), X(t)]$ and functional mean $\mathfrak{m}(t) := \operatorname{E}[X(t)] \in L^2[0,1]$. The covariance operator \mathcal{K} , defined by $\mathcal{K}f(t) := \int_0^1 a(s,t)f(s)ds$ for $f \in L^2[0,1]$, is symmetric, positive, compact, and hence diagonalizable by the spectral theorem via the eigenvector-eigenvalue pairs $(e_n, \lambda_n)_{n=1}^{\infty}$ with non-negative eigenvalues. The naive definition of the functional Mahalanobis distance d_{FM} reads

$$d_{FM}(f;X) := \|\mathcal{K}^{-\frac{1}{2}}f\|_{L^{2}[0,1]},\tag{2}$$

and the problem we face in this infinite-dimensional setting is the non-invertibility of $\mathcal{K}^{\frac{1}{2}}$. When the inverse exists, it is given by $\mathcal{K}^{-\frac{1}{2}}f = \sum_{n=1}^{\infty} \frac{1}{\sqrt{\lambda_n}} \langle e_n, f \rangle e_n$, but since \mathcal{K} is of trace class, we have that $\sum_{n=1}^{\infty} \lambda_n < \infty$. This restricts the set of elements for which the inverse is well defined. In fact, if X is a Gaussian process and f is a sample path of X, then a classical result from Gaussian probability theory states that $\mathcal{K}^{-\frac{1}{2}}f$ will almost surely not exist, see e.g. [Hai23, Proposition 4.45]. The first paper to use d_{FM} resolved this issue by approximating \mathcal{K} via its M biggest eigenvalues, where M was determined via cross-validation [PL15]. Further theoretical advances were later made to the functional theory under the assumption that \mathcal{K} is injective [BBC20], using the RKHS $\mathcal{H}(\mathcal{K}) := \mathcal{K}^{1/2}(L^2[0,1])$ to regularize d_{FM} by considering the minimization problem

$$f_{\alpha} := \underset{h \in \mathcal{H}(\mathcal{K})}{\operatorname{argmin}} \|f - h\|^{2} + \alpha \|\mathcal{K}^{-\frac{1}{2}}h\|^{2}$$
$$= (\mathcal{K} + \alpha I)^{-1}\mathcal{K}f$$
$$= \sum_{n=1}^{\infty} \frac{\lambda_{n}}{\lambda_{n} + \alpha} \langle f, e_{n} \rangle e_{n},$$

for some $\alpha > 0$. The regularized functional Mahalanobis distance is defined by replacing f with f_{α} in (2), or equivalently by considering Tikhonov regularization on the operator $\mathcal{K}^{\frac{1}{2}}$. This effectively bypasses the previous invertibility issues, allowing for a well-behaved anomaly distance on $L^2[0,1]$ with theoretical guarantees like consistency of the sample estimator, and well-understood distributional properties under Gaussian assumptions on X [BBC20].

A key limitation of the current functional Mahalanobis theory is that the sample estimator of d_{FM} is special to the $L^2[0, 1]$ setting, and essentially reverts to the finite-dimensional Euclidean theory. The procedure involves discretizing *d*-dimensional sample paths on a grid of *T* time steps, and computing the Mahalanobis distance in \mathbb{R}^{Td} [PL15; RS05]. While this method works for $L^2[0, 1]$, it fails for other Hilbert spaces since the construction relies on special properties of $L^2[0, 1]$. Our aim in this paper is to derive computational formulas and theoretical results for the Mahalanobis distance in the most general infinite-dimensional setting. The current functional approach has several limitations, including the injectivity assumption of \mathcal{K} . In practical, data-driven settings, the rank of \mathcal{K} is finite and bounded by the number of data points, making the injectivity assumption problematic. Our unified framework does not require injectivity, and overcomes these issues by showing that the sample Mahalanobis estimator arises naturally by considering Cameron-Martin spaces with respect to empirical measures. This framework encompasses non-injective, general Hilbert space, and finite-dimensional cases within a single theory, eliminating the need for separate constructions.

1.2 Overview of the unified framework

In our unified framework we work on a Banach space $(V, \|\cdot\|)$ with continuous dual denoted by V^* . Our main object of study is Borel probability measures μ on V of finite second moment, denoted $\mu \in \mathcal{M}_V$ as per Definition 2.2. For such measures μ , we are able to define the vector-valued mean

$$\mathfrak{m}:=\int_V x d\mu(x),$$

and covariance operator $\mathcal{K}: V^* \to V$

$$\mathcal{K}f := \int_V x f(x) d\mu_{\mathfrak{m}}(x),$$

through the use of Bochner integrals, where $\mu_{\mathfrak{m}}$ is the measure μ shifted by \mathfrak{m} . The fundamental object we will work with is the μ -variance norm defined by

$$\|x\|_{\mu\text{-cov}} := \sup_{\substack{\text{Cov}^{\mu}[f, f] \le 1, \\ f \in V^*}} f(x),$$
(3)

which is well-defined for all $x \in V$, but is allowed to be infinite. This measure-theoretic notion was first suggested in the original pre-print of [Sha+23], but the authors provided no formal theory for the infinite dimensional case, and the idea was subsequently reworked into a variance norm with respect to a finite sample only, without the use of probability measures or laws. Our construction presented in this paper extends the data-driven finitedimensional variance norm, to a variance norm with respect to the law μ . In particular, when μ is a Gaussian measure $||x||_{\mu-\text{cov}}$ coincides with the Cameron-Martin norm of x. Crucially, our construction relies only on the fact that μ has second finite moment, and does not use any Gaussian probability theory. In this extended setting, the Banach space Mahalanobis distance is subsequently defined as

$$d_M(x;\mu) := \|x - \mathfrak{m}\|_{\mu - \operatorname{cov}_2}$$

which is well-defined for all $x \in V$, and coincides with the classical \mathbb{R}^d , functional, and kernelized Mahalanobis distances, with the added benefit that our definition supports the case of non-injective covariance operator \mathcal{K} .

By introducing Tikhonov regularization on the operator $\mathcal{K}^{\frac{1}{2}}$ when V is a Hilbert space, we are able to define a regularized variance norm similar to the functional $L^2[0,1]$ setting [BBC20]. In our case, the α -regularized μ -variance norm with $\alpha > 0$ is given by

$$|x_{\alpha}\|_{\mu\text{-cov}} = \left\| (\mathcal{K} + \alpha I)^{-1} \mathcal{K}^{\frac{1}{2}} x \right\|$$
$$= \sum_{n=1}^{\infty} \frac{\lambda_n}{(\lambda_n + \alpha)^2} \langle e_n, x \rangle, \tag{4}$$

where $(e_n, \lambda_n)_{n=1}^{\infty}$ are the eigenvector-eigenvalue pairs of the covariance operator \mathcal{K} (the sum in (4) is a finite sum if dim $V < \infty$). We prove in Theorem 2.33 that a consistent estimator of (4) is given by $||x_{\alpha}||_{\mu^N - cov}$, where $\mu^N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$ is the empirical measure of a finite sample $\{x_1, \dots, x_N\} \subset V$ assumed to be drawn from μ . More importantly, this entails that we no longer need one theory for the covariance operator of a random process, and a different theory for the finite sample estimator. Our framework allows to use the same results, theorems, and definitions for the sample estimator and the underlying random process by considering the variance norm with respect to the empirical measure μ^N .

When μ is a Gaussian measure, the exact distribution of the Mahalanobis distance is known and is equal to that of an infinite sum of independent χ_1^2 random variables. This generalizes the work of [BBC20] to the non-injective and general Hilbert case. The expression (4) for empirical measures can be efficiently computed through an SVD decomposition of the inner product Gram matrix of the data, as detailed in Theorem 3.3. This procedure, fully outlined in Algorithms 1 to 3, has strong connections to kernel principal component analysis [SSM98]. Specifically, this framework allows us to define a kernelized nearest-neighbour Mahalanobis distance, which we show can be computed with the same time complexity as the classical kernelized Mahalanobis distance. This is $\mathcal{O}(N^3 + N^2K)$ time for fitting the model, and $\mathcal{O}(NK + NM)$ time for inference, where N is the number of data points, K is the time complexity of a single inner product evaluation, and M is the number of eigenvalues considered.

Our work was motivated by the lack of theory surrounding anomaly detection on Banach spaces, and more specifically work on anomaly detection for time series data using the signature transform [Sha+23; Aky+22; Arr+24], an object originating from the theory of rough paths [Lyo98]. Existing methods for signature-based anomaly detection have been limited to very low-dimensional time series due to the exponential $\mathcal{O}(Td^m)$ time complexity in the path dimension d when computing m-level truncated signatures of time series of length T. Our unified framework addresses this bottleneck, allowing for efficient computations of signature Mahalanobis distances in linear time with respect to d through the use of signature kernels [KO19; Sal+21a]. This improvement in time complexity enables these methods to be applied to high-dimensional time series data.

1.3 Organization of the paper

This paper is organized as follows: Section 2 presents results on the theoretical foundation of variance norms and Mahalanobis distance on Banach and Hilbert spaces. We begin by constructing the variance norm using ideas originating from the theory of Cameron-Martin spaces, an approach that classically involves Gaussian probability measures. However, our construction is done purely functional-analytically, without relying on any Gaussian assumptions. Additionally, we present some initial theoretical results for nearest-neighbour Mahalanobis distances by developing concentration inequalities in the finite-dimensional Gaussian case via Lipschitz functions on the Stiefel manifold.

In Section 3, we derive algorithms for computing the variance norm with respect to empirical measures of observed data in the Hilbert and Banach space settings, including the kernelized setting. We show that the variance norm — and hence also the Mahalanobis distance — depends solely on the choice of inner product, and hence that it can naturally be kernelized in the sense of kernel learning by working on RKHSs. The full procedure is detailed in Algorithms 1 to 3.

We conclude the paper with Section 4, where we conduct an extensive empirical study of kernelized multivariate time series semi-supervised anomaly detection using 12 real-world datasets from the multivariate UEA archive [Bag+18]. We demonstrate how the kernelized nearest-neighbor variance norm outperforms the kernelized Mahalanobis distance on both PR-AUC and ROC-AUC on average, including the linear non-kernelized case. We find that the non-kernelized anomaly distances perform well on average but are often outperformed by dynamic-time kernels such as the signature kernel, the global alignment kernel, and the Volterra reservoir kernel, which comparatively have the most first-place rankings across all datasets. Inspired by the $L^2[0, 1]$ inner product, we also experiment with a class of positive definite integral-type kernels, which can be connected to the family of linear time warping kernels. Our code is publicly available at https://github.com/nikitazozoulenko/kern el-timeseries-anomaly-detection.

2 Theoretical Foundations of Variance Norms

Throughout this section, we consider a Banach space $(V, |\cdot|)$ with its continuous dual V^* , and a Borel probability measure μ defined on V. The primary objective of this section is to develop a comprehensive theory of variance norms on Banach spaces by extending the concepts of Cameron-Martin spaces and norms to non-Gaussian measures. For the classical treatment of Gaussian measures on Banach spaces we refer to [Lif12; Bog15; Hai23].

This section is organized as follows: Section 2.1 introduces the necessary setup, assumptions on μ , and the covariance operator associated with μ . In Section 2.2, we define the Cameron-Martin space with respect to a potentially non-Gaussian measure and prove that the variance norm coincides with the extended Cameron-Martin norm. Section 2.2 presents specialized results for the Hilbert space case, expressed in terms of the eigenvectors and eigenvalues of the covariance operator. Section 2.4 defines the Mahalanobis distance and the nearest-neighbor Mahalanobis distance in terms of the variance norm. Section 2.5 presents a method for obtaining concentration inequalities for the nearest-neighbor Mahalanobis distance in the finite-dimensional Gaussian case. In Section 2.6, we introduce the concept of Tikhonov-regularized variance norms, and we prove that they can be consistently estimated using empirical measures of a sample. We also provide distributional results for the case of a Gaussian measure. Finally, Section 2.7 discusses the kernelized variance norm, obtained by letting V be a RKHS.

2.1 Covariance operators

In order to extend the classical theory beyond Gaussian measures, we first need to identify suitable conditions on the underlying measures to obtain a well-defined Cameron-Martin space. Note that we will not assume that our underlying Banach space is separable. We do this in order to consider possibly non-separable reproducing kernel Hilbert spaces in the sequel. We will consider the following class of measures: **Definition 2.1.** We say that a measure μ on a Banach space V has finite p-th moment if $\|\cdot\| \in L^p(V,\mu)$.

Definition 2.2. Let V be a Banach space. We denote by \mathcal{M}_V the set of all Borel probability measures μ of finite second moment such that the identity map $V \ni x \mapsto x \in V$ is strongly μ -measurable in the Bochner sense, see e.g. [AB07, Definition 11.36].

The assumptions in Definition 2.2 allow us to consider non-separable Banach spaces V. In particular, empirical measures of the form $\frac{1}{N}\sum_{i=1}^{N} \delta_{x_i}$ always belong to \mathcal{M}_V . On the other hand, if V were to be separable, then measurability on V would be equivalent to strong measurability, and in that case \mathcal{M}_V would be the set of all Borel probability measures on V with finite second moment.

Definition 2.3. Suppose that $\mu \in \mathcal{M}_V$ for a Banach space V. Using the Bochner integral, we define

$$\mathfrak{m} := \int_V x d\mu(x) = \mathbf{E}^{x \sim \mu} \big[x \big]$$

to be the mean of the measure μ . This integral is well defined since the identity was assumed to be strongly measurable, and it is finite since $\|\cdot\| \in L^2(V,\mu) \subset L^1(V,\mu)$. On the continuous dual V^* we define the covariance quadratic form $q: V^* \times V^* \to \mathbb{R}$ by

$$q(f,g) := \operatorname{Cov}^{\mu}[f,g]$$

= $\operatorname{E}^{x \sim \mu} \left[f(x - \mathfrak{m})g(x - \mathfrak{m}) \right]$
= $\langle i^* f, i^* g \rangle_{L^2(V,\mu_{\mathfrak{m}})},$

for $f, g \in V^*$. Here $\mu_{\mathfrak{m}}$ is the measure obtained by shifting μ by the mean \mathfrak{m} , i.e. the pushforward of $x \mapsto x + \mathfrak{m}$, and $i^* : V^* \to L^2(V, \mu_{\mathfrak{m}})$ is the inclusion map sending $f \in V^*$ to its equivalence class in $L^2(V, \mu_{\mathfrak{m}})$.

Note that the inclusion map $i^* : V^* \to L^2(V, \mu_{\mathfrak{m}})$ is not necessarily injective. The quadratic form q and the map i^* are well-defined since

$$\int_{V} |f(x-y)|^{2} d\mu(x) \leq \|f\|_{V^{*}}^{2} \int_{V} \left(\|x\|^{2} + 2\|y\| \|x\| + \|y\|^{2} \right) d\mu(x) < \infty,$$

for all $f \in V^*$ and all shifts $y \in V$. Moreover, we have that $\mu \in \mathcal{M}_V \iff \mu_y \in \mathcal{M}_V$. We observe that q defines a positive quadratic form on V^* , but that it may fail to be an inner product if we have that q(f, f) = 0 for $f \neq 0$. We will sometimes refer to the quadratic form q as the functional covariance of μ .

An alternative characterization of the quadratic form q is through the so-called covariance operator of μ . This turns out to be a natural functional-analytic object to study when we no longer have access to the Gaussian tools from the classical theory of Cameron-Martin spaces.

Definition 2.4. Let $\mu \in \mathcal{M}_V$ for a Banach space V. We define the covariance operator of μ to be the bounded linear operator $\mathcal{K}: V^* \to V$ defined via

$$\mathcal{K}f := \int_V x f(x) d\mu_{\mathfrak{m}}(x),$$

for $f \in V^*$.

Note that the map $x \mapsto xf(x)$ is strongly $\mu_{\mathfrak{m}}$ -measurable for all $f \in V^*$ since both $x \mapsto x$ and $x \mapsto f(x)$ are strongly $\mu_{\mathfrak{m}}$ -measurable, and hence their product also is. The same map is also Bochner integrable since we have the bound $||xf(x)|| \leq ||f||_{V^*} ||x||^2$, which additionally implies that \mathcal{K} indeed is a bounded linear operator.

Remark 2.5. An alternative way to define \mathcal{K} is by using the quadratic form q to first define a linear operator $\mathcal{K}: V^* \to V^{**}$ via $(\mathcal{K}f)(g) = q(f,g)$. One then realizes, through Lemma 2.6 below, that $\mathcal{K}f$ actually is an evaluation functional of the vector $\int_V xf(x)d\mu_{\mathfrak{m}}(x) \in V$.

Lemma 2.6. Let $\mu \in \mathcal{M}_V$ for some Banach space V. The covariance operator $\mathcal{K} : V^* \to V$ is the unique operator satisfying

$$q(f,g) = f(\mathcal{K}g)$$

for all $f, g \in V^*$.

Proof Suppose that \mathcal{K} is the covariance operator of μ , and fix $f, g \in V^*$. Using the Bochner integral representation of \mathcal{K} we obtain that

$$\begin{split} q(f,g) &= \int_{V} f(x)g(x)d\mu_{\mathfrak{m}}(x) = \int_{V} f(xg(x))d\mu_{\mathfrak{m}}(x) \\ &= f\bigg(\int_{V} xg(x)d\mu_{\mathfrak{m}}(x)\bigg) = f(\mathcal{K}g), \end{split}$$

where the second to last equality follows from the fact that bounded operators commute with Bochner integrals, see e.g. [AB07, Lemma 11.45].

Conversely, if $q(f,g) = f(\widetilde{\mathcal{K}}g)$ for all $f,g \in V^*$ and some operator $\widetilde{\mathcal{K}}$, then $0 = f(\widetilde{\mathcal{K}}g - \mathcal{K}g)$. Consequently $\widetilde{\mathcal{K}}g = \mathcal{K}g$ by the Hahn-Banach theorem, for each $g \in V^*$.

In the sequel we will need the fact that \mathcal{K} is a compact operator. This will be used in the Hilbert space setting to diagonalize \mathcal{K} (where \mathcal{K} will additionally be of trace class) to obtain a computable expression for the variance norm. We choose to prove the result in the Banach case for full generality.

Theorem 2.7. Suppose that $\mu \in \mathcal{M}_V$ for a Banach space V. Then the covariance operator $\mathcal{K}: V^* \to V$ is a compact operator.

Proof Suppose that f_n is a bounded sequence in V^* , say $||f_n||_{V^*} \leq 1$. By Alaoglu's Theorem [Lax14, Theorem 12.3] there exists a weak*-convergent subsequence f_{n_k} converging to some $f \in V^*$, that is $\lim_k f_{n_k} = f$ pointwise. Since $||xf_{n_k}(x)|| \leq ||x||^2$ for all $x \in V$, it follows by the Dominated Convergence Theorem for Bochner integrals [AB07, Theorem 11.46] that

$$\lim_{k} \mathcal{K} f_{n_{k}} = \int_{V} \lim_{k} x f_{n_{k}}(x) d\mu_{\mathfrak{m}}(x) = \mathcal{K} f$$

which concludes the proof that \mathcal{K} is compact.

Example 2.8 (Case $V = \mathbb{R}^d$). Let μ be a measure on \mathbb{R}^d with well-defined covariance matrix $\Sigma = \mathbb{E}^{x \sim \mu}[(x - \mathfrak{m})(x - \mathfrak{m})^T]$ and mean $\mathfrak{m} = \mathbb{E}^{x \sim \mu}[x]$. The measure μ can be viewed as the push-forward measure of some d-dimensional random vector X with covariance matrix Σ and mean \mathfrak{m} . In this finite dimensional setting, we have that $\mathbb{R}^d = V = V^*$, and

$$q(a,b) = \mathbf{E}^{x \sim \mu} \left[\langle a, x - m \rangle \langle b, x - m \rangle \right]$$
$$= \mathbf{E}^{x \sim \mu} \left[\sum_{i=1}^{n} \sum_{j=1}^{n} a_i b_j (x - m)_i (x - m)_j \right]$$
$$= \langle a, \Sigma b \rangle$$

for all $a, b \in \mathbb{R}^d$. Lemma 2.6 tells us that the covariance operator \mathcal{K} simply is the linear mapping associated with the covariance matrix Σ .

Example 2.9 (Case V = C[0,1]). Suppose that V = C[0,1] is the Banach space of continuous functions on [0,1] equipped with the sup norm. Then $V^* = \mathcal{M}[0,1]$ is the space of all finite signed Borel measures on [0,1], and each $\nu \in \mathcal{M}[0,1]$ acts on V as a continuous linear functional via the pairing

$$\langle \nu, x \rangle = \int_0^1 x(t) d\nu(t) d\nu(t)$$

Consider $\mu \in \mathcal{M}_V$. Denote the covariance function by $a(s,t) := E^{x \sim \mu_{\mathfrak{m}}} [x(t)x(s)]$ for $s, t \in [0,1]$, which is well-defined since $\mu \in \mathcal{M}_V$. We then see that the covariance operator $\mathcal{K} : V^* \to V$ is given by

$$\begin{split} (\mathcal{K}\nu)(t) &= \int_{V} x(t) \langle \nu, x \rangle d\mu_{\mathfrak{m}}(x) \\ &= \int_{V} x(t) \int_{0}^{1} x(s) d\nu(s) d\mu_{\mathfrak{m}}(x) \\ &= \int_{0}^{1} a(s,t) d\nu(s), \end{split}$$

where the change of order of integration is permitted by Fubini's theorem, since we have that $\int_{V} ||x||^2 d\mu_{\mathfrak{m}}(x) < \infty$ by assumption, where $\|\cdot\|$ in this case is the supremum norm.

The following example demonstrates how we recover the functional Mahalanobis setting in $L^2[0,1]$ [PL15; BBC20] as a special case in our framework, without any assumptions on the continuity of the covariance function $a(s,t) := E^{x \sim \mu_{\mathfrak{m}}}[x(t)x(s)]$, or injectivity of the covariance operator \mathcal{K} .

Example 2.10 (Case $V = L^2[0,1]$). Consider the case when $\mu \in \mathcal{M}_V$ and $V = L^2[0,1]$ is the Hilbert space of square integrable functions on [0,1]. Using that μ has finite second moment, one can show that $a(\cdot,t) \in L^2[0,1]$ for a.e. t. The covariance operator $\mathcal{K} : L^2[0,1] \to L^2[0,1]$ can then be written as

$$(\mathcal{K}f)(t) = \int_{V} x(t) \int_{0}^{1} f(s)x(s)dsd\mu_{\mathfrak{m}}(x)$$
$$= \int_{0}^{1} a(s,t)f(s)ds,$$

which we see is a Hilbert-Schmidt integral operator, since

$$\begin{split} \int_{0}^{1} \int_{0}^{1} |a(s,t)| ds dt &\leq \int_{0}^{1} \int_{0}^{1} \int_{V} |x(s)x(t)| d\mu_{\mathfrak{m}}(x) ds dt \\ &= \int_{V} \|x\|_{L^{1}[0,1]}^{2} d\mu_{\mathfrak{m}}(x) \\ &\leq \int_{V} \|x\|^{2} d\mu_{\mathfrak{m}}(x) < \infty, \end{split}$$

where $\|\cdot\|$ is the $L^2[0,1]$ -norm.

2.2 Extended Cameron-Martin spaces

A fundamental challenge when studying the covariance operator $\mathcal{K} : V^* \to V$ is that \mathcal{K} may be non-injective. We address this issue by essentially quotienting out the degenerate part of μ by taking the closure of V^* in the space $L^2(V, \mu_m)$ with respect to the L^2 -norm, which is intrinsically related to the quadratic form q. This is a classical approach used to study Gaussian measures on Banach spaces, and it will allow us to extend \mathcal{K} to an injective operator defined on a closed subspace of $L^2(V, \mu_m)$.

Definition 2.11. Let V be a Banach space. For $\mu \in \mathcal{M}_V$ we define the space \mathcal{R}_{μ} to be the closure of $i^*(V^*)$ in the Hilbert space $L^2(V, \mu_{\mathfrak{m}})$. Here i^* is the map sending $f \in V^*$ to its equivalence class in $L^2(V, \mu_{\mathfrak{m}})$.

The space \mathcal{R}_{μ} plays a crucial role throughout this section. As a closed subset of a Hilbert space, \mathcal{R}_{μ} inherits a Hilbert space structure under the $L^2(V, \mu_{\mathfrak{m}})$ -norm. Our goal is to extend \mathcal{K} to an operator $\mathcal{C} : \mathcal{R}_{\mu} \to V$, where the image $H_{\mu} = \mathcal{C}(\mathcal{R}_{\mu})$ will be defined as the Cameron-Martin space of μ . Furthermore, \mathcal{K} will be a symmetric compact operator and an isometry between these two spaces. This is what in the sequel will allow us to consider empirical measures of a sample, which by definition will give rise to finite-rank, and in particular non-injective, covariance operators.

Proposition 2.12. The covariance operator $\mathcal{K} : V^* \to V$ extends to a linear operator $\mathcal{C} : \mathcal{R}_{\mu} \to V$, where \mathcal{R}_{μ} is the $L^2(V, \mu_{\mathfrak{m}})$ -closure of $i^*(V^*)$, via the limit

$$\mathcal{C}k := \lim_{n \to \infty} \mathcal{K}f_n = \int_V xk(x)d\mu_{\mathfrak{m}}(x),$$

where $(i^*f_n)_{n=1}^{\infty}$ is any sequence converging to $k \in \mathcal{R}_{\mu} \subset L^2(V, \mu_{\mathfrak{m}})$.

Remark 2.13. Note that $Ci^*f = \mathcal{K}f$ for all $f \in V^*$.

Proof Let $k \in \mathcal{R}_{\mu}$. Since $i^*(V^*)$ is dense in \mathcal{R}_{μ} , there exists a sequence $f_n \in V^*$ such that $\|i^*f_n - k\|_{L^2(V,\mu_{\mathfrak{m}})} \to 0$ as $n \to \infty$. By Hölders inequality we have that

$$\begin{aligned} \left\| \mathcal{K}f_n - \int_V xk(x)d\mu_{\mathfrak{m}}(x) \right\| &\leq \int_V \|x\| \left| (k - f_n)(x) \right| d\mu_{\mathfrak{m}}(x) \\ &= \left(\int_V \|x\|^2 d\mu_{\mathfrak{m}}(x) \right)^{\frac{1}{2}} \left(\int_V \left| (k - f_n)(x) \right|^2 d\mu_{\mathfrak{m}}(x) \right)^{\frac{1}{2}}, \end{aligned}$$

which goes to 0 as $n \to \infty$. This holds for any such sequence.

We can now define the Cameron-Martin space of a general (possibly non-Gaussian) measure μ . The Cameron-Martin space will be a Hilbert space isometrically isomorphic to \mathcal{R}_{μ} , whose norm will naturally be given by the covariance-adjusted distance through the extended covariance operator.

Definition 2.14. Let V be a Banach space, and let $\mu \in \mathcal{M}_V$ with extended covariance operator $\mathcal{C} : \mathcal{R}_{\mu} \to V$. We define the Cameron-Martin space H_{μ} of μ to be the set $H_{\mu} := \mathcal{C}(R_{\mu})$.

Proposition 2.15. The operator $C : \mathcal{R}_{\mu} \to H_{\mu}$ is invertible. Hence H_{μ} is a Hilbert space under the norm

$$\|h\|_{H_{\mu}} := \|\mathcal{C}^{-1}h\|_{L^{2}(V,\mu_{\mathfrak{m}})},$$

$$\langle h, l \rangle_{H_{\mu}} := \langle \mathcal{C}^{-1}h, \mathcal{C}^{-1}l \rangle_{L^{2}(V,\mu_{\mathfrak{m}})},$$

where $h, l \in H_{\mu}$. The spaces H_{μ} and \mathcal{R}_{μ} are isometrically isomorphic via the map \mathcal{C} .

Proof We need to prove that C is injective. To this end, assume that Ck = 0 for some $k \in \mathcal{R}_{\mu}$. By definition there exists a sequence $f_n \in V^*$ such that $i^*f_n \to k$ in \mathcal{R}_{μ} . Lemma 2.6 then implies that

$$0 = \lim_{n} g(\mathcal{K}f_n) = \lim_{n} \langle i^*g, i^*f_n \rangle_{L^2(V,\mu_{\mathfrak{m}})} = \langle i^*g, k \rangle_{L^2(V,\mu_{\mathfrak{m}})}$$

for all $g \in V^*$. By continuity we obtain that $\langle g, k \rangle_{L^2(V,\mu_{\mathfrak{m}})} = 0$ for all $g \in \mathcal{R}_{\mu}$. Consequently we find that k = 0 ($\mu_{\mathfrak{m}}$ -a.e.), which shows that $\mathcal{C} : \mathcal{R}_{\mu} \to H$ is injective. The latter statement of the Theorem follows from the Hilbert space structure of $\mathcal{R}_{\mu} \subset L^2(V,\mu_{\mathfrak{m}})$ and the linearity of \mathcal{C} .

The following fundamental theorem shows that $\|\cdot\|_{\mu\text{-cov}}$ is a genuine norm on a subspace of V, and infinite otherwise. More precisely, this subspace is the Cameron-Martin space $H_{\mu} \subset V$, and the μ -variance norm coincides with the Cameron-Martin Hilbert norm when restricted to this space. We begin by formally defining the μ -variance norm.

Definition 2.16. Let $\mu \in \mathcal{M}_V$ for a Banach space V. The μ -variance norm for $x \in V$ is defined as

$$||x||_{\mu\text{-}cov} := \sup_{f \in V^*, q(f,f) \le 1} f(x).$$

Theorem 2.17. Suppose that $\mu \in \mathcal{M}_V$ for a Banach space V. Then the Cameron-Martin space of μ is characterized by

$$H_{\mu} = \{ h \in V : \|h\|_{\mu - cov} < \infty \}.$$

Furthermore, the Cameron-Martin norm $\|\cdot\|_{H_{\mu}}$ and the variance norm $\|\cdot\|_{\mu-cov}$ coincide on H_{μ} , or in other words

$$||h||_{H_{\mu}} := ||\mathcal{C}^{-1}h||_{L^{2}(V,\mu_{\mathfrak{m}})} = \sup_{f \in V^{*}, q(f,f) \le 1} f(h) =: ||h||_{\mu\text{-}cov}$$

for all $h \in H_{\mu}$.

Proof Suppose that h = Ck for some $k \in \mathcal{R}_{\mu}$. We want to show that the variance norm $\|h\|_{\mu-\text{cov}}$ is finite and equal to $\|h\|_{H_{\mu}}$. To this end, observe that

$$\begin{split} \|h\|_{\mu\text{-cov}} &= \sup_{f \in V^*, q(f, f) \le 1} f(\mathcal{C}k) \\ &= \sup_{f \in V^*, q(f, f) \le 1} \langle i^* f, k \rangle_{L^2(V, \mu_{\mathfrak{m}})} \\ &= \sup_{l \in \mathcal{R}_{\mu}, \|l\|_{L_2} \le 1} \langle l, k \rangle_{L^2(V, \mu_{\mathfrak{m}})} \\ &= \|\mathcal{C}^{-1}h\|_{L^2(V, \mu_{\mathfrak{m}})}, \end{split}$$

where the third equality follows by the fact that $i^*(V^*)$ is dense in \mathcal{R}_{μ} .

Conversely, assume that $||x||_{\mu\text{-cov}} < \infty$ for some $x \in V$. Let $T_x : V^* \to \mathbb{R}$ denote the evaluation functional $T_x g = g(x)$. For $g \in V^*$ we have the bound

$$|T_xg| = |g(x)| \le \|i^*g\|_{L^2(V,\mu_{\mathfrak{m}})} \sup_{f \in V^*, q(f,f) \le 1} f(x),$$

hence T_x extends to a linear operator $\mathcal{T}_x : \mathcal{R}_\mu \to \mathbb{R}$ by continuity. More specifically, $\mathcal{T}_x k$ for $k \in \mathcal{R}_\mu$ can be defined via $\mathcal{T}_x k := \lim_n T_x f^{(n)} = \lim_n f^{(n)}(x)$ where $f^{(n)} \in V^*$ is any sequence such that $||k - i^* f^{(n)}||_{L^2(V,\mu_m)} \to 0$. The operator norm for a general bounded operator $\mathcal{T} \in \mathcal{R}^*_\mu$ is given by

$$\begin{aligned} \|\mathcal{T}\|_{\mathcal{R}^*} &:= \sup_{k \in \mathcal{R}_{\mu}, \, \|k\|_{L^2(V,\mu_{\mathfrak{m}})} \leq 1} \mathcal{T}k \\ &= \sup_{f \in V^*, \, q(f,f) \leq 1} \mathcal{T}i^*f, \end{aligned}$$

where the last equality follows by the fact that $i^*(V^*)$ is dense in \mathcal{R}_{μ} . Restricting this to extended evaluation functionals \mathcal{T}_x we obtain that

$$\|\mathcal{T}_x\|_{\mathcal{R}^*} = \sup_{f \in V^*, q(f,f) \le 1} f(x) = \|x\|_{\mu\text{-cov}}.$$

Since $\mathcal{T}_x \in \mathcal{R}^*_{\mu}$ if and only if the operator norm is finite, we may use the fact that \mathcal{R}_{μ} is a Hilbert space to identify \mathcal{T}_x with an element of \mathcal{R}_{μ} itself, say k_x , such that $\mathcal{T}_x l = \langle l, k_x \rangle$ for all $l \in \mathcal{R}_{\mu}$. If $f \in V^*$, then

$$f(\mathcal{C}k_x - x) = \langle k_x, i^*f \rangle_{L^2(V,\mu_{\mathfrak{m}})} - f(x) = f(x) - f(x) = 0,$$

and it follows by Hahn-Banach that $Ck_x = h$. This concludes the proof.

Remark 2.18. In the above theorem we proved that $h \in H_{\mu}$ if and only if the evaluation functional T_h extends to a continuous linear functional on $\mathcal{R}_{\mu} \subset L^2(V, \mu_{\mathfrak{m}})$. This also proves that H_{μ} is a reproducing kernel Hilbert space.

2.3 Hilbert space setting

We now specialize to the case where V is a Hilbert space with inner product $\langle \cdot, \cdot \rangle$. In this setting, we have that $V = V^*$, allowing us to diagonalize the covariance operator $\mathcal{K} : V \to V$ and express the μ -variance norm in terms of the eigenvalues of \mathcal{K} , irrespective of injectivity.

Recall from Lemma 2.6 that

$$\langle g, \mathcal{K}f \rangle = q(g, f) = q(f, g) = \langle f, \mathcal{K}g \rangle$$

for all $f, g \in V$. This, combined with Theorem 2.7 and the fact that $\langle f, \mathcal{K}f \rangle \geq 0$ for all $f \in V$, implies that \mathcal{K} is a symmetric, positive, compact operator. Consequently, the spectral theorem [Lax14, Theorem 28.3] implies that there exists an orthonormal sequence of eigenvectors $(e_n)_{n=1}^{\infty}$ and corresponding non-negative eigenvalues $(\lambda_n)_{n=1}^{\infty}$ such that

$$\mathcal{K}f = \sum_{n=1}^{\infty} \lambda_n \langle f, e_n \rangle e_n, \quad \forall f \in V.$$
(5)

Our ultimate goal is to derive an explicit computational formula for the variance norm $\|\cdot\|_{H_{\mu}}$. To achieve this, we first need to characterize the Cameron-Martin space H_{μ} , as Theorem 2.17 states that this is the subspace of V where the variance norm is finite. Understanding H_{μ} will also be crucial when we later consider empirical measures constructed from observed data in Section 3.

Theorem 2.19. Let $\mu \in \mathcal{M}_V$ for a Hilbert space V, and let $(e_n)_{n=1}^{\infty}$ and $(\lambda_n)_{n=1}^{\infty}$ be orthonormal eigenvectors and eigenvalues that diagonalizes the covariance operator $\mathcal{K} : V \to V$. Then the Cameron-Martin space H_{μ} is given by

$$\begin{split} H_{\mu} &= \left\{ h \in V : \sum_{n=1, \lambda_n \neq 0}^{\infty} \frac{\langle h, e_n \rangle^2}{\lambda_n} < \infty \text{ and } \left(\forall n \ge 1, \lambda_n = 0 \implies \langle h, e_n \rangle = 0 \right) \right\} \\ &= \mathcal{K}^{\frac{1}{2}}(V), \end{split}$$

where $\mathcal{K}^{\frac{1}{2}}$ is the square root operator of \mathcal{K} . The variance norm is given by

$$\|h\|_{H_{\mu}}^{2} = \begin{cases} \|\mathcal{K}^{-\frac{1}{2}}h\|^{2} = \sum_{n=1,\,\lambda_{n}\neq0}^{\infty} \frac{\langle h,e_{n}\rangle^{2}}{\lambda_{n}} & \text{if } h \in \mathcal{K}^{\frac{1}{2}}(V), \\ \infty & \text{otherwise,} \end{cases}$$
(6)

for all $h \in V$.

Proof Let $h \in H_{\mu}$, that is $h = Ck = \lim_{n \to \infty} \mathcal{K}f^{(n)}$ for some $k \in \mathcal{R}_{\mu}$ and a sequence $i^*f^{(n)} \in i^*(V^*)$ such that $||k - i^*f^{(n)}||_{L^2(V,\mu_{\mathfrak{m}})} \to 0$. Using the symmetry of \mathcal{K} we obtain that

$$\langle h, e_j \rangle = \lim_n \langle \mathcal{K} f^{(n)}, e_j \rangle = \lim_n \langle f^{(n)}, \lambda_j e_j \rangle$$

for all $j \ge 1$, and hence $\langle h, e_j \rangle = 0$ whenever $\lambda_j = 0$. Moreover, Lemma 2.6 implies that $\|i^* f^{(n)}\|_{L^2(V,\mu_{\mathfrak{m}})} = \|\mathcal{K}^{\frac{1}{2}} f^{(n)}\|$ for all $n \ge 1$, and consequently we find that

$$\begin{split} \|h\|_{H_{\mu}}^{2} &= \|k\|_{L^{2}(V,\mu_{\mathfrak{m}})}^{2} = \lim_{n} \|i^{*}f^{(n)}\|_{L^{2}(V,\mu_{\mathfrak{m}})}^{2} = \lim_{n} \|\mathcal{K}^{\frac{1}{2}}f^{(n)}\|^{2} \\ &= \lim_{n} \sum_{j=1}^{\infty} \lambda_{j} \langle f^{(n)}, e_{j} \rangle^{2} \geq \sum_{j=1}^{\infty} \lim_{n} \lambda_{j} \langle f^{(n)}, e_{j} \rangle^{2} \\ &= \sum_{j=1, \lambda_{j} \neq 0}^{\infty} \lim_{n} \frac{\langle \mathcal{K}f^{(n)}, e_{j} \rangle^{2}}{\lambda_{j}} = \sum_{j=1, \lambda_{j} \neq 0}^{\infty} \frac{\langle h, e_{j} \rangle^{2}}{\lambda_{j}}, \end{split}$$

where the inequality follows by Fatou's lemma. This shows that $h \in \mathcal{K}^{\frac{1}{2}}(V)$ since $||h||^2_{H_{\mu}} < \infty$.

Conversely, let $h \in \mathcal{K}^{\frac{1}{2}}(V)$. We want to show that $h = \mathcal{C}k$ for some $k \in \mathcal{R}_{\mu}$. We do this by defining the sequence

$$f^{(n)} := \sum_{j=1,\lambda_j \neq 0}^n \frac{\langle h, e_j \rangle}{\lambda_j} e_j,$$

and noting that $\mathcal{K}f^{(n)} = \sum_{j=1}^{n} \langle h, e_j \rangle e_j \to h$ as $n \to \infty$. Furthermore, $i^*f^{(n)}$ converges in $L^2(V, \mu_{\mathfrak{m}})$ to some element $k \in R_{\mu}$ since we have the bound

$$\|i^*f^{(n)} - i^*f^{(m)}\|_{L^2(V,\mu_{\mathfrak{m}})} = \sum_{j=n,\,\lambda_j\neq 0}^m \frac{\langle h, e_j \rangle^2}{\lambda_j} \le \sum_{j=n,\,\lambda_j\neq 0}^\infty \frac{\langle h, e_j \rangle^2}{\lambda_j}$$

for all n < m. Consequently we obtain that $h = Ck = \lim_{n \to \infty} \mathcal{K}(f^{(n)})$, from which (6) follows by definition.

Remark 2.20. Note that the expression for the variance norm is $||h||_{H_{\mu}} = \sqrt{\langle h, \mathcal{K}^{-1}h \rangle} = ||\mathcal{K}^{-\frac{1}{2}}h||$, analogous to the finite dimensional \mathbb{R}^d case.

In the functional Mahalanobis distance literature, [BBC20] argues that the naive Mahalanobis distance $\|\mathcal{K}^{-\frac{1}{2}}h\|$ fails to be defined due to the non-invertibility of the square root operator $\mathcal{K}^{\frac{1}{2}}$ in the $L^2[0,1]$ setting. However, when viewed through the lens of variance norms and Cameron-Martin spaces as per Theorem 2.19, we can see how such a notion can still be made precise despite the difficulties present in the infinite dimensional and singular settings by allowing the anomaly distance to be infinite if the covariance structure of the underlying distribution does not match the new samples. This point of view was for instance taken in [Sha+23] for their conformance score anomaly distance in the finite dimensional setting.

2.4 Mahalanobis distance and conformance score

Having introduced the necessary theoretical background in the previous subsections, we are now ready to define the Mahalanobis distance on any general Banach space V. We claim that a natural definition of an anomaly distance on V is the μ -variance norm of a new sample x against the mean \mathfrak{m} , as outlined in the following definition:

Definition 2.21. Let $\mu \in \mathcal{M}_V$ for a Banach space V. We define the Mahalanobis distance $d_M(x;\mu)$ of the element $x \in V$ with respect to the measure μ to be

$$d_M(x;\mu) = \|x - \mathfrak{m}\|_{\mu\text{-}cov},$$

where \mathfrak{m} is the mean of μ .

Theorem 2.17 tells us that the Mahalanobis distance is finite for all x such that $x - \mathfrak{m} \in H_{\mu}$, where H_{μ} is the Cameron-Martin space of μ . When μ is an empirical measure of the form $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$, we will show in Section 3 that the sample Mahalanobis distance $d(x; \mu^N)$ is finite if and only if $x \in span(x_1, \dots, x_n)$, and infinite otherwise. This point of view has previously not been addressed in the infinite-dimensional setting, likely due to the restrictive assumption that the covariance operator \mathcal{K} must be injective [BBC20]. Such an assumption excludes both the sample covariance operator, which is of finite rank, and the case of a singular covariance matrix.

The following examples demonstrate that our definition coincides with, and in fact extends the Mahalanobis distance in \mathbb{R}^d to random variables with possibly degenerate co-variance matrices Σ . Furthermore, the estimator of the Mahalanobis distance will be given by the case where μ is the empirical measure of the underlying data. Example 2.23 is equivalent to the setting presented in [Sha+23].

Example 2.22 (Finite Dimensional Case). Let μ be a measure on \mathbb{R}^d with well-defined covariance matrix $\Sigma = \mathbb{E}^{x \sim \mu}[(x - \mathfrak{m})(x - \mathfrak{m})^T]$ and mean $\mathfrak{m} = \mathbb{E}^{x \sim \mu}[x]$. We have that

$$q(a,b) = \mathbf{E}^{x \sim \mu} \left[\langle a, x - \mathfrak{m} \rangle \langle b, x - \mathfrak{m} \rangle \right]$$

= $\mathbf{E}^{x \sim \mu} \left[\sum_{i=1}^{n} \sum_{j=1}^{n} a_i b_j (x - \mathfrak{m})_i (x - \mathfrak{m})_j \right]$
= $\sum_{i=1}^{n} \sum_{j=1}^{n} a_i b_j \Sigma_{i,j} = \langle a, \Sigma b \rangle = \langle \Sigma a, b \rangle = a^T \Sigma b$

for all $a, b \in \mathbb{R}^d$. Lemma 2.6 tells us that the covariance operator of μ is simply Σ . Furthermore, Theorem 2.19 implies that the Cameron-Martin space is given by $H_{\mu} = Im(\Sigma^{\frac{1}{2}}) = Im(\Sigma)$. This, together with Theorem 2.17, gives that the variance norm is given by

$$\|x\|_{H_{\mu}}^{2} = \sup_{a \in \mathbb{R}^{d}} \frac{\langle a, x \rangle^{2}}{a^{T} \Sigma a} = \begin{cases} x^{T} \Sigma^{-1} x & \text{if } x \in Im(\Sigma), \\ +\infty & \text{otherwise.} \end{cases}$$

Note also that we can write $x^T \Sigma^{-1} x = \langle \Sigma^{-\frac{1}{2}} x, \Sigma^{-\frac{1}{2}} x \rangle = |\Sigma^{-\frac{1}{2}} x|^2$. Replacing the inverse with the pseudo-inverse amounts to calculating the variance norm of the projection of x onto H_{μ} . The Mahalanobis distance is then given by $||x - \mathfrak{m}||_{H_{\mu}}$.

Example 2.23 (Empirical measure on \mathbb{R}^d). Let $\mu = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ be the empirical measure associated with a corpus of n observed data points $\{x_1, \dots, x_n\} \subset \mathbb{R}^d$. Then

$$\hat{\mathfrak{m}} = \mathrm{E}^{x \sim \mu} [x]$$

and

$$\hat{\Sigma} = \mathbf{E}^{x \sim \mu} [(x - \hat{\mathfrak{m}})(x - \hat{\mathfrak{m}})^T]$$

become the sample mean and empirical covariance matrix, and the Cameron-Martin norm is given by

$$\|x\|_{H_{\mu}}^{2} = \begin{cases} x^{T}\hat{\Sigma}^{-1}x & \text{if } x \in Im(\hat{\Sigma}), \\ +\infty & \text{otherwise.} \end{cases}$$

If $\hat{\Sigma}$ is invertible, then the Cameron-Martin norm evaluated at the point $x - \hat{\mathfrak{m}}$ coincides with the classical Mahalanobis distance, which is given by $d_M(x;\mu) = (x - \hat{\mathfrak{m}})^T \hat{\Sigma}^{-1}(x - \hat{\mathfrak{m}})$. This illustrates how $\|x - \hat{\mathfrak{m}}\|_{H_{\mu}}$, when interpreted through the Cameron-Martin norm, is the natural choice for a generalization of the Mahalanobis distance to any Banach space V.

The Gaussian Cameron-Martin literature is rich in examples of Cameron-Martin spaces and norms. A classical example is the Wiener measure on C[0, 1], the space of continuous functions, where the Cameron-Martin space is given by the space of absolutely continuous functions with square integrable derivative, with corresponding Cameron-Martin norm

$$\|h\|_{H_{\mu}} = \int_0^1 |\dot{h}(t)|^2 dt$$

More generally, there exist expressions for Cameron-Martin spaces and norms for Gaussian measures on C[0, 1] in the case where the underlying Gaussian process can be written as an integral with respect to Gaussian white noise. We refer to [Lif12] for further details.

While the Mahalanobis distance is a widely used anomaly distance for anomaly detection in the \mathbb{R}^d setting, it may perform poorly when the underlying data is not Gaussian. For non-Gaussian data, the mean element might be a poor choice of reference point when calculating anomaly distances, and so the k-nearest-neighbor (kNN) distance to the normal corpus points may serve as a better alternative [HKF04; VF11; Sha+23]. This approach requires choosing a metric for calculating the nearest-neighbours. In the finite dimensional setting the Euclidean, Minkowski, Manhattan, or even the Mahalanobis distance itself are commonly used.

Shao et al. [Sha+23] coined the term *conformance score* for the case when the Mahalanobis distance is used in conjunction with the 1-nearest-neighbour Mahalanobis distance, see also [VF11]. We aim to generalize this notion to the general setting of a Banach space V and a law $\mu \in \mathcal{M}_V$. Note that the notion of variance norm in [Sha+23] is restricted to a finite sample only, while our unified framework considers any law μ .

Definition 2.24. Let $\mu \in \mathcal{M}_V$ for a Banach space V, and let $\{x_1, \dots, x_N\} \subset V$ be a corpus of observed data. We define the conformance score $d_C(x; \mu)$ of x with respect to μ and the corpus as

$$d_C(x;\mu) = \min_{1 \le i \le N} \|x - x_i\|_{\mu - cov}$$

In other words, the conformance score is the nearest-neighbour μ -variance distance to the corpus, and in particular is new for the infinite dimensional setting covering laws μ not necessarily restricted to empirical measures. Furthermore, while Shao et al. [Sha+23] do provide a definition of conformance score for infinite dimensional V, the authors were only able to prove results and derive computational formulas for the case when V is finite dimensional, since the infinite dimensional definition depends non-trivially on V^{*}. Our framework presented in this paper overcomes this limitation. In Section 3, we will derive formulas for how to compute variance norms in the general infinite dimensional setting.

2.5 Finite-dimensional concentration inequalities

Despite the growing use of conformance scores for anomaly detection [Sha+23; Arr+24], their probabilistic properties remain underdeveloped and unexplored. To address this gap, we aim to provide concentration inequalities for the conformance score in the finitedimensional setting under the assumption that the data is Gaussian. The closest previous work is [SP16], which studied the finite sample properties of k-nearest-neighbour distances for an i.i.d. sample with a fixed reference point, using independence and binomial concentration. However, this approach is incompatible with our setting, as the conformance score is a minimum over a dependent sample. Our goal is to derive a test statistic for the conformance score at a given significance level, analogous to Hotelling's T^2 test for the classical Mahalanobis distance.

We will focus on the special case of Gaussian data, where the new sample is assumed to be contained in the normal corpus. This approach is motivated by the fact that the resulting random variable will be distributed according to the Haar measure on the Stiefel manifold, or equivalently, as the first d rows of a Haar-distributed random variable over the orthogonal group $\mathbb{O}(n)$. The setting is as follows. Consider an i.i.d. sample X_1, \dots, X_n drawn from $N_d(0, \Sigma)$ where $\Sigma \in \mathbb{R}^{d \times d}$ is non-singular. Let $X^T = (X_1 | \dots | X_n) \in \mathbb{R}^{d \times n}$ be the data matrix. We approximate the nearest-neighbour Mahalanobis distance by assuming that the data is mean-centered with known mean, and by including the new sample point in the normal corpus, say X_1 . We are interested in studying the concentration properties of

$$C(n,d) := \min_{2 \le i \le n} \|\sqrt{n} (X^T X)^{-\frac{1}{2}} (X_1 - X_i)\|$$
$$= \sqrt{n} \min_{2 \le i \le n} \|U(e_1 - e_i)\|.$$

Here $U := (X^T X)^{-\frac{1}{2}} X^T$ is distributed according to the Haar measure on the Stiefel manifold, and the e_i are the standard unit basis vectors in \mathbb{R}^n . It is important to note that C(n,d) as defined above is not the exact conformance score, but rather the *'in-sample'* conformance since X_1 is assumed to belong to the normal corpus when estimating the covariance matrix. This approach simplifies our analysis while still providing valuable insights into the behaviour of the conformance scores. Notably, this formulation allows for the use of classical results of random matrix theory on the Stiefel manifold. **Definition 2.25.** Let $d \leq n$ be natural numbers. The Stiefel manifold $S_{d,n}$ is the subset of $d \times n$ matrices defined by $S_{d,n} := \{P \in \mathbb{R}^{d \times n} : PP^T = I_d\}$ with the subspace topology inherited from the normed space $\mathbb{R}^{d \times n}$ and

$$\|P\|_{HS} := \sqrt{tr(PP^T)}$$

We let $\mu_{d,n}$ denote the Haar measure on $\mathcal{S}_{d,n}$, and μ_n the Haar measure on $\mathbb{O}(n)$.

If n = d then $S_{d,n} = \mathbb{O}(n)$ coincides with the orthogonal group. In this case we have that $\mu_{d,n} = \mu_n$. If W is a random variable with distribution μ_n , then the $S_{d,n}$ -valued random variable obtained by taking the first d rows of W has distribution $\mu_{d,n}$.

Recall that $X^T = (X_1 | \cdots | X_n) \in \mathbb{R}^{d \times n}$ is the data matrix of the sample. Thus, $U := (X^T X)^{-\frac{1}{2}} X^T \in \mathcal{S}_{d,n}$ represents the first *d* rows of a Haar random matrix $W \in \mathbb{O}(n)$. Write

$$W = (W_1 | \cdots | W_n) = \begin{pmatrix} U_1 \\ V_1 \end{pmatrix} | \cdots | \begin{pmatrix} U_n \\ V_n \end{pmatrix}$$

and note that $||W_i - W_j||^2 = 2$ for $i \neq j$. Consequently, we see that the minimum of $\mu_{d,n}$ -distributed random variables can equivalently be written as a maximum over $\mu_{n-d,n}$ -distributed random variables:

$$\min_{2 \le i \le n} \|U_1 - U_i\|^2 = 2 - \max_{2 \le i \le n} \|V_1 - V_i\|^2$$

Our goal is to study the concentration properties of the conformance score from the perspective of concentration of measure results of Lipschitz functions on the Stiefel manifold. The analysis relies on the following classical concentration of measure result on the Stiefel manifold, see e.g. [Led01, p. 27] for a standard reference.

Theorem 2.26. If $f : S_{d,n} \to \mathbb{R}$ is L-Lipschitz with respect to $\|\cdot\|_{HS}$, with mean $m_f := \int f d\mu_{d,n}$, then

$$\mu_{d,n}(m_f - f \ge t) \le e^{-\frac{(n-1)t^2}{8L^2}}$$

for all t > 0.

We are interested in the following Lipschitz functions, defined on the Stiefel manifold.

Lemma 2.27. Let $k \leq n$ and let $f : S_{k,n} \to \mathbb{R}$ be the function

$$f: (V_1|\cdots|V_n) \mapsto \max_{2 \le i \le n} \|V_1 - V_i\|.$$

Then f and f^2 are $\sqrt{2}$ and 4-Lipschitz w.r.t $\|\cdot\|_{HS}$, respectively.

Proof Let $A = (A_1 | \cdots | A_n) \in \mathcal{S}_{k,n}$ and $B = (B_1 | \cdots | B_n) \in \mathcal{S}_{k,n}$. Then

$$|f(A) - f(B)| = \left| \max_{2 \le i \le n} ||A_1 - A_i|| - \max_{2 \le i \le n} ||B_1 - B_i|| \right|$$

$$\leq \max_{2 \le i \le n} ||A_1 - B_1 + A_i - B_i||$$

$$\leq \sqrt{2} ||A - B||_{op} \le \sqrt{2} ||A - B||_{HS},$$

where $\|\cdot\|_{op}$ is the operator norm. Similarly, for f^2 we find that

$$\begin{aligned} \left| f(A)^2 - f(B)^2 \right| &= \left| f(A) - f(B) \right| \left| f(A) + f(B) \right| \\ &\leq 2\sqrt{2} \left| f(A) - f(B) \right| \\ &\leq 4 \|A - B\|_{HS}, \end{aligned}$$

where we used the fact that $||A_1 - A_i|| = \sqrt{2}$ for all $2 \le i \le n$.

By applying Theorem 2.26 to the Lipschitz function f^2 in Lemma 2.27 we obtain that

$$P(C(n,d)^{2} < c^{2}) = P\left(\min_{2 \le i \le n} n \|U_{1} - U_{i}\|^{2} < c^{2}\right)$$

$$= P\left(2 - \max_{2 \le i \le n} \|V_{1} - V_{i}\|^{2} < \frac{c^{2}}{n}\right)$$

$$= P\left(m_{f^{2}} - \max_{2 \le i \le n} \|V_{1} - V_{i}\|^{2} < m_{f^{2}} - (2 - \frac{c^{2}}{n})\right)$$

$$\ge 1 - e^{-\frac{(n-1)}{128}(m_{f^{2}} - 2 + \frac{c^{2}}{n})^{2}}$$
(7)

for all c such that $2 - \frac{c^2}{n} < m_{f^2}$. For a given $\alpha \in (0, 1)$, we can set (7) equal to $1 - \alpha$ to obtain a test statistic at the significance level α . Similar analysis can be done using f and m_f instead of f^2 and m_{f^2} .

Example 2.28. A simple lower bound on the mean m_{f^2} reads

$$m_{f^2} = \mathbb{E}[\max_{2 \le i \le n} \|V_1 - V_i\|^2] \ge \mathbb{E}[\|V_1 - V_i\|^2] = 2\frac{n-d}{n} =: l(n,d),$$
(8)

which is obtained from the identity $W = (W_1 | \cdots | W_n) = \begin{pmatrix} U_1 | \cdots | U_n \\ V_1 | \cdots | V_n \end{pmatrix} \in \mathbb{O}(n)$. As a simple application of (7), we can use the lower bound l(n,d) in (8) to obtain the following test statistic: To choose a valid c we need to use the lower bound on m_{f^2} , and to bound the actual probability involving the term $2 - \frac{c^2}{n} - m_{f^2}$ we must again use the lower bound. To this end, let $0 < \varepsilon \leq 1$ be a free variable controlling the threshold. We choose c^2 via

$$2 - \frac{c^2}{n} = (1 - \varepsilon)l(n, d) \iff c^2 = 2d + 2\varepsilon(n - d).$$

The term $m_{f^2} - 2 + rac{c^2}{n}$ can then be bounded by

$$m_f - 2 + \frac{c^2}{n} = m_f - (1 - \varepsilon)l(n, d)$$
$$\geq \varepsilon l(n, d) = 2\varepsilon \frac{n - d}{n}.$$

The concentration of measure result then reads

$$P\left(Conf(n,d)^{2} < (1-\varepsilon)2d + \varepsilon 2n\right) \ge 1 - e^{-\frac{(n-1)}{128}(m_{f}-2+\frac{\varepsilon^{2}}{n})^{2}}$$
$$= 1 - e^{-\frac{\varepsilon^{2}(n-d)^{2}}{n32}\frac{n-1}{n}}.$$

A test statistic is obtained by equating the RHS to $1 - \alpha$ for some significance level α , which will determine ε . The lower bound (8) that we used is not sharp, and to obtain an informative test statistic a sharper bound is most likely required. A sharper bound could for instance be achieved by comparing the spherical expectation m_{f^2} with a corresponding Gaussian expectation using spherical coordinates. However, obtaining such a bound for the maximum is highly nontrivial since it involves an expectation of a maximum of dependent Erlang or chi-squared distributions. We leave this to future work.

2.6 Regularized variance norms

A classical result from Cameron-Martin theory states that $\mu(H_{\mu}) = 0$ whenever μ is a Gaussian measure and dim $(H_{\mu}) = \infty$, see e.g. [Hai23, Proposition 4.45]. This implies that the sample outcomes of a V-valued random variable will almost surely not lie in the Cameron-Martin space H_{μ} , making the variance norm infinite with probability one. This issue was addressed in the functional data analysis literature for $V = L^2[0, 1]$ under the assumptions of a continuous covariance function and an injective covariance operator [BBC20]. Using our unified framework based on μ -variance norms, we extend these results to the general Hilbert setting without requiring continuity of the underlying stochastic process or injectivity of the covariance operator.

Let V be a Hilbert space. Recall by Theorem 2.19 that the μ -variance norm of $x \in V$ is given by $||x||_{H_{\mu}} = ||\mathcal{K}^{-\frac{1}{2}}x||$ if $x \in Im(\mathcal{K}^{\frac{1}{2}})$, and infinity otherwise. One natural way to extend this to a finite semi-norm defined on all of V, is to replace the inverse $\mathcal{K}^{-\frac{1}{2}}$ with the *Tikhonov regularized operator* $R_{\alpha} = (\mathcal{K} + \alpha I)^{-1}\mathcal{K}^{\frac{1}{2}}$ with smoothing parameter α . Tikhonov regularization is a classical tool used in statistic (e.g. ridge regression) and functional analysis to deal with ill-posed equations, see e.g. [Kre13]. In contrast to the inverse $\mathcal{K}^{-\frac{1}{2}}$, the Tikhonov operator R_{α} is defined on all of V and is an approximation of the pseudo-inverse of $\mathcal{K}^{\frac{1}{2}}$.

Definition 2.29. Let V be a Hilbert space, and let $\mu \in \mathcal{M}_V$. We define the Tikhonov regularized μ -variance norm with smoothing parameter $\alpha > 0$ as

$$\|x_{\alpha}\|_{\mu\text{-}cov} := \|(\mathcal{K} + \alpha I)^{-1} \mathcal{K}^{\frac{1}{2}} x\|$$
(9)

for $x \in V$, where \mathcal{K} is the covariance operator of μ .

We will sometimes refer to (9) as the α -regularized variance norm. Since \mathcal{K} is a positive symmetric compact operator, it can be diagonalized by some orthonormal sequence of eigenvectors $(e_n)_{n=1}^{\infty}$ with non-negative eigenvalues $(\lambda_n)_{n=1}^{\infty}$. Furthermore, we have that $(\mathcal{K} + \alpha I)^{-1} \mathcal{K}^{\frac{1}{2}} x = \sum_{n=1}^{\infty} \frac{\sqrt{\lambda_n}}{\lambda_n + \alpha} \langle e_n, x \rangle e_n$ for all $x \in V$, and hence the α -regularized variance norm $||x_{\alpha}||_{\mu\text{-cov}}$ can be computed via Parseval's identity, as summarized below.

Proposition 2.30. Let V be a Hilbert space, and let $\mu \in \mathcal{M}_V$. The α -regularized μ -variance norm is given by

$$||x_{\alpha}||_{\mu\text{-}cov}^{2} = \sum_{n=1}^{\infty} \frac{\lambda_{n}}{(\lambda_{n} + \alpha)^{2}} \langle e_{n}, x \rangle$$

for all $x \in V$.

An equivalent approach to the above, and what will motivate the notation x_{α} in (9), was first considered in [BBC20] in the special case $V = L^2[0, 1]$. We will now present an equivalent construction in our general setting without the assumption of injectivity on the covariance operator: The idea is to approximate each $x \in V$ via an element $x_{\alpha} \in H_{\mu}$ where α is a smoothing parameter, and then take the μ -variance norm of x_{α} which always will be finite since $x_{\alpha} \in H_{\mu}$ by construction. Since no closest element in H_{μ} to x exists in the infinite dimensional case (since H_{μ} might not closed), x_{α} is chosen by minimizing

$$x_{\alpha} := \underset{h \in H_{\mu}}{\operatorname{argmin}} \|x - h\|^{2} + \alpha \|h\|_{H_{\mu}}^{2}.$$
 (10)

Since H_{μ} is a reproducing kernel Hilbert space as discussed in the remark proceeding Theorem 2.17, it follows from [CZ07, Theorem 8.4] that the unique solution x_{α} to (10) is given by

$$x_{\alpha} = (\mathcal{K} + \alpha I)^{-1} \mathcal{K} x = \sum_{n=1}^{\infty} \frac{\lambda_n}{\lambda_n + \alpha} \langle e_n, x \rangle e_n,$$

where the second equality is due to diagonalization of \mathcal{K} , analogous to the Tikhonov case. Moreover, we have that the μ -variance norm of x_{α} is

$$||x_{\alpha}||_{\mu-\text{cov}} = ||\mathcal{K}^{\frac{1}{2}}x_{\alpha}||_{V} = ||(\mathcal{K}+\alpha I)^{-1}\mathcal{K}^{\frac{1}{2}}x||_{2}$$

and hence this approach coincides with Tikhonov perspective in Definition 2.29 in our generalized setting.

Now that we know that the regularized variance norm $||x_{\alpha}||$ is finite for all $x \in V$, we may proceed analogously as in [BBC20] and find the distribution of the Mahalanobis distance in the Gaussian setting, but without assumptions on the continuity of the stochastic process or injectivity of the covariance operator. We do this by using results from the classical theory of Gaussian measures on Banach spaces, as well as our own intermediate result presented in Proposition 2.32.

Theorem 2.31. Let μ be a Gaussian measure on a Hilbert space V, and let $\alpha > 0$. If $x \sim \mu$ is drawn from the measure μ , then the squared α -regularized Mahalanobis distance has distribution

$$\|(x-\mathfrak{m})_{\alpha}\|^{2} \stackrel{d}{=} \sum_{n=1}^{\infty} \left(\frac{\lambda_{n}}{\lambda_{n}+\alpha}\right)^{2} Y_{n},$$

where \mathfrak{m} is the mean of μ , λ_n are the eigenvalues of the covariance operator \mathcal{K} of μ , and Y_1, Y_2, \cdots is a sequence of *i.i.d.* standard χ_1^2 random variables.

Proof Let $(e_n)_{n=1}^{\infty}$ be an orthonormal sequence of eigenvectors of \mathcal{K} . It follows by Proposition 2.30 that

$$\|(x-\mathfrak{m})_{\alpha}\|^{2} = \sum_{n=1}^{\infty} \frac{\lambda_{n}}{(\lambda_{n}+\alpha)^{2}} \langle e_{n}, x-\mathfrak{m} \rangle^{2}.$$

Since μ is a Gaussian measure, the vectors e_n are by definition Gaussian distributed when acting as continuous linear functionals on V. Moreover, we have that

$$\mathbf{E}^{\mu} \big[\langle e_n, x - \mathfrak{m} \rangle \langle e_m, x - \mathfrak{m} \rangle \big] = \langle e_n \mathcal{K} e_m \rangle = \begin{cases} \lambda_n & \text{if } n = m, \\ 0 & \text{otherwise}, \end{cases}$$

$$\mathbf{E}^{\mu} \big[\langle e_n, x - \mathfrak{m} \rangle \big] = \langle e_n, \mathbf{E}^{\mu} [x] \rangle - \langle e_n, \mathfrak{m} \rangle = 0,$$

from which the result follows.

In practice we will not know the exact form of the measure μ , but rather we will work with empirical measures $\mu^N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$ based on a finite sample $\{x_1, \dots, x_N\}$ drawn from some unknown μ . Hence it would be useful to know that the sample variance norm $\|x_{\alpha}\|_{\mu^{N-\text{cov}}}$ is a consistent estimator of the actual variance norm $\|x_{\alpha}\|_{\mu-\text{cov}}$. Before we are able to prove consistency in our generalized framework, we first need an intermediate result deviating from the work in [BBC20] since their arguments for the case $V = L^2[0, 1]$ are special for the Hilbert-Schmidt covariance operator with continuous covariance function.

Proposition 2.32. Let V be a separable Hilbert space, and let $\mu^N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$ be an empirical measure of an *i.i.d.* sample drawn from $\mu \in \mathcal{M}_V$. Then we have that

$$\|\mathcal{K} - \mathcal{K}_N\|_{op} \to 0$$

almost surely as $N \to \infty$, where \mathcal{K} and \mathcal{K}_N are the covariance operators of μ and μ^N respectively, and $\|\cdot\|_{op}$ is the operator norm.

Proof First, note that all covariance operators \mathcal{K} of some measure $\mu \in \mathcal{M}_V$ are of trace class since

$$\operatorname{Tr}(|\mathcal{K}|) = \sum_{n=1}^{\infty} \langle \mathcal{K}e_n, e_n \rangle = \sum_{n=1}^{\infty} \int_V \langle x, e_n \rangle^2 d\mu_{\mathfrak{m}}(x)$$
$$= \int_V \|x\|^2 d\mu_{\mathfrak{m}}(x) = \int_V \|x\|^2 d\mu(x) - \|\mathfrak{m}\|^2$$
(11)

for any orthonormal basis $(e_n)_{n=1}^{\infty}$. Furthermore, since the empirical measure μ^N converges weakly to μ [Var58, Theorem 3], it follows from (11) together with the strong law of large numbers that $\operatorname{Tr}(|\mathcal{K}_N|) \to \operatorname{Tr}(|\mathcal{K}|)$ as $N \to \infty$ almost surely. Similarly, for all $y, z \in V$ and all covariance operators \mathcal{K} we have that

$$\langle z, \mathcal{K}y \rangle = \int_V \langle z, x \rangle \langle y, x \rangle d\mu(x) - \langle z, \mathfrak{m} \rangle \langle y, \mathfrak{m} \rangle$$

almost surely. Hence by weak convergence of measures and the strong law of large numbers we obtain that $\langle x, \mathcal{K}_N y \rangle \rightarrow \langle x, \mathcal{K} y \rangle$ almost surely, that is $\mathcal{K}_N \rightarrow \mathcal{K}$ in the weak operator topology. Since we have convergence in both trace class and weak operator topology, [Ara81, Theorem II] implies that $\|\mathcal{K} - \mathcal{K}_N\|_{op} \rightarrow 0$ almost surely.

The following theorem proves that the regularized μ -variance norm can be consistently estimated though the use of empirical measures, or equivalently through the use of the empirical covariance operator. The convergence in Theorem 2.33 can in fact be strengthened to uniform convergence on bounded sets, using the same arguments outlined in the proof.

and

Theorem 2.33. Let V be a separable Hilbert space, and let $\mu^N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$ be an empirical measure of an *i.i.d.* sample drawn from $\mu \in \mathcal{M}_V$. Then the regularized μ_N -variance norm converges pointwise to the regularized μ -variance norm, or in other words

$$||x_{\alpha}||_{\mu^{N}-cov} \to ||x_{\alpha}||_{\mu-cov}$$

almost surely as $N \to \infty$ for all $x \in V$ and all $\alpha > 0$.

Proof Fix $x \in V$ and $\alpha > 0$. For brevity we write $T_N^{\alpha} = (\mathcal{K}_N + \alpha I)^{-1}$ and $T^{\alpha} = (\mathcal{K} + \alpha I)^{-1}$. By the reverse triangle inequality we obtain that

$$\begin{aligned} \left| \|x_{\alpha}\|_{\mu_{N}} - \|x_{\alpha}\|_{\mu} \right| &= \left| \|T_{N}^{\alpha} \mathcal{K}_{N}^{\frac{1}{2}} x\| - \|T^{\alpha} \mathcal{K}^{\frac{1}{2}} x\| \right| \\ &\leq \left\| T_{N}^{\alpha} \mathcal{K}_{N}^{\frac{1}{2}} x - T^{\alpha} \mathcal{K}^{\frac{1}{2}} x \right\| \\ &\leq \|T_{N}^{\alpha}\|_{op} \|\mathcal{K}_{N}^{\frac{1}{2}} x - \mathcal{K}^{\frac{1}{2}} x\| + \|T_{N}^{\alpha} - T^{\alpha}\|_{op} \|\mathcal{K}^{\frac{1}{2}} x\|. \end{aligned}$$
(12)

First note that T_N^{α} and T^{α} are bounded by $||T_N^{\alpha}||_{op} \leq \frac{1}{\alpha}$. Proposition 2.32 implies that $||\mathcal{K}_N - \mathcal{K}||_{op} \to 0$ almost surely, from which it follows that the first term of (12) tends to 0 as $N \to \infty$. The second term also tends to 0 as $N \to \infty$ by [GGK12, Corollary 8.3] since $\mathcal{K} - \mathcal{K}_N = (\mathcal{K} + \alpha I) - (\mathcal{K}_N + \alpha I)$, which concludes the proof.

2.7 Kernelized variance norms

In this section we introduce the concept of a kernelized variance norm in the spirit of *kernel learning* using reproducing kernel Hilbert spaces (RKHS) [CS01; Sch09]. We begin by recalling the basic definitions used in RKHS theory.

Definition 2.34. A positive definite (p.d.) k on a set \mathcal{X} is defined as a symmetric function $k: \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ satisfying

$$\sum_{i=1}^{N}\sum_{j=1}^{N}a_{i}a_{j}k(x_{i},x_{j})\geq 0$$

for all $x_1, \dots x_N \in \mathcal{X}$, all $a_1, \dots, a_N \in \mathbb{R}$ and all integers $N \geq 1$.

Definition 2.35. Let \mathcal{X} be a set, and let \mathcal{H} be a Hilbert space of functions of \mathcal{X} . A reproducing kernel is defined as a function $k : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$ satisfying

- (i) $\forall x \in \mathcal{X}, \ k(\cdot, x) \in \mathcal{H},$
- (*ii*) $\forall x \in \mathcal{X}, \ \forall f \in \mathcal{H}, \ f(x) = \left\langle f, k(\cdot, x) \right\rangle_{\mathcal{H}}.$

Furthermore, \mathcal{H} is said to be a reproducing kernel Hilbert space (RKHS) if there exists a reproducing kernel for \mathcal{H} . This kernel is unique by Riesz representation theorem.

All reproducing kernels are p.d. kernels. Moreover, the Moore-Aronszajn theorem [Aro50] states that all p.d. kernels define a unique RKHS, with reproducing kernel given by the p.d. kernel. From a machine learning standpoint it may be advantageous to view RKHSs through the lens of *feature maps*. A feature map is defined as a function ϕ : $\mathcal{X} \to \mathcal{F}$ where \mathcal{F} is a Hilbert space. Every feature map defines a unique p.d. kernel via $k(x, y) := \langle \phi(x), \phi(y) \rangle_{\mathcal{F}}$. Conversely, given a RKHS with reproducing kernel k, the feature map $\phi(x) := k(\cdot, x)$ reproduces k since $\langle \phi(x), \phi(y) \rangle_{\mathcal{H}} = \langle k(\cdot, x), k(\cdot, y) \rangle_{\mathcal{H}} = k(x, y)$. Hence we may simply consider feature maps in our subsequent analysis.

A kernelized variance norm can be constructed by lifting an initial measure μ using a feature map ϕ via $\nu := \mu \circ \phi^{-1}$, and then calculating the ν -variance norm on the RKHS corresponding to ϕ . From a machine learning standpoint we will mostly be interested in empirical measures of the form $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$, hence we may simple let the kernelized measure be $\nu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$. As we will show in Section 3, the eigenvalues and eigenvectors of the covariance oper-

As we will show in Section 3, the eigenvalues and eigenvectors of the covariance operator of an empirical measure is uniquely determined by the inner product Gram matrix of the data. This in turn uniquely determines the variance norm by Theorem 2.19 and Proposition 2.30. Consequently we will be able to use the *kernel trick* to avoid explicitly computing the (possibly infinite dimensional) feature map, and thus computing the variance norm solely through the use of kernel evaluations.

Note additionally that a RKHS need not necessarily be separable. It is for this reason we defined the class of measures \mathcal{M}_V to be compatible with the non-separable case, to be able to consider the kernelized setting in full generality.

3 Computing Variance Norms

For most machine learning applications, the underlying probability measure μ is not explicitly known, and we must base our models on finite samples assumed to be drawn from μ . A natural estimator of the underlying distribution is the empirical measure $\mu^N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$, whose variance norm we showed converges almost surely to the actual μ -variance norm as per Theorem 2.33. In this section, we derive computational formulas for the variance norm with respect to μ^N .

First, we consider the Hilbert case in Section 3.1, where the variance norm is determined by the eigenvectors of the covariance operator of μ^N via Theorem 2.19. These eigenvectors are obtained using an SVD decomposition of the inner product Gram matrix of our data in V, or equivalently, via the kernel Gram matrix when V is an RKHS. Section 3.2 outlines the full procedure for calculating the Mahalanobis distance and the nearest-neighbour Mahalanobis distance. To conclude, we also derive expressions for $||x||_{\mu^N-\text{cov}}$ in the general Banach case in Section 3.3, extending the results of [Sha+23] to the infinite-dimensional setting.

3.1 Hilbert Case

By recognizing that the covariance operator \mathcal{K} is a finite rank operator when $\mu = \frac{1}{N} \sum_{n=1}^{N} \delta_{x_i}$ is an empirical measure, we can turn the problem of calculating the variance norm $\|h\|_{H_{\mu}}$

into a finite-dimensional problem. This approach yields an explicit solution in terms of the covariance operator's eigenvalues and eigenvectors, as motivated by Theorem 2.19.

Proposition 3.1. Let V be a Hilbert space, and let $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$ be an empirical measure. Then $\mu \in \mathcal{M}_V$, and the covariance operator $\mathcal{K} : V \to V$ of μ is given by

$$\mathcal{K}y = \frac{1}{N} \sum_{i=1}^{N} (x_i - \mathfrak{m}) \langle y, x_i - \mathfrak{m} \rangle$$

for $y \in V$, and the Cameron-Martin space H_{μ} is

$$H_{\mu} = span\{x_1 - \mathfrak{m}, \cdots, x_N - \mathfrak{m}\} = span\{e_1, \cdots, e_M\},\$$

where $\mathfrak{m} = \frac{1}{N} \sum_{i=1}^{N} x_i$ is the empirical mean of the data, and e_1, \dots, e_M are the eigenvectors of \mathcal{K} with positive eigenvalues.

Proof The expression for the covariance operator \mathcal{K} follows directly from the Bochner integral representation in Definition 2.4. Theorem 2.19 says that the Cameron-Martin space is given by $H_{\mu} = \mathcal{K}^{\frac{1}{2}}(V)$, and the result follows since \mathcal{K} is a finite rank operator.

Corollary 3.2. Let V be a Hilbert space, and let $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$ be an empirical measure. If $(e_m)_{m=1}^{M}$ is an orthonormal set of all the eigenvectors of K possessing positive eigenvalues $(\lambda_m)_{m=1}^{M}$, then the variance norm of μ is given by

$$\|h\|_{\mu\text{-}cov}^2 = \sum_{m=1}^M \frac{\langle h, e_m \rangle^2}{\lambda_m}$$

for $h \in H_{\mu} = Im(\mathcal{K})$, and $+\infty$ otherwise.

Proof The result follows directly from Theorem 2.19 and Proposition 3.1.

The following theorem presents an algorithm for computing the eigenvalues and eigenvectors of the empirical covariance operator, based on an SVD decomposition of the inner product Gram matrix. This concept is closely related to the techniques used for kernel PCA [SSM98]. We however give our own original proof, and relate the results back to the Cameron-Martin space of the empirical measure.

Theorem 3.3. Let V be a Hilbert space, and let $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$ be an empirical measure. Let $A \in \mathbb{R}^{N \times N}$ be defined by $A_{i,j} = \langle f_i, f_j \rangle$, where $f_i = \frac{x_i - \mathfrak{m}}{\sqrt{N}}$, with SVD decomposition $A = U \Sigma U^T$. Let $v^{(n)}$ be the n-th column of U, and $\lambda_n = \Sigma_{n,n}$. Define $M = \max\{m \leq N : \lambda_m > 0\}$. Then the elements defined by

$$e_n = \sum_{i=1}^N v_i^{(n)} f_i,$$
(13)

are orthogonal eigenvectors of \mathcal{K} with corresponding eigenvalue λ_n for $n \leq M$, and $e_n = 0$ for $M < n \leq N$. Moreover, we have that $span\{e_1, \dots, e_M\} = H_{\mu}$. **Proof** To prove that e_m is an eigenvector of \mathcal{K} with eigenvalue λ_m for $m \in \{1, \dots, M\}$, observe that

$$\mathcal{K}\left(\sum_{i=1}^{N} v_i^{(m)} f_i\right) = \sum_{j=1}^{N} f_j \left\langle \sum_{i=1}^{N} v_i^{(m)} f_i, f_j \right\rangle$$
$$= \sum_{j=1}^{N} f_j \sum_{i=1}^{N} v_i^{(m)} \left\langle f_i, f_j \right\rangle$$
$$= \sum_{j=1}^{N} f_j \lambda_m v_j^{(m)},$$

where we used that $\sum_{i=1}^{N} v_i^{(m)} \langle f_i, f_j \rangle = (Av^{(m)})_j = \lambda_m v_j^{(m)}$. Next, we check that the vectors $\{e_1, ..., e_N\}$ are linearly independent. We see that

$$\langle e_n, e_m \rangle = \left\langle \sum_{i=1}^N v_i^{(m)} f_i, \sum_{j=1}^N v_j^{(n)} f_j \right\rangle = \sum_{i=1}^N \sum_{j=1}^N v_i^{(m)} v_j^{(n)} \langle f_i, f_j \rangle$$
$$= \langle v^{(n)}, Av^{(m)} \rangle_{\mathbb{R}^N} = \lambda_m \langle v^{(n)}, v^{(m)} \rangle_{\mathbb{R}^N}$$

for $n, m \in \{1, \dots, N\}$. When $n \neq m$ we have that $\langle v^{(n)}, v^{(m)} \rangle_{\mathbb{R}^N} = 0$, hence $\langle e_n, e_m \rangle = 0$. For $1 \leq m \leq M$ the element e_m is non-zero since $\langle e_m, e_m \rangle = \lambda_m \langle v^{(m)}, v^{(m)} \rangle_{\mathbb{R}^N} > 0$. On the other hand, if $M < n \le N$ then $\lambda_n = 0$, hence $\langle e_n, e_n \rangle = 0$ and consequently $e_n = 0$.

Finally, we want to use Proposition 3.1 to conclude that $H_{\mu} = span\{e_1, \dots, e_M\}$. To this end, define $f = (f_1, \dots, f_N)$ and $e = (e_1, \dots, e_N)$ as column vectors. Using this notation, (13) can be written as $f = Ue \iff e = U^T f$, from which it follows that $span\{f_1, \dots, f_N\} = span\{e_1, \dots, e_N\} = span\{e_1, \dots, e_M\}$. This concludes the proof.

The generality of Theorem 3.3, which only assumes V is a Hilbert space, allows us to extend its application to compute eigenvalues of a kernelized covariance operator. Consider the setting where we have observations $\{x_1, \cdots, x_N\} \subset \mathcal{X}$ where \mathcal{X} is any set, and a feature map $\phi : \mathcal{X} \to \mathcal{F}$ where \mathcal{F} is a Hilbert space. By forming the empirical measure $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$ on the feature space, we can let our underlying space V be the RKHS associated with ϕ . In this case, the inner product Gram matrix becomes the Gram matrix of the reproducing kernel. This gives us an alternative theoretical foundation for the kernelized Mahalanobis distance, first introduced in [RL01]. Furthermore, if a kernel trick is known for ϕ , then we can avoid expensive direct computations of ϕ when computing the kernelized variance norm.

3.2 Algorithms

To handle the fact that the μ -variance norm $||x||_{\mu-cov}$ is only finite for vectors in the Cameron-Martin space H_{μ} , we can consider three alternative approaches:

(i) Use the Tikhonov α -regularized variance norm as per Definition 2.29 which always is finite.

- (ii) Project x onto to the space H_{μ} before calculating the variance norm of the projection. This is equivalent to using the Tikhonov formula with $\alpha = 0$, or in the finite dimensional \mathbb{R}^d setting replacing the inverse with the pseudo-inverse.
- (iii) Define the anomaly distance as infinite if the distance from x to H_{μ} exceeds a predetermined threshold. Otherwise, compute the distance using either approach (ii) or (iii).

We will present our algorithms from the perspective of (i) and (ii) above, in the setting using the kernelized variance norm, e.g. empirical measures of the form $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$ with respect to data $\{x_1, \dots, x_N\} \subset \mathcal{X}$, where \mathcal{X} is any set, and $\phi : \mathcal{X} \to \mathcal{F}$ is a feature map for some Hilbert space \mathcal{F} . We can without loss of generality consider algorithms for the variance norm in the RKHS setting, since if we let $\mathcal{X} = V$ be a Hilbert space, and $\phi = Id$ the identity, then our original non-kernelized setting is recovered. In this sense, the kernelized version serves as a strict generalization of the standard Hilbert setting. We will sometimes refer to the non-kernelized setting as the linear kernel case.

Due to possible non-linearities of the feature map, we might not be able to directly compute the inner product of the normalized elements $\langle f_n, f_m \rangle$. We address this by expressing the inner products of the scaled vectors f_i as linear combinations of $\langle x_n, x_m \rangle$. Furthermore, since we are mapping the data into a possibly infinite dimensional Hilbert space, we might not be able to store the eigenvectors in memory. To overcome this, we instead directly compute the inner products of our corpus elements with the eigenvectors of \mathcal{K} via the following: Let N, f_i , M, $v_i^{(m)}$, e_m , and λ_m be given as per Theorem 3.3. Then by bilinearity of the inner product, we obtain that

$$\langle f_m, f_n \rangle = \frac{1}{N} \bigg(\langle x_m, x_n \rangle - \frac{1}{N} \sum_{j=1}^N \langle x_m, x_j \rangle - \frac{1}{N} \sum_{j=1}^N \langle x_n, x_j \rangle + \frac{1}{N^2} \sum_{j=1}^N \sum_{i=1}^N \langle x_i, x_j \rangle \bigg),$$

and

$$\langle \frac{e_m}{\sqrt{\lambda_m}}, h \rangle = \sum_{i=1}^N \frac{v_i^{(m)}}{\sqrt{\lambda_m N}} \bigg(\langle x_i, h \rangle - \frac{1}{N} \sum_{j=1}^N \langle x_j, h \rangle \bigg),$$

for all $i \in \{1, \dots, N\}$ and $m \in \{1, \dots, M\}$.

Next, we present algorithms for computing the kernelized variance norm for the empirical measure $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$ with respect to a corpus $\{\phi(x_1), \dots, \phi(x_N)\}$, where ϕ is a feature map corresponding to a reproducing kernel k, with known kernel trick. Recall that the non-kernelized setting is recovered by setting $\phi = Id$ to be the identity, or equivalently by using the linear kernel $k(x, y) = \langle x, y \rangle$. The model fitting procedure is detailed in Algorithm 1, which has time complexity $\mathcal{O}(N^2(K+N))$ where K is the time complexity of a single kernel evaluation. Inference using the Mahalanobis distance and conformance score anomaly distances is detailed by Algorithm 2 and Algorithm 3, respectively. The time complexity of both methods is $\mathcal{O}(N(K+M))$ where M is the number of eigenvalues used in the computation.

Algorithm 1: Kernelized Gram matrix w.r.t. $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$.

Input: Data $\{x_1, \dots, x_N\}$. // Compute normalized Gram matrix via kernel trick 1 $B_{i,j} \leftarrow \langle \phi(x_i), \phi(x_j) \rangle = k(x_i, x_j)$ for $i, j \in \{1, \dots, N\}$ 2 Column mean $a_i \leftarrow \frac{1}{N} \sum_{j=1}^{N} B_{i,j}$ for $i \in \{1, \dots, N\}$ 3 Matrix mean $b \leftarrow \frac{1}{N} \sum_{i=1}^{N} a_i$ 4 $A_{i,j} \leftarrow \frac{1}{N} (B_{i,j} - a_i - a_j + b)$ for $i, j \in \{1, \dots, N\}$ // $A_{i,j} = \langle f_i, f_j \rangle$ // Compute inner products of eigenvectors and data 5 Compute SVD decomposition $U\Sigma U^t = A$ 6 Set $M \leftarrow \max\{m \le N : \Sigma_{m,m} > \lambda\}$ 7 for n = 1 to N do 8 $\begin{bmatrix} \text{for } m = 1 \text{ to } M \text{ do} \\ 9 \end{bmatrix} \begin{bmatrix} E_{n,m} \leftarrow \sum_{i=1}^{N} \frac{U_{i,m}}{\sqrt{N\Sigma_{m,m}}} (B_{i,n} - a_n) \\ // E_{n,m} = \langle e_m, \phi(x_n) \rangle \end{bmatrix}$

Output: Matrix E, and SVD decomposition $A = U\Sigma U^t$.

Algorithm 2: Kernelized Mahalanobis distance w.r.t. $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$. Input: Data $\{x_1, \dots, x_N\}$. SVD decomposition matrices $U, \Sigma \in \mathbb{R}^{N \times N}$ and $E \in \mathbb{R}^{N \times M}$ as per Algorithm 1. A new sample y. // Compute inner product of eigenvectors and sample 1 Use kernel trick $s_i \leftarrow \langle \phi(y), \phi(x_i) \rangle = k(y, x_i)$ for $i \in \{1, \dots, N\}$ 2 Average $r \leftarrow \frac{1}{N} \sum_{i=1}^{N} s_i$ 3 $p_m \leftarrow \frac{1}{\sqrt{N\Sigma_{m,m}}} \sum_{i=1}^{N} U_{i,m}(s_i - r)$ for $m \in \{1, \dots, M\}$ // $p_m = \langle e_m, \phi(y) \rangle$ // Calculate Mahalanobis distance 4 Average $c_m \leftarrow \frac{1}{N} \sum_{i=1}^{N} E_{i,m}$ for $m \in \{1, \dots, M\}$ // $c_m = \langle e_m, \frac{1}{N} \sum_{n=1}^{N} \phi(x_n) \rangle$ 5 $d \leftarrow \sqrt{\sum_{m=1}^{M} \frac{(p_m - c_m)^2}{\Sigma_{m,m}}}$ // $d = \|\phi(y) - \frac{1}{N} \sum_{n=1}^{N} \phi(x_n)\|_{H_{\mu}}^2$ Output: Kernelized Mahalanobis distance $d_M(y; \mu)$.

Algorithm 3: Kernelized conformance score w.r.t. $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$.

Input: Data $\{x_1, \dots, x_N\}$. SVD decomposition matrices $U, \Sigma \in \mathbb{R}^{N \times N}$ and $E \in \mathbb{R}^{N \times M}$ as per Algorithm 1. A new sample y. // Compute inner product of eigenvectors and sample 1 Use kernel trick $s_i \leftarrow \langle \phi(y), \phi(x_i) \rangle = k(y, x_i)$ for $i \in \{1, \dots, N\}$ 2 Average $r \leftarrow \frac{1}{N} \sum_{i=1}^{N} s_i$ 3 $p_m \leftarrow \frac{1}{\sqrt{N\Sigma_{m,m}}} \sum_{i=1}^{N} U_{i,m}(s_i - r)$ for $m \in \{1, \dots, M\}$ // $p_m = \langle e_m, \phi(y) \rangle$ // Calculate nearest-neighbour Mahalanobis distance 4 $d_n \leftarrow \sum_{m=1}^{M} \frac{(p_m - E_{n,m})^2}{\Sigma_{m,m}}$ for $n \in \{1, \dots, N\}$ // $d_n = \|\phi(y) - \phi(x_n)\|_{H_{\mu}}^2$ 5 $c \leftarrow \sqrt{\min_n d_n}$ Output: Kernelized conformance score c.

3.3 Banach case

In this section we prove a formula for the μ -variance norm when the underlying space V is any general Banach space not necessarily a Hilbert space, and μ is an empirical measure. Shao et al. [Sha+23] proved a similar theorem for the case when V is finite dimensional, for their non-measure-theoretic definition of variance norms using elementary linear algebra. The difficulty in the general infinite dimensional setting stems from the fact that the variance norm depends non-trivially on all of V^* , which might be infinite dimensional, despite the fact that the Cameron-Martin space H_{μ} is finite dimensional.

Theorem 3.4. Let $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$ be an empirical measure on a Banach space V. Write $y_i := x_i - \mathfrak{m}, i \in \{1, \dots, N\}$, for the centered data, where $\mathfrak{m} = \frac{1}{N} \sum_{i=1}^{N} x_i$ is the empirical mean. Then the following statements hold:

- (i) The Cameron-Martin space is given by $H_{\mu} = span\{y_1, \dots, y_N\}$.
- (ii) Suppose that $\{e_1, \dots, e_M\}$ is a basis of H_{μ} , and denote by $a^{(x)} \in \mathbb{R}^M$ the coordinates of a fixed projection of x onto H_{μ} . Then the μ -variance norm is given by

$$\|x\|_{\mu\text{-}cov} = \begin{cases} (a^{(x)})^T \Sigma^{-1} a^{(x)} & \text{if } a^{(x)} \in Im(\Sigma), \\ +\infty & \text{otherwise,} \end{cases}$$

for $x \in V$, where $\Sigma \in \mathbb{R}^{M \times M}$ is the empirical covariance matrix of the coordinates $a^{(y_1)}, \dots, a^{(y_N)}$.

Proof We begin by proving (*ii*), from which (*i*) will follow. Since the covariance operator $\mathcal{K}: V^* \to V$ is a finite rank operator, we have that $\mathcal{C}(\mathcal{R}_{\mu}) = \mathcal{K}(V^*)$, where $\mathcal{C}: \mathcal{R}_{\mu} \to V$ is the extended covariance operator. Hence it follows from the expression

$$\mathcal{K}f = \frac{1}{N}\sum_{i=1}^{N} (x_i - \mathfrak{m})f(x_i - \mathfrak{m})$$

that $H_{\mu} = \mathcal{K}(V^*) \subset span\{y_1, \dots, y_N\}$, which by Theorem 2.17 implies that $||x||_{\mu-cov}$ is infinite for all $x \notin span\{y_1, \dots, y_N\}$. Consequently we will only need to consider this finite span in the subsequent analysis.

Next, observe that by writing $y_i = \sum_{m=1}^{M} a_m^{(y_i)} e_m$, we obtain the following expression for the functional quadratic form of μ for all $f \in V^*$

$$q(f,f) = \frac{1}{N} \sum_{i=1}^{N} f(y_i)^2$$
$$= \frac{1}{N} \sum_{i=1}^{N} \left(\sum_{m=1}^{M} a_m^{(y_i)} f(e_m) \right)^2$$
$$= \frac{1}{N} \sum_{i=1}^{N} \langle a^{(y_i)}, b \rangle_{\mathbb{R}^M}^2$$
$$= \langle b, \Sigma b \rangle_{\mathbb{R}^M},$$

with b given by $b_m = f(e_m)$, and where $\Sigma \in \mathbb{R}^{M \times M}$ is the empirical covariance matrix of the coordinates $a^{(y_1)}, \dots, a^{(y_N)}$. Conversely, if $b \in \mathbb{R}^M$ is fixed, then $f_b(x) := \langle a^{(x)}, b \rangle_{\mathbb{R}^M}$ defines a continuous linear functional on $span\{y_1, \dots, y_N\}$, which extends continuously to V via Hahn-Banach. Consequently we obtain that

$$\begin{aligned} \|x\|_{\mu\text{-cov}}^2 &= \sup_{f \in V^*} \frac{f(x)^2}{q(f,f)} \\ &= \sup_{b \in \mathbb{R}^M} \frac{\langle a^{(x)}, b \rangle_{\mathbb{R}^M}^2}{\langle b, \Sigma b \rangle_{\mathbb{R}^M}} \\ &= \begin{cases} (a^{(x)})^T \Sigma^{-1} a^{(x)} & \text{if } a^{(x)} \in Im(\Sigma), \\ +\infty & \text{otherwise,} \end{cases} \end{aligned}$$

where the last equality follows from Example 2.22. The fact that we have equality in $H_{\mu} = span\{y_1, \dots, y_N\}$ follows by Theorem 2.17 since $a^{(x)} \in Im(\Sigma)$ if and only if $x \in span\{y_1, \dots, y_N\}$, which proves (i).

We want to stress that the choice of basis and projection map is purely for computational convenience. Different choices of basis and projection map will lead to the same expression for the variance norm, since it is independent of basis by Definition 2.16.

Corollary 3.5. If V is a Banach space, then for all $x \in V$ we have that

 $||x||_{\mu\text{-}cov} = ||a^{(x)}||_{\nu\text{-}cov},$

where $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$ and $a^{(x)}$ are defined as in Theorem 3.4, and $\nu = \frac{1}{N} \sum_{\delta_a(y_i)}$ is an empirical measure on \mathbb{R}^M .

While Corollary 3.5 gives us a way to calculate the variance norm in terms of the coordinates of the data with respect to some basis of the Cameron-Martin space H_{μ} , the method of obtaining said basis will depend heavily on the structure of V. A priori it is not clear how one would obtain a computable algorithm that works for any general Banach space V without imposing a Hilbert space structure, contrary to the Hilbert case in Section 3.2.

4 Application to Multivariate Time Series

In this section, we apply the theory developed in Section 2 and Section 3 to semi-supervised multivariate time series anomaly detection. Given a collection of non-anomalous multivariate time series, referred to as the normal data corpus, we are presented with new time series samples that we want to classify as either belonging to the normal class or as anomalies. We achieve this by defining an anomaly distance with respect to the corpus using either the Mahalanobis distance or the conformance score (see Definitions 2.21 and 2.24), assuming that our data originates from some suitable Hilbert space. For this particular problem, a natural choice for the Hilbert space is $V = (L^2[0,1])^d$. However, our framework is agnostic to the specific Hilbert space. For a given normal corpus $\{x_1, \dots, x_N\} \subset V$ we form either the standard empirical measure $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{x_i}$, or the kernelized empirical measure $\mu = \frac{1}{N} \sum_{i=1}^{N} \delta_{\phi(x_i)}$ where ϕ is a feature map corresponding to a positive definite kernel. The empirical measure μ can be interpreted as an estimator of the underlying distribution of the normal corpus. Given a new sample y, we proceed by calculating either the Mahalanobis distance

$$d_M(y;\mu) = \|y - \mathfrak{m}\|_{\mu\text{-cov}},$$

where $\mathfrak{m} = \frac{1}{N} \sum_{i=1}^{N} x_i$ is the mean of the normal corpus, or the conformance score

$$d_C(y;\mu) = \min_{1 \le n \le N} \|y - x_n\|_{\mu-cov},$$

using Algorithms 1 to 3, which defines an anomaly distance to the normal corpus. We then classify each new sample y as being anomalous or non-anomalous based on a threshold $\gamma > 0$.

The threshold γ can be determined through various approaches. A data-driven method involves splitting the normal corpus into training and validation sets, then choosing γ as an empirical quantile of the anomaly distances in the validation set. Alternatively, a theoretical approach uses the guarantees for the Mahalanobis distance and conformance score, as outlined in Theorem 2.31 and the concentration results of Section 2.5, assuming the data follows a Gaussian distribution. If a labelled subset of outliers is available, a supervised optimization method can be employed, using k-fold cross-validation to determine an optimal threshold; however, this requires access to a supervised dataset of anomalies. In our experiments, we evaluate each anomaly distance using Precision-Recall AUC and ROC-AUC metrics, which consider sensitivity across all positive thresholds, thus eliminating the need to select a fixed threshold explicitly.

Since Theorem 3.3 implies that the μ -variance norm depends solely on the Gram matrix $\langle x_i, x_j \rangle$ of the normal corpus, our anomaly distances depend only on how we choose to measure the similarity of two elements x_i and x_j via some inner product. We could for instance keep to the linear inner product given by our choice of Hilbert space, or we could choose to kernelize our method which corresponds to working with non-linear inner products given by a positive definite kernel k via $\langle \phi(x_i), \phi(x_j) \rangle = k(x_i, x_j)$ using a kernel trick. Here we consider both alternatives.

Although semi-supervised anomaly detection using the nearest-neighbour Mahalanobis distance, as opposed to the classical Mahalanobis distance, has been successfully employed in the finite-dimensional \mathbb{R}^d setting [VF11; SK20; Sha+23; Arr+24], to our knowledge no comprehensive comparison of these two anomaly distances has been carried out in the literature. In this section we carry out an extensive comparison of our newly introduced kernelized conformance score (including the non-kernelized linear case) against the kernelized Mahalanobis distance for the task of semi-supervised time series anomaly detection, using the infinite-dimensional framework developed in the previous sections.

4.1 Time series kernels

We begin by giving a brief summary of the time series kernels we will consider in our experimentation. These will consist of the linear kernel given by the $(L^2[0,1])^d$ inner product, a family of generalized integral-class kernels related to linear time warping, and a collection of time-dynamic state-of-the-art time series kernels including the global alignment kernel, the Volterra reservoir kernel, and the signature kernel.

4.1.1 Static and integral class kernels

Consider first the natural Hilbert space $(L^2[0,1])^d$, where the inner product is given by

$$\langle x, y \rangle = \int_{[0,1]} \langle x_t, y_t \rangle_{\mathbb{R}^d} dt \tag{14}$$

for $x, y \in (L^2[0,1])^d$. If x and y are discretized on a regular time grid of size T, and consequently can be viewed as d-dimensional time series of length T, then the inner product (14) can simply be computed by flattening x and y into vectors in \mathbb{R}^{Td} , and then calculating their Euclidean dot product. Instead of using the Euclidean dot product, or in other words the linear kernel, one could just as well replace this with any static kernel defined on \mathbb{R}^{dT} as per Definition 4.1 below. Similarly, we can also replace the linear static kernel in (14) to obtain a class of integral-type kernels with respect to a static kernel. In our experimentation we will consider both flattened and integral-type kernels. We give the following definitions:

Definition 4.1. A static kernel on \mathbb{R}^d is a defined as a positive definite kernel $k : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$.

Definition 4.2. Let k be a static kernel on \mathbb{R}^d . We define the time series integral class kernel of k to be the kernel

$$K_k(x,y) = \int_{[0,1]} k(x_t, y_t) dt$$

defined for d-dimensional time series x and y.

The positive definiteness of K_k follows trivially from that of k. By replacing the Euclidean dot product with a possibly non-linear static kernel, an algorithm may be able to take certain non-linearities of the data into account to increase classification accuracies. The integral type kernels can in fact be seen as a variant of linear time warping kernels, which was first introduced in [Shi+01]. The static kernels we consider in this paper are:

- (i) The linear kernel $k_{linear}(x, y) = \langle x, y \rangle$, which does not have any hyperparameters.
- (ii) The polynomial kernel $k_{poly}(x, y) = (c + \langle x, y \rangle)^p$ with hyperparameters $c \in \mathbb{R}$ and $p \in \mathbb{Z}$.
- (iii) The RBF kernel $k_{RBF}(x, y) = e^{-\frac{|x-y|^2}{2\sigma^2}}$ with hyperparameter $\sigma > 0$.

We thus have two distinct classes of time series kernels parametrized by static kernels: One is to consider flattened time series and static kernels in \mathbb{R}^{Td} , and the other is integral class kernels with respect to static kernels on \mathbb{R}^d . Note that the integral and static class kernels coincide when k is the linear kernel, and are distinct otherwise. These time series kernels can be computed in $\mathcal{O}(Td)$ time.

4.1.2 GLOBAL ALIGNMENT KERNEL

The global alignment kernel (GAK) [Cut+07; Cut11] is a dynamic-time kernel which is able to take non-linear time lags into account when measuring the similarity of two time series via dynamic time warping. While the classical dynamic time warping fails to define positive definite kernels due to failing to satisfy the triangle inequality, the global alignment kernel is able to overcome this by summing over all possible global alignments of the time series.

Definition 4.3. Let $x = (x_1, \dots, x_T)$ and $y = (y_1, \dots, x_L)$ be two time series of length Tand L respectively. An alignment π , denoted $\pi \in \mathcal{A}(T, L)$, is defined as a pair $\pi = (\pi_1, \pi_2)$ of vectors of length $p \leq T + L - 1$ such that $1 = \pi_1(1) \leq \dots \leq \pi_1(p) = T$ and $1 = \pi_2(1) \leq \dots \leq \pi_2(p) = L$. Given a similarity measure $\varphi : \mathbb{R}^d \times \mathbb{R}^d \to [0, \infty)$, the cost $D_{x,y}(\pi)$ is defined as

$$D_{x,y}(\pi) := \sum_{i=1}^{|\pi|} \varphi(x_{\pi_1(i)}, y_{\pi_2(i)}),$$

and the global alignment kernel is defined as

$$K_{GA}(x,y) = \sum_{\pi \in \mathcal{A}(T,L)} e^{-D_{x,y}(\pi)}$$
$$= \sum_{\pi \in \mathcal{A}(T,L)} \prod_{i=1}^{|\pi|} \kappa(x_{\pi_1(i)}, y_{\pi_2(i)}),$$
(15)

where $\kappa = e^{-\varphi}$ is the local similarity.

Cuturi et al. [Cut+07] proved that K_{GA} defined via a local kernel κ is positive definite if $\frac{\kappa}{1+\kappa}$ is positive definite. A sufficient condition for this is for κ to be geometrically or infinitely divisible. In practice the local kernel $\kappa = \frac{k_{RBF}}{2-k_{RBF}}$ is often used, and due to the exponential nature of (15) the GAK kernel is always made to be normalized in feature space via $\frac{K_{GA}(x,y)}{\sqrt{K_{GA}(x,x)K_{GA}(y,y)}}$. The GAK kernel has a single hyperparameter $\sigma > 0$ inherited from the static RBF kernel, and $K_{GA}(x,y)$ can be computed in $\mathcal{O}(TLd)$ time using dynamic programming.

4.1.3 VOLTERRA RESERVOIR KERNEL

The Volterra Reservoir Kernel (VRK) [GGO22] is a universal dynamic kernel designed for sequences of arbitrary length. The kernel is built by constructing a state-space representation of the classical Volterra series expansions [Wie58; San83; BC85], a series representation for analytic maps between sequences. As discussed in detail in [GGO22; Cuc+22] this idea is closely related to the principle of reservoir computing [MNM02; JH04] and associated kernels [GO21]. The VRK kernel was recently shown to outperform the RBF, GAK, and signature kernels in a market forecasting task [GGO22].

For sequences of length T the VRK kernel with hyperparameters $\tau \in \mathbb{R}$ and $\lambda \in (0, 1)$ is defined as

$$K^{\text{VOLT}}(x,y) = 1 + \sum_{k=1}^{T} \lambda^{2k} \prod_{t=0}^{k-1} \frac{1}{1 - \tau^2 \langle x_{T-t}, y_{T-t} \rangle}$$

for time series x and y of equal length such that $\tau^2 ||x|| ||y|| < 1$, where $\langle \cdot, \cdot \rangle$ is the Euclidean inner product on \mathbb{R}^d . The solution can be computed in $\mathcal{O}(Td)$ time using a recursive relation between the kernel at different time steps.

4.1.4 Signature kernels

The signature kernel [KO19; Sal+21a] is a positive definite kernel for sequential data based on tools originating from stochastic analysis and rough path theory [Lyo98]. It has many desirable theoretical properties such as invariance to time-reparametrization, universality, and characteristicness on compact sets. Algorithms using signature kernels have successfully been applied to a wide variate of fields since their inception, for instance in Bayesian forecasting [TO20], hypothesis testing [Sal+21b], and for support vector machines [Sal+21a; TOS23] achieveing state-of-the-art accuracies.

Below we give a very brief construction of the signature kernel, which is defined as an inner product in the extended tensor algebra via the so-called *signature transform*. For a detailed introduction to signature kernels we refer to the seminal papers [KO19; Sal+21a], the review article [LO23], and the lecture notes [CS24].

Definition 4.4. Let \mathcal{H} be a Hilbert space. The *m*-fold iterated integral of a bounded variation path $x \in BV([0,1], \mathcal{H})$ is recursively defined as

$$S_0(x) := 1, \qquad S_{m+1}(x) = \int_0^1 S_m(x) \otimes dx_t.$$

We define the signature transform as the map

$$S: BV([0,1], \mathcal{H}) \to \prod_{m=0}^{\infty} \mathcal{H}^{\otimes m}$$
$$x \mapsto (S_m(x))_{m=0}^{\infty},$$

and similarly, we define the truncated signature as the map $S_{0:n}(x) := (S_m(x))_{m=0}^n$. Here we use the convention that $\mathcal{H}^{\otimes 0} = \mathbb{R}$.

Given a path $x \in BV([0,1], \mathbb{R}^d)$ and a static kernel k on \mathbb{R}^d , we may canonically lift kto a path k_x taking values in its RKHS \mathcal{H} via $t \mapsto k(x_t, \cdot) \in \mathcal{H}$ using the reproducing kernel property of k. If k is the linear kernel $k(x, y) = \langle x, y \rangle_{\mathbb{R}^d}$, then k_x is simply the original path x. However, if we choose k to be a non-linear kernel such as the RBF kernel, then k_x would genuinely be different to x, and in this particular case k_x would take values in an infinite dimensional Hilbert space where direct computations of truncated signature features are impossible. The main idea behind the signature kernel is to define the sequential kernel k^{sig} w.r.t. a static kernel k as the inner product of signature transforms $S(k_x)$ and $S(k_y)$ given two paths x and y. **Definition 4.5.** Let k be a static kernel on \mathbb{R}^d . We define the k-lifted signature kernel as the mapping $k^{sig}: BV([0,1],\mathbb{R}^d) \times BV([0,1],\mathbb{R}^d) \to \mathbb{R}$,

$$k^{sig}(x,y) = \sum_{m=0}^{\infty} \left\langle S_m(k_x), S_m(k_y) \right\rangle_{\mathcal{H}^{\otimes m}},$$

where $\langle \cdot, \cdot \rangle_{\mathcal{H}^{\otimes m}}$ is the Hilbert-Schmidt inner product defined as $\langle a, b \rangle_{\mathcal{H}^{\otimes m}} = \sum_{i=1}^{n} \langle a_i, b_i \rangle_{\mathcal{H}}$ for elements $a = a_1 \otimes \cdots \otimes a_m$ and $b = b_1 \otimes \cdots \otimes b_m$. The truncated singuture kernel $k_{0:n}^{sig}$ is defined similarly using the truncated signature.

There are currently three main algorithms for computing signature kernels, each of which come with their separate advantages and disadvantages. We list the methods below:

- 1. If dim $\mathcal{H} = d < \infty$, then the truncated signature $S_{0:m}(x)$ can be computed exactly in $\mathcal{O}(Td^m)$ time, when treating the time series x as a piecewise linear path of length T. The truncated signature can then be computed by taking the inner product of the truncated signature. In practice this method is only applicable when k is the trivial linear kernel, and when d is very small (e.g. d < 5) due to the exponential time complexity.
- 2. The second method takes advantage of the kernel trick to compute $k_{0:m}^{sig}(x, y)$ via a Horner-type scheme [KO19]. This can be computed in $\mathcal{O}(LTmd)$ time using a nongeometric approximation of $k^{sig}(x, y)$, or exactly in $\mathcal{O}(LT(md + m^3))$ when viewing x and y as piecewise linear paths of lengths T and L with state-space \mathbb{R}^d . We refer the reader to [KO19, Algorithms 3 and 6] for more details.
- 3. The last method is due to Salvi et al. [Sal+21a], who proved that the signature kernel $k^{sig}(x|_{[0,s]}, y|_{[0,t]})$ solves the Goursat PDE

$$k^{sig}(x|_{[0,s]}, y|_{[0,t]}) = 1 + \int_0^s \int_0^t k^{sig}(x|_{[0,u]}, y|_{[0,v]}) \langle dk_{x_u}, dk_{x_v} \rangle_{\mathbb{R}^d},$$
(16)

where $x|_{[0,x]}$ denotes the restriction of x to the interval [0,s]. Equation (16) can be solved for piecewise linear paths using numerical PDE methods in $\mathcal{O}(LTd)$ time [Sal+21a], but is often much slower than the truncated approaches.

Generally the RBF-lifted signature kernel is preferred over the vanilla signature kernel. This is partly due to the latter having a tendency to blow up when the underlying time series are not properly normalized, something which is particularly pronounced for the PDE signature kernel which essentially acts as an inner product of tensor exponentials. In our experimentation we will use the truncated signature kernel with the linear and RBF static kernels, as well as the RBF-lifted PDE signature kernel.

Another recently introduced variant of the signature and its signature kernel is the so-called *randomized signature* [Cuc+21], which we now define.

Definition 4.6. Let $M \geq 1$ be an integer. Fix an initial condition $z_0 \in \mathbb{R}^M$, random matrices $A_1, \dots, A_d \in \mathbb{R}^{M \times M}$, random biases $b_1, \dots, b_d \in \mathbb{R}^M$ and an activation function σ . The randomized signature Z of $x \in BV([0,1], \mathbb{R}^d)$ is defined as the solution of the controlled differential equation (CDE)

$$dZ_t = \sum_{i=1}^d \sigma(A_i Z_t + b_i) dx_t^{(i)}, \qquad Z_0 = z_0,$$
(17)

where $x^{(i)}$ denotes the *i*'th component of x.

The randomized signature was first constructed by Cuchiero et al. [Cuc+21] as a random projection of the signature, with an argument based on a non-trivial application of the Johnson-Lindenstrauss lemma. It is a reservoir system and, in theory, has the same expressive power as the signature transform in the limit as $M \to \infty$. Randomized signatures have recently been successfully used for market anomaly detection [Aky+22], graph conversion [Sch+23], optimal portfolio selection [Aky+23; CM24], generative time series modelling [BGW24], and for learning rough dynamical systems [Com+23]. The CDE (17) has since been studied from the perspective of randomly initialized ResNets [CLS23], and path developments on compact Lie groups [LLN22; LLN23; CT24]. In our experiments, we will use Gaussian random matrices and biases, with tanh activation function.

4.2 Experiments

In this section we present an empirical study comparing the (potentially kernelized) Mahalanobis distance to the conformance score for semi-supervised multivariate time series anomaly detection. Our primary objective is to validate Algorithms 1 to 3 presented in this paper for this infinite dimensional setting. For comparisons of the finite-dimensional conformance score against other established methods like isolation forests, shapelets, and local outlier factors, we refer readers to [Sha+23]. Within the functional data analysis literature, the $(L^2[0,1])^d$ Mahalanobis distance has been evaluated against other common functional anomaly detection methodologies such as boxplots, outliergrams, and depth-based trimming [AR14; BBC20].

We will use UEA multivariate time series repository [Bag+18; Rui+21] in our experimentation, which in recent years has become a standard benchmark for multivariate time series classification. The repository contains 30 real world datasets consisting of multivariate time series, 26 of which are of equal lengths ranging from 8 to 2500 time steps, with state-space dimension ranging from 2 to 1345, see Table 1 for a summary. For the task of semi-supervised anomaly detection task we employ a one-versus-rest approach. In each experiment, we designate a single class label as the normal corpus, while considering all other classes as anomalous. We evaluate the performance using both precision-recall AUC (PR AUC) and ROC AUC. The results are then averaged across all class labels for a comprehensive assessment. Our experiment code is publically available at https://github.com/nikitazozoulenko/kernel-timeseries-anomaly-detection.

Code	Name	Train size	Test size	Dims	Length	Classes	Avg. Corpus Size
AWR	ArticularyWordRecognition	275	300	9	144	25	11
\mathbf{AF}	AtrialFibrillation	15	15	2	640	3	5
BM	BasicMotions	40	40	6	100	4	10
\mathbf{CR}	Cricket	108	72	6	1197	12	9
DDG	DuckDuckGeese	50	50	1345	270	5	10
\mathbf{EW}	EigenWorms	128	131	6	17,984	5	26
\mathbf{EP}	Epilepsy	137	138	3	206	4	34
\mathbf{EC}	EthanolConcentration	261	263	3	1751	4	65
\mathbf{ER}	ERing	30	270	4	65	6	5
FD	FaceDetection	5890	3524	144	62	2	2945
\mathbf{FM}	FingerMovements	316	100	28	50	2	158
HMD	HandMovementDirection	160	74	10	400	4	40
HW	Handwriting	150	850	3	152	26	6
HB	Heartbeat	204	205	61	405	2	102
LIB	Libras	180	180	2	45	15	12
LSST	LSST	2459	2466	6	36	14	176
MI	MotorImagery	278	100	64	3000	2	139
NATO	NATOPS	180	180	24	51	6	30
PD	PenDigits	7494	3498	2	8	10	749
PEMS	PEMS-SF	267	173	963	144	7	38
\mathbf{PS}	PhonemeSpectra	3315	3353	11	217	39	85
RS	RacketSports	151	152	6	30	4	38
SRS1	SelfRegulationSCP1	268	293	6	896	2	134
SRS2	SelfRegulationSCP2	200	180	7	1152	2	100
SWJ	StandWalkJump	12	15	4	2500	3	4
UW	UWaveGestureLibrary	120	320	3	315	8	15

Table 1: Summary of the 26 equal length UEA multivariate time series datasets. In our empirical study we consider all datasets of total size less than 8000, where the average corpus size is greater than 30.

4.2.1 EXPERIMENTAL SETUP

In total we will consider two different anomaly distances, namely the Mahalanobis distance and the conformance score, together with 11 different time series kernels. This includes the linear Euclidean kernel corresponding to the non-kernelized $(L^2[0,1])^d$ setting. In our open-source code we provide efficient PyTorch implementations of each kernel on both GPU and CPU, as well as an implementation of Algorithms 1 to 3 for computing the kernelized Mahalanobis distance and the kernelized conformance score. We will consider the following time series kernels in our experimentation, all of which were defined in Section 4.1:

- 1. The family of time series kernels obtained by flattening a given time series of length T into a vector in \mathbb{R}^{Td} , and then applying a static kernel. We will use the RBF, polynomial, and linear kernels as our choices of static kernels, the latter of which corresponds to the $(L^2[0,1])^d$ inner product.
- 2. The family of integral-class kernels (linear time warping), with RBF and polynomial static kernels.
- 3. The global alignment kernel (GAK).
- 4. The Volterra reservoir kernel (VRK).
- 5. Four different variants of the signature kernel: The truncated signature kernel, the RBF-lifted truncated signature kernel, the RBF-lifted PDE signature kernel, and randomized signatures with tanh activation.

Our work adds to the growing body of literature on anomaly detection using signature features, which was first studied in [Sha+23]. This was done by explicitly computing *m*-level truncated signature features, which has $\mathcal{O}(Td^m)$ time complexity. Truncated signatures were later successfully used for market anomaly detection [Aky+22], and radio astronomy [Arr+24]. The use of signatures has however been limited to very low-dimensional time series due to the exponential time complexity of explicitly computing truncated signatures. Our unified framework addresses this bottleneck, allowing for efficient computations of both signature conformance scores and signature Mahalanobis distances in $\mathcal{O}(T^2d)$ time. This significant improvement in *d* opens up the use of these methods for high-dimensional time series data.

4.2.2 Hyper-parameter selection

For each kernel, dataset, and class label, we run an extensive grid search on the designated training set using repeated k-fold cross-validation with 4 folds and 10 repeats to find the optimal kernel hyper-parameters. Let \mathbb{R}^d be the state-space, and let T be the length of the time series for a given dataset. For each method using the RBF static kernel we use the range $\sigma \in \frac{1}{\sqrt{d}} \{e^{-2}, e^{-1}, 1, e^1, e^2\}$, and similarly for the polynomial kernel we use $p \in \{2, 3, 4\}$, and $c \in \{\frac{1}{4}, \frac{1}{2}, 1, 2, 4\}$. For the GAK kernel we use the previously specified σ without the \sqrt{d} term, multiplied by $\sqrt{T} \cdot \text{med}(||x - y||)$ as is recommended in [Cut11]. For the VRK kernel we use $\tau \in \frac{1}{\sqrt{d}} \{\frac{1}{8}, \frac{1}{4}, \frac{1}{2}, 1\}$, and we let λ vary from 0.25 to 0.999 on an inverse logarithmic grid of size 10.

The signature kernels inherit hyper-parameters from their respective static kernels. We additionally scale the kernel-lifted paths by $s \in \frac{1}{\sqrt{d}} \{\frac{1}{4}, \frac{1}{2}, 1, 2, 4\}$ for the truncated signature, and by $s \in \frac{1}{\sqrt{d}} \{\frac{1}{8}, \frac{1}{4}, \frac{1}{2}, 1\}$ for the untruncated PDE kernel. We use lower values for the PDE signature kernel since untruncated signatures essentially can be viewed as tensor exponentials, whose inner products will blow up if the input values are too big. For the truncated signature kernel we let the truncation level be in $\{1, 2, 3, 4, 5, 6, 7\}$. For the randomized signature we use the tanh activation function with number of features in $\{10, 25, 50, 100, 200\}$, and random matrix variances taken from a logarithmic grid of size 8 from 0.00001 to 1. Since the randomized signature is a randomized kernel, we perform the cross validation with 5 different random seeds for the random matrix initializations, and take the best performing model (using the training set only), as is common practice for randomized methods.

Furthermore, for each method we also cross-validate over the Tikhonov regularization parameter $\alpha \in \{10^{-8}, 10^{-5}, 10^{-2}\}$, whether to concatenate time as an additional dimension to each time series, and the eigenvalue threshold λ in Algorithm 1. We set an upper limit of 50 eigenvalues for the computation of the variance norm. For numerical stability, and to make the choices of α and λ be comparable across all kernels and datasets, we normalize all time series kernels K in feature space via $\frac{K(x,y)}{\sqrt{K(x,x)K(y,y)}}$.

4.2.3 Pre-processing

For each dataset and each class label, we normalize the data to have mean zero and standard deviation one, using the statistics of the normal corpus. Average-pooling is then performed to reduce the maximum length of all time series to 100 time steps. After this, we concatenate the zero vector to each time series to allow each dynamic kernel to be translation-sensitive, and we clip all values to be in [-5, 5] for additional numerical stability. Furthermore, in our cross-validation we also include the choice of adding time as an additional dimension to all time series. For the VRK kernel specifically, we perform further clipping of the data based on the τ hyper-parameter, which is required to make the VRK kernel well-defined.

4.2.4 Data and results

Due to the high computational cost of evaluating 11 time series kernels on all 26 UEA datasets, with up to 40 experiments per dataset-kernel combination, each of which goes through an extensive repeated k-fold cross-validation, we focus our analysis on UEA datasets with a total size under 8,000 entries (see Table 1). This excluded PENDIGITS and FACEDE-TECTION. Additionally, to ensure a sufficient statistical sample size, we only considered datasets where the average corpus size exceeded 30 entries, resulting in a final selection of 12 datasets.

The anomaly distances were computed as described by Algorithms 1 to 3. The optimal kernel hyper-parameters were obtained separately for the Mahalanobis distance and the conformance score, via a 10 times repeated 4-fold cross-validation on the training data for each dataset. The objective score used in the cross-validation was the sum of ROC-AUC and PR-AUC. When calculating the precision-recall metric, we let the non-anomalous class be the positive class. The final model was then evaluated on the out-of-sample test set to obtain the final results, presented in Table 2 and Table 3.

Dataset		ROC AUC												
Dataset		linear	RBF	poly	$I_{\rm RBF}$	$I_{\rm poly}$	GAK	VRK	$S_{\rm lin}$	$S_{\rm RBF}$	$S^{\infty}_{\mathrm{RBF}}$	$S_{ anh}^{ m rand}$		
EP	С	.89	.95	.91	.94	.91	.97	.94	.98	.98	.92	.94		
ĽГ	Μ	.70	.81	.80	.80	.81	.88	.90	.98	.97	.91	.95		
EC	С	.55	.56	.55	.55	.55	.56	.58	.59	.55	.56	.55		
EC	Μ	.57	.56	.58	.55	.57	.57	.60	.56	.56	.56	.56		
FM	С	.58	.52	.54	.54	.53	.60	.53	.49	.49	.48	.54		
1, 101	Μ	.58	.51	.53	.50	.52	.54	.48	.49	.48	.51	.56		
HMD	С	.55	.43	.54	.53	.49	.46	.52	.53	.48	.50	.57		
	Μ	.45	.50	.47	.46	.51	.55	.54	.50	.50	.52	.52		
HB	С	.63	.64	.60	.61	.61	.59	.67	.70	.72	.62	.61		
	Μ	.61	.59	.61	.62	.59	.61	.62	.69	.60	.59	.60		
LSST	С	.54	.61	.53	.61	.56	.68	.53	.57	.67	.62	.62		
1001	Μ	.62	.68	.66	.67	.66	.67	.67	.67	.65	.63	.67		
MI	С	.51	.54	.57	.57	.53	.50	.54	.43	.57	.60	.45		
1011	Μ	.51	.52	.47	.46	.49	.50	.54	.47	.49	.43	.46		
PEMS	С	.91	.92	.90	.91	.89	.93	.90	.93	.92	.93	.87		
1 121015	Μ	.48	.69	.53	.66	.52	.77	.90	.80	.79	.71	.72		
\mathbf{PS}	С	.62	.65	.65	.64	.63	.66	.67	.70	.69	.56	.67		
15	Μ	.65	.67	.65	.65	.64	.65	.68	.71	.70	.54	.69		
\mathbf{RS}	С	.79	.73	.74	.80	.81	.77	.46	.73	.77	.68	.76		
105	Μ	.34	.58	.48	.60	.42	.83	.61	.79	.73	.75	.69		
SRS1	С	.68	.81	.79	.80	.81	.77	.81	.61	.77	.71	.77		
	Μ	.73	.60	.70	.59	.58	.62	.72	.77	.77	.77	.75		
SRS2	С	.57	.51	.53	.53	.53	.54	.50	.49	.48	.53	.50		
	Μ	.57	.55	.54	.55	.59	.55	.53	.54	.50	.50	.52		
Avg. AUC	С	.65	.66	.65	.67	.65	.67	.64	.65	.67	.64	.66		
	Μ	.57	.60	.58	.59	.58	.65	.65	.66	.65	.62	.64		
Avg. Rank	С	11.0	10.7	10.9	10.0	11.8	8.8	9.9	10.4	8.8	12.4	11.1		
Avg. nalik	Μ	13.5	13.2	14.1	14.8	14.8	9.6	10.2	8.8	12.1	15.0	11.3		

Table 2: One-versus-rest ROC-AUC for the semi-supervised anomaly detection experiments on the UEA multivariate time series repository. The conformance and Mahalanobis methods are denoted by C and M, respectively. The symbols I, S, S^{∞} and S^{rand} represent the integral, truncated signature, PDE signature, and randomized signature kernels, respectively.

4.2.5 DISCUSSION

For the Mahalanobis distance, there seems to be a clear advantage to working in the kernelized setting, as the results show that the linear $(L^2[0,1])^d$ inner product achieves the lowest average test scores out of all methods, with ROC-AUC and PR-AUC scores of 0.57 and 0.39, respectively. The GAK, VRK and truncated signature kernels on the other hand perform best in this regard, obtaining AUC scores of 0.65-0.66 and 0.46-0.49, respectively.

The average test scores for the conformance score (nearest-neighbour Mahalanobis distance) do not differ much between the choices of kernels, but can have significant differences

Dataset		Precision-Recall AUC										
		linear	RBF	poly	$I_{\rm RBF}$	$I_{\rm poly}$	GAK	VRK	$S_{ m lin}$	$S_{\rm RBF}$	$S_{\mathrm{RBF}}^{\infty}$	$S_{ anh}^{ ext{rand}}$
EP	C	.79	.88	.81	.87	.75	.92	.86	.96	.96	.80	.85
EF	M	.42	.58	.54	.57	.58	.73	.73	.95	.94	.78	.89
EC	С	.28	.28	.29	.28	.31	.29	.32	.32	.29	.30	.28
EC	M	.30	.32	.31	.30	.32	.32	.33	.28	.28	.30	.30
FM	С	.56	.52	.55	.57	.56	.60	.54	.54	.52	.53	.54
1, 111	M	.55	.51	.52	.50	.52	.52	.51	.52	.49	.53	.55
HMD	C	.30	.24	.31	.30	.28	.26	.27	.28	.29	.27	.29
IIIVID	M	.25	.29	.26	.28	.27	.31	.33	.29	.29	.33	.27
HB	С	.58	.60	.53	.56	.55	.55	.63	.63	.63	.58	.60
IID	M	.58	.56	.57	.61	.55	.59	.61	.64	.55	.56	.58
LSST	C	.12	.12	.10	.15	.11	.14	.10	.12	.19	.14	.10
1001	M	.14	.15	.16	.17	.14	.14	.15	.17	.17	.14	.15
MI	C	.54	.54	.57	.57	.54	.49	.55	.47	.56	.60	.49
1011	M	.55	.54	.53	.49	.53	.53	.57	.51	.50	.46	.48
PEMS	C	.79	.82	.79	.81	.79	.83	.83	.83	.81	.82	.72
1 121015	M	.34	.41	.33	.39	.31	.48	.64	.50	.52	.38	.40
PS	С	.05	.06	.06	.05	.05	.07	.06	.07	.07	.03	.06
15	M	.05	.06	.05	.05	.05	.05	.06	.07	.08	.03	.07
RS	C	.65	.64	.66	.72	.70	.62	.40	.56	.67	.52	.55
105	Μ	.20	.33	.26	.34	.23	.66	.36	.61	.53	.57	.44
SRS1	C	.69	.78	.76	.78	.77	.74	.76	.63	.76	.70	.75
51(51	Μ	.77	.67	.69	.66	.64	.69	.68	.75	.73	.74	.74
SRS2	С	.57	.51	.53	.54	.52	.55	.52	.51	.50	.54	.51
5162	Μ	.55	.55	.54	.56	.58	.55	.54	.55	.51	.51	.52
Avg. AUC	С	.49	.50	.50	.52	.49	.51	.49	.49	.52	.49	.48
Avg. AUC	Μ	.39	.41	.40	.41	.39	.46	.46	.49	.47	.44	.45
Avg. Rank	С	11.0	11.1	10.6	8.0	11.4	10.3	9.8	10.6	7.7	11.2	12.8
	Μ	13.2	13.1	15.2	13.8	15.1	10.5	10.0	8.8	12.4	14.5	12.0

Table 3: One-versus-rest PR-AUC for the semi-supervised anomaly detection experiments on the UEA multivariate time series repository. The conformance and Mahalanobis methods are denoted by C and M, respectively. The symbols I, S, S^{∞} and S^{rand} represent the integral, truncated signature, PDE signature, and randomized signature kernels, respectively.

within a single dataset. The average ROC and PR AUC scores calculated over all datasets range from 0.64-0.67 and 0.49-0.52, respectively, with the best results obtained from the RBF integral, RBF signature, and GAK kernel.

When it comes to average rank, the Mahalanobis linear truncated signature and conformance RBF truncated signature takes the number one spot, with the VRK and GAK kernels as close second place contenders. These kernels also have the most number of first places across all datasets, especially the linear truncated signature kernel. However, since the results are very dataset dependent, the best performing model and kernel combination will vary on a case-by-case basis.

Fig. 2 shows a pairwise scatter plot of the Mahalanobis distance and conformance score test results for all kernels and all datasets. The results suggest that most of the time there is no significant advantage to using one anomaly distance over the other, except for a few cases seen in the upper left quadrant where the conformance score greatly outperforms the Mahalanobis distance. The difference in performance seem to be more pronounced for the simple flattened and integral-class kernels, where the average difference is 0.07 points, as opposed to the dynamic-time kernels where the average difference is 0.02 points. This difference is more pronounced for the PR-AUC metric, and two interesting examples are RACKETSPORTS and PEMS-SF where the PR-AUC doubles for select kernels when using the conformance method.

When it comes to computing the variance norm according to Algorithms 1 to 3, both the Mahalanobis and conformance methods on average obtained their highest cross validation scores using a low number of eigenvalues, as seen in Fig. 1. Furthermore, we see that both methods in general preferred a low regularization parameter α , with $\alpha = 1e-08$ being most commonly used.

Acknowledgements

TC has been supported by the EPSRC Programme Grant EP/S026347/1 and acknowledges the support of the Erik Ellentuck Fellowship at the Institute for Advanced Study. NZ has been supported by the Roth Scholarship at Imperial College London. For the purpose of open access, the author(s) has applied a Creative Commons Attribution (CC BY) licence (where permitted by UKRI, 'Open Government Licence' or 'Creative Commons Attribution No-derivatives (CC BY-ND) licence' may be stated instead) to any Author Accepted Manuscript version arising.

References

- [AB07] C. Aliprantis and K. Border. Infinite Dimensional Analysis: A Hitchhiker's Guide. Springer, 2007.
- [AGA13] M. Amer, M. Goldstein, and S. Abdennadher. "Enhancing one-class Support Vector Machines for unsupervised anomaly detection". In: Proceedings of the ACM SIGKDD Workshop on Outlier Detection and Description, ODD 2013. 2013, pp. 8–15.




Figure 1: Optimal hyper-parameters for computing the anomaly distance as per Algorithms 1 to 3, sampled across all datasets and all kernels, normalized by the number of classes per dataset. The results were obtained via a repeated k-fold cross-validation on the train set.




Figure 2: Pairwise comparison of one-versus-rest test scores for the Mahalanobis distance and the Conformance score. Each point represents one kernel and one dataset.

- [Aky+22] E. Akyildirim, M. Gambara, J. Teichmann, and S. Zhou. "Applications of Signature Methods to Market Anomaly Detection". In: Preprint, arXiv 2201.02441 (2022).
- [Aky+23] E. Akyildirim, M. Gambara, J. Teichmann, and S. Zhou. "Randomized Signature Methods in Optimal Portfolio Selection". In: *Preprint*, arXiv 2312.16448 (2023).
- [AR14] A. Arribas-Gil and J. Romo. "Shape outlier detection and visualization for functional data: the outliergram". In: *Biostatistics* 15.4 (2014), pp. 603–619.
- [Ara81] J. Arazy. "More on Convergence in Unitary Matrix Spaces". In: Proceedings of the American Mathematical Society 83.1 (1981), pp. 44–48.
- [Aro50] N. Aronszajn. "Theory of Reproducing Kernels". In: Transactions of the American Mathematical Society 68.3 (1950), pp. 337–404.
- [Arr+24] P. Arrubarrena, M. Lemercier, B. Nikolic, T. Lyons, and T. Cass. "Novelty Detection on Radio Astronomy Data using Signatures". In: *Preprint, arXiv* 2402.14892 (2024).
- [Bag+18] A. Bagnall, H. A. Dau, J. Lines, M. Flynn, J. Large, A. Bostrom, P. Southam, and E. Keogh. "The UEA multivariate time series classification archive, 2018". In: *Preprint, arXiv 1811.00075* (2018).
- [BBC20] J. R. Berrendero, B. Bueno-Larraz, and A. Cuevas. "On Mahalanobis distance in functional settings". In: J. Mach. Learn. Res. 21.1 (2020). ISSN: 1532-4435.
- [BC85] S. Boyd and L. Chua. "Fading memory and the problem of approximating nonlinear operators with Volterra series". In: *IEEE Transactions on Circuits* and Systems 32.11 (1985), pp. 1150–1161.
- [BGW24] F. Biagini, L. Gonon, and N. Walter. "Universal randomised signatures for generative time series modelling". In: *Preprint, arXiv 2406.10214* (2024).
- [Bog15] V. Bogachev. *Gaussian Measures*. Mathematical Surveys and Monographs. American Mathematical Society, 2015.
- [Bre+00] M. M. Breunig, H.-P. Kriegel, R. T. Ng, and J. Sander. "LOF: identifying density-based local outliers". In: Proceedings of the 2000 ACM SIGMOD International Conference on Management of Data. Dallas, Texas, USA: Association for Computing Machinery, 2000, pp. 93–104.
- [Bro+22] P. O. Brown, M. C. Chiang, S. Guo, Y. Jin, C. K. Leung, E. L. Murray, A. G. M. Pazdor, and A. Cuzzocrea. "Mahalanobis Distance Based K-Means Clustering". In: *Big Data Analytics and Knowledge Discovery*. Ed. by R. Wrembel, J. Gamper, G. Kotsis, A. M. Tjoa, and I. Khalil. Cham: Springer International Publishing, 2022, pp. 256–262.
- [Cha+20] Z. Chang, W. Chen, Y. Gu, and H. Xu. "Mahalanobis-Taguchi System for Symbolic Interval Data Based on Kernel Mahalanobis Distance". In: *IEEE Access* 8 (2020), pp. 20428–20438.

[CLS23]	N. M. Cirone, M. Lemercier, and C. Salvi. "Neural signature kernels as infinite-
	width-depth-limits of controlled ResNets". In: Proceedings of the 40th Interna-
	tional Conference on Machine Learning. 2023.

- [CM24] C. Cuchiero and J. Möller. "Signature Methods in Stochastic Portfolio Theory". In: Preprint, arXiv 2310.02322 (2024).
- [Com+23] E. M. Compagnoni, A. Scampicchio, L. Biggio, A. Orvieto, T. Hofmann, and J. Teichmann. "On the effectiveness of Randomized Signatures as Reservoir for Learning Rough Dynamics". In: *IJCNN*. 2023, pp. 1–8.
- [CS01] F. Cucker and S. Smale. "On the mathematical foundations of learning". In: Bulletin of the American Mathematical Society 39 (2001), pp. 1–49.
- [CS24] T. Cass and C. Salvi. "Lecture notes on rough paths and applications to machine learning". In: *Preprint, arXiv 2404.06583* (2024).
- [CT24] T. Cass and W. F. Turner. "Free probability, path developments and signature kernels as universal scaling limits". In: *Preprint, arXiv 2402.12311* (2024).
- [Cuc+21] C. Cuchiero, L. Gonon, L. Grigoryeva, J.-P. Ortega, and J. Teichmann. "Expressive Power of Randomized Signature". In: Advances in Neural Information Processing Systems. Workshop on the Symbiosis of Deep Learning and Differential Equations. 2021.
- [Cuc+22] C. Cuchiero, L. Gonon, L. Grigoryeva, J.-P. Ortega, and J. Teichmann. "Discretetime signatures and randomness in reservoir computing". In: *IEEE Trans. Neu*ral Netw. Learn. Syst. 33.11 (2022), pp. 6321–6330. ISSN: 2162-237X,2162-2388.
- [Cut+07] M. Cuturi, J.-P. Vert, O. Birkenes, and T. Matsui. "A Kernel for Time Series Based on Global Alignments". In: 2007 IEEE International Conference on Acoustics, Speech and Signal Processing. Vol. 2. 2007, pp. II-413-II-416.
- [Cut11] M. Cuturi. "Fast global alignment kernels". In: *Proceedings of the 28th International Conference on Machine Learning*. Omnipress, 2011, pp. 929–936.
- [CZ07] F. Cucker and D. X. Zhou. Learning Theory: An Approximation Theory Viewpoint. Cambridge Monographs on Applied and Computational Mathematics. Cambridge University Press, 2007.
- [Dan+20] S. Daneshgadeh Çakmakçı, T. Kemmerich, T. Ahmed, and N. Baykal. "Online DDoS attack detection using Mahalanobis distance and Kernel-based learning algorithm". In: Journal of Network and Computer Applications 168 (2020), p. 102756.
- [DJM00] R. De Maesschalck, D. Jouan-Rimbaud, and D. Massart. "The Mahalanobis distance". In: Chemometrics and Intelligent Laboratory Systems 50.1 (2000), pp. 1–18.
- [GGK12] I. Gohberg, S. Goldberg, and M. Kaashoek. Basic Classes of Linear Operators. Birkhäuser Basel, 2012.
- [GGO22] L. Gonon, L. Grigoryeva, and J.-P. Ortega. "Reservoir kernels and Volterra series". In: *Preprint, arXiv 2212.14641* (2022).

- [GO21] L. Grigoryeva and J.-P. Ortega. "Dimension reduction in recurrent networks by canonicalization". In: *Journal of Geometric Mechanics* 13.4 (2021), pp. 647–677.
- [Hai23] M. Hairer. An Introduction to Stochastic PDEs. 2023. URL: https://www.hai rer.org/notes/SPDEs.pdf.
- [HKF04] V. Hautamaki, I. Karkkainen, and P. Franti. "Outlier detection using k-nearest neighbour graph". In: Proceedings of the 17th International Conference on Pattern Recognition. Vol. 3. 2004, 430–433 Vol.3.
- [HXD03] Z. He, X. Xu, and S. Deng. "Discovering Cluster Based Local Outliers". In: Pattern Recognition Letters 24 (2003), pp. 1641–1650.
- [JH04] H. Jaeger and H. Haas. "Harnessing Nonlinearity: Predicting Chaotic Systems and Saving Energy in Wireless Communication". In: Science 304.5667 (2004), pp. 78–80.
- [KO19] F. J. Kiraly and H. Oberhauser. "Kernels for Sequentially Ordered Data". In: Journal of Machine Learning Research 20.31 (2019), pp. 1–45.
- [Kre13] R. Kress. Linear Integral Equations. Applied Mathematical Sciences. Springer New York, 2013.
- [KSZ08] H.-P. Kriegel, M. Schubert, and A. Zimek. "Angle-based outlier detection in high-dimensional data". In: Proceedings of the 14th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. Las Vegas, Nevada, USA: Association for Computing Machinery, 2008, pp. 444–452.
- [Lah+17] H. Lahdhiri, O. Taouali, I. Elaissi, I. Jaffel, M. F. Harakat, and H. Messaoud. "A new fault detection index based on Mahalanobis distance and kernel method". In: *The International Journal of Advanced Manufacturing Technology* 91.5 (2017), pp. 2799–2809.
- [Lax14] P. Lax. Functional Analysis. Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts. Wiley, 2014.
- [Led01] M. Ledoux. *The Concentration of Measure Phenomenon*. Mathematical surveys and monographs. American Mathematical Society, 2001.
- [Lif12] M. Lifshits. *Lectures on Gaussian Processes*. Graduate Texts in Mathematics. Springer Berlin, Heidelberg, 2012.
- [LKK10] R. Lin, E. Khalastchi, and G. A. Kaminka. "Detecting anomalies in unmanned vehicles using the Mahalanobis distance". In: 2010 IEEE International Conference on Robotics and Automation. 2010, pp. 3038–3044.
- [LLN22] H. Lou, S. Li, and H. Ni. "Path Development Network with Finite-dimensional Lie Group Representation". In: Preprint, arXiv 2204.00740 (2022).
- [LLN23] H. Lou, S. Li, and H. Ni. "PCF-GAN: generating sequential data via the characteristic function of measures on the path space". In: *Thirty-seventh Conference* on Neural Information Processing Systems. 2023.
- [LO23] D. Lee and H. Oberhauser. "The Signature Kernel". In: *Preprint, arXiv 2305.04625* (2023).

[LTZ08]	F. T. Liu, K. M. Ting, and ZH. Zhou. "Isolation Forest". In: 2008 Eighth IEEE International Conference on Data Mining. 2008, pp. 413–422.
[Lyo98]	T. J. Lyons. "Differential equations driven by rough signals." In: <i>Revista Matemática Iberoamericana</i> 14.2 (1998), pp. 215–310.
[Mah36]	P. C. Mahalanobis. "On the generalized distance in statistics". In: <i>Proceedings</i> of the National Institute of Sciences (Calcutta) 2 (1936), pp. 49–55.
[MNM02]	W. Maass, T. Natschläger, and H. Markram. "Real-time computing without stable states: a new framework for neural computation based on perturbations". In: <i>Neural Computation</i> 14 (2002), pp. 2531–2560.
[Pan+21]	G. Pang, C. Shen, L. Cao, and A. V. D. Hengel. "Deep Learning for Anomaly Detection: A Review". In: <i>ACM Comput. Surv.</i> 54.2 (2021).
[PH09]	E. Pekalska and B. Haasdonk. "Kernel discriminant analysis for positive definite and indefinite kernels". In: <i>IEEE Trans Pattern Anal Mach Intell</i> 31.6 (2009), pp. 1017–1032.
[PL15]	E. J. Pedro Galeano and R. E. Lillo. "The Mahalanobis Distance for Functional Data With Applications to Classification". In: <i>Technometrics</i> 57.2 (2015), pp. 281–291.
[RL01]	A. Ruiz and P. E. López-de-Teruel. "Nonlinear kernel-based statistical pattern analysis". In: <i>IEEE Trans Neural Netw</i> 12.1 (2001), pp. 16–32.
[RS05]	J. O. Ramsay and B. W. Silverman. Functional Data Analysis. Springer, 2005.
[Ruf+20]	L. Ruff, R. A. Vandermeulen, N. Görnitz, A. Binder, E. Müller, K. Müller, and M. Kloft. "Deep Semi-Supervised Anomaly Detection". In: 8th International Conference on Learning Representations. 2020.
[Ruf+21]	L. Ruff, J. Kauffmann, R. Vandermeulen, G. Montavon, W. Samek, M. Kloft, T. Dietterich, and KR. Müller. "A Unifying Review of Deep and Shallow Anomaly Detection". In: <i>Proceedings of the IEEE</i> PP (2021), pp. 1–40.
[Rui+21]	A. P. Ruiz, M. Flynn, J. Large, M. Middlehurst, and A. Bagnall. "The great multivariate time series classification bake off: a review and experimental evaluation of recent algorithmic advances". In: <i>Data Mining and Knowledge Discovery</i> 35.2 (2021), pp. 401–449.
[Sal+21a]	C. Salvi, T. Cass, J. Foster, T. Lyons, and W. Yang. "The Signature Kernel Is the Solution of a Goursat PDE". In: <i>SIAM Journal on Mathematics of Data Science</i> 3.3 (2021), pp. 873–899.
[Sal+21b]	C. Salvi, M. Lemercier, C. Liu, B. Horvath, T. Damoulas, and T. Lyons. "Higher Order Kernel Mean Embeddings to Capture Filtrations of Stochas- tic Processes". In: <i>Advances in Neural Information Processing Systems</i> . Ed. by M. Ranzato, A. Beygelzimer, Y. Dauphin, P. Liang, and J. W. Vaughan. Vol. 34. Curran Associates, Inc., 2021, pp. 16635–16647.
[San83]	I. W. Sandberg. "Series expansions for nonlinear systems". In: <i>Circuits, Systems and Signal Processing</i> 2.1 (1983), pp. 77–87.
	48

- [Sch+23] B. Schäfl, L. Gruber, J. Brandstetter, and S. Hochreiter. "G-Signatures: Global Graph Propagation With Randomized Signatures". In: *Preprint, arXiv 2302.08811* (2023).
- [Sch09] A. J. Schölkopf Bernhard;Smola. Learning with kernels: support vector machines, regularization, optimization and beyond. Adaptive computation and machine learning. MIT Press, 2009.
- [SCZ18] J. Shang, M. Chen, and H. Zhang. "Fault detection based on augmented kernel Mahalanobis distance for nonlinear dynamic processes". In: Computers & Chemical Engineering 109 (2018), pp. 311–321.
- [Sha+23] Z. Shao, R. S.-Y. Chan, T. Cochrane, P. Foster, and T. Lyons. "Dimensionless Anomaly Detection on Multivariate Streams with Variance Norm and Path Signature". In: *Preprint, arXiv 2006.03487* (2023).
- [Shi+01] H. Shimodaira, K.-i. Noma, M. Nakai, and S. Sagayama. "Dynamic Time-Alignment Kernel in Support Vector Machine". In: Advances in Neural Information Processing Systems. Ed. by T. Dietterich, S. Becker, and Z. Ghahramani. Vol. 14. MIT Press, 2001.
- [SK20] H. Sarmadi and A. Karamodin. "A novel anomaly detection method based on adaptive Mahalanobis-squared distance and one-class kNN rule for structural health monitoring under environmental effects". In: *Mechanical Systems and Signal Processing* 140 (2020), p. 106495.
- [SP16] S. Singh and B. Poczos. "Finite-Sample Analysis of Fixed-k Nearest Neighbor Density Functional Estimators". In: Advances in Neural Information Processing Systems. Ed. by D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett. Vol. 29. Curran Associates, Inc., 2016.
- [SSM98] B. Schölkopf, A. Smola, and K.-R. Müller. "Nonlinear Component Analysis as a Kernel Eigenvalue Problem". In: Neural Computation 10.5 (1998), pp. 1299– 1319.
- [TO20] C. Toth and H. Oberhauser. "Bayesian Learning from Sequential Data using Gaussian Processes with Signature Covariances". In: Proceedings of the 37th International Conference on Machine Learning. Vol. 119. 2020, pp. 9548–9560.
- [TOS23] C. Toth, H. Oberhauser, and Z. Szabo. "Random Fourier Signature Features". In: Preprint, arXiv 2311.12214 (2023).
- [Var58] V. S. Varadarajan. "On the Convergence of Sample Probability Distributions". In: Sankhyā: The Indian Journal of Statistics (1933-1960) 19.1/2 (1958), pp. 23–26. ISSN: 00364452.
- [VF11] G. Verdier and A. Ferreira. "Adaptive Mahalanobis Distance and k -Nearest Neighbor Rule for Fault Detection in Semiconductor Manufacturing". In: *IEEE Transactions on Semiconductor Manufacturing* 24.1 (2011), pp. 59–68.
- [Wan+11] P.-C. Wang, C.-T. Su, K.-H. Chen, and N.-H. Chen. "The application of rough set and Mahalanobis distance to enhance the quality of OSA diagnosis". In: *Expert Systems with Applications* 38.6 (2011), pp. 7828–7836.

- [Wie58] N. Wiener. Nonlinear Problems in Random Theory. The Technology Press of MIT, 1958, p. 131.
- [WYT07] D. Wang, D. S. Yeung, and E. C. C. Tsang. "Weighted Mahalanobis Distance Kernels for Support Vector Machines". In: *IEEE Transactions on Neural Net*works 18.5 (2007), pp. 1453–1462.
- [XNZ08] S. Xiang, F. Nie, and C. Zhang. "Learning a Mahalanobis distance metric for data clustering and classification". In: *Pattern recognition* 41.12 (2008), pp. 3600–3612.