CLASSIFICATION OF QUASI-FREE COMPACT GROUP ACTIONS ON CUNTZ ALGEBRAS

JAMES GABE

ABSTRACT. Quasi-free actions of finite groups on Cuntz algebras \mathcal{O}_n for $n \geq 2$ are classified up to conjugacy by data in the representation ring. Partial results are obtained for quasi-free actions by compact groups.

Let G be a 2nd countable compact group and $\pi: G \to \mathcal{U}(\mathbb{C}^n)$ a faithful unitary representation for $n \geq 2$. This induces a G-action α^{π} on the Cuntz algebra \mathcal{O}_n (the C*-algebra generated by isometries s_1, \ldots, s_n with $\sum_{j=1}^n s_j s_j^* = 1$) called the *quasi-free* action, given by $\alpha_g^{\pi}(s_j) = \sum_{k=1}^n \pi(g)_{k,j} s_k$ for $g \in G$. Such actions and their fixed point algebras (known as Doplicher–Roberts algebras) have been studied for instance in [5, 3, 4, 8, 9, 10]. Goldstein and Izumi showed [7] that quasi-free outer G-actions on \mathcal{O}_{∞} (defined analogously) for G finite are all conjugate, and Izumi recently proved the same for compact groups [10, Corollary 7.3] under the same condition as in the following theorem. The following theorem classifies quasi-free actions on \mathcal{O}_n by using a recent classification theorem [6].

In the following, $\mathcal{F}(\mathbb{C}^n)$ denotes the full Fock space $\bigoplus_{k=0}^{\infty} (\mathbb{C}^n)^{\otimes k}$, and $\mathcal{F}(\pi) \colon G \to \mathcal{U}(\mathcal{F}(\mathbb{C}^n))$ denotes the induced representation. The statement for finite groups below follows from a result of Burnside [1, Chapter XV, Theorem IV].

Theorem. Let G be a 2^{nd} countable compact group, let $n \geq 2$, and suppose $\pi_1, \pi_2: G \rightarrow \mathcal{U}(\mathbb{C}^n)$ are representations so that every irreducible representation of G is equivalent to a subrepresentation of $\mathcal{F}(\pi_1)$ and $\mathcal{F}(\pi_2)$ (this is automatic for faithful representations if G is finite). Then α^{π_1} and α^{π_2} are conjugate (i.e. there exists $\beta \in \operatorname{Aut}(\mathcal{O}_n)$ with $\beta \alpha^{\pi_1} = \alpha^{\pi_2} \beta$) if and only if $1 - [\pi_1]$ and $1 - [\pi_2]$ are associated in the representation ring R(G).

Proof. As seen for instance in [7, Section 8], the G-equivariant Cuntz-Toeplitz extension

$$0 \to (\mathcal{K}(\mathcal{F}(\mathbb{C}^n)), \operatorname{Ad} \mathcal{F}(\pi_j)) \to (\mathcal{E}_n, \operatorname{Ad} \mathcal{F}(\pi_j)) \to (\mathcal{O}_n, \alpha^{\pi_j}) \to 0$$

has a *G*-equivariant completely positive splitting, and hence induces an exact triangle in KK^G (see [12, Appendix A] for details on KK^G as a triangulated category). By results of Pimsner [15, Lemma 4.7, Remark 4.10(2)], there are canonical KK^G -equivalences $(\mathcal{K}(\mathcal{F}(\mathbb{C}^n)), \operatorname{Ad} \mathcal{F}(\pi_j)) \sim_{KK^G} \mathbb{C}$ and $(\mathcal{E}_n, \operatorname{Ad} \mathcal{F}(\pi_j)) \sim_{KK^G} \mathbb{C}$ which induce an isomorphism in KK^G of this exact triangle and a triangle of the form

$$\Sigma(\mathcal{O}_n, \alpha^{\pi_j}) \to \mathbb{C} \xrightarrow{1-[\pi_j]} \mathbb{C} \xrightarrow{\iota_j} (\mathcal{O}_n, \alpha^{\pi_j})$$

where $\iota_j \in KK^G(\mathbb{C}, (\mathcal{O}_n, \alpha^{\pi_j}))$ is induced by the unital inclusion $\mathbb{C} \hookrightarrow \mathcal{O}_n$, and $1 - [\pi_j] \in KK^G(\mathbb{C}, \mathbb{C}) \cong R(G)$ (see [11, Remark 2.15(1)] for this isomorphism). Consider the following diagram in KK^G whose rows are exact triangles

 $^{2020\} Mathematics\ Subject\ Classification.\ 46L35,\ 46L55,\ 46L80,\ 20C15.$

This was supported by DFF grants 1054-00094B and 1026-00371B.

JAMES GABE

By the axioms of triangulated categories and [14, Proposition 1.1.20], there exists a KK^{G} -equivalence x making the right square commute if and only if there exists a KK^{G} -equivalence u making the middle square commute. As $\mathcal{F}(\pi_j)$ contains every irreducible representation, it follows that α^{π_j} is isometrically shift-absorbing by [10, Example 7.1]. By [6, Corollary 6.4], the existence of an x as above is equivalent to the existence of a conjugacy between α^{π_1} and α^{π_2} , while existence of such a $u \in KK^G(\mathbb{C}, \mathbb{C}) \cong R(G)$ simply means that $1 - [\pi_1]$ and $1 - [\pi_2]$ are associated in R(G).

A similar result for C^* -algebras generated by vector bundles was obtained by Dadarlat in [2, Theorem 1.1].

I suspect it is possible to take a conjugacy between α^{π_1} and α^{π_2} and induce an invertible element in R(G) implementing the associatedness of $1 - [\pi_1]$ and $1 - [\pi_2]$, but a proof of this eludes me. I strongly doubt that the converse is possible.

In the above theorem, the assumption that $\mathcal{F}(\pi_j)$ contains every irreducible representation was to ensure that the action α^{π_j} is isometrically shift-absorbing. By [10, Theorem 1.2] (one implication is proved in [13]) this is equivalent to the fixed-point algebra $\mathcal{O}_n^{\alpha^{\pi_j}}$ (the Doplicher-Roberts algebra) being simple and purely infinite (since α^{π_j} is minimal). It is suggested in [10, Example 7.1] that the condition in the above theorem might also be necessary for α^{π_j} being isometrically shift-absorbing.

I thank Gábor Szabó for useful conversations, Wojciech Szymański for helpful comments and for pointing out the reference [2], and Masaki Izumi for suggesting generalising the result from finite to compact groups.

References

- [1] W. Burnside. Theory of groups of finite order. Dover Publications, Inc., New York, xxiv+512, 1955.
- [2] M. Dadarlat. The C^* -algebra of a vector bundle. J. Reine Angew. Math., 670:121–143, 2012.
- S. Doplicher and J. Roberts. Endomorphisms of C*-algebras, cross products and duality for compact groups. Ann. of Math. (2) 130(1):75–119, 1989.
- S. Doplicher and J. Roberts. A new duality theory for compact groups. Invent. Math. 98(1):157–218, 1989.
- [5] D. Evans. On \mathcal{O}_n . Publ. Res. Inst. Math. Sci. 16(3):915–927, 1980.
- [6] J. Gabe and G. Szabó. The dynamical Kirchberg–Phillips theorem. Acta Math., 232(1):1–77, 2024.
- [7] P. Goldstein and M. Izumi. Quasi-free actions of finite groups on the Cuntz algebra \mathcal{O}_{∞} . Tohoku Math. J. (2), 63(4):729–749, 2011.
- [8] M. Izumi. Finite group actions on C^{*}-algebras with the Rohlin property. I. Duke Math. J., 122(2):233–280, 2004.
- M. Izumi. Finite group actions on C*-algebras with the Rohlin property. II. Adv. Math., 184(1):119– 160, 2004.
- [10] M. Izumi. Minimal compact group actions on C^* -algebras with simple fixed point algebras. arXiv:2405.03231.
- [11] G. G. Kasparov. Equivariant KK-theory and the Novikov conjecture. Invent. Math., 91(1):147–201, 1988.
- [12] R. Meyer and R. Nest. The Baum-Connes conjecture via localisation of categories. *Topology*, 45(2):209–259, 2006.
- [13] M. Mukohara. Inclusions of simple C^* -algebras arising from compact group actions. arXiv:2401.13989.
- [14] A. Neeman. Triangulated categories. Ann. of Math. Stud., 148. Princeton University Press, Princeton, NJ, viii+449, 2001.
- [15] M. V. Pimsner. A class of C^{*}-algebras generalizing both Cuntz-Krieger algebras and crossed products by Z. In Free probability theory (Waterloo, ON, 1995), volume 12 of Fields Inst. Commun., pages 189–212. Amer. Math. Soc., Providence, RI, 1997.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF SOUTHERN DENMARK, 5230 ODENSE, DENMARK

Email address: gabe@imada.sdu.dk