

# Decoding *Dmrt1*: Insights into vertebrate sex determination and gonadal sex differentiation

Barbora Augstenová<sup>1</sup>, Wen-Juan Ma<sup>1\*</sup>

1. Research group of Ecology, Evolution and Genetics, Biology Department, Vrije Universiteit Brussel, Brussels, Belgium

\*Corresponding: wen-juan.ma@vub.be

## Abstract

The *Dmrt* gene family is characterized by a conserved DM domain. It includes nine genes in vertebrates and is crucial to sex determination and sexual differentiation. *Dmrt1* is pivotal in testis formation and function by interacting with genes crucial for Sertoli cell differentiation, such as *Sox9*. It represses female-determining pathways and ovarian formation by silencing *Foxl2*. Across 127 vertebrate species, *Dmrt1* exhibits sexually dimorphic expression, prior to and during gonadal sex differentiation and in adult testes, implicating its role in master regulation of sex determination and gonadal sex differentiation. *Dmrt1* emerges as a master sex-determining gene in one fish, frog, chicken and reptile, with candidacy in 12 other vertebrate species. Recent studies suggest epigenetic regulation of *Dmrt1* in its promoter methylation and transposable element insertion introducing epigenetic modification to *cis*-regulatory elements of *Dmrt1*, alongside non-coding RNA involvement, in a wide spectrum of sex-determining mechanisms involving genetic, to genetic interactions with environment, to environmental factors. Additionally, alternative splicing of *Dmrt1* was found in all vertebrate groups except amphibians. *Dmrt1* has evolved many lineage-specific isoforms (ranging from 2 to 10) but has no sex-specific splicing variants in any taxa, which is in sharp contrast to the constitutional sex-specific splicing of *Dsx* in insects. Future research should focus on understanding the molecular basis of environmental sex determination from a broader taxon, and the molecular basis of epigenetic regulation. It is also essential to understand why and how multiple alternative splicing variants of *Dmrt1* evolve and the specific roles each isoform plays in sex determination and gonadal sex differentiation, as well as the significant differences in the molecular mechanisms and functions of alternative splicing between *Dmrt1* in vertebrates and sex-specific splicing of *Dsx* in insects. Understanding the differences could provide deeper insights into the evolution of sex-determining mechanisms between vertebrates and insects.

## Key words


*Dmrt1*, DM domain, sex determination, gonadal sex differentiation, master sex-determining gene, GSD, ESD, epigenetic regulation, alternative splicing, vertebrates

## 1. The structure and evolution of *Dmrt* genes

The name of the *Dmrt* genes derives from the fruit fly *Drosophila melanogaster doublesex* (*Dsx*) gene and the nematode worm *Caenorhabditis elegans male abnormal 3* (*Mab-3*) gene Related Transcriptional factors (Raymond *et al.*, 1998). *Dmrt* genes are characterized by the presence of the DM domain, DNA-binding motif, an unusual cysteine-rich zinc DNA binding motif composed of two intertwined zinc fingers (Bellefroid *et al.*, 2013). Genes, such as *Mab-3*, *Dsx*, and *Dmrt* genes with the DM domain are quite conserved in their role of regulating sex determination and sexual differentiation across metazoans, including nematodes, insects, and vertebrates (Zarkower, 2013).

Different taxa have evolved different numbers of *Dmrt* genes (Bellefroid *et al.*, 2013; Wang *et al.*, 2023). In vertebrates, there are overall recognized nine *Dmrt* genes *Dmrt1* – *Dmrt8* and *Dmrt2b* (Figure 1). The gene *Dmrt2b* is restricted to fishes. Earlier studies suggested the *Dmrt6* was missing in fishes (Veith *et al.*, 2006a), however recent studies reported *Dmrt6* in certain fishes including coelacanth, tilapia, and Southern catfish (Zhang *et al.*, 2014; Wang *et al.*, 2023). The genes *Dmrt7* and *Dmrt8* are only present in mammals (Figure 1) (Bellefroid *et al.*, 2013; Dong *et al.*, 2020). *Dmrt2*, *Dmrt3*, *Dmrt4*, *Dmrt5*, and *Dsx*-like genes were suggested to widely distribute in vertebrates (Wang *et al.*, 2023). DMRT proteins have been classified into different subfamilies, based on additional conserved protein domains and on the exon-intron structures (Figure 1). The nomenclature of the *Dmrt* genes is not always clear, e.g. in mouse the genes *Dmrt3*, *Dmrt4*, *Dmrt5* are also known as *Dmrta3*, *Dmrta1*, *Dmrta2* respectively (Bellefroid *et al.*, 2013). *Dmrt3*, *Dmrt4* and *Dmrt5* belong to one subfamily characterized by the presence of a conserved DMA domain at their C-terminus, and each gene has two coding exons, with the DM domain encoded by the first coding exon and the DMA domain encoded by the second one (Ottolenghi *et al.*, 2002). The structure of *Dmrt5* can be characterized additionally by a conserved DMB domain near the C-terminus (Figure 1) (Guo *et al.*, 2004; Sheng *et al.*, 2014). The DM domain of *Dmrt8* is not so conserved across mammals in comparison to other *Dmrt* genes, and contains many mutations, and in some cases the functional DM domain was lost altogether (Figure 1) (Bellefroid *et al.*, 2013).

Regarding the evolutionary origin of *Dmrt* genes, no conclusive phylogenetic relationship among all DMRT proteins can be drawn at the level of metazoans, because there is little sequence similarity outside the DM and DMA domains (Bellefroid *et al.*, 2013), and the length of DM domain (~70 amino acids) was too short to provide phylogenetic certainty. However, a reduced dataset revealed eight monophyletic groups containing *Dmrt* genes, and most clades have limited phylogenetic distribution (Wexler *et al.*, 2014). *Dmrt1* orthologs were mainly found in vertebrates such as mammals, birds, amphibians and teleost fishes (Wexler *et al.*, 2014). *Dmrt2*, *Dmrt3*, *Dmrt4* and *Dmrt5* orthologs of each of these genes were already present in the ancestor of deuterostomes, containing both DM and DMA domains (Figure 1). Comparison between DM and DMA domain phylogenies suggests that vertebrate *Dmrt2* have lost their DMA domain since related sequences in cephalochordates and hemichordates contain this domain (Bellefroid *et al.*, 2013). The *Dmrt2* ortholog group, containing sequences from different taxa, appear to have diverse roles in various metazoan taxa such as somiogenesis and regulation of left-right axial patterning. *Dmrt8* show strong similarity over their entire sequence to *Dmrt7*. Both have been found in mammals but not in other vertebrate species. In *Dmrt8* from human, chimp, orangutan and possibly other mammals, the open reading frame is disrupted by a stop codon 5' of the DM domain, hence they could not encode a functional DM domain (Veith *et al.*, 2006b).



**Figure 1.** Domain architecture of *Dmrt* gene family and their main distribution across vertebrate groups. All *Dmrt* genes share the highly conserved protein motif DM domain. *Dmrt3*, *Dmrt4* and *Dmrt5* share a highly conserved DMA domain at their C-terminus. *Dmrt5* has additional conserved DMB domain. The conserved DM domain on *Dmrt8* was disrupted and thus produced non-functional protein. Two important key paralogs of *Dmrt1*, *DmY* and *DmW*, are also included.

## 2. *Dmrt1* and its regulatory network

*Dmrt1*, the *Dsx* and *Mab3*-related transcription factor 1, is a pivotal gene within the *Dmrt* family, extensively studied for its role in transcriptional regulation by DMRT proteins. *Dmrt1* is indispensable for the maintenance of testis development and function, and the suppression of female-determining pathways in males across various vertebrate species (Hong *et al.*, 2007). Its crucial role in sex determination has been documented in chicken, frogs, fishes, and reptiles, where *Dmrt1* and its paralogs have been identified as the master, or candidate master, sex-determining gene (Table S1) (Matsuda *et al.*, 2002; Ferguson-Smith, 2006; Yoshimoto *et al.*, 2008; Smith *et al.*, 2009; Rhen & Schroeder, 2010; Herpin & Schartl, 2011; Ma *et al.*, 2016; Cui *et al.*, 2017; Ge *et al.*, 2017; Mustapha *et al.*, 2018; Wang *et al.*, 2022b). The role conservation of *Dmrt1* across vertebrates highlights its fundamental importance in the genetic regulatory networks that govern sex determination and gonadal sex differentiation.

The genetic regulatory network involving *Dmrt1* is complex and essential for the proper development of gonads and sexual differentiation in vertebrates. Central to this network is the interaction between *Dmrt1* and *Sox9* (Zarkower & Murphy, 2022). *Sox9*, another critical transcription factor for testis development, work synergistically with *Dmrt1* to promote the differentiation and maintenance of Sertoli cells, which are essential for testis formation in vertebrates (Raymond *et al.*, 2000; Huang *et al.*, 2005b). The interaction between *Dmrt1*

and *Sox9* is critical for the upregulation of genes involved in testis development and the repression of ovarian pathways. In addition, *Dmrt1* is involved in feedback loops that regulate its own expression as well as the expression of other genes pivotal for sexual differentiation. For instance, *Dmrt1* expression is suppressed by the forkhead transcription factor 2 gene *Foxl2*, a gene crucial for ovarian development, ensuring the suppression of male-specific pathways in females (Matson *et al.*, 2011). Additionally, *Dmrt1* can stabilize its expression patterns with autoregulatory feedback mechanisms, which is essential for maintaining sexual differentiation (Murphy *et al.*, 2010).

The activity of *Dmrt1* is modulated by various signaling pathways. The Notch signaling pathway plays a crucial role in the development and function of various tissues including the gonads. The pathway components interact with *Dmrt1* to fine-tune the sexual fate balance between male and female-determining pathways during embryonic gonadal development. This interaction is critical for the precise regulation of *Dmrt1* expression, resulting in the differentiation and proliferation of Sertoli cells essential for testis formation and spermatogenesis (Nef *et al.*, 2005). Beyond Notch signaling, *Dmrt1* is also involved in crosstalk with the *Wnt* signaling pathway that regulates sexual differentiation. The *Wnt* signaling pathway is crucial for the formation and function of Sertoli cells and Leydig cells in mouse, which are secreted by steroid and located in the interstitial region between testis cords (Chassot *et al.*, 2008; Tang *et al.*, 2008). *Wnt4* was shown to suppress male gonadal development. The *Wnt* pathway interacts with *Dmrt1* and *Sox9* to regulate the balance between female and male gonadal differentiation (Kim *et al.*, 2006; Chassot *et al.*, 2008).

### **3. *Dmrt1* expression during development and adulthood in vertebrates**

*Dmrt1* is crucially involved in gonadal sex differentiation across a wide range of taxa in vertebrates (Mawaribuchi *et al.*, 2012; Ayers *et al.*, 2013; Huang *et al.*, 2017). The role of *Dmrt1* in testes development and functionality in vertebrates was first described in humans, where the deletion of autosomal chromosome 9 region containing *Dmrt1* (placed in the distal 9p region - 9p24.3) led to complete or partial gonadal dysgenesis, and hemizygoty of this region led to testicular defects and further to XY sex reversals (Raymond *et al.*, 1999; Ledig *et al.*, 2010).

*Dmrt1* has diverse expression dynamics in the somatic and gonadal tissues during development in vertebrates. In Table S1 with 127 vertebrate species (75 in fishes, 17 in amphibians, 16 in mammals and 19 in reptiles including avians), we have summarized the *Dmrt1* temporal and spatial expression pattern in gonads prior to and during gonadal sex differentiation, at sub-adult and adult stages, as well as various adult somatic tissues. Several general patterns can be drawn from Table S1. First, *Dmrt1* expression was examined in undifferentiated gonads in a handful of species with either similar expression level or higher expression in XY individuals or gonads. Second, for the vast majority of species, *Dmrt1* showed a sexually dimorphic expression pattern during the gonadal sex differentiation process with increased expression in males and decreased in females, highlighting its conserved role involving testis formation during development. However, in certain frog species, there was no detected difference in *Dmrt1* expression during gonadal sex differentiation (Table S1) (Duarte-Guterman *et al.*, 2012; Xu *et al.*, 2015). Third, in sub-adult or adult gonads, *Dmrt1* is exclusively expressed in testes in 27 species, has significant higher expression in testes and weaker in ovaries in about 52 species, and inconclusive expression pattern in 11 species due to expression data in testes but no data examined in ovaries (Table S1). *Dmrt1* expression is also detected in ovotestes with either

weak expression (protandrous hermaphrodite fish *Acanthopagrus latus*) or stronger expression than either testes or ovaries (protogynous hermaphrodite fish *Epinephelus coioides*). This is consistent with *Dmrt1*'s role in formation and maintenance of testicular development. Fourth, *Dmrt1* has no expression or very weak expression in adult somatic tissues such as brain, liver, kidney, mussel, heart, spleen, eye, among others (Table S1).

*Dmrt1* expression change can be associated with sex change in certain fish species with sequential hermaphroditism. The *Dmrt1* expression in protandrous clownfish *Amphiprion bicinctus* (Pomacentridae) decreased in gonads during and after the transformation from male to female (Casas *et al.*, 2016). In the closely related clownfish species *Amphiprion clarkii*, *Dmrt1* was expressed in nonbreeder and male gonads (Wang *et al.*, 2022a). However, *Dmrt1* expression level was retained and, surprisingly, at even higher levels in ovaries after the sex transformation, suggesting a potential role in promoting ovarian development and require further functional studies in this species (Casas *et al.*, 2016; Wang *et al.*, 2022a). In the fish species with protogynous hermaphroditism such as *Monopterus albus* (Synbranchidae), *Epinephelus merra* (Serranidae) or *Halichoeres tenuispinis* (Labridae), the expression of *Dmrt1* increased in gonads at the time of the transformation from females to males, suggesting role of *Dmrt1* in testis formation and development (Huang *et al.*, 2005b; Alam *et al.*, 2008; Miyake *et al.*, 2012).

As summarized in Table S1, studies have shown that *Dmrt1* at adult stage is predominantly expressed in testis (Shibata *et al.*, 2002; Hong *et al.*, 2007). Interestingly, *Dmrt1* expression levels vary with breeding season condition. Various studies across vertebrates show that *Dmrt1* expression differs between the breeding and non-breeding season with higher expression in testis during breeding season, suggesting *Dmrt1* expression is associated with the active status of spermatogenesis. The higher expression of *Dmrt1* in testis of males in breeding season has been found in the Iberian Mole *Talpa occidentalis* (Talpidae) (Dadhich *et al.*, 2010), in various rodent species (Massoud *et al.*, 2021), in reptiles such as the Italian wall lizard *Podarcis siculus* (Lacertidae) (Capriglione *et al.*, 2010), as well as the rainbow trout fish *Oncorhynchus mykiss* (Salmonidae) (Marchand *et al.*, 2000).

#### **4. *Dmrt1*'s role in gonadal sex differentiation in vertebrates**

The transcriptional regulation function of *Dmrt1* is best studied in mouse. Gene/allelic gain and loss studies demonstrated that at least one functional copy of *Dmrt1* was required for the proliferation and radial migration of Sertoli cells and germ cells, and the development of testes (Raymond *et al.*, 2000; Fahrioglu *et al.*, 2006; Kim *et al.*, 2007). Furthermore, functional studies in mouse showed that *Dmrt1* determined whether male germ cells had undergone mitosis and spermatogonia differentiation, or meiosis. Loss of *Dmrt1* in the undifferentiated spermatogonia disrupted cyclical gene expression in Sertoli cells, activated transcription of the spermatogonia differentiation factor, which prevented meiosis and promoted spermatogonia development (Matson *et al.*, 2011). *Dmrt1* was not required for the development and differentiation of ovaries in the mouse (Raymond *et al.*, 2000). However, *Dmrt1* was expressed in the genital ridge, the gonadal primordium, in both sexes of the mouse, and was required for normal meiotic prophase in the embryonic ovaries and formation of primordial follicles. Loss of *Dmrt1* in the embryonic ovary resulted in the reduction of the number of primordial follicles in the juvenile ovary, although the females were still fertile (Krentz *et al.*, 2011). In rabbits, recent study shows *Dmrt1* was involved in testis determination, and yet it was also essential for female fertility.

The XX and XY individuals with knockout of *Dmrt1* did not undergo meiosis and folliculogenesis, and XX adult females without *Dmrt1* were sterile (Dujardin *et al.*, 2023). Furthermore, *Dmrt1* seems to be indispensable in postnatal testes differentiation in therian mammals overall (Raymond *et al.*, 2000; Kim *et al.*, 2007). Finally, *Dmrt1* plays a role in the origin of germ cell cancers such as testicular teratoma in several mammals including humans (Looijenga *et al.*, 2006; Turnbull *et al.*, 2010; Kratz *et al.*, 2011; Krentz *et al.*, 2011).

Functional studies of *Dmrt1* in non-mammals have been constrained to only a handful of species (Table S1). For instance, *Dmrt1* or its paralogues were identified as candidate master sex-determining genes in 7 fishes with knockdown or knockout functional assay, such as medaka (*Oryzias latipes*), *Astyanax scabripinnis*, Chinese tongue sole (*Cynoglossus semilaevis*), Siamese fighting fish (*Betta splendens*), Spotted scat (*Scatophagus argus*), African scat (*Scatophagus tetracanthus*), Spotbanded scat (*Selenotoca multifasciata*) (Table 1). For the rest of fishes, most species focused on expression profile during gonadal sex differentiation and at adult stages. In many fishes, such as Nile tilapia (*Oreochromis niloticus*), black porgy (*Acanthopagrus schlegelii*), Spotted scat (*Scatophagus argus*), Chinese tongue sole (*Cynoglossus semilaevis*), Spotted knifejaw (*Oplegnathus punctatus*), Japanese eel (*Anguilla japonica*), *Dmrt1* showed sexually dimorphic expression with higher expression in males during gonadal sex differentiation throughout development and was predominantly expressed in adult testis (He *et al.*, 2003; Li *et al.*, 2013; Chen *et al.*, 2014; Cui *et al.*, 2017; Mustapha *et al.*, 2018; Qi *et al.*, 2024). In the zebrafish (*Danio rerio*), knockdown of *Dmrt1* at early embryonic stage resulted in fertile females, or in rare cases resulted in males with testicular dysgenesis, suggesting a role for male germ cell maintenance and proliferation, and no requirement for ovary development (Lin *et al.*, 2017; Webster *et al.*, 2017). In the Siamese fighting fish (*Betta splendens*) with XX/XY sex chromosome system, *Dmrt1* showed male-biased expression in undifferentiated gonads and adult testis, and the knockout of *Dmrt1* resulted in ovarian development in XY individuals. Transposable element insertion led to reduced *Dmrt1* expression, resulting in sex reversals (XY with ovaries), suggesting a role of epigenetic regulation of *Dmrt1* in association with testis development (Wang *et al.*, 2022b) (see details in section ‘Role of epigenetic regulation of *Dmrt1* in sexual differentiation’).

In many studied frog species such as *Xenopus laevis*, *Bombina bombina*, *Bufo viridis*, *Hyla arborea*, *Rana arvalis* and *Rana temporaria*, *Dmrt1* showed strong sexually dimorphic, male-biased expression in gonads either throughout development and/or at adult stage (Table S1) (Shibata *et al.*, 2002; Piprek *et al.*, 2013; Ma *et al.*, 2018c; b). The paralog of *Dmrt1*, *DmW*, was an ovary-determining gene in the African claw frog *Xenopus tropicalis* as knockdown of *DmW* led to testis formation (Yoshimoto *et al.*, 2008) (see details in section ‘*DmW* as master sex-determining gene in the African clawed frog’). *Dmrt1* had low expression in undifferentiated gonads at early embryogenesis, and increased dramatically during gonadal sex differentiation in the testis but was low in the developing ovaries (Piprek *et al.*, 2013), suggesting a role involving in testis formation and function. In the European common frog *Rana temporaria*, *Dmrt1* is located on the sex chromosome. Population genetics and pedigree analysis showed that  $F_{st}$  between sexes was significantly elevated at the *Dmrt1* region, and *Dmrt1* polymorphism (*Dmrt1* Y-chromosome haplotypes) covaried with sex determination in various populations across Europe (Ma *et al.*, 2016). Furthermore, *Dmrt1* Y-chromosome haplotype was also associated with varying level of masculinization, with evidence of association with male testis formation and development throughout development (Rodrigues *et al.*, 2017; Ma *et al.*, 2018a; b). All

evidence together points out *Dmrt1* is a strong candidate master sex-determining gene in this species (Table 1). In salamander *Hynobius retardatus*, high temperature led to downregulation of *Dmrt1* expression, leading to sex reversal from males to females (Sakata *et al.*, 2006).

Most reptiles have genetic sex determination (GSD). All avians are possessing GSD with ZZ/ZW system, and Squamata mostly have GSD with either XY/XX or ZW/ZZ systems (Sarre *et al.*, 2011). However, sex, in certain lineages such as turtles and crocodiles, is predominantly determined by temperature (TSD) in thermal sensitive period during embryogenesis, which is one primary trigger of environmental sex-determining mechanism (ESD) (Table S1) (Valenzuela & Lance, 2004; Bista & Valenzuela, 2020). In a few species including the Australian bearded dragon, GSD is sensitive to temperature and was overturned by temperature within one generation. Study showed instant creation of a lineage of ZZ temperature-sensitive bearded dragons when altering the incubation temperature during thermal sensitive period, and W chromosome was eliminated in these animals (Holleley *et al.*, 2015). In the red-eared slider turtle (*Trachemys scripta*) with TSD, gene gain and loss analysis showed that *Dmrt1* was required for testes differentiation throughout development and was considered a strong master sex-determining gene (Ge *et al.*, 2017). In most studied TSD species, e.g. *Alligator mississippiensis*, *Lepidochelys olivacea*, *Chelydra serpentina*, *Trachemys scripta*, *Dmrt1* was expressed and increased during temperature-sensitive embryogenesis period at male producing temperature (Table S1) (Rhen *et al.*, 2007; Urushitani *et al.*, 2011; Bieser & Wibbels, 2014; Ge *et al.*, 2017; Díaz-Hernández *et al.*, 2020), which was consistently with the role of *Dmrt1* involving in testis formation and function. Furthermore, *Dmrt1* was shown to play an important role in proliferation and maintenance of spermatogonia and spermatocytes in the Italian wall lizard *Podarcis sicula* (Capriglione *et al.*, 2010).

## **5. *Dmrt1* and its paralogs' role as master sex-determining gene**


The DM domain has a conserved role in sexual development across both vertebrates and invertebrates (Hong *et al.*, 2007; Zarkower & Murphy, 2022), which makes *Dmrt1* a good candidate for the master sex-determining gene and is expected to be a crucial component of the sex-determining pathway in many vertebrates. So far *Dmrt1*, or its paralogs *DmY* and *DmW* (Figure 1), are demonstrated to be the master sex-determining genes in chicken, fish and frog respectively (Matsuda *et al.*, 2002; Nanda *et al.*, 2002; Yoshimoto *et al.*, 2008; Smith *et al.*, 2009; Yoshimoto *et al.*, 2010), and are candidate master sex-determining genes in 12 other vertebrates (Table 1).

### **5.1 *Dmrt1bY/DmY* as master sex-determining gene in medaka fish**

Duplication of *Dmrt1*, *Dmrt1bY* or *DmY*, was identified as master sex-determining gene in the medaka fish (*Oryzias latipes*) (Figure 2) (Matsuda *et al.*, 2002; Nanda *et al.*, 2002). The same gene was later found to be involved in sex determination in one closely related species *O. curvinotus*, but not in any other closely related fish species (Matsuda *et al.*, 2002; Zhang, 2004). Both species possess XX/XY sex chromosomes and the master sex-determining gene *Dmrt1bY/DmY* originates from *Dmrt1* by duplication in the common ancestor ca 4-10 million years ago (Mya) (Nanda *et al.*, 2002; Kondo *et al.*, 2004).

The *Dmrt1bY/DmY* gene plays a key role in testis development, and it alone is sufficient to lead to testes differentiation (Figure 2) (Kobayashi *et al.*, 2004; Murakami *et al.*, 2007). *Dmrt1bY/DmY* plays an important

role in regulating the primordial germ cells proliferation during embryonic gonad development and was expressed at gonadal undifferentiated stage (Figure 2). The *Dmrt1* itself was expressed only after the testis differentiation was completed (Kobayashi *et al.*, 2004; Herpin *et al.*, 2007). In mature testes both *Dmrt1* and *Dmrt1bY/DmY* were expressed in Sertoli cells. *Dmrt1bY/DmY* expression was lower in mature testes, yet *Dmrt1* expression was relatively higher and was also detected in the testis of XX sexual reversals, indicating *Dmrt1* is crucial in spermatogenesis in medaka fish. Together it suggests separate roles for each gene, *Dmrt1bY/DmY* for testis formation and differentiation and *Dmrt1* for spermatogenesis (Kobayashi *et al.*, 2004). Interestingly, examination of one chimeric fish where germ cells consisted of XX cells but the somatic cells of the testis contained both female (XX) and male (XY) cells, indicating that XY cells with *Dmrt1bY/DmY* can be sufficient for testis development (Shinomiya *et al.*, 2002; Murakami *et al.*, 2007). Neither *Dmrt1* nor *Dmrt1bY/DmY* was expressed in ovaries (Kobayashi *et al.*, 2004).



**Figure 2.** Diagram of the key sex-determining pathway of *Dmrt1* and its paralog *DmY*, *DmW*, being characterized as the master sex-determining gene in medaka fish, African clawed frog, chicken and red-eared slider turtle. In vertebrates, all gonads start developing as undifferentiated gonads and then gonadal sex differentiation progresses. The master sex-determining gene *Dmrt1* and its paralogs initiate their expression either prior or at the onset of gonadal sex differentiation during early embryogenesis. The main gene network directly regulated by *Dmrt1* and its paralogs are illustrated in each case which ultimately leads to differentiated gonads at the adult stage. Gene regulator network illustration is adapted from previous studies for fish, frog chicken and turtle respectively (Miura *et al.*, 1998; Hediger Niessen *et al.*, 2004; Sánchez & Chaouiya, 2018; Li *et al.*, 2022).



Spontaneous XX sex-reversed males are not uncommon in medaka fish. XX individuals can develop as fully functional males even without presence of *Dmrt1bY/DmY* due to the presence of yet-to-be identified autosomal modifiers (Nanda *et al.*, 2003). Noticeably, at 34°C all XX individuals would develop as XX sex-reversed males, indicating temperature as an important factor in inducing XX sex reversals in medaka (Nanda *et al.*, 2003). Interestingly, no XY sex reversals (XY females) were found at low temperature treatment, suggesting *Dmrt1bY/DmY* expression was not affected by temperature. On the other hand, high temperature also led to expression of *Dmrt1* during early embryonic development (Hattori *et al.*, 2007), indicating an important role of *Dmrt1* in the testis development in the XX sex-reverted individuals. Taken together, *Dmrt1bY/DmY* is sufficient, however not essential, to induce male pathway development. On the other hand, autosomal copy of *Dmrt1* seems crucial for the development and maintenance of functional testis (Masuyama *et al.*, 2012).

## 5.2 *DmW* as master sex-determining gene in the African clawed frog

A paralog of *Dmrt1*, *DmW*, located on the W chromosome, was identified as the master sex-determining gene in the African clawed frog *Xenopus laevis* (Figure 2) (Yoshimoto *et al.*, 2008, 2010). Due to allopolyploidization in the ancestral species, *X. laevis* possesses two paralogs of *Dmrt1*, *Dmrt1.L* (*DMRT1 $\alpha$* ) and *Dmrt1.S* (*DMRT1 $\beta$* ), located on the autosomal chromosomal pairs L1 (1) and S1 (2), respectively (Yoshimoto *et al.*, 2008; Bewick *et al.*, 2011). *DmW* was the result of partial duplication of *Dmrt1.S* (*Dmrt1 $\beta$* ), specifically exon 2 and 3, and it lacked the transactivation domain-coding region (Yoshimoto *et al.*, 2008, 2010; Mawaribuchi *et al.*, 2017a). The *Dmrt1* duplication occurred approximately 47 Mya after allotetraploidization, and *DmW* was found in several closely related *Xenopus* species (Bewick *et al.*, 2011; Mawaribuchi *et al.*, 2017b; Cauret *et al.*, 2020, 2023). Other than *X. laevis*, *DmW* was not female-specific in most *Xenopus* species except in *X. gilli* (Cauret *et al.*, 2020, 2023). Most likely it was inherited from the most recent common ancestor of *X. laevis* and *X. gilli* and later it got lost in other species (Cauret *et al.*, 2020, 2023). The role of *DmW* in female gonad differentiation of *X. gilli* is yet to be investigated.

*DmW* is required for ovary development in females in *X. laevis*, and it functions as an antagonistic competitor of *Dmrt1* (Figure 2) (Yoshimoto *et al.*, 2010). Both *Dmrt1* and *DmW* bind to the same regulatory regions of target genes (yet-to-be-discovered) that are crucial in testes development. During the embryonic gonad development, *DmW* mRNA was more abundant than *Dmrt1* mRNA in the primordial gonads of ZW tadpoles early in sex determination, which could suppress the male-determining pathway by antagonizing *Dmrt1* transcriptional activity to prevent the target gene interacting with *Dmrt1*, leading to *Dmrt1* transcriptional repression (Figure 2). As a result, *DmW* inhibits the gene cascades for testis formation, resulting in ovary formation. *DmW* is missing in ZZ embryos and *Dmrt1* could bind to the target gene to initiate the transcriptional activity, which directs the sex-determining pathway towards testes development (Yoshimoto *et al.*, 2010; Yoshimoto & Ito, 2011). Interestingly, the WW individuals evolve as phenotypically normal females, suggesting that the Z-specific region on the Z chromosome is not required for development of females (Mawaribuchi *et al.*, 2017b), which further confirms the major role of *DmW* in sex determination of *X. laevis*. However, it is possible that there are factors functioning upstream to *DmW* that have yet to be discovered (Yoshimoto *et al.*, 2008; Cauret *et al.*, 2023).

### 5.3 *Dmrt1* as master sex-determining gene in chicken

*Dmrt1* is on the Z chromosome of chicken (*Gallus gallus*), which led to the speculation of the role of *Dmrt1* in bird sex determination for years (Smith *et al.*, 1999, 2007; Nanda *et al.*, 2000; Ellegren, 2001). The function of *Dmrt1* in gonadal differentiation was finally confirmed with *Dmrt1* knockdown in ZZ (normally male) chicken embryos, which led to feminization with ovarian development instead (Smith *et al.*, 2009). *Dmrt1* is thus required for proper testis formation and function in chicken (Smith *et al.*, 2009). Over-expression of *Dmrt1* in ZW females led to increased *Sox9* (possibly *Amh* too) expression and induced male sex-determining pathway (Figure 2). This further antagonised the female pathway by reducing the expression of the critical feminizing enzyme aromatase in embryonic chicken gonads (Lambeth *et al.*, 2014).

Because birds lack global Z chromosome-level dosage compensation, and chicken embryos need two functional copies of *Dmrt1* in order to develop into males with functional testes (Smith *et al.*, 2009; Lambeth *et al.*, 2014), it was proposed that *Dmrt1* was an essential sex-linked regulator of gonadal differentiation in the chicken and probably across avians, and it likely acts with a dosage mechanism (Lambeth *et al.*, 2014). Indeed, knockout of one *Dmrt1* allele in ZZ embryos, with CRISPR-Cas9-based monoallelic targeting approach, led to developing ovaries in ZZ male chicken with a single functional copy of *Dmrt1*. This demonstrated the avian sex-determining mechanism is based on *Dmrt1* dosage (Ioannidis *et al.*, 2021). Additionally, *Dmrt1* knockout in ZZ male chicken embryos led to incomplete feminization of the gonadal tissue, leading to development of ovotestis, and the genome-wide gene expression were not completely altered to ovary-like expression. Knockout of one *Dmrt1* allele also led to lower fitness and higher mortality in both ZZ and ZW individuals, and germ cells were not fully developed in the knockout individuals (Ioannidis *et al.*, 2021; Lee *et al.*, 2021), which together suggests crucial role of *Dmrt1* in germ cell development and thus reproduction of both sexes. Since all birds possess homologous sex chromosomes and higher expression of *Dmrt1* prior to gonad differentiation was described not only in chicken but also in zebra finch and emu (Hirst *et al.*, 2017), it is thus possible that *Dmrt1* is required for gonadal sex differentiation in most birds (Table S1). Furthermore, it is possible that there is a female-specific gene on the W chromosome that functions as an upstream regulator to suppress *Dmrt1* gene that is yet to be discovered.

### 5.4 *Dmrt1* as master regulator of temperature sex determination in turtle

The molecular mechanism of ESD has been a long-standing mystery until a recent discovery in the red-eared slider turtle *Trachemys scripta*. Prior to the study, the thermosensitive genetic triggers for gonadal sex differentiation remained largely unknown. Recent studies found the *Dmrt1* expression was temperature-dependent and showed sexually dimorphic pattern preceding gonadal sex differentiation (Ge *et al.*, 2017). The study showed *Dmrt1* expression responded rapidly to temperature change and aromatase inhibitor treatment. Gene function analysis demonstrated *Dmrt1* was necessary and sufficient for initiation of testes formation and development (Kettlewell *et al.*, 2000; Ge *et al.*, 2017). The findings suggest that *Dmrt1* is candidate master regulator of temperature-dependent sex determination (Ge *et al.*, 2017). *Dmrt1* expression was further regulated by the methylation in its promoter region (Ge *et al.*, 2018), and it is repressed in female producing temperatures by phosphorylated signal transducer and activator of transcription 3 (*STAT3*), the phosphorylation is induced by calcium signalling pathways induced by high temperatures (Weber *et al.*, 2020).

## 6. Role of epigenetic regulation of *Dmrt1* in sexual differentiation

Epigenetics is the study of changes in gene function that cannot be explained by changes in DNA sequences (Russo *et al.*, 1996). There are three main epigenetic regulation mechanisms: DNA methylation, modification of histones and the involvement of non-coding RNAs (ncRNAs) (Piferrer, 2013). Epigenetic regulation allow organisms to integrate environmental and genetic factors to modify their gene activities for a particular phenotype, and play an essential role in various biological processes across the tree of life, such as germline imprinting, X-chromosome inactivation in mammals, mating-type silencing in yeast, temperature-dependent vernalization in plants (Piferrer, 2013, 2021; Gunes *et al.*, 2016; Tachibana, 2016).


Sex determination is typically categorized as genetic (GSD) or environmental mechanism (ESD), where genetic (gene or genes) or environmental (temperature, PH, social cues, salt etc.) factors functioning as the primary signal of the pathway to develop as a female or a male (Beukeboom & Perrin, 2014). However, sex determination in many species depends on the contribution of both genetic and environmental factors. It has been widely accepted that sex determination is a spectrum between pure GSD and ESD, where GSD and ESD only represent the opposite ends of continuum (Beukeboom & Perrin, 2014; Straková *et al.*, 2020). Epigenetic regulation of sex determination strictly is not equal to ESD, because the epigenetic mechanisms such as DNA methylation, histone modification and ncRNAs can be involved in the whole spectrum of sex determination from GSD to ESD regardless of the primary triggering factors. For instance, epigenetic regulation can take place in GSD, or in systems where GSD interacts with ESD, or pure ESD species (Figure 3). The molecular basis of sex-determining pathway of GSD has been extensively studied and in some cases is well understood (Figure 2), however, the underlying genetics of ESD are largely unknown. Furthermore, the role of epigenetic regulation on both GSD and ESD has not been well studied and more efforts thus should be carried on in such directions.

In this section, we delve into the emerging role of epigenetic regulation in sex determination, particularly focusing on the key sex-determining gene *Dmrt1*. While *Dmrt1*'s canonical function in orchestrating gonadal sex differentiation is well-established, the epigenetic regulation of *Dmrt1* and its implications for sex determination represent a novel frontier in the field. Here, we synthesize current knowledge and recent discoveries regarding the epigenetic regulation of *Dmrt1*, offering insights into its multifaceted role in sexual differentiation across diverse vertebrate taxa. By exploring the epigenetic landscape of sex determination, we aim to shed light on the dynamic interplay between genetic and environmental factors underlying sex-determining pathways.

Regarding epigenetic regulation of *Dmrt1*, comparing to other epigenetic forms such as modification of histones and ncRNAs involvement, methylation of *Dmrt1* DNA sequence (i.e. mostly at the promoter region) in the nucleus is a bit more explored and so far has been the main characterized epigenetic modification (Figure 3). Methylation occurs when a methyl group replaces carbon 5' of a deoxycytidine adjacent to a guanidine (CpG), which islands are genomic regions with elevated CpG content that are typically associated with promoter or regulatory regions (Li *et al.*, 2015; Mugal *et al.*, 2015). Changes in methylation levels in the CpG islands at the promoter regions can regulate the level of *Dmrt1* expression.

### 6.1 Epigenetic regulation of *Dmrt1* in species with ESD

To our best knowledge, the only study that demonstrates the molecular mechanism of ESD is in the red-eared slider turtle (*Trachemys scripta*) (Figure 2 and 3) (Ge *et al.*, 2018; Georges & Holleley, 2018). This species has temperature-dependent sex determination (TSD), where embryos exposed at low temperature at thermal sensitive developmental period develop as males while at high temperature they develop as females. Studies have shown that the histone H3K27 (H3 lysine 27) demethylase KDM6B had sexually dimorphic expression, which was temperature-dependent during the early embryonic stage before the gonad showed sexual differentiation (Figure 2) (Ge *et al.*, 2017, 2018). Knockdown of the *Kdm6b* gene at low temperature (i.e. 26°C typically develop as males) led to male-to-female sex reversals in most surviving embryos. In addition, the authors found KDM6B directly regulated the transcription of *Dmrt1* by removing the trimethylation of H3K27 near the promoter of *Dmrt1* (Ge *et al.*, 2018). *Kdm6b* disruption led to an overexpression of *Dmrt1*, which rescued the sex reversal and developed as males. Finally, a recent study identified a link between temperature and the activation of signal transducer and activator of transcription 3 (STAT3), which regulated the expression of *Kdmtb* (Weber *et al.*, 2020). They found that at warmer, female-producing temperature, an influx of Ca<sup>2+</sup> initiated, which led STAT3 to be phosphorylated that silenced *Kdm6b* transcription. The silencing of *Kdm6b* resulted in suppressing of *Dmrt1* expression, which ultimately repressed testis development (Figure 2) (Weber *et al.*, 2020). Taken together, the study found the epigenetic mechanism is responsible for TSD by regulating *Dmrt1* expression at the early embryonic stages. The next step would be to investigate more ESD species to understand how broad this epigenetic mechanism is, before a general molecular mechanism for ESD can be drawn.



**Figure 3.** The model of epigenetic regulation of *Dmrt1* involving in sex determination and gonadal sex differentiation across the spectrum of sex determination, from GSD to ESD, is illustrated. The primary signals in sex determination range from GSD to ESD. Epigenetic regulation mechanism has been found in all spectrum of sex-determining systems, GSD, GSD interacts with ESD and ESD. In the GSD in the chicken, it was shown that MHM region (male hypermethylated region) is located next to the *Dmrt1*. MHM is hypermethylated that blocks the ncRNA which leads to upregulation of *Dmrt1*, resulting in testis development in males. In females, MHM is hypomethylated that activates the ncRNA which downregulates *Dmrt1*, resulting in ovary development in females. In many fishes with GSD interaction with ESD, the current epigenetic regulation has found methylation of *Dmrt1* promoter region to suppress its expression or involvement of transposable elements associated methylation mechanism which is yet to discover the exact molecular basis.

## 6.2 Epigenetic regulation of *Dmrt1* in species where genetic factors interact with environmental factors

### 6.2.1 Via methylation of *Dmrt1* promoter region

Epigenetic mechanism of sex determination has been also beautifully illustrated in a flatfish with sex determination involving interaction between genetic and environmental factors (Figure 3). In a marine flatfish, the Chinese half-smooth tongue sole (*Cynoglossus semilaevis*), there is a female heterogametic (ZZ/ZW) sex chromosome system, the Z-linked *Dmrt1* was a putative male-determining gene, and the spermatogenesis of *Dmrt1* mutant ZZ male was disrupted and developed ovary-like testis, suggesting *Dmrt1* was an essential male sex-determining gene (Chen *et al.*, 2014; Zhou & Chen, 2016; Cui *et al.*, 2017; Wang *et al.*, 2019). Furthermore, elevated temperature was shown to lead to hypomethylation of the *Dmrt1* promoter in ZW embryos, which increased the *Dmrt1* expression at the time of sex determination in early embryogenesis, resulting in ZW pseudo-male development. Further crossing the ZW pseudo-male with standard ZW female revealed extremely male-biased offspring (94% of all offspring), suggesting the methylation change can be inherited across generations (Chen *et al.*, 2014). All these together demonstrate that genetic factor interacts with environmental factor such as temperature to determine sex in this species, and temperature functions as a stimuli to alter *Dmrt1* gene expression via epigenetic methylation mechanism to lead to development of sex reversals. In the Nile tilapia (*Oreochromis niloticus*) with male heterogametic (XX/XY) sex chromosome system, functional studies show that once *Dmrt1* was mutated, the gonad could not be rescued to develop functional testis (Qi *et al.*, 2024). This suggests *Dmrt1* is the only male sex-determining pathway gene indispensable for sex determination and functional testis development in this species. Furthermore, high temperature at crucial early embryogenesis leads to male development in this species with methylation assay (Wang *et al.*, 2018).

Other than epigenetic regulation of *Dmrt1* directly affects sex determination and leads to sex reversals in the aforementioned studies, it also shows effects on gonadal sex differentiation. For instance, in the Asian ray-finned fish (*Culter alburnus*), one study found that CpG methylation pattern of the *Dmrt1* promoter was highly sexually dimorphic, where it was hypermethylated in the ovaries but not in the testis. This leads to very high *Dmrt1* expression in testis and almost non-detected in ovaries and several other somatic tissues (Jia *et al.*, 2019), suggesting epigenetic modifications may play an important role in the gonad differentiation in this fish. In the Japanese flounder (*Paralichthys olivaceus*), the 57-69% *Dmrt1* promoter CpGs was methylated in the ovary but not in testis, leading to *Dmrt1* expression 70 times higher in the testis than in the ovary (Wen *et al.*, 2014). This suggests that epigenetic modification affects the *Dmrt1* gene expression which associates with gonadal differentiation. In the European sea bass (*Dicentrarchus labrax*), the sex is determined by genetic and temperature factors. Studies show sex-specific methylation and temperature response for *Dmrt1* and *cyp19a1a* genes in juvenile gonads, suggesting epigenetic regulation is possibly involved in gonadal

differentiation in this species (Anastasiadi *et al.*, 2018). In the blotched snakehead (*Channa maculata*), the CpG methylation level of the *Dmrt1* promoter was correlated negatively with its gene expression in the testis and ovary, with *Dmrt1* being expressed 15 times higher in the testis than in the ovary during gonadal development. Steroid treatment affected *Dmrt1* expression and could induce male to female sex reversal. This study suggests *Dmrt1* is involved in testis differentiation and spermatogenesis in this species and epigenetic regulation can explain the gonadal differentiation throughout development (Ou *et al.*, 2022).

### 6.2.2 Involving transposable elements (TEs)

In the Siamese fighting fish (*Betta splendens*), association mapping study shows that *Dmrt1* was located in the identified small sex-determining region and demonstrated a functional role where knockout of *Dmrt1* led to ovarian development in individuals with XY genotypes (Wang *et al.*, 2022b). Furthermore, this study found that a transposable element *Drbx1* inserted in the fourth intron of X-linked *Dmrt1* copy, resulting in epigenetic modifications to the neighboring *cis*-regulatory elements in the *Dmrt1* promoter region, confirmed with methylation assays (Figure 3) (Wang *et al.*, 2022b). Similarly like in other fishes described above, this study again shows methylation on *Dmrt1* promoter regions regulate the gene expression and leads to sex reversal.

### 6.3 Epigenetic regulation of *dmrt1* in species with GSD

*Dmrt1* was identified as master sex-determining gene in chickens, and two copies of *Dmrt1* expression were required during early embryogenesis to initiate testis development (Smith *et al.*, 2009). A region neighboring *Dmrt1* is differentially methylated between males and females. In males, this region is hypermethylated, male hypermethylated region (MHM), and is transcriptionally inactive, whereas this region in females is hypomethylated and transcribed into a long ncRNA (Teranishi *et al.*, 2001). MHM is hypothesized to inhibit *Dmrt1* expression in females, as the MHM transcript accumulates at the site of transcription adjacent to *Dmrt1*. A functional study with injection of MHM into rooster testicles led to a strong downregulation of *Dmrt1* and a paling of the comb color, suggesting a change in sex hormones (Figure 3). Other studies show other sex-determining genes such as *Amh* and *Sox9* were unaffected at transcriptional levels, suggesting that MHM specifically regulates *Dmrt1* (Figure 3) (Yang *et al.*, 2010).

## 7. Role of alternative splicing of *Dmrt1* in gonadal sex differentiation

Alternative splicing refers to the mechanism where different exons or a fraction of an exon are joined in different combinations, to produce unique but related mRNAs transcripts, from a single pre-mRNA (Wang *et al.*, 2015). Alternative splicing leads to an increase in the coding capacity of genes and therefore an increase of proteome complexity, constituting a ubiquitous regulatory mechanism of gene expression. Thus alternative splicing was considered one key mechanism to contribute to phenotypic diversity among animal and plant taxa (Wright *et al.*, 2022). One of the most fundamental developmental processes which alternative splicing is contributing to is sex determination (Wright *et al.*, 2022). The essential role of alternative splicing on sex determination has been extensively studied in a shared common pathway composed of the *Transformer* (*Tra*) gene and its downstream target *Doublesex* (*Dsx*) (Suzuki *et al.*, 2001; Pomiankowski *et al.*, 2004; Cho *et al.*, 2007; Beukeboom & Zande, 2010). Female spliced *tra* transcripts give rise to functional TRA proteins that directly regulate splicing of female *Dsx* mRNAs, producing female DSX proteins and result in female development. When *Tra* is spliced into the male variant, no TRA proteins are produced. This results in splicing


of male isoform *Dsx* mRNAs, male DSX proteins and male development (Figure 4A) (Verhulst *et al.*, 2010; Gempe & Beye, 2011). Remarkably, sex-specific splicing of *Tra* and *Dsx*, key regulation genes in sex-determining and sexual differentiation in *Drosophila melanogaster*, is conserved across diverse insects, including the silk moth *Bombyx mori*, red flour beetle *Tribolium castaneum*, love fruit flies *Bactrocera oleae*, the housefly *M. domestica*, the medfly *Ceratitis capitata*, the honeybee *Apis mellifera* and the jewelry wasp *Nasonia vitripennis*, suggesting an ancient origin and role in sex determination and sexual differentiation (Cho *et al.*, 2007; Giebel *et al.*, 2009; Verhulst *et al.*, 2010; Geuverink & Beukeboom, 2014; Van De Zande & Verhulst, 2014). *Dmrt1* is the ortholog gene of *Dsx* in vertebrates, suggesting the strong conservative role in sex determination in eukaryotes (Vicoso, 2019).

In sharp contrast to the well-studied sex-specific splicing role of *Dsx* in sex determination in insects, sex-specific splicing role of *Dmrt1* in vertebrates have recently been explored, mostly focusing on adult or developmental expression studies and no functional study on isoform level has been done yet. We have compiled in total 16 vertebrate species with alternative splicing variants (Table 2), with 10 fishes, 2 mammals (human and mouse), 4 reptiles including avians (1 chicken, 1 turtle, 1 crocodile and 1 alligator). There are clear patterns we could observe from these studies. First, it is unclear whether alternative splicing is directly involved in the genetic mechanism of sex determination in vertebrates. For most of these species, master sex-determining gene has not yet been identified or verified. For chicken with *Dmrt1* being the master sex-determining gene, it is unclear whether and how splicing variants play a role in sex determination. Second, unlike the constitutional sex-specific splicing variants of *Dsx* which generates female and male-specific DSX protein in insects, there were no sex-specific splicing variants in any of the studied 16 species (Table 2). Third, there are, however, dynamics of temporal expression prior or during gonadal sex differentiation and adult of sex-limited or sex-biased expression for one or all of the *Dmrt1* isoforms, as well as variation in expression breadth across adult tissues (Table 2, Figure 4B). One study showed exclusive expression in testis in both adult and gonadal sex differentiation stages, both showed male-limited expression, suggesting a role in testis formation and function (Raghuvver & Senthilkumaran, 2009). Several (6) species showed one or all *Dmrt1* isoforms were exclusively expressed in adult testes or developmental testes (Huang *et al.*, 2005a; Lu *et al.*, 2007; Zhao *et al.*, 2007a; Deloffre *et al.*, 2009; Su *et al.*, 2015; Domingos *et al.*, 2018). The rest 10 species showed significant higher expression in adult testis, and during gonadal sex differentiation or transformation to males from females (Table 2). All these expression dynamics suggest a crucial role of *Dmrt1* isoforms in testicular formation and/or function. Lastly, to our best knowledge, so far there have been none reported alternative splicing in any amphibians. Finally, a broader spectrum of taxa and functional studies are needed to better understand its role in sex determination and sexual differentiation.

### 7.1 Alternative splicing of *Dmrt1* in fishes

Sex-determining mechanism in fishes is extremely diverse and labile, governed by genetic, environmental factors, or their interactions. Certain teleost fishes undergo natural sequential sex change, potentially an ideal system to study sex determination and sexual differentiation on which one sex is transformed into the other. Among the 10 fishes with alternative splicing of *Dmrt1*, the isoform number ranges from 2 to 6 (Table 2, Figure 4C). Half of all such species do not show, at any developmental stage, male-specific splicing of *Dmrt1*, and all isoforms showed higher expression in adult testes or developmental testes (Figure B). For instance, Zebrafish (*Danio rerio*) is shown to have ZZ/ZW sex chromosome system with a

polygenic sex-determining mechanism with varying contributing loci including *Dmrt1*, *Amh*, *Sox9*, *Cyp19a1* and *Foxl2a* (Kossack & Draper, 2019). In this species, alternative splicing has been found in *Dmrt1* with three different isoforms (*Dmrt1a*, *Dmrt1b*, *Dmrt1c*) which encoded proteins with 267, 246 and 132 amino acids. All three isoforms are expressed in both testis and ovary but significantly highly expressed in testis (Guo *et al.*, 2005).



**Figure 4.** Sex-specific splicing of *Dsx* in insects and alternative splicing of *Dmrt1* in vertebrates. A. Sex-specific splicing of *Dsx* in *Drosophila melanogaster* is illustrated and can be found in all insects. Illustration is adapted from (Hediger Niessen *et al.*, 2004). The female isoform of *Dsx* mRNAs can produce functional female DSX protein and splicing of male isoform *Dsx* mRNAs generates male DSX protein. B. Summary of expression dynamics of *Dmrt1* alternative splicing isoforms, illustrated by example of 2 splicing isoforms from several vertebrates (Table 2). *Dmrt1* does not have female or male-specific splicing isoforms as insects, instead different isoform (or all isoforms) could have stage-specific male-limited expression, as illustrated here isoform B has male-limited expression at gonadal sex differentiation stage and isoform A has male-limited expression in adult testis. All isoforms typically have strong male-biased expression during gonadal sex differentiation and in testis than ovary, and often they either do not have or have very weak expression in somatic tissues such as brain (B), kidney (K) and liver (L). In some studies, isoforms also start



expressing and with male-biased expression in undifferentiated gonads during embryogenesis. C. One example of illustration of alternative-splicing isoforms from each major vertebrate group except amphibian, and the species from each group was chosen to have the highest number of isoforms found in that group. \*: the isoform number range found in that group. Both within and across vertebrate groups, the alternative splicing isoform variant do not have the fixed mechanism or pattern, and it seems to be rather random. For example, isoform b in fish is splicing out of exon 2 and exon 3, in reptile is splicing out of exon 5 and retaining a bit of intron region, in chicken is splicing out of exon 5 and retaining both intron 4 and intron 5 region, in mouse is splicing out exon 5 and exon 4 is joining between the entire exon 4 and a fraction of exon 5.

In certain species all isoforms showed testis-specific expression at adult stage but were all expressed in both sexes during gonadal sex differentiation process. For instance, in the rice field eel (*Monopterus albus*) with protogynous hermaphrodite, the 4 isoforms of *Dmrt1* (encoding proteins with 301, 196, 300 and 205 amino acids) were exclusively expressed in adult gonads. Two of the splice forms, *Dmrt1a* and *Dmrt1b*, had an expression from ovary to testis during the gonadal transformation that occurs in the sex change process (Huang *et al.*, 2005b). In the European sea bass (*Dicentrarchus labrax*), a teleost fish with polygenic sex determination induced by temperature, two alternative splicing of *Dmrt1* were detected (*Dmrt1a* and *Dmrt1b*), with *Dmrt1a* splicing out exon4 (Deloffre *et al.*, 2009). Both transcripts were expressed at a similar level and firstly detected at 150 days post-hatch coinciding with the start of sex differentiation. With development, the expression increases in testis and decreases in ovary, and only expressed in testis tissues in adults. Both transcripts were suggested to be required for testicular function (Deloffre *et al.*, 2009).

In the fish barramundi (*Lates calcarifer*), a protandrous hermaphrodite species, two isoforms were identified, *Dmrt1a* and *Dmrt1b*. Interestingly, different isoforms showed different sex-limited expression (Domingos *et al.*, 2018). The isoform *Dmrt1a*, which composes the DM domain, showed exclusive expression in adult testis, while *Dmrt1b* (lacking this DM domain) was detected in testis and ovaries, with higher expression in testis (Domingos *et al.*, 2018).

### 7.2 Alternative splicing of *Dmrt1* in mammals

There are 3 and 4 alternative splicing variants of *Dmrt1* identified in human and mouse respectively (Figure 4C). Their predominant expression in testis indicates a certain role in testis function, yet whether and how each isoform functions are unclear. In human, all 3 *Dmrt1* isoforms were expressed in testis (Cheng *et al.*, 2006). However, sex-specific splicing was not concluded due to expression was only conducted in the testis tissue. In mouse, all 4 *Dmrt1* isoforms showed testis-limited expression at adult stage, but all were expressed during gonadal sex differentiation process with higher expression in males (Huang *et al.*, 2005a).

### 7.3 Alternative splicing of *Dmrt1* in reptiles

Reptiles have 2-10 isoforms with a highest number 10 in the alligator species *Alligator sinensis* (Figure 4C) (Table 2). In many reptile species, *Dmrt1* expression is upregulated in male embryos. In the painted turtle (*Chrysemys picta*) with TSD, *Dmrt1* is upregulated in males during gonadal development. Study further found that alternative splicing of *Dmrt1* were differentially expressed by temperature at stage 19 and 22 in developing gonads after the onset of sex determination. Both splicing variants displayed a significant male-biased expression pattern when exposed to male-producing temperatures (Mizoguchi & Valenzuela, 2020). In the crocodile *Crocodylus palustris* and the alligator *Alligator sinensis*, they have 2 and 10 *Dmrt1* isoforms respectively (Figure 4C). No isoform was sex-specific as they were expressed in both testis and ovary tissues, as well as during gonadal sex differentiation process.

All birds have genetic sex determination with female heterogametic (ZZ/ZW) sex chromosome system. The *Dmrt1* gene has been identified as master sex-determining gene in chicken with dosage effects (Smith *et al.*, 2009). Chicken *Dmrt1* generates 6 different splicing variants during gonadal sex differentiation and their expression were testis specific in adults. The splicing variants had differential expression along the different stages, suggesting stage-specific splicing variants. *Dmrt1b* was exclusively expressed in embryo gonads and

higher in male than female gonads at stage 31 which was the key time of gonadal sexual differentiation. *Dmrt1c* was highly expressed in female than male gonads at stage 31. *Dmrt1f* was detectable only in male gonads at stage 31 (Zhao *et al.*, 2007a).

## 8. Conclusion

Nine *Dmrt* genes with conserved DM domain with DNA-binding motif distribute across vertebrate groups. *Dmrt1* is the most studied gene within *Dmrt* family regarding transcription regulation of DMRT protein and plays a crucial role in regulating sex determination and gonadal differentiation across vertebrates. From our compiled vertebrate species, it showed that *Dmrt1* was expressed in the undifferentiated gonads in certain species. Mostly *Dmrt1* shows sexually dimorphic expression, being significantly higher or exclusively expressed, in males during gonadal sex differentiation and predominantly expressed in adult testes, with minimal or no expression in ovaries and other somatic tissues. Notably, *Dmrt1* has been identified as a master sex-determining gene in chicken, frog, fish, and reptile species, and proposed as a candidate in additional 12 species (6 fishes, 2 amphibians, 4 birds).

Recent studies highlight the role of epigenetic regulation, involving methylation of the *Dmrt1* promoter region, transposable element insertion and ncRNA involvement, in sex determination and gonadal sex differentiation. Furthermore, alternative splicing of *Dmrt1*, akin to the sex-specific role of *Dsx* in insects, showed no sex-specific splicing pattern in all studied species but with sex-limited expression in males in certain species, highlighting a major difference in the regulatory mechanism between *Dmrt1* in vertebrates and *Dsx* in insects.

One open question is how generalizable the molecular mechanisms identified in the red-eared slider turtle are across vertebrate taxa. Future research should focus on investigating the molecular basis of ESD in a broader spectrum of species. Furthermore, it is crucial to explore the role of epigenetic regulation in the continuum of sex-determining systems, ranging from GSD to systems where GSD interacts with ESD, to pure ESD. Detailed molecular studies are needed to understand these underlying mechanisms. Finally, the alternative splicing of *Dmrt1* requires deeper understanding. It is essential to understand why and how multiple (up to 10) alternative splicing variants of *Dmrt1* evolve and the specific roles each isoform plays in sex determination and gonadal sex differentiation. Another interesting aspect is the significant differences in the molecular mechanisms and functions of alternative splicing between *Dmrt1* in vertebrates and sex-specific splicing of *Dsx* in insects. Understanding the differences could provide deeper insights into the evolution of sex-determining mechanism between vertebrates and insects.

**Author Contributions:** W.-J. M. designed the study. B. A. and W.-J. M. collected data and synthesized ideas from literature. B. A. visualized the synthesized ideas in figures with significant input from W.-J. M. W.-J. M. and B. A. wrote the manuscript. All authors have read and agreed to the submitted version of the manuscript.

**Data Availability Statement:** NA.

**Acknowledgments:** We thank Paris Veltsos for insightful discussion and comments on the manuscript. This work is funded by the ERC starting grant (FrogWY, 101039501) and a starting grant from Vrije Universiteit Brussel (OZR4049) to Wen-Juan Ma. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

**Conflicts of Interest**

The authors declare no conflict of interest.

Table 1. List of species across various vertebrate groups which *Dmrt1* was identified as master, or candidate master, sex-determining gene, and the expression pattern throughout gonadal development and at adult stages.

Class	Species	SD system	Master SD gene	Expression during gonadal sex differentiation	Expression in gonads of sub-adult/adult	Expression in adult somatic tissue	Method	Gene function	Reference
Actinopterygii	<i>Oryzias latipes</i>	GSD (XX/XY)	yes ( <i>Dmrt1bY/DmY</i> )	testes ( <i>Dmrt1a</i> , <i>Dmrt1bY/DmY</i> )	testes ( <i>Dmrt1bY/DmY</i> ; <i>Dmrt1a</i> ), ovaries <i>Dmrt1a</i>	<i>Dmrt1bY/DmY</i> expressed: pituitary, heart, liver, spleen; <i>Dmrt1bY/DmY</i> very weak expression (intestine, eye); <i>Dmrt1a</i> expressed: spleen (males); <i>Dmrt1a</i> very weak expression: spleen (females); not expressed: brain, muscle, gills	PCR, histology	SD, gonad differentiation ( <i>Dmrt1bY/DmY</i> ), spermatogenesis ( <i>Dmrt1a</i> )	Ohmuro-Matsuyama <i>et al.</i> , 2003; Kobayashi <i>et al.</i> , 2004
Actinopterygii	<i>Astyanax scabripinnis</i>	GSD (ZZ/ZW)	candidate	NA	testes	not examined	qPCR	NA	Castro <i>et al.</i> , 2019
Actinopterygii	<i>Cynoglossus semilaevis</i>	GSD (ZZ/ZW)	candidate	testes	testes	not examined	qPCR	testes differentiation, spermatogenesis	(Cui <i>et al.</i> , 2017)
Actinopterygii	<i>Betta splendens</i>	GSD (XX/XY)	candidate	males (testes)	testes	not examined	qPCR	gonad differentiation	Wang <i>et al.</i> , 2022a
Actinopterygii	<i>Scatophagus argus</i>	GSD (XX/XY)	candidate	NA	testes	not examined	PCR	NA	Mustapha <i>et al.</i> , 2018
Actinopterygii	<i>Scatophagus tetracanthus</i>	GSD (XX/XY)	candidate ( <i>Dmrt1Y</i> )	NA	testes ( <i>Dmrt1X</i> , <i>Dmrt1Y</i> ), ovaries ( <i>Dmrt1X</i> weak)	not examined	qPCR, PCR	NA	Peng <i>et al.</i> , 2023
Actinopterygii	<i>Selenotoca multifasciata</i>	GSD (XX/XY)	candidate ( <i>Dmrt1Y</i> )	NA	testes ( <i>Dmrt1X</i> , <i>Dmrt1Y</i> ), ovaries ( <i>Dmrt1X</i> )	not examined	PCR, RNAseq	NA	Jiang <i>et al.</i> , 2022
Amphibia	<i>Hyla arborea</i>	GSD (XX/XY)	candidate	testes	NA	not examined	histology	NA	(Pipek <i>et al.</i> , 2013)
Amphibia	<i>Xenopus laevis</i>	GSD (ZZ/ZW)	yes ( <i>DmW</i> )	testes ( <i>Dmrt1</i> ), ovaries ( <i>Dmrt1</i> weak, <i>DmW</i> )	testes ( <i>Dmrt1</i> ), ovaries ( <i>DmW</i> weak, <i>Dmrt1</i> weak)	<i>DmW</i> expressed: liver; very weak expression mesonephros; <i>DmW</i> not expressed: intestine, brain, stomach, spleen, heart	qPCR, PCR, histology, Southern blot	NA	Yoshimoto <i>et al.</i> , 2008a, 2010 (Fujitani <i>et al.</i> , 2016; Mawaribuchi <i>et al.</i> , 2017b)
Amphibia	<i>Rana temporaria</i>	GSD (XX/XY)	candidate	testes (strong), ovaries (weak)	NA	not examined	histology	NA	Pipek <i>et al.</i> 2014

Aves	<i>Cairina moschata</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR, RNAseq	SD, gonad differentiation	Wang <i>et al.</i> , 2015; Bai <i>et al.</i> , 2020
Aves	<i>Dromaius novaehollandiae</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR	SD, gonad differentiation	Hirst <i>et al.</i> , 2017
Aves	<i>Taeniopygia guttata</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR	SD, gonad differentiation	Hirst <i>et al.</i> , 2017
Aves	<i>Gallus gallus domesticus</i>	GSD (ZZ/ZW)	yes ( <i>Dmrt1</i> )	testes (strong), ovaries (weak)	testes	expressed: heart; not expressed: liver, spleen, lung, kidney	qPCR, histology	SD, gonad differentiation, germ cell proliferation	Zhao <i>et al.</i> , 2007b; Lambeth <i>et al.</i> , 2014; Zhao <i>et al.</i> , 2023
Aves	<i>Coturnix japonica</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	testes	not examined	RNAseq	SD, gonad differentiation	Okuno <i>et al.</i> , 2020
Reptilia	<i>Trachemys scripta</i>	TSD	yes ( <i>Dmrt1</i> )	testes (strong), ovaries (weak)	NA	not examined	qPCR, histology	gonad differentiation	Bieser & Wibbels, 2014; Gu & Walters, 2017

Table 2. List of vertebrates with alternative splicing of *Dmrt1* and each of their expression throughout development.

Class	Species	SD system	Isoform number	Sex-limited expression	Expression breath in adult	Expression during gonadal sex differentiation or sex change	Reference
Actinopterygii	<i>Clarias gariepinus</i>	GSD (XX/XY)	3	yes	testis	all isoforms are only expressed in developing males	Raghuveer & Senthilkumaran, 2009
	<i>Danio rerio</i>	GSD (ZZ/ZW), polygenic	3	No	<i>Dmrt1a</i> is highly expressed in testis; weaker expression of <i>Dmrt1b</i> and <i>Dmrt1c</i> testis	NA	Guo <i>et al.</i> , 2005; Kossack & Draper, 2019
	<i>Megalobrama amblycephala</i>	GSD (XX/XY)	4	yes (adult only)	testis	all isoforms are expressed in males and only one isoform is expressed in initial stage during transformation into female	Su <i>et al.</i> , 2015
	<i>Halichoeres tenuispinis</i>	Protogynous hermaphrodite	2	No	testis and ovary (higher expression in testis)	both isoforms are expressed but higher in males than females	Miyake <i>et al.</i> 2012
	<i>Lates calcarifer</i>	Protandrous hermaphrodites	2	Yes ( <i>Dmrt1a</i> ), no ( <i>Dmrt1b</i> )	<i>Dmrt1a</i> is only expressed in testis; <i>Dmrt1b</i> in both	NA	Domingos <i>et al.</i> , 2018
	<i>Dicentrarchus labrax</i>	GSD	2	yes (only at adult stage)	not in somatic tissues, but in testis	both isoforms are expressed, both are highly expressed in testis	Deloffre <i>et al.</i> , 2009
	<i>Epinephelus merra</i>	Protogynous hermaphrodite	2	No	testis and ovary	all isoforms are expressed in both sexes but higher in transformation into males	Alam <i>et al.</i> 2008
	<i>Monopterus albus</i>	Protogynous hermaphrodite	4	yes (only at adult stage)	not in somatic tissues, only in testis	all isoforms are expressed in developing gonads in both sexes, significantly higher in testis	Huang <i>et al.</i> 2005
	<i>Solea senegalensis</i>	GSD (XX/XY)	6	No	testis and ovary (higher in testis)	NA	Cross <i>et al.</i> 2020
	<i>Acipenser schrenckii</i>	GSD (ZZ/ZW)	2	No	testis and ovary (higher in testis)	both isoforms are expressed but higher in developing females	Okada <i>et al.</i> 2017
Mammalia	<i>Homo sapiens</i>	GSD (XX/XY)	3	Inconclusive	only studied in testis	NA	Cheng <i>et al.</i> 2006
	<i>Mus musculus</i>	GSD (XX/XY)	4	yes (only at adult stage)	testis	all isoforms are expressed in developing gonads in both sexes, significantly higher in males	Lu <i>et al.</i> 2007
Reptilia	<i>Crocodylus palustris</i>	TSD	8	No	testis, kidney, brain	all isoforms are expressed throughout temperature sensitive period, higher in males than females	Anand <i>et al.</i> , 2008
	<i>Chrysemys picta</i>	TSD	2	No	testis and ovary (higher in testis)	both isoforms are expressed but higher in males than females	Mizoguchi & Valenzuela, 2020
	<i>Alligator sinensis</i>	TSD	10	No	testis and ovary	NA	Wan <i>et al.</i> 2013
Aves	<i>Gallus gallus</i>	GSD (ZZ/ZW)	6	yes (only testis at developing stage 31)	testis and heart	<i>Dmrt1a</i> is not expressed; <i>Dmrt1b</i> , <i>Dmrt1c</i> , <i>Dmrt1d</i> , <i>Dmrt1e</i> are expressed in both sexes; <i>Dmrt1f</i> is only expressed in testis	Zhao <i>et al.</i> 2007; Smith <i>et al.</i> 2009

## Reference

- Alam, M.A., Kobayashi, Y., Horiguchi, R., Hirai, T. & Nakamura, M. 2008. Molecular cloning and quantitative expression of sexually dimorphic markers *Dmrt1* and *Foxl2* during female-to-male sex change in *Epinephelus merra*. *Gen. Comp. Endocrinol.* **157**: 75–85.
- Anastasiadi, D., Vandeputte, M., Sánchez-Baizán, N., Allal, F. & Piferrer, F. 2018. Dynamic epimarks in sex-related genes predict gonad phenotype in the European sea bass, a fish with mixed genetic and environmental sex determination. *Epigenetics* **13**: 988–1011.
- Ayers, K.L., Davidson, N.M., Demiyah, D., Roeszler, K.N., Grützner, F., Sinclair, A.H., *et al.* 2013. RNA sequencing reveals sexually dimorphic gene expression before gonadal differentiation in chicken and allows comprehensive annotation of the W-chromosome. *Genome Biol.* **14**: R26.
- Bai, D.P., Chen, Y., Hu, Y.Q., He, W.F., Shi, Y.Z., Fan, Q.M., *et al.* 2020. Transcriptome analysis of genes related to gonad differentiation and development in Muscovy ducks. *BMC Genomics* **21**: 1–17. BMC Genomics.
- Bellefroid, E.J., Leclère, L., Saulnier, A., Keruzore, M., Sirakov, M., Vervoort, M., *et al.* 2013. Expanding roles for the evolutionarily conserved *Dmrt* sex transcriptional regulators during embryogenesis. *Cell. Mol. Life Sci.* **70**: 3829–3845.
- Beukeboom, L.E.O.W. & Zande, L.V.A.N.D.E. 2010. Genetics of sex determination in the haplodiploid wasp *Nasonia vitripennis* (Hymenoptera : Chalcidoidea). *J. Genet.* **89**: 333–339.
- Beukeboom, L.W. & Perrin, N. 2014. The Evolution of Sex Determination. In: *The Evolution of Sex Determination*. Oxford University Press, Oxford, New York.
- Bewick, A.J., Anderson, D.W. & Evans, B.J. 2011. Evolution of the closely related, sex-related genes *dm-w* and *dmrt1* in african clawed frogs (*xenopus*). *Evolution.* **65**: 698–712.
- Bieser, K.L. & Wibbels, T. 2014. Chronology, magnitude and duration of expression of putative sex-determining/differentiation genes in a turtle with temperature-dependent sex determination. *Sex. Dev.* **8**: 364–375.
- Bista, B. & Valenzuela, N. 2020. Turtle insights into the evolution of the reptilian karyotype and the genomic architecture of sex determination. *Genes.* **11**: 416.
- Capriglione, T., Vaccaro, M.C., Morescalchi, M.A., Tammaro, S. & De Iorio, S. 2010. Differential *DMRT1* expression in the gonads of *podarcis sicula* (Reptilia: Lacertidae). *Sex. Dev.* **4**: 104–109.
- Casas, L., Saborido-Rey, F., Ryu, T., Michell, C., Ravasi, T. & Irigoien, X. 2016. Sex Change in Clownfish: Molecular Insights from Transcriptome Analysis. *Sci. Rep.* **6**: 1–19.
- Castro, J.P., Hattori, R.S., Yoshinaga, T.T., Silva, D.M.Z.D.A., Foresti, F., Santos, M.H., *et al.* 2019. Differential Expression of *dmrt1* in *Astyanax scabripinnis* (Teleostei, Characidae) Is Correlated with B Chromosome Occurrence. *Zebrafish* **16**: 182–188.
- Cauret, C.M.S., Gansauge, M.T., Tupper, A.S., Furman, B.L.S., Knytl, M., Song, X.Y., *et al.* 2020. Developmental Systems Drift and the Drivers of Sex Chromosome Evolution. *Mol. Biol. Evol.* **37**: 799–810.
- Cauret, C.M.S., Jordan, D.C., Kukoly, L.M., Burton, S.R., Anele, E.U., Kwiecien, J.M., *et al.* 2023. Functional dissection and assembly of a small, newly evolved, W chromosome-specific genomic



- region of the African clawed frog *Xenopus laevis*. *PLoS Genet.* **19**: 1–27.
- Chassot, A.A., Ranc, F., Gregoire, E.P., Roepers-Gajadien, H.L., Taketo, M.M., Camerino, G., *et al.* 2008. Activation of  $\beta$ -catenin signaling by *Rspo1* controls differentiation of the mammalian ovary. *Hum. Mol. Genet.* **17**: 1264–1277.
- Chen, S., Zhang, G., Shao, C., Huang, Q., Liu, G., Zhang, P., *et al.* 2014. Whole-genome sequence of a flatfish provides insights into ZW sex chromosome evolution and adaptation to a benthic lifestyle. *Nat. Genet.* **46**: 253–260.
- Cheng, P.L., Dumas, G.A., Smith, J.T., Leger, A., Plamondon, A., McGrath, M.J., *et al.* 2006. Transcriptional diversity of DMRT1 (*dsx*- and *mab3*-related transcription factor 1) in human testis. *Cell Res.* **26**: 389–397.
- Cho, S., Huang, Z.Y. & Zhang, J. 2007. Sex-specific splicing of the honeybee doublesex gene reveals 300 million years of evolution at the bottom of the insect sex-determination pathway. *Genetics* **177**: 1733–1741.
- Cui, Z., Liu, Y., Wang, W., Wang, Q., Zhang, N., Lin, F., *et al.* 2017. Genome editing reveals *dmrt1* as an essential male sex-determining gene in Chinese tongue sole (*Cynoglossus semilaevis*). *Sci. Rep.* **7**: 42213.
- Dadhich, R.K., Real, F.M., Zurita, F., Barrionuevo, F.J., Burgos, M. & Jiménez, R. 2010. Role of apoptosis and cell proliferation in the testicular dynamics of seasonal breeding mammals: A study in the Iberian mole, *Talpa occidentalis*. *Biol. Reprod.* **83**: 83–91.
- Deloffre, L.A.M., Martins, R.S.T., Mylonas, C.C. & Canario, A.V.M. 2009. Alternative transcripts of DMRT1 in the European sea bass: Expression during gonadal differentiation. *Aquaculture* **293**: 89–99. Elsevier B.V.
- Diáz-Hernández, V., Dominguez-Mora, P., Chino-Palomo, L., Marmolejo-Valencia, A., Harfush, M. & Merchant-Larios, H. 2020. Spatiotemporal Expression of *Foxl2* and *Dmrt1* before, during, and after Sex Determination in the Sea Turtle *Lepidochelys olivacea*. *Sex. Dev.* **13**: 286–296.
- Domingos, J.A., Budd, A.M., Banh, Q.Q., Goldsbury, J.A., Zenger, K.R. & Jerry, D.R. 2018. Sex-specific *dmrt1* and *cyp19a1* methylation and alternative splicing in gonads of the protandrous hermaphrodite barramundi. *PLoS One* **13**: 1–19.
- Dong, J., Li, J., Hu, J., Sun, C., Tian, Y., Li, W., *et al.* 2020. Comparative Genomics Studies on the *dmrt* Gene Family in Fish. *Front. Genet.* **11**: 1–17.
- Duarte-Guterman, P., Ryan, M.J. & Trudeau, V.L. 2012. Developmental expression of sex steroid- and thyroid hormone-related genes and their regulation by triiodothyronine in the gonad-mesonephros of a Neotropical frog, *Physalaemus pustulosus*. *Gen. Comp. Endocrinol.* **177**: 195–204. Elsevier Inc.
- Dujardin, E., Andre, M., Dewaele, A., Mandon-Pepin, B., Poulat, F., Frambourg, A., *et al.* 2023. DMRT1 is a Testis Determining Gene in Rabbits and is Also Essential for Female Fertility. *eLift* **12**: RP89284.
- Ellegren, H. 2001. Hens, cocks and avian sex determination A quest for genes on Z or W? *EMBO Rep.* **21**: 192–196.
- Fahrioglu, U., Murphy, M.W., Zarkower, D. & Bardwell, V.J. 2006. mRNA expression analysis and the molecular basis of neonatal testis defects in *Dmrt1* mutant mice. *Sex. Dev.* **1**: 42–58.
- Ferguson-Smith, M. 2006. The evolution of sex chromosomes and sex determination in vertebrates and the

- key role of DMRT1. *Sex. Dev.* **1**: 2–11.
- Fujitani, K., Otomo, A., Wada, M., Takamatsu, N. & Ito, M. 2016. Sexually dimorphic expression of *Dmrt1* and *c H2AX* in germ stem cells during gonadal development in *Xenopus laevis*. *FEBS Open Bio* **6**: 276–284.
- Ge, C., Ye, J., Weber, C., Sun, W., Zhang, H., Zhou, Y., *et al.* 2018. The histone demethylase KDM6B regulates temperature-dependent sex determination in a turtle species. *Science*. **360**: 645–648.
- Ge, C., Ye, J., Zhang, H., Zhang, Y., Sun, W., Sang, Y., *et al.* 2017. *Dmrt1* induces the male pathway in a turtle species with temperature-dependent sex determination. *Dev.* **144**: 2222–2233.
- Gempe, T. & Beye, M. 2011. Function and evolution of sex determination mechanisms, genes and pathways in insects. *BioEssays* **33**: 52–60.
- Georges, A. & Holleley, C.E. 2018. How does temperature determine sex? *Science*. **360**: 601–602.
- Geuverink, E. & Beukeboom, L.W. 2014. Phylogenetic distribution and evolutionary dynamics of the sex determination genes *doublesex* and *transformer* in insects. *Sex. Dev.* **8**: 38–49.
- Giebel, J.D., Kamping, A., Beukeboom, L.W. & Zande, L. Van De. 2009. Identification and characterization of the *doublesex* gene of *Nasonia*. *Insect Mol. Biol.* **18**: 315–324.
- Gu, L. & Walters, J.R. 2017. Evolution of sex chromosome dosage compensation in animals: A beautiful theory, undermined by facts and bedeviled by details. *Genome Biol. Evol.* **9**: 2461–2476.
- Gunes, S.O., Metin Mahmutoglu, A. & Agarwal, A. 2016. Genetic and epigenetic effects in sex determination. *Birth Defects Res. Part C - Embryo Today Rev.* **108**: 321–336.
- Guo, Y., Cheng, H., Huang, X., Gao, S., Yu, H. & Zhou, R. 2005. Gene structure, multiple alternative splicing, and expression in gonads of zebrafish *Dmrt1*. *Biochem. Biophys. Res. Commun.* **330**: 950–957.
- Guo, Y., Li, Q., Gao, S., Zhou, X., He, Y., Shang, X., *et al.* 2004. Molecular cloning, characterization, and expression in brain and gonad of *Dmrt5* of zebrafish. *Biochem. Biophys. Res. Commun.* **324**: 569–575.
- Hattori, R.S., Gould, R.J., Fujioka, T., Saito, T., Kurita, J., Strüssmann, C.A., *et al.* 2007. Temperature-dependent sex determination in Hd-rR medaka *Oryzias latipes*: Gender sensitivity, thermal threshold, critical period, and DMRT1 expression profile. *Sex. Dev.* **1**: 138–146.
- He, C.L., Du, J.L., Wu, G.C., Lee, Y.H., Sun, L.T. & Chang, C.F. 2003. Differential *Dmrt1* transcripts in gonads of the protandrous black porgy, *Acanthopagrus schlegelii*. *Cytogenet. Genome Res.* **101**: 309–313.
- Hediger Niessen, M., Burghardt, G., Siegenthaler, C., Buser, N., Hilfiker-Kleiner, D., Dübendorfer, A., *et al.* 2004. Sex determination in *Drosophila melanogaster* and *Musca domestica* converges at the level of the terminal regulator *doublesex*. *Dev. Genes Evol.* **214**: 29–42.
- Herpin, A. & Scharf, M. 2011. *Dmrt1* genes at the crossroads: A widespread and central class of sexual development factors in fish. *FEBS J.* **278**: 1010–1019.
- Herpin, A., Schindler, D., Kraiss, A., Hornung, U., Winkler, C. & Scharf, M. 2007. Inhibition of primordial germ cell proliferation by the medaka male determining gene *Dmrt1bY*. *BMC Dev. Biol.* **7**: 1–15.
- Hirst, C.E., Major, A.T., Ayers, K.L., Brown, R.J., Mariette, M., Sackton, T.B., *et al.* 2017. Sex reversal and comparative data undermine the W chromosome and support Z-linked DMRT1 as the regulator of

- gonadal sex differentiation in birds. *Endocrinology* **158**: 2970–2987.
- Holleley, C.E., O’Meally, D., Sarre, S.D., Marshall Graves, J.A., Ezaz, T., Matsubara, K., *et al.* 2015. Sex reversal triggers the rapid transition from genetic to temperature-dependent sex. *Nature* **523**: 79–82.
- Hong, C.S., Park, B.Y. & Saint-Jeannet, J.P. 2007. The function of Dmrt genes in vertebrate development: It is not just about sex. *Dev. Biol.* **310**: 1–9.
- Huang, S., Ye, L. & Chen, H. 2017. Sex determination and maintenance: The role of DMRT1 and FOXL2. *Asian J. Androl.* **19**: 619–624.
- Huang, X., Guo, Y., Shui, Y., Gao, S., Yu, H., Cheng, H., *et al.* 2005a. Multiple alternative splicing and differential expression of dmrt1 during gonad transformation of the rice field eel. *Biol. Reprod.* **73**: 1017–1024.
- Huang, X., Guo, Y., Shui, Y., Gao, S., Yu, H., Cheng, H., *et al.* 2005b. Multiple Alternative Splicing and Differential Expression of dmrt1 During Gonad Transformation of the Rice Field Eel. *Biol. Reprod.* **73**: 1017–1024.
- Ioannidis, J., Taylor, G., Zhao, D., Liu, L., Idoko-Akoh, A., Gong, D., *et al.* 2021. Primary sex determination in birds depends on DMRT1 dosage, but gonadal sex does not determine adult secondary sex characteristics. *Proc. Natl. Acad. Sci. U. S. A.* **118**: 1–10.
- Jia, Y., Zheng, J., Chi, M., Liu, S., Jiang, W., Cheng, S., *et al.* 2019. Molecular identification of dmrt1 and its promoter CpG methylation in correlation with gene expression during gonad development in *Culter alburnus*. *Fish Physiol. Biochem.* **45**: 245–252.
- Jiang, D.N., Huang, Y.Q., Zhang, J.M., Mustapha, U.F., Peng, Y.X., Huang, H., *et al.* 2022. Establishment of the Y-linked Dmrt1Y as the candidate sex determination gene in spotbanded scat (*Selenotoca multifasciata*). *Aquac. Reports* **23**: 101085.
- Kettlewell, J.R., Raymond, C.S. & Zarkower, D. 2000. Temperature-dependent expression of turtle Dmrt1 prior to sexual differentiation. *Genesis* **26**: 174–178.
- Kim, S., Bardwell, V.J. & Zarkower, D. 2007. Cell type-autonomous and non-autonomous requirements for Dmrt1 in postnatal testis differentiation. *Dev. Biol.* **307**: 314–327.
- Kim, Y., Kobayashi, A., Sekido, R., DiNapoli, L., Brennan, J., Chaboissier, M.C., *et al.* 2006. Fgf9 and Wnt4 Act as Antagonistic Signals to Regulate Mammalian Sex Determination. *PLoS Biol.* **4**: e187.
- Kobayashi, T., Matsuda, M., Kajiura-Kobayashi, H., Susuki, A., Saito, N., Nakamoto, M., *et al.* 2004. Two DM domain genes, DMY and DMRT1, involved in testicular differentiation and development in the Medaka, *Oryzias latipes*. *Dev. Dyn.* **231**: 518–526.
- Kondo, M., Nanda, I., Hornung, U., Schmid, M. & Schartl, M. 2004. Evolutionary Origin of the Medaka Y Chromosome Mariko. *Curr. Biol.* **14**: 1664–1669.
- Kossack, M.E. & Draper, B.W. 2019. Genetic regulation of sex determination and maintenance in zebrafish (*Danio rerio*). *Curr Top Dev Biol* **134**: 119–149.
- Kratz, C.P., Han, S.S., Rosenberg, P.S., Berndt, S.I., Burdett, L., Yeager, M., *et al.* 2011. Variants in or near KITLG, BAK1, DMRT1, and TERT-CLPTM1L predispose to familial testicular germ cell tumour. *J. Med. Genet.* **48**: 473–476.
- Krentz, A.D., Murphy, M.W., Sarver, A.L., Griswold, M.D., Bardwell, V.J. & Zarkower, D. 2011. DMRT1

- promotes oogenesis by transcriptional activation of *Stra8* in the mammalian fetal ovary. *Dev. Biol.* **356**: 63–70.
- Lambeth, L.S., Raymond, C.S., Roeszler, K.N., Kuroiwa, A., Nakata, T., Zarkower, D., *et al.* 2014. Over-expression of *DMRT1* induces the male pathway in embryonic chicken gonads. *Dev. Biol.* **389**: 160–172.
- Ledig, S., Hiort, O., Scherer, G., Hoffmann, M., Wolff, G., Morlot, S., *et al.* 2010. Array-CGH analysis in patients with syndromic and non-syndromic XY gonadal dysgenesis: Evaluation of array CGH as diagnostic tool and search for new candidate loci. *Hum. Reprod.* **25**: 2637–2646.
- Lee, H.J., Seo, M., Choi, H.J., Rengaraj, D., Jung, K.M., Park, J.S., *et al.* 2021. *DMRT1* gene disruption alone induces incomplete gonad feminization in chicken. *FASEB J.* **35**: e21876.
- Li, M.H., Yang, H.H., Li, M.R., Sun, Y.L., Jiang, X.L., Xie, Q.P., *et al.* 2013. Antagonistic roles of *Dmrt1* and *Foxl2* in sex differentiation via estrogen production in tilapia as demonstrated by TALENs. *Endocrinology* **154**: 4814–4825.
- Li, X.Y., Mei, J., Ge, C.T., Liu, X.L. & Gui, J.F. 2022. Sex determination mechanisms and sex control approaches in aquaculture animals. *Sci. China Life Sci.* **65**: 1091–1122.
- Li, Z., Dai, H., Martos, S.N., Xu, B., Gao, Y., Li, T., *et al.* 2015. Distinct roles of DNMT1-dependent and DNMT1-independent methylation patterns in the genome of mouse embryonic stem cells. *Genome Biol.* **16**: 1–15. *Genome Biology*.
- Lin, Q., Mei, J., Li, Z., Zhang, X., Zhou, L. & Gui, J. 2017. Self-Renewal and Differentiation of Male Germ Cells. *Genetics* **207**: 1007–1022.
- Looijenga, L.H.J., Hersmus, R., Gillis, A.J.M., Pfundt, R., Stoop, H.J., Van Gurp, R.J.H.L.M., *et al.* 2006. Genomic and expression profiling of human spermatocytic seminomas: Primary spermatocyte as tumorigenic precursor and *DMRT1* as candidate chromosome 9 gene. *Cancer Res.* **66**: 290–302.
- Lu, H., Huang, X., Zhang, L., Guo, Y., Cheng, H. & Zhou, R. 2007. Multiple alternative splicing of mouse *Dmrt1* during gonadal differentiation. *Biochem. Biophys. Res. Commun.* **352**: 630–634.
- Ma, W.-J., Rodrigues, N., Sermier, R., Brelsford, A. & Perrin, N. 2016. *Dmrt1* polymorphism covaries with sex-determination patterns in *Rana temporaria*. *Ecol. Evol.* 5107–5117.
- Ma, W.-J., Veltsos, P., Sermier, R., Parker, D.J. & Perrin, N. 2018a. Evolutionary and developmental dynamics of sex-biased gene expression in common frogs with proto-Y chromosomes. *Genome Biol.* **19**: 156. *Genome Biology*.
- Ma, W.-J., Veltsos, P., Toups, M.A., Rodrigues, N., Sermier, R., Jeffries, D.L., *et al.* 2018b. Tissue specificity and dynamics of sex-biased gene expression in a common frog population with differentiated, yet homomorphic, sex chromosomes. *Genes.* **9**: 294.
- Ma, W.J., Veltsos, P., Sermier, R., Parker, D.J. & Perrin, N. 2018c. Evolutionary and developmental dynamics of sex-biased gene expression in common frogs with proto-Y chromosomes. *Genome Biol.* **19**: 1–17. *Genome Biology*.
- Marchand, O., Govoroun, M., D’Cotta, H., McMeel, O., Lareyre, J.J., Bernot, A., *et al.* 2000. *DMRT1* expression during gonadal differentiation and spermatogenesis in the rainbow trout, *Oncorhynchus mykiss*. *Biochim. Biophys. Acta - Gene Struct. Expr.* **1493**: 180–187.
- Massoud, D., Lao-Pérez, M., Ortega, E., Burgos, M., Jiménez, R. & Barrionuevo, F.J. 2021. Divergent

- seasonal reproductive patterns in syntopic populations of two murine species in southern Spain, *Mus spretus* and *Apodemus sylvaticus*. *Animals* **11**: 1–16.
- Masuyama, H., Yamada, M., Kamei, Y., Fujiwara-Ishikawa, T., Todo, T., Nagahama, Y., *et al.* 2012. Dmrt1 mutation causes a male-to-female sex reversal after the sex determination by Dmy in the medaka. *Chromosom. Res.* **20**: 163–176.
- Matson, C.K., Murphy, M.W., Sarver, A.L., Griswold, M.D., Bardwell, V.J. & Zarkower, D. 2011. DMRT1 prevents female reprogramming in the postnatal mammalian testis. *Nature* **476**: 101–105.
- Matsuda, M., Nagahama, Y., Shinomiya, A., Sato, T., Matsuda, C., Kobayashi, T., *et al.* 2002. DMY is a Y-specific DM-domain gene required for male development in the medaka fish. *Nature* **417**: 559–563.
- Mawaribuchi, S., Musashijima, M., Wada, M., Izutsu, Y., Kurakata, E., Park, M.K., *et al.* 2017a. Molecular evolution of two distinct dmrt1 promoters for germ and somatic cells in vertebrate gonads. *Mol. Biol. Evol.* **34**: 724–733.
- Mawaribuchi, S., Takahashi, S., Wada, M., Uno, Y., Matsuda, Y., Kondo, M., *et al.* 2017b. Sex chromosome differentiation and the W- and Z-specific loci in *Xenopus laevis*. *Dev. Biol.* **426**: 393–400. Elsevier Inc.
- Mawaribuchi, S., Yoshimoto, S., Ohashi, S., Takamatsu, N. & Ito, M. 2012. Molecular evolution of vertebrate sex-determining genes. *Chromosom. Res.* **20**: 139–151.
- Miura, I., Takase, M. & Nakamura, M. 1998. Sex determination and differentiation in amphibians. *Tanpakushitsu Kakusan Koso.* **43**: 462–469.
- Miyake, Y., Sakai, Y. & Kuniyoshi, H. 2012. Molecular cloning and expression profile of sex-specific genes, figla and dmrt1, in the protogynous hermaphroditic fish, *Halichoeres poecilopterus*. *Zoolog. Sci.* **29**: 690–701.
- Mizoguchi, B. & Valenzuela, N. 2020. Alternative splicing and thermosensitive expression of Dmrt1 during urogenital development in the painted turtle, *Chrysemys picta*. *PeerJ* **2020**: 1–25.
- Mugal, C.F., Arndt, P.F., Holm, L. & Ellegren, H. 2015. Evolutionary Consequences of DNA Methylation on the GC Content in Vertebrate Genomes. *Genes|Genomes|Genetics* **5**: 441–447.
- Murakami, T., Maki, N., Nishida-Umehara, C., Matsuda, Y. & Agata, K. 2007. Establishment of high-resolution FISH mapping system and its application for molecular cytogenetic characterization of chromosomes in newt, *Cynops pyrrhogaster* (Urodela, Amphibia). *Chromosom. Res.* **15**: 471–484.
- Murphy, M.W., Sarver, A.L., Rice, D., Hatzi, K., Ye, K., Melnick, A., *et al.* 2010. Genome-wide analysis of DNA binding and transcriptional regulation by the mammalian Doublesex homolog DMRT1 in the juvenile testis. *Proc. Natl. Acad. Sci. U. S. A.* **107**: 13360–13365.
- Mustapha, U.F., Jiang, D.N., Liang, Z.H., Gu, H.T., Yang, W., Chen, H.P., *et al.* 2018. Male-specific Dmrt1 is a candidate sex determination gene in spotted scat (*Scatophagus argus*). *Aquaculture* **495**: 351–358. Elsevier.
- Nanda, I., Hornung, U., Kondo, M., Schmid, M. & Schartl, M. 2003. Common spontaneous sex-reversed XX males of the medaka *Oryzias latipes*. *Genetics* **163**: 245–251.
- Nanda, I., Kondo, M., Hornung, U., Asakawa, S., Winkler, C., Shimizu, A., *et al.* 2002. A duplicated copy of DMRT1 in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*. *Proc. Natl. Acad. Sci.* **99**: 11778–11783.

- Nanda, I., Zend-Ajusich, E., Shan, Z., Grützner, F., Scharl, M., Burt, D.W., *et al.* 2000. Conserved synteny between the chicken Z sex chromosome and human chromosome 9 includes the male regulatory gene DMRT1: A comparative (re)view on avian sex determination. *Cytogenet. Cell Genet.* **89**: 67–78.
- Nef, S., Schaad, O., Stallings, N.R., Cederroth, C.R., Pitetti, J.L., Schaer, G., *et al.* 2005. Gene expression during sex determination reveals a robust female genetic program at the onset of ovarian development. *Dev. Biol.* **287**: 361–377.
- Ohmuro-Matsuyama, Y., Matsuda, M., Kobayashi, T., Ikeuchi, T. & Nagahama, Y. 2003. Expression of DMY and DMRT1 in the various tissues of the medaka, *Oryzias latipes*. *Fish Physiol. Biochem.* **28**: 171.
- Okuno, M., Miyamoto, S., Itoh, T., Seki, M., Suzuki, Y., Mizushima, S., *et al.* 2020. Expression profiling of sexually dimorphic genes in the Japanese quail, *Coturnix japonica*. *Sci. Rep.* **10**: 1–11.
- Ottolenghi, C., Fellous, M., Barbieri, M. & McElreavey, K. 2002. Novel paralogy relations among human chromosomes support a link between the phylogeny of doublesex-related genes and the evolution of sex determination. *Genomics* **79**: 333–343.
- Ou, M., Chen, K., Gao, D., Wu, Y., Luo, Q., Liu, H., *et al.* 2022. Characterization, expression and CpG methylation analysis of *Dmrt1* and its response to steroid hormone in blotched snakehead (*Channa maculata*). *Comp. Biochem. Physiol. Part - B Biochem. Mol. Biol.* **257**: 1–10.
- Peng, Y.X., Huang, Y.Q., Zhong, J., Jiang, Z.T., Fan, S., Shi, H.J., *et al.* 2023. Corrigendum to Identification of sex-linked marker and candidate sex determination gene in ornamental fish, African scat (*Scatophagus tetracanthus*). *Aquaculture* **569**: 739386. Elsevier B.V.
- Piferrer, F. 2021. Epigenetic mechanisms in sex determination and in the evolutionary transitions between sexual systems. *Philos. Trans. R. Soc. B Biol. Sci.* **376**.
- Piferrer, F. 2013. Epigenetics of sex determination and gonadogenesis. *Dev. Dyn.* **242**: 360–370.
- Piprek, R.P., Pecio, A., Laskowska-Kaszub, K., Kubiak, J.Z. & Szymura, J.M. 2013. Sexual dimorphism of AMH, DMRT1 and RSPO1 localization in the developing gonads of six anuran species. *Int. J. Dev. Biol.* **57**: 871–875.
- Pomiankowski, A., Nöthiger, R. & Wilkins, A. 2004. The Evolution of the Drosophila Sex-Determination Pathway. *Genetics* **166**: 1761–1773.
- Qi, S., Dai, S., Zhou, X., Wei, X., Chen, P., He, Y., *et al.* 2024. *Dmrt1* is the only male pathway gene tested indispensable for sex determination and functional testis development in tilapia. *PLoS Genet.* **20**: 1–32.
- Raghuvver, K. & Senthilkumaran, B. 2009. Identification of multiple *dmrt1*s in catfish: Localization, dimorphic expression pattern, changes during testicular cycle and after methyltestosterone treatment. *J. Mol. Endocrinol.* **42**: 437–448.
- Raymond, C.S., Murphy, M.W., O’Sullivan, M.G., Bardwell, V.J. & Zarkower, D. 2000. *Dmrt1*, a gene related to worm and fly sexual regulators, is required for mammalian testis differentiation. *Genes Dev.* **14**: 2587–2595.
- Raymond, C.S., Parker, E.D., Kettlewell, J.R., Brown, L.G., Page, D.C., Kusz, K., *et al.* 1999. A region of human chromosome 9p required for testis development contains two genes related to known sexual regulators. *Hum. Mol. Genet.* **8**: 989–996.
- Raymond, C.S., Shamu, C.E., Shen, M.M., Seifert, K.J., Hirsch, B., Hodgkin, J., *et al.* 1998. Evidence for

- evolutionary conservation of sex-determining genes. *Nature* **391**: 691–695.
- Rhen, T., Metzger, K., Schroeder, A. & Woodward, R. 2007. Expression of putative sex-determining genes during the thermosensitive period of gonad development in the snapping turtle, *Chelydra serpentina*. *Sex. Dev.* **1**: 255–270.
- Rhen, T. & Schroeder, A. 2010. Molecular mechanisms of sex determination in reptiles. *Sex. Dev.* **4**: 16–28.
- Rodrigues, N., Studer, T., Dufresnes, C., Ma, W., Veltsos, P. & Perrin, N. 2017. Dmrt1 polymorphism and sex-chromosome differentiation in *Rana temporaria*. *Mol. Ecol.* **26**: 4897–4905.
- Russo, V.E.A., Martienssen, R.A. & Riggs, A.D. (eds). 1996. *Epigenetic mechanisms of gene regulation*. V.E.A. Russo, R.A. Martienssen, A.D. Riggs. 1996. Cold Spring Harbor Laboratory Press, New York, USA. Cold Spring Harbor Laboratory Press, New York.
- Sakata, N., Miyazaki, K. & Wakahara, M. 2006. Up-regulation of P450arom and down-regulation of Dmrt-1 genes in the temperature-dependent sex reversal from genetic males to phenotypic females in a salamander. *Dev. Genes Evol.* **216**: 224–228.
- Sánchez, L. & Chaouiya, C. 2018. Logical modelling uncovers developmental constraints for primary sex determination of chicken gonads. *J. R. Soc. Interface* **15**: 20180165.
- Sarre, S.D., Ezaz, T. & Georges, A. 2011. Transitions Between Sex-Determining Systems in Reptiles and Amphibians. *Annu. Rev. Genomics Hum. Genet.* **12**: 391–406.
- Sheng, Y., Chen, B., Zhang, L., Luo, M., Cheng, H. & Zhou, R. 2014. Identification of Dmrt genes and their up-regulation during gonad transformation in the swamp eel (*Monopterus albus*). *Mol. Biol. Rep.* **41**: 1237–1245.
- Shibata, K., Takase, M. & Nakamura, M. 2002. The Dmrt1 expression in sex-reversed gonads of amphibians. *Gen. Comp. Endocrinol.* **127**: 232–241.
- Shinomiya, A., Shibata, N., Sakaizumi, M. & Hamaguchi, S. 2002. Sex reversal of genetic females (XX) induced by the transplantation of XY somatic cells in the medaka, *Oryzias latipes*. *Int. J. Dev. Biol.* **46**: 711–717.
- Smith, C.A., McClive, P.J., Reed, K.J., Western, P.S. & Sinclair, A.H. 1999. Conservation of a sex-determining gene. *Nature* **402**: 601.
- Smith, C.A., Roeszler, K.N., Hudson, Q.J. & Sinclair, A.H. 2007. Avian sex determination: What, when and where? *Cytogenet. Genome Res.* **117**: 165–173.
- Smith, C.A., Roeszler, K.N., Ohnesorg, T., Cummins, D.M., Farlie, P.G., Doran, T.J., *et al.* 2009. The avian Z-linked gene DMRT1 is required for male sex determination in the chicken. *Nature* **461**: 267–271.
- Straková, B., Rovatsos, M., Kubička, L. & Kratochvíl, L. 2020. Evolution of Sex Determination in Amniotes: Did Stress and Sequential Hermaphroditism Produce Environmental Determination? *BioEssays* **42**: 1–9.
- Su, L., Zhou, F., Ding, Z., Gao, Z., Wen, J., Wei, W., *et al.* 2015. Transcriptional variants of Dmrt1 and expression of four Dmrt genes in the blunt snout bream, *Megalobrama amblycephala*. *Gene* **573**: 205–215.
- Suzuki, M.G., Ohbayashi, F., Mita, K. & Shimada, T. 2001. The mechanism of sex-specific splicing at the doublesex gene is different between *Drosophila melanogaster* and *Bombyx mori*. *Insect Biochem. Mol.*

*Biol.* **31**: 1201–1211.

- Tachibana, M. 2016. Epigenetics of sex determination in mammals. *Reprod. Med. Biol.* **15**: 59–67.
- Tang, H., Brennan, J., Karl, J., Hamada, Y., Raetzman, L. & Capel, B. 2008. Notch signaling maintains Leydig progenitor cells in the mouse testis. *Development* **135**: 3745–3753.
- Teranishi, M., Shimada, Y., Hori, T., Nakabayashi, O., Kikuchi, T., Macleod, T., *et al.* 2001. Transcripts of the MHM region on the chicken Z chromosome accumulate as non-coding RNA in the nucleus of female cells adjacent to the *hw??I* locus. *Chromosom. Res.* **9**: 147–165.
- Turnbull, C., Rapley, E.A., Seal, S., Pernet, D., Renwick, A., Hughes, D., *et al.* 2010. Variants near DMRT1, TERT and ATF7IP are associated with testicular germ cell cancer. *Nat. Genet.* **42**: 604–607.
- Urushitani, H., Katsu, Y., Miyagawa, S., Kohno, S., Ohta, Y., Guillette, L.J., *et al.* 2011. Molecular cloning of anti-Müllerian hormone from the American alligator, *Alligator mississippiensis*. *Mol. Cell. Endocrinol.* **333**: 190–199.
- Valenzuela, N. & Lance, V.A. 2004. *Temperature-Dependent Sex Determination in Vertebrates* (N. Valenzuela & V. A. Lance, eds). Smithsonian Books.
- Van De Zande, L. & Verhulst, E.C. 2014. Genomic imprinting and maternal effect genes in haplodiploid sex determination. *Sex. Dev.* **8**: 74–82.
- Veith, A., Schäfer, M., Klüver, N., Schmidt, C., Schultheis, C., Schartl, M., *et al.* 2006a. Tissue-specific expression of *dmrt* genes in embryos and adults of the platyfish *Xiphophorus maculatus*. *Zebrafish* **3**: 325–337.
- Veith, A.M., Klattig, J., Dettai, A., Schmidt, C., Englert, C. & Volff, J.N. 2006b. Male-biased expression of X-chromosomal DM domain-less *Dmrt8* genes in the mouse. *Genomics* **88**: 185–195.
- Verhulst, E.C., van de Zande, L. & Beukeboom, L.W. 2010. Insect sex determination: It all evolves around transformer. *Curr. Opin. Genet. Dev.* **20**: 376–383.
- Vicoso, B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome turnover. *Nat. Ecol. Evol.* **3**: 1632–1641.
- Wang, F.L., Yan, L.X., Shi, H.J., Liu, X.Y., Zheng, Q.Y., Sun, L.N., *et al.* 2018. Genome-wide identification, evolution of DNA methyltransferases and their expression during gonadal development in Nile tilapia. *Comp. Biochem. Physiol. Part - B Biochem. Mol. Biol.* **226**: 73–84.
- Wang, H., Fang, Y., Wang, L., Zhu, W., Ji, H., Wang, H., *et al.* 2015. Heterosis and differential gene expression in hybrids and parents in *Bombyx mori* by digital gene expression profiling. *Sci. Rep.* **5**: 8750.
- Wang, H., Qu, M., Tang, W., Liu, S. & Ding, S. 2022a. Transcriptome Profiling and Expression Localization of Key Sex-Related Genes in a Socially-Controlled Hermaphroditic Clownfish, *Amphiprion clarkii*. *Int. J. Mol. Sci.* **23**: 9085.
- Wang, L., Sun, F., Wan, Z.Y., Yang, Z., Tay, Y.X., Lee, M., *et al.* 2022b. Transposon-induced epigenetic silencing in the X chromosome as a novel form of *dmrt1* expression regulation during sex determination in the fighting fish. *BMC Biol.* **20**: 1–16. BMC Biology.
- Wang, Q., Cao, T., Wang, Y., Li, X. & Wang, Y. 2023. Genome-wide identification and comparative analysis of *Dmrt* genes in echinoderms. *Sci. Rep.* **13**: 1–9.



- Wang, Q., Cui, Z., Guo, H., Zhang, N., Xu, W., Yang, Y., *et al.* 2019. Functional analysis of the promoter of the *dmrt1* gene in Chinese tongue sole, *Cynoglossus semilaevis*. *J. Oceanol. Limnol.* **37**: 1333–1341.
- WANG, Y., LIU, J., HUANG, B., XU, Y.-M., LI, J., HUANG, L.-F., *et al.* 2015. Mechanism of alternative splicing and its regulation. *Biomed. Reports* **3**: 152–158.
- Weber, C., Zhou, Y., Lee, J.G., Looger, L.L., Qian, G., Ge, C., *et al.* 2020. Temperature-dependent sex determination is mediated by pSTAT3 repression of *Kdm6b*. *Science*. **368**: 303–306.
- Webster, K.A., Schach, U., Ordaz, A., Steinfeld, J.S., Draper, B.W. & Siegfried, K.R. 2017. *Dmrt1* is necessary for male sexual development in zebrafish. *Dev. Biol.* **422**: 33–46.
- Wen, A.Y., You, F., Sun, P., Li, J., Xu, D.D., Wu, Z.H., *et al.* 2014. CpG methylation of *dmrt1* and *cyp19a* promoters in relation to their sexual dimorphic expression in the Japanese flounder *Paralichthys olivaceus*. *J. Fish Biol.* **84**: 193–205.
- Wexler, J.R., Plachetzki, D.C. & Kopp, A. 2014. Pan-metazoan phylogeny of the DMRT gene family: A framework for functional studies. *Dev. Genes Evol.* **224**: 175–181.
- Wright, C.J., Smith, C.W.J. & Jiggins, C.D. 2022. Alternative splicing as a source of phenotypic diversity. *Nat. Rev. Genet.* **23**: 697–710.
- Xu, W., Li, Y.Y., Lou, Q.Q., Chen, X.R., Qin, Z.F. & Wie, W.J. 2015. Low concentrations of dihydrotestosterone induce female-to-male sex reversal in the frog *Pelophylax nigromaculatus*. *Environ. Toxicol. Chem.* **34**: 2370–2377.
- Yang, X., Zheng, J., Xu, G., Qu, L., Chen, S., Li, J., *et al.* 2010. Exogenous cMHM regulates the expression of DMRT1 and ER $\alpha$  in avian testes. *Mol. Biol. Rep.* **37**: 1841–1847.
- Yoshimoto, S., Ikeda, N., Izutsu, Y., Shiba, T., Takamatsu, N. & Ito, M. 2010. Opposite roles of DMRT1 and its W-linked paralogue, DM-W, in sexual dimorphism of *Xenopus laevis*: Implications of a ZZ/ZW-type sex-determining system. *Development* **137**: 2519–2526.
- Yoshimoto, S. & Ito, M. 2011. A ZZ/ZW-type sex determination in *xenopus laevis*. *FEBS J.* **278**: 1020–1026.
- Yoshimoto, S., Okada, E., Umemoto, H., Tamura, K., Uno, Y., Nishida-umehara, C., *et al.* 2008. A W-linked DM-domain gene, DM-W, participates in primary ovary development in *Xenopus laevis*. *Proc. Natl. Acad. Sci.* **105**: 2469–2474.
- Zarkower, D. 2013. *DMRT Genes in Vertebrate Gametogenesis*, 1st ed. Elsevier Inc.
- Zarkower, D. & Murphy, M.W. 2022. DMRT1: An Ancient Sexual Regulator Required for Human Gonadogenesis. *Sex. Dev.* **16**: 112–125.
- Zhang, J. 2004. Evolution of DMY, a Newly Emergent Male Sex-Determination Gene of Medaka Fish. *Genetics* **166**: 1887–1895.
- Zhang, X., Wang, H., Li, M., Cheng, Y., Jiang, D., Sun, L., *et al.* 2014. Isolation of Doublesex- and Mab-3-related transcription factor 6 and its involvement in spermatogenesis in tilapia. *Biol. Reprod.* **91**: 1–10.
- Zhao, Y., Lu, H., Yu, H., Cheng, H. & Zhou, R. 2007a. Multiple alternative splicing in gonads of chicken DMRT1. *Dev. Genes Evol.* **217**: 119–126.
- Zhao, Y., Lu, H., Yu, H., Cheng, H. & Zhou, R. 2007b. Multiple alternative splicing in gonads of chicken DMRT1. *Dev. Genes Evol.* **217**: 119–126.



**Table S1.** List of species in which *Dmrt1* expression was studied throughout gonadal development and at adult stages, *Dmrt1*'s role in sex determination and its gene function across various vertebrate groups.

Class	Family	Species	Sex determination	Master SD gene	Expression during gonadal sex differentiation	Expression in gonads of sub-adults/adults	Expression in adult somatic tissues	Methods	Gene function	Reference
Actinistia	Latimeriidae	<i>Latimeria chalumnae</i>	NA	NA	NA	testes, (no ovaries data)	not expressed: liver	RNAseq	NA	Forconi et al. 2013
Actinopterygii	Acipenseridae	<i>Acipenser baeri</i>	NA	NA	NA	testes (strong), ovaries (weak)	very low expression: muscle; not expressed: brain, liver, gills, kidney	qPCR	NA	Berbellio et al. 2013
Actinopterygii	Acipenseridae	<i>Acipenser fulvescens</i>	NA	NA	NA	testes (strong), ovaries (weak)	not examined	RNAseq	NA	Hale et al. 2010
Actinopterygii	Acipenseridae	<i>Acipenser gulfstenstedtii</i>	NA	NA	testes, ovotestes (strongest), ovaries (weak)	NA	not examined	qPCR	NA	Fajkowski et al. 2016
Actinopterygii	Acipenseridae	<i>Acipenser ruthenus</i>	NA	NA	testes (strong), ovaries (weak)	NA	not examined	qPCR	NA	Wang et al. 2019, Wang et al. 2017
Actinopterygii	Acipenseridae	<i>Acipenser schrenkii</i>	NA	NA	testes (weak), ovaries (weak)	testes (strong), ovaries (weak)	not expressed: brain, pituitary, gill, heart, head kidney, posterior kidney, liver, spleen, caecum	qPCR, PCR	NA	Okada et al. 2017
Actinopterygii	Acipenseridae	<i>Acipenser sinensis</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR, histc	NA	Leng et al. 2015, Hu et al. 2024
Actinopterygii	Acipenseridae	<i>Acipenser stellatus</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR	NA	Burcea et al. 2018
Actinopterygii	Acipenseridae	<i>Scaphiynchus platyrhynchus</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR	NA	Amberg et al. 2013
Actinopterygii	Actinopterygii	<i>Tetraodon nigroviridis</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not expressed: liver, eye, brain, skin, heart, gills, head, kidney	PCR	NA	Brunner et al. 2001
Actinopterygii	Adrianchthyidae	<i>Oryzias latipes</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR, RNA	NA	Dong et al. 2021
Actinopterygii	Adrianchthyidae	<i>Oryzias latipes</i>	NA	( <i>Dmrt1b1/Dmrt1</i> )	testes (Dmrt1a, Dmrt1b1/Dmrt1)	testes (Dmrt1b1/Dmrt1; Dmrt1a), ovaries Dmrt1a	Dmrt1b1/Dmrt1 expressed: pituitary, heart, liver, spleen; Dmrt1b1/Dmrt1 very weak expression (intestine, eye); Dmrt1a expressed: testis	PCR, histol	SD, gonad differentiation (Dmrt1b1/Dmrt1), spermatogenesis (Dmrt1a)	Ohmura-Matsuyama et al. 2003, Kobayashi et al. 2004
Actinopterygii	Anguillidae	<i>Anguilla japonica</i>	NA	NA	testes, ovaries	testes (strong), ovaries (weak)	not examined	qPCR, histc	NA	Jeng et al. 2019
Actinopterygii	Anoplopomatidae	<i>Anoplopoma fimbria</i>	NA	NA	testes (strong), ovaries (weak)	testes	not expressed: brain, pituitary, gill, heart, spleen, liver, intestine	PCR, semi-	NA	Smith et al. 2013
Actinopterygii	Atherinopsidae	<i>Odontesthes bonariensis</i>	NA	NA	testes	testes	not expressed: eye, gill, liver, kidney, spleen, epithelium olfactory, heart, muscle, intestine, brain-pituitary	PCR, South	NA	Fernandino et al. 2006
Actinopterygii	Bagridae	<i>Pseudobagrus ussuriensis</i>	NA	NA	testes	testes	not examined	qPCR	NA	Pan et al. 2021
Actinopterygii	Batrachoidae	<i>Halobatrachus didactylus</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	very low expression: heart, liver, spleen, kidney, swim bladder, intestine, muscle, gill	qPCR, histc	NA	Ubeda-Manzanaro et al. 2014, Li et al. 2014
Actinopterygii	Butidae	<i>Oxyeleotris marmorata</i>	NA	NA	testes (strong), ovaries (weak)	testes	very low expression: heart, liver, spleen, kidney, muscle, pituitary; not expressed: brain	qPCR	NA	Liu et al. 2020
Actinopterygii	Centrarchidae	<i>Micropterus salmoides</i>	NA	NA	testes	testes (strong), ovaries (weak)	very low expression: Heart, Liver, Brain, Intestine, Gill, Kidney	qPCR	NA	Yan et al. 2019
Actinopterygii	Channidae	<i>Channa maculata</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not expressed: gills, liver, spleen, intestines, middle kidney, muscle, head kidney, heart, hypothalamus, brain	qPCR	NA	Ou et al. 2020
Actinopterygii	Characidae	<i>Astyanax altiparanae</i>	NA	NA	testes	testes (strong), ovaries (weak)	not expressed: liver, gills, muscle, heart, kidney, gut, brain	qPCR	NA	Adolfi et al. 2015
Actinopterygii	Characidae	<i>Astyanax scabripinnis</i>	GSD (ZZ/ZW)	candidate	NA	testes	not examined	qPCR	NA	Castro et al. 2019
Actinopterygii	Cichlidae	<i>Oreochromis niloticus</i>	GSD (XX/XY)	NA	testes	NA	not examined	semi-qPCR, testes differentiation, function	NA	Kobayashi et al. 2007, Qi et al. 2024
Actinopterygii	Clariidae	<i>Clarias fuscus</i>	GSD (XX/XY)	NA	testes (strong), ovaries	testes (strong), ovaries	not examined	qPCR, RNA	NA	Lin et al. 2021
Actinopterygii	Clariidae	<i>Clarias galiepinus</i>	GSD	NA	testes	testes	not expressed: heart, muscle, liver, kidney, spleen, brain	PCR, histol	NA	Raghuvver and Senthikumar 2009
Actinopterygii	Cynoglossidae	<i>Cynoglossus semilaevis</i>	GSD (ZZ/ZW)	candidate	testes	testes	not examined	qPCR	testes differentiation, spermatogenesis	Cui et al. 2017
Actinopterygii	Cyprinidae	<i>Carassius auratus gibelio</i>	GSD (XX/XY)	NA	NA	testes (strong), ovaries (weak)	not expressed: heart, liver, hypothalamus, pituitary, kidney, spleen	qPCR	NA	Wang et al. 2022a
Actinopterygii	Cyprinidae	<i>Culter alburnus</i>	NA	NA	testes	testes (strong), ovaries (weak)	very low expression: eye, brain, heart, kidney, muscle; no expression: liver, spleen	qPCR	NA	Jia et al. 2018
Actinopterygii	Cyprinidae	<i>Cyprinus carpio</i>	GSD (XX/XY)	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR, histc	NA	Tao et al. 2022, Wang et al. 2023
Actinopterygii	Cyprinidae	<i>Cyprinus carpio haematopterus</i>	GSD (XX/XY)	NA	testes (strong), ovaries (weak)	NA	not examined	qPCR	NA	Jiang et al. 2020
Actinopterygii	Cyprinidae	<i>Danio rerio</i>	GSD (ZZ/ZW)	NA	testes, ovaries	testes (strong), ovaries (weak)	not examined	qPCR, PCR	male germ cells proliferation, maintenance	Guo et al. 2005, Webster et al. 2017
Actinopterygii	Cyprinidae	<i>Gnathopogon caerulescens</i>	GSD (XX/XY)	NA	NA	testes (strong), ovaries (weak)	not expressed: brain, pituitary, eye, gills, heart, liver, gall bladder, spleen, kidney, air-bladder	PCR	NA	Ashida et al. 2013
Actinopterygii	Cyprinidae	<i>Gobiocypris rarus</i>	GSD	NA	testes	testes	not expressed: brain, heart, liver, muscle	PCR	NA	Zhang et al. 2009
Actinopterygii	Cyprinidae	<i>Labeo rohita</i>	NA	NA	NA	testes (strong), ovaries	very weak: brain, heart, kidney, liver; not expressed: muscle	qPCR	NA	Sahoo et al. 2019
Actinopterygii	Cyprinidae	<i>Megalobrama amblycephala</i>	NA	NA	males (strong), females (weak)	testes	not expressed: brain, gills, kidney, liver, muscle, spleen	qPCR	NA	Su et al. 2015
Actinopterygii	Cyprinidae	<i>Puntius sophore</i>	NA	NA	testes, ovaries	testes, ovaries	not examined	qPCR	NA	Mahanty et al. 2019
Actinopterygii	Cyprinidae	<i>Schizothorax kozlovi</i>	NA	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not expressed: gills, muscle, spleen, liver, kidney, heart, brain	qPCR	NA	He et al. 2020
Actinopterygii	Diodontidae	<i>Diodon hystrix</i>	NA	NA	testes, ovaries NA	testes, ovaries NA	not examined	qPCR, RNA	NA	Chen et al. 2021
Actinopterygii	Gadidae	<i>Gadus morhua</i>	GSD (XX/XY)	NA	testes	testes (strong), ovaries (weak)	not expressed: brain, eye, head kidney, gills, heart, muscle, pyloric caeca, spleen, liver	PCR	NA	Johnsen et al. 2010
Actinopterygii	Ictaluridae	<i>Ictalurus punctatus</i>	GSD (XX/XY)	NA	testes	testes (strong), ovaries (weak)	not expressed: brain, kidney, head kidney, liver	qPCR	NA	Xu et al. 2022
Actinopterygii	Labridae	<i>Haliclaeres trimaculatus</i>	Protogynous hermaphrodite	no	NA	gonads during sex transformation, testes	not expressed: brain, gill, intestine, liver, gallbladder, muscle, fin	qPCR	possibly sex transition	Miyake et al. 2012
Actinopterygii	Labridae	<i>Haliclaeres trimaculatus</i>	Protogynous hermaphrodite	no	NA	testes (strong), ovaries (weak)	not examined	qPCR, histc	possibly sex transition	Nozu et al. 2015
Actinopterygii	Labridae	<i>Lates calcarifer</i>	Protandrous hermaphrodite	no	NA	testes (Dmrt1a), testes and ovaries (Dmrt1b)	not examined	qPCR	NA	Domingos et al. 2018
Actinopterygii	Mastacembelidae	<i>Mastacembelus armatus</i>	Protogynous hermaphrodite	no	NA	testes (strong), ovotestes (weak) ovaries (weak)	not expressed: spleen, tail kidney, muscle, pectoral fin, heart, gill, eye, brain, liver	qPCR, RNA	NA	Xue et al. 2021
Actinopterygii	Moronidae	<i>Dicentrarchus labrax</i>	NA	NA	testes (strong), ovaries (weak)	testes	not expressed: brain, liver, heart, kidney, muscle, intestine, branchial filament	PCR	NA	Deloffre et al. 2009
Actinopterygii	Oplegnathidae	<i>Oplegnathus fasciatus</i>	GSD (X1X1X2X2/X1X2Y)	NA	NA	testes	very weak expression: liver; not expressed: heart, muscle, kidney, gills, intestine, brain	qPCR	NA	Li et al. 2021
Actinopterygii	Oplegnathidae	<i>Oplegnathus punctatus</i>	GSD (X1X1X2X2/X1X2Y)	no	males (strong), females (weak)	testes (strong), ovaries (weak)	expressed: stomach; very weak expression: liver, intestine, gill; not expressed: brain, kidney, muscle, skin, heart, spleen, e	qPCR	regulation of genes involved in SD, gonad differentiation	Zhao et al. 2024a
Actinopterygii	Ophronemidae	<i>Betta splendens</i>	GSD (XX/XY)	candidate	males (testes)	testes	not examined	qPCR	gonad differentiation	Wang et al. 2022b
Actinopterygii	Osteoglossidae	<i>Scleropages formosus</i>	GSD (ZZ/ZW)	NA	NA	testes (strong), ovaries (weak)	not examined	RNAseq	NA	Zhao et al. 2024b
Actinopterygii	Oxudercidae	<i>Acanthopagrus amnatorus</i>	NA	NA	NA	testes (strong), ovaries (weak)	not examined	qPCR	NA	Kong et al. 2024
Actinopterygii	Paralichthyidae	<i>Paralichthys olivaceus</i>	GSD (XX/XY)	NA	NA	testes (strong), ovaries (weak)	not examined	qPCR	NA	Wan et al. 2013, Shu et al. 2022
Actinopterygii	Plecoglossidae	<i>Plecoglossus altivelis</i>	GSD (XX/XY)	NA	NA	testes (strong), ovaries (weak)	very weak expression: intestine, spleen; not expressed: brain, heart, kidney, liver, muscle	qPCR	NA	Wang et al. 2014
Actinopterygii	Poeciliidae	<i>Poecilia reticulata</i>	GSD (XX/XY)	NA	testes	testes, ovaries	not examined	histology	NA	Kamaszewski et al. 2020
Actinopterygii	Poeciliidae	<i>Xiphophorus maculatus</i>	GSD (XX/XY/WY/Y)	NA	NA	testes	not expressed: brain, eye, gills, liver, white muscle, skin	PCR	NA	Veth et al. 2006
Actinopterygii	Pomacentridae	<i>Amphiprion bicinctus</i>	Protandrous hermaphrodite	no	NA	testes (strong), ovaries (weak)	not examined	qPCR	NA	Casas et al. 2016
Actinopterygii	Pomacentridae	<i>Amphiprion clarkii</i>	Protandrous hermaphrodite	no	NA	testes (weak), ovaries (strong)	not examined	qPCR, RNA	NA	Wang et al. 2022c
Actinopterygii	Rivulidae	<i>Austrolebias charrua</i>	GSD	NA	testes	testes	not examined	PCR	NA	Arezo et al. 2014
Actinopterygii	Salmonidae	<i>Brachymystax tsilingensis</i>	NA	NA	NA	testes	not examined	qPCR	NA	Huang et al. 2023
Actinopterygii	Salmonidae	<i>Oncorhynchus mykiss</i>	GSD (XX/XY)	NA	undifferentiated gonads in XY individual (strong)	testes (strong), ovaries (weak)	not expressed: skin, heart, liver, kidney	semi-qPCR	NA	Murchand et al. 2020
Actinopterygii	Scatophagidae	<i>Scatophagus argus</i>	GSD (XX/XY)	candidate	NA	testes	not examined	PCR	NA	Mustapha et al. 2018
Actinopterygii	Scatophagidae	<i>Scatophagus tetracanthus</i>	GSD (XX/XY)	candidate (Dmrt1Y)	NA	testes (Dmrt1ΔX, Dmrt1Y), ovaries (Dmrt1ΔX weak)	not examined	qPCR, PCR	NA	Peng et al. 2023
Actinopterygii	Scatophagidae	<i>Selenotoca multifasciata</i>	GSD (XX/XY)	candidate (Dmrt1Y)	NA	testes (Dmrt1X, Dmrt1Y), ovaries (Dmrt1X)	not examined	PCR, RNAs	NA	Jiang et al. 2022
Actinopterygii	Sciaenidae	<i>Larimichthys crocea</i>	NA	NA	NA	testes	not examined	qPCR, RNA	NA	Xiao et al. 2019, Wan et al. 2020
Actinopterygii	Sciaenidae	<i>Nbeo albiflora</i>	NA	NA	NA	testes (strong), ovaries (weak)	not examined	qPCR, RNA	NA	Zhu et al. 2022
Actinopterygii	Scorpaenidae	<i>Sebastes schlegelii</i>	GSD	NA	NA	testes (strong), ovaries (weak)	not expressed: liver, heart, brain, kidney, spleen, gill, muscle, intestine	qPCR	NA	Ma et al. 2014
Actinopterygii	Serranidae	<i>Epinephelus coioides</i>	Protogynous hermaphrodite	no	NA	testes, ovotestes (strong), ovaries (weak)	not expressed: liver, kidney, spleen, fat, heart, muscle, pituitary, brain, stomach, intestine, gills	qPCR, PCR	possibly sex transition	Xia et al. 2007, Chen et al. 2019, Lyu et al. 2019
Actinopterygii	Serranidae	<i>Epinephelus merra</i>	Protogynous hermaphrodite	no	NA	testes (strong), ovaries	not expressed: brain, gills, heart, intestine, kidney, liver, muscle, ovary, pituitary, spleen	qPCR, PCR	possibly sex transition	Alam et al. 2008
Actinopterygii	Sillaginidae	<i>Sillago sihama</i>	NA	NA	NA	testes (strong), ovaries	not examined	RNAseq	NA	Tian et al. 2019
Actinopterygii	Siniperidae	<i>Siniperca chuatsi</i>	GSD (XX/XY)	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	very weak expression: optic-tectum, cerebellum, hypothalamus, medulla oblongata, pituitary, heart, liver, intestine, stomi	qPCR	NA	Han et al. 2021
Actinopterygii	Sparidae	<i>Acanthopagrus latus</i>	Protandrous hermaphrodite	no	NA	testes (strong), ovotestes (weak), ovaries (weak)	not expressed: liver, spleen, kidney, pituitary, brain	qPCR	NA	Li et al. 2020

Actinopterygii	Sparidae	<i>Acanthopogon schlegelii</i>	Protandrous hermaphrodite	no	NA	testes (strong), ovaries (weak)	very weak expression: seminal vesicle; not expressed: brain, pituitary, heart, head kidney, kidney, liver, spleen, intestine	qPCR, hist: testes differentiation; germ cell maintenance	He et al. 2003, Wu et al. 2012
Actinopterygii	Sparidae	<i>Sparus auratus</i>	Protandrous hermaphrodite	no	NA	testes, ovaries NA	not examined	qPCR NA	Liarte et al. 2007
Actinopterygii	Synbranchidae	<i>Monopterus albus</i>	Protogynous hermaphrodite	no	NA	testes (strong), ovotestes, ovaries (weak)	not examined	qPCR, PCR possibly sex transition	Huang et al. 2005
Actinopterygii	Tetraodontidae	<i>Takifugu bimaculatus</i>	GSD	NA	testes	NA	not examined	semi-qPCR NA	Zhong et al. 2022
Actinopterygii	Tetraodontidae	<i>Takifugu rubripes</i>	GSD (XX/XY)	NA	males (strong), females (weak)	NA	not examined	qPCR, RNA NA	Yan et al. 2021
Amphibia	Bombinatoridae	<i>Bombina orientalis</i>	GSD	NA	testes (strong), ovaries (weak)	NA	not examined	histology NA	Piprek et al. 2014
Amphibia	Bufo	<i>Bufo viridis</i>	GSD (XX/XY)	NA	testes (strong), ovaries (weak)	NA	not examined	histology NA	Piprek et al. 2014
Amphibia	Bufo	<i>Bufo marinus</i>	GSD (ZZ/ZW)	NA	testes, ovaries	NA	not examined	qPCR NA	Abramyan et al. 2009
Amphibia	Cryptobranchidae	<i>Andrias davidianus</i>	GSD	NA	testes	NA	not examined	qPCR, RNA NA	Yang et al. 2024
Amphibia	Hylidae	<i>Hyla arborea</i>	GSD (XX/XY)	candidate	testes	NA	not examined	histology NA	Piprek et al. 2014
Amphibia	Hynobiidae	<i>Hynobius retardatus</i>	GSD	NA	testes	NA	not examined	PCR NA	Sakata et al. 2006
Amphibia	Leptodactylidae	<i>Physalaemus pustulosus</i>	GSD	NA	testes, ovaries	NA	not examined	qPCR	Duarte-Guterman et al. 2012
Amphibia	Pipidae	<i>Xenopus laevis</i>	GSD (ZZ/ZW)	yes (DmW)	testes (Dmrt1), ovaries (Dmrt1 weak, DmW)	testes (Dmrt1), ovaries (DmW weak, Dmrt1 weak)	DmW expressed: liver; very weak expression mesonephros; DmW not expressed: intestine, brain, stomach, spleen, heart	qPCR, PCR NA	Yoshimoto et al. 2008, 2010, Piprek et al. 2018, Fujitani et al. 2016, Mawaribuchi et al. 2017
Amphibia	Pipidae	<i>Xenopus tropicalis</i>	GSD (VZ, WY, ZZ/ZW, WW)	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR NA	Duarte-Guterman et al. 2011
Amphibia	Ranidae	<i>Glandirana rugosa</i>	GSD (XX/XY; ZZ/ZW)	no	testes	testes	not expressed: brain, heart, lung, liver, muscle, kidney, pancreas, spleen	PCR NA	Shibata et al. 2002,
Amphibia	Ranidae	<i>Odorrana tormota</i>	GSD	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR, RNA NA	Shu et al. 2019
Amphibia	Ranidae	<i>Pelophylax nigromaculatus</i>	GSD	NA	testes, ovaries	testes, ovaries	not examined	qPCR NA	Xu et al. 2015
Amphibia	Ranidae	<i>Rana arvalis</i>	GSD (XX/XY)	NA	testes (strong), ovaries (weak)	NA	not examined	histology NA	Piprek et al. 2014
Amphibia	Ranidae	<i>Rana chensinensis</i>	GSD	NA	undifferentiated gonads and testes	testes	not expressed: brain, liver, kidney	PCR NA	Li et al. 2019
Amphibia	Ranidae	<i>Rana temporaria</i>	GSD (XX/XY)	candidate	testes (strong), ovaries (weak)	NA	not examined	histology NA	Piprek et al. 2014
Amphibia	Salamandridae	<i>Cynops orientalis</i>	GSD	NA	testes	testes	not examined	qPCR, RNA NA	Biscotti et al. 2020
Amphibia	Salamandridae	<i>Pleurodeles waltl</i>	GSD (ZZ/ZW)	NA	testes, ovaries NA	testes, ovaries NA	not examined	PCR NA	Flament et al. 2009
Aves	Anatidae	<i>Cairina moschata</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR, RNA SD, gonad differentiation	Wang et al. 2015, Bai et al. 2020
Aves	Casuaridae	<i>Dromolus novaehollandiae</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR SD, gonad differentiation	Hirst et al. 2017
Aves	Estrilidae	<i>Taeniopygia guttata</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	NA	not examined	qPCR SD, gonad differentiation	Hirst et al. 2017
Aves	Phasianidae	<i>Coturnix japonica</i>	GSD (ZZ/ZW)	candidate	testes (strong), ovaries (weak)	testes	not examined	RNAseq SD, gonad differentiation	Okuno et al. 2020
Aves	Phasianidae	<i>Gallus gallus domesticus</i>	GSD (ZZ/ZW)	yes (Dmrt1)	testes (strong), ovaries (weak)	testes	expressed: heart; not expressed: liver, spleen, lung, kidney	qPCR, hist: SD, gonad differentiation, germ cell proliferation	Zhao et al. 2007, Lambeth et al. 2014, Clinton and Zhao et al. 2023
Mammalia	Bovidae	<i>Bos taurus</i>	GSD (XX/XY)	no	testes (strong), ovaries (weak)	NA	not examined	qPCR NA	Pianelli et al. 2020
Mammalia	Bovidae	<i>Capra hircus</i>	GSD (XX/XY)	no	testes, ovaries NA	testes, ovaries NA	not examined	qPCR, hist: testes integrity, reduction of immune response	Wei et al. 2021
Mammalia	Bovidae	<i>Ovis aries</i>	GSD (XX/XY)	NA	testes, ovaries NA	testes, ovaries NA	not examined	qPCR NA	Li et al. 2018
Mammalia	Hominidae	<i>Homo sapiens sapiens</i>	GSD (XX/XY)	no	testes	testes	not examined	qPCR, PCR, gonad differentiation	Raymond et al. 1999; Moniot et al. 2000, Cheng et al. 2006, Ledig et al. 2010
Mammalia	Leporidae	<i>Oryctolagus cuniculus</i>	GSD (XX/XY)	no	testes, ovaries	NA	not examined	qPCR, hist: NA	Khali et al. 2018, Dujardin et al. 2023
Mammalia	Macropodidae	<i>Notamacropus eugenii</i>	GSD (XX/XY)	no	testes, ovaries	testes, ovaries	not examined	histology NA	Pask et al. 2003
Mammalia	Muridae	<i>Apodemus sylvaticus</i>	GSD (XX/XY)	no	NA	testes, ovaries NA	not examined	histology NA	Massoud et al. 2021
Mammalia	Muridae	<i>Microtus duodecimcostatus</i>	GSD (XX/XY)	no	NA	testes, ovaries NA	not examined	histology NA	Lao-Pérez et al. 2021
Mammalia	Muridae	<i>Mus musculus</i>	GSD (XX/XY)	no	testes	testes	not examined	qPCR, hist: NA	Rahmoun et al. 2014
Mammalia	Muridae	<i>Mus musculus</i>	GSD (XX/XY)	no	testes, ovaries	testes, ovaries	not examined	histology, gonad differentiation, spermatogenesis	Pask et al. 2003, Lei et al. 2007, Kim et al. 2007
Mammalia	Muridae	<i>Mus spretus</i>	GSD (XX/XY)	no	testes, ovaries NA	testes, ovaries NA	not examined	histology NA	Massoud et al. 2021
Mammalia	Ornithorhynchidae	<i>Ornithorhynchus anatinus</i>	GSD (X1X2X3X4X5X1X2X3X4X5/X1X2X3X4X5Y1Y2Y3Y4Y5)	no	NA	testes	not expressed: liver, kidney, brain, muscle	PCR, histol NA	Tsend-Ayush et al. 2009
Mammalia	Soricidae	<i>Crocidura russula</i>	GSD (XX/XY)	no	testes, ovaries NA	NA	not examined	histology NA	Massoud et al. 2014
Mammalia	Suidae	<i>Sus domesticus</i>	GSD (XX/XY)	no	testes, ovaries NA	NA	not examined	qPCR NA	Stenhouse et al. 2022
Mammalia	Tachyglossidae	<i>Tachyglossus aculeatus</i>	GSD (X1X2X3X4X5X1X2X3X4X5/X1X2X3X4X5Y1Y2Y3Y4)	no	testes, ovaries	testes, ovaries	not examined	PCR, histol NA	Shearwin-Whitatt et al. 2024
Mammalia	Talpidae	<i>Talpa occidentalis</i>	GSD (XX/XY)	no	testes, ovaries	testes, ovaries NA	not examined	qPCR, hist: NA	Zurita et al. 2005, Carmona et al. 2009, Dadhich et al. 2011
Reptilia	Agamidae	<i>Papano vitticeps</i>	GSD (ZZ/ZW)	NA	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	RNAseq NA	Wagner et al. 2023
Reptilia	Alligatoridae	<i>Alligator mississippiensis</i>	TSD	no	testes (strong), ovaries (weak)	NA	not examined	qPCR NA	Urushitani et al. 2011
Reptilia	Alligatoridae	<i>Alligator sinensis</i>	TSD	no	NA	testes (strong), ovaries (weak)	not examined	RNAseq NA	Wan et al. 2013
Reptilia	Chelonidae	<i>Lepidochelys olivacea</i>	TSD	no	undifferentiated gonads in MPT, testes (stro)	NA	not examined	qPCR NA	Diaz-Hernández et al. 2020
Reptilia	Chelydridae	<i>Chelydra serpentina</i>	TSD	no	testes, ovaries (weak)	NA	not examined	qPCR NA	Rhen et al. 2007
Reptilia	Crocodylidae	<i>Crocodylus palustris</i>	TSD	no	testes (strong), ovaries (weak)	NA	not examined	qPCR NA	Anand et al. 2008
Reptilia	Emydidae	<i>Chrysemys picta</i>	TSD	no	testes (strong), ovaries (weak)	NA	not examined	qPCR, RNA NA	Mizoguchi and Valenzuela 2020, Mizoguchi et al. 2022
Reptilia	Emydidae	<i>Testudo scripta</i>	TSD	yes (Dmrt1)	testes (strong), ovaries (weak)	NA	not examined	qPCR, hist: gonad differentiation	Bieser and Wübeler 2014, Ge et al. 2017
Reptilia	Eublepharidae	<i>Eublepharis macularius</i>	TSD	no	testes (strong), ovaries (weak)	NA	not examined	qPCR, PCR NA	Pallotta et al. 2022
Reptilia	Eublepharidae	<i>Mauermys mutica</i>	TSD	no	testes (strong), ovaries (weak)	NA	not examined	histology NA	Liu et al. 2021
Reptilia	Geomydidae	<i>Chinemys reevesii</i>	TSD	no	NA	testes, ovaries NA	not expressed: liver, spleen, brain, kidney	PCR NA	Hoshi and Nakao et al. 2008
Reptilia	Geomydidae	<i>Mauermys reevesii</i>	TSD	no	testes (strong), ovaries (weak)	testes (strong), ovaries (weak)	not examined	qPCR RACE NA	Tang et al. 2018, Shu et al. 2020
Reptilia	Lacertidae	<i>Podarcis sicula</i>	GSD (ZZ/ZW)	NA	NA	testes	not expressed: liver, heart	semi-qPCR NA	Capriglione et al. 2010
Reptilia	Trionychidae	<i>Pelodiscus sinensis</i>	GSD (ZZ/ZW)	NA	testes	testes	not expressed: muscle, kidney, lung, spleen, liver; very weak expression: heart	qPCR, PCR, gonad differentiation	Sun et al. 2017

- Abramyan, J., Feng, C.W. & Koopman, P. 2009. Cloning and expression of candidate *Sex. Dev.* genes in the cane toad (*Bufo marinus*). *Dev. Dynam.* 238: 2430–2441.
- Adolfi, M.C., Carreira, A.C.O., Jesus, L.W.O., Bogerd, J., Funes, R.M., Schartl, M., et al. 2015. Molecular cloning and expression analysis of *Dmrt1* and *Sox9* during gonad development and male reproductive cycle in the lambari fish, *Astyanax Altiparanae*. *Reprod. Biol. Endocrinol.* 13: 1–15.
- Alam, M.A., Kobayashi, Y., Horiguchi, R., Hirai, T. & Nakamura, M. 2008. Molecular cloning and quantitative expression of sexually dimorphic markers *Dmrt1* and *Foxl2* during female-to-male sex change in *Epinephelus merra*. *Gen. Comp. Endocrinol.* 157: 75–85.
- Amberg, J.J., Goforth, R.R. & Sepúlveda, M.S. 2013. Antagonists to the *Wnt* cascade exhibit sex-specific expression in gonads of sexually mature shovelnose sturgeon. *Sex. Dev.* 7: 308–315.
- Anand, A., Patel, M., Lalremruata, A., Singh, A.P., Agrawal, R., Singh, L., et al. 2008. Multiple alternative splicing of *Dmrt1* during gonadogenesis in Indian mugger, a species exhibiting temperature-dependent sex determination. *Gene* 425: 56–63.
- Arezo, M.J., Papa, N., Gutterierrez, V., Garcia, G. & Berois, N. 2014. Sex determination in annual fishes: searching for the master sex-determining gene in *Austrolebias charrua* (Cyprinodontiformes, Rivulidae). *Genet. Mol. Biol.* 37: 364–374.
- Ashida, H., Ueyama, N., Kinoshita, M. & Kobayashi, T. 2013. Molecular identification and expression of *Foxl2* and *Dmrt1* genes from willow minnow *Gnathopogon caeruleus*. *Reprod. Biol.* 13: 317–324.
- Bai, D.P., Chen, Y., Hu, Y.Q., He, W.F., Shi, Y.Z., Fan, Q.M., et al. 2020. Transcriptome analysis of genes related to gonad differentiation and development in Muscovy ducks. *BMC Genomics* 21: 1–17.
- Berbejillo, J., Martínez-Bengochea, A., Bedó, G. & Vizziano-Cantonnet, D. 2013. Expression of *Dmrt1* and *Sox9* during gonadal development in the Siberian sturgeon (*Acipenser baerii*). *Fish Physiol. Biochem.* 39: 91–94.
- Bieser, K.L. & Wibbels, T. 2015. Chronology, magnitude and duration of expression of putative sex-determining/differentiation genes in a turtle with temperature-dependent sex determination. *Sex. Dev.* 8: 364–375. S.
- Biscotti, M.A., Carducci, F., Barucca, M., Gerdol, M., Pallavicini, A., Schartl, M., et al. 2020. The transcriptome of the newt *Cynops orientalis* provides new insights into evolution and function of sexual gene networks in sarcopterygians. *Sci. Rep.* 10: 1–14.
- Brunner, B., Hornung, U., Shan, Z., Nanda, I., Kondo, M., Zend-Ajusch, E., et al. 2001. Genomic organization and expression of the Doublesex-related gene cluster in vertebrates and detection of putative regulatory regions for *Dmrt1*. *Genomics* 77: 8–17.
- Barucca, A., Popa, G.-O., Florescu, I.E., Maereanu, M., Dudu, A., Georgescu, S.E., et al. 2018. Expression characterization of six genes possibly involved in gonad development for stellate sturgeon individuals (*Acipenser stellatus*, Pallas 1771). *Int. J. Genomics* 1: 7835637.
- Capriglione, T., Vaccaro, M.C., Morescalchi, M.A., Tammara, S. & De Iorio, S. 2010. Differential *Dmrt1* expression in the gonads of *Podarcis sicula* (Reptilia: Lacertidae). *Sex. Dev.* 4: 104–109. S.
- Carmona, F.D., Lupiáñez, D.G., Martín, J.E., Burgos, M., Jiménez, R., & Zurita, F. 2009. The spatio-temporal pattern of testis organogenesis in mammals—insights from the mole. *Int. J. Dev. Biol.* , 53: 1035–1044.
- Casas, L., Saborido-Rey, F., Ryu, T., Michell, C., Ravasi, T. & Irigoien, X. 2016. Sex change in clownfish: Molecular insights from transcriptome analysis. *Sci. Rep.* 2016:6:16: 1–19.
- Castro, J.P., Hattori, R.S., Yoshinaga, T.T., Silva, D.M.Z.D.A., Foresti, F., Santos, M.H., et al. 2019. Differential expression of *Dmrt1* in *Astyanax scabripinnis* (Teleostei, Characidae) is correlated with B chromosome occurrence. *Zebrafish* 16: 182–188.
- Chen, C.Y., Tsai, Y.J. & Chang, C.F. 2019. The roles of *Cyp19a1a* and *Dmrt1* during gonadal sex differentiation and sex change in orange-spotted grouper, *Epinephelus coioides*. *J. Mar. Sci. Tech.* 27: 282–291.
- Chen, H., Li, Z., Wang, Y., Huang, H., Yang, X., Li, S., et al. 2021. Comparison of gonadal transcriptomes uncovers reproduction-related genes with sexually dimorphic expression patterns in diodon hystrix. *Animals* 11: 1042.
- Cheng, P.L., Dumas, G.A., Smith, J.T., Leger, A., Plamondon, A., McGrath, M.J., et al. 2006. Transcriptional diversity of *Dmrt1* (dsx- and mab3-related transcription factor 1) in human testis. *Cell Res.* 16: 389–393.
- Clinton, M. & Zhao, D. 2023. Avian sex determination: A chicken and egg conundrum. *Sex. Dev.* 17: 120–133. S.
- Cui, Z., Liu, Y., Wang, W., Wang, Q., Zhang, N., Lin, F., et al. 2017. Genome editing reveals *Dmrt1* as an essential male sex-determining gene in Chinese tongue sole (Cynoglossus semilaevis). *Sci. Rep.* 7: 1–10.
- Dadhich, R.K., Barriounevo, F.J., Lupiáñez, D.G., Real, F.M., Burgos, M. & Jiménez, R. 2011. Expression of genes controlling testicular development in adult testis of the seasonally breeding Iberian mole. *Sex. Dev.* 5: 77–88.
- Deloffre, L.A.M., Martins, R.S.T., Mylonas, C.C. & Canario, A.V.M. 2009. Alternative transcripts of *Dmrt1* in the European sea bass: Expression during gonadal differentiation. *Aquaculture* 293: 89–99.
- Díaz-Hernández, V., Domínguez-Mora, P., Chino-Palomo, L., Marmolejo-Valencia, A., Harfush, M. & Merchant-Larios, H. 2020. Spatiotemporal expression of *Foxl2* and *Dmrt1* before, during, and after sex determination in the sea turtle *Lepidochelys olivacea*. *Sex. Dev.* 13: 286–296.
- Domingos, J.A., Budd, A.M., Banh, Q.Q., Goldsbury, J.A., Zenger, K.R. & Jerry, D.R. 2018. Sex-specific *Dmrt1* and *Cyp19a1* methylation and alternative splicing in gonads of the protandrous hermaphrodite barramundi. *PLoS One* 13: e0204182.
- Dong, Z., Li, X., Yao, Z., Wang, C., Guo, Y., Wang, Q., et al. 2021. *Oryzias curvirostris* in Sanya does not contain the male sex-determining gene *DmY*. *Animals* 11: 1327
- Duarte-Guterman, P. & Trudeau, V.L. 2011. Transcript profiles and triiodothyronine regulation of sex steroid- and thyroid hormone-related genes in the gonad–mesonephros complex of *Silurana tropicalis*. *Mol. Cell. Endocrinol.* 331: 143–149.
- Duarte-Guterman, P., Ryan, M.J. & Trudeau, V.L. 2012. Developmental expression of sex steroid- and thyroid hormone-related genes and their regulation by triiodothyronine in the gonad–mesonephros of a Neotropical frog, *Physalaemus pustulosus*. *Gen. Comp. Endocrinol.* 177: 195–204.
- Dujardin, E., André, M., Dewaele, A., Mandon-Pépin, B., Poulat, F., Frambourg, A., et al. 2023. *Dmrt1* is a testis-determining gene in rabbits and is also essential for female fertility. *Life* 12: RP89284.
- Fajkowska, M., Rzepkowska, M., Adamek, D., Ostaszewska, T. & Szczepkowski, M. 2016. Expression of *Dmrt1* and vtg genes during gonad formation, differentiation and early maturation in cultured Russian sturgeon *Acipenser gueldenstaedtii*. *J. Fish. Biol.* 89: 1441–1449.
- Fernandino, J.I., Guilgur, L.G. & Somoza, G.M. 2006. *Dmrt1* expression analysis during spermatogenesis in pejerrey, *Odontesthes bonariensis*. *Fish. Physiol. Biochem.* 32: 231–240.
- Flament, S., Dumond, H., Chardard, D. & Chesnel, A. 2009. Lifelong testicular differentiation in *Pleurodeles waltl* (Amphibia, Caudata). *Reprod. Biol. Endocrinol.* 7: 1–12.
- Forconi, M., Canapa, A., Barucca, M., Biscotti, M.A., Capriglione, T., Buonocore, F., et al. 2013. Characterization of sex determination and sex differentiation genes in *Latimeria*. *PLoS One.* 8.
- Fujitani, K., Otomo, A., Wada, M., Takamatsu, N. & Ito, M. 2016. Sexually dimorphic expression of *Dmrt1* and *γH2AX* in germ stem cells during gonadal development in *Xenopus laevis*. *FEBS Open Bio.* 6: 276–284.
- Ge, C., Ye, J., Zhang, H., Zhang, Y., Sun, W., Sang, Y., et al. 2017. *Dmrt1* induces the male pathway in a turtle species with temperature-dependent sex determination. *Development* 144: 2222–2233.
- Guo, Y., Cheng, H., Huang, X., Gao, S., Yu, H. & Zhou, R. 2005. Gene structure, multiple alternative splicing, and expression in gonads of zebrafish *Dmrt1*. *Biochem. Biophys. Res. Commun.* 330: 950–957.
- Hale, M.C., Jackson, J.R. & DeWoody, J.A. 2010. Discovery and evaluation of candidate sex-determining genes and xenobiotics in the gonads of lake sturgeon (*Acipenser fulvescens*). *Genetica* 138: 745–756.
- Han, C., Wang, C., Ouyang, H., Zhu, Q., Huang, J., Han, L., et al. 2021. Characterization of dmrts and their potential role in gonadal development of mandarin fish (*Siniperca chuatsi*). *Aquac. Rep.* 21: 100802.
- He, C.L., Du, J.L., Wu, G.C., Lee, Y.H., Sun, L.T. & Chang, C.F. 2003. Differential *Dmrt1* transcripts in gonads of the protandrous black porgy, *Acanthopagrus schlegelii*. *Cytogenet. Genome Res.* 101: 309–313. S.
- He, Y., Wu, X., Zhu, Y. & Yang, D. 2020. Expression Profiles of *Dmrt1* in *Schizothorax kozlovi*, and their relation to CpG methylation of its promoter and temperature. *Zoolog. Sci.* 37: 140.
- Hirst, C.E., Major, A.T., Ayers, K.L., Brown, R.J., Mariette, M., Sackton, T.B., et al. 2017. Sex reversal and comparative data undermine the W chromosome and support Z-linked *Dmrt1* as the regulator of gonadal sex differentiation in birds. *Endocrinology* 158: 2970–2987.
- Hoshi, H. & Nakao, A. 2008. Molecular cloning of full-length *Dmrt1* cDNA of reeves turtle (*Chinemys reevesii*). *J. Vet. Med. Sci.* 70: 687–692.
- Hu, Y., Tan, R., Zhu, X., Wang, B., Wang, J., Guo, B., et al. 2024. Genome-wide identification, phylogeny and expressional profile of the *Dmrt* gene family in Chinese sturgeon (*Acipenser sinensis*). *Sci. Rep.* 14: 4231.
- Huang, L., Ye, H., Yue, H., Leng, X., Ruan, R., Du, H., et al. 2023. Comparative transcriptome analysis reveals sexually dimorphic gene expression in the gonads of *Brachymystax tsinlingensis*. *Animals* 13: 3690.
- Huang, X., Guo, Y., Shui, Y., Gao, S., Yu, H., Cheng, H., et al. 2005. Multiple alternative splicing and differential expression of *Dmrt1* during gonad transformation of the rice field eel. *Biol. Reprod.* 73: 1017–1024.
- Jeng, S.R., Wu, G.C., Yueh, W.S., Kuo, S.F., Dufour, S. & Chang, C.F. 2019. *Dmrt1* (doublesex and mab-3-related transcription factor 1) expression during gonadal development and spermatogenesis in the Japanese eel. *Gen. Comp. Endocrinol.* 279: 154–163.
- Jia, Y., Zheng, J., Chi, M., Liu, S., Jiang, W., Cheng, S., et al. 2019. Molecular identification of *Dmrt1* and its promoter CpG methylation in correlation with gene expression during gonad development in *Culter alburnus*. *Fish Physiol. Biochem.* 45: 245–252.
- Jiang, D.N., Huang, Y.Q., Zhang, J.M., Mustapha, U.F., Peng, Y.X., Huang, H., et al. 2022. Establishment of the Y-linked *Dmrt1Y* as the candidate sex determination gene in spotbanded scat (*Selenotoca multifasciata*). *Aquac. Rep.* 23: 101085.
- Jiang, M., Jia, S., Chen, J., Chen, K., Ma, W., Wu, X., et al. 2020. Timing of gonadal development and dimorphic expression of sex-related genes in gonads during early sex differentiation in the yellow river carp. *Aquaculture* 518: 734825.
- Johnsen, H., Seppola, M., Torgersen, J.S., Delghandi, M. & Andersen, Ø. 2010. Sexually dimorphic expression of *Dmrt1* in immature and mature Atlantic cod (*Gadus morhua* L.). *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 156: 197–205.
- Kamaszewski, M., Skrobisz, M., Wójcik, M., Kawalski, K., Szczepański, A., Bujarski, P., et al. 2020. The role of transcription factors in gonad development and sex differentiation of a teleost model fish—guppy (*Poecilia reticulata*). *Animals* 10: 2401.
- Khalil, I., Eman K., Metwally, M.A., El-Shafey, A.A. & Eshra, E.A. 2018. Expression pattern of *Dmrt1* and *Stra8* genes during postnatal development of rabbit testes. *J. Vet. Anat.* 11: 15–29.
- Kim, S., Bardwell, V.J. & Zarkower, D. 2007. Cell type-autonomous and non-autonomous requirements for *Dmrt1* in postnatal testis differentiation. *Dev. Biol.* 307: 314–327.
- Kobayashi, T., Kajiura-Kobayashi, H., Guan, G. & Nagahama, Y. 2008. Sexual dimorphic expression of *Dmrt1* and *Sox9a* during gonadal differentiation and hormone-induced sex reversal in the teleost fish Nile tilapia (*Oreochromis niloticus*). *Dev. Dynam.* 237: 297–306.

Kobayashi, T., Matsuda, M., Kajjura-Kobayashi, H., Susuki, A., Saito, N., Nakamoto, M., et al. 2004. Two DM domain genes, *DmY* and *Dmrt1*, involved in testicular differentiation and development in the medaka, *Oryzias latipes*. *Dev. Dynam.* 231: 518–526.

Kong, D., Zhao, X., Pan, Y. & Song, N. 2024. Gonadal transcriptome analysis of sex-biased gene and genome-wide investigation of Dmrt gene family in *Acanthogobius ommaturus*. *Comp. Biochem. Physiol. Part D Genomics Proteomics* 50: 101204.

Lambeth, L.S., Raymond, C.S., Roeszler, K.N., Kuroiwa, A., Nakata, T., Zarkower, D., et al. 2014. Over-expression of *Dmrt1* induces the male pathway in embryonic chicken gonads. *Dev. Biol.* 389: 160–172.

Lao-Pérez, M., Massoud, D., Real, F.M., Hurtado, A., Ortega, E., Burgos, M., et al. 2021. Mediterranean pine vole, *Microtus duodecimcostatus*: A paradigm of an opportunistic breeder. *Animals* 11: 1639.

Ledig, S., Hiort, O., Scherer, G., Hoffmann, M., Wolff, G., Morlot, S., et al. 2010. Array-CGH analysis in patients with syndromic and non-syndromic XY gonadal dysgenesis: evaluation of array CGH as diagnostic tool and search for new candidate loci. *Hum. Reprod.* 25: 2637–2646.

Lei, N., Hornbaker, K.I., Rice, D.A., Karpova, T., Agbor, V.A. & Heckert, L.L. 2007. Sex-specific differences in mouse *Dmrt1* expression are both cell type- and stage-dependent during gonad development. *Biol. Reprod.* 77: 466–475.

Leng, X.Q., Du, H.J., Li, C.J. & Cao, H. 2016. Molecular characterization and expression pattern of *Dmrt1* in the immature Chinese sturgeon *Acipenser sinensis*. *J. Fish. Biol.* 88: 567–579.

Li, H., Zhu, Q., Chen, R., Liu, M. & Xu, D. 2021. Identification and characterization of dimorphic expression of sex-related genes in rock bream, a fish with multiple sex chromosomes. *Front. Genet.* 12: 791179.

Li, S., Lin, G., Fang, W., Huang, P., Gao, D., Huang, J., et al. 2020. Gonadal transcriptome analysis of sex-related genes in the protandrous yellowfin seabream (*Acanthopagrus latus*). *Front. Genet.* 11: 551435.

Li, T., Lu, Z., Luo, R., Gao, J., Zhao, X. & Ma, Y. 2018. Expression and cellular localization of double sex and mab-3 related transcription factor 1 in testes of postnatal Small-Tail Han sheep at different developmental stages. *Gene* 642: 467–473.

Li, X.-Y., Jin, W.-J., Lei, L. & Zhang, Y.-H. 2019. Identification and expression of *Dmrt1* and *Sox9* during the gonadal differentiation of *Rana chensinensis*. *Turk. J. Zool.* 43: 319–330.

Li, X.Y., Li, Z., Zhang, X.J., Zhou, L. & Gui, J.F. 2014. Expression characterization of testicular *Dmrt1* in both Sertoli cells and spermatogenic cells of polyploid gibel carp. *Gene* 548: 119–125.

Liarte, S., Chaves-Pozo, E., García-Alcazar, A., Mulero, V., Meseguer, J. & García-Ayala, A. 2007. Testicular involution prior to sex change in gilthead seabream is characterized by a decrease in *Dmrt1* gene expression and by massive leukocyte infiltration. *Reprod. Biol. Endocrinol.* 5: 1–15.

Lin, X., Zhou, D., Zhang, X., Li, G., Zhang, Y., Huang, C., et al. 2021. A first insight into the gonad transcriptome of hong kong catfish (*Clarias fuscus*). *Animals* 11: 1131.

Liu, W., Zhang, H., Xiang, Y., Jia, K., Luo, M. & Yi, M. 2020. A novel germ line and somatic cell expression of two sexual differentiation genes, *Dmrt1* and *Foxl2* in marbled goby (*Oxyeleotris marmorata*). *Aquaculture* 516: 734619.

Liu, X., Zhu, Y., Zhao, Y., Wang, Y., Li, W., Hong, X., et al. 2021. *Vasa* expression is associated with sex differentiation in the Asian yellow pond turtle, *Mauremys mutica*. *J. Exp. Zool. B Mol. Dev. Evol.* 336: 431–442.

Lyu, Q., Hu, J., Yang, X.K., Liu, X.C., Chen, Y., Bin, Xiao, L., et al. 2019. Expression profiles of dmrts and foxls during gonadal development and sex reversal induced by 17 $\alpha$ -methyltestosterone in the orange-spotted grouper. *Gen. Comp. Endocrinol.* 274: 26–36.

Ma, L., Wang, W., Xiang, X., Jiang, J., Song, H., Jiang, H., et al. 2014. Characterization of the *Dmrt1* gene in the black rockfish *Sebastes schlegelii* revealed a remarkable sex-dimorphic expression. *Fish Physiol. Biochem.* 40: 1263–1274.

Mahanty, A., Purohit, G.K., Mohanty, S. & Mohanty, B.P. 2019. Heat stress-induced alterations in the expression of genes associated with gonadal integrity of the teleost *Puntius sophore*. *Fish. Physiol. Biochem.* 45: 1409–1417.

Marchand, O., Govoroun, M., D’Cotta, H., McMeel, O., Lareyre, J.J., Bernot, A., et al. 2000. *Dmrt1* expression during gonadal differentiation and spermatogenesis in the rainbow trout, *Oncorhynchus mykiss*. *Biochimica et Biophysica Acta (BBA) - Gene Structure and Expression* 1493: 180–187.

Massoud, D., Barriounevo, F.J., Ortega, E., Burgos, M. & Jiménez, R. 2014. The testis of greater white-toothed shrew *Crocodyra russula* in Southern European populations: A case of adaptive lack of seasonal involution? *J. Exp. Zool. B Mol. Dev. Evol.* 322: 304–315.

Massoud, D., Lao-Pérez, M., Ortega, E., Burgos, M., Jiménez, R., Barriounevo, F.J., et al. 2021. Divergent seasonal reproductive patterns in syntopic populations of two murine species in southern Spain, *Mus spretus* and *Apodemus sylvaticus*. *Animals* 11: 243.

Mawaribuchi, S., Takahashi, S., Wada, M., Uno, Y., Matsuda, Y., Kondo, M., et al. 2017. Sex chromosome differentiation and the W- and Z-specific loci in *Xenopus laevis*. *Dev. Biol.* 426: 393–400.

Miyake, Y., Sakai, Y. & Kuniyoshi, H. 2012. Molecular cloning and expression profile of sex-specific genes, *Figla* and *Dmrt1*, in the protogynous hermaphroditic fish, *Halichoeres poecilopterus*. *Zool. Sci.* 29: 690–710.

Mizoguchi, B. & Valenzuela, N. 2020. Alternative splicing and thermosensitive expression of *Dmrt1* during urogenital development in the painted turtle, *Chrysemys picta*. *PeerJ* 2020: e8639.

Mizoguchi, B., Topping, N.E., Lavin, A.M. & Valenzuela, N. 2022. Cadmium ecotoxic effects on embryonic *Dmrt1* and aromatase expression in *Chrysemys picta* turtles may implicate changes in DNA methylation. *Genes* 13: 1318.

Moniot, B., Berta, P., Scherer, G., Südbeck, P. & Poulat, F. 2000. Male specific expression suggests role of *Dmrt1* in human sex determination. *Mech. Dev.* 91: 323–325.

Mustapha, U.F., Jiang, D.N., Liang, Z.H., Gu, H.T., Yang, W., Chen, H.P., et al. 2018. Male-specific *Dmrt1* is a candidate sex determination gene in spotted scat (*Scatophagus argus*). *Aquaculture* 495: 351–358.

Nozu, R., Horiguchi, R., Kobayashi, Y. & Nakamura, M. 2015. Expression profile of doublesex/male abnormal-3-related transcription factor-1 during gonadal sex change in the protogynous wrasse, *Halichoeres trimaculatus*. *Mol. Reprod. Dev.* 82: 859–866.

Ohmuro-Matsuyama, Y., Matsuda, M., Kobayashi, T., Ikeuchi, T. & Nagahama, Y. 2003. Expression of *DmY* and *Dmrt1* in various tissues of the medaka (*Oryzias latipes*). *Zool. Sci.* 20: 1395–1398.

Okada, H., Hagihara, S., Yamashita, K., Ijiri, S. & Adachi, S. 2017. Expression pattern of *Foxl2* and *Dmrt1* in gonad of Amur sturgeon *Acipenser schrenckii* in relation to sex differentiation. *Aquaculture* 479: 712–720.

Okuno, M., Miyamoto, S., Itoh, T., Seki, M., Suzuki, Y., Mizushima, S., et al. 2020. Expression profiling of sexually dimorphic genes in the Japanese quail, *Coturnix japonica*. *Sci. Rep.* 10: 1–11.

Ou, M., Chen, K., Gao, D., Wu, Y., Luo, Q., Liu, H., et al. 2022. Characterization, expression and CpG methylation analysis of *Dmrt1* and its response to steroid hormone in blotched snakehead (*Channa maculata*). *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 257: 110672.

Pallotta, M.M., Fogliano, C. & Carotenuto, R. 2022. Temperature incubation influences gonadal gene expression during leopard gecko development. *Animals* 12: 3186.

Pan, Z.J., Zhu, C.K., Chang, G.L., Wu, N., Ding, H.Y. & Wang, H. 2021. Differential expression analysis and identification of sex-related genes by gonad transcriptome sequencing in estradiol-treated and non-treated Ussuri catfish *Pseudobagrus ussuriensis*. *Fish. Physiol. Biochem.* 47: 565–581.

Pask, A.J., Behringer, R.R. & Renfree, M.B. 2003. Expression of *Dmrt1* in the mammalian ovary and testis – from marsupials to mice. *Cytogenet. Genome Res.* 101: 229–236. S.

Peng, Y.X., Huang, Y.Q., Zhong, J., Jiang, Z.T., Fan, S., Shi, H.J., et al. 2023. Identification of sex-linked marker and candidate sex determination gene in ornamental fish, African scat (*Scatophagus tetracanthus*). *Aquaculture* 563: 739023.

Piprek, R.P., Dambulewicz, M., Kloc, M. & Kubiak, J.Z. 2018. Transcriptome analysis identifies genes involved in sex determination and development of *Xenopus laevis* gonads. *Differentiation* 100: 46–56.

Piprek, R.P., Pecio, A., Laskowska-Kaszub, K., Kubiak, J.Z. & Szymura, J.M. 2014. Sexual dimorphism of *Amh*, *Dmrt1* and *Rspo1* localization in the developing gonads of six anuran species. *Int. J. Dev. Biol.* 57: 891–895.

Planells, B., Gómez-Redondo, I., Sánchez, J.M., McDonald, M., Cánovas, Á., Lonergan, P., et al. 2020. Gene expression profiles of bovine genital ridges during sex determination and early differentiation of the gonads. *Biol. Reprod.* 102: 38–52.

Qi, S., Dai, S., Zhou, X., Wei, X., Chen, P., He, Y., et al. 2024. *Dmrt1* is the only male pathway gene tested indispensable for sex determination and functional testis development in tilapia. *PLoS Genet.* 20: e1011210.

Raghuvver, K. & Senthilkumaran, B. 2009. Identification of multiple *Dmrt1* s in catfish: localization, dimorphic expression pattern, changes during testicular cycle and after methyltestosterone treatment. *J. Mol. Endocrinol.* 42: 437–448.

Rahmoun, M., Perez, J., Saunders, P.A., Boizet-Bonhoure, B., Wilhelm, D., Poulat, F., et al. 2015. Anatomical and molecular analyses of XY ovaries from the African Pygmy mouse *Mus minutoides*. *Sex. Dev.* 8: 356–363.

Raymond, C.S., Parker, E.D., Kettlewell, J.R., Brown, L.G., Page, D.C., Kuz, K., et al. 1999. A region of human chromosome 9p required for testis development contains two genes related to known sexual regulators. *Hum. Mol. Genet.* 8: 989–996

Rhen, T., Metzger, K., Schroeder, A. & Woodward, R. 2007. Expression of putative sex-determining genes during the thermosensitive period of gonad development in the snapping turtle, *Chelydra serpentina*. *Sex. Dev.* 1: 255–270.

Sahoo, L., Sahoo, S., Mohanty, M., Sankar, M., Dixit, S., Das, P., et al. 2021. Molecular characterization, computational analysis and expression profiling of *Dmrt1* gene in Indian major carp, *Labeo rohita* (Hamilton 1822). *Anim. Biotechnol.* 32: 413–426.

Sakata, N., Miyazaki, K. & Wakahara, M. 2006. Up-regulation of P450arom and down-regulation of *Dmrt-1* genes in the temperature-dependent sex reversal from genetic males to phenotypic females in a salamander. *Dev. Genes. Evol.* 216: 224–228.

Shearwin-Whyatt, L., Fenelon, J., Yu, H., Major, A., Qu, Z., Zhou, Y., et al. 2024. *AmhY* and sex determination in egg-laying mammals. *bioRxiv* 2024.04.18.588952.

Shibata, K., Takase, M. & Nakamura, M. 2002. The *Dmrt1* expression in sex-reversed gonads of amphibians. *Gen. Comp. Endocrinol.* 127: 232–241.

Shu, C., Wang, L., Zou, C., Tan, X., Zou, Y., Kong, L., et al. 2022. Function of *Foxl2* and *Dmrt1* proteins during gonadal differentiation in the olive flounder *Paralichthys olivaceus*. *Int. J. Biol. Macromol.* 215: 141–154.

Shu, O., Xiao-Li, L., Li Wei, Y.-K. & Xin-Ping, Z. 2020. Cloning and expression characteristics of *Dmrt1* gene of *Mauremys mutica*. *J. Hydrobiol.* 44.

Shu, Y., Zhang, H., Cai, Q., Tang, D., Wang, G., Liu, T., et al. 2019. Integrated mRNA and miRNA expression profile analyses reveal the potential roles of sex-biased miRNA-mRNA pairs in gonad tissues of the Chinese concave-eared torrent frog (*Odorrana tormota*). *J. Exp. Zool. B Mol. Dev. Evol.* 332: 69–80.

Smith, E.K., Guzmán, J.M. & Luckenbach, J.A. 2013. Molecular cloning, characterization, and sexually dimorphic expression of five major sex differentiation-related genes in a Scorpaeniform fish, sablefish (*Anoplopoma fimbria*). *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 165: 125–137.

Stenhouse, C., Cortes-Araya, Y., Donadeu, F.X. & Ashworth, C.J. 2022. Associations between testicular development and fetal size in the pig. *J. Anim. Sci. Biotechnol.* 13: 1–11.

Su, L., Zhou, F., Ding, Z., Gao, Z., Wen, J., Wei, W., et al. 2015. Transcriptional variants of *Dmrt1* and expression of four *Dmrt* genes in the blunt snout bream, *Megalobrama amblycephala*. *Gene* 573: 205–215.

Sun, W., Cai, H., Zhang, G., Zhang, H., Bao, H., Wang, L., et al. 2017. *Dmrt1* is required for primary male sexual differentiation in Chinese soft-shelled turtle *Pelodiscus sinensis*. *Sci. Rep.* 2017 7:17: 1–14.

Tang, W.Q., Mu, Y., Valenzuela, N. & Du, W.G. 2018. Effects of incubation temperature on the expression of sex-related genes in the chinese pond turtle, *Mauremys reevesii*. *Sex. Dev.* 11: 307–319.

Tao, B., Liao, X., Chen, L., Li, Y., Chen, K., Jia, S., et al. 2022. Germ cells are not essential for sexual dimorphism of gonads in common carp, *C. carpio* L. *Aquaculture* 547: 737501.

Tian, C., Li, Z., Dong, Z., Huang, Y., Du, T., Chen, H., et al. 2019. Transcriptome analysis of male and female mature gonads of silver sillago (*Sillago sihama*). *Genes* 129 10: 129.

Tsend-Ayush, E., Lim, S.L., Pask, A.J., Hamdan, D.D.M., Renfree, M.B. & Grützner, F. 2009. Characterisation of *Atrx*, *Dmrt1*, *Dmrt7* and *Wt1* in the platypus (*Ornithorhynchus anatinus*). *Reprod. Fertil. Dev.* 21: 985–991.

Úbeda-Manzanaro, M., Merto, M.A., Ortiz-Delgado, J.B., Rebordinos, L. & Sarasquete, C. 2014. Expression profiling of the sex-related gene *Dmrt1* in adults of the Lusitanian toadfish *Halobatrachus didactylus* (Bloch and Schneider, 1801). *Gene* 535: 255–265.

Urushitani, H., Katsu, Y., Miyagawa, S., Kohno, S., Ohta, Y., Guillelte, L.J., et al. 2011. Molecular cloning of anti-Müllerian hormone from the American alligator, *Alligator mississippiensis*. *Mol. Cell Endocrinol.* 333: 190–199.

Veith, A.M., Schäfer, M., Klüver, N., Schmidt, C., Schultheis, C., Schartl, M., et al. 2006. Tissue-specific expression of *Dmrt* genes in embryos and adults of the platyfish *Xiphophorus maculatus*. *Zebrafish* 3: 325–337.

Wagner, S., Whiteley, S.L., Castelli, M., Patel, H.R., Deveson, I.W., Blackburn, J., et al. 2023. Gene expression of male pathway genes *Sox9* and *Amh* during early sex differentiation in a reptile departs from the classical amniote model. *BMC Genomics* 24: 1–18.

Wan, H.F., Zhong, Z.W., Jiang, Y.H., Zou, P.F., Zhang, Z.P. & Wang, Y.L. 2020. Genome-wide investigation of *Dmrt* gene family in large yellow croaker (*Larimichthys crocea*). *Theriogenology* 156: 272–282.

Wan, Q.H., Pan, S.K., Hu, L., Zhu, Y., Xu, P.W., Xia, J.Q., et al. 2013. Genome analysis and signature discovery for diving and sensory properties of the endangered Chinese alligator. *Cell Res.* 23: 1091–1105.

Wang, H., Qu, M., Tang, W., Liu, S. & Ding, S. 2022c. Transcriptome profiling and expression localization of key sex-related genes in a socially-controlled hermaphroditic clownfish, *Amphiprion clarkii*. *Int. J. Mol. Sci.* 23: 9085.

Wang, J.H., Miao, L., Li, M.Y., Guo, X.F., Pan, N., Chen, Y.Y., et al. 2014. Cloning the *Dmrt1* and *DmrtA2* genes of ayu (*Plecoglossus altivelis*) and mapping their expression in adult, larval, and embryonic stages. *Zool. Res.* 35: 99.

Wang, L., Sun, F., Wan, Z.Y., Yang, Z., Tay, Y.X., Lee, M., et al. 2022b. Transposon-induced epigenetic silencing in the X chromosome as a novel form of *Dmrt1* expression regulation during sex determination in the fighting fish. *BMC Biol.* 20: 1–16.

Wang, M., Chen, L., Zhou, Z., Xiao, J., Chen, B., Huang, P., et al. 2023. Comparative transcriptome analysis of early sexual differentiation in the male and female gonads of common carp (*Cyprinus carpio*). *Aquaculture* 563: 738984.

Wang, M.T., Li, Z., Ding, M., Yao, T.Z., Yang, S., Zhang, X.J., et al. 2022a. Two duplicated *Gsdf* homeologs cooperatively regulate male differentiation by inhibiting *Cyp19a1a* transcription in a hexaploid fish. *PLoS Genet.* 18: e1010288.

Wang, Q., Weng, H., Chen, Y., Wang, C., Lian, S., Wu, X., et al. 2015. Early development of gonads in Muscovy duck embryos. *Br. Poult. Sci.* 56: 390–397.

Wang, W., Zhu, H., Dong, Y., Dong, T., Tian, Z. & Hu, H. 2019. Identification and dimorphic expression of sex-related genes during gonadal differentiation in sterlet *Acipenser ruthenus*, a primitive fish species. *Aquaculture* 500: 178–187.

Wang, W., Zhu, H., Dong, Y., Tian, Z., Dong, T., Hu, H., et al. 2017. Dimorphic expression of sex-related genes in different gonadal development stages of sterlet, *Acipenser ruthenus*, a primitive fish species. *Fish Physiol. Biochem.* 43: 1557–1569.

Webster, K.A., Schach, U., Ordaz, A., Steinfeld, J.S., Draper, B.W. & Siegfried, K.R. 2017. *Dmrt1* is necessary for male sexual development in zebrafish. *Dev. Biol.* 422: 33–46.

Wei, Y.D., Du, X.M., Yang, D.H., Ma, F.L., Yu, X.W., Zhang, M.F., et al. 2021. *Dmrt1* regulates the immune response by repressing the *Tlr4* signaling pathway in goat male germline stem cells. *Zool. Res.* 42: 14.

Wen, A.Y., You, F., Sun, P., Li, J., Xu, D.D., Wu, Z.H., et al. 2014. CpG methylation of *Dmrt1* and *Cyp19a* promoters in relation to their sexual dimorphic expression in the Japanese flounder *Paralichthys olivaceus*. *J. Fish. Biol.* 84: 193–205.

Wu, G.C., Chiu, P.C., Lin, C.J., Lyu, Y.S., Lan, D.S. & Chang, C.F. 2012. Testicular *Dmrt1* is involved in the sexual fate of the ootestis in the protandrous black porgy. *Biol. Reprod.* 86: 41–42.

Xia, W., Zhou, L., Yao, B., Li, C.J. & Gui, J.F. 2007. Differential and spermatogenic cell-specific expression of *Dmrt1* during sex reversal in protogynous hermaphroditic groupers. *Mol. Cell Endocrinol.* 263: 156–172.

Xiao, J., Cao, K., Zou, Y., Xiao, S., Wang, Z. & Cai, M. 2019. Sex-biased gene discovery from the gonadal transcriptomes of the large yellow croaker (*Larimichthys crocea*). *Aquac. Fish* 4: 9–16.

Xu, S., Zhang, S., Zhang, W., Liu, H., Wang, M., Zhong, L., et al. 2022. Genome-Wide Identification, phylogeny, and expression profile of the *Dmrt* (doublesex and mab-3 related transcription factor) gene family in channel catfish (*Ictalurus punctatus*). *Front. Genet.* 13: 891204.

Xu, W., Li, Y.Y., Lou, Q.Q., Chen, X.R., Qin, Z.F. & Wie, W.J. 2015. Low concentrations of dihydrotestosterone induce female-to-male sex reversal in the frog *Pelophylax nigromaculatus*. *Environ. Toxicol. Chem.* 34: 2370–2377.

Xue, L., Jia, D., Xu, L., Huang, Z., Fan, H., Chen, B., et al. 2021. Bulk and single-cell RNA-seq reveal the sexually dimorphic expression pattern of *Dmrtb1* in zig-zag eel (*Mastacembelus armatus*). *Aquaculture* 545: 737194.

Yan, H., Shen, X., Jiang, J., Zhang, L., Yuan, Z., Wu, Y., et al. 2021. Gene expression of *Takifugu rubripes* gonads during Al- or MT-induced masculinization and E2-induced feminization. *Endocrinology* 162: 1–20.

Yan, N., Hu, J., Li, J., Dong, J., Sun, C. & Ye, X. 2019. Genomic organization and sexually dimorphic expression of the *Dmrt1* gene in largemouth bass (*Micropterus salmoides*). *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 234: 68–77.

Yang, S., Tang, X., Yan, F., Yang, H., Xu, L., Jian, Z., et al. 2024. A time-course transcriptome analysis revealing the potential molecular mechanism of early gonadal differentiation in the Chinese giant salamander. *Comp. Biochem. Physiol. Part D Genomics Proteomics* 50: 101200.

Yoshimoto, S., Ikeda, N., Izutsu, Y., Shiba, T., Takamatsu, N. & Ito, M. 2010. Opposite roles of *Dmrt1* and its W-linked paralogue, DM-W, in sexual dimorphism of *Xenopus laevis*: implications of a ZZ/ZW-type sex-determining system. *Development* 137: 2519–2526.

Yoshimoto, S., Okada, E., Umamoto, H., Tamura, K., Uno, Y., Nishida-Umehara, C., et al. 2008. AW-linked DM-domain gene, DM-W, participates in primary ovary development in *Xenopus laevis*. *Proc. Natl. Acad. Sci. USA* 105: 2469–2474.

Zhang, X., Zha, J. & Wang, Z. 2008. Influences of 4-nonylphenol on doublesex- and mab-3-related transcription factor 1 gene expression and vitellogenin mRNA induction of adult rare minnow (*Gobiocypris rarus*). *Environ. Toxicol. Chem.* 27: 196–205.

Zhao, C., Bian, C., Mu, X., Zhang, X. & Shi, Q. 2024b. Gonadal transcriptome sequencing reveals sexual dimorphism in expression profiling of sex-related genes in Asian arowana (*Scleropages formosus*). *Front. Genet.* 15: 1381832.

Zhao, H., Xiao, Y., Xiao, Z., Wu, Y., Ma, Y. & Li, J. 2024a. Genome-wide investigation of the *Dmrt* gene family sheds new insight into the regulation of sex differentiation in spotted knifejaw (*Oplegnathus punctatus*) with fusion chromosomes (Y). *Int. J. Biol. Macromol.* 257: 128638.

Zhao, Y., Lu, H., Yu, H., Cheng, H. & Zhou, R. 2007. Multiple alternative splicing in gonads of chicken *Dmrt1*. *Dev. Genes Evol.* 217: 119–126.

Zhong, Z., Wang, Y., Feng, Y., Xu, Y., Zhao, L., Jiang, Y., et al. 2022. The molecular regulation mechanism of *Dmrt1*—based on the establishment of the testis cell line derived from two-spot puffer *Takifugu bimaculatus*. *Fish Physiol. Biochem.* 48: 1475–1494.

Zhu, Q., Qin, Z., Chen, R., Wang, L., Tan, P. & Xu, D. 2022. Comparative transcriptome analyses characterize expression signatures among males, females, neo-males, and gynogenetic females in the yellow drum (*Nibea albiflora*). *Front. Genet.* 13: 872815.

Zurita, F., Carmona, F.D., Lupiáñez, D.G., Barrionuevo, F.J., Guioli, S., Burgos, M., et al. 2006. Meiosis onset is postponed to postnatal stages during ootestis development in female moles. *Sex. Dev.* 1: 66–76. S.