

# Spin polarization of fermions at local equilibrium: Second-order gradient expansion

Xin-Li Sheng,<sup>1,\*</sup> Francesco Becattini,<sup>2,†</sup> Xu-Guang Huang,<sup>3,4,5,‡</sup> and Zhong-Hua Zhang<sup>3,§</sup>

<sup>1</sup>*INFN Sezione di Firenze, Florence, Italy*

<sup>2</sup>*Università di Firenze and INFN Sezione di Firenze, Florence, Italy*

<sup>3</sup>*Physics Department and Center for Field Theory and Particle Physics, Fudan University, Shanghai 200438, China*

<sup>4</sup>*Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Fudan University, Shanghai 200433, China*

<sup>5</sup>*Shanghai Research Center for Theoretical Nuclear Physics,  
National Natural Science Foundation of China and Fudan University, Shanghai 200438, China*

We present a calculation of the spin polarization of spin-1/2 fermions in a relativistic fluid at local thermodynamic equilibrium at the second order in the gradient expansion, including second-order derivatives. The second-order derivative terms vanish if the local equilibrium hypersurface is the hyperplane  $t = \text{const}$  in the collision center-of-mass frame. However, since the freeze-out hypersurface has a non-trivial space-time structure, these terms may result in a non-vanishing contribution to the spin polarization, whose magnitude needs to be assessed with numerical computations.

## I. INTRODUCTION

Spin polarization in a relativistic fluid is a subject of major phenomenological relevance in relativistic heavy-ion collisions, where the Quark Gluon Plasma (QGP) is formed; for recent reviews on spin polarization, see, e.g., [1–3].

Global and local polarization of  $\Lambda$  hyperons have been measured by the STAR collaboration at RHIC energies [4–6], and by the ALICE collaboration at the LHC energy [7, 8]. The hydrodynamic model with statistical hadronic freeze-out proved to be successful in reproducing the data for several observables. Global spin polarization in heavy ion collision at high energy was quantitatively and successfully predicted by this model [9–11] with the spin-thermal vorticity coupling [12], whereas its momentum dependence could not be reproduced [6, 13, 14]. This motivated further theoretical investigations on other possible sources of spin polarization in heavy-ion collisions, which led to the finding that also the (thermal) shear tensor and the gradient of chemical potentials contribute [15–17] at the linear order in the gradients of the hydro-thermodynamic fields (see also studies in refs. [18–22]). By including the spin polarization induced by the thermal shear tensor, the models can reproduce the experimental data [23, 24] with a noteworthy sensitivity on the initial hydrodynamic conditions as well as on the transport coefficients [19, 24].

In hydrodynamics and non-equilibrium statistical mechanics, any physical observable can be expressed in terms of the hydro-thermodynamic fields through a so-called gradient expansion [25, 26], which provides a good approximation when these fields are slowly varying in spacetime. More precisely, the expansion in gradients is appropriate when the Knudsen number  $K_n$ , that is the ratio between the interaction length and the typical scale of the inhomogeneity of the fields satisfies  $K_n \ll 1$ , the so-called hydrodynamic limit<sup>1</sup>. However, there are two kinds of terms in the general gradient expansion which are conceptually distinct: those stemming from the expansion of the local equilibrium part of the density operator, which are non-dissipative, and those stemming from the expansion of the dissipative part of the density operator, which are associated to the so-called transport coefficients [27], such as shear viscosity.

It is known that the gradient expansion of the local equilibrium density operator gives rise to terms proportional to thermal vorticity, thermal shear tensor and gradient of chemical potential for the spin polarization, which are first order in the gradient expansion [15]. Nevertheless, we do not have any knowledge on possible second order gradient terms in the same expansion. In this work, we aim at filling this gap by calculating the linear response of the spin polarization to the second order gradient of hydrodynamic quantities at local thermodynamic equilibrium. We will not delve into the possible dissipative terms (i.e. arising from the expansion in the dissipative part of the density operator) at the same order of expansion. In heavy-ion collisions, numerical simulations show that the components of thermal vorticity tensor  $\varpi_{\mu\nu}$  and thermal shear tensor  $\xi_{\mu\nu}$  are small enough [9, 10, 23, 28–33] to ensure the validity of the hydrodynamic limit, hence of the gradient expansion. However, some components, such as the  $\varpi_{tz}$  component as shown in ref. [10], could have significant variations over the freeze-out hypersurface, implying that second order gradient terms  $\partial_\rho \varpi_{\mu\nu}$  and  $\partial_\rho \xi_{\mu\nu}$  could provide a relevant contribution.

The paper is organized as follows: sec. II features a general discussion on the mean values of operators at local equilibrium. In sec. III, we derive the mean value of the Wigner function for spin-1/2 fermions. In sec. IV we

\* sheng@fi.infn.it

† becattini@fi.infn.it

‡ huangxuguang@fudan.edu.cn

§ zhzhang22@m.fudan.edu.cn

<sup>1</sup> The Knudsen number can be defined as  $K_n = \lambda/|\beta/\partial\beta|$  where  $\lambda$  is the interaction length and  $\beta$  the four-temperature field.




FIG. 1: Spacetime diagram of a relativistic heavy-ion collision. Here  $\Sigma_{\text{eq}}$  (black line) is the hypersurface where LTE is achieved and  $\Sigma_{\text{FO}}$  (blue line) is the hypersurface where the quark-gluon plasma decouples.

calculate the mean spin polarization vector up to second order in gradient expansion. Finally, section V presents the summary of this work and the related outlook.

### Notation

In this paper we use the natural units, with  $\hbar = c = K = 1$ . The Minkowskian metric tensor is  $\text{diag}(1, -1, -1, -1)$ ; for the Levi-Civita symbol we use the convention  $\epsilon^{0123} = 1$ . We will use the relativistic notation with repeated indices assumed to be saturated. Operators in Hilbert space will be denoted by a wide upper hat, e.g.  $\widehat{H}$ , except for the Dirac field operator  $\psi(x)$ .

## II. LOCAL THERMODYNAMIC EQUILIBRIUM AND ITS EXPANSION

For a relativistic fluid at Local Thermodynamic Equilibrium (LTE), the density operator is given by [34–36]

$$\widehat{\rho}_{\text{LE}} = \frac{1}{Z_{\text{LE}}} \exp \left[ - \int_{\Sigma} d\Sigma_{\mu} \left( \widehat{T}^{\mu\nu}(y) \beta_{\nu}(y) - \frac{1}{2} \widehat{S}^{\mu\nu\lambda}(y) \Omega_{\nu\lambda}(y) - \widehat{j}^{\mu}(y) \zeta(y) \right) \right], \quad (1)$$

where  $Z_{\text{LE}}$  is the normalization factor ensuring  $\text{Tr} \widehat{\rho}_{\text{LE}} = 1$ ,  $\beta_{\nu} = u_{\nu}/T$  is the four-temperature vector,  $\Omega_{\nu\lambda}$  is the spin potential,  $\zeta = \mu/T$  is the ratio between a chemical potential and temperature, and  $y$  is the point on the hypersurface  $\Sigma$ . For heavy ion collisions, we are mostly interested on calculations at the freeze-out, so the hypersurface is the  $\Sigma_{\text{FO}}$  featured in fig. 1. The formula (1) includes the possibility of a spin potential  $\Omega_{\nu\lambda}$  not coinciding with thermal vorticity:

$$\varpi_{\nu\lambda} = \frac{1}{2} (\partial_{\lambda} \beta_{\nu} - \partial_{\nu} \beta_{\lambda}),$$

which makes it certainly pseudo-gauge dependent [37]. Thus, the operator (1) pertains to a particular choice of the pseudo-gauge, for instance the so-called canonical one where  $\widehat{T}^{\mu\nu}$  and  $\widehat{S}^{\mu\nu\lambda}$  are obtained from the Lagrangian of the theory by applying the Nöther theorem.

When calculating the mean value of a local operator  $\widehat{O}(x)$  at LTE:

$$\langle \widehat{O}(x) \rangle_{\text{LE}} = \text{Tr} \left( \widehat{O}(x) \widehat{\rho}_{\text{LE}} \right)$$

in a relativistic fluid with slowly varying thermodynamic fields  $\beta, \zeta$  (small Knudsen number), it is expected that the above mean value will be mostly determined by the value of those fields at the point  $x$  while their values far from  $x$  will give rise to corrections. One can then expand the fields  $\beta_{\nu}(y)$ ,  $\Omega_{\nu\lambda}(y)$ , and  $\zeta(y)$  in Taylor series around the point

$x$  where the operator  $\widehat{O}$  is to be evaluated, thereby generating a gradient expansion. At the leading order, one can approximate:

$$\beta_\nu(y) - \beta_\nu(x) \simeq (y-x)^\lambda \partial_\lambda \beta_\nu(x), \quad \zeta(y) - \zeta(x) \simeq (y-x)^\lambda \partial_\lambda \zeta(x),$$

implying ( $\varpi = \Omega$  for the sake of simplicity) [36]:

$$\widehat{\rho}_{\text{LE}} \simeq \frac{1}{Z_{\text{LE}}} \exp \left[ -\beta_\nu(x) \widehat{P}^\nu + \frac{1}{2} \varpi_{\nu\lambda}(x) \widehat{J}_x^{\nu\lambda} - \frac{1}{2} \xi_{\nu\lambda}(x) \widehat{Q}_x^{\nu\lambda} + \zeta(x) \widehat{Q} + \partial_\nu \zeta(x) \widehat{D}_x^\nu \right], \quad (2)$$

where  $\xi$  is the thermal shear tensor:

$$\xi_{\nu\lambda} = \frac{1}{2} (\partial_\lambda \beta_\nu + \partial_\nu \beta_\lambda),$$

$\widehat{P}^\mu$  are the four-momentum operators,  $\widehat{J}_x^{\lambda\nu}$  are the angular momentum-boost operators with respect to the point  $x$ , and the operators  $\widehat{Q}_x$  and  $\widehat{D}_x$  read:

$$\widehat{Q}_x^{\lambda\nu} = \int_\Sigma d\Sigma_\mu(y) \left[ (y-x)^\lambda \widehat{T}^{\mu\nu} + (y-x)^\nu \widehat{T}^{\mu\lambda} \right], \quad \widehat{D}_x^\lambda = \int_\Sigma d\Sigma_\mu(y) (y-x)^\lambda \widehat{j}^\mu. \quad (3)$$

The operators  $\widehat{Q}_x$  and  $\widehat{D}_x$  are responsible for the shear-induced polarization [15, 16] and spin Hall effect [17] respectively and, unlike  $\widehat{P}$  and  $\widehat{J}$ , are dependent on the hypersurface  $\Sigma$ ; this is a crucial difference because the corrections stemming from  $\widehat{Q}_x$  and  $\widehat{D}_x$  will, in general, be dependent on the normal vector field to the hypersurface besides the thermo-hydrodynamic fields.

It is the purpose of this paper to go beyond the leading order Taylor expansion of the thermodynamic fields and to include the second derivatives of  $\beta$  and  $\zeta$  in the expansion (2), which will eventually be associated to integral operators of the same sort of  $\widehat{Q}_x$  and  $\widehat{D}_x$ . To maintain a general viewpoint, it is convenient to recast  $\widehat{\rho}_{\text{LE}}$  as:

$$\widehat{\rho}_{\text{LE}} = \frac{1}{Z_{\text{LE}}} \exp \left[ \widehat{A}_x + \widehat{B}_x \right], \quad (4)$$

where the operators  $\widehat{A}_x$  and  $\widehat{B}_x$  are defined as:

$$\begin{aligned} \widehat{A}_x &\equiv -\beta_\nu(x) \widehat{P}^\nu + \zeta(x) \widehat{Q}, \\ \widehat{B}_x &\equiv -\int_\Sigma d\Sigma_\mu \widehat{T}^{\mu\nu}(y) [\beta_\nu(y) - \beta_\nu(x)] + \frac{1}{2} \int_\Sigma d\Sigma_\mu \widehat{S}^{\mu\nu\lambda}(y) \Omega_{\nu\lambda}(y) \\ &\quad + \int_\Sigma d\Sigma_\mu \widehat{j}^\mu(y) [\zeta(y) - \zeta(x)], \end{aligned} \quad (5)$$

with  $\widehat{P}^\nu \equiv \int_\Sigma d\Sigma_\mu \widehat{T}^{\mu\nu}$  being the four-momentum operator and  $\widehat{Q} \equiv \int_\Sigma d\Sigma_\mu \widehat{j}^\mu$  being the total charge. Due to energy-momentum conservation and current conservation, the operator  $\widehat{A}_x$  is independent of the hypersurface  $\Sigma$  unlike  $\widehat{B}_x$ . In eq. (5) we have not extracted the spin potential at the point  $x$ . Indeed, since the spin potential coincides with the thermal vorticity at global equilibrium, we have considered the spin potential  $\Omega_{\nu\lambda}(y) \approx \Omega_{\nu\lambda}(x)$  as a first order term. In this power counting scheme, the contribution from  $\widehat{B}_x$  is at least  $\mathcal{O}(\partial)$  and therefore is assumed to be much smaller than the contribution from  $\widehat{A}_x$ , allowing us to expand the exponential in (4) up to the quadratic term in  $\widehat{B}_x$  using the Kubo identity,

$$\begin{aligned} e^{\widehat{A}_x + \widehat{B}_x} &\approx e^{\widehat{A}_x} + \int_0^1 dz e^{z\widehat{A}_x} \widehat{B}_x e^{-z\widehat{A}_x} \\ &\quad + \int_0^1 dz_1 \int_0^{z_1} dz_2 e^{z_1\widehat{A}_x} \widehat{B}_x e^{(z_1-z_2)\widehat{A}_x} \widehat{B}_x e^{-z_1\widehat{A}_x} + \mathcal{O}(\widehat{B}^3). \end{aligned}$$

For the local operator  $\widehat{O}(x)$ , the mean value at LTE is thus given by:

$$\begin{aligned} \langle \widehat{O}(x) \rangle_{\text{LE}} &\approx \langle \widehat{O}(0) \rangle_A + \int_0^1 dz \langle \widehat{O}(0), e^{z\widehat{A}_x} \widehat{B}_x e^{-z\widehat{A}_x} \rangle_{A,C} \\ &\quad + \int_0^1 dz_1 \int_0^{z_1} dz_2 \langle \widehat{O}(0), e^{z_2\widehat{A}_x} \widehat{B}_x e^{-z_2\widehat{A}_x}, e^{z_1\widehat{A}_x} \widehat{B}_x e^{-z_1\widehat{A}_x} \rangle_{A,C} + \mathcal{O}(\widehat{B}^3), \end{aligned} \quad (6)$$

where  $\langle \rangle_A$  denotes the mean value at the homogeneous global thermodynamic equilibrium described by  $\hat{A}$ :

$$\langle \hat{O} \rangle_A \equiv \frac{\text{Tr} [\hat{O} e^{\hat{A}_x}]}{\text{Tr} [e^{\hat{A}_x}]}.$$

In eq. (6) we have shifted the operator  $\hat{O}(x)$  by means of the identity  $\hat{O}(0) \equiv e^{-ix \cdot \hat{P}} \hat{O}(x) e^{ix \cdot \hat{P}}$  and  $\hat{B}_x$  likewise. The connected mean values are defined as:

$$\begin{aligned} \langle \hat{O}_1, \hat{O}_2 \rangle_{A,C} &\equiv \langle \hat{O}_1 \hat{O}_2 \rangle_A - \langle \hat{O}_1 \rangle_A \langle \hat{O}_2 \rangle_A, \\ \langle \hat{O}_1, \hat{O}_2, \hat{O}_3 \rangle_{A,C} &\equiv \langle \hat{O}_1 \hat{O}_2 \hat{O}_3 \rangle_A - \langle \hat{O}_1 \hat{O}_2 \rangle_A \langle \hat{O}_3 \rangle_A - \langle \hat{O}_2 \hat{O}_3 \rangle_A \langle \hat{O}_1 \rangle_A \\ &\quad - \langle \hat{O}_1 \hat{O}_3 \rangle_A \langle \hat{O}_2 \rangle_A + 2 \langle \hat{O}_1 \rangle_A \langle \hat{O}_2 \rangle_A \langle \hat{O}_3 \rangle_A, \end{aligned} \quad (7)$$

for arbitrary operators  $\hat{O}_{1,2,3}$ .

We now focus on free spin-1/2 fermions, for which the canonical energy-stress tensor, canonical spin tensor, and the current are given by [38],

$$\begin{aligned} \hat{T}^{\mu\nu}(x) &= \frac{i}{2} \bar{\psi}(x) \gamma^\nu (\overrightarrow{\partial}^\mu - \overleftarrow{\partial}^\mu) \psi(x), \\ \hat{S}^{\mu\nu\lambda}(x) &= -\frac{1}{2} \epsilon^{\mu\nu\lambda\rho} \bar{\psi}(x) \gamma_\rho \gamma^5 \psi(x), \\ \hat{j}^\mu(x) &= \bar{\psi}(x) \gamma^\mu \psi(x), \end{aligned} \quad (8)$$

where  $\psi(x)$  is the Dirac field. By substituting (8) into eq. (5) and taking the 4D Fourier transforms, it is possible to express  $\hat{B}_0$  as follows:

$$\hat{B}_0 = e^{-ix \cdot \hat{P}} \hat{B}_x e^{ix \cdot \hat{P}} = \frac{1}{(2\pi)^5} \int d^4 k_1 d^4 k_2 \bar{\psi}(k_2) \mathcal{B}(k_2, k_1) \psi(k_1), \quad (9)$$

where  $\psi(k)$  is the Fourier component of the Dirac field,

$$\psi(k) \equiv \int d^4 x \psi(x) e^{ik \cdot x}, \quad \bar{\psi}(k) \equiv \psi^\dagger(k) \gamma^0 = \int d^4 x \bar{\psi}(x) e^{-ik \cdot x}. \quad (10)$$

Since  $\hat{B}_0$  is Hermitian,  $\mathcal{B}(k_2, k_1)$  is a  $4 \times 4$  Dirac matrix fulfilling  $\gamma^0 \mathcal{B}^\dagger(k_2, k_1) \gamma^0 = \mathcal{B}(k_1, k_2)$ , whose explicit form reads:

$$\mathcal{B}(k_2, k_1) = -\frac{1}{(2\pi)^3} \int_\Sigma d\Sigma_\mu e^{-iq \cdot (y-x)} \left\{ \gamma^\mu k^\nu [\beta_\nu(y) - \beta_\nu(x)] + \frac{1}{4} \epsilon^{\mu\nu\lambda\sigma} \gamma_\sigma \gamma^5 \Omega_{\nu\lambda}(y) - \gamma^\mu [\zeta(y) - \zeta(x)] \right\}, \quad (11)$$

where  $k \equiv (k_1 + k_2)/2$  is the average momentum and  $q \equiv k_1 - k_2$  is the relative momentum, respectively. The fields  $\beta_\nu(y)$ ,  $\Omega_{\nu\lambda}(y)$ , and  $\zeta(y)$  can be expressed as Taylor series with respect to  $y - x$ . For instance:

$$\beta_\nu(y) = \beta_\nu(x) + \sum_{n=1}^{\infty} \frac{1}{n!} \left[ \prod_{j=1}^n (y^{\lambda_j} - x^{\lambda_j}) \partial_{\lambda_j} \right] \beta_\nu \Big|_{y=x},$$

where  $\partial_{\lambda_j}$  acts on  $\beta_\nu$ . Then:

$$\begin{aligned} e^{-iq \cdot (y-x)} [\beta_\nu(y) - \beta_\nu(x)] &= e^{-iq \cdot (y-x)} \sum_{n=1}^{\infty} \frac{1}{n!} \left[ \prod_{j=1}^n (y^{\lambda_j} - x^{\lambda_j}) \partial_{\lambda_j} \right] \beta_\nu \Big|_{y=x} \\ &= \sum_{n=1}^{\infty} \frac{1}{n!} \left[ \prod_{j=1}^n \partial_{\lambda_j} \right] \beta_\nu \Big|_{y=x} \left[ \prod_{j=1}^n i \frac{\partial}{\partial q_j} \right] e^{-iq \cdot (y-x)} \equiv [\beta_\nu(x + i\partial_q) - \beta_\nu(x)] \left( e^{-iq \cdot (y-x)} \right). \end{aligned}$$

The other terms in (11) involving  $\Omega$  and  $\zeta$  can be expressed likewise. Finally, all the functions  $\beta(x + i\partial_q)$  etc. can be moved out of the integral, so that eq. (11) turns into:

$$\mathcal{B}(k_2, k_1) = -\frac{1}{(2\pi)^3} \left\{ \gamma^\mu k^\nu [\beta_\nu(x + i\partial_q) - \beta_\nu(x)] + \frac{1}{4} \epsilon^{\mu\nu\lambda\sigma} \gamma_\sigma \gamma^5 \Omega_{\nu\lambda}(x + i\partial_q) - \gamma^\mu [\zeta(x + i\partial_q) - \zeta(x)] \right\} \int_\Sigma d\Sigma_\mu e^{-iq \cdot (y-x)}, \quad (12)$$

where it is understood that the term within the curl brackets acts on the integral over  $\Sigma$ . Now, with the help of the Gauss's theorem, that integral can be converted to an integral over a portion of a hyperplanes  $\Sigma_B$  with constant normal vector  $\hat{t}^\mu = (1, 0, 0, 0)$  and an integral over a 4D region  $\Omega_B$  enclosed by two space-like hyperplanes and a part of the hypersurface  $\Sigma$  (see fig. 2 in Appendix A). Assuming that the space extent of  $\Sigma_B$  and  $\Omega_B$  are sufficiently large, we obtain (see Appendix A for the derivation):

$$\begin{aligned} \int_\Sigma d\Sigma_\mu e^{-iq \cdot (y-x)} &\simeq (2\pi)^3 \left[ \hat{t}_\mu e^{-iq^0(y_B^0 - x^0)} - iq_\mu \int_{y_B^0}^{y_\Sigma^0(q)} dy^0 e^{-iq^0(y^0 - x^0)} \right] \delta^3(\mathbf{q}) \\ &= (2\pi)^3 e^{-iq^0(y_\Sigma^0(q) - x^0)} \delta^3(\mathbf{q}) \hat{t}_\mu, \end{aligned} \quad (13)$$

where  $y_B^0$  denotes the constant time for  $\Sigma_B$  and  $y_\Sigma^0(q)$  denotes a constant time, dependent on  $q$ , which can be considered as an average time for the space-like part of the hypersurface  $\Sigma$  (see again Appendix A for the precise definition). By plugging eq. (13) into eq. (12), we can finally express  $\mathcal{B}(k_2, k_1)$  as follows:

$$\begin{aligned} \mathcal{B}(k_2, k_1) &\simeq -\hat{t}_\mu \left\{ \gamma^\mu k^\nu [\beta_\nu(x + i\partial_q) - \beta_\nu(x)] + \frac{1}{4} \epsilon^{\mu\nu\lambda\rho} \gamma_\rho \gamma^5 \Omega_{\nu\lambda}(x + i\partial_q) \right. \\ &\quad \left. - \gamma^\mu [\zeta(x + i\partial_q) - \zeta(x)] \right\} \left[ e^{-iq^0(y_\Sigma^0(q) - x^0)} \delta^3(\mathbf{q}) \right], \end{aligned} \quad (14)$$

with the same proviso as for eq. (12).

### III. THE WIGNER FUNCTION AT LOCAL EQUILIBRIUM

The general formula to calculate the mean spin vector of spin 1/2 fermions reads [39]:

$$S^\mu(p) = \frac{1}{2} \frac{\int_\Sigma d\Sigma \cdot p_+ \text{tr}[\gamma^\mu \gamma^5 W(x, p)]}{\int_\Sigma d\Sigma \cdot p_+ \text{tr}[W(x, p)]}, \quad (15)$$

where  $\text{tr}$  stands for the trace over the spinorial indices and  $p_+^\mu \equiv p^\mu \theta(p^0)$  selects the positive-energy part of the Wigner function, removing antiparticle contribution. The covariant Wigner function is defined as [40, 41]:

$$W_{ab}(x, p) \equiv \frac{1}{(2\pi)^4} \int d^4y e^{-ip \cdot y} \left\langle \bar{\psi}_b \left( x + \frac{y}{2} \right) \psi_a \left( x - \frac{y}{2} \right) \right\rangle, \quad (16)$$

where  $a, b$  are spinorial Dirac indices and  $\langle (\dots) \rangle \equiv \text{Tr}[\hat{\rho}(\dots)]$  denotes the mean value with a specific density matrix. It should be stressed that, even though the argument  $p$  of the Wigner function is not necessarily on-shell, the formula (15), for a free field, makes  $p$  an on-shell vector and it holds for any choice of the hypersurface  $\Sigma$  [39].

The goal of this section is the calculation of the Wigner function at LTE by using the expansion in eq. (6). For this purpose, we first make use of the Fourier decomposition in eq. (10) to convert the definition (16) to the momentum space:

$$W_{ab}(x, p) = \frac{1}{(2\pi)^8} \int d^4k_1 d^4k_2 \delta^4 \left( p - \frac{k_1 + k_2}{2} \right) e^{-i(k_1 - k_2) \cdot x} \langle \bar{\psi}_b(k_2) \psi_a(k_1) \rangle, \quad (17)$$

then we apply eq. (6) by setting  $\hat{O} = \bar{\psi}_b(k_2) \psi_a(k_1)$ . The result can be in general written as a series in powers of  $\hat{B}$ :

$$W_{\text{LE}}(x, p) = W_0(x, p) + W_{\text{lin}}(x, p) + W_{\text{quad}}(x, p) + \mathcal{O}(\hat{B}^3). \quad (18)$$

We are now going to determine the form of the Wigner function up to the quadratic term.

### A. Main term

The  $\widehat{B}$ -independent part of the Wigner function is given by:

$$W_{0,ab}(x,p) = \frac{1}{(2\pi)^8} \int d^4q e^{-iq \cdot x} \langle \bar{\psi}_b(p_2) \psi_a(p_1) \rangle_A, \quad (19)$$

where

$$p \equiv (p_1 + p_2)/2, \quad q \equiv p_1 - p_2 \implies p_{1,2} \equiv p \pm q/2 \quad (20)$$

and  $\widehat{A}_x$  is given by eq. (5). Noting that the field operators in coordinate space obey the equal-time anticommutating relation  $\{\psi_a(t, \mathbf{x}), \psi_b^\dagger(t, \mathbf{x}')\} = i\delta^3(\mathbf{x} - \mathbf{x}')\delta_{ab}$ , one can readily derive the commutators  $[\psi_a(p), \widehat{P}^\mu] = p^\mu \psi_a(p)$  and  $[\psi_a(p), \widehat{Q}] = \psi_a(p)$ . It then follows that the expectation value in eq. (19) is given by

$$\langle \bar{\psi}_b(p_2) \psi_a(p_1) \rangle_A = n_F(x, p_1) \{\bar{\psi}_b(p_2), \psi_a(p_1)\}, \quad (21)$$

where  $n_F$  is the Fermi-Dirac distribution parameterized by local variables  $\beta^\mu(x)$  and  $\zeta(x)$ ,

$$n_F(x, p) \equiv \frac{1}{1 + \exp[\beta(x) \cdot p - \zeta(x)]}. \quad (22)$$

By quantizing the Dirac field in a standard way, one can show that the anticommutator in eq. (21) reads:

$$\{\bar{\psi}_b(p_2), \psi_a(p_1)\} = (2\pi)^5 \delta(p^2 - m^2) \text{sgn}(p^0) (\not{p} + m)_{ab} \delta^4(q), \quad (23)$$

where  $\text{sgn}$  is the sign function. Substituting eqs. (21) and (23) into eq. (19), we obtain

$$W_0(x, p) = \frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{(2\pi)^3} (\not{p} + m) n_F(x, p), \quad (24)$$

which is exactly the known Wigner function for free fermions [41–43] at global thermodynamic equilibrium. We note that  $p^0$  in eq. (24) could be either positive or negative, indicating that  $W_0$  contains contributions from both particles and antiparticles. We also find that  $W_0$  does not have axial-vector components, indicating that particles are not polarized at this order.

### B. Linear term

The linear term in the expansion of the LTE density operator reads:

$$\begin{aligned} W_{\text{lin},ab}(x,p) &= \frac{1}{(2\pi)^{13}} \int d^4k_1 d^4k_2 d^4k_3 d^4k_4 \delta^4\left(p - \frac{k_1 + k_2}{2}\right) \mathcal{B}_{cd}(k_4, k_3) \\ &\quad \times \int_0^1 dz e^{z\beta \cdot (k_3 - k_4)} \langle \bar{\psi}_b(k_2) \psi_a(k_1), \bar{\psi}_d(k_4) \psi_c(k_3) \rangle_{A,C}, \end{aligned} \quad (25)$$

where  $a, b, c, d$  are spinorial Dirac indices and the connected mean value is defined in eq. (7). One can check that:

$$\langle \bar{\psi}_b(k_2) \psi_a(k_1), \bar{\psi}_d(k_4) \psi_c(k_3) \rangle_{A,C} = n_F(x, k_2) [1 - n_F(x, k_1)] \{\bar{\psi}_b(k_2), \psi_c(k_3)\} \{\bar{\psi}_d(k_4), \psi_a(k_1)\}, \quad (26)$$

where the anticommutators are given by eq. (23). Substituting eqs. (23) and (26) into (25), we obtain:

$$\begin{aligned} W_{\text{lin}}(x,p) &= \frac{1}{(2\pi)^3} \int d^4q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \\ &\quad \times n_F(x, p_1) [1 - n_F(x, p_2)] \int_0^1 dz e^{z\beta \cdot q} (\not{p}_2 + m) \mathcal{B}(p_2, p_1) (\not{p}_1 + m), \end{aligned} \quad (27)$$

where  $p_1, p_2, p, q$  are like in eq. (20).

### C. Quadratic term

By applying eq. (6) to the Wigner function in eq. (17) and expressing  $\widehat{B}$  in terms of the Dirac field operators as shown in the eq. (9), one can derive the quadratic term in eq. (18):

$$\begin{aligned}
W_{\text{quad},ab}(x,p) &= \frac{1}{(2\pi)^{18}} \int d^4p_1 d^4p_2 d^4k_1 d^4k_2 d^4k_3 d^4k_4 \delta^4\left(p - \frac{p_1 + p_2}{2}\right) \\
&\times \int_0^1 dz_1 \int_0^{z_1} dz_2 e^{z_2\beta \cdot (k_1 - k_2) + z_1\beta \cdot (k_3 - k_4)} \mathcal{B}_{cd}(k_2, k_1) \mathcal{B}_{ef}(k_4, k_3) \\
&\times \langle \bar{\psi}_b(p_2) \psi_a(p_1), \bar{\psi}_c(k_2) \psi_d(k_1), \bar{\psi}_e(k_4) \psi_f(k_3) \rangle_{A,C}, \tag{28}
\end{aligned}$$

where the subscripts  $a, b, c, d, e, f$  are spinorial indices. The connected mean value turns out to be:

$$\begin{aligned}
&\langle \bar{\psi}_b(p_2) \psi_a(p_1), \bar{\psi}_c(k_2) \psi_d(k_1), \bar{\psi}_e(k_4) \psi_f(k_3) \rangle_{A,C} \\
&= (1 - n_F(x, p_1)) n_F(x, p_2) [(1 - n_F(x, k_1)) \{\bar{\psi}_b(p_2), \psi_f(k_3)\} \{\bar{\psi}_c(k_2), \psi_a(p_1)\} \{\bar{\psi}_e(k_4), \psi_d(k_1)\} \\
&\quad - n_F(x, k_2) \{\bar{\psi}_b(p_2), \psi_d(k_1)\} \{\bar{\psi}_c(k_2), \psi_f(k_3)\} \{\bar{\psi}_e(k_4), \psi_a(p_1)\}] ,
\end{aligned}$$

where the Fermi-Dirac distributions are defined in (22) and the anticommutators are given in eq. (23). After completing some momentum integrals, the equation (28) simplifies to the following form:

$$\begin{aligned}
W_{\text{quad}}(x,p) &= \frac{1}{(2\pi)^3} \int d^4q d^4q' \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \text{sgn}(p_3^0) n_F(p_1) [1 - n_F(p_3)] \\
&\times \int_0^1 dz_1 \int_0^{z_1} dz_2 \left\{ -e^{z_2\beta \cdot q + z_1\beta \cdot q'} n_F(p_2) + e^{z_2\beta \cdot q' + z_1\beta \cdot q} [1 - n_F(p_2)] \right\} \\
&\times (\not{p}_3 + m) \mathcal{B}(p_3, p_2) (\not{p}_2 + m) \mathcal{B}(p_2, p_1) (\not{p}_1 + m), \tag{29}
\end{aligned}$$

where we have defined  $p_1 \equiv p + (q + q')/2$ ,  $p_2 \equiv p + (-q + q')/2$ , and  $p_3 \equiv p - (q + q')/2$ .

### IV. SPIN POLARIZATION AT LOCAL EQUILIBRIUM

We can now use the expansion (18) in eq. (15) to determine the spin polarization vector. First, though, we will rearrange the terms of the expansion (18) in increasing order of gradients:

$$W_{\text{LE}}(x,p) = W^{(0)}(x,p) + W^{(1)}(x,p) + W^{(2)}(x,p) + \mathcal{O}(\partial^3). \tag{30}$$

where

$$W^{(0)} = W_0, \quad W^{(1)} = W_{\text{lin}}^{(1)}, \quad W^{(2)} = W_{\text{lin}}^{(2)} + W_{\text{quad}}, \tag{31}$$

and the superscripts denote their orders in the gradient expansion. By first order in the gradient expansion we mean terms including  $\partial_\mu \beta_\nu(x)$ ,  $\partial_\mu \zeta(x)$ , and  $\Omega_{\mu\nu}(x)$ , while for second order we mean second order derivatives such as  $\partial_\mu \partial_\lambda \beta_\nu(x)$ ,  $\partial_\mu \partial_\lambda \zeta(x)$ ,  $\partial_\lambda \Omega_{\mu\nu}(x)$  as well as quadratic terms in the first order derivatives such as  $[\partial_\mu \beta_\nu(x)][\partial_\lambda \beta_\sigma(x)]$ ,  $[\partial_\mu \zeta(x)][\partial_\nu \zeta(x)]$ . Hence, up to second order in gradient, the spin polarization is given by:

$$\begin{aligned}
S^{(0)\mu}(p) &= 0, \\
S^{(1)\mu}(p) &= \frac{\int d\Sigma \cdot p_+ \text{tr} [\gamma^\mu \gamma^5 W^{(1)}(x,p)]}{2 \int d\Sigma \cdot p_+ \text{tr} [W^{(0)}(x,p)]}, \\
S^{(2)\mu}(p) &= \frac{\int d\Sigma \cdot p_+ \text{tr} [\gamma^\mu \gamma^5 W^{(2)}(x,p)]}{2 \int d\Sigma \cdot p_+ \text{tr} [W^{(0)}(x,p)]} - \frac{\left\{ \int d\Sigma \cdot p_+ \text{tr} [\gamma^\mu \gamma^5 W^{(1)}(x,p)] \right\} \left\{ \int d\Sigma \cdot p_+ \text{tr} [W^{(1)}(x,p)] \right\}}{2 \left\{ \int d\Sigma \cdot p_+ \text{tr} [W^{(0)}(x,p)] \right\}^2}, \tag{32}
\end{aligned}$$

where the zeroth order part  $S^{(0)\mu}(p)$  vanishes because  $\text{tr}[\gamma^\mu \gamma^5 W^{(0)}] = 0$ . Also, the denominator in eq. (32) can be straightforwardly obtained from eq. (24):

$$\text{tr} [W_0(x,p)] = 4m \frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{(2\pi)^3} n_F(x,p). \tag{33}$$

### A. First order in the gradient expansion

At the first order in the gradient expansion, the Wigner function is obtained by taking the first-order term in the expansion of  $\mathcal{B}$  in eq. (14):

$$\mathcal{B}^{(1)}(p_2, p_1) = -\hat{t}_\mu \left\{ \gamma^\mu [p^\nu \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x)] i\partial_q^\lambda + \frac{1}{4} \epsilon^{\mu\nu\lambda\rho} \gamma_\rho \gamma^5 \Omega_{\nu\lambda}(x) \right\} \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \quad (34)$$

and plugging it into eq. (27). The result reads:

$$W^{(1)}(x, p) = \frac{1}{(2\pi)^3} \int d^4q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) [f_A^\lambda(p_2, p_1) i\partial_\lambda^q + f_B(p_2, p_1)] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right]. \quad (35)$$

where the auxiliary functions  $f_{A,B}$  are defined as

$$\begin{aligned} f_A^\lambda(p_2, p_1) &\equiv -\hat{t}_\mu \left\{ \partial^\lambda [p_\nu \beta^\nu(x) - \zeta(x)] \right\} (\not{p}_2 + m) \gamma^\mu (\not{p}_1 + m) n_F(x, p_1) [1 - n_F(x, p_2)] \int_0^1 dz e^{z\beta \cdot q}, \\ f_B(p_2, p_1) &\equiv -\frac{1}{4} \hat{t}_\mu \epsilon^{\mu\nu\lambda\sigma} \Omega_{\nu\lambda} (\not{p}_2 + m) \gamma_\sigma \gamma^5 (\not{p}_1 + m) n_F(x, p_1) [1 - n_F(x, p_2)] \int_0^1 dz e^{z\beta \cdot q}. \end{aligned}$$

and  $p, q, p_1, p_2$  are defined like in eq. (20). Noting that the product of the two delta functions can be rewritten as:

$$\delta(p_1^2 - m^2) \delta(p_2^2 - m^2) = \frac{1}{2|p^0|} \delta\left(p^2 + \frac{q^2}{4} - m^2\right) \delta\left(q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0}\right), \quad (36)$$

the integral in  $d^4q$  in eq. (35) can be carried out by parts, so to obtain:

$$W^{(1)}(x, p) = \frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} [D_\lambda^q f_A^\lambda(p_2, p_1) + f_B(p_2, p_1)], \quad (37)$$

where the operator  $D_\lambda^q$  is defined as:

$$D_\lambda^q \equiv -i\partial_\lambda^q + \frac{p_\lambda}{p^0} (i\partial_0^q + y_\Sigma^0(q) - x^0). \quad (38)$$

Somewhat lengthy but simple calculations involving the trace of multiple  $\gamma$  matrices give the scalar and axial-vector components of the Wigner function at this order:

$$\begin{aligned} \text{tr} [W^{(1)}(x, p)] &= -\frac{\delta(p^2 - m^2)}{(2\pi)^3 |p^0|} n_F(x, p) [1 - n_F(x, p)] (y_\Sigma^0 - x^0) 4mp^\lambda \partial_\lambda [p^\sigma \beta_\sigma(x) - \zeta(x)], \\ \text{tr} [\gamma^\mu \gamma^5 W^{(1)}(x, p)] &= -\frac{\delta(p^2 - m^2)}{(2\pi)^3 |p^0|} n_F(x, p) [1 - n_F(x, p)] \\ &\quad \times \left\{ 2\epsilon^{\mu\nu\rho\lambda} p_\nu \hat{t}_\rho \partial_\lambda [p^\tau \beta_\tau(x) - \zeta(x)] + (p^\mu p_\tau - g_\tau^\mu m^2) \hat{t}_\rho \epsilon^{\rho\nu\lambda\tau} \Omega_{\nu\lambda}(x) \right\}. \end{aligned} \quad (39)$$

Substituting the second equation of (39) and eq. (33) in eq. (32), the polarization at first order in gradient expansion can be obtained:

$$\begin{aligned} S^{(1)\mu}(p) &= -\frac{1}{8mN} \int d\Sigma \cdot p_+ n_F(x, p) [1 - n_F(x, p)] \\ &\quad \times \left\{ \epsilon^{\mu\nu\lambda\sigma} \Omega_{\nu\lambda} p_\sigma - \frac{2}{p \cdot \hat{t}} \hat{t}_\nu \epsilon^{\mu\nu\lambda\sigma} p_\lambda [(\xi_{\sigma\rho} + \Omega_{\sigma\rho} - \varpi_{\sigma\rho}) p^\rho - \partial_\sigma \zeta] \right\}, \end{aligned} \quad (40)$$

where

$$N \equiv \int d\Sigma \cdot p_+ n_F(x, p) \quad (41)$$

and  $\xi_{\sigma\rho} \equiv (\partial_\sigma \beta_\rho + \partial_\rho \beta_\sigma)/2$  is the thermal shear tensor. The first term reduces to the polarization induced by thermal vorticity when  $\Omega_{\nu\lambda} = \varpi_{\nu\lambda}$ , which is the major contribution to the  $\Lambda$  hyperons' global polarization. The contribution proportional to  $\xi_{\sigma\rho}$  is the shear-induced polarization that coincides with the result of ref. [15] and, essentially, with



that of ref. [44]<sup>2</sup>. The contribution from  $\partial_\sigma \zeta$  is the so-called spin Hall effect [17] and again, our result essentially coincides with the result reported in [44]. Finally, the contribution from the difference between spin potential and thermal vorticity coincides with the calculation in ref. [45] and, essentially, with that in ref. [44].

A key feature of eq. (40) is its dependence on the vector  $\hat{t}$  which is the direction of time in the centre-of-mass frame of the colliding nuclei (see fig. 1). As it was pointed out in ref. [15], the appearance of this vector stems from the essential dependence of the operator  $\widehat{Q}_x^{\lambda\nu}$  in eq. (3) on the hypersurface  $\Sigma$  (unlike  $\widehat{J}_x^{\lambda\nu}$ ); in this respect,  $\hat{t}$  can be considered as the average normal vector to the hypersurface  $\Sigma$ .

## B. Second order in the gradient expansion

At the second order in the gradient expansion, the Wigner function contains both linear and quadratic contributions, as expressed by eqs. (31) and (32). We first focus on  $W_{\text{lin}}^{(2)}$ , which is obtained by plugging the second order term of the expansion of eq. (14):

$$\mathcal{B}^{(2)}(p_2, p_1) = \hat{t}_\mu \left\{ \frac{1}{2} \gamma^\mu [p^\nu \partial_\lambda \partial_\sigma \beta_\nu(x) - \partial_\lambda \partial_\sigma \zeta(x)] \partial_q^\lambda \partial_q^\sigma + \frac{i}{4} \epsilon^{\mu\nu\lambda\rho} \gamma_\rho \gamma^5 [\partial_\sigma \Omega_{\nu\lambda}(x)] \partial_q^\sigma \right\} \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right]$$

into  $W_{\text{lin}}$  in eq. (27). The result reads (see Appendix B):

$$W_{\text{lin}}^{(2)}(x, p) = \frac{1}{(2\pi)^3} \int d^4q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \\ \times [f_C^{\lambda\sigma}(p_2, p_1) \partial_\lambda^q \partial_\sigma^q + f_D^\sigma(p_2, p_1) i \partial_\sigma^q] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right], \quad (42)$$

where the auxiliary functions  $f_{C,D}$  are defined as:

$$f_C^{\lambda\sigma}(p_2, p_1) \equiv \frac{1}{2} \hat{t}_\mu \{ \partial^\lambda \partial^\sigma [p_\nu \beta^\nu(x) - \zeta(x)] \} (\not{p}_2 + m) \gamma^\mu (\not{p}_1 + m) n_F(x, p_1) [1 - n_F(x, p_2)] \int_0^1 dz e^{z\beta \cdot q}, \\ f_D^\sigma(p_2, p_1) \equiv \frac{1}{4} \hat{t}_\mu \epsilon^{\mu\nu\lambda\rho} [\partial^\sigma \Omega_{\nu\lambda}(x)] (\not{p}_2 + m) \gamma_\rho \gamma^5 (\not{p}_1 + m) n_F(x, p_1) [1 - n_F(x, p_2)] \int_0^1 dz e^{z\beta \cdot q}. \quad (43)$$

Integrating by parts, we can carry out the integral over  $d^4q$  in eq. (42) and obtain:

$$W_{\text{lin}}^{(2)}(x, p) = -\frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} [D_\lambda^q D_\sigma^q f_C^{\lambda\sigma}(p_2, p_1) - D_\sigma^q f_D^\sigma(p_2, p_1)] \\ + \frac{\delta'(p^2 - m^2)}{(2\pi)^3 4|p^0|} \left[ g_{\lambda\sigma} - \frac{2p_\lambda}{p^0} g_{\sigma 0} + \frac{p_\lambda p_\sigma}{(p^0)^2} \right] \lim_{q^\mu \rightarrow 0} f_C^{\lambda\sigma}(p_2, p_1), \quad (44)$$

where the operator  $D^q$  is defined in eq. (38). We note that at the second order in gradient, the Wigner function features a term proportional to  $\delta'(p^2 - m^2)$ , including an off-shell contribution. It should be stressed that off-shell contributions must vanish once integrated over the hypersurface [39]. In this case, by using the explicit expression of  $f_C^{\lambda\sigma}$  in (43), one can show that the term proportional to  $\delta'(p^2 - m^2)$  only contributes to the scalar and vector components of the Wigner function, but not to the axial-vector one, which is relevant for the spin vector calculation.

Altogether, by using eq. (33) and eq. (44) in the last equation of eq. (32), after working out all traces one obtains:

$$S_{\text{lin}}^{(2)\mu}(p) = \frac{1}{4m(p \cdot \hat{t})^2 N} \int d\Sigma \cdot p_+ n_F(x, p) [1 - n_F(x, p)] (y_\Sigma(0) - x) \cdot \hat{t} \\ \times \hat{t}_\alpha p_\rho \left[ \epsilon^{\mu\sigma\alpha\rho} p^\lambda p^\nu \partial_\sigma \xi_{\nu\lambda} + \left( \frac{1}{2} p^\alpha \epsilon^{\mu\nu\lambda\rho} - \epsilon^{\mu\alpha\lambda\rho} p^\nu \right) p^\sigma \partial_\sigma \varpi_{\nu\lambda} \right. \\ \left. - \epsilon^{\mu\sigma\alpha\rho} p^\lambda \partial_\sigma \partial_\lambda \zeta + \frac{1}{2} \epsilon^{\alpha\nu\lambda\sigma} \partial^\rho (\Omega_{\nu\lambda} - \varpi_{\nu\lambda}) (p^\mu p_\sigma - m^2 g_\sigma^\mu) \right], \quad (45)$$

where  $N$  is shown in eq. (41). The most striking feature of eq. (45) is its explicit dependence on the average time  $y_\Sigma^0(0) = y_\Sigma(0) \cdot \hat{t}$  of the space-like part of the hypersurface  $\Sigma$ . This finding, surprising as it seems, is indeed in full

<sup>2</sup> In ref. [44] the normalization of the mean spin polarization vector slightly differs from the most commonly used.

agreement with the expectation that a local equilibrium term, which is not supposed to survive at global equilibrium, shows a dependence on the shape of the hypersurface, as it was discussed at the end of the previous subsection. It now appears that eq. (45) is a pure relativistic effect: indeed, if  $\Sigma$  is a flat 3D hypersurface,  $x^0 = y_\Sigma^0$  for any point  $x$  on  $\Sigma$ , resulting in a vanishing  $S_{\text{lin}}^{(2)\mu}$ . The feature that  $S_{\text{lin}}^{(2)\mu}(p)$  is proportional to  $y_\Sigma^0(0) - x^0$  can also be understood as a consequence of space and time reversal symmetry (P and T symmetry). Since  $S^\mu(p, x^0) \rightarrow -S^\mu(p, -x^0)$  (where, for clarity, the dependence on  $x^0$  is made explicit) under PT transformation, it must be constructed using PT-odd building blocks. At second order in gradients, these are  $\partial_\mu \partial_\nu \beta_\rho$ ,  $\partial_\mu \partial_\nu \zeta$ , and  $\partial_\mu \Omega_{\nu\rho}$  (which are all PT-even) multiplied by arbitrary powers of  $p$  and odd powers of  $y_\Sigma(0) - x$  (such combination is due to time translation symmetry). However, the power of  $y_\Sigma(0) - x$  must be smaller than the order in gradients, implying that only the first power of  $y_\Sigma(0) - x$  can appear in  $S_{\text{lin}}^{(2)\mu}(p)$ . Furthermore, the request of like transformations under P and T separately of both sides of equation (45), implies that only the time component  $y_\Sigma^0(0) - x^0$  can appear.

We now come to the spin polarization induced by quadratic terms in the gradient expansion. According to eq. (32), this reads:

$$S_{\text{quad}}^{(2)\mu}(p) = \frac{1}{2} \frac{\int d\Sigma \cdot p_+ \text{tr} [\gamma^\mu \gamma^5 W_{\text{quad}}(x, p)]}{\int d\Sigma \cdot p_+ \text{tr} [W^{(0)}(x, p)]} - S^{(1)\mu}(p) \frac{\int d\Sigma \cdot p_+ \text{tr} [W^{(1)}(x, p)]}{\int d\Sigma \cdot p_+ \text{tr} [W^{(0)}(x, p)]}, \quad (46)$$

where the second term arises because also the particle number is modified at the first order in the gradient expansion. The second term on the right hand side of eq. (46) which can be obtained from eqs. (33), (39), and (40) does not need further explanation. On the other hand, the function  $W_{\text{quad}}$  is obtained by plugging  $\mathcal{B}^{(1)}$  in eq. (34) into eq. (29). The expression of  $W_{\text{quad}}$  is too complicated to be listed here, however we have found a compact formula for the axial-vector component of  $W_{\text{quad}}$  as a function of already known quantities, whose derivation is reported in the Appendix C:

$$\text{tr} [\gamma^\mu \gamma^5 W_{\text{quad}}(x, p)] = \frac{[1 - 2n_F(x, p)] \text{tr} [\gamma^\mu \gamma^5 W^{(1)}(x, p)] \text{tr} [W^{(1)}(x, p)]}{[1 - n_F(x, p)] \text{tr} [W^{(0)}(x, p)]}. \quad (47)$$

Plugging the above expression into  $S_{\text{quad}}^{(2)\mu}$  in eq. (46), one can obtain the spin polarization induced by quadratic terms of first order quantities. Note that, according to eq. (39), if  $\Sigma$  is a flat 3D hypersurface with normal vector  $\hat{t}^\mu = (1, 0, 0, 0)$ ,  $\text{tr} [W^{(1)}(x, p)] = 0$  because of vanishing  $y_\Sigma^0(0) - x^0$  for any point  $x$  on  $\Sigma$ , leading to vanishing  $S_{\text{quad}}$ . Therefore, for such a flat hypersurface, the total spin polarization at second order, given by the sum  $S^{(2)\mu} = S_{\text{lin}}^{(2)\mu} + S_{\text{quad}}^{(2)\mu}$ , vanishes. This is in agreement with calculations at the global equilibrium, which are notably independent of the hypersurface, which show that there is no quadratic contribution in thermal vorticity for the spin polarization vector [46].

## V. SUMMARY

In summary, we have derived an expression of the average spin polarization vector of spin-1/2 fermions at local thermodynamic equilibrium up to the second order in the gradient expansion of hydro-thermodynamic fields, upgrading previous results at the leading order. We have calculated both the contributions from the quadratic term in the gradients and the linear in the second order derivatives.

The calculations in this work are based on the local thermodynamic equilibrium density operator and therefore have an expected dependence on the freeze-out hypersurface  $\Sigma$ . Under some geometric approximation, we find that the second order correction  $S^{(2)\mu}$  depends on the time coordinate  $x^0$  of the centre-of-mass frame for any point  $x$  on  $\Sigma$ . Notably, for the case of  $\Sigma$  coinciding with a hyperplane with a constant normal vector  $\hat{t}^\mu = (1, 0, 0, 0)$ , the term  $S^{(2)\mu}$  vanishes. Nevertheless, in relativistic heavy ion collision  $\Sigma$  is neither flat nor purely space-like, resulting in a non-vanishing  $S^{(2)\mu}$ .

In relativistic heavy ion collisions, while the quadratic term is expected to be a small correction to the leading order in view of the smallness of  $\varpi$  and  $\xi$  at the freeze-out, as shown by numerical simulations, the impact of the second order derivative term is not easy to estimate and requires numerical computations.

## ACKNOWLEDGMENTS

X.L.S. and F. B. are partly supported by ICSC - *Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing*, funded by European Union - NextGenerationEU and by the project PRIN2022 *Advanced Probes of the Quark Gluon Plasma* funded by “Ministero dell’Università e della Ricerca”. Z.-H.Z. and X.-G.H. are supported by the National Key Research and Development Program of China (Grant No. 2022YFA1604900),

the Natural Science Foundation of Shanghai (Grant No. 23JC1400200), and the National Natural Science Foundation of China (Grant No. 12147101, No. 12225502, and No. 12075061).

- 
- [1] X.-G. Huang, J. Liao, Q. Wang, and X.-L. Xia, *Lect. Notes Phys.* **987**, 281 (2021), 2010.08937.
- [2] J.-H. Gao, Z.-T. Liang, Q. Wang, and X.-N. Wang, *Lect. Notes Phys.* **987**, 195 (2021), 2009.04803.
- [3] F. Becattini, M. Buzzegoli, T. Niida, S. Pu, A.-H. Tang, and Q. Wang (2024), 2402.04540.
- [4] L. Adamczyk et al. (STAR), *Nature* **548**, 62 (2017), 1701.06657.
- [5] J. Adam et al. (STAR), *Phys. Rev. C* **98**, 014910 (2018), 1805.04400.
- [6] J. Adam et al. (STAR), *Phys. Rev. Lett.* **123**, 132301 (2019), 1905.11917.
- [7] S. Acharya et al. (ALICE), *Phys. Rev. C* **101**, 044611 (2020), [Erratum: *Phys.Rev.C* 105, 029902 (2022)], 1909.01281.
- [8] S. Acharya et al. (ALICE), *Phys. Rev. Lett.* **128**, 172005 (2022), 2107.11183.
- [9] F. Becattini, G. Inghirami, V. Rolando, A. Beraudo, L. Del Zanna, A. De Pace, M. Nardi, G. Pagliara, and V. Chandra, *Eur. Phys. J. C* **75**, 406 (2015), [Erratum: *Eur.Phys.J.C* 78, 354 (2018)], 1501.04468.
- [10] I. Karpenko and F. Becattini, *Eur. Phys. J. C* **77**, 213 (2017), 1610.04717.
- [11] B. Fu, K. Xu, X.-G. Huang, and H. Song, *Phys. Rev. C* **103**, 024903 (2021), 2011.03740.
- [12] F. Becattini, V. Chandra, L. Del Zanna, and E. Grossi, *Annals Phys.* **338**, 32 (2013), 1303.3431.
- [13] T. Niida (STAR), *Nucl. Phys. A* **982**, 511 (2019), 1808.10482.
- [14] F. Becattini and M. A. Lisa, *Ann. Rev. Nucl. Part. Sci.* **70**, 395 (2020), 2003.03640.
- [15] F. Becattini, M. Buzzegoli, and A. Palermo, *Phys. Lett. B* **820**, 136519 (2021), 2103.10917.
- [16] S. Y. F. Liu and Y. Yin, *JHEP* **07**, 188 (2021), 2103.09200.
- [17] S. Y. F. Liu and Y. Yin, *Phys. Rev. D* **104**, 054043 (2021), 2006.12421.
- [18] B. Fu, S. Y. F. Liu, L. Pang, H. Song, and Y. Yin, *Phys. Rev. Lett.* **127**, 142301 (2021), 2103.10403.
- [19] C. Yi, S. Pu, and D.-L. Yang, *Phys. Rev. C* **104**, 064901 (2021), 2106.00238.
- [20] X.-Y. Wu, C. Yi, G.-Y. Qin, and S. Pu, *Phys. Rev. C* **105**, 064909 (2022), 2204.02218.
- [21] S. Alzhrani, S. Ryu, and C. Shen, *Phys. Rev. C* **106**, 014905 (2022), 2203.15718.
- [22] N. Weickgenannt, D. Wagner, E. Speranza, and D. H. Rischke, *Phys. Rev. D* **106**, 096014 (2022), 2203.04766.
- [23] F. Becattini, M. Buzzegoli, G. Inghirami, I. Karpenko, and A. Palermo, *Phys. Rev. Lett.* **127**, 272302 (2021), 2103.14621.
- [24] A. Palermo, E. Grossi, I. Karpenko, and F. Becattini (2024), 2404.14295.
- [25] P. Kovtun, *J. Phys. A* **45**, 473001 (2012), 1205.5040.
- [26] P. Romatschke and U. Romatschke, *Relativistic Fluid Dynamics In and Out of Equilibrium*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2019), ISBN 978-1-108-48368-1, 978-1-108-75002-8, 1712.05815.
- [27] F. Becattini, M. Buzzegoli, and E. Grossi, *Particles* **2**, 197 (2019), 1902.01089.
- [28] O. Teryaev and R. Usubov, *Phys. Rev. C* **92**, 014906 (2015).
- [29] W.-T. Deng and X.-G. Huang, *Phys. Rev. C* **93**, 064907 (2016), 1603.06117.
- [30] Y. Jiang, Z.-W. Lin, and J. Liao, *Phys. Rev. C* **94**, 044910 (2016), [Erratum: *Phys.Rev.C* 95, 049904 (2017)], 1602.06580.
- [31] H. Li, L.-G. Pang, Q. Wang, and X.-L. Xia, *Phys. Rev. C* **96**, 054908 (2017), 1704.01507.
- [32] D.-X. Wei, W.-T. Deng, and X.-G. Huang, *Phys. Rev. C* **99**, 014905 (2019), 1810.00151.
- [33] X.-G. Deng, X.-G. Huang, Y.-G. Ma, and S. Zhang, *Phys. Rev. C* **101**, 064908 (2020), 2001.01371.
- [34] D. N. Zubarev, A. V. Prozorkevich, and S. A. Smolyanskii, *Theor. Math. Phys.* **40**, 821 (1979).
- [35] C. van Weert, *Annals of Physics* **140**, 133 (1982), ISSN 0003-4916, URL <https://www.sciencedirect.com/science/article/pii/0003491682903384>.
- [36] F. Becattini, L. Bucciattini, E. Grossi, and L. Tinti, *Eur. Phys. J. C* **75**, 191 (2015), 1403.6265.
- [37] F. Becattini, W. Florkowski, and E. Speranza, *Phys. Lett. B* **789**, 419 (2019), 1807.10994.
- [38] C. Itzykson and J. B. Zuber, *Quantum Field Theory*, International Series In Pure and Applied Physics (McGraw-Hill, New York, 1980), ISBN 978-0-486-44568-7.
- [39] F. Becattini, *Lect. Notes Phys.* **987**, 15 (2021), 2004.04050.
- [40] U. W. Heinz, *Phys. Rev. Lett.* **51**, 351 (1983).
- [41] D. Vasak, M. Gyulassy, and H. T. Elze, *Annals Phys.* **173**, 462 (1987).
- [42] R.-H. Fang, L.-G. Pang, Q. Wang, and X.-N. Wang, *Phys. Rev. C* **94**, 024904 (2016), 1604.04036.
- [43] N. Weickgenannt, X.-L. Sheng, E. Speranza, Q. Wang, and D. H. Rischke, *Phys. Rev. D* **100**, 056018 (2019), 1902.06513.
- [44] Y.-C. Liu and X.-G. Huang, *Sci. China Phys. Mech. Astron.* **65**, 272011 (2022), 2109.15301.
- [45] M. Buzzegoli, *Phys. Rev. C* **105**, 044907 (2022), 2109.12084.
- [46] A. Palermo and F. Becattini, *Eur. Phys. J. Plus* **138**, 547 (2023), 2304.02276.
- [47] V. Shtabovenko, R. Mertig, and F. Orellana (2023), 2312.14089.

### Appendix A: Integral over the hypersurface

The aim of this section is to calculate the integral over the freeze-out hypersurface:

$$\int_{\Sigma} d\Sigma_{\mu} e^{-iq \cdot (y-x)}. \quad (\text{A1})$$

The hypersurface  $\Sigma$  can be split into a space-like part  $\Sigma_S$  and a time-like part  $\Sigma_T$  (there is also a light-like part, but it is 2D surface with vanishing measure). It is convenient to introduce two auxiliary hypersurfaces  $\Sigma_B$  and  $\Sigma'$ , which are portions of hyperplanes with constant times  $y_B^0$  and  $y_{\Sigma}^0$  in the centre-of-mass frame, see fig. 2. The hypersurface  $\Sigma_B$  is defined as the portion of hyperplane joining the intersections between the initial local equilibrium hypersurface and the freeze-out, while  $\Sigma'$  is a portion of hyperplane, with  $\hat{t}$  as normal vector, which intersects the hypersurface  $\Sigma$  at some 2D boundary. Obviously, for any point  $y^{\mu}$  on  $\Sigma'$ , the time component  $y^0$  is a constant. The idea is that, in general, it is possible to approximate:

$$\int_{\Sigma'} d\Sigma_{\mu} e^{-iq \cdot (y-x)} \simeq \int_{\Sigma_S} d\Sigma_{\mu} e^{-iq \cdot (y-x)}. \quad (\text{A2})$$

and that  $y^0$  can be chosen on  $\Sigma'$  so as to make this approximation an equality for the time component of the equation (A2). Any point on  $\Sigma_S$  has a time component which is a function of the space Cartesian coordinates  $\mathbf{y}$ , that is  $y^0 = y^0(\mathbf{y})$ . Thus, the  $\mu = 0$  component of eq. (A2) can be rewritten as:

$$\int d^3\mathbf{y} e^{-iq^0(y_{\Sigma}^0 - x^0)} e^{i\mathbf{q} \cdot (\mathbf{y} - \mathbf{x})} = \int d^3\mathbf{y} e^{-iq^0(y^0(\mathbf{y}) - x^0)} e^{i\mathbf{q} \cdot (\mathbf{y} - \mathbf{x})} \quad (\text{A3})$$

with unknown  $y_{\Sigma}^0$ . In the equation (A3) it can be shown that, by using  $y^i$ ,  $i = 1, 2, 3$  as variables to describe the hypersurface  $\Sigma_S$ , it turns out that  $d\Sigma_0 = d^3\mathbf{y}$  without any other multiplicative factor. In general, the integration domains in  $\mathbf{y}$  can be different, yet this equation should have a solution for each  $q$ , that is  $y_{\Sigma}^0(q)$ . In the limit of  $q^{\mu} \rightarrow 0$ , and assuming that the domains of  $\mathbf{y}$  are approximately the same, one can rewrite eq. (A3) as:

$$\int d^3\mathbf{y} \left( e^{-iq^0 y_{\Sigma}^0} - e^{-iq^0 y^0(\mathbf{y})} \right) \simeq q^0 \int d^3\mathbf{y} (y_{\Sigma}^0 - y^0(\mathbf{y})) = 0$$

giving as average time for  $\Sigma_S$ :

$$y_{\Sigma}^0(0) \simeq \frac{\int_{\Sigma_S} d\Sigma y^0(\mathbf{y})}{\int_{\Sigma_S} d\Sigma}.$$

By using eq. (A2), the integral in eq. (A1) can be now cast in the following form:

$$\int_{\Sigma} d\Sigma_{\mu} e^{-iq \cdot (y-x)} \simeq \int_{\Sigma' \cup \Sigma_L} d\Sigma_{\mu} e^{-iq \cdot (y-x)},$$

where  $\Sigma_L$  is the part of  $\Sigma$  joining  $\Sigma'$  and  $\Sigma_B$  (which is approximate to  $\Sigma_T$ , see fig. 2). Using the Gauss' theorem, one can convert the above 3D integration to a 3D integration over  $\Sigma_B$  plus a 4D integral over the region encompassed by  $\Sigma'$ ,  $\Sigma_L$ , and  $\Sigma_B$ . Furthermore, if the hypersurface  $\Sigma_B$  is large enough, we can approximate the integrals with their infinite space extension limits and obtain:

$$\begin{aligned} \int_{\Sigma} d\Sigma_{\mu} e^{-iq \cdot (y-x)} &\simeq \int_{\Sigma' \cup \Sigma_L} d\Sigma_{\mu} e^{-iq \cdot (y-x)} \simeq \hat{t}_{\mu} e^{-iq^0(y_B^0 - x^0)} \int d^3\mathbf{y} e^{i\mathbf{q} \cdot (\mathbf{y} - \mathbf{x})} + \int_{y_B^0}^{y_{\Sigma}^0(q)} dy^0 \int d^3\mathbf{y} \frac{\partial}{\partial y^{\mu}} e^{-iq \cdot (y-x)} \\ &\simeq (2\pi)^3 \left[ \hat{t}_{\mu} e^{-iq^0(y_B^0 - x^0)} - iq_{\mu} \int_{y_B^0}^{y_{\Sigma}^0(q)} dy^0 e^{-iq^0(y^0 - x^0)} \right] \delta^3(\mathbf{q}) = (2\pi)^3 e^{-iq^0(y_{\Sigma}^0 - x^0)} \delta^3(\mathbf{q}) \hat{t}_{\mu}, \end{aligned}$$

which is eq. (13) in the main text.

### Appendix B: Calculation of $W_{\text{lin}}^{(2)}$

In this section we show some of most important steps of the evaluation of  $W_{\text{lin}}^{(2)}$  in eq. (42). We first focus on the contribution from  $f_D^{\sigma}$ , which is given by

$$I_D = \frac{1}{(2\pi)^3} \int d^4q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) f_D^{\sigma}(p_2, p_1) i \partial_{\sigma}^q \left[ e^{-iq^0(y_{\Sigma}^0 - x^0)} \delta^3(\mathbf{q}) \right].$$




FIG. 2: Spacetime diagram of a relativistic heavy-ion collision. Here  $\Sigma_{\text{eq}}$  (black line) is the hypersurface where LTE is achieved and  $\Sigma_{\text{FO}}$  is the hypersurface where the quark-gluon plasma decouples, which consists of the space-like part  $\Sigma_S$  (blue line) and the time-like parts  $\Sigma_T$  (magenta lines). For calculating the integral over  $\Sigma_{\text{FO}}$ , we introduce two flat hypersurfaces  $\Sigma_B$  and  $\Sigma'$  (red dashed lines) with constant times  $y_B^0$  and  $y_\Sigma^0$ , respectively.

Integrating by parts, the above formula can be turned into:

$$I_D = -\frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \left\{ [i\partial_\sigma^q f_D^\sigma(p_2, p_1)] \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) + f_D^\sigma(p_2, p_1) i\partial_\sigma^q [\delta(p_1^2 - m^2) \delta(p_2^2 - m^2)] \right\}. \quad (\text{B1})$$

In this transformation, we have neglected derivatives of the sign functions, because they yield delta functions of an energy equation that in general cannot be fulfilled due to the on-shell conditions enforced by other delta functions. Rewriting the product of two delta functions as in eq. (36), we can calculate their momentum derivative as follows:

$$\begin{aligned} i\partial_\sigma^q [\delta(p_1^2 - m^2) \delta(p_2^2 - m^2)] &= i \frac{q_\sigma}{4|p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) + \frac{ip_\sigma}{2|p^0|p^0} \delta \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta' \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \\ &= \frac{p_\sigma}{p^0} i\partial_0^q \left[ \frac{1}{2|p^0|} \delta \left( p^2 + \frac{q^2}{2} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] + i \frac{q_\sigma}{4|p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \\ &\quad - \frac{ip_\sigma q^0}{4|p^0|p^0} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right). \end{aligned}$$

When plugging the above formula back into  $I_D$ , we find that  $\delta(q^0 - \mathbf{p} \cdot \mathbf{q}/p^0)$  and  $\delta(\mathbf{q})$  involve the restriction  $q^\mu = 0$ , implying that the last two terms on the right-hand side do not contribute to  $I_D$ . Therefore, the (B1) becomes:

$$I_D = -\frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \left\{ [i\partial_\sigma^q f_D^\sigma(p_2, p_1)] \frac{1}{2|p^0|} \delta \left( p^2 + \frac{q^2}{2} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) + f_D^\sigma(p_2, p_1) \frac{p_\sigma}{p^0} i\partial_0^q \left[ \frac{1}{2|p^0|} \delta \left( p^2 + \frac{q^2}{2} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] \right\}. \quad (\text{B2})$$

For the derivative with respect to  $q^0$  in the (B2), we integrate by parts again:

$$I_D = -\frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \frac{1}{2|p^0|} \delta \left( p^2 + \frac{q^2}{2} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \times \left[ i\partial_\sigma^q - \frac{p_\sigma}{p^0} (i\partial_0^q + y_\Sigma^0 - x^0) \right] f_D^\sigma(p_2, p_1).$$

to finally get:

$$I_D = \frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} D_\sigma^q f_D^\sigma(p_2, p_1), \quad (\text{B3})$$

where the operator  $D_\sigma^q$  is defined in eq. (38).

We now turn to the contribution from  $f_C^{\lambda\sigma}$ , which is given by:

$$I_C = \frac{1}{(2\pi)^3} \int d^4q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) f_C^{\lambda\sigma}(p_2, p_1) \partial_\lambda^q \partial_\sigma^q \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right].$$

Similarly for the foregoing case, the above integral can be carried out by parts to obtain:

$$I_C = \frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \left\{ [\partial_\lambda^q \partial_\sigma^q f_C^{\lambda\sigma}(p_2, p_1)] \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \right. \\ \left. + 2 [\partial_\lambda^q f_C^{\lambda\sigma}(p_2, p_1)] [\partial_\sigma^q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2)] + f_C^{\lambda\sigma}(p_2, p_1) [\partial_\lambda^q \partial_\sigma^q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2)] \right\},$$

where the momentum derivatives of sign functions have been neglected for the same reason as for the previous case, and the symmetry  $f_C^{\lambda\sigma} = f_C^{\sigma\lambda}$  has been used. The first and second terms on the right-hand-side of the above formula can be simplified in a similar way as for the previous calculation of  $I_D$ . Then  $I_C$  is converted to:

$$I_C = \frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} \left[ \partial_\lambda^q \partial_\sigma^q f_C^{\lambda\sigma}(p_2, p_1) + 2i \frac{p_\sigma}{p^0} (i\partial_0^q + y_\Sigma^0 - x^0) \partial_\lambda^q f_C^{\lambda\sigma}(p_2, p_1) \right] \\ + \frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) f_C^{\lambda\sigma}(p_2, p_1) [\partial_\lambda^q \partial_\sigma^q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2)]. \quad (\text{B4})$$

For the last term on the right hand side, we have:

$$\partial_\lambda^q \partial_\sigma^q \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) = \frac{q_\lambda q_\sigma}{8|p^0|} \delta'' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) + \frac{g_{\lambda\sigma}}{4|p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \\ + \frac{p_\lambda p_\sigma}{2(p^0)^2 |p^0|} \delta \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta'' \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) + \frac{q_\lambda p_\sigma + p_\lambda q_\sigma}{4p^0 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta' \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right). \quad (\text{B5})$$

In turn, the third and last terms on the right hand side of eq. (B5) can be worked out as follows:

$$\frac{p_\lambda p_\sigma}{2(p^0)^2 |p^0|} \delta \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta'' \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) = \partial_q^0 \partial_q^0 \left[ \frac{p_\lambda p_\sigma}{2(p^0)^2 |p^0|} \delta \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] \\ - \partial_q^0 \left[ \frac{q^0 p_\lambda p_\sigma}{2(p^0)^2 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] + \frac{(q^0)^2 p_\lambda p_\sigma}{8(p^0)^2 |p^0|} \delta'' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \\ + \frac{p_\lambda p_\sigma}{4(p^0)^2 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right), \quad (\text{B6})$$

and

$$\frac{q_\lambda p_\sigma + p_\lambda q_\sigma}{4p^0 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta' \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) = \partial_q^0 \left[ \frac{q_\lambda p_\sigma + p_\lambda q_\sigma}{4p^0 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] \\ - \frac{g_{\lambda 0} p_\sigma + p_\lambda g_{\sigma 0}}{4p^0 |p^0|} \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) - \frac{q^0 (q_\lambda p_\sigma + p_\lambda q_\sigma)}{8p^0 |p^0|} \delta'' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \quad (\text{B7})$$

Substituting eqs. (B5)-(B7) into eq. (B4) and using the fact that  $\delta(q^0 - \mathbf{p} \cdot \mathbf{q}/p^0)$  and  $\delta^3(\mathbf{q})$  restricts  $q^\mu = 0$ , we arrive at:

$$I_C = \frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} \left[ \partial_\lambda^q \partial_\sigma^q f_C^{\lambda\sigma}(p_2, p_1) + 2i \frac{p_\sigma}{p^0} (i\partial_0^q + y_\Sigma^0 - x^0) \partial_\lambda^q f_C^{\lambda\sigma}(p_2, p_1) \right] \\ + \frac{1}{(2\pi)^3} \int d^4q e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) f_C^{\lambda\sigma}(p_2, p_1) \left\{ \partial_q^0 \partial_q^0 \left[ \frac{p_\lambda p_\sigma}{2(p^0)^2 |p^0|} \delta \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right] \right. \\ \left. + \frac{1}{4|p^0|} \left[ g_{\lambda\sigma} - \frac{g_{\lambda 0} p_\sigma + p_\lambda g_{\sigma 0}}{p^0} + \frac{p_\lambda p_\sigma}{(p^0)^2} \right] \delta' \left( p^2 + \frac{q^2}{4} - m^2 \right) \delta \left( q^0 - \frac{\mathbf{p} \cdot \mathbf{q}}{p^0} \right) \right\}.$$

Finally, by integrating by parts again and carrying out the integral over  $d^4q$ , we obtain

$$I_C = \frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} \left[ \partial_\lambda^q \partial_\sigma^q + 2i \frac{p_\sigma}{p^0} (i\partial_0^q + y_\Sigma^0 - x^0) \partial_\lambda^q - \frac{p_\lambda p_\sigma}{(p^0)^2} (i\partial_0^q + y_\Sigma^0 - x^0)^2 \right] f_C^{\lambda\sigma}(p_2, p_1) \\ + \frac{\delta'(p^2 - m^2)}{(2\pi)^3 4|p^0|} \lim_{q^\mu \rightarrow 0} \left[ g_{\lambda\sigma} - \frac{g_{\lambda 0} p_\sigma + p_\lambda g_{\sigma 0}}{p^0} + \frac{p_\lambda p_\sigma}{(p^0)^2} \right] f_C^{\lambda\sigma}(p_2, p_1).$$

which can be expressed in a compact form using the operator  $D_\sigma^q$  in eq. (39),

$$I_C = -\frac{\delta(p^2 - m^2)}{(2\pi)^3 2|p^0|} \lim_{q^\mu \rightarrow 0} D_\lambda^q D_\sigma^q f_C^{\lambda\sigma}(p_2, p_1) + \frac{\delta'(p^2 - m^2)}{(2\pi)^3 4|p^0|} \lim_{q^\mu \rightarrow 0} \left[ g_{\lambda\sigma} - \frac{2p_\lambda}{p^0} g_{\sigma 0} + \frac{p_\lambda p_\sigma}{(p^0)^2} \right] f_C^{\lambda\sigma}(p_2, p_1). \quad (\text{B8})$$

The Wigner function  $W_{\text{lin}}^{(2)}$  in (44) is then obtained by summing eqs. (B8) and (B3).

### Appendix C: Calculation of $W_{\text{quad}}$

To make notations more compact, we first define:

$$\begin{aligned} \Gamma_{1,\lambda}(p) &\equiv \hat{t}_\mu \gamma^\mu [p^\nu \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x)], \\ \Gamma_2 &\equiv \frac{1}{4} \hat{t}_\mu \epsilon^{\mu\nu\lambda\sigma} \gamma_\sigma \gamma^5 \Omega_{\nu\lambda}(x), \end{aligned}$$

where the dependence of  $\Gamma_{1,\lambda}$  and  $\Gamma_2$  on  $x$  is understood; also,  $\Gamma_{1,\lambda}$  depends on  $p$  whereas  $\Gamma_2$  is independent of  $p$ . Thereby, the first order part  $\mathcal{B}^{(1)}$  in eq. (34) can be rewritten as:

$$\mathcal{B}^{(1)}(p_2, p_1) = -[\Gamma_{1,\lambda}(p) i\partial_q^\lambda + \Gamma_2] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right].$$

Substituting it into  $W_{\text{quad}}$  in eq. (29), one obtains:

$$\begin{aligned} W_{\text{quad}}(x, p) &= \int dQ (\not{p}_3 + m) \left\{ [\Gamma_{1,\lambda}(p - q/2) i\partial_{q'}^\lambda + \Gamma_2] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right] \right\} \\ &\quad \times (\not{p}_2 + m) \left\{ [\Gamma_{1,\sigma}(p + q'/2) i\partial_q^\sigma + \Gamma_2] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \right\} (\not{p}_1 + m), \end{aligned} \quad (\text{C1})$$

where  $p_1 \equiv p + (q + q')/2$ ,  $p_2 \equiv p + (-q + q')/2$ ,  $p_3 \equiv p - (q + q')/2$ , and

$$\begin{aligned} dQ &\equiv \frac{1}{2(2\pi)^3} d^4 q d^4 q' \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \text{sgn}(p_3^0) \\ &\quad \times n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] \left[ 1 + \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot (q - q') \right]. \end{aligned}$$

The  $dQ$  in the above equation has been obtained through the Taylor expansion in  $q^\mu$  or  $q'^\mu$  of the following expression:

$$\begin{aligned} &n_F(p_1) [1 - n_F(p_3)] \int_0^1 dz_1 \int_0^{z_1} dz_2 \left\{ -e^{z_2 \beta \cdot q + z_1 \beta \cdot q'} n_F(p_2) + e^{z_2 \beta \cdot q' + z_1 \beta \cdot q} [1 - n_F(p_2)] \right\} \\ &\simeq \frac{1}{2} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] \left[ 1 + \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot (q - q') + \mathcal{O}(q^2, q'^2, q \cdot q') \right], \end{aligned} \quad (\text{C2})$$

where  $\mathcal{O}(q^2, q'^2, q \cdot q')$  terms can be dropped because they will not contribute to later calculations. In eq. (C1),  $\Gamma_{1,\lambda}$  is multiplied with a momentum derivative acting on  $\delta^3(\mathbf{q})$  or  $\delta^3(\mathbf{q}')$ , while  $\Gamma_2$  does not. The momentum derivative gives rise to complicated expressions when trying to integrate out the delta functions, thus we separate  $W_{\text{quad}}$  into three parts:

$$W_{\text{quad}} = W_{\text{quad},11} + W_{\text{quad},12} + W_{\text{quad},22},$$

with:

$$\begin{aligned} W_{\text{quad},11}(x, p) &= \int dQ (\not{p}_3 + m) \left\{ [\Gamma_{1,\lambda}(p - q/2) i\partial_{q'}^\lambda] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right] \right\} \\ &\quad \times (\not{p}_2 + m) \left\{ [\Gamma_{1,\sigma}(p + q'/2) i\partial_q^\sigma] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \right\} (\not{p}_1 + m), \\ W_{\text{quad},12}(x, p) &= \int dQ (\not{p}_3 + m) \left\{ [\Gamma_{1,\lambda}(p - q/2) i\partial_{q'}^\lambda] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right] \right\} (\not{p}_2 + m) \Gamma_2 (\not{p}_1 + m) \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \\ &\quad + \int dQ (\not{p}_3 + m) \Gamma_2 (\not{p}_2 + m) \left\{ [\Gamma_{1,\sigma}(p + q'/2) i\partial_q^\sigma] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \right\} (\not{p}_1 + m) \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right], \\ W_{\text{quad},22}(x, p) &= \int dQ (\not{p}_3 + m) \Gamma_2 (\not{p}_2 + m) \Gamma_2 (\not{p}_1 + m) \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}') \right] \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right]. \end{aligned}$$

It is hard to obtain explicit formulae for the various  $W_{\text{quad}}$ , especially  $W_{\text{quad},11}$ . Notwithstanding, calculating the axial-vector component is relatively easy with the help of FeynCalc [47].

We first focus on the first contribution to the  $W_{\text{quad}}$ . Working out the trace of Dirac matrices, we find:

$$\begin{aligned} \text{tr} [\gamma^\mu \gamma^5 W_{\text{quad},11}(x, p)] &= 8p^0 \epsilon^{\mu\nu\alpha\rho} p_\nu \int dQ \left\{ \hat{t}_\rho(q_\alpha + q'_\alpha) - \frac{1}{2p^0} [q_\alpha q'_\rho + \hat{t} \cdot (q - q')(q_\alpha + q'_\alpha) \hat{t}_\rho] \right\} \\ &\quad \times \left[ (p^{\nu'} - q^{\nu'}/2) \partial_{\lambda'} \beta_{\nu'}(x) - \partial_{\lambda'} \zeta(x) \right] \left[ (p^\nu + q^\nu/2) \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x) \right] \\ &\quad \times \left\{ \partial_q^\lambda \partial_{q'}^{\lambda'} \sin[(q^0 + q'^0)(y_\Sigma^0 - x^0)] \delta^3(\mathbf{q}') \delta^3(\mathbf{q}) \right\}, \end{aligned} \quad (\text{C3})$$

taking advantage of the fact that (C3) is a real function. Integrating by parts, we can rewrite it as:

$$\text{tr} [\gamma^\mu \gamma^5 W_{\text{quad},11}(x, p)] = \frac{4p^0}{(2\pi)^3} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] \epsilon^{\mu\nu\alpha\rho} p_\nu (I_{\alpha\rho}^{(1)} + I_{\alpha\rho}^{(2)} + I_{\alpha\rho}^{(3)} + I_{\alpha\rho}^{(4)}), \quad (\text{C4})$$

where:

$$\begin{aligned} I_{\alpha\rho}^{(1)} &\equiv \int d^4q d^4q' g_1(p; q, q') g_2(p; q, q') \left[ \partial_q^\lambda \partial_{q'}^{\lambda'} g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \right], \\ I_{\alpha\rho}^{(2)} &\equiv \int d^4q d^4q' g_1(p; q, q') \left[ \partial_q^\lambda \partial_{q'}^{\lambda'} g_2(p; q, q') \right] g_{3,\alpha\rho\lambda\lambda'}(p; q, q'), \\ I_{\alpha\rho}^{(3)} &\equiv \int d^4q d^4q' g_1(p; q, q') \left[ \partial_q^\lambda g_2(p; q, q') \right] \left[ \partial_{q'}^{\lambda'} g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \right], \\ I_{\alpha\rho}^{(4)} &\equiv \int d^4q d^4q' g_1(p; q, q') \left[ \partial_{q'}^{\lambda'} g_2(p; q, q') \right] \left[ \partial_q^\lambda g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \right], \end{aligned}$$

with auxiliary functions

$$\begin{aligned} g_1(p; q, q') &\equiv \sin[(q^0 + q'^0)(y_\Sigma^0 - x^0)] \delta^3(\mathbf{q}') \delta^3(\mathbf{q}) \text{sgn}(p_1^0) \text{sgn}(p_2^0) \text{sgn}(p_3^0), \\ g_2(p; q, q') &\equiv \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2), \\ g_{3,\alpha\rho\lambda\lambda'}(p; q, q') &\equiv \left[ 1 + \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot (q - q') \right] \\ &\quad \times \left\{ \hat{t}_\rho(q_\alpha + q'_\alpha) - \frac{1}{2p^0} [q_\alpha q'_\rho + \hat{t} \cdot (q - q')(q_\alpha + q'_\alpha) \hat{t}_\rho] \right\} \\ &\quad \times \left[ (p^{\tau'} - q^{\tau'}/2) \partial_{\lambda'} \beta_{\tau'}(x) - \partial_{\lambda'} \zeta(x) \right] \left[ (p^\tau + q^\tau/2) \partial_\lambda \beta_\tau(x) - \partial_\lambda \zeta(x) \right]. \end{aligned}$$

We note that we have neglected momentum derivatives of the sign functions because they will give terms like  $\delta(p_1^0)$ , a constraint that in general cannot be fulfilled.

In the term  $I_{\alpha\rho}^{(1)}$ , the delta functions in  $g_1$  and  $g_2$  yield:

$$\delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2) \delta^3(\mathbf{q}) \delta^3(\mathbf{q}') = \frac{\delta(p^2 - m^2)}{4(p^0)^2} \delta^4(q) \delta^4(q'), \quad (\text{C5})$$

forcing  $q^\mu$  and  $q'^\mu$  to be vanishing vectors. Therefore  $I_{\alpha\rho}^{(1)}$  vanishes because:

$$I_{\alpha\rho}^{(1)} \propto \lim_{q^\mu, q'^\mu \rightarrow 0} \sin[(q^0 + q'^0)(y_\Sigma^0 - x^0)] = 0.$$

In the term  $I_{\alpha\rho}^{(2)}$ , the momentum derivatives act on  $g_2$ , which can be expressed as:

$$g_2(p; q, q') = \frac{1}{4} \delta \left( p^2 - m^2 + \frac{q^2 + q'^2}{4} + \frac{q \cdot q'}{2} \right) \delta \left[ q \cdot \left( p + \frac{q'}{2} \right) \right] \delta \left[ q' \cdot \left( p - \frac{q}{2} \right) \right].$$

If both  $\partial_q^\lambda$  and  $\partial_{q'}^{\lambda'}$  act on  $\delta \left( p^2 - m^2 + \frac{q^2 + q'^2}{4} + \frac{q \cdot q'}{2} \right)$ , the remaining two delta functions in  $g_2$ , with the delta functions in  $g_1$ , will lead to  $\delta^4(q) \delta^4(q')$  and thus  $g_1 = 0$  because:

$$\lim_{q^\mu, q'^\mu \rightarrow 0} \sin[(q^0 + q'^0)(y_\Sigma^0 - x^0)] = 0.$$



If  $\partial_q^\lambda$  or  $\partial_{q'}^{\lambda'}$  (or both of them) acts on  $\delta \left[ q \cdot \left( p + \frac{q'}{2} \right) \right]$  or  $\delta \left[ q' \cdot \left( p - \frac{q}{2} \right) \right]$ , one can convert it to a derivative with respect to the time component  $q^0$  or  $q'^0$  and then integrate by parts. The only term that survives in our calculation reads:

$$I_{\alpha\rho}^{(2)} = \frac{1}{4} \int d^4q d^4q' g_1(p; q, q') \left( p^\lambda + \frac{q'^\lambda}{2} \right) \left( p^{\lambda'} - \frac{q'^{\lambda'}}{2} \right) g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \\ \times \delta \left( p^2 - m^2 + \frac{q^2 + q'^2}{4} + \frac{q \cdot q'}{2} \right) \delta' \left[ q \cdot \left( p + \frac{q'}{2} \right) \right] \delta' \left[ q' \cdot \left( p - \frac{q}{2} \right) \right].$$

Integrating again by parts, we find:

$$I_{\alpha\rho}^{(2)} \propto \lim_{q, q' \rightarrow 0} \frac{\partial}{\partial q^0} \frac{\partial}{\partial q'^0} \sin [(q^0 + q'^0)(y_\Sigma^0 - x^0)] \hat{t}_\rho(q_\alpha + q'_\alpha) \propto \hat{t}_\alpha \hat{t}_\rho,$$

which is symmetric with respect to its indices  $\alpha, \rho$ .

The calculation of  $I_{\alpha\rho}^{(3)}$  is similar to that of  $I_{\alpha\rho}^{(2)}$ , and the only term that survive reads:

$$I_{\alpha\rho}^{(3)} = \frac{1}{4} p^\lambda \int d^4q d^4q' g_1(p; q, q') \delta \left( p^2 - m^2 + \frac{q^2 + q'^2}{4} + \frac{q \cdot q'}{2} \right) \\ \times \delta' \left[ q \cdot \left( p + \frac{q'}{2} \right) \right] \delta \left[ q' \cdot \left( p - \frac{q}{2} \right) \right] \left[ \partial_{q'}^{\lambda'} g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \right].$$

Then,  $\delta' \left[ q \cdot \left( p + \frac{q'}{2} \right) \right]$  gives rise to a derivative with respect to  $q^0$ , and after integrating by parts, it turns out:

$$I_{\alpha\rho}^{(3)} = -\frac{1}{4} p^\lambda \lim_{q^\mu, q'^\mu \rightarrow 0} \frac{\partial}{\partial q^0} \sin [(q^0 + q'^0)(y_\Sigma^0 - x^0)] \text{sgn}(p_1^0) \text{sgn}(p_2^0) \text{sgn}(p_3^0) \\ \times \frac{\delta \left[ p^2 - m^2 + \frac{q^2 + q'^2}{4} + \frac{q \cdot q'}{2} \right]}{\left( p^0 + \frac{q'^0}{2} \right) \left| p^0 - \frac{q^0}{2} \right| \left| p^0 + \frac{q'^0}{2} \right|} \left[ \partial_{q'}^{\lambda'} g_{3,\alpha\rho\lambda\lambda'}(p; q, q') \right] \\ = -\frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{4(p^0)^3} (y_\Sigma^0 - x^0) \hat{t}_\rho g_\alpha^{\lambda'} \left[ p^\tau \partial_{\lambda'} \beta_{\tau'}(x) - \partial_{\lambda'} \zeta(x) \right] p^\lambda \left[ p^\tau \partial_\lambda \beta_\tau(x) - \partial_\lambda \zeta(x) \right].$$

Finally, the term  $I_{\alpha\rho}^{(4)}$  yields an expression which turns out to be the same as that of  $I_{\alpha\rho}^{(3)}$  because of the symmetry in the exchange of  $q^\mu$  with  $q'^\mu$ . Substituting  $I_{\alpha\rho}^{(i)}$  with  $i = 1, 2, 3, 4$  back into eq. (C4), we obtain:

$$\text{tr} \left[ \gamma^\mu \gamma^5 W_{\text{quad},11}(x, p) \right] = -\frac{2\delta(p^2 - m^2) \text{sgn}(p^0)}{(2\pi)^3 (p^0)^2} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] (y_\Sigma^0 - x^0) \\ \times \epsilon^{\mu\nu\alpha\rho} p_\nu \hat{t}_\rho g_\alpha^{\lambda'} \left[ p^\tau \partial_\lambda \beta_\tau(x) - \partial_\lambda \zeta(x) \right] p^{\lambda'} \left[ p^{\tau'} \partial_{\lambda'} \beta_{\tau'}(x) - \partial_{\lambda'} \zeta(x) \right].$$

Now we can proceed to the calculation of  $W_{\text{quad},12}$ . First, we rewrite the delta functions to the following form:

$$\delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2) \delta^3(\mathbf{q}) = \frac{1}{4 \left| p^0 + \frac{q'^0}{2} \right|} \delta \left[ p^2 + \frac{(q')^2}{4} - m^2 \right] \delta(p \cdot q') \delta^{(4)}(q), \\ \delta(p_1^2 - m^2) \delta(p_2^2 - m^2) \delta(p_3^2 - m^2) \delta^3(\mathbf{q}') = \frac{1}{4 \left| p^0 - \frac{q^0}{2} \right|} \delta \left[ p^2 + \frac{q^2}{4} - m^2 \right] \delta(p \cdot q) \delta^{(4)}(q').$$

Plugging them into  $W_{\text{quad},12}$  and integrating out  $q^\mu$  or  $q'^\mu$ :

$$\begin{aligned}
W_{\text{quad},12}(x,p) &= \frac{1}{32(2\pi)^3} \int d^4q \delta \left[ p^2 + \frac{q^2}{4} - m^2 \right] \delta(p \cdot q) n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] \\
&\quad \times \text{sgn} \left( p^0 - \frac{q^0}{2} \right) \text{sgn} \left( p^0 + \frac{q^0}{2} \right) [p^\nu \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x)] [\gamma \cdot (p - q/2) + m] \\
&\quad \times \left\{ \frac{1}{p^0 + \frac{q^0}{2}} \left[ 1 - \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot q \right] (\hat{t} \cdot \gamma) [\gamma \cdot (p + q/2) + m] \gamma_\sigma \gamma^5 \right. \\
&\quad \left. + \frac{1}{p^0 - \frac{q^0}{2}} \left[ 1 + \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot q \right] \gamma_\sigma \gamma^5 [\gamma \cdot (p - q/2) + m] (\hat{t} \cdot \gamma) \right\} \\
&\quad \times [\gamma \cdot (p + q/2) + m] \hat{t}_\tau \epsilon^{\tau\nu\alpha\sigma} \Omega_{\nu\alpha}(x) \left\{ i \partial_q^\lambda \left[ e^{-iq^0(y_\Sigma^0 - x^0)} \delta^3(\mathbf{q}) \right] \right\}.
\end{aligned}$$

Integrating by parts, we can rewrite it as follows:

$$\begin{aligned}
W_{\text{quad},12}(x,p) &= \frac{\delta(p^2 - m^2)}{32(2\pi)^3 |p^0|} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] [p^\nu \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x)] \hat{t}_\tau \epsilon^{\tau\nu\alpha\sigma} \Omega_{\nu\alpha}(x) \\
&\quad \times \lim_{q \rightarrow 0} D_q^\lambda [\gamma \cdot (p - q/2) + m] \left\{ \frac{1}{p^0 + \frac{q^0}{2}} \left[ 1 - \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot q \right] \hat{t} \cdot \gamma [\gamma \cdot (p + q/2) + m] \gamma_\sigma \gamma^5 \right. \\
&\quad \left. + \frac{1}{p^0 - \frac{q^0}{2}} \left[ 1 + \frac{1 - 4e^{\beta \cdot p} + e^{2\beta \cdot p}}{6(e^{2\beta \cdot p} - 1)} \beta \cdot q \right] \gamma_\sigma \gamma^5 [\gamma \cdot (p - q/2) + m] (\hat{t} \cdot \gamma) \right\} [\gamma \cdot (p + q/2) + m],
\end{aligned}$$

where the operator  $D_q^\lambda$  is given in eq. (38). The axial-vector component is given by:

$$\begin{aligned}
\text{tr} [\gamma^\mu \gamma^5 W_{\text{quad},12}(x,p)] &= - \frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{(2\pi)^3 (p^0)^2} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] (y_\Sigma^0 - x^0) \\
&\quad \times p^\lambda [p^\nu \partial_\lambda \beta_\nu(x) - \partial_\lambda \zeta(x)] (g_\sigma^\mu m^2 - p^\mu p_\sigma) \hat{t}_\tau \epsilon^{\tau\nu\alpha\sigma} \Omega_{\nu\alpha}(x).
\end{aligned}$$

The calculation of  $W_{\text{quad},22}$  is quite straightforward by using the relation (C5). Integrating out the delta functions for  $q^\mu$  and  $q'^\mu$ , we arrive at:

$$\begin{aligned}
W_{\text{quad},22}(x,p) &= \frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{8(2\pi)^3 (p^0)^2} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] \\
&\quad \times (\not{p} + m) \Gamma_2(\not{p} + m) \Gamma_2(\not{p} + m),
\end{aligned}$$

which has a vanishing axial-vector component:

$$\text{tr} [\gamma^\mu \gamma^5 W_{\text{quad},22}(x,p)] = 0.$$

Hence, the whole quadratic contribution to axial-vector component of the Wigner function at the second order reads:

$$\begin{aligned}
\text{tr} [\gamma^\mu \gamma^5 W_{\text{quad}}(x,p)] &= \frac{\delta(p^2 - m^2) \text{sgn}(p^0)}{(2\pi)^3 (p^0)^2} n_F(p) [1 - n_F(p)] [1 - 2n_F(p)] (y_\Sigma^0 - x^0) \\
&\quad \times p^{\lambda'} \partial_{\lambda'} [p^{\tau'} \beta_{\tau'}(x) - \zeta(x)] \\
&\quad \times \{ 2\epsilon^{\mu\nu\rho\lambda} p_\nu \hat{t}_\rho \partial_\lambda [p^\tau \beta_\tau(x) - \zeta(x)] + (p^\mu p_\tau - g_\tau^\mu m^2) \hat{t}_\rho \epsilon^{\rho\nu\lambda\tau} \Omega_{\nu\lambda}(x) \}. \tag{C6}
\end{aligned}$$

Comparing eq. (C6) with eqs. (24) and (39) one can finally prove eq. (47).