

# TOPOLOGICAL COMPLEXITY OF IDEAL LIMIT POINTS

MAREK BALCERZAK, SZYMON GŁĄB, AND PAOLO LEONETTI

ABSTRACT. Given an ideal  $\mathcal{I}$  on the nonnegative integers  $\omega$  and a Polish space  $X$ , let  $\mathcal{L}(\mathcal{I})$  be the family of subsets  $S \subseteq X$  such that  $S$  is the set of  $\mathcal{I}$ -limit points of some sequence taking values in  $X$ . First, we show that  $\mathcal{L}(\mathcal{I})$  may attain arbitrarily large Borel complexity. Second, we prove that if  $\mathcal{I}$  is a  $G_{\delta\sigma}$ -ideal then all elements of  $\mathcal{L}(\mathcal{I})$  are closed. Third, we show that if  $\mathcal{I}$  is a simply coanalytic ideal and  $X$  is first countable, then every element of  $\mathcal{L}(\mathcal{I})$  is simply analytic. Lastly, we studied certain structural properties and the topological complexity of minimal ideals  $\mathcal{I}$  for which  $\mathcal{L}(\mathcal{I})$  contains a given set.

## 1. INTRODUCTION

Let  $\mathcal{I}$  be an ideal on the nonnegative integers  $\omega$ , that is, a subset of  $\mathcal{P}(\omega)$  closed under taking subsets and finite unions. Unless otherwise stated, it is assumed that  $\mathcal{I}$  is admissible, namely,  $\omega \notin \mathcal{I}$  and that  $\mathcal{I}$  contains the family  $\text{Fin}$  of finite subsets of  $\omega$ . Intuitively, the ideal  $\mathcal{I}$  represents the family of “small” subsets of  $\omega$ . An important example is the family of asymptotic density zero sets

$$\mathcal{Z} := \left\{ S \subseteq \omega : \lim_{n \rightarrow \infty} \frac{|S \cap [0, n]|}{n} = 0 \right\}.$$

Define  $\mathcal{I}^+ := \mathcal{P}(\omega) \setminus \mathcal{I}$ . Ideals are regarded as subsets of the Cantor space  $\{0, 1\}^\omega$ , hence we can speak about their topological complexity. For instance,  $\text{Fin}$  is a  $F_\sigma$ -ideal, and  $\mathcal{Z}$  is a  $F_{\sigma\delta}$ -ideal which is not  $F_\sigma$ .

Pick also a sequence  $\mathbf{x} = (x_n)$  be taking values in a topological space  $X$ . Then, we denote by  $\Lambda_{\mathbf{x}}(\mathcal{I})$  the set of  $\mathcal{I}$ -limit points of  $\mathbf{x}$ , that is, the set of all  $\eta \in X$  for which there exists a subsequence  $(x_{n_k})$  such that

$$\lim_{k \rightarrow \infty} x_{n_k} = \eta \quad \text{and} \quad \{n_k : k \in \omega\} \in \mathcal{I}^+.$$

---

2010 *Mathematics Subject Classification*. Primary: 40A35, 54H05. Secondary: 11B05, 28A05, 54A20.

*Key words and phrases*. Ideal limit points; Borel and analytic ideals; simply analytic set; Fubini product; pruned trees; Wadge reduction.

It is well known that, even in the case where  $\mathbf{x}$  is a real bounded sequence, it is possible that  $\Lambda_{\mathbf{x}}(\mathcal{Z})$  is the empty set, see [13, Example 4]. The topological nature of the sets of  $\mathcal{I}$ -limits points  $\Lambda_{\mathbf{x}}(\mathcal{I})$  and their relationship with the slightly weaker variant of  $\mathcal{I}$ -cluster points have been studied in [2], cf. also [16, 17].

In this work, we continue along this line of research. To this aim, we introduce our main definition:

**Definition 1.1.** Let  $X$  be a topological space and  $\mathcal{I}$  be an ideal on  $\omega$ . We denote by  $\mathcal{L}_X(\mathcal{I})$  the family of sets of  $\mathcal{I}$ -limit points of sequences  $\mathbf{x}$  taking values in  $X$  together with the emptyset, that is,

$$\mathcal{L}_X(\mathcal{I}) := \{A \subseteq X : A = \Lambda_{\mathbf{x}}(\mathcal{I}) \text{ for some sequence } \mathbf{x} \in X^\omega\} \cup \{\emptyset\}.$$

If the topological space  $X$  is understood, we write simply  $\mathcal{L}(\mathcal{I})$ .

A remark is in order about the the addition of  $\{\emptyset\}$  in the above definition: it has been proved by Meza-Alcántara in [28, Section 2.7] that, if  $X = [0, 1]$ , then there exists a  $[0, 1]$ -valued sequence  $\mathbf{x}$  such that  $\Lambda_{\mathbf{x}}(\mathcal{I}) = \emptyset$  if and only if there exists a function  $\phi : \omega \rightarrow \mathbb{Q} \cap [0, 1]$  such that  $\phi^{-1}[A] \in \mathcal{I}$  for every set  $A \subseteq \mathbb{Q} \cap [0, 1]$  with at most finitely many limit points; cf. also [3, Proposition 6.4] and, more generally, [10] for analogues in compact uncountable spaces. On the other hand, if  $X$  is not compact, it is easy to see that, for every ideal  $\mathcal{I}$ , there is a sequence with no  $\mathcal{I}$ -limit points. Thus, the addition of  $\{\emptyset\}$  in the above definition avoids the repetition of known results in the literature and to add further subcases based on the topological structure of the underlying space  $X$ .

We summarize in Theorem 1.2 and Theorem 1.3 below the known results from [2, 16, 17] about the families  $\mathcal{L}_X(\mathcal{I})$ .

For, given a topological space  $X$  and an ordinal  $1 \leq \alpha < \omega_1$ , we use the standard Borel pointclasses notations  $\Sigma_\alpha^0(X)$  and  $\Pi_\alpha^0(X)$ , so that  $\Sigma_1^0(X)$  stands for the open sets of  $X$ ,  $\Pi_1^0(X)$  for the closed sets,  $\Sigma_2^0(X)$  for the  $F_\sigma$ -sets, etc.; we denote by  $\Delta_\alpha^0(X) := \Sigma_\alpha^0(X) \cap \Pi_\alpha^0(X)$  the ambiguous classes; also, if  $X$  is a Polish space,  $\Sigma_1^1(X)$  stands for the analytic sets,  $\Pi_1^1(X)$  for the coanalytic sets, etc., see e.g. [25, Section 11.B] or [31, Section 3.6]. Again, we suppress the reference to the underlying space  $X$  if it is clear from the context.

Recall that an ideal  $\mathcal{I}$  is a *P-ideal* if it is  $\sigma$ -directed modulo finite sets, that is, for every sequence  $(A_n)$  with values in  $\mathcal{I}$  there exists  $A \in \mathcal{I}$  such that  $A_n \setminus A$  is finite for all  $n \in \omega$ . Important examples include the  $\Sigma_1^1$  P-ideals, which are known to be necessarily  $\Pi_3^0$ . A topological space  $X$  is *discrete* if it contains only isolated points.

**Theorem 1.2.** *Let  $X$  be a nondiscrete first countable Hausdorff space. Then:*

- (i)  $\mathcal{L}(\mathcal{I}) \subseteq \Pi_1^0$  if and only if  $\mathcal{I}$  is a  $\Sigma_2^0$  ideal, provided that  $\mathcal{I}$  is a  $\Sigma_1^1$   $P$ -ideal.
- (ii)  $\mathcal{L}(\mathcal{I}) \subseteq \Sigma_2^0$ , provided that  $\mathcal{I}$  is a  $\Sigma_1^1$   $P$ -ideal.

*If, in addition, all closed subsets of  $X$  are separable, then:*

- (iii)  $\mathcal{L}(\mathcal{I}) = \Pi_1^0$ , provided that  $\mathcal{I}$  is a  $\Sigma_2^0$  ideal.
- (iv)  $\mathcal{L}(\mathcal{I}) = \Sigma_2^0$ , provided that  $\mathcal{I}$  is  $\Sigma_1^1$   $P$ -ideal which is not  $\Sigma_2^0$ .

*Proof.* It follows by [2, Theorems 2.2, 2.5, 2.7, and 3.4]. □

To state the next result, we recall some further definitions. An ideal  $\mathcal{I}$  has the *hereditary Baire property* if the restriction  $\mathcal{I} \upharpoonright A := \{S \cap A : S \in \mathcal{I}\}$  has the Baire property for every  $A \in \mathcal{I}^+$ . Note that all analytic ideals have the hereditary Baire property: indeed, the proof goes verbatim as in [24, Theorem 3.13], considering that analytic sets are closed under continuous preimages, and that they have the Baire property, see e.g. [31, Proposition 4.1.2 and Theorem 4.3.2]. In addition, there exist ideals with the Baire property but without the hereditary Baire property, see e.g. [32, Proposition 2.1].

Also, an ideal  $\mathcal{I}$  on  $\omega$  is said to be a  $P^+$ -ideal if, for every decreasing sequence  $(A_n)$  with values in  $\mathcal{I}^+$ , there exists  $A \in \mathcal{I}^+$  such that  $A \setminus A_n$  is finite for all  $n \in \omega$ . It is known that all  $F_\sigma$ -ideals are  $P^+$ -ideals, see [23] and [22, Observation 2.2]. We remark that  $P^+$ -ideal may have arbitrarily high Borel complexity, as it has been proved in [21, p. 2031] and [22, Example 2.6], cf. also [14].

Moreover, an ideal  $\mathcal{I}$  is called a *Farah ideal* if there exists a sequence  $(K_n)$  of hereditary compact subsets in  $\mathcal{P}(\omega)$  such that  $S \in \mathcal{I}$  if and only if for all  $n \in \omega$  there exists  $k \in \omega$  such that  $S \setminus [0, k] \in K_n$ , see [8, 19, 20]. It is known that all analytic  $P$ -ideals are Farah and that all Farah ideals are  $F_{\sigma\delta}$ . On the other hand, it is still unknown whether the converse holds, namely, all  $F_{\sigma\delta}$  ideals are Farah ideals, see [7] and [19, p. 60].

**Theorem 1.3.** *Let  $X$  be a first countable Hausdorff space. Then:*

- (i)  $\mathcal{L}(\mathcal{I}) \subseteq \Pi_1^0$ , provided that  $\mathcal{I}$  is  $P^+$ -ideal.
- (ii)  $\mathcal{L}(\mathcal{I}) \subseteq \Pi_1^0$  if and only if  $\mathcal{I}$  is a  $P^+$ -ideal, provided that  $\mathcal{I}$  has the hereditary Baire property and  $X$  is nondiscrete metrizable.
- (iii)  $\mathcal{L}(\mathcal{I}) \subseteq \Sigma_2^0$ , provided that  $\mathcal{I}$  is a Farah ideal.
- (iv)  $\mathcal{L}(\mathcal{I}) = \{\{\eta\} : \eta \in X\}$  if and only if  $\mathcal{I}$  is maximal, provided that  $|X| \geq 2$ .

*If, in addition,  $X$  is second countable, then:*

- (v)  $\Pi_1^0 \subseteq \mathcal{L}(\mathcal{I})$ , provided that  $\mathcal{I}$  has the hereditary Baire property.

- (vi)  $\Sigma_2^0 \subseteq \mathcal{L}(\mathcal{I})$ , provided that  $\mathcal{I}$  has the hereditary Baire property and is not a  $P^+$ -ideal.

*Proof.* It follows by [16, Proposition 1.4, Corollary 3.9, and Theorem 4.5] and [17, Corollary 2.5 and Theorems 2.4, 2.8, and 2.10]. (We added the ONLY IF part of item (iv), which is straightforward.)  $\square$

In what follows, we divide our main results into four sections. First, we show in Section 2 that  $\mathcal{L}(\mathcal{I})$  can be equal to families of arbitrarily high Borel complexity. Also, we prove that we cannot have the equality  $\mathcal{L}(\mathcal{I}) = \Pi_2^0$ : more precisely,  $\mathcal{L}(\mathcal{I}) \subseteq \Pi_2^0$  if and only if  $\mathcal{L}(\mathcal{I}) \subseteq \Pi_1^0$ . Then, we show that, if  $X$  is a first countable space and  $\mathcal{I}$  is a simply coanalytic ideal (see Section 3 for details), then every  $\Lambda_{\mathbf{x}}(\mathcal{I})$  is simply analytic. Lastly, we study structural and topological properties of “smallest” ideals  $\mathcal{I}$  such that  $\mathcal{L}(\mathcal{I})$  contains a given subsets of  $X$ .

## 2. LARGE AND SMALL BOREL COMPLEXITIES

Our first main result computes explicitly some families  $\mathcal{L}(\mathcal{I})$ , proving that they may attain arbitrarily large Borel complexity (Theorem 2.5 below). This is somehow related to [16, Question 3.11], which asks about the existence of a Borel ideal  $\mathcal{I}$  such that  $\mathcal{L}(\mathcal{I})$  contains sets with large Borel complexities.

For, given (possibly nonadmissible) ideals  $\mathcal{I}$  and  $\mathcal{J}$  on two countably infinite sets  $Z$  and  $W$ , respectively, we define their *Fubini product* by

$$\mathcal{I} \times \mathcal{J} := \{S \subseteq Z \times W : \{n \in Z : \{k \in W : (n, k) \in S\} \notin \mathcal{J}\} \in \mathcal{I}\},$$

which is an ideal on the countably infinite set  $Z \times W$ , see e.g. [7, Chapter 1]. Hence, recursively,  $\text{Fin}^\alpha := \text{Fin} \times \text{Fin}^{\alpha-1}$  for all integers  $\alpha \geq 2$  is an ideal on  $\omega^\alpha$ .

**Theorem 2.1.** *Let  $\mathcal{I}, \mathcal{J}$  be (possibly nonadmissible) ideals on  $\omega$  such that  $\mathcal{I} \times \mathcal{J}$  is an admissible ideal on  $\omega^2$ . Fix also a bijection  $h : \omega^2 \rightarrow \omega$  and let  $\mathbf{x} = (x_n)$  be a sequence with values in a first countable space  $X$ . For each  $n \in \omega$ , define the sequence  $\mathbf{x}^{(n)} = (x_k^{(n)} : k \in \omega)$  by  $x_k^{(n)} := x_{h(n,k)}$  for all  $k \in \omega$ . Then*

$$\Lambda_{\mathbf{x}}(h[\mathcal{I} \times \mathcal{J}]) = \{\eta \in X : \{n \in \omega : \eta \in \Lambda_{\mathbf{x}^{(n)}}(\mathcal{J})\} \notin \mathcal{I}\}, \quad (1)$$

where  $h[\mathcal{I} \times \mathcal{J}]$  stands for the family  $\{h[S] : S \in \mathcal{I} \times \mathcal{J}\}$ .

*Proof.* Let  $A$  and  $B$  be the left and right hand side of (1), respectively.

INCLUSION  $A \subseteq B$ . The inclusion is clear if  $A = \emptyset$ . Otherwise fix a point  $\eta \in A$ . Hence there exists a set  $S \subseteq \omega^2$  such that  $S \notin \mathcal{I} \times \mathcal{J}$  and the subsequence  $(x_{h(s)} : s \in S)$  is convergent to  $\eta$ . By the definition of Fubini product  $\mathcal{I} \times \mathcal{J}$ ,

$$N := \{n \in \omega : K_n \in \mathcal{J}^+\} \in \mathcal{I}^+, \quad \text{where } K_n := \{k \in \omega : (n, k) \in S\}.$$

At this point, for each  $n \in N$ , the subsequence  $(x_k^{(n)} : k \in K_n)$  is convergent to  $\eta$  and  $K_n \in \mathcal{J}^+$ . Since  $N \in \mathcal{I}^+$ , we obtain that  $\eta \in B$ .

INCLUSION  $B \subseteq A$ . The inclusion is clear if  $B = \emptyset$ . Otherwise fix a point  $\eta \in B$  and let  $(U_n)$  be a decreasing local base of neighborhoods at  $\eta$ . Hence there exists  $N \in \mathcal{I}^+$  such that  $\eta$  is a  $\mathcal{J}$ -limit point of  $\mathbf{x}^{(n)}$  for each  $n \in N$ , let us say  $\lim_{k \in K_n} x_k^{(n)} = \eta$  for some  $K_n \in \mathcal{J}^+$ . Upon removing finitely many elements, we can suppose without loss of generality that

$$\forall n \in N, \forall k \in K_n, \quad x_k^{(n)} \in U_n.$$

Now, set  $S := \{(n, k) \in \omega^2 : n \in N, k \in K_n\}$  and note that  $S \notin \mathcal{I} \times \mathcal{J}$ . It follows that the subsequence  $(x_{h(s)} : s \in S)$  is convergent to  $\eta$ : indeed there are only finitely many elements of the subsequence outside each  $U_n$ . Therefore  $\eta \in A$ , which concludes the proof.  $\square$

**Remark 2.2.** The above result holds for every topological space  $X$  if  $\mathcal{I} = \{\emptyset\}$ . Indeed, in the second part of the proof it is enough to let  $N$  be a singleton.

It is worth noting that if  $\mathcal{I}$  is an ideal on a countably infinite set  $Z$ ,  $\phi : Z \rightarrow \omega$  is a bijection, and  $\mathbf{x} \in X^Z$  is a  $Z$ -indexed sequence with values in a topological space  $X$ , then  $\phi[\mathcal{I}] := \{\phi[S] : S \in \mathcal{I}\}$  is an ideal on  $\omega$  and  $\Lambda_{\mathbf{x}}(\mathcal{I}) = \Lambda_{\mathbf{y}}(\phi[\mathcal{I}])$ , where  $\mathbf{y} \in X^\omega$  is the sequence defined by  $y_n := x_{\phi^{-1}(n)}$  for all  $n \in \omega$ . Hence we may use interchangeably  $\mathcal{L}(\mathcal{I})$  or  $\mathcal{L}(\phi[\mathcal{I}])$ . In particular, Theorem 2.1 allows to compute explicitly families of the type  $\mathcal{L}(\mathcal{I} \times \mathcal{J})$ . For, we state two consequence of Theorem 2.1:

**Corollary 2.3.** *Let  $X$  be a topological space and  $\mathcal{I}$  be an ideal on  $\omega$ . Then*

$$\mathcal{L}(\emptyset \times \mathcal{I}) = \left\{ \bigcup_n A_n : A_0, A_1, \dots \in \mathcal{L}(\mathcal{I}) \right\}.$$

*Proof.* It follows by Theorem 2.1 and Remark 2.2 that

$$\begin{aligned} \mathcal{L}(\emptyset \times \mathcal{I}) &= \{\Lambda_{\mathbf{x}}(\emptyset \times \mathcal{I}) : \mathbf{x} \in X^\omega\} \\ &= \left\{ \bigcup_n \Lambda_{\mathbf{x}^{(n)}}(\mathcal{I}) : \mathbf{x} \in X^\omega \right\} \\ &= \left\{ \bigcup_n \Lambda_{\mathbf{x}^{(n)}}(\mathcal{I}) : \mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots \in X^\omega \right\} \\ &= \left\{ \bigcup_n A_n : A_0, A_1, \dots \in \mathcal{L}(\mathcal{I}) \right\}, \end{aligned}$$

completing the proof.  $\square$

**Corollary 2.4.** *Let  $X$  be a first countable space and  $\mathcal{I}$  be an ideal on  $\omega$ . Then*

$$\mathcal{L}(\text{Fin} \times \mathcal{I}) = \{\limsup_n A_n : A_0, A_1, \dots \in \mathcal{L}(\mathcal{I})\}.$$

*Proof.* It follows by Theorem 2.1 that

$$\begin{aligned} \mathcal{L}(\text{Fin} \times \mathcal{I}) &= \{\Lambda_{\mathbf{x}}(\text{Fin} \times \mathcal{I}) : \mathbf{x} \in X^\omega\} \\ &= \{ \{\eta \in X : \exists^\infty n \in \omega, \eta \in \Lambda_{\mathbf{x}^{(n)}}(\mathcal{I})\} : \mathbf{x} \in X^\omega \} \\ &= \left\{ \bigcap_n \bigcup_{k \geq n} \Lambda_{\mathbf{x}^{(k)}}(\mathcal{I}) : \mathbf{x}^{(0)}, \mathbf{x}^{(1)}, \dots \in X^\omega \right\} \\ &= \{\limsup_n A_n : A_0, A_1, \dots \in \mathcal{L}(\mathcal{I})\}, \end{aligned}$$

completing the proof.  $\square$

At this point, recall that  $\text{Fin}$  is a  $F_\sigma$ -ideal, and that  $\emptyset \times \text{Fin}$  is an analytic  $P$ -ideal which is not  $F_\sigma$ , see [7, Example 1.2.3] and [25, Exercise 23.1]. Thus, it follows by Theorem 1.2 that

$$\mathcal{L}(\text{Fin}) = \Pi_1^0 \quad \text{and} \quad \mathcal{L}(\emptyset \times \text{Fin}) = \Sigma_2^0. \quad (2)$$

With the above premises, we are able to extend (2) to certain ideals with large Borel complexity families  $\mathcal{L}(\mathcal{I})$ .

**Theorem 2.5.** *Let  $X$  be a complete metrizable space. Then, for each positive integer  $\alpha$ , we have:*

- (i)  $\text{Fin}^\alpha$  is a  $\Sigma_{2^\alpha}^0$ -ideal and  $\mathcal{L}(\text{Fin}^\alpha) = \Pi_{2^{\alpha-1}}^0$ .
- (ii)  $\emptyset \times \text{Fin}^\alpha$  is a  $\Pi_{2^{\alpha+1}}^0$ -ideal and  $\mathcal{L}(\emptyset \times \text{Fin}^\alpha) = \Sigma_{2^\alpha}^0$ .

*Proof.* (i) The complexity of  $\text{Fin}^\alpha$  is obtained applying recursively [28, Proposition 1.6.16], while the computation of  $\mathcal{L}(\text{Fin}^\alpha)$  is obtained putting together the base case (2), Corollary 2.4, and [25, Exercise 23.5(i)].

The proof of (ii) goes similarly, replacing Corollary 2.4 with Corollary 2.3.  $\square$

In particular, if  $X$  is a complete metrizable space,  $\mathcal{L}_X(\text{Fin}^2) = \Pi_3^0$ . This provides a generalization of [2, Example 4.2], where it is proved constructively that there exists a *real* sequence  $\mathbf{x}$  such that  $\Lambda_{\mathbf{x}}(\text{Fin}^2)$  is equal to  $[0, 1] \setminus \mathbb{Q}$  (note that the latter is not a  $F_\sigma$ -set, hence  $\mathcal{L}_{\mathbb{R}}(\text{Fin}^2) \cap (\Pi_2^0 \setminus \Sigma_2^0)$  is nonempty).

Our second main result deals with ideals with small topological complexity. Suppose that  $X$  is a first countable space, and recall that

$$\mathcal{L}(\mathcal{I}) \subseteq \Pi_1^0, \quad (3)$$

provided  $\mathcal{I}$  is a  $\Sigma_2^0$ -ideal, see [2, Theorem 2.3]; note that the Hausdorffness hypothesis is not needed here. In the next result, we are going to show that the same conclusion holds if  $\mathcal{I}$  is a  $\Sigma_3^0$ -ideal.

**Theorem 2.6.** *Let  $X$  be a first countable space, and suppose that  $\mathcal{I}$  is a  $\Sigma_3^0$ -ideal. Then inclusion (3) holds.*

It is worth noting that Theorem 2.6 is not a consequence of the former result as it really includes new cases: indeed, as remarked also by Solecki in [30, p. 345], there exists a  $\Delta_3^0$ -ideal on  $\omega$  (hence, both  $\Sigma_3^0$  and  $\Pi_3^0$ ) which is neither  $\Sigma_2^0$  nor  $\Pi_2^0$ , see [34]; cf. also [4, 5] and [28, Proposition 1.2.1].

In addition, Theorem 2.6 allows us to prove a generalization of the folklore result that every  $\Sigma_2^0$ -ideal is a  $P^+$ -ideal, see [23]. For a different proof of the second part, see also [9, Proposition 10.1].

**Corollary 2.7.** *Let  $\mathcal{I}$  be a  $\Sigma_3^0$ -ideal on  $\omega$ . Then  $\mathcal{I}$  is a  $P^+$ -ideal, and there exists a  $\Sigma_2^0$ -ideal  $\mathcal{J}$  such that  $\mathcal{I} \subseteq \mathcal{J}$ .*

*Proof.* The first part follows putting together Theorem 2.6 and Theorem 1.3.(ii) (note that  $\mathcal{I}$  is Borel, hence with the hereditary Baire property). The second part follows by the known fact that a Borel ideal on  $\omega$  is contained in a  $\Sigma_2^0$ -ideal if and only if it is contained in a  $P^+$ -ideal, see [23].  $\square$

At this point, we divide the proof of Theorem 2.6 into two intermediate steps. To this aim, we recall that properties of ideals can be often expressed by finding *critical* ideals with respect to some preorder, cf. e.g. the survey [19]. To this aim, let  $\mathcal{I}$  and  $\mathcal{J}$  be two ideals on two countably infinite sets  $Z$  and  $W$ , respectively. Then we say that  $\mathcal{I}$  is below  $\mathcal{J}$  in the Rudin–Blass ordering, shortened as

$$\mathcal{I} \leq_{\text{RB}} \mathcal{J},$$

if there is a finite-to-one map  $\phi : W \rightarrow Z$  such that  $S \in \mathcal{I}$  if and only if  $\phi^{-1}[S] \in \mathcal{J}$  for all subsets  $S \subseteq Z$ . The restriction of these orderings to maximal ideals  $\mathcal{I}$ , and the Borel cardinality of the quotients  $\mathcal{P}(Z)/\mathcal{I}$  have been extensively studied, see e.g. [6, 18, 26] and references therein.

**Lemma 2.8.** *Let  $X$  be a first countable Hausdorff space and  $\mathcal{I}$  be an ideal on  $\omega$  with the hereditary Baire property. Suppose that inclusion (3) does not hold. Then  $\emptyset \times \text{Fin} \leq_{\text{RB}} \mathcal{I}$ .*

*Proof.* Since inclusion (3) fails and  $X$  is first countable, there exists a sequence  $\mathbf{x}$  such that  $S := \Lambda_{\mathbf{x}}(\mathcal{I})$  is not closed, hence not sequentially closed. Therefore there exists a sequence  $\mathbf{y}$  taking values in  $S$  which is convergent to some limit  $\eta \in X \setminus S$ . Since  $X$  is Hausdorff, we may suppose without loss of generality that  $y_n \neq y_m$  for all distinct  $n, m \in \omega$ . Now, for each  $n \in \omega$ , there exists  $A_n \in \mathcal{I}^+$  such that  $\lim_{k \in A_n} x_k = y_n$ . Define  $B_0 := A_0 \cup (\omega \setminus \bigcup_n A_n)$  and, recursively,  $B_{n+1} := A_{n+1} \setminus \bigcup_{k \leq n} B_k$  for all  $n \in \omega$ . Hence  $\{B_n : n \in \omega\}$  is a partition of  $\omega$  into  $\mathcal{I}$ -positive sets, for each  $n \in \omega$ , the restriction  $\mathcal{I} \upharpoonright B_n$  is an ideal on  $B_n$  with the Baire property.

It follows by Talagrand's characterization of meager ideals that  $\text{Fin} \leq_{\text{RB}} \mathcal{I} \upharpoonright B_n$  for each  $n \in \omega$ . More explicitly, for each  $n \in \omega$  there exists a finite-to-one map  $\phi_n : B_n \rightarrow \omega$  such that

$$\forall W \subseteq \omega, \quad \phi_n^{-1}[W] \in \mathcal{I} \quad \text{if and only if} \quad W \in \text{Fin}, \quad (4)$$

see [33, Theorem 2.1]; cf. also [1] for further characterizations of meager ideals based of  $\mathcal{I}$ -limit points of sequences. Define the map  $\phi : \omega \rightarrow \omega^2$  by

$$\forall n \in \omega, \forall k \in B_n, \quad \phi(k) = (n, \phi_n(k)).$$

We claim that  $\phi$  is a witnessing function for  $\emptyset \times \text{Fin} \leq_{\text{RB}} \mathcal{I}$ . For, suppose that  $W \subseteq \omega^2$  belongs to  $\emptyset \times \text{Fin}$ . Then

$$\phi^{-1}[W] = \bigcup_{n \in \omega} \phi_n^{-1}[\{k \in \omega : (n, k) \in W\}] \quad (5)$$

Since each  $\phi_n$  is finite-to-one,  $\phi^{-1}[W]$  has finite intersection with each  $B_n$ . Then either  $\phi^{-1}[W]$  is finite or the subsequence  $(x_n : n \in \phi^{-1}[W])$  is convergent to  $\eta$ , while  $\eta$  is not an  $\mathcal{I}$ -limit point of  $\mathbf{x}$ . Hence, in both cases,  $\phi^{-1}[W] \in \mathcal{I}$ . Conversely, suppose that  $W$  does not belong to  $\emptyset \times \text{Fin}$ , so that there exists  $n_0 \in \omega$  such that  $\{k \in \omega : (n_0, k) \in W\} \notin \text{Fin}$ . It follows by (4) and (5) that  $\phi^{-1}[W]$  contains

$$\phi_{n_0}^{-1}[\{k \in \omega : (n_0, k) \in W\}] \in \mathcal{I}^+.$$



Therefore  $\phi^{-1}[W] \in \mathcal{I}$  if and only if  $W \in \emptyset \times \text{Fin}$  for each  $W \subseteq \omega^2$ .  $\square$

**Corollary 2.9.** *Under the same hypotheses of Lemma 2.8,  $\Sigma_2^0 \subseteq \mathcal{L}(\mathcal{I})$ .*

*Proof.* Thanks to Lemma 2.8 and [16, Proposition 3.8], we have  $\mathcal{L}(\emptyset \times \text{Fin}) \subseteq \mathcal{L}(\mathcal{I})$ . The claim follows by Equation (2).  $\square$

For the next intermediate result, given topological spaces  $X, Y$  and subsets  $A \subseteq X$  and  $B \subseteq Y$ , we say that  $A$  is *Wadge reducible to  $B$* , shortened as

$$A \leq_W B,$$

if there exists a continuous map  $\Phi : X \rightarrow Y$  such that  $\Phi^{-1}[B] = A$  (or, equivalently,  $x \in A$  if and only if  $\Phi(x) \in B$  for all  $x \in X$ ), see e.g. [25, Definition 21.13]. If, in addition,  $X$  and  $Y$  are Polish spaces with  $X$  zero-dimensional, then  $B$  is said to be  $\Pi_3^0$ -hard if  $A \leq_W B$  for some  $A \in \Pi_3^0(X)$ . Lastly, if  $B$  is a  $\Pi_3^0(Y)$  set which is also  $\Pi_3^0$ -hard, then it is called  $\Pi_3^0$ -complete. (Analogous definitions can be given for other classes of sets in Polish spaces, see [25, Definition 22.9].)

**Lemma 2.10.** *Let  $\mathcal{I}$  be an ideal on  $\omega$  such that  $\emptyset \times \text{Fin} \leq_{\text{RB}} \mathcal{I}$ . Then  $\mathcal{I}$  is  $\Pi_3^0$ -hard.*

*Proof.* First, recall that  $\emptyset \times \text{Fin}$  is a  $\Pi_3^0$ -complete subset of  $\mathcal{P}(\omega^2)$ , see e.g. [25, Exercise 23.1]. Hence, to complete the proof, it is sufficient to show that

$$\emptyset \times \text{Fin} \leq_W \mathcal{I}.$$

By hypothesis, there exists a finite-to-one function  $\phi : \omega \rightarrow \omega^2$  such that  $S \in \emptyset \times \text{Fin}$  if and only if  $\phi^{-1}[S] \in \mathcal{I}$ . Now, define the map  $\Phi : \mathcal{P}(\omega^2) \rightarrow \mathcal{P}(\omega)$  by  $\Phi(S) := \phi^{-1}[S]$ , so that  $S \in \emptyset \times \text{Fin}$  if and only if  $\Phi(S) \in \mathcal{I}$ . Hence, we only need to show that  $\Phi$  is continuous. For, fix  $n \in \omega$  and define  $S := \phi[\{0, \dots, n\}]$ . It follows that, for all  $A, B \subseteq \omega^2$  with  $A \cap S = B \cap S$ ,

$$\begin{aligned} \Phi(A) \cap [0, n] &= \{k \in \omega : \phi(k) \in A \text{ and } k \leq n\} \\ &= \{k \in \omega : \phi(k) \in A \cap S\} = \Phi(B) \cap [0, n], \end{aligned}$$

which proves the continuity of  $\Phi$ .  $\square$

*Proof of Theorem 2.6.* Suppose that  $X$  is a first countable space and  $\mathcal{I}$  is an ideal on  $\omega$  such that inclusion (3) fails. We claim that  $\mathcal{I}$  is not a  $\Sigma_3^0$ -ideal.

If  $\mathcal{I}$  is not a Borel ideal, then the claim is trivial. Hence, let us suppose hereafter that  $\mathcal{I}$  is Borel. In particular,  $\mathcal{I}$  has the hereditary Baire property. At this point, it follows by Lemma 2.8 and Lemma 2.10 that  $\mathcal{I}$  is a  $\Pi_3^0$ -hard. To sum up,  $\mathcal{I}$  is a Borel subset of a zero-dimensional Polish space and it is  $\Pi_3^0$ -hard. We conclude by [25, Theorem 22.10] that the  $\mathcal{I}$  is not a  $\Sigma_3^0$ -ideal.  $\square$

## 3. SIMPLY ANALYTICITY

In [2, Proposition 4.1], the first and last-named authors proved the following:

**Proposition 3.1.** *Let  $\mathbf{x}$  be a sequence taking values in a Hausdorff regular first countable space  $X$ . Let also  $\mathcal{I}$  be a coanalytic ideal on  $\omega$ . Then  $\Lambda_{\mathbf{x}}(\mathcal{I})$  is analytic.*

However, the classical definition of analytic sets  $A \subseteq X$  as projections of Borel subsets of  $X \times X$  is usually considered in *Polish spaces*  $X$ , cf. e.g. [31, Chapter 4] or [25, Chapter 14]. In addition, the above result has been re-proved in [17, Theorem 4.1] for zero-dimensional Polish spaces  $X$ , so that

$$\mathcal{L}(\mathcal{I}) \subseteq \Sigma_1^1.$$

whenever  $\mathcal{I}$  is a  $\Pi_1^1$ -ideal (notice that the above notation is meaningful).

In this Section, our aim is to reformulate and clarify the statement and the proof of Proposition 3.1, extending in turn both the latter and the special case treated in [17, Theorem 4.1].

Note that there are several papers in which the notion of analytic set is adapted to more general topological spaces, see e.g. [15, 29]. The theory of the latter extension, called *K-analytic sets*, is important and nontrivial, and it does not proceed verbatim as in the classical one of analytic sets.

For our purposes, we generalize in a straight way one of possible definitions of analytic sets to the case of arbitrary topological spaces. Hereafter,  $\pi_X$  stands for the usual projection on  $X$ .

**Definition 3.2.** Let  $X$  be a topological space. A subset  $A \subseteq X$  is said to be *simply analytic*, shortened as *s-analytic*, if there exists an uncountable Polish space  $Y$  and a Borel subset  $B \subseteq X \times Y$  such that  $A = \pi_X[B]$ .

Note that Definition 3.2 is coherent with the classical notion of analytic set. Indeed, it is well known that a subset  $A$  of a Polish space  $X$  is analytic if and only if there exists an uncountable Polish space  $Y$  and a Borel  $B \subseteq X \times Y$  such that  $A = \pi_X[B]$ , cf. [31, Proposition 4.1.1]. At this point, let  $Y_1, Y_2$  be two uncountable Polish spaces. Then there exists a Borel isomorphism  $h : Y_2 \rightarrow Y_1$ , as it follows by [31, Theorem 3.3.13]. Let  $A := \pi_X[B_1]$  be a s-analytic subset for a Borel set  $B_1 \subseteq X \times Y_1$ . Then  $B_2 := \{(x, y) \in X \times Y_2 : (x, h(y)) \in B_1\}$  is a Borel subset of  $X \times Y_2$  by Lemma 3.4 below and, clearly,  $\pi_X[B_2] = A$ . Therefore, in Definition 3.2 one may assume without loss of generality that  $Y = \omega^\omega$ .

Lastly, a subset  $A \subseteq X$  is said to be *s-coanalytic* if  $A^c := X \setminus A$  is s-analytic. Observe that every Borel set in  $X$  is both s-analytic and s-coanalytic.

These observations enable us to consider Proposition 3.1 valid as it was stated in the original version in [2]. They allow, in addition, to remove in its statement the hypothesis of regularity (and also the property of being Hausdorff; however, the latter one has been used in [2] only because a first countable space is Hausdorff if and only if every sequence has at most one limit):

**Theorem 3.3.** *Let  $\mathbf{x}$  be a sequence taking values in a first countable space  $X$ . Let also  $\mathcal{I}$  be a  $s$ -coanalytic ideal on  $\omega$ . Then  $\Lambda_{\mathbf{x}}(\mathcal{I})$  is  $s$ -analytic.*

For the proof of Theorem 3.3, we will need some intermediate lemmas.

**Lemma 3.4.** *Let  $f : X \rightarrow Z$  and  $g : Y \rightarrow W$  be Borel functions, where  $X, Y, Z, W$  are topological spaces. Then the map  $h : X \times Y \rightarrow Z \times W$  defined by*

$$\forall (x, y) \in X \times Y, \quad h(x, y) := (f(x), g(y))$$

*is Borel.*

*Proof.* Let  $U \subseteq Z$  and  $V \subseteq W$  be arbitrary open sets. It is enough to show that  $h^{-1}[U \times V]$  is a Borel subset of  $X \times Y$ . The latter set is equal to  $f^{-1}[U] \times g^{-1}[V]$ , which belongs to  $\mathcal{B}(X) \otimes \mathcal{B}(Y)$ . The claim follows by the inclusion  $\mathcal{B}(X) \otimes \mathcal{B}(Y) \subseteq \mathcal{B}(X \times Y)$ , see e.g. [11, Proposition 251E]. (For the converse inclusion of the latter, which does not hold for every  $X$  and  $Y$ , see [12].)  $\square$

**Lemma 3.5.** *Let  $f : X \rightarrow Y$  be a Borel function, where  $X$  is a topological space and  $Y$  is an uncountable Polish space. If  $A \subseteq Y$  is an analytic set, then  $f^{-1}[A]$  is a  $s$ -analytic subset of  $X$ .*

*Proof.* Since  $A \subseteq Y$  is analytic, there exists a Borel subset  $B \subseteq Y \times Y$  such that  $A$  is the projection on the first coordinate of  $B$ . At this point, define  $C := \{(x, y) \in X \times Y : (f(x), y) \in B\}$ . Then  $C$  is Borel set by Lemma 3.4 and  $\pi_X[C] = f^{-1}[A]$ . Therefore  $f^{-1}[A]$  is a  $s$ -analytic set, which concludes the proof.  $\square$

**Lemma 3.6.** *Let  $A \subseteq X \times Y$  be a  $s$ -analytic set, where  $X$  is a topological space and  $Y$  is an uncountable Polish space. Then  $C := \pi_X[A]$  is a  $s$ -analytic set.*

*Proof.* Let  $Z$  be an uncountable Polish space and  $B \subseteq X \times Y \times Z$  be a Borel set such that  $A = \pi_{X \times Y}[B]$ . Now, observe that  $C = \pi_X[B]$  and obviously  $Y \times Z$  is an uncountable Polish space.  $\square$

**Lemma 3.7.** *Let  $(A_n)$  be a sequence of  $s$ -analytic subsets of a topological space  $X$ . Then both  $\bigcup_n A_n$  and  $\bigcap_n A_n$  are  $s$ -analytic.*

*Proof.* The proof proceeds verbatim as in [31, Proposition 4.1.2].  $\square$

We are ready for the proof of Theorem 3.3.

*Proof of Theorem 3.3.* Note that  $\mathcal{I}^+$  is s-analytic and let  $\mathcal{N}$  be the set of strictly increasing sequences  $(n_k)$  of nonnegative integers. Since  $\mathcal{N}$  can be regarded as a closed subset of  $\omega^\omega$ , it is a Polish space by Alexandrov's theorem, see e.g. [31, Theorem 2.2.1]. For each  $\eta \in X$ , fix a decreasing local base  $(U_{\eta,m} : m \in \omega)$  of open neighbourhoods of  $\eta$ . Then  $\eta$  is an  $\mathcal{I}$ -limit point of  $\mathbf{x}$  if and only if there exists a sequence  $(n_k) \in \mathcal{N}$  such that  $\{n_k : k \in \omega\} \in \mathcal{I}^+$  and  $\{k \in \omega : x_{n_k} \notin U_{\eta,m}\} \in \text{Fin}$  for all  $m \in \omega$ .

At this point, define the continuous function

$$\psi : \mathcal{N} \rightarrow \{0, 1\}^\omega : (n_k) \mapsto \chi_{\{n_k : k \in \omega\}},$$

where  $\chi_S$  stands for the characteristic function of a set  $S \subseteq \omega$ . In addition, for each  $m \in \omega$ , define the function  $\zeta_m : \mathcal{N} \times X \rightarrow \{0, 1\}^\omega$  by

$$\forall (n_k) \in \mathcal{N}, \forall \eta \in X, \quad \zeta_m((n_k), \eta) := \chi_{\{t \in \omega : x_{n_t} \notin U_{\eta,m}\}}$$

Identifying each  $S \subseteq \omega$  with its characteristic function  $\chi_S$ , it follows that

$$\Lambda_{\mathbf{x}}(\mathcal{I}) = \pi_X \left[ (\psi^{-1}[\mathcal{I}^+] \times X) \cap \bigcap_{m \in \omega} \zeta_m^{-1}[\text{Fin}] \right]. \quad (6)$$

**CLAIM 1.**  $\psi^{-1}[\mathcal{I}^+] \times X$  is a s-analytic subset of  $\mathcal{N} \times X$ .

*Proof.* Thanks to Lemma 3.5 and the fact that  $\psi$  is continuous,  $\psi^{-1}[\mathcal{I}^+]$  is an (ordinary) analytic subset of  $\mathcal{N}$ . Hence there exists an uncountable Polish space  $Y$  and a Borel set  $B \subseteq \mathcal{N} \times Y$  such that  $\psi^{-1}[\mathcal{I}^+] = \pi_{\mathcal{N}}[B]$ . The claim follows from the fact that

$$B \times X \in \mathcal{B}(\mathcal{N} \times Y) \times \mathcal{B}(X) \subseteq \mathcal{B}(\mathcal{N} \times Y \times X),$$

see e.g. [11, Proposition 251E], and that  $\pi_{\mathcal{N} \times X}[B \times X] = \pi_{\mathcal{N}}[B] \times X$ .  $\square$

**CLAIM 2.** For each  $\eta \in X$  and  $m \in \omega$ , the section  $\zeta_m(\cdot, \eta)$  is continuous.

*Proof.* Fix  $\eta \in X$  and  $m \in \omega$ . It is enough to show that the section  $\zeta_m(\cdot, \eta)$  is sequentially continuous. For, pick a sequence  $(n_k^{(p)} : p \in \omega)$  of elements of  $\mathcal{N}$  which is convergent to some  $(n_k) \in \mathcal{N}$ . Then, for each  $t \in \omega$ , there exists  $p_t \in \omega$  such that  $n_t^{(p)} = n_t$  for all  $p \geq p_t$ . Hence  $\zeta_m((n_k^{(p)}), \eta)(t) = \zeta_m((n_k), \eta)(t)$  for all  $p \geq p_t$ . This proves that  $\lim_p \zeta_m((n_k^{(p)}), \eta) = \zeta_m((n_k), \eta)$ .  $\square$

**CLAIM 3.** For each  $(n_k) \in \mathcal{N}$  and  $m \in \omega$ , the section  $\zeta_m((n_k), \cdot)$  is Borel measurable.

*Proof.* Fix  $(n_k) \in \mathcal{N}$  and  $m \in \omega$  and define for convenience  $\xi : X \rightarrow \{0, 1\}^\omega$  by  $\xi(\eta) := \zeta_m((n_k), \eta)$  for all  $\eta \in X$ . It is enough to show that preimage  $\xi^{-1}[U]$  is Borel in  $X$  for every basic open set  $U \subseteq \mathcal{N}$ . This is clear if  $U = \emptyset$ . Otherwise there exist  $p \in \omega$  and  $r_0, r_1, \dots, r_p \in \{0, 1\}$  such that  $U = \{a \in \{0, 1\}^\omega : a(t) = r_t \text{ for all } t = 0, 1, \dots, p\}$ . Set  $A := \{t \in \{0, 1, \dots, p\} : r_t = 0\}$  and  $B := \{t \in \{0, 1, \dots, p\} : r_t = 1\}$ . It follows that

$$\begin{aligned} \xi^{-1}[U] &= \{\eta \in X : \zeta_m((n_k), \eta)(t) = r_t \text{ for all } t = 0, 1, \dots, p\} \\ &= \left( \bigcap_{t \in A} \{\eta \in X : \zeta_m((n_k), \eta)(t) = 0\} \right) \cap \left( \bigcap_{t \in B} \{\eta \in X : \zeta_m((n_k), \eta)(t) = 1\} \right) \\ &= \left( \bigcap_{t \in A} \{\eta \in X : x_{n_t} \in U_{\eta, m}\} \right) \cap \left( \bigcap_{t \in B} \{\eta \in X : x_{n_t} \notin U_{\eta, m}\} \right) \\ &= \left( \bigcap_{t \in A} \bigcup_{\eta \in X : x_{n_t} \in U_{\eta, m}} U_{\eta, m} \right) \cap \left( \bigcap_{t \in B} \bigcap_{\eta \in X : x_{n_t} \notin U_{\eta, m}} U_{\eta, m}^c \right). \end{aligned}$$

Since each  $U_{\eta, m}$  is open, we conclude that  $\xi^{-1}[U]$  is a Borel set.  $\square$

**CLAIM 4.** For each  $m \in \omega$ , the set  $\zeta_m^{-1}[\text{Fin}]$  is Borel.

*Proof.* Thanks to Claim 2, Claim 3, and [31, Theorem 3.1.30], each map  $\zeta_m$  is Borel measurable. The conclusion follows since  $\text{Fin}$  is a  $F_\sigma$ -set.  $\square$

To conclude the proof of Proposition 3.3, we have by Claim 1 and Claim 4 that both  $\psi^{-1}[\mathcal{I}^+] \times X$  and  $\sigma_m^{-1}[\text{Fin}]$  are s-analytic subsets of  $\mathcal{N} \times X$ , hence also their intersection by Lemma 3.7. By the identity (6),  $\Lambda_{\mathbf{x}}(\mathcal{I})$  is the projection on  $X$  of a s-analytic subset of  $\mathcal{N} \times X$ , which is s-analytic by Lemma 3.6.  $\square$

#### 4. MINIMAL IDEALS $\mathcal{I}_W$ AND THEIR COMPLEXITIES

In this last Section, following the line of research initiated in [16, 17], we recall the definition of certain ideals and we study their structural and topological complexity. For, given a sequence  $\mathbf{x}$  taking values in a first countable Hausdorff space  $X$  and a subset  $W \subseteq X$ , define the ideal

$$\mathcal{I}_W := \{A \subseteq \omega : \mathbb{L}_{\mathbf{x}|A} \cap W = \emptyset\},$$

where  $L_{\mathbf{x}\upharpoonright A} := \Lambda_{\mathbf{x}\upharpoonright A}(\text{Fin})$ , see [16, Section 2]. More explicitly,  $A \subseteq \omega$  belongs to  $\mathcal{I}_W$  if and only if  $A$  is finite or, in the opposite,  $A$  is infinite and there are no infinite subsets  $B \subseteq A$  such that  $(x_n : n \in B)$  is convergent to some element of  $W$ . Note that  $\mathcal{I}_W$  may be not admissible: for instance, if  $W = \emptyset$  then  $\mathcal{I}_W = \mathcal{P}(\omega)$ .

The main reason of its introduction is the following:

**Theorem 4.1.** *Let  $\mathbf{x}$  be a sequence taking values in a first countable Hausdorff space  $X$  and fix a subset  $W \subseteq X$ . Then*

$$\Lambda_{\mathbf{x}}(\mathcal{I}_W) = W \cap L_{\mathbf{x}}. \quad (7)$$

*In particular, if  $X$  is separable, then  $W \in \mathcal{L}(\mathcal{I}_W)$ .*

*Proof.* The first part follows by [16, Theorem 2.2(i)]. This implies, if  $W \subseteq L_{\mathbf{x}}$ , then  $\Lambda_{\mathbf{x}}(\mathcal{I}_W) = W$ , so that  $W \in \mathcal{L}(\mathcal{I}_W)$ . Hence, the second part is obtained by choosing a sequence  $\mathbf{x}$  with dense image.  $\square$

Note that that Theorem 4.1 proves that, even if  $X = \mathbb{R}$ , the family  $\mathcal{L}(\mathcal{I})$  may contain sets which are not Borel, which answered an open question in [2].

It is also worth to remark that ideals  $\mathcal{I}_W$  are not the only ones for which (7) holds, see [16, Theorem 2.7]; cf. also [17, Theorems 3.4 and 3.5] for further refinements of Theorem 4.1 in regular and Polish spaces. However, we show that they are the smallest ideals with such property. For, we say that  $\mathcal{I}$  contains an isomorphic copy of  $\mathcal{J}$  if there exists a bijection  $\phi$  on  $\omega$  such that  $\phi[\mathcal{J}] \subseteq \mathcal{I}$ , where  $\phi[\mathcal{J}]$  stands for the family  $\{\phi[S] : S \in \mathcal{J}\}$ .

**Proposition 4.2.** *Let  $\mathbf{x}$  be a sequence taking values in a compact metric space  $X$  and fix a dense subset  $W \subseteq X$ . Let also  $\mathcal{I}$  be an ideal on  $\omega$  such that  $W \in \mathcal{L}(\mathcal{I})$ . Suppose also that  $\mathbf{x}$  has dense image, and that*

$$\forall A \in \mathcal{I}^+, \forall \mathbf{y} \in X^\omega, \quad \Lambda_{\mathbf{y}\upharpoonright A}(\mathcal{I} \upharpoonright A) \neq \emptyset. \quad (8)$$

*Then  $\mathcal{I}$  contains an isomorphic copy of  $\mathcal{I}_W$ .*

*Proof.* By hypothesis, there exists a sequence  $\mathbf{y}$  taking values in  $X$  such that  $\Lambda_{\mathbf{y}}(\mathcal{I}) = W$ . Let  $d$  be a compatible metric on  $X$  and note that the denseness of  $W$  implies that  $L_{\mathbf{y}} = X$ . Since  $X$  is compact, there exists a strictly increasing sequence of integers  $(\iota_n : n \in \omega)$  such that  $\iota_0 := 0$  and, for each  $n \in \omega$ , the family of open balls  $\{B(x_k, 2^{-n}) : k \in [\iota_n, \iota_{n+1}]\}$  is an open cover of  $X$ .

At this point, define recursively the map  $\phi : \omega \rightarrow \omega$  as it follows:

- (i)  $\phi(0)$  is the smallest integer  $k \in \omega$  such that  $d(y_{\phi(k)}, x_0) < 1$ ;

- (ii) for each integer  $m \in (1, \iota_1)$ ,  $\phi(m)$  is the smallest integer  $k \in \omega$  such that  $d(y_{\phi(k)}, x_m) < 1$  and  $k \notin \phi[\{0, 1, \dots, m-1\}]$ ;
- (iii) for each  $n, m \in \omega$  with  $n \geq 1$  and  $m \in [\iota_n, \iota_{n+1})$ ,  $\phi(m)$  is the smallest integer  $k \in \omega$  such that  $d(y_{\phi(k)}, x_m) < 2^{-n}$  and  $k \notin \phi[\{0, 1, \dots, m-1\}]$ .

It follows by construction that  $\phi$  is a bijection on  $\omega$  which satisfies

$$\lim_{n \rightarrow \infty} d(y_{\phi(n)}, x_n) = 0. \quad (9)$$

Fix  $A \in \mathcal{I}^+$  and note that the set of  $\mathcal{I} \upharpoonright A$ -limit points of  $\mathbf{y} \upharpoonright A$  is contained both in  $L_{\mathbf{y} \upharpoonright A}$  and in  $\Lambda_{\mathbf{y}}(\mathcal{I})$ . Since the latter is equal to  $W$ , we obtain by (8) and (9) that

$$\emptyset \neq \Lambda_{\mathbf{y} \upharpoonright A}(\mathcal{I} \upharpoonright A) \subseteq L_{\mathbf{y} \upharpoonright A} \cap W = L_{\mathbf{x} \upharpoonright \phi^{-1}[A]} \cap W.$$

By the definition of  $\mathcal{I}_W$ , we get  $\phi^{-1}[A] \notin \mathcal{I}_W$ . Therefore  $\phi[\mathcal{I}_W] \subseteq \mathcal{I}$ .  $\square$

As remarked in the Introduction, the technical condition (8) has been already studied in the literature. For instance, if  $X$  is compact, all  $F_\sigma$ -ideals  $\mathcal{I}$  satisfy (8), taking into account [2, Theorem 2.3] and [27, Lemma 3.1(vi)].

The following result, due to He et al. [16], deals with the topological complexity of ideals  $\mathcal{I}_W$ . Hereafter, we assume for simplicity that  $X$  is the Cantor space and  $\mathbf{x}$  is an enumeration of the rationals (i.e., finitely supported sequences) of  $X$ , with  $x_0 := (0, 0, \dots)$ .

**Theorem 4.3.** *Let  $W$  be a subset of the Cantor space  $X = \{0, 1\}^\omega$ . Then:*

- (i) if  $W$  is closed then  $\mathcal{I}_W$  is a  $\Sigma_2^0$ -ideal;
- (ii) if  $W$  is  $\Sigma_2^0$  then  $\mathcal{I}_W$  is a  $\Pi_3^0$ -ideal;
- (iii) if  $W$  is open then  $\mathcal{I}_W$  is an analytic  $P$ -ideal.

*Proof.* It follows by [16, Theorem 2.2(ii)] by choosing a sequence  $\mathbf{x}$  with dense image (the case  $W = \emptyset$  holds as well).  $\square$

Taking into account also the results obtained in Section 2, we prove some characterizations for the [non]closedness of the subset  $W$ :

**Theorem 4.4.** *Let  $W$  be a nonempty subset of the Cantor space  $X = \{0, 1\}^\omega$  such that  $\mathcal{I}_W$  has the hereditary Baire property. Then the following are equivalent:*

- (i)  $W$  is not closed;
- (ii)  $\emptyset \times \text{Fin} \leq_{\text{RB}} \mathcal{I}_W$ ;
- (iii)  $\mathcal{I}_W$  is  $\Pi_3^0$ -hard;
- (iv)  $\Sigma_2^0 \subseteq \mathcal{L}(\mathcal{I}_W)$ .

*Proof.* (i)  $\implies$  (ii). Thanks to Theorem 4.1,  $W \in \mathcal{L}(\mathcal{I}_W)$ , hence the inclusion  $\mathcal{L}(\mathcal{I}_W) \subseteq \Pi_1^0$  fails. The claim follows by Lemma 2.8.

(ii)  $\implies$  (iii). It follows by Lemma 2.10.

(ii)  $\implies$  (iv). It follows by Corollary 2.9.

(iii)  $\implies$  (i). If  $W$  is closed, then  $\mathcal{I}_W$  would be a  $\Sigma_2^0$ -ideal by Theorem 4.3.

(iv)  $\implies$  (i). As in the previous implication, if  $W$  is closed, then  $\mathcal{I}_W$  would be  $\Sigma_2^0$ . Hence  $\mathcal{L}(\mathcal{I}_W) = \Pi_1^0$  by Theorem 1.2. However, since  $X$  is separable, there exists a countable dense subset, which is  $\Sigma_2^0$  and not closed.  $\square$

On the same lines, we are able to characterize the openness of the set  $W$  and the  $P$ -property of the ideal  $\mathcal{I}_W$ , improving Theorem 4.3(iii):

**Theorem 4.5.** *Let  $W$  be a subset of the Cantor space  $X = \{0, 1\}^\omega$ . Then the following are equivalent:*

- (i)  $W$  is open;
- (ii)  $\mathcal{I}_W$  is an analytic  $P$ -ideal;
- (iii)  $\mathcal{I}_W$  is a  $P$ -ideal.

*Proof.* (i)  $\implies$  (ii). It follows by Theorem 4.3(iii).

(ii)  $\implies$  (iii). This is obvious.

(iii)  $\implies$  (i). Let us suppose that  $W$  is not open, i.e.,  $W^c$  is not closed. Since  $X$  is metrizable,  $W^c$  is not sequentially closed. Hence it is possible to pick a sequence  $\mathbf{y}$  taking values in  $W^c$  which is convergent to some  $\eta \in W$ . Let us denote by  $d$  a compatible metric on  $X$ , so that  $\lim_k d(y_k, \eta) = 0$ . Since  $\mathbf{x}$  is dense, for each  $k \in \omega$  there exists an infinite set  $A_k \subseteq \omega$  such that  $\lim \mathbf{x} \upharpoonright A_k = y_k$ . Considering that  $y_k \in W^c$ , it follows that  $L_{\mathbf{x} \upharpoonright A_k} \cap W = \emptyset$ , hence  $A_k \in \mathcal{I}_W$ .

At this point, pick a set  $A \subseteq \omega$  such that  $A_k \setminus A \in \text{Fin}$  for all  $k \in \omega$ . Set for convenience  $b_{-1} := 0$  and define recursively a sequence of integers  $(b_k : k \in \omega)$  such that  $b_k$  is the smallest element of  $A \cap A_k$  for which  $d(y_k, x_{b_k}) \leq d(y_k, \eta)$  and  $b_k > b_{k-1}$  (note that this is well defined). It follows by the triangle inequality that

$$\forall k \in \omega, \quad d(x_{b_k}, \eta) \leq d(x_{b_k}, y_k) + d(y_k, \eta) \leq 2d(y_k, \eta).$$

Therefore  $B := \{b_k : k \in \omega\}$  is an infinite subset of  $A$  such that  $\lim \mathbf{x} \upharpoonright B = \eta$ . Since  $\eta \in W$ , this proves that  $L_{\mathbf{x} \upharpoonright A} \cap W \neq \emptyset$ . Hence, by the definition of  $\mathcal{I}_W$ , we conclude that  $A \notin \mathcal{I}_W$ . To sum up,  $(A_k : k \in \omega)$  is a witnessing sequence of sets in  $\mathcal{I}_W$  which fails the  $P$ -property for the ideal  $\mathcal{I}_W$ .  $\square$

As a consequence, we obtain that there are only three possibilities for the complexity of ideals  $\mathcal{I}_W$ :



**Corollary 4.6.** *Let  $W$  be a subset of the Cantor space  $X = \{0, 1\}^\omega$ . Then exactly one of the following cases occurs:*

- (i)  $\mathcal{I}_W$  is a  $\Sigma_2^0$ -ideal;
- (ii)  $\mathcal{I}_W$  is a Borel ideal which is not  $\Sigma_3^0$ ;
- (iii)  $\mathcal{I}_W$  is not a Borel ideal.

*Proof.* First, if  $W = \emptyset$ , then  $\mathcal{I}_W = \mathcal{P}(\omega)$ , which satisfies only (i). Hence suppose hereafter that  $W$  is nonempty. Suppose also that  $\mathcal{I}_W$  is Borel, so that (iii) fails and  $\mathcal{I}_W$  has the hereditary Baire property. If  $W$  is closed then  $\mathcal{I}_W$  is  $\Sigma_2^0$  by Theorem 4.3, hence (i) holds. If  $W$  is not closed then  $W$  is  $\Pi_3^0$ -hard by Theorem 4.4, which is equivalent to be not  $\Sigma_3^0$  by [25, Theorem 22.10], hence (ii) holds.  $\square$

In light of Theorem 4.3, one may ask whether the third case in Corollary 4.6 really occurs. We answer in the affirmative with our last main result:

**Theorem 4.7.** *Let  $W$  be a Borel subset of the Cantor space  $X = \{0, 1\}^\omega$  which is not  $\Sigma_2^0$ . Then  $\mathcal{I}_W$  is not analytic, hence not Borel.*

Putting together the above results, we have the following consequence:

**Corollary 4.8.** *Let  $W$  be a Borel subset of the Cantor space  $X = \{0, 1\}^\omega$ . Then the following are equivalent:*

- (i)  $W$  is  $\Sigma_2^0$ ;
- (ii)  $\mathcal{I}_W$  is a  $\Pi_3^0$ -ideal;
- (iii)  $\mathcal{I}_W$  is a Borel ideal;
- (iv)  $\mathcal{I}_W$  is an analytic ideal.

*Proof.* (i)  $\implies$  (ii). It follows by Theorem 4.3(i).

(ii)  $\implies$  (iii)  $\implies$  (iv). They are obvious.

(iv)  $\implies$  (i). It follows by Theorem 4.7.  $\square$

We recall now some definition about trees, which will be needed in the proof of Theorem 4.7.

Denote by  $\{0, 1\}^{<\omega}$  the set of finite  $\{0, 1\}$ -sequences. We say that  $a = (a_0, \dots, a_n) \in \{0, 1\}^{<\omega}$  is an extension of  $b = (b_0, \dots, b_m) \in \{0, 1\}^{<\omega}$ , shortened as  $a \subseteq b$ , if  $n \leq m$  and  $a_k = b_k$  for all  $k \leq n$ . Given an infinite sequence  $z \in \{0, 1\}^\omega$ , we define  $z \upharpoonright n := (z_0, \dots, z_n) \in \{0, 1\}^{<\omega}$  for each  $n \in \omega$ . A *tree on  $\{0, 1\}$*  is a subset  $T \subseteq \{0, 1\}^{<\omega}$  with the property that,  $a \in T$  for all  $a, b \in \{0, 1\}^{<\omega}$  such that  $a \subseteq b \in T$ . The *body* of a tree  $T$  on  $\{0, 1\}$ , denoted by  $[T]$ , is the set of all its

infinite branches, that is, the set of all sequences  $z \in \{0, 1\}^\omega$  such that  $z \upharpoonright n \in T$  for all  $n \in \omega$ .

A tree  $T$  on  $\{0, 1\}$  is said to be *pruned* if every  $a \in T$  has a proper extension, that is, for all  $a \in T$  there exists  $b \in T$  such that  $a \neq b$  and  $a \subseteq b$ , see [25, Definition 2.1]. Accordingly, we define

$$\text{PTr}_2 := \{T \subseteq \{0, 1\}^{<\omega} : T \text{ is a pruned tree}\}.$$

Identifying a pruned tree with its characteristic function, the set  $\text{PTr}_2$  can be regarded as a closed subset of the Polish space  $\{0, 1\}^{\{0, 1\}^{<\omega}}$ , see [25, Exercise 4.32]. Thanks to Alexandrov's theorem [25, Theorem 3.11],  $\text{PTr}_2$  is a Polish space.

With the above premises, let  $\text{WF}_2^*$  be the set of pruned trees on  $\{0, 1\}$  for which every infinite branch contains finitely many ones, that is,

$$\text{WF}_2^* := \{T \in \text{PTr}_2 : \forall z \in [T], \forall^\infty n, z_n = 0\},$$

see [25, Section 33.A]. It is known that  $\text{WF}_2^*$  is a  $\Pi_1^1$ -complete subset of  $\text{PTr}_2$ , hence it is conanalytic but not analytic, see [25, Exercise 27.3].

More generally, given a nonempty subset  $W$  of the Cantor space  $\{0, 1\}^\omega$ , let  $\mathcal{T}(W)$  be the set of pruned trees  $T$  on  $\{0, 1\}$  which contains an infinite branch in  $W$ , that is,

$$\mathcal{T}(W) := \{T \in \text{PTr}_2 : [T] \cap W \neq \emptyset\}.$$

It is immediate to check that  $\mathcal{T}(W_{\text{irr}})$  coincides with  $\text{WF}_2^*$ , where  $W_{\text{irr}}$  is the set of irrationals  $\{z \in \{0, 1\}^\omega : \exists^\infty n, z_n = 1\}$ .

With the above premises, we are ready for the proof of Theorem 4.7.

*Proof of Theorem 4.7.* We divide the proof in some intermediate steps. Hereafter, as in the statement of the result,  $W$  is a given Borel subset of the Cantor space which is not  $\Sigma_2^0$ .

**CLAIM 5.** Fix nonempty subsets  $A, B \subseteq \{0, 1\}^\omega$  such that  $A \leq_W B$ . Then

$$\mathcal{T}(A) \leq_W \mathcal{T}(B). \tag{10}$$

*Proof.* Since  $A$  is Wadge reducible to  $B$ , there exists a continuous map  $\phi : \{0, 1\}^\omega \rightarrow \{0, 1\}^\omega$  such that  $\phi^{-1}[B] = A$ , i.e.,  $z \in A$  if and only if  $\phi(z) \in B$ . Let  $\mathcal{K}$  be the family of compact subsets of  $\{0, 1\}^\omega$ , endowed with the Vietoris topology. It follows by [25, Exercise 4.32] that the map  $\psi : \text{PTr}_2 \rightarrow \mathcal{K}$  defined by

$$\forall T \in \text{PTr}_2, \quad \psi(T) := [T].$$

is a well-defined homeomorphism. In addition, its inverse map is given by  $\psi^{-1}(K) = \{z \upharpoonright n : z \in K, n \in \omega\}$  for all  $K \in \mathcal{K}$ . Thanks to [25, Exercise 4.29(vi)], the map  $\varphi : \mathcal{K} \rightarrow \mathcal{K}$  defined by  $\Phi(K) := \{\phi(z) : z \in K\}$  is continuous.

At this point, define the function  $f : \text{PTr}_2 \rightarrow \text{PTr}_2$  by

$$f := \psi^{-1} \circ \Phi \circ \psi,$$

so that  $f(T) = \{\phi(z) \upharpoonright n : z \in [T], n \in \omega\}$  for each pruned tree  $T$ . Now, it is sufficient to show that  $f$  is a witnessing map for the claimed Wadge reduction (10). For, note that  $f$  is a composition of three continuous maps, hence it is continuous. Moreover, the following chain of equivalences holds for each pruned tree  $T$  on  $\{0, 1\}$ :

$$\begin{aligned} T \in \mathcal{T}(A) & \quad \text{if and only if} \quad [T] \cap A = \emptyset, \\ & \quad \text{if and only if} \quad \Phi([T]) \cap B = \emptyset, \\ & \quad \text{if and only if} \quad [f(T)] \cap B = \emptyset, \\ & \quad \text{if and only if} \quad f(T) \in \mathcal{T}(B). \end{aligned}$$

Therefore  $f^{-1}[\mathcal{T}(B)] = \mathcal{T}(A)$ , which completes the proof.  $\square$

**CLAIM 6.**  $\text{WF}_2^* \leq_W \mathcal{T}(W)$ .

*Proof.* Thanks to [25, Theorem 22.10], the Borel set  $W$  is  $\Pi_2^0$ -hard, that is,  $A \leq_W W$  for all  $A \in \Pi_2^0$ . In particular,  $W_{\text{irr}} \leq_W W$ . Hence by Claim 5, we conclude that  $\text{WF}_2^* = \mathcal{T}(W_{\text{irr}}) \leq_W \mathcal{T}(W)$ .  $\square$

**CLAIM 7.**  $\mathcal{T}(W) \leq_W \mathcal{I}_W$ .

*Proof.* We need to show that there exists a continuous map  $h : \text{PTr}_2 \rightarrow \mathcal{P}(\omega)$  such that

$$T \in \mathcal{T}(W) \quad \text{if and only if} \quad h(T) \in \mathcal{I}_W$$

for all pruned trees  $T$  on  $\{0, 1\}$ .

First, define the map  $f : \{0, 1\}^{<\omega} \rightarrow \omega$  as follows: for each  $s = (s_0, \dots, s_k) \in \{0, 1\}^{<\omega}$ , let  $f(s)$  be the unique nonnegative integer such that

$$z_{f(s)} = (s_0, \dots, s_k, 1, 0, 0, \dots).$$

Of course,  $f$  is injective. Recalling that  $x_0 = (0, 0, \dots)$  is the unique sequence with empty support, it is easy to see the image of  $f$  is the set of positive integers.

At this point, define the map  $h : \text{PTr}_2 \rightarrow \mathcal{P}(\omega)$  by  $h(T) := \{f(s) : s \in T\}$  for all  $T \in \text{PTr}_2$ . Since  $T$  is an infinite and  $f$  is injective, then  $h(T)$  is infinite as well. It is also not difficult to show that  $h$  is continuous: indeed, a basic clopen set containing  $h(T)$  is of the type

$$V := \{S \in \omega : f(A) \subseteq S \text{ and } f(B) \cap S = \emptyset\}$$

for some finite sets  $A, B \subseteq \{0, 1\}^{<\omega}$  such that every restriction of each  $s \in A$  does not belong to  $B$ . Then, the set  $U := \{T \in \text{PTr}_2 : A \subseteq T \text{ and } B \cap T = \emptyset\}$  is an open set such that  $h[U] \subseteq V$ .

Next, we claim that

$$[T] = L_{\mathbf{x} \upharpoonright h(T)}. \quad (11)$$

On the one hand, fix  $a \in L_{\mathbf{x} \upharpoonright h(T)}$ . Then there exists a strictly increasing sequence of positive integers  $(p_k)$  such that  $\lim_k z_{p_k} = a$ . For each  $k \in \omega$ , set  $s^k := f^{-1}(p_k)$  and define  $\ell_k \in \omega$  so that  $s^k = (s_0^k, s_1^k, \dots, s_{\ell_k}^k)$ . Hence

$$z_{p_k} = z_{f(s^k)} = (s_0^k, s_1^k, \dots, s_{\ell_k}^k, 1, 0, 0, \dots).$$

Fix  $n \in \omega$ . Then there exists  $k_0 \in \omega$  such that  $a \upharpoonright n = z_{p_k} \upharpoonright n$  for all  $k \geq k_0$ . Since  $f$  is injective, there exists  $k_1 \geq \max\{n, k_0\}$  such that  $\ell_{k_1} \geq n$ . It follows that

$$a \upharpoonright n = z_{f(s^{k_1})} \upharpoonright n = (s_0^{k_1}, s_1^{k_1}, \dots, s_n^{k_1}) = s^{k_1} \upharpoonright n.$$

Since  $T$  is a tree and  $s^{k_1} \in T$ , we obtain  $a \upharpoonright n \in T$ . By the arbitrariness of  $n$ , it follows that  $a$  is an infinite branch of  $T$ . Therefore  $L_{\mathbf{x} \upharpoonright h(T)} \subseteq [T]$ .

On the other hand, fix  $a \in [T]$ , so that  $a \upharpoonright n \in T$  for all  $n \in \omega$ . It follows that  $f(a \upharpoonright n) \in h(T)$  and  $z_{f(a \upharpoonright n)} = (a_0, \dots, a_n, 1, 0, 0, \dots)$  for all  $n \in \omega$ . It is clear that  $\lim_n z_{f(a \upharpoonright n)} = a$ . Therefore the opposite inclusion  $[T] \subseteq L_{\mathbf{x} \upharpoonright h(T)}$  holds.

Recall that  $T \in \mathcal{T}(W)$  if and only if  $[T] \cap W = \emptyset$ . Thanks to (11), this is equivalent to  $L_{\mathbf{x} \upharpoonright h(T)} \cap W = \emptyset$ , that is,  $h(T) \in \mathcal{I}_W$ .  $\square$

To conclude the proof, thanks to Claim 6, Claim 7, and the transitivity of  $\leq_W$ , we get  $\text{WF}_2^* \leq_W \mathcal{I}_W$ . Hence there exists a continuous map  $g : \text{PTr}_2 \rightarrow \mathcal{P}(\omega)$  such that

$$g^{-1}[\mathcal{I}_W] = \text{WF}_2^*.$$

The conclusion follows by the facts that  $\text{WF}_2^*$  is  $\Pi_1^1$ -complete (hence, not analytic) and that analytic sets are closed under continuous preimages, see e.g. [31, Proposition 4.1.2].  $\square$

It is possible to show that  $\mathcal{I}_{W_{\text{irr}}}$  is coanalytic, hence  $\Pi_1^1$ -complete. We leave as open question for the interested reader to check whether  $\mathcal{I}_W$  is always coanalytic whenever  $W \subseteq \{0, 1\}^\omega$  is Borel and not  $\Sigma_2^0$ .

## 5. ACKNOWLEDGMENTS

The authors are grateful to Rafal Filipów and Adam Kwela (University of Gdansk, PL) for several discussions related to the results of the manuscript.

## REFERENCES

1. M. Balcerzak, S. Głąb, and P. Leonetti, *Another characterization of meager ideals*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **117** (2023), no. 2, Paper No. 90, 10.
2. M. Balcerzak and P. Leonetti, *On the relationship between ideal cluster points and ideal limit points*, Topology Appl. **252** (2019), 178–190.
3. P. Barbarski, R. Filipów, N. Mrożek, and P. Szuca, *When does the Katětov order imply that one ideal extends the other?*, Colloq. Math. **130** (2013), no. 1, 91–102.
4. J. Calbrix, *Classes de Baire et espaces d'applications continues*, C. R. Acad. Sci. Paris Sér. I Math. **301** (1985), no. 16, 759–762.
5. ———, *Filtres boréliens sur l'ensemble des entiers et espaces des applications continues*, Rev. Roumaine Math. Pures Appl. **33** (1988), no. 8, 655–661.
6. W. W. Comfort and S. Negrepontis, *The theory of ultrafilters*, Die Grundlehren der mathematischen Wissenschaften, Band 211, Springer-Verlag, New York-Heidelberg, 1974.
7. I. Farah, *Analytic quotients: theory of liftings for quotients over analytic ideals on the integers*, Mem. Amer. Math. Soc. **148** (2000), no. 702, xvi+177.
8. ———, *Luzin gaps*, Trans. Amer. Math. Soc. **356** (2004), no. 6, 2197–2239.
9. R. Filipów, K. Kowitz, and A. Kwela, *Characterizing existence of certain ultrafilters*, Ann. Pure Appl. Logic **173** (2022), no. 9, Paper No. 103157, 31.
10. R. Filipow, A. Kwela, and P. Leonetti, *Borel complexity of sets of ideal limit points*, manuscript.
11. D. H. Fremlin, *Measure theory. Vol. 2*, Torres Fremlin, Colchester, 2003, Broad foundations, Corrected second printing of the 2001 original.
12. D. H. Fremlin, R. A. Johnson, and E. Wajch, *Countable network weight and multiplication of Borel sets*, Proc. Amer. Math. Soc. **124** (1996), no. 9, 2897–2903.
13. J. A. Fridy, *Statistical limit points*, Proc. Amer. Math. Soc. **118** (1993), no. 4, 1187–1192.
14. S. Garcia-Ferreira and O. Guzmán, *More on MAD families and P-points*, Topology Appl. **305** (2022), Paper No. 107871, 10.
15. R. W. Hansell, J. E. Jayne, and C. A. Rogers, *K-analytic sets*, Mathematika **30** (1983), no. 2, 189–221 (1984).
16. X. He, H. Zhang, and S. Zhang, *The Borel complexity of ideal limit points*, Topology Appl. **312** (2022), Paper No. 108061, 12.
17. ———, *More on ideal limit points*, Topology Appl. **322** (2022), Paper No. 108324.

18. G. Hjorth and A. S. Kechris, *New dichotomies for Borel equivalence relations*, Bull. Symbolic Logic **3** (1997), no. 3, 329–346.
19. M. Hrušák, *Combinatorics of filters and ideals*, Set theory and its applications, Contemp. Math., vol. 533, Amer. Math. Soc., Providence, RI, 2011, pp. 29–69.
20. M. Hrušák and D. Meza-Alcántara, *Comparison game on Borel ideals*, Comment. Math. Univ. Carolin. **52** (2011), no. 2, 191–204.
21. M. Hrušák, D. Meza-Alcántara, E. Thümmel, and C. Uzcátegui, *Ramsey type properties of ideals*, Ann. Pure Appl. Logic **168** (2017), no. 11, 2022–2049.
22. M. Hrušák and J. L. Verner, *Adding ultrafilters by definable quotients*, Rend. Circ. Mat. Palermo (2) **60** (2011), no. 3, 445–454.
23. W. Just and A. Krawczyk, *On certain Boolean algebras  $\mathcal{P}(\omega)/I$* , Trans. Amer. Math. Soc. **285** (1984), no. 1, 411–429.
24. V. Kadets, D. Seliutin, and J. Tryba, *Conglomerated filters and statistical measures*, J. Math. Anal. Appl. **509** (2022), no. 1, Paper No. 125955, 17.
25. A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.
26. ———, *Rigidity properties of Borel ideals on the integers*, vol. 85, 1998, 8th Prague Topological Symposium on General Topology and Its Relations to Modern Analysis and Algebra (1996), pp. 195–205.
27. P. Leonetti and F. Maccheroni, *Characterizations of ideal cluster points*, Analysis (Berlin) **39** (2019), no. 1, 19–26.
28. D. Meza-Alcántara, *Ideals and filters on countable sets*, (2009), PhD Thesis, UNAM México.
29. C. A. Rogers, J. E. Jayne, C. Dellacherie, F. Topsoe, J. Hoffmann-Jorgensen, D. A. Martin, A. S. Kechris, and A. H. Stone, *Analytic sets*, Academic Press, London, 1980.
30. S. Solecki, *Analytic ideals*, Bull. Symbolic Logic **2** (1996), no. 3, 339–348.
31. S. M. Srivastava, *A course on Borel sets*, Graduate Texts in Mathematics, vol. 180, Springer-Verlag, New York, 1998.
32. M. Staniszewski, *On ideal equal convergence II*, J. Math. Anal. Appl. **451** (2017), no. 2, 1179–1197.
33. M. Talagrand, *Compacts de fonctions mesurables et filtres non mesurables*, Studia Math. **67** (1980), no. 1, 13–43.
34. F. van Engelen, *On Borel ideals*, Ann. Pure Appl. Logic **70** (1994), no. 2, 177–203.

INSTITUTE OF MATHEMATICS, LODZ UNIVERSITY OF TECHNOLOGY, AL. POLITECHNIKI 8,  
93-590 LODZ, POLAND

*Email address:* `marek.balcerzak@p.lodz.pl`

INSTITUTE OF MATHEMATICS, LODZ UNIVERSITY OF TECHNOLOGY, AL. POLITECHNIKI 8,  
93-590 LODZ, POLAND

*Email address:* `szymon.glab@p.lodz.pl`

DEPARTMENT OF ECONOMICS, UNIVERSITÀ DEGLI STUDI DELL'INSUBRIA, VIA MONTE  
GENEROSO 71, 21100 VARESE, ITALY

*Email address:* `leonetti.paolo@gmail.com`