Estimating invertible processes in Hilbert spaces, with applications to functional ARMA processes

Sebastian Kühnert^{*1}, Gregory Rice^{†2}, and Alexander Aue^{‡3}

¹Fakultät für Mathematik, Ruhr-Universität Bochum, Bochum, DE ²Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, CN ³Department of Statistics, University of California, Davis, Davis, US

July 16, 2024

Abstract

Invertible processes naturally arise in many aspects of functional time series analysis, and consistent estimation of the infinite dimensional operators that define them are of interest. Asymptotic upper bounds for the estimation error of such operators for processes in the Hilbert space $L^2[0,1]$ have been considered in recent years. This article adds to the theory in this area in several ways. We derive consistent estimates for the operators defining an invertible representation of a stationary process in a general separable Hilbert space under mild conditions that hold for many classes of functional time series. Moreover, based on these results, we derive consistency results with explicit rates for related operator estimates for Hilbert space-valued causal linear processes, as well as functional MA, AR and ARMA processes.

MSC 2020 subject classifications: 47B38, 60G10, 62F12 Keywords: ARMA; functional time series; linear processes; invertible processes

1 Introduction

Over the past three decades, the field *functional data analysis* (fDA) has experienced substantial growth, likely driven by increasing interest in examining high-dimensional data that arises from continuous observations across various domains, such as time, space, and frequency; see Horváth and Kokoszka (2012) and Ramsay and Silverman (2005) for text-book reviews of fDA. In many cases, functional dara are collected over time. For instance, one might gather continuous observations of electricity prices in a specific region and transform them into daily electricity price curves, see Liebl (2013). These collections of functional data objects are commonly known as *functional time series* (fTS) and there has been progress in developing methods for their analysis and modeling in recent years. For an overview of fTS analysis (fTSA), we refer to Bosq (2000)

^{*}sebastian.kuehnert@ruhr-uni-bochum.de

[†]grice@uwaterloo.ca

[‡]aaue@ucdavis.edu

and Kokoszka and Reimherr (2017). A typical and general setting for fTSA is to assume that the data attain their values in a separable, infinite dimensional Hilbert space \mathcal{H} , which we consider throughout this paper.

As is the case with scalar time series, the most oft used models in fTSA are linear models. These include \mathcal{H} -valued autoregressive (AR) processes, often described as functional autoregressive (fAR) process, and to a lesser extent in the literature \mathcal{H} -valued moving average (fMA) and fARMA processes. Such processes typically admit under regularity conditions stationary solutions that may be represented as \mathcal{H} -valued linear and invertible processes. For testing stationarity of general, fTS, we refer the reader to Horváth et al. (2014), Aue and van Delft (2020) and van Delft et al. (2021), and the monograph Bosq (2000) remains an excellent reference for linear processes in function spaces. Such processes also arise in the study of many non-linear fTS processes. For instance under regularity conditions the solutions for the volatility process in \mathcal{H} -valued versions of the functional (generalized) autoregressive conditionally heteroscedastic models (f(G)ARCH), see Hörmann et al. (2013), Aue et al. (2017) and Cerovecki et al. (2019), can be represented as (weak) \mathcal{H} -valued linear and invertible processes; see e.g. Kühnert (2020) where the processes attained values in the separable Hilbert space $\mathcal{H} = L^2[0,1]$ of squared Lebesgue-integrable functions with domain [0, 1].

When it comes to estimating the infinite dimensional parameters defining such processes, early references such as Bosq (2000) put forward consistent estimators of the operators defining fAR processes. Limit theorems for \mathcal{H} -valued linear processes can for instance be found in Merlevède (1996), Merlevède et al. (1997) and Düker (2018), and in Banach spaces in Račkauskas and Suguet (2010). Further, Aue and Klepsch (2017) and Klepsch and Klüppelberg (2017) approached the estimation of the operators in the linear and inverted representation pointwise, without stating explicit estimation rates. In Kühnert (2020), the complete sequences of operators in the linear and inverted representation of invertible linear processes originating from fGARCH models with values in the specific Hilbert space $L^{2}[0, 1]$. For fAR operators, Caponera and Panaretos (2022) provided explicit rates under quite mild conditions, and Kara Terki and Mourid (2024) derived both exponential bounds and convergence rates based on sieve estimates. The MA and ARMA case, though, has received substantially less attention in the literature. However, Turbillon et al. (2008) discussed estimating the fMA(1) operator under the quite strict condition that it commutes with the error processes covariance operator, and Kuenzer (2024) recently derived consistent estimators of the operators defining an H-valued ARMA process based on a two step approach involving sequentially esitmating the AR and MA components, and employing dimension reduction using an increasing number of principal components. The best achievable rate of estimation under fairly general conditions for the operator defining an fAR process of order one is Mas and Pumo (2009), who establish central limit theorems for such estimators.

This article formulates consistency results with quantifiable rates for the operators defining a general, \mathcal{H} -valued linear and invertible process based on functional Yule-Walker equations and a Tychonoff-regularized estimator, without assuming that the processes stem from specific models. These results are established under a Sobolev type condition and the well-known and practical weak dependence concept for fTSA called L^p -m-approximibility introduced in Hörmann and Kokoszka (2010). These results are then applied to obtain consistency results for the operators defining fAR, fMA and fARMA processes for arbitrary AR and MA orders.

The rest of the paper is organized as follows. Section 2 contains our notation and basic definitions and features in functional time series analysis. Section 3 establishes our operator estimates and asymptotic results in the infinite-dimensional setting. In Section 4, we demonstrate applications of these results to obtain consistent estimators of the operators in fARMA models. Section 5 summarizes the paper, and the proofs of all technical and auxiliary finite-dimensional results are collected in Appendices A-B.

Throughout this article, all asymptotic statements are meant for the sample size $N \to \infty$, unless otherwise noted.

2 Preliminaries

2.1 Notation

The additive identity of linear spaces is denoted by 0, and the identity map by I. On Cartesian product spaces V^n , with $n \in \mathbb{N}$, scalar multiplication and addition are defined component-wise. W let $(\mathcal{B}, \|\cdot\|)$ denote a Banach space, and $(\mathcal{H}, \langle \cdot, \cdot \rangle), (\mathcal{H}_\star, \langle \cdot, \cdot \rangle_\star)$ be Hilbert spaces with their respective norms and inner products. Then, \mathcal{B}^n is a Banach space and \mathcal{H}^n a Hilbert space, if they are endowed with the norm $\|x\|^2 = \sum_{i=1}^n \|x_i\|^2$ and the inner product $\langle x, y \rangle = \sum_{i=1}^n \langle x_i, y_i \rangle$, respectively, where $x \coloneqq (x_1, \ldots, x_n)^{\mathsf{T}}, y \coloneqq (y_1, \ldots, y_n)^{\mathsf{T}}$. By $\mathcal{L}_{\mathcal{H},\mathcal{H}_\star}, \mathcal{S}_{\mathcal{H},\mathcal{H}_\star} \subsetneq \mathcal{L}_{\mathcal{H},\mathcal{H}_\star}$, and $\mathcal{N}_{\mathcal{H},\mathcal{H}_\star} \subseteq \mathcal{S}_{\mathcal{H},\mathcal{H}_\star}$ we denote the spaces of linear, bounded; Hilbert-Schmidt (H-S); and nuclear (trace-class) operators $A: \mathcal{H} \to \mathcal{H}_\star$, respectively, with operator norm $\|\cdot\|_{\mathcal{L}}$; H-S inner product $\langle \cdot, \cdot \rangle_{\mathcal{S}}$ and H-S norm $\|\cdot\|_{\mathcal{S}}$; and nuclear norm $\|\cdot\|_{\mathcal{N}}$, respectively, with $\mathcal{T}_{\mathcal{H}} \coloneqq \mathcal{T}_{\mathcal{H},\mathcal{H}}$ for $\mathcal{T} \in \{\mathcal{L}, \mathcal{S}, \mathcal{N}\}$. For a given *complete orthonormal system* (CONS) $(a_j)_{j\in\mathbb{N}}$ of $\mathcal{H}, \prod_{a_m}^{a_n}: \mathcal{H} \to \text{span}\{a_m, a_{m+1}, \ldots, a_n\} \subset \mathcal{H}$, with n > m, denotes the projection operator onto the closed linear subspace spanned by a_m, \ldots, a_n . All random variables are defined on a common probability space $(\Omega, \mathfrak{A}, \mathbb{P})$. We write $X_n = o_{\mathbb{P}}(c_n)$ and $X_n = O_{\mathbb{P}}(c_n)$ (for $n \to \infty$) for X_n/c_n converging in probability and being stochastically bounded, respectively, for some sequence $(c_n)_{n\in\mathbb{N}} \subset (0,\infty)$. For $p \in [1,\infty), L_{\mathcal{H}}^p = L_{\mathcal{H}}^p(\Omega,\mathfrak{A},\mathbb{P})$ is the space of (classes of) random variables $X \in \mathcal{H}$ with $\mathbb{E}\|X\|^p < \infty, (X_k)_{k\in\mathbb{Z}}$ is an $L_{\mathcal{H}}^p$ -process if $(X_k)_{k\in\mathbb{Z}} \subset L_{\mathcal{H}}^p$, and centered if $\mathbb{E}(X_k) = 0$ for all k, with expectation understood as a Bochner-integral.

2.2 Stationarity and lagged (cross-)covariance operators

Let $(\mathcal{H}, \langle \cdot, \cdot \rangle), (\mathcal{H}_{\star}, \langle \cdot, \cdot \rangle_{\star})$ be separable Hilbert spaces with their respective inner products. The cross-covariance operator of $X \in L^2_{\mathcal{H}}$ and $Y \in L^2_{\mathcal{H}_{\star}}$ is defined by $\mathscr{C}_{X,Y} := \mathbb{E}[X - \mathbb{E}(X)] \otimes [Y - \mathbb{E}(Y)]$, where $x \otimes y := \langle x, \cdot \rangle y$ denotes the tensorial product of $x \in \mathcal{H}, y \in \mathcal{H}_{\star}$, and the covariance operator of X by $\mathscr{C}_X = \mathscr{C}_{X,X}$. If $\mathbb{E}||X||^2 < \infty$, $\mathscr{C}_X \in \mathcal{N}_{\mathcal{H}}$ is positive semi-definite and self-adjoint with $||\mathscr{C}_X||_{\mathcal{N}} = \mathbb{E}||X - \mathbb{E}(X)||^2$. Further, $\mathscr{C}_{X,Y} \in \mathcal{N}_{\mathcal{H},\mathcal{H}_{\star}}$, for the adjoint holds $\mathscr{C}^*_{X,Y} = \mathscr{C}_{Y,X} \in \mathcal{N}_{\mathcal{H}_{\star},\mathcal{H}}$, and $||\mathscr{C}_{X,Y}||_{\mathcal{N}} \leq \mathbb{E}||X - \mathbb{E}(X)||||Y - \mathbb{E}(Y)||_{\star}$.

A process $\mathbf{X} = (X_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ is *(strictly) stationary* if $(X_{t_1}, \ldots, X_{t_n}) \stackrel{d}{=} (X_{t_1+h}, \ldots, X_{t_n+h})$ for all $h, t_1, \ldots, t_n \in \mathbb{Z}, n \in \mathbb{N}$, and weakly stationary if it is an $L^2_{\mathcal{H}}$ -process with $\mathbb{E}(X_k) = \mu$ for all k for some $\mu \in \mathcal{H}$, and if $\mathscr{C}_{X_k, X_\ell} = \mathscr{C}_{X_{k+h}, X_{\ell+h}}$ for all $h, k, \ell \in \mathbb{Z}$. We call $(X_k)_{k \in \mathbb{Z}}$ a white noise *(WN)* if $(X_k)_{k \in \mathbb{Z}}$ is a centered, weakly stationary process with $\mathbb{E}||X_k||^2 > 0$ for all k, and $\mathscr{C}_{X_k, X_\ell} = 0$ for $k \neq l$. Provided the process $\mathbf{X} \subset L^2_{\mathcal{H}}$ is weakly stationary, the lag-h-covariance operators of \mathbf{X} are defined by $\mathscr{C}^h_{\mathbf{X}} \coloneqq \mathscr{C}_{X_0, X_h}$, for $h \in \mathbb{Z}$, where $(\mathscr{C}^h_{\mathbf{X}})^* = \mathscr{C}^{-h}_{\mathbf{X}}$. We call $\mathscr{C}_{\mathbf{X}} \coloneqq \mathscr{C}^0_{\mathbf{X}}$ the covariance operator of \mathbf{X} . If two processes $\mathbf{X} = (X_k)_{k \in \mathbb{Z}} \subset L^2_{\mathcal{H}}, \mathbf{Y} = (Y_k)_{k \in \mathbb{Z}} \subset L^2_{\mathcal{H}'}$ satisfy $\mathscr{C}_{X_k, Y_\ell} = \mathscr{C}_{X_{k+h}, Y_{\ell+h}}$ for all h, k, ℓ , the lag-h-cross-covariance operators of \mathbf{X}, \mathbf{Y} are defined by $\mathscr{C}^h_{\mathbf{X}, \mathbf{Y}} \coloneqq \mathscr{C}_{X_0, Y_h}$ for $h \in \mathbb{Z}$, where $(\mathscr{C}^h_{\mathbf{X}, \mathbf{Y}})^* = \mathscr{C}^{-h}_{\mathbf{Y}, \mathbf{X}}$. Further, we write $\mathscr{D}_{\mathbf{X}, \mathbf{Y}} \coloneqq \mathscr{C}^1_{\mathbf{X}, \mathbf{Y}}$.

3 Estimation of the operators in \mathcal{H} -valued linear and invertible processes

This section demonstrates an estimation procedure for operators of *invertible* processes $\mathbf{X} = (X_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ and derives asymptotic results. The use of these results for estimating the operators of invertible, causal *linear processes* is also discussed. Throughout we let $\boldsymbol{\varepsilon} = (\varepsilon_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ be a stationary and ergodic \mathcal{H} -valued WN.

3.1 Estimation in the inverted representation

We call a centered process $\mathbf{X} = (X_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ invertible if it is stationary and satisfies

$$X_k = \sum_{j=1}^{\infty} \psi_j(X_{k-j}) + \varepsilon_k, \quad k \in \mathbb{Z},$$
(1)

where $(\psi_j)_{j \in \mathbb{N}} \subset \mathcal{L}_{\mathcal{H}}$ is a sequence of linear operators. Stationary processes satisfying (1) arise in many different situations. Examples include invertible linear processes, which are discussed at length in Chapter 7 of Bosq (2000). This class of time series obviously also includes fAR processes, but also fMA and fARMA processes under natural conditions; see Spangenberg (2013) and Kuenzer (2024). Moreover, estimating the fAR(MA) parameters derived from invertible processes is also beneficial when dealing with f(G)ARCH processes, as point-wise squared (point-wise) f(G)ARCH models are fAR(MA) models, see Hörmann et al. (2013) and Kühnert (2020).

Assumption 3.1. $X \subset L^2_{\mathcal{H}}$ is a centered, invertible process with representation (1), and $\varepsilon \subset L^4_{\mathcal{H}}$.

Assumption 3.2. The covariance operator $\mathscr{C}_{\varepsilon} \colon \mathcal{H} \to \mathcal{H}$ is injective.

Remark 3.1. Injectivity of the covariance operator $\mathscr{C}_{\varepsilon}$ holds if and only if $\overline{\operatorname{im}}(\mathscr{C}_{\varepsilon}) = \mathcal{H}$, see *Hsing and Eubank (2015)*. It is also equivalent to the condition that there is no affine, closed, and proper subspace $U \subsetneq \mathcal{H}$ with $\mathbb{P}(\varepsilon_0 \in U) = 1$.

The goal of this section is to define, based on a sample X_1, \ldots, X_N from \mathbf{X} , consistent estimators of the operators $(\psi_j)_{j \in \mathbb{N}}$ in (1). The estimators we propose are based on estimates for lagged (cross-) covariance operators of processes

$$X_k^{[L]} \coloneqq (X_k, X_{k-1}, \dots, X_{k-L+1})^\top, \quad k \in \mathbb{Z},$$
(2)

taking value in the Cartesian product Hilbert space \mathcal{H}^L . We assume that the lag parameter $L = L_N \in \mathbb{N}$ grows as a function of the sample size. We define the empirical version $\hat{\mathscr{C}}_{\mathbf{X}^{[L]}}$ of the covariance operator $\mathscr{C}_{\mathbf{X}^{[L]}}$ through

$$\hat{\mathscr{C}}_{\mathbf{X}^{[L]}} \coloneqq \frac{1}{N-L+1} \sum_{k=L}^{N} X_k^{[L]} \otimes X_k^{[L]}, \quad L \le N.$$
(3)

We also define $\hat{\mathscr{D}}_{\mathbf{X}^{[L]},\mathbf{X}}$, the empirical version of $\mathscr{D}_{\mathbf{X}^{[L]},\mathbf{X}} = \mathscr{C}^{1}_{\mathbf{X}^{[L]},\mathbf{X}} \in \mathcal{N}_{\mathcal{H}^{L},\mathcal{H}}$, by

$$\hat{\mathscr{D}}_{\boldsymbol{X}^{[L]},\boldsymbol{X}} \coloneqq \frac{1}{N-L} \sum_{k=L}^{N-1} X_k^{[L]} \otimes X_{k+1}, \quad L < N.$$
(4)

For all $L, \mathscr{C}_{\mathbf{X}^{[L]}}$ and $\mathscr{D}_{\mathbf{X}^{[L]}, \mathbf{X}}$ are unbiased estimators for $\mathscr{C}_{\mathbf{X}^{[L]}}$ and $\mathscr{D}_{\mathbf{X}^{[L]}, \mathbf{X}}$, respectively. To derive consistency results for estimates of $(\psi_j)_{j \in \mathbb{N}}$ in (1), we assume that these operator estimators have consistency rates for a growing $L = L_N$ as follows.

Assumption 3.3. For some sequence $L = L_N \to \infty$ with $L = o(N^{1/3})$ holds

$$\|\hat{\mathscr{C}}_{\boldsymbol{X}^{[L]}} - \mathscr{C}_{\boldsymbol{X}^{[L]}}\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(L^{3/2}N^{-1/2}),\tag{5}$$

$$\|\widehat{\mathscr{D}}_{\mathbf{X}^{[L]},\mathbf{X}} - \mathscr{D}_{\mathbf{X}^{[L]},\mathbf{X}}\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(LN^{-1/2}).$$
(6)

- **Remark 3.2.** (a) Assumption 3.3 holds for centered, $L^4_{\mathcal{H}}$ -m-approximable processes, see Kühnert (2024), where the well-known concept $L^p_{\mathcal{H}}$ -m-approximability was introduced by Hörmann and Kokoszka (2010). For $p \geq 1$, a process $(X_k)_{k\in\mathbb{Z}} \subset L^p_{\mathcal{H}}$ is called $L^p_{\mathcal{H}}$ -m-approximable if for each k it holds $X_k = f(\epsilon_k, \epsilon_{k-1}, \ldots)$ for some i.i.d. process $(\epsilon_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ and a measurable function $f: \mathcal{H}^{\infty} \to \mathcal{H}$ (i.e. $(X_k)_k$ is causal (w.r.t. $(\epsilon_k)_k$)) such that $\sum_{m=1}^{\infty} \nu_p(X_m - X_m^{(m)}) < \infty$, where $\nu_p(\cdot) \coloneqq (\mathbb{E} \|\cdot\|^p)^{1/p}$, and where $X^{(m)}_k \coloneqq f(\epsilon_k, \epsilon_{k-1}, \ldots, \epsilon_{k-m+1}, \epsilon^{(m)}_{k-m}, \epsilon^{(m)}_{k-m-1}, \ldots)$ for all k, m, with $(\epsilon_j^{(m)})_j$ being independent copies of $(\epsilon_j)_j$ for all m. We would like to point out that Assumption 3.3 can also be satisfied for π -dependent processes, where π -dependence is a recently introduced weak dependence concept for metric space-valued processes by Kutta (2024), as L^p -m-approximibility implies π -dependence.
 - (b) Other centered processes that satisfy Assumption 3.3 are those that fulfill the rather strict summability condition in Kühnert (2022), and if they are i.i.d. with finite fourth moments, see, e.g. Horváth and Kokoszka (2012) for covariance operator estimation of i.i.d. processes.
 - (c) It is worthwhile to note that Assumption 3.3 can even be satisfied for not necessarily causal processes, for instance linear processes (LP) in H, that is

$$X_k = \sum_{\ell = -\infty}^{\infty} \phi_\ell(\varepsilon_{k-\ell}),\tag{7}$$

with $\sum_{\ell=-\infty}^{\infty} \|\phi_{\ell}\|_{\mathcal{L}} < \infty$. This is the case if the LP's operators fulfill the stronger condition $\sum_{k=1}^{\infty} \sum_{\ell=k}^{\infty} [\|\phi_{\ell}\|_{\mathcal{L}} + \|\phi_{-\ell}\|_{\mathcal{L}}] < \infty$.

We also make use of estimates of functional principal components. Before we do, we make several useful observations. The covariance operator of an LP can be described in terms of the linear operators in (15) and error covariance operator $\mathscr{C}_{\varepsilon}$ through

$$\mathscr{C}_{\boldsymbol{X}} = \sum_{i=0}^{\infty} \phi_i \mathscr{C}_{\boldsymbol{\varepsilon}} \phi_i^*.$$
(8)

The following are then immediate consequences of Assumptions 3.1–3.2. Now, let $(\lambda_j, c_j)_{j \in \mathbb{N}} = (\lambda_j(L), c_j(L))_{j \in \mathbb{N}}$, and $(\hat{\lambda}_j, \hat{c}_j)_{j \in \mathbb{N}} = (\hat{\lambda}_j(L), \hat{c}_j(L))_{j \in \mathbb{N}}$ be the sequence of eigenvalue/eigenfunction pairs of the covariance operators $\mathscr{C}_{\mathbf{X}^{[L]}}$, and $\hat{\mathscr{C}}_{\mathbf{X}^{[L]}} \in \mathcal{N}_{\mathcal{H}^L}$, respectively. According to Bosq (2000), Lemma 4.2, it holds that

$$\sup_{j\in\mathbb{N}} |\hat{\lambda}_j - \lambda_j| \le \|\hat{\mathscr{C}}_{\mathbf{X}^{[L]}} - \mathscr{C}_{\mathbf{X}^{[L]}}\|_{\mathcal{L}}.$$
(9)

Suitable estimators for the eigenfunctions c_j , which are unambiguous except for their sign, are $\hat{c}'_j := \operatorname{sgn}\langle \hat{c}_j, c_j \rangle \hat{c}_j$, with $\operatorname{sgn}(x) := \mathbb{1}_{[0,\infty)}(x) - \mathbb{1}_{(-\infty,0)}(x), x \in \mathbb{R}$. As covariance operators are positive semi-definite, injectivity of $\mathscr{C}_{\mathbf{X}^{[L]}} \in \mathcal{N}_{\mathcal{H}^L}$ after Proposition 3.1 is equivalent to strict positivity. Hence, the eigenvalues satisfy $\lambda_1 \geq \lambda_2 \geq \cdots > 0$. If the eigenspaces associated to each eigenvalues λ_j are one-dimensional, according to Bosq (2000), Lemma 4.3, it holds that

$$\|\hat{c}_j' - c_j\| \leq \frac{2\sqrt{2}}{\alpha_j} \|\hat{\mathscr{C}}_{\mathbf{X}^{[L]}} - \mathscr{C}_{\mathbf{X}^{[L]}}\|_{\mathcal{L}},$$

where $\alpha_1 \coloneqq \lambda_1 - \lambda_2$, and $\alpha_j \coloneqq \min(\lambda_{j-1} - \lambda_j, \lambda_j - \lambda_{j+1})$ for j > 1. Moreover, for all $k \in \mathbb{N}$ it holds with $\Lambda_k \coloneqq \sup_{1 \le j \le k} (\lambda_j - \lambda_{j+1})^{-1}$ that

$$\sup_{1 \le j \le k} \|\hat{c}_j' - c_j\| \le 2\sqrt{2} \Lambda_k \|\hat{\mathscr{C}}_{\boldsymbol{X}^{[L]}} - \mathscr{C}_{\boldsymbol{X}^{[L]}}\|_{\mathcal{L}}.$$
(10)

We note $\Lambda_k = (\lambda_k - \lambda_{k+1})^{-1}$ if there is a convex function $\kappa \colon \mathbb{R} \to \mathbb{R}$ with $\kappa(j) = \lambda_j$ for all j. We remind the reader that below we will be taking $L = L_N$ with $L_N \to \infty$, and $\Lambda_k = \Lambda_{k,N}$.

The estimates for the operators in the inverted representation are derived based on the approximate Yule-Walker equation

$$\mathscr{D}_{\boldsymbol{X}^{[L]},\boldsymbol{X}} = \Psi_L \mathscr{C}_{\boldsymbol{X}^{[L]}} + \sum_{\ell > L} \psi_\ell \mathscr{C}_{\boldsymbol{X}^{[L]},\boldsymbol{X}}^{1-\ell},$$
(11)

where the process $\mathbf{X}^{[L]} = (X_k^{[L]})_{k \in \mathbb{Z}} \subset \mathcal{H}^L$ is defined in (2), and the operator $\Psi_L \in \mathcal{L}_{\mathcal{H}^L, \mathcal{H}}$ is given by

$$\Psi_L \coloneqq (\psi_1 \cdots \psi_L). \tag{12}$$

Thereby, $\sum_{\ell>L} \psi_{\ell} \mathscr{C}_{\mathbf{X}^{[L]},\mathbf{X}}^{1-\ell}$ exists for all fixed $L \in \mathbb{N}$, as $\sum_{\ell\geq 1} \psi_{\ell} \mathscr{C}_{\mathbf{X}^{[L]},\mathbf{X}}^{1-\ell}$ converges in the operator norm topology under Assumption 3.1.

Based on a sample X_1, \ldots, X_N from X, we focus on estimating Ψ_L . This can be achieved by using Tychonoff regularization, see, e.g., Hall and Horowitz (2005), with an estimator $\hat{\Psi}_L = \hat{\Psi}_L(K, N)$ of the form

$$\hat{\Psi}_{L} \coloneqq \hat{\mathscr{D}}_{\boldsymbol{X}^{[L]},\boldsymbol{X}} \hat{\mathscr{C}}_{\boldsymbol{X}^{[L]}}^{\dagger} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} = \hat{\mathscr{D}}_{\boldsymbol{X}^{[L]},\boldsymbol{X}} (\hat{\mathscr{C}}_{\boldsymbol{X}^{[L]}} + \theta_{N} \mathbb{I})^{-1} \prod_{\hat{c}_{1}}^{\hat{c}_{K}},$$
(13)

where $\mathbb{I}: \mathcal{H}^L \to \mathcal{H}^L$ is the identity map, $\hat{\mathscr{C}}_{\mathbf{X}^{[L]}}$ and $\hat{\mathscr{D}}_{\mathbf{X}^{[L]},\mathbf{X}}$ are the estimates in (3)–(4), $K = K_N \in \mathbb{N}, (\theta_N)_{N \in \mathbb{N}} \subset (0, \infty)$ are tuning parameter sequences satisfying $K_N \to \infty, \theta_N \to 0$, and $\hat{c}_1, \ldots, \hat{c}_K$ are the eigenfunctions associated to the first K largest eigenvalues $\hat{\lambda}_1 \geq \cdots \geq \hat{\lambda}_K$ of $\hat{\mathscr{C}}_{\mathbf{X}^{[L]}}$.

We are able to establish the consistency of $\hat{\Psi}_L$ under the following assumptions.

Assumption 3.4. The eigenvalues $\lambda_j = \lambda_j(L)$ of $\mathscr{C}_{\mathbf{X}^{[L]}}$ satisfy $\lambda_j \neq \lambda_{j+1}$ for each $L = L_N$ and j. Assumption 3.5. Each of the operators in (1) satisfy that $\psi_i \in \mathcal{S}_{\mathcal{H}}$, and $\sum_{i=1}^{\infty} \|\psi_i\|_{\mathcal{S}}^2 < \infty$. Assumption 3.4, which implies that each eigenspace of the covariance operator $\mathscr{C}_{\mathbf{X}^{[L]}}$ is onedimensional, seems to be strict. For example when \mathbf{X} is an independent and identically distributed process the eigenvalues of $\mathscr{C}_{\mathbf{X}^{[L]}} = \text{diag}(\mathscr{C}_{\mathbf{X}}, \ldots, \mathscr{C}_{\mathbf{X}})$ have multiplicity at least L. However, for nondegenerate linear processes in which at least some of the operators ϕ_j in (15) are non-zero, this condition is more plausible. This condition is weaker than many related works in which a specific form of the eigenvalue gaps $f(j) = \lambda_j - \lambda_{j-1}$ are prescribed; see for instance the equation 3.2 in Hall and Horowitz (2007).

The method we use to quantify the error between $\hat{\Psi}_L$ and Ψ_L entails first decomposing Ψ_L into a suitable finite dimensional representation and a remainder term, and then evaluating the error of $\hat{\Psi}_L$ in estimating each component. In order to quantify the estimation of the remainder term, we employ a Sobolev condition akin to that used in Hall and Meister (2007) to derive precise asymptotic results for the estimation errors of densities in the deconvolution setting. To state our Sobolev type condition, we let $(\mu_j, d_j)_{j \in \mathbb{N}}$ denote the eigenvalue/eigenfunction pairs corresponding to the covariance operator $\mathscr{C}_{\mathbf{X}} \in \mathcal{N}_{\mathcal{H}}$. Then, $(c_i \otimes d_j)_{i,j \in \mathbb{N}} = (c_i(L) \otimes d_j)_{i,j \in \mathbb{N}}$ is a CONS of $\mathcal{S}_{\mathcal{H}^L,\mathcal{H}}$, so, if $\Psi_L \in \mathcal{S}_{\mathcal{H}^L,\mathcal{H}}$, which holds if the ψ_i are H-S operators, $\Psi_L = \sum_i \sum_j \langle \Psi_L, c_i \otimes d_j \rangle_{\mathcal{S}} (c_i \otimes d_j)$.

Assumption 3.6. For some $\beta > 0$, the limit $S_{\Psi}(\beta) \coloneqq \lim_{N \to \infty} S_{\Psi_{L_N}}(\beta) < \infty$ exists, where for each $L = L_N \in \mathbb{N}$ holds,

$$S_{\Psi_L}(\beta) = S_{\Psi_L}(\beta, (c_i \otimes d_j)_{i,j}) = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \langle \Psi_L, c_i \otimes d_j \rangle_{\mathcal{S}}^2 \left(1 + i^{2\beta} + j^{2\beta}\right) < \infty.$$

Theorem 3.1. Let Assumptions 3.1–3.6 hold, and let $K = K_N \to \infty$ and $\theta_N \to 0$ be further sequences with $\theta_N = o(\lambda_K)$ and also

$$K^{1+\beta}\lambda_{K}^{-1}\Lambda_{K}L^{2} = \mathcal{O}(N^{1/2}), \quad \sum_{\ell > L} \|\psi_{\ell}\|_{\mathcal{L}} = \mathcal{O}(K\Lambda_{K}L^{3/2}N^{-1/2}), \quad \theta_{N} = \mathcal{O}(K\Lambda_{K}L^{2}N^{-1/2}),$$

where $\beta > 0$ is defined in Assumption 3.6. Then, we have

$$\|\hat{\Psi}_L - \Psi_L\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(K^{-\beta}).$$
(14)

Remark 3.3. (a) Theorem 3.1 improves on comparable results in the literature in several ways. In contrast to Kühnert (2020), our result is stated for general, invertible linear processes without assuming that they originate from a specific model, and where the processes are allowed to attain values in arbitrary, separable Hilbert space rather than only in L²[0,1]. Most notably, we analyze the complete operators and state explicit asymptotic upper bounds, rather than either focusing on finite-dimensional projections which is commonly done in operator estimation, see Hörmann et al. (2013) and Aue et al. (2017), or omitting stating the rate when completely observing the operators, see Cerovecki et al. (2019) for fGARCH models, and Aue and Klepsch (2017) and Klepsch and Klüppelberg (2017) for invertible linear processes. Further, we measured the estimation errors even with the H-S- instead of the weaker operator norm as in Aue and Klepsch (2017) and Klepsch and Klüppelberg (2017). The Sobolev type condition we utilized in our estimation, is, as far as we are aware, in the context of operator estimation only used for fGARCH models in Kühnert (2020). Moreover, we refrained from a simplifying convexity assumption for the eigenvalues.

- (b) Although Λ_K is defined via the eigenvalues λ_K , we can only simplify the sequences' requirements in Theorem 3.1 (and in other results) in certain situations. For example, if for each L there are convex functions $\kappa_L \colon \mathbb{R} \to \mathbb{R}$ so that for the eigenvalues holds $\kappa_L(j) = \lambda_j(L)$ for all j, and if $\lambda_j(L)$ decays for $j \to \infty$ exponentially fast for each L, and if $\lambda_j(L) = c_L e^{-j}$ for some constant $c_L > 0$. Then, we obtain the explicit form $\Lambda_K = (\lambda_K - \lambda_{K+1})^{-1} = d_L \lambda_K^{-1}$, with $d_L := (c_L(1 - e^{-1}))^{-1} > 0$.
- (c) By following the lines of the proof of Theorem 3.1 (and all other asymptotic results), it seems like our result could have been even formulated in sense of L^2 -convergence instead of stochastic boundedness with a certain rate, but our inequalities contain reciprocals of eigenvalue estimates which only converge in the stochastic sense.

3.2 Estimating the operators in the causal linear process representation

We now suppose $\mathbf{X} = (X_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ follows a causal LP, see (7) for general LP's, so

$$X_k = \varepsilon_k + \sum_{j=1}^{\infty} \phi_j(\varepsilon_{k-j}), \quad k \in \mathbb{Z},$$
(15)

where $(\phi_j)_{j \in \mathbb{N}} \subset \mathcal{L}_{\mathcal{H}}$ are linear operators with $\sum_{j=1}^{\infty} \|\phi_j\|_{\mathcal{L}} < \infty$. Such a LP is invertible if it may be expressed as in (1). The monograph Bosq (2000) was devoted to this class of processes in its entirety. For general existence theorems of LPs and their invertibility, see Section 7 of this book. We now take up the estimation of the operators ϕ_i in (15) of an invertible LP. Estimating these operators are of use in deriving prediction and confidence sets when applying \mathcal{H} -valued LP models.

Relating the estimates from an inverted representation of X to its causal linear representation can be done using the following proposition.

Proposition 3.1. Let Assumptions 3.1–3.2 hold. Then

- (a) The covariance operator $\mathscr{C}_{\mathbf{X}} : \mathcal{H} \to \mathcal{H}$ is injective.
- (b) The covariance operators $\mathscr{C}_{\mathbf{X}^{[L]}} \colon \mathcal{H}^L \to \mathcal{H}^L$ are injective for all L.
- (c) If $\sum_{\ell=0}^{\infty} A_{\ell}(\varepsilon_{i-\ell}) = \sum_{\ell=0}^{\infty} B_{\ell}(\varepsilon_{i-\ell})$ for some $i \in \mathbb{N}$, then, $A_{\ell} = B_{\ell}$ for all $\ell \in \mathbb{N}$.
- (d) If \mathbf{X} is a causal linear process, and if $\sum_{\ell=0}^{\infty} A_{\ell}(X_{i-\ell}) = \sum_{\ell=0}^{\infty} B_{\ell}(X_{i-\ell})$ for some $i \in \mathbb{N}$, then $A_{\ell} = B_{\ell}$ for all $\ell \in \{0, 1, 2, \}$.

For an invertible LP X in Assumption 3.1 such that

$$X_k = \varepsilon_k + \sum_{i=1}^{\infty} \psi_i(X_{k-i}), \text{ and } X_{k-i} = \varepsilon_{k-i} + \sum_{j=1}^{\infty} \phi_j(\varepsilon_{k-i-j}),$$

for all $k, i \in \mathbb{Z}$, we have with $\phi_0 \coloneqq \mathbb{I}$ that

$$X_k = \varepsilon_k + \sum_{i=1}^{\infty} \left[\sum_{j=1}^{i} \psi_j \phi_{i-j} \right] (\varepsilon_{k-i}), \quad k \in \mathbb{Z}.$$

This representation, the linear representation (15), Assumption 3.2 and Proposition 3.1 (c) yield

$$\phi_i = \sum_{j=1}^i \psi_j \phi_{i-j}, \quad i \in \mathbb{N}.$$

Hence, with $\hat{\psi}_j = \hat{\Psi}_L^{(j)}$ denoting the *j*'th component of the estimator $\hat{\Psi}_L$ in (13) for $\Psi_L = (\psi_1, \ldots, \psi_L)$, a reasonable estimate for ϕ_i is $\hat{\phi}_i = \hat{\phi}_i(L, K)$, defined iteratively with $\hat{\phi}_0 \coloneqq \mathbb{I}$, and

$$\hat{\phi}_i \coloneqq \sum_{j=1}^i \hat{\psi}_j \hat{\phi}_{i-j}, \quad i \in \mathbb{N}.$$
(16)

By assuming that the operators satisfy a Sobolev condition, we also obtain the following consistency result for the complete operators in the linear representation.

Proposition 3.2. Under the assumptions of Theorem 3.1 holds

$$\|\hat{\phi}_i - \phi_i\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(K^{-\beta}), \quad i \in \mathbb{N}.$$

Remark 3.4. The novelties of Proposition 3.2 are similar to those in Theorem 3.1, see Remark 3.3 (a). Here, too, in the linear representation, the processes can attain their values in a general, separable Hilbert space, they are not constrained to stem from a certain type of time series, and we estimated the complete operators with stating an explicit asymptotic upper bound for the estimation errors in the H-S norm. Such a result is to the best of our knowledge new. However, consistent estimates for the operators in the linear representation, measured in the operator norm, and without convergence rate, can be found in Aue and Klepsch (2017) and Klepsch and Klüppelberg (2017).

4 Consequences for fARMA processes

The asymptotic results for the operator estimators in the linear and in the invertible representations in Section 3 can be applied to estimate the parameters of many linear functional time series processes with stationary and WN innovations, including fAR, fMA, and fARMA processes. Such processes are thoroughly discussed in Bosq (2000), where the focus is mainly on fAR processes. Spangenberg (2013) establishes general sufficient conditions under which fARMA processes in Banach spaces have causal and invertible linear process representations as in equations (15) and (1). Further, Klepsch et al. (2017) discusses prediction methods for fARMA(p,q) processes, and Kuenzer (2024) derived consistent operator estimates for fMA(1) and fARMA(1,1) processes based on the Hannan–Rissanen method, and making using of functional principal component analysis. Herein, by making use of operator estimates for the operators defining fAR, fMA and fARMA processes, with explicit asymptotic upper bounds.

4.1 fAR processes

We start by demonstrating the implications of our general results to fAR processes. Formally, a centered process $(X_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is a functional autoregressive process with order $p \in \mathbb{N}$ (fAR(p)) if

$$X_k = \varepsilon_k + \sum_{i=1}^p \alpha_i(X_{k-i}) \text{ a.s.}, \quad k \in \mathbb{Z},$$
(17)

where $(\varepsilon_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is a WN, and where $\alpha_i \in \mathcal{L}_{\mathcal{H}}$ are operators with $\alpha_p \neq 0$. With

$$A_p = (\alpha_1 \cdots \alpha_p) \in \mathcal{L}_{\mathcal{H}^p, \mathcal{H}},$$

and Ψ_p in (12), we have $A_p = \Psi_p$, and the exact Yule-Walker equation $\mathscr{C}^1_{X^{[p]},X} = A_p \mathscr{C}_{X^{[p]}}$. Then,

$$\hat{A}_p \coloneqq \hat{\Psi}_p$$

with $\hat{\Psi}_p = \hat{\Psi}_{p,K}$ being the estimator in (13) for L = p, and some $K \in \mathbb{N}$, and with $\hat{\alpha}_i \coloneqq \hat{A}_{p,K}^{(i)}$, $i \in \{1, \ldots, p\}$ being its components, we obtain the following corollary of Theorem 3.1.

Corollary 4.1. Let $(X_k)_{k \in \mathbb{Z}} \subset \mathcal{H}$ be the fAR(p) process in (17), and suppose the conditions of Theorem 3.1 are satisfied. Then, we have

$$\|\hat{\alpha}_i - \alpha_i\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(K^{-\beta}), \quad i = 1, \dots, p.$$
(18)

Remark 4.1. To the best of our knowledge, the result in Corollary 4.1 which states explicit asymptotic rates for the estimation errors for the operators of fAR processes in general, separable Hilbert space for any fAR order, and where the errors are only required to be stationary, ergodic white noises instead of i.i.d., is new. Explicit rates for fAR(1) processes with i.i.d. errors can also be found in Caponera and Panaretos (2022), where their results require a specific relationship between the lag-1-cross covariance to the covariance operator. Further, Bosq (2000) states consistency results for the fAR(1) operator projected on a finite-dimensional sup-space, and also the complete operator without stating an explicit rate.

4.2 fMA processes

Formally, a centered process $(X_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is a functional moving-average process with order $q \in \mathbb{N}$ (fMA(q)) if

$$X_k = \varepsilon_k + \sum_{j=1}^q \beta_j(\varepsilon_{k-j}) \text{ a.s.}, \quad k \in \mathbb{Z},$$

where $(\varepsilon_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is a WN, and where $\beta_j \in \mathcal{L}_{\mathcal{H}}$ are operators with $\beta_q \neq 0$. By putting

$$\hat{\beta}_j = \hat{\phi}_j,\tag{19}$$

with $\hat{\phi}_j$ in (16), we get the following result.

Corollary 4.2. Let $(X_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ be the fMA(q) process in (19) with $q \in \mathbb{N}$, and suppose the conditions of Theorem 3.1 are satisfied. Then,

$$\|\hat{\beta}_j - \beta_j\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(K^{-\beta}), \quad j = 1, \dots, q.$$

Remark 4.2. As far as we are aware, explicit asymptotic rates for the estimation errors for the operators of fMA processes in general, separable Hilbert spaces for any order as formulated in Corollary 4.2 have not been stated in the literature so far. Nonetheless, Turbillon et al. (2008) already estimated the fMA(1) operator under the very limiting condition that the fMA operator and the error processes' covariance operator commute. Moreover, for a simulation study for fMA(1) processes, see Kuenzer (2024).

4.3 fARMA processes

We finally turn to the estimation of fARMA processes. A centered process $(X_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is called a functional autoregressive moving-average process with orders $p, q \in \mathbb{N}$ (fARMA(p, q)) if

$$X_k = \varepsilon_k + \sum_{i=1}^p \alpha_i(X_{k-i}) + \sum_{j=1}^q \beta_j(\varepsilon_{k-j}) \text{ a.s.}, \quad k \in \mathbb{Z},$$
(20)

where $(\varepsilon_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ is a WN, and $\alpha_i, \beta_j \in \mathcal{L}_{\mathcal{H}}$ are operators with $\alpha_p \neq 0, \beta_q \neq 0$. We combine the fARMA(p,q) representation of the process $(X_k)_{k\in\mathbb{Z}}$ in (20) and its inverted representation (1). We note that, as with multivariate ARMA processes, in general the operators α_i and β_j are not identifiable. Sufficient conditions for the identifiability of fARMA operators are put forward in equation (10) and Propositions 1-3 of Kuenzer (2024). Spangenberg (2013) establishes conditions under which such processes may be represented using (15) and (1). We assume below that the fARMA models that we consider are identifiable and admit solutions that are ISLPs.

4.3.1 fARMA(1,1) processes

To describe the main idea, we first discuss estimation for fARMA(1,1) processes of the form

$$X_k = \varepsilon_k + \alpha_1(X_{k-1}) + \beta_1(\varepsilon_{k-1}) \text{ a.s.}, \quad k \in \mathbb{Z}.$$
 (21)

From this identity, Proposition 3.1(c,d) and $\varepsilon_{k-1} = X_{k-1} - \sum_{j=1}^{\infty} \psi_j(X_{k-1-j})$ it follows with $\psi_0 = \mathbb{I}$,

$$\psi_i = \begin{cases} \alpha_1 + \beta_1, & \text{if } i = 1, \\ -\beta_1 \psi_{i-1}, & \text{if } i > 1. \end{cases}$$
(22)

Although we may use the estimator $\hat{\Psi}_L$ in (13) to asymptotically estimate all operators ψ_i consistently, we cannot immediately obtain consistent estimates for the ARMA(1,1) operators, as $\psi_1 \in S_{\mathcal{H}}$ does not have a bounded inverse. However, with $\hat{\psi}_i = \hat{\Psi}_L^{(i)}$ being the *i*'th component of the estimator $\hat{\Psi}_L$ in (13), equation (22) suggests the following estimators $\hat{\alpha}_1 = \hat{\alpha}_1(L, K, M)$ for α_1 and $\hat{\beta}_1 = \hat{\beta}_1(L, K, M)$ for β_1 :

$$\hat{\alpha}_{1} \coloneqq \hat{\psi}_{1} - \hat{\beta}_{1}, \qquad \hat{\beta}_{1} \coloneqq -\hat{\psi}_{2} \hat{\psi}_{1}^{\dagger} \prod_{\hat{f}_{1}}^{\hat{f}_{M}} = -\hat{\psi}_{2} \hat{\psi}_{1}^{*} (\hat{\psi}_{1} \hat{\psi}_{1}^{*} + \gamma_{N} \mathbb{I})^{-1} \prod_{\hat{f}_{1}}^{\hat{f}_{M}}, \qquad (23)$$

where $(K_N)_N, (L_N)_N, (M_N)_N \subset \mathbb{N}, (\gamma_N)_{N \in \mathbb{N}} \subset (0, \infty)$ are tuning parameter sequences such that $\min\{K_N, L_N, M_N\} \to \infty$, and $\gamma_N \to 0$ as $N \to \infty$, $\hat{f}_1, \ldots, \hat{f}_M$ are the eigenfunctions associated to the eigenvalues $\hat{\rho}_1 \geq \cdots \geq \hat{\rho}_M \geq 0$ of $\hat{\psi}_1 \hat{\psi}_1^* \colon \mathcal{H} \to \mathcal{H}$, and f_1, \ldots, f_M are the eigenfunctions associated to the eigenvalues $\rho_1 \geq \cdots \geq \rho_M \geq 0$ of $\psi_1 \psi_1^* \colon \mathcal{H} \to \mathcal{H}$. These estimators are consistent under the following assumptions.

Assumption 4.1. $\alpha_1, \beta_1 \in S_H$ are H-S operators.

Assumption 4.2. The image of the operator $\psi_1 : \mathcal{H} \to \mathcal{H}$ lies dense.

Assumption 4.3. The eigenvalues of $\psi_1\psi_1^*: \mathcal{H} \to \mathcal{H}$ satisfy $\rho_j \neq \rho_{j+1}$ for all $j \in \mathbb{N}$.

Assumption 4.4. For some $\gamma > 0$ holds

$$\sum_{i=1}^{\infty}\sum_{j=1}^{\infty}\langle\beta_1, f_i\otimes f_j\rangle_{\mathcal{S}}^2(1+i^{2\gamma}+j^{2\gamma})<\infty.$$

Theorem 4.1. Let the assumptions of Theorem 3.1, Assumptions 4.1–4.4, and $M^{1+\gamma}\rho_M^{-1}P_M = O(K^{\beta})$ and $\gamma_N = o(\rho_M M^{-\gamma})$ hold. Then, the fARMA(1,1) operators satisfy

$$\max\{\|\hat{\alpha}_1 - \alpha_1\|_{\mathcal{S}}, \|\hat{\beta}_1 - \beta_1\|_{\mathcal{S}}\} = \mathcal{O}_{\mathbb{P}}(M^{-\gamma}).$$

- **Remark 4.3.** (a) Rather than using a Sobolev condition and Tychonoff regularization in Theorem 4.1 twice, Kuenzer (2024) derived consistent estimates for the fARMA(1,1) operators based on the Hannan-Rissanen method and multiple other technical conditions.
 - (b) Assumptions 4.1, 4.3–4.4 are of the same type as the assumptions for the operators in the inverted representation in Theorem 3.1. The additional listed Assumption 4.2 is needed so that identifiability of β_1 we estimate based on the identity $\psi_2 = -\beta_1\psi_1$ in (22) is guaranteed.

4.3.2 fARMA(p,q) processes

The procedure for centered fARMA(p,q) processes $(X_k)_{k\in\mathbb{Z}} \subset \mathcal{H}$ in (20) for arbitrary p,q is similar as for fARMA(1,1) processes. Similar to the transformations that led to (22), we obtain for any pand q, with $\alpha_i = \beta_j = 0$ for i > p, j > q,

$$\sum_{i=1}^{\infty} \psi_i(X_{k-i}) = \sum_{i=1}^{\max(p,q)} \left[\alpha_i + \beta_i - \sum_{j=1}^{i-1} \beta_j \psi_{i-j} \right] (X_{k-i}) - \sum_{i=\max(p,q)+1}^{\infty} \left[\sum_{j=1}^q \beta_j \psi_{i-j} \right] (X_{k-i}).$$

Comparing the left and right hand sides of the above and applying Proposition 3.1 (d), we see that

$$\psi_{i} = \begin{cases} \alpha_{i} + \beta_{i} - \sum_{j=1}^{i-1} \beta_{j} \psi_{i-j}, & \text{if } 1 \le i \le \max(p,q), \\ -\sum_{j=1}^{q} \beta_{j} \psi_{i-j}, & \text{if } i > \max(p,q). \end{cases}$$
(24)

We proceed to obtain estimates for β_i by using (24) for $i > \max(p,q)$, and then subsequently obtaining estimates for the α_i . Since the right hand side of (24) involves sums of compositions of the β_j with ψ_{i-j} for $i > \max(p,q)$, we cannot immediately retrieve estimates for β_j by applying Tychonoff regularized versions of the estimates ψ_{i-j} . For i = p + q, with

$$B_q \coloneqq (\beta_1 \cdots \beta_q) \in \mathcal{L}_{\mathcal{H}^q, \mathcal{H}},$$

$$\Psi'_{[i]} = \Psi'_{[i]}(p, q) \coloneqq (\psi_{p+q+i-1} \cdots \psi_{p+i}) \in \mathcal{L}_{\mathcal{H}^q, \mathcal{H}}, \quad i \ge 0,$$
(25)

the identity (24) becomes $\psi_{p+q} = -B_q \Psi_{[0]}^{\prime \top}$. Identifiability of B_q , and thus of β_1, \ldots, β_q , are given if the image of $\Psi_{[0]}^{\prime \top} \in \mathcal{L}_{\mathcal{H},\mathcal{H}^q}$ lies dense. Identifiability of B_q can be established using the relationship

$$\Psi'_{[q]} = -B_q \prod , \qquad (26)$$

where $\prod = \prod(p,q) \in \mathcal{L}_{\mathcal{H}^q}$ is the operator-valued matrix defined by

$$\prod \coloneqq \begin{bmatrix} \Psi'_{[q-1]} \\ \Psi'_{[q-2]} \\ \vdots \\ \Psi'_{[0]} \end{bmatrix} = \begin{bmatrix} \psi_{p+2q-2} & \psi_{p+2q-3} & \cdots & \psi_{p+q-1} \\ \psi_{p+2q-3} & \psi_{p+2q-4} & \cdots & \psi_{p+q-2} \\ \vdots & \vdots & \cdots & \vdots \\ \psi_{p+q-1} & \psi_{p+q-2} & \cdots & \psi_{p} \end{bmatrix}.$$
(27)

For $\Psi'_{[i]}$ and $\prod = \prod(p,q)$, we use the estimates $\hat{\Psi}'_{[i]} = \hat{\Psi}'_{[i]}(L,K)$ and $\hat{\prod} = \hat{\prod}(p,q;L,K)$, respectively, defined by

$$\hat{\Psi}'_{[i]} \coloneqq (\hat{\psi}_{p+q+i-1} \cdots \hat{\psi}_{p+i}) \in \mathcal{L}_{\mathcal{H}^q, \mathcal{H}}, \quad i \ge 0, \text{ and}$$

$$(28)$$

$$\hat{\Pi} \coloneqq \begin{bmatrix} \Psi'_{[q-1]} \\ \hat{\Psi}'_{[q-2]} \\ \vdots \\ \hat{\Psi}'_{[0]} \end{bmatrix} = \begin{bmatrix} \hat{\psi}_{p+2q-2} & \hat{\psi}_{p+2q-3} & \cdots & \hat{\psi}_{p+q-1} \\ \hat{\psi}_{p+2q-3} & \hat{\psi}_{p+2q-4} & \cdots & \hat{\psi}_{p+q-2} \\ \vdots & \vdots & \cdots & \vdots \\ \hat{\psi}_{p+q-1} & \hat{\psi}_{p+q-2} & \cdots & \hat{\psi}_{p} \end{bmatrix},$$
(29)

where $\hat{\psi}_i = \hat{\Psi}_{L,K}^{(i)}$ denotes the *i*th component of $\hat{\Psi}_L$ in (13). Due to (37), the operator-valued vectors $\Psi'_{[i]}, \hat{\Psi}'_{[i]}$ and matrices $\prod, \hat{\prod}$ are under Assumption 3.5 elements of $\mathcal{S}_{\mathcal{H}^q,\mathcal{H}}$ and $\mathcal{S}_{\mathcal{H}^q}$, respectively, thus $\prod\prod^*, \hat{\prod} \hat{\prod}^* \in \mathcal{N}_{\mathcal{H}^q}$ are nuclear. As a result of (24) and (26), the estimators $\hat{\alpha}_i = \hat{\alpha}_i(L, K, M)$ and $\hat{\beta}_j = \hat{\beta}_j(L, K, M)$ for α_1 and β_1 , respectively, are

$$\hat{\alpha}_{i} \coloneqq \hat{\psi}_{i} + \hat{B}_{[i]} \hat{\Psi}_{[i]}^{\prime\prime\prime} , \ 1 \le i \le p, \qquad \hat{\beta}_{j} \coloneqq \begin{cases} \hat{B}_{q}^{(j)}, & \text{if } 1 \le j \le q, \\ 0, & \text{if } j > q. \end{cases}$$
(30)

In these definitions, $\hat{B}_q = \hat{B}_q(L, K, M)$ stands for the estimator for B_q defined by

$$\hat{B}_q \coloneqq -\hat{\Psi}'_{[q]} \hat{\Pi}^{\dagger} \prod_{\hat{h}_1}^{\hat{h}_M} = -\hat{\Psi}'_{[q]} \hat{\Pi}^{*} \left(\hat{\Pi} \hat{\Pi}^{*} + \gamma_N \mathbb{I} \right)^{-1} \prod_{\hat{h}_1}^{\hat{h}_M},$$

where $(K_N)_N, (L_N)_N, (M_N)_N \subset \mathbb{N}, (\gamma_N)_{N \in \mathbb{N}} \subset (0, \infty)$ are sequences with $\min\{K_N, L_N, M_N\} \to \infty$ and $\gamma_N \to 0$. Further, $\hat{h}_1, \ldots, \hat{h}_M$ and h_1, \ldots, h_M are the eigenfunctions associated to the eigenvalues $\hat{\zeta}_1 \geq \cdots \geq \hat{\zeta}_M \geq 0$ and $\zeta_1 \geq \cdots \geq \zeta_M \geq 0$ of $\prod \prod^* : \mathcal{H}^q \to \mathcal{H}^q$ and $\prod \prod^* : \mathcal{H}^q \to \mathcal{H}^q$, respectively, and $(h_i \otimes d_j)_{i,j}$ is a CONS of $\mathcal{S}_{\mathcal{H}^q,\mathcal{H}}$, where d_j are the eigenfunctions of $\mathscr{C}_{\mathbf{X}}$. Further, we define $\hat{B}_{[i]} = \hat{B}_{[i]}(q; L, K), B_{[i]} = B_{[i]}(q), \hat{\Psi}_{[i]}^{'''} = \hat{\Psi}_{[i]}^{'''}(p, q; L, K), \Psi_{[i]}^{'''} = \Psi_{[i]}^{'''}(p, q) \in \mathcal{S}_{\mathcal{H}^{\max(i,q)},\mathcal{H}}$ for any i, q by

$$\hat{B}_{[i]} \coloneqq \begin{cases} \hat{B}_i, & \text{if } 1 \le i < q, \\ \hat{B}_q, & \text{if } i \ge q, \end{cases} \quad \text{and} \quad B_{[i]} \coloneqq \begin{cases} B_i, & \text{if } 1 \le i < q, \\ B_q, & \text{if } i \ge q, \end{cases}$$
(31)

$$\hat{\Psi}_{[i]}^{\prime\prime\prime} \coloneqq \begin{cases} \hat{\Psi}_{[0]}^{\prime\prime}, & \text{if } 1 \le i \le q, \\ \hat{\Psi}_{[i-q]}^{\prime\prime}, & \text{if } i > q, \end{cases} \quad \text{and} \quad \Psi_{[q]}^{\prime\prime\prime} \coloneqq \begin{cases} \Psi_{[0]}^{\prime\prime}, & \text{if } 1 \le i \le q, \\ \Psi_{[i-q]}^{\prime\prime}, & \text{if } i > q, \end{cases} \tag{32}$$

with $\hat{\Psi}''_{[i]}$ and $\hat{\Psi}''_{[i]}$ being identical to $\hat{\Psi}'_{[i]}$ in (28) and $\Psi'_{[i]}$ in (25) for all *i*, respectively, except for a reversed sign in the first component.

Assumption 4.5. $\alpha_1, \ldots, \alpha_p, \beta_1, \ldots, \beta_q \in S_H$ are *H-S* operators.

Assumption 4.6. The image of the operator-valued matrix $\prod : \mathcal{H}^q \to \mathcal{H}^q$ lies dense.

Assumption 4.7. The eigenvalues of $\prod \prod^* : \mathcal{H}^q \to \mathcal{H}^q$ satisfy $\zeta_j \neq \zeta_{j+1}$ for all $j \in \mathbb{N}$.

The following Sobolev condition, which is similar to the one in the ARMA(1,1) case, enables us also to deduce a consistency result for the estimation errors for the complete ARMA(p,q) operators.

Assumption 4.8. For some $\gamma > 0$ holds

$$\sum_{i=1}^{\infty}\sum_{j=1}^{\infty} \langle B_q, h_i \otimes d_j \rangle_{\mathcal{S}}^2 \left(1 + i^{2\gamma} + j^{2\gamma}\right) < \infty.$$

Theorem 4.2. Let the assumptions of Theorem 3.1, Assumptions 4.5–4.8, and $M^{1+\gamma}\zeta_M^{-1}Z_M = O(K^{\beta})$ and $\gamma_N = o(\zeta_M M^{-\gamma})$ hold. Then, for the fARMA(p,q) operators holds

$$\max_{1 \le i \le p, 1 \le j \le q} \left\{ \|\hat{\alpha}_i - \alpha_i\|_{\mathcal{S}}, \|\hat{\beta}_j - \beta_j\|_{\mathcal{S}} \right\} = \mathcal{O}_{\mathbb{P}}(M^{-\gamma}).$$

Remark 4.4. To the best of our knowledge, Theorem 4.2 which states explicit asymptotic upper bounds for the estimation errors for all operators of fARMA(p,q) processes attaining values in arbitrary separable Hilbert spaces for arbitrary orders is an entirely new result in the literature.

5 Conclusion

5.1 Summary

This article establishes consistent Yule-Walker type estimates using Tychonoff-regularization for the operators of (functional) invertible linear processes satisfying Sobolev and mild weak dependence conditions in separable Hilbert spaces. Building on these results, we also establish consistent Yule-Walker estimates for the operators of functional AR, MA and ARMA processes with arbitrary orders. The estimates for the complete operators in all of these models are derived on the basis of asymptotic consistency results for finite-dimensional projections of a growing dimension, which are of use in their own right. Our results represent innovations in the current literature in numerous ways: The invertible linear and the functional AR, MA and ARMA processes are allowed to attain their values in general, separable Hilbert spaces, and we derive explicit asymptotic upper bounds for estimation errors of all operators in the inverted and linear representation of our invertible linear process, as well as for the AR, MA and ARMA processes with arbitrary orders. Further, the definition of all our models requires the errors only to be strictly stationary, ergodic white noises instead of independent and identically distributed, which is of use when studying solutions of non-linear function valued time series processes.

5.2 A promising concept of invertibility

It is worth to point out that Granger and Andersen (1978) established a more general notion of invertibility as the series representation (1) for real-valued processes. They called a real-valued

process $(X_k)_k$ invertible if $\mathbb{E}(\hat{\varepsilon}_k - \varepsilon_k)^2 \to 0$ for $k \to \infty$, with

$$\varepsilon_k = X_k - f(X_{k-1}, X_{k-2}, \dots, X_{k-p}, \varepsilon_{k-1}, \varepsilon_{k-2}, \dots, \varepsilon_{k-q}), \quad k \in \mathbb{N},$$
(33)

$$\hat{\varepsilon}_k = X_k - f(X_{k-1}, X_{k-2}, \dots, X_{k-p}, \hat{\varepsilon}_{k-1}, \hat{\varepsilon}_{k-2}, \dots, \hat{\varepsilon}_{k-q}), \quad k \in \mathbb{N},$$
(34)

where $\hat{\varepsilon}_0, \hat{\varepsilon}_{-1}, \ldots, \hat{\varepsilon}_{1-q}$ and $X_0, X_{-1}, \ldots, X_{1-p}$ are given for some $p, q \in \mathbb{N}$, where $(\varepsilon_k)_k$ is i.i.d. and centered with finite second moments, and where f is a measurable function. In the case that the function f is unknown it was replaced by its estimate \hat{f} (if given). Hallin (1980, 1981) generalized this concept of invertibility even further by using $\hat{\varepsilon}_0, \hat{\varepsilon}_{-1}, \ldots$ rather than letting time tend to infinity which enabled also the analysis of time-dependent processes.

The more general concept of invertibility described above can also be established for processes in a general, separable Banach space \mathcal{B} endowed with the norm $\|\cdot\|$. Namely, we call a process $(X_k)_k \subset \mathcal{B}$ mild Granger-Andersen invertible if

$$\|\hat{\varepsilon}_{k_N} - \varepsilon_{k_N}\| = \mathbf{o}_{\mathbb{P}}(1),$$

where N is the sample size (as usual), $k_N \to \infty$, $(\varepsilon_k)_k$ is i.i.d., and $\hat{\varepsilon}_{k_N}$ and ε_{k_N} have the representations (33)–(34), respectively. This notion of invertibility is based on Granger and Andersen (1978) but it is milder (hence its name), because finite second moments of the innovations are not required, the convergence is in a weaker sense (in probability rather than in the L^2 -sense), and, for technical reasons, $(k_n)_n$ can be explicitly chosen. It can be shown that invertibility in the common sense (1) implies mild Granger-Andersen invertibility. Moreover, as mentioned in Granger and Andersen (1978) for real-valued processes, there are also functional time series which are mild Granger-Andersen invertible but not invertible as in the sense (1), and there are processes that are not invertible in both senses: Under certain conditions bilinear functional time series in a general, separable Hilbert space as stated in Hörmann and Kokoszka (2010) can be mild Granger-Andersen invertible but not invertible in the sense (1), and, e.g., the non-linear MA(1) process $X_k = \alpha_1(\varepsilon_{k-1}^2) + \varepsilon_k$ a.s. in $L^2[0, 1]$ can be strictly stationary but not invertible in both senses.

5.3 Future research

Although mild Granger-Andersen invertibility discussed in the previous section is useful in situations where invertibility in the sense (1) does not hold, our proof technique generally cannot be used as one not necessarily can truncate a series, and let the number of summands go to infinity. As a result, the entire estimation method must be modified and appropriate assumptions must be made. Another research topic is the derivation of estimators and consistency rates in general Banach, possibly even in metric spaces.

Acknowledgements

The authors would like to note that they did not receive any funds for this project. Furthermore, the majority of Sebastian Kühnert's work took place during his employment at University of California, Davis, and the remaining work at Ruhr University Bochum.

References

- Aue, A. and A. van Delft (2020). Testing for stationarity of functional time series in the frequency domain. Annals of Statistics 48(5), 2505–2547.
- Aue, A., L. Horváth, and D. Pellatt (2017). Functional generalized autoregressive conditional heteroskedasticity. *Journal of Time Series Analysis 38*, 3–21.
- Aue, A. and J. Klepsch (2017). Estimating functional time series by moving average model fitting. arXiv:1701.00770. https://arxiv.org/abs/1701.00770.
- Bosq, D. (2000). *Linear Processes in Function Spaces*. Lecture Notes in Statistics. New York: Springer.
- Caponera, A. and V. Panaretos (2022). On the rate of convergence for the autocorrelation operator in functional autoregression. *Statistics & Probability Letters 189*(109575).
- Cerovecki, C., C. Francq, S. Hörmann, and J.-M. Zakoïan (2019). Functional GARCH models: The quasi-likelihood approach and its applications. *Journal of Econometrics* 209(2), 353–375.
- van Delft, A., V. Characiejus, and H. Dette (2021). A nonparametric test for stationarity in functional time series. *Statistica Sinica* 31(3), pp. 1375–1395.
- Düker, M.-C. (2018). Limit theorems for Hilbert space-valued linear processes under long-range dependence. *Stochastic Processes and their Applications* 128(5), 1439–1465.
- Granger, C. W. J. and A. Andersen (1978). On the invertibility of time series models. *Stochastic Processes and their Applications* 8(1), 87–92.
- Hall, P. and J. Horowitz (2005). Nonparametric methods for inference in the presence of instrumental variables. *Annals of Statistics* 33(6), 2904–2929.
- Hall, P. and J. Horowitz (2007). Methodology and convergence rates for functional linear regression. Annals of Statistics 35(1), 70–91.
- Hall, P. and A. Meister (2007). A ridge-parameter approach to deconvolution. Annals of Statistics 35(4), 1535–1558.
- Hallin, M. (1980). Invertibility and generalized invertibility of time series models. Journal of the Royal Statistical Society Series B: Statistical Methodology 42(2), 210–212.
- Hallin, M. (1981). Addendum to: "invertibility and generalized invertibility of time series models". Journal of the Royal Statistical Society Series B: Statistical Methodology 43(1), 103.
- Hörmann, S., L. Horváth, and R. Reeder (2013). A functional version of the ARCH model. Econometric Theory 29(2), 267–288.
- Hörmann, S. and P. Kokoszka (2010). Weakly dependent functional data. Annals of Statistics 38, 1845–1884.

- Hörmann, S. and P. Kokoszka (2010). Weakly dependent functional data. Ann. Statist. 38, 1845– 1884.
- Horváth, L. and P. Kokoszka (2012). Inference for functional data with applications. New York: Springer.
- Horváth, L., P. Kokoszka, and G. Rice (2014). Testing stationarity of functional time series. Journal of Econometrics 179(1), 66–82.
- Hsing, T. and R. Eubank (2015). Theoretical Foundations of Functional Data Analysis, with an Introduction to Linear Operators. West Sussex: Wiley.
- Kara Terki, N. and T. Mourid (2024). Exponential bounds and convergence rates of sieve estimators for functional autoregressive processes. *Sankhya A 86*(1), 364–391.
- Klepsch, J. and C. Klüppelberg (2017). An innovations algorithm for the prediction of functional linear processes. *Journal of Multivariate Analysis* 155, 252–271.
- Klepsch, J., C. Klüppelberg, and T. Wei (2017). Prediction of functional ARMA processes with an application to traffic data. *Econometrics and Statistics* 1, 128–149.
- Kokoszka, P. and M. Reimherr (2017). *Introduction to Functional Data Analysis*. Chapman and Hall/CRC.
- Kuenzer, T. (2024). Estimation of functional ARMA models. Bernoulli 30(1), 117–142.
- Kühnert, S. (2019). Über funktionale ARCH- und GARCH-Zeitreihen. Ph. D. thesis, University of Rostock.
- Kühnert, S. (2020). Functional ARCH and GARCH models: A Yule-Walker approach. *Electronic Journal of Statistics* 14(2), 4321–4360.
- Kühnert, S. (2022). Lagged covariance and cross-covariance operators of processes in Cartesian products of abstract Hilbert spaces. *Electronic Journal of Statistics* 16(2), 4823–4862.
- Kühnert, S. (2024+). Estimating lagged (cross-)covariance operators of L^p-m-approximable processes in Cartesian product Hilbert spaces. In revision. arXiv:2402.08110. https://https://arxiv.org/pdf/2402.08110.
- Kutta, T. (2024+). Approximately mixing time series. https://papers.ssrn.com/sol3/papers.cfm?abstract_i
- Liebl, D. (2013). Modeling and forecasting electricity spot prices: A functional data perspective. Annals of Applied Statistics 7(3), 1562–1592.
- Mas, A. and B. Pumo (2009). Linear processes for functional data.
- Merlevède, F. (1996). Central limit theorem for linear processes with values in a Hilbert space. Stochastic Processes and their Applications 65(1), 103–114.
- Merlevède, F., M. Peligrad, and S. Utev (1997). Sharp conditions for the CLT of linear processes in a Hilbert space. *Journal of Theoretical Probability* 10, 681–693.

- Račkauskas, A. and C. Suquet (2010). On limit theorems for Banach-space-valued linear processes. Lithuanian Mathematical Journal 50, 71–87.
- Ramsay, J. and B. Silverman (2005). *Functional data analysis*. Springer Series in Statistics. New York: Springer.
- Spangenberg, F. (2013). Strictly stationary solutions of ARMA equations in Banach spaces. Journal of Multivariate Analysis 121, 127–138.
- Turbillon, C., D. Bosq, J.-M. Marion, and B. Pumo (2008). Estimation du paramètre des moyennes mobiles Hilbertiennes. Comptes Rendus. Mathématique 346(5-6), 347–350.

A Some operator norm (in-)equalities

In various proofs, we make use of the following operator norm (in-)equalities.

Lemma A.1. Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a Hilbert space with respective inner product, and let $A_i, B_{ij} \in \mathcal{L}_{\mathcal{H}}$ and $S_{ij} \in \mathcal{S}_{\mathcal{H}}$ be bounded and H-S operators, respectively, where $i = 1, \ldots, m \in \mathbb{N}, j = 1, \ldots, n \in \mathbb{N}$. Then, the following holds.

(a) The operator-valued vector $\mathbf{A} \coloneqq (A_1 \cdots A_m)$ satisfies $\mathbf{A} \in \mathcal{L}_{\mathcal{H}^m, \mathcal{H}}$, with

$$\|\boldsymbol{A}\|_{\mathcal{L}} \le \left\|\sum_{i=1}^{m} A_i\right\|_{\mathcal{L}}.$$
(35)

(b) For the operator-valued matrix $\mathbf{B} \coloneqq (B_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$ holds $\mathbf{B} \in \mathcal{L}_{\mathcal{H}^m, \mathcal{H}^n}$, with

$$\|\boldsymbol{B}\|_{\mathcal{L}} \le \sum_{i=1}^{m} \|(B_{i1} \cdots B_{in})\|_{\mathcal{L}}.$$
 (36)

(c) For the operator-valued matrix $\mathbf{S} \coloneqq (S_{ij})_{1 \leq i \leq m, 1 \leq j \leq n}$ holds $\mathbf{S} \in S_{\mathcal{H}^m, \mathcal{H}^n}$, with

$$\|\boldsymbol{S}\|_{\mathcal{S}}^{2} = \sum_{i=1}^{m} \sum_{j=1}^{n} \|S_{ij}\|_{\mathcal{S}}^{2}.$$
(37)

Proof. (a) For any $x := (x_1, \ldots, x_m)^{\top} \in \mathcal{H}^m$ holds

$$\|\mathbf{A}(x)\| = \left\|\sum_{i=1}^{m} A_i(x_i)\right\| \le \left\|\sum_{i=1}^{m} A_i\right\|_{\mathcal{L}} \|x_i\|.$$

Thus, as $||x_i|| \le ||x||$ for all *i*, the assertion follows from the definition of the norm $|| \cdot ||_{\mathcal{L}}$.

(b) Elementary conversions and the definition of our norms lead indeed to

$$\|\boldsymbol{B}\|_{\mathcal{L}} = \left(\sup_{\|\boldsymbol{x}\| \le 1} \sum_{i=1}^{m} \|(B_{i1} \cdots B_{in})(\boldsymbol{x})\|^{2}\right)^{1/2}$$

$$\leq \sum_{i=1}^{m} \sup_{\|\boldsymbol{x}\| \le 1} \|(B_{i1} \cdots B_{in})(\boldsymbol{x})\| = \sum_{i=1}^{m} \|(B_{i1} \cdots B_{in})\|_{\mathcal{L}}.$$

(c) See Kühnert (2019), Lemma 2.16 (b).

B Proofs of the results in Sections 3-4

B.1 Proofs of results in Section 3

Proof of Proposition 3.1. (a) Due to (8) holds for any $x \in \mathcal{H}$ with $x \neq 0$,

$$\langle \mathscr{C}_X(x), x \rangle = \sum_{i=0}^{\infty} \langle \phi_i \mathscr{C}_{\varepsilon} \phi_i^*(x), x \rangle = \sum_{i=0}^{\infty} \left\langle \mathscr{C}_{\varepsilon}^{1/2} \phi_i^*(x), \mathscr{C}_{\varepsilon}^{1/2} \phi_i^*(x) \right\rangle = \sum_{i=0}^{\infty} \left\| \mathscr{C}_{\varepsilon}^{1/2} \phi_i^*(x) \right\|^2$$

Due to $\phi_0^* = \phi_0 = \mathbb{I}$, and as $\mathscr{C}_{\varepsilon}^{1/2}$ is injective after Assumption 3.2, we have $\|\mathscr{C}_{\varepsilon}^{1/2}\phi_i^*(x)\| > 0$. Thus, for any $x \in \mathcal{H}$ with $x \neq 0$ holds $\langle \mathscr{C}_X(x), x \rangle > 0$, in other words, \mathscr{C}_X is a strictly positive operator which is as such injective.

(b) Suppose by way of contradiction that there exists a $\mathbf{v} = (v_1, \ldots, v_L)^\top \in \mathcal{H}^L$ such that $\mathbf{v} \neq 0$, and

$$0 = \langle \mathscr{C}_{\mathbf{X}^{[L]}}(\mathbf{v}), \mathbf{v} \rangle = \operatorname{var} \Big(\sum_{i=0}^{L-1} \langle X_{-i}, v_i \rangle \Big)$$

If this is so there exists an $r \leq L - 1$ so that $v_r \neq 0$, and almost surely

$$\langle X_{-r}, v_r \rangle = -\sum_{i=r+1}^{L-1} \langle X_{-i}, v_i \rangle.$$
(38)

Notice the right hand side of the above is measurable with respect to $\sigma(\varepsilon_i, i \leq -r-1)$. Due to the representation (15), the left hand side of the above is equal to

$$\sum_{\ell=0}^{\infty} \langle \phi_{\ell}(\varepsilon_{-r-\ell}), v_r \rangle,$$

where $\phi_0 = \mathbb{I}$. Multiplying each side of (38) with $\langle \varepsilon_{-r}, v_r \rangle$ and taking expectations gives that $\langle \mathscr{C}_{\varepsilon}(v_r), v_r \rangle = 0$, with $v_r \neq 0$, which contradicts the injectivity of $\mathscr{C}_{\varepsilon}$.

(c) According to Fubini's theorem, and as linear operators commute with the expected value, we have for any i, j, k,

$$\mathbb{E}\langle c_k, \varepsilon_{i-j} \rangle \sum_{\ell=0}^{\infty} A_\ell(\varepsilon_{i-\ell}) = \sum_{\ell=0}^{\infty} A_\ell(\mathbb{E}\langle c_k, \varepsilon_{i-j} \rangle \varepsilon_{i-\ell}).$$
(39)

We note that $\mathbb{E}\langle c_k, \varepsilon_{i-j} \rangle \varepsilon_{i-\ell} = 0$ when $j \neq \ell$, and when $j = \ell$,

$$\mathbb{E}\langle c_k, \varepsilon_{i-j}\rangle \varepsilon_{i-j} = \sum_{r=0}^{\infty} \mathbb{E}\langle c_k, \varepsilon_{i-j}\rangle \langle c_r, \varepsilon_{i-j}\rangle c_r = \lambda_k c_k$$

after Bosq (2000), equation (1.36). Combining with (39), we have that

$$\mathbb{E}\langle c_k, \varepsilon_{i-j} \rangle \sum_{\ell=0}^{\infty} A_\ell(\varepsilon_{i-\ell}) = \lambda_k A_j(c_k).$$

Similarly,

$$\mathbb{E}\langle c_k, \varepsilon_{i-j} \rangle \sum_{\ell=0}^{\infty} B_\ell(\varepsilon_{i-\ell}) = \lambda_k B_j(c_k).$$

Hence, due the hypothesis $\lambda_k A_j(c_k) = \lambda_k B_j(c_k)$ for all $j, k \in \mathbb{N}$, and as C_{ε} is injective, $\lambda_k > 0$, thus $A_j(c_k) = B_j(c_k)$ for all $j, k \in \mathbb{N}$. Consequently, as $(c_j)_j$ forms a CONS of \mathcal{H} , for all j holds indeed $A_j = B_j$.

(d) We note that if X is an SLP, then $Z_i = \sum_{\ell=0}^{\infty} A_\ell(X_{i-\ell})$ is also an SLP, since

$$Z_i = \sum_{\ell=0}^{\infty} A_\ell(X_{i-\ell}) = \sum_{\ell=0}^{\infty} D_\ell(\varepsilon_{i-\ell}),$$

where $D_{\ell} = \sum_{j=0}^{\ell} A_j \phi_{\ell-j}$. By hypothesis we also have that

$$Z_i = \sum_{\ell=0}^{\infty} B_\ell(X_{i-\ell}) = \sum_{\ell=0}^{\infty} F_\ell(\varepsilon_{i-\ell}),$$

where $F_{\ell} = \sum_{j=0}^{\ell} B_j \phi_{\ell-j}$. According to part (c), $F_{\ell} = D_{\ell}$ for all $\ell \in \mathbb{N}$. Since $\phi_0 = \mathbb{I}$ we have then that $A_0 = D_0 = F_0 = B_0$. Now the result follows by induction.

In several places, we make use of the following upper bounds derived from Bosq (2000). From the definition of $\mathbf{X}^{[L]} = (X_k^{[L]})_{k \in \mathbb{Z}}$, and stationarity of $\mathbf{X} = (X_k)_{k \in \mathbb{Z}}$ follows

$$\|\mathscr{C}_{\mathbf{X}^{[L]}}\|_{\mathcal{N}} = \mathbb{E}\|X_0^{[L]}\|^2 = L \mathbb{E}\|X_0\|^2.$$
(40)

Further, due to the definition of $X^{[L]}$, stationarity and Cauchy-Schwarz inequality, holds,

$$\begin{aligned} \|\mathscr{C}_{\boldsymbol{X}^{[L]}}^{h}\|_{\mathcal{N}} &\leq \mathbb{E} \|X_{0}^{[L]}\|^{2} = L \mathbb{E} \|X_{0}\|^{2}, \quad h \in \mathbb{Z}, \\ \|\mathscr{C}_{\boldsymbol{X}^{[L]},\boldsymbol{X}}^{h}\|_{\mathcal{N}} &\leq \sqrt{L} \mathbb{E} \|X_{0}\|^{2}, \quad h \in \mathbb{Z}. \end{aligned}$$
(41)

Moreover, the eigenvalues $\lambda_k = \lambda_k(L)$ of $\mathscr{C}_{\mathbf{X}^{[L]}}$ satisfy $\lambda_1 \ge \lambda_2 \ge \cdots > 0$ after Proposition 3.1 (b) and Assumption 3.4, and $\|\mathscr{C}_{\mathbf{X}^{[L]}}\|_{\mathcal{N}} = \sum_{j=1}^{\infty} \lambda_j$ and (40) yield

$$\lambda_k < \frac{1}{k} \sum_{j=1}^k \lambda_j < k^{-1} L \mathbb{E} ||X_0||^2, \quad k, L \in \mathbb{N},$$

and consequently, with $\Lambda_k = \Lambda_k(L)$ in (10), provided $\mathbb{E} ||X_0||^2 > 0$, we obtain

$$\Lambda_k \ge \lambda_k^{-1} > kL^{-1}(\mathbb{E}||X_0||^2)^{-1}, \quad k, L \in \mathbb{N}.$$

At first, towards proving Theorem 3.1, we focus on an idealized case in which Ψ_L is finitedimensional, and may be diagonalized with respect to the CONS $(c_i \otimes d_j)_{i,j} = (c_i(L) \otimes d_j)_{i,j}$ of $\mathcal{S}_{\mathcal{H}^{L},\mathcal{H}}$, where $(\lambda_{j}, c_{j})_{j \in \mathbb{N}} = (\lambda_{j}(L), c_{j}(L))_{j \in \mathbb{N}}$ and $(\mu_{j}, d_{j})_{j \in \mathbb{N}}$ are the eigenpair sequences of the covariance operators $\mathscr{C}_{\mathbf{X}^{[L]}} \in \mathcal{N}_{\mathcal{H}^{L}}$ and $\mathscr{C}_{\mathbf{X}} \in \mathcal{N}_{\mathcal{H}}$, respectively. Thus, if $\Psi_{L} \in \mathcal{S}_{\mathcal{H}^{L},\mathcal{H}}$, which is given if all ψ_{i} are H-S operators, we have

$$\Psi_L = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \langle \Psi_L, c_i \otimes d_j \rangle_{\mathcal{S}}(c_i \otimes d_j).$$

Hereinafter, we make use of this identity, but state some technnical assumptions first.

Assumption B.1. For fixed $K \in \mathbb{N}$ and almost all $L \in \mathbb{N}$, there are constants $p_{i,j,L} \in \mathbb{R}$ so that

$$\Psi_L = \Psi_L(K) = \sum_{i=1}^K \sum_{j=1}^K p_{i,j,L}(c_i \otimes d_j).$$
(42)

Theorem B.1. Let Assumptions 3.1–3.5, B.1 hold. Further, let $K \in \mathbb{N}$ be fixed, and suppose that $\sum_{\ell>L} \|\psi_\ell\|_{\mathcal{L}} = o(\Lambda_K L^{3/2} N^{-1/2})$, and also $\theta_N = o(\lambda_K)$ and $\theta_N = o(\Lambda_K L^2 N^{-1/2})$, where $\Lambda_K = \sup_{1 \leq j \leq K} (\lambda_j - \lambda_{j+1})^{-1}$. Then, it holds

$$\|\hat{\Psi}_L - \Psi_L(K)\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}\left(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2}\right), \quad K \in \mathbb{N}.$$
(43)

Proof. To reiterate, $(\lambda_j, c_j)_{j \in \mathbb{N}} = (\lambda_j(L), c_j(L))_{j \in \mathbb{N}}, (\hat{\lambda}_j, \hat{c}_j)_{j \in \mathbb{N}} = (\hat{\lambda}_j(L), \hat{c}_j(L))_{j \in \mathbb{N}}$ and $(\mu_j, d_j)_{j \in \mathbb{N}}$ are the eigenpair sequences of the covariance operators $\mathscr{C}_{\mathbf{X}^{[L]}}, \mathscr{C}_{\mathbf{X}^{[L]}} \in \mathcal{N}_{\mathcal{H}^L}$ and $\mathscr{C}_{\mathbf{X}} \in \mathcal{N}_{\mathcal{H}}$, respectively, where $\mathbf{X}^{[L]} = (X_k^{[L]})_{k \in \mathbb{Z}} \subset \mathcal{H}^L$, with $L \in \mathbb{N}$, is the process in (2) being defined via our ISLP $\mathbf{X} = (X_k)_{k \in \mathbb{Z}}$ in Assumption 3.1. Further, $(c_i \otimes d_j)_{i,j} = (c_i(L) \otimes d_j)_{i,j}$ is a CONS of $\mathcal{S}_{\mathcal{H}^L,\mathcal{H}}$, and due to Assumptions 3.5, B.1, $\Psi_L \in \mathcal{S}_{\mathcal{H}^L,\mathcal{H}}$ is an H-S operator which has for some $K \in \mathbb{N}$ and almost all $L \in \mathbb{N}$ the representation $\Psi_L = \Psi_L(K) = \sum_{i=1}^K \sum_{j=1}^K p_{i,j,L}(c_i \otimes d_j)$. Moreover, for the eigenvalues of $\mathscr{C}_{\mathbf{X}^{[L]}}$ holds $\lambda_1 > \cdots > \lambda_K > 0$ for all L after Assumptions 3.1–3.4. Throughout, we write $\hat{\mathcal{D}}, \mathcal{D}$ for $\hat{\mathcal{O}}_{\mathbf{X}^{[L]},\mathbf{X}} = \mathscr{C}_{\mathbf{X}^{[L]},\mathbf{X}}^1 = \mathscr{C}_{\mathbf{X}^{[L]},\mathbf{X}}^1 \in \mathcal{S}_{\mathcal{H}^L,\mathcal{H}}$, and \mathscr{C}, \mathscr{C} for $\mathscr{C}_{\mathbf{X}^{[L]}}, \mathscr{C}_{\mathbf{X}^{[L]}} \in \mathcal{N}_{\mathcal{H}^L}$, respectively. From the definition of Ψ_L , due to $\mathscr{C}^{\dagger} = (\mathscr{C} + \theta_N \mathbb{I})^{-1}$, the approximate Yule-Walker equation (11), and $\mathscr{C}^{\ddagger} := \mathscr{C} \mathscr{C}^{\dagger}$, follows

$$\begin{split} \hat{\Psi}_{L} - \Psi_{L} &= (\hat{\mathscr{D}} - \mathscr{D}) \hat{\mathscr{C}^{\dagger}} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} + \mathscr{D} \Big(\hat{\mathscr{C}^{\dagger}} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} - \mathscr{C^{\dagger}} \prod_{c_{1}}^{c_{K}} \Big) + \mathscr{D} \mathscr{C^{\dagger}} \prod_{c_{1}}^{c_{K}} - \Psi_{L} \\ &= (\hat{\mathscr{D}} - \mathscr{D}) \hat{\mathscr{C}^{\dagger}} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} + \mathscr{D} \Big(\hat{\mathscr{C}^{\dagger}} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} - \mathscr{C^{\dagger}} \prod_{c_{1}}^{c_{K}} \Big) + \Big(\sum_{\ell > L} \psi_{\ell} \mathscr{C}_{\boldsymbol{X}^{[L]}, \boldsymbol{X}}^{1-\ell} \Big) \mathscr{C^{\dagger}} \prod_{c_{1}}^{c_{K}} + \Psi_{L} \Big(\mathscr{C^{\dagger}} \prod_{c_{1}}^{c_{K}} - \mathbb{I} \Big). \end{split}$$

Consequently, due to triangle inequality and operator-valued Hölder's inequality, we have

$$\|\hat{\Psi}_{L} - \Psi_{L}\|_{\mathcal{S}} \leq \|\hat{\mathscr{D}} - \mathscr{D}\|_{\mathcal{S}} \|\hat{\mathscr{C}}^{\dagger} \prod_{\hat{c}_{1}}^{\hat{c}_{K}}\|_{\mathcal{L}} + \|\mathscr{D}\|_{\mathcal{S}} \|\hat{\mathscr{C}}^{\dagger} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} - \mathscr{C}^{\dagger} \prod_{c_{1}}^{c_{K}}\|_{\mathcal{L}} + \left\|\sum_{\ell > L} \psi_{\ell} \mathscr{C}^{1-\ell}_{\mathbf{X}^{[L]}, \mathbf{X}}\right\|_{\mathcal{S}} \|\mathscr{C}^{\dagger} \prod_{c_{1}}^{c_{K}}\|_{\mathcal{L}} + \left\|\Psi_{L} \left(\mathscr{C}^{\dagger} \prod_{c_{1}}^{c_{K}} - \mathbb{I}\right)\right\|_{\mathcal{S}}.$$

$$(44)$$

Thereby, according to the definition of $\hat{\mathscr{C}}^{\dagger}, \mathscr{C}^{\dagger}$ and the operator norm, for all K, L, N holds

$$\left\| \hat{\mathscr{C}}^{\dagger} \prod_{\hat{c}_1}^{\hat{c}_K} \right\|_{\mathcal{L}} = \left\| (\hat{\mathscr{C}} + \theta_N \mathbb{I})^{-1} \prod_{\hat{c}_1}^{\hat{c}_K} \right\|_{\mathcal{L}} = \sup_{1 \le j \le K} (\hat{\lambda}_j + \theta_N)^{-1} = (\hat{\lambda}_K + \theta_N)^{-1} =: \hat{\ell}_{K,N}, \quad (45)$$

$$\left\| \mathscr{C}^{\dagger} \prod_{c_1}^{c_K} \right\|_{\mathcal{L}} = \left\| (\mathscr{C} + \theta_N \mathbb{I})^{-1} \prod_{c_1}^{c_K} \right\|_{\mathcal{L}} = \sup_{1 \le j \le K} (\lambda_j + \theta_N)^{-1} = (\lambda_K + \theta_N)^{-1} =: \ell_{K,N}.$$
(46)

Further, due to (41),

$$\|\mathscr{D}\|_{\mathcal{S}} \le \sqrt{L} \,\mathbb{E} \|X_0\|^2. \tag{47}$$

From the definition of the operator norm, the projection operators and $\hat{\mathscr{C}}^{\dagger}$ and \mathscr{C}^{\dagger} follows with $\hat{x}_j \coloneqq \langle x, \hat{c}'_j \rangle, x_j \coloneqq \langle x, c_j \rangle, \hat{\ell}_{j,N} \coloneqq (\hat{\lambda}_j + \theta_N)^{-1}$ and $\ell_{j,N} \coloneqq (\lambda_j + \theta_N)^{-1}$,

$$\begin{aligned} \left\| \hat{\mathscr{C}}^{\dagger} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} - \mathscr{C}^{\dagger} \prod_{c_{1}}^{c_{K}} \right\|_{\mathcal{L}} &= \sup_{\|\boldsymbol{x}\| \leq 1} \left\| \sum_{j=1}^{\infty} \hat{x}_{j} \hat{\mathscr{C}}^{\dagger} \prod_{\hat{c}_{1}}^{\hat{c}_{K}} \hat{c}_{j}' - x_{j} \mathscr{C}^{\dagger} \prod_{c_{1}}^{c_{K}} c_{j} \right\| \\ &= \sup_{\|\boldsymbol{x}\| \leq 1} \left\| \sum_{j=1}^{K} \hat{x}_{j} \hat{\mathscr{C}}^{\dagger} (\hat{c}_{j}') - x_{j} \mathscr{C}^{\dagger} (c_{j}) \right\| \\ &\leq \sup_{\|\boldsymbol{x}\| \leq 1} \sum_{j=1}^{K} \left\| \hat{x}_{j} \hat{\ell}_{j,N} (\hat{c}_{j}' - c_{j}) \right\| + \sup_{\|\boldsymbol{x}\| \leq 1} \sum_{j=1}^{K} \left\| (\hat{x}_{j} \hat{\ell}_{j,N} - x_{j} \ell_{j,N}) (c_{j}) \right\|. \end{aligned}$$
(48)

For the first term in (48) holds due to elementary conversions and (10),

$$\sup_{\|\boldsymbol{x}\| \le 1} \sum_{j=1}^{K} \|\hat{x}_{j}\hat{\ell}_{j,N}(\hat{c}_{j}' - c_{j})\| \le K \sup_{1 \le j \le K} \hat{\ell}_{j,N} \|\hat{c}_{j}' - c_{j}\| \le 2\sqrt{2}K\hat{\ell}_{K,N}\Lambda_{K} \|\hat{\mathscr{C}} - \mathscr{C}\|_{\mathcal{L}}.$$
 (49)

For the second term in (49) holds due to similar conversions as above,

$$\sup_{\|\boldsymbol{x}\| \le 1} \sum_{j=1}^{K} \left\| \left(\hat{x}_{j} \hat{\ell}_{j,N} - x_{j} \ell_{j,N} \right) (c_{j}) \right\| \le \sup_{\|\boldsymbol{x}\| \le 1} \sum_{j=1}^{K} \left\| (\hat{x}_{j} - x_{j}) \hat{\ell}_{j,N} (c_{j}) \right\| + \sup_{\|\boldsymbol{x}\| \le 1} \sum_{j=1}^{K} \left\| x_{j} (\hat{\ell}_{j,N} - \ell_{j,N}) (c_{j}) \right\|$$

$$\le K \hat{\ell}_{K,N} \sup_{1 \le j \le K} \left\| \hat{c}_{j}' - c_{j} \right\| + \sup_{1 \le j \le K} \left\| (\hat{\ell}_{j,N} - \ell_{j,N}) (c_{j}) \right\|$$

$$\le 2\sqrt{2} K \hat{\ell}_{K,N} \Lambda_{K} \| \hat{\mathscr{C}} - \mathscr{C} \|_{\mathcal{L}} + \sup_{1 \le j \le K} | \hat{\ell}_{j,N} - \ell_{j,N} |.$$
(50)

Due to the definition of $\hat{\ell}_{j,N}$ and $\ell_{j,N}$, and (9), we have

$$\sup_{1 \le j \le K} |\hat{\ell}_{j,N} - \ell_{j,N}| \le \sup_{1 \le j \le K} \hat{\ell}_{j,N} \ell_{j,N} |\hat{\lambda}_j - \lambda_j| \le \hat{\ell}_{K,N} \ell_{K,N} \|\hat{\mathscr{C}} - \mathscr{C}\|_{\mathcal{L}}.$$
(51)

Hence, by combining (48)–(51), and due to $\ell_{K,N} \leq \lambda_K^{-1} \leq \Lambda_K$, we obtain

$$\left\|\hat{\mathscr{C}}^{\dagger}\prod_{\hat{c}_{1}}^{\hat{c}_{K}}-\mathscr{C}^{\dagger}\prod_{c_{1}}^{c_{K}}\right\|_{\mathcal{L}} \leq \|\hat{\mathscr{C}}-\mathscr{C}\|_{\mathcal{L}}\,\hat{\ell}_{K,N}\Lambda_{K}\big(4\sqrt{2}K+1\big).$$
(52)

Also, by using triangle inequality, operator-valued Hölder's inequality and (41), we get for any L,

$$\left\|\sum_{\ell>L}\psi_{\ell}\mathscr{C}_{\boldsymbol{X}^{[L]},\boldsymbol{X}}^{1-\ell}\right\|_{\mathcal{S}} \leq \sqrt{L}\,\mathbb{E}\|X_{0}\|^{2}\sum_{\ell>L}\|\psi_{\ell}\|_{\mathcal{L}},\tag{53}$$

where the series exists for all L after Assumption 3.1. At last, after the definition of the H-S norm, with $\Psi_L = \Psi_L(K)$ in (42), $c_i \otimes d_j$, with $c_i = c_i(L)$ and d_j being the eigenfunctions of $\mathscr{C} = \mathscr{C}_{\mathbf{X}^{[L]}}$ and $\mathscr{C}_{\mathbf{X}}$, respectively, $\mathscr{C}^{\ddagger} = \mathscr{C}(\mathscr{C} + \theta_N \mathbb{I})^{-1}$, with δ_{jk} being the *Kronecker-Delta*, so $\delta_{jk} = 1$ if j = k, and $\delta_{jk} = 0$ if $j \neq k$, with $\Psi_L = \Psi_L(K)$, we get

$$\begin{split} \left\| \Psi_{L} \left(\mathscr{C}^{\dagger} \prod_{c_{1}}^{C_{K}} - \mathbb{I} \right) \right\|_{\mathcal{S}}^{2} &= \sum_{i=1}^{\infty} \left\| \Psi_{L} \left(\mathbb{1}_{[1,K]}(i) \mathscr{C}^{\dagger} - \mathbb{I} \right) (c_{i}) \right\|^{2} \\ &= \sum_{i=1}^{\infty} \left(\mathbb{1}_{[1,K]}(i) \lambda_{i} (\lambda_{i} + \theta_{N})^{-1} - 1 \right)^{2} \left\| \sum_{k=1}^{K} \sum_{\ell=1}^{K} p_{k,\ell,L} (c_{k} \otimes d_{\ell}) (c_{i}) \right\|^{2} \\ &= \sum_{i=1}^{K} \left(\lambda_{i} (\lambda_{i} + \theta_{N})^{-1} - 1 \right)^{2} \left\| \sum_{k=1}^{K} \sum_{\ell=1}^{K} p_{k,\ell,L} \delta_{i,k} d_{\ell} \right\|^{2} \\ &= \theta_{N}^{2} \left\| \sum_{k=1}^{K} \sum_{\ell=1}^{K} (\lambda_{k} + \theta_{N})^{-1} p_{k,\ell,L} d_{\ell} \right\|^{2} \\ &\leq \theta_{N}^{2} \ell_{K,N}^{2} \left\| \sum_{k=1}^{K} \sum_{\ell=1}^{K} p_{k,\ell,L} d_{\ell} \right\|^{2} = \theta_{N}^{2} \ell_{K,N}^{2} \left\| \Psi_{L} (K) \right\|_{\mathcal{S}}^{2} . \end{split}$$
 (54)

Thereby, due to the definition of $\Psi_L(K)$ and Ψ_L , and Assumption 3.5, we have

$$\lim_{L \to \infty} \|\Psi_L(K)\|_{\mathcal{S}}^2 \le \lim_{L \to \infty} \|\Psi_L\|_{\mathcal{S}}^2 = \lim_{L \to \infty} \sum_{i=1}^L \|\psi_i\|_{\mathcal{S}}^2 = \sum_{i=1}^\infty \|\psi_i\|_{\mathcal{S}}^2 < \infty.$$
(55)

Altogether, by plugging (45)-(47) and (52)-(54) in (44), we obtain for all K, L, N,

$$\begin{aligned} \|\hat{\Psi}_L - \Psi_L(K)\|_{\mathcal{S}} &\leq \hat{\ell}_{K,N} \|\hat{\mathscr{D}} - \mathscr{D}\|_{\mathcal{S}} + \sqrt{L} \mathbb{E} \|X_0\|^2 \|\hat{\mathscr{C}} - \mathscr{C}\|_{\mathcal{L}} \hat{\ell}_{K,N} \Lambda_K (4\sqrt{2}K + 1) \\ &+ \ell_{K,N} \sqrt{L} \mathbb{E} \|X_0\|^2 \sum_{\ell > L} \|\psi_\ell\|_{\mathcal{L}} + \theta_N \ell_{K,N} \|\Psi_L(K)\|_{\mathcal{S}}. \end{aligned}$$

Hence, due to (5)–(6), $(\hat{\lambda}_K + \theta_N)^{-1} \xrightarrow{\mathbb{P}} \lambda_K^{-1}$ after $\hat{\lambda}_K \xrightarrow{\mathbb{P}} \lambda_K$, $\theta_N = o(\lambda_K)$ and the continuous mapping theorem, and because of $\sum_{\ell > L} \|\psi_\ell\|_{\mathcal{L}} = o(\Lambda_K L^{3/2} N^{-1/2})$, $\theta_N = o(\Lambda_K L^2 N^{-1/2})$ and (55), our claim is proven.

To prove the convergence result for the estimators in the case of the complete operators, we use the following auxiliary result.

Lemma B.1. Let $C, \hat{C} \in \mathcal{L}_{\mathcal{H}}$ be positive semi-definite, self-adjoint, compact operators with eigenvalues $\varphi_1 > \varphi_2 > \cdots > 0$, and $\hat{\varphi}_1 \ge \hat{\varphi}_2 \ge \cdots \ge 0$, respectively, where \hat{C} denotes an estimate for C which is defined based on a sample with sample size $N \in \mathbb{N}$ such that $\|\hat{C} - C\|_{\mathcal{L}} = O_{\mathbb{P}}(c_N)$, where $(c_N)_{N \in \mathbb{N}} \subset (0, \infty)$ is a sequence with $c_N \to 0$. Further, let $(a_N)_{N \in \mathbb{N}}, (b_N)_{N \in \mathbb{N}} \subset (0, \infty)$ be sequences with $a_N \to \infty$ and $b_N \to 0$ such that both $b_N = o(\varphi_{a_N})$ and $c_N = o(\varphi_{a_N})$. Then,

$$(\hat{\varphi}_{a_N} + b_N)^{-1} = \mathcal{O}_{\mathbb{P}}(\varphi_{a_N}^{-1}).$$

Proof. Evidently, for any a_N holds $|\hat{\varphi}_{a_N} - \varphi_{a_N}| \leq ||\hat{C} - C||_{\mathcal{L}}$, and thus

$$|\hat{\varphi}_{a_N}/\varphi_{a_N}-1| \leq \frac{\|\hat{C}-C\|_{\mathcal{L}}}{\varphi_{a_N}}$$

where the right-hand side goes to zero because $\|\hat{C} - C\|_{\mathcal{L}} = \mathcal{O}_{\mathbb{P}}(c_N)$ and $c_N = \mathcal{O}(\varphi_{a_N})$. Hence, $\hat{\varphi}_{a_N}/\varphi_{a_N} \xrightarrow{\mathbb{P}} 1$, and $(\hat{\varphi}_{a_N} + b_N)/\varphi_{a_N} \xrightarrow{\mathbb{P}} 1$ as $b_N = \mathcal{O}(\varphi_{a_N})$. Further, the continuous mapping theorem implies $\varphi_{a_N}/(\hat{\varphi}_{a_N} + b_N) \xrightarrow{\mathbb{P}} 1$ from which directly follows $(\hat{\varphi}_{a_N} + b_N)^{-1} = \mathcal{O}_{\mathbb{P}}(\varphi_{a_N}^{-1})$.

Proof of Theorem 3.1. In this proof, we make use of the notation in the proof of Theorem B.1. Firstly, $\Psi_L(K) = \sum_{i=1}^K \sum_{j=1}^K \langle \Psi_L, c_i \otimes d_j \rangle_{\mathcal{S}}(c_i \otimes d_j)$ and Assumption 3.6 imply

$$\|\Psi_L(K) - \Psi_L\|_{\mathcal{S}}^2 = \sum_{\substack{i=1\\\max(i,j)>K}}^{\infty} \langle \Psi_L, c_i \otimes d_j \rangle_{\mathcal{S}}^2 \le K^{-2\beta} S_{\Psi_L}(\beta), \quad K, L \in \mathbb{N}.$$
 (56)

Further, due to arguments in the proof of Theorem B.1, $\theta_N = o(\lambda_K)$ and $K^{1+\beta}\lambda_K^{-1}\Lambda_K L^2 = O(N^{1/2})$ which give $\hat{\ell}_{K,N} = (\hat{\lambda}_K + \theta_N)^{-1} = O_{\mathbb{P}}(\lambda_K^{-1})$ after Lemma B.1, and by using the triangle inequality, $\sum_{\ell>L} \|\psi_\ell\|_{\mathcal{L}} = o(K\Lambda_K L^{3/2} N^{-1/2})$, and $\theta_N = o(K\Lambda_K L^2 N^{-1/2})$, we get indeed

$$\begin{split} \| \tilde{\Psi}_L - \Psi_L \|_{\mathcal{S}} &\leq \| \tilde{\Psi}_L - \Psi_L(K) \|_{\mathcal{S}} + \| \Psi_L(K) - \Psi_L \|_{\mathcal{S}} \\ &= \mathcal{O}_{\mathbb{P}}(K \lambda_K^{-1} \Lambda_K L^2 N^{-1/2}) + \mathcal{O}(K^{-\beta}) \\ &= \mathcal{O}_{\mathbb{P}}(K^{-\beta}). \end{split}$$

We now turn towards proving Proposition 3.2. Under Assumption B.1 it follows for any ℓ and K and L sufficiently large that

$$\psi_{\ell} = \psi_{\ell}(K) \coloneqq \left[\Psi_{L}(K)\right]^{(\ell)} = \sum_{i=1}^{K} \sum_{j=1}^{K} p_{i,j,L}(c_{i} \otimes d_{j})^{(\ell)} = \sum_{i=1}^{K} \sum_{j=1}^{K} p_{i,j,L}(c_{i}^{(\ell)} \otimes d_{j}).$$
(57)

The finite-dimensional representation of the operators ϕ_i are not necessarily as simple. For instance for ϕ_2 holds due to $\phi_1 = \psi_1$ and $\phi_2 = \psi_1 \phi_1$ under Assumption B.1 for sufficiently large L,

$$\phi_2(K) = \psi_1^2(K) = \sum_{i=1}^K \sum_{i'=1}^K \sum_{j=1}^K \sum_{j'=1}^K p_{i,j,L} p_{i',j',L} \underbrace{\left(c_i^{(\ell)} \otimes d_j\right) \left(c_{i'}^{(\ell)} \otimes d_{j'}\right)}_{= \langle c_i^{(\ell)}, d_{j'} \rangle c_{i'}^{(\ell)} \otimes d_j}$$

Nevertheless, we can use the recursively defined finite-dimensional approximations

$$\phi_i = \phi_i(K) \coloneqq \sum_{j=1}^i \psi_j(K)\phi_{i-j}(K), \quad i \in \mathbb{N}, \quad \text{with } \phi_0(K) \coloneqq \mathbb{I}.$$
(58)

Proposition B.1. Let the assumptions of Theorem B.1 hold. Then,

$$\|\hat{\phi}_i - \phi_i(K)\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2}), \quad i, K \in \mathbb{N}.$$

Proof. With $\phi'_i := \|\phi_i(K)\|_{\mathcal{L}}, \hat{\psi}'_j := \|\hat{\psi}_j\|_{\mathcal{L}}$ for all $i, j, \hat{\phi}_0 = \phi_0(K) = \mathbb{I}$, (16), (57)–(58), elementary conversions and $\|\hat{\psi}_j - \psi_j(K)\|_{\mathcal{S}} \le \|\hat{\Psi}_L - \Psi_L(K)\|_{\mathcal{S}}$, we obtain for any $i \in \mathbb{N}$,

$$\begin{aligned} \|\hat{\phi}_{i} - \phi_{i}(K)\|_{\mathcal{S}} &= \left\|\sum_{j=1}^{i} \hat{\psi}_{j} \hat{\phi}_{i-j} - \psi_{j}(K) \phi_{i-j}(K)\right\|_{\mathcal{S}} \\ &\leq \sum_{j=1}^{i} \|\hat{\psi}_{j}\|_{\mathcal{L}} \|\hat{\phi}_{i-j} - \phi_{i-j}(K)\|_{\mathcal{S}} + \|\phi_{i-j}(K)\|_{\mathcal{L}} \|\hat{\psi}_{j} - \psi_{j}(K)\|_{\mathcal{S}} \\ &\leq \sum_{j=1}^{i-1} \hat{\psi}_{j}' \|\hat{\phi}_{i-j} - \phi_{i-j}(K)\|_{\mathcal{S}} + \sum_{j=0}^{i-1} \|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} \phi_{j}'. \end{aligned}$$

Since $\|\hat{\phi}_1 - \phi_1(K)\|_{\mathcal{S}} = \|\hat{\psi}_1 - \psi_1(K)\|_{\mathcal{S}} \le \|\hat{\Psi}_L - \Psi_L(K)\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2})$ after Theorem B.1, we obtain an asymptotic upper bound for $\|\hat{\phi}_i - \phi_i(K)\|_{\mathcal{S}}$ by doing (i-1) iterations of the above inequality. By using $\phi'_0 = 1$ and the convention that factors depending on j_k for some k in the multi-sums below are set 1 if $\sum_{j_k=1}^0$ is given, due to $\phi'_i = \|\phi_i(K)\|_{\mathcal{L}}$ and $\hat{\psi}'_j = \|\hat{\psi}_j\|_{\mathcal{L}}$ for all i, j, after Theorem B.1, implying also $\|\hat{\psi}_j\|_{\mathcal{L}} = \|\psi_j(K)\|_{\mathcal{L}} + o_{\mathbb{P}}(1)$ for all j, with

$$\Theta(i,K) \coloneqq 1 + \sum_{k=1}^{i-1} \sum_{j_1=1}^{i-j_0-1} \cdots \sum_{j_k=1}^{i-j_{k-1}-1} \left(\prod_{\ell=1}^{k-1} \|\psi_{j_k}(K)\|_{\mathcal{L}}\right) \Big[\|\phi_{j_k}(K)\|_{\mathcal{L}} + \|\psi_{j_k}(K)\|_{\mathcal{L}} \Big],$$

it holds for each i, K,

$$\begin{split} \|\hat{\phi}_{i} - \phi_{i}(K)\|_{\mathcal{S}} \\ &\leq \|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} \\ &\times \left[\sum_{j_{1}=0}^{i-1} \phi_{j_{1}}' + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=0}^{i-j_{1}-1} \hat{\psi}_{j_{1}}' \phi_{j_{2}}' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-2}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-2} \hat{\psi}_{j_{k}}'\right) \phi_{j_{i-1}}'\right] \\ &+ \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-1} \hat{\psi}_{j_{k}}'\right) \underbrace{\|\hat{\phi}_{i-\sum_{k=1}^{i-1} j_{k}} - \phi_{i-\sum_{k=1}^{i-1} j_{k}}(K)\|_{\mathcal{S}}}_{&= \|\hat{\psi}_{1} - \psi_{1}(K)\|_{\mathcal{S}} \end{split}$$

$$&\leq \|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} \\ &\times \left[1 + \sum_{j_{1}=1}^{i-1} \phi_{j_{1}}' + \sum_{j_{1}=1}^{i-j_{1}-1} \sum_{j_{2}=1}^{i-j_{1}-1} \hat{\psi}_{j_{1}}' \phi_{j_{2}}' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-2} \hat{\psi}_{j_{k}}'\right) \phi_{j_{i-1}}' \\ &+ \sum_{j_{1}=1}^{i-1} \hat{\psi}_{j_{1}}' + \sum_{j_{1}=1}^{i-j_{1}-1} \sum_{j_{2}=1}^{i-j_{1}-1} \hat{\psi}_{j_{1}}' \hat{\psi}_{j_{2}}' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-1} \hat{\psi}_{j_{k}}'\right) \right] \end{split}$$

$$= \|\hat{\Psi}_L - \Psi_L(K)\|_{\mathcal{S}} \big[\Theta(i, K) + o_{\mathbb{P}}(1) \big].$$

Consequently, after Theorem B.1, our claim is shown.

Proof of Proposition 3.2. By using the arguments in the proof of Proposition B.1, with $\phi'_i := \|\phi_j(K)\|_{\mathcal{L}}, \phi''_i := \|\phi_j\|_{\mathcal{L}}, \hat{\psi}'_j := \|\hat{\psi}_j\|_{\mathcal{L}}$ and $\psi'_j := \|\psi_j(K)\|_{\mathcal{L}}$ for all i, j, K, and $\|\psi_j(K) - \psi_j\|_{\mathcal{S}} \leq |\psi_j|_{\mathcal{L}}$

 $\|\Psi_L(K) - \Psi_L\|_{\mathcal{S}}$ for $L \ge j$, we get for $L \ge i$,

$$\begin{split} \|\hat{\phi}_{i} - \phi_{i}\|_{\mathcal{S}} &\leq \|\hat{\phi}_{i} - \phi_{i}(K)\|_{\mathcal{S}} + \|\phi_{i}(K) - \phi_{i}\|_{\mathcal{S}} \\ &\leq \Big\|\sum_{j=1}^{i} \hat{\psi}_{j} \hat{\phi}_{i-j} - \psi_{j}(K) \phi_{i-j}(K)\Big\|_{\mathcal{S}} + \Big\|\sum_{j=1}^{i} \psi_{j}(K) \phi_{i-j}(K) - \psi_{j} \phi_{i-j}\Big\|_{\mathcal{S}} \\ &\leq \sum_{j=1}^{i-1} \hat{\psi}_{j}' \|\hat{\phi}_{i-j} - \phi_{i-j}(K)\|_{\mathcal{S}} + \|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} \sum_{j=0}^{i-1} \phi_{j}' \\ &\quad + \sum_{j=1}^{i-1} \psi_{j}' \|\phi_{i-j}(K) - \phi_{i-j}\|_{\mathcal{S}} + \|\Psi_{L}(K) - \Psi_{L}\|_{\mathcal{S}} \sum_{j=0}^{i-1} \phi_{j}''. \end{split}$$

Following the lines in the proof of Proposition B.1, $\phi'_i = \|\phi_i(K)\|_{\mathcal{L}} \to \phi''_i = \|\phi_i\|_{\mathcal{L}}$ for all i, and $\hat{\psi}'_j = \|\hat{\psi}_j\|_{\mathcal{L}} = \psi''_j + o_{\mathbb{P}}(1)$ for all j after Theorem 3.1, where $\psi''_j = \|\psi_j\|_{\mathcal{L}}$, and $\psi''_0 = 1$, with

$$\Theta'(i) \coloneqq \lim_{K \to \infty} \Theta(i, K) = 1 + \sum_{k=1}^{i-1} \sum_{j_1=1}^{i-j_0-1} \cdots \sum_{j_k=1}^{i-j_{k-1}-1} \left(\prod_{\ell=1}^{k-1} \|\psi_{j_k}\|_{\mathcal{L}}\right) \Big[\|\phi_{j_k}\|_{\mathcal{L}} + \|\psi_{j_k}\|_{\mathcal{L}} \Big],$$

for each i, we obtain indeed

$$\begin{split} \|\hat{\phi}_{i} - \phi_{i}\|_{\mathcal{S}} &\leq \|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} \\ &\times \left[1 + \sum_{j_{1}=1}^{i-1} \phi_{j_{1}}' + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \hat{\psi}_{j_{1}}' \phi_{j_{2}}' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-2} \hat{\psi}_{j_{k}}'\right) \phi_{j_{i-1}}' \right. \\ &+ \sum_{j_{1}=1}^{i-1} \hat{\psi}_{j_{1}}' + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \hat{\psi}_{j_{1}}' \hat{\psi}_{j_{2}}' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{1}-2} \left(\prod_{k=1}^{i-1} \hat{\psi}_{j_{k}}'\right)\right] \\ &+ \|\Psi_{L}(K) - \Psi_{L}\|_{\mathcal{S}} \\ &\times \left[1 + \sum_{j_{1}=1}^{i-1} \phi_{j_{1}}'' + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \psi_{j_{1}}' \phi_{j_{2}}'' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-2} \psi_{j_{k}}'\right) \phi_{j_{i-1}}'' \right. \\ &+ \sum_{j_{1}=1}^{i-1} \psi_{j_{1}}'' + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \psi_{j_{1}}' \psi_{j_{2}}'' + \dots + \sum_{j_{1}=1}^{i-1} \sum_{j_{2}=1}^{i-j_{1}-1} \dots \sum_{j_{i-1}=1}^{i-j_{i-2}-1} \left(\prod_{k=1}^{i-1} \psi_{j_{k}}'\right)\right] \\ &= O_{\mathbb{P}}(K^{-\beta}) \Theta'(i) = O_{\mathbb{P}}(K^{-\beta}). \\ \\ \\ \Box = \left(\sum_{j=1}^{i-j_{1}-1} (i-j_{j_{1}-1}) - \sum_{j=1}^{i-j_{1}-1} (i-j_{j_{1}-1}) - \sum_{j=i-1}^{i-j_{1}-1} (i-j_{1}-1) - \sum_{j=i-1}^{i-j_{1}-1} (i-j_{1}-1) -$$

B.2 Proofs of results in Section 4

Proof of Corollary 4.1. As already indicated, part (a) and (b) follow immediately from Theorems B.1–3.1, respectively, by putting L = p, and due to $A_p = \Psi_p$ and $\hat{A}_{p,K} = \hat{\Psi}_{p,K}$.

Proof of Corollary 4.2. The parts (a) and (b) result from Propositions B.1, 3.2, respectively, by putting $\beta_j = \phi_j$ and $\hat{\beta}_j = \hat{\phi}_j$.

Assumption B.2. For fixed $M \in \mathbb{N}$, there are constants $b_{i,j,1} \in \mathbb{R}$ so that

$$\beta_1 = \beta_1(M) = \sum_{i=1}^M \sum_{j=1}^M b_{i,j,1}(f_i \otimes f_j).$$
(59)

To assume a finite-dimensional representation of α_1 is not necessary, as the Assumptions B.1–B.2 entail together with (23) that for α_1 holds with $\psi_1(K)$ in (57),

$$\alpha_1 = \alpha_1(K, M) \coloneqq \psi_1(K) + \beta_1(M)$$

Theorem B.2. Let $(X_k)_{k\in\mathbb{Z}}$ be the fARMA(1,1) process in (21), let the assumptions of Theorem B.1 and Assumptions 4.1–4.3, B.2 hold, and the sequence in (23) satisfies $\gamma_N = o(\lambda_K^{-1} \Lambda_K L^2 N^{-1/2})$. Then, for fixed K, M, and $L = L_N \to \infty$ in Theorem B.1 holds

$$\max\{\|\hat{\alpha}_{1} - \alpha_{1}(K, M)\|_{\mathcal{S}}, \|\hat{\beta}_{1} - \beta_{1}(M)\|_{\mathcal{S}}\} = O_{\mathbb{P}}(\lambda_{K}^{-1}\Lambda_{K}L^{2}N^{-1/2}).$$

Proof. The conversions in this proof are similar to those in the proof of Theorem B.1. We remind that $(\rho_j, f_j)_{j \in \mathbb{N}}, (\hat{\rho}_j, \hat{f}_j)_{j \in \mathbb{N}}$ are the eigenpair sequences of $\psi_1 \psi_1^*, \hat{\psi}_1 \hat{\psi}_1^* \in \mathcal{N}_{\mathcal{H}}$, respectively, where $\psi_i \in S_{\mathcal{H}}$ are the H-S operators in the inverted representation (1), and $\hat{\psi}_i = \hat{\Psi}_{L,K}^{(i)} \in S_{\mathcal{H}}$ its estimators, the *i*th components of $\hat{\Psi}_L$ in (13). Similar as in the proofs above, we have

$$\sup_{j \in \mathbb{N}} |\hat{\rho}_j - \rho_j| \le \|\hat{\psi}_1 \hat{\psi}_1^* - \psi_1 \psi_1^*\|_{\mathcal{L}}.$$
(60)

Further, the eigenvalues satisfy $\rho_1 > \rho_2 > \cdots > 0$ due to Assumptions 4.2–4.3 and since $\psi_1 \psi_1^*$ is injective which is given as the image of ψ_1 lies dense. Thus,

$$\sup_{1 \le j \le M} \|\hat{f}_j' - f_j\| \le 2\sqrt{2} P_M \|\hat{\psi}_1 \hat{\psi}_1^* - \psi_1 \psi_1^*\|_{\mathcal{L}}, \quad M \in \mathbb{N},$$
(61)

where $\hat{f}'_j \coloneqq \operatorname{sgn}\langle \hat{f}_j, f_j \rangle \hat{f}_j$ and $P_M \coloneqq \operatorname{sup}_{1 \le j \le M} (\rho_j - \rho_{j+1})^{-1}$. Thereby, due to triangle inequality and sub-multiplicativity of the operator norm holds,

$$\|\hat{\psi}_1\hat{\psi}_1^* - \psi_1\psi_1^*\|_{\mathcal{L}} \le \|\hat{\psi}_1 - \psi_1\|_{\mathcal{L}} \big(\|\psi_1\|_{\mathcal{L}} + \|\hat{\psi}_1\|_{\mathcal{L}}\big).$$
(62)

Moreover, $(\gamma_N)_{N \in \mathbb{N}} \subset (0, \infty)$ is a sequence with $\gamma_N \to 0$, and for β_1 holds $\beta_1 = \beta_1(M) = \sum_{i=1}^M \sum_{j=1}^M b_{i,j,1}(f_i \otimes f_j)$ after Assumption B.2.

As $\hat{\beta}_1 \coloneqq -\hat{\psi}_2 \hat{\psi}_1^{\dagger} \coprod_{\hat{f}_1}^{\hat{f}_M}$, where $\hat{\psi}_1^{\dagger} = \hat{\psi}_1^* (\hat{\psi}_1 \hat{\psi}_1^* + \gamma_N \mathbb{I})^{-1}$, with $\psi_1^{\dagger} \coloneqq \psi_1^* (\psi_1 \psi_1^* + \gamma_N \mathbb{I})^{-1}$, $\psi_1^{\ddagger} \coloneqq \psi_1^* \psi_1^{\dagger}$, and $\psi_2 = -\beta_1 \psi_1$ from (22), due to triangle and operator-valued Hölder's inequality holds

$$\|\hat{\beta}_{1} - \beta_{1}\|_{\mathcal{S}} \leq \|\hat{\psi}_{2} - \psi_{2}\|_{\mathcal{S}} \left\|\hat{\psi}_{1}^{\dagger}\prod_{\hat{f}_{1}}^{\hat{f}_{M}}\right\|_{\mathcal{L}} + \|\psi_{2}\|_{\mathcal{S}} \left\|\hat{\psi}_{1}^{\dagger}\prod_{\hat{f}_{1}}^{\hat{f}_{M}} - \psi_{1}^{\dagger}\prod_{f_{1}}^{f_{M}}\right\|_{\mathcal{L}} + \left\|\beta_{1}\left(\psi_{1}^{\dagger}\prod_{f_{1}}^{f_{M}} - \mathbb{I}\right)\right\|_{\mathcal{S}}.$$
 (63)

Due to the definition of $\hat{\psi}_1^{\dagger}$ and the operator norm, $\|A\|_{\mathcal{L}}^2 = \|A^*A\|_{\mathcal{L}}$ for any $A \in \mathcal{L}_{\mathcal{H}}$, as $\hat{\psi}_1\hat{\psi}_1^*$ commutes with $(\hat{\psi}_1\hat{\psi}_1^* + \gamma_N\mathbb{I})^{-1}$, the projection operator $\coprod_{\hat{f}_1}^{\hat{f}_M}$ with $\hat{\psi}_1\hat{\psi}_1^*$ and $(\hat{\psi}_1\hat{\psi}_1^* + \gamma_N\mathbb{I})^{-1}$, and as $(\coprod_{\hat{f}_1}^{\hat{f}_M})^* = (\coprod_{\hat{f}_1}^{\hat{f}_M})^2 = \coprod_{\hat{f}_1}^{\hat{f}_M},$ it holds

$$\left\| \hat{\psi}_{1}^{\dagger} \prod_{\hat{f}_{1}}^{\hat{f}_{M}} \right\|_{\mathcal{L}}^{2} = \left\| \prod_{\hat{f}_{1}}^{\hat{f}_{M}} (\hat{\psi}_{1}^{\dagger})^{*} \hat{\psi}_{1}^{\dagger} \prod_{\hat{f}_{1}}^{\hat{f}_{M}} \right\|_{\mathcal{L}} = \left\| \hat{\psi}_{1} \hat{\psi}_{1}^{*} (\hat{\psi}_{1} \hat{\psi}_{1}^{*} + \gamma_{N} \mathbb{I})^{-2} \prod_{\hat{f}_{1}}^{\hat{f}_{M}} \right\|_{\mathcal{L}}$$
$$= \sup_{1 \le j \le M} \frac{\hat{\rho}_{j}}{(\hat{\rho}_{j} + \gamma_{N})^{2}} \le (\hat{\rho}_{M} + \gamma_{N})^{-1} =: \hat{p}_{M,N}.$$
(64)

We get as in the proof of Theorem B.1, from the definition of the projection operators, $\hat{\psi}_1^{\dagger}$ and ψ_1^{\dagger} , with $\hat{f}'_j \coloneqq \operatorname{sgn}\langle f_j, \hat{f}_j \rangle \hat{f}_j, \hat{x}_j \coloneqq \langle x, \hat{f}'_j \rangle, x_j \coloneqq \langle x, f_j \rangle, \hat{p}_{j,N} \coloneqq (\hat{\rho}_j + \gamma_N)^{-1}$ and $p_{j,N} \coloneqq (\rho_j + \gamma_N)^{-1}$:

$$\begin{aligned} \left\| \hat{\psi}_{1}^{\dagger} \prod_{\hat{f}_{1}}^{\hat{f}_{M}} - \psi_{1}^{\dagger} \prod_{f_{1}}^{\hat{f}_{M}} \right\|_{\mathcal{L}} \\ &= \sup_{\|\boldsymbol{x}\| \leq 1} \left\| \sum_{j=1}^{M} \hat{x}_{j} \hat{p}_{j,N} \hat{\psi}_{1}^{*}(\hat{f}_{j}') - x_{j} \hat{p}_{j,N} \psi_{1}^{*}(f_{j}) \right\| \\ &\leq \sup_{\|\boldsymbol{x}\| \leq 1} \sum_{j=1}^{M} \left\| \hat{x}_{j} \hat{p}_{j,N} \hat{\psi}_{1}^{*}(\hat{f}_{j}' - f_{j}) \right\| + \left\| (\hat{x}_{j} - x_{j}) \hat{p}_{j,N} \hat{\psi}_{1}^{*}(f_{j}) \right\| + \left\| x_{j} \left[\hat{p}_{j,N} \hat{\psi}_{1}^{*} - p_{j,N} \psi_{1}^{*} \right](f_{j}) \right\|. \end{aligned}$$
(65)

Due to elementary conversions and (61)-(62) holds

$$\sup_{\|\boldsymbol{x}\| \le 1} \sum_{j=1}^{M} \left\| \hat{x}_{j} \hat{p}_{j,N} \hat{\psi}_{1}^{*} (\hat{f}_{j}' - f_{j}) \right\| \le M \sup_{1 \le j \le M} \hat{p}_{j,N} \left\| \hat{\psi}_{1}^{*} (\hat{f}_{j}' - f_{j}) \right\| \le 2\sqrt{2} M \hat{p}_{M,N} P_{M} \left\| \hat{\psi}_{1} \right\|_{\mathcal{L}} \left\| \hat{\psi}_{1} \hat{\psi}_{1}^{*} - \psi_{1} \psi_{1}^{*} \right\|_{\mathcal{L}}.$$
(66)

Further, due to similar conversions as above,

$$\sup_{\|\boldsymbol{x}\| \leq 1} \sum_{j=1}^{M} \left\| (\hat{x}_{j} - x_{j}) \hat{p}_{j,N} \hat{\psi}_{1}^{*}(f_{j}) \right\| \\
\leq M \sup_{1 \leq j \leq M} \hat{p}_{j,N} \| \hat{f}_{j}' - f_{j} \| \| \hat{\psi}_{1}^{*}(f_{j}) \| \\
\leq 2\sqrt{2} M \hat{p}_{M,N} P_{M} \| \hat{\psi}_{1} \hat{\psi}_{1}^{*} - \psi_{1} \psi_{1}^{*} \|_{\mathcal{L}} \sup_{1 \leq j \leq M} \left(\| \psi_{1}^{*}(f_{j}) \| + \| [\hat{\psi}_{1}^{*} - \psi_{1}^{*}](f_{j}) \| \right) \\
\leq 2\sqrt{2} M \hat{p}_{M,N} P_{M} \| \hat{\psi}_{1} \hat{\psi}_{1}^{*} - \psi_{1} \psi_{1}^{*} \|_{\mathcal{L}} \left(\rho_{M}^{1/2} + \| \hat{\psi}_{1} - \psi_{1} \|_{\mathcal{L}} \right).$$
(67)

Moreover, we get

$$\begin{split} \sup_{\|\boldsymbol{x}\| \leq 1} & \sum_{j=1}^{M} \left\| x_{j} \left[\hat{p}_{j,N} \hat{\psi}_{1}^{*} - p_{j,N} \psi_{1}^{*} \right](f_{j}) \right\| \\ &= \sup_{1 \leq j \leq M} \left\| \left[\hat{p}_{j,N} \hat{\psi}_{1}^{*} - p_{j,N} \psi_{1}^{*} \right](f_{j}) \right\| \\ &\leq \sup_{1 \leq j \leq M} \left\| (\hat{p}_{j,N} - p_{j,N}) \hat{\psi}_{1}^{*}(f_{j}) \right\| + \sup_{1 \leq j \leq M} \left\| p_{j,N} \left[\hat{\psi}_{1}^{*} - \psi_{1}^{*} \right](f_{j}) \right\| \end{split}$$

$$\leq \sup_{1 \leq j \leq M} |\hat{p}_{j,N} - p_{j,N}| \sup_{1 \leq j \leq M} \left(\|\psi_1^*(f_j)\| + \|[\hat{\psi}_1^* - \psi_1^*](f_j)\|\right) + p_{M,N}\|\hat{\psi}_1 - \psi_1\|_{\mathcal{L}}$$

$$\leq P_M \Big[\hat{p}_{M,N} \|\hat{\psi}_1\hat{\psi}_1^* - \psi_1\psi_1^*\|_{\mathcal{L}} \big(\rho_M^{1/2} + \|\hat{\psi}_1 - \psi_1\|_{\mathcal{L}}\big) + \|\hat{\psi}_1 - \psi_1\|_{\mathcal{L}} \Big],$$
(68)

where we utilized $p_{M,N} \leq \rho_M^{-1} \leq P_M$. By combining (65)–(68), using (62), and $\|\hat{\psi}_1 - \psi_1\|_{\mathcal{L}} = o_{\mathbb{P}}(1)$ and thus $\|\hat{\psi}_1\|_{\mathcal{L}} = \|\psi_1\|_{\mathcal{L}} + o_{\mathbb{P}}(1)$ after Theorem B.1, for fixed K, M, and $L = L_N \to \infty$ holds

$$\begin{aligned} \left\| \hat{\psi}_{1}^{\dagger} \prod_{\hat{f}_{1}}^{f_{M}} - \psi_{1}^{\dagger} \prod_{f_{1}}^{f_{M}} \right\|_{\mathcal{L}} &\leq 2\sqrt{2}M\hat{p}_{M,N}P_{M} \| \hat{\psi}_{1}\hat{\psi}_{1}^{*} - \psi_{1}\psi_{1}^{*} \|_{\mathcal{L}} \Big[\| \hat{\psi}_{1} \|_{\mathcal{L}} + \rho_{M}^{1/2} + \| \hat{\psi}_{1} - \psi_{1} \|_{\mathcal{L}} \Big] \\ &+ P_{M} \Big[\hat{p}_{M,N} \| \hat{\psi}_{1}\hat{\psi}_{1}^{*} - \psi_{1}\psi_{1}^{*} \|_{\mathcal{L}} \Big(\rho_{M}^{1/2} + \| \hat{\psi}_{1} - \psi_{1} \|_{\mathcal{L}} \Big) + \| \hat{\psi}_{1} - \psi_{1} \|_{\mathcal{L}} \Big] \\ &\leq P_{M} \| \hat{\psi}_{1} - \psi_{1} \|_{\mathcal{L}} \Big[2\| \psi_{1} \|_{\mathcal{L}} \hat{p}_{M,N} \Big(2\sqrt{2}M \Big[\| \psi_{1} \|_{\mathcal{L}} + \rho_{M}^{1/2} \Big] + \rho_{M}^{1/2} \Big) + 1 + o_{\mathbb{P}}(1) \Big]. \end{aligned}$$

$$\tag{69}$$

Furthermore, similar as in the proof of Theorem B.1, we get with $\beta_1 = \beta_1(M)$ in (59), because f_j are the eigenfunctions of $\psi_1\psi_1^*$, and $\psi_1^{\ddagger} \coloneqq \psi_1\psi_1^*(\psi_1\psi_1^* + \gamma_N\mathbb{I})^{-1}$,

$$\left\|\beta_1\left(\psi_1^{\ddagger} \prod_{f_1}^{f_M} - \mathbb{I}\right)\right\|_{\mathcal{S}}^2 \le \gamma_N^2 p_{M,N}^2 \left\|\sum_{k=1}^M \sum_{\ell=1}^M b_{k,\ell,1} f_\ell\right\|^2 = \gamma_N^2 p_{M,N}^2 \|\beta_1(M)\|_{\mathcal{S}}^2.$$
(70)

By plugging the inequalities (64), (69)–(70) into (63), and afterwards using that for all *i* holds $\|\hat{\psi}_i - \psi_i\|_{\mathcal{L}} \leq \|\hat{\psi}_i - \psi_i\|_{\mathcal{S}} \leq \|\hat{\Psi}_L - \Psi_L\|_{\mathcal{S}}$, we get

$$\begin{aligned} \|\hat{\beta}_{1} - \beta_{1}(M)\|_{\mathcal{S}} \\ &\leq \hat{p}_{M,N}^{1/2} \|\hat{\psi}_{2} - \psi_{2}\|_{\mathcal{S}} + \|\psi_{2}\|_{\mathcal{S}} P_{M} \|\hat{\psi}_{1} - \psi_{1}\|_{\mathcal{L}} \\ &\times \left[2\|\psi_{1}\|_{\mathcal{L}} \,\hat{p}_{M,N} \left(2\sqrt{2}M \left[\|\psi_{1}\|_{\mathcal{L}} + \rho_{M}^{1/2} \right] + \rho_{M}^{1/2} \right) + 1 + o_{\mathbb{P}}(1) \right] + \gamma_{N} \, p_{M,N} \|\beta_{1}(M)\|_{\mathcal{S}} \\ &\leq \|\hat{\Psi}_{L} - \Psi_{L}\|_{\mathcal{S}} \left(\hat{p}_{M,N}^{1/2} + \|\psi_{2}\|_{\mathcal{S}} P_{M} \left[2\|\psi_{1}\|_{\mathcal{L}} \,\hat{p}_{M,N} \left(2\sqrt{2}M \left[\|\psi_{1}\|_{\mathcal{L}} + \rho_{M}^{1/2} \right] + \rho_{M}^{1/2} \right) + 1 + o_{\mathbb{P}}(1) \right] \right) \\ &+ \gamma_{N} \, p_{M,N} \|\beta_{1}(M)\|_{\mathcal{S}} \,. \end{aligned}$$

$$(71)$$

Since K, M are fixed, $L = L_N \to \infty$, $\hat{p}^a_{M,N} \xrightarrow{\mathbb{P}} \rho_M^{-a}$ after (60), (62) and Theorem B.1 for any $a \neq 0$, and $\gamma_N = o(\lambda_K^{-1} \Lambda_K L^2 N^{-1/2})$, we get $\|\hat{\beta}_1 - \beta_1(M)\|_{\mathcal{S}} = O_{\mathbb{P}}(\lambda_K^{-1} \Lambda_K L^2 N^{-1/2})$. Further, $\hat{\alpha}_1 = \hat{\psi}_1 - \hat{\beta}_1$ and $\alpha_1(K, M) = \psi_1(K) - \beta_1(M)$ entail

$$\|\hat{\alpha}_{1} - \alpha_{1}(K, M)\|_{\mathcal{S}} \leq \|\bar{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} + \|\beta_{1} - \beta_{1}(M)\|_{\mathcal{S}}$$
$$= O_{\mathbb{P}} (\lambda_{K}^{-1} \Lambda_{K} L^{2} N^{-1/2}).$$

Hence, our claim is proven.

Proof of Theorem 4.1. By combining the inequalities in the proofs of Theorems 3.1, B.1–B.2, for $\min\{K_N, L_N, M_N\} \to \infty$ holds because of $\hat{p}_{M,N} = \mathcal{O}_{\mathbb{P}}(\rho_M^{-1})$ according to Lemma B.1 with $\|\hat{\psi}_1\hat{\psi}_1^* - \psi_1\psi_1^*\|_{\mathcal{L}} = \mathcal{O}_{\mathbb{P}}(K^{-\beta}), M^{1+\gamma}\rho_M^{-1}P_M = \mathcal{O}(K^{\beta}) \text{ and } \gamma_N = \mathcal{O}(\rho_M M^{-\gamma}), \text{ and Assumption 4.4}$ indeed $\|\hat{\beta}_1 - \beta_1\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(M^{-\gamma}), \text{ thus } \|\hat{\alpha}_1 - \alpha_1\|_{\mathcal{S}} \le \|\hat{\Psi}_L - \Psi_L\|_{\mathcal{S}} + \|\hat{\beta}_1 - \beta_1\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(M^{-\gamma}).$

Assumption B.3. For fixed $M \in \mathbb{N}$, there are constants $b'_{i,j,q} \in \mathbb{R}$ so that

$$B_q = B_q(M) = \sum_{i=1}^M \sum_{j=1}^M b'_{i,j,q}(h_i \otimes d_j).$$

Due to the identity (24), which is equivalent to $\alpha_i = \psi_i + B_{[i]} \Psi_{[i]}^{\prime\prime\prime}$ for $i = 1, \ldots, p$, we get similar as in Section 3.2, a finite-dimensional representation of α_i via the finite-dimensional representations of the operators ψ_i and β_j . Namely, it holds

$$\alpha_i = \alpha_i(K, M) \coloneqq \psi_i(K) + B_{[i]}(M) \Psi_{[i]}^{\prime\prime}(K), \quad 1 \le i \le p.$$

Theorem B.3. Let $(X_k)_{k\in\mathbb{Z}}$ be the fARMA(p,q) process in (20), let the assumptions of Theorem B.1 and Assumptions 4.5–4.7 and B.3 hold, and let $\gamma_N = o(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2})$. Then, for fixed K, M, and $L = L_N \to \infty$ as in Theorem B.1, it holds

$$\max_{1 \le i \le p, 1 \le j \le q} \left\{ \| \hat{\alpha}_i - \alpha_i(K, M) \|_{\mathcal{S}}, \| \hat{\beta}_j - \beta_j(M) \|_{\mathcal{S}} \right\} = \mathcal{O}_{\mathbb{P}} \left(\lambda_K^{-1} \Lambda_K L^2 N^{-1/2} \right).$$
(72)

Proof. This proof proceeds as the proof of Theorem B.2. Herein, $(\zeta_j, h_j)_{j \in \mathbb{N}}$, $(\hat{\zeta}_j, \hat{h}_j)_{j \in \mathbb{N}}$ are the eigenpair sequences of $\prod \prod^*, \hat{\prod} \hat{\prod}^* \in \mathcal{N}_{\mathcal{H}^q}$, respectively, with $\prod, \hat{\prod} \in \mathcal{S}_{\mathcal{H}^q}$ defined in (27) and (29), respectively, and where $\prod \in \mathcal{S}_{\mathcal{H}^q}$ and $\prod \prod^* \in \mathcal{N}_{\mathcal{H}^q}$ hold after Assumption 3.5 and (37). Further,

$$\sup_{j\in\mathbb{N}} |\hat{\zeta}_j - \zeta_j| \le \left\| \hat{\prod} \hat{\prod}^* - \prod \hat{\prod}^* \right\|_{\mathcal{L}},\tag{73}$$

with $\zeta_1 > \zeta_2 > \cdots > 0$ after Assumptions 4.6–4.7, and

$$\sup_{1 \le j \le M} \|\hat{h}_j' - h_j\| \le 2\sqrt{2}Z_M \left\| \hat{\prod} \hat{\prod}^* - \prod M^* \right\|_{\mathcal{L}}, \quad M \in \mathbb{N},$$
(74)

with $\hat{h}'_j \coloneqq \operatorname{sgn}\langle \hat{h}_j, h_j \rangle \hat{h}_j$ and $Z_M \coloneqq \operatorname{sup}_{1 \le j \le M} (\zeta_j - \zeta_{j+1})^{-1}$. Due to the definition of the operatorvalued vectors and matrices, and (35), holds

$$\left\|\hat{\Pi} - \Pi\right\|_{\mathcal{L}} \le \sum_{i=0}^{q-1} \|\hat{\Psi}'_{[i]} - \Psi'_{[i]}\|_{\mathcal{L}} .$$
(75)

Consequently, as $\|\hat{\Psi}'_{[i]} - \Psi'_{[i]}\|_{\mathcal{S}} \le \|\hat{\Psi}_L - \Psi_L\|_{\mathcal{S}}$ for $L \ge p + q + i - 1$, we obtain for $L \ge p + 2q - 2$,

$$\left\|\hat{\Pi}\hat{\Pi}^{*}-\Pi\Pi^{*}\right\|_{\mathcal{L}} \leq \left\|\hat{\Pi}-\Pi\right\|_{\mathcal{L}}\left(\left\|\Pi\right\|_{\mathcal{L}}+\left\|\hat{\Pi}\right\|_{\mathcal{L}}\right)$$
$$\leq q\|\hat{\Psi}_{L}-\Psi_{L}\|_{\mathcal{S}}\left(\left\|\Pi\right\|_{\mathcal{L}}+\left\|\hat{\Pi}\right\|_{\mathcal{L}}\right).$$
(76)

At first, we show the claimed consistency results for the operators β_j . For $\hat{B}_q = -\hat{\Psi}'_{[q]} \hat{\Pi}^{\dagger} \coprod_{\hat{h}_1}^{\hat{h}_M}$, with $\hat{\Pi}^{\dagger} = \hat{\Pi}^* (\hat{\Pi} \hat{\Pi}^* + \gamma_N \mathbb{I})^{-1}$ and $\Pi^{\dagger} = \Pi^* (\Pi \Pi^* + \gamma_N \mathbb{I})^{-1}$, where $(\gamma_N)_{N \in \mathbb{N}} \subset (0, \infty)$ is a sequence with $\gamma_N \to 0$, and with $\Pi^{\ddagger} := \Pi \Pi^{\dagger}$ and $\Psi'_{[q]} = -B_q \prod$ in (26), holds

$$\hat{B}_{q} - B_{q} = \left(\Psi'_{[q]} - \hat{\Psi}'_{[q]}\right) \hat{\prod}^{\dagger} \prod_{\hat{h}_{1}}^{\hat{h}_{M}} + \Psi'_{[q]} \left(\prod^{\dagger} \prod_{h_{1}}^{h_{M}} - \hat{\prod}^{\dagger} \prod_{\hat{h}_{1}}^{\hat{h}_{M}}\right) + B_{q} \left(\prod^{\ddagger} \prod_{h_{1}}^{h_{M}} - \mathbb{I}\right)$$

Subsequently, by using triangle inequality and operator-valued Hölder's inequality, the definition of the operators and the norms below, and ideas from the proof of Theorem B.2, with $B_q = B_q(M)$ in Assumption B.3 holds for sufficiently large L with $\hat{z}_{M,N} \coloneqq (\hat{\zeta}_M + \gamma_N)^{-1}$ and $z_{M,N} \coloneqq (\zeta_M + \gamma_N)^{-1}$:

$$\|\hat{B}_{q} - B_{q}\|_{\mathcal{S}} \leq \hat{z}_{M,N}^{1/2} \|\hat{\Psi}_{L} - \Psi_{L}\|_{\mathcal{S}} + \|\Psi'_{[q]}\|_{\mathcal{S}} \left\|\prod^{\dagger}\prod_{h_{1}}^{h_{M}} - \prod^{\dagger}\prod_{\hat{h}_{1}}^{\hat{h}_{M}} \right\|_{\mathcal{L}} + \gamma_{N} z_{M,N} \|B_{q}(M)\|_{\mathcal{S}}.$$
 (77)

Thereby, similar as in the proof of Theorem B.2, by using (73)–(76) and $\|\cdot\|_{\mathcal{L}} \leq \|\cdot\|_{\mathcal{S}}$, we get

$$\begin{split} &\|\Pi^{\dagger} \prod_{h_{1}}^{h_{M}} - \hat{\Pi}^{\dagger} \prod_{\hat{h}_{1}}^{h_{M}} \|_{\mathcal{L}} \\ &\leq 2\sqrt{2}M\hat{z}_{M,N}Z_{M} \|\hat{\Pi} \prod^{*} - \Pi\Pi^{*} \|_{\mathcal{L}} \left(\|\hat{\Pi}\|_{\mathcal{L}} + \zeta_{M}^{1/2} + \|\hat{\Pi} - \Pi\|_{\mathcal{L}} \right) \\ &+ \zeta_{M}^{-1} \Big[\hat{z}_{M,N} \|\hat{\Pi} \prod^{*} - \Pi\Pi^{*} \|_{\mathcal{L}} \left(\zeta_{M}^{1/2} + \|\hat{\Pi} - \Pi\|_{\mathcal{L}} \right) + \|\hat{\Pi} - \Pi\|_{\mathcal{L}} \Big] \\ &= qZ_{M} \|\hat{\Psi}_{L} - \Psi_{L} \|_{\mathcal{S}} \Big[2 \|\Pi\|_{\mathcal{L}} \hat{z}_{M,N} \Big(2\sqrt{2}M \Big[\|\Pi\|_{\mathcal{L}} + \zeta_{M}^{1/2} \Big] + \zeta_{M}^{1/2} \Big) + 1 + o_{\mathbb{P}}(1) \Big]. \end{split}$$
(78)

By plugging (78) into (77), for fixed K, M, and $L = L_N \to \infty$, since $\hat{z}^a_{M,N} \xrightarrow{\mathbb{P}} \zeta^{-a}_M$ after (73), (76) and Theorem B.1 for any $a \neq 0$, and $\gamma_N = o(\lambda_K \Lambda_K L^{3/2} N^{-1/2})$, we get $\|\hat{B}_q - B_q(M)\|_{\mathcal{S}} = O_{\mathbb{P}}(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2})$, and subsequently $\|\hat{\beta}_j - \beta_j(M)\|_{\mathcal{S}} = O_{\mathbb{P}}(\lambda_K^{-1}\Lambda_K L^2 N^{-1/2})$ for each j.

Moreover, due to $\hat{\alpha}_i = \hat{\psi}_i + \hat{B}_{[i]} \hat{\Psi}_{[i]}^{\prime\prime\prime\top}, \ \alpha_i(K,M) = \psi_i(K) + \hat{B}_{[i]}(M) \Psi_{[i]}^{\prime\prime\prime\top}(K), \ \alpha_i = \beta_j = \hat{\beta}_j = 0$ for i > p and j > q, and elementary conversions, we obtain for $L \ge p + q + i - 1$ for any i, p, q:

$$\begin{aligned} \|\hat{\alpha}_{i} - \alpha_{i}(K, M)\|_{\mathcal{S}} &\leq \|\hat{\psi}_{i} - \psi_{i}(K)\|_{\mathcal{S}} + \|\hat{B}_{[i]}(M)\|_{\mathcal{L}}\|\hat{\Psi}_{[i]}^{\prime\prime\prime} - \Psi_{[i]}^{\prime\prime\prime}(K)\|_{\mathcal{S}} + \|\Psi_{[i]}^{\prime\prime\prime}(K)\|_{\mathcal{L}}\|\hat{B}_{[i]} - B_{[i]}(M)\|_{\mathcal{S}} \\ &\leq (1 + \|\hat{B}_{[i]}(M)\|_{\mathcal{L}})\|\hat{\Psi}_{L} - \Psi_{L}(K)\|_{\mathcal{S}} + \|\Psi_{[i]}^{\prime\prime\prime\prime}(K)\|_{\mathcal{L}}\|\hat{B}_{q} - B_{q}(M)\|_{\mathcal{S}}. \end{aligned}$$

Hence, due to $\|\hat{B}_{[i]}(M)\|_{\mathcal{L}} \leq \|B_{[i]}(M)\|_{\mathcal{L}} + \|\hat{B}_{[i]} - B_{[i]}(M)\|_{\mathcal{L}}$ and $\|\hat{B}_{[i]} - B_{[i]}(M)\|_{\mathcal{L}} \leq \|\hat{B}_q - B_q(M)\|_{\mathcal{L}}$, we have $\|\hat{\alpha}_i - \alpha_i(K, M)\|_{\mathcal{S}} = \mathcal{O}_{\mathbb{P}}(\lambda_K^{-1}\Lambda_K L^{3/2}N^{-1/2})$ for any *i*. Therefore, (72) is shown. \Box

Proof of Theorem 4.2. Based on the arguments in the proofs of Theorems 4.1, B.3, Assumption 4.8, for min{ K_N, L_N, M_N } $\rightarrow \infty$ holds after $z_{M,N} = O_{\mathbb{P}}(\zeta_M^{-1})$ according to Lemma B.1 with $\|\hat{\Pi}\hat{\Pi}^* - \prod_{i=1}^{n} \|_{\mathcal{L}} = O_{\mathbb{P}}(K^{-\beta}), M^{1+\gamma}\zeta_M^{-1}Z_M = O(K^{\beta})$ and $\gamma_N = o(\zeta_M M^{-\gamma})$, that $\|\hat{B}_q - B_q\|_{\mathcal{S}} = O_{\mathbb{P}}(M^{-\gamma})$, so $\|\hat{\beta}_j - \beta_j\|_{\mathcal{S}} = O_{\mathbb{P}}(M^{-\gamma})$ for each j. Further, due to $\hat{\alpha}_i = \hat{\psi}_i + \hat{B}_{[i]}\hat{\Psi}_{[i]}^{\prime\prime\prime} + in$ (30), $\alpha_i = \psi_i + B_{[i]}\Psi_{[i]}^{\prime\prime\prime} + in$, and similar conversions as in Theorem B.3 give for sufficiently large L for each i,

$$\|\hat{\alpha}_{i} - \alpha_{i}\|_{\mathcal{S}} \leq (1 + \|B_{[i]}\|_{\mathcal{L}}) \|\Psi_{L} - \Psi_{L}\|_{\mathcal{S}} + \|\Psi_{[i]}^{\prime\prime\prime}\|_{\mathcal{L}} \|B_{q} - B_{q}\|_{\mathcal{S}} + o_{\mathbb{P}}(M^{-\gamma})$$

= $O_{\mathbb{P}}(M^{-\gamma}).$