

Noncommutative Lightcones from Quantum $SO(2,1)$ Conformal Groups

Martina Adamo^{*}, Angel Ballesteros[†] and Flavio Mercati[‡]

Departamento de Física, Universidad de Burgos, 09001 Burgos, Spain

July 18, 2024

Abstract

Five new families of noncommutative lightcones in $2+1$ dimensions are presented as the quantizations of the inequivalent Poisson homogeneous structures that emerge when the lightcone is constructed as a homogeneous space of the $SO(2,1)$ conformal group. Each of these noncommutative lightcones maintains covariance under the action of the respective quantum deformation of the $SO(2,1)$ conformal group. We discuss the role played by $SO(2,1)$ automorphisms in the classification of inequivalent Poisson homogeneous lightcones, as well as the geometric aspects of this construction. The localization properties of the novel quantum lightcones are analyzed and shown to be deeply connected with the geometric features of the Poisson homogeneous spaces.

KEYWORDS: Noncommutative spacetimes, quantum groups, Poisson–Lie groups, homogeneous spaces, quantization, conformal symmetry, localization

1 Introduction

Quantum Field Theory (QFT) on noncommutative spacetimes was initially proposed in 1947 [1] as a way to cut off the ultraviolet divergences of quantum electrodynamics. The idea gained new momentum after the limitations of describing quantum gravity as a perturbative QFT were highlighted. In particular, the emergence of the Planck length as the natural scale at which quantum gravity breaks down as an effective field theory strongly suggests that there are limitations to the localizability of spacetime events [2], and the standard smooth structure of spacetime might have to be replaced with a new mathematical description. After dramatic progress in the 1970s–80s on the foundations of such a description [3–5], we now have noncommutative formulations of all the basic tools of geometry and topology, which allow us, in principle, to study QFT on noncommutative spaces [6]. Quantum groups [7–9] are the noncommutative generalization of Lie groups and play a central role in the description of the symmetries of noncommutative spaces [10–12]. The ‘semiclassical’ counterpart of quantum groups are Poisson–Lie (hereafter PL) groups, which are standard Lie groups equipped with a Poisson bracket that is compatible with the group structure [9, 13]. A PL group contains all the essential structural information concerning the noncommutativity properties induced by the associated quantum group, with the exception of ordering ambiguities that may arise in the quantization process. The advantage of working with PL groups instead of their quantum counterparts is their computational simplicity, while still capturing most of the nonclassical aspects of a noncommutative space.

A key open question in the physics of noncommutative spaces is the fate of causality. These theories inherently possess a weakened notion of locality (see, *e.g.*, [14, 15]), which is bound to interact in unpredictable ways with the notion of causality, another pillar of modern QFT and classical relativistic theories such as General Relativity. In particular, the possibility of fuzzy/noncommutative lightcones emerging in noncommutative QFTs has been contemplated in [16–19]. Moreover, recent works have explicitly highlighted that the difficulties of defining a sharp notion of causal relations between ‘points’ in certain noncommutative spacetimes can be circumvented by making use of their associated noncommutative spaces of worldlines [20–23].

^{*}marti.adamo@gmail.com, madamo@ubu.es

[†]angelb@ubu.es

[‡]flavio.mercati@gmail.com

In this paper, we present a novel approach to this problem by explicitly constructing noncommutative lightcones in three spacetime dimensions, seen as quantum homogeneous spaces of a quantum $\text{SO}(2,1)$ group. In this framework, the noncommutative lightcone is constructed with no reference to any ambient noncommutative space, thus avoiding any causality issues arising from the fuzziness of the embedding. In this way, five new noncommutative lightcones with quantum conformal group invariance will be presented and analyzed.

In the next section, the classical (both future or past) lightcones \mathcal{L}_{\pm} will be constructed as the homogeneous spaces that are obtained by quotienting the $\text{SO}(2,1)$ group by the stabilizing subgroup of a lightlike vector, which we will call K . Under the appropriate $\text{SO}(2,1)$ parametrization, the orbits of a lightlike vector under the (Lorentz) group action coincide with the future- or past-oriented half lightcones. The conformal group $\text{SO}(2,1)$ then acts as isometry group on the lightcone, and a geometric picture involving the underlying conformal symmetries emerges.

Afterwards, in order to define an intrinsic noncommutative structure on the lightcone, we consider its Poisson homogeneous (hereafter PH) structures. To this aim, we introduce in Section 3 the PL structures on $\text{SO}(2,1)$, which fall into three distinct families (up to Lie algebra automorphisms), characterized by three classes of classical r -matrices [24–26]. At this point, a deeper insight into $\text{SO}(2,1)$ automorphisms turns out to be needed, since we realized that some of them do not preserve the quotient structure $\mathcal{L}_{\pm} = \text{SO}(2,1)/K$ defining the lightcones, a fact that – to the best of our knowledge – has not been previously considered in the literature. The complete study of $\text{SO}(2,1)$ automorphisms acting on the lightcones \mathcal{L}_{\pm} is presented in Section 4. As a consequence, two of the families of $\text{SO}(2,1)$ r -matrices split into two inequivalent classes.

Therefore, we have to consider five different families of PL structures on $\text{SO}(2,1)$, which give rise to five nonisomorphic PH space structures on the lightcones that are explicitly presented in Section 5. This is a consequence of the property, shared by all five families, of coisotropy with respect to the subgroup K (see, *e.g.*, [27–30]).

The associated noncommutative lightcones \mathcal{L}_{\pm}^{ℓ} can then be obtained in Section 6 as the quantization of the associated PH spaces, defining the algebras generating the corresponding quantum homogeneous spaces [31]. Following the approach introduced in [14, 15, 32], the limitations to the localization of regions on the lightcones imposed by such noncommutative geometries are studied in terms of the representation theory of the corresponding noncommutative algebras, and they turn out to be directly connected with the geometric properties of the PH lightcones they come from. Finally, some remarks and open problems close the paper.

2 The lightcone in 2+1 dimensions as a homogeneous space

The Lie algebra $\mathfrak{so}(2,1)$ of the pseudo-orthogonal group in 2+1 dimensions, expressed in terms of the antisymmetric generators $\mathbb{M}^{IJ} = -\mathbb{M}^{JI}$, is given by

$$[\mathbb{M}_{IJ}, \mathbb{M}_{KL}] = \eta_{IL} \mathbb{M}_{JK} + \eta_{JK} \mathbb{M}_{IL} - \eta_{IK} \mathbb{M}_{JL} - \eta_{JL} \mathbb{M}_{IK}, \quad (1)$$

where $\eta_{IJ} = \text{diag}(-1, 1, 1)$ is the metric of the three-dimensional ambient Minkowski space, with coordinates labeled as $I, J, K, \dots = 0, 1, 2$.

The orbits of this group in the 3D ambient Minkowski space belong to three classes: spacelike two-sheeted ‘mass’ hyperboloids, one-sheeted timelike hyperboloids, and the future and past lightcones, from which the origin $(0, 0, 0)$ can be singled out as a fixed point under the action of the group. In this paper, we are interested in the lightcones. Each of them can be defined as the homogeneous space obtained as the quotient of $\text{SO}(2,1)$ by the isotropy subgroup of a lightlike vector. If the vector is future-oriented, one gets the future lightcone, and if it is past-oriented, the past lightcone is obtained.

To explicitly build the lightcones, we need to introduce the defining representation of the $\mathfrak{so}(2,1)$ Lie algebra as three-dimensional real matrices $(a, b, c, \dots = 0, 1, 2)$

$$\rho(\mathbb{M}_{IJ})^a{}_b = \delta_I^a \eta_{Jb} - \delta_J^a \eta_{Ib}. \quad (2)$$

Without loss of generality, we can choose the following lightlike stabilizing vectors as

$$v_{\pm}^a = \pm(l, 0, -l), \quad l > 0, \quad (3)$$

where v_{+}^a generates the future lightcone and v_{-}^a generates the past one. The positive quantity l is a dimensional constant that gives the unit of length to the ambient coordinates. Both v_{+}^a and v_{-}^a have the same isotropy subgroup, which is generated by $\mathbb{K} = \mathbb{M}_{21} - \mathbb{M}_{01}$.

Note that the the Lorentz group of $(d+1)$ -dimensional Minkowski space and the conformal group of the $(d-1)$ -dimensional Euclidean space, $\text{Conf}(d-1)$, are isomorphic. The Lie algebras of the two groups coincide, and the natural basis of the conformal algebra $\mathfrak{conf}(1)$, made of the generators of translations \mathbb{T} , dilations \mathbb{D} , and special conformal transformations \mathbb{K} , is related to the basis (1) of $\mathfrak{so}(2,1)$ by the following linear redefinition:

$$\mathbb{D} = \mathbb{M}_{02}, \quad \mathbb{T} = \mathbb{M}_{01} + \mathbb{M}_{21}, \quad \mathbb{K} = \mathbb{M}_{21} - \mathbb{M}_{01}. \quad (4)$$

The generator \mathbb{K} is the only one that stabilizes the vector (3), since

$$\rho(\mathbb{D})^a{}_b v_{\pm}^b = v_{\pm}^a, \quad \rho(\mathbb{T})^a{}_b v_{\pm}^b = \pm 2l \delta^a{}_1, \quad \rho(\mathbb{K}_i)^a{}_b v_{\pm}^b = 0, \quad (5)$$

which suggests a local factorization of the connected component to the identity of $\text{SO}(2,1)$ (the orthochronous component $\text{SO}_+(2,1)$) that exploits this fact and reads

$$e^{\tau \rho(\mathbb{T})} = \begin{pmatrix} 1 + \frac{\tau^2}{2} & \tau & -\frac{\tau^2}{2} \\ \tau & 1 & -\tau \\ \frac{\tau^2}{2} & \tau & 1 - \frac{\tau^2}{2} \end{pmatrix}, \quad e^{\lambda \rho(\mathbb{D})} = \begin{pmatrix} \text{ch } \lambda & 0 & \text{sh } \lambda \\ 0 & 1 & 0 \\ \text{sh } \lambda & 0 & \text{ch } \lambda \end{pmatrix}, \quad e^{\sigma \rho(\mathbb{K})} = \begin{pmatrix} 1 + \frac{\sigma^2}{2} & -\sigma & \frac{\sigma^2}{2} \\ -\sigma & 1 & -\sigma \\ -\frac{\sigma^2}{2} & \sigma & 1 - \frac{\sigma^2}{2} \end{pmatrix}. \quad (6)$$

Now, a generic $\text{SO}_+(2,1)$ element is given by

$$g(\tau, \lambda, \sigma) = e^{\tau \rho(\mathbb{T})} e^{\lambda \rho(\mathbb{D})} e^{\sigma \rho(\mathbb{K})} \\ = \begin{pmatrix} \frac{1}{2}(e^{\lambda-2\tau\sigma+e^{-\lambda}(1+\tau^2)}(1+\sigma^2)) & \tau - e^{-\lambda}(1+\tau^2)\sigma & \frac{1}{2}(e^{\lambda-2\tau\sigma+e^{-\lambda}(1+\tau^2)}(-1+\sigma^2)) \\ -\sigma + e^{-\lambda}\tau(1+\sigma^2) & 1 - 2e^{-\lambda}\tau\sigma & -\sigma + e^{-\lambda}\tau(-1+\sigma^2) \\ \frac{1}{2}(e^{\lambda-2\tau\sigma+e^{-\lambda}(-1+\tau^2)}(1+\sigma^2)) & \tau - e^{-\lambda}(-1+\tau^2)\sigma & \frac{1}{2}(e^{\lambda-2\tau\sigma+e^{-\lambda}(-1+\tau^2)}(-1+\sigma^2)) \end{pmatrix}. \quad (7)$$

This factorization is appropriate for constructing the future and past lightcones as $\text{SO}(2,1)$ homogeneous spaces in the form $\mathcal{L}_{\pm} = \text{Conf}(1)/\mathbb{K}$ since by acting with a generic $\text{SO}_+(2,1)$ element (7) on the lightlike vector (3), the action of \mathbb{K} , factorized to the right, leaves v_{\pm}^a invariant, thus generating a pair of two-dimensional surfaces parametrized by the group parameters τ and λ only.

Given the local coordinates τ , λ , and σ , the group axioms of $\text{SO}(2,1)$ can be encoded into a binary multiplication law

$$g(\tau_1, \lambda_1, \sigma_1) g(\tau_2, \lambda_2, \sigma_2) = g\left(\tau_1 + \frac{\tau_2 e^{\lambda_1}}{1 - \tau_2 \sigma_1}, \lambda_1 + \lambda_2 - 2 \log(\tau_2 \sigma_1 - 1), \sigma_2 + \frac{\sigma_1 e^{\lambda_2}}{1 - \tau_2 \sigma_1}\right), \quad (8)$$

an inverse map

$$g^{-1}(\tau, \lambda, \sigma) = g\left(\frac{\tau}{\tau\sigma - e^{\lambda}}, \lambda - 2 \log(e^{\lambda} - \tau\sigma), \frac{\sigma}{\tau\sigma - e^{\lambda}}\right), \quad (9)$$

and the specification of the coordinates of the identity element $g(0, 0, 0) = e$. The multiplication map (8) allows the calculation of the left- and right-invariant vector fields associated with the Lie algebra generators \mathbb{T} , \mathbb{D} , and \mathbb{K} in the coordinates τ , λ , σ

$$\begin{aligned} X_{\mathbb{T}}^L &= e^{\lambda} \partial_{\tau} + 2\sigma \partial_{\lambda} + \sigma^2 \partial_{\sigma}, & X_{\mathbb{T}}^R &= \partial_{\tau}, \\ X_{\mathbb{D}}^L &= \partial_{\lambda} + \sigma \partial_{\sigma}, & X_{\mathbb{D}}^R &= \tau \partial_{\tau} + \partial_{\lambda}, \\ X_{\mathbb{K}}^L &= \partial_{\sigma}, & X_{\mathbb{K}}^R &= \tau^2 \partial_{\tau} + 2\tau \partial_{\lambda} + e^{\lambda} \partial_{\sigma}. \end{aligned} \quad (10)$$

It is important to realize that the coordinates τ , λ , σ do not cover all of $\text{SO}_+(2,1)$. In fact, the spatial rotations by π given by

$$e^{\pi \mathbb{M}_{12}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad (11)$$

can only be reached through (7) with a singular choice of coordinates: $\tau \rightarrow \infty$, $\sigma \rightarrow \infty$, and $\lambda \rightarrow \infty$, in such a way that $e^{-\lambda\tau^2} \rightarrow 1$ and $e^{-\lambda\sigma^2} \rightarrow 1$. Therefore, all transformations that involve such a π rotation will belong to the boundary of our coordinate patch, and we can reach them by multiplying a regular group element of the form (7) by the matrix (11).

Notice also that $\text{SO}(2,1)$ has two connected components: $\text{SO}_+(2,1)$, the orthochronous component, and $\text{SO}_-(2,1)$, the non-orthochronous one. The three-dimensional representation of the elements of the latter can be obtained from (7) by multiplying $g(\tau, \lambda, \sigma)$ by the matrix

$$PT = \text{diag}(-1, 1, -1), \quad \text{where} \quad P = \text{diag}(1, 1, -1), \quad T = \text{diag}(-1, 1, 1), \quad (12)$$

which has unit determinant, and therefore belongs to $\text{SO}(2,1)$, but is not connected to the identity. This matrix implements a parity transformation and a time reversal, and connects the two stabilizing vectors

$$v_+^a = (PT)^a_b v_-^b, \quad (13)$$

so that the past lightcone can also be seen as generated by the application of non-orthochronous transformations to the future-oriented vector v_+^a .

The two lightcones can be described in terms of embedding coordinates x_\pm^I within the ambient (2+1)-dimensional Minkowski space. To avoid confusion between the coordinates on the group (τ , λ , and σ) and those on the homogeneous space, we will replace τ with y and λ with z , namely

$$x_\pm^I(y, z) = g(y, z, \sigma) v_\pm = \pm e^{-z} l \begin{pmatrix} y^2 + 1 \\ 2y \\ y^2 - 1 \end{pmatrix}. \quad (14)$$

Notice how both coordinate y and z are dimensionless (they are combinations of angles and rapidities, and therefore dimensionless in natural units). It is the constant l that gives the dimensions of a length (or time, which is the same in natural units) to the embedding coordinates, which satisfy the conditions

$$\eta_{IJ} x_\pm^I(y, z) x_\pm^J(y, z) = 0, \quad \text{sgn}(x_\pm^0(y, z)) = \pm 1, \quad (15)$$

indicating that $x_+^I(y, z)$ defines the future lightcone \mathcal{L}_+ , and $x_-^I(y, z)$ the past one, \mathcal{L}_- .

The action of a generic $\text{SO}_+(2,1)$ conformal group element (7) on ambient coordinates is

$$(x'_\pm)^I(y, z) = g(\tau, \lambda, \sigma)^I_J x^J(y, z) = g(\tau, \lambda, \sigma)^I_J g(y, z, \sigma)^J_K v_\pm^K = x_\pm^I(y', z'), \quad (16)$$

which means that coordinates y and z transform according to (8), namely

$$y' = \tau + \frac{ye^\lambda}{1-\sigma y}, \quad z' = z + \lambda - 2 \log(\sigma y - 1). \quad (17)$$

Notice that the ray $x^0 = x^2$ lies on the boundary of the coordinate patch covered by y and z , since to reach this light ray, we must simultaneously take the limits $y \rightarrow \infty$ and $z \rightarrow \infty$, in such a way that $e^{-z}y^2$ remains finite. In fact, the ray $x^0 = x^2$ is connected to the origin of the homogeneous space by the transformation (11).

It is worth stressing that the geometrical interpretation of $\text{SO}(2,1) \simeq \text{Conf}(1)$ as the group of conformal isometries of the real line is recovered by further quotienting by the dilations, which move the point on the lightcone along a light ray, either towards the origin or towards infinity (see Fig. 1). Quotienting the future lightcone by the action of dilations then yields the *celestial sphere*, which is the space of future-directed (or past-directed) light rays, that is, the future or past boundary of the lightcone. This space can be coordinatized with the polar angle $\theta = \arctan(x^2/x^1)$, and then mapped to the real line via a stereographic projection (excluding one point, the point at infinity, which coincides with the light ray $x^0 = x^2$ that is not covered by the coordinates y, z)

$$\frac{\cos \theta}{1 - \sin \theta} = \frac{x^2 + \sqrt{(x^1)^2 + (x^2)^2}}{x^1} = y, \quad (18)$$

which provides a geometrical interpretation of the coordinate y . As shown in (17), the action onto y of translations and dilations is linear: $y' = y + \tau$, $y' = e^\lambda y$, while special conformal transformations coincide with an inversion, followed by a translation, followed by a second inversion, $y' = y/(1 - \sigma y)$.




Figure 1: The future and past lightcones in the three-dimensional Minkowski ambient space. The coordinate lines of y (in blue) are obtained through pure translations and lie on the intersection of the cone with 45° planes. Those of z (in red) correspond to dilations and lie on the intersections of the cone with vertical planes (parallel to the time axis x^0). The black dots represent the origin $(1, 0, -1)$ of the future lightcone and the origin $(-1, 0, 1)$ of the past lightcone when constructed as the homogeneous spaces \mathcal{L}_\pm . Notice that the boundary of our coordinate patch is the light ray $x^0 = x^2$ (black dashed line).

3 Poisson–Lie groups and Poisson homogeneous spaces

As it is well known, a PL structure on a Lie group is the semiclassical limit of a quantum group: a Hopf algebra [7, 9, 33, 34] that represents a noncommutative deformation of the algebra of functions on a Lie group. Conversely, a quantum group is a Hopf algebra quantization of a given PL group.

The aim of this paper is to construct noncommutative analogues of the lightcones \mathcal{L}_\pm , which must be covariant under the corresponding quantum $\text{SO}(2,1)$ conformal groups. To achieve this, we adopt the ‘semiclassical’ approach used for constructing the noncommutative Minkowski and (A)dS spacetimes in [35, 36], as well as the noncommutative spaces of worldlines given in [20–23] as quantizations of PH spaces that are covariant under the action of the corresponding PL groups.

Therefore, we have to begin by selecting a (coboundary) PL structure Π on the Lie group $\text{SO}(2,1)$ with Lie algebra $\mathfrak{so}(2,1)$ given by a classical r -matrix defined on $\mathfrak{so}(2,1) \wedge \mathfrak{so}(2,1)$. The Poisson bivector defining Π is explicitly provided by the Sklyanin bracket

$$\{f, g\} = r^{IJ} (X_I^L f X_J^L g - X_I^R f X_J^R g), \quad f, g \in \mathcal{C}^\infty(\text{SO}(2,1)), \quad (19)$$

where X_I^L and X_I^R are the left- and right-invariant vector fields on $\text{SO}(2,1)$ given by (10).

Moreover, the tangent counterpart of a coboundary PL group $(\text{SO}(2,1), \Pi)$ is a coboundary Lie bialgebra $(\mathfrak{so}(2,1), \delta)$, where the Lie bialgebra cocommutator map $\delta : \mathfrak{so}(2,1) \rightarrow \mathfrak{so}(2,1) \wedge \mathfrak{so}(2,1)$ is obtained from the r -matrix through the expression

$$\delta(\mathbb{X}) = [\mathbb{X} \otimes 1 + 1 \otimes \mathbb{X}, r], \quad \forall \mathbb{X} \in \mathfrak{so}(2,1). \quad (20)$$

The theory of PH spaces (see [27–30] and references therein) states that PL groups $(\text{SO}(2,1), \Pi)$ will lead to PH structures (\mathcal{L}_\pm, π) on the lightcones provided that the associated Lie bialgebra $(\mathfrak{so}(2,1), \delta)$ satisfies the so-called coisotropy condition for the cocommutator δ with respect to the

isotropy subalgebra of the homogeneous space, in this case generated by \mathbb{K}

$$\delta(\mathbb{K}) \subseteq \mathbb{K} \wedge \mathfrak{so}(2, 1). \quad (21)$$

Then the Poisson structure π on the homogeneous space is obtained by canonical projection from the corresponding Π on $\text{SO}(2,1)$ given by Eq. (19). In that case, we have a PH lightcone, in which the additional Poisson bracket π is compatible with the action of the conformal group. Moreover, when the stronger condition

$$\delta(\mathbb{K}) \subseteq \mathbb{K} \wedge \mathbb{K}, \quad (22)$$

is met, \mathbb{K} becomes a sub-Lie bialgebra, leading to a PH space of Poisson-subgroup type. In our case, since \mathbb{K} is one-dimensional, this situation is only realized when $\delta(\mathbb{K}) = 0$.

It is also well known that the Lie algebra $\mathfrak{sl}(2, \mathbb{R}) \simeq \mathfrak{so}(2, 1)$ admits three families of inequivalent Lie bialgebra structures, which correspond to three distinct families of PL structures (see [24, 25] and also [26], where the explicit construction of the corresponding quantum groups and algebras is provided). Expressed in terms of the $\text{SL}(2, \mathbb{R})$ generators \mathbb{I}_+ , \mathbb{I}_- , \mathbb{I}_3 with commutation relations

$$[\mathbb{I}_3, \mathbb{I}_\pm] = \pm 2 \mathbb{I}_\pm, \quad [\mathbb{I}_+, \mathbb{I}_-] = \mathbb{I}_3, \quad (23)$$

the three classical r -matrices generating these PL structures are

$$r_I = \alpha \mathbb{I}_+ \wedge \mathbb{I}_-, \quad r_{II} = \beta \mathbb{I}_3 \wedge (\mathbb{I}_+ + \mathbb{I}_-), \quad r_{III} = \mathbb{I}_3 \wedge \mathbb{I}_+, \quad (24)$$

where α and β are real numbers that parametrize each family of Lie bialgebras (different values of α and β correspond to inequivalent r -matrices, not related to each other by Lie algebra automorphisms). The r -matrix r_{III} does not have a corresponding parameter, because its rescaling by a real number is induced by an automorphism.

In this paper, we are interested in expressing the $\text{SO}(2,1)$ group and its r -matrices in terms of the conformal generators \mathbb{T} , \mathbb{D} , and \mathbb{K} . The isomorphism between $\mathfrak{sl}(2, \mathbb{R})$ and $\mathfrak{so}(2, 1)$

$$\mathbb{I}_+ = -\mathbb{M}_{01} - \mathbb{M}_{12} = \mathbb{K}, \quad \mathbb{I}_- = -\mathbb{M}_{01} + \mathbb{M}_{12} = -\mathbb{T}, \quad \mathbb{I}_3 = -2\mathbb{M}_{02} = -2\mathbb{D}, \quad (25)$$

leads to

$$r_I = \alpha \mathbb{T} \wedge \mathbb{K}, \quad r_{II} = 2\beta \mathbb{D} \wedge (\mathbb{T} - \mathbb{K}), \quad r_{III} = -2\mathbb{D} \wedge \mathbb{K}. \quad (26)$$

The three r -matrices above are representatives of three automorphism classes at the PL group level, but it will be necessary to analyze in detail the role of such group automorphisms at the level of the coisotropic PH spaces. To the best of our knowledge, this aspect has not been previously treated in the literature, and it will turn out to be essential, since in doing so we will show that more than three inequivalent PH lightcones exist and have to be considered.

First, we recall that the automorphisms of the group $\text{SO}(2,1)$ are all inner, meaning that they can all be obtained by conjugation by an element of the group

$$g(\tau, \lambda, \sigma) \rightarrow h g(\tau, \lambda, \sigma) h^{-1}, \quad h \in \text{SO}(2, 1). \quad (27)$$

To discuss the effect of automorphisms on the PL groups and PH spaces, we need to study their action on the r -matrices, which in turn requires the action on the Lie algebra generators. Given a basis of linearly independent generators $\mathbb{X}_i \in \mathfrak{so}(2, 1)$, $i = 1, 2, 3$, the action of h from (27) on each element \mathbb{X}_i is given by

$$\mathbb{X}_i \rightarrow h(\phi_1, \phi_2, \psi) \mathbb{X}_i h^{-1}(\phi_1, \phi_2, \psi), \quad (28)$$

where ϕ_1 , ϕ_2 , ψ are three parameters on the group. The three PL groups coming from (26) are invariant under automorphisms of $\text{SO}(2,1)$, so any automorphism of the form (27) induces an ‘unphysical’ recoordination of the group manifold that in turn induces a map between equivalent PL structures.

Nevertheless, the situation changes when PH spaces associated to certain automorphism-equivalent PL groups are considered. As we showed in the previous section, these homogeneous spaces are obtained, geometrically, by applying the group transformations to a point of the ambient Minkowski

space in which the representation (7) of $SO(2,1)$ lives. Only the coordinates τ and λ act nontrivially on the lightlike point we chose as origin of the homogeneous space, and the σ coordinate does not contribute. This implies that the automorphisms on the $SO(2,1)$ group that just mix the τ and λ coordinates project onto coordinate changes on the homogeneous space and can be said to preserve said homogeneous space (or to preserve the quotient structure). However, $SO(2,1)$ automorphisms that mix the σ coordinate with τ and λ cannot be expressed as maps from the homogeneous space onto itself and, therefore, they do not commute with the quotient $SO(2,1)/K$. The former type of automorphism is then an unphysical change of coordinates of the PH space, while the latter changes the Poisson structure on the homogeneous space, despite not changing the PL group.

4 $SO(2,1)$ r -matrices and quotient-preserving automorphisms

To make the above considerations explicit, we describe the most generic automorphism map (27) on $SO(2,1)$ as follows: when belonging to the connected component of the identity, $h \in SO_+(2,1)$, it will be written for convenience as

$$h = e^{\phi_2 \mathbb{K}} e^{\phi_1 \mathbb{D}} e^{\psi \mathbb{T}}, \quad (29)$$

which corresponds, through Eq. (27), to the following transformation rule on the group coordinates:

$$\begin{aligned} \tau &\rightarrow \left(\frac{e^{-\phi_1(1+\sigma\psi)}}{(\tau+\psi)(1+\sigma\psi)-e^\lambda\psi} - \phi_2 \right)^{-1}, \\ \lambda &\rightarrow \lambda - 2 \log \left(1 + \sigma\psi + e^{\phi_1}\phi_2 \left(e^\lambda\psi - (\tau+\psi)(1+\sigma\psi) \right) \right), \\ \sigma &\rightarrow \frac{e^\lambda\phi_2(\psi e^{\phi_1}\phi_2 - 1) - (e^{\phi_1}\phi_2(\sigma\psi + 1) - \sigma)(\phi_2(\tau + \psi) - e^{-\phi_1})}{(\sigma\psi + 1)(e^{\phi_1}\phi_2(\tau + \psi) - 1) - \psi\phi_2 e^{\lambda+\phi_1}}. \end{aligned} \quad (30)$$

The parametrization chosen for h does not cover all of $SO_+(2,1)$, because (just like the parametrization (7)) it excludes the rotation (11). The effect on the coordinates of conjugating $g(\tau, \lambda, \sigma)$ with the matrix (11) is found to be

$$\tau \rightarrow \frac{\sigma}{e^\lambda - \sigma\tau}, \quad \lambda \rightarrow \lambda - 2 \log \left(e^\lambda - \sigma\tau \right), \quad \sigma \rightarrow \frac{\tau}{e^\lambda - \sigma\tau}. \quad (31)$$

Finally, the automorphisms generated by the non-orthochronous component $SO_-(2,1)$ can be obtained from (27) by further conjugating by the PT matrix (12). The conjugation $g(\tau, \lambda, \sigma) \rightarrow (PT)g(\tau, \lambda, \sigma)(PT)^{-1}$ acts on the group coordinates as follows:

$$\tau \rightarrow -\tau, \quad \lambda \rightarrow \lambda, \quad \sigma \rightarrow -\sigma. \quad (32)$$

A quick inspection of Eq. (30) reveals that the automorphisms that do not mix the σ coordinate with τ and λ are those such that $\psi = 0$, and only those. The conjugation by the PT matrix (32) preserves the quotient as well. The only transformations that do not preserve the quotient, and therefore are physical and can generate inequivalent PH spaces, are those of the form (30) that involve a nonzero ψ parameter, or those that involve the rotation (31) of π of the 1 and 2 axes. Therefore, we will need only to focus on these ‘physical’ automorphisms. We may as well set the parameters ϕ_1 and ϕ_2 to zero, because, as can be verified, the effect of a transformation of the form (30) on the PH space is the same as that of a transformation with the same ψ and $\phi_1 = \phi_2 = 0$, namely

$$\tau \rightarrow \tau + \psi \left(1 - \frac{e^\lambda}{1+\sigma\psi} \right), \quad \lambda \rightarrow \lambda - 2 \log(1 + \sigma\psi), \quad \sigma \rightarrow \frac{\sigma}{1 + \sigma\psi}, \quad (33)$$

modulo changes of coordinates on the homogeneous space. In conclusion, from now on we will only focus on automorphisms generated by (11) and (29) with $\phi_1 = \phi_2 = 0$.

The action (28) of these physical automorphisms on the generators in the conformal basis, $\mathbb{X}_1 = \mathbb{T}$, $\mathbb{X}_2 = \mathbb{D}$, $\mathbb{X}_3 = \mathbb{K}$, induce the following transformation rules:

$$\mathbb{T} \rightarrow \mathbb{K}, \quad \mathbb{D} \rightarrow -\mathbb{D}, \quad \mathbb{K} \rightarrow \mathbb{T}, \quad (34)$$

and

$$\mathbb{T} \rightarrow \mathbb{T}, \quad \mathbb{D} \rightarrow \mathbb{D} - \psi\mathbb{T}, \quad \mathbb{K} \rightarrow \mathbb{K} + \psi^2\mathbb{T} - 2\psi\mathbb{D}, \quad (35)$$

respectively. In particular, the latter introduce an additional parameter in the r -matrices (26):

$$r_I = \alpha(\mathbb{T} \wedge \mathbb{K} - 2\psi\mathbb{T} \wedge \mathbb{D}) \equiv \alpha_1\mathbb{T} \wedge (\mathbb{K} + \alpha_2\mathbb{D}) = r_I(\alpha_1, \alpha_2), \quad (36a)$$

$$\begin{aligned} r_{II} &= -2\beta(\mathbb{D} \wedge \mathbb{K} - \psi\mathbb{T} \wedge \mathbb{K} + (1 + \psi^2)\mathbb{T} \wedge \mathbb{D}) \\ &\equiv \beta_1((\mathbb{D} \wedge \beta_2\mathbb{T}) \wedge \mathbb{K} + (1 + \beta_2^2)\mathbb{T} \wedge \mathbb{D}) = r_{II}(\beta_1, \beta_2), \end{aligned} \quad (36b)$$

$$\begin{aligned} r_{III} &= -2(\mathbb{D} \wedge \mathbb{K} - \psi\mathbb{T} \wedge \mathbb{K} + \psi^2\mathbb{T} \wedge \mathbb{D}) \\ &\equiv -2((\mathbb{D} - \gamma\mathbb{T}) \wedge \mathbb{K} + \gamma^2\mathbb{T} \wedge \mathbb{D}) = r_{III}(\gamma), \end{aligned} \quad (36c)$$

where we relabeled our parameters $\alpha_1, \alpha_2, \beta_1, \beta_2$ and γ for ease of reading.

Although the transformation we used to introduce a dependence of the r -matrices on the parameter ψ does not correspond to a recoordination on the lightcones, some pure recoordination automorphisms of the form (29) with $\psi = 0$ could relate some of the r -matrices in (36) with each other. This happens for the following subsets of r -matrices: in the family r_I (36a), we can connect two r -matrices of the family with the same α_1 but different non-zero $\alpha_2 \neq \alpha'_2$ with recoordination parameters:

$$\phi_1 = \log \frac{\alpha'_2}{\alpha_2}, \quad \phi_2 = 0, \quad \Rightarrow \quad r_I(\alpha_1, \alpha_2) \xrightarrow[\phi_1, \phi_2]{} r_I(\alpha_1, \alpha'_2). \quad (37)$$

This means that all the r -matrices of this family with a non-zero α_2 are actually connected by a recoordination, so we can simply choose a representative for them. The r -matrices with $\alpha_2 = 0$ are disconnected from the ones with $\alpha_2 \neq 0$. Therefore, we split this family into two uniparametric families

$$r_I(\alpha_1, \alpha_2 = 0) = \alpha_1\mathbb{T} \wedge \mathbb{K}, \quad r_I(\alpha_1, \alpha_2 = 1) = \alpha_1\mathbb{T} \wedge (\mathbb{D} + \mathbb{K}). \quad (38)$$

The same happens for the second family r_{II} (36b):

$$\phi_1 = \log \frac{1+\beta'_2}{1+\beta_2}, \quad \phi_2 = \frac{-\beta_2+\beta'_2}{1+\beta_2}, \quad \Rightarrow \quad r_{II}(\beta_1, \beta_2) \xrightarrow[\phi_1, \phi_2]{} r_{II}(\beta_1, \beta'_2), \quad (39)$$

with the difference that this transformation is allowed for any value of the parameters, so we have a single uniparametric family

$$r_{II}(\beta_1, \beta_2 = 0) = \beta_1\mathbb{D} \wedge (\mathbb{K} - \mathbb{T}). \quad (40)$$

Finally, in the case of the third family r_{III} (36c), for non-zero values of the parameter $\gamma \neq \gamma'$, we can connect two r -matrices of the family with the following recoordination parameters:

$$\phi_1 = 2 \log \frac{\gamma'}{\gamma}, \quad \phi_2 = \frac{\gamma - \gamma'}{\gamma^2}, \quad \Rightarrow \quad r_{III}(\gamma) \xrightarrow[\phi_1, \phi_2]{} r_{III}(\gamma'). \quad (41)$$

This shows that all non-zero values of the parameter γ are connected by a recoordination. Therefore, we again have a splitting into two subcases:

$$r_{III}(\gamma = 0) = -2\mathbb{D} \wedge \mathbb{K}, \quad r_{III}(\gamma = 1) = 2(\mathbb{T} - \mathbb{D}) \wedge \mathbb{K} - 2\mathbb{T} \wedge \mathbb{D}. \quad (42)$$

Thus, the five classes of inequivalent structures obtained so far are the following:

$$r_I^A(\alpha_A) = \alpha_A\mathbb{T} \wedge \mathbb{K}, \quad (43a)$$

$$r_I^B(\alpha_B) = \alpha_B\mathbb{T} \wedge (\mathbb{D} + \mathbb{K}), \quad (43b)$$

$$r_{II}(\beta) = \beta\mathbb{D} \wedge (\mathbb{K} - \mathbb{T}), \quad (43c)$$

$$r_{III}^A = -2\mathbb{D} \wedge \mathbb{K}, \quad (43d)$$

$$r_{III}^B = 2(\mathbb{T} - \mathbb{D}) \wedge \mathbb{K} - 2\mathbb{T} \wedge \mathbb{D}, \quad (43e)$$

where r_I^B and r_{III}^B arise as new ones. We have three uniparametric families of r -matrices depending on the real parameters $\alpha_A, \alpha_B, \beta \in \mathbb{R}$, and two additional isolated r -matrices.

Finally, the effect of the π rotation automorphism (34) on the families (43) leads to

$$\begin{aligned}
r_I^A(\alpha_A) &= -\alpha_A \mathbb{T} \wedge \mathbb{K} = r_I^A(-\alpha_A), \\
r_I^B(\alpha_B) &= \alpha_B \mathbb{K} \wedge (\mathbb{T} - \mathbb{D}), \\
r_{II}(\beta) &= \beta \mathbb{D} \wedge (\mathbb{K} - \mathbb{T}) = r_{II}(\beta), \\
r_{III}^A &= 2\mathbb{D} \wedge \mathbb{T}, \\
r_{III}^B &= -2(\mathbb{K} + \mathbb{D}) \wedge \mathbb{T} + 2\mathbb{K} \wedge \mathbb{D}.
\end{aligned} \tag{44}$$

The families r_I^A and r_{II} did not change (the automorphism just sends the parameters α_A and β to $-\alpha_A$ and β), while the other three families are modified. However, upon a closer inspection, these changes correspond to the following recoordinatizations of the families we already have:

$$\begin{aligned}
\forall \phi_1 \in \mathbb{R}, \quad \phi_2 = -\frac{1}{2}, & \quad \Rightarrow \quad r_I^A(-\alpha_B) \xrightarrow{\phi_1, \phi_2} r_I^B(\alpha_B), \\
\phi_1 = 0, \quad \phi_2 = 1, & \quad \Rightarrow \quad r_{III}^B \xrightarrow{\phi_1, \phi_2} r_{III}^A, \\
\phi_1 = 0, \quad \phi_2 = 2, & \quad \Rightarrow \quad r_{III}^B \xrightarrow{\phi_1, \phi_2} r_{III}^B.
\end{aligned} \tag{45}$$

Therefore, Eq. (43) provides the full set of families of r -matrices to be considered in order to construct noncommutative lightcones with quantum $\text{SO}(2,1)$ invariance.

5 Poisson homogeneous structures on the lightcone

The five families of r -matrices (43) correspond to the following five cocommutators that define, through (20), the associated Lie bialgebra structures on the $\mathfrak{so}(2,1)$ algebra:

$$\begin{aligned}
r_I^A: \quad & \delta(\mathbb{T}) = -2\alpha_A \mathbb{T} \wedge \mathbb{D}, \quad \delta(\mathbb{D}) = 0, \quad \delta(\mathbb{K}) = 2\alpha_A \mathbb{D} \wedge \mathbb{K}, \\
r_I^B: \quad & \delta(\mathbb{T}) = -2\alpha_B \mathbb{T} \wedge \mathbb{D}, \quad \delta(\mathbb{D}) = \alpha_B \mathbb{T} \wedge \mathbb{D}, \quad \delta(\mathbb{K}) = \alpha_B (2\mathbb{D} + \mathbb{T}) \wedge \mathbb{K}, \\
r_{II}: \quad & \delta(\mathbb{T}) = -2\beta \mathbb{T} \wedge \mathbb{K}, \quad \delta(\mathbb{D}) = -\beta \mathbb{D} \wedge (\mathbb{K} + \mathbb{T}), \quad \delta(\mathbb{K}) = \beta \mathbb{T} \wedge \mathbb{K}, \\
r_{III}^A: \quad & \delta(\mathbb{T}) = 2\mathbb{T} \wedge \mathbb{K}, \quad \delta(\mathbb{D}) = 2\mathbb{D} \wedge \mathbb{K}, \quad \delta(\mathbb{K}) = 0, \\
r_{III}^B: \quad & \delta(\mathbb{T}) = 2\mathbb{T} \wedge (\mathbb{K} - 2\mathbb{D}), \quad \delta(\mathbb{D}) = 2\mathbb{D} \wedge (\mathbb{K} + \mathbb{T}), \quad \delta(\mathbb{K}) = 2(2\mathbb{D} - \mathbb{T}) \wedge \mathbb{K}.
\end{aligned} \tag{46}$$

The coisotropy condition (21) holds for all five cases above, while only the matrix r_{III}^A (43d) satisfies the Poisson subgroup condition (22), which makes \mathbb{K} into a sub-Lie bialgebra.

Now, by using the Sklyanin bracket (19), we get five PL structures Π on $\text{SO}(2,1)$

$$\begin{aligned}
r_I^A: \quad & \{\tau, \lambda\} = -2\alpha_A \tau, \quad \{\sigma, \tau\} = 0, \quad \{\sigma, \lambda\} = -2\alpha_A \sigma, \\
r_I^B: \quad & \{\tau, \lambda\} = \alpha_B (e^\lambda - 2\tau - 1), \quad \{\sigma, \tau\} = -\alpha_B e^\lambda \sigma, \quad \{\sigma, \lambda\} = -\alpha_B \sigma (2 + \sigma), \\
r_{II}: \quad & \{\tau, \lambda\} = \beta (e^\lambda - \tau^2 - 1), \quad \{\sigma, \tau\} = -\beta e^\lambda (\sigma - \tau), \quad \{\sigma, \lambda\} = \beta (e^\lambda - \sigma^2 - 1), \\
r_{III}^A: \quad & \{\tau, \lambda\} = 2\tau^2, \quad \{\sigma, \tau\} = -2e^\lambda \tau, \quad \{\sigma, \lambda\} = 2(1 - e^\lambda), \\
r_{III}^B: \quad & \{\tau, \lambda\} = -2(e^\lambda - (\tau - 1)^2), \quad \{\sigma, \tau\} = 2e^\lambda (\sigma - \tau), \quad \{\sigma, \lambda\} = -2(e^\lambda - (\sigma - 1)^2).
\end{aligned} \tag{47}$$

Notice that, at this PL group level, the inequivalent structures are just three (or rather two uniparametric families plus a single, isolated case): $r_I^{A,B}$, r_{II} , and $r_{III}^{A,B}$, since the ‘ A ’ and ‘ B ’ subfamilies of families I and III are related by automorphisms. Casimir functions for all these Poisson structures (47) have been determined and are as follows:

$$\begin{aligned}
C_I^A &= \frac{\tau}{\sigma}, \quad C_I^B = \frac{1 - e^\lambda + \sigma\tau + 2\tau}{2\sigma}, \quad C_{II} = \frac{1 - e^\lambda + \sigma\tau}{\sigma - \tau}, \\
C_{III}^A &= \frac{1 - e^\lambda + \sigma\tau}{\tau}, \quad C_{III}^B = \frac{1 - e^\lambda + \sigma\tau - \sigma - \tau}{\sigma - \tau},
\end{aligned} \tag{48}$$

The level sets of these Casimir functions identify the symplectic leaves of the Poisson structure.

Now, the Poisson brackets π of the PH spaces of each of the five families (43) can be simply obtained by projecting the PL brackets (47) to the first two coordinates, τ and λ , and replacing τ with y and λ with z

$$r_I^A : \{y, z\} = -2\alpha_A y, \quad (49a)$$

$$r_I^B : \{y, z\} = \alpha_B (e^z - 2y - 1), \quad (49b)$$

$$r_{II} : \{y, z\} = \beta (e^z - y^2 - 1), \quad (49c)$$

$$r_{III}^A : \{y, z\} = 2y^2, \quad (49d)$$

$$r_{III}^B : \{y, z\} = -2 (e^z - (y - 1)^2). \quad (49e)$$

$$(49f)$$

We have found a set of Darboux coordinates $\{q, p\} = 1$ for each case

$$\begin{aligned} r_I^A : z = p, \quad y = \pm e^{-2\alpha_A(q-q_0)}, \\ r_I^B : z = p, \quad y = \frac{1}{2}(e^p - 1) + e^{-2\alpha_B(q-q_0)}, \\ r_{II} : z = p, \quad y = -\sqrt{1 - e^p} \tan(\sqrt{1 - e^p} \beta (q - q_0)), \\ r_{III}^A : z = p, \quad y = -(2(q - q_0))^{-1}, \\ r_{III}^B : z = p, \quad y = 1 - e^{p/2} \tanh(2e^{p/2}(q - q_0)), \end{aligned} \quad (50)$$

where q_0 is an integration constant and the \pm symbol refers to two possible choices of Darboux coordinates.

The coordinates y and z appearing in (49) are equally valid on the future and the past lightcone, \mathcal{L}_+ and \mathcal{L}_- , depending on which of the two embeddings (14) is used. The Poisson brackets (49) induce the following Poisson brackets on the ambient Minkowski space:

$$\{x_\pm^I, x_\pm^J\} = (u_\pm x_\pm^L + w_\pm) \varepsilon^{IJ}{}_{\mathcal{K}} x_\pm^{\mathcal{K}}, \quad (51)$$

with the following choices for u_\pm and w_\pm for the five families:

$$\begin{aligned} r_I^A : u_\pm = -2\alpha_A \delta_\pm^1, \quad w_\pm = 0, \\ r_I^B : u_\pm = \alpha_B (\delta_\pm^0 + 2\delta_\pm^1 - \delta_\pm^2), \quad w_\pm = \mp 2l, \\ r_{II} : u_\pm = 2\beta \delta_\pm^0, \quad w_\pm = \mp l, \\ r_{III}^A : u_\pm = -2(\delta_\pm^0 + \delta_\pm^2), \quad w_\pm = 0, \\ r_{III}^B : u_\pm = 4(\delta_\pm^1 - \delta_\pm^0), \quad w_\pm = \pm l, \end{aligned} \quad (52)$$

where l is the positive constant chosen for the stabilizing vectors (3). These brackets reduce to (49) on the lightcones \mathcal{L}_\pm , when the conditions (15) are satisfied.

Notice how all the right-hand sides of Eqs. (51) are proportional to $\varepsilon^{IJ}{}_{\mathcal{K}} x_\pm^{\mathcal{K}}$, where $\varepsilon^{IJ}{}_{\mathcal{K}}$ is the 3D Levi-Civita tensor with one index lowered with the 2+1-dimensional Minkowski metric. Under inner automorphisms, the embedding coordinates transform like contravariant vectors under classical Lorentz transformations. Thus, the effect of an automorphism on the commutation relations (51) is to Lorentz-transform the constants u_\pm as if they formed a covariant vector, while the constants w_\pm are left invariant. Therefore, two models whose u_\pm are related by a Lorentz transformation are equivalent, in the sense that they are related by a coordinate change, as long as their w_\pm values coincide.

The constant u_\pm is spacelike in both cases of family *I*, timelike in family *II*, and lightlike in both cases of family *III*. Consequently, apart from a rescaling, one can connect cases r_I^A and r_I^B , as well as r_{III}^A and r_{III}^B , using an automorphism. However, the ‘A’ cases have $w_\pm = 0$, while the ‘B’ cases do not. The conclusion is that the ‘A’ cases are limiting cases of the ‘B’ ones, where $l \rightarrow 0$. Given that l is the length scale of the ambient Minkowski space, sending l to zero corresponds to observing




Figure 2: The regions of the lightcones where the five Poisson brackets vanish are given by the black curves defined by the intersection of the lightcones with the planes shown in this figure.

the model from infinitely far away, where the features relevant in a neighbourhood of the origin of size $\sim l$ cannot be appreciated.

This rewriting makes it easier to identify the submanifolds of the lightcones where the Poisson brackets vanish in each model, which will be relevant in the quantum case concerning localization properties. These are all one-dimensional submanifolds obtained by intersecting the lightcones (15) with the planes $u_{\pm} x_{\pm}^1 + w_{\pm} = 0$, that is

$$\begin{aligned}
 r_I^A : \quad & x_{\pm}^1 = 0, && \text{timelike plane through the origin,} \\
 r_I^B : \quad & x_{\pm}^0 + 2x_{\pm}^1 - x_{\pm}^2 \mp 2l = 0, && \text{timelike planes separated by } 2l, \\
 r_{II} : \quad & x_{\pm}^0 \mp l = 0, && \text{spacelike planes separated by } 2l, \\
 r_{III}^A : \quad & x_{\pm}^0 + x_{\pm}^2 = 0, && \text{lightlike plane through the origin,} \\
 r_{III}^B : \quad & x_{\pm}^1 - x_{\pm}^0 \pm l = 0, && \text{lightlike planes separated by } 2l,
 \end{aligned} \tag{53}$$

where the \pm symbol refers to the future and past lightcones. We plotted these intersecting planes, for each of the five models, in Fig. 2. Notice how the parameter l appearing in three of the expressions above simply sets the scale for the whole picture without changing anything substantial: for r_I^B , r_{II} , and r_{III}^B , it determines the vertical distance between the origin and the two intersecting planes. Independently of the value of l , the intersection remain consistent: two light rays in the case of r_I^A , two hyperbolas for r_I^B , two circles for r_{II} , one light ray for r_{III}^A , and two parabolas for r_{III}^B .

Cases r_I^A and r_{III}^A can be seen as degenerate limits of r_I^B and r_{III}^B , respectively, as $l \rightarrow 0$. This further clarifies the sense in which the ‘A’ and ‘B’ subcases are not mere recoordinatizations of each other. Instead, the ‘A’ cases are just the ‘B’ cases viewed from infinitely far away, making the failure of the two planes to meet at the origin not appreciable (and, in the $r_I^B \rightarrow r_I^A$ case, further subject to a Lorentz transform that makes the intersecting planes vertical). Similarly, the Poisson brackets between embedding coordinates (51) show that the two ‘A’ models can be obtained from the ‘B’ models via an automorphism of the x_{\pm}^1 followed by the $l \rightarrow 0$ limit. In other words, the ‘A’ models are essentially the ‘B’ models seen from afar (and from a Lorentz-transformed perspective).

6 Representations of noncommutative lightcones and their localization properties

Introducing a fuzziness parameter $\ell > 0$, which plays a similar role as \hbar in ordinary Quantum Mechanics (see [32] for a more detailed discussion), and promoting the coordinate functions y, z to

operators \hat{y} , \hat{z} , we can quantize Eq. (49):

$$r_I^A : [\hat{y}, \hat{z}] = -2i\ell\alpha_A\hat{y}, \quad (54a)$$

$$r_I^B : [\hat{y}, \hat{z}] = i\ell\alpha_B(e^{\hat{z}} - 2\hat{y} - 1), \quad (54b)$$

$$r_{II} : [\hat{y}, \hat{z}] = i\ell\beta(e^{\hat{z}} - \hat{y}^2 - 1), \quad (54c)$$

$$r_{III}^A : [\hat{y}, \hat{z}] = 2i\ell\hat{y}^2, \quad (54d)$$

$$r_{III}^B : [\hat{y}, \hat{z}] = -2i\ell(e^{\hat{z}} - (\hat{y} - 1)^2), \quad (54e)$$

with no ordering issues arising. This ℓ -quantization should be understood in the sense that its semiclassical limit

$$\{\cdot, \cdot\} = \lim_{\ell \rightarrow 0} \frac{[\cdot, \cdot]}{i\ell}, \quad (55)$$

recovers the five PH lightcones (49). The commutation relations (54) are supplemented by the Hermiticity conditions on the ‘quantum coordinates’ \hat{y} and \hat{z}

$$\hat{y}^\dagger = \hat{y}, \quad \hat{z}^\dagger = \hat{z}, \quad (56)$$

which are compatible with the commutators, as can be easily verified. Notice how the fuzziness parameter ℓ is dimensionless; it determines the magnitude of fuzziness implied by the commutation relations of the dimensionless coordinates \hat{y} and \hat{z} .

The irreducible regular representations of an algebra isomorphic to case r_I^A in Eq. (54a) have been described in [37] and further analyzed in [14, 15]. The representation is based on the Darboux coordinates (50), and the operators take the form

$$\hat{z} = -i\ell\partial_q, \quad \hat{y} = 0, \pm e^{-2\alpha_A(q-q_0)}, \quad (57)$$

where the three cases, in which \hat{y} is respectively positive, negative, or zero, are the three Hermitian irreducible representations, and the integration constant q_0 just shifts the origin of the representation horizontally. The Hilbert space is that of square-integrable functions $L^2(\mathbb{R})$ of $q \in \mathbb{R}$.

A similar analysis applies also to the algebras in Eq. (54b) and Eq. (54d):

$$\begin{aligned} r_I^B : \hat{z} &= -i\ell\partial_q, \quad \hat{y} = \frac{1}{2} \left(e^{-i\ell\partial_q} - 1 \right) + e^{-2\alpha_B(q-q_0)}, \\ r_{III}^A : \hat{z} &= -i\ell\partial_q, \quad \hat{y} = -(2(q-q_0))^{-1}. \end{aligned} \quad (58)$$

In the case of r_{III}^A , the singularity at $q = 0$ forces us to change the Hilbert space for a space of functions that vanish at the origin. For model r_{III}^B in Eq. (54e), we modify the representation used for algebra r_{III}^A :

$$\hat{z} = -\log\left(4(q-q_0)^2 + e^{p_0+i\ell\partial_q}\right), \quad \hat{y} = 1 - (2(q-q_0))^{-1}, \quad (59)$$

with a Hilbert space consisting again of square-integrable functions that vanish at the origin.

The model r_{II} in Eq. (54c) is special because one of the variables of the canonical pair we find has a compact domain, indicating that the canonically conjugate operator should have a discrete spectrum. This compactness is related to the fact, observed above, that in this model the region where the Poisson brackets vanish is a compact circle. Rather than using the Darboux variables found in (50), which depend on products of q and p and make quantization difficult, it is convenient to consider the cylindrical coordinates θ , t in the ambient space (already used in (18) for the geometrical interpretation of y):

$$y = \frac{\cos\theta}{1 - \sin\theta}, \quad z = -\log\left(\frac{\pm t(1 - \sin\theta)}{2}\right), \quad (60)$$

where the \pm symbol refers to the future and past lightcones. In these coordinates, the Poisson brackets (49c) of case r_{II} turn into $\{\theta, t\} = 2\beta(t \mp 1)$, and a set of Darboux coordinates can be found as $\theta = q$, $t = \pm(e^{2\beta(p-p_0)} + 1)$, which imply:

$$y = \frac{\cos q}{1 - \sin q}, \quad z = -\log(1 - \sin q) - \log\left(\frac{e^{2\beta(p-p_0)} + 1}{2}\right). \quad (61)$$

We then found a choice of Darboux coordinates that is suitable for representing case r_{II} of Eq. (54c), because there is no ordering ambiguity in the map that connects them to \hat{y} , \hat{z} :

$$\hat{y} = \frac{\cos \theta}{1 - \sin \theta}, \quad \hat{z} = -\log(1 - \sin \theta) - \log\left(\frac{e^{2\beta(\hat{L}_\theta - p_0)} + 1}{2}\right). \quad (62)$$

The Hilbert space will be that of square-integrable periodic functions of $\theta \in [0, 2\pi)$, and the operator \hat{L}_θ in the expression (62) for \hat{z} is the conjugate momentum of the angle θ , which has a discrete spectrum and can be obtained as a self-adjoint extension of the derivative operator $-i\ell\partial_\theta$.

We now turn to the problem of localizability, *i.e.*, how well points can be resolved in noncommutative spaces such as (54). This problem can be discussed in terms of the states on the algebras (54). These states are positive linear functionals of norm one on the algebra, and maximally-localized states can be described as limits of well-normalized states. The maximally-localized states determine how well a region can be resolved. For all models in Eq. (54) except r_{II} (54c), we can discuss the perfectly-localized states in the Darboux variables, which close the Heisenberg algebra $[\hat{q}, \hat{p}] = i\ell$ and have a real spectrum. The coordinates \hat{q} or the momenta \hat{p} can be arbitrarily well localized around a value if we consider normalized states that are increasingly more peaked around a particular value. In the limit of perfect localization, these turn into delta functions, which are not normalizable and therefore called improper states, but admit the physical interpretation of an idealization, as one can push the localization limit arbitrarily far while always maintaining a normalizable state. Thus, \hat{p} or \hat{q} can be separately localized arbitrarily well, but there are limits on their simultaneous localization, in accordance with the uncertainty relation

$$\delta\hat{q} \delta\hat{p} \geq \frac{\ell}{2}. \quad (63)$$

The optimally-localized states in both \hat{p} and \hat{q} are the coherent states, whose wavefunctions are Gaussians given by

$$\psi(q) = \frac{1}{\sqrt{\sigma\sqrt{2\pi}}} e^{-\left(\frac{q-q_0}{2\sigma}\right)^2 + i\frac{p_0 q_0}{\ell}} \Rightarrow \langle\hat{q}\rangle = q_0, \quad \langle\hat{p}\rangle = p_0, \quad \delta\hat{q} = \sigma, \quad \delta\hat{p} = \frac{\ell}{2\sigma}. \quad (64)$$

The commutation relations of an algebra impose that, on any state, the uncertainty product is larger than the expectation value of the right-hand side. In the case of the Heisenberg algebra, the right-hand side is proportional to the identity operator so its expectation value is a constant, and one can never perfectly localize a state in phase space. When the commutators are non-constant, as in (54), the situation might be different. The uncertainty relations implied by our commutators are

$$\begin{aligned} r_I^A: & \quad \delta\hat{y} \delta\hat{z} \geq -\ell |\alpha_A \langle\hat{y}\rangle|, \\ r_I^B: & \quad \delta\hat{y} \delta\hat{z} \geq \frac{\ell}{2} \left| \alpha_B \left\langle e^{\hat{z}} - 2\hat{y} - 1 \right\rangle \right|, \\ r_{II}: & \quad \delta\hat{y} \delta\hat{z} \geq \frac{\ell}{2} \left| \beta \left\langle e^{\hat{z}} - \hat{y}^2 - 1 \right\rangle \right|, \\ r_{III}^A: & \quad \delta\hat{y} \delta\hat{z} \geq \frac{\ell}{2} |\langle\hat{y}^2\rangle|, \\ r_{III}^B: & \quad \delta\hat{y} \delta\hat{z} \geq \ell \left| \left\langle e^{\hat{z}} - (\hat{y} - 1)^2 \right\rangle \right|, \end{aligned} \quad (65)$$

and in all five cases there are (limits of) well-normalized states that annihilate the expectation values on the right-hand sides above.

As a first exploration of the locality structure of our noncommutative spacetimes, we can identify the regions of the lightcone where the right-hand sides of (65) vanish. Improper states defined as delta functions localized on a point of these regions will be perfectly localized, as both the uncertainties $\delta\hat{z}$ and $\delta\hat{y}$ will vanish on those states, and they will be obtainable as limits of sequences of well-normalized states on the corresponding algebras. These regions have been discussed already in Section 5, and are depicted in Fig. 2 as intersections between the lightcones and certain planes.

It is worth stressing that in the special case r_{III}^A , in which the subgroup K is a PL subgroup, its associated noncommutative spacetime (54d) is just quadratic in \hat{y}^2 , while all the other commutators

always possess a linear term when a formal power series expansion is considered. After quantization, this makes the noncommutativity of this special case milder, because one needs to localize a state further from the line $y = 0$ in order to see the fuzziness, compared to the other models.

To conclude, we stress that the approach here presented can be used with no modification for the case of noncommutative 3+1-dimensional lightcones, which can be constructed as quantizations of coisotropic PH structures coming from the PL structures of the $SO(3,1)$ Lorentz group when interpreted in a conformal setting. We believe that the new noncommutative lightcones (54) presented here offer a novel twist on an important issue in noncommutative physics: the interplay between the relaxed notion of locality implied by noncommutativity and the notion of causality, which is fundamental for QFT. Indeed, a number of studies in the past have focused on the possibility of fuzzy lightcones emerging from the spacetime correlations of noncommutative QFTs [16–19]; however, the noncommutativity we describe here stays on the lightcone. The lightcone itself is sharp, but there are limitations to localizing points *on the lightcone*. This feature might be interesting from a phenomenological point of view, in light of studies of propagation of massless particles in a quantum spacetime (see [38, 39]). This approach complements nicely the studies of noncommutative spaces of worldlines initiated recently [20–23], and it would be particularly interesting to compare the results of the present paper with those focusing on the spaces of noncommutative lightlike worldlines [21]. Work on all these lines is in progress and will be presented elsewhere.

Acknowledgements

M.A., A.B., and F.M. would like to acknowledge support from the Agencia Estatal de Investigación (Spain) under grants PID2019-106802GB-I00/AEI/10.13039/501100011033, the Q-CAYLE Project funded by the Regional Government of Castilla y León (Junta de Castilla y León), and the Ministry of Science and Innovation MICIN through NextGenerationEU (PRTR C17.I1). F.M. acknowledges support from the Agencia Estatal de Investigación (Spain) under grant CNS2023-143760.

References

- [1] H. S. Snyder, “Quantized Space-Time,” *Phys. Rev.* **71** (Jan, 1947) 38–41.
- [2] S. Hossenfelder, “Minimal Length Scale Scenarios for Quantum Gravity,” *Living Rev. Rel.* **16** (2013) 2.
- [3] A. Connes, “Noncommutative Differential Geometry,” *Inst. Hautes Études Sci. Publ. Math.* **62** (1985) 257.
- [4] S. L. Woronowicz, “Twisted $SU(2)$ Group. An Example of a Non-Commutative Differential Calculus,” *PRIMS* **23** no. 1, (1987) 117–181.
- [5] S. L. Woronowicz, “Compact matrix pseudogroups,” *Commun. Math. Phys.* **111** (1987) 613–665.
- [6] P. Vitale, M. Adamo, R. Dekhil, and D. Fernández-Silvestre, “Introduction to noncommutative field and gauge theory,” *PoS QG-MMSchools* (2024) 007.
- [7] S. Majid, *Foundations of Quantum Group Theory*. Cambridge University Press, Cambridge, 1995.
- [8] S. Majid, *A Quantum Groups Primer*. London Mathematical Society Lecture Note Series. Cambridge University Press, 2002.
- [9] A. N. P. Vyjayanthi Chari, *A guide to quantum groups*. Cambridge University Press, 1994.
- [10] R. J. Szabo, “Quantum field theory on noncommutative spaces,” *Phys. Rept.* **378** (2003) 207–299.
- [11] P. Aschieri, M. Dimitrijevic, P. Kulish, F. Lizzi, and J. Wess, *Noncommutative Spacetimes: Symmetries in Noncommutative Geometry and Field Theory*. Lecture Notes in Physics. Springer Berlin Heidelberg, 2011.
- [12] A. Chamseddine, C. Consani, N. Higson, M. Khalkhali, H. Moscovici, and G. Yu, *Advances in Noncommutative Geometry: On the Occasion of Alain Connes’ 70th Birthday*. Springer International Publishing, 2020.

- [13] V. G. Drinfel'd, "Hamiltonian structures of lie groups, lie bialgebras and the geometric meaning of the classical Yang-Baxter equations," *Sov. Math. Dokl.* **27** (1983) 68–71.
- [14] F. Lizzi, M. Manfredonia, F. Mercati, and T. Poulain, "Localization and Reference Frames in κ -Minkowski Spacetime," *Phys. Rev. D* **99** no. 8, (2019) 085003.
- [15] F. Lizzi, M. Manfredonia, and F. Mercati, "Localizability in κ -Minkowski spacetime," *Int. J. Geom. Meth. Mod. Phys.* **17** no. supp01, (2020) 2040010.
- [16] M. Arzano and L. T. Consoli, "Signal propagation on κ -Minkowski spacetime and nonlocal two-point functions," *Phys. Rev. D* **98** no. 10, (2018) 106018, [arXiv:1808.02241 \[hep-th\]](#).
- [17] F. Mercati and M. Sergola, "Light Cone in a Quantum Spacetime," *Phys. Lett.* **B787** (2018) 105–110.
- [18] F. Mercati and M. Sergola, "Pauli–Jordan function and scalar field quantization in κ -Minkowski noncommutative spacetime," *Phys. Rev.* **D98** no. 4, (2018) 045017.
- [19] N. Franco, K. Hersent, V. Maris, and J.-C. Wallet, "Quantum causality in κ -Minkowski and related constraints," *Class. Quant. Grav.* **40** no. 16, (2023) 164001.
- [20] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, "Noncommutative spaces of worldlines," *Phys. Lett. B* **792** (2019) 175–181.
- [21] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, "The noncommutative space of light-like worldlines," *Phys. Lett. B* **829** (2022) 137120.
- [22] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, "All noncommutative spaces of κ -Poincaré geodesics," *J. Phys. A: Math. Theor.* **55** no. 43, (Nov, 2022) 435205.
- [23] A. Ballesteros, G. Gubitosi, I. Gutierrez-Sagredo, and F. J. Herranz, " κ -Galilean and κ -Carrollian noncommutative spaces of worldlines," *Phys. Lett. B* **838** (Mar., 2023) 137735.
- [24] A. G. Reyman, "Poisson Structures Related to Quantum Groups," in *Quantum Groups and its Applications in Physics*, L. Castellani and J. Wess, eds., p. 407. IOS, Amsterdam, 1996.
- [25] X. Gomez, "Classification of three-dimensional Lie bialgebras," *J. Math. Phys.* **41** (2000) 4939.
- [26] J. Lukierski and V. N. Tolstoy, "Quantizations of $D = 3$ Lorentz symmetry," *Eur. Phys. J. C* **77** (2017) 226.
- [27] J.-H. Lu, *Multiplicative and affine Poisson structures on Lie groups*. PhD thesis, University of California, Berkeley, 1990.
- [28] N. Ciccoli and F. Gavarini, "A quantum duality principle for coisotropic subgroups and Poisson quotients," *Adv. Math.* **199** (2006) 104–135.
- [29] A. Ballesteros, C. Meusburger, and P. Naranjo, "AdS Poisson homogeneous spaces and Drinfel'd doubles," *J. Phys. A: Math. Theor.* **50** (2017) 395202.
- [30] A. Ballesteros, I. Gutierrez-Sagredo, and F. Mercati, "Coisotropic Lie bialgebras and complementary dual Poisson homogeneous spaces," *J. Phys. A Math. Theor.* **54** (2021) 315203.
- [31] M. S. Dijkuizen and T. H. Koornwinder, "Quantum homogeneous spaces, duality and quantum 2-spheres," *Geom. Dedicata* **52** (1994) 291–315.
- [32] A. Ballesteros, G. Gubitosi, I. Gutierrez-Sagredo, and F. Mercati, "Fuzzy worldlines with κ -Poincaré symmetries," *J. High Energ. Phys.* **2021** (2021) 80.
- [33] V. G. Drinfel'd, "Quantum Groups," in *Proc. Int. Congr. Math. (Berkeley 1986)*, A. Gleason, ed., pp. 798–820. American Mathematical Society, Providence RI, 1987.
- [34] E. Abe, *Hopf Algebras*. Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 2004.
- [35] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, "The κ -(A)dS noncommutative spacetime," *Phys. Lett. B* **796** (2019) 93–101.
- [36] A. Ballesteros, I. Gutierrez-Sagredo, and F. J. Herranz, "Noncommutative (A)dS and Minkowski spacetimes from quantum Lorentz subgroups," *Class. Quantum Grav.* **39** (2022) 015018.
- [37] L. Dabrowski and G. Piacitelli, "Canonical κ -Minkowski Spacetime," [arXiv:1004.5091 \[math-ph\]](#).
- [38] A. Addazi, "Quantum gravity phenomenology at the dawn of the multi-messenger era – A review," *Prog. Part. Nucl. Phys.* **125** (2022) 103948.
- [39] R. Alves Batista *et al.*, "White Paper and Roadmap for Quantum Gravity Phenomenology in the Multi-Messenger Era," [arXiv:2312.00409 \[gr-qc\]](#).