Hypergeometric sheaves and extraspecial groups in even characteristic

Lee Tae Young

Department of Mathematics, Rutgers University, Piscataway, NJ 08854, USA tae.young@binghamton.edu

July 18, 2024

Abstract

We determine precisely which irreducible hypergeometric sheaves have an extraspecial normalizer in characteristic 2 as their geometric monodromy groups. This resolves the last open case of the determination of local monodromy at 0 of irreducible hypergeometric sheaves with finite geometric monodromy group.

2010 Mathematics Subject Classification: 20C15, 11T23, 20D15 **Keywords:** Local systems, Monodromy groups, Extraspecial groups, Exponential Sums

Contents

1	Introduction	1
2	Preliminaries	2
3	m2sp-elements of extraspecial normalizers	3
4	Local monodromy at ∞ of hypergeometric sheaves	6
5	Determination of the finiteness of the monodromy	14

1 Introduction

Let p be a prime. The finite quotient groups of the étale fundamental group $\pi_1^{et}(\mathbb{G}_m/\overline{\mathbb{F}_p})$ are the finite groups generated by its Sylow p-subgroups together with at most one other element; this was conjectured by Abhyankar [1] and proved by Harbater [4]. In their recent series of papers, Katz, Rojas-León and Tiep realized many pairs of such finite group G and its

faithful complex representation V as the geometric monodromy groups of explicitly written $\overline{\mathbb{Q}_{\ell}}$ -local systems, usually hypergeometric sheaves, on $\mathbb{G}_m/\overline{\mathbb{F}_p}$.

In [7–9], they determined which pairs (G, V) can be realized using irreducible hypergeometric sheaves. It would be desirable to achieve a complete classification of irreducible hypergeometric sheaves in terms of their monodromy groups. As hypergeometric sheaves are completely determined by its local monodromies, one can first ask which local monodromy at 0 can be realized. Specifically, we want to see which triples (G, V, g), where $g \in G$ is an element with certain properties which must be satisfied by the local monodromy at 0 of a hypergeometric sheaf, can be realized by irreducible hypergeometric sheaves. This was also completed in [7–9] for all but one type of groups: the so-called extraspecial normalizers in characteristic 2.

The goal of this paper is to completely answer this question, not only about the local monodromy at 0 but also at ∞ , and write down a complete classification of irreducible hypergeometric sheaves whose geometric monodromy group is a finite extraspecial normalizer in characteristic 2. In Theorem 3.1, we classify the elements of extraspecial normalizers in characteristic 2 which has odd order and has no eigenvalues of multiplicity exceeding 2 on an 2^n -dimensional representation extending the unique irreducible representation of the normal extraspecial 2-subgroup of order 2^n of that dimension. This is one of several properties that must be satisfied by the local monodromies at 0 and ∞ of irreducible hypergeometric sheaves. After taking some properties of hypergeometric sheaves into account, in Theorem 4.4 we get a more restrictive list of possible local monodromies. We then study the hypergeometric sheaves defined by each combination of the local monodromies at 0 and ∞ , and determine precisely when it has finite geometric monodromy group. This is our main result Theorem 5.7, which says that these local systems are precisely those discovered and studied earlier by Pink, Sawin, Katz and Tiep in [8,9], where the isomorphism type of their geometric monodromy groups were determined. This completes the classification of the local systems in question.

2 Preliminaries

In this section, we recall some definitions and basic properties of the objects we will study. For more definitions and properties, we refer to [5] and [8].

For $x \in \{0, \infty\}$, we fix a choice of an inertia group $I(x) < \pi_1^{et}(\mathbb{G}_m/\overline{\mathbb{F}_p})$, and we denote by P(x) the wild inertial group in I(x). We also fix an element γ_x of order prime to p in I(x) whose image in I(x)/P(x) is a topological generator of this pro-cyclic group.

For a nontrivial additive character ψ of \mathbb{F}_p and multiplicative characters χ_1, \ldots, χ_D and ρ_1, \ldots, ρ_M of a finite extension of \mathbb{F}_p , there exists a hypergeometric sheaf of type (D, M)

$$\mathcal{H} = \mathcal{H}yp_{\psi}(\chi_1, \dots, \chi_D; \rho_1, \dots, \rho_M).$$

It is tame at 0, and its local monodromy at ∞ (that is, the restriction to $I(\infty)$) is given by Tame \oplus Wild, where Tame is a *M*-dimensional tame representation and Wild is a (D - M)-dimensional irreducible representation of Swan conductor 1. The geometric monodromy

group $G_{\text{geom}} \leq \operatorname{GL}_D(\overline{\mathbb{Q}_\ell})$ of \mathcal{H} is the Zariski closure of the image of $\pi_1^{et}(\mathbb{G}_m/\overline{\mathbb{F}_p})$ on \mathcal{H} . We denote by $J, Q \leq G_{\text{geom}}$ and $g_x \in G_{\text{geom}}$ the images of $I(\infty)$, $P(\infty)$ and γ_x on \mathcal{H} when the local system is clear from the context. The restriction Wild $|_Q$ (or equivalently Wild $|_{P(\infty)}$) decomposes into a direct sum of W_0 pairwise nonisomorphic irreducible representations, where W_0 is the p'-part of dim Wild = D - M.

The characters χ_i are called the *upstairs characters*, and ρ_j are called the *downstairs characters* of \mathcal{H} . \mathcal{H} is irreducible if and only if no $\{\chi_1, \ldots, \chi_D\} \cap \{\rho_1, \ldots, \rho_M\} = \emptyset$. Also, G_{geom} can be finite only when χ_1, \ldots, χ_D are pairwise distinct and so are ρ_1, \ldots, ρ_M . For all hypergeometric sheaves in the rest of this paper, we will always assume that these properties are satisfied. In particular, the element $g_0 \in G_{\text{geom}} < \operatorname{GL}_D(\overline{\mathbb{Q}_\ell})$ must be an **ssp**-element and g_∞ must be an **m2sp**-element, as defined below.

Definition. An element $g \in GL_n(\mathbb{C})$ is said to be

- (i) an ssp-element if each eigenvalue of g has multiplicity 1, and
- (ii) an m2sp-element if each eigenvalue of g has multiplicity at most 2.

Katz and Tiep [9, Theorem 5.2.9] [10] showed that most irreducible primitive hypergeometric sheaves (as representations of their G_{geom}) satisfy the following condition.

Definition. A group G and its faithful complex representation V (or equivalently, a subgroup $G \leq \operatorname{GL}_n(\mathbb{C})$) is said to satisfy *condition* (**S**) if V is irreducible, primitive, tensor indecomposable, and not tensor induced. If in addition $\mathbf{Z}(G)$ is finite, then we say (G, V)satisfies *condition* (**S**+).

By a result of Guralnick and Tiep [3, Proposition 2.8], such finite groups, in particular the G_{geom} of (S+)-hypergeometric sheaves, must be either almost quasisimple or an extraspecial normalizer as defined below.

Definition. An extraspecial normalizer in characteristic p is a finite group $G < \operatorname{GL}(V)$ which contains a normal subgroup $R = \mathbb{Z}(R)E$, where $V = \mathbb{C}^{p^n}$ for some $1 \leq n \in \mathbb{Z}$, E is an extraspecial p-group of order p^n which acts irreducibly on V, and either R = E or $\mathbb{Z}(R) \cong C_4$, the cyclic group of order 4.

In [7–9], Katz, Rojas-León and Tiep determined whether each triple (G, V, g) of a finite group G which is either almost quasisimple or an extraspecial normalizer, its faithful complex representation V, and an element $g \in G$ which acts as an **ssp**-element on V can be realized as the G_{geom} and g_0 of an irreducible hypergeometric sheaf. The only case which was not completely covered by their results is the case of extraspecial normalizers in characteristic 2.

3 m2sp-elements of extraspecial normalizers

To find the candidates for g_{∞} , we first find all m2sp-elements of odd order in extraspecial normalizers in characteristic 2. This is just an extension of [8, Theorem 8.5], which classifies the ssp-elements.

Theorem 3.1. Let $V = \mathbb{C}^{2^n}$ and let G be a finite irreducible subgroup of $\operatorname{GL}(V)$ that satisfies $(\mathbf{S}+)$ and is an extraspecial normalizer, so that $G \triangleright R = \mathbf{Z}(R)E$ for some normal subgroup R which contains an extraspecial 2-group $E = 2_{\epsilon}^{1+2n}$ that acts irreducibly on V, and either R = E or $\mathbf{Z}(R) \cong C_4$. Suppose that an element $g \in G$ of odd order has no eigenvalues with multiplicity larger than 2 on V and that $2^n \ge 16$. Then there exists integers a_1, \ldots, a_t such that $a_1 + \cdots + a_t = n$, integers r_1, \ldots, r_t such that $0 \le r_i < 2^{a_i} - \epsilon_i$ where $\epsilon_i = -1$ for $2 \le i \le t$ and $\epsilon_1 = (-1)^{t-1}\epsilon$, so that the image of g in $G/\mathbf{Z}(G)R$ lies in a maximal torus

$$C_{2^{a_1}-\epsilon_1}\times\cdots\times C_{2^{a_t}-\epsilon_t}<\mathbf{O}_{2a_1}^{\epsilon_1}\times\cdots\times\mathbf{O}_{2a_t}^{\epsilon_t}\leq\mathbf{O}_{2n}^{\epsilon}.$$

Moreover, one of the following holds:

- (a) $2^{a_1} + 1, \ldots, 2^{a_t} + 1$ are pairwise coprime, and $gcd(r_i, 2^{a_i} + 1) = 1$ for all *i*. In this case $\overline{o}(g) = \prod_{i=1}^{t} (2^{a_i} + 1)$. (This is precisely the situation of [8, Theorem 8.5(ii)].)
- (b) $2^{a_1}+1,\ldots,2^{a_{t-1}}+1$ are pairwise coprime, $gcd(r_i,2^{a_i}+1)=1$ for $1 \le i \le t-1$, $a_t=1$, and $r_t=0$. In this case $\overline{o}(g) = \prod_{i=1}^{t-1} (2^{a_i}+1) = (\prod_{i=1}^t (2^{a_i}+1))/3$.
- (c) $2^{a_1} + 1, \ldots, 2^{a_{t-1}} + 1$ are pairwise coprime, $gcd(r_i, 2^{a_i} + 1) = 1$ for all $i, a_t = 1$, and a_i is odd for some (unique) i < t. In this case, n is even and $\overline{o}(g) = \prod_{i=1}^{t-1} (2^{a_i} + 1) = (\prod_{i=1}^{t} (2^{a_i} + 1))/3$.
- (d) $2^{a_1} 1, 2^{a_2} + 1, \dots, 2^{a_t} + 1$ are pairwise coprime, $gcd(r_i, 2^{a_i} + 1) = 1$ for $2 \le i \le t$, and $gcd(r_1, 2^{a_1} 1) = 1$.

In cases (a), (b) and (c), the spectrum of g on V is

$$\left\{\xi\prod_{i=1}^{t}\zeta_{i}^{r_{i}}\mid\zeta_{i}\in\mu_{2^{a_{i}}+1}\setminus\{1\}\right\}$$

as a multiset, where ξ is a root of unity of odd order. In case (d), the spectrum of g on V is

$$\left\{ \xi \prod_{i=1}^{t} \zeta_{i}^{r_{i}} \mid \zeta_{1} \in \mu_{2^{a_{1}}-1} \sqcup \{1\}, \zeta_{i} \in \mu_{2^{a_{i}}+1} \setminus \{1\} \text{ for } i \geq 2 \right\}$$

as a multiset, where ξ is again a root of unity of odd order.

Proof. We use the notations in the proof of [8, Theorem 8.5]. By that proof we know that $V = V_1 \otimes V_2 \otimes \cdots \otimes V_t$ and $E = E_1 \circ E_2 \circ \cdots \circ E_t$ where $V_i = \mathbb{C}^{2^{a_i}}$ and $E_i \cong 2^{1+2a_i}_{\epsilon_i}$ for $\epsilon_i = \pm (\epsilon_i = \pm 1$ when interpreted as a number). Also, g = zh for some $z \in \mathbb{Z}(GL(V))$ and $h \in \langle s_1, \ldots, s_t \rangle$, where $s_i \in \operatorname{Sp}^{\epsilon_i}(V_i) < \operatorname{Sp}^{\epsilon}(V)$ is an element of order $2^{a_i} - \epsilon_i$ whose spectrum on V_i is $\mu_{2^{a_i}+1} \setminus \{1\}$ if $\epsilon_i = -$, and $\mu_{2^{a_i}-1}$ (with 1 having multiplicity 2) if $\epsilon_i = +$. We may write $h = s_1^{r_1} s_2^{r_2} \cdots s_t^{r_t}$ for $0 \leq r_i < 2^{a_i} - \epsilon_i$.

Suppose that g has no eigenvalues of multiplicity larger than 2 on V. Since G acts irreducibly on V, z acts as a scalar. Hence $h = gz^{-1}$, whose spectrum is given by the

product of the spectra of $s_i^{r_i}$'s as described above, also has no eigenvalue of multiplicity larger than 2. If $\epsilon_{i_1} = \epsilon_{i_2} = +$ for some $i_1 \neq i_2$, then the spectrum of $s_{i_1}s_{i_2}$ on $V_{i_1} \otimes V_{i_2}$ has eigenvalue 1 with multiplicity 4, whence the spectrum of h must also have eigenvalue with multiplicity at least 4. Therefore, ϵ_i can be + for at most one i.

Suppose that $\epsilon_i = -$ for every *i*. Then the spectrum of *h* on *V* is

$$\left\{\prod_{i=1}^{t} \zeta_i^{r_i} \mid \zeta_i \in \mu_{2^{a_i}+1} \setminus \{1\}\right\} \text{ (as a multiset)}$$

In order to have no eigenvalue of multiplicity larger than 2, we must have $gcd(r_i, 2^{a_i} + 1) = 1$ for all *i* with at most one possible exception $i = i_0$ for which $(r_{i_0}, a_{i_0}) = (0, 1)$. Also, if $d := gcd(2^{a_{i_1}} + 1, 2^{a_{i_2}} + 1) > 1$ for some $i_1 \neq i_2$ and $gcd(r_{i_1}, 2^{a_{i_1}} + 1) = gcd(r_{i_2}, 2^{a_{i_2}} + 1) = 1$, then we may choose $(\zeta_i, \zeta_j) = (\zeta, \zeta^{-1}), (\zeta^2, \zeta^{-2}), \ldots, (\zeta^{-1}, \zeta)$ for some primitive *d*th root of unity ζ to get d - 1 copies of some eigenvalue. The only possibility is d = 3. Moreover, if both $2^{a_{i_1}} + 1$ and $2^{a_{i_2}} + 1$ are larger than 3, then we can instead choose $(\zeta_{i_1}, \zeta_{i_2}) =$ $(\xi, \eta), (\xi\zeta, \eta\zeta^{-1}), (\xi\zeta^{-1}, \eta\zeta)$ for some primitive $2^{a_{i_1}} + 1$ th and $2^{a_{i_2}} + 1$ th roots of unity ξ and η to get 3 copies of same eigenvalue. Therefore, either $a_i = 1$ or $a_j = 1$. Of course, at most one such i_0 and at most one such pair (i_1, i_2) can exist, and if they both exist then $i_0 \in \{i_1, i_2\}$. After reindexing if necessary, we can assume that $a_1 > a_2 > \cdots > a_{t-1} \ge a_t$, and we are in one of the situations (a), (b) and (c).

Next, suppose that $\epsilon_1 = +$ and $\epsilon_2 = \cdots = \epsilon_t = -$. The spectrum of h on V is now

$$\left\{\prod_{i=1}^{t} \zeta_{i}^{r_{i}} \mid \zeta_{1} \in \mu_{2^{a_{1}}-1} \sqcup \{1\}, \zeta_{i} \in \mu_{2^{a_{i}}+1} \setminus \{1\} \text{ for } i \geq 2\right\} \text{ (as a multiset).}$$

Since 1 has multiplicity 2 in $\mu_{2^{a_1}-1} \sqcup \{1\}$, the numbers $\prod_{i=2}^{t} \zeta_i^{r_i}$ should be all distinct, and no two of them differ by a factor of a $(2^{a_1}-1)$ th root of unity. Therefore we are in the situation (d).

In the above theorem, we had the condition that $2^{a_i} + 1$ are pairwise coprime. It might be convenient to note the following easy equivalent condition.

Lemma 3.2. $2^{a} + 1$ and $2^{b} + 1$ are coprime if and only if a and b have different 2-parts (the highest power of 2 dividing it).

Proof. Let a_2 and b_2 be the 2-parts of a and b. Note that $2^a + 1 = (2^{a_2} + 1)(2^{a-a_2} - 2^{a-2a_2} + \cdots - 2^{a_2} + 1)$. The "only if" direction is immediate from this. For the converse, assume that $a_2 > b_2$, and observe that

$$\gcd(2^{a}+1,2^{b}+1) \mid \gcd(2^{2a}-1,2^{2b}-1) = 2^{2\gcd(a,b)} - 1 = (2^{\gcd(a,b)}-1)(2^{\gcd(a,b)}+1).$$

Since $2^{\operatorname{gcd}(a,b)} - 1$ divides $2^a - 1$, it is coprime to $2^a + 1$. Also, $\operatorname{gcd}(a,b)$ is not divisible by a_2 , so $a/\operatorname{gcd}(a,b)$ is even. Hence $2^{\operatorname{gcd}(a,b)} + 1$ also divides $2^a - 1$, so it is coprime to $2^a + 1$. Therefore $\operatorname{gcd}(2^a + 1, 2^b + 1) = 1$.

4 Local monodromy at ∞ of hypergeometric sheaves

Let \mathcal{H} be an irreducible hypergeometric sheaf over a field \mathbb{F}_q of characteristic 2 which is of type (D, M) with W := D - M > 0 and whose geometric monodromy group $G := G_{\text{geom}}$ is an extraspecial normalizer of characteristic 2. By [8, Theorem 9.19(ii)], G has a normal extraspecial 2-subgroup $E \cong 2^{1+2n}_{\pm}$ where $2^n = D$, and $\mathbf{Z}(E)$ is the Sylow 2-subgroup of $\mathbf{Z}(G)$; in other words, R = E.

In this section, we determine which of the elements in Theorem 3.1 can appear as g_{∞} of \mathcal{H} . We must take into account the structure of the image Q of $P(\infty)$.

Proposition 4.1. At least one of the following holds:

- (a) \mathcal{H} is Kloosterman,
- (b) $Q \cap E = 1$, so the image of Q in $G/\mathbf{Z}(G)E$ is isomorphic to Q, or
- (c) $(Q \cap E)\mathbf{Z}(E)$ is elementary abelian, each irreducible constituent of Wild $|_{Q \cap E}$ is nontrivial, and (dim Tame) $\cdot |\{Q \cap Q^e \cap E \mid e \in E\}| = D = 2^n$.

Proof. Suppose that \mathcal{H} is not in the cases (a) and (b), so that $Q \cap E$ is nontrivial and dim Tame > 0. By [8, Proposition 4.8], $Q \cap \mathbf{Z}(G) = 1$ and $|\mathbf{Z}(G)|_2 \leq 2$. Since Q is a 2-group and $\mathbf{Z}(E) \leq \mathbf{Z}(G)$, we have $Q \cap \mathbf{Z}(E) = 1$. Therefore, $Q \cap E$ is isomorphic to its image in $E/\mathbf{Z}(E)$, so $(Q \cap E)\mathbf{Z}(E)$ is an elementary abelian 2-group.

Let Φ be the monodromy representation of \mathcal{H} . Consider its restrictions to the subgroups $Q \cap E \triangleleft (Q \cap E)\mathbf{Z}(E) \trianglelefteq E$. Since $Q \cap E \trianglelefteq Q \trianglelefteq J$, the restriction $\Phi|_{Q \cap E}$ decomposes into $\mathsf{Tame}|_{Q \cap E} \oplus \mathsf{Wild}|_{Q \cap E} = (\dim \mathsf{Tame}) \cdot \mathbb{1}_{Q \cap E} \oplus \mathsf{Wild}|_{Q \cap E}$. Hence, there exists some irreducible constituent Ψ of $\Phi|_{(Q \cap E)\mathbf{Z}(E)}$ (which is one-dimensional since $(Q \cap E)\mathbf{Z}(E)$ is abelian as we saw above) whose restriction to $Q \cap E$ is trivial. On the other hand, since $\Phi|_E$ is irreducible and faithful, by Clifford theory, no irreducible constituent of its restriction to the normal subgroup $(Q \cap E)\mathbf{Z}(E)$ can contain a normal subgroup of E in the kernel, so in particular $\Psi \neq \mathbb{1}_{(Q \cap E)\mathbf{Z}(E)}$. Therefore ker $\Psi = Q \cap E$, and each irreducible constituent of $\Phi|_{(Q \cap E)\mathbf{Z}(E)}$ has kernel $Q^e \cap E$ for some $e \in E$. In particular, no irreducible constituent of $\Phi|_{(Q \cap E)\mathbf{Z}(E)}$ other than Ψ restricts to $\mathbb{1}_{Q \cap E}$, so the multiplicity of Ψ in $\Phi|_{(Q \cap E)\mathbf{Z}(E)}$ is exactly the multiplicity of $\mathbb{1}_{Q \cap E}$ in $\Phi|_{Q \cap E}$, and it divides dim Φ .

Now consider the restriction of the irreducible representation Wild of J to the normal subgroups $Q \cap E \trianglelefteq Q \trianglelefteq J$. If Wild $|_{Q \cap E}$ has $\mathbb{1}_{Q \cap E}$ as an irreducible constituent, then there exists an irreducible constituent Θ of Wild $|_Q$ with $Q \cap E \le \ker \Theta$. Since $Q \cap E \trianglelefteq J$, it follows that $Q \cap E \le \bigcap_{j \in J} (\ker \Theta)^j = \ker \operatorname{Wild} |_Q$. This is impossible since Φ is faithful and $Q \le \ker \operatorname{Tame}$. Therefore, Wild $|_{Q \cap E}$ cannot have $\mathbb{1}_{Q \cap E}$ as an irreducible constituent, so the multiplicity of $\mathbb{1}_{Q \cap E}$ in $\Phi|_{Q \cap E}$ is exactly dim Tame. By the previous paragraph, this number divides dim $\Phi = D = 2^n$.

As we now know what are the possible intersections $Q \cap E$, it is natural to ask what are the possible quotients $Q/(Q \cap E)$. As g_{∞} normalizes Q, we can simply find all 2-subgroups of $G/\mathbb{Z}(G)E$ which are normalized by the elements described in Theorem 3.1. **Proposition 4.2.** Let a_1, \ldots, a_t be positive integers such that $a_1 + \cdots + a_t = n$, and let $g = (g_1, \ldots, g_t)$ be an element of $\operatorname{Sp}_{2a_1}(2) \times \cdots \times \operatorname{Sp}_{2a_t}(2) \leq \operatorname{Sp}_{2n}(2)$. Suppose that P is a 2-subgroup of $\operatorname{Sp}_{2n}(2)$ such that $g \in \mathbf{N}_{\operatorname{Sp}_{2n}(2)}(P)$.

- (a) Suppose that g_1, \ldots, g_t are irreducible over \mathbb{F}_2 , have distinct orders $o(g_i)$, and every nonzero orbit of the action of g_i on $\mathbb{F}_2^{2a_i}$ has length $o(g_i)$. Then P must be trivial.
- (b) Suppose that g₁,..., g_{t-1} are irreducible over F₂, have distinct orders o(g_i), and every nonzero orbit of the action of g_i on F₂^{2a_i} has length o(g_i). Also suppose that V_t = V_{t1}⊕V_{t2} for some a_t-dimensional subspaces V_{t1} and V_{t2}, g_t = g_{t,1} ⊕ g_{t,2} ∈ (GL(V_{t1}) × GL(V_{t2})) ∩ Sp_{2at}(2) for some g_{t1} and g_{t2} which are irreducible over F₂, has same order o(g_t) which is not equal to any other o(g_i), and every nonzero orbit of the actions of g_{t,1} and g_{t,2} on F₂^{at} has length o(g_t). Then P must be elementary abelian of order not exceeding 2^{a²}.

Proof. Let
$$A_m$$
 be the $m \times m$ matrix $\begin{pmatrix} & 1 \\ & \ddots & \\ 1 & & \end{pmatrix}$. Let $V = V_1 \oplus \cdots \oplus V_t$ be the natural

module of $\operatorname{Sp}_{2a_1}(2) \times \cdots \times \operatorname{Sp}_{2a_t}(2)$ with the symplectic form $A_{2a_1} \oplus \cdots \oplus A_{2a_t}$.

(a) We first show that V_1, \ldots, V_t are the minimal $\langle g \rangle$ -submodules of V. Since each g_i is irreducible, it is clear that each V_i is a minimal $\langle g \rangle$ -submodule of V. Conversely, suppose that a minimal submodule U contains a vector $v = (v_1, \ldots, v_t) \in V_1 \oplus \cdots \oplus V_t$ with at least two nonzero components v_i, v_j . We may assume that $o(g_i) < o(g_j)$. Then $g^{o(g_i)}v - v =$ $(g_1^{o(g_i)}v_1 - v_1, \ldots, g_t^{o(g_t)}v_t - v_t) \in M$. The *i*th component of this vector is $g_i^{o(g_i)}v_i - v_i = 0$, while the *j*th component is nonzero. We may repeat this until we get a vector with only one nonzero component. Therefore, M intersects at least one of V_1, \ldots, V_t nontrivially, so by the minimality, M is one of them.

Since g_1, \ldots, g_t are irreducible over \mathbb{F}_2 and have pairwise coprime orders, the minimal $\langle g \rangle$ -submodules of V are V_1, \ldots, V_t . Since P is a 2-subgroup of $\operatorname{GL}_{2n}(2)$ normalized by g, $\mathbf{C}_V(P)$ is a nontrivial $\langle g \rangle$ -submodule of V, so it must contain some V_i . After relabeling if necessary, we may assume that $V_1 \subseteq \mathbf{C}_V(P)$. Now the action of P on V/V_1 fixes a nontrivial $\langle g \rangle$ -submodule by the same reason. Repeating this, we can see that after relabeling,

$$P \leq \left\{ \begin{pmatrix} I_{2a_1} & X_{1,2} & \cdots & X_{1,t} \\ 0 & I_{2a_2} & \cdots & X_{2,t} \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & 0 & I_{2a_t} \end{pmatrix} \middle| X_{i,j} \in M_{2a_i \times 2a_j}(\mathbb{F}_2) \right\}.$$

On the other hand, the elements of P preserves the symplectic form $A_{2a_1} \oplus \cdots \oplus A_{2a_t}$, so

$$\begin{pmatrix} A_{2a_{1}} & & \\ & A_{2a_{2}} & & \\ & & \ddots & \\ & & & A_{2a_{t}} \end{pmatrix}$$

$$= \begin{pmatrix} I_{2a_{1}} & & \\ X_{1,2}^{T} & I_{2a_{2}} & & \\ \vdots & \vdots & \ddots & \\ X_{1,t}^{T} & X_{2,t}^{T} & \cdots & I_{2a_{t}} \end{pmatrix} \begin{pmatrix} A_{2a_{1}} & & \\ & A_{2a_{2}} & & \\ & & \ddots & \\ & & A_{2a_{t}} \end{pmatrix} \begin{pmatrix} I_{2a_{1}} & X_{1,2} & \cdots & X_{1,t} \\ 0 & I_{2a_{2}} & \cdots & X_{2,t} \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & 0 & I_{2a_{t}} \end{pmatrix}$$

$$= \begin{pmatrix} A_{2a_{1}} & & \\ X_{1,2}^{T}A_{2a_{1}} & X_{2,t}^{T}A_{2a_{2}} & & \\ \vdots & \vdots & \ddots & \\ X_{1,t}^{T}A_{2a_{1}} & X_{1,2}^{T}A_{2a_{2}} & \cdots & A_{2a_{t}} \end{pmatrix} \begin{pmatrix} I_{2a_{1}} & X_{1,2} & \cdots & X_{1,t} \\ 0 & I_{2a_{2}} & \cdots & X_{2,t} \\ 0 & 0 & \ddots & \vdots \\ 0 & 0 & 0 & I_{2a_{t}} \end{pmatrix}$$

$$= \begin{pmatrix} A_{2a_{1}} & & A_{2a_{1}}X_{1,2} & \cdots & A_{2a_{1}}X_{1,t} \\ A_{2a_{1}} & X_{1,2}^{T}A_{2a_{1}} & X_{1,2}^{T}A_{2a_{1}}X_{1,2} + A_{2a_{2}} & \cdots & X_{1,t}^{T}A_{2a_{1}}X_{1,t} + A_{2a_{1}}X_{2,t} \\ \vdots & \vdots & \ddots & \vdots \\ X_{1,t}^{T}A_{2a_{1}} & X_{1,t}^{T}A_{2a_{1}}X_{1,2} + X_{2,t}^{T}A_{2a_{2}} & \cdots & X_{1,t}^{T}A_{2a_{1}}X_{1,t} + \cdots + A_{2a_{t}} \end{pmatrix}$$

which shows that $X_{i,j} = 0$ for all pairs (i, j). Therefore P is trivial.

(b) By the same argument as in (a), we see that the minimal g_{∞} -invariant subspaces of V are V_1, \ldots, V_{t-1} and certain subspaces of V_t . Moreover, the minimal polynomials of $g_{t,1}$ and $g_{t,2}$ are both irreducible of degree a_t , so their least common multiple is the minimal polynomial of g_t . Hence, every minimal g_{∞} -invariant subspace of V_t has dimension a_t . Therefore, for a suitable choice of basis, we have

$$P \leq \left\{ \begin{pmatrix} I_{2a_1} & X_{1,2} & \cdots & X_{1,t-1} & X_{1,t} & X_{1,t+1} \\ 0 & I_{2a_2} & \cdots & X_{2,t-1} & X_{2,t} & X_{2,t+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & I_{2a_{t-1}} & X_{t-1,t} & X_{t-1,t+1} \\ X_{t,1} & X_{t,2} & \cdots & X_{t,t-1} & I_{a_t} & X_{t,t+1} \\ X_{t+1,1} & X_{t+1,2} & \cdots & X_{t+1,t-1} & 0 & I_{a_t} \end{pmatrix} \right\}$$

where the symplectic form in this basis becomes $A_{2a_1} \oplus \cdots \oplus A_{2a_{t-1}} \oplus B$ for some symplectic form $B = \begin{pmatrix} B_1 & B_2 \\ B_2^T & B_3 \end{pmatrix}$ on $\mathbb{F}_2^{2a_t}$, where B_1, B_2, B_3 are $a_t \times a_t$ matrices. Also, there exists $0 \leq i_1 \leq i_2 \leq t-1$ such that $X_{i,t} = 0$ for all $i > i_1, X_{i,t+1} = 0$ for all $i > i_2, X_{t,i} = 0$ for all $i \leq i_1$, and $X_{t+1,i} = 0$ for all $i \leq i_2$. In particular, for each $i = 1, \ldots, t-1$, at least one of $X_{i,t}$ and $X_{t,i}$ is 0, and at least one of $X_{i,t+1}$ and $X_{t+1,i}$ is 0. Since each element of P preserves the symplectic form, we have the following equality:

$$\begin{pmatrix} A_{2a_1} & & & \\ & \ddots & & \\ & & B_1 & B_2 \\ & & B_2^T & B_3 \end{pmatrix}$$

$$= \begin{pmatrix} I_{2a_1} & & X_{t,1}^T & X_{t+1,1}^T \\ X_{1,2}^T & I_{2a_2} & & X_{t,2}^T & X_{t+1,2}^T \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ X_{1,t-1}^T & X_{2,t-1}^T & I_{2a_{t-1}} & X_{t,t-1}^T & X_{t+1,t-1}^T \\ X_{1,t+1}^T & X_{2,t+1}^T & X_{t-1,t+1}^T & X_{t-1,t+1}^T & I_{at} \end{pmatrix} \begin{pmatrix} A_{2a_1} & & \\ & A_{2a_2} \\ & \ddots & \\ & & A_{2a_{t-1}} \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & &$$

If $i_2 = 0$, then $X_{i,t} = X_{i,t+1} = 0$ for all $i = 1, \ldots, t-1$. If $i_1 > 0$, then $X_{t,1} = X_{t+1,1} = 0$, so by computing the first row of the product above, we get $X_{1,2} = \cdots = X_{1,t+1} = 0$. If $i_1 = 0 < i_2$, then $X_{i,t} = X_{t+1,1} = 0$ for $i = 1, \ldots, t-1$, so by computing the (1, t+1)- and (t, t+1)-entries, we get $A_{2a_1}X_{1,t+1} + X_{t,1}^TB_1X_{t,t+1} + X_{t,1}^TB_2 = 0$ and $B_1X_{t,t+1} + B_2 = 0$, so that $X_{1,t+1} = 0$. Therefore, regardless of the numbers i_1 and i_2 and the symplectic form B, we always have $X_{1,t} = X_{1,t+1} = 0$.

Similarly, assume that $X_{j,t} = X_{j,t+1} = 0$ for all j < i. If $i_2 < i$, then $X_{i,t} = X_{i,t+1} = 0$. If $i_1 \ge i$, then $X_{t,j} = X_{t+1,j} = 0$ for j = 1, ..., i, so by computing the (i, t)- and (i, t + 1)entry, we get $A_{2a_i}X_{i,t} = 0$ and $A_{2a_i}X_{i,t+1} = 0$. If $i_1 < i \le i_2$, then $X_{k,t} = X_{t+1,j} = 0$ for k = i, ..., t - 1 and j = 1, ..., i, so by computing the (i, t + 1) and (t, t + 1)-entries, we get $A_{2a_i}X_{i,t+1} + X_{t,i}^TB_1X_{t,t+1} + X_{t,i}^TB_2 = 0$ and $B_1X_{t,t+1} + B_2 = 0$, so that $A_{2a_i}X_{i,t+1} = 0$. Therefore, in any case, we get $X_{i,t} = X_{i,t+1} = 0$. By induction we get $X_{i,t} = X_{i,t+1} = 0$ for all $i \in \{1, ..., t - 1\}$. The above matrix now becomes

$$\begin{pmatrix} A_{2a_1} & 0 & \cdots & 0 & X_{t,1}^T B_1 + X_{t+1,1}^T B_2^T & X_{t,1}^T B_2 + X_{t+1,1}^T B_3 \\ X_{1,2}^T A_{2a_1} & A_{2a_2} & \cdots & 0 & X_{t,2}^T B_1 + X_{t+1,2}^T B_2^T & X_{t,2}^T B_2 + X_{t+1,2}^T B_3 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ X_{1,t-1}^T A_{2a_1} & X_{2,t-1}^T A_{2a_2} & \cdots & A_{2a_{t-1}} & X_{t,t-1}^T B_1 + X_{t+1,t-1}^T B_2^T & X_{t,t-1}^T B_2 + X_{t+1,t-1}^T B_3 \\ 0 & 0 & \cdots & 0 & B_1 & B_2 \\ 0 & 0 & \cdots & 0 & X_{t,t+1}^T B_1 + B_2^T & X_{t,t+1}^T B_2 + B_3 \end{pmatrix} \begin{pmatrix} I_{2a_1} & X_{1,2} & \cdots & X_{1,t-1} & 0 & 0 \\ 0 & I_{2a_2} & \cdots & X_{2,t-1} & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & I_{2a_{t-1}} & 0 & 0 \\ X_{t,1} & X_{t,2} & \cdots & X_{t,t-1} & I_{a_t} & X_{t,t+1} \\ X_{t+1,1} & X_{t+1,2} & \cdots & X_{t+1,t-1} & 0 & I_{a_t} \end{pmatrix}$$

By computing the *t*th column, we immediately see that $X_{t,i}^T B_1 + X_{t+1,i}^T B_2^T = 0$ for $i = 1, \ldots, t-1$ and $X_{t,t+1}^T B_1 = 0$. Then we can also compute the t + 1th column and see that

$$\begin{split} X_{t,i}^T B_2 + X_{t+1,i}^T B_3 &= 0 \text{ for } i = 1, \dots, t-1 \text{ and } B_2^T X_{t,t+1} + X_{t,t+1}^T B_2 = 0. \text{ The matrix is now} \\ \begin{pmatrix} A_{2a_1} & 0 & \cdots & 0 & 0 & 0 \\ X_{1,2}^T A_{2a_1} & A_{2a_2} & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ X_{1,t-1}^T A_{2a_1} & X_{2,t-1}^T A_{2a_2} & \cdots & A_{2a_{t-1}} & 0 & 0 \\ 0 & 0 & \cdots & 0 & B_1 & B_2 \\ 0 & 0 & \cdots & 0 & B_2^T & X_{t,t+1}^T B_2 + B_3 \end{pmatrix} \begin{pmatrix} I_{2a_1} & X_{1,2} & \cdots & X_{1,t-1} & 0 & 0 \\ 0 & I_{2a_2} & \cdots & X_{2,t-1} & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & I_{2a_{t-1}} & 0 & 0 \\ X_{t,1} & X_{t,2} & \cdots & X_{t,t-1} & I_{a_t} & X_{t,t+1} \\ X_{t+1,1} & X_{t+1,2} & \cdots & X_{t+1,t-1} & 0 & I_{a_t} \end{pmatrix} \end{split}$$

so it is now straightforward to check that $X_{i,j} = 0$ for $1 \le i < j \le t - 1$. Therefore, we get

$$P \leq \left\{ \begin{pmatrix} I_{2a_{1}} & & & \\ & I_{2a_{2}} & & & \\ & & \ddots & & \\ & & & I_{2a_{t-1}} & \\ & & & & I_{a_{t}} & X_{t,t+1} \\ & & & & & I_{a_{t}} & X_{t,t+1} \\ & & & & & I_{a_{t}} \end{pmatrix} \middle| \begin{array}{c} X_{t,t+1} \in M_{a_{t},a_{t}}(2), \\ X_{t,t+1}^{T} B_{1} = 0, \\ B_{2}^{T} X_{t,t+1} = (B_{2}^{T} X_{t,t+1})^{T} \\ \end{array} \right\}$$

so P is an elementary abelian 2-group of order at most $2^{a_t^2}$.

Proposition 4.3. $W/D \ge (2 - \sqrt{2})/4 > 0.146$. If W is even, then $W/D \ge 7(2 - \sqrt{2})/16 > 0.256$. In general, $W/D > ((2 - \sqrt{2})/2)(1 - 1/W_0)$, where W_0 is the 2'-part of W.

Proof. This is almost identical to the first part of the proof of [8, Theorem 7.4]. The only thing that has to be changed is that in the notation of that proof, we now have p = r = 2 so that $|Q| \ge 2$, and \overline{g} is now a 2-element of $\operatorname{Sp}_{2n}(2)$. In this setting, we still get the bound $W/D \ge (1 - 1/\sqrt{2})(1 - 1/2) = (2 - \sqrt{2})/4 > 0.146$. If W is even, then Q has an irreducible representation of even dimension, so it cannot be abelian. Therefore $|Q| \ge 8$ and we get $W/D \ge (1 - 1/\sqrt{2})(1 - 1/8) = 7(1 - 1/\sqrt{2})/8 = 7(2 - \sqrt{2})/16 > 0.256$. For the last statement, recall that Wild $|_Q$ is a sum of W_0 distinct nontrivial irreducible constituents. Therefore $|Q| \ge |\operatorname{Irr}(Q)| > W_0$.

Now we determine the local monodromy of \mathcal{H} at ∞ and the downstairs characters.

Theorem 4.4. Suppose that an irreducible hypergeometric sheaf \mathcal{H} has an extraspecial normalizer as its G_{geom} . Then g_{∞} is as in one of the cases described in Theorem 3.1, with the following additional conditions:

- (a) It is as in Theorem 3.1(a), with t = 1. \mathcal{H} is Kloosterman in this case.
- (b) It is as in Theorem 3.1(d), $t = 2, 2 \mid na_2$, and the downstairs characters are $Char(2^{a_2} + 1) \setminus \{1\}$.
- (c) It is as in Theorem 3.1(d), t = 1, and the downstairs character is $\{1\}$.
- (d) It is as in Theorem 3.1(b), n = 2, and the downstairs characters are $Char(3) \setminus \{1\}$.

Proof. (1) Suppose that g is as in Theorem 3.1(a). By Proposition 4.2(a) we know that $Q/(Q \cap E) = 1$, so $Q \leq E$. First suppose that \mathcal{H} is Kloosterman. Then dim Wild $= D = 2^n$, so the spectrum of g on Wild is $\eta(\mu_{2^n+1} \setminus \{1\})$ for some root of unity η of odd order. Since there is no Tame in this case, this is the entire spectrum of g, so t = 1.

Suppose now that \mathcal{H} is not Kloosterman. Since Q is nontrivial, we are in the situation of Proposition 4.1(c) with $Q = Q \cap E$. In particular, Q is abelian. Therefore, dim Tame = D - dim Wild is odd, and it divides $D = 2^n$, so dim Tame = 1 and dim Wild = $2^n - 1$. The spectrum of g on Wild is $\eta \mu_{2^n-1}$ for some root of unity η of odd order. This does not happen for g as in Theorem 3.1(a).

(2) Now consider the case (b) of Theorem 3.1. Here, the spectrum of g on V is

$$\left\{\xi\prod_{i=1}^{t-1}\zeta_i \mid \zeta_i \in \mu_{2^{a_i}+1} \setminus \{1\}\right\}$$

with every eigenvalue having multiplicity 2. Therefore, if an irreducible hypergeometric sheaf has such g as its $g_{\infty} \in G_{\text{geom}}$, then the spectrum of g_{∞} on Wild must be the above set, with every eigenvalue having multiplicity 1. Hence dim Wild $= D/2 = 2^{n-1}$. On the other hand, since $Q \cap E$ is an abelian normal subgroup of Q and $Q/(Q \cap E)$ is either trivial or has order 2 by Proposition 4.2(b), the irreducible characters of Q must have degree 1 or 2. Therefore, the 2-part of dim Wild is 1 or 2. This forces n = 1 or 2. The downstairs characters must be $\{1\}$ if n = 1 and $\text{Char}(3) \setminus \{1\}$ if n = 2. However, if n = 1, then the only hypergeometric sheaf which can have such g_{∞} together with g_0 as in Theorem 3.1(a) is $\mathcal{H}yp_{\psi}(\text{Char}(3) \setminus \{1\}; 1)$. The G_{geom} of this sheaf has an element of order 3 (namely g_0) and a nontrivial 2-subgroup. By the proof of [8, Theorem 9.3(i)], we get $G_{\text{geom}} = S_3$. Therefore $n \neq 1$.

(3) Next, consider the case Theorem 3.1(c). Let $i_{odd} < t$ be the unique index such that $a_{i_{odd}}$ is odd. Then the spectrum is given by

$$\{\xi\omega^{j}\prod_{i=1}^{t-1}\zeta_{i}^{r_{i}} \mid \zeta_{i} \in \mu_{2^{a_{i}}+1} \setminus \{1\}, j \in \{1,2\}\}$$

=2 $\{\xi\prod_{i=1}^{t-1}\zeta_{i} \mid \zeta_{i} \in \mu_{2^{a_{i}}+1} \setminus \{1\} \text{ for } i \neq i_{odd}, \zeta_{i_{odd}} \in \mu_{2^{a_{i}}_{odd}+1} \setminus \{\omega, \omega^{2}\}\}$
 $\cup \{\xi\prod_{i=1}^{t-1}\zeta_{i} \mid \zeta_{i} \in \mu_{2^{a_{i}}+1} \setminus \{1\} \text{ for } i \neq i_{odd}, \zeta_{i_{odd}} \in \{\omega, \omega^{2}\}\}.$

where ξ is a root of unity of odd order and ω is a primitive 3rd root of unity. Those with multiplicity 2 must appear exactly once in both Tame and Wild.

Suppose that $a_{i_{odd}} = 1$. By the proof of Proposition 4.2(b) with $B_1 = A_2 = B_3$ and

 $B_2 = 0$, we see that $Q/(Q \cap E) = 1$. The spectrum of g is

$$2\left\{\xi\prod_{i=1}^{t-1}\zeta_{i}\mid\zeta_{i}\in\mu_{2^{a_{i+1}}}\setminus\{1\}\text{ for }i\neq i_{odd},\zeta_{i_{odd}}=1\right\}$$
$$\cup\left\{\xi\prod_{i=1}^{t-1}\zeta_{i}\mid\zeta_{i}\in\mu_{2^{a_{i+1}}}\setminus\{1\}\text{ for }i\neq i_{odd},\zeta_{i_{odd}}\in\{\omega,\omega^{2}\}\right\}.$$

Since $Q \leq E$ and Q is nontrivial, we are in the situation of Proposition 4.1(c) with abelian Q. Hence, dim Tame is an odd number dividing D. Therefore dim Tame = 1. Since dim Tame is at least the number of eigenvalues of g with multiplicity 2, we must have t = 2 and $n = a_{i_{odd}} + a_2 = 2$. However, in this case, E is of type +, so g_0 must be as in Theorem 3.1(a) with even t, which is impossible.

Now suppose that $a_{i_{odd}} > 1$. Then from Proposition 4.2(a), we can see that $Q/(Q \cap E)$ is trivial, so that $1 \neq Q \leq E$. Also, since \mathcal{H} has nonzero tame part, it is not Kloosterman. Therefore, we are again in the situation of Proposition 4.1(c) with $Q = Q \cap E$. Therefore dim Tame = 1, but this is impossible since dim Tame must be at least the number of eigenvalues of g with multiplicity 2, and there are at least $|\mu_{2^{a_{i_{odd}+1}}} \setminus {\omega, \omega^{2}}| \geq 3$ of them.

(4) For Theorem 3.1(d), the eigenvalues of multiplicity 2 are

$$\{\xi\zeta_2\zeta_3\cdots\zeta_t \mid \zeta_i \in \mu_{2^{a_i+1}} \setminus \{1\}\}$$
 if $t \ge 2$, and $\{\xi\}$ if $t = 1$.

Therefore, the spectra on Tame and Wild both contain them, so \mathcal{H} is not Kloosterman. Also, Proposition 4.2(b) shows that $Q/(Q \cap E)$ is elementary abelian of order not exceeding $2^{a_1^2}$.

If we are in the situation of Proposition 4.1(b), then Q is elementary abelian and dim Wild is odd. Hence, the eigenvalues of g on Wild are the dim Wildth roots of $(\xi\zeta_2\zeta_3\cdots\zeta_t)^{\dim Wild}$ for $\zeta_i \in \mu_{2^{a_i}+1}$ if $t \geq 2$. Hence $\prod_{i=2}^t (2^{a_i}+1)$ divides dim Wild, but not all $\prod_{i=2}^t (2^{a_i}+1)$ th roots of $(\xi\zeta_2\zeta_3\cdots\zeta_t)^{\dim Wild}$ are in the spectrum of g. Therefore t = 1, $a_1 = n$ and dim Wild divides $2^n - 1$. By Proposition 4.3, $(2^n - 1)/\dim Wild \in \{1,3\}$.

Suppose that dim Wild = $(2^n - 1)/3$. Let χ be the character of \mathcal{H} as a representation of G, and $\chi_{\mathsf{Tame}} = (\dim \mathsf{Tame}) \mathbb{1}_P$ and χ_{Wild} be that of Tame and Wild as a representation of Q, so that $\chi|_Q = \chi_{\mathsf{Tame}} + \chi_{\mathsf{Wild}}$. Also let $x \in Q$ be any nonidentity element. Then x has order 2, so the eigenvalues of the action of x on \mathcal{H} are 1 or -1. Therefore,

$$\mathbb{Z} \ni \chi(x) = \chi_{\mathsf{Tame}}(x) + \chi_{\mathsf{Wild}}(x) \ge \dim \mathsf{Tame} - \dim \mathsf{Wild} = 1 + (2^n - 1)/3 > 2^{n-2}.$$

On the other hand, by [2, Lemma 2.4], $|\chi(x)|$ is a power of $\sqrt{2}$, and since χ is faithful, $\chi(x) < 2^n$. Therefore $\chi(x) = 2^{n-1}$ for all $x \in Q \setminus \{1\}$. But then

$$2^{n} - \frac{2^{n} - 1}{3} = \dim \mathsf{Tame} = (\chi|_{Q}, 1_{Q}) = \frac{1}{|Q|} \sum_{x \in Q} \chi(x) = \frac{2^{n} + (|Q| - 1)2^{n-1}}{|Q|} = 2^{n-1} + \frac{2^{n-1}}{|Q|} + \frac{2^{n-1}}{|Q|} + \frac{2^$$

so that

$$|Q| = \frac{2^{n-1}}{2^{n-1} - \frac{2^n - 1}{3}} = \frac{2^{n-1}}{\frac{2^{n-1} + 1}{3}}$$

This forces $(2^{n-1}+1)/3 = 1$, so n = 2, t = 2, and $a_1 = a_2 = 1$. Hence *E* must be of type -, and by Theorem 3.1(a), g_0 must have order 5. Therefore, the hypergeometric sheaf must be $\mathcal{H}yp_{\psi}(\mathsf{Char}(5) \setminus \{1\}; \mathsf{Char}(3))$ up to tensoring by a Kummer sheaf. However, this sheaf has $G_{\text{geom}} = \mathsf{S}_5$ as studied in the proof of [8, Theorem 9.3].

Suppose now that Proposition 4.1(c) holds, so that dim Tame divides $D = 2^n$. If dim Tame = 1, then since there must be at most one eigenvalue of g of multiplicity 2, we see that t = 1. The spectrum of g on Wild is $\xi \mu_{2^n-1}$.

Now assume dim Tame > 1. Then dim Tame = 2^m for some integer 0 < m < n, and dim Wild = $2^m(2^{n-m}-1)$. The spectrum of g on Wild is then $\eta \mu_{2^{n-m}-1}(\mu_{2^m+1} \setminus \{1\})$ for some root of unity η of odd order. Therefore, the $(2^{n-m}-1)(2^m+1)$ th powers of $\zeta_2\zeta_3\cdots\zeta_t$ are same for all $\zeta_i \in \mu_{2^{a_i}+1} \setminus \{1\}$, so $\prod_{i=2}^t (2^{a_i}+1)$ divides $(2^{n-m}-1)(2^m+1)$. Also, multiplication by a primitive $(2^{n-m}-1)$ th root does not change the spectrum of g on Wild. From the description of the spectrum of g, one immediately sees that if a root of unity which is not 1 has this property, then its order divides $2^{a_1}-1$. Therefore $2^{n-m}-1$ divides $2^{a_1}-1$, and $\prod_{i=2}^t (2^{a_i}+1)$ divides $2^m + 1$. On the other hand, if we take an eigenvalue of g on Wild and multiply by $(2^m + 1)$ th roots of unity, then all but one of the $2^m + 1$ numbers are also contained in the spectrum. But if $t \geq 3$, then strictly more than one multiple of an eigenvalue by $(2^{a_2} + 1)(2^{a_3} + 1)$ th roots of unity are missing from the spectrum of g. Therefore we must have $t \leq 2$.

If t = 2, then we must have $2^{a_2} + 1 = 2^m + 1$, since among the multiples of an eigenvalue by the $(2^m + 1)$ th roots of unity, only one must be missing, but $2^{a_1} + 1$ divides $2^m + 1$ and one from the multiples by $(2^{a_1} + 1)$ th roots of unity is already missing. Hence $m = a_2$, $a_1 = n - m$, dim Tame = 2^{a_2} , and dim Wild = $(2^{a_1} - 1)2^{a_2}$. Since $2^{n-m} - 1$ and $2^m + 1$ are relatively prime, we also see that $2 \mid nm$.

If t = 1, then the spectrum of g on Wild is

$$\{\eta\zeta_{1}\zeta_{2} \mid \zeta_{1} \in \mu_{2^{n-m}-1}, \zeta_{2} \in \mu_{2^{m}+1} \setminus \{1\}\}\$$

for some root of unity η of odd order, and this must be contained in $\xi\mu_{2^n-1}$. Therefore $(2^m + 1)(2^{n-m} - 1) = 2^n + 2^{n-m} - 2^m - 1$ divides $2^n - 1$, which forces n = 2m. Note that $(\chi|_Q, 1_Q) = \dim \mathsf{Tame} = 2^m$. Also recall that $Q \cap E$ is normalized by Q and g_{∞}, χ is faithful, and the nontrivial irreducible constituents of $\chi|_Q$ has $1_{Q\cap E}$ as a constituent when restricted to $Q \cap E$. Therefore $(\chi|_{Q\cap E}, 1_{Q\cap E}) = 2^m$. On the other hand, since $\chi|_E$ is the unique irreducible character of E of degree $2^n, \chi|_{Q\cap E}$ is a multiple of the regular character $\rho_{Q\cap E}$ of $Q \cap E$. Since $(\rho_{Q\cap E}, 1_{Q\cap E}) = 1$, we get $\chi|_{Q\cap E} = 2^m \rho_{Q\cap E}$. In particular, $|Q \cap E| = \rho_{Q\cap E}(1) = \chi|_{Q\cap E}(1)/2^m = 2^m$. Hence, the g-invariant subspace $(Q \cap E)\mathbf{Z}(E)/\mathbf{Z}(E)$ of $E/\mathbf{Z}(E)$ has dimension m, but the g-invariant subspaces of $E/\mathbf{Z}(E)$ have dimension 0, n or 2n. Therefore this case is impossible.

5 Determination of the finiteness of the monodromy

From Theorem 3.1(a), we know the possible upstairs characters of \mathcal{H} , and the possible downstairs characters were determined in Theorem 4.4. However, not every pair of these upstairs and downstairs characters have the desired G_{geom} ; in fact, G_{geom} is not even finite for some of them. In this section, we determine the pairs with finite G_{geom} .

By [5, 8.14.6], the finiteness of G_{geom} is equivalent to saying that the values of trace function after a Tate twist are algebraic integers, and it is enough to check the *p*-adic valuations. This can be done by using the so-called *V*-test of Kubert [6, Proposition 13.2] [5, Section 8.16].

The definition of the function V can be found in [6, Discussion after Proposition 13.1]; for an alternative definition, for a given prime p and integers $k > 0, 0 \le a < p^k - 1$ and z, we have

$$V\left(\frac{a}{p^k - 1} + z\right) = \frac{\text{sum of digits of } a \text{ in base } p}{k(p - 1)},$$

cf. [5, Section 8.16] [6, Theorem 13.4]. For the rest of this paper, we will use the following basic properties of the function V, often without mentioning. See [6, Discussion before Proposition 13.2] for the proofs.

- (1) V(x) = 0 if and only if $x \in \mathbb{Z}$.
- (2) V(x) + V(-x) = 1 for any $x \notin \mathbb{Z}$.

(3)
$$V(1/2) = V(1/2)$$

- (4) V(x) = V(px) for any x.
- (5) $V(x) + V(y) \ge V(x+y)$ for any x, y.
- (6) (Hasse-Davenport relation) $\sum_{i=1}^{r} V(x+i/r) = V(rx) + \frac{r-1}{2}$ for any x and $p \nmid r \in \mathbb{Z}$.

In practice, instead of the original inequality stated in [6, Proposition 13.2], the following simplified version is used. Although this version is already known to experts, it seems to be not written explicitly in literature to my knowledge, so here I include the statement and proof for the convenience of readers.

Lemma 5.1. Let χ be a multiplicative character of order $p^d - 1$, and let \mathcal{G} be the hypergeometric sheaf $\mathcal{H}yp_{\psi}(\chi^{a_1}, \ldots, \chi^{a_D}; \chi^{b_1}, \ldots, \chi^{b_M})$ where $a_1, \ldots, a_D, b_1, \ldots, b_M \in \{0, \ldots, p^d - 2\}$ are pairwise distinct. Then the geometric monodromy group G_{geom} of \mathcal{G} is finite if and only if for every $x \in \mathbb{Q}/\mathbb{Z}$ whose denominator is not divisible by p and for every $N \in \{1, \ldots, p^d - 2\}$ which is relatively prime to $p^d - 1$, we have the inequality

$$\sum_{i=1}^{D} V(Na_i + x) + \sum_{j=1}^{M} V(-Nb_j - x) \ge \frac{D + M - 1}{2}$$

Proof. The only difference between this and [6, Proposition 13.2] is that the term $(1/D) \sum_{i,j} V(Na_i - Nb_j)$ is replaced by M/2. Suppose that the inequality in this lemma holds for all x and N. Let N_0 be any possible value of N, and choose $(N, x) = (N_0, -N_0b_j)$ and $(-N_0, N_0b_j)$ and take the sum of the resulting inequalities. Since V(r) + V(-r) = 1 if $r \notin \mathbb{Z}$ and 0 otherwise, the sum of these two inequalities becomes $D + M - 1 \ge D + M - 1$. Therefore, the equality must hold for both pairs $(N, x) = (N_0, -N_0b_j)$ and $(-N_0, N_0b_j)$. Now take the sum of the equalities for the pairs $(N, x) = (N_0, -N_0b_1), (N_0, -N_0b_2), \dots, (N_0, -N_0b_M)$. Then we get

$$\sum_{i,j} V(N_0 a_i - N_0 b_j) + \frac{M(M-1)}{2} = \frac{M(D+M-1)}{2} = \frac{DM + M(M-1)}{2}$$

so that $(1/D) \sum_{i,j} V(N_0 a_i - N_0 b_j) = M/2.$

Conversely, suppose that the inequality in [6, Proposition 13.2] holds for all x and N. By an entirely analogous argument, we get

$$\sum_{i,j} V(Na_i - Nb_j) + \frac{M(M-1)}{2} = \frac{M(D-1)}{2} + \frac{M}{D} \sum_{i,j} V(Na_i - Nb_j)$$

$$(1/D) \sum_{i,j} V(Na_i - Nb_j) = (1/(D-M))(\frac{M(D-1)}{2} - \frac{M(M-1)}{2}) = M/2$$

so that $(1/D) \sum_{i,j} V(Na_i - Nb_j) = (1/(D - M))(\frac{m(D - D)}{2} - \frac{m(M - D)}{2}) = M/2.$

From now on, given a hypergeometric sheaf \mathcal{G} as in Lemma 5.1, we will call

$$V_{\mathcal{G},\mathrm{up}}(N,x) := \sum_{i=1}^{D} V(Na_i + x) - D/2$$

the "upstairs part" of the V-test, and

$$V_{\mathcal{G},\text{down}}(N,x) := \sum_{j=1}^{M} V(-Nb_j - x) - M/2$$

the "downstairs part" of the test. Thus the inequality for the V-test becomes

$$V_{\mathcal{G},\mathrm{up}}(N,x) + V_{\mathcal{G},\mathrm{down}}(N,x) + \frac{1}{2} \ge 0.$$

Now we return to the notations in the previous section, so \mathcal{H} is an irreducible hypergeometric sheaf of rank $D = 2^n$ in characteristic 2 whose G_{geom} is an extraspecial normalizer. The element g_0 is as described in Theorem 3.1(a); let $a_1 > \cdots > a_t$ be the positive integers in this description, so that $a_1 + \cdots + a_t = n$, and the numbers $A_i := 2^{a_i} + 1$ are pairwise coprime. The upstairs characters are the multiplicative characters of order dividing $A_1A_2 \cdots A_t$ but not dividing $A_1A_2 \cdots A_t/A_i$ for any i, tensored with a common multiplicative character. Therefore, the V-test of this \mathcal{H} has the upstairs part

$$V_{\mathcal{H},\mathrm{up}}(N,x) = \sum_{i_1=1}^{2^{a_1}} \sum_{i_2=1}^{2^{a_2}} \cdots \sum_{i_t=1}^{2^{a_t}} V\left(N\left(\frac{i_1}{A_1} + \dots + \frac{i_t}{A_t}\right) + x\right) - \frac{D}{2}$$

=
$$\sum_{i_1,\dots,i_{t-1}} \left(V\left(A_t N\left(\frac{i_1}{A_1} + \dots + \frac{i_{t-1}}{A_{t-1}}\right) + A_t x\right) + 2^{a_t - 1} - V\left(N\left(\frac{i_1}{A_1} + \dots + \frac{i_{t-1}}{A_{t-1}}\right) + x\right)\right) - \frac{D}{2}$$

=
$$\cdots$$

=
$$\sum_{i=0}^{t} \left((-1)^{t-i} \sum_{1 \le t_1 \le \dots \le t_i \le t} V(A_{t_1} A_{t_2} \cdots A_{t_i} x)\right)$$

where the equalities follow from the Hasse-Davenport relation stated above.

Let us write

$$V(2; r_1, \dots, r_t; x) := \sum_{i=0}^{t} \left((-1)^{t-i} \sum_{1 \le t_1 < \dots < t_i \le t} V((2^{r_{t_1}} + 1) \cdots (2^{r_{t_i}} + 1)x) \right)$$

so that $V_{\mathcal{H},up}(N,x) = V(2;a_1,\ldots,a_t;x)$. We now prove several inequalities involving this function.

Lemma 5.2. Let $r \in \mathbb{Z}_{>0}$. In characteristic 2, the following hold.

(a) $V((2^r + 1)x) < 2V(x) < V((2^r + 1)x) + 1.$ (b) If $V((2^r + 1)x) = 0$, then V(x) = 0 or 1/2.

 $1 \le t_1 \le \cdots \le t_i \le t$

Proof. (a) If $x \in \mathbb{Z}$, then $V((2^r+1)x) = V(x) = 0$. If $x \notin \mathbb{Z}$, then we have $V((2^r+1)x) \leq V(x) = 0$. $V(2^{r}x) + V(x) = 2V(x) = 2 - 2V(-x) \le 2 - V(-(2^{r}+1)x) = V((2^{r}+1)x) + 1.$

(b) Since $V((2^r+1)x) = 0$, we must have $(2^r+1)x \in \mathbb{Z}$. If $x \in \mathbb{Z}$, then V(x) = 0. So suppose $x \notin \mathbb{Z}$. Then $V(x) = 1 - V(-x) = 1 - V((2^r + 1)x - x) = 1 - V(2^r x) = 1 - V(x)$, so V(x) = 1/2.

Corollary 5.3. In characteristic 2, $V(2; r_1; x) = V((2^r+1)x) - V(x) \in [-1/2, 1/2)$ for every positive integer r. It is -1/2 if and only if $x = i/(2^r + 1)$ for some integer i not divisible by $2^r + 1.$

Proof. If $V((2^r + 1)x) = 0$ then $V(2; r_1; x) = 0$ or -1/2 by Lemma 5.2(b). So assume $0 < V((2^r + 1)x) < 1$. Then

$$-V(x) < V((2^{r} + 1)x) - V(x) < 1 - V(x).$$

On the other hand, from Lemma 5.2(a) we get

$$V(x) - 1 \le V((2^r + 1)x) - V(x) \le V(x).$$

Combining these two inequalities, we get $V(2; r_1; x) \in (-1/2, 1/2)$.

Corollary 5.4. (a) If g_{∞} is as in Theorem 4.4(a), then $V_{\mathcal{H},down}(N,x) = 0$.

- (b) If g_{∞} is as in Theorem 4.4(b), then $V_{\mathcal{H},down}(N,x) = V(-(2^{a_2}+1)x) V(-x) \in [-1/2, 1/2)$. Also, $V_{\mathcal{H},down}(N,x) = -1/2$ precisely when $x = i/(2^{a_2}+1)$ for some integer i not divisible by $2^{a_2}+1$.
- (c) If g_{∞} is as in Theorem 4.4(c), then $V_{\mathcal{H},down}(N,x) = V(-x) 1/2 \in [-1/2, 1/2).$

Proof. (a) and (c) are obvious. For (b), the downstairs part is $V(-(2^{a_2}+1)x) - V(-x) = V(2; a_2; -x)$, so we can apply Corollary 5.3.

Lemma 5.5. In characteristic 2, $V(2; r_1, r_2; x) = V((2^{r_1} + 1)(2^{r_2} + 1)x) - V((2^{r_1} + 1)x) - V((2^{r_2} + 1)x) + V(x) \in [-3/4, 3/4)$ for every positive integers r_1, r_2 such that $2^{r_1} + 1$ and $2^{r_2} + 1$ are coprime. It is -3/4 if and only if $(2^{r_1} + 1)(2^{r_2} + 1)x \in \mathbb{Z}$ and V(x) = 1/4.

Proof. Let $X := V(2; r_1, r_2; x) = V((2^{r_1}+1)(2^{r_2}+1)x) - V((2^{r_1}+1)x) - V((2^{r_2}+1)x) + V(x)$. First note that

$$V((2^{r_1} + 1)(2^{r_2} + 1)x)$$

$$\leq \frac{1}{2}(V(2^{r_1}(2^{r_2} + 1)x) + V((2^{r_2} + 1)x) + V(2^{r_2}(2^{r_1} + 1)x) + V((2^{r_1} + 1)x))$$

$$= V((2^{r_2} + 1)x) + V((2^{r_1} + 1)x)$$

so that $X \leq V(x)$.

If $(2^{r_1} + 1)(2^{r_2} + 1)x \in \mathbb{Z}$, then $V((2^{r_1} + 1)x), V((2^{r_2} + 1)x) \in \{0, 1/2\}$. If both of them are 0, then $x \in \mathbb{Z}$, so X = 0. If one of them is 0, then V(x) = 1/2 and X = 0. If both of them are 1/2, then

$$\frac{1}{4} = \frac{1}{2}V((2^{r_1} + 1)x) \le \frac{1}{2}(V(2^{r_1}x) + V(x)) = V(x) = 1 - V(-x)$$
$$= 1 - \frac{1}{2}(V(-x) + V(-2^{r_1}x)) \le 1 - \frac{1}{2}V(-(2^{r_1} + 1)x) = \frac{3}{4}$$

so $X \in [-3/4, -1/4]$.

Suppose that $(2^{r_1}+1)(2^{r_2}+1)x \notin \mathbb{Z}$. Then

$$V((2^{r_2}+1)x) + V((2^{r_1}+1)x) = 2 - V(-(2^{r_2}+1)x) - V(-(2^{r_1}+1)x)$$

$$\leq 2 - V(-(2^{r_1}+1)(2^{r_2}+1)x) = 1 + V((2^{r_1}+1)(2^{r_2}+1)x)$$

so that $X \ge V(x) - 1$.

Also,

$$2V(x) - 1 = 1 - V(-2^{r_1}x) - V(-x) \le 1 - V(-(2^{r_1} + 1)x)$$
$$= V((2^{r_1} + 1)x) \le V(2^{r_1}x) + V(x) = 2V(x)$$

and similarly $2V(x) - 1 \le V((2^{r_2} + 1)x) \le 2V(x)$, so -3V(x) < X < -3V(x) + 3. Combining these results, we can see that $X \in [-3/4, 3/4)$, and X = -3/4 happens only when $(2^{r_1} + 1)(2^{r_2} + 1)x \in \mathbb{Z}$ and V(x) = 1/4.

For a_1, \ldots, a_t such that $2^{a_j} + 1$ are pairwise coprime, we will write

$$x_{a_1,\dots,a_t}(i_1,\dots,i_t) := \sum_{j=1}^t \frac{i_j}{2^{a_j}+1}.$$

If the numbers a_1, \ldots, a_t are clear from the context, we will omit the subscript and simply write $x(i_1, \ldots, i_t)$.

Lemma 5.6. For any positive integers a_1, \ldots, a_t with $t \ge 2$ such that $(2^{a_j} + 1)$ are pairwise coprime, at least one of the following holds.

- (i) There exist integers i_1, \ldots, i_t such that $V(2; a_1, \ldots, a_t; x(i_1, \ldots, i_t)) \neq -1/2$ and for every $j, (2^{a_j} + 1) \nmid i_j$.
- (ii) There exist some $j_0 \in \{1, ..., t\}$ and integers $i_1, ..., i_t$ such that $(2^{a_{j_0}} + 1) \mid i_{j_0}$ and $V(2; a_1, ..., a_t; x(i_1, ..., i_t)) \notin [-1/2, 1/2].$

Proof. We use induction on t.

(1) First assume that t = 2. We show that (i) holds in this case by finding explicit integers i_1, i_2 . Let b_j be the 2-part of a_j . Then $2^{a_j} + 1 = (2^{b_j} + 1)(2^{a_j-b_j} - 2^{a_j-2b_j} + \cdots - 2^{b_j} + 1)$. Since $2^{a_j} + 1$ are pairwise coprime, we must have $b_1 \neq b_2$. We may assume that $b_1 > b_2$. Let

$$i_1 = \frac{2^{a_1} + 1}{2^{b_1} + 1} (2^{b_1/2} - 1)(2^{b_1/4} - 1) \cdots (2^{2b_2} - 1)(2^{b_2} - 1) \text{ and } i_2 = \frac{2^{a_2} + 1}{2^{b_2} + 1}.$$

Then $2^{a_j} + 1$ does not divide i_j for both j = 1, 2, and

$$\begin{split} V\left(\frac{i_1}{2^{a_1}+1} + \frac{i_2}{2^{a_2}+1}\right) &= V\left(\frac{(2^{b_1/2}-1)(2^{b_1/4}-1)\cdots(2^{2b_2}-1)(2^{b_2}-1)}{2^{b_1}+1} + \frac{1}{2^{b_2}+1}\right) \\ &= V\left(\frac{(2^{b_1}-1)(2^{b_1/2}-1)\cdots(2^{b_2}-1) + (2^{b_1}+1)(2^{b_1/2}+1)\cdots(2^{2b_2}+1)(2^{b_2}-1)}{2^{2b_1}-1}\right) \\ &= V\left(\frac{(2^{b_2}-1)B}{2^{2b_1}-1}\right) \end{split}$$

where

$$B := (2^{b_1} - 1)(2^{b_1/2} - 1) \cdots (2^{2b_2} - 1) + (2^{b_1} + 1)(2^{b_1/2} + 1) \cdots (2^{2b_2} + 1).$$

This number is just the sum of all numbers of the form $2^{1+(2^{k_1}+\cdots+2^{k_r})b_2}$ for some $1 \leq k_r < \cdots < k_1 \leq \log_2(b_1/b_2)$ with $r \equiv \log_2(b_1/b_2) \mod 2$. These numbers are distinct for different choice of numbers k_1, \ldots, k_r , and the ratio of two of them is at least 2^{b_2} . Therefore, if we write B in base 2, the digits 1 are always followed by $b_2 - 1$ digits 0; in other words, there are at least $b_2 - 1$ 0's between any two 1's. Thus multiplying B with $2^{b_2} - 1$, which is just b_2 consecutive 1's when written in base 2, is the same as replacing each occurrence of b_2 consecutive base 2 digits $0 \ldots 01$ in the base 2 representation of B by $1 \ldots 1$. Since there are

exactly $2^{\log_2(b_1/b_2)-1} = (b_1/2b_2)$ distinct choices of k_1, \ldots, k_r , there are the same number of b_2 consecutive digits $0 \ldots 01$. Therefore $(2^{b_2} - 1)B$ has exactly $b_1/2$ nonzero base 2 digits, and since $(2^{b_2} - 1)B < 2^{2b_1} - 1$, we have

$$V\left(\frac{(2^{b_2}-1)B}{2^{2b_{t-1}}-1}\right) = \frac{b_1/2}{2b_1} = \frac{1}{4}, \text{ so that } V(2;a_1,a_2;x(i_1,i_2))$$

Therefore, (i) holds when t = 2.

(2) Now assume that the statement holds for $t = t_0 - 1$ so that at least one of (i) and (ii) holds, and consider the case $t = t_0$. Assume that $a_1 > a_2 > \cdots > a_{t_0}$, and suppose that both (i) and (ii) are not true. Then for any i_2, \ldots, i_{t_0} such that $(2^{a_j} + 1) \nmid i_j$ for every $j \in \{2, \ldots, t_0\}$, we have

$$2^{a_1}V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, \dots, i_{t_0})) = (2^{a_1} + 1)V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, \dots, i_{t_0})) - V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, \dots, i_{t_0})) = \sum_{i=0}^{2^{a_1}} V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(i, i_2, \dots, i_{t_0})) - \sum_{i=0}^{2^{a_1}} V(2; a_2, \dots, a_{t_0}; x(i, i_2, \dots, i_{t_0})) = \sum_{i=0}^{2^{a_1}} V(2; a_1, \dots, a_{t_0}; x(i, i_2, \dots, i_{t_0})) = (-2^{a_1-1} - \frac{1}{2}, -2^{a_1-1} + \frac{1}{2}].$$

Here, the second equality follows from the Hasse-Davenport relation stated in the beginning of this section, and the last equality and containment follow from the assumption that both (i) and (ii) are not true.

On the other hand, $2^{2 \operatorname{lcm}(a_2,\ldots,a_{t_0})} - 1$ is a common denominator of $\{i_j/(2^{a_j}+1) \mid j = 2,\ldots,t_0\}$ and $x(0,i_2,\ldots,i_{t_0})$ is the sum of them, so by the definition of the function V, we have

$$V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, \dots, i_{t_0})) \in \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_{t_0})}\right) \mathbb{Z}$$

Note that since $2^{a_1} + 1, \ldots, 2^{a_{t_0}} + 1$ are pairwise coprime, as we saw above, the 2-parts of a_j are distinct, so in particular $a_1 \ge 2^{t_0-1}$. Hence if $a_1 \ge 16$, then we have

$$2^{a_1} \ge 2^{(\log_2 a_1)^2} \ge 2^{(t_0 - 1)\log_2 a_1} = (a_1)^{t_0 - 1} > \prod_{j=2}^{t_0} a_{t_0} \ge \operatorname{lcm}(a_2, \dots, a_{t_0}).$$

One can also easily check that $2^{a_1} > \text{lcm}(a_2, \ldots, a_{t_0})$ when $a_1 < 16$. Therefore,

$$V(2; a_2, \dots, a_{t_0}; (2^{a_1}+1)x(0, i_2, \dots, i_{t_0})) \in \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_{t_0})}\right) \mathbb{Z} \cap \left[-\frac{1}{2} - \frac{1}{2^{a_1+1}}, -\frac{1}{2} + \frac{1}{2^{a_1+1}}\right] = \left\{\frac{1}{2}\right\}$$

Therefore, a_2, \ldots, a_{t_0} does not satisfy (i).

For any $j_0 \in \{2, \ldots, t_0\}$ and i_2, \ldots, i_{t_0} such that i_j is not divisible by $2^{a_j} + 1$ if and only if $j_0 \neq j$, we have

$$(2^{a_1} + 1)V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, i_3, \dots, i_{t_0}))$$

$$= \sum_{i=0}^{2^{a_1}} V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(i, i_2, i_3, \dots, i_{t_0}))$$

$$= \sum_{i=0}^{2^{a_1}} (V(2; a_1, \dots, a_{t_0}; x(i, i_2, i_3 \dots, i_{t_0})) + V(2; a_2, \dots, a_{t_0}; x(i, i_2, i_3, \dots, i_{t_0})))$$

$$= \sum_{i=0}^{2^{a_1}} V(2; a_1, \dots, a_{t_0}; x(i, i_2, i_3 \dots, i_{t_0})) + V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, i_3, \dots, i_{t_0})).$$

As before, we get

$$V(2; a_2, \dots, a_{t_0}; (2^{a_1} + 1)x(0, i_2, i_3, \dots, i_{t_0})) \in \left[-\frac{2^{a_1} + 1}{2^{a_1 + 1}}, \frac{2^{a_1} + 1}{2^{a_1 + 1}}\right] \cap \left(\frac{1}{2\operatorname{lcm}(a_3, \dots, a_{t_0})}\right) \mathbb{Z}$$
$$= \left[-\frac{1}{2}, \frac{1}{2}\right].$$

Therefore (ii) also fails for a_2, \ldots, a_{t_0} . This contradicts the induction hypothesis that at least one of (i) and (ii) holds for $t = t_0 - 1$. Therefore, at least one of (i) and (ii) must also hold for a_1, \ldots, a_{t_0} .

Now we are ready to prove the main result of this paper.

Theorem 5.7. The irreducible hypergeometric sheaves in characteristic 2 of type (D, M) with D > M whose geometric monodromy group is an extraspecial normalizer are the following, up to tensoring with a Kummer sheaf.

- (a) $\mathcal{H}yp_{\psi}(\mathsf{Char}(2^{a}+1)\setminus\{1\};\emptyset)$ for some positive integer a.
- (b) $\mathcal{H}yp_{\psi}(\mathsf{Char}(2^{a}+1) \setminus \{\mathbb{1}\}; \mathsf{Char}(2^{b}+1) \setminus \{\mathbb{1}\})$ for some positive integers a > b with different 2-parts.
- (c) $\mathcal{H}yp_{\psi}((\mathsf{Char}(2^{a}+1) \setminus \{\mathbb{1}\}) \times (\mathsf{Char}(2^{b}+1) \setminus \{\mathbb{1}\}); \mathbb{1})$ for some positive integers a and b with different 2-parts, and $a+b \geq 5$.

Proof. Let g_0 be as in Theorem 3.1(a), and let t and $a_1 > \cdots > a_t$ be as defined there. First suppose that $t \ge 3$.

(1) If g_{∞} is as in Theorem 4.4(a), then the V-test inequality is

$$V(2; a_1, \dots, a_t; x) + \frac{1}{2} \ge 0.$$
 (5.7.1)

Suppose that this inequality holds for all x. As before, let $x(i_1, \ldots, i_t) := \frac{i_1}{2^{a_1}+1} + \cdots + \frac{i_t}{2^{a_t}+1}$ for integers i_j . From the properties (1) and (2) of the function V stated before Lemma 5.1, we see that

$$V(2; a_1, \dots, a_t; x(i_1, \dots, i_t)) + V(2; a_1, \dots, a_t; -x(i_1, \dots, i_t)) = -1 \text{ if } \forall j, \ 2^{a_j} + 1 \nmid i_j. \ (5.7.2)$$

As we assumed that (5.7.1) holds for all x, it follows that

$$V(2; a_1, \dots, a_t; x(i_1, \dots, i_t)) = -\frac{1}{2}$$
 if no i_j is divisible by $2^{a_j} + 1$.

If i_j is divisible by $2^{a_j} + 1$ for some j, then we instead get

$$V(2; a_1, \dots, a_t; x(i_1, i_2, \dots, i_t)) + V(2; a_1, \dots, a_t; x(-i_1, -i_2, \dots, -i_t)) = 0,$$

so from (5.7.1) we get

$$V(2; a_1, \dots, a_t; x(i_1, i_2, \dots, i_t)) \in \left[-\frac{1}{2}, \frac{1}{2}\right].$$
(5.7.3)

However, since we are assuming $t \ge 3$, by Lemma 5.6 at least one of (5.7.2) and (5.7.3) must fail. Therefore (5.7.1) cannot hold for all x.

(2) Suppose that g_{∞} is as in Theorem 4.4(b). Write dim Tame = 2^{b} for some integer $1 \leq b < n$. The V-test inequality is

$$V(2; a_1, \dots, a_t; x) + V(2; b; -x) + \frac{1}{2} \ge 0.$$
(5.7.4)

By taking the sum of (5.7.4) for $x = x(i_1, \ldots, i_t)$ and $x(-i_1, \ldots, -i_t)$ defined as before, we again see that

$$V(2; a_1, \dots, a_t; x(i_1, \dots, i_t)) + V(2; b; -x(i_1, \dots, i_t)) \begin{cases} = -\frac{1}{2} \text{ if } (2^{a_j} + 1) \nmid i_j \forall j, \\ \in [-1/2, 1/2] \text{ otherwise} \end{cases}$$

and that $(2^b + 1)x(i_1, \ldots, i_t) \notin \mathbb{Z}$ if $(2^{a_j} + 1) \nmid i_j$ for all j. Hence there exists at least one j such that $2^{a_j} + 1$ does not divide $2^b + 1$. If $(2^{a_j} + 1) \nmid i_j$ for all j, then the sum of the left-hand side of the above equalities for $i_1 = 0, \ldots, 2^{a_1}$ can be written as

$$\sum_{i=0}^{2^{a_1}} \left(V(2; a_1, \dots, a_t; x(i, i_2, \dots, i_t)) + V(2; b; -x(i, i_2, \dots, i_t)) \right)$$

=2^{a_1}V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) + V(2; b; -(2^{a_1} + 1)x(0, i_2, \dots, i_t)).

By Lemma 5.5 applied to $V(2; b, a_1; -x(i, i_2, ..., i_t)) = V(2; b; -(2^{a_1} + 1)x(0, i_2, ..., i_t)) - V(2; b; -x(0, i_2, ..., i_t)),$

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t))$$

= $-\frac{1}{2} + 2^{-a_1}V(2; a_1, \dots, a_t; x(0, i_2, \dots, i_t)) - 2^{-a_1}V(2; b, a_1; -x(0, i_2, \dots, i_t))$
 $\in \left[-\frac{1}{2} - \frac{1}{2^{a_1+1}} - \frac{3}{2^{a_1+2}}, -\frac{1}{2} + \frac{1}{2^{a_1+1}} + \frac{3}{2^{a_1+2}}\right] = \left[-\frac{1}{2} - \frac{5}{2^{a_1+2}}, -\frac{1}{2} + \frac{5}{2^{a_1+2}}\right].$

By a similar argument as in the proof of Lemma 5.6, we see that $\operatorname{lcm}(a_2, \ldots, a_t) < 2^{(\log_2 a_1)^2} < 2^{a_1+2-\log_2 5}$ for $a_1 \geq 18$, and one can check that for $a_1 < 18$ we also have $\operatorname{lcm}(a_2, \ldots, a_t) < 2^{a_1+2}/5$. Therefore we have

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) \\ \in \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_t)}\right) \mathbb{Z} \cap \left[-\frac{1}{2} - \frac{5}{2^{a_1+2}}, -\frac{1}{2} + \frac{5}{2^{a_1+2}}\right] = \left\{-\frac{1}{2}\right\}.$$
(5.7.5)

Also, if $(2^{a_j} + 1) \mid i_j$ for some $j \in \{2, \ldots, t\}$, then as before we have

$$2^{a_1}V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, i_3, \dots, i_t)) + V(2; b; -(2^{a_1} + 1)x(0, i_2, i_3, \dots, i_t))$$

$$= \sum_{i=0}^{2^{a_1}} \left(V(2; a_1, \dots, a_t; x(i, i_2, i_3, \dots, i_t)) + V(2; b; -x(i, i_2, i_3, \dots, i_t)) \right)$$

$$\in \left[-\frac{2^{a_1} + 1}{2}, \frac{2^{a_1} + 1}{2} \right]$$

so by Lemma 5.5 applied to $V(2; b, a_1; -x(i, i_2, ..., i_t)) = V(2; b; -(2^{a_1} + 1)x(0, i_2, ..., i_t)) - V(2; b; -x(0, i_2, ..., i_t))$, we get

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, i_3, \dots, i_t)) \\\in \left[-\frac{1}{2} - \frac{5}{2^{a_1+2}}, \frac{1}{2} + \frac{5}{2^{a_1+2}}\right] \cap \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_t)}\right) \mathbb{Z} = \left[-\frac{1}{2}, \frac{1}{2}\right].$$
(5.7.6)

As before, by Lemma 5.6, one of (5.7.5) and (5.7.6) must fail, so (5.7.4) fails for some x.

(3) Finally suppose that g_{∞} is as in Theorem 4.4(c). In this case E is of type +, so t is even. The V-test inequality is

$$V(2; a_1, \dots, a_t; x) + V(-x) \ge 0.$$
(5.7.7)

We again get

$$V(2; a_1, \dots, a_t; x(i_1, \dots, i_t)) + V(-x(i_1, \dots, i_t)) - \frac{1}{2} \begin{cases} = -\frac{1}{2} \text{ if } (2^{a_j} + 1) \nmid i_j \forall j, \\ \in [-1/2, 1/2] \text{ otherwise.} \end{cases}$$

If $(2^{a_j}+1) \nmid i_j$ for $j = 2, \ldots, t$, then the sum of the left-hand side of the above inequalities for $i_1 = 0, \ldots, 2^{a_1}$ is

$$2^{a_1}V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) + V(-(2^{a_1} + 1)x(0, i_2, \dots, i_t)) - \frac{1}{2}$$

= $\sum_{i=0}^{2^{a_1}} \left(V(2; a_1, \dots, a_t; x(i, \dots, i_t)) + V(-x(i, \dots, i_t)) - \frac{1}{2} \right) \in \left[-2^{a_1-1} - \frac{1}{2}, -2^{a_1-1} + \frac{1}{2} \right].$

Therefore we get

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) \in \left(-\frac{1}{2} - \frac{1}{2^{a_1}}, -\frac{1}{2} + \frac{1}{2^{a_1}}\right] \cap \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_t)}\right) \mathbb{Z}.$$

If $a_1 \ge 20$, then $lcm(a_2, \ldots, a_t) < 2^{(\log_2 a_1)^2} < 2^{a_1-1}$, and one can check that we still have $lcm(a_2, \ldots, a_t) < 2^{a_1-1}$ for $a_1 < 20$, unless t = 4 and $(a_1, a_2, a_3, a_4) = (8, 7, 6, 4)$. Therefore, assuming $(a_1, a_2, a_3, a_4) \ne (8, 7, 6, 4)$, we must have

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) = -1/2.$$
(5.7.8)

For i_2, \ldots, i_t with $(2^{a_j} + 1) \mid i_j$ for some j, we have

$$2^{a_1}V(2;a_2,\ldots,a_t;(2^{a_1}+1)x(0,i_2,\ldots,i_t)) + V(-(2^{a_1}+1)x(0,i_2,\ldots,i_t)) - \frac{1}{2}$$
$$= \sum_{i=0}^{2^{a_1}} \left(V(2;a_1,\ldots,a_t;x(i,\ldots,i_t)) + V(-x(i,\ldots,i_t)) - \frac{1}{2} \right) \in \left[-2^{a_1-1} - \frac{1}{2}, 2^{a_1-1} + \frac{1}{2} \right].$$

so if $(a_1, a_2, a_3, a_4) \neq (8, 7, 6, 4)$, then

$$V(2; a_2, \dots, a_t; (2^{a_1} + 1)x(0, i_2, \dots, i_t)) \\ \in \left(-\frac{1}{2} - \frac{1}{2^{a_1}}, \frac{1}{2} + \frac{1}{2^{a_1}}\right] \cap \left(\frac{1}{2\operatorname{lcm}(a_2, \dots, a_t)}\right) \mathbb{Z} = \left[-\frac{1}{2}, \frac{1}{2}\right].$$
(5.7.9)

By Lemma 5.6, not both (5.7.8) and (5.7.9) can be true at the same time. Therefore (5.7.7) fails unless t = 4 and $(a_1, a_2, a_3, a_4) = (8, 7, 6, 4)$. For this case, one can quickly find values of x for which (5.7.7) fails using computers.

(4) We proved that t < 3. For t = 1, E is of type -, so g_{∞} must be as in Theorem 4.4(a) (the Pink-Sawin Kloosterman sheaf [9, Theorem 7.3.8]) or Theorem 4.4(b) (studied in [9, Theorem 8.5.5] when $a_1 \ge 4$ and in [11, Theorem 4.4] when $a_1 < 4$).

For t = 2, E is of type +, so g_{∞} must be as in Theorem 4.4(c) (studied in [9, Proposition 9.1.9] when $a + b \ge 4$) or Theorem 4.4(d). For g_{∞} as in Theorem 4.4(c), the condition $a_1 + a_2 \ge 4$ is necessary here, because the case $a_1 + a_2 = 3$ has $G_{\text{geom}} = 2A_8$ as shown in [9, Theorem 9.1.8]. For Theorem 4.4(d), we must have $a_1 = a_2 = 1$, which is impossible.

Acknowledgment

I would like to thank Pham Huu Tiep for suggesting this problem and giving helpful advice.

References

- [1] S. S. Abhyankar, Coverings of algebraic curves, Amer. J. Math. 79 (1957), 825–856. MR0094354
- [2] R. M. Guralnick and P. H. Tiep, Cross characteristic representations of even characteristic symplectic groups, Trans. Amer. Math. Soc. 356 (2004), no. 12, 4969–5023. MR2084408

- R. M. Guralnick and P. H. Tiep, Symmetric powers and a problem of Kollár and Larsen, Invent. Math. 174 (2008), no. 3, 505–554. MR2453600
- [4] D. Harbater, Abhyankar's conjecture on Galois groups over curves, Invent. Math. 117 (1994), no. 1, 1–25. MR1269423
- [5] N. M. Katz, Exponential sums and differential equations, Annals of Mathematics Studies, 124, Princeton University Press, Princeton, NJ, 1990. MR1081536
- [6] N. M. Katz, G₂ and hypergeometric sheaves, Finite Fields Appl. 13 (2007), no. 2, 175–223. MR2307123
- [7] N. M. Katz, A. Rojas-León and P. H. Tiep, *Rigid local systems and Sporadic Simple groups*, Mem. Amer. Math. Soc. (to appear).
- [8] N. M. Katz and P. H. Tiep, Monodromy groups of Kloosterman and hypergeometric sheaves, Geom. Funct. Anal. 31 (2021), no. 3, 562–662. MR4311580
- [9] N. M. Katz and P. H. Tiep, *Exponential sums, hypergeometric sheaves, and monodromy groups,* Ann. of Math. Stud. (to appear).
- [10] N. M. Katz and P. H. Tiep, Condition (S+) in ranks 4, 8, and 9, J. Pure Appl. Algebra 228 (2024), no. 9, Paper No. 107679, 13 pp. MR4728133
- [11] N. M. Katz and P. H. Tiep, *Moments, exponential sums, and monodromy groups,* preprint.