

THE REVISED BOOMERANG CONNECTIVITY TABLES AND THEIR CONNECTION TO THE DIFFERENCE DISTRIBUTION TABLE

KIRPA GARG, SARTAJ UL HASAN, CONSTANZA RIERA, AND PANTELIMON STĂNICĂ*

ABSTRACT. It is well-known that functions over finite fields play a crucial role in designing substitution boxes (S-boxes) in modern block ciphers. In order to analyze the security of an S-box, recently, three new tables have been introduced: the Extended Boomerang Connectivity Table (EBCT), the Lower Boomerang Connectivity Table (LBCT), and the Upper Boomerang Connectivity Table (UBCT). In fact, these tables offer improved methods over the usual Boomerang Connectivity Table (BCT) for analyzing the security of S-boxes against boomerang-style attacks. Here, we put in context these new EBCT, LBCT, and UBCT concepts by connecting them to the DDT for a differentially δ -uniform function and also determine the EBCT, LBCT, and UBCT entries of three classes of differentially 4-uniform power permutations, namely, Gold, Kasami and Bracken-Leander. We also determine the Double Boomerang Connectivity Table (DBCT) entries of the Gold function. As byproducts of our approach, we obtain some previously published results quite easily.

1. INTRODUCTION

In symmetric cryptography, vectorial Boolean functions over a finite field play a significant role, particularly in designing block ciphers' S-boxes. There are various kinds of cryptanalytic attacks possible on these S-boxes. One well-known attack on S-boxes is the differential attack, introduced by Biham and Shamir [1]. This attack exploits the non-uniformity in the distribution of output differences corresponding to a given input difference. To quantify the degree of resistance of S-Boxes against differential attacks, Nyberg [15] introduced the notion of Difference Distribution Table (DDT) and differential uniformity.

Let p be a prime number and n be a positive integer. We denote by \mathbb{F}_q the finite field with $q = p^n$ elements, by \mathbb{F}_q^* the multiplicative cyclic group of non-zero elements of \mathbb{F}_q and by $\mathbb{F}_q[X]$ the ring of polynomials in one variable X with coefficients in \mathbb{F}_q . For a function $F : \mathbb{F}_q \rightarrow \mathbb{F}_q$, and for any $a \in \mathbb{F}_q$, the derivative of F in the direction a is defined as $D_F(X, a) := F(X + a) - F(X)$ for all $X \in \mathbb{F}_q$. For any $a, b \in \mathbb{F}_q$, the Difference Distribution Table (DDT) entry $DDT_F(a, b)$ at point (a, b) is the number of solutions $X \in \mathbb{F}_q$ of the equation $D_F(X, a) = b$. Further, the differential uniformity of F , denoted by Δ_F , is given by $\Delta_F := \max \{DDT_F(a, b) \mid a \in \mathbb{F}_q^*, b \in \mathbb{F}_q\}$. We call a function F a perfect nonlinear (PN) function and an almost perfect nonlinear (APN) function if $\Delta_F = 1$ and $\Delta_F = 2$, respectively. Functions with low differential uniformity are more resistant to differential attacks, making them desirable for cryptographic applications.

2020 *Mathematics Subject Classification.* 12E20, 11T06, 94A60.

Key words and phrases. Finite fields, permutation polynomials, almost perfect non-linear functions, Upper Boomerang Connectivity Table, Lower Boomerang Connectivity Table, Extended Boomerang Connectivity Table, Double Boomerang Connectivity Table.

The work of K. Garg is supported by the University Grants Commission (UGC), Government of India. The work of S. U. Hasan is partially supported by Core Research Grant CRG/2022/005418 from the Science and Engineering Research Board, Government of India. The work of P. Stănică (corresponding author) is partially supported by a grant from the NPS Foundation.

In 1999, Wagner [16] introduced a new attack on block ciphers, called the boomerang attack, which can be seen as an extension of the differential attack. It allows the cryptanalyst to use two unrelated differential characteristics to attack the same cipher by using one differential to defeat the first half and another to defeat the second half of the cipher. Further, Cid et al. [7] introduced the notion of the Boomerang Connectivity Table (BCT), which can be used to more accurately evaluate the probability of generating a right quartet in boomerang-style attacks such that the boomerang incompatibility, ladder switch and S-box switch can easily be detected. For effectively computing the entries in the BCT, Li et al. [13] proposed an equivalent formulation as described below, which avoids computing the inverse of the S-box, and consequently, it can be defined even for non-permutation. For any $a, b \in \mathbb{F}_{2^n}$, the BCT entry at $(a, b) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$, denoted as $\mathcal{B}_F(a, b)$, is the number of solutions in $\mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$ of the following system

$$\begin{cases} F(X) + F(Y) = b \\ F(X + a) + F(Y + a) = b. \end{cases}$$

The boomerang uniformity of F is defined as $\mathcal{B}_F := \max \{ \mathcal{B}_F(a, b) \mid a, b \in \mathbb{F}_{2^n}^* \}$. However, it turns out that the BCT has limitations when evaluating boomerang switches in multiple rounds. This was observed by Wang and Peyrin [17], who showed that the BCT is not applicable effectively in some cases involving multiple rounds by giving an incompatibility example on two rounds of AES. To address this challenge, they introduced a new tool known as the Boomerang Difference Table (BDT) along with its variant BDT'. Later, in 2020, Delaune et al. [8] renamed these tables as the Upper Boomerang Connectivity Table (UBCT) and the Lower Boomerang Connectivity Table (LBCT), respectively, to highlight the fact that UBCT and LBCT emphasize the upper and lower characteristic, respectively. The Feistel Boomerang Extended Table (FBET), or Boomerang Extended Table (BET) for SPN ciphers, proposed by Boukerrou et al. [3], has been renamed as the Extended Boomerang Connectivity Table (EBCT) by the authors in [8]. The entries in this table count the number of values such that the boomerang will return on a single S-box, with all the differences fixed.

Further, to study the behavior of two consecutive S-boxes in the boomerang attack, Hadipour et al. [11] introduced the notion of Double Boomerang Connectivity Table (DBCT). It may be noted that Yang et al. [19] illustrated the effect of an S-box on the probability of the 7-round boomerang distinguisher by comparing the DDT, BCT and DBCT entries of different S-boxes. Their findings highlight the importance of investigating the DBCT, in addition to the DDT and BCT, for evaluating an S-box's resistance to a boomerang attack. However, determining the DBCT entries for an S-box (or equivalently, for a vectorial Boolean function) is quite challenging, and rather limited research has been done in this direction.

Recently, Eddahmani and Mesnager [10] studied various properties of EBCT, LBCT and UBCT. They also determined these entries for the inverse function. In a separate work, Man et al. [14] also gave the DBCT, LBCT and UBCT entries for the inverse function. In this work, we discuss the DBCT entries of the Gold function, and we provide a connection between these new EBCT, LBCT and UBCT concepts and the DDT entries for a differentially δ -uniform function. As a consequence, it turns out that our result covers a particular case of a result of Eddahmani et al. [10] and Man et al. [14] for the inverse function over \mathbb{F}_{2^n} . Moreover, we explicitly compute EBCT, LBCT and UBCT entries of three classes of differentially-4 uniform power permutations. We shall now give the structure of the paper. We first recall some definitions and results in Section 2. In Section 3, we provide some general results on EBCT. We also give a generalization for the definitions of EBCT, LBCT and UBCT for any function over \mathbb{F}_{2^n} . In Section 4, we find a connection between the EBCT, LBCT and UBCT entries of a differentially δ -uniform function and also express them in terms of the DDT

entries. Section 5 and Section 6 deal with the EBCT, LBCT and UBCT entries of APN functions, as well as some other well-known functions. We also computed the DBCT entries for the Gold function in Section 7. In Section 8, we discuss the invariance of EBCT, LBCT and UBCT under extended affine equivalence. Finally, we conclude the paper in Section 9.

2. PRELIMINARIES

We will first recall several definitions and lemmas used in the subsequent sections.

Definition 2.1. [8] Let F be a permutation of \mathbb{F}_{2^n} . The Extended Boomerang Connectivity Table (EBCT) of F is a $2^n \times 2^n \times 2^n \times 2^n$ table where the entry at $a, b, c, d \in \mathbb{F}_{2^n}$ is

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \left| \begin{cases} F(X) + F(X + a) = b \\ F(X) + F(X + c) = d \\ F^{-1}(F(X) + d) + F^{-1}(F(X + a) + d) = a \end{cases} \right. \right\} \right|.$$

Definition 2.2. [17] Let F be a permutation of \mathbb{F}_{2^n} . The Lower Boomerang Connectivity Table (LBCT) of F is defined as a $2^n \times 2^n \times 2^n$ table where the entry at $a, b, c \in \mathbb{F}_{2^n}$ is

$$\text{LBCT}_F(a, b, c) = \left| \left\{ X \in \mathbb{F}_{2^n} \left| \begin{cases} F(X) + F(X + b) = c \\ F^{-1}(F(X) + c) + F^{-1}(F(X + a) + c) = a \end{cases} \right. \right\} \right|.$$

Definition 2.3. [17] Let F be a permutation of \mathbb{F}_{2^n} . The Upper Boomerang Connectivity Table (UBCT) of F is defined as a $2^n \times 2^n \times 2^n$ table where the entry at $a, b, c \in \mathbb{F}_{2^n}$ is

$$\text{UBCT}_F(a, b, c) = \left| \left\{ X \in \mathbb{F}_{2^n} \left| \begin{cases} F(X) + F(X + a) = b \\ F^{-1}(F(X) + c) + F^{-1}(F(X + a) + c) = a \end{cases} \right. \right\} \right|.$$

Recently, Eddahmani and Mesnager [10] and, Man et al. [14] redefined the notions of LBCT and UBCT for a permutation F without involving its compositional inverse, as stated in the following lemmas.

Lemma 2.4. [10] *Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c \in \mathbb{F}_{2^n}$, we have*

$$\text{LBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} X + Y = b \\ F(X + a) + F(Y + a) = c \\ F(X) + F(Y) = c \end{cases} \right. \right\} \right|.$$

Lemma 2.5. [10] *Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c \in \mathbb{F}_{2^n}$, we have*

$$\text{UBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} F(X + a) + F(Y + a) = c \\ F(X) + F(Y) = c \\ F(X) + F(X + a) = b \end{cases} \right. \right\} \right|.$$

Definition 2.6. (Feistel Boomerang Connectivity Table) [3] Let $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ and $a, b \in \mathbb{F}_{2^n}$. The *Feistel Boomerang Connectivity Table* (FBCT) of F is given by a $2^n \times 2^n$ table, of entry at $(a, b) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$ defined by

$$\text{FBCT}_F(a, b) = \left| \left\{ X \in \mathbb{F}_{2^n} \mid F(X + a + b) + F(X + b) + F(X + a) + F(X) = 0 \right\} \right|.$$

This is also denoted as $\nabla_F(a, b)$. The maximum of the entries of the FBCT of F , for $a \neq b, ab \neq 0$, is also known as the *second-order zero differential uniformity* of F , denoted as ∇_F , which is defined over any finite field characteristic.

Remark 2.7. If F is a permutation of \mathbb{F}_{2^n} , then it is immediate that

$$\text{FBCT}_F(a, b) = \sum_{c \in \mathbb{F}_{2^n}} \text{LBCT}_F(a, b, c).$$

Note that this implies that $\text{LBCT}_F(a, b, c) \leq \text{FBCT}_F(a, b)$, $\forall a, b, c \in \mathbb{F}_{2^n}$.

Definition 2.8. (Double Difference Distribution Table) [10] Let F be a permutation of \mathbb{F}_{2^n} . The *double difference distribution table* (DDDT) of F is a $2^n \times 2^n \times 2^n$ table where the entry at $(a, b, c) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$ is given by

$$\text{DDDT}_F(a, b, c) = |\{X \in \mathbb{F}_{2^n} \mid F(X + a + b) + F(X + b) + F(X + a) + F(X) = c\}|.$$

Note that $\text{FBCT}_F(a, b) = \text{DDDT}_F(a, b, 0)$.

Moreover, in [10] it was shown that for $a, b, c, d \in \mathbb{F}_{2^n}$ with $abcd = 0$, the values of $\text{EBCT}(a, b, c, d)$ are trivial to compute as are the values of $\text{LBCT}_F(a, b, c)$ and $\text{UBCT}_F(a, b, c)$, for $abc = 0$. We recall these results in the following three lemmas.

Lemma 2.9. [10] Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c, d \in \mathbb{F}_{2^n}$ with $abcd = 0$, we have

$$\text{EBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \mid \begin{cases} 2^n & \text{if } a = b = c = d = 0 \\ \text{DDT}_F(c, d) & \text{if } a = b = 0, c \neq 0, d \neq 0 \\ \text{DDT}_F(a, b) & \text{if } a \neq 0, b \neq 0, c = d = 0 \\ 0 & \text{otherwise} \end{cases} \right\} \right|.$$

Lemma 2.10. [10] Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c \in \mathbb{F}_{2^n}$ with $abc = 0$, we have

$$\text{LBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \mid \begin{cases} 2^n & \text{if } b = c = 0 \\ 0 & \text{if } b \neq 0, c = 0 \\ \text{DDT}_F(b, c) & \text{if } a = 0 \\ 0 & \text{if } b = 0, c \neq 0 \end{cases} \right\} \right|.$$

Lemma 2.11. [10] Let F be a permutation of \mathbb{F}_{2^n} . Then, for $a, b, c \in \mathbb{F}_{2^n}$ with $abc = 0$, we have

$$\text{UBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \mid \begin{cases} 2^n & \text{if } a = b = 0 \\ 0 & \text{if } a = 0, b \neq 0, c \neq 0 \\ \text{DDT}_F(a, b) & \text{if } c = 0 \\ 0 & \text{if } a \neq 0, b = 0, c \neq 0 \end{cases} \right\} \right|.$$

Also, [16] $\text{UBCT}_F(a, b, b) = \text{DDT}_F(a, b)$ and $\text{BCT}_F(a, c) = \sum_{b \in \mathbb{F}_{2^n}} \text{UBCT}_F(a, b, c)$. Addi-

tionally, it is easy to see that $\text{DDT}_F(a, c) = \text{LBCT}_F(a, a, c)$. Next, we recall the notion of DBCT as follows.

Definition 2.12. [19] Let $F(X)$ be a mapping from \mathbb{F}_{2^n} to itself. The Double Boomerang Connectivity Table (DBCT) is a $2^n \times 2^n$ table defined for $(a, d) \in \mathbb{F}_{2^n}^2$ by

$$\text{DBCT}_F(a, d) = \sum_{b, c} \text{dbct}(a, b, c, d),$$

where $\text{dbct}(a, b, c, d) = \text{UBCT}_F(a, b, c) \text{LBCT}_F(b, c, d)$. For $a = 0$ or $d = 0$, it can be easily obtained that

$$\text{DBCT}_F(0, d) = \sum_c \text{UBCT}_F(0, 0, c) \text{LBCT}_F(0, c, d) = 2^{2n}, \text{ and}$$

$$\text{DBCT}_F(a, 0) = \sum_b \text{UBCT}_F(a, b, 0) \text{LBCT}_F(b, 0, 0) = 2^{2n}.$$

3. GENERAL RESULTS ON THE EBCT

In a similar way as the results in [10] and [14], we can redefine the EBCT for a permutation F without involving its compositional inverse, as stated in the following lemma.

Lemma 3.1. *Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c, d \in \mathbb{F}_{2^n}$, we have*

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \mid \begin{cases} F(X) + F(X + a) = b \\ F(X) + F(X + c) = d \\ F(X + a + c) + F(X + a) = d \end{cases} \right\} \right|.$$

Proof.

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \mid \begin{cases} 1) F(X) + F(X + a) = b \\ 2) F(X) + F(X + c) = d \\ 3) F^{-1}(F(X) + d) + F^{-1}(F(X + a) + d) = a \end{cases} \right\} \right|.$$

Equation 2) implies that $F(X) + d = F(X + c)$. Equation 3) is then equivalent to

$$\begin{aligned} F^{-1}(F(X + c)) + F^{-1}(F(X + a) + d) &= a \\ F^{-1}(F(X + a) + d) &= X + a + c \\ F(X + a) + d &= F(X + a + c), \end{aligned}$$

completing the proof of the lemma. \square

As a consequence, we can derive the connection between the EBCT of a function and the EBCT of its compositional inverse.

Corollary 3.2. *Let F be a permutation of \mathbb{F}_{2^n} . Then for $a, b, c, d \in \mathbb{F}_{2^n}$, we have $\text{EBCT}_F(a, b, c, d) = \text{EBCT}_{F^{-1}}(b, a, d, c)$.*

Proof. We have by Lemma 3.1 that $\text{EBCT}_F(a, b, c, d)$ is given by the following equations:

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \mid \begin{cases} 1) F(X) + F(X + a) = b \\ 2) F(X) + F(X + c) = d \\ 3) F(X + a + c) + F(X + a) = d \end{cases} \right\} \right|.$$

Let $Y = F(X)$. Then, equation 1) gives $Y + F(X + a) = b$, which is equivalent to $F(X + a) = Y + b$, and then to $X + a = F^{-1}(Y + b)$, rendering the equation $F^{-1}(Y + b) + F^{-1}(Y) = a$.

Similarly, Equation 2) is equivalent to $F^{-1}(Y + d) + F^{-1}(Y) = c$.

Lastly, inserting Equation 1) in Equation 3), we obtain $F(X + a + c) = Y + d + b$, which is equivalent to $F^{-1}(Y + b + d) = X + a + c$. Since $F^{-1}(Y + b) + F^{-1}(Y) = F^{-1}(Y + b) + X = a$, this gives in turn $F^{-1}(Y + b + d) = F^{-1}(Y + b) + c$, equivalent to $F^{-1}(Y + b + d) + F^{-1}(Y + b) = c$.

From these three equations, the result follows. \square

Lemmas 2.5, 2.4 and 3.1 also suggest a generalization of these concepts for the set of all functions over \mathbb{F}_{2^n} (not necessarily permutations).

Definition 3.3. Let F be a function over \mathbb{F}_{2^n} . Then, for $a, b, c, d \in \mathbb{F}_{2^n}$, we can define the EBCT, LBCT and UBCT as, respectively:

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \mid \begin{cases} F(X) + F(X + a) = b \\ F(X) + F(X + c) = d \\ F(X + a + c) + F(X + a) = d \end{cases} \right\} \right|.$$

$$\text{LBCT}_F(a, b, c) = \left| \left\{ X \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} F(X+a) + F(X+a+b) = c \\ F(X) + F(X+b) = c \end{cases} \right. \right\} \right|.$$

$$\text{UBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} F(X+a) + F(Y+a) = c \\ F(X) + F(Y) = c \\ F(X) + F(X+a) = b \end{cases} \right. \right\} \right|.$$

Remark 3.4. Note that if F is a permutation, these definitions are equivalent to the original definitions of these concepts. Therefore, we will refer to this definition for all functions.

4. EBCT, LBCT AND UBCT OF A FUNCTION IN TERMS OF ITS DDT

In this section, we present a general result that gives an intriguing connection between the EBCT, LBCT and UBCT entries of a differentially δ -uniform function F with its DDT entries.

Theorem 4.1. Let F be a differentially δ -uniform function on \mathbb{F}_{2^n} . Let $k \in \{1, 2, \dots, \frac{\delta}{2}\}$. If $\text{DDT}_F(b, c) = 2k$, we let $\{x_0, x_0+b, x_1, x_1+b, \dots, x_{k-1}, x_{k-1}+b\}$ denote the solutions of the equation $F(X+b)+F(X) = c$; if $\text{DDT}_F(a, b) = 2k$, we let $\{y_0, y_0+a, y_1, y_1+a, \dots, y_{k-1}, y_{k-1}+a\}$ denote the solutions of the equation $F(X+a) + F(X) = b$; if $\text{DDT}_F(c, d) = 2k$, let $\{z_0, z_0+c, z_1, z_1+c, \dots, z_{k-1}, z_{k-1}+c\}$ be the solutions of the equation $F(X+c) + F(X) = d$. Then, for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have

$$\text{EBCT}_F(a, b, c, d) = \begin{cases} \text{DDT}_F(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_F(c, d) = 2k, a = z_i + z_j, i \neq j, b = F(z_i) + F(z_j), \\ & k \in \{2, \dots, \frac{\delta}{2}\}, \text{ and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \text{ or} \\ & \text{DDT}_F(c, d) = 2k, a = z_i + z_j + c, b = F(z_i) + F(z_j) + d \\ & k \in \{2, \dots, \frac{\delta}{2}\}, \text{ and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover for $a, b, c \in \mathbb{F}_{2^n}^*$, we have

$$\text{LBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(b, c) & \text{if } \text{DDT}_F(b, c) = 2, a = b, \text{ or} \\ & \text{DDT}_F(b, c) = 2k, a \in \{b, x_i + x_j, x_i + x_j + b\}, \\ & k \in \{2, \dots, \frac{\delta}{2}\}, \text{ and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise,} \end{cases}$$

and,

$$\text{UBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(a, b) & \text{if } \text{DDT}_F(a, b) = 2 \text{ and } c = b, \text{ or} \\ & \text{DDT}_F(a, b) = 2k, k \in \{2, \dots, \frac{\delta}{2}\} \text{ and} \\ & c \in \{b, F(y_i) + F(y_j), F(y_i) + F(y_j) + b\} \text{ for} \\ & i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By Lemma 3.1 (or Definition 3.3, in general),

$$\text{EBCT}_F(a, b, c, d) = \left| \left\{ X \in \mathbb{F}_{2^n} \left| \begin{cases} F(X) + F(X+a) = b \\ F(X) + F(X+c) = d \\ F(X+a+c) + F(X+a) = d \end{cases} \right. \right\} \right|.$$

Also, by Lemma 2.4 (or Definition 3.3, in general),

$$\text{LBCT}_F(a, c, d) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} X + Y = c \\ F(X + a) + F(Y + a) = d \\ F(X) + F(Y) = d \end{cases} \right. \right\} \right|.$$

For both EBCT and LBCT, the equations above imply that $F(X + a + c) + F(X + a) = F(X) + F(X + c) = d$. Since F is differentially δ -uniform, the equation $F(X) + F(X + c) = d$ can have zero, two or $2k$ solutions, for $k \in \{2, \dots, \frac{\delta}{2}\}$. We can then divide the proof into three cases:

- If $\text{DDT}_F(c, d) = 0$, then the equation has no solutions, and therefore $\text{LBCT}_F(a, c, d) = \text{EBCT}_F(a, b, c, d) = 0$.
- If $\text{DDT}_F(c, d) = 2$, then the solutions are of the form $z_0, z_0 + c$. This implies that $a = c$, and then, in that case, $\text{LBCT}_F(a, c, d) = \text{DDT}_F(c, d)$. For EBCT, from the condition $F(X) + F(X + a) = b$, we obtain that $b = d$.
- If $\text{DDT}_F(c, d) = 2k$, then the solutions are of the form $z_0, z_0 + c, z_1, z_1 + c, \dots, z_{k-1}, z_{k-1} + c$, where $k \in \{2, \dots, \frac{\delta}{2}\}$. Then, in order for the above equation to have a solution, $a \in \{0, c, z_i + z_j, z_i + z_j + c, i \neq j\}$. Since $a \neq 0$, then, when $a = c, z_i + z_j, z_i + z_j + c$, $\text{LBCT}_F(a, c, d) = \text{DDT}_F(c, d)$. If $a \notin \{c, z_i + z_j, z_i + z_j + c\}$, $\text{LBCT}_F(a, c, d) = \text{EBCT}_F(a, b, c, d) = 0$. A change of indices for LBCT gives the stated result. Now, we give $\text{EBCT}_F(a, b, c, d)$ when $a \in \{c, z_i + z_j, z_i + z_j + c\}$.
 - If $a = c$, then as before, $b = d$, in which case $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d)$. If $b \neq d$, there are no solutions.
 - If $a = z_i + z_j, i \neq j$, then inserting the solution z_i in the equation $F(X + a) + F(X) = b$ gives $F(z_i + a) + F(z_i) = F(z_j) + F(z_i) = b$, while the other equations are satisfied. If $b = F(z_j) + F(z_i)$, then, $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d)$. If $b \neq F(z_j) + F(z_i)$, then there are no solutions.
 - If $a = z_i + z_j + c$, then inserting the solution z_i in the equation $F(X + a) + F(X) = b$ gives $F(z_i + a) + F(z_i) = F(z_j + c) + F(z_i) = b$, while the other equations are satisfied. If $b = F(z_i) + F(z_j + c) = F(z_i) + F(z_j) + d$, then, $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d)$. If $b \neq F(z_i) + F(z_j) + d$, then there are no solutions.

This proves the result for EBCT and LBCT.

Lastly, we have by Lemma 2.11, or by definition 3.3 in general, that:

$$\text{UBCT}_F(a, b, c) = \left| \left\{ (X, Y) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n} \left| \begin{cases} F(X + a) + F(Y + a) = c \\ F(X) + F(Y) = c \\ F(X) + F(X + a) = b \end{cases} \right. \right\} \right|.$$

From the first two equations, we obtain $F(X) + F(X + a) = F(Y) + F(Y + a) = b$. Since F is differentially δ -uniform, both this equation and the equation $F(X) + F(X + a) = b$ can have zero, two or $2k$ solutions, for $k \in \{2, \dots, \frac{\delta}{2}\}$. We can then divide the proof in three cases:

- If $\text{DDT}_F(a, b) = 0$, then the equation $F(X) + F(X + a) = b$ has no solutions, and therefore $\text{UBCT}_F(a, b, c) = 0$.
- If $\text{DDT}_F(a, b) = 2$, then the solutions of $F(X) + F(X + a) = F(Y) + F(Y + a) = b$ will imply that either $Y = X$ or $Y = X + a$. Clearly $Y \neq X$ as $c \neq 0$. Thus, we have $Y = X + a$ and in that case $\text{UBCT}_F(a, b, c) \neq 0$ only if $c = b$. Therefore, $\text{UBCT}_F(a, b, c) = \text{DDT}_F(a, b)$ if $c = b$.
- If $\text{DDT}_F(a, b) = 2k$, then the solutions of $F(X) + F(X + a) = b$ are of the form $y_0, y_0 + a, y_1, y_1 + a, \dots, y_{k-1}, y_{k-1} + a$, where $k \in \{2, \dots, \frac{\delta}{2}\}$. Then, setting $X = y_i$

and $Y = y_j$ in the first two equations, where $i, j \in \{0, 1, \dots, k-1\}$. We obtain that $F(y_i) + F(y_i + a) = F(y_j) + F(y_j + a)$, so $y_i \in \{y_j, y_j + a, y_l, y_l + a\}$ where $l (\neq i, j) \in \{0, 1, \dots, k-1\}$. If $y_i \neq y_j$ we get $c = 0$, a contradiction; if $y_i = y_j + a$, we obtain that $b = c$. If $y_i = y_l$, then $c = F(y_i) + F(y_l)$; if $y_i = y_l + a$, we obtain $c = F(y_i) + F(y_l + a) = F(y_i) + F(y_l) + b$.

This proves the result for UBCT. \square

Next, we provide a characterisation of EBCT entries of a differentially δ -uniform function F in terms of UBCT and LBCT entries of F , and vice versa.

Theorem 4.2. *Let F be a differentially δ -uniform function. Then for $a, b, c, d \in \mathbb{F}_{2^n}^*$, $\text{EBCT}_F(a, b, c, d) = 2$ if and only if $\text{LBCT}_F(a, c, d) = \text{UBCT}_F(c, d, b) = 2$. Furthermore, if $\delta \geq 4$, for $k \in \{2, \dots, \frac{\delta}{2}\}$, let $A_k = \{(a, c, d) \mid \exists b, \text{ with } \text{EBCT}_F(a, b, c, d) = 2k\}$ and $B_k = \{(c, d, b) \mid \text{EBCT}_F(a, b, c, d) = 2k, \text{ for some } a\}$. Then, $(a, c, d) \in A_k$ if and only if $\text{LBCT}_F(a, c, d) = 2k$, and $(c, d, b) \in B_k$ if and only if $\text{UBCT}_F(c, d, b) = 2k$.*

Proof. Let F be a differentially δ -uniform function. Then from the Theorem 4.1, for $a, b, c, d \in \mathbb{F}_{2^n}^*$,

$$\text{EBCT}_F(a, b, c, d) = \begin{cases} \text{DDT}_F(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_F(c, d) = 2k, a = z_i + z_j, i \neq j, b = F(z_i) + F(z_j), \text{ or} \\ & \text{DDT}_F(c, d) = 2k, a = z_i + z_j + c, b = F(z_i) + F(z_j) + d \\ & k \in \{2, \dots, \frac{\delta}{2}\} \text{ and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise,} \end{cases}$$

where $\{z_0, z_0 + c, z_1, z_1 + c, \dots, z_{k-1}, z_{k-1} + c\}$ are the $2k$ solutions of the equation $F(X + c) + F(X) = d$ for $k = \{1, 2, \dots, \frac{\delta}{2}\}$ if $\text{DDT}_F(c, d) = 2k$.

On the other hand, we have

$$\text{LBCT}_F(a, c, d) = \begin{cases} \text{DDT}_F(c, d) & \text{DDT}_F(c, d) = 2, a = c, \text{ or} \\ & a \in \{c, z_i + z_j, z_i + z_j + c\}, k \in \{2, \dots, \frac{\delta}{2}\}, \\ & \text{and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{UBCT}_F(c, d, b) = \begin{cases} \text{DDT}_F(c, d) & \text{DDT}_F(c, d) = 2, b = d, \text{ or} \\ & b \in \{d, F(z_i) + F(z_j), F(z_i) + F(z_j) + d\}, k \in \{2, \dots, \frac{\delta}{2}\}, \\ & \text{and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise.} \end{cases}$$

If $\text{EBCT}_F(a, b, c, d) = 2$, then $a = c, b = d$ and $\text{DDT}_F(c, d) = 2$. Then from Theorem 4.1, $\text{LBCT}_F(a, c, d) = \text{UBCT}_F(c, d, b) = \text{DDT}_F(c, d) = 2$. On the other hand, if $\text{LBCT}_F(a, c, d) = \text{UBCT}_F(c, d, b) = 2$, then again using Theorem 4.1, we have $a = c, b = d$ and $\text{DDT}_F(c, d) = 2$, so $\text{EBCT}_F(a, b, c, d) = 2$.

If $(a, c, d) \in A_k$, where $k = \{2, \dots, \frac{\delta}{2}\}$ then $\text{DDT}_F(c, d) = 2k$ and $a \in \{c, z_i + z_j, z_i + z_j + c\}$, $i, j \in \{0, 1, \dots, k-1\}, i \neq j$, which implies that $\text{LBCT}_F(a, c, d) = 2k$. On the other hand, if $\text{LBCT}_F(a, c, d) = 2k$, then $\text{DDT}_F(c, d) = 2k$ and $a \in \{c, z_i + z_j, z_i + z_j + c\}$, $i, j \in \{0, 1, \dots, k-1\}, i \neq j$. We consider the cases:

- If $a = c$, then $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d) = 2k$ if $b = d$, so $(a, c, d) \in A_k$ for $b = d$.
- If $a = z_i + z_j$, then $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d) = 2k$ if $b = F(z_i) + F(z_j)$, so we get $(a, c, d) \in A_k$.

- If $a = z_i + z_j + c$, then $\text{EBCT}_F(a, b, c, d) = \text{DDT}_F(c, d) = 2k$ if $b = F(z_i) + F(z_j + c) = F(z_i) + F(z_j) + d$, so $(a, c, d) \in A_k$.

Similarly, we can prove the result for the connection between the EBCT and the UBCT. \square

Remark 4.3. Note that there can be a, c, d values such that $\text{EBCT}_F(a, b, c, d) = 0$, but $\text{LBCT}_F(a, c, d) \neq 0$, and b, c, d values with $\text{EBCT}_F(a, b, c, d) = 0$, and $\text{UBCT}_F(c, d, b) \neq 0$.

Remark 4.4. Note that, if F is a permutation, the result for the connection between the EBCT and the LBCT can be easily deduced from Definitions 2.1 and 2.2.

Remark 4.5. Note that the theorem implies that $(\text{EBCT}_F(a, b, c, d))^2 \leq \text{LBCT}_F(a, c, d) \cdot \text{UBCT}_F(c, d, b)$. Furthermore, the equality is met for every $a, b, c, d \in \mathbb{F}_{2^n}$ if and only if F is APN.

5. CONSEQUENCES OF OUR RESULTS FOR APN FUNCTIONS

We first state the EBCT, LBCT and UBCT entries of an APN function F and give these entries in terms of the DDT entries of F . It is known that $H(X) = X^{-1}$ over \mathbb{F}_{2^n} , where n is odd, is an APN function and the authors in [10, 14] have computed its EBCT, LBCT and UBCT entries, which turns out to be a special case of the following result.

Corollary 5.1. If F is an APN function over \mathbb{F}_{2^n} , then for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have

$$\text{EBCT}_F(a, b, c, d) = \begin{cases} \text{DDT}_F(a, b) & \text{if } a = c \text{ and } b = d \\ 0 & \text{otherwise.} \end{cases}$$

$$\text{LBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(a, c) & \text{if } a = b \\ 0 & \text{otherwise,} \end{cases}$$

and,

$$\text{UBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(a, b) & \text{if } c = b \\ 0 & \text{otherwise,} \end{cases}$$

Proof. The proof of this corollary is straightforward from Theorem 4.1, by setting $\delta = 2$. \square

Remark 5.2. For an APN function F , one can compute the explicit entries of all the three tables, $\text{EBCT}_F(a, b, c, d)$, $\text{LBCT}_F(a, b, c)$ and $\text{UBCT}_F(a, b, c)$, for all $a, b, c, d \in \mathbb{F}_{2^n}$ using Lemmas 2.11–2.9 and our aforementioned corollary.

Remark 5.3. For an APN function F , Corollary 5.1, Lemma 2.10 and Remark 2.7 imply the result from [3][Theorem 1]:

$$\text{FBCT}_F(a, b) = \begin{cases} 0 & \text{if } ab \neq 0, a \neq b, \\ 2^n & \text{otherwise.} \end{cases}$$

where $a, b \in \mathbb{F}_{2^n}$.

6. CONSEQUENCES OF OUR RESULTS FOR SOME WELL-KNOWN FUNCTIONS

In this section, we will give the EBCT, LBCT and UBCT entries of the Gold, Kasami and Bracken-Leander functions for those parameters such that the functions are differentially 4-uniform permutations; in the case of the Gold function, we also state and prove the EBCT, LBCT and UBCT entries for any parameters. We can also give a very short proof for the EBCT, LBCT and UBCT entries of the inverse function for n even, which is the main result in [10, 14] (for n odd, see the previous section). It may be noted that for those $a, b, c \in \mathbb{F}_{2^n}$, for

which $abc = 0$, the explicit values of LBCT and UBCT can be obtained by using Lemma 2.4 and Lemma 2.10, and, for those $a, b, c, d \in \mathbb{F}_{2^n}$, for which $abcd = 0$, the explicit values of EBCT can be obtained by using Lemma 2.9. Thus, throughout this section, we shall determine the LBCT and UBCT entries for all those $a, b, c \in \mathbb{F}_{2^n}$, for which $abc \neq 0$ and EBCT entries for all those $a, b, c, d \in \mathbb{F}_{2^n}$, for which $abcd \neq 0$.

For clarity, we first state the results we will use to compute EBCT, LBCT and UBCT entries of differentially 4-uniform functions using Theorem 4.1 and Theorem 4.2, which we will need for the functions in this section.

Corollary 6.1. *Let F be a differentially 4-uniform function over \mathbb{F}_{2^n} . Then, for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have*

$$\text{EBCT}_F(a, b, c, d) = \begin{cases} \text{DDT}_F(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_F(c, d) = 4, a = z_0 + z_1, b = F(z_0) + F(z_1), \text{ or} \\ & \text{DDT}_F(c, d) = 4, a = z_0 + z_1 + c, b = F(z_0) + F(z_1 + c) \\ 0 & \text{otherwise,} \end{cases}$$

where $z_0, z_0 + c, z_1, z_1 + c$ are the four solutions of the equation $F(X + c) + F(X) = d$ if $\text{DDT}_F(c, d) = 4$. Next,

$$\text{LBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(b, c) & \text{if } \text{DDT}_F(b, c) = 2 \text{ and } a = b, \text{ or} \\ & \text{DDT}_F(b, c) = 4 \text{ and } a \in \{b, x_0 + x_1, x_0 + x_1 + b\}, \\ 0 & \text{otherwise,} \end{cases}$$

where $x_0, x_0 + b, x_1, x_1 + b$ are the four solutions of the equation $F(X + b) + F(X) = c$ if $\text{DDT}_F(b, c) = 4$. Further,

$$\text{UBCT}_F(a, b, c) = \begin{cases} \text{DDT}_F(a, b) & \text{if } \text{DDT}_F(a, b) = 2 \text{ and } c = b, \text{ or} \\ & \text{DDT}_F(a, b) = 4 \text{ and } c \in \{b, F(y_0) + F(y_1), F(y_0) + F(y_1) + b\}, \\ 0 & \text{otherwise,} \end{cases}$$

where $y_0, y_0 + a, y_1, y_1 + a$ are the four solutions of the equation $F(X + a) + F(X) = b$ if $\text{DDT}_F(a, b) = 4$.

Proof. The proof directly follows from Theorem 4.1 by taking $\delta = 4$. \square

Corollary 6.2. *Let F be a differentially 4-uniform function over \mathbb{F}_{2^n} . Let $A = \{(a, c, d) \mid \exists b, \text{ with } \text{EBCT}_F(a, b, c, d) = 4\}$ and $B = \{(c, d, b) \mid \text{EBCT}_F(a, b, c, d) = 4, \text{ for some } a\}$. Then for $a, b, c \in \mathbb{F}_{2^n}^*$, $\text{EBCT}_F(a, b, c, d) = 2$ if and only if $\text{LBCT}_F(a, c, d) = \text{UBCT}_F(c, d, b) = 2$. Furthermore, $(a, c, d) \in A$ if and only if $\text{LBCT}_F(a, c, d) = 4$, and $(c, d, b) \in B$ if and only if $\text{UBCT}_F(c, d, b) = 4$.*

Proof. By setting $\delta = 4$, one can see that the proof directly follows from Theorem 4.2. \square

We will here use Corollary 6.1– 6.2 to compute the concrete values for three infinite classes of differentially 4-uniform power permutations over \mathbb{F}_{2^n} (Table 1) having the best known nonlinearity.

We first consider the Gold function $F_1(X) = X^{2^k+1}$, a differentially 2^ℓ -uniform function over \mathbb{F}_{2^n} , where $\ell = \gcd(k, n)$ and compute its EBCT, LBCT and UBCT entries. We will use the relative trace in the following way. For a general k , we do not have in general that $\mathbb{F}_{2^k} \subseteq \mathbb{F}_{2^n}$. However, we can naturally embed the elements of \mathbb{F}_{2^n} in $\mathbb{F}_{2^{km}}$, since $m = \frac{n}{\gcd(k, n)}$. Then, $\sum_{i=0}^{m-1} \alpha^{2^{ki}} = \text{Tr}_k^{km}(\alpha)$.

Family	d	condition	LBCT/UBCT
Inverse	$2^n - 2$	$n = 2k, k > 1$	[10, 14]
Gold	$2^k + 1$	$n = 2k, k \text{ odd}, \gcd(i, n) = 2$	This paper
Kasami	$2^{2i} - 2^i + 1$	$n = 2k, k \text{ odd}, \gcd(i, n) = 2$	This paper
Bracken-Leander	$2^{2k} + 2^k + 1$	$n = 4k, k \text{ odd}$	This paper

Table 1. Differentially 4-uniform permutations X^d over \mathbb{F}_{2^n} .

Corollary 6.3. Let $F_1(X) = X^{2^k+1}$, the general Gold function. Let $\ell = \gcd(k, n)$, and $m = \frac{n}{\ell}$. Then,

$$\text{EBCT}_{F_1}(a, b, c, d) = \begin{cases} 2^\ell & \text{if } \text{Tr}_k^{km} \left(\frac{d}{c^{2^k+1}} \right) = \text{Tr}_1^m(1) \text{ and } a = c, b = d, \text{ or} \\ & \text{Tr}_k^{km} \left(\frac{d}{c^{2^k+1}} \right) = \text{Tr}_1^m(1) \text{ and } a = uc, \text{ and } b = (u + u^2)c^{2^k+1} + ud, \\ & \text{where } u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{LBCT}_{F_1}(a, b, c) = \begin{cases} 2^\ell & \text{if } \text{Tr}_k^{km} \left(\frac{c}{b^{2^k+1}} \right) = \text{Tr}_1^m(1) \text{ and } a \in b\mathbb{F}_{2^\ell}^*, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{UBCT}_{F_1}(a, b, c) = \begin{cases} 2^\ell & \text{if } \text{Tr}_k^{km} \left(\frac{b}{a^{2^k+1}} \right) = \text{Tr}_1^m(1), c \in \{b, (u + u^2)a^{2^k+1} + ub, u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}\}, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. By Theorem 4.1, $\text{EBCT}_{F_1}(a, b, c, d)$ equals

$$\begin{cases} \text{DDT}_{F_1}(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_{F_1}(c, d) = 2k, a = z_i + z_j, i \neq j, b = F_1(z_i) + F_1(z_j), \text{ or} \\ & \text{DDT}_{F_1}(c, d) = 2k, a = z_i + z_j + c, b = F_1(z_i) + F_1(z_j) + d \\ & k \in \{2, \dots, \frac{\delta}{2}\}, \text{ and } i, j \in \{0, 1, \dots, k-1\}, i \neq j, \\ 0 & \text{otherwise,} \end{cases}$$

where $\{z_0, z_0 + c, z_1, z_1 + c, \dots, z_{k-1}, z_{k-1} + c\}$ are the $2k$ solutions of the equation $F(X + c) + F(X) = d$ for $\text{DDT}_F(c, d) = 2k$, where $k \in \{1, 2, \dots, \frac{\delta}{2}\}$.

It is known from [2, Lemma 4] that $X^{2^k+1} + (X + c)^{2^k+1} = d$ has either zero or 2^ℓ distinct solutions if and only if $\sum_{i=0}^{m-1} D^{2^{ki}} = m \equiv \text{Tr}_1^m(1) \pmod{2}$, where $D = \frac{d}{c^{2^k+1}}$ (zero solutions otherwise), $\ell = \gcd(k, n)$ and $m = \frac{n}{\ell}$. The solutions are of the form $X + cu$, where $u \in \mathbb{F}_{2^\ell}$. Then, the set of the sum of two different solutions $z_i, z_j, i \neq j, z_j \neq z_i + c$, of the equation $(X + c)^{2^k+1} + X^{2^k+1} = d$ is cu where $u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}$. For given any $z_i + z_j = cu$, where $u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}$, we need to compute the value of $F(z_i) + F(z_j)$. We have

$$\begin{aligned} (uc)^{2^k+1} &= (z_0 + z_1)^{2^k+1} = z_0^{2^k+1} + z_1^{2^k} z_0 + z_1 z_0^{2^k} + z_1^{2^k+1} \\ &= F_1(z_0) + F_1(z_1) + (z_0 + uc)^{2^k} z_0 + (z_0 + uc) z_0^{2^k} \\ &= F_1(z_0) + F_1(z_1) + z_0(uc)^{2^k} + uc z_0^{2^k}. \end{aligned}$$

On the other hand, since $(z_0 + c)^{2^k+1} + z_0^{2^k+1} = d$, we get that $c^{2^k} z_0 + c z_0^{2^k} + c^{2^k+1} = d$. Given that $\ell|k$, then $u^{2^k} = u$, and therefore $z_0(uc)^{2^k} + uc z_0^{2^k} = u(c^{2^k} z_0 + c z_0^{2^k}) = u(d + c^{2^k+1})$. Substituting above, it gives that $F_1(z_0) + F_1(z_1) = (uc)^{2^k+1} + u(d + c^{2^k+1}) = (u + u^2)c^{2^k+1} + ud$.

If $\ell = 1$, we get that F_1 is APN, and using Corollary 5.1, we are done. Next, let us suppose that $\ell > 1$ or equivalently, $\text{DDT}_{F_1}(c, d) \neq 2$, then we will compute the LBCT and UBCT entries via (and using the notations of) Theorem 4.2. Obviously, $\text{LBCT}_{F_1}(a, c, d) \neq 2$, for any $(a, c, d) \in \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^*$ and, $\text{UBCT}_{F_1}(c, d, b) \neq 2$, for any $(c, d, b) \in \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^*$. Notice that

$$A = \left\{ (c, c, d), (uc, c, d) \mid \text{DDT}_{F_1}(c, d) = 2^\ell \text{ and } u \in \mathbb{F}_{2^\ell}^* \setminus \{1\} \right\},$$

as, in the case $\text{DDT}_{F_1}(c, d) = 2^\ell$, $(c, c, d) \in A$, for $b = d$ and $(uc, c, d) \in A$, for $b = (u + u^2)c^{2^k+1} + ud$. Thus, from Corollary 6.2, we have $\text{LBCT}_{F_1}(a, c, d) = 2^\ell$, for $a \in \{c, uc\}$, where $u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}$. This completes the LBCT case.

Similarly, one can write

$$B = \left\{ (c, d, d), (c, d, (u + u^2)c^{2^k+1} + ud) \mid \text{DDT}_{F_1}(c, d) = 2^\ell \text{ and } u \in \mathbb{F}_{2^\ell}^* \setminus \{1\} \right\},$$

as, in the case $\text{DDT}_{F_1}(c, d) = 4$, $(c, d, d) \in B$, for $a = c$ and $(c, d, (u + u^2)c^{2^k+1} + ud) \in B$, for $a = uc$. This gives us the UBCT of the Gold function. \square

It is known that the Gold function is a differentially 4-uniform permutation over \mathbb{F}_{2^n} , when $\gcd(k, n) = 2$, $n = 2t$ and t is odd. By using the above Corollary 6.3, one can see that either $u = \omega$ or $u = \omega^2$, thus giving the EBCT, LBCT and UBCT entries of differentially 4-uniform Gold permutation.

We will derive here the FBCT for the Gold function for any parameters (found also in [9], but our proof is significantly shorter).

Corollary 6.4. *Let $F_1(X) = X^{2^k+1}$, with $\gcd(k, n) = \ell$ and $m = \frac{n}{\ell}$, where $ab \neq 0$. Without loss of generality, we take $b = 1$. Then:*

$$\text{FBCT}_{F_1}(a, 1) = \begin{cases} 2^n & \text{if } a \in \mathbb{F}_{2^\ell}^*, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Observe that

$$\text{LBCT}_{F_1}(a, 1, c) = \begin{cases} 2^\ell & \text{if } \sum_{i=0}^{m-1} c^{2^{ki}} = 1 \text{ and } a \in \mathbb{F}_{2^\ell}^*, \\ 0 & \text{otherwise.} \end{cases}$$

This implies that, if $a \notin \mathbb{F}_{2^\ell}^*$, then $\text{FBCT}_{F_1}(a, 1) = 0$. Let us now assume that $a \in \mathbb{F}_{2^\ell}^*$. Note that the condition $\sum_{i=0}^{m-1} c^{2^{ki}} = 1$ is equivalent to $\text{DDT}(1, c) = 2^\ell$. Denoting as is usual

$$\omega_i = |\{c \mid \text{DDT}(1, c) = i\}|, \text{FBCT}_{F_1}(a, 1) = \sum_{c \in \mathbb{F}_{2^n}} \text{LBCT}_{F_1}(a, 1, c) = 2^\ell \omega_{2^\ell} = \sum_{i=0}^{2^\ell} i \omega_i = 2^n.$$

This proves our claim. \square

Remark 6.5. *This prior result can be generalized. For any function F , $\text{FBCT}_F(1, 1) = 2^n$. If $x_i + x_j$, $i \neq j$, is independent of c , then, by a similar argument, for $a \in \{x_i + x_j, x_i + x_j + 1\}$ we have $\text{FBCT}_F(a, 1) = 2^n$. If, for every $i \neq j$ we have that $x_i + x_j$ is independent of c , then, for any $a \neq 0$:*

$$\text{FBCT}_{F_1}(a, 1) = \begin{cases} 2^n & \text{if } a \in \{1, x_i + x_j, x_i + x_j + 1\} \\ 0 & \text{otherwise.} \end{cases}$$

Example 6.6. *In Table 2, Corollary 6.3 is illustrated by giving explicit values of the EBCT, LBCT and UBCT entries of the Gold function over $\mathbb{F}_{2^{2t}} = \langle g \rangle$, where g is a primitive element of $\mathbb{F}_{2^{2t}}$.*

t	k	a	b	c	d	ω	EBCT $_{F_1}(a, b, c, d)$	LBCT $_{F_1}(a, b, c)$	UBCT $_{F_1}(a, b, c)$
3	2	g^{44}	g^{23}	g^{16}	for any d	g^{21}	0	4	0
3	2	g^2	g^{26}	g^{53}	for any d	g^{21}	0	0	4
5	6	g^{351}	g^{692}	g^2	for any d	g^{341}	0	0	0
5	4	g^2	g^{359}	g^{11}	for any d	g^{682}	0	0	4
3	2	g^{44}	g^8	g^{23}	g^{16}	g^{21}	4	0	0
5	4	g^{684}	g^{11}	g^2	g^{359}	g^{682}	4	0	0

Table 2. EBCT, LBCT and UBCT entries for the Gold function over $\mathbb{F}_{2^{2t}}$

The Kasami function is a well-known nonlinear function over \mathbb{F}_{2^n} , given by $F_2(X) = X^{2^{2k}-2^k+1}$. Hertel and Pott [12] showed that for $\gcd(k, n) = 2$, where $n = 2t$, t is odd and $3 \nmid t$, it is a differentially 4-uniform permutation. We next investigate the EBCT, LBCT and UBCT entries of F_2 over \mathbb{F}_{2^n} .

Corollary 6.7. *Let $F_2(X) = X^{2^{2k}-2^k+1}$, where $\gcd(k, n) = 2$, $n = 2t$ and t is odd and $3 \nmid t$. Then for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have*

$$\text{EBCT}_{F_2}(a, b, c, d) = \begin{cases} 4 & \text{if } \text{DDT}_{F_2}(c, d) = 4 \text{ and } a = c, b = d, \text{ or} \\ & \text{DDT}_{F_2}(c, d) = 4, a = \omega c + \alpha^{2^k+1}, \text{ and } b = \omega d + \alpha^{2^{3k}+1}, \text{ or} \\ & \text{DDT}_{F_2}(c, d) = 4, a = \omega^2 c + \alpha^{2^k+1}, \text{ and } b = \omega^2 d + \alpha^{2^{3k}+1} \\ 0 & \text{otherwise,} \end{cases}$$

for $\alpha \in \mathbb{F}_{2^n}^*$ with $c^{2^{2k}} + c^{2^k} \alpha^{2^{3k}-2^k} + c \alpha^{2^{3k}+2^{2k}-2^k-1} + d \alpha^{2^{2k}-1} = 0$ and $\text{Tr}_k^{kt} \left(1 + \frac{c}{\alpha^{2^k+1}} \right) = 0$.

$$\text{LBCT}_{F_2}(a, b, c) = \begin{cases} 4 & \text{if } \text{DDT}_{F_2}(b, c) = 4 \text{ and } a \in \left\{ b, b\omega + \alpha^{2^k+1}, b\omega^2 + \alpha^{2^k+1} \right\}, \\ 0 & \text{otherwise,} \end{cases}$$

for $\alpha \in \mathbb{F}_{2^n}^*$ with $b^{2^{2k}} + b^{2^k} \alpha^{2^{3k}-2^k} + b \alpha^{2^{3k}+2^{2k}-2^k-1} + c \alpha^{2^{2k}-1} = 0$ and $\text{Tr}_k^{kt} \left(1 + \frac{b}{\alpha^{2^k+1}} \right) = 0$.

$$\text{UBCT}_{F_2}(a, b, c) = \begin{cases} 4 & \text{if } \text{DDT}_{F_2}(a, b) = 4 \text{ and } c \in \left\{ b, b\omega + \alpha^{2^{3k}+1}, b\omega^2 + \alpha^{2^{3k}+1} \right\}, \\ 0 & \text{otherwise,} \end{cases}$$

for $\alpha \in \mathbb{F}_{2^n}^*$ satisfying $a^{2^{2k}} + a^{2^k} \alpha^{2^{3k}-2^k} + a \alpha^{2^{3k}+2^{2k}-2^k-1} + b \alpha^{2^{2k}-1} = 0$, $\text{Tr}_k^{kt} \left(1 + \frac{a}{\alpha^{2^k+1}} \right) = 0$ and ω is the cube root of unity.

Proof. By Corollary 6.1, $\text{EBCT}_{F_2}(a, b, c, d)$ equals

$$\begin{cases} \text{DDT}_{F_2}(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_{F_2}(c, d) = 4, a = z_0 + z_1, b = F_2(z_0) + F_2(z_1), \text{ or} \\ & \text{DDT}_{F_2}(c, d) = 4, a = z_0 + z_1 + c, b = F_2(z_0) + F_2(z_1 + c) \\ 0 & \text{otherwise,} \end{cases}$$

where $z_0, z_0 + c, z_1, z_1 + c$ are the four solutions of the equation $F_2(X + c) + F_2(X) = d$ if $\text{DDT}_{F_2}(c, d) = 4$. We write $F_2(X + c) + F_2(X) = d$ as the system

$$(6.1) \quad \begin{cases} Y + X = c, \\ F_2(Y) + F_2(X) = d. \end{cases}$$

Now by using the argument given in [6], and the fact that $\gcd(2^k + 1, 2^n - 1) = 1$, we assume that $X = u^{2^k+1}, Y = v^{2^k+1} = (u + \alpha)^{2^k+1}$, for some $\alpha \in \mathbb{F}_{2^n}^*$ such that $v = u + \alpha$. Then, after simplifying System (6.1), we have

$$\begin{cases} \left(\frac{u}{\alpha}\right)^{2^k} + \frac{u}{\alpha} + 1 + \frac{c}{\alpha^{2^k+1}} = 0, \\ \left(\frac{u}{\alpha}\right)^{2^{3k}} + \frac{u}{\alpha} + 1 + \frac{d}{\alpha^{2^{3k}+1}} = 0, \end{cases}$$

or equivalently,

$$\begin{cases} \left(\frac{u}{\alpha}\right)^{2^k} + \frac{u}{\alpha} + 1 + \frac{c}{\alpha^{2^k+1}} = 0, \\ c^{2^{2k}} + c^{2^k} \alpha^{2^{3k}-2^k} + c \alpha^{2^{3k}+2^{2k}-2^k-1} + d \alpha^{2^{2k}-1} = 0. \end{cases}$$

Notice that the first equation of the above system has either 0 or 4 solutions in \mathbb{F}_{2^n} , depending on whether $\sum_{i=0}^{t-1} \left(1 + \frac{c}{\alpha^{2^k+1}}\right)^{2^{ki}}$ is zero or not, respectively. This first equation is simply the equation for the DDT entry of X^{2^k+1} at (α, c) , and if $\text{DDT}_{X^{2^k+1}}(\alpha, c) = 4$, then the four solutions of $\left(\frac{u}{\alpha}\right)^{2^k} + \frac{u}{\alpha} + 1 + \frac{c}{\alpha^{2^k+1}} = 0$ are in $\{u, u + \alpha, u + \alpha\omega, u + \alpha\omega^2\}$. This will give us $X \in \left\{u^{2^k+1}, (u + \alpha)^{2^k+1}, (u + \alpha\omega)^{2^k+1}, (u + \alpha\omega^2)^{2^k+1}\right\}$ as solutions of $F(X) + F(X + c) = d$, or equivalently $\{z_0, z_0 + b, z_1, z_1 + b\} \subseteq \left\{u^{2^k+1}, (u + \alpha)^{2^k+1}, (u + \alpha\omega)^{2^k+1}, (u + \alpha\omega^2)^{2^k+1}\right\}$.

Then for $z_0 = u^{2^k+1}$, we have the following two cases:

Case 1. Let $z_1 = (u + \alpha\omega)^{2^k+1}$ and $z_2 = (u + \alpha\omega^2)^{2^k+1}$. Then $z_0 + z_1 = a$ and $F_2(z_0) + F_2(z_1) = b$ can be written as

$$\begin{cases} u^{2^k+1} + (u + \alpha\omega)^{2^k+1} = a \\ u^{2^{3k}+1} + (u + \alpha\omega)^{2^{3k}+1} = b. \end{cases}$$

Also, by using $u^{2^{3k}+1} + (u + \alpha)^{2^{3k}+1} = d$, and the second equation of above system will give us $b = d\omega + \alpha^{2^{3k}+1}$. To compute a , we use $u^{2^k+1} + (u + \alpha)^{2^k+1} = c$ and the first equation of the above system and simplify them to get $a = c\omega + \alpha^{2^k+1}$.

Case 2. Let $z_1 = (u + \alpha\omega^2)^{2^k+1}$ and $z_2 = (u + \alpha\omega)^{2^k+1}$. By using the same argument as in the above case, we get $a = c\omega^2 + \alpha^{2^k+1}$ and $b = d\omega^2 + \alpha^{2^{3k}+1}$.

Summarizing both cases, we get the EBCT of the Kasami function.

We next compute the LBCT and UBCT entries via (and using the notations of) Corollary 6.2. As $\text{DDT}_{F_2}(c, d)$ is either 0 or 4, then, $\text{LBCT}_{F_2}(a, c, d) \neq 2$, for any $(a, c, d) \in \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^*$ and, $\text{UBCT}_{F_2}(c, d, b) \neq 2$, for any $(c, d, b) \in \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^* \times \mathbb{F}_{2^n}^*$.

Now, one can write set

$$A = \left\{ (c, c, d), (\omega c + \alpha^{2^k+1}, c, d), (\omega^2 c + \alpha^{2^k+1}, c, d) \mid \text{DDT}_{F_2}(c, d) = 4 \right\},$$

for $c, d \in \mathbb{F}_{2^n}^*$. This is because, when $\text{DDT}_{F_2}(c, d) = 4$, $(c, c, d) \in A$, for $b = d$, $(\omega c + \alpha^{2^k+1}, c, d) \in A$, for $b = d\omega + \alpha^{2^{3k}+1}$ and $(\omega^2 c + \alpha^{2^k+1}, c, d) \in A$, for $b = d\omega^2 + \alpha^{2^{3k}+1}$. Finally, from Corollary 6.2, we have $\text{LBCT}_{F_2}(a, c, d) = 4$, for $a \in \left\{c, \omega c + \alpha^{2^k+1}, \omega^2 c + \alpha^{2^k+1}\right\}$. Thus, by reordering the indexes, we complete the LBCT case.

Similarly, one can write the set

$$B = \left\{ (c, d, d), (c, d, d\omega + \alpha^{2^{3k}+1}), (c, d, d\omega^2 + \alpha^{2^{3k}+1}) \mid \text{DDT}_{F_2}(c, d) = 4 \right\},$$

as, in the case $\text{DDT}_{F_2}(c, d) = 4$, $(c, d, d) \in B$, for $a = c$, $(c, d, d\omega + \alpha^{2^{3k}+1}) \in B$, for $a = \omega c + \alpha^{2^k+1}$ and $(c, d, d\omega^2 + \alpha^{2^{3k}+1}) \in B$, for $a = \omega^2 c + \alpha^{2^k+1}$. Hence, one can get the UBCT of the Kasami function by reordering the indexes. \square

Example 6.8. *The examples given in Table 3 (g denotes a primitive element of $\mathbb{F}_{2^{2t}}$) illustrate Corollary 6.7.*

t	k	a	b	c	d	α	ω	$\text{EBCT}_{F_2}(a, b, c, d)$	$\text{LBCT}_{F_2}(a, b, c)$	$\text{UBCT}_{F_2}(a, b, c)$
5	2	g^6	g	g^{605}	for any d	g^{50}	g^{341}	0	0	4
5	6	g^5	g	g^{774}	for any d	g^{994}	g^{682}	0	0	0
5	2	g^{401}	g^{605}	g^6	g	g^{50}	g^{341}	4	0	0
5	6	g^{84}	g^{24}	g^2	for any d	g^{681}	g^{682}	0	4	0

Table 3. EBCT, LBCT and UBCT entries for Kasami function over $\mathbb{F}_{2^{2t}}$

Bracken and Leander [4] studied the differential uniformity of $F_3(X) = X^{2^{2k}+2k+1}$ over \mathbb{F}_{2^n} where $n = 4k$, which is often known as the Bracken-Leander function. They showed that F_3 is differentially 4-uniform, and its nonlinearity is $2^{n-1} - 2^{\frac{n}{2}}$. Further, Calderini and Villa [5] investigated the BCT entries and showed that the boomerang uniformity of the Bracken-Leander function is upper bounded by 24. Here, we first recall some results from [18] that will be used to compute EBCT, LBCT and UBCT entries for the Bracken-Leander function.

Lemma 6.9. [18, Lemma 1] *For any $X \in \mathbb{F}_{2^n}$ and $c \in \mathbb{F}_{2^n}$, let $t = \text{Tr}_k^n(c + 1) \in \mathbb{F}_{2^k}$, where $n = 4k$,*

- (1) *If $F_3(X) + F_3(X + 1) = c$, then $\text{Tr}_k^n(x) = t$.*
- (2) *Let $Z = X + X^{2^k}$. Then $F_3(x) + F_3(X + 1) = c$ holds if and only if $(X, Z) \in \mathbb{F}_{2^n} \times \mathbb{F}_{2^n}$ satisfies the below equations:*

$$(6.2) \quad Z^2 + (t + 1)Z + c^{2^k} + c^{2^{2k}} = 0$$

$$(6.3) \quad X^2 + X + Z^{2^k+1} + Z^{2^k} + z^2 + c + 1 = 0.$$

Next, we define W to be the set of all $X \in \mathbb{F}_{2^n}$ such that there exists $X' \in \mathbb{F}_{2^n} \setminus \{X, X + 1\}$ with $X^{2^{2k}+2k+1} + (X + 1)^{2^{2k}+2k+1} = X'^{2^{2k}+2k+1} + (X' + 1)^{2^{2k}+2k+1}$. We now recall a result describing W explicitly.

Lemma 6.10. [18, Theorem 2] *Given elements $u \in \mathbb{F}_{2^{2k}} \setminus \{\mathbb{F}_{2^k}\}$ and $v \in \mathbb{F}_{2^n} \setminus \{\mathbb{F}_{2^{2k}}\}$ such that*

$$u + u^{2^k} = 1, v + v^{2^{2k}} = 1,$$

which always exist, then $X \in W$ if and only if

$$X = (tu + \beta)v + \tau,$$

where $t \in \mathbb{F}_{2^k} \setminus \{1\}$, $\tau \in \mathbb{F}_{2^{2k}}$, and $\beta \in \mathbb{F}_{2^k}$ satisfies

$$\beta = (t + 1)^{-1}(\alpha^2 + \alpha) + (t + 1)(u^2 + u) + 1,$$

for some $\alpha \in \mathbb{F}_{2^k}$.

Remark 6.11. *The proof of the above lemma also shows that $X' = X + (t + 1)u + \alpha$.*

Next, we determine the EBCT, LBCT and UBCT entries for the Bracken-Leander function.

Corollary 6.12. *Let $F_3(X) = X^{2^{2k}+2^k+1}$, where $n = 4k$. Then for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have*

$$\text{EBCT}_{F_3}(a, b, c, d) = \begin{cases} \text{DDT}_{F_3}(c, d) & \text{if } a = c, b = d, \\ & \text{or } \text{DDT}_{F_3}(c, d) = 4, a = c((t+1)u + \alpha), \text{ and} \\ & \quad b = d(\alpha + (t+1)u^{2^k}) + F_3(c)\gamma, \\ & \text{or } \text{DDT}_{F_3}(c, d) = 4, a = c((t+1)u + \alpha + 1), \text{ and} \\ & \quad b = d(\alpha + (t+1)u^{2^k} + 1) + F_3(c)\gamma, \\ 0 & \text{otherwise,} \end{cases}$$

where $t = \text{Tr}_k^n\left(\frac{d}{F_3(c)}\right)$, and $F_3(tu + \beta)v + \tau + F_3(tu + \beta)v + \tau + 1 = \frac{d}{F_3(c)}$,

$$\text{LBCT}_{F_3}(a, b, c) = \begin{cases} \text{DDT}_{F_3}(b, c) & \text{if } \text{DDT}_{F_3}(b, c) = 2 \text{ and } a = b, \\ & \text{or } \text{DDT}_{F_3}(b, c) = 4, \text{ and} \\ & \quad a \in \{b, b((t+1)u + \alpha), b((t+1)u + \alpha + 1)\}, \\ 0 & \text{otherwise,} \end{cases}$$

where $t = \text{Tr}_k^n\left(\frac{c}{F_3(b)}\right)$, and $F_3(tu + \beta)v + \tau + F_3(tu + \beta)v + \tau + 1 = \frac{c}{F_3(b)}$, and,

$$\text{UBCT}_{F_3}(a, b, c) = \begin{cases} \text{DDT}_{F_3}(a, b) & \text{if } \text{DDT}_{F_3}(a, b) = 2 \text{ and } c = b, \\ & \text{or } \text{DDT}_{F_3}(a, b) = 4 \text{ and} \\ & \quad c \in \{b, b(\alpha + (t+1)u^{2^k}) + F_3(a)\gamma, \text{ and} \\ & \quad b(\alpha + (t+1)(1+u) + 1) + F_3(a)\gamma\}, \\ 0 & \text{otherwise,} \end{cases}$$

where $t = \text{Tr}_k^n\left(\frac{b}{F_3(a)}\right)$, and $F_3(tu + \beta)v + \tau + F_3(tu + \beta)v + \tau + 1 = \frac{b}{F_3(a)}$.

In all the tables computed, $\gamma = ((t+1)^{-1}(\alpha^4 + \alpha^2) + (t+1)^3(u^4 + u^2) + \alpha^2 t + \alpha t + t(t+1)(u + u^2))$, $\tau \in \mathbb{F}_{2^{2k}}$, $u \in \mathbb{F}_{2^{2k}} \setminus \{\mathbb{F}_{2^k}\}$, $v \in \mathbb{F}_{2^n} \setminus \{\mathbb{F}_{2^{2k}}\}$ satisfying $u + u^{2^k} = 1$, $v + v^{2^{2k}} = 1$ and $\beta \in \mathbb{F}_{2^k}$ satisfying $\beta = (t+1)^{-1}(\alpha^2 + \alpha) + (t+1)(u^2 + u) + 1$, for some $\alpha \in \mathbb{F}_{2^k}$.

Proof. By Corollary 6.1, $\text{EBCT}_{F_3}(a, b, c, d)$ is equal to

$$\begin{cases} \text{DDT}_{F_3}(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_{F_3}(c, d) = 4, a = z_0 + z_1, b = F_3(z_0) + F_3(z_1), \text{ or} \\ & \text{DDT}_{F_3}(c, d) = 4, a = z_0 + z_1 + c, b = F_3(z_0) + F_3(z_1 + c) \\ 0 & \text{otherwise,} \end{cases}$$

where $z_0, z_0 + c, z_1, z_1 + c$ are the four solutions of the equation $F_3(X + c) + F_3(X) = d$ if $\text{DDT}_{F_3}(c, d) = 4$. This is equivalent to $F_3\left(\frac{X}{c} + 1\right) + F_3\left(\frac{X}{c}\right) = \frac{d}{c^{2^{2k}+2^k+1}}$ having solutions $\left\{\frac{z_0}{c}, \frac{z_0}{c} + 1, \frac{z_1}{c}, \frac{z_1}{c} + 1\right\}$. We can simplify $F_3\left(\frac{X}{c} + 1\right) + F_3\left(\frac{X}{c}\right) = \frac{d}{F_3(c)}$ as

$$\left(\frac{X}{c}\right)^{2^{2k}+2^k} + \left(\frac{X}{c}\right)^{2^{2k}+1} + \left(\frac{X}{c}\right)^{2^k+1} + \left(\frac{X}{c}\right)^{2^{2k}} + \left(\frac{X}{c}\right)^{2^k} + \frac{X}{c} + 1 = \frac{d}{F_3(c)}.$$

Let $\text{Tr}_k^n\left(\frac{d}{F_3(c)} + 1\right) = \text{Tr}_k^n\left(\frac{d}{F_3(c)}\right) = t$, where Tr_k^n denotes the relative trace map from \mathbb{F}_{2^n} to \mathbb{F}_{2^k} .

First assume that $t = 1$, then from the proof of [4, Theorem 1], we know that $F_3\left(\frac{X}{c} + 1\right) + F_3\left(\frac{X}{c}\right) = \frac{d}{F_3(c)}$ can have at most two solutions, a contradiction. Next, for $t \neq 1$, from Lemma 6.10, for $u \in \mathbb{F}_{2^{2k}} \setminus \{\mathbb{F}_{2^k}\}$ and $v \in \mathbb{F}_{2^n} \setminus \{\mathbb{F}_{2^{2k}}\}$ such that

$u + u^{2^k} = 1$, $v + v^{2^k} = 1$, we know that $\frac{z_0}{c} \in W$ if and only if $\frac{z_0}{c} = (tu + \beta)v + \tau$ and $\frac{z_1}{c} \in \left\{ \frac{z_0}{c} + (t+1)u + \alpha, \frac{z_0}{c} + (t+1)u + \alpha + 1 \right\}$, where $\tau \in \mathbb{F}_{2^{2k}}$, and $\beta \in \mathbb{F}_{2^k}$ satisfies the relation $\beta = (t+1)^{-1}(\alpha^2 + \alpha) + (t+1)(u^2 + u) + 1$, for some $\alpha \in \mathbb{F}_{2^k}$. Thus, $F_3\left(\frac{X}{c} + 1\right) + F_3\left(\frac{X}{c}\right) = \frac{d}{F_3(c)}$ can be written as $F_3((tu + \beta)v + \tau) + F_3((tu + \beta)v + \tau + 1) = \frac{d}{F_3(c)}$.

Again, we divide our analysis into the following two cases:

Case 1. Let $\frac{z_1}{c} = \frac{z_0}{c} + (t+1)u + \alpha$, then we have $a = c((t+1)u + \alpha)$. Now, $b = F_3(z_0) + F_3(z_1) = z_0^{2^{2k}+2^k+1} + z_1^{2^{2k}+2^k+1} = c^{2^{2k}+2^k+1}(F_3\left(\frac{z_0}{c}\right) + F_3\left(\frac{z_0}{c} + (t+1)u + \alpha\right))$. For $y_0 = \frac{z_0}{c}$ and $D = \frac{d}{F_3(c)}$, we have

$$\begin{aligned}
\frac{b}{F_3(c)} &= F_3(y_0) + F_3(y_0 + (t+1)u + \alpha) \\
&= (\alpha + (t+1)u)(y_0^{2^{2k}+2^k} + y_0^{2^{2k}+1} + y_0^{2^k+1}) + (\alpha + (t+1)u)^2(y_0^{2^{2k}} + y_0^{2^k} + y_0) \\
&\quad + (t+1)(y_0^{2^{2k}+1} + (\alpha + (t+1)u)(y_0 + y_0^{2^k})) + (\alpha + (t+1)u)^d \\
&= ((\alpha + (t+1)u) + (\alpha + (t+1)u)^2)(y_0^{2^{2k}} + y_0^{2^k} + y_0) + (\alpha + (t+1)u)(1 + D) \\
&\quad + (t+1)(y_0^{2^{2k}+1} + (\alpha + (t+1)u)(y_0 + y_0^{2^k})) + (\alpha + (t+1)u)^d \\
&= (t+1)(tu + \beta + 1)(y_0^{2^{2k}} + y_0^{2^k} + y_0) + (\alpha + (t+1)u)(1 + D) \\
&\quad + (t+1)(y_0^{2^{2k}+1} + (\alpha + (t+1)u)(y_0 + y_0^{2^k})) + (\alpha + (t+1)u)^d \\
&= (t+1)(tu + \beta + 1)(tu + \beta + (tu^{2^k} + \beta)v^{2^k} + \tau^{2^k}) + (\alpha + (t+1)u)(1 + D) \\
&\quad + (t+1)((tu + \beta)^2 v^{2^{2k}+1} + \tau(tu + \beta) + \tau^2) \\
&\quad + (\alpha + (t+1)u)(tu + \beta) + (\alpha + (t+1)u)^d \\
&= (t+1)(D + tu + \beta + 1) + (t+1)(tu + \beta + 1)(tu + \beta) + (\alpha + (t+1)u)(1 + D) \\
&\quad + (t+1)(\alpha + (t+1)u)(tu + \beta) + (\alpha + (t+1)u)^d \\
&= (\alpha + (t+1)u^{2^k})D + (t+1)^{-1}(\alpha^4 + \alpha^2) + (t+1)^3(u^4 + u^2) \\
&\quad + \alpha^2 t + \alpha t + t(t+1)(u + u^2).
\end{aligned}$$

The above expression will give us $b = d(\alpha + (t+1)u^{2^k}) + F_3(c)((t+1)^{-1}(\alpha^4 + \alpha^2) + (t+1)^3(u^4 + u^2) + \alpha^2 t + \alpha t + t(t+1)(u + u^2)) = d(\alpha + (t+1)u^{2^k}) + F_3(c)\gamma$ (say), where $\gamma = ((t+1)^{-1}(\alpha^4 + \alpha^2) + (t+1)^3(u^4 + u^2) + \alpha^2 t + \alpha t + t(t+1)(u + u^2))$.

Case 2. Let $\frac{z_1}{c} = \frac{z_0}{c} + (t+1)u + \alpha + 1$, then by same argument as in Case 1, we have $a = c((t+1)u + \alpha + 1)$ and $b = d(\alpha + (t+1)u^{2^k} + 1) + F_3(c)((t+1)^{-1}(\alpha^4 + \alpha^2) + (t+1)^3(u^4 + u^2) + \alpha^2 t + \alpha t + t(t+1)(u + u^2))$.

Combining both of these cases, we get the EBCT entries for the Bracken-Leander function.

We again use (also its notations) Corollary 6.2 to compute the LBCT and UBCT entries for the Bracken-Leander function. As we have computed $\text{EBCT}_{F_3}(a, b, c, d) = 2$, when $a = c, b = d$ and $\text{DDT}_{F_3}(c, d) = 2$, then $\text{LBCT}_{F_3}(a, c, d) = 2$, for $a = c$ when $\text{DDT}_{F_3}(c, d) = 2$ and, $\text{UBCT}_{F_3}(c, d, b) = 2$, for $b = d$ when $\text{DDT}_{F_3}(c, d) = 2$.

Also, $A = \{(c, c, d), (c((t+1)u + \alpha), c, d), (c((t+1)u + \alpha + 1), c, d) \mid \text{DDT}_{F_3}(c, d) = 4\}$, as, in the case $\text{DDT}_{F_4}(c, d) = 4$, $(c, c, d) \in A$, for $b = d$, $(c((t+1)u + \alpha), c, d) \in A$, for $b = d(\alpha + (t+1)u^{2^k}) + F_3(c)\gamma$ and $(c((t+1)u + \alpha + 1), c, d) \in A$, for $b = d(\alpha + (t+1)u^{2^k} + 1) + F_3(c)\gamma$. Thus, from Corollary 6.2, we have $\text{LBCT}_{F_3}(a, c, d) = 4$, for $a \in \{c, c((t+1)u + \alpha), c((t+1)u + \alpha + 1)\}$. This completes the LBCT case.

Similarly, one can write $B = \{(c, d, d), (c, d, d(\alpha + (t + 1)u^{2^k}) + F_3(c)\gamma), (c, d, d(\alpha + (t + 1)u^{2^k} + 1) + F_3(c)\gamma) \mid \text{DDT}_{F_3}(c, d) = 4\}$, as, in the case of $\text{DDT}_{F_3}(c, d) = 4$, $(c, d, d) \in B$, for $a = c$, $(c, d, d(\alpha + (t + 1)u^{2^k}) + F_3(c)\gamma) \in B$, for $a = c((t + 1)u + \alpha)$ and $(c, d, d(\alpha + (t + 1)u^{2^k} + 1) + F_3(c)\gamma) \in B$, for $a = c((t + 1)u + \alpha + 1)$.

Hence, we get the LBCT and UBCT of the Bracken-Leander function by just reordering the indexes. \square

Example 6.13. To illustrate Corollary 6.12, Table 4 gives explicit values of the LBCT and UBCT entries of Bracken-Leander function over $\mathbb{F}_{2^{4k}}^* = \langle g \rangle$ (g is a primitive element of $\mathbb{F}_{2^{4k}}$).

k	a	b	c	d	α	u	v	τ	$\text{EBCT}_{F_3}(a, b, c, d)$	$\text{LBCT}_{F_3}(a, b, c)$	$\text{UBCT}_{F_3}(a, b, c)$
2	g	g	g	d	$-$	$-$	$-$	$-$	2	2	2
2	g^{71}	g^3	g^{32}	d	g^{255}	g^{17}	g^{123}	g^{17}	0	4	0
2	g^{54}	g^{37}	g^{26}	d	g^{85}	g^{34}	g^{224}	g^{17}	0	4	0
2	g^{70}	g^3	g^{103}	d	g^{85}	g^{34}	g^{224}	g^{17}	0	0	4
2	g^{36}	g^{103}	g^{70}	g^3	g^{85}	g^{34}	g^{224}	g^{17}	4	0	0

Table 4. EBCT, LBCT and UBCT entries for Bracken-Leander function over $\mathbb{F}_{2^{4k}}^* = \langle g \rangle$

Via our Corollary 6.1, we can also give a very short proof for the EBCT, LBCT and UBCT entries of the inverse function for n even, which is the main result in [10, 14] (for n odd, see the previous section).

Corollary 6.14. Let $F_4(X) = X^{2^n - 2}$ where n is even. Then for $a, b, c, d \in \mathbb{F}_{2^n}^*$, we have

$$\text{EBCT}_{F_4}(a, b, c, d) = \begin{cases} 4 & \text{if } a = c = \frac{1}{b} = \frac{1}{d}, \\ & \text{or } b = \frac{1}{a}, d = \frac{1}{c}, \text{ and } (ad)^2 + ad + 1 = 0, \\ 2 & \text{if } a = c, b = d, d \neq \frac{1}{c} \text{ and } \text{Tr}\left(\frac{1}{cd}\right) = 0, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{LBCT}_{F_4}(a, b, c) = \begin{cases} 4 & \text{if } a = b = \frac{1}{c}, \\ & \text{or } b = \frac{1}{c}, (ac)^2 + ac + 1 = 0, \\ 2 & \text{if } a = b \neq \frac{1}{c}, \text{Tr}\left(\frac{1}{bc}\right) = 0, \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{UBCT}_{F_4}(a, b, c) = \begin{cases} 4 & \text{if } b = c = \frac{1}{a}, \\ & \text{or } b = \frac{1}{a}, (ac)^2 + ac + 1 = 0, \\ 2 & \text{if } c = b \neq \frac{1}{a}, \text{Tr}\left(\frac{1}{ab}\right) = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. For all three results, we have to study the spectrum of the inverse function, which can be found, for instance, implied in the proofs in [10, 14], but we recall it below for easy reference.

The equation $(X + A)^{2^n - 2} + X^{2^n - 2} = B$, for $A, B \neq 0$, and n even, gives:

- If $X = 0$ or $X = A$, then $B = \frac{1}{A}$, otherwise there are no solutions.
- If $X \neq 0, A$, then the equation is equivalent to $X^2 + AX + \frac{A}{B} = 0$, which has two solutions if and only if $\text{Tr}\left(\frac{1}{AB}\right) = 0$, and zero solutions, otherwise.

By Corollary 6.1, $\text{EBCT}_{F_4}(a, b, c, d)$ is equal to

$$\begin{cases} \text{DDT}_{F_4}(c, d) & \text{if } a = c, b = d, \text{ or} \\ & \text{DDT}_{F_4}(c, d) = 4, a = z_0 + z_1, b = F_4(z_0) + F_4(z_1), \text{ or} \\ & \text{DDT}_{F_4}(c, d) = 4, a = z_0 + z_1 + c, b = F_4(z_0) + F_4(z_1 + c) \\ 0 & \text{otherwise,} \end{cases}$$

where $z_0, z_0 + c, z_1, z_1 + c$ are the four solutions of the equation $F(X + c) + F(X) = d$, if $\text{DDT}_F(c, d) = 4$.

If $a = c, b = d, d \neq \frac{1}{c}$ and $\text{Tr}\left(\frac{1}{cd}\right) = 0$, then $\text{EBCT}_{F_4}(a, b, c, d) = \text{DDT}_{F_4}(c, d) = 2$.

If $d = \frac{1}{c}$, and without loss of generality, we let $z_0 = 0, z_1 = \omega c$. If $a = c, b = d$, then $\text{EBCT}_{F_4}(a, b, c, d) = \text{DDT}_{F_4}(c, d) = 4$. If $a = z_0 + z_1 = \omega c$ and $b = F_4(z_0) + F_4(z_1) = \frac{1}{\omega c}$, then $\text{EBCT}_{F_4}(a, b, c, d) = \text{DDT}_{F_4}(c, d) = 4$. If $a = z_0 + z_1 + c = \omega^2 c$ and $b = F_4(z_0) + F_4(z_1 + c) = \frac{1}{\omega^2 c}$, then $\text{EBCT}_{F_4}(a, b, c, d) = \text{DDT}_{F_4}(c, d) = 4$. Note that these last two cases can be summarized as $b = \frac{1}{a}, d = \frac{1}{c}$ and $(ad)^2 + ad + 1 = 0$. In all other cases, $\text{EBCT}_{F_4}(a, b, c, d) = 0$.

We now compute the LBCT and UBCT of the inverse function using Corollary 6.2. As we have seen $\text{EBCT}_{F_4}(a, b, c, d) = 2$, when $a = c, b = d, d \neq \frac{1}{c}$ and $\text{Tr}\left(\frac{1}{cd}\right) = 0$, we infer that $\text{LBCT}_{F_4}(a, c, d) = 2$ if $a = c \neq \frac{1}{d}$ and $\text{Tr}\left(\frac{1}{cd}\right) = 0$. Similarly, $\text{UBCT}_{F_4}(c, d, b) = 2$ if $b = d \neq \frac{1}{a}$ and $\text{Tr}\left(\frac{1}{cd}\right) = 0$.

Clearly, when $d = \frac{1}{c}$, we can write $A = \{(c, c, d), (\omega c, c, d), (\omega^2 c, c, d) \mid \text{DDT}_{F_4}(c, d) = 4\}$, as, in the case $\text{DDT}_{F_4}(c, d) = 4$, $(c, c, d) \in A$, for $b = d$, $(\omega c, c, d) \in A$, for $b = \frac{1}{\omega c}$ and $(\omega^2 c, c, d) \in A$, for $b = \frac{1}{\omega^2 c}$. Thus, from Corollary 6.2, we have $\text{LBCT}_{F_4}(a, c, d) = 4$, for $a \in \{c, \omega c, \omega^2 c\}$ and $d = \frac{1}{c}$. This is equivalent to saying $\text{LBCT}_{F_4}(a, c, d) = 4$ when $d = \frac{1}{c}$ and $(ad)^2 + ad = 1 = 0$.

In a similar way, one can write $B = \{(c, d, d), (c, d, \frac{1}{\omega c}), (c, d, b = \frac{1}{\omega c}) \mid \text{DDT}_{F_4}(c, d) = 4\}$, as, in the case $\text{DDT}_{F_4}(c, d) = 4$, $(c, d, d) \in B$, for $a = c$, $(c, d, \frac{1}{\omega c}) \in B$, for $a = \omega c$ and $(c, d, \frac{1}{\omega c}) \in B$, for $a = \omega^2 c$.

This completes the proof. \square

Using 2.7 and our results for the inverse function, we can easily compute its FBCT (also found in [9]).

Corollary 6.15. *Let $F_4(X) = X^{2^n - 2}$ over \mathbb{F}_{2^n} , with n even. Then, for $a, b \in \mathbb{F}_{2^n}^*$,*

$$\text{FBCT}_{F_4}(a, b) = \begin{cases} 2^n, & \text{if } a = b, \\ 4, & \text{if } \left(\frac{a}{b}\right)^2 + \frac{a}{b} + 1 = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If $a = b$, then trivially $\text{FBCT}_F(a, b) = 2^n$ for any function F . If $a \neq b$, by Corollary 6.14, $\text{LBCT}(a, b, c)$ is nonzero if and only if $b = \frac{1}{c}, (ac)^2 + ac + 1 = 0$, which is equivalent to $b = \frac{1}{c}, \left(\frac{a}{b}\right)^2 + \frac{a}{b} + a = 0$. In that case, $\text{LBCT}(a, b, c) = 4$. The result follows. \square

7. DBCT OF GOLD FUNCTION

In this section, we study the DBCT entries of the Gold function. To this end, we need the LBCT and UBCT entries of the Gold function, which we have already computed in Corollary 6.3. Also, if $k \mid n$, then for any $\alpha \in \mathbb{F}_{2^n}$, we have $\sum_{i=0}^{t-1} \alpha^{2^{ki}} = \text{Tr}_k^n(\alpha)$, the relative trace of \mathbb{F}_{2^n} over \mathbb{F}_{2^k} . If $k \nmid n$, then the elements of \mathbb{F}_{2^n} can be embedded in $\mathbb{F}_{2^{km}}$, since $m = \frac{n}{\gcd(k, n)}$.

Theorem 7.1. *Let $F_1(X) = X^{2^k+1}$, over \mathbb{F}_{2^n} where $\gcd(k, n) = \ell$ and $m = \frac{n}{\ell}$ is odd. Then,*

$$\text{DBCT}_{F_1}(a, d) = \begin{cases} 2^{2\ell} ((2^\ell - 2) + |N(a, d)|) & \text{if } \text{Tr}_k^{km} \left(\frac{d}{(a^{2^k+1})^{2^k+1}} \right) = u^4, u \in \mathbb{F}_{2^\ell}^* \setminus \{1\} \\ 2^{2\ell} \cdot |N(a, d)| & \text{otherwise,} \end{cases}$$

where $|N(a, d)|$ denotes the cardinality of $\left\{ b \in \mathbb{F}_{2^n}^* \mid \text{Tr}_k^{km} \left(\frac{b}{a^{2^k+1}} \right) = \text{Tr}_k^{km} \left(\frac{d}{b^{2^k+1}} \right) = m \right\}$.

Proof. First, assume that $a = 0$ or $d = 0$, then using definition of DBCT, we can write $\text{DBCT}_{F_1}(a, d) = 2^{2n}$. For $a, d \in \mathbb{F}_{2^n}^*$, we can write

$$\text{DBCT}_F(a, d) = \sum_{b=c} \text{UBCT}_F(a, b, c) \text{LBCT}_F(b, c, d) + \sum_{b \neq c} \text{UBCT}_F(a, b, c) \text{LBCT}_F(b, c, d).$$

Recall, for $a, d \in \mathbb{F}_{2^n}^*$, $\ell = \gcd(k, n)$ and $m = \frac{n}{\ell}$,

$$\text{LBCT}_{F_1}(b, c, d) = \begin{cases} 2^\ell & \text{if } \text{Tr}_k^{km} \left(\frac{d}{c^{2^k+1}} \right) = \text{Tr}_1^m(1) \text{ and } b \in c\mathbb{F}_{2^\ell}^* \text{ for } b, c, d \in \mathbb{F}_{2^n}^* \\ \text{DDT}_{F_1}(c, d) & \text{if } b = 0 \\ 0 & \text{otherwise,} \end{cases}$$

$$\text{UBCT}_{F_1}(a, b, c) = \begin{cases} 2^\ell & \text{if } \text{Tr}_k^{km} \left(\frac{b}{a^{2^k+1}} \right) = \text{Tr}_1^m(1) \text{ and for } a, b, c \in \mathbb{F}_{2^n}^*, \\ & c \in \left\{ b, (u+u^2)a^{2^k+1} + ub, u \in \mathbb{F}_{2^\ell}^* \setminus \{1\} \right\} \\ \text{DDT}_{F_1}(a, b) & \text{if } c = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Clearly for $b = 0$, we have $\text{DDT}_F(a, b) = \text{DDT}_{F_1}(a, 0) = 0$ as $a \neq 0$, thus, giving us $\text{UBCT}_{F_1}(a, b, c) = 0$ and similarly for $c = 0$, we get $\text{DDT}_{F_1}(c, d) = \text{DDT}_{F_1}(0, d) = 0$ as $d \neq 0$ giving us $\text{LBCT}_{F_1}(a, b, c) = 0$.

When $b \neq c$, then $\text{UBCT}_{F_1}(a, b, c) = 2^\ell$ only if $c = (u+u^2)a^{2^k+1} + ub$ and $\text{DDT}_{F_1}(a, b) = 2^\ell$, where $u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}$. Also, $\text{LBCT}_{F_1}(b, c, d) = 2^\ell$ only if $b = vc$ for some $v \in \mathbb{F}_{2^\ell}^*$ and $\text{DDT}_{F_1}(c, d) = 2^\ell$. After combining them, we have $c = (u+u^2)a^{2^k+1} + uvc$, and $\text{DDT}_{F_1}(a, b) = \text{DDT}_{F_1}(c, d) = 2^\ell$. Notice that $v \neq u^{-1}$, because otherwise $a = 0$. Thus, we have $c = \frac{(u+u^2)}{1+uv} a^{2^k+1}$ and therefore, $b = \frac{v(u+u^2)}{1+uv} a^{2^k+1}$. Then $\text{UBCT}_{F_1}(a, b, c) = \text{LBCT}_{F_1}(b, c, d) = 2^\ell$ if $\text{DDT}_{F_1}(a, b) = \text{DDT}_{F_1}(c, d) = 2^\ell$ which is the same as $\sum_{i=0}^{m-1} \frac{b^{2^{ki}}}{a^{2^{ki}(2^k+1)}} = \sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{c^{2^{ki}(2^k+1)}} = m$, where $m = \frac{n}{\ell}$. This is equivalent to $\sum_{i=0}^{m-1} \left(\frac{v(u+u^2)}{1+uv} \right)^{2^{ki}} = \sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{\left(\frac{(u+u^2)}{1+uv} a^{2^k+1} \right)^{2^{ki}(2^k+1)}} = m$.

One can see that the first equality $\sum_{i=0}^{m-1} \left(\frac{v(u+u^2)}{1+uv} \right)^{2^{ki}} = m$, is nothing but $m \left(\frac{v(u+u^2)}{1+uv} \right) = m$, as $u, v \in \mathbb{F}_{2^\ell} \subseteq \mathbb{F}_{2^k}$. This is the same as saying $m \frac{u^2v+1}{uv+1} = 0$, or equivalently we have $\frac{u^2v+1}{uv+1} = 0$ or equivalently, $v = \frac{1}{u^2}$, as m is odd. Then we have $\frac{u^2v+1}{uv+1} = 0$ or equivalently, $v = \frac{1}{u^2}$. This will give us $c = u^2(a^{2^k+1})$ and $b = a^{2^k+1}$. Thus, for $b = a^{2^k+1}$ and $c = u^2(a^{2^k+1})$, we have $\sum_{i=0}^{m-1} \frac{b^{2^{ki}}}{a^{2^{ki}(2^k+1)}} = 1 (\equiv m \pmod{2})$. Also, $\sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{(u^2 a^{2^k+1})^{2^{ki}(2^k+1)}} = m$ for some $a, d \in \mathbb{F}_{2^\ell}$. Obviously, such a and d always exist because $\sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{(u^2 a^{2^k+1})^{2^{ki}(2^k+1)}} = 1$ if and only if there exists an $A \in \mathbb{F}_{km}$ such that $\frac{d^{2^{ki}}}{(u^2 a^{2^k+1})^{2^{ki}(2^k+1)}} = 1 + A + A^2$. Finally, we can conclude that for all $u \in \mathbb{F}_{2^\ell}$, choosing $c = u^2(a^{2^k+1})$ and $b = a^{2^k+1}$, for

$\text{UBCT}_{F_1}(a, b, c) = \text{LBCT}_{F_1}(b, c, d) = 2^\ell$, we should have $\sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{u^4(a^{2^k+1})^{2^{ki}(2^k+1)}} = m$. Therefore, $\text{DBCT}_{F_1}(a, d) = 2^{2\ell}(2^\ell - 2)$ if $\text{Tr}_k^{km} \left(\frac{d}{(a^{2^k+1})^{2^k+1}} \right) = u^4$, for some $u \in \mathbb{F}_{2^\ell}^* \setminus \{1\}$.

Let us now have $b = c$, then

$$\sum_{b=c} \text{UBCT}_F(a, b, c) \text{LBCT}_F(b, c, d) = \sum_b \text{DDT}_{F_1}(a, b) \text{DDT}_{F_1}(b, d),$$

when $\sum_{i=0}^{m-1} \frac{b^{2^{ki}}}{a^{2^{ki}(2^k+1)}} = \sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{b^{2^{ki}(2^k+1)}} = m$. If any of these sums is different from m , then the product of the DDT's is 0. Hence, we need to compute $|N(a, d)|$, where

$$\begin{aligned} N(a, d) &= \left\{ b \in \mathbb{F}_{2^n}^* \mid \sum_{i=0}^{m-1} \frac{b^{2^{ki}}}{a^{2^{ki}(2^k+1)}} = \sum_{i=0}^{m-1} \frac{d^{2^{ki}}}{b^{2^{ki}(2^k+1)}} = m \right\} \\ &= \left\{ b \in \mathbb{F}_{2^n}^* \mid \text{Tr}_k^{km} \left(\frac{b}{a^{2^k+1}} \right) = \text{Tr}_k^{km} \left(\frac{d}{b^{2^k+1}} \right) = m \right\}, \end{aligned}$$

which completes the proof. \square

Example 7.2. To illustrate Theorem 7.1, Table 5 gives explicit values of the DBCT entries of Gold function over $\mathbb{F}_{2^n}^* = \langle g \rangle$ (g is a primitive element of \mathbb{F}_{2^n}).

m	k	a	d	u	$\text{DBCT}_{F_1}(a, d)$
3	2	g^{25}	g^{22}	$g^3 + g^2 + g + 1$	160
3	2	g^{63}	g^{56}	–	64
5	2	g^4	g^{186}	–	1024
5	4	g^2	g^{868}	$g^5 + g^3 + g + 1$	1024

Table 5. DBCT entries for the Gold function over \mathbb{F}_{2^n}

It would be interesting to investigate the DBCT entries of other functions, such as Kasami and Bracken-Leander functions.

8. INVARIANCE UNDER EXTENDED AFFINE EQUIVALENCE

It is easy to check that, for any function F , and a function $G = F_2 \circ F \circ F_1$, where $F_i(X) = A_i X + B_i$ are affine functions (we say that F is Extended Affine Equivalent (EA) to G), then $\text{DDT}_G(\alpha, \beta) = \text{DDT}_F \left(\alpha, \frac{\beta}{A_2} \right)$. Therefore, EBCT, LBCT and UBCT are not preserved under EA. However, from Theorem 4.1, we can see that $\sum_{a,c} \text{LBCT}_F(a, b, c)$, $\sum_{b,c} \text{LBCT}_F(a, b, c)$ and $\sum_{a,b,c} \text{EBCT}_F(a, b, c)$ are equivalent under EA.

Also, if $x_i + x_j$ is independent of c for every $i \neq j$, then $\sum_c \text{LBCT}_F(a, b, c)$ is preserved under EA.

9. CONCLUSION

We have determined the entries of the newly proposed Extended Boomerang Connectivity Table (EBCT), Lower Boomerang Connectivity Table (LBCT) and Upper Boomerang Connectivity Table (UBCT) of a δ -uniform function by establishing connections with the entries of the well-known Difference Distribution Table (DDT), which we consider to be a significant observation. We also give the EBCT, LBCT and UBCT entries of three classes of differentially 4-uniform power permutations, namely, Gold, Kasami and Bracken-Leander. Moreover, it turns out that one of our results covers the result of Eddahmani et al. [10] and

Man et al. [14] for the inverse function over \mathbb{F}_{2^n} . We also compute the DBCT entries for the Gold function. We further challenge the community to determine the DBCT entries of other classes of interesting functions, such as Kasami and Bracken-Leander functions over \mathbb{F}_{2^n} .

REFERENCES

- [1] E. Biham, A. Shamir, *Differential cryptanalysis of DES-like cryptosystems*, J. Cryptol. 4(1) (1991), 3–72.
- [2] C. Blondeau, A. Canteaut, P. Charpin, *Differential properties of power functions*, Int. J. Inf. Coding Theory 1(2) (2010), 149–170.
- [3] H. Boukerrou, P. Huynh, V. Lallemand, B. Mandal, M. Minier, *On the Feistel counterpart of the boomerang connectivity table*, IACR Trans. Symmetric Cryptol. 1 (2020), 331–362.
- [4] C. Bracken, G. Leander, *A highly nonlinear differentially 4 uniform power mapping that permutes fields of even degree*, Finite Fields Their Appl. 16(4) (2010), 231–242.
- [5] M. Calderini, I. Villa *On the boomerang uniformity of some permutation polynomials* Cryptogr. Commun. 12 (2020), 1161–1178.
- [6] C. Carlet, K. H. Kim, S. Mesnager, *A direct proof of APN-ness of the Kasami functions*, Des. Codes Cryptogr. 89 (2021), 441–446.
- [7] C. Cid, T. Huang, T. Peyrin, Y. Sasaki, and L. Song, *Boomerang connectivity table: a new cryptanalysis tool*. In: Nielsen J., Rijmen V. (eds.), Advances in Cryptology, EUROCRYPT 2018, LNCS 10821, Springer, Cham, pp. 683–714, 2018.
- [8] S. Delaune, P. Derbez, M. Vavrille, *Catching the Fastest Boomerangs: Application to SKINNY*, IACR Trans. Symmetric Cryptol. 4 (2020), 104–129.
- [9] S. Eddahmani, S. Mesnager, *Explicit values of the DDT, the BCT, the FBCT, and the FBDT of the inverse, the gold, and the Bracken-Leander S-boxes*, Cryptogr. Commun. 14 (2022), 1301–1344.
- [10] S. Eddahmani, S. Mesnager, *Determination of cryptographic tables and properties related to the revised boomerang and its application to a fundamental S-box*, Cryptology ePrint Archive (2024), <https://eprint.iacr.org/2024/577>.
- [11] H. Hadipour, N. Bagheri, L. Song, *Improved rectangle attacks on SKINNY and CRAFT*, IACR Trans. Symmetric Cryptol. 2 (2021), 140–198.
- [12] D. Hertel, A. Pott *Two results on maximum nonlinear functions*, Des. Codes Cryptogr., 47 (2008), 225–235.
- [13] K. Li, L. Qu, B. Sun, C. Li, *New results about the boomerang uniformity of permutation polynomials*, IEEE Trans. Inf. Theory 65(11) (2019), 7542–7553.
- [14] Y. Man, N. Li, Z. Liu, X. Zeng, *The Explicit values of the UBCT, the LBCT and the DBCT of the inverse function*, (2024), [arXiv:2404.12208](https://arxiv.org/abs/2404.12208).
- [15] K. Nyberg, *Differentially uniform mappings for cryptography*. In: Helleseht T. (eds.), Advances in Cryptology–EUROCRYPT 1993, LNCS 765, Springer, Berlin, Heidelberg, pp. 55–64, 1994.
- [16] D. Wagner, *The boomerang attack*, In: L. R. Knudsen (ed.) Fast Software Encryption-FSE 1999. LNCS 1636, Springer, Berlin, Heidelberg, pp. 156–170, 1999.
- [17] H. Wang, T. Peyrin, *Boomerang Switch in Multiple Rounds. Application to AES Variants and Deoxys*, IACR Trans. Symmetric Cryptol. 1 (2019), 142–169.
- [18] M. Xiong, H. Yan, *A note on the differential spectrum of a differentially 4-uniform power function*, Finite Fields Their Appl. 48 (2017), 117–125.
- [19] Q. Yang, L. Song, S. Sun, D. Shi, L. Hu, *New properties of the double boomerang connectivity table*, IACR Trans. Symmetric Cryptol. 4 (2022), 208–242.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY JAMMU, JAMMU 181221, INDIA
Email address: kirpa.garg@gmail.com

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY JAMMU, JAMMU 181221, INDIA
Email address: sartaj.hasan@iitjammu.ac.in

DEPARTMENT OF COMPUTER SCIENCE, ELECTRICAL ENGINEERING AND MATHEMATICAL SCIENCES, WESTERN NORWAY UNIVERSITY OF APPLIED SCIENCES, 5020 BERGEN, NORWAY
Email address: csr@hvl.no

APPLIED MATHEMATICS DEPARTMENT, NAVAL POSTGRADUATE SCHOOL, MONTEREY, CA 93943, USA
Email address: pstanica@nps.edu