

New Integrable Chiral Cosmological Models with Two Scalar Fields

Vsevolod R. Ivanov,^{1,*} and Sergey Yu. Vernov^{2,†}

¹*Physics Department, Lomonosov Moscow State University*

Leninskie Gory 1, 119991, Moscow, Russia

²*Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University,*

Leninskie Gory 1, 119991, Moscow, Russia

We construct integrable chiral cosmological models with two scalar fields and potentials represented in terms of hyperbolic functions. Using the conformal transformation of the metric and the corresponding models with induced gravity terms, we obtain the general solutions in the spatially flat, open and closed Friedmann universes and the corresponding integrals of motion. The obtained general solutions can be written in terms of the Jacobi elliptic functions of the conformal time.

PACS numbers: 04.20.Jb, 04.50.Kd, 98.80.-k

Keywords: Chiral cosmological model, integrability, modified gravity

I. INTRODUCTION

Most cosmological models, which describe the global evolution of the Universe, include scalar fields. To describe only one epoch of the Universe evolution, single field models can be used. Using two-field models, one can describe multiple epochs or multiple physical phenomenons. Models with a single scalar field nonminimally coupled to gravity can be transformed to models with a minimally coupled scalar field with a canonical kinetic term by the metric and scalar field transformations. On the other hand, it is not possible to transform a model with two scalar fields nonminimally coupled to gravity to a model with two minimally coupled scalar fields and a standard kinetic part of the Lagrangian in the most general case [1]. After the metric transformation, one obtains a general relativity model with nonstandard kinetic terms of scalar fields, a so-called chiral cosmological model (CCM) [2].

In the Friedmann–Lemaître–Robertson–Walker (FLRW) metric, the Einstein’s equations reduce to a system of ordinary differential equations. The problem of searching for integrable cosmological models with scalar fields is being actively investigated, essentially in the case of single-field

* vsvd.ivanov@gmail.com

† svernov@theory.sinp.msu.ru

cosmological models, but such models have been found only for a few specific forms of the scalar field potential [3–11].

A two-field CCM with a constant potential is trivially integrated not only in the spatially-flat FLRW metric, but also in the Bianchi I metric [12, 13]. At the same time, finding integrable cosmological models with multiple scalar fields is, in general, an extremely complicated task. We know only a few examples of integrable cosmological models with two or more fields [14–18]. The goal of this paper is to construct new integrable CCM with two fields and potentials written in terms of hyperbolic functions.

There are a few methods to integrate evolution equations in the FLRW metric. The integrability can be proven by using a suitable parametric time [6], but it is not immediately clear as to how to find this parametric time. The superpotential method, which is also known as the Hamilton-Jacobi equation approach, is useful to integrate not only single-field models with the exponential potential [3], but also two-field models with the exponential potential [18]. Note that this method is actively used to study inflation and to construct cosmological models with particular exact solutions [3, 18–35].

Another method for constructing one-field integrable models is based on the conformal metric transformation and the construction of the corresponding model in the Jordan frame [5, 8, 10, 36]. In Ref. [5], the authors construct two-field models with one ordinary scalar field, one phantom field and induced gravity terms to prove the integrability of a single-field cosmological model with a potential written in terms of hyperbolic functions. Note that integrability of this model has been proven in the closed and open Friedmann universes as well [37]. In Refs. [5, 38, 39], the proposed cosmological model has been intensively studied in the context of the early Universe evolution, with special emphasis on inflation and the possibility of crossing the Big Bang-Big Crunch singularity. In Ref. [10], the authors have proven integrability of the same model in a simpler way by using a single-field integrable model with non-minimal coupling constructed in Ref. [9]. Note that this model belongs to a wide class of integrable models with minimally coupled scalar fields and potentials in terms of hyperbolic functions. Their integrability have been proven by using a suitable parametric time in Ref. [6] (see also Ref. [10]).

To construct a new integrable CCM with two fields, we start from models with two nonminimally coupled scalar fields. The single-field integrable model proposed in Ref. [9] has an interesting feature: the Ricci scalar is an integral of motion. Recently, N -field cosmological models with the same property have been found and their integrability in the spatially flat FLRW metric have been proven [40]. In this paper, we find general solutions of evolution equations in the Friedmann

universe with arbitrary spatial curvature for such two-field integrable models with non-minimal coupling. After finding such solutions, we obtain two-field chiral cosmological models in the Einstein frame by the conformal transformation of the metric. For a few two-field potentials, we get general solutions. In the conformal time, we get analytic expressions of the general solutions in terms of the Jacobi elliptic functions.

The structure of the paper is as follows. In Section II, we consider two-field models with induced gravity terms. In Section III, we construct the corresponding CCMs by the conformal transformation of the metric and the introduction of new fields. In Section IV, we find solutions in the FLRW metric for the model with the nonminimal coupling. Examples of integrated CCMs and their general solutions in the conformal time are given in Section V. The results are summarized in Section VI.

II. TWO-FIELD MODEL WITH NONMINIMAL COUPLING

Let us consider models with two nonminimally coupled scalar fields, described by an action

$$S = \int d^4x \sqrt{-g} \left[U(\sigma^1, \sigma^2) R - \frac{1}{2} g^{\mu\nu} (C_1 \partial_\mu \sigma^1 \partial_\nu \sigma^1 + C_2 \partial_\mu \sigma^2 \partial_\nu \sigma^2) - V(\sigma^1, \sigma^2) \right], \quad (1)$$

where the functions U and V are differentiable, R is the Ricci scalar, and C_A are constants. Positive values of C_A correspond to the case of standard scalar fields.

By varying the action (1), we obtain evolution equations

$$U \left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right) = \nabla_\mu \nabla_\nu U - g_{\mu\nu} \square U + \frac{1}{2} T_{\mu\nu}, \quad (2)$$

where

$$T_{\mu\nu} = C_1 \partial_\mu \sigma^1 \partial_\nu \sigma^1 + C_2 \partial_\mu \sigma^2 \partial_\nu \sigma^2 - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} (C_1 \partial_\alpha \sigma^1 \partial_\beta \sigma^1 + C_2 \partial_\alpha \sigma^2 \partial_\beta \sigma^2) + V \right).$$

and the field equations are

$$C_A \square \sigma^A - V_{,\sigma^A} + R U_{,\sigma^A} = 0, \quad (3)$$

$A = 1, 2$; $F_{,\sigma^A} \equiv \frac{dF}{d\sigma^A}$ for any function $F(\sigma^1, \sigma^2)$, and there is no summation in A .

From Eq. (2), it follows

$$3\square U - UR = \frac{1}{2} g^{\mu\nu} T_{\mu\nu} = -\frac{1}{2} g^{\alpha\beta} \sum_{A=1}^2 C_A \partial_\alpha \sigma^A \partial_\beta \sigma^A - 2V. \quad (4)$$

Our goal is to find such functions U and V that Eq. (4) has an integral of motion. Using the field equations (3), we get

$$\square U = \sum_{A=1}^2 \frac{1}{C_A} \left(U_{,\sigma^A} V_{,\sigma^A} - R (U_{,\sigma^A})^2 \right) + \sum_{A,B=1}^2 U_{,\sigma^A \sigma^B} g^{\alpha\beta} \partial_\alpha \sigma^A \partial_\beta \sigma^B. \quad (5)$$

Substituting Eq. (5) into Eq. (4), we obtain

$$2V + \sum_{A=1}^2 \frac{3}{C_A} U_{,\sigma^A} V_{,\sigma^A} = R \left(U + \sum_{A=1}^2 \frac{3}{C_A} [U_{,\sigma^A}]^2 \right) + \frac{g^{\alpha\beta}}{2} \sum_{A,B=1}^2 [6U_{,\sigma^A \sigma^B} + C_A \delta_{AB}] \partial_\alpha \sigma^A \partial_\beta \sigma^B. \quad (6)$$

To solve Eq. (6), we choose such a function $U(\sigma^1, \sigma^2)$ that

$$6U_{,\sigma^A \sigma^B} + C_A \delta_{AB} = 0, \quad (7)$$

for all values of A and B , and

$$U + \sum_{A=1}^2 \frac{3}{C_A} [U_{,\sigma^A}]^2 = U_0, \quad (8)$$

where U_0 is a constant.

The solution of Eqs. (7) and (8) is

$$U(\sigma^1, \sigma^2) = U_0 - \frac{C_1}{12} (\sigma^1 - \sigma_0^1)^2 - \frac{C_2}{12} (\sigma^2 - \sigma_0^2)^2, \quad (9)$$

where σ_0^A are constants.

With such a function U , Eq. (6) is

$$2V + \sum_{A=1}^2 \frac{1}{2} \sigma^A V_{,\sigma^A} = R U_0. \quad (10)$$

If R is a constant, $R = R_0$, then Eq. (10) is a linear first-order partial differential equation with $V(\sigma^1, \sigma^2)$ as an unknown function. Equation (10) has the following general solution:

$$V = \frac{R_0 U_0}{2} + (\sigma^1 - \sigma_0^1)^4 f \left(\frac{\sigma^2 - \sigma_0^2}{\sigma^1 - \sigma_0^1} \right), \quad (11)$$

where f is an arbitrary differentiable function.

Note that for models with functions U and V given by Eqs. (9) and (11), the Ricci scalar R is the integral of motion in an arbitrary metric. This result can be generalized on models with an arbitrary number of scalar fields [40].

We consider the case of $U_0 > 0$, when the function $U > 0$ for some values of σ^A . The case of $U_0 < 0$ corresponds to antigravity. If $U_0 = 0$, then Eq. (10) restricts the potential without the assumption that R is a constant. It means that nontrivial solutions of the Friedmann equations can exist only for potential (11) with $U_0 = 0$. In the case of single-field models, an analogous restriction on the potential has been obtained in Ref. [41].

III. THE CORRESPONDING MODELS IN THE EINSTEIN FRAME

Now we construct models in the Einstein frame that correspond to the previously chosen functions U and V . To get a model with two standard scalar fields, we take $C_A = 1$. Also, we can take $\sigma_0^A = 0$ without loss of generality.

By doing a conformal transformation of the initial metric $g_{\mu\nu}$ of the form

$$g_{\mu\nu}^E = \frac{2U}{M_{\text{Pl}}^2} g_{\mu\nu}, \quad (12)$$

one arrives at the following action in the Einstein frame:

$$S_E = \int d^4x \sqrt{-g^E} \left(\frac{M_{\text{Pl}}^2}{2} R_E - \frac{g_E^{\mu\nu}}{2} \sum_{A,B=1}^2 K_{AB} \partial_\mu \sigma^A \partial_\nu \sigma^B - V_E \right), \quad (13)$$

where M_{Pl} is the reduced Planck mass,

$$K_{AB} = \frac{M_{\text{Pl}}^2}{2U} \left(\delta_{AB} + \frac{3}{U} U_{,\sigma^A} U_{,\sigma^B} \right), \quad (14)$$

$$V_E = \frac{M_{\text{Pl}}^4}{4U^2} V. \quad (15)$$

For the chosen function $U(\sigma^1, \sigma^2)$ the matrix K_{AB} is not diagonal:

$$\begin{aligned} K_{11} &= \frac{M_{\text{Pl}}^2}{2U^2} \left(U + \frac{1}{12} (\sigma^1)^2 \right), \\ K_{12} &= K_{21} = \frac{M_{\text{Pl}}^2}{24U^2} \sigma^1 \sigma^2, \\ K_{22} &= \frac{M_{\text{Pl}}^2}{2U^2} \left(U + \frac{1}{12} (\sigma^2)^2 \right). \end{aligned} \quad (16)$$

It is possible to diagonalize the matrix K_{AB} by introducing new fields χ^1 and χ^2 . In general, one has

$$K_{CD} \partial_\mu \sigma^C \partial_\nu \sigma^D = K_{CD} \frac{\partial \sigma^C}{\partial \chi^A} \frac{\partial \sigma^D}{\partial \chi^B} \partial_\mu \chi^A \partial_\nu \chi^B = G_{AB} \partial_\mu \chi^A \partial_\nu \chi^B,$$

where G_{AB} is the new ‘‘mass’’ matrix.

In our particular case, we take

$$\sigma^1 = \sqrt{12U_0} \tanh \frac{\chi^2}{\sqrt{6}M_{\text{Pl}}}, \quad \sigma^2 = \frac{\sqrt{12U_0}}{\cosh \frac{\chi^2}{\sqrt{6}M_{\text{Pl}}}} \tanh \frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}, \quad (17)$$

and get the following matrix G_{AB} :

$$\begin{aligned} G_{11} &= 1, \\ G_{12} &= G_{21} = 0, \\ G_{22} &= \cosh^2 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right). \end{aligned} \quad (18)$$

In terms of the new fields, the function U takes the form

$$U(\chi^1, \chi^2) = \frac{U_0}{\cosh^2\left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}\right) \cosh^2\left(\frac{\chi^2}{\sqrt{6}M_{\text{Pl}}}\right)}. \quad (19)$$

Action (13) can be rewritten as

$$S_{\text{E}} = \int d^4x \sqrt{-g^{(\text{E})}} \left(\frac{M_{\text{Pl}}^2}{2} R^{(\text{E})} - \frac{g^{(\text{E})\mu\nu}}{2} \left[\partial_\mu \chi^1 \partial_\nu \chi^1 + \cosh^2\left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}\right) \partial_\mu \chi^2 \partial_\nu \chi^2 \right] - V_{\text{E}} \right), \quad (20)$$

where

$$V_{\text{E}} = \frac{M_{\text{Pl}}^4 \left(\cosh \frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \cosh \frac{\chi^2}{\sqrt{6}M_{\text{Pl}}} \right)^2}{4U_0} \left(\frac{R_0}{2} + 144U_0 \left[\tanh \frac{\chi^2}{\sqrt{6}M_{\text{Pl}}} \right]^4 f \left(\frac{\tanh \frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}}{\sinh \frac{\chi^2}{\sqrt{6}M_{\text{Pl}}}} \right) \right). \quad (21)$$

By varying the action (20), we get the following equations:

$$R_{\mu\nu}^{(\text{E})} - \frac{1}{2} g_{\mu\nu}^{(\text{E})} R^{(\text{E})} = \frac{1}{M_{\text{Pl}}^2} T_{\mu\nu}^{\text{E}}, \quad (22)$$

$$\square \chi^1 - \frac{\sqrt{6}}{12M_{\text{Pl}}} \sinh\left(\frac{\chi^1}{3\sqrt{6}M_{\text{Pl}}}\right) g^{(\text{E})\mu\nu} \partial_\mu \chi^2 \partial_\nu \chi^2 - \frac{\partial V_{\text{E}}}{\partial \chi^1} = 0; \quad (23)$$

$$\cosh^2\left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}\right) \square \chi^2 + \frac{\sqrt{6}}{6M_{\text{Pl}}} \sinh\left(\frac{\chi^1}{3\sqrt{6}M_{\text{Pl}}}\right) g^{(\text{E})\mu\nu} \partial_\mu \chi^1 \partial_\nu \chi^2 - \frac{\partial V_{\text{E}}}{\partial \chi^2} = 0, \quad (24)$$

where

$$\begin{aligned} T_{\mu\nu}^{\text{E}} &= \partial_\mu \chi^1 \partial_\nu \chi^1 + \cosh^2\left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}\right) \partial_\mu \chi^2 \partial_\nu \chi^2 \\ &- g_{\mu\nu}^{(\text{E})} \left(\frac{1}{2} g^{(\text{E})\alpha\beta} \partial_\alpha \chi^1 \partial_\beta \chi^2 + \frac{1}{2} \cosh^2\left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}}\right) g^{(\text{E})\alpha\beta} \partial_\alpha \chi^2 \partial_\beta \chi^2 + V_{\text{E}}(\chi^1, \chi^2) \right). \end{aligned} \quad (25)$$

In the Einstein frame, the d'Alembert operator \square acting on a scalar is

$$\square = \frac{1}{\sqrt{g^{(\text{E})}}} \partial_\mu \left(\sqrt{g^{(\text{E})}} g^{(\text{E})\mu\nu} \partial_\nu \right). \quad (26)$$

IV. INTEGRABLE COSMOLOGICAL MODELS WITH NONMINIMAL COUPLING

A. The FLRW metric with conformal time

To get cosmological solutions in the analytic form, we consider the FLRW metric with conformal time

$$ds^2 = a^2(\tau) \left(-d\tau^2 + \frac{dr^2}{1 - Kr^2} + r^2 d\theta^2 + r^2 \sin^2(\theta) d\phi^2 \right), \quad (27)$$

where $a(\tau)$ is the scale function. A positive curvature index K is associated with a closed Universe, $K = 0$ with a flat Universe and a negative K with an open one.

In this metric, Eqs. (2) and (3) with $C_A = 1$ have the following form:

$$6U [h^2 + K] + 6h\dot{U} = \frac{1}{2} [(\dot{\sigma}^1)^2 + (\dot{\sigma}^2)^2] + a^2V, \quad (28)$$

$$U [2\dot{h} + h^2 + K] + \ddot{U} + H\dot{U} = \frac{1}{2}a^2V - \frac{1}{4} [(\dot{\sigma}^1)^2 + (\dot{\sigma}^2)^2], \quad (29)$$

$$\ddot{\sigma}^A + 2h\dot{\sigma}^A - 6U'_{,\sigma^A}a^2R + a^2V_{,\sigma^A} = 0, \quad (30)$$

where $h = \dot{a}/a$, and “dots” denote derivatives with respect to the parametric time τ .

The Ricci scalar is

$$R = \frac{6}{a^2} \left(\frac{\ddot{a}}{a} + K \right). \quad (31)$$

Integrating equation $R = R_0$, namely,

$$\ddot{a} + Ka = \frac{R_0}{6}a^3, \quad (32)$$

we obtain

$$\dot{a}^2 + Ka^2 = \frac{R_0}{12}a^4 + C, \quad (33)$$

where C is an integration constant.

Equation (33) has the general solution in terms of the Jacobi elliptic function:

$$a(\tau) = \frac{\sqrt{6C}}{\sqrt{3K + \sqrt{9K^2 - 3R_0C}}} \times \operatorname{sn} \left(\frac{1}{6} \sqrt{18K + 6\sqrt{9K^2 - 3R_0C}} (\tau - \tau_0) \left| \frac{\sqrt{R_0C (6K^2 - R_0C + 2K\sqrt{9K^2 - 3R_0C})}}{R_0C - 6K^2 - 2K\sqrt{9K^2 - 3R_0C}} \right. \right), \quad (34)$$

if $R_0 > 0$. For $R_0 = 0$, we get

$$a(\tau) = \begin{cases} a_1 \sin(\sqrt{K}\tau) + a_2 \cos(\sqrt{K}\tau), & K > 0; \\ a_1\tau + a_0, & K = 0; \\ a_1 \sinh(\sqrt{K}\tau) + a_2 \cosh(\sqrt{K}\tau), & K < 0; \end{cases} \quad (35)$$

where a_0 , a_1 and a_2 are arbitrary constants.

To find solutions for scalar fields, we substitute $\sigma^A = y^A/a$ into Eq. (30) and obtain a system

$$\ddot{y}^A + Ky^A + a^3 V_{,\phi^A} \left(\frac{y^1}{a}, \frac{y^2}{a} \right) = 0. \quad (36)$$

For V given by (11), we get the following system of two equations:

$$\ddot{y}^A + Ky^A + \frac{\partial \tilde{V}}{\partial y^A} = 0, \quad (37)$$

where $A = 1, 2$ and

$$\tilde{V}(y^1, y^2) = (y^1)^4 f\left(\frac{y^2}{y^1}\right).$$

System (37) has the first integral

$$\frac{1}{2}(\dot{y}^1)^2 + \frac{1}{2}(\dot{y}^2)^2 + \frac{K}{2}[(y^1)^2 + (y^2)^2] + \tilde{V} = E, \quad (38)$$

where E is an integration constant.

To analyze the cosmological evolution we consider equations in the cosmic time, which is defined by the equation $dt = a(\tau)d\tau$. Using

$$h = aH, \quad \dot{U} = a \frac{dU}{dt}, \quad \dot{y}^A = a^2 \frac{d\sigma^A}{dt} + a^2 H \sigma^A, \quad (39)$$

where H is the Hubble parameter, we get Eq. (28) in the following form:

$$6U \left[H^2 + \frac{K}{a^2} \right] + 6H \frac{dU}{dt} = \frac{1}{2} \left[\frac{d\sigma^1}{dt} \right]^2 + \frac{1}{2} \left[\frac{d\sigma^2}{dt} \right]^2 + V. \quad (40)$$

Using Eq. (38), we get from Eq. (40) the following equation:

$$6U_0 H^2 + 6U_0 \frac{K}{a^2} = \frac{1}{2} U_0 R_0 + \frac{E}{a^4}. \quad (41)$$

Equation (41) has the following solutions for $R_0 > 0$:

$$t - t_0 = \pm \frac{\sqrt{3}}{\sqrt{R_0}} \ln \left(\frac{R_0 U_0 a^2 - 6U_0 K}{\sqrt{R_0}} + \sqrt{R_0 U_0^2 a^4 - 12K U_0^2 a^2 + 2E U_0} \right). \quad (42)$$

At $R_0 = 0$, we obtain the following nonconstant solutions:

$$a(t) = \begin{cases} \pm \frac{\sqrt{K(E - 6K^2 U_0^3 (t - t_0)^2)}}{\sqrt{6U_0 K}}, & K \neq 0, \\ \pm \frac{\sqrt{3}}{3U_0} \sqrt{\pm \sqrt{6E U_0} U_0 (t - t_0)}, & K = 0. \end{cases} \quad (43)$$

Equation (41) coincides with the equation obtained in Ref. [9] for $K = 0$ and in Ref. [10] for an arbitrary K in the case of the corresponding single-field model. Thus, the analysis of the Universe

evolution in depending on the values of the constants K , R_0 and E , including conditions of the existence of bounce solutions given in Ref. [10], is also valid for the considered two-field models. For this reason, we do not repeat it in this paper. Note only that in the case $K = 0$, the Hubble parameter has the following simple form:

$$H(t) = \begin{cases} \frac{\sqrt{3R_0} \left(e^{2\sqrt{3R_0}t/3} + B \right)}{6 \left(e^{2\sqrt{3R_0}t/3} - B \right)}, & R_0 > 0 \\ \frac{1}{2(t - t_0)}, & R_0 = 0. \end{cases} \quad (44)$$

where B is a constant.

Note that explicit form of the $a(\tau)$ does not depend on the form of the potential and is given by Eq. (34) for $R_0 > 0$ and Eq. (35) for $R_0 = 0$. To get the general solutions in the conformal time, so it is suitable to write out expressions for $H(\tau)$ and $a(\tau)$ in explicitly real forms. We present formulas for the case of $K = 0$ and $R_0 > 0$ only.

For $K = 0$ and $R_0 > 0$, we obtain

1. if $E > 0$, then

$$H(\tau) = \sqrt{\frac{R_0}{3}} \frac{\operatorname{dn} \left(a_0 \sqrt{\frac{R_0}{3}} (\tau - \tau_0) \middle| \frac{1}{2} \right)}{1 - \operatorname{cn} \left(a_0 \sqrt{\frac{R_0}{3}} (\tau - \tau_0) \middle| \frac{1}{2} \right)}, \quad (45)$$

$$a(\tau) = \frac{a_0 \operatorname{sn} \left(a_0 \sqrt{\frac{R_0}{3}} (\tau - \tau_0) \middle| \frac{1}{2} \right)}{1 + \operatorname{cn} \left(a_0 \sqrt{\frac{R_0}{3}} (\tau - \tau_0) \middle| \frac{1}{2} \right)}, \quad a_0 = \left(\frac{2E}{U_0 R_0} \right)^{\frac{1}{4}}, \quad (46)$$

$$\tau(t) = \tau_0 + \frac{1}{a_0} \sqrt{\frac{3}{R_0}} F \left(\arccos \left(\frac{a_0^2 - a^2}{a_0^2 + a^2} \right) \middle| \frac{1}{2} \right),$$

$$a_0 \sqrt{\frac{R_0}{12}} (\tau - \tau_0) \in \left(0, K \left(\frac{1}{2} \right) \right);$$

2. if $E < 0$, then

$$a(\tau) = \frac{a_0}{\operatorname{cn} \left(a_0 \sqrt{\frac{R_0}{6}} (\tau - \tau_0) \middle| \frac{1}{2} \right)}, \quad a_0 = \left(\frac{-2E}{U_0 R_0} \right)^{\frac{1}{4}}, \quad (47)$$

$$H(\tau) = \sqrt{\frac{R_0}{6}} \operatorname{sn} \left(a_0 \sqrt{\frac{R_0}{6}} (\tau - \tau_0) \middle| \frac{1}{2} \right) \operatorname{dn} \left(a_0 \sqrt{\frac{R_0}{6}} (\tau - \tau_0) \middle| \frac{1}{2} \right), \quad (48)$$

$$\tau(t) = \tau_0 + \frac{1}{a_0} \sqrt{\frac{6}{R_0}} F \left(\arcsin \left(\sqrt{\frac{a^2 - a_0^2}{a^2}} \right) \middle| \frac{1}{2} \right),$$

$$a_0 \sqrt{\frac{R_0}{6}} (\tau - \tau_0) \in \left(-K \left(\frac{1}{2} \right), K \left(\frac{1}{2} \right) \right);$$

3. if $E = 0$, then

$$H(\tau) = \pm \sqrt{\frac{R_0}{12}}, \quad a(\tau) = \frac{a_0}{1 \mp a_0 \sqrt{R_0/12} (\tau - \tau_0)}, \quad (49)$$

$$\tau(t) = \tau_0 \pm \frac{1}{a_0} \sqrt{\frac{12}{R_0}} \left(1 - e^{\mp \sqrt{R_0/12} (t-t_0)} \right) = \tau_0 \pm \frac{1}{a_0} \sqrt{\frac{12}{R_0}} \left(1 - \frac{a_0}{a} \right),$$

$$\pm a_0 \sqrt{\frac{R_0}{12}} (\tau - \tau_0) \in (-\infty, 1).$$

Here F is the incomplete elliptic integral of the first kind, defined as

$$F(\varphi | m) = \int_0^\varphi \frac{d\theta}{\sqrt{1 - m \sin^2 \theta}}, \quad (50)$$

$K(m) \equiv F(\pi/2 | m)$ is the complete elliptic integral of the first kind, and the Jacobi elliptic functions sn , cn and dn have been used.

In the Einstein frame, the FLRW metric (27) becomes

$$ds^2 = a_{\text{E}}^2(\tau) \left(-d\tau^2 + \frac{dr^2}{1 - Kr^2} + r^2 d\theta^2 + r^2 \sin^2(\theta) d\varphi^2 \right), \quad (51)$$

where

$$a_{\text{E}}(\tau) = \sqrt{\frac{U}{U_0}} a(\tau). \quad (52)$$

V. POTENTIALS AND SOLUTIONS OF THE FIELD EQUATIONS

A. The way to get the general cosmological solution of the CCM

In the FLRW metric with the conformal time, Eqs. (22)–(24) have the following form:

$$3M_{\text{Pl}}^2 (h_{\text{E}}^2 + K) = \frac{1}{2} (\dot{\chi}^1)^2 + \frac{1}{2} G_{22} (\dot{\chi}^2)^2 + a_{\text{E}}^2 V_{\text{E}}, \quad (53)$$

$$-M_{\text{Pl}}^2 \left(2\dot{h}_{\text{E}} + h_{\text{E}}^2 + K \right) = \frac{1}{2} (\dot{\chi}^1)^2 + \frac{1}{2} G_{22} (\dot{\chi}^2)^2 - a_{\text{E}}^2 V_{\text{E}}, \quad (54)$$

$$\ddot{\chi}^1 + 2h_E \dot{\chi}^1 - \frac{1}{2} \frac{dG_{22}}{d\chi^1} (\dot{\chi}^2)^2 + a_E^2 \frac{\partial V_E}{\partial \chi^1} = 0, \quad (55)$$

$$G_{22} \ddot{\chi}_2 + 2h_E G_{22} \dot{\chi}_2 + \frac{dG_{22}}{d\chi^1} \dot{\chi}_1 \dot{\chi}_2 + a_E^2 \frac{\partial V_E}{\partial \chi^2} = 0. \quad (56)$$

Here “dots” denote $d/d\tau$, and $h_E \equiv \dot{a}_E/a_E = a_E H_E$.

Our goal is to get the general solutions of these equations for the potentials given by the formula (21). If we know the general solutions for the corresponding model in the Jordan frame, or, in other words, if the functions $a(\tau)$, $\sigma^1(\tau)$, and $\sigma^2(\tau)$ are known, then the scale function a_E is given by Eq. (52) and the Hubble parameter

$$H_E(\tau) = \frac{M_{\text{Pl}}}{\sqrt{2U(\sigma^1, \sigma^2)}} \left(H(\tau) + \frac{1}{2a(\tau)U(\sigma^1, \sigma^2)} \frac{dU}{d\tau} \right). \quad (57)$$

Here we use the fact that conformal times for both frames coincide (up to an additive constant, which can be set to be zero without loss of generality), since $dt/a = dt_E/a_E$. Having these expressions for a_E , H_E ,

To get $\chi^1(\tau)$ and $\chi^2(\tau)$, we use the relations

$$\chi^1 = \sqrt{6} M_{\text{Pl}} \operatorname{arctanh} \left(\frac{\sigma^2}{\sqrt{12U_0 - (\sigma^1)^2}} \right), \quad \chi^2 = \sqrt{6} M_{\text{Pl}} \operatorname{arctanh} \left(\frac{\sigma^1}{\sqrt{12U_0}} \right). \quad (58)$$

It is not always possible to integrate Eq. (37). To integrate field equations in the Jordan frame we must firstly choose a particular form of the potential V . Below we present the general solutions for three types of the potential V . The obtained solutions allow us to get the general solutions for the corresponding CCM.

B. Potential $V = \frac{1}{2} R_0 U_0$

We start from the case of a constant potential

$$V_0 = \frac{1}{2} R_0 U_0. \quad (59)$$

The corresponding potential in the Einstein frame is

$$V_E = \frac{M_{\text{Pl}}^4 R_0}{8U_0} \cosh^2 \left(\frac{\chi^1}{\sqrt{6} M_{\text{Pl}}} \right) \cosh^2 \left(\frac{\chi^2}{\sqrt{6} M_{\text{Pl}}} \right). \quad (60)$$

For $\tilde{V} \equiv 0$, Eq. (37) has the following solutions:

$$y^A = \begin{cases} B_1^A \sin(\sqrt{K}\tau) + B_0^A \cos(\sqrt{K}\tau), & K > 0; \\ B_1^A \tau + B_0^A, & K = 0; \\ B_1^A \sinh(\sqrt{K}\tau) + B_0^A \cosh(\sqrt{K}\tau), & K < 0; \end{cases} \quad (61)$$

where B_0^A and B_1^A are integration constants.

Using the expression (34), we get $\sigma^A(\tau)$. It allows us to get the general solution for the CCM with the potential (21) by Eqs. (52) and (58).

$$\text{C. Potential } V = \frac{1}{2}R_0U_0 + c_1 (\sigma^1)^4 + c_2 (\sigma^2)^4$$

Let us choose the potential V as

$$V_1(\sigma^1, \sigma^2) = \frac{R_0U_0}{2} + c_1 (\sigma^1)^4 + c_2 (\sigma^2)^4, \quad (62)$$

where c_1 and c_2 are constant.

For this potential V , system (37) is completely separable, and we have 2 independent integrals of motion:

$$\frac{1}{2} \left(\frac{dy^A}{d\tau} \right)^2 + \frac{K}{2} (y^A)^2 + c_A (y^A)^4 = E_A, \quad (63)$$

where E_A are integration constants.

The form of solutions of (63) depends on the signs of c_A and E_A . For $K = 0$, we obtain:

1. $E_A > 0, \quad c_A > 0$:

$$y^A(\tau) = \left(\frac{E_A}{c_A} \right)^{\frac{1}{4}} \text{cn} \left(\pm 2 (E_A c_A)^{\frac{1}{4}} (\tau - \tau_i) + \text{cn}^{-1} \left(u_n \left| \frac{1}{2} \right| \right) \left| \frac{1}{2} \right| \right); \quad (64)$$

2. $E_A < 0, \quad c_A < 0$:

$$y^A(\tau) = \frac{\left(\frac{E_A}{c_A} \right)^{\frac{1}{4}}}{\text{cn} \left(\pm 2 (E_A c_A)^{\frac{1}{4}} (\tau - \tau_i) + \text{cn}^{-1} \left(\frac{1}{u_A} \left| \frac{1}{2} \right| \right) \left| \frac{1}{2} \right| \right)}; \quad (65)$$

3. $E_A > 0, \quad c_A < 0$:

$$y^A(\tau) = \left| \frac{E_A}{c_A} \right|^{\frac{1}{4}} \frac{\text{sn} \left(\pm 2\sqrt{2} |E_A c_A|^{\frac{1}{4}} (\tau - \tau_i) + \text{sn}^{-1} \left(\frac{2u_A}{1+u_A^2} \left| \frac{1}{2} \right| \right) \left| \frac{1}{2} \right| \right)}{1 + \text{cn} \left(\pm 2\sqrt{2} |E_A c_A|^{\frac{1}{4}} (\tau - \tau_i) + \text{cn}^{-1} \left(\frac{1-u_A^2}{1+u_A^2} \left| \frac{1}{2} \right| \right) \left| \frac{1}{2} \right| \right)}. \quad (66)$$

Here $u_A = |c_A/E_A|^{1/4} y^A(\tau_i)$, and the choice of sign in “ \pm ” is determined by initial conditions.

Obviously, there are no solutions when $E_A < 0$ and $c_A > 0$.

For $c_A = 0$, E_A has to be non-negative, and we have $y^A(\tau)$ given by Eq. (61).

For the potential V given by (62), the corresponding potential V_E has the following form in terms of the new fields:

$$V_E = \frac{M_{\text{Pl}}^4}{4U_0^2} \left[\frac{1}{2} R_0 U_0 \cosh^4 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right) \cosh^4 \left(\frac{\chi^2}{\sqrt{6}M_{\text{Pl}}} \right) + (12U_0)^2 c_1 \cosh^4 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right) \sinh^4 \left(\frac{\chi^2}{\sqrt{6}M_{\text{Pl}}} \right) + (12U_0)^2 c_2 \sinh^4 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right) \right]. \quad (67)$$

If we put $\chi^2 = 0$, then V_E coincides with the potential from single-field model proposed by Bars and Chen in Ref. [5]:

$$V_{\text{BC}} = \frac{M_{\text{Pl}}^4}{4U_0^2} \left[\frac{1}{2} R_0 U_0 \cosh^4 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right) + (12U_0)^2 c_2 \sinh^4 \left(\frac{\chi^1}{\sqrt{6}M_{\text{Pl}}} \right) \right]. \quad (68)$$

So, the proposed integrable model with potential (67) is two-field generalization of the known single-field integrable model [5].

$$\mathbf{D. \quad Potential} \quad V = \frac{1}{2} R_0 U_0 + c \left((\sigma^1)^2 + (\sigma^2)^2 \right)^2$$

For the potential

$$V_2 = \frac{R U_0}{2} + c \left((\sigma^1)^2 + (\sigma^2)^2 \right)^2, \quad (69)$$

where c is a nonzero constant, general solutions can also be explicitly written in terms of the Jacobi elliptic functions. In this case, Eq. (38) with $K = 0$ takes the form

$$\frac{1}{2} \left(\frac{dy^1}{d\tau} \right)^2 + \frac{1}{2} \left(\frac{dy^2}{d\tau} \right)^2 + c \left((y^1)^2 + (y^2)^2 \right)^2 = E. \quad (70)$$

By introducing ‘‘polar coordinates’’,

$$y^1(\tau) = \rho(\tau) \cos(\theta(\tau)), \quad y^2(\tau) = \rho(\tau) \sin(\theta(\tau)), \quad (71)$$

it can be rewritten as

$$\frac{1}{2} \left(\frac{d\rho}{d\tau} \right)^2 + \frac{1}{2} \rho^2 \left(\frac{d\theta}{d\tau} \right)^2 + c\rho^4 = E. \quad (72)$$

In analogy with classical mechanics, one may observe that θ is a cyclic coordinate, which means that

$$\rho^2 \frac{d\theta}{d\tau} = L = \text{const}, \quad (73)$$

and the equation for ρ reads

$$\frac{1}{2} \left(\frac{d\rho}{d\tau} \right)^2 + \frac{L^2}{2\rho^2} + c\rho^4 = E. \quad (74)$$

In what follows, we restrict ourselves to the case of $c > 0$ only. In such a case, the solution for ρ reads

$$\rho(\tau) = \sqrt{\rho_0^2 - (\rho_0^2 - \rho_1^2) \operatorname{sn}^2 \left(\frac{\sqrt{v_0 - v_2}}{2\beta} \tau + C_\rho \left| \frac{v_0 - v_1}{v_0 - v_2} \right. \right)}. \quad (75)$$

Here, C_ρ is a constant of integration,

$$\rho_{0,1}^2 = 2\sqrt{\frac{E}{3c}} v_{0,1}, \quad v_k = \cos \left(\frac{1}{3} \arccos \left(- \left(\frac{E_{min}}{E} \right)^{3/2} \right) - \frac{2\pi k}{3} \right), \quad k = 0, 1, 2,$$

$$\beta = \frac{1}{4} \left(\frac{3}{cE} \right)^{1/4}, \quad E_{min} = \frac{3}{2} \left(\frac{cL^4}{2} \right)^{1/3}.$$

Note that since $E \geq E_{min}$, we always have $v_0 \geq v_1 > 0 > v_2$. The equality $v_0 = v_1$ holds when $E = E_{min}$. The solution for θ is

$$\theta(\tau) = \frac{\operatorname{sgn}(L)}{2v_0\sqrt{v_0 - v_2}} \left(\frac{E_{min}}{E} \right)^{3/4} \Pi \left(1 - \frac{v_1}{v_0}; \frac{\sqrt{v_0 - v_2}}{2\beta} \tau + C_\rho \left| \frac{v_0 - v_1}{v_0 - v_2} \right. \right) + C_\theta. \quad (76)$$

Here

$$\Pi(n; u | m) \equiv \int_0^u \frac{dw}{1 - n \operatorname{sn}^2(w | m)}$$

is the incomplete elliptic integral of the third kind, and C_θ is a constant of integration.

If $E = E_{min}$, then $\rho = \rho_{min} = (L^2/4c)^{1/6} = \text{const}$, and

$$\theta(\tau) = \frac{L}{\rho_{min}^2} \tau + C_\theta. \quad (77)$$

For the potential V_2 , the corresponding potential V_E takes the form

$$V_E = \frac{M_{\text{Pl}}^4}{4U_0^2} \left[\frac{1}{2} R_0 U_0 \cosh^4 \left(\frac{\chi^1}{\sqrt{6} M_{\text{Pl}}} \right) \cosh^4 \left(\frac{\chi^2}{\sqrt{6} M_{\text{Pl}}} \right) + (12U_0)^2 c \left(\cosh^2 \left(\frac{\chi^1}{\sqrt{6} M_{\text{Pl}}} \right) \sinh^2 \left(\frac{\chi^2}{\sqrt{6} M_{\text{Pl}}} \right) + \sinh^2 \left(\frac{\chi^1}{\sqrt{6} M_{\text{Pl}}} \right) \right)^2 \right]. \quad (78)$$

The general solution for the CCM with this potential can be found by Eqs. (52) and (58).

VI. CONCLUSIONS

In this paper, we have constructed new integrable chiral cosmological models with two-fields and potentials in terms of hyperbolic functions. These potentials are presented in Eqs. (60), (67), and (78). The general solutions have been found in terms of the Jacobi elliptic functions of the conformal time.

To get these solutions we have used the corresponding models in the Jordan frame, for which the Ricci scalar is an integral of motion. The integration of the initial system of equations reduces to the integration of system (37) of two second-order equations, that has one integral of motion for any potential V given by Eq. (11). In this paper, we present examples of the potential V , for which system (37) has two independent integrals of motion and, hence, is integrable.

Two-field CCMs actively investigated in cosmology to describe the evolution of the observable Universe, including inflation, primordial black hole formation and dark energy [42–49]. The constructed models can be considered as two-field generalizations of the single-field integrable model proposed in Ref. [5], which has been actively investigated as a possible description of the evolution of the early Universe. We hope that the proposed integrable models will be useful to describe the Universe evolution.

ACKNOWLEDGEMENTS

V.R.I. is supported by the “BASIS” Foundation grant No. 22-2-2-6-1.

REFERENCES

-
- [1] D. I. Kaiser, “Conformal Transformations with Multiple Scalar Fields,” *Phys. Rev. D* **81** (2010) 084044, [arXiv:1003.1159 \[gr-qc\]](#).
 - [2] S. V. Chervon, “On the chiral model of cosmological inflation,” *Russ. Phys. J.* **38** (1995) 539–543.
 - [3] D. S. Salopek and J. R. Bond, “Nonlinear evolution of long wavelength metric fluctuations in inflationary models,” *Phys. Rev. D* **42** (1990) 3936–3962.
 - [4] A. J. Maciejewski, M. Przybylska, T. Stachowiak, and M. Szydłowski, “Global integrability of cosmological scalar fields,” *J. Phys. A* **41** (2008) 465101, [arXiv:0803.2318 \[math-ph\]](#).
 - [5] I. Bars and S.-H. Chen, “The Big Bang and Inflation United by an Analytic Solution,” *Phys. Rev. D* **83** (2011) 043522, [arXiv:1004.0752 \[hep-th\]](#).
 - [6] P. Fré, A. Sagnotti, and A. S. Sorin, “Integrable Scalar Cosmologies I. Foundations and links with String Theory,” *Nucl. Phys. B* **877** (2013) 1028–1106, [arXiv:1307.1910 \[hep-th\]](#).
 - [7] P. Fré, A. S. Sorin, and M. Trigiante, “Integrable Scalar Cosmologies II. Can they fit into Gauged Extended Supergravity or be encoded in N=1 superpotentials?,” *Nucl. Phys. B* **881** (2014) 91–180, [arXiv:1310.5340 \[hep-th\]](#).

- [8] A. Y. Kamenshchik, E. O. Pozdeeva, A. Tronconi, G. Venturi, and S. Y. Vernov, “Integrable cosmological models with non-minimally coupled scalar fields,” *Class. Quant. Grav.* **31** (2014) 105003, [arXiv:1312.3540 \[hep-th\]](#).
- [9] B. Boisseau, H. Giacomini, D. Polarski, and A. A. Starobinsky, “Bouncing Universes in Scalar-Tensor Gravity Models admitting Negative Potentials,” *JCAP* **07** (2015) 002, [arXiv:1504.07927 \[gr-qc\]](#).
- [10] A. Y. Kamenshchik, E. O. Pozdeeva, A. Tronconi, G. Venturi, and S. Y. Vernov, “Interdependence between integrable cosmological models with minimal and non-minimal coupling,” *Class. Quant. Grav.* **33** no. 1, (2016) 015004, [arXiv:1509.00590 \[gr-qc\]](#).
- [11] A. Y. Kamenshchik, E. O. Pozdeeva, A. A. Starobinsky, A. Tronconi, G. Venturi, and S. Y. Vernov, “Induced gravity, and minimally and conformally coupled scalar fields in Bianchi-I cosmological models,” *Phys. Rev. D* **97** no. 2, (2018) 023536, [arXiv:1710.02681 \[gr-qc\]](#).
- [12] V. R. Ivanov and S. Y. Vernov, “Integrable cosmological models with an additional scalar field,” *Eur. Phys. J. C* **81** no. 11, (2021) 985, [arXiv:2108.10276 \[gr-qc\]](#).
- [13] V. R. Ivanov and S. Y. Vernov, “Anisotropic Solutions for R^2 Gravity Model with a Scalar Field,” *Phys. Atom. Nucl.* **86** no. 6, (2023) 1526–1532, [arXiv:2301.06836 \[gr-qc\]](#).
- [14] L. P. Chimento, “General solution to two-scalar field cosmologies with exponential potentials,” *Class. Quant. Grav.* **15** (1998) 965–974.
- [15] P. Christodoulidis, “General solutions to \mathcal{N} -field cosmology with exponential potentials,” *Eur. Phys. J. C* **81** no. 5, (2021) 471, [arXiv:1811.06456 \[astro-ph.CO\]](#).
- [16] A. Giacomini, E. González, G. Leon, and A. Paliathanasis, “Variational symmetries and superintegrability in multifield cosmology,” *Phys. Rev. D* **105** no. 4, (2022) 044010, [arXiv:2104.13649 \[gr-qc\]](#).
- [17] P. Christodoulidis and A. Paliathanasis, “ \mathcal{N} -field cosmology in hyperbolic field space: stability and general solutions,” *JCAP* **05** (2021) 038, [arXiv:2101.09582 \[gr-qc\]](#).
- [18] J. G. Russo, “New exact solutions in multi-scalar field cosmology,” *JCAP* **07** (2023) 066, [arXiv:2304.12360 \[gr-qc\]](#).
- [19] A. G. Muslimov, “On the Scalar Field Dynamics in a Spatially Flat Friedman Universe,” *Class. Quant. Grav.* **7** (1990) 231–237.
- [20] A. R. Liddle, P. Parsons, and J. D. Barrow, “Formalizing the slow roll approximation in inflation,” *Phys. Rev. D* **50** (1994) 7222–7232, [arXiv:astro-ph/9408015](#).
- [21] W. H. Kinney, “A Hamilton-Jacobi approach to nonslow roll inflation,” *Phys. Rev. D* **56** (1997) 2002–2009, [arXiv:hep-ph/9702427](#).
- [22] I. Y. Aref’eva, A. S. Koshelev, and S. Y. Vernov, “Exact solution in a string cosmological model,” *Theor. Math. Phys.* **148** (2006) 895–909, [arXiv:astro-ph/0412619](#).
- [23] D. Bazeia, C. B. Gomes, L. Losano, and R. Menezes, “First-order formalism and dark energy,” *Phys. Lett. B* **633** (2006) 415–419, [arXiv:astro-ph/0512197](#).

- [24] I. Y. Aref'eva, A. S. Koshelev, and S. Y. Vernov, "Crossing of the $w = -1$ barrier by D3-brane dark energy model," *Phys. Rev. D* **72** (2005) 064017, [arXiv:astro-ph/0507067](#).
- [25] K. Skenderis and P. K. Townsend, "Hamilton-Jacobi method for curved domain walls and cosmologies," *Phys. Rev. D* **74** (2006) 125008, [arXiv:hep-th/0609056](#).
- [26] S. Y. Vernov, "Construction of Exact Solutions in Two-Fields Models and the Crossing of the Cosmological Constant Barrier," *Teor. Mat. Fiz.* **155** (2008) 47–61, [arXiv:astro-ph/0612487](#).
- [27] P. K. Townsend, "Hamilton-Jacobi mechanics from pseudo-supersymmetry," *Class. Quant. Grav.* **25** (2008) 045017, [arXiv:0710.5178 \[hep-th\]](#).
- [28] I. Y. Aref'eva, N. V. Bulatov, and S. Y. Vernov, "Stable Exact Solutions in Cosmological Models with Two Scalar Fields," *Theor. Math. Phys.* **163** (2010) 788–803, [arXiv:0911.5105 \[hep-th\]](#).
- [29] A. Y. Kamenshchik, A. Tronconi, G. Venturi, and S. Y. Vernov, "Reconstruction of Scalar Potentials in Modified Gravity Models," *Phys. Rev. D* **87** no. 6, (2013) 063503, [arXiv:1211.6272 \[gr-qc\]](#).
- [30] T. Harko, F. S. N. Lobo, and M. K. Mak, "Arbitrary scalar field and quintessence cosmological models," *Eur. Phys. J. C* **74** (2014) 2784, [arXiv:1310.7167 \[gr-qc\]](#).
- [31] P. Binetruy, E. Kiritsis, J. Mabillard, M. Pieroni, and C. Rosset, "Universality classes for models of inflation," *JCAP* **04** (2015) 033, [arXiv:1407.0820 \[astro-ph.CO\]](#).
- [32] S. Y. Vernov, V. R. Ivanov, and E. O. Pozdeeva, "Superpotential Method for $F(R)$ Cosmological Models," *Phys. Part. Nucl.* **51** no. 4, (2020) 744–749, [arXiv:1912.07049 \[gr-qc\]](#).
- [33] S. V. Chervon, I. V. Fomin, E. O. Pozdeeva, M. Sami, and S. Y. Vernov, "Superpotential method for chiral cosmological models connected with modified gravity," *Phys. Rev. D* **100** no. 6, (2019) 063522, [arXiv:1904.11264 \[gr-qc\]](#).
- [34] I. Fomin and S. Chervon, "Exact and Slow-Roll Solutions for Exponential Power-Law Inflation Connected with Modified Gravity and Observational Constraints," *Universe* **6** no. 11, (2020) 199.
- [35] A. Y. Kamenshchik, E. O. Pozdeeva, A. Tribolet, A. Tronconi, G. Venturi, and S. Y. Vernov, "The Superpotential Method and the Amplification of Inflationary Perturbations," [arXiv:2406.19762 \[gr-qc\]](#).
- [36] A. Y. Kamenshchik, E. O. Pozdeeva, A. Tronconi, G. Venturi, and S. Y. Vernov, "Integrable cosmological models in the Einstein and in the Jordan frames and Bianchi-I cosmology," *Phys. Part. Nucl.* **49** no. 1, (2018) 1–4, [arXiv:1606.04260 \[gr-qc\]](#).
- [37] I. Bars, S.-H. Chen, P. J. Steinhardt, and N. Turok, "Complete Set of Homogeneous Isotropic Analytic Solutions in Scalar-Tensor Cosmology with Radiation and Curvature," *Phys. Rev. D* **86** (2012) 083542, [arXiv:1207.1940 \[hep-th\]](#).
- [38] I. Bars, S.-H. Chen, and N. Turok, "Geodesically Complete Analytic Solutions for a Cyclic Universe," *Phys. Rev. D* **84** (2011) 083513, [arXiv:1105.3606 \[hep-th\]](#).
- [39] I. Bars, S.-H. Chen, P. J. Steinhardt, and N. Turok, "Antigravity and the Big Crunch/Big Bang Transition," *Phys. Lett. B* **715** (2012) 278–281, [arXiv:1112.2470 \[hep-th\]](#).

- [40] V. R. Ivanov and S. Y. Vernov, “Integrable Cosmological Models with an Arbitrary Number of Scalar Fields,” [arXiv:2407.05002](#) [gr-qc].
- [41] I. Y. Aref’eva, N. V. Bulatov, R. V. Gorbachev, and S. Y. Vernov, “Non-minimally coupled cosmological models with the Higgs-like potentials and negative cosmological constant,” *Class. Quant. Grav.* **31** (2014) 065007, [arXiv:1206.2801](#) [gr-qc].
- [42] S. V. Chervon, “Chiral Cosmological Models: Dark Sector Fields Description,” *Quant. Matt.* **2** (2013) 71–82, [arXiv:1403.7452](#) [gr-qc].
- [43] G. K. Karananas and J. Rubio, “On the geometrical interpretation of scale-invariant models of inflation,” *Phys. Lett. B* **761** (2016) 223–228, [arXiv:1606.08848](#) [hep-ph].
- [44] P. Christodoulidis, D. Roest, and E. I. Sfakianakis, “Attractors, Bifurcations and Curvature in Multi-field Inflation,” *JCAP* **08** (2020) 006, [arXiv:1903.03513](#) [gr-qc].
- [45] A. Gundhi, S. V. Ketov, and C. F. Steinwachs, “Primordial black hole dark matter in dilaton-extended two-field Starobinsky inflation,” *Phys. Rev. D* **103** no. 8, (2021) 083518, [arXiv:2011.05999](#) [hep-th].
- [46] M. Braglia, D. K. Hazra, F. Finelli, G. F. Smoot, L. Sriramkumar, and A. A. Starobinsky, “Generating PBHs and small-scale GWs in two-field models of inflation,” *JCAP* **08** (2020) 001, [arXiv:2005.02895](#) [astro-ph.CO].
- [47] S. R. Geller, W. Qin, E. McDonough, and D. I. Kaiser, “Primordial black holes from multifield inflation with nonminimal couplings,” *Phys. Rev. D* **106** no. 6, (2022) 063535, [arXiv:2205.04471](#) [hep-th].
- [48] M. Braglia, A. Linde, R. Kallosh, and F. Finelli, “Hybrid α -attractors, primordial black holes and gravitational wave backgrounds,” *JCAP* **04** (2023) 033, [arXiv:2211.14262](#) [astro-ph.CO].
- [49] E. O. Pozdeeva and S. Y. Vernov, “Primordial Black Holes in Induced Gravity Inflationary Models with Two Scalar Fields,” [arXiv:2407.00999](#) [gr-qc].