# New Laboratory Constraints on Neutrinophilic Mediators

P. S. Bhupal Dev,<sup>1,\*</sup> Doojin Kim,<sup>2,†</sup> Deepak Sathyan,<sup>3,‡</sup> Kuver Sinha,<sup>4,§</sup> and Yongchao Zhang<sup>5,¶</sup>

<sup>1</sup>Department of Physics and McDonnell Center for the Space Sciences,

Washington University, St. Louis, MO 63130, USA

<sup>2</sup>Mitchell Institute for Fundamental Physics and Astronomy,

Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

<sup>3</sup>Maryland Center for Fundamental Physics, Department of Physics,

University of Maryland, College Park, MD 20742, USA

<sup>4</sup>Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019, USA

<sup>5</sup>School of Physics, Southeast University, Nanjing 211189, China

Neutrinophilic mediators are well-motivated messenger particles that can probe some of the least known sectors of fundamental physics involving nonstandard interactions of neutrinos with themselves and potentially with dark matter. In particular, light mediators coupling to the active neutrinos will induce new decay modes of the Standard Model mesons (e.g.,  $\pi^{\pm}, K^{\pm} \rightarrow \ell^{\pm} + \stackrel{(-)}{\nu} + \phi$ ), charged leptons (e.g.,  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu} + \phi$ ), and gauge bosons (e.g.,  $Z \rightarrow \nu + \bar{\nu} + \phi$ ). A common lore is that these decays suffer from infrared divergences in the limit of the vanishing mediator mass, i.e.,  $m_{\phi} \rightarrow 0$ . Here, we show for the first time that including the 1-loop contributions of these mediators to the standard 2-body decays (e.g.,  $\pi^{\pm}, K^{\pm} \rightarrow \ell^{\pm} + \stackrel{(-)}{\nu}$ , etc.), the infrared divergence from the 3-body decay cancels out exactly by virtue of the Kinoshita–Lee–Nauenberg theorem. Including these cancellation effects, we then update the existing laboratory constraints on neutrinophilic scalar mediators, thereby extending the limits far beyond the decaying parent particle mass and excluding a wider range of parameter space. These new "physical" limits derived here have significant implications for the future detection prospects of nonstandard neutrino (self-)interactions.

Introduction. – Neutrinos are the least understood out of the Standard Model (SM) particles. In particular, they can have potentially large nonstandard interactions and can serve as a portal to beyond the SM (BSM) physics. While nonstandard neutrino interactions with charged SM fermions are readily probed with neutrino scattering and oscillation experiments [1, 2], neutrino self-interactions [3] and possible connections to dark matter [4] can be effectively probed by studying the interactions of neutrinophilic mediators. In fact, it is common to have (light) scalar or vector bosons mediating the selfinteractions among active neutrinos in many BSM scenarios. For instance, a light leptonic scalar  $\phi$  can couple to neutrinos in the form of  $\phi \nu \bar{\nu}$  or  $\phi \nu \nu^c$  depending on the lepton number carried by  $\phi$ , which induces neutrino self-interactions and gives rise to interesting signals at both low-energy experiments [5-8] and high-energy colliders [9-11], as well as from astrophysical [12-18] and cosmological [19–25] observables; see Ref. [3] for a recent review. Another example of neutrinophilic mediators is the so-called Majoron particle J with interaction structure  $J\bar{\nu}i\gamma_5\nu$  originating from global symmetry breaking in seesaw models [26-28]. In some scenarios, the scalar might also couple to neutrino and dark (matter) particle  $\chi$ , e.g., in the form of  $\phi \bar{\nu} \chi$  [29–38]. Such couplings may contribute to neutrino self-interactions at the 1-loop level [39] or radiatively generate nontrivial electromagnetic properties of neutrinos [40]. There are also some seesaw models with a scalar coupling to the active neutrinos and heavy neutrino N via  $\phi N \nu$  [41–53].

Such nonstandard interactions of neutrinos via neu-



FIG. 1. Feynman diagrams for (a) the meson decay  $M^{\pm} \rightarrow \ell^{\pm} + {}^{(-)}_{\nu} + \phi$ , (b) the SM process  $M^{\pm} \rightarrow \ell^{\pm} + {}^{(-)}_{\nu}$ , and (c) the 1-loop correction due to the interaction in Eq. (1).

trinophilic mediators induce new decay modes of SM particles, e.g.,  $\pi^{\pm}, K^{\pm} \to \ell^{\pm} + {}^{(-)}_{\nu}/\chi + \phi$  (see Fig. 1a) and  $Z \rightarrow \nu + \bar{\nu}/\chi + \phi$  [6, 54, 55]. Then the corresponding experimental data, e.g., the decay widths and the spectra of charged leptons  $\ell^{\pm}$  from meson decays can be used to set limits on these decay rates as a function of the mediator mass  $m_{\phi}$ , as done by the PIENU [56] and NA62 [57] experiments using charged pion and kaon decays, respectively. However, it is a common lore that these decay channels are potentially subject to the infrared (IR) divergences; the corresponding partial widths approach infinity as  $m_{\phi} \rightarrow 0$  (see e.g., Refs. [6, 9, 58, 59]). This is clearly unphysical. We show that the IR divergence is removed by including the interference between the 1-loop contribution (Fig. 1c) and the 2-body decay (Fig. 1b). This is reminiscent of the standard calculations of quantum electrodynamics [60] (see also, e.g., Ref. [61]) and expected as a natural consequence of the Kinoshita–Lee–Nauenberg (KLN) theorem [62, 63].

For illustration purposes, we focus on the following

decay processes in this letter: exotic charged-meson decays  $\mathsf{M}^{\pm} \to \ell^{\pm} + \stackrel{(-)}{\nu} + \phi$  with  $\mathsf{M} = \pi, K$  and  $\ell = e, \mu$ , hadronic tau decays  $\tau^{\pm} \to \pi^{\pm} + \stackrel{(-)}{\nu} + \phi$ , and Z boson decays  $Z \to \nu + \bar{\nu} + \phi$ . More general cases involving dark matter  $\chi$  or heavy neutrino N in the final state such as  $\mathsf{M}^{\pm} \to \ell^{\pm} + \chi/N + \phi$  with nonzero mass  $m_{\chi/N}$  are also of great interest, e.g., for DM phenomenology and heavy neutrino searches, and will be reported in our forthcoming work [38].

We point out that summing up the tree and 1-loop contributions will not only give "physical" constraints on the associated decays in the IR limit of small  $\phi$  mass but also, in general, improve the constraints at large  $\phi$ mass. When the mediator is heavy, the tree-level process is kinematically suppressed or forbidden and the BSM effects are dominated by the virtual mediator in the loop. This has far-reaching implications for the experimental limits on  $m_{\phi}$  and its couplings.

**Meson decays.** – Let us first consider the meson decays  $M^{\pm} \rightarrow \ell^{\pm} + {}^{(-)}_{\nu} + \phi$  with the light scalar  $\phi$  emitted from the neutrino line (Fig. 1a). We consider the generic coupling in the form of

$$\mathcal{L} = g_{\nu} \phi \bar{\nu} \nu \,. \tag{1}$$

The couplings of  $\phi$  can be either flavor-diagonal or flavoroff-diagonal. Possible ultraviolet (UV) completions of this effective operator can be found in Refs. [6, 7, 9]. It is well known that the partial widths for the 2-body leptonic decays  $\mathsf{M}^{\pm} \to \ell^{\pm} + {}^{(\overline{\nu})}$  are helicity-suppressed in the SM, i.e., proportional to the charged-lepton mass squared  $m_{\ell}^2$ . This is not the case for the 3-body decay; see the Supplement for details. In the small  $m_{\phi}$  limit, the partial width  $\Gamma(\mathsf{M}^{\pm} \to \ell^{\pm} + {}^{(\overline{\nu})} + \phi)$  can be written as

$$\Gamma \simeq \frac{G_F^2 m_{\mathsf{M}}^3 f_{\mathsf{M}}^2 |V|^2 g_{\nu}^2}{128\pi^3} \left[ -x_{\ell\mathsf{M}} (1 - x_{\ell\mathsf{M}})^2 \log x_{\phi\mathsf{M}} + C_2(x_{\ell\mathsf{M}}) \right]$$
(2)

where  $G_F$  is the Fermi constant,  $m_{\rm M}$  and  $f_{\rm M}$  are respectively the charged-meson mass and decay constant, V is the CKM matrix element ( $V_{ud}$  for pions and  $V_{us}$ for kaons),  $x_{ab} \equiv m_a^2/m_b^2$ , and  $C_2(x_{\ell M})$  is a dimensionless function of the mass ratio  $x_{\ell M}$ , given in Eq. (S7) in the Supplement. It is apparent that the first term in Eq. (2) is IR-divergent, i.e., goes to infinity in the limit of  $m_{\phi} \to 0$ , or equivalently  $x_{\phi M} \to 0$ . For concreteness, we neglect the neutrino mass, which will not affect our results in the  $\phi$  mass range of interest here.

The coupling in Eq. (1) also induces a self-energy correction to the neutrino line, as shown in Fig. 1c. The amplitude  $\mathcal{M}^{(0)}$  of the tree-level SM decay  $\mathsf{M}^{\pm} \to \ell^{\pm} + {\overset{(-)}{\nu}}$ , shown in Fig. 1b, interferes with the amplitude  $\mathcal{M}^{(1)}$  for the 1-loop diagram, when we calculate the partial width  $\Delta\Gamma^{\text{loop}}(\mathsf{M}^{\pm} \to \ell^{\pm} + {\overset{(-)}{\nu}})$ . In particular, the interference  $\operatorname{term}$ 

$$\operatorname{Re}\left[\mathcal{M}^{(0)*}\mathcal{M}^{(1)}\right] \propto g_{\nu}^{2}, \qquad (3)$$

which is at the same order in  $g_{\nu}$  as the partial width in Eq. (2). The full expression for  $\Delta\Gamma^{\text{loop}}(\mathsf{M}^{\pm} \to \ell^{\pm} + \overset{(-)}{\nu})$  is given in Eq. (S8) of the Supplement. The IR-divergent part is  $+x_{\ell\mathsf{M}}(1-x_{\ell\mathsf{M}})^2 \log x_{\phi\mathsf{M}}$  that exactly cancels out the first term of Eq. (2), as expected from KLN theorem.

Taking into account the loop effects, we show in Fig. 2 the updated limits from  $\pi^{\pm}$  and  $K^{\pm}$  decays in the left and right panels, respectively. We first report conservative limits, taking 90% C.L. uncertainty ranges of the partial widths based on the information in the PDG data [64]; more details can be found in the Supplement. The blue and red lines are for the muon and electron decay modes, respectively.

In the case of  $\ell = e$ , the logarithmic divergence is heavily suppressed by  $x_{eM} \equiv m_e^2/m_{\pi,K}^2$ , hence the  $C_2$ term becomes more important in Eq. (2) for  $m_{\phi} \gtrsim eV$ . In other words, the IR divergence does not dominate the decay rate of  $M^{\pm} \rightarrow e^{\pm} + \stackrel{(-)}{\nu} + \phi$ . The limits on  $M^{\pm} \rightarrow e^{\pm} + \stackrel{(-)}{\nu} + \phi$  from the  $M^{\pm} \rightarrow e^{\pm} + \stackrel{(-)}{\nu}$  data (red lines) are flat even in the limit of  $m_{\phi} \rightarrow 0$ , showing little differences between the tree-level and tree+loop-level limits, as expected. When  $m_{\phi} \ll m_{\rm M}$ , the pion and kaon decay limits on  $g_{\nu}$  in the electron channel are respectively  $5.9 \times 10^{-3}$  and  $2.2 \times 10^{-3}$ .

By contrast, in the muon case  $\ell = \mu$ , as  $m_{\mu}$  is comparable to  $m_{\pi,K}$ , the divergent behavior is noticeable in the  $m_{\phi} \rightarrow 0$  limit, which is clear from the dashed blue lines in Fig. 2 (the label "tree"). The corresponding IRfree limits with the loop corrections are presented by the solid blue lines, labeled as "tree+loop". Notably, while the tree-level decay  $\mathsf{M}^{\pm} \to \ell^{\pm} + \overset{(-)}{\nu} + \phi$  is kinematically forbidden when  $m_{\phi} \geq m_{\mathsf{M}} - m_{\ell}$ , the 1-loop contribution to the decay  $M^{\pm} \rightarrow \ell^{\pm} + \tilde{\nu}$  still exists. Consequently, the solid blue lines in Fig. 2 can extend to large  $m_{\phi}$ , even beyond the parent particle mass  $m_{\rm M}$ , whereas the dashed blue lines quickly vanish as  $m_{\phi}$  gets closer to  $m_{\mathsf{M}} - m_{\mu}$ . With the loop contributions included,  $\pi^{\pm} \rightarrow \mu^{\pm} + \nu^{(-)} + \phi$ gives  $g_{\nu} < 0.061$  in the massless  $\phi$  limit. For heavy  $\phi$ , this decay constrains  $m_{\phi}$  up to ~ 14 GeV for  $g_{\nu} < 1$ . The IR limit for the kaon decay is relatively weaker, i.e.,  $q_{\nu} < 0.53$  is allowed, while the UV limit is ~ 15 GeV for  $g_{\nu} < 1$ . The dip feature of the solid blue line at around 200 MeV in the right panel of Fig. 2 is due to the substantial cancellation of the tree and 1-loop contributions.

These limits can be further improved by a dedicated shape analysis of the final decay products. The PIENU experiment has provided limits on the branching ratio (BR) of  $\pi^{\pm} \rightarrow e^{\pm}/\mu^{\pm} + \stackrel{(-)}{\nu} + X$  as a function of the invisible X mass [56]. We reinterpret them with our updated partial width calculations in both electron and muon channels, which are shown respectively by the or-



FIG. 2. Limits on meson decays  $\pi^{\pm}$ ,  $K^{\pm} \to e^{\pm}/\mu^{\pm} + {}^{(-)}\nu^{+} + \phi$  at 90% C.L., in the plane of  $m_{\phi}$  and  $g_{\nu}$ . The dashed lines are the limits with only the tree-level contribution, while the solid lines are for those including the 1-loop contributions (cf. Fig. 1). The red and blue lines are respectively for the  $g_{\nu_e}$  and  $g_{\nu_{\mu}}$  couplings. The purple and orange lines are respectively the corresponding limits from PIENU (left) [56] and NA62 (right) [57]. The gray-shaded regions are excluded by other constraints [3]. See the text for more details.

ange and purple lines in the left panel of Fig. 2. In the electron channel, we find that the limit of  $g_{\nu}$  is improved by ~ 13% for small  $m_{\phi}$ . In contrast, the loop-included result (solid) differs from the tree-level one (dashed) in the muon channel. Qualitatively, when  $m_{\phi}$  approaches the kinematic threshold,  $m_{\pi} - m_{\mu} \simeq 34$  MeV, the tree-level contribution is highly suppressed by phase space and the loop contribution becomes the dominant BSM effect. Therefore, the "tree+loop" limit gets much stronger at  $m_{\phi} \gtrsim 10$  MeV. For  $m_{\phi} \rightarrow 0$ , the PIENU limit on  $g_{\nu_{\mu}}$  is 0.029, while at  $m_{\phi} \sim 10$  MeV, it is improved to  $7.2 \times 10^{-3}$ .

Similarly, the NA62 experiment has reported the shape-analysis-based limits on BR( $K^{\pm} \rightarrow \mu^{\pm} + \stackrel{(-)}{\nu} + X$ ) with X being a scalar [57]. They are of the order of  $\mathcal{O}(10^{-6})$  for 10 MeV  $< m_X < 370$  MeV, roughly three orders of magnitude stronger than the limits from the partial widths above. The resultant "tree" and "tree+loop" limits are shown respectively by the dashed and solid purple lines in the right panel of Fig. 2. Again, with the 1-loop contribution included, the NA62 limits get much stronger, especially when  $m_{\phi}$  is close to the threshold; at  $m_{\phi} = 370$  MeV, the limit can reach down to  $2.1 \times 10^{-3}$ .

We note that due to the existence of an off-shell neutrino propagator in the 3-body decay, the energy/angular distribution of the charged lepton from meson decays might be (mildly) affected, and the corresponding limits should be interpreted accordingly; see e.g., Fig. 7 of Ref. [57]. Therefore, more dedicated analyses of the PIENU and NA62 data may improve to some extent the limits on  $m_{\phi}$  and  $g_{\nu}$  reported here. We will examine this aspect in future work.

Speaking of the existing limits, all the gray-shaded regions in both panels of Fig. 2 show the exclusions by current terrestrial, astrophysical, and cosmological data [3], i.e., those from the cosmic microwave background [20, 65], big bang nucleosynthesis [23], SN1987A [66, 67], IceCube High Energy Starting Events [17], the highenergy neutrinos detected by IceCube from the blazar TXS 0506+056 [16], and double-beta decays (only for  $\nu_e$ ) [68]. Other existing limits, e.g., those from stellar cooling [69], are relatively weaker for the parameter space of our interest and hence not shown in Fig. 2. Moreover, the coupling in Eq. (1) induces 1-loop couplings of  $\phi$  to the quarks and charged leptons [26], which would give additional limits from neutrino-electron and neutrinonucleus scattering [70], e.g., those from Borexino [71] and COHERENT [72]. However, they are highly suppressed by the loop factor and the heavy W and Z particles in the loop, and are therefore not shown here.

Finally, as natural extensions, we have also calculated the partial widths for other cases and the corresponding meson decay limits: (i) The scalar is replaced by a pseudoscalar J with couplings to neutrinos in the form of  $J\bar{\nu}i\gamma_5\nu$ ; we find that the results are the same as the scalar case above. (ii) The scalar couples to the charged leptons, i.e.,  $g_{\ell}\phi\bar{\ell}\ell$  or  $g_{\ell}\phi\bar{\ell}i\gamma_5\ell$  (with  $\ell=e,\mu$ ) [58, 59, 73], where the same cancellation happens. However, such couplings contribute to the anomalous magnetic moments of electron [74] and muon [75], which give rise to more stringent limits than the meson decay limits under consideration [76]. Therefore we do not pursue this case further. (iii) The analysis above can also be applied to the charged D meson decays, i.e.,  $D^{\pm} \rightarrow \ell^{\pm} + \overset{(-)}{\nu} + \phi$ , and also to the semileptonic B-meson decays. However, the corresponding limits are weaker than the ones shown here [6, 9]. (iv) If the scalar  $\phi$  is replaced by a vector boson Z', the corresponding partial width  $\Gamma(\mathsf{M}^{\pm} \to \ell^{\pm} + \overset{(-)}{\nu} + Z')$ is dominated by the term  $m_{\rm M}^4/m_{Z'}^2$  originating from the longitudinal polarization of Z'; this is much larger



FIG. 3. Limits on tau decay  $\tau^{\pm} \rightarrow \pi^{\pm} + {}^{(-)}_{\nu} + \phi$  at 90% C.L. in the plane of  $m_{\phi}$  and  $g_{\nu,\tau}$ . The red and blue lines are respectively for  $g_{\nu}$  and  $g_{\tau}$ . The existing limit from SN1987A is relevant only to  $g_{\nu}$ . The other notations are the same as in Fig. 2.

than the IR divergent term  $m_{\ell}^2 \log(m_{Z'}^2/m_{\rm M}^2)$ , see e.g., Refs. [58, 59, 70, 77, 78]. We delve into this intriguing case in forthcoming work [38].

**Tau decays.** – One of the dominant tau-lepton decay channels in the SM is  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu}$ , which is closely related to the charged-pion decays  $\pi^{\pm} \rightarrow \ell^{\pm} + \stackrel{(-)}{\nu}$ . Calculations of the decay channel  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu} + \phi$  are similar to those for the charged-meson decays, and the details are given in the Supplement. The resultant "tree" and "tree+loop" limits on  $m_{\phi}$  and  $g_{\nu}$  estimated with the  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu}$  partial width measurement are presented in Fig. 3 respectively by the dashed and solid red lines. The most important limit is from SN1987A [66, 67], shown by the gray-shaded region.

Here we also consider the case of  $\phi$  coupling to  $\tau$  via  $g_{\tau}\phi\tau^{+}\tau^{-}$ . The corresponding limits from  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu}$  are shown by the (dashed) blue lines. The existing limits on  $g_{\tau}$  are much weaker, mainly from the measurement of the anomalous  $\tau$  magnetic moment. The current ATLAS constraint of  $-0.057 < a_{\tau} < 0.024$  [79] leads to the exclusion bound of  $g_{\tau} > 1.1$  [76], and is out of the presentation range in Fig. 3.

We find that, once the 1-loop contribution is included, the allowed values of  $g_{\nu,\tau}$  are smaller than respectively 0.48 and 0.23 in the  $m_{\phi} \rightarrow 0$  limit. For  $g_{\nu,\tau} < 1$ ,  $m_{\phi}$  is constrained up to 22 GeV and 58 GeV, respectively, which are well beyond the  $\tau$  mass and the existing SN1987A limit.

The pure leptonic decay channel  $\tau^{\pm} \rightarrow \ell^{\pm} + \nu_{\ell} + \nu_{\tau}$ (with  $\ell = e, \mu$ ) can also be used to set limits on exotic decays, i.e.,  $\tau^{\pm} \rightarrow \ell^{\pm} + \nu_{\ell} + \nu_{\tau} + \phi$  with  $\phi$  emitted from the neutrino or charged-lepton lines [8, 55]. However, with respect to the 3-body decay  $\tau^{\pm} \rightarrow \pi^{\pm} + {}^{(-)}_{\nu} + \phi$  considered above, these 4-body leptonic decays are both phase-space



FIG. 4. Limits on the decay  $Z \rightarrow \nu + \bar{\nu} + \phi$  (orange) at 90% C.L. in the plane of  $m_{\phi}$  and  $g_{\nu}$ . The brown line shows the LHC Run-3 prospect (for  $\nu_{\mu}$ ). The other notations are the same as in Fig. 2.

and BR-suppressed. Similarly, we expect that the 4-body decay channels from muon, i.e.,  $\mu^{\pm} \rightarrow e^{\pm} + \nu_e + \nu_{\mu} + \phi$ , will not give competitive limits. Nevertheless, we will examine these 4-body decays in future work [38] for completeness.

Z boson decays. – The invisible Z decay data can be utilized to set limits on our neutrinophilic mediator  $\phi$ through the  $Z \to \nu + \bar{\nu} + \phi$  decay channel. The calculational details are given in the Supplement. Just like in our previous cases, the tree-level contribution shows IR divergence which is removed by including the 1-loop contributions. But here the 1-loop corrections come from the neutrino self-energy, as well as from the  $Z\nu\bar{\nu}$  vertex, unlike the meson and tau cases above. While we observe this cancellation even with (almost) massless neutrinos, it is interesting to compare our findings with the results in Ref. [8], where the IR divergence was regulated by the neutrino mass, which becomes relevant only in the regime  $m_{\phi} \lesssim m_{\nu}$ . Our result is more general in this sense. The orange lines in Fig. 4 show the resulting limits from invisible Z data;  $g_{\nu} < 1.4$  is constrained for  $m_{\phi} \ll m_Z$  with 1-loop contributions included. Due to the cancellation of the tree and loop contributions, the "tree+loop" limit gets much weaker at around  $m_{\phi} \sim 30 \text{ GeV}$  (as shown by the "gap").

The current limit mainly comes from SN1987A [66, 67] within the presentation range, indicated by the grayshaded region in Fig. 4. The search prospect of  $\phi$  at the large hadron collider (LHC) Run-3 in the  $W^{\pm} \rightarrow \mu^{\pm} +$ MET channel with an integrated luminosity of 300 fb<sup>-1</sup> and 0.1% systematics is shown by the brown line [11]. It is clear from the solid orange lines in Fig. 4 that, for the  $\phi$  mediator, the invisible Z-decay data have excluded a sizable range of parameter space beyond the SN1987A limit, complementing the prospect at the LHC.

One can also derive limits from the exotic W boson decays, i.e.,  $W^{\pm} \rightarrow \ell^{\pm} + \overset{(-)}{\nu} + \phi$ , and the calculations

are very similar to the Z boson case. However, the uncertainty  $\Delta BR(W^{\pm} \rightarrow \ell^{\pm} + \overset{(-)}{\nu}) \simeq 3.6 \times 10^{-3}$ , is much larger than that from the invisible Z data,  $7.3 \times 10^{-4}$  [64]. The resulting exclusions on the coupling are  $g_{\nu} > 1$  and are thus not shown in Fig. 4.

The invisible Z data can also be used for other rare Z decay channels; e.g.,  $Z \rightarrow \nu + \bar{\nu} + Z'$  with a neutrinophilic vector mediator Z'. This certainly carries nontrivial physics implications and features in mitigating the associated IR divergence, which are quite different from the case of  $M^{\pm} \rightarrow \ell^{\pm} + {}^{(-)}_{\nu} + Z'$ . We will defer the detailed discussion of the vector case for future publication [38].

**Discussions and conclusions.** – In this letter, we have studied the exotic decays of charged mesons, tau lepton, and Z gauge boson in the presence of a (light) neutrinophilic scalar  $\phi$ . We particularly focused on the IR divergence arising in the  $m_{\phi} \rightarrow 0$  limit which is shown to be removed with the 1-loop contributions included. The methodology here can also be applied to other decay channels, e.g.,  $\pi^0 \rightarrow \gamma + \gamma + \phi$  with  $\phi$  coupling to photons or the channel  $\pi^{\pm} \rightarrow e^{\pm} + \stackrel{(-)}{\nu} + a$  with pseudoscalar *a* coupling to the *W* mediator or the valence quarks of  $\pi^{\pm}$  [80–82]. One may also constrain the hadronic couplings of a (light) scalar from meson and tau decays, e.g., in the channel of  $\tau^{\pm} \rightarrow \pi^{\pm} + \stackrel{(-)}{\nu} + \phi$  with  $\phi$  coupling to  $\pi^{\pm}$  instead of  $\stackrel{(-)}{\nu}$  or  $\tau^{\pm}$ .

Several implications of the 1-loop corrections are worth mentioning. (i) The 1-loop contributions are important in not only removing the IR divergence but also, in general, bringing new limits in the region of parameter space that is kinematically "forbidden" to constrain at the tree level (see the solid lines in Figs. 2 through 4). (ii) When conducting similar phenomenological studies, one should carefully include loop contributions to perform the theory calculations more accurately and place experimental bounds more robustly without the unphysical IR divergence. (iii) Some past experimental limits should be revisited accordingly, e.g., the PIENU and NA62 limits re-interpreted in our study.

In summary, the SM should be IR-finite, as stated by the KLN theorem. This holds even in the presence of BSM couplings. We have demonstrated this general feature with a scalar  $\phi$  interacting with the active neutrinos and  $\tau^{\pm}$ .

**Acknowledgments.**– We thank Kaladi Babu, Bhaskar Dutta, Sudip Jana, Lorenzo Ricci, and Oleksandr Tomalak for useful discussions and comments on the draft. BD is supported by the U.S. Department of Energy grant No. DE-SC 0017987. The work of DK is supported by the DOE Grant No. DE-SC0010813. DS is supported by NSF Grant No. PHY-2210361 and by the Maryland Center for Fundamental Physics. KS is supported by the U.S. Department of Energy grant DE-SC0009956. YZ is supported by the National Natural Science Foundation of China under grant No. 12175039, the 2021 Jiangsu Shuangchuang (Mass Innovation and Entrepreneurship) Talent Program No. JSSCBS20210144, and the "Fundamental Research Funds for the Central Universities". BD, DK, and KS acknowledge the Center for Theoretical Underground Physics and Related Areas (CETUP\* 2024) and the Institute for Underground Science at SURF for hospitality and for providing a stimulating environment, where this work was finalized.

- \* bdev@wustl.edu
- <sup>†</sup> doojin.kim@tamu.edu
- <sup>‡</sup> dsathyan@umd.edu
- <sup>§</sup> kuver.sinha@ou.edu
- <sup>¶</sup> zhangyongchao@seu.edu.cn
- Y. Farzan and M. Tortola, Neutrino oscillations and Non-Standard Interactions, Front. in Phys. 6 (2018) 10, [1710.09360].
- [2] P. S. B. Dev et al., Neutrino Non-Standard Interactions: A Status Report, SciPost Phys. Proc. 2 (2019) 001, [1907.00991].
- [3] J. M. Berryman et al., Neutrino self-interactions: A white paper, Phys. Dark Univ. 42 (2023) 101267, [2203.01955].
- [4] M. Blennow, E. Fernandez-Martinez, A. Olivares-Del Campo, S. Pascoli, S. Rosauro-Alcaraz and A. V. Titov, *Neutrino Portals to Dark Matter*, *Eur. Phys. J. C* 79 (2019) 555, [1903.00006].
- [5] P. S. Pasquini and O. L. G. Peres, Bounds on Neutrino-Scalar Yukawa Coupling, Phys. Rev. D 93 (2016) 053007,
   [1511.01811]. [Erratum: Phys.Rev.D 93, 079902 (2016)].
- [6] J. M. Berryman, A. De Gouvêa, K. J. Kelly and Y. Zhang, Lepton-Number-Charged Scalars and Neutrino Beamstrahlung, Phys. Rev. D 97 (2018) 075030, [1802.00009].
- [7] K. J. Kelly and Y. Zhang, Mononeutrino at DUNE: New Signals from Neutrinophilic Thermal Dark Matter, Phys. Rev. D 99 (2019) 055034, [1901.01259].
- [8] V. Brdar, M. Lindner, S. Vogl and X.-J. Xu, *Revisiting neutrino self-interaction constraints from Z and τ decays*, *Phys. Rev. D* 101 (2020) 115001, [2003.05339].
- [9] A. de Gouvêa, P. S. B. Dev, B. Dutta, T. Ghosh, T. Han and Y. Zhang, *Leptonic Scalars at the LHC*, *JHEP* 07 (2020) 142, [1910.01132].
- [10] P. S. B. Dev, B. Dutta, T. Ghosh, T. Han, H. Qin and Y. Zhang, Leptonic scalars and collider signatures in a UV-complete model, JHEP 03 (2022) 068, [2109.04490].
- [11] K. Agashe, S. Airen, R. Franceschini, D. Kim, A. V. Kotwal, L. Ricci et al., "Unification" of BSM Searches and SM Measurements: the case of lepton+MET and m<sub>W</sub>, 2404.17574.
- K. C. Y. Ng and J. F. Beacom, Cosmic neutrino cascades from secret neutrino interactions, Phys. Rev. D 90 (2014) 065035, [1404.2288]. [Erratum: Phys.Rev.D 90, 089904 (2014)].
- [13] I. M. Shoemaker and K. Murase, Probing BSM Neutrino Physics with Flavor and Spectral Distortions: Prospects

for Future High-Energy Neutrino Telescopes, Phys. Rev. D 93 (2016) 085004, [1512.07228].

- [14] L. Heurtier and Y. Zhang, Supernova Constraints on Massive (Pseudo)Scalar Coupling to Neutrinos, JCAP 02 (2017) 042, [1609.05882].
- [15] A. Das, A. Dighe and M. Sen, New effects of non-standard self-interactions of neutrinos in a supernova, JCAP 05 (2017) 051, [1705.00468].
- [16] K. J. Kelly and P. A. N. Machado, Multimessenger Astronomy and New Neutrino Physics, JCAP 10 (2018) 048, [1808.02889].
- [17] M. Bustamante, C. Rosenstrøm, S. Shalgar and I. Tamborra, Bounds on secret neutrino interactions from highenergy astrophysical neutrinos, Phys. Rev. D 101 (2020) 123024, [2001.04994].
- [18] I. Esteban, S. Pandey, V. Brdar and J. F. Beacom, Probing secret interactions of astrophysical neutrinos in the high-statistics era, Phys. Rev. D 104 (2021) 123014, [2107.13568].
- [19] F.-Y. Cyr-Racine and K. Sigurdson, *Limits on Neutrino-Neutrino Scattering in the Early Universe*, *Phys. Rev. D* 90 (2014) 123533, [1306.1536].
- [20] M. Archidiacono and S. Hannestad, Updated constraints on non-standard neutrino interactions from Planck, JCAP 07 (2014) 046, [1311.3873].
- [21] G.-y. Huang, T. Ohlsson and S. Zhou, Observational Constraints on Secret Neutrino Interactions from Big Bang Nucleosynthesis, Phys. Rev. D 97 (2018) 075009, [1712.04792].
- [22] G. Barenboim, P. B. Denton and I. M. Oldengott, Constraints on inflation with an extended neutrino sector, Phys. Rev. D 99 (2019) 083515, [1903.02036].
- [23] N. Blinov, K. J. Kelly, G. Z. Krnjaic and S. D. McDermott, Constraining the Self-Interacting Neutrino Interpretation of the Hubble Tension, Phys. Rev. Lett. 123 (2019) 191102, [1905.02727].
- [24] K.-F. Lyu, E. Stamou and L.-T. Wang, Self-interacting neutrinos: Solution to Hubble tension versus experimental constraints, Phys. Rev. D 103 (2021) 015004, [2004.10868].
- [25] A. Das and S. Ghosh, Flavor-specific interaction favors strong neutrino self-coupling in the early universe, JCAP 07 (2021) 038, [2011.12315].
- [26] Y. Chikashige, R. N. Mohapatra and R. D. Peccei, Are There Real Goldstone Bosons Associated with Broken Lepton Number?, Phys. Lett. B 98 (1981) 265–268.
- [27] G. B. Gelmini and M. Roncadelli, Left-Handed Neutrino Mass Scale and Spontaneously Broken Lepton Number, Phys. Lett. B 99 (1981) 411–415.
- [28] J. Schechter and J. W. F. Valle, Neutrino Decay and Spontaneous Violation of Lepton Number, Phys. Rev. D 25 (1982) 774.
- [29] C. Boehm and P. Fayet, Scalar dark matter candidates, Nucl. Phys. B 683 (2004) 219–263, [hep-ph/0305261].
- [30] J. Barranco, O. G. Miranda, C. A. Moura, T. I. Rashba and F. Rossi-Torres, *Confusing the extragalactic neutrino* flux limit with a neutrino propagation limit, *JCAP* 10 (2011) 007, [1012.2476].
- [31] Y. Farzan, S. Pascoli and M. A. Schmidt, *Recipes and Ingredients for Neutrino Mass at Loop Level*, *JHEP* 03 (2013) 107, [1208.2732].
- [32] A. Olivares-Del Campo, C. Bœhm, S. Palomares-Ruiz and S. Pascoli, Dark matter-neutrino interactions through the lens of their cosmological implications, Phys.

Rev. D 97 (2018) 075039, [1711.05283].

- [33] S. Ghosh, R. Khatri and T. S. Roy, Dark neutrino interactions make gravitational waves blue, Phys. Rev. D 97 (2018) 063529, [1711.09929].
- [34] C. Hagedorn, J. Herrero-García, E. Molinaro and M. A. Schmidt, *Phenomenology of the Generalised Scotogenic* Model with Fermionic Dark Matter, JHEP 11 (2018) 103, [1804.04117].
- [35] M. Hufnagel and X.-J. Xu, Dark matter produced from neutrinos, JCAP 01 (2022) 043, [2110.09883].
- [36] C. Alvarado, C. Bonilla, J. Leite and J. W. F. Valle, Phenomenology of fermion dark matter as neutrino mass mediator with gauged B-L, Phys. Lett. B 817 (2021) 136292, [2102.07216].
- [37] J. Herms, S. Jana, V. P. K. and S. Saad, Light neutrinophilic dark matter from a scotogenic model, Phys. Lett. B 845 (2023) 138167, [2307.15760].
- [38] P. S. B. Dev, D. Kim, D. Sathyan, K. Sinha and Y. Zhang, "New Constraints on Neutrino-Dark Matter Interactions: A Comprehensive Analysis." in preparation, 2024.
- [39] I. Bischer, W. Rodejohann and X.-J. Xu, Loop-induced Neutrino Non-Standard Interactions, JHEP 10 (2018) 096, [1807.08102].
- [40] G. Herrera and I. M. Shoemaker, Dark Neutrino Moments From Light Loops, 2406.08663.
- [41] E. Ma, Pathways to naturally small neutrino masses, Phys. Rev. Lett. 81 (1998) 1171–1174, [hep-ph/9805219].
- [42] E. Ma, Verifiable radiative seesaw mechanism of neutrino mass and dark matter, Phys. Rev. D 73 (2006) 077301, [hep-ph/0601225].
- [43] C. Boehm, Y. Farzan, T. Hambye, S. Palomares-Ruiz and S. Pascoli, Is it possible to explain neutrino masses with scalar dark matter?, Phys. Rev. D 77 (2008) 043516, [hep-ph/0612228].
- [44] T. Hambye, F. S. Ling, L. Lopez Honorez and J. Rocher, Scalar Multiplet Dark Matter, JHEP 07 (2009) 090, [0903.4010]. [Erratum: JHEP 05, 066 (2010)].
- [45] Y. Farzan, S. Pascoli and M. A. Schmidt, AMEND: A model explaining neutrino masses and dark matter testable at the LHC and MEG, JHEP 10 (2010) 111, [1005.5323].
- [46] A. Ibarra and C. Simonetto, Understanding neutrino properties from decoupling right-handed neutrinos and extra Higgs doublets, JHEP 11 (2011) 022, [1107.2386].
- [47] Y. Farzan and E. Ma, Dirac neutrino mass generation from dark matter, Phys. Rev. D 86 (2012) 033007, [1204.4890].
- [48] Y. Farzan and S. Palomares-Ruiz, Dips in the Diffuse Supernova Neutrino Background, JCAP 06 (2014) 014, [1401.7019].
- [49] T. Li and J. Liao, Loop effect in the coherent neutrinonucleus scattering, JHEP 02 (2021) 099, [2008.00743].
- [50] M. Chianese, B. Fu and S. F. King, Interplay between neutrino and gravity portals for FIMP dark matter, JCAP 01 (2021) 034, [2009.01847].
- [51] S.-F. Ge, P. Pasquini and J. Sheng, Solar neutrino scattering with electron into massive sterile neutrino, Phys. Lett. B 810 (2020) 135787, [2006.16069].
- [52] S.-F. Ge, P. Pasquini and J. Sheng, Solar active-sterile neutrino conversion with atomic effects at dark matter direct detection experiments, JHEP 05 (2022) 088, [2112.05560].
- [53] M. Chianese, B. Fu and S. F. King, Dark Matter

in the Type Ib Seesaw Model, JHEP **05** (2021) 129, [2102.07780].

- [54] V. D. Barger, W.-Y. Keung and S. Pakvasa, Majoron Emission by Neutrinos, Phys. Rev. D 25 (1982) 907.
- [55] A. P. Lessa and O. L. G. Peres, Revising limits on neutrino-Majoron couplings, Phys. Rev. D 75 (2007) 094001, [hep-ph/0701068].
- [56] PIENU collaboration, A. Aguilar-Arevalo et al., Search for three body pion decays  $\pi^+ \rightarrow l^+ \nu X$ , Phys. Rev. D 103 (2021) 052006, [2101.07381].
- [57] NA62 collaboration, E. Cortina Gil et al., Search for K<sup>+</sup> decays to a muon and invisible particles, Phys. Lett. B 816 (2021) 136259, [2101.12304].
- [58] C. E. Carlson and B. C. Rislow, New Physics and the Proton Radius Problem, Phys. Rev. D 86 (2012) 035013, [1206.3587].
- [59] B. Dutta, D. Kim, A. Thompson, R. T. Thornton and R. G. Van de Water, Solutions to the MiniBooNE Anomaly from New Physics in Charged Meson Decays, Phys. Rev. Lett. **129** (2022) 111803, [2110.11944].
- [60] F. Bloch and A. Nordsieck, Note on the Radiation Field of the electron, Phys. Rev. 52 (1937) 54–59.
- [61] M. E. Peskin and D. V. Schroeder, An Introduction to quantum field theory. Addison-Wesley, Reading, USA, 1995, 10.1201/9780429503559.
- [62] T. Kinoshita, Mass singularities of Feynman amplitudes, J. Math. Phys. 3 (1962) 650–677.
- [63] T. D. Lee and M. Nauenberg, Degenerate Systems and Mass Singularities, Phys. Rev. 133 (1964) B1549–B1562.
- [64] PARTICLE DATA GROUP collaboration, R. L. Workman et al., *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [65] M. Escudero and S. J. Witte, A CMB search for the neutrino mass mechanism and its relation to the Hubble tension, Eur. Phys. J. C 80 (2020) 294, [1909.04044].
- [66] E. W. Kolb and M. S. Turner, Supernova SN 1987a and the Secret Interactions of Neutrinos, Phys. Rev. D 36 (1987) 2895.
- [67] S. Shalgar, I. Tamborra and M. Bustamante, Corecollapse supernovae stymic secret neutrino interactions, Phys. Rev. D 103 (2021) 123008, [1912.09115].
- [68] T. Brune and H. Päs, Massive Majorons and constraints on the Majoron-neutrino coupling, Phys. Rev. D 99 (2019) 096005, [1808.08158].
- [69] M. Escudero, D. Hooper, G. Krnjaic and M. Pierre, Cosmology with A Very Light  $L_{\mu} L_{\tau}$  Gauge Boson, JHEP 03 (2019) 071, [1901.02010].

- [70] R. Laha, B. Dasgupta and J. F. Beacom, Constraints on New Neutrino Interactions via Light Abelian Vector Bosons, Phys. Rev. D 89 (2014) 093025, [1304.3460].
- [71] BOREXINO collaboration, G. Alimonti et al., The Borexino detector at the Laboratori Nazionali del Gran Sasso, Nucl. Instrum. Meth. A 600 (2009) 568–593, [0806.2400].
- [72] COHERENT collaboration, D. Akimov et al., Observation of Coherent Elastic Neutrino-Nucleus Scattering, Science 357 (2017) 1123–1126, [1708.01294].
- [73] G. Krnjaic, G. Marques-Tavares, D. Redigolo and K. Tobioka, Probing Muonphilic Force Carriers and Dark Matter at Kaon Factories, Phys. Rev. Lett. **124** (2020) 041802, [1902.07715].
- [74] X. Fan, T. G. Myers, B. A. D. Sukra and G. Gabrielse, Measurement of the Electron Magnetic Moment, Phys. Rev. Lett. 130 (2023) 071801, [2209.13084].
- [75] MUON G-2 collaboration, D. P. Aguillard et al., Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm, Phys. Rev. Lett. 131 (2023) 161802, [2308.06230].
- [76] M. Lindner, M. Platscher and F. S. Queiroz, A Call for New Physics : The Muon Anomalous Magnetic Moment and Lepton Flavor Violation, Phys. Rept. 731 (2018) 1– 82, [1610.06587].
- [77] V. Barger, C.-W. Chiang, W.-Y. Keung and D. Marfatia, Constraint on parity-violating muonic forces, Phys. Rev. Lett. 108 (2012) 081802, [1109.6652].
- [78] P. Bakhti and Y. Farzan, Constraining secret gauge interactions of neutrinos by meson decays, Phys. Rev. D 95 (2017) 095008, [1702.04187].
- [79] ATLAS collaboration, G. Aad et al., Observation of the γγ→ττ Process in Pb+Pb Collisions and Constraints on the τ-Lepton Anomalous Magnetic Moment with the ATLAS Detector, Phys. Rev. Lett. 131 (2023) 151802, [2204.13478].
- [80] W. A. Bardeen, R. D. Peccei and T. Yanagida, CON-STRAINTS ON VARIANT AXION MODELS, Nucl. Phys. B 279 (1987) 401–428.
- [81] W. Altmannshofer, S. Gori and D. J. Robinson, Constraining axionlike particles from rare pion decays, Phys. Rev. D 101 (2020) 075002, [1909.00005].
- [82] M. Bauer, M. Neubert, S. Renner, M. Schnubel and A. Thamm, *Flavor probes of axion-like particles*, *JHEP* 09 (2022) 056, [2110.10698].

# Supplemental Material

#### A Data used for the limits

Here we collect in Table S1 all the data used in the Letter for the meson, tau, and Z decay limits. For instance, from the  $\pi \to e\nu$  data in the table, the  $1\sigma$  range uncertainty is

$$\Delta\Gamma(\pi \to e\nu) = \frac{\mathrm{BR}(\pi \to e\nu)}{\tau_{\pi^{\pm}}} \left[ \frac{\Delta\tau_{\pi^{\pm}}}{\tau_{\pi^{\pm}}} + \frac{\Delta\mathrm{BR}(\pi \to e\nu)}{\mathrm{BR}(\pi \to e\nu)} \right].$$
 (S1)

To apply for the limits at the 90% C.L., we multiply the  $1\sigma$  uncertainties by a factor of 1.64.

TABLE S1. Data used for the limits, with  $\tau$  (or  $\Gamma$ ) being the lifetime (or width) of the decaying parent particle, BR the corresponding BR for the channels in the first column, and  $\Delta\Gamma$  the  $1\sigma$  uncertainties for the partial widths. The last column is the experimental limits from the spectrum analysis of the charged leptons involved. Taken from PDG [64] unless otherwise specified.

Channel	Data	$1\sigma$ limits	Relevant expt.
$\pi \to e \nu$	$\tau = (2.6033 \pm 0.0005) \times 10^{-8} \text{ sec}$ BR = (1.230 ± 0.004) × 10 <sup>-4</sup>	$\Delta \Gamma = 1.07 \times 10^{-20} \text{ MeV}$	PIENU [56]
$\pi  ightarrow \mu  u$	$\tau = (2.6033 \pm 0.0005) \times 10^{-8} \text{ sec}$ BR = 0.9998770 ± 0.0000004	$\Delta \Gamma = 4.85 \times 10^{-18} \text{ MeV}$	PIENU [56]
$K \to e \nu$	$\tau = (1.2380 \pm 0.0020) \times 10^{-8} \text{ sec}$ BR = (1.582 ± 0.007) × 10 <sup>-5</sup>	$\Delta \Gamma = 5.06 \times 10^{-21} \ {\rm MeV}$	
$K \to \mu \nu$	$\tau = (1.2380 \pm 0.0020) \times 10^{-8} \text{ sec}$ BR = 0.6356 ± 0.0011	$\Delta \Gamma = 1.13 \times 10^{-16} \ {\rm MeV}$	NA62 [57]
$ au  o \pi  u$	$\tau = (2.093 \pm 0.005) \times 10^{-13} \text{ sec}$ BR = 0.1082 ± 0.0005	$\Delta \Gamma = 2.38 \times 10^{-12} \text{ MeV}$	
$Z  o \nu \bar{\nu}$	$\Gamma = (2.4955 \pm 0.0023) \text{ GeV} BR = 0.20000 \pm 0.00055$	$\Delta \Gamma = 1.83~{\rm MeV}$	

### **B** Meson decay $M \rightarrow \ell + \nu + \phi$

For the meson decay  $\mathsf{M}(p) \to \ell(p_\ell) + \nu(p_\nu) + \phi(p_\phi)$ , with  $\phi$  coupling to neutrinos with the strength  $g_\nu$ , the squared amplitude is given by

$$\sum |\mathcal{M}(\mathsf{M} \to \ell + \nu + \phi)|^2 = \frac{8g_{\nu}^2 f_{\mathsf{M}}^2 G_F^2 |V|^2}{q^4} \left\{ q^4 (p_{\ell} \cdot p_{\nu}) + m_{\ell}^2 \Big[ 2(q \cdot p_{\nu})(q^2 + (q \cdot p_{\ell})) - q^2(p_{\ell} \cdot p_{\nu}) \Big] \right\},$$
(S2)

with q being the momentum of the neutrino mediator. After a lengthy calculation, one can find the total partial width to be

$$\Gamma(\mathsf{M} \to \ell + \nu + \phi) = \frac{g_{\nu}^2 G_F^2 m_\mathsf{M}^3 f_\mathsf{M}^2 |V|^2}{128\pi^3} f_1(x_{\phi\mathsf{M}}, x_{\ell\mathsf{M}}), \qquad (S3)$$

with  $x_{ab} \equiv m_a^2/m_b^2$ , and the dimensionless function  $f_1(x_1, x_2)$  is given by

$$f_{1}(x_{1}, x_{2}) = \frac{1}{3} \frac{\lambda^{1/2}(1, x_{1}, x_{2})}{1 - x_{2}} \left( 1 + 10x_{1} + x_{1}^{2} - x_{2}(9 - 6x_{1} + x_{1}^{2}) + x_{2}^{2}(18 - 19x_{1}) - 10x_{2}^{3} \right) \\ + \left( 2x_{1}(1 + x_{1}) - x_{2}\left(1 - 3x_{1}^{2}\right) - 2x_{2}^{2}(1 - 3x_{1}) + x_{2}^{3} \right) \arctan \frac{\lambda^{1/2}(1, x_{1}, x_{2})}{1 - x_{1} + x_{2}} \\ - \left[ x_{1}(1 - 3x_{2}^{2}) - x_{2}(1 - x_{2})^{2} + (2 - x_{2} + 4x_{2}^{2} - 3x_{2}^{3}) \frac{x_{1}^{2}}{(1 - x_{2})^{2}} \right] \operatorname{arctanh} \frac{(1 - x_{2})\lambda^{1/2}(1, x_{1}, x_{2})}{(1 - x_{2})^{2} + x_{1}(1 + x_{2})}$$
(S4)

where

$$\lambda(a, b, c) \equiv a^2 + b^2 + c^2 - 2ab - 2ac - 2bc.$$
(S5)

For sufficiently small  $x_1$ ,

$$f_1(x_1, x_2) \simeq -x_2(1 + 2x_2 - x_2^2) \operatorname{arctanh} \frac{1 - x_2}{1 + x_2} + \frac{1}{6}(1 - x_2) \left[ 2 - 4x_2(4 - 5x_2) - 3x_2(1 - x_2) \log \frac{x_1^2 x_2}{(1 - x_2)^4} \right].$$
(S6)

Then we get the IR-divergent part, i.e., the first term in Eq. (2), and the rest of  $f_1(x_1, x_2)$  is the finite function

$$C_{2}(x_{2}) \simeq -x_{2}(1+2x_{2}-x_{2}^{2})\operatorname{arctanh}\frac{1-x_{2}}{1+x_{2}} + \frac{1}{6}(1-x_{2})\left[2-4x_{2}(4-5x_{2})-3x_{2}(1-x_{2})\frac{x_{2}}{(1-x_{2})^{4}}\right].$$
(S7)

For interference term between Figs. 1 (b) and (c), its contribution to the width of the decay  $M \rightarrow \ell + \nu$  is

$$\Delta\Gamma^{\text{loop}}(\mathsf{M} \to \ell + \nu) = -\frac{g_{\nu}^2 G_F^2 m_\mathsf{M} m_{\ell}^2 f_\mathsf{M}^2 |V|^2}{128\pi^3} \left(1 - \frac{m_{\ell}^2}{m_\mathsf{M}^2}\right)^2 f_1^{\text{loop}}(x_{\phi\mathsf{M}}, x_{\ell\mathsf{M}}), \tag{S8}$$

where we have removed the UV divergence, which can always be done by adding counterterms in a UV-complete theory, and the dimensionless function

$$f_1^{\text{loop}}(x_1, x_2) = \frac{5}{2} - \log \frac{x_1(1-x_2)^2}{16\pi^2}, \qquad (S9)$$

leading to the IR term proportional to  $x_{\ell M}(1 - x_{\ell M})^2 \log x_{\phi M}$ , which cancels exactly with the first term in Eq. (2) from the 3-body decay  $M \rightarrow \ell + \nu + \phi$ , as expected.

## $\mathbf{C} \quad \ \mathbf{Tau \ decay} \ \tau \to \pi + \nu + \phi$

In the SM, the matrix element for the semileptonic decay  $\tau \to \pi + \nu$  is closely correlated with that for  $\pi \to \ell + \nu$ . For the case of  $\phi$  coupling to the neutrino, we can easily obtain the squared amplitude for the decay  $\tau(p) \to \pi(p_{\pi}) + \nu(p_{\nu}) + \phi(p_{\phi})$ :

$$\frac{1}{2}\sum |\mathcal{M}_{\nu}(\tau \to \pi + \nu + \phi)|^2 = \frac{4g_{\nu}^2 f_{\pi}^2 G_F^2 |V_{ud}|^2}{q^4} \left\{ q^4(p \cdot p_{\nu}) + m_{\tau}^2 \Big[ 2(q \cdot p_{\nu})((q \cdot p) - q^2) - q^2(p \cdot p_{\nu}) \Big] \right\},$$
(S10)

where the factor of 1/2 is for averaging over the spins of tau in the initial state. Then the partial width reads

$$\Gamma(\tau \to \pi + \nu + \phi) = \frac{g_{\nu}^2 G_F^2 m_{\tau}^3 f_{\pi}^2 |V_{ud}|^2}{256\pi^3} f_2(x_{\phi\tau}, x_{\pi\tau}), \qquad (S11)$$

with the dimensionless function

$$f_{2}(x_{1}, x_{2}) = -\frac{\lambda^{1/2}(1, x_{1}, x_{2})}{3} \left[ (10 + x_{1}^{2} - 8x_{2} + x_{2}^{2}) + \frac{x_{1}}{1 - x_{2}} (19 - 6x_{2} - 10x_{2}^{2}) \right] \\ - \left( 1 - x_{2}(2 + x_{2}) + 2x_{1} \left( 3 + x_{2}^{2} \right) + x_{1}^{2} (3 + 2x_{2}) \right) \operatorname{arctanh} \frac{\lambda^{1/2}(1, x_{1}, x_{2})}{1 - x_{1} + x_{2}} \\ + \left[ (1 - x_{2})^{2} + 2x_{1}(3 - x_{2}^{2}) + \frac{x_{1}^{2}}{(1 - x_{2})^{2}} \left( 3 - x_{2}(4 - x_{2} + 2x_{2}^{2}) \right) \right] \operatorname{arctanh} \frac{(1 - x_{2})\lambda^{1/2}(1, x_{1}, x_{2})}{(1 - x_{2})^{2} + x_{1}(1 + x_{2})}.$$
(S12)

In the limit of  $x_1 \to 0$ , we have

$$f_2(x_1, x_2) \simeq -\frac{1}{3}(1 - x_2)(10 - 8x_2 + x_2^2) - (1 - x_2)^2 \log \frac{x_1}{(1 - x_2)^2} - x_2^2 \log x_2.$$
(S13)

The loop contribution to the decay  $\tau \to \pi + \nu$  is

$$\Delta\Gamma^{\text{loop}}(\tau \to \pi + \nu) = -\frac{g_{\nu}^2 G_F^2 m_{\tau}^3 f_{\pi}^2 |V_{ud}|^2}{256\pi^3} \left(1 - \frac{m_{\pi}^2}{m_{\tau}^2}\right)^2 f_1^{\text{loop}}(x_{\phi\tau}, x_{\pi\tau}).$$
(S14)

For the case of  $\phi$  coupling to the tau with the strength  $g_{\tau}$ , the amplitude square is very similar to Eq. (S10):

$$\frac{1}{2}\sum |\mathcal{M}_{\tau}(\tau \to \pi + \nu + \phi)|^2 = \frac{4g_{\nu}^2 f_{\pi}^2 G_F^2 |V_{ud}|^2}{(q^2 - m_{\tau}^2)^2} \left\{ q^4(p \cdot p_{\nu}) + m_{\tau}^2 \Big[ 2(q \cdot p_{\nu})((q \cdot p) + q^2) - q^2(p \cdot p_{\nu}) \Big] \right\},$$
(S15)

with q being the momentum of the tau propagator. For the partial width, one only needs to replace the coupling  $g_{\nu}$ 

by  $g_{\tau}$  in Eq. (S11), and the corresponding dimensionless function is

$$f_{3}(x_{1}, x_{2}) = -\frac{\lambda^{1/2}(1, x_{1}, x_{2})}{6} \Big[ 119 - 115x_{2} + 2x_{2}^{2} - x_{1}(61 - 26x_{2}) + 2x_{1}^{2} \Big] \\ + \sqrt{x_{1}(4 - x_{1})}(1 - x_{2}) \Big( 2(9 - x_{2}) - x_{1}(3 + x_{2}) \Big) \left[ \frac{\pi}{2} + \arctan \frac{x_{1}(3 - x_{1} + x_{2})\lambda^{-1/2}(1, x_{1}, x_{2})}{\sqrt{x_{1}(4 - x_{1})}} \right] \\ + \Big( 8 - 16x_{2} + 7x_{2}^{2} - 2x_{1} \left( 18 - 12x_{2} + x_{2}^{2} \right) + x_{1}^{2} (3 - 2x_{2}) \Big) \arctan \frac{\lambda^{1/2}(1, x_{1}, x_{2})}{1 - x_{1} + x_{2}} \\ + x_{2}^{2}(1 - x_{1})^{2} \operatorname{arctanh} \frac{(1 - x_{1})\lambda^{1/2}(1, x_{1}, x_{2})}{(1 - x_{1})^{2} - x_{2}(1 + x_{1})}.$$
(S16)

In the limit of  $x_1 \to 0$ ,

$$f_3(x_1, x_2) \simeq -\frac{1}{6}(1 - x_2)(119 - 115x_2 + 2x_2^2) - 4(1 - x_2)^2 \log \frac{x_1}{(1 - x_2)^2} - x_2^2 \log x_2.$$
(S17)

The loop contribution is the same as in Eq. (S14) with  $g_{\nu}$  replaced by  $g_{\tau}$  and multiplied by a factor of 4.

### **D** Z boson decay $Z \rightarrow \nu + \bar{\nu} + \phi$

For the decay  $Z(p) \rightarrow \nu(p_{\nu}) + \bar{\nu}(p_{\bar{\nu}}) + \phi(p_{\phi})$ , the amplitude square is

$$\frac{1}{3} \sum |\mathcal{M}(Z \to \nu + \bar{\nu} + \phi)|^2 = \frac{2\sqrt{2}g_{\nu}^2 m_Z^2 G_F}{3q^4} \left[ 2(q_1 \cdot p_{\nu})(q_1 \cdot p_{\bar{\nu}}) + 2(q_2 \cdot p_{\nu})(q_2 \cdot p_{\bar{\nu}}) - (q_1^2 + q_2^2)(p_{\nu} \cdot p_{\bar{\nu}}) + \frac{2}{m_Z^2} \left( 2(p \cdot p_{\bar{\nu}})(p \cdot q_1)(q_1 \cdot p_{\nu}) + 2(p \cdot p_{\nu})(p \cdot q_2)(q_2 \cdot p_{\bar{\nu}}) - (q_1^2 + q_2^2)(p \cdot p_{\nu})(p \cdot p_{\bar{\nu}}) \right) \right],$$
(S18)

where the factor of 1/3 is for averaging the spins of Z boson, and  $q_{1,2} = p_{\phi} + p_{\nu,\bar{\nu}}$  is the momentum of the (anti)neutrino propagator. Then the corresponding partial decay width reads

$$\Gamma(Z \to \nu \bar{\nu} \phi) = \frac{g_{\nu}^2 G_F m_Z^3}{96\sqrt{2}\pi^3} f_4(x_{\phi Z}) \,, \tag{S19}$$

with the dimensionless function

$$f_4(x) = -\frac{1}{6}(1-x)(17+8x-x^2) - (1+3x)\log x.$$
(S20)

The dimensionless function for the corresponding loop contribution to  $Z \to \nu \bar{\nu}$  is, with the prefactors the same as in Eq. (S19):

$$f_{2}^{\text{loop}}(x) = -\frac{3}{2} + \log x + 2(1 - \log x) - \frac{2 + 3x + 2x \log x}{(1 + x)} - x(3 + 2\log x) \log\left(\frac{x}{1 + x}\right) \\ + \frac{1}{1 + x} {}_{2} f_{1}^{(0,0,1,0)} \left(1, 1, 3, -\frac{1}{x}\right) + \frac{2}{x} {}_{2} f_{1}^{(0,0,1,0)} \left(1, 2, 3, -\frac{1}{x}\right) + \frac{1}{x} {}_{2} f_{1}^{(0,1,0,0)} \left(1, 2, 3, -\frac{1}{x}\right) \\ + 2 \int_{0}^{1} db \left[ {}_{2} F_{1}^{(0,1,0,0)} \left(1, 0, 2, \frac{b}{x(1 - b)}\right) - {}_{2} F_{1}^{(0,0,1,0)} \left(1, 0, 2, \frac{b}{x(1 - b)}\right) \right],$$
(S21)

where  $_{2}f_{1}$  and  $_{2}F_{1}$  are respectively the hyper-geometric and regularized hyper-geometric functions.