

State-dependent mobility edge in kinetically constrained models

Manthan Badbaria,^{1,*} Nicola Pancotti,^{2,3} Rajeev Singh,¹ Jamir Marino,⁴ and Riccardo J. Valencia-Tortora⁴

¹*Department of Physics, Indian Institute of Technology (Banaras Hindu University), Varanasi 221005, India*

²*AWS Center for Quantum Computing, Pasadena, CA 91125, USA*

³*California Institute of Technology, Pasadena, CA, USA*

⁴*Institut für Physik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany*

(Dated: July 19, 2024)

In this work, we show that the kinetically constrained quantum East model lies between a quantum scarred and a many-body localized system featuring an unconventional type of mobility edge in the spectrum. We name this scenario *state-dependent* mobility edge: while the system does not exhibit a sharp separation in energy between thermal and non-thermal eigenstates, the abundance of non-thermal eigenstates results in slow entanglement growth for *many* initial states, such as product states, below a finite energy density. We characterize the state-dependent mobility edge by looking at the complexity of classically simulating dynamics using tensor network for system sizes well beyond those accessible via exact diagonalization. Focusing on initial product states, we observe a qualitative change in the dynamics of the bond dimension needed as a function of their energy density. Specifically, the bond dimension typically grows *polynomially* in time up to a certain energy density, where we locate the state-dependent mobility edge, enabling simulations for long times. Above this energy density, the bond dimension typically grows *exponentially* making the simulation practically unfeasible beyond short times, as generally expected in interacting theories. We correlate the polynomial growth of the bond dimension to the presence of many non-thermal eigenstates around that energy density, a subset of which we compute via tensor network.

In the era of noisy intermediate scale quantum (NISQ) devices [1], a natural task that could display quantum advantage over classical computers is the simulation of quantum many-body dynamics itself [2–5]. Indeed, the simulation of quantum systems is typically expected to be hard for the same reason quantum computers are believed to be powerful, namely entanglement. However, displaying provable quantum advantage is a challenging task in itself due to the noisy nature of current quantum computers [6–11]. Indeed, while quantum computers are developed so are also classical algorithm, making the quantum advantage effectively a moving target. Even more surprisingly, also assuming the ability to implement large-scale error-correcting protocols and possess a fault-tolerant quantum computer, there are strongly interacting quantum systems whose simulation is not guaranteed to be more efficient using a quantum computer over a classical one. Indeed, there exist systems whose properties make them amenable to being efficiently simulated on a classical computer. Examples include Hamiltonians exhibiting non-thermal behavior at finite energy density, e.g. area-law excited eigenstates, such as quantum many-body scarred systems [12–20] and many-body localized systems (MBL) systems [21, 22], or mechanisms hindering the propagation of quantum correlations, such as dynamical confinement [23–28]. Indeed, as these systems are characterized by a slow growth of entanglement, typically considered a measure of ‘classical complexity,’ they are amenable to be efficiently simulated using classical algorithms (e.g., via tensor networks) with modest computational resources up to long times [29–31]. The presence of exceptions

to the naively expected quantum advantage in computing dynamics raises an intriguing possibility: can non-thermal features be diagnosed and characterized by examining how efficiently dynamics can be computed using classical algorithms?

To this aim, we focus on one-dimensional non-integrable systems. A key point of our approach is the choice of an appropriate candidate for efficient classical simulations susceptible to the complexity due to entanglement. Tensor-network methods fulfill such a requirement. Indeed, the resources required—the bond dimension—typically scale exponentially with time in far-from-equilibrium dynamics since finite-energy volume-law eigenstates of the Hamiltonian participate predominantly in the dynamics [32]. However, the presence of non-thermal eigenstates of the Hamiltonian could significantly reduce complexity, enabling efficient simulations for ‘long’ times [29–31]. Having selected a classical method, we need to define complexity. We categorize a computational task as ‘hard’ (or ‘easy’) if the required computational resources scale (sub-)exponentially. We distinguish complexity along the space and time domain. For the time domain, we adopt the bond dimension χ of the matrix product state representation of the evolved states as a measure of complexity. We choose χ as it encodes the actual computational time needed in performing operations, such as dynamics, in tensor network methods. Moreover, χ is closely linked to entanglement, widely regarded as an indicator of ‘hardness’ in representing quantum states on a classical computer. If χ grows (sub-)exponentially in time the task is said to be (‘easy’) ‘hard’ in the time-domain. Instead, for the complexity along the space domain, we look at the degree of separability of the evolved state, which directly reflects on the

* manthan.badbaria.phy19@itbhu.ac.in

dimension of the potentially accessible Hilbert space. Specifically, we distinguish whether the state describing the whole system can or cannot be written as product states of (potentially entangled) states describing smaller disconnected subsystems. In case it can, we name the state as separable and the space-complexity is said to be ‘easy.’ Otherwise, we say that is inseparable and space-complexity is ‘hard.’ If the state is separable, the accessible Hilbert space is effectively reducible in smaller disconnected Hilbert spaces each describing different subsystems, as the degrees of freedom defined on such subsystems are not entangled. Instead, if the state is inseparable, the accessible Hilbert space is strictly irreducible. Combining space and time complexity in simulating dynamics of initial product states, we then aim to infer the properties of non-thermal eigenstates of the Hamiltonian.

For the sake of concreteness, we test our complexity-oriented proxy on the kinetically constrained quantum East model [33–41]. From a condensed matter and statistical physics perspective, it attracts interest because it exhibits both fast and slow thermalizing dynamical phases alongside with localization, despite being translationally invariant and non-integrable [33–36, 39–41]. This behavior is markedly different from MBL systems, where localization arises due to many-body wave function interference caused by quenched disorder, while instead it arises by making transport a higher-order process in the quantum East model. Moreover, it features extreme slowdown of thermalization as well as dynamical heterogeneity similarly to structural glasses [33, 42], from which its classical counterpart is inspired [42–46]. From a quantum information perspective, the digital (i.e. Floquet) version hosts special points where it reduces to a Clifford circuit [37], making it a candidate for investigating deviations from the latter, and has been shown to still display localization despite lack of energy conservation [38]. Additionally, its localized nature has been shown to aid in passively protecting quantum information against certain types of coupling to an external environment [39].

In our work we combine a quantum information and statistical physics oriented approach to characterize the quantum East model at finite energy density. Specifically, we aim to inspect spectral properties by using the complexity of simulating dynamics of initial product states via tensor network, going beyond the small system sizes accessible via exact diagonalization. By doing so, we can identify parameter regimes where *many* initial states are easily simulable. Then, by closely inspecting the properties of such states, we discover and compute, via DMRG-X [47], a family of non-thermal finite-energy density eigenstates of the Hamiltonian mostly responsible for the observed slow dynamics.

Interestingly, the easily simulable states are mostly clustered in their energies, and so it is tempting to identify a sort of mobility edge in the spectrum – a sepa-




FIG. 1. Cartoon summarizing the complexity phase diagram of simulating dynamics of initial product states in the quantum East model as a function of their energy density $\epsilon \in [0, 1]$ and the physical time t reached. Each box is split into two triangles which encode the complexity either along the space domain (upper triangle) or along the time domain (lower triangle). A simulation is (easy) hard in the time domain if the computational resources needed, i.e. the bond dimension of the tensor network, grows (sub-)exponentially in time. A simulation is easy (hard) in the space domain if the evolved state is (in)separable. Simulations of initial product states are typically easy in the time domain if the energy density is below a certain energy density ϵ^* , dependent on the parameters of the Hamiltonian, indicating the existence of a *state-dependent* mobility edge in the spectrum (cf. Sec. III C). The sharp differentiation between active and inactive regions, typical of glassy-like systems, makes the space-complexity undergo a transition from a ‘easy’ to ‘hard’ as a function of time t : for $t < t^*$ the entanglement between the different active regions remains negligible and the state can be written as a tensor product of (potentially entangled) states describing the different active regions, i.e. it is separable; for $t > t^*$, the entanglement between the different active regions is no longer negligible and the state is inseparable. In the separable case, the accessible Hilbert space \mathcal{H} appears as if it is reducible in smaller disconnected subspaces \mathcal{H}_i , each describing the i -th active region, similarly to *fragmented* systems [19, 48, 49].

ration of a thermal region and a localized one in the spectrum of the Hamiltonian—challenging the common belief that a mobility edge is exclusive to disordered systems [50–52]. However, small-scale exact diagonalization calculations suggest that the spectrum of the quantum East model does not exhibit a distinct separation between non-thermal eigenstates and thermal ones [34] (see Appendix A). Instead, its spectrum displays features more reminiscent of quantum many-body scarred (QMBS) systems, even though the non-thermal eigenstates are not evenly spaced in energy, are exponentially many in the system size, and are nearly product states [34] (see Appendix A). This contrasts with QMBS systems, where the non-thermal eigenstates are evenly spaced in energy and constitute a zero fraction of the spectrum in the thermodynamic limit [19]. All these ingredients open up an intriguing possibility: the existence of a *state-dependent* mobility edge. In other words, the system behaves akin to having a many-body mobility edge for a large class of initial states, such as product states. To draw an analogy, our scenario resembles QMBS systems, where the impact of the non-thermal eigenstates significantly influ-

ences the dynamics of some initial states. However, differently from QMBS systems, the exponential abundance of non-thermal eigenstates suggests that the non-thermal behavior becomes prevalent across a wide range of initial states, particularly among product states. Under this lens, we expect, and numerically confirm, that simulating dynamics of initial product states can be carried out efficiently via tensor network up to a finite energy density, while instead being computationally hard above it.

As a byproduct of our analysis, the existence of parameters and many initial states whose dynamics can be carried out efficiently suggests that kinetically constrained inspired circuits can be easily simulable on classical computers, alongside Clifford, MBL [53] and fractonic [48, 54] inspired. Additionally, identifying regimes where dynamics of product states are classically challenging holds practical relevance since they are the easily preparable initial states in current NISQ devices [35, 55–61].

A. Summary of results

In Sec. I and Sec. II we introduce our measure of complexity and the model, respectively. In Sec. III we discuss our results, summarized in Fig. 1, which are:

- (i) While time-complexity is typically ‘hard’ in the delocalized phase, as expected for generic non-integrable models, we observe both ‘easy’ and ‘hard’ regimes in the localized phase, indicating the existence of a *state-dependent* mobility edge (see Sec. III C);
- (ii) The sharp differentiation between active and inactive regions, typical of kinetically constrained models and glassy systems [33, 34, 45, 62, 63], makes the initial distribution of excitations matter in dictating time-complexity (see Sec. III D) and space-complexity (see Sec. III E). In the latter, we observe a transition from ‘easy’ to ‘hard’ space-complexity as a function of time. Specifically, in the ‘easy’ regime, the state is approximately separable and its dynamics can be faithfully computed by the dynamics of suitably chosen non-overlapping subsystems. Instead, in the ‘hard’ regime the state is inseparable;
- (iii) We connect the points above by computing a novel family of non-thermal localized eigenstates at finite energy density via tensor network (see Sec. III F). We use such states to explain the extremely slow dynamics observed for a large class of initial states (see Sec. III G).

In Sec. IV we conclude by listing possible fruitful directions in the context of random quantum circuits [64].

I. COMPLEXITY ORIENTED PROXY OF NON-THERMAL EIGENSTATES

To overcome the limitations faced by exact diagonalization, here we propose a complexity-oriented proxy based on tensor-networks for detecting non-thermal eigenstates. Our approach is based on the observation that typical thermalizing systems display an exponential growth of the bond dimension during dynamics, as opposed to systems displaying non-thermal behavior (e.g. MBL systems) where instead the bond dimension grows polynomially in time [29, 30]. An intermediate scenario is constituted by systems with a mobility edge, i.e. systems displaying non-thermal eigenstates in a certain energy window, and thermal ones in the others. Specifically, in such system we envision that the complexity of simulating the dynamics depends on the energy of the specific state at hand, potentially allowing a distinction between the two regions.

To this end, we consider the dynamics of initial product states. Specifically, we first select initial states with a small energy variance so that the eigenstates of the Hamiltonian participating in the dynamics are mostly within a small energy window. Then, we use the way the bond dimension χ grows in time as a measure of time-complexity. In typical non-integrable systems, we expect an exponential growth of χ after a quench, as the bipartite entanglement entropy $S \propto \log \chi$ grows linearly in time [32]. Instead, a sub-exponential growth of χ could indicate the existence of non-thermal eigenstates in the spectrum. In the following, we label it as an (‘easy’) ‘hard’ task if the bond dimension needed scales (sub-)exponentially in time. In such a manner, we can potentially locate a mobility edge within the spectrum at fixed parameters of the Hamiltonian.

Before continuing, we also mention Krylov complexity used as a measure for characterizing how operators explore the available space [65, 66], which has been recently applied also to the quantum East model here discussed [67], and dynamics of operators using tensor networks [68]. However, computing the dynamics of operators generally involves democratically the full spectrum of the Hamiltonian, and so lacks the desired energy resolution needed for detecting a mobility edge.

II. MODEL

We study the quantum East model [33, 34] in open boundary conditions with Hamiltonian

$$\hat{H} = -\frac{1}{2} \sum_{j=0}^N \hat{n}_j (e^{-s} \hat{\sigma}_{j+1}^x - 1), \quad (1)$$

where $\hat{\sigma}_j^\alpha$ is the Pauli- α matrix on site j ; $\hat{n}_j = (1 - \hat{\sigma}_j^z)/2$ is the projector onto the state $|1\rangle$ in the local z basis. The term \hat{n}_j in Eq. (1) is the kinetic constraint, which

translates to a nontrivial action of the Hamiltonian solely to the right ('East') of a previously excited ($|1\rangle$) site. Consequently, the Hamiltonian acts trivially on empty strings without any excited sites to its left, making the location and occupation of the first excited spin a conserved quantity. Thus, the Hilbert space splits into N dynamically disconnected sectors indexed with the position of the first occupied site $|1\rangle$; i.e. the k -th sector has $(k-1)$ zeros preceding $|1\rangle$ on the k -th site. This feature is in stark contrast with systems undergoing Hilbert space fragmentation, where the number of disconnected subspaces is $\mathcal{O}(\exp(N))$ [19, 48, 49] (cf. Sec. II B for a comparison with Hilbert space fragmentation).

Because of the trivial action of Hamiltonian on empty sites, the results do not depend on the sector considered in the thermodynamic limit. Thus, we fix $k=1$ throughout our work, without loss of generality. Once the 'East symmetry' sector is fixed, any product state can be dynamically accessed by any other. In other words, the sector is irreducible, as the Hamiltonian does not possess any other nontrivial (excluding the energy) conserved quantity.

A. Localization

Despite being non-integrable and translational invariant, numerical evidence mostly based on each diagonalization indicates that the quantum East model displays a dynamical transition separating a fast and slow thermalizing phase as a result of the competition of the kinetic term, controlled by e^{-s} , and the potential one $\propto \sum_j \hat{n}_j$ [33, 34]. Intuitively, when the kinetic term dominates ($s \lesssim 0$), excitations propagate ballistically, making the details of the initial state rapidly lost, while instead when it is small $s \gtrsim 0$, excitations propagate slowly, making the details of the initial state *potentially* matter up to long times. In Ref. [34] it was shown that such dynamical transition is linked to the delocalization-localization transition occurring in the ground state. When $s < 0$ the ground state is delocalized; namely, the wave function is spread along the lattice with homogeneous probability and amplitude of finding an occupied site. Instead, the ground state is localized for $s > 0$, namely the corresponding wave functions contain nontrivial excitations only on a small region of the lattice around the first excitation fixing the East symmetry, while it is approximately in the vacuum state everywhere else. In other words, for $s > 0$ the probability of finding an occupied site in the ground state decays exponentially as [34]

$$\langle \hat{n}_j \rangle \sim e^{-j/\xi}, \quad (2)$$

where ξ is the localization length, parametrically small in s , beyond which the ground state can be approximated as a product state of empty sites. In turn, the localized ground state can be used, together with empty strings, as a basis of area-law states arbitrarily close to true eigenstates of the Hamiltonian, partially explaining the possi-

ble slow thermalization in the localized phase [34, 39]. However, such construction is mostly limited to low-energies and so there is still a lack of a complete understanding of the dynamical phase transition at finite-energy density. Here, we aim to investigate such dynamical phase transition more closely, focusing on whether the system hosts a mobility edge despite being disorder-free. However, from exact diagonalization calculations, a sharp separation of a thermal and non-thermal region looks unlikely (see Appendix A) [34]. Nonetheless, since many non-thermal eigenstates have a large overlap with product states (see Appendix A), there is the possibility that the system behaves akin to having a many-body mobility edge for a large class of initial states, such as product states, and displays a 'state-dependent mobility edge.' We will address such open questions using the complexity-oriented proxy introduced in Sec. I.

B. Comparison with Hilbert space fragmentation

Before presenting our results, we briefly summarize the differences between the quantum East model and systems displaying Hilbert space fragmentation. We do so by briefly summarizing the main properties of fragmented systems, while we refer to, e.g., Ref. [19] for a more comprehensive and complete discussion.

Given a Hamiltonian \hat{H} , the Hilbert space \mathcal{H} on which it acts can be generally decomposed into dynamically disconnected subspaces $\{\mathcal{H}_n\}$, referred to as Krylov subspaces, as

$$\mathcal{H} = \bigoplus_{n=1}^Q \mathcal{H}_n, \quad \mathcal{H}_n = \text{span}_t \{e^{-i\hat{H}t} |\psi_n\rangle\}, \quad (3)$$

where Q is the number of Krylov subspaces and

$$\text{span}_t \{e^{-i\hat{H}t} |\psi_n\rangle\} \equiv \text{span} \{|\psi_n\rangle, \hat{H}|\psi_n\rangle, \hat{H}^2|\psi_n\rangle, \dots\} \quad (4)$$

denotes the subspace spanned by the time evolution of the state $|\psi_n\rangle$. The states $|\psi_n\rangle$ are chosen so that they are not eigenstates of the Hamiltonian and their Krylov subspaces are distinct. More concretely, $|\psi_n\rangle$ are typically chosen to be product states, as they are the ones more easily accessible experimentally, although recent works are investigating the case where they are entangled states [41, 49, 69]. In doing such decomposition, a key question concerns how many Krylov subspaces the system displays, as well as the properties of each of them. Concerning the number Q of Krylov subspaces, we can distinguish two main scenarios depending on how Q scales with the system size N . Specifically, we could have

$$Q = \begin{cases} \mathcal{O}(N^p), & \text{with } p \geq 0 \\ \mathcal{O}(\exp(N)) \end{cases} \quad (5)$$

The first scenario occurs for systems exhibiting no or 'conventional' abelian or non-abelian symmetries, where

with ‘conventional’ we mean that they can be written as the sum of local terms (e.g. $U(1)$ symmetry), or products of one-site unitary (e.g. \mathbb{Z}_2 symmetry). In this case, the different Krylov subspaces are labeled by the quantum numbers associated with such symmetries. Instead, when $Q = \mathcal{O}(\exp(N))$ the system is said to be fragmented, and it occurs when the system displays non-local conserved quantities [49] (e.g. the presence of bit strings which are invariant under the action of the Hamiltonian) associated with non-conventional symmetries. Once the different subspaces are labeled, a relevant question concerns investigating their properties, such as whether the system thermalizes or not within each Krylov subspace [70–72].

Given these definitions, it is evident that the quantum East model does not display Hilbert space fragmentation. Indeed, it conserves (trivially) the energy and the location of the first excited site. Given that in a system of size N there are N different ways in which we can locate the first excited site, we can label N different Krylov subspaces. Furthermore, each of the Krylov subspaces cannot be further reduced, namely, it is ergodic, as seeding any product state belonging to such subspace in Eq. (4) allows us to explore the whole subspace. However, the remarkable aspect is that despite having the possibility to explore an exponentially large Hilbert space, the way such exploration occurs undergoes a dynamical phase transition from a regime where exploration occurs quickly to a regime where it happens extremely slowly, impeding thermalization for exponentially long times in the system size. Additionally, we envision the scenario where the efficiency with which the Hilbert space is explored depends not only on the Hamiltonian but also on the initial state seeded $|\psi_n\rangle$. Indeed, there could be, at fixed Hamiltonian parameters, a product state $|\psi_n\rangle$ for which the connectivity effectively results higher than others, allowing to explore a larger portion of the accessible Hilbert space faster, which could be enough to make the system thermalize for that specific initial state [73]. This point of view is equivalent to the original question we posed, about the existence of a *state-dependent* mobility edge in the model at hand, and further justifies why we look at the dynamics of initial product states.

III. STATE-DEPENDENT MOBILITY EDGE

In this section, we investigate the time-complexity and space-complexity of evolving initial product states as a function of their properties, i.e. energy density and spatial structure, and the Hamiltonian parameters. Based on this, we discover a family of localized eigenstates with finite energy density responsible for the observed behavior for many initial states considered, corroborating the sensitivity of our complexity-oriented proxy to non-thermal eigenstates of the Hamiltonian.

A. Initial states

As we are interested in investigating the existence of a *state-dependent* mobility edge, we need to specify the class of states of our interest. To our ends, we investigate the dynamics of initial product states in the computational basis

$$|\psi\rangle = |1\rangle \bigotimes_{k=1}^N |0/1\rangle_k, \quad (6)$$

where we keep the first site fixed to $|1\rangle$, making the dynamics occur in the largest irreducible ‘East’ symmetry sector of our model, while $|0/1\rangle$ could be either $|0\rangle$ or $|1\rangle$. As discussed in Sec. I, a quench protocol could be used to probe information about the eigenstates when the energy variance of the initial state is small. Product states as the one in Eq. (6), with M excitations ($|1\rangle$), have

$$\begin{aligned} \langle\psi|\hat{H}|\psi\rangle &= \frac{M}{2}, & \langle\psi|\hat{H}^2|\psi\rangle &= \frac{e^{-2s}}{4}M + \frac{M^2}{4}, \\ \frac{\sqrt{\Delta H}}{\langle\psi|\hat{H}|\psi\rangle} &= \frac{e^{-s}\sqrt{M}}{M} \sim \frac{1}{\sqrt{M}}, \end{aligned} \quad (7)$$

where $\Delta H \equiv \langle\psi|\hat{H}^2|\psi\rangle - \langle\psi|\hat{H}|\psi\rangle^2$. Since we are interested in making statements at finite energy density (i.e. we want $\langle\hat{H}\rangle/N$ finite for $N \rightarrow \infty$), we set $M = mN$, with $m \in [0, 1]$ the density of excitations. In such a manner, the energy density is finite while the relative fluctuation around the mean goes to zero in the thermodynamic limit. Thus, such states are good candidates for analyzing the spectrum of the quantum East model. Notice that the average energy depends solely on m and not on their location, allowing us to isolate the impact of the spatial structure in the dynamics keeping the energy fixed.

B. Details on the numerical methods

For a fixed value of s and m , we sample up to 100 random product states to mitigate sample biases, and we simulate their dynamics using the Time Evolving Block Decimation algorithm [74]. We keep the Schmidt singular values larger than 10^{-14} and we set the timestep $\Delta t = 10^{-3}$. We stop the simulation when either the maximum bond dimension reaches 512 or the time reached is enough to compute the quantities of interest. For each simulation, we investigate how the max bond dimension $\chi_{\max}(t) = \max_{j \in [1, N-1]} \chi_j(t)$ grows in time by fitting either a polynomial ($\propto t^\alpha$) or exponential ($\propto e^{rt}$) function, where α and r are positive constants, depending on which one better approximates the data. Then, we link such behavior to the spectral properties of the Hamiltonian, as detailed in Sec. I. We highlight that trotterizing the continuous-time dynamics induces undesired errors tied to the finite time step. However, we do not expect the quantity of interest (i.e., the way the bond dimension




FIG. 2. (a) Fraction of sampled initial product states (cf. Eq. (6)) for which time-complexity is easy $f_{\text{easy}} \in [0, 1]$ as a function of the normalized energy density $\epsilon = (\langle H \rangle - E_{\text{min}})/(E_{\text{max}} - E_{\text{min}})$ and s . The grey region indicates the lack of data, as the chosen initial states (cf Eq. (6)) do not span the whole energy spectrum. The yellow line serves as guidance to indicate where $f_{\text{easy}} = 0.5$. It is dashed if it changes as the system size N increases (for $0 \lesssim s \lesssim 0.3$), and continuous if it does not (see Fig. 3). Deep in the localized ($s > 1$) and delocalized phase ($s < 0$), the behavior displays typicality, namely either $f_{\text{easy}} \approx 0$ (typically ‘hard’) or $f_{\text{easy}} \approx 1$ (typically ‘easy’) weakly depending on ϵ . Near the transition point on the localized side ($0 \lesssim s \lesssim 0.5$) f_{easy} highly depends on ϵ , giving evidence of the existence of a mobility edge along the spectrum. (b-c) Dynamics of $\chi_{\text{max}}(t)$ for all the sampled states for different values of s and ϵ , marked in (a). The dashed line represents a fit for a representative state. Deep in the ‘easy’ and ‘hard’ regime the growth of $\chi_{\text{max}}(t)$ weakly depends on the parameters of the initial states. Instead, in the regions separating typically ‘hard’ or ‘easy’ we observe different behaviors for states with the same ϵ , signaling a role played by other features of the initial state, such as its spatial structure.

grows) to be qualitatively affected by such errors, provided that we use a small enough time step. We present results on system size $N = 30$, as our statements are not appreciably affected upon increasing N apart from near the transition point $0 \leq s \lesssim 0.3$, as we will show (cf. Appendix B for details on the scaling analysis in N). All the results were obtained using the python package quimb [75].

C. Role of energy density

Here we focus on the impact of the energy density of the initial states in Eq. (6) in dictating the time-complexity of simulating the dynamics. While time-complexity is typically ‘hard’ in the delocalized phase ($s < 0$), as expected for generic non-integrable models, we remarkably observe both ‘easy’ and ‘hard’ regimes in the localized phase ($s > 0$) also at finite energy density, indicating the existence of a *state-dependent* mobility edge.

To extract the behavior in the thermodynamic limit, we measure the energy with respect to the ground state and most excited state energies, namely the normalized energy density $\epsilon = (\langle \hat{H} \rangle - E_{\text{min}})/(E_{\text{max}} - E_{\text{min}}) \in [0, 1]$, where E_{min} and E_{max} are the energy of the ground state and the most excited state (which can be computed via DMRG minimizing the energy of $-\hat{H}$), respectively. Despite the advantage of initializing product states (cf. Eq. (6)) in isolating the interplay of their properties, as we will discuss, they have a drawback: they do not

always allow an efficient sampling over ϵ . Specifically, as s decreases, it is not possible to sample from the extremes of the spectrum, as the ground state and the most excited states are ‘far’ from the singly occupied state and the completely filled state, respectively. Nonetheless, as we are mostly interested in the central region of the spectrum, associated with high-temperature, such limitation does not play a major role in our results. Finally, we highlight that the many-body spectrum is not symmetric around $\epsilon = 0.5$ ($\hat{H} \neq -\hat{H}$ up to a unitary transformation) and so there are no reasons to expect a symmetric mobility edge.

In Fig. 2(a), we show the fraction $f_{\text{easy}} \in [0, 1]$ of states for which time-complexity is ‘easy’, i.e. $\chi_{\text{max}}(t)$ grows sub-exponentially in time, at a given ϵ and s . Specifically, f_{easy} is the average, over the sampled initial states at a given energy density ϵ , of a binomial variable which takes the value 0 if the time-complexity is hard, or 1 if it is easy. As expected, in the delocalized phase ($s < 0$) the system mostly displays exponential growth of the bond dimension ($f_{\text{easy}} \approx 0$) as we move towards the middle of the spectrum ϵ (cf. Fig. 2a), reflecting the thermal nature of the whole spectrum. On the other hand, for $s > 0$ we observe regions where f_{easy} is large not only near the extreme of the spectrum but also at finite energy density ϵ (cf. Fig. 2(b)). For $0 \lesssim s \lesssim 0.5$ (cf. Fig. 2(c)), we observe as ϵ increases an ‘inversion’ of f_{easy} , namely for small ϵ the dynamics is typically ‘simple’ to be simulated ($f_{\text{easy}} \approx 1$), while moving towards the center of the spectrum dynamics is typically ‘hard’ to be simulated ($f_{\text{easy}} \approx 0$). While for $0 \leq s \lesssim 0.3$ we argue that the existence of such inversion




FIG. 3. Fraction of randomly sampled initial product states for which time-complexity is easy (f_{easy}) as a function of the energy density ϵ at fixed $s = \{0.3, 0.4\}$ and different system sizes $N = \{15, 30, 60, 90\}$ (from light to dark curves). Referring to Fig. 2(a), the data correspond to two vertical cuts. Deep in the localized phase, the results are not affected by N and simulations are always easy (not shown). Instead, the closer s is to the localization point ($s = 0$), the more f_{easy} is affected by N . The bars correspond to the statistical uncertainty due to the finite number of initial states sampled. Inset: ϵ^* , where $f_{\text{easy}}(\epsilon^*) = 0.5$, as a function of N . The error bars come from the statistical uncertainty in f_{easy} which makes multiple ϵ compatible, within a standard deviation, with our definition of $f(\epsilon^*) = 0.5$. For $0 \leq s \lesssim 0.3$ the easy-to-hard inversion occurs at an energy density that decreases in N , providing evidence that solely the ground state is likely localized in the $N \rightarrow \infty$ limit. Instead, for $0.3 \lesssim s \lesssim 0.5$ we observe that upon increasing N the inversion occurs at a size-independent ϵ , indicating the existence of a *state-dependent* mobility edge.

is a finite-size effect, i.e. the energy density at which it occurs decreases in N indicating that solely the ground state is localized, for $0.3 \lesssim s \lesssim 0.5$ we observe that upon increasing N the inversion occurs at the same ϵ (see Fig. 3). Such size-independent inversion provides evidence of a *state-dependent* mobility edge in the spectrum. Instead, deep in the localized phase $s \gtrsim 0.5$, $f_{\text{easy}} \approx 1$ at any ϵ indicating that a large fraction of highly excited eigenstates becomes non-thermal, with the fraction increasing with s .

Based on our results, the *state-dependent* mobility edge is reminiscent more of a crossover than a sharp transition. Indeed, we observe parameter regimes where f_{easy} assumes value far from its extremes ($f_{\text{easy}} \neq \{0, 1\}$), indicating that other features apart from the energy density are playing a role in the dynamics (see e.g. Fig. 2(c) where at fixed $\epsilon = 0.1$ dynamics of $\chi_{\text{max}}(t)$ changes based on the initial state). We further investigate such dependence in the following section.

D. Role of spatial structure

Here, we investigate the role of the initial state structure in the time-complexity. In doing so, we show how the size of excited regions and their distance plays a crucial role in dictating the time-complexity of simulating their dynamics [33, 34]. Additionally, we show that the localization length of the ground state serves as a length scale in the system useful in understanding finite-energy density phenomena.

Long-time dynamics of slowly thermalizing systems can depend on features of the initial state beyond the conserved quantities of the system. In kinetically constrained models (KCMs), the spatial structure of the initial state plays a key role up to extremely long times, in a manner reminiscent of glassy systems [33, 34, 42, 62, 63]. Such a dependence stems from the sharp differentiation between active and inactive regions in KCMs. Typically, as in the quantum East model, active regions are constituted by $|1\rangle$, while inactive regions by $|0\rangle$. As a result, a natural parameter to capture the spatial structure is the initial size w of inactive regions (number of consecutive $|0\rangle$) as it controls when active regions will entangle [33]. However, this parameter does not contain information regarding the size of the active regions. To capture such information, we look at the initial size of active regions k (number of consecutive $|1\rangle$), which is also in a one-to-one correspondence with their average energy (cf. Eq. (7)) for initial product states in the computational basis.

As we are dealing with initial states with multiple active regions, we trade k and w with their averages over the system k_{avg} and w_{avg} , respectively. Inspecting the time-complexity as a function of these two parameters, we observe that such quantities are not so effective in predicting the time-complexity of simulating the states at hand (see Appendix D). However, by including the localization length ξ of the ground state at the corresponding s (we restrict to $s > 0$), we observe that the predictive power improves (see Fig. 4(a)). The localization length ξ serves as a length scale as in the localized phase active regions expand exponentially slowly and remain within few ξ up to long times. As k_{avg} increases the simulations are typically harder, while instead, the opposite occurs as excitations are farther apart. Such behavior could be understood as the interplay of dynamics within each cluster of excitations and between different ones. Intuitively, if clusters are far from each other on average ($w_{\text{avg}}/\xi \gg 1$), the time-complexity is mostly dictated by the dynamics within each cluster since the propagating front generates little entanglement. In turn, the larger the cluster is, and so is its energy, the more it is typically hard to simulate, in agreement with Fig. 2(a).

A possible analogy to explain such behavior is given by looking at the system as a collection of subsystems with a certain temperature, directly linked to




FIG. 4. (a) The same data as presented in Fig. 2 labeling the states based on the initial average size of clusters of consecutive excitations ($|1\rangle$) normalized by N (k_{avg}/N) and average distance between clusters w_{avg} in units of ξ (w_{avg}/ξ), with ξ the localization length of the ground state at the respective value of s . The red dots correspond to ‘hard’ time-complexity ($\chi_{\text{max}}(t) \propto e^{rt}$) with exponential rate r . The blue dots correspond to ‘easy’ time-complexity ($\chi_{\text{max}}(t) \propto t^\alpha$) with power-law exponent α . Simulations are typically hard if either k_{avg}/N increases and w_{avg}/ξ decreases. (b-c) dynamics of the bond dimension χ_i along each cut for two representative states. In (b), clusters of excitations have started to entangle beyond a transient time. Whereas in (c), clusters of excitations haven’t entangled yet due to the presence of large islands of consecutive $|0\rangle$.

the number of excitations, separated by completely inactive regions at zero temperature (the empty state $|00\dots 0\rangle$ is the true ground state of the quantum East model): if the inert regions are too extended, the hottest source dominates the hardness in simulating the dynamics. Such a picture is supported by inspecting more closely how the bond dimension, or equivalently entanglement, at each possible bipartition, evolves in time (cf. Fig 4(b,c)). In Fig. 4(b) islands of excitations have started to appreciably entangle beyond a transient time. Instead, in Fig. 4(c) clusters of excitations are not strongly entangled due to the presence of large inactive regions between them.

The distinction between inactive and active regions opens up the possibility of understanding the dynamics of generic product states in terms of concatenated and weakly entangled clusters in the localized phase. We further investigate this observation in the next sections.

E. Transition in space-complexity

We have observed that the structure of the initial state plays a leading role in dictating the time-complexity in the localized phase. Such a dependence stems from the sharp differentiation between active and inactive regions in kinetically constrained models (KCMs). Here, we go a step further showing that such distinction also leads to a crossover in the space-complexity from a fragmented regime, where the state is separable, to a fully ergodic one, where the state is non-separable.

Heterogeneity is a hallmark of classical glassy systems, which manifests also in their quantum counterpart as ob-

served in other studies [42, 63]. The quantum East model makes no exception, as it is evident from the dynamics of various observables, such as occupation number and entanglement entropy. This feature opens up the possibility of distinguishing two timescales: one where dynamics occurs *mostly* within each cluster of excitations (intra-cluster), and another when it also *appreciably* involves different clusters of excitations (inter-cluster). Specifically, we could define a time t^* up to which the system is approximately separable, since the entanglement between the different clusters is negligible, and the whole dynamics is encoded in the dynamics of each cluster separately. Such separability in the evolved state could be formalized using the Lieb-Robinson bound [76]: quantum correlations propagate at most ballistically with exponentially small corrections in systems with a finite local Hilbert space and short-range interactions. In KCMs, empty regions are completely inactive, and the finite velocity of propagation of entanglement in the system (upon neglecting the exponentially small correction in the distance) implies the separability just mentioned up to time t^* where the active regions appreciably merge. As a consequence, an initial product state (cf. Eq. (6)) with \mathcal{N} cluster evolves as

$$|\psi(t)\rangle \approx \bigotimes_{n=1}^{\mathcal{N}} |\psi_n(t)\rangle \quad t \lesssim t^* \quad (8)$$

where $|\psi_n(t)\rangle$ is the time-evolved state describing the n -th cluster. In other words, the evolved state is given by a product state of each cluster, i.e. it is separable (up to exponentially small corrections coming from the propagating fronts). The transition from separability to fully ergodic is accompanied by a change in the dimension




FIG. 5. Dynamics of the entanglement entropy S , computed on the last bond, upon initializing the state $|111\rangle \otimes |0\rangle^w \otimes |1\rangle$ with $w \in [1, 8]$ in the delocalized ($s < 0$) and localized phase ($s > 0$). The long-time values of S are independent of w as expected in the fastly thermalizing regime ($s = -0.5$), whereas it decreases as w increases for $s = 0.5$ indicating localization and slow thermalization. (b) Time t^* at which the two clusters appreciably entangled, i.e. when $S > \Delta$, as a function of w . We set $\Delta = 10^{-2}$ for the sake of concreteness (see text for a more compelling discussion in terms of Lieb-Robinson bound). In the delocalized phase ($s < 0$), t^* grows linearly in w as excitations propagate ballistically. Instead, t^* grows exponentially in w in the localized phase ($s > 0$), due to the extreme slowdown in dynamics. The results were obtained via exact diagonalization.

of the accessible Hilbert space \mathcal{D} . Specifically,

$$\mathcal{D}(t) = \begin{cases} \sim \mathcal{N} \times 2^{N/\mathcal{N}} & t \lesssim t^*, \\ 2^N & t \gtrsim t^*, \end{cases} \quad (9)$$

which in words translates, keeping N/\mathcal{N} fixed, to a transition in the dimension of the accessible Hilbert space from *polynomial* to *exponential* in the system size N . Such transition could be interpreted as a transition between a fragmented regime [19, 48, 49] to a fully ergodic one. However, we once more highlight that such a transition is present in the quantum East model upon neglecting the exponential small corrections coming from the Lieb-Robinson bound. Instead, in exactly fragmented systems the different Hilbert space sectors are completely disconnected, and the time t^* is formally infinite.

As the Lieb-Robinson bound constitutes an upper bound, it is in principle possible to observe slower than ballistic propagation. This is the case for the quantum East model, where the propagation could be exponentially slow in the localized phase [33]. As a consequence, the dependence of the time t^* with the distance w , defined as the number of empty sites $|0\rangle$ between two clusters, strongly depends on whether the system is localized or delocalized. To test this, we compute the evolution of the entanglement entropy between two clusters, of size 3 and 1 respectively, at a distance w , i.e.

$|\psi(t=0)\rangle = |111\rangle \otimes |0\rangle^w \otimes |1\rangle$. Specifically, we look at the entanglement entropy S on the last bond, and we define a threshold Δ so that we consider the two clusters not entangled if $S < \Delta$. The specific value of Δ is arbitrary and chosen just to show the dependence of t^* on w (cf. Fig. 5). In the delocalized phase, t^* grows linearly in w as excitations propagate ballistically. Instead, t^* grows exponentially in w in the localized phase, due to the extreme slowdown in dynamics (cf. Fig. 5(b)). Moreover, the value reached by S in the long time limit is independent on w for $s < 0$, as expected for systems evolving into a volume law state, differently from the localized phase ($s > 0$), indicating non-thermal behavior at least up to the times reached. Since t^* could be very large in the system size, our observation could prove valuable in different directions:

- (i) For time $t < t^*$ the dynamics of the whole system can be efficiently simulated as a collection of its subsystems making a negligible error [77, 78].
- (ii) Separability up to time t^* justifies the investigation of smaller system sizes in order to grasp the behavior or larger ones. Specifically, from Eq. (8), the dynamics of a product state given by \mathcal{N} cluster is completely encoded looking at the dynamics of \mathcal{N} single-cluster states up to time t^* .

Based on (ii), we now investigate the time-complexity in simulating the time evolution of single-cluster states, namely kink states. We will show that such investigation not only will confirm the observed time-complexity in the previous sections, but also allow us to identify a class of area-law states responsible for such behavior. This will provide evidence of the sensitivity of our complexity-oriented proxy to non-thermal eigenstates.

F. Localization at finite energy-density

Following the conclusions of the previous section, we investigate the time-complexity of simulating the dynamics of the kink states

$$|\mathbf{k}\rangle = |1\rangle^k \otimes |00\dots 0\rangle. \quad (10)$$

As always for product states in the computational basis, we have a one-to-one correspondence between k and the energy density ϵ . Since we now have a single state for each point in the parameter space ϵ vs s , we show in Fig. 6 the relative error $\delta_{p/e}$ between the polynomial and the exponential fit performed on the time evolution of the max bond dimension $\chi_{\max}(t)$. If $\delta_{p/e} > 1$, the exponential fit better approximates the data, while instead vice versa if $\delta_{p/e} < 1$. We observe that the time-complexity of simulating kink states follows a similar trend as the one observed in Fig. 2(a). This signals that, as expected, the hardest part to simulate is given by regions densely excited, or in other terms the ‘hottest’ regions.




FIG. 6. Time-complexity upon initializing kink states $|\mathbf{k}\rangle = \otimes_{j=1}^k |1\rangle \otimes_{j=k+1}^N |0\rangle$ with $k \in [1, N/2]$ with $N = 30$. Polynomial and exponential fits are performed to determine the complexity of simulating these states. (a) Relative error $\delta_{e/p}$ between the exponential and polynomial fits as a function of energy density ϵ and s . For $\delta_{e/p} < 1$, the state is ‘hard’ to simulate. Conversely, for $\delta_{e/p} > 1$, the state is ‘easy’ to simulate. We see a complexity trend similar to the one observed in Fig. 2(a), signaling that most of the complexity of simulating product states could be understood in terms of the evolution of kink states. (b) Dynamics of χ_{\max} for kink states with varying k , in the delocalized ($s = -0.5$) regime and in the localized ($s = 0.5$) regime. In the delocalized regime, states are ‘hard’ to simulate irrespective of k . Instead, in the localized regime, k has a huge impact on the time-complexity.

To better understand our observations, we more closely investigate the eigenstates of the Hamiltonian. In particular, from Exact Diagonalization calculations, we observe that the kink states in Eq. (10) have a large overlap with a limited number of eigenstates of the Hamiltonian deep in the localized phase. Inspecting such eigenstates, we observe that they have a density profile similar to a kink state with an additional localized tail on the right edge, which we term as *localized kink states*. Due to the limited system sizes accessible via Exact Diagonalization, we further characterize such states resorting to DMRG-X, a variation of the standard DMRG, which allows finding area-law eigenstates which are ‘near’ to the initial state seeded to the algorithm [47]. In our case, we seed the kink states (cf. Eq. (10)) and let the algorithm find the best approximating eigenstate, setting a maximal bond dimension of 100. For each of these states, the algorithm could either fail to converge, which we interpret as the absence of a localized kink state, or converge.

In Fig. 7 we show the localization length ξ of the states for which the algorithm was able to converge (we set as convergence criteria a variance of the Hamiltonian $< 10^{-5}$). We observe that the algorithm can find a state with $\epsilon = 0$, faithfully reproducing the results from DMRG, as it corresponds to the ground state of the Hamiltonian. Additionally, the algorithm can find other localized states at finite energy density as s increases, corresponding to the localized kink states above




FIG. 7. (a) Inverse of the localization length ξ of localized kink eigenstates (variance of energy $< 10^{-5}$) obtained via DMRG-X [47], as a function of their energy density $\epsilon = (\langle H \rangle - E_{\min}) / (E_{\max} - E_{\min})$ in a system of size $N = 30$. The algorithm converges not only at zero energy density ($\epsilon = 0$), corresponding to the ground state, but also at finite energy $\epsilon > 0$, indicating the existence of localized kink-states along the spectrum for $s > 0$. (b) Density profile of some localized kink states found via DMRG-X for different ϵ at fixed $s = 1.4$. The dashed lines are the exponential fit used to extract ξ .

introduced. We observe that ξ is parametrically small in s , while instead it is parametrically large in ϵ . In Appendix E we provide further details on the results obtained via DMRG-X. This further provides evidence of the effectiveness of our complexity-oriented quantity for detecting the presence of thermal and non-thermal eigenstates states.

G. A unifying picture

The localized kink states found could be used, together with empty strings, as building blocks for defining a large class of states close to eigenstates of the Hamiltonian for larger system sizes. To show this, let us name $|\tilde{k}^N\rangle$ the localized kink-states, with k the number of excitations in the kink (excluding the localized tail) and N the system size. First of all, we observe that the state

$$|\Psi_k^L\rangle \equiv |\tilde{k}^N\rangle \otimes |0\rangle^{\otimes(L-N)}, \quad (11)$$

with support on $L > N$ sites, has energy variance

$$\langle \Psi_k^L | \Delta \hat{H} | \Psi_k^L \rangle \sim e^{-s} \langle \tilde{k}^N | \hat{n}_N | \tilde{k}^N \rangle \sim e^{-s} e^{-(N-k)/\xi} \quad (12)$$

which depend solely on the occupation on the last site of the state $|\tilde{k}^N\rangle$, since the only contribution to the variance comes from the boundary term between $|\tilde{k}^N\rangle$ and the subsequent string of empty sites. Since the variance in Eq. (12) is independent on L , it is small if $(N - k) \gg \xi$, i.e. $|\tilde{k}^N\rangle$ is localized. Since ξ does not scale with N , such a condition is satisfied if $k/N \ll 1$ for large N , which can be fulfilled for $k < cN$ with $c \ll 1$

a small constant and $k < N^p$ with $p < 1$. Recalling the correspondence between the number of excitations and the average energy, the former case corresponds to states with an extensive energy, while the latter to states with a sub-extensive one. The bound in k imposed is quite strict in practice and could be relaxed by observing the exponential dependence of the variance on $(N - k)/\xi$. Specifically, by setting a target variance ε in Eq. (12), it would be enough to have $(N - k) \sim \xi \log(1/\varepsilon)$.

Having proven that the localized kink states found at finite N are quasi-eigenstates of larger systems, we can do a step further. Specifically, we could use them as a basis for states in a system of size $L > N$. Allowing $k \in [0, k_{\max}]$, where $|\tilde{0}\rangle \equiv |0\rangle^{\otimes N}$ and k_{\max} corresponds to the maximum number of excitations for which the system display localization, the number of states we can define is $(k_{\max} + 1)^{L/N}$. Let us call $|\Psi^L\rangle \in \{|\tilde{0}^N\rangle, |\tilde{1}^N\rangle, \dots, |\tilde{k}_{\max}^N\rangle\}^{\otimes L/N}$. Since $|\tilde{0}^N\rangle$ blocks do not contribute to the variance, the only contribution comes from the states $|\tilde{k}^N\rangle$, giving

$$\langle \Psi^L | \Delta \hat{H} | \Psi^L \rangle \sim e^{-s} \sum_{j=1}^{\mathcal{M}} e^{-(N-k_j)/\xi} < \mathcal{M} e^{-(N-k_{\max})/\xi}, \quad (13)$$

where \mathcal{M} is the number of states $|\tilde{k}\rangle$ with $k > 1$, and k_j is the size of the j -th concatenated localized kink state. We highlight that \mathcal{M} is potentially unbounded in the thermodynamic limit, making the variance inevitably large. The energy is given by

$$\langle \Psi^L | \hat{H} | \Psi^L \rangle \sim \sum_{j=1}^{\mathcal{M}} \langle \tilde{k}_j | \hat{H} | \tilde{k}_j \rangle \sim \sum_{j=1}^{\mathcal{M}} k_j. \quad (14)$$

Given these results, we now aim to find a trade-off between small variance and finite energy density. To this end, let us embed the maximum number of localized states possible, namely $\mathcal{M} = L/N$, and consider that the kink states embedded have $k \sim cN$, with $c \ll 1$. Let us consider the case where $N = L^\alpha$, with $\alpha < 1$, to which corresponds $\mathcal{M} = L^{1-\alpha}$ and $k \sim cL^\alpha$. For this parametrization, we have $\langle \Psi^L | \hat{H} | \Psi^L \rangle \sim cL$ and $\langle \Psi^L | \Delta \hat{H} | \Psi^L \rangle < L^{1-\alpha} e^{-L^\alpha/\xi}$, implying that we can have finite energy density states with small variance and so slowly evolving.

The existence of localized kink eigenstates, which include the ground state [34], allows us to at least partially explain the ease of simulating the dynamics of a large class of initial states deep in the localized phase. Indeed, the product states in the computational basis considered are very close (in overlap) to concatenated localized kink states, if they exist, which we have shown can have small variance and are thus slowly evolving. However, the localized states here identified do not allow us to explain the dynamics of *all* the product states simulated. Indeed, based on our numerical results, we do not find localized kink states at large energy density for $s \lesssim 1$, while instead

looking at time-complexity as well as small system size exact diagonalization (cf. Appendix A) we would argue that non-thermal eigenstates should exist. This opens up the possibility that non-thermal eigenstates qualitatively different from localized kink states exist.

IV. DISCUSSION AND PERSPECTIVES

In this work, we have characterized the kinetically constrained quantum East model by inspecting the complexity of simulating the dynamics via tensor networks. In doing so, we have distinguished complexity along the time (time-complexity) and space (space-complexity) domain. We have linked the ease of time-complexity to the presence of numerous non-thermal eigenstates at finite energy densities, a subset of which we have computed via DMRG-X. The abundance of non-thermal eigenstates causes the system to exhibit behavior akin to having a mobility edge when initializing product states, a scenario we term the *state-dependent* mobility edge. Space-complexity, on the other hand, is susceptible to the separability of the evolved state. We have shown that the sharp distinction between active and inactive regions, characteristic of kinetically constrained models, induces a transition in space-complexity during the dynamics of initial product states, as summarized by Eq. (8).

Intriguingly, the transition encoded in Eq. (8) is reminiscent of the dynamical transition in the complexity of performing the so-called sampling task [79, 80]. The sampling problem involves the extraction of events according to the probability distribution provided by the many-body quantum state, and it is widely regarded as a leading contender in demonstrating provable quantum advantage. Additionally, and more relevant for us, the sampling problem has gathered attention also for defining novel kinds of dynamical phase transitions linked to an easy to hard transition during dynamics, which could already happen in free bosons systems in a lattice [81]. Specifically, in Ref. [81], it was considered free bosons sparsely located along a lattice: for times smaller than the time at which particle interfere, controlled by the Lieb-Robinson bound, the task is easy as quantum particles behave as distinguishable classical ones; for larger times than the interference time the task becomes hard, as interference between particles can no longer be disregarded. This strongly resembles the transition encoded in Eq. (8).

A. (Random) unitary circuits and thermal bubbles

The results obtained in the quantum East model open up the intriguing possibility that the corresponding quantum circuit could be efficiently simulable as well. More concretely, the generator of the dynamics could be Trot-

terized as

$$e^{-i\hat{H}T} = \prod_{n=1}^{T/\Delta t} e^{-i\hat{H}\Delta t} \approx \prod_{n=1}^{T/\Delta t} \left(\prod_{j=1}^{N-1} e^{-i\hat{h}_{j,j+1}\Delta t} \right) \quad (15)$$

where the two-sites operator $e^{-i\hat{h}_{j,j+1}\Delta t}$ for the quantum East model results equal to

$$e^{-i\hat{h}_{j,j+1}\Delta t} = e^{iJ\hat{n}_j\hat{\sigma}_{j+1}^x\Delta t/2} e^{-i\hat{n}_j\Delta t/2} \quad (16)$$

where $J \equiv e^{-s}$. By using $\hat{n}_j^k = \hat{n}_j$ for $k > 0$, and $(\hat{\sigma}_j^x)^k = 1$ for k odd, we can write the evolution operator exactly as a

$$e^{-i\hat{h}_{j,j+1}\Delta t} = U_j + U_{j,j+1} \\ U_j = 1 - \hat{n}_j + \hat{n}_j \cos(J\Delta t/2) e^{-i\Delta t/2} \quad (17) \\ U_{j,j+1} = i\hat{n}_j\hat{\sigma}_{j+1}^x \sin(J\Delta t/2) e^{-i\Delta t/2},$$

which can be seen as a *tilted*-CNOT, i.e. a CNOT which does not necessarily perform conditioned perfect spin-flips. This opens up the possibility that a subgroup of such a family of circuits, which does not belong to Clifford circuits (apart for fine tuned parameters [37]), could be efficiently simulated via classical algorithms. In such direction, the Floquet version of the quantum East model [38] has been shown to display localization starting from a single excitation, despite the absence of energy conservation. However, analogously to what is here discussed in the continuous time version ($\Delta t \rightarrow 0$), also in the Trotterized one the initial state could play a role in dynamics. A detailed analysis of such a circuit could lead to the discovery of a novel family of *tilted*-CNOTs that can be simulated efficiently, alongside Clifford, MBL inspired [53], as well as fractonic random circuits [48, 54].

Going beyond a deterministic evolution, an intriguing direction could be the addition of randomness in the circuit. We can envision different ways to do that. For instance, by randomly applying gates belonging to the one listed in Eq. (17) with random J . In such a scenario, we envision that the system could display an entanglement transition between volume law to area law (if J is sampled mostly from the localized side, namely $J < 1$). Another possibility is applying Eq. (17) deterministically for time τ (even by keeping Δt small so that we are approximating the continuous time evolution), and then applying some gates to each site with probability p , and repeat. In such a protocol, notice that the density of random gates in the total space-time volume is finite. By doing so, we could naively envision that the addition of gates preserving the directional character together with the sharp differentiation between active (given by $|1\rangle$) and inactive (given by $|0\rangle$) regions should not spoil the main features of the quantum East model. However, we observe that such a picture is challenged




FIG. 8. Snapshots at different times $te^{-s} = \{15, 25, \dots, 75\}$ (from light to dark curves) of the entanglement entropy on each cut under the quantum East model deep in the localized phase ($s = 1.5$) with Z gates applied with probability p on each site after a cycling time $\tau = 0.5$. For $p = 0$, entanglement remains localized near the initially seeded excitation. For $p > 0$, the active region heats up, as indicated by the linear growth of S_j on each j (until saturating its upper bound given by an infinite temperature thermal state), and expands diffusively towards the inactive region (see inset). Results averaged over 80 random realization for $p = 0.10$ (shaded areas are statistical uncertainties). The initial state is $|1\rangle \otimes_{j=2}^N |0\rangle$. The value of τ does not qualitatively change the observed behavior.

also by the random application of the Z -gate. Indeed, even an infinitesimal p , leads to the disruption of the slow entanglement growth, making it ballistic, and locally heats up the system to infinite temperature as a result of the non-commutativity of Z -gates with the kinetic constraint terms. To show this, we initialize a state with a single excitation. We observe that the Z -gate heats the system solely in the already active regions as expected, which in turn heats the surroundings. Intriguingly, the application of Z gates turns the spread of entanglement (see Fig. 8) from subdiffusive (for $p = 0$, corresponding to the deterministic dynamics) to diffusive (for $p > 0$). Such phenomenon is reminiscent of avalanches in MBL systems [82–86], and it could potentially serve as a toy model for their investigation.

Acknowledgements We thank Juan P. Garrahan and Shane P. Kelly for insightful discussions. This project has been supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Project-ID 429529648 – TRR 306 QuCoLiMa (“Quantum Cooperativity of Light and Matter”); and by the Dynamics and Topology Center funded by the State of Rhineland Palatinate. Parts of this research were conducted using the Mogon supercomputer and/or advisory services offered by Johannes Gutenberg University Mainz (hpc.uni-mainz.de), which is a member of the AHRP (Alliance for High Performance Computing in Rhineland Palatinate, www.ahrp.info) and the Gauss Alliance e.V. We gratefully acknowledge the computing time granted on the Mogon supercomputer at Johannes Gutenberg University Mainz (hpc.uni-mainz.de) through the project “DysQCorr”. M.B. acknowledges the support and the re-

sources provided by PARAM Shivay Facility under the National Supercomputing Mission, Government of India at the Indian Institute of Technology, Varanasi

Appendix A: Full spectrum of the Hamiltonian for small system size

Here we reproduce the results from Ref [34] showing the presence of exponentially large number of non-thermal eigenstates with large overlap with product states. To do so, we compute the full spectrum of the quantum East model (cf. Eq. (1)) for small system sizes via exact diagonalization. We fix the first site in the $|1\rangle$ state. In the upper panel of Fig. 9 we show the half-cut entanglement entropy S of the eigenstates $|\psi_\epsilon\rangle$ in the delocalized regime ($s = -0.5$) and the localized regime ($s = 0.5$) as a function of the normalized energy $\epsilon = (E_n - E_{\min})/(E_{\max} - E_{\min})$, where E_n is the energy of the n -th eigenstate, E_{\max} is the maximum energy, and E_{\min} is the ground state energy. In the delocalized ($s < 0$) phase the spectrum appears to be thermal as expected, while instead in the localized regime ($s > 0$) there are many non-thermal eigenstates with a large overlap with the product states (see lower-panel). The product states $\{|\psi_p\rangle\}$ where obtained by writing the eigenstates $\{|\psi_\epsilon\rangle\}$ as MPS and then truncate their bond-dimension to 1 on each bond.




FIG. 9. **Upper panel:** half-cut entanglement entropy of the eigenstates $|\psi_\epsilon\rangle$ of the quantum East model (cf. Eq. (1)) as a function of the normalized energy $\epsilon = (E - E_{\min})/(E_{\max} - E_{\min})$. **Lower panel:** overlap between $|\psi_\epsilon\rangle$ and its best approximating product state $|\psi_p\rangle$ as a function of ϵ . In the delocalized phase ($s < 0$) the spectrum appears to be thermal, as typical for non-integrable systems. Instead, in the localized phase ($s > 0$) there are (exponentially) many non-thermal eigenstates with large overlap with product states. We show data for $N = 13$ sites.




FIG. 10. Dynamics of the maximum entanglement entropy of ~ 100 product states (cf. Eq. (6)) at fixed normalized energy $\epsilon = (\langle H \rangle - E_{\min})/(E_{\max} - E_{\min}) = 0.4$ and different values of s . Deep in the delocalized phase ($s < 0$), we observe a linear growth of S_{\max} as expected for thermal systems. Whereas in the localized phase ($s > 0$), we have logarithmic growth of S_{\max} due to localization.

Appendix B: Entanglement entropy dynamics

In Fig. 2(a) we have located a state-dependent mobility edge for initial product states using the way $\chi_{\max}(t)$ grows in time, distinguishing a ‘hard’ and ‘easy’ time-complexity region. Here we present the data for the dynamics of max entanglement entropy $S_{\max}(t)$ along the system, to support the arguments presented in Sec. III A.

For states represented in the form of MPS with bond dimension χ , the maximal entanglement entropy is limited to $\log \chi$. Thus, if $S_{\max}(t) \propto t$, the bond-dimension needed grows exponentially. Instead, if S_{\max} grows sub-linearly, e.g. logarithmically as occurs for localized systems, the needed bond dimension scales polynomially. In Fig. 10, we show the data for the growth of S_{\max} for a sample of product states at a fixed $\epsilon = 0.4$ and different s . For states deep in the delocalized regime (e.g. $s = -0.3$), we have a linear growth of $S_{\max}(t)$, as expected in the thermal regime. In the localized phase $s > 0$, we observe $S_{\max}(t) \propto \log t$, indicating non-thermal properties.

Appendix C: Finite size scaling

In the section III A, we uncover a mobility edge in the system for a class of product states. We studied the model on a system size of $N = 30$, and the growth rate of χ was used as an indicator to classify ‘hard’ or ‘easy’ regimes. Here, we perform a finite-size scaling analysis in the system size N . To this end, we perform two different types of finite-size scaling. The first is to show that boundary effects are irrelevant up to the times reached. The second instead, is to properly address sampling biases.




FIG. 11. Dynamics of product states (cf. Eq. (C1)) with similar spatial structure and same ϵ for different system sizes N . (a) Dynamics of $\chi_{\max}(t)$ for different ϵ and system size $N \in [30, 60, 90, 120, 150]$. The curves for different N cannot be distinguished as they overlap, proving the absence of finite size effects. (b-c) exponential rate r and power-law exponent α as a function of N for different ϵ . Both r and α are independent of the system size N .

In the first scaling analysis, we compare states with similar local spatial structures. We do so, as we have established in the main text that spatial structure plays a vital role in dictating the time-complexity. In practice, we consider a randomly sampled product state $|\psi_{N=30}\rangle$ for system size $N = 30$ with energy density ϵ , and we obtain states for larger N by simply concatenating copies of them

$$|\psi_N\rangle = |\psi_{N=30}\rangle^{[N/30]} \quad (\text{C1})$$

where we restrict to system sizes N multiple of 30. We highlight that these states have comparable same energy density ϵ (apart for $1/N$ corrections). We choose three representative values of $\epsilon = \{0.22, 0.54, 0.8\}$ at fixed $s = 0.5$. Then, we simulate the dynamics of such states up to time $T = 64$ via TEBD. In Fig. 11(a) we show the dynamics of $\chi_{\max}(t)$ for different values of energy density ϵ . Each curve with a specific color consists of five overlapping curves for system size $N = \{30, 60, 90, 120, 150\}$. The overlap gives a clear indication that growth dynamics and hence our results are unaffected by boundary effects. As an additional test, in Fig. 11(b-c) we show the computed exponential rates r and power-law exponents α extracted by fitting $\chi_{\max}(t)$ for different ϵ and N . Also here, there are no signatures of finite-size effects.

In the second scaling analysis (see Fig. 3), we instead compare results obtained via randomly sampling initial states that do not necessarily share the same spatial structure. Specifically, we perform the same sampling procedure discussed in Sec. III B for different system sizes N . For the sake of clarity, we show results along two vertical cuts of Fig. 2(a) fixing s and swiping the energy density ϵ . Deep in the localized phase, results are not affected by changing the system size N , and dynamics are always easily computable. Instead, the more we get near the transition point $s = 0$ the more the results can be affected by changing the system size N . Based on our scaling analysis, we observe that for $0 \lesssim s \lesssim 0.3$ the different curves obtained for different N do not tend to collapse, and the value at which the transition looks to decrease to




FIG. 12. Same data presented in Fig. 4(a) without rescaling w_{avg} with the ground state localization length ξ . The red dots correspond to hard time-complexity with r the corresponding rate. The blue dots correspond to easy time-complexity with α the power-law. Both hard and easy simulations overlap as the effect of ξ on growth dynamics is not considered, highlighting its key role also out-of-equilibrium.

zero. We give two possible interpretations of such a result: (a) no mobility edge exists in such parameter regime and solely the ground state is localized; (b) the number of initial states sampled is not large enough to obtain convergent results for s small, as other features of the initial state play a key role in dictating the dynamics, such as the initial distribution of excitations (see Sec. III D). Instead, for $0.3 \lesssim s \lesssim 0.5$ we observe that upon increasing N the inversion occurs at a size-independent ϵ , indicating the existence of a state-dependent mobility edge.

Appendix D: Role of the equilibrium localization length ξ in dictating time-complexity

In Sec. III D we have discussed how the localization length ξ of the ground state provides a length scale. To prove its predictive power, in Fig. 12 we show the same data as in Fig. 4 without rescaling the average distance between excitations w_{avg} with ξ . We see that the ‘hard’ and the ‘easy’ simulations completely overlap and we are not able to differentiate between them.

Appendix E: Properties of the localized kink states found via DMRG-X

Here we provide some additional results on the states $|\psi_X\rangle$ obtained via DMRG-X. We refer to Ref. [47] for the interested reader on the technical details. In Fig. 13 we show the energy variance ΔH computed on $|\psi_X\rangle$ (setting a maximal bond dimension $\chi = 100$) in a system of size $N = 30$. We only show the data points for which $\Delta H \leq 10^{-5}$. In Fig. 13 we show the overlap between the states $|\psi_X\rangle$ and the kink states $|\mathbf{k}\rangle = |1\rangle^{\otimes k} \otimes |0\rangle^{\otimes (N-k)}$ seeded




FIG. 13. **Left panel:** energy variance $\Delta E_{\mathbf{x}} = (\langle \psi_{\mathbf{x}} | \hat{H}^2 | \psi_{\mathbf{x}} \rangle - \langle \psi_{\mathbf{x}} | \hat{H} | \psi_{\mathbf{x}} \rangle^2)$ computed on the states found via DMRG-X upon seeding kink states $|\mathbf{k}\rangle = |1\rangle^{\otimes k} \otimes |0\rangle^{\otimes (N-k)}$. We consider the algorithm converged solely if $\Delta E_{\mathbf{x}} \leq 10^{-5}$. **Right panel:** modulus of the overlap between the states obtained via DMRG-X $|\psi_{\mathbf{x}}\rangle$ and the initially seeded kink states $|\mathbf{k}\rangle = |1\rangle^{\otimes k} \otimes |0\rangle^{\otimes (N-k)}$. Results obtained for system size $N = 30$.

in the variational approach. We observe large overlap (way larger than the typical overlap $1/\mathcal{D}$ with $\mathcal{D} = 2^N$ the Hilbert space dimension), indicating that such states are highly relevant in dictating the dynamical features of the kink states $|\mathbf{k}\rangle$.

-
- [1] J. Preskill, Quantum computing in the NISQ era and beyond, *Quantum* **2**, 79 (2018).
- [2] R. P. Feynman, Simulating physics with computers, *International Journal of Theoretical Physics* **21**, 467–488 (1982).
- [3] A. J. Daley, I. Bloch, C. Kokail, S. Flannigan, N. Pearson, M. Troyer, and P. Zoller, Practical quantum advantage in quantum simulation, *Nature* **607**, 667–676 (2022).
- [4] A. Miessen, P. J. Ollitrault, F. Tacchino, and I. Tavernelli, Quantum algorithms for quantum dynamics, *Nature Computational Science* **3**, 25–37 (2022).
- [5] Y. Kim, A. Eddins, S. Anand, K. X. Wei, E. van den Berg, S. Rosenblatt, H. Nayfeh, Y. Wu, M. Zaletel, K. Temme, and A. Kandala, Evidence for the utility of quantum computing before fault tolerance, *Nature* **618**, 500–505 (2023).
- [6] J. Tindall, M. Fishman, E. M. Stoudenmire, and D. Sels, Efficient tensor network simulation of ibm’s eagle kicked ising experiment, *PRX Quantum* **5**, 010308 (2024).
- [7] S. Patra, S. S. Jahromi, S. Singh, and R. Orús, Efficient tensor network simulation of ibm’s largest quantum processors, *Phys. Rev. Res.* **6**, 013326 (2024).
- [8] H.-J. Liao, K. Wang, Z.-S. Zhou, P. Zhang, and T. Xiang, Simulation of ibm’s kicked ising experiment with projected entangled pair operator (2023), arXiv:2308.03082 [quant-ph].
- [9] S. Anand, K. Temme, A. Kandala, and M. Zaletel, Classical benchmarking of zero noise extrapolation beyond the exactly-verifiable regime (2023), arXiv:2306.17839 [quant-ph].
- [10] T. Begušić, J. Gray, and G. K.-L. Chan, Fast and converged classical simulations of evidence for the utility of quantum computing before fault tolerance, *Science Advances* **10**, 10.1126/sciadv.adk4321 (2024).
- [11] M. S. Rudolph, E. Fontana, Z. Holmes, and L. Cincio, Classical surrogate simulation of quantum systems with lowesa (2023), arXiv:2308.09109 [quant-ph].
- [12] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing many-body dynamics on a 51-atom quantum simulator, *Nature* **551**, 579 (2017).
- [13] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Weak ergodicity breaking from quantum many-body scars, *Nature Physics* **14**, 745 (2018).
- [14] S. Choi, C. J. Turner, H. Pichler, W. W. Ho, A. A. Michailidis, Z. Papić, M. Serbyn, M. D. Lukin, and D. A. Abanin, Emergent su(2) dynamics and perfect quantum many-body scars, *Phys. Rev. Lett.* **122**, 220603 (2019).
- [15] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum many-body scars and weak breaking of ergodicity, *Nature Physics* **17**, 675 (2021).
- [16] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Quantum scarred eigenstates in a rydberg atom chain: Entanglement, breakdown of thermalization, and stability to perturbations, *Phys. Rev. B* **98**, 155134 (2018).
- [17] V. Khemani, C. R. Laumann, and A. Chandran, Signatures of integrability in the dynamics of rydberg-blockaded chains, *Phys. Rev. B* **99**, 161101(R) (2019).
- [18] C. J. Turner, J.-Y. Desaulles, K. Bull, and Z. Papić, Correspondence principle for many-body scars in ultracold rydberg atoms, *Phys. Rev. X* **11**, 021021 (2021).
- [19] S. Moudgalya, B. A. Bernevig, and N. Regnault, Quantum many-body scars and hilbert space fragmentation: a review of exact results, *Reports on Progress in Physics* **85**, 086501 (2022).
- [20] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, Quantum many-body scars: A quasiparticle perspective, *Annual Review of Condensed Matter Physics* **14**, 443 (2023).
- [21] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, *Annual Review of Condensed Matter Physics* **6**, 15 (2015), <https://doi.org/10.1146/annurev-conmatphys-031214-014726>.
- [22] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, *Rev. Mod. Phys.* **91**, 021001 (2019).
- [23] M. Kormos, M. Collura, G. Takács, and P. Calabrese, Real-time confinement following a quantum quench to a non-integrable model, *Nature Physics* **13**, 246 (2017).

- [24] A. Lerose, F. M. Surace, P. P. Mazza, G. Perfetto, M. Collura, and A. Gambassi, Quasilocalized dynamics from confinement of quantum excitations, *Phys. Rev. B* **102**, 041118(R) (2020).
- [25] F. M. Surace, P. P. Mazza, G. Giudici, A. Lerose, A. Gambassi, and M. Dalmonte, Lattice gauge theories and string dynamics in rydberg atom quantum simulators, *Phys. Rev. X* **10**, 021041 (2020).
- [26] F. Liu, R. Lundgren, P. Titum, G. Pagano, J. Zhang, C. Monroe, and A. V. Gorshkov, Confined quasiparticle dynamics in long-range interacting quantum spin chains, *Phys. Rev. Lett.* **122**, 150601 (2019).
- [27] R. Javier Valencia Tortora, P. Calabrese, and M. Collura, Relaxation of the order-parameter statistics and dynamical confinement, *Europhysics Letters* **132**, 50001 (2020).
- [28] O. Pomponio, M. A. Werner, G. Zarand, and G. Takacs, Bloch oscillations and the lack of the decay of the false vacuum in a one-dimensional quantum spin chain, *SciPost Phys.* **12**, 061 (2022).
- [29] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded growth of entanglement in models of many-body localization, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [30] M. Serbyn, Z. Papić, and D. A. Abanin, Universal slow growth of entanglement in interacting strongly disordered systems, *Phys. Rev. Lett.* **110**, 260601 (2013).
- [31] A. Ehrenberg, A. Deshpande, C. L. Baldwin, D. A. Abanin, and A. V. Gorshkov, Simulation complexity of many-body localized systems, arXiv preprint arXiv:2205.12967 (2022).
- [32] H. Kim and D. A. Huse, Ballistic spreading of entanglement in a diffusive nonintegrable system, *Phys. Rev. Lett.* **111**, 127205 (2013).
- [33] M. van Horssen, E. Levi, and J. P. Garrahan, Dynamics of many-body localization in a translation-invariant quantum glass model, *Phys. Rev. B* **92**, 100305 (2015).
- [34] N. Pancotti, G. Giudice, J. I. Cirac, J. P. Garrahan, and M. C. Bañuls, Quantum east model: Localization, non-thermal eigenstates, and slow dynamics, *Phys. Rev. X* **10**, 021051 (2020).
- [35] R. J. Valencia-Tortora, N. Pancotti, M. Fleischhauer, H. Bernien, and J. Marino, Rydberg platform for non-ergodic chiral quantum dynamics, *Phys. Rev. Lett.* **132**, 223201 (2024).
- [36] T. Bhore, J.-Y. Desaulles, and Z. Papić, Deep thermalization in constrained quantum systems, *Phys. Rev. B* **108**, 104317 (2023).
- [37] B. Bertini, C. De Fazio, J. P. Garrahan, and K. Klobas, Exact quench dynamics of the floquet quantum east model at the deterministic point, *Phys. Rev. Lett.* **132**, 120402 (2024).
- [38] B. Bertini, P. Kos, and T. c. v. Prosen, Localized dynamics in the floquet quantum east model, *Phys. Rev. Lett.* **132**, 080401 (2024).
- [39] R. J. Valencia-Tortora, N. Pancotti, and J. Marino, Kinetically constrained quantum dynamics in superconducting circuits, *PRX Quantum* **3**, 020346 (2022).
- [40] A. Geißler and J. P. Garrahan, Slow dynamics and nonergodicity of the bosonic quantum east model in the semiclassical limit, *Physical Review E* **108**, 034207 (2023).
- [41] P. Brighi, M. Ljubotina, and M. Serbyn, Hilbert space fragmentation and slow dynamics in particle-conserving quantum east models, *SciPost Physics* **15**, 093 (2023).
- [42] Z. Lan, M. van Horssen, S. Powell, and J. P. Garrahan, Quantum slow relaxation and metastability due to dynamical constraints, *Phys. Rev. Lett.* **121**, 040603 (2018).
- [43] J. P. Garrahan, R. L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk, and F. van Wijland, First-order dynamical phase transition in models of glasses: an approach based on ensembles of histories, *Journal of Physics A: Mathematical and Theoretical* **42**, 075007 (2009).
- [44] P. Chleboun, A. Faggionato, and F. Martinelli, Time scale separation in the low temperature east model: rigorous results, *J. Stat. Mech.* **2013**, L04001 (2013).
- [45] J. P. Garrahan, Aspects of non-equilibrium in classical and quantum systems: Slow relaxation and glasses, dynamical large deviations, quantum non-ergodicity, and open quantum dynamics, *Physica A: Statistical Mechanics and its Applications* **504**, 130 (2018).
- [46] L. Causser, I. Lesanovsky, M. C. Bañuls, and J. P. Garrahan, Dynamics and large deviation transitions of the xor-fredrickson-andersen kinetically constrained model, *Phys. Rev. E* **102**, 052132 (2020).
- [47] V. Khemani, F. Pollmann, and S. L. Sondhi, Obtaining highly excited eigenstates of many-body localized hamiltonians by the density matrix renormalization group approach, *Phys. Rev. Lett.* **116**, 247204 (2016).
- [48] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, Ergodicity breaking arising from hilbert space fragmentation in dipole-conserving hamiltonians, *Phys. Rev. X* **10**, 011047 (2020).
- [49] S. Moudgalya and O. I. Motrunich, Hilbert space fragmentation and commutant algebras, *Phys. Rev. X* **12**, 011050 (2022).
- [50] I. Mondragon-Shem, A. Pal, T. L. Hughes, and C. R. Laumann, Many-body mobility edge due to symmetry-constrained dynamics and strong interactions, *Phys. Rev. B* **92**, 064203 (2015).
- [51] D. J. Luitz, N. Laflorencie, and F. Alet, Many-body localization edge in the random-field heisenberg chain, *Phys. Rev. B* **91**, 081103 (2015).
- [52] P. Naldesi, E. Ercolessi, and T. Roscilde, Detecting a many-body mobility edge with quantum quenches, *SciPost Physics* **1**, 10.21468/scipostphys.1.1.010 (2016).
- [53] C. Sünderhauf, D. Pérez-García, D. A. Huse, N. Schuch, and J. I. Cirac, Localization with random time-periodic quantum circuits, *Phys. Rev. B* **98**, 134204 (2018).
- [54] S. Pai, M. Pretko, and R. M. Nandkishore, Localization in fractonic random circuits, *Phys. Rev. X* **9**, 021003 (2019).
- [55] I. Lesanovsky and J. P. Garrahan, Kinetic constraints, hierarchical relaxation, and onset of glassiness in strongly interacting and dissipative rydberg gases, *Phys. Rev. Lett.* **111**, 215305 (2013).
- [56] C. Pérez-Espigares, I. Lesanovsky, J. P. Garrahan, and R. Gutiérrez, Glassy dynamics due to a trajectory phase transition in dissipative rydberg gases, *Phys. Rev. A* **98**, 021804(R) (2018).
- [57] M. M. Valado, C. Simonelli, M. D. Hoogerland, I. Lesanovsky, J. P. Garrahan, E. Arimondo, D. Ciampini, and O. Morsch, Experimental observation of controllable kinetic constraints in a cold atomic gas, *Phys. Rev. A* **93**, 040701(R) (2016).
- [58] I. Lesanovsky and J. P. Garrahan, Out-of-equilibrium structures in strongly interacting rydberg gases with dissipation, *Phys. Rev. A* **90**, 011603(R) (2014).
- [59] D. Gribben, I. Lesanovsky, and R. Gutiérrez, Quench dynamics of a dissipative rydberg gas in the classical and quantum regimes, *Phys. Rev. A* **97**, 011603(R) (2018).
- [60] M. Ostmann, M. Marcuzzi, J. P. Garrahan, and I. Lesanovsky, Localization in spin chains with facilitation

- tion constraints and disordered interactions, Phys. Rev. A **99**, 060101(R) (2019).
- [61] M. Marcuzzi, J. c. v. Minář, D. Barredo, S. de Léséleuc, H. Labuhn, T. Lahaye, A. Browaeys, E. Levi, and I. Lesanovsky, Facilitation dynamics and localization phenomena in rydberg lattice gases with position disorder, Phys. Rev. Lett. **118**, 063606 (2017).
- [62] Z. Lan, M. van Horssen, S. Powell, and J. P. Garrahan, Quantum slow relaxation and metastability due to dynamical constraints, Physical review letters **121**, 040603 (2018).
- [63] L. Zadnik and J. P. Garrahan, Slow heterogeneous relaxation due to constraints in dual xxz models, Physical Review B **108**, L100304 (2023).
- [64] M. P. Fisher, V. Khemani, A. Nahum, and S. Vijay, Random quantum circuits, Annual Review of Condensed Matter Physics **14**, 335–379 (2023).
- [65] D. E. Parker, X. Cao, A. Avdoshkin, T. Scaffidi, and E. Altman, A universal operator growth hypothesis, Phys. Rev. X **9**, 041017 (2019).
- [66] E. Rabinovici, A. Sánchez-Garrido, R. Shir, and J. Sonner, Krylov complexity from integrability to chaos, Journal of High Energy Physics **2022**, 1 (2022).
- [67] H. G. Menzler and R. Jha, Krylov localization as a probe for ergodicity breaking, arXiv preprint arXiv:2403.14384 (2024).
- [68] M. C. Bañuls, N. Y. Yao, S. Choi, M. D. Lukin, and J. I. Cirac, Dynamics of quantum information in many-body localized systems, Phys. Rev. B **96**, 174201 (2017).
- [69] Y. Li, P. Sala, and F. Pollmann, Hilbert space fragmentation in open quantum systems, Phys. Rev. Res. **5**, 043239 (2023).
- [70] G. De Tomasi, D. Hetterich, P. Sala, and F. Pollmann, Dynamics of strongly interacting systems: From fock-space fragmentation to many-body localization, Phys. Rev. B **100**, 214313 (2019).
- [71] L. Herviou, J. H. Bardarson, and N. Regnault, Many-body localization in a fragmented hilbert space, Phys. Rev. B **103**, 134207 (2021).
- [72] S. Moudgalya, A. Prem, R. Nandkishore, N. Regnault, and B. A. Bernevig, Thermalization and its absence within krylov subspaces of a constrained hamiltonian, in *Memorial Volume for Shoucheng Zhang* (WORLD SCIENTIFIC, 2021) p. 147–209.
- [73] Y. Han, X. Chen, and E. Lake, Exponentially slow thermalization and the robustness of hilbert space fragmentation, arXiv preprint arXiv:2401.11294 (2024).
- [74] G. Vidal, Efficient classical simulation of slightly entangled quantum computations, Phys. Rev. Lett. **91**, 147902 (2003).
- [75] J. Gray, quimb: A python package for quantum information and many-body calculations, Journal of Open Source Software **3**, 819 (2018).
- [76] E. H. Lieb and D. W. Robinson, The finite group velocity of quantum spin systems, Communications in mathematical physics **28**, 251 (1972).
- [77] J. Haah, M. B. Hastings, R. Kothari, and G. H. Low, Quantum algorithm for simulating real time evolution of lattice hamiltonians, SIAM Journal on Computing **52**, FOCS18 (2021).
- [78] C.-F. (Anthony) Chen, A. Lucas, and C. Yin, Speed limits and locality in many-body quantum dynamics, Reports on Progress in Physics **86**, 116001 (2023).
- [79] S. Aaronson and A. Arkhipov, The computational complexity of linear optics (2010), arXiv:1011.3245 [quant-ph].
- [80] A. P. Lund, M. J. Bremner, and T. C. Ralph, Quantum sampling problems, bosonsampling and quantum supremacy, npj Quantum Information **3**, 10.1038/s41534-017-0018-2 (2017).
- [81] A. Deshpande, B. Fefferman, M. C. Tran, M. Foss-Feig, and A. V. Gorshkov, Dynamical phase transitions in sampling complexity, Phys. Rev. Lett. **121**, 030501 (2018).
- [82] W. De Roeck and F. m. c. Huveneers, Stability and instability towards delocalization in many-body localization systems, Phys. Rev. B **95**, 155129 (2017).
- [83] T. Thiery, F. m. c. Huveneers, M. Müller, and W. De Roeck, Many-body delocalization as a quantum avalanche, Phys. Rev. Lett. **121**, 140601 (2018).
- [84] D. J. Luitz, F. m. c. Huveneers, and W. De Roeck, How a small quantum bath can thermalize long localized chains, Phys. Rev. Lett. **119**, 150602 (2017).
- [85] A. Morningstar, L. Colmenarez, V. Khemani, D. J. Luitz, and D. A. Huse, Avalanches and many-body resonances in many-body localized systems, Phys. Rev. B **105**, 174205 (2022).
- [86] D. Sels, Bath-induced delocalization in interacting disordered spin chains, Phys. Rev. B **106**, L020202 (2022).