ON THE CONFORMAL GROUP OF A GLOBALLY HYPERBOLIC SPACETIME

ALI BLEYBEL

ABSTRACT. The Groups of causal and conformal automorphisms of globally hyperbolic spacetimes were studied.

In two dimensions, we prove that all globally hyperbolic spacetimes that are directed and connected are causally isomorphic. We work out the consequences of this fact and obtain a (partial) classification of the causal automorphism and conformal groups of a two-dimensional globally hyperbolic space.

Finally, we present a generalization for locally conformally flat globally hyperbolic spacetimes.

1. INTRODUCTION

In General Relativity, conformal mappings are important because they preserve the causal structure (up to time orientation) and light-like geodesics up to parametrization.

Zeeman, [11], showed that the group of causal automorphisms of Minkowski spacetime (of dimension > 2) coincides with the group G generated by orthochronous Lorentz transformations, space-time translations, and dilatations. Later, in [9], Levichev generalized these results to smooth time-orientable Lorentzian manifolds M with dimensions > 2. More precisely, he showed that any causal automorphism of M is automatically conformal, provided that it is past and future distinguishing.

Although there are many results concerning the conformal group of a compact Lorentzian manifold (e.g., [10]), results concerning the group of conformal automorphisms of spacetime are difficult to find in the litterature (especially physics litterature).

This study addresses this problem using the back-and-forth method (reminiscent of model theory). We obtain a partial result, namely for two-dimensional globally hyperbolic spacetimes. In particular, we isolate an order-theoretic condition satisfied by spacetimes causally isomorphic to two-dimensional Minkowski spacetime. These results might be of interest especially in the light of the celestial holography program.

While the results obtained using our method can probably be shown via more standard methods (especially in the 2-dimensional case), it is worthwhile to note that the importance of this work lies in its potential for generalization to higher dimensional spacetimes. Furthermore, in the 2-dimensional case, this paper is largely self contained.

The rest of this paper is organized as follows. Section 2 outlines the background material necessary for the rest of the paper and the notation and introductory

results. In Section 3, we introduce and prove the main theorem. Section 4 provides a partial classification of the causal and conformal automorphism groups. Section 5 generalizes the results obtained in the case of locally conformally flat globally hyperbolic spacetimes.

2. Preliminaries

We assume the reader is familiar with elementary first-order logic and model theory.

2.1. Order theory. Let (X, \prec) be a partially ordered set, with the order relation denoted by \prec . Recall that \prec , a binary relation, is a subset of the Cartesian product $X \times X$. Being an order relation, \prec is a reflexive, transitive, and antisymmetric relation on X. If A is a subset of X, the *induced order* on A is the binary relation $\prec |_A := (A \times A) \cap \prec \ldots \prec |_A$ is an order relation on A.

Given (partially) ordered sets X, Y, a partial isomorphism $X \rightarrow Y$ is an order isomorphism $A \rightarrow B$ with $A \subset X, B \subset Y$ are equipped with the induced orders from X and Y respectively.

The set $\{x \in \mathfrak{C} | a \prec x \prec b\}$ will be denoted by [a, b] and is called the Alexandrov interval with endpoints a, b. For a spacetime \mathcal{M} , an interval [a, b] is also called a (closed) causal diamond.

By a *pattern*, we mean a finite set equipped with a strict order relation.

2.2. The setting. We will be using the following signature

$$L := \{\twoheadrightarrow, \ll\},$$

where \rightarrow , \ll are two binary relation symbols.

2.2.1. Theory. We consider the theory T, which consists of all statements which are logical consequences of the following axioms:

- (1) $\forall x (x \ll x)$ (2) $\forall x \forall y \forall z ((x \ll y \& y \ll z) \rightarrow x \ll z)$
- $(3) \ \forall x (x \twoheadrightarrow x)$
- (4) $\forall x \forall y ((x \twoheadrightarrow y \& y \twoheadrightarrow x) \to x = y)$
- (5) .not. $(x \ll y \& x \twoheadrightarrow y)$.

And for each natural number n > 0, the following axiom: (6)_n

$$\forall x_1 \forall x_2 \dots \forall x_n \left[\left(\bigwedge_{i=1}^{n-1} x_i \twoheadrightarrow x_{i+1} \& x_1 \twoheadrightarrow x_n \right) \longrightarrow \bigwedge_{j,k,1 \le j \le k \le n} x_j \twoheadrightarrow x_k \right].$$

2.2.2. Types and quantifier-free types. Let n > 0 and let $\bar{x} := (x_1, \ldots, x_n)$ be a tuple of *n*-variables.

An *n*-type $t = t(\bar{x})$ in T is a set of formulas $\phi(\bar{x})$, such that for each formula $\psi(\bar{x}) \in t(\bar{x})$, there exists a model M of T and a tuple of elements $(a_1, \ldots, a_n) \in M^n$ such that $\psi(a_1, \ldots, a_n)$ holds in M.

The case where n = 1 corresponds to 1-types.

We say that $t(\bar{x})$ is a quantifier-free type if all formulas in t are quantifier-free.

For a tuple $\bar{a} \in M^n$, n > 0, we denote by $\operatorname{tp}(\bar{a})$ the set of all formulas $\phi(\bar{x})$ such that $\phi(\bar{a})$ holds in M. Let X be a set of elements of M, and let \bar{b} be a listing of the elements of X (so \bar{b} can be of infinite length). The type of \bar{a} over X, denoted by $\operatorname{tp}(\bar{a}/\bar{b})$ (or, equivalently, $\operatorname{tp}(\bar{a}/X)$) is the set of all formulas $\phi(\bar{x}, \bar{y})$ (where \bar{y} is a tuple of variables), such that $\phi(\bar{a}, \bar{b}')$ holds in M. Here \bar{b}' is some tuple of elements of X of the same length as \bar{y} .

The quantifier-free type of \bar{a} over X, denoted by $qftp(\bar{a}/X)$ is the set of all quantifier-free formulas $\phi(\bar{x}, \bar{y})$ such that $\phi(\bar{a}, \bar{b}')$ holds in M.

2.3. Corresponding notions in the context of spacetime physics. The causality relation in the context of spacetime physics is denoted by $\prec: x \prec y$ iff $x \in J^-(y)$ where $J^-(y)$ is the past causal cone of y; note that $y \in J^-(y)$. We will also make use of the relation $\ll: x \ll y$ (Chronology) iff $x \in I^-(y)$ (x belongs to the *open* past lightcone of y). For a globally hyperbolic spacetime, the relation \ll is transitive and irreflexive; it follows that \ll is antisymmetric. The relation \ll is then a strict order.

Let $X \subset \mathcal{M}$ be a finite set. Then X is a *strict pattern* if for all $x, y \in X, x \neq y$, $x \ll y$ or $x \not\prec y$. The point of this definition is to exclude non-open relations. The statements $x \ll y$ and $I^{-}(y) \cap I^{+}(x) \neq \emptyset$ are equivalent. Other equivalent statements are $I^{+}(y) \subsetneq I^{+}(x)$ and $I^{-}(x) \subsetneq I^{-}(y)$.

The expression $x \to y$ (used earlier in the context of the Theory T) is defined here as $x \prec y \& x \ll y$.

We denote by x^{\perp} the set of elements causally unrelated to x, i.e. $\{y \in \mathcal{M} | x \not\prec y \& y \not\prec x\}$. The relation $y \in x^{\perp}$ is also denoted by x || y.

2.4. Assertion. Assume $a \rightarrow b$. Then, for any c, we have

$$\begin{aligned} a \prec c \& c \twoheadrightarrow b &\to a \twoheadrightarrow c \\ b \ll c &\to a \ll c \\ b ||c &\to c \not\prec a \end{aligned}$$

2.5. *Remark.* The relation $x \prec y$ can be expressed as (modulo T):

$$x \prec y \equiv x \ll y$$
 .or. $x \twoheadrightarrow y$

where we used the following axiom (5):

.not.
$$(x \ll y \& x \twoheadrightarrow y)$$
.

To see this, observe that $x \to y \leftrightarrow (x \prec y \& x \ll y)$, and the required equivalence follows.

We have:

2.6. **Theorem.** Let \mathcal{M} be a globally hyperbolic spacetime. Then \mathcal{M} is a model of the theory T in the language \mathcal{L} .

Proof. It is a standard fact that the relations \rightarrow and \ll defined as in subsection 2.3, satisfy the axioms (1), through (5), and (6)_n of 2.2.1 for all n > 0. For details, please see [8].

2.7. Conformal transformations. A conformal diffeomorphism $\phi : \mathcal{M} \to \mathcal{N}$ where $(\mathcal{M}, g_{\mathcal{M}})$ and $(\mathcal{N}, g_{\mathcal{N}})$ are two *n*-dimensional pseudo-Riemannian manifolds of signature $(p, q), 0 \leq p \leq q, p + q = n$ is a C^{∞} -diffeomorphism $\varphi : \mathcal{M} \to \mathcal{N}$ such that

$$\varphi^*(g_{\mathcal{N}}) = e^{\psi}g_{\mathcal{M}},$$

for some arbitrary C^{∞} -function $\psi : \mathcal{M} \to \mathbb{R}$.

3. Causal isomorphisms of globally hyperbolic spacetimes

3.1. General considerations. Let \mathcal{C}, \mathcal{D} be models of T. A causal isomorphism $f : \mathcal{C} \to \mathcal{D}$ is a bijective map satisfying

$$\forall x, y \in \mathcal{C}, x \prec y \leftrightarrow f(x) \prec f(y)$$

The above definition applies to globally hyperbolic spacetimes \mathcal{M}, \mathcal{N} .

Let a, b be points in \mathcal{M}, \mathcal{N} respectively.

We have:

3.2. **Theorem.** Let \mathcal{M} and \mathcal{N} be globally hyperbolic spacetimes. Assume that \mathcal{M} and \mathcal{N} are upward and downward directed.

A partial map $\{a\} \to \{b\}$ can be extended to a causal isomorphism $\mathcal{M} \to \mathcal{N}$ if and only if for any finite tuples $(a_0 = a, a_2, \ldots, a_n)$ and $(b_0 = b, b_2, \ldots, b_n)$ with $qftp(a_1, \ldots, a_n) = qftp(b_1, \ldots, b_n)$ (in the language $L := \{\ll, \twoheadrightarrow\}$) the following conditions hold: and

$$(3.2.1) \qquad \mathcal{O}_0 \cap \mathcal{O}_1 \cap \dots \cap \mathcal{O}_n \neq \emptyset \longleftrightarrow \mathcal{O}'_0 \cap \mathcal{O}'_1 \cap \dots \cap \mathcal{O}'_n \neq \emptyset$$

where \mathcal{O}_i is one of $I^+(a_i), I^-(a_i), a_i^{\perp}$ (respectively \mathcal{O}'_i is one of $I^+(b_i), I^-(b_i), b_i^{\perp}$), and, for all $I, J \subset \{0, 1, \ldots, n\}$,

$$(3.2.2) \qquad \exists x \left[\bigwedge_{i \in I} a_i \twoheadrightarrow x \& \bigwedge_{j \in J} x \twoheadrightarrow a_j \right] \longleftrightarrow \exists y \left[\bigwedge_{i \in I} b_i \twoheadrightarrow y \& \bigwedge_{j \in J} y \twoheadrightarrow b_j \right]$$

Before proceeding to the proof of this Theorem, let us show the following:

3.3. Lemma. Keep the above notation and hypotheses (conditions 3.2.1 and 3.2.2). Assume that $c \in \mathcal{M}$ satisfies $[\bigwedge_{i \in I} a_i \twoheadrightarrow c \& \bigwedge_{j \in J} c \twoheadrightarrow a_j]$ for some $I, J \subset \{0, 1, \ldots, n\}$, and that, furthermore:

$$c \in \mathcal{O}_{i_0} \cap \mathcal{O}_{i_1} \cap \cdots \cap \mathcal{O}_{i_k},$$

for some open sets $\mathcal{O}_{i_{\ell}}$ where \mathcal{O}_{i} is one of $I^{+}(a_{i_{\ell}}), I^{-}(a_{i_{\ell}}), a_{i_{\ell}}^{\perp}, \ell = 0, 1, \dots, k, i_{\ell} \in \{0, 1, \dots, n\}.$

Then, we have:

$$\exists d[\bigwedge_{i\in I} b_i \twoheadrightarrow d \& \bigwedge_{j\in J} d \twoheadrightarrow b_j] \& d \in \mathcal{O}'_{i_0} \cap \mathcal{O}'_{i_1} \cap \dots \cap \mathcal{O}'_{i_k},$$

with $\mathcal{O}'_{i_{\ell}}, \ \ell = 0, 1, \dots, k$ are such that

$$\mathcal{O}_{i_{\ell}} = I^{\pm}(a_{i_{\ell}}) \leftrightarrow \mathcal{O}'_{i_{\ell}} = I^{\pm}(b_{i_{\ell}}) \& \mathcal{O}_{i_{\ell}} = a^{\perp}_{i_{\ell}} \leftrightarrow \mathcal{O}'_{i_{\ell}} = b^{\perp}_{i_{\ell}}.$$

Proof. We may suppose that $I \cap J = \emptyset$ (otherwise $c = a_m$ for some $m \in I \cap J$ and it suffices to take $d = b_m$).

Observe then that I, J and $\{i_0, i_1, \ldots, i_k\}$ are disjoint.

Recall that $\{x \in \mathcal{M} | x \twoheadrightarrow e \text{ Or } e \twoheadrightarrow x\} = \mathcal{M} \setminus (I^+(e) \cup I^-(e) \cup e^{\perp})\}.$

Let us consider the case $|I| = 1, |J| = 0, I = \{a_s\}$ for simplicity; the other case can be treated similarly.

Then, as $a_s \to c$, $c \in \mathcal{M} \setminus (I^+(a_s) \cup I^-(a_s) \cup a_s^{\perp}) =: \mathcal{F}$; it follows that $c \in \mathcal{F} \cap \mathcal{O}_{i_0} \cap \mathcal{O}_{i_1} \cap \cdots \cap \mathcal{O}_{i_k}$.

On the other hand, if $\mathcal{F}' \cap \mathcal{O}'_{i_0} \cap \mathcal{O}'_{i_1} \cap \cdots \cap \mathcal{O}'_{i_k} = \emptyset$ where $\mathcal{F}' = \mathcal{N} \setminus (I^+(b_s) \cup I^-(b_s) \cup b_s^{\perp})$ we get $\mathcal{O}'_{i_0} \cap \mathcal{O}'_{i_1} \cap \cdots \cap \mathcal{O}'_{i_k} \subset (I^+(b_s) \cup I^-(b_s) \cup b_s^{\perp})$. It can now be seen that the latter implies that there are relations among $\{b_{i_\ell}, b_s, \ell = 0, 1, \dots, n\}$ which are nonexistent among $\{a_{i_\ell}, a_s, \ell = 0, 1, \dots, n\}$ contradicting the hypotheses, hence the claim. \Box

3.3.1. Proof of Lemma. Let $\mathfrak{C}, \mathfrak{D}$ be dense subsets of \mathcal{M}, \mathcal{N} respectively. These sets are equipped with the induced relations \ll (Chronology) and \twoheadrightarrow (Horismos). Back & Forth:

A finite partial isomorphism $\mathcal{M} \dashrightarrow \mathcal{N}$ is an order isomorphism $p : A \to B$, with A, B finite subsets of \mathcal{M} and \mathcal{N} respectively, equipped with the corresponding induced order relations.

We will construct, inductively, a family of partial isomorphisms $p_i : A_i \to B_i$, for $i \in \mathbb{N}$ where A_i, B_i are finite subsets of $\mathfrak{C}', \mathfrak{D}'$ (with $\mathfrak{C} \subset \mathfrak{C}' \subset \mathcal{M}, \mathfrak{D} \subset \mathfrak{D}' \subset \mathcal{N}$ and $\mathfrak{C}', \mathfrak{D}'$ are countable), such that

$$(3.3.1) A_i \subset A_{i+1}, B_i \subset B_{i+1}$$

$$(3.3.2) \qquad \qquad \bigcup_{i \in \mathbb{N}} A_i = \mathfrak{C}$$

$$(3.3.3) \qquad \qquad \bigcup_{i\in\mathbb{N}} B_i = \mathfrak{D}',$$

(3.3.4)
$$qftp(a_n/\hat{a}) = qftp(b_n/\hat{b})$$

The union $\bigcup_{i\in\mathbb{N}} p_i$ will then be an order isomorphism $\mathfrak{C}' \to \mathfrak{D}'$.

3.4. Back & forth. Let p_n be a partial isomorphism mapping a_i to b_i for $i = 0, 1, \ldots, n$.

<u>Forth:</u> (n is even) Let c be an element of $\mathfrak{C} \setminus A$.

Let c_{\downarrow} be the set $\{x \in A | x \ll c\}$. Similarly, $c_{\uparrow} := \{x \in A | c \ll x\}$, and $c_{\parallel} := \{x \in A | c \ll x\}$.

 $A|x||c\}.$

The sets c_{\rightarrow} and c_{\leftarrow} are defined as

$$c_{\twoheadrightarrow}:=\{x\in A|c\twoheadrightarrow x\},\quad c_{\leftarrow}:=\{x\in A|x\twoheadrightarrow c\}.$$

We will assume that $c_{\rightarrow} = c_{\leftarrow} = \emptyset$. It follows that

$$\bigcap_{x \in c_{\downarrow}} I^{+}(x) \cap \bigcap_{x \in c_{\uparrow}} I^{-}(x) \cap \bigcap_{x \in c_{\parallel}} x^{\perp} \neq \emptyset$$

By the assumptions of the Lemma, we have

$$\bigcap_{x \in c_{\downarrow}} I^{+}(p_{n}(x)) \cap \bigcap_{x \in c_{\uparrow}} I^{-}(p_{n}(x)) \cap \bigcap_{x \in c_{\parallel}} p_{n}(x)^{\perp} \neq \emptyset$$

and it suffices to choose a d in the above intersection.

Denote $a_{n+1} := c$ and $b_{n+1} := d$.

<u>Back:</u> (*n* is odd) Let *d* be an element of $\mathcal{N} \setminus B$. Let d_{\downarrow} be the set $\{x \in b | x \ll d\}$. Similarly, $d_{\uparrow} := \{x \in B | d \ll x\}$, and $d_{\parallel} := \{x \in B | x \parallel d\}$.

The sets d_{\rightarrow} and d_{\leftarrow} are defined as

We will assume that $d_{\rightarrow} = d_{\leftarrow} = \emptyset$.

It follows that

$$\bigcap_{x \in d_{\downarrow}} I^{+}(x) \cap \bigcap_{x \in d_{\uparrow}} I^{-}(x) \cap \bigcap_{x \in d_{\parallel}} x^{\perp} \neq \emptyset$$

By the assumptions of the Lemma, we have

$$\bigcap_{x \in d_{\downarrow}} I^+(p_n^{-1}(x)) \cap \bigcap_{x \in d_{\uparrow}} I^-(p_n^{-1}(x)) \cap \bigcap_{x \in d_{\parallel}} p_n^{-1}(x)^{\perp} \neq \emptyset$$

and it suffices to choose a c in the above intersection.

Denote $a_{n+1} := c$ and $b_{n+1} := d$.

The case where c_{\rightarrow} (say) is non-empty is handled similarly by using condition (3.2.2). More precisely, assume that c satisfies

$$\left[\bigwedge_{i\in I} a_i \twoheadrightarrow c \& \bigwedge_{j\in J} c \twoheadrightarrow a_j\right]$$

for some $I, J \subset \{0, 1, ..., n\}$. Then by condition 3.2.2, there exists some $d \in \mathcal{N}$ satisfying

$$\left[\bigwedge_{i\in I} b_i \twoheadrightarrow d \& \bigwedge_{j\in J} d \twoheadrightarrow b_j\right].$$

Using Lemma 3.3, we can assume that $\operatorname{tp}(d/\overline{b}) = \operatorname{tp}(c/\overline{a})$. However, might be the case that $d \notin \mathfrak{D}$, in which case we let $\mathfrak{D}_n := \mathfrak{D}_{n-1} \cup \{d\}$, where \mathfrak{D}_{n-1} was defined at the earlier stages.

If, in the back stage, $d_{\rightarrow} \neq \emptyset$ (say), then a similar reasoning as above can be applied, and we define a set $\mathfrak{C}_n := \mathfrak{C}_{n-1} \cup \{c\}$, where c satisfies $\operatorname{tp}(c/\bar{a}) = \operatorname{tp}(d/\bar{b})$.

Define

$$\mathfrak{C}' := igcup_{n\in\mathbb{N}} \mathfrak{C}_n, \quad \mathfrak{D}' := igcup_{n\in\mathbb{N}} \mathfrak{D}_n$$

and

$$p := \bigcup_{k \in \mathbb{N}} p_k.$$

Then:

3.5. Lemma. The map p is a total causal isomorphism $\mathfrak{C}' \to \mathfrak{D}'$.

Proof. The map p is total:

let x be an element of \mathfrak{C}' . Then $x = a_\ell$ for some $\ell \in \mathbb{N}$. Hence, $x \in A_\ell$. It follows that $p|_{A_\ell} = p_\ell$ is defined on $x = a_\ell$.

To show that p is bijective, we argue similarly: let, for the sake of a contradiction, $x, y \in \mathfrak{C}'$ be elements such that p(x) = p(y). By the above reasoning, there exists some i such that $p_i(x) = p(x) = p(y) = p_i(y)$, contradicting the injectivity of p_i . Similarly, let $z \in \mathfrak{C}'$. Then $z = a_\ell$ for some sufficiently large ℓ ; then $z = a_\ell \in B_k$ (for some natural number k). We have $p_k(A_k) = B_k$.

Let us define

$$\Phi: \mathcal{M} \to \mathcal{N}, x \mapsto \lim_{k} p(x_k), \text{where } x_k \to x \text{ as } k \to \infty, x_k \in \mathbf{dom}(p).$$

This is well defined as otherwise p ceases to be a causal morphism:

3.6. Lemma. The map Φ is a well-defined non-trivial causal isomorphism $\mathcal{M} \to \mathcal{N}$.

Proof of Lemma:

First we show that Φ is a well defined map: let $y, z \in \mathcal{N}$ be two spacetime points, such that $\lim_k p(x_k) = y$ and $\lim_\ell p(x'_\ell) = z$, where $(x_k)_k, (x'_\ell)_\ell$ both converge to $x \in \mathcal{M}$.

Let \mathcal{V} be a neighborhood $\mathcal{V} = [a, b]$ of x, with $a, b \in \mathfrak{C}$; then

$$x_k, x'_\ell \in \mathcal{V} \ \forall k, \ell > N_1, \& p(\mathcal{V} \cap \mathfrak{C}') \subset [p(a), p(b)]$$

for some $N_1 \in \mathbb{N}$. In particular $p(x_k), p(x'_{\ell}) \in [p(a), p(b)]$ for all $k, \ell > N_1$.

Considering a sequence $\mathcal{V}_n := [a_n, b_n], a_n, b_n \in \mathfrak{C}$ of nested causal diamonds $\mathcal{V}_{n+1} \subset \mathcal{V}_n$, such that $a_n, b_n \to x$ as $n \to \infty$ we obtain $\lim_{k\to\infty} p(x_k) = \lim_{\ell\to\infty} p(x'_\ell)$, thus y = z and Φ is well defined.

Similarly, Φ is a causal morphism.

Finally, to see that Φ is bijective, we observe that p is bijective, and that each point $z \in \mathcal{N}$ is the limit of some sequence $(z_k)_k$: the sequence $(p^{-1}(z_k))_k$ being contained in a compact subset of \mathcal{M} it converges to some limit $\Phi^{-1}(z)$.

3.7. *Remark.* While back & forth can be used (in general) to show that two countable structures are isomorphic, the above result extends the isomorphism $\mathfrak{C}' \to \mathfrak{D}'$ to $\Phi : \mathcal{M} \to \mathcal{N}$. In reality, the two countable structures $\mathfrak{C}', \mathfrak{D}'$ that were shown

isomorphic by p are both dense in \mathcal{M}, \mathcal{N} respectively, and that furthermore, \mathcal{M}, \mathcal{N} are *standard* pseudo-riemannian manifolds (hence \mathcal{M} is rigid over \mathfrak{C}' , and similarly for \mathcal{N}).

4. Partial classification of $Aut(\mathcal{M})$ for \mathcal{M} a two-dimensional globally hyperbolic spacetime

This section considers the case of a two-dimensional globally hyperbolic spacetime.

We have:

4.1. **Theorem.** Let \mathcal{M} be a two-dimensional globally hyperbolic spacetime having non-compact Cauchy surfaces.

Assume furthermore that \mathcal{M} is upward and downward directed, i.e. \mathcal{M} satisfies the following

(†) For all $x, y \in \mathcal{M}$, there exist $u, v \in \mathcal{M}$ such that $x, y \prec u \& v \prec x, y$.

Then a causal isomorphism exists $i : \mathcal{M} \to \mathbb{M}^2$. Consequently, the group of causal automorphisms of \mathcal{M} is $Aut(\mathbb{M}^2)$.

Let \mathcal{C} be a countable dense subset of \mathcal{M} .

4.2. Lemma. Let $A, B \subset C, A \cap B = \emptyset$ be finite. Assume that $A \ll B$; then there exists some $z \in C$ such that $A \ll \{z\} \ll B$.

Proof. This is done by induction on |A| and |B|.

- a) If $A = B = \emptyset$, then the statement holds since \mathcal{M} is non-empty.
- b) If |A| = 0 then by induction on |B|:
 - i. |B| = 1: the statement follows since \mathcal{M} is globally hyperbolic with no boundary;
 - ii. |B| = 2: the statement follows by the assumption that \mathcal{M} is directed.
 - iii. |B| > 2: the statement follows easily by induction.
- c) The case |B| = 0 is treated similarly.
- d) |A| = |B| = 1: follows by the denseness of \mathcal{C} .
- e) |A| = 1, |B| = 2: write $A = \{a\}, B = \{b_1, b_2\}$ with $a \ll b_1, b_2$. Then b_1, b_2 are contained in the forward light cone with vertex a. Similarly, a is contained in the past light cones of b_1, b_2 .

The past light cones of vertices b_1, b_2 have nonempty intersection $I_{12} := I^-(b_1) \cap I^-(b_2)$ (since *a* belongs to this intersection).

Let $z \in I_{12} \cap I^+(a)$. Then, z satisfies the required property.

- f) A similar argument applies when |A| = 2, |B| = 1.
- g) |A| = 2, |B| = 2: This is the base step for induction; Let $A = \{a_1, a_2\}, B := \{b_1, b_2\}$. Then $a_1, a_2 \ll b_1, b_2$. As $a_1, a_2 \ll b_1$, two null curves emanating from a_1, a_2 intersect at a point p. Then $p \ll b_1, b_2$.

Similarly, two null curves containing b_1, b_2 respectively intersect at some point q. We have $p \ll q$; for some $z \in [p,q]$, $p \ll z \ll q$. Hence, z satisfies the requirements of the Lemma.





Assume the induction hypothesis holds for all sets A, B satisfying |A| = m, |B| = n.

For |A| = m + 1, |B| = n: let $A' \subset A$, |A'| = m, and let $\{a\} := A \setminus A'$. Then, by the induction hypothesis, there exists some z such that $A' \ll \{z\} \ll B$.

If $a \ll z$, we are done. Otherwise, to conclude, apply the result for the sets $\{a, z\}, B$. For |A| = m, |B| = n + 1 a similar reasoning applies.

It follows immediately:

4.3. Lemma. Keep the above notation. Let $\{a_0, a_1, \ldots, a_n\} \subset \mathcal{M}$, and $\{b_0, b_1, \ldots, b_n\} \subset \mathbb{M}^2$ with $qftp(\bar{a}) = qftp(\bar{b})$. Then

$$(4.3.1) \qquad \mathcal{O}_1 \cap \mathcal{O}_2 \cap \dots \cap \mathcal{O}_n \neq \emptyset \leftrightarrow \mathcal{O}'_1 \cap \mathcal{O}'_2 \cap \dots \cap \mathcal{O}'_n \neq \emptyset$$

4.4. **Proof of Theorem.** The result follows from the above Lemmas, together with the observation that the condition $qftp(\bar{a}) = qftp(\bar{b})$ holds for the tuples $\bar{a} = a_0, \bar{b} = b_0$ with arbitrary $a_0 \in \mathcal{M}$ and $b_0 \in \mathbb{M}^2$.

4.5. Two-dimensional spacetimes with compact Cauchy surfaces. To handle the case where a spacetime \mathcal{M} has compact Cauchy surfaces, we use the following result:

4.6. Theorem ([6]). Let M be a Lorentzian manifold (or a semi-Riemannian manifold) with universal covering $\pi : \overline{M} \to M$. Denote by Γ the group $\pi_1(\mathcal{M})$. Then, $\Gamma \subset Aut(\mathcal{M}) =: G$ (respectively $\Gamma \subset Conf(\mathcal{M}) =: G_1$). Also, Aut(M) (or, Conf(M)) is isomorphic to $\mathcal{N}(\Gamma)/\Gamma$ in which $\mathcal{N}(\Gamma)$ is the normalizer of Γ in $Aut(\overline{M})$

We obtain the following:

(or, Conf(M)).

4.7. Theorem. Let \mathcal{M} be a two-dimensional globally hyperbolic spacetime having compact Cauchy surfaces.

Assume furthermore that \mathcal{M} (the universal covering space of \mathcal{M}) is directed. Then the group $Aut(\mathcal{M})$ of causal automorphisms of \mathcal{M} is given by

 $\mathcal{N}(\mathbb{Z})/\mathbb{Z}$

where $\mathcal{N}(\mathbb{Z})$ is the normalizer of \mathbb{Z} in $Aut(\mathbb{M}^2)$.

4.8. Non-directed two-dimensional globally hyperbolic spacetime. Let us define

$$\mathcal{D} := \{ (x, t) \in \mathbb{M}^2 | |x| + |t| < 1 \}.$$

This is the diamond $I^+((0,-1)) \cap I^-((0,+1))$. Note that \mathcal{D} is causally isomorphic to \mathbb{M}^2 .

4.9. **Theorem** ([5]). Any two-dimensional spacetime with non-compact Cauchy surfaces can be causally isomorphically embedded into \mathcal{D} .

We denote the above embedding by ι . Let us recall the statement for $\mathcal{M} = \mathbb{M}^2$:

4.10. **Theorem** ([3]). The group $Aut(\mathbb{M}^2)$ of causal automorphisms of the Minkowski plane is given by

$$(Homeo_{\leq}(\mathbb{R}))^2 \rtimes S_2$$

We may characterize causal automorphisms of non-directed globally hyperbolic spacetimes by considering the embedding ι .

4.11. **Theorem.** Let \mathcal{M} be a two-dimensional globally hyperbolic spacetime. If \mathcal{M} is not directed, then any causal automorphism of \mathcal{M} induces a homeomorphism $\partial_{\iota}\mathcal{M} \rightarrow \partial_{\iota}\mathcal{M}$, where $\partial_{\iota}\mathcal{M}$ is equipped with the standard topology.

By analyzing the structure of non-directed spacetimes, we obtain a more precise classification of the automorphism group:



FIGURE 3.

4.12. **Theorem.** Let \mathcal{M} be a two-dimensional globally hyperbolic spacetime. Then $Aut(\mathcal{M})$ is one of the following groups:

- 1. $\Gamma := (Homeo_{\leq}(\mathbb{R}))^2 \rtimes S_2$;
- 2. $\mathcal{N}_{\Gamma}(\mathbb{Z})/\mathbb{Z}$;
- 3. $\Gamma' := a$ subgroup of Γ stabilizing a countable set;
- 4. $\mathcal{N}_{\Gamma'}(\mathbb{Z})/\mathbb{Z}$;

5.
$$Homeo_{\leq}(\mathbb{R}) \rtimes S_2;$$

6.
$$\mathcal{N}_{Homeo_{\leq}(\mathbb{R})\rtimes S_2}(\mathbb{Z})/\mathbb{Z}$$

Proof. Items (1) and (2) have already been investigated (they correspond to directed spacetimes with non-compact and compact Cauchy surfaces, respectively). Assume that \mathcal{M} has non-compact Cauchy surfaces.

Assume furthermore that \mathcal{M} is upward but not downward-directed, and let $a_1, a_2 \in \partial \mathcal{M}$ be such that

$$(\ddagger) \qquad \exists b(a_1, a_2 \prec b \& \nexists x (x \in \mathcal{M} \& x \prec a_1, a_2)).$$

Then any automorphism of \mathcal{D} must preserve the type $tp(a_1, a_2)$. We have three possible cases:

- 1. There exist timelike curves γ_1, γ_2 containing a_1, a_2 respectively such that for any $b_1 \in \gamma_1, b_2 \in \gamma_2$ satisfying $a_1 \ll b_1 \ll b, a_2 \ll b_2 \ll b$ we have (‡) holds for b_1, b_2 .
- 2. There exist null curves γ_1, γ_2 containing a_1, a_2 respectively such that for any $b_1 \in \gamma_1, b_2 \in \gamma_2$ satisfying $a_1 \twoheadrightarrow b_1 \twoheadrightarrow b, a_2 \twoheadrightarrow b_2 \twoheadrightarrow b$ we have (\ddagger) holds for b_1, b_2 .
- 3. For all causal curves γ_1, γ_2 with $a_1 \in \gamma_1, a_2 \in \gamma_2$, there exist some $b_1 \in \gamma_1, b_2 \in \gamma_2$ satisfying $a_1 \prec b_1 \prec b, a_2 \prec b_2 \prec b$ and (‡) does not hold for b_1, b_2 .

Cases 1. and 2. imply the existence of a small interval in $\partial \mathcal{M}$ (i.e., the image of an interval of \mathbb{R} by a homeomorphism $\mathbb{R} \to \partial \mathcal{M}$). By paracompactness of $\partial \mathcal{M}$, we conclude the existence of a countable set of b (as in (‡) above), and hence Aut(\mathcal{M}) stabilizes a countable subset of \mathcal{M} .

In case 3., the group of causal automorphisms stabilizing $\partial \mathcal{M}$ is seen to be $\operatorname{Homeo}_{\leq}(\mathbb{R}) \rtimes S_2$: let $\partial \mathcal{M}$ be defined by $x^- = \varphi(x^+)$, and let $\Phi : \mathcal{M} \to \mathcal{M}, (x^+, x^-) \mapsto (\phi(x^+), \psi(x^-))$ be a causal automorphism of \mathcal{M} , where ϕ, ψ are increasing homeomorphisms of \mathbb{R} . Denoting the extension of Φ to \mathcal{D} by Φ' , let $\Phi' : \mathcal{D} \to \mathcal{D}, (x^+, x^-) \mapsto (\phi(x^+), \psi(x^-))$ (with ϕ, ψ defined over \mathbb{R}). Then, on $\partial \iota \mathcal{M}$: $\psi(\varphi(x^+)) = \varphi(\phi(x^+))$; let u be a parameter for $\partial \mathcal{M}, x^+ = f_+(u), x^- = f_-(u)$, then $\phi(f_+(u)) = f_+(\varphi(u)), \psi(f_-(u)) = f_-(\varphi(u))$. We may assume that f_+ or f_- is monotone (otherwise, it can be seen that for some $x \in \partial \mathcal{M}$, we have $I^{\pm}(x) \cap \iota \mathcal{M} \neq \emptyset$, impossible by Theorem 4.1 of [6]). It follows that φ is monotone, as required.

The case where \mathcal{M} is downward but not upward directed is treated similarly. If \mathcal{M} is neither upward nor downward directed, then the same analysis still applies (with minor changes), and we also get that $\operatorname{Aut}(\mathcal{M})$ is one of the groups listed in the Theorem.

The other cases (i.e., \mathcal{M} has compact Cauchy surfaces) follow by the application of Theorem 4.7.

4.13. Group of conformal automorphisms of \mathcal{M} . Let us denote by Diffeo $(\mathbb{R})_{\leq}$ the group of increasing C^{∞} -diffeomorphisms of the real line.

In [5], a characterization of the conformal automorphism group of a two-dimensional globally hyperbolic spacetime with a non-compact Cauchy surface is given. We obtain a simpler characterization of C^{∞} -conformal diffeomorphisms of the Minkowski plane using null coordinates $x^+ = t + x, x^- = t - x$ (the standard metric on \mathbb{M}^2 written in terms of null coordinates is given by $-dx^+dx^-$):

4.14. **Theorem.** The group $Conf(\mathbb{M}^2)$ of conformal diffeomorphisms of the plane is given by

$$Diffeo_{\leq}(\mathbb{R})^2 \rtimes D_2,$$

where D_2 is the dihedral group, $D_2 \simeq C_2 \times C_2$.

Proof. Let $\Phi : \mathbb{M}^2 \to \mathbb{M}^2, (x^+, x^-) \mapsto (X^+, X^-)$ be a conformal diffeomorphism of the plane.

We have:

$$-dX^{+}dX^{-} = -e^{\varphi(x^{+},x^{-})}dx^{+}dx^{-},$$

as Φ is a conformal map, with $\varphi(x^+, x^-)$ being an arbitrary function. Writing $X^+ = X^+(x^+, x^-)$ and $X^- = X^-(x^+, x^-)$ we obtain

$$-\left(\frac{\partial X^{+}}{\partial x^{+}}dx^{+} + \frac{\partial X^{+}}{\partial x^{-}}dx^{-}\right)\left(\frac{\partial X^{-}}{\partial x^{+}}dx^{+} + \frac{\partial X^{-}}{\partial x^{-}}dx^{-}\right) = -e^{\varphi(x^{+},x^{-})}dx^{+}dx^{-},$$

hence

$$X_{+}^{+}X_{+}^{-} = 0$$

$$X_{-}^{+}X_{-}^{-} = 0$$

$$X_{+}^{+}X_{-}^{-} + X_{-}^{+}X_{+}^{-} > 0$$

Finally,

$$\begin{split} X^+ &= X^+(x^+), X^- = X^-(x^-), \quad \text{and} \quad X^+_+ X^-_- > 0 \\ & \text{Or} \\ X^+ &= X^+(x^-), X^- = X^-(x^+), \quad \text{and} \quad X^+_- X^-_+ > 0. \end{split}$$

The rest of the proof now follows [3]: any conformal diffeomorphism of \mathbb{M}^2 is of one the forms $(x^+, x^-) \mapsto (f(x^+), g(x^-))$ or $(x^+, x^-) \mapsto (f(x^-), g(x^+))$, where f, g are diffeomorphisms of the real line, which are both increasing or decreasing. It follows that the group $\operatorname{Conf}(\mathbb{M}^2)$ is generated by elements in $\operatorname{Diffeo}_{\leq}(\mathbb{R})^2$, space reflections $(x^+, x^-) \mapsto (x^-, x^+)$ and time reflections $(x^+, x^-) \mapsto (-x^-, -x^+)$.

We still need to see that the subgroup $\text{Diffeo}_{\leq}(\mathbb{R})^2$ is normal in $\text{Conf}(\mathbb{M}^2)$. Let p_1 denote the transposition $(x^+, x^-) \mapsto (x^-, x^+)$; for any $n \in \text{Diffeo}_{\leq}(\mathbb{R})^2$, $n : (x^+, x^-) \mapsto (f(x^+), g(x^-))$ we have $p_1 n p_1 : (x^+, x^-) \mapsto (g(x^+), f(x^-)$ where g, f are increasing diffeomorphisms of \mathbb{R} . Hence $p_1 n p_1 \in \text{Diffeo}_{\leq}(\mathbb{R})^2$ as required. Let p_2 denote the map $(x^+, x^-) \mapsto (-x^-, -x^+)$; then we have

$$p_2np_2: (x^+, x^-) \mapsto (-x^-, -x^+) \mapsto (f(-x^-), g(-x^+)) \mapsto (-g(-x^+), -f(-x^-)).$$

Defining f_1, g_1 by $f_1(x^+) = -g(-x^+), g_1 = -f(-x^-)$ it can be seen that both f_1 and g_1 are increasing diffeomorphisms of \mathbb{R} . Let now $p := p_1 p_2 : (x^+, x^-) \mapsto (-x^+, -x^-)$; a similar reasoning shows that pnp is in $\text{Diffeo}_{\leq}(\mathbb{R})^2$, whence the result follows. This proves the Theorem. \Box

Applying the above considerations allows us to obtain the following:

4.15. **Theorem.** Let \mathcal{M} be a two-dimensional globally hyperbolic spacetime. Then $Conf(\mathcal{M})$ is one of the following groups:

1. $\Gamma_1 := (Diffeo_{\leq}(\mathbb{R}))^2 \rtimes D_2$; 2. $\mathcal{N}_{\Gamma_1}(\mathbb{Z})/\mathbb{Z}$; 3. $\Gamma'_1 := a \ subgroup \ of \ \Gamma_1 \ stabilizing \ a \ countable \ set;$ 4. $\mathcal{N}_{\Gamma'_1}(\mathbb{Z})/\mathbb{Z}$; 5. $\Gamma_2 := Diffeo_{\leq}(\mathbb{R}) \rtimes D_2;$ 6. $\mathcal{N}_{\Gamma_2}(\mathbb{Z})/\mathbb{Z};$ 7. $\Gamma'_2 := a \ subgroup \ of \ \Gamma_2 \ stabilizing \ a \ countable \ set;$ 8. $\mathcal{N}_{\Gamma'_2}(\mathbb{Z})/\mathbb{Z}.$

Proof. The same proof method for Theorem 4.11 still applies after making the necessary changes at the appropriate places (extra care should be taken for conformal anti-causal diffeomorphisms). \Box

5. Locally conformally flat globally hyperbolic spacetimes

While a generic spacetime of dimension ≥ 3 has very few causal (or conformal) automorphisms, the locally conformally flat case is more interesting.

5.1. **Theorem.** Let \mathcal{M} be globally hyperbolic spacetime. Assume that \mathcal{M} is

- (1) simply connected ;
- (2) locally conformally flat;
- (3) directed;
- (4) has non-compact Cauchy surfaces;
- (5) $\dim(\mathcal{M}) \geq 3$;

Then \mathcal{M} is globally conformally flat.

Proof. It suffices to check that the conditions (3.2.1, 3.2.2) are met. Let $\mathcal{D} = [p, q]$ be an interval in \mathcal{M} .

By compactness of \mathcal{D} , a finite cover exists \mathcal{D}_i , $i = 1, \ldots, \ell$ by conformally flat opens. The metric restricted to \mathcal{D}_i has the form

$$g_i := g|_{\mathcal{D}_i} = \Omega_i (-dt^2 + \sum_{i=1}^{n-1} dx_i^2)$$

Then

$$g|_{\mathcal{D}} = \Omega g_{\mathbb{M}^n}$$

where Ω is obtained from the Ω_i using partition of unity. For any finite set $A \subset \mathcal{D}$, we can then use the causal embedding $\mathcal{D} \hookrightarrow \mathbb{M}^n$ to deduce the required result.

For a given spacetime \mathcal{M} , we denote by $\overline{\mathcal{M}}$ its universal covering space.

5.2. **Theorem.** Let \mathcal{M} be a globally hyperbolic spacetime. Assume that $\overline{\mathcal{M}}$ is

(1) locally conformally flat; (2) directed; (3) dim(\mathcal{M}) ≥ 3 ; Let $\Gamma := \pi_1(\mathcal{M})$. Then Conf(\mathcal{M}) = $\mathcal{N}_{Conf(\mathbb{M}^n)}(\Gamma)/\Gamma$.

Combining this with the result by A. V. Levichev ([9]), we obtain:

5.3. Theorem. Let \mathfrak{M} be a globally hyperbolic spacetime having dimension ≥ 3 . Assume that \mathfrak{M} is locally conformally flat and that (\mathfrak{M}, \ll) is directed. Then \mathfrak{M} is conformally homogeneous.

References

- Peter J. Cameron, Deborah C. Lockett, Posets, homomorphisms and homogeneity, Discrete Mathematics 310 (2010) 604-613.
- P.J. Cameron, J. Nešetřil, Homomorphism-homogeneous relational structures, Combinatorics, Probability and Computing 15 (2006) 91-103.
- A García-Parrado Gómez-Lobo and E Minguzzi, Product posets and causal automorphisms of the plane, Class. Quant. Gravity, vol. 28. no. 14 (2011).
- 4. Hawking, S. R., King, A. R. and McCarthy, P. J., J. Math. Phys. 17, 174 (1976).
- 5. Kim, D.-H., J. Geom. Phys. 96 (2015) 146-154.
- Kim, D.-H., A classification of two-dimensional spacetimes with non-compact Cauchy surfaces, J. Geom. Phys. 73 (2013) 252–259.

- Kim, D.-H., A note on non-compact Cauchy surfaces, Class. Quantum. Grav. 25, (2008) pp.238002.
- E. H. Kronheimer and R. Penrose (1967). On the structure of causal spaces, Math. Proc. Cambridge Philosophical Society, 63, pp 481501 doi:10.1017/S030500410004144X
- 9. A.V. Levichev, Prescribing the conformal geometry of a Lorentz manifold by means of its causal structure, Sov. Math. Dokl. 35, (1987) pp.452.
- 10. Melnick, K. & Pecastaing, V. The conformal group of a compact simply connected Lorentzian manifold, arXiv:1911.06251v2
- 11. Zeeman, E. C., J. Math. Phys. 5, 490 (1964); doi: 10.1063/1.1704140

FACULTY OF SCIENCES (I), LEBANESE UNIVERSITY, BEIRUT, LEBANON