CONCENTRATION AND MOMENT INEQUALITIES FOR HEAVY-TAILED RANDOM MATRICES

MORITZ JIRAK, STANISLAV MINSKER, YIQIU SHEN, AND MARTIN WAHL

ABSTRACT. We prove Fuk-Nagaev and Rosenthal-type inequalities for the sums of independent random matrices, focusing on the situation when the norms of the matrices possess finite moments of only low orders. Our bounds depend on the "intrinsic" dimensional characteristics such as the effective rank, as opposed to the dimension of the ambient space. We illustrate the advantages of such results in several applications, including new moment inequalities for sample covariance matrices and the corresponding eigenvectors of heavy-tailed random vectors. Moreover, we demonstrate that our techniques yield sharpened versions of the moment inequalities for empirical processes.

1. INTRODUCTION

Fuk-Nagaev inequalities (Fuk and Nagaev, 1971; Nagaev, 1979) generalize exponential deviation inequalities for the sums of independent random variables, such as Berstein's, Prokhorov's and Bennett's inequalities, to case when the random variables satisfy minimal integrability conditions. For example, a corollary of Fuk and Nagaev's results is the following bound: for a sequence of independent, centered random variables X_1, \ldots, X_n such that $\max_k \mathbb{E}|X_k|^p < \infty$ for some $p \ge 2$,

$$\mathbb{P}\left(\left|\sum_{k=1}^{n} X_{k}\right| \geq t\right) \leq 2\exp\left(-C_{1}(p)\frac{t^{2}}{\sum_{k=1}^{n} \mathbb{E}X_{k}^{2}}\right) + \mathbb{P}\left(\max_{k}|X_{k}| > t/4\right) + C_{2}(p)\left(\frac{\sum_{k=1}^{n} \mathbb{E}|X_{k}|^{p}}{t^{p}}\right)^{2}$$

Nagaev (1979) describes the applications of such results to the laws of large numbers and moment inequalities. Later, Einmahl and Li (2008), Rio (2017) and Bakhshizadeh et al.

²⁰¹⁰ Mathematics Subject Classification. 60B20, 60E15, 62H25, 15A42.

Key words and phrases. Fuk-Nagaev inequality, Rosenthal's inequality, Covariance estimation, Principal Components Analysis.

S. Minsker and Y. Shen acknowledge support by the National Science Foundation grant DMS CAREER-2045068.

(2023), among others, improved the original estimates by Fuk and Nagaev in several ways: first, the inequalities were extended to martingales and Banach-space valued random variables, and second, the constants were sharpened. For example, inequalities due to Rio (2017); Bakhshizadeh et al. (2023) hold with $C_1(p) = 1/2 + \delta$ for any $\delta > 0$ and $C_2(p)$ of order p^p . The latter fact is important as the order of growth of these constants translates into the tail behavior of $|\sum_{k=1}^{n} X_k|$.

The goal of this work is to prove a version of the Fuk-Nagaev inequality for the sums of independent random matrices and use it to sharpen existing moment inequalities. Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of independent self-adjoint¹ random matrices such that $\mathbb{E}W_k = 0_{d \times d}$ for all k and where the expectation is taken element-wise. Assume that for all k, $||W_k|| \leq U$ with probability 1 where $|| \cdot ||$ stands for the operator (spectral) norm. A line of work by Ahlswede and Winter (2002); Oliveira (2010); Tropp (2012) culminated in the following version of the so-called matrix Bernstein's inequality: for all t > 0,

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| \ge t\right) \le 2d \exp\left(-\frac{t^{2}/2}{\sigma^{2} + Ut/3}\right)$$

where $\sigma^2 = \|\sum_{k=1}^n \mathbb{E}W_k^2\|$. An attractive feature of this inequality (as opposed to, say, Talagrand's concentration inequality (Talagrand, 1996)) is that it yields a bound for $\mathbb{E}\|\sum_{k=1}^n W_k\|$, namely, that $\mathbb{E}\|\sum_{k=1}^n W_k\| \leq K(\sigma\sqrt{\log(d)} + U\log(d))$ for some absolute constant K > 0. Tropp's results have been extended in two directions: first, it was shown by Minsker (2017); Tropp (2015) that the dimension factor d can essentially be replaced by the so-called *effective rank* $r\left(\sum_{k=1}^n \mathbb{E}W_k^2\right)$ where

$$r(A) := \frac{\operatorname{trace}(A)}{\|A\|}.$$

for a positive definite matrix A. In particular, this version of Bernstein's inequality is applicable in the context of random Hilbert-Schmidt operators acting on Hilbert spaces. Second, the boundedness assumption was relaxed by Koltchinskii (2011) to the requirement that $\max_k || || W_k || ||_{\psi_1} < \infty$ where the ψ_1 norm of a random variable Z is defined via $|| Z ||_{\psi_1} = \inf\{r > 0 : \mathbb{E}e^{|Z/r|} \leq 2\}$. Finally, Klochkov and Zhivotovskiy (2020) showed that

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_k\right\| \ge t\right) \le c_1 r(V_n^2) \exp\left(-c_2 \frac{t^2}{\|V_n^2\| + Rt}\right)$$

¹It is well-known that the case of general rectangular matrices reduces to this one via the so-called "Hermitian dilation," see (Tropp, 2015, section 2.1.17).

for all $t \ge c_3(\|V_n^2\|^{1/2} + M)$ where V_n^2 is any matrix satisfying $V_n^2 \succeq \sum_{k=1}^n \mathbb{E}W_k^2$ and $R = \|\max_k \|W_k\|\|_{\psi_1}$. Our results allow to relax the integrability assumptions even further and cover the case of heavy-tailed random matrices, namely, random matrices such that $\mathbb{E}\|W\|^p = \infty$ for some p > 0 (however, we are still able to recover the known bounds for the "light-tailed" random matrices). For example, Theorem 2.2 below implies that for all $p \ge 1$,

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\|V_{n}^{2}\|^{1/2}\sqrt{q} + q\mathbb{E}\max_{k}\|W_{k}\| + \frac{p}{\log(ep)}(\mathbb{E}\max_{k}\|W_{k}\|^{p})^{1/p}\right)$$

where $q = \log(r(V_n^2)) \lor p$ and K is an absolute constant. This inequality sharpens previously known results of this type by Junge and Zeng (2013); Dirksen (2011); Chen et al. (2012): for example, a version of Rosenthal's inequality by Chen et al. (2012) states that

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\|V_{n}^{2}\|^{1/2}\sqrt{r} + r\mathbb{E}^{1/p}\max_{k}\|W_{k}\|^{p}\right)$$

where $r = \log(d) \lor p$. The fact that our bound depends on $r(V_n^2)$ instead of d immediately allows one to extend it to Hilbert-Schmidt operators acting on Hilbert spaces.

Finally, let us remark that the order of the constants in the inequality stated above is optimal: in the scalar case d = 1, it is known (Johnson et al., 1985) that without any additional assumptions, the best order of C(p) in the inequality

$$\mathbb{E}^{1/p} \Big| \sum_{k=1}^{n} W_k \Big|^p \le C(p) \left(\left(\sum_{k=1}^{n} \mathbb{E} W_k^2 \right)^{1/2} + \left(\sum_{k=1}^{n} \mathbb{E} |W_k|^p \right)^{1/p} \right)$$

is $C(p) = K \frac{p}{\log(p)}$ while Pinelis and Utev (1985) showed that $C_1(p) = K \sqrt{p}$ and $C_2(p) = Kp$ are the best possible in the inequality of the form

(1)
$$\mathbb{E}^{1/p} \Big| \sum_{k=1}^{n} W_k \Big|^p \le C_1(p) \left(\sum_{k=1}^{n} \mathbb{E} W_k^2 \right)^{1/2} + C_2(p) \mathbb{E}^{1/p} \max_k |W_k|^p.$$

It is clear that our results yield a sharper version the inequality (1) for large p whenever $\mathbb{E} \max_k |W_k|$ is much smaller than $\mathbb{E}^{1/p} \max_k |W_k|^p$.

4 MORITZ JIRAK, STANISLAV MINSKER, YIQIU SHEN, AND MARTIN WAHL

1.1. Organization of the paper. The rest of the exposition is organized as follows: we present the main results and their proofs in sections 2.1 and 2.2. Applications of the developed techniques to the empirical processes is discussed in section 2.3 while implications for the problems of matrix subsampling, covariance estimation and eigenvector estimation are described in sections 3.1, 3.2 and 3.3 respectively. Finally, section 4 contains the required background and proofs of the lemmas that were omitted from the main exposition.

2. Main results

In this section we state the new concentration and moment bounds - Theorems 2.1 and 2.2. The required notation will be introduced on demand. Let us remark that throughout the paper, the values of constants K, c, $C(\cdot)$ is often unspecified and can change from line to line; we use K and c to denote absolute constants and $C(\cdot)$ to denote constants whose value depends on the parameters in brackets.

2.1. Fuk-Nagaev-type inequality. The following proposition is the key technical ingredient that will serve as the starting point for the derivation of the main results. Therefore, we state it separately. Everywhere below, $M = \max_{k=1,...,n} ||W_k||$, $Q_{1/2}(Z)$ stands for the median of a real-valued random variable Z, and $\varepsilon_1, \ldots, \varepsilon_n$ denote independent symmetric Bernoulli random variables that are independent from W_1, \ldots, W_n . Finally, $\|\cdot\|_2$ is the Euclidean norm.

Proposition 2.1. Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of centered, independent, self-adjoint random matrices. Let U > 0, and assume that V_n^2 satisfies $V_n^2 \succeq \sum_k \mathbb{E} W_k^2 I\{\|W_k\| \leq U\}$. Finally, set $\sigma_U^2 = \|V_n^2\|$. Then, whenever

(2)
$$\frac{t}{2} \ge \sigma_U \lor U/3 \lor \sup_{\|v\|_2=1} Q_{1/2} \left(\left\langle \left(\sum_{k=1}^n W_k \right) v, v \right\rangle \right),$$

the following inequality holds:

(3)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > 12t\right) \leq 64r\left(V_{n}^{2}\right)\exp\left[-\frac{(t/2)^{2}}{\sigma_{U}^{2} + tU/6}\right] + 16\mathbb{P}\left(\left\|\sum_{k=1}^{n} \varepsilon_{k}W_{k}\mathbf{1}\left\{\|W_{k}\| > U\right\}\right\| > t/2\right)\mathbb{P}\left(\left\|\sum_{k=1}^{n} \varepsilon_{k}W_{k}\right\| > t\right) + 4\mathbb{P}\left(M > t\right).$$

If the random matrices are symmetrically distributed (that is, W_j and $-W_j$ are equidistributed for all j), then

(4)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > 3t\right) \leq 16r\left(V_{n}^{2}\right) \exp\left[-\frac{(t/2)^{2}}{\sigma^{2} + tU/6}\right] + 4\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\mathbf{1}\left\{\|W_{k}\| > U\right\}\right\| > t/2\right) \mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > t\right) + \mathbb{P}\left(M > t\right)$$

under the assumption that $\frac{t}{2} \geq \sigma_U \vee U/3$.

Remark 2.1. If W_1, \ldots, W_n have symmetric distribution, then $\mathbb{P}(\|\sum_{k=1}^n W_k\| > t) \geq \frac{1}{2}\mathbb{P}(M > t)$ in view of Lévy's inequality (Ledoux and Talagrand, 2013, Proposition 2.3). This shows that the quantile U is necessary in the lower bound for t. The term σ_U is also known to be necessary – for instance, take $W_j = \xi_j A_j$ where ξ_1, \ldots, ξ_n are *i.i.d.* N(0,1) random variables and A_1, \ldots, A_n are fixed self-adjoint matrices (Tropp, 2012).

Proof. Let us reduce the general case to the situation when W_1, \ldots, W_n are symmetric. To this end, it suffices to apply Lemma 2.3.7 in the book by van der Vaart and Wellner (1996): it implies that whenever $6t \ge \sup_{\|v\|_2=1} Q_{1/2} \left(\left(\sum_{k=1}^n W_k \right) v, v \right) \right)$,

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > 12t\right) \leq 4\mathbb{P}\left(\left\|\sum_{k=1}^{n} \varepsilon_{k} W_{k}\right\| > 3t\right).$$

Obviously, this inequality holds without any assumptions if W_1, \ldots, W_k are symmetrically distributed. Next, in view of Hoffmann-Jørgensen inequality (Proposition 4.1),

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > 3t\right) \le 4\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > t\right)^{2} + \mathbb{P}\left(M > t\right).$$

Given U > 0, we define, for each $k = 1, \ldots, n$,

$$\overline{W}_k := \varepsilon_k W_k \mathbf{1}\{ \|W_k\| \le U \} \text{ and } \Delta_k := \varepsilon_k W_k \mathbf{1}\{ \|W_k\| > U \}.$$

Clearly, $\|\sum_{k=1}^{n} \varepsilon_k W_k\| \leq \|\sum_{k=1}^{n} \widetilde{W}_k\| + \|\sum_{k=1}^{n} \Delta_k\|$, all the random matrices \widetilde{W}_k , Δ_k are symmetric, and

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > t\right) \le \mathbb{P}\left(\left\|\sum_{k=1}^{n}\widetilde{W_{k}}\right\| > t/2\right) + \mathbb{P}\left(\left\|\sum_{k=1}^{n}\Delta_{k}\right\| > t/2\right) := A_{1} + A_{2}.$$

Therefore,

(5)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > 3t\right) \le 4A_{1} + 4A_{2}\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > t\right) + \mathbb{P}\left(M > t\right).$$

The first term on the right-hand side of inequality (5) can be bounded directly via Theorem 4.1. Let V_n^2 satisfy $V_n^2 \succeq \sum_k \mathbb{E} W_k^2 I\{\|W_k\| \le U\}$. Then for $\sigma_U^2 = \|V_n^2\|$ and t such that $\frac{t}{2} \ge \sigma_U + U/3$,

(6)
$$4A_1 \le 16r(V_n^2) \exp\left[-\frac{(t/2)^2}{\sigma_U^2 + tU/6}\right].$$

The result follows.

We are now ready to deduce the first main result of the paper.

Theorem 2.1. Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of centered, independent, selfadjoint random matrices, and assume that $\mathbb{E}M^p < \infty$ for some p > 1. Moreover, suppose that V_n^2 satisfies $V_n^2 \succeq \sum_k \mathbb{E}W_k^2$, and set $\sigma^2 = ||V_n^2||$. Then

(7)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > 12t\right) \leq K\left(r(V_{n}^{2})\exp\left[-\frac{(t/2)^{2}}{\sigma^{2} + 4t\mathbb{E}M}\right] + \mathbb{P}\left(M \geq t\right) + \left(\left(\frac{p}{\log(ep)}\right)^{p}\frac{\mathbb{E}M^{p}}{t^{p}}\right)^{2}\right)$$

whenever $t \geq 2(\sigma \vee \mathbb{E}M/3)$ and where K is an absolute constant.

Proof. We will continue using the notation introduced in the proof of Proposition 2.1. First of all, note that $\sup_{\|v\|_{2}=1} Q_{1/2}\left(\left\langle \left(\sum_{j=1}^{n} W_{j}\right) v, v\right\rangle\right) \leq \sigma\sqrt{2}$ for the choice of σ stated above. Next, plugging the inequality

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\| > t\right) \leq \mathbb{P}\left(\left\|\sum_{k=1}^{n}\widetilde{W_{k}}\right\| > t/2\right) + \mathbb{P}\left(\left\|\sum_{k=1}^{n}\Delta_{k}\right\| > t/2\right)$$

into relation (3) with V_n^2 and σ^2 specified in the conditions of the theorem, we deduce that

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{k}\right\| > 12t\right) \leq 128r\left(V_{n}^{2}\right)\exp\left[-\frac{(t/2)^{2}}{\sigma^{2} + tU/6}\right] + 4\mathbb{P}\left(M > t\right) + 16\left(\mathbb{P}\left(\left\|\sum_{k=1}^{n} \Delta_{k}\right\| > t/2\right)\right)^{2}.$$

Next, we apply Markov's inequality to get the bound

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\| > t/2\right) \leq \frac{\mathbb{E}\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\|^{p}}{(t/2)^{p}}.$$

Theorem 4.2 implies that for all $p \ge 2$,

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\|^{p}\right)^{1/p} \leq K \frac{p}{\log(p)} \left(\mathbb{E}\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\| + \left(\mathbb{E}\max_{k\leq n} \|\mathbf{\Delta}_{k}\|^{p}\right)^{1/p}\right).$$

Moreover, if we set $U := 24\mathbb{E}M$, then Proposition 4.2 applies with q = 1 and $t_0 = 0$, the latter due to the inequality

(8)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} \Delta_{k}\right\| > 0\right) \le \mathbb{P}\left(M > U\right) \le \frac{\mathbb{E}M}{U} \le 1/24.$$

Therefore,

(9)
$$\mathbb{E}\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\| \leq 6\mathbb{E}\max_{k \leq n} \|\mathbf{\Delta}_{k}\| \leq 6\mathbb{E}\max_{k \leq n} \|W_{k}\| \leq 6\left(\mathbb{E}\max_{k \leq n} \|W_{k}\|^{p}\right)^{1/p},$$

implying that

(10)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} \Delta_{k}\right\| > t/2\right) \leq \left(K\frac{p}{\log(ep)}\right)^{p} \frac{\mathbb{E}M^{p}}{t^{p}}$$

Note that whenever 0 , the same result still holds since

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} \boldsymbol{\Delta}_{k}^{(1)}\right\|^{p}\right)^{1/p} \leq K \mathbb{E}^{1/p} \max_{k \leq n} \|W_{k}\|^{p}$$

in view of Proposition 4.2. The conclusion follows.

Remark 2.2. In principle, one can increase the power in the last term on the righthand side of (7) from 2 to 2^k by applying Hoffmann-Jørgensen inequality k times.

2.2. Moment inequalities. Now we will establish moment inequalities by integrating the tail estimates of Proposition 2.1. As before, assume that V_n^2 satisfies $V_n^2 \succeq \sum_k \mathbb{E}W_k^2$ and let $\sigma^2 = ||V_n^2||$.

Theorem 2.2. Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of centered, independent, selfadjoint random matrices, and let $Q_p = \inf\{s > 0 : \mathbb{P}(\|\sum_{k=1}^n \Delta_k\| > s/2) \leq \frac{1}{8}3^{-p}\}$. Then for all $p \geq 1$,

(11)
$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\sigma\sqrt{q} + qQ_{1} + Q_{p} + \mathbb{E}^{1/p}M^{p}\right),$$

where $q = \log(r(V_n^2)) \lor p$ and K > 0 is an absolute constant. In particular, we have the following "closed-form" Rosenthal-type moment inequalities:

(12)
$$\left(\mathbb{E} \left\| \sum_{k=1}^{n} W_k \right\|^p \right)^{1/p} \leq K \left(\sigma \sqrt{q} + q \mathbb{E} M + \frac{p}{\log(ep)} (\mathbb{E} M^p)^{1/p} \right) \text{ and}$$

(13)
$$\left(\mathbb{E} \left\| \sum_{k=1}^{n} W_k \right\|^p \right)^{1/p} \leq K \left(\sigma \sqrt{q} + \log(r(V_n^2)) \mathbb{E} M + p \|M\|_{\psi_1} \right).$$

Remark 2.3. The well-known relation $||M||_{\psi_1} \leq K \log(n) \max_{k \leq n} ||X_k||_{\psi_1}$ (van der Vaart and Wellner, 1996) could be useful when combined with the inequality (13).

Proof. Observe that

(14)
$$\mathbb{E} \left\| \sum_{k=1}^{n} W_{k} \right\|^{p} \leq 2^{p} \mathbb{E} \left\| \sum_{k=1}^{n} \varepsilon_{k} W_{k} \right\|^{p}$$
$$= 2^{p} p \int_{0}^{\infty} t^{p-1} \mathbb{P} \left(\left\| \sum_{k=1}^{n} \varepsilon_{k} W_{k} \right\| \geq t \right) dt = 6^{p} \cdot p \int_{0}^{\infty} t^{p-1} \mathbb{P} \left(\left\| \sum_{k=1}^{n} \varepsilon_{k} W_{k} \right\| \geq 3t \right) dt,$$

where we used the symmetrization inequality (van der Vaart and Wellner, 1996, Lemma 2.3.6) on the first step and the integration by parts formula on the second step, and the linear change of variables on the third step. To estimate the last integral, we choose $U = Q_1/2$ and apply inequality (4) in the range $t \ge t_0 := 2(\sigma \lor U/3)$ to

deduce that

$$\mathbb{E} \left\| \sum_{k=1}^{n} W_{k} \right\|^{p} \leq 12^{p} (\sigma \vee U/3)^{p} + 6^{p} p \left(\int_{0}^{\infty} t^{p-1} \left(16r \left(V_{n}^{2} \right) \exp \left[-\frac{(t/2)^{2}}{\sigma^{2} + tU/6} \right] \wedge 1 \right) dt + \int_{0}^{\infty} t^{p-1} \mathbb{P} \left(M \geq t \right) dt + \int_{0}^{\infty} 4t^{p-1} \mathbb{P} \left(\left\| \sum_{k=1}^{n} \Delta_{k} \right\| > t/2 \right) \mathbb{P} \left(\left\| \sum_{k=1}^{n} \varepsilon_{k} W_{k} \right\| > t \right) dt \right)$$

Recalling the definition of Q_p , one easily checks that

$$6^{p} p \int_{0}^{\infty} 4t^{p-1} \mathbb{P}\left(\left\|\sum_{k=1}^{n} \mathbf{\Delta}_{k}\right\| > t/2\right) \mathbb{P}\left(\left\|\sum_{k=1}^{n} \varepsilon_{k} W_{k}\right\| > t\right) dt$$
$$\leq 4 \cdot 6^{p} Q_{p}^{p} + 2^{p-1} \mathbb{E}\left\|\sum_{k=1}^{n} \varepsilon_{k} W_{k}\right\|^{p}.$$

Combined with the first line of display (14), this inequality implies that

$$2^{p-1}\mathbb{E}\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\|^{p} \leq 12^{p}(\sigma \vee U/3)^{p} + 6^{p}\mathbb{E}M^{p} + 4 \cdot 6^{p}Q_{p}^{p} + 6^{p}p\int_{0}^{\infty}t^{p-1}\left(16r\left(V_{n}^{2}\right)\exp\left[-\frac{(t/2)^{2}}{\sigma^{2} + tU/6}\right] \wedge 1\right)dt.$$

Application of Corollary 4.1 to the last integral yields the inequality (11). Finally, let us prove the inequalities (12) and (13). To this end, we need to obtain the upper bounds for the quantities Q_1 and Q_p . To estimate Q_1 , recall the inequality (8) which implies that $Q_1/2 \leq 24\mathbb{E}M$. Similary, by (10) and Markov's inequality, we deduce that $Q_p \leq K \frac{p}{\log(ep)} \mathbb{E}^{1/p} M^p$. If, on the other hand, $\|M\|_{\psi_1} < \infty$, then the second inequality of Theorem 4.2 combined with the bound (9) and the well-known estimate $\mathbb{E}M \leq K \|M\|_{\psi_1}$ imply that $\|\sum_{k=1}^n \Delta_k\|_{\psi_1} \leq \|M\|_{\psi_1}$, whence

$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} \Delta_{k}\right\| > t\right) \le e^{-C\frac{t}{\|M\|_{\psi_{1}}}}$$

and $Q_p \leq C'p \|M\|_{\psi_1}$. Let us remark that the inequality (12) can be also obtained by integrating the tail bound (7) directly.

10 MORITZ JIRAK, STANISLAV MINSKER, YIQIU SHEN, AND MARTIN WAHL

Next, we deduce a version of the previous result that holds for sums of nonnegative definite random matrices.

Corollary 2.1. Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of independent, nonnegative definite matrices and let $M = \max_{j=1,\ldots,n} ||W_j||$. Moreover, let $A_n := \sum_{j=1}^n \mathbb{E}W_j$. Then for all $p \ge 1$,

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\|A_{n}\| + q \mathbb{E}M + \frac{p}{\log(ep)}(\mathbb{E}M^{p})^{1/p}\right) \text{ and}$$
$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\|A_{n}\| + \log(r(A_{n}))\mathbb{E}M + p\|M\|_{\psi_{1}}\right),$$

where $q = \log(r(A_n)) \lor p$ and K > 0 is an absolute constant.

Proof. In view of Minkowski's inequality followed by the symmetrization inequality,

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq \|A_{n}\| + \left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k} - \mathbb{E}W_{k}\right\|^{p}\right)^{1/p} \\ \leq \|A_{n}\| + 2\mathbb{E}\left(\mathbb{E}\left\|\sum_{k=1}^{n} \varepsilon_{k} W_{k}\right\|^{p}\right)^{1/p},$$

where $\varepsilon_1, \ldots, \varepsilon_n$ are i.i.d random signs independent from W_1, \ldots, W_n . To estimate the second term in the sum above, we will apply the inequality (12) together with the following choice of V_n^2 : recall that in (12), we set $U = 24\mathbb{E}M$ and note that for all j,

$$\mathbb{E}\left[W_j^2 I\{\|W_j\| \le 24\mathbb{E}M\}\right] \preceq 24\mathbb{E}M \cdot \mathbb{E}W_j$$

since $W_j \succeq 0$ with probability 1. This relation implies that we can set $V_n^2 = 24\mathbb{E}M \cdot A_n$, whence $r(V_n^2) = r(A_n)$. Moreover, $\sigma\sqrt{q} = \sqrt{24q\|A_n\|\mathbb{E}M} \leq \|A_n\| + 6q\mathbb{E}M$, hence (12) yields the bound

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n}\varepsilon_{k}W_{k}\right\|^{p}\right)^{1/p} \leq K'\left(\|A_{n}\| + q\mathbb{E}M + \frac{p}{\log(ep)}\left(\mathbb{E}M^{p}\right)^{1/p}\right),$$

implying the claim. The second inequality is obtained in a similar manner where the inequality (13) is used in place of (12). \Box

2.3. Inequalities for empirical processes. The only part of the previous arguments that exploits the "non-commutative" nature of the random variables is the application of Matrix Bernstein's inequality. In this section, we state the results produced by our method for general empirical processes. The only required modification is the application of Bousquet's version of Talagrand's concentration inequality (38) in place of Bernstein's inequality. We state only the versions of Theorems 2.1 and (11) and remark on the key differences. The required changes to the proofs are minimal hence we avoid the details.

Let \mathcal{F} be a set of measurable real-valued functions defined on some measurable space S and let X_1, \ldots, X_n be i.i.d copies of an S-valued random variable X. Assume that $\mathbb{E}f(X) = 0$ for all $f \in \mathcal{F}$. Let us set $F(x) := \sup_{f \in \mathcal{F}} |f(x)|$, $M = \max_{k \leq n} F(X_k)$, and suppose that $\mathbb{E}M^p < \infty$ for some $p \geq 2$. Denote $Z = \sup_{f \in \mathcal{F}} \sum_{k=1}^n f(X_k)$; for simplicity, we will assume that Z is measurable. Finally, let σ_* satisfy $\sigma_*^2 \geq n \sup_{f \in \mathcal{F}} \mathbb{E}f^2(X)$. For example, in the main case of interest of this paper, $\|\sum_{k=1}^n W_k\| = \sup_{\|v\|_2 = 1} |\operatorname{tr}\left((\sum_{k=1}^n W_k)vv^T\right)|$ corresponding to $\mathcal{F} = \{f_v(\cdot) = |\operatorname{tr}\left((\cdot)vv^T\right)|, \|v\|_2 = 1\}$. The following result can be viewed as an extension of Adamczak's inequality (Adamczak, 2008) to the heavy-tailed case.

Theorem 2.3. For all $t \ge \sqrt{2}\sigma_*$,

$$\mathbb{P}\left(Z > 24\left(\mathbb{E}Z + t\right)\right) \le K \left(\exp\left(-\frac{t^2}{2\sigma_*^2 + 64t\mathbb{E}M}\right) + \mathbb{P}\left(M \ge t\right) + \left(\frac{p}{\log(ep)}\right)^{2p} \left(\frac{\mathbb{E}M^p}{t^p}\right)^2\right)$$

where K is an absolute constant.

Integrating this tail bound, we obtain the following moment inequalities.

Theorem 2.4. For all $p \ge 1$,

(15)
$$(\mathbb{E}Z^p)^{1/p} \le K \left(\mathbb{E}Z + \sigma_* \sqrt{p} + p \mathbb{E}M + \frac{p}{\log(ep)} (\mathbb{E}M^p)^{1/p} \right).$$

Let us compare this result with the bound of Theorem 2.2. When applied to the sums of random matrices, we get the inequality

$$\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\|^{p}\right)^{1/p} \leq K\left(\mathbb{E}\left\|\sum_{k=1}^{n} W_{k}\right\| + \sigma_{*}\sqrt{p} + p\mathbb{E}M + \frac{p}{\log(ep)}(\mathbb{E}M^{p})^{1/p}\right)$$

12 MORITZ JIRAK, STANISLAV MINSKER, YIQIU SHEN, AND MARTIN WAHL

The main difference is that this bounds includes $\mathbb{E} \|\sum_{k=1}^{n} W_k\|$ on the right-hand side. However, for large values of p, it is better than (12) since σ_*^2 can be much smaller than $\|\sum_k \mathbb{E} W_k^2\|$. Moreover, Theorem 2.4 improves upon the inequality proved by (Boucheron et al., 2005, Theorem 12): the latter states that for all $p \geq 2$,

$$\left(\mathbb{E}Z^p\right)^{1/p} \le K \left(\mathbb{E}Z + \sigma_* \sqrt{p} + p(\mathbb{E}M^p)^{1/p}\right).$$

The estimate provided by (15) is better for large values of p if $\mathbb{E}M$ is smaller than $\mathbb{E}^{1/p}M^p$.

3. Applications

In this section, we apply the inequalities to get improved bounds to two classical problems - matrix subsampling and covariance estimation.

3.1. Norms of random submatrices. Let B be a self-adjoint matrix, and let $\delta_1, \ldots, \delta_d$ be i.i.d. Bernoulli random variables with $\mathbb{E}\delta_1 = \delta \in (0, 1)$. Define $R = \text{diag}(\delta_1, \ldots, \delta_d)$. We are interested in the spectral norm of the matrix BR formed by the columns B_i , $i \in I$ of B with indices corresponding to the random set $I = \{1 \leq i \leq d : \delta_i = 1\}$. This problem has previously been studied by Rudelson and Vershynin (2007) and Tropp (2015) who showed that

(16)
$$\mathbb{E}\|BR\|^2 \le K\left(\delta\|B\|^2 + \frac{\log(n\delta)}{\lfloor\delta^{-1}\rfloor} \sum_{k=1}^{\lfloor\delta^{-1}\rfloor} \|B_{(k)}\|_2^2\right)$$

and

$$\mathbb{E}\|BR\|^{2} \leq 1.72 \left(\delta\|B\|^{2} + \log\left(2\frac{\|B\|_{\mathrm{F}}^{2}}{\|B\|^{2}}\right) \|B_{(1)}\|_{2}^{2}\right)$$

respectively, where $B_{(j)}$ denotes the column with the *j*-th largest norm and *K* is a numerical constant. Note that $||B_{(1)}||_2^2 \ge \frac{1}{\lfloor \delta^{-1} \rfloor} \sum_{k=1}^{\lfloor \delta^{-1} \rfloor} ||B_{(k)}||_2^2$ but it is possible that $\log(n\delta) > \log\left(2\frac{||B||_F^2}{||B||^2}\right)$ when the matrix *B* has small "stable rank" srank(*B*) := $\frac{||B||_F^2}{||B||^2}$. We will show below that Tropp's bound can be improved, and that (16) holds with $\log(n\delta)$ replaced with $\log(n\delta) \wedge \log(\operatorname{srank}(B))$. To this end, let e_1, \ldots, e_d denote the standard Euclidean basis, and observe that

$$||BR||^2 = \left\|\sum_{k=1}^d \delta_k B_k e_k^T\right\|^2 = \left\|\sum_{k=1}^d \delta_k B_k B_k^T\right\|.$$

We will apply Corollary 2.1 to the last expression with $W_k = \delta_k B_k B_k^T$. Note that $A_n = \delta B B^T$ so that $||A_n|| = \delta ||B||^2$, and that $\mathbb{E}M = \mathbb{E}(\max_{k \leq d} \delta_k ||B_k||_2^2)$. According to Lemma 5.1 in (Rudelson and Vershynin, 2007) or Proposition 2.3 in (Tropp, 2008),

$$\mathbb{E} \max_{k=1,\dots,d} \delta_k \|B_k\|_2^2 \le \frac{2}{\lfloor \delta^{-1} \rfloor} \sum_{k=1}^{\lfloor \delta^{-1} \rfloor} \|B_{(k)}\|_2^2.$$

Finally, the effective rank $r(A_n) = \frac{\|B\|_F^2}{\|B\|^2}$ coincides with the stable rank of *B*. We record the following bound.

Corollary 3.1. The inequalities

$$\mathbb{E}||BR||^{2} \leq K\left(\delta||B||^{2} + \log\left(\frac{||B||_{\mathrm{F}}^{2}}{||B||^{2}}\right) \frac{1}{\lfloor\delta^{-1}\rfloor} \sum_{k=1}^{\lfloor\delta^{-1}\rfloor} ||B_{(k)}||_{2}^{2}\right)$$

and

$$\mathbb{E}\|BR - \delta B\|^{2} \le K(1 - \delta) \left(\delta \|B\|^{2} + \log\left(\frac{\|B\|_{\mathrm{F}}^{2}}{\|B\|^{2}}\right) \frac{1}{\lfloor \delta^{-1} \rfloor} \sum_{k=1}^{\lfloor \delta^{-1} \rfloor} \|B_{(k)}\|_{2}^{2} \right)$$

hold for all $\delta \in (0,1)$ and a numerical constant K > 0.

Proof. The first inequality has already been established above. The proof of the second bound is quite similar: it suffices to note that

$$\mathbb{E} \|BR - \delta B\|^2 = \mathbb{E} \left\| \sum_{k=1}^d (\delta_k - \delta)^2 B_k B_k^T \right\|,$$

and that $\left\| \mathbb{E} \sum_{k=1}^d (\delta_k - \delta)^2 B_k B_k^T \right\| = \delta (1 - \delta) \|B\|^2$ and
 $\mathbb{E} \max_{k=1 \le d} (\delta_k - \delta)^2 \|B_k\|_2^2 \le \frac{2(1 - \delta)}{\lfloor \delta^{-1} \rfloor} \sum_{k=1}^{\lfloor \delta^{-1} \rfloor} \|B_{(k)}\|_2^2.$

3.2. Covariance estimation. In this section, we consider applications of our results to the covariance estimation problem. Let $X \in \mathbb{R}^d$ be a random vector such that $\mathbb{E}X = 0$ and $\mathbb{E}XX^{\top} = \Sigma$. Given a sequence $X_1, \ldots, X_n \in \mathbb{R}^d$ of i.i.d. copies of X, what is an upper bound for the error of the sample covariance matrix? In other words, we would like to estimate $r_n := \left\| \frac{1}{n} \sum_{j=1}^n X_j X_j^{\top} - \Sigma \right\|$. One of the long-standing open questions asks for the minimal assumptions on the distribution of X

such that $n = C(\varepsilon)d$ suffices to guarantee that $\mathbb{E}r_n \leq \varepsilon \|\Sigma\|$, or that $r_n \leq \varepsilon \|\Sigma\|$ with high probability. Results of this type are often referred to as the "quantitative versions of the Bai-Yin theorem," after (Bai and Yin, 2008). Let us give an (incomplete) overview of the rich history of the problem. It has long been known that sub-Gaussian distributions satisfy the required conditions (Vershynin, 2012). Moreover, very general and precise characterization of the behavior of the sample covariance of Gaussian random vectors with values in Banach spaces has been found by Koltchinskii and Lounici (2017) and, very recently, further sharpened by Han (2022) in the finite-dimensional case. For the log-concave and the sub-exponential distributions, the problem was first considered by Kannan et al. (1997), and the bounds were significantly improved and refined by Bourgain (1996) and Rudelson (1999). It took much longer to eliminate the unnecessary logarithmic factors, until the problem was finally solved by Adamczak et al. (2010, 2011). Finally, the case of heavy-tailed distributions was investigated by Vershynin (2011); Srivastava and Vershynin (2013); Mendelson and Paouris (2014); Guédon et al. (2017) and Tikhomirov (2018), who showed that $4 + \varepsilon$ moments are sufficient to get the desired bound. Specifically, Tikhomirov's results imply that if $\Sigma = I_d$ and $\sup_{\|v\|_2=1} \mathbb{E} |\langle X, v \rangle|^p = T < \infty$ for some p > 4, then

$$r_n \le C(p) \left(T^{2/p} \sqrt{\frac{d}{n}} + \frac{\max_j \|X_j\|_2^2}{n} \right)$$

with probability at least 1-1/n. Abdalla and Zhivotovskiy (2024) refined Tikhomirov's estimates and essentially showed that $n = C(\varepsilon)r(\Sigma)$ samples suffices to get the desired guarantees in expectation, although they considered the sample covariance based on properly truncated random vectors. Next, we show that the results by Abdalla and Zhivotovskiy (2024) can be combined with the moment inequalities developed in this paper to get a sharp moment inequality for r_n .

Theorem 3.1. Let $X \in \mathbb{R}^d$ be a random vector such that $\mathbb{E}X = 0$ and $\mathbb{E}[XX^\top] = \Sigma$. Let $X_1, \ldots, X_n \in \mathbb{R}^d$ be i.i.d. copies of X. Assume that $\frac{r(\Sigma)}{n} \leq c$ for a sufficiently small positive constant c, and that for some p > 4

(17)
$$\sup_{\|v\|_{2}=1} \frac{\mathbb{E}^{1/p} \left| \langle X, v \rangle \right|^{p}}{\mathbb{E}^{1/2} \langle X, v \rangle^{2}} = \kappa < \infty.$$

Then

$$\left(\mathbb{E}\left\|\frac{1}{n}\sum_{j=1}^{n}X_{j}X_{j}^{\top}-\Sigma\right\|^{2}\right)^{1/2} \leq C(\kappa,p)\left(\|\Sigma\|\sqrt{\frac{r(\Sigma)}{n}}+\frac{\mathbb{E}^{2/p}\max_{j\leq n}\|X_{j}\|_{2}^{p}}{n}\right)$$

Proof. Our proof builds on the results by Abdalla and Zhivotovskiy (2024) which in turn sharpen the inequality due to Tikhomirov (2018). Before we dive into the details, let us mention that the "hypercontractivity" condition (17) implies in particular that $\|\mathbb{E}\|X\|^2 X X^T \| \leq \kappa^2 \operatorname{tr}(\Sigma) \|\Sigma\|$; the proof of this fact can be found in (Wei and Minsker, 2017, Lemma 2.3). Note that

$$\left\|\frac{1}{n}\sum_{j=1}^{n}X_{j}X_{j}^{\top}-\Sigma\right\| = \sup_{\|v\|_{2}=1}\left|\frac{1}{n}\sum_{j=1}^{n}\langle X_{j},v\rangle^{2}-\mathbb{E}\langle X,v\rangle^{2}\right|.$$

Let us state the following decomposition of the error into "peaky" and "spread" parts (Tikhomirov, 2018) that holds for arbitrary $\lambda > 0$:

(18)
$$\sup_{\|v\|_{2}=1} \left| \frac{1}{n} \sum_{j=1}^{n} \langle X_{j}, v \rangle^{2} - \mathbb{E} \langle X, v \rangle^{2} \right| \leq \underbrace{\sup_{\|v\|_{2} \leq 1} \frac{1}{n} \sum_{j=1}^{n} \langle X_{j}, v \rangle^{2} \mathbf{1} \{\lambda \langle X_{i}, v \rangle^{2} > 1\}}_{\text{Peaky part}} + \underbrace{\sup_{\|v\|_{2} \leq 1} \left| \frac{1}{\lambda n} \sum_{j=1}^{n} \psi(\lambda \langle X_{j}, v \rangle^{2}) - \mathbb{E} \langle X, v \rangle^{2} \right|}_{\text{Spread part}},$$

where

$$\psi(x) = \begin{cases} x, & \text{for } x \in [-1, 1];\\ \operatorname{sign}(x) & \text{for } |x| > 1. \end{cases}$$

We will estimate the two terms separately, starting with the "spread" part. To this end, we will apply Proposition 4 in (Abdalla and Zhivotovskiy, 2024) which implies that for $\lambda = \frac{1}{\kappa^2 \|\Sigma\|} \sqrt{\frac{r(\Sigma)}{n}}$,

$$\sup_{\|v\|_{2} \le 1} \left| \frac{1}{\lambda n} \sum_{j=1}^{n} \psi(\lambda \langle X_{j}, v \rangle^{2}) - \mathbb{E} \langle X, v \rangle^{2} \right| \le C \kappa^{2} \|\Sigma\| \left(\sqrt{\frac{r(\Sigma)}{n}} + \frac{t}{\sqrt{r(\Sigma)n}} \right)$$

with probability at least $1 - e^{-t}$. For $t = r(\Sigma)$, we get in particular that

(19)
$$\sup_{\|v\|_{2} \le 1} \left| \frac{1}{\lambda n} \sum_{j=1}^{n} \psi(\lambda \langle X_{j}, v \rangle^{2}) - \mathbb{E} \langle X, v \rangle^{2} \right| \le C \kappa^{2} \|\Sigma\| \sqrt{\frac{r(\Sigma)}{n}}$$

with probability at least $1 - e^{-r(\Sigma)}$. Next, we will estimate the "peaky" term in the inequality (18). Equation (5) in the work by Abdalla and Zhivotovskiy (2024) states

that for all subsets $\mathcal{J} \subseteq [n]$ of cardinality at most k,

(20)
$$\left\|\frac{1}{n}\sum_{j\in\mathcal{J}}X_jX_j^{\top}\right\| \le \frac{f(k,[n])}{n}$$

for the function f(k, [n]) defined via

$$f(k, [n]) = \sup_{\|y\|_2 = 1, \|y\|_0 \le k, \operatorname{supp}(y) \subseteq [n]} \left\| \sum_{j=1}^n y_j X_j \right\|_2^2$$

and the bounds holds uniformly over all such subsets J. The following result provides a bound for f(k, [n]).

Theorem 3.2. (Abdalla and Zhivotovskiy, 2024, Theorem 3) Assume that $\frac{r(\Sigma)}{n} \leq c'$ for a sufficiently small positive constant c'. Then

$$f(k, [n]) \le C(p, \kappa) \left(\max_{j \le n} \|X_j\|_2^2 + \|\Sigma\| k \left(\frac{n}{k}\right)^{4/(4+p)} \log^4 \frac{n}{k} \right),$$

with probability at least $1 - \frac{C(p)}{n}$, and the bound holds simultaneously for all integers k satisfying $r(\Sigma) \leq k \leq c'n$.

Since p > 4, this result implies that

(21)
$$f(k, [n]) \le C'(p, \kappa) \left(\max_{j \le n} \|X_j\|_2^2 + \|\Sigma\|\sqrt{nk} \right).$$

Next, let $\lambda = \frac{1}{\kappa^2 \|\Sigma\|} \sqrt{\frac{r(\Sigma)}{n}}$. Following Tikhomirov (2018); Abdalla and Zhivotovskiy (2024), let us define the random set

$$I_v = \left\{ j \in [n] : \langle X_j, v \rangle^2 > 1/\lambda \right\}$$

and $m = \sup_{\|v\|_2 \le 1} |I_v|$. Then, in view of the inequality (20), we see that

(22)
$$\frac{m}{n\lambda} \leq \sup_{\|v\|_2 \leq 1} \frac{1}{n} \sum_{j=1}^n \langle X_j, v \rangle^2 \mathbf{1} \{\lambda \langle X_i, v \rangle^2 > 1\} \leq \frac{f(m, [n])}{n}$$

Now, if $r(\Sigma) \leq m \leq c'n$, then we can employ Theorem 3.2 to derive the following bound that holds with probability at least $1 - \frac{C(p)}{n}$:

$$m \leq \lambda f(m, [n]) \leq C'(p) \left(\max_{j \leq n} \|X_j\|_2^2 + \|\Sigma\|\sqrt{nm} \right) \cdot \frac{1}{\kappa^2 \|\Sigma\|} \sqrt{\frac{r(\Sigma)}{n}}$$
$$\leq \frac{C'(p)}{\kappa^2} \left(\frac{\max_{j \leq n} \|X_j\|^2}{\|\Sigma\|} \sqrt{\frac{r(\Sigma)}{n}} + \sqrt{m}\sqrt{r(\Sigma)} \right).$$

Solutions to the inequality $x \leq a\sqrt{x} + b$ satisfy $x \leq 2\max(a^2, b)$. Therefore, with probability at least $1 - \frac{C(p)}{n}$,

(23)
$$m \le C_1(p) \max\left(r(\Sigma), \frac{\max_{j\le n} \|X_j\|^2}{\kappa^2 \|\Sigma\|} \sqrt{\frac{r(\Sigma)}{n}}\right)$$

It remains to show that $m \leq c'n$ if $r(\Sigma) < cn$ for c small enough (clearly, if $m < r(\Sigma)$, then (23) holds). By the definition of I_v , for any $v \in \mathbb{R}^d$,

.

$$(24) \quad |I_v| = \sum_{j=1}^n \mathbf{1}\left\{\langle X_j, v \rangle^2 > \kappa^2 \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}}\right\} = \sum_{j=1}^n \mathbf{1}\left\{\frac{|\langle X_j, v \rangle|}{\left(\kappa^2 \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}}\right)^{1/2}} > 1\right\}$$
$$\leq \sum_{j=1}^n \rho\left(\frac{|\langle X_j, v \rangle|}{\left(\kappa^2 \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}}\right)^{1/2}}\right),$$

where

$$\rho(x) = \begin{cases} 0 & x \le 1/2 \\ 2x - 1 & x \in (1/2, 1] \\ 1 & x > 1 \end{cases}$$

is such that $\mathbf{1}\{x \ge 1/2\} \ge \rho(x) \ge \mathbf{1}\{x \ge 1\}$. For brevity, set

$$Z_j(v) := \frac{|\langle X_j, v \rangle|}{\left(\kappa^2 \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}}\right)^{1/2}} \quad \text{and} \quad S = \sup_{\|v\|_2 \le 1} \left(\sum_{j=1}^n \rho(Z_j(v)) - \mathbb{E}\rho(Z_j(v))\right).$$

In view of Markov's inequality and assumption (17),

(25)
$$\mathbb{E}\rho(Z_j(v)) \le \mathbb{P}\left(\langle X, v \rangle^2 > \frac{\kappa^2}{2} \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}}\right) \le \frac{4\mathbb{E}\langle X, v \rangle^4}{\kappa^4 \|\Sigma\|^2} \cdot \frac{r(\Sigma)}{n} \le \frac{4r(\Sigma)}{n}$$

Denoting $\sigma^2 = \sup_{||v||_2 \le 1} \operatorname{Var}(\rho(Z_1(v)))$, we deduce that

$$\mathbb{P}\left(\sup_{\|v\|_{2}\leq 1}|I_{v}|\geq c'n\right)\leq \mathbb{P}\left(\sup_{\|v\|_{2}=1}\sum_{j=1}^{n}\rho(Z_{j})>c'n\right)\leq \mathbb{P}\left(S>c'n-4r(\Sigma)\right).$$

We will apply Theorem 4.3 to estimate the right hand side in the display above. Specifically, $\mathbb{P}\left(S > c'n - 4r(\Sigma)\right) \leq e^{-t}$ whenever

$$c'n - 4r(\Sigma) - 2\mathbb{E}S - \sigma\sqrt{2tn} - 4t/3 \ge 0.$$

To prove that this relation holds for suitable choices of parameters c and t (where $r(\Sigma)/n \leq c$), first observe that

$$\sigma^2 \le \mathbb{E}(\rho(Z)^2 \le \mathbb{E}\rho(Z) \le \frac{4r(\Sigma)}{n},$$

where the last inequality follows from the bound (25). Next we will estimate $\mathbb{E}S$. Let $\varepsilon_1, \ldots \varepsilon_n$ be a sequence of independent random signs. The standard argument based on the symmetrization and contraction inequalities (Ledoux and Talagrand, 2013, Theorem 4.4), together with the fact that $\rho(x)$ is Lipschitz continuous with Lipschitz constant equal to 2, yields that

ī

ī

$$\mathbb{E}S \leq 2\mathbb{E}\sup_{\|v\|_{2}\leq 1} \left|\sum_{j=1}^{n} \varepsilon_{j}\rho(Z_{j})\right| \leq 8\mathbb{E}\sup_{\|v\|_{2}\leq 1} \left|\sum_{j=1}^{n} \varepsilon_{j}\frac{|\langle X_{j}, v\rangle|}{\left(\kappa^{2}\|\Sigma\|\sqrt{\frac{n}{r(\Sigma)}}\right)^{1/2}}\right|$$
$$\leq \frac{16}{\left(\kappa^{2}\|\Sigma\|\sqrt{\frac{n}{r(\Sigma)}}\right)^{1/2}} \left(\mathbb{E}\sup_{\|v\|_{2}\leq 1}\left(\sum_{j=1}^{n}\langle X_{j}, v\rangle\right)^{2}\right)^{1/2}.$$

Since

$$\mathbb{E}\sup_{\|v\|_{2}\leq 1}\left(\sum_{j=1}^{n}\langle X_{j},v\rangle\right)^{2} = \mathbb{E}\left\|\sum_{j=1}^{n}X_{j}\right\|^{2}\leq\kappa^{2}n\|\Sigma\|r(\Sigma)$$

by assumption (17), we conclude that

(26)
$$\mathbb{E}\sup_{\|v\|_2 \le 1} \left(\sum_{j=1}^n \rho(Z_j) - \mathbb{E}\rho(Z_j) \right) \le 16n^{1/4} (r(\Sigma))^{3/4}.$$

As a consequence, we have to choose c and t such that

(27)
$$4r(\Sigma) + 32n^{1/4}(r(\Sigma))^{3/4} + 2\sqrt{2t}\sqrt{r(\Sigma)} + 4t/3 \le c'n,$$

which is satisfied if both $r(\Sigma)$ t do not exceed a constant times n. When (27) holds, we conclude that $m \leq c'n$ with probability at least $1 - e^{-t}$, and that the inequality (23) holds with probability at least 1 - c'(p)/n. Combining this result with the estimates (21), (22) and Theorem 3.2, we deduce that with probability at least 1 - c'(p)/n, the "peaky" term admits the upper bound of the form

(28)
$$\sup_{\|v\|_{2} \leq 1} \frac{1}{n} \sum_{j=1}^{n} \langle X_{j}, v \rangle^{2} \mathbf{1} \left\{ \langle X_{i}, v \rangle^{2} > \kappa^{2} \|\Sigma\| \sqrt{\frac{n}{r(\Sigma)}} \right\}$$
$$\leq \frac{f(m, [n])}{n} \leq C_{2}(p, \kappa) \left(\frac{\max_{j \leq n} \|X_{j}\|_{2}^{2}}{n} + \|\Sigma\| \frac{r(\Sigma)}{n} \right)$$

Combining the estimates (19) and (28) with the decomposition (18), we conclude that with probability at least $1 - e^{-r(\Sigma)} - c'(p)/n$,

(29)
$$\left\|\frac{1}{n}\sum_{j=1}X_jX_j^{\top} - \Sigma\right\| \le C(p,\kappa)\left(\frac{\max_{j\le n}\|X_j\|_2^2}{n} + \|\Sigma\|\sqrt{\frac{r(\Sigma)}{n}}\right).$$

To obtain the desired bound in expectation, let us define the event

$$A := \left\{ \left\| \frac{1}{n} \sum_{j=1} X_j X_j^\top - \Sigma \right\| \le C(p,\kappa) \left(\frac{\max_{j \le n} \|X_j\|_2^2}{n} + \|\Sigma\| \sqrt{\frac{r(\Sigma)}{n}} \right) \right\}.$$

Then $\mathbb{P}(A^c) \leq e^{-r(\Sigma)} + c'(p)/n$ and

$$\mathbb{E}^{1/2} \left\| \frac{1}{n} \sum_{j=1}^{n} X_j X_j^{\top} - \Sigma \right\|^2 \le C_1(p,\kappa) \left(\frac{\mathbb{E}^{1/2} \max_{j \le n} \|X_j\|_2^4}{n} + \|\Sigma\| \sqrt{\frac{r(\Sigma)}{n}} \right) \\ + \mathbb{E}^{1/2} \left[\left\| \frac{1}{n} \sum_{j=1}^{n} X_j X_j^{\top} - \Sigma \right\|^2 \mathbf{1}(A^c) \right].$$

Hölder's inequality implies that

$$\mathbb{E}^{1/2}\left[\left\|\frac{1}{n}\sum_{j=1}X_jX_j^{\top} - \Sigma\right\|^2 \mathbf{1}(A^c)\right] \le \mathbb{E}^{2/p}\left\|\frac{1}{n}\sum_{j=1}X_jX_j^{\top} - \Sigma\right\|^{p/2} \left(\frac{c'(p)}{n} \vee e^{-r(\Sigma)}\right)^{\frac{p-4}{2p}}$$

Finally, we invoke Rosenthal's inequality (12) to deduce that

$$\mathbb{E}^{2/p} \left\| \frac{1}{n} \sum_{j=1}^{\infty} X_j X_j^{\top} - \Sigma \right\|^{p/2} \le C(p) \left[\sqrt{\frac{r(\Sigma)}{n}} \|\Sigma\| \sqrt{\log(er(\Sigma))} + \frac{\log(er(\Sigma))}{n} \mathbb{E} \max_{j \le n} \|X_j\|_2^2 + \frac{1}{n} \mathbb{E}^{2/p} \max_{j \le n} \|X_j\|_2^p \right].$$

Since $r(\Sigma) < cn$ by assumption, $\log(er(\Sigma)) < C(p) (n \lor e^{r(\Sigma)})^{(p-4)/2p}$, implying the final form of the bound. Let us remark that in the course of the proof, we obtained a slightly stronger inequality

$$\left(\mathbb{E}\left\|\frac{1}{n}\sum_{j=1}^{n}X_{j}X_{j}^{\top}-\Sigma\right\|^{2}\right)^{1/2} \leq C(\kappa,p)\left(\|\Sigma\|\sqrt{\frac{r(\Sigma)}{n}}\right) + \frac{\mathbb{E}^{1/2}\max_{j\leq n}\|X_{j}\|_{2}^{4}}{n}\bigvee\frac{\mathbb{E}^{2/p}\max_{j\leq n}\|X_{j}\|_{2}^{p}}{n}\left(\frac{1}{n}+e^{-r(\Sigma)}\right)^{\frac{p-4}{2p}}\right).$$

3.3. Empirical eigenvector estimation. In this section, we continue the considerations of section 3.2. Let $X \in \mathbb{R}^d$ be a random vector such that $\mathbb{E}X = 0$ and $\mathbb{E}XX^{\top} = \Sigma$, and let $\lambda_1 \geq \cdots \geq \lambda_d$ and u_1, \ldots, u_d be the eigenvalues and the eigenvectors of Σ , respectively. Moreover, let $g_1 = \lambda_1 - \lambda_2$ and $g_j = \min(\lambda_{j-1} - \lambda_j, \lambda_j - \lambda_{j+1})$ for $j = 2, \ldots, d$ be the different spectral gaps, meaning that if $g_j > 0$, then the eigenvector u_j is uniquely determined up to the sign. Given a sequence $X_1, \ldots, X_n \in \mathbb{R}^d$ of i.i.d. copies of X, let $\hat{\lambda}_1 \geq \cdots \geq \hat{\lambda}_d$ and $\hat{u}_1, \ldots, \hat{u}_d$ be the eigenvalues and eigenvectors of the empirical covariance matrix $\hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n X_k X_k^{\top}$, respectively. A question that has been studied for decades asks for perturbation bounds for the empirical spectral characteristics, we refer for instance to Anderson (1963); Horváth and Kokoszka (2012); Hsing and Eubank (2015) for some classical and more recent results and applications. For the special case of spectral projectors, the Davis-Kahan inequality (cf. Davis and Kahan (1970)) is among the most prominent tools, see (Jirak and Wahl, 2020; Yu et al., 2015) and the references therein for some recent context. Combining the Davis-Kahan inequality with Theorem 3.1 we get that

(30)
$$\mathbb{E}^{1/2} \left\| \hat{u}_{j} \hat{u}_{j}^{\top} - u_{j} u_{j}^{\top} \right\|_{2}^{2} \leq C(\kappa, p) \left(\frac{\|\Sigma\|}{g_{j}} \sqrt{\frac{r(\Sigma)}{n}} + \frac{\mathbb{E}^{2/p} \max_{k \leq n} \|X_{k}\|_{2}^{p}}{ng_{j}} \right),$$

provided that $g_j > 0$. Note that the requirement $\frac{r(\Sigma)}{n} \leq c$ can be dropped in this case because the left-hand side is always bounded by $\sqrt{2}$. Although prominent, estimates of the type (30) are often sub-optimal, we refer to (Jirak and Wahl, 2020, 2023) for a detailed discussion. It turns out that the complexity of the problem is captured by the *relative rank*

(31)
$$r_j(\Sigma) = \sum_{i \neq j} \frac{\lambda_i}{|\lambda_i - \lambda_j|} + \frac{\lambda_j}{g_j},$$

in contrast to the effective rank $r(\Sigma)$. The following result improves upon the bounds in (Jirak and Wahl, 2023, Corollary 5) as it requires less moments.

Theorem 3.3. Assume that for some p > 4 condition (17) holds. Let $j \in \{1, \ldots, d\}$ be such that $g_j > 0$. Then

$$\mathbb{E}^{1/2} \|\hat{u}_j - u_j\|_2^2 \le \mathbb{E}^{1/2} \|\hat{u}_j \hat{u}_j^\top - u_j u_j^\top\|_2^2 \le C(\kappa, p) \left(\sqrt{\frac{\lambda_j}{g_j}} \sqrt{\frac{r_j(\Sigma)}{n}} + \frac{r_j(\Sigma)}{n^{1-2/p}}\right),$$

where $\|\cdot\|_2$ also denotes the Frobenius norm for matrices.

Remark 3.1. The random vector X can be expressed in terms of the Karhunen-Loève decomposition $X = \sum_{i=1}^{d} \sqrt{\lambda_i} \eta_i u_i$, where the Karhunen-Loève coefficients η_1, \ldots, η_d are uncorrelated with $\mathbb{E}\eta_i^2 = 1$ and defined by $\eta_i = \langle X, u_i \rangle / \sqrt{\lambda_i}$. In this case, (17) holds if η_1, \ldots, η_d form a martingale difference sequence with $\max_{i \leq d} \mathbb{E}^{1/p} |\eta_i|^p \leq \kappa / \sqrt{p-1}$, as can be seen by an application of Burkholder's inequality (Theorem 2.1 in Rio (2009)).

Remark 3.2. Since $\lambda_j \leq ||\Sigma||$ and $|\lambda_k - \lambda_j| \geq g_j$ for $k \neq j$, we have the inequality

$$\sqrt{\frac{\lambda_j}{g_j}}\sqrt{\frac{r_j(\Sigma)}{n}} \le \frac{\|\Sigma\|}{g_j}\sqrt{\frac{r(\Sigma)}{n}},$$

meaning that the first term on the right-hand side of Theorem 3.3 is always an improvement over the corresponding one in (30).

Proof of Theorem 3.3. Suppose that the sign of \hat{u}_j is chosen such that $\langle \hat{u}_j, u_j \rangle \geq 0$. By display (5.24) in Hsing and Eubank (2015) combined with Proposition 1 in Jirak and Wahl (2024) (see also Lemma 2 in Jirak and Wahl (2023)), we have

(32)
$$\|\hat{u}_j - u_j\|_2 \le \|\hat{u}_j \hat{u}_j^\top - u_j u_j^\top\|_2 \le 4\sqrt{2} \|T_j (\hat{\Sigma} - \Sigma) T_j\|$$

with

$$T_j = |R_j|^{1/2} + g_j^{-1/2} u_j u_j^{\top}, \qquad |R_j|^{1/2} = \sum_{i \neq j} \frac{1}{\sqrt{|\lambda_i - \lambda_j|}} u_i u_i^{\top}.$$

Since T_j is symmetric,

$$T_j\left(\hat{\Sigma}-\Sigma\right)T_j = \frac{1}{n}\sum_{k=1}^n (T_jX_k)(T_jX_k)^\top - \mathbb{E}(T_jX)(T_jX)^\top.$$

Thus, estimating $||T_j(\hat{\Sigma} - \Sigma)T_j||$ is again a covariance estimation problem with random vector given by

$$T_j X = \sum_{i \neq j} \left(\frac{\lambda_i}{|\lambda_i - \lambda_j|} \right)^{1/2} \eta_i u_i + \left(\frac{\lambda_j}{g_j} \right)^{1/2} \eta_j u_j$$

with Karhunen-Loève coefficients η_1, \ldots, η_d introduced in Remark 3.1. We now explore the fact that assumption (17) is invariant under linear transformations. Indeed, for any $u \neq 0$, we have

$$\frac{\mathbb{E}^{1/p}|\langle T_j X, u \rangle|^p}{\mathbb{E}^{1/2}\langle T_j X, u \rangle^2} = \frac{\mathbb{E}^{1/p}|\langle X, T_j u \rangle|^p}{\mathbb{E}^{1/2}\langle X, T_j u \rangle^2} \le \sup_{\|v\|_2 = 1} \frac{\mathbb{E}^{1/p}|\langle X, v \rangle|^p}{\mathbb{E}^{1/2}\langle X, v \rangle^2} \le \kappa,$$

implying that (17) also holds for $T_j X$. In addition, setting $v = u_j$ in (17), we get

$$\max_{i \le d} \mathbb{E}^{1/p} |\eta_i|^p \le \kappa.$$

Using the triangle inequality, it follows that

$$\mathbb{E}^{2/p} \|T_j X\|^p = \mathbb{E}^{2/p} \left(\sum_{i \neq j} \frac{\lambda_i}{|\lambda_i - \lambda_j|} \eta_i^2 + \frac{\lambda_j}{g_j} \eta_j^2 \right)^{p/2}$$
$$\leq \sum_{i \neq j} \frac{\lambda_i}{|\lambda_i - \lambda_j|} \mathbb{E}^{2/p} |\eta_i|^p + \frac{\lambda_j}{g_j} \mathbb{E}^{2/p} |\eta_j|^p \leq r_j(\Sigma) \kappa^2$$

This in turn yields the estimate

(33)
$$\mathbb{E}^{2/p} \max_{i \le n} \|T_j X_i\|^p \le n^{2/p} r_j(\Sigma) \kappa^2.$$

Moreover, we have the relations

(34)
$$\operatorname{trace}\left(T_{j}\Sigma T_{j}\right) = r_{j}(\Sigma), \qquad \|T_{j}\Sigma T_{j}\| = \left(\max_{i\neq j}\frac{\lambda_{i}}{|\lambda_{i} - \lambda_{j}|}\right)\bigvee\frac{\lambda_{j}}{g_{j}} \le \frac{2\lambda_{j}}{g_{j}}$$

Theorem 3.1, together with displays (33) and (34), now yields the inequality

$$\begin{split} \mathbb{E}^{1/2} \left\| \hat{u}_j \hat{u}_j^\top - u_j u_j^\top \right\|_2^2 &\leq 4\sqrt{2} \mathbb{E}^{1/2} \|T_j (\hat{\Sigma} - \Sigma) T_j\|^2 \\ &\leq C(\kappa, p) \left(\|T_j \Sigma T_j\| \sqrt{\frac{r(T_j \Sigma T_j)}{n}} + \frac{\mathbb{E}^{2/p} \max_{i \leq n} \|T_j X_i\|_2^p}{n} \right) \\ &\leq C'(\kappa, p) \left(\sqrt{\frac{\lambda_j}{g_j}} \sqrt{\frac{r_j(\Sigma)}{n}} + \frac{r_j(\Sigma)}{n^{1-2/p}} \right), \end{split}$$

provided that $\frac{r_j(\Sigma)}{n} \leq c$. Finally, in case that $\frac{r_j(\Sigma)}{n} > c$ we trivially have

$$\mathbb{E}^{1/2} \left\| \hat{u}_{j} \hat{u}_{j}^{\top} - u_{j} u_{j}^{\top} \right\|_{2}^{2} \le \sqrt{2} \le \frac{\sqrt{2}}{c} \frac{r_{j}(\Sigma)}{n^{1-2/p}}$$

The conclusion follows.

4. AUXILIARY RESULTS

In this section, we collect the background material and technical results that our arguments rely on.

Theorem 4.1. (Minsker, 2017, Theorem 3.1) and (Tropp, 2015, Theorem 7.7.1) Let $W_1, \ldots, W_n \in \mathbb{C}^{d \times d}$ be a sequence of independent, centered, self-adjoint random matrices such that $||W_k|| \leq U$, $k = 1, \ldots, n$ almost surely. Assume that $V_n^2 \succeq \sum_k \mathbb{E}W_k^2$ and let $\sigma^2 = ||V_n^2||$. Then for any $t \geq \sigma + U/3$,

(35)
$$\mathbb{P}\left(\left\|\sum_{k=1}^{n} W_{i}\right\| > t\right) \leq 4r\left(V_{n}^{2}\right)\exp\left[-\frac{t^{2}/2}{\sigma^{2} + tU/3}\right]$$

Corollary 4.1. Let $p \ge 1$. Under the assumptions of Theorem 4.1,

(36)
$$\mathbb{E}^{1/p} \left\| \sum_{k=1}^{n} W_k \right\|^p \le K \left(\sigma \sqrt{q} + Uq \right)$$

where $q = \log(er(V_n^2)) \lor p$ and K > 0 is a numerical constant.

Proof. Inequalities of this type are well known. See for instance (Wainwright, 2019, exercise 2.8) or (Talagrand, 2014, Lemma 2.3.2) where the case p = 1 is considered.

Proposition 4.1. (Hoffmann-Jørgensen inequality: Ledoux and Talagrand, 2013, Proposition 6.7) Let X_1, \ldots, X_n be independent, symmetrically distributed random variables with values in a separable Banach space with norm $\|\cdot\|_B$. Set $S_k = \sum_{i=1}^k X_i$, $k \leq N$. Then for any s, t > 0,

$$\mathbb{P}(\|S_N\|_B > 2t + s) \le 4(\mathbb{P}(\|S_N\|_B > t))^2 + \mathbb{P}\left(\max_{i \le N} \|X_i\|_B > s\right)$$

Proposition 4.2. (Ledoux and Talagrand, 2013, Proposition 6.8) Let $0 < q < \infty$ and let X_1, \ldots, X_n be independent, symmetrically distributed random variables with values in a separable Banach space with norm $\|\cdot\|_B$. Set $S_k = \sum_{i=1}^k X_i$, $k \leq N$. Then for $t_0 = \inf\{t > 0 : \mathbb{P}(\|S_N\|_B > t) \leq (2 \cdot 3^p)^{-1}\}$,

(37)
$$\mathbb{E} \|S_N\|_B^p \le 2 \cdot 3^p \mathbb{E} \max_{i \le N} \|X_i\|_B^p + 2 (3t_0)^p.$$

Theorem 4.2. (Ledoux and Talagrand, 2013, Theorems 6.20 and 6.21) Let X_1, \ldots, X_n be independent random variables with values in a separable Banach space with norm $\|\cdot\|_B$. There exists a numerical constant K such that for all p > 1,

$$\mathbb{E}^{1/p} \left\| \sum_{k=1}^{n} X_k \right\|_B^p \le K \frac{p}{\log(ep)} \left(\mathbb{E} \left\| \sum_{k=1}^{n} X_k \right\|_B + \mathbb{E}^{1/p} \max_k \|X_k\|_B^p \right) \text{ and} \\ \left\| \left\| \sum_{k=1}^{n} X_k \right\|_B \right\|_{\psi_1} \le K \left(\mathbb{E} \left\| \sum_{k=1}^{n} X_k \right\|_B + \left\| \max_k \|X_k\|_B \right\|_{\psi_1} \right).$$

Theorem 4.3. (Bousquet, 2003, Theorem 7.3) Let \mathcal{F} be a countable set of measurable real-valued functions and let X_1, \ldots, X_n be i.i.d. Assume that $\mathbb{E}f(X_1) = 0$ for all $f \in \mathcal{F}$ and that $\sup_{f \in \mathcal{F}} |f(X_1)| \leq U$ with probability 1. Denote $Z = \sup_{f \in \mathcal{F}} \sum_{k=1}^n f(X_k)$. Assume that $\sigma_*^2 \geq n \sup_{f \in \mathcal{F}} \mathbb{E}f^2(X_1)$ and set $v = \sigma_*^2 + 2\mathbb{E}[Z]$. Then for all $t \geq 0$,

(38)
$$\mathbb{P}\left(Z \ge \mathbb{E}Z + \sqrt{2tv} + \frac{tU}{3}\right) \le e^{-t}.$$

The inequality (38) immediately implies that with probability at least $1 - e^{-t}$,

$$Z \le 2\mathbb{E}Z + \sigma_*\sqrt{2t} + 4tU/3$$

References

Abdalla, P. and Zhivotovskiy, N. (2024). Covariance estimation: Optimal dimensionfree guarantees for adversarial corruption and heavy tails. *To appear in the Journal* of the European Mathematical Society.

- Adamczak, R. (2008). A tail inequality for suprema of unbounded empirical processes with applications to markov chains. *Electronic Journal of Probability*, 13:1000– 1034.
- Adamczak, R., Litvak, A., Pajor, A., and Tomczak-Jaegermann, N. (2010). Quantitative estimates of the convergence of the empirical covariance matrix in log-concave ensembles. *Journal of the American Mathematical Society*, 23(2):535–561.
- Adamczak, R., Litvak, A. E., Pajor, A., and Tomczak-Jaegermann, N. (2011). Sharp bounds on the rate of convergence of the empirical covariance matrix. *Comptes Rendus. Mathématique*, 349(3-4):195–200.
- Ahlswede, R. and Winter, A. (2002). Strong converse for identification via quantum channels. *IEEE Transactions on Information Theory*, 48(3):569–579.
- Anderson, T. W. (1963). Asymptotic theory for principal component analysis. Ann. Math. Statist., 34:122–148.
- Bai, Z.-D. and Yin, Y.-Q. (2008). Limit of the smallest eigenvalue of a large dimensional sample covariance matrix. In Advances In Statistics, pages 108–127.
- Bakhshizadeh, M., Maleki, A., and De La Pena, V. H. (2023). Sharp concentration results for heavy-tailed distributions. *Information and Inference: A Journal of the IMA*, 12(3):1655–1685.
- Boucheron, S., Bousquet, O., Lugosi, G., and Massart, P. (2005). Moment inequalities for functions of independent random variables. *The Annals of Probability*, 33(2):514–560.
- Bourgain, J. (1996). Random points in isotropic convex sets. *Convex geometric* analysis, 34:53–58.
- Bousquet, O. (2003). Concentration inequalities for sub-additive functions using the entropy method. In *Stochastic inequalities and applications*, pages 213–247. Springer.
- Chen, R. Y., Gittens, A., and Tropp, J. A. (2012). The masked sample covariance estimator: an analysis using matrix concentration inequalities. *Information and Inference: A Journal of the IMA*, 1(1):2–20.
- Davis, C. and Kahan, W. M. (1970). The rotation of eigenvectors by a perturbation. III. SIAM Journal of Numerical Analysis, 7:1–46.
- Dirksen, S. (2011). Noncommutative and vector-valued Rosenthal inequalities. *Ph.D.* thesis.
- Einmahl, U. and Li, D. (2008). Characterization of LIL behavior in Banach space. Transactions of the American Mathematical Society, 360(12):6677–6693.
- Fuk, D. K. and Nagaev, S. V. (1971). Probability inequalities for sums of independent random variables. *Theory of Probability & Its Applications*, 16(4):643–660.

- Guédon, O., Litvak, A. E., Pajor, A., and Tomczak-Jaegermann, N. (2017). On the interval of fluctuation of the singular values of random matrices. *Journal of the European Mathematical Society*, 19(5).
- Han, Q. (2022). Exact spectral norm error of sample covariance. arXiv preprint arXiv:2207.13594.
- Horváth, L. and Kokoszka, P. (2012). *Inference for functional data with applications*. Springer Series in Statistics. Springer, New York.
- Hsing, T. and Eubank, R. (2015). Theoretical foundations of functional data analysis, with an introduction to linear operators. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester.
- Jirak, M. and Wahl, M. (2020). Perturbation bounds for eigenspaces under a relative gap condition. Proceedings of the American Mathematical Society, 148(2):479–494.
- Jirak, M. and Wahl, M. (2023). Relative perturbation bounds with applications to empirical covariance operators. Adv. Math., 412:Paper No. 108808, 59.
- Jirak, M. and Wahl, M. (2024). Quantitative limit theorems and bootstrap approximations for empirical spectral projectors. *Probability Theory and Related Fields*.
- Johnson, W. B., Schechtman, G., and Zinn, J. (1985). Best constants in moment inequalities for linear combinations of independent and exchangeable random variables. *The Annals of Probability*, pages 234–253.
- Junge, M. and Zeng, Q. (2013). Noncommutative Bennett and Rosenthal inequalities. The Annals of Probability, 41(6):4287–4316.
- Kannan, R., Lovász, L., and Simonovits, M. (1997). Random walks and an $O^*(n^5)$ volume algorithm for convex bodies. *Random Structures & Algorithms*, 11(1):1–50.
- Klochkov, Y. and Zhivotovskiy, N. (2020). Uniform Hanson-Wright type concentration inequalities for unbounded entries via the entropy method. *Electronic Journal* of *Probability*, 25(22):1–30.
- Koltchinskii, V. (2011). Von Neumann entropy penalization and low-rank matrix estimation. *The Annals of Statistics*, 39(6):2936–2973.
- Koltchinskii, V. and Lounici, K. (2017). Concentration inequalities and moment bounds for sample covariance operators. *Bernoulli*, 23(1):110–133.
- Ledoux, M. and Talagrand, M. (2013). Probability in Banach Spaces: isoperimetry and processes. Springer Science & Business Media.
- Mendelson, S. and Paouris, G. (2014). On the singular values of random matrices. Journal of the European Mathematical Society, 16(4).
- Minsker, S. (2017). On some extensions of Bernstein's inequality for self-adjoint operators. *Statistics & Probability Letters*, 127:111–119.
- Nagaev, S. V. (1979). Large deviations of sums of independent random variables. *The Annals of Probability*, pages 745–789.

- Oliveira, R. I. (2010). Sums of random Hermitian matrices and an inequality by rudelson. *Electronic Communications in Probability*, 15:203–212.
- Pinelis, I. and Utev, S. (1985). Estimates of the moments of sums of independent random variables. *Theory of Probability & Its Applications*, 29(3):574–577.
- Rio, E. (2009). Moment inequalities for sums of dependent random variables under projective conditions. *Journal of Theoretical Probability*, 22(1):146–163.
- Rio, E. (2017). About the constants in the Fuk-Nagaev inequalities. *Electronic Communications in Probability*, 22(28):12p.
- Rudelson, M. (1999). Random vectors in the isotropic position. Journal of Functional Analysis, 164(1):60–72.
- Rudelson, M. and Vershynin, R. (2007). Sampling from large matrices: An approach through geometric functional analysis. *Journal of the ACM*, 54(4).
- Srivastava, N. and Vershynin, R. (2013). Covariance estimation for distributions with $2 + \varepsilon$ moments. The Annals of Probability, 41(5):3081–3111.
- Talagrand, M. (1996). New concentration inequalities in product spaces. Inventiones mathematicae, 126(3):505–563.
- Talagrand, M. (2014). Upper and lower bounds for stochastic processes, volume 60. Springer.
- Tikhomirov, K. (2018). Sample covariance matrices of heavy-tailed distributions. International Mathematics Research Notices, 2018(20):6254–6289.
- Tropp, J. A. (2008). Norms of random submatrices and sparse approximation. Comptes Rendus. Mathématique, 346(23-24):1271–1274.
- Tropp, J. A. (2012). User-friendly tail bounds for sums of random matrices. Foundations of computational mathematics, 12:389–434.
- Tropp, J. A. (2015). An introduction to matrix concentration inequalities. Foundations and Trends (R) in Machine Learning, 8(1-2):1–230.
- van der Vaart, A. W. and Wellner, J. A. (1996). Weak convergence and empirical processes. Springer Series in Statistics. Springer-Verlag, New York.
- Vershynin, R. (2011). Approximating the moments of marginals of high-dimensional distributions. The Annals of Probability, 39(4):1591–1606.
- Vershynin, R. (2012). How close is the sample covariance matrix to the actual covariance matrix? *Journal of Theoretical Probability*, 25(3):655–686.
- Wainwright, M. J. (2019). High-dimensional statistics: A non-asymptotic viewpoint, volume 48. Cambridge university press.
- Wei, X. and Minsker, S. (2017). Estimation of the covariance structure of heavytailed distributions. Advances in neural information processing systems, 30.
- Yu, Y., Wang, T., and Samworth, R. J. (2015). A useful variant of the Davis-Kahan theorem for statisticians. *Biometrika*, 102(2):315–323.

28 MORITZ JIRAK, STANISLAV MINSKER, YIQIU SHEN, AND MARTIN WAHL

UNIVERSITÄT WIEN, AUSTRIA Email address: moritz.jirak@univie.ac.at

UNIVERSITY OF SOUTHERN CALIFORNIA Email address: minsker@usc.edu

UNIVERSITY OF SOUTHERN CALIFORNIA Email address: yiqiushe@usc.edu

UNIVERSITÄT BIELEFELD, GERMANY Email address: martin.wahl@math.uni-bielefeld.de