

Series over Bessel functions as series in terms of Riemann's zeta function

Slobodan B. Tričković¹, Miomir S. Stanković²

¹University of Niš, Department of Mathematics,
ORCID: 0000-0002-3766-8579, e-mail: sbt@junis.ni.ac.rs

²Mathematical Institute of the Serbian Academy of
Sciences and Arts, Belgrade, Serbia

July 19, 2024

Abstract

Relying on the Hurwitz formula, we find sums of the series over sine and cosine functions through the Hurwitz zeta function. Using another summation formula for these trigonometric series, we find finite sums of some series over the Riemann zeta function.

Keywords: Dirichlet Riemann's zeta function, Hurwitz's zeta function, Gamma function, Bessel functions, spherical Bessel functions.

Subject classification: 11M35; 33B15; 33E20.

1 Main results

Apostol [2] derived Hurwitz's formula

$$\zeta(1-s, a) = \frac{\Gamma(s)}{(2\pi)^s} \left(e^{-i\pi s/2} \sum_{n=1}^{\infty} \frac{e^{2i\pi na}}{n^s} + e^{i\pi s/2} \sum_{n=1}^{\infty} \frac{e^{-2i\pi na}}{n^s} \right),$$

where $\zeta(1-s, a)$ is the Hurwitz zeta function with $0 < a \leq 1$. We can write it in the equivalent form

$$\zeta(1-s, a) = \frac{2\Gamma(s)}{(2\pi)^s} \sum_{n=1}^{\infty} \frac{1}{n^s} \cos\left(\frac{\pi}{2}s - 2n\pi a\right), \quad (1.1)$$

which enables us to find the limiting value

$$\lim_{s \rightarrow 2m-1} (\zeta(1-s, a) + \zeta(1-s, 1-a)) = \lim_{s \rightarrow 2m-1} \left(\frac{4\Gamma(s)}{(2\pi)^s} \sum_{n=1}^{\infty} \frac{\cos \frac{\pi s}{2} \cos 2n\pi a}{n^s} \right) = 0,$$

where $m \in \mathbb{N}$.

Theorem 1.1 For $\operatorname{Re} s > 1$ and $0 < a \leq 1$ there hold the following formulas

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\sin(nx+y)}{n^s} &= \frac{(2\pi)^s}{2\Gamma(s)\sin\pi s} \left(\cos\left(y - \frac{\pi s}{2}\right) \zeta\left(1-s, \frac{x}{2\pi}\right) \right. \\ &\quad \left. - \cos\left(y + \frac{\pi s}{2}\right) \zeta\left(1-s, 1 - \frac{x}{2\pi}\right) \right), \\ \sum_{n=1}^{\infty} \frac{\cos(nx+y)}{n^s} &= \frac{(2\pi)^s}{2\Gamma(s)\sin\pi s} \left(\sin\left(y + \frac{\pi s}{2}\right) \zeta\left(1-s, 1 - \frac{x}{2\pi}\right) \right. \\ &\quad \left. - \sin\left(y - \frac{\pi s}{2}\right) \zeta\left(1-s, \frac{x}{2\pi}\right) \right), \end{aligned} \quad (1.2)$$

where $y \in \mathbb{R}$.

Proof. We replace a first with $1 - x/2\pi$, then with $x/2\pi$. Further, we multiply the first equality by $e^{i(y+\pi s/2)}$, and the latter by $e^{i(y-\pi s/2)}$. After subtracting, we obtain

$$\begin{aligned} &e^{i(y+\pi s/2)} \zeta\left(1-s, 1 - \frac{x}{2\pi}\right) - e^{i(y-\pi s/2)} \zeta\left(1-s, \frac{x}{2\pi}\right) \\ &= \frac{\Gamma(s)}{(2\pi)^s} (e^{i\pi s} - e^{-i\pi s}) \sum_{n=1}^{\infty} \frac{e^{i(nx+y)}}{n^s} \\ &= \frac{2\Gamma(s)\sin\pi s}{(2\pi)^s} \left(i \sum_{n=1}^{\infty} \frac{\cos(nx+y)}{n^s} - \sum_{n=1}^{\infty} \frac{\sin(nx+y)}{n^s} \right). \end{aligned}$$

Hence, comparing the corresponding real and imaginary parts, we arrive at (1.2). \square

For $y = 0$, we obtain

$$\sum_{n=1}^{\infty} \frac{\cos nx}{n^s} = \frac{(2\pi)^s}{4\Gamma(s)} \frac{\zeta\left(1-s, 1 - \frac{x}{2\pi}\right) + \zeta\left(1-s, \frac{x}{2\pi}\right)}{\cos\frac{\pi s}{2}}.$$

Thus, we evaluate

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\cos nx}{n^{2m-1}} &= \frac{(2\pi)^{2m-1}}{4(2m-2)!} \lim_{s \rightarrow 2m-1} \frac{\zeta\left(1-s, 1 - \frac{x}{2\pi}\right) + \zeta\left(1-s, \frac{x}{2\pi}\right)}{\cos\frac{\pi s}{2}} \\ &= \frac{(-1)^{m-1}(2\pi)^{2m-2}}{(2m-2)!} \left(\zeta'\left(2-2m, 1 - \frac{x}{2\pi}\right) + \zeta'\left(2-2m, \frac{x}{2\pi}\right) \right). \end{aligned} \quad (1.3)$$

We shall use (1.3) to obtain a finite sum of the series over the zeta functions.

Theorem 1.2 For $m \in \mathbb{N}$, there holds

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{\zeta(2k)}{(2k)_{2m-1}} \left(\frac{x}{2\pi}\right)^{2k} &= \frac{\log x - H_{2m-2}}{2(2m-2)!} + \frac{\left(\frac{2\pi}{x}\right)^{2m-2}}{2(2m-2)!} \left(\zeta'\left(2-2m, 1 - \frac{x}{2\pi}\right) \right. \\ &\quad \left. + \zeta'\left(2-2m, \frac{x}{2\pi}\right) \right) + (-1)^m \sum_{k=0}^{m-2} \frac{(-1)^k x^{2k-2m+2} \zeta(2m-2k-1)}{2(2k)!}. \end{aligned} \quad (1.4)$$

Proof. Taking account of [7]

$$\sum_{n=1}^{\infty} \frac{\cos nx}{n^s} = \frac{\pi x^{s-1}}{2\Gamma(s) \cos \frac{\pi s}{2}} + \sum_{k=0}^{\infty} \frac{(-1)^k \zeta(s-2k)}{(2k)!} x^{2k},$$

after letting $s \rightarrow 2m-1$, it follows

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\cos nx}{n^{2m-1}} &= \sum_{k=0}^{m-2} \frac{(-1)^k \zeta(2m-2k-2)}{(2k)!} x^{2k} + \lim_{s \rightarrow 2m-1} \left(\frac{\pi x^{s-1}}{2\Gamma(s) \cos \frac{\pi s}{2}} \right. \\ &\quad \left. + \frac{(-1)^{m-1} \zeta(s-2m+2)}{(2m-2)!} x^{2m-2} \right) + \sum_{k=m}^{\infty} \frac{(-1)^k \zeta(2m-2k-1)}{(2k)!} x^{2k}. \end{aligned} \quad (1.5)$$

By bringing the fractions in brackets to the same denominator and applying L'Hôpital's rule, we determine the above limiting value

$$\begin{aligned} \lim_{s \rightarrow 2m-1} \left(\frac{\pi x^{s-1}}{2\Gamma(s) \cos \frac{\pi s}{2}} + \frac{(-1)^{m-1} \zeta(s-2m+2)}{(2m-2)!} x^{2m-2} \right) \\ = \frac{(-1)^m x^{2m-2}}{(2m-2)!} (\log x - H_{2m-2}). \end{aligned}$$

As for the remainder in (1.5), we introduce a new running index by $k = m+j-1$, then reverting to k and recalling the relation

$$\zeta(1-2n) = (-1)^n \frac{2(2n-1)! \zeta(2n)}{(2\pi)^{2n}} = (-1)^n \frac{2\Gamma(2n) \zeta(2n)}{(2\pi)^{2n}}, \quad (1.6)$$

we come to the formula

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{\cos nx}{n^{2m-1}} &= \frac{(-1)^m x^{2m-2}}{(2m-2)!} (\log x - H_{2m-2}) + \sum_{k=0}^{m-2} \frac{(-1)^k x^{2k} \zeta(2m-2k-1)}{(2k)!} \\ &\quad + 2(-1)^{m-1} x^{2m-2} \sum_{k=1}^{\infty} \frac{\zeta(2k)}{(2k)_{2m-1}} \left(\frac{x}{2\pi} \right)^{2k}. \end{aligned}$$

Using (1.3), we obtain (1.4). \square

We are dealing now with a more general series.

Theorem 1.3 *For $m, p \in \mathbb{N}$ and $0 < x < 2\pi$, there holds*

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(m+n)_p}{(2n)_{2p+2m}} \zeta(2n) \left(\frac{x}{2\pi} \right)^{2n} \\ = (-1)^{m-1} 4x^{p+2m-2} \sum_{k=1}^p C_k \sum_{j=0}^{2k-1} (-1)^j \prod_{i=1}^j (2k-i) \sum_{n=1}^{\infty} \frac{\zeta(2n) \left(\frac{x}{2\pi} \right)^{2n}}{(2n)_{2m+j+1}}. \end{aligned} \quad (1.7)$$

Proof. We consider a decomposition of the fraction of two Pochhammer symbols in (1.7)

$$\begin{aligned} \frac{(m+n)_p}{(2n)_{2p+2m}} &= \frac{(m+n) \cdots (m+n+p-1)}{(2n)_{2m}(2n+2m) \cdots (2n+2m+2p-1)} \\ &= \frac{1}{2^p(2n)_{2m}(2n+2m+1)(2n+2m+3) \cdots (2n+2m+2p-1)}. \end{aligned}$$

By applying Heaviside's method, in the next step, we make a decomposition of the rational function

$$\frac{P(n)}{Q(n)} = \frac{1}{(2n+2m+1)(2n+2m+3) \cdots (2n+2m+2p-1)},$$

with $P(n) = 1$, $Q(n) = (2n+2m+1)(2n+2m+3) \cdots (2n+2m+2p-1)$. Thus, we have

$$\frac{P(n)}{Q(n)} = \frac{C_1}{n-a_1} + \frac{C_2}{n-a_2} + \cdots + \frac{C_p}{n-a_p},$$

where the constants C_k , $k = 1, \dots, p$, are to be determined, and $Q(a_k) = 0$. So, we have

$$C_k = \frac{P(a_k)}{Q'(a_k)}, \quad a_k = -(2m+2k-1)/2, \quad k = 1, \dots, p,$$

which means that the left-hand side series in (1.7) we can express as follows

$$\sum_{n=1}^{\infty} \frac{(m+n)_p}{(2n)_{2p+2m}} \zeta(2n) \left(\frac{x}{2\pi}\right)^{2n} = \sum_{k=1}^p \frac{C_k}{2^{p-1}} \sum_{n=1}^{\infty} \frac{\zeta(2n) \left(\frac{x}{2\pi}\right)^{2n}}{(2n)_{2m}(2n+2m+2k-1)}. \quad (1.8)$$

The whole idea consists in finding constants c_1, \dots, c_{2k} in the representation

$$\frac{1}{2n+2m+2k-1} = \frac{c_1}{2n+2m} + \frac{c_2}{(2n+2m)_2} + \cdots + \frac{c_{2k}}{(2n+2m)_{2k}},$$

where $k = 1, \dots, p$. We multiply the numerator and denominator of the left-hand side fraction by the missing factors between $(2n+2m+2k-1)$ and $(2n+2m)$, including the latter. Also, we bring the right-hand side fractions to the same denominator. Thus, we obtain the equality of the numerators

$$\begin{aligned} (2n+2m) \cdots (2n+2m+2k-2) &= c_1(2n+2m+1) \cdots (2n+2m+2k-1) \\ &+ c_2(2n+2m+2) \cdots (2n+2m+2k-1) + \cdots + c_{2k}, \quad k = 1, \dots, p. \end{aligned}$$

The highest power on the left-hand side and the product at c_1 is $(2n)^{2k-1}$. There follows $c_1 = 1$. After rearrangements, we have

$$\begin{aligned} &- (2k-1)(2n+2m+1) \cdots (2n+2m+2k-2) \\ &= c_2(2n+2m+2) \cdots (2n+2m+2k-1) + c_3(2n+2m+3) \cdots \times \quad (1.9) \\ &\cdots (2n+2m+2k-1) + \cdots + c_{2k-1}(2n+2m+2k-1) + c_{2k}. \end{aligned}$$

Regarding both sides as polynomial functions in n of the degree $2k - 2$, we take the $(2k - 2)$ th derivative. As a result, we have

$$-(2k - 1)2^{2k-2}(2k - 2)! = c_2 2^{2k-2}(2k - 2)!.$$

Hence, we find $c_2 = -(2k - 1)$. Replacing this value in (1.9), we can determine c_3 similarly

$$\begin{aligned} & (2k - 1)(2k - 2)(2n + 2m + 2) \cdots (2n + 2m + 2k - 2) \\ &= c_3(2n + 2m + 3) \cdots (2n + 2m + 2k - 1) + \\ & c_4(2n + 2m + 4) \cdots (2n + 2m + 2k - 1) + \cdots + c_{2k-1}(2n + 2m + 2k - 1) + c_{2k}. \end{aligned}$$

Now, the polynomials on both sides are of the degree $2k - 3$, and we take the $(2k - 3)$ th derivative, and obtain

$$(2k - 1)(2k - 2)2^{2k-3}(2k - 3)! = c_3 2^{2k-3}(2k - 3)!,$$

which implies $c_3 = (2k - 1)(2k - 2)$. By repeating this procedure, we arrive at the relation

$$\frac{1}{(2n)_{2m}(2n + 2m + 2k - 1)} = \sum_{j=0}^{2k-1} \frac{(-1)^j}{(2n)_{2m+j+1}} \prod_{i=1}^j (2k - i).$$

Together with (1.8), this gives rise to (1.7). \square

Thus, the procedure to evaluate the left-hand side sum of (1.7) reduces to a multiple application of (1.4).

2 Applications to some series over Bessel functions

Bessel functions $J_\nu(x)$ defined by the Swiss mathematician Daniel Bernoulli, then generalized and developed by Friedrich Bessel while studying the dynamics of gravitational systems in the second decade of the 19th century, are canonical solutions of homogeneous Bessel's differential equation

$$x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + (x^2 - \nu^2)y = 0 \quad (2.1)$$

for an arbitrary complex number ν . The particular solution

$$J_\nu(x) = \sum_{m=0}^{\infty} \frac{(-1)^m \left(\frac{x}{2}\right)^{2m+\nu}}{m! \Gamma(m + \nu + 1)}, \quad (2.2)$$

is called Bessel's function of the first kind order ν [5]. Apart from that, it can be represented by Poisson's integral

$$J_\nu(z) = \frac{2 \left(\frac{z}{2}\right)^\nu}{\Gamma\left(\frac{1}{2}\right) \Gamma\left(\nu + \frac{1}{2}\right)} \int_0^{\pi/2} \sin^{2\nu} \theta \cos(z \cos \theta) d\theta, \quad \operatorname{Re} \nu > -\frac{1}{2}, \quad (2.3)$$

proved by Poisson [4] and Lommel [3] to be a solution of homogenous Bessel's differential equation for $2\nu \in \mathbb{N}_0$, relying on the summation of trigonometric series, in [6], we derived summation formulas for the series

$$\sum_{n=1}^{\infty} \frac{J_{\nu}(nx)}{n^{\alpha}} = \frac{\pi(\frac{x}{2})^{\alpha-1} \sec \frac{\pi(\alpha-\nu)}{2}}{2\Gamma(\frac{\alpha-\nu+1}{2})\Gamma(\frac{\alpha+\nu+1}{2})} + \sum_{k=0}^{\infty} \frac{(-1)^k (\frac{x}{2})^{\nu+2k} \zeta(\alpha-\nu-2k)}{k! \Gamma(\nu+k+1)}, \quad (2.4)$$

where $\alpha > 0$, $\alpha > \nu > -\frac{1}{2}$, $0 < x < 2\pi$.

Providing $\alpha - \nu = 2m$, $m \in \mathbb{N}$, the right-hand side series in (2.4) truncates because Riemann's zeta function equals zero if the argument is a negative even integer. So, setting $\alpha = \nu + 2m$ in (2.4) brings the series in closed form

$$\sum_{n=1}^{\infty} \frac{J_{\nu}(nx)}{n^{\nu+2m}} = \frac{m!(-1)^m x^{\nu+2m-1} \sqrt{\pi}}{2^{\nu}(2m)!\Gamma(\nu+m+\frac{1}{2})} + \sum_{k=0}^m \frac{(-1)^k \zeta(2m-2k)(\frac{x}{2})^{\nu+2k}}{k! \Gamma(\nu+k+1)}, \quad (2.5)$$

where $0 < x < 2\pi$, $\operatorname{Re} \nu > -2m - \frac{1}{2}$.

In the case $\alpha - \nu = 2m - 1$, $m \in \mathbb{N}$, we cannot immediately place this in the right-hand side of the relation (2.4) because one encounter singularities, i.e. in $\sec \frac{\pi(\alpha-\nu)}{2}$ within the numerator of the first term and in the member of the right-hand series for the index $k = m - 1$.

Theorem 2.1 For $\alpha = \nu + 2m - 1$, $m \in \mathbb{N}$, there holds

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{J_{\nu}(nx)}{n^{\nu+2m-1}} &= \frac{(-1)^{m-1} (\frac{x}{2})^{\nu+2m-2}}{2\Gamma(m)\Gamma(\nu+m)} \left(H_{m-1} + H_{\nu+m-1} - 2 \ln \frac{x}{2} \right) \\ &+ \sum_{k=0}^{m-2} \frac{(-1)^k \zeta(2m-2k-1)(\frac{x}{2})^{\nu+2k}}{k! \Gamma(\nu+k+1)} + \sum_{k=1}^{\infty} \frac{\Gamma(2k)\zeta(2k)(\frac{x}{4\pi})^{2k}}{\Gamma(m+k)\Gamma(\nu+m+k)}. \end{aligned} \quad (2.6)$$

Proof. Because of what is said above, it is necessary to take the limiting value in (2.4) when $\alpha \rightarrow \nu + 2m - 1$

$$\lim_{\alpha \rightarrow \nu+2m-1} \left(\frac{\pi(\frac{x}{2})^{\alpha-1} \sec \frac{\pi(\alpha-\nu)}{2}}{\Gamma(\frac{\alpha-\nu+1}{2})\Gamma(\frac{\alpha+\nu+1}{2})} + \sum_{k=0}^{m-1} \frac{(-1)^k \zeta(\alpha-\nu-2k)(\frac{x}{2})^{\nu+2k}}{k! \Gamma(\nu+k+1)} \right).$$

For $k = 0, 1, \dots, m-2$ all the terms have no singularities if $\alpha = \nu + 2m - 1$, so it suffices to deal only with the term for $k = m - 1$, then we find

$$\begin{aligned} \lim_{\alpha \rightarrow \nu+2m-1} &\left(\frac{\pi(\frac{x}{2})^{\alpha-1} \sec \frac{\pi(\alpha-\nu)}{2}}{\Gamma(\frac{\alpha-\nu+1}{2})\Gamma(\frac{\alpha+\nu+1}{2})} + \frac{(-1)^{m-1} (\frac{x}{2})^{\nu+2m-2} \zeta(\alpha-\nu-2m+2)}{(m-1)! \Gamma(\nu+m)} \right) \\ &= \frac{(-1)^{m-1} (\frac{x}{2})^{\nu+2m-2}}{2\Gamma(m)\Gamma(\nu+m)} \left(H_{m-1} + H_{\nu+m-1} - 2 \log \frac{x}{2} \right), \end{aligned}$$

where γ is Euler-Mascheroni's constant. Thus, we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{J_{\nu}(nx)}{n^{\nu+2m-1}} &= \frac{(-1)^{m-1} \left(\frac{x}{2}\right)^{\nu+2m-2}}{2\Gamma(m)\Gamma(\nu+m)} \left(H_{m-1} + H_{\nu+m-1} - 2\log \frac{x}{2}\right) \\ &+ \sum_{k=0}^{m-2} \frac{(-1)^k \zeta(2m-2k-1) \left(\frac{x}{2}\right)^{\nu+2k}}{k!\Gamma(\nu+k+1)} + \sum_{k=m}^{\infty} \frac{(-1)^k \zeta(2m-2k-1) \left(\frac{x}{2}\right)^{\nu+2k}}{k!\Gamma(\nu+k+1)}. \end{aligned}$$

As for the remainder, we transform it by shifting the index, so we introduce the substitution $k = m + j - 1$, then revert to k instead of using j , i.e.

$$\begin{aligned} \sum_{k=m}^{\infty} \frac{(-1)^k \zeta(2m-2k-1) \left(\frac{x}{2}\right)^{\nu+2k}}{k!\Gamma(\nu+k+1)} \\ = (-1)^{m-1} \left(\frac{x}{2}\right)^{\nu+2m-2} \sum_{k=1}^{\infty} \frac{(-1)^k \zeta(1-2k) \left(\frac{x}{2}\right)^{2k}}{\Gamma(m+k)\Gamma(\nu+m+k)}. \end{aligned}$$

Using (1.6) substituting there k for n , the last series becomes

$$2(-1)^{m-1} \left(\frac{x}{2}\right)^{\nu+2m-2} \sum_{k=1}^{\infty} \frac{\Gamma(2k)\zeta(2k) \left(\frac{x}{4\pi}\right)^{2k}}{\Gamma(m+k)\Gamma(\nu+m+k)},$$

whereby we arrive at (2.6). \square

In a special case, setting $\nu = \frac{1}{2}$ in (2.6), we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{J_{1/2}(nx)}{n^{2m-1/2}} &= \frac{(-1)^{m-1} \left(\frac{x}{2}\right)^{2m-3/2}}{2\Gamma(m)\Gamma(\frac{1}{2}+m)} \left(H_{m-1} + H_{m-1/2} - 2\log \frac{x}{2}\right) \quad (2.7) \\ &+ \sqrt{\frac{x}{2}} \sum_{k=0}^{m-2} \frac{(-1)^k \zeta(2m-2k-1) \left(\frac{x}{2}\right)^{2k}}{k!\Gamma(\frac{1}{2}+k+1)} + \sum_{k=1}^{\infty} \frac{\Gamma(2k)\zeta(2k) \left(\frac{x}{4\pi}\right)^{2k}}{\Gamma(m+k)\Gamma(\frac{1}{2}+m+k)}. \end{aligned}$$

After some modifications and referring to (1.4), we bring the right-hand series in (2.7) to the closed form, i.e.

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{\Gamma(2k)\zeta(2k) \left(\frac{x}{4\pi}\right)^{2k}}{\Gamma(m+k)\Gamma(\frac{1}{2}+m+k)} &= \frac{(-1)^{m-1}}{\sqrt{2\pi}} 8x^{2m-3/2} \sum_{k=1}^{\infty} \frac{\zeta(2k)}{(2k)_{2m}} \left(\frac{x}{2\pi}\right)^{2k} \\ &= \frac{(-1)^m 4x^{2m-3/2}}{(2m-1)!\sqrt{2\pi}} (H_{2m-1} - \log 2\pi) - \frac{4}{\sqrt{2x\pi}} \sum_{k=0}^{m-2} \frac{(-1)^k \zeta(2m-2k-1) x^{2k+1}}{(2k+1)!} \\ &+ \frac{(-1)^m 4(2\pi)^{2m-3/2}}{(2m-1)!\sqrt{x}} \left(\zeta'\left(1-2m, 1 - \frac{x}{2\pi}\right) - \zeta'\left(1-2m, 1 + \frac{x}{2\pi}\right) \right). \quad (2.8) \end{aligned}$$

By differentiating the basic property of Hurwitz's function

$$\zeta(s, a) = a^{-s} + \zeta(s, a+1)$$

with respect to s and putting then $a = \frac{x}{2\pi}$ and $s = 1 - 2m$, for the last term of (2.8), we find

$$-\zeta'\left(1 - 2m, 1 + \frac{x}{2\pi}\right) = -\left(\frac{x}{2\pi}\right)^{2m-1} \log \frac{x}{2\pi} - \zeta'\left(1 - 2m, \frac{x}{2\pi}\right).$$

We can rewrite now (2.8) and use it to give the left-hand side series of (2.7) the closed form

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{J_{1/2}(nx)}{n^{2m-1/2}} &= \frac{(-1)^m 4(2\pi)^{2m-1}}{(2m-1)! \sqrt{2\pi x}} \left(\frac{1}{2} (H_{2m-1} - \log x) \left(\frac{x}{2\pi}\right)^{2m-1} \right. \\ &\left. + \zeta'\left(1 - 2m, 1 - \frac{x}{2\pi}\right) - \zeta'\left(1 - 2m, \frac{x}{2\pi}\right) \right) + \sqrt{\frac{2x}{\pi}} \sum_{k=0}^{m-2} \frac{(-x^2)^k \zeta(2m-2k-1)}{(2k+1)!}. \end{aligned} \quad (2.9)$$

2.1 Series over spherical Bessel functions

We can generalize (2.9) by considering spherical Bessel functions $j_k(z)$, linked to Bessel functions of the first kind by the equality

$$j_p(x) = \sqrt{\frac{\pi}{2x}} J_{p+\frac{1}{2}}(x), \quad p \in \mathbb{N}_0.$$

So, multiplying both sides of (2.4) by $\sqrt{\frac{\pi}{2x}}$ and setting $\nu = p + \frac{1}{2}$, $p \in \mathbb{N}_0$, we have

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{j_p(nx)}{n^\alpha} &= \frac{\pi \sqrt{\pi} \left(\frac{x}{2}\right)^{\alpha-\frac{3}{2}}}{2 \Gamma\left(\frac{\alpha-p}{2} + \frac{1}{4}\right) \Gamma\left(\frac{\alpha+p}{2} + \frac{3}{4}\right) \cos\left(\frac{\pi(\alpha-p)}{2} - \frac{\pi}{4}\right)} \\ &\quad + \frac{\sqrt{\pi}}{2} \sum_{k=0}^{\infty} \frac{(-1)^k \zeta\left(\alpha - p - \frac{1}{2} - 2k\right) \left(\frac{x}{2}\right)^{p+2k}}{k! \Gamma\left(p + \frac{1}{2} + k + 1\right)}. \end{aligned} \quad (2.10)$$

Here, we are referring to Euler's reflection formula

$$\Gamma(1-s)\Gamma(s) = \frac{\pi}{\sin \pi s}, \quad s \notin \mathbb{Z},$$

with $s = \frac{\alpha-p}{2} + \frac{1}{4}$, and obtain

$$\Gamma\left(\frac{3}{4} - \frac{\alpha-p}{2}\right) \Gamma\left(\frac{\alpha-p}{2} + \frac{1}{4}\right) = \frac{\pi}{\cos\left(\frac{\pi}{4} - \frac{\pi(\alpha-p)}{2}\right)} = \frac{\pi}{\cos\left(\frac{\pi(\alpha-p)}{2} - \frac{\pi}{4}\right)}.$$

Replacing this in the first term of (2.10), and using the property

$$\Gamma\left(n + \frac{1}{2}\right) = \frac{\sqrt{\pi} \Gamma(2n)}{2^{2n-1} \Gamma(n)} = \frac{(2n)!}{2^{2n} n!} \sqrt{\pi},$$

we obtain the summation formula for the series over spherical Bessel functions

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(s)^{n-1} j_p((an-b)x)}{(an-b)^\alpha} &= \frac{c\sqrt{\pi}x^{\alpha-\frac{3}{2}}\Gamma(\frac{3}{4}-\frac{\alpha-p}{2})}{2^{\alpha+\frac{1}{2}}\Gamma(\frac{3}{4}+\frac{\alpha+p}{2})} \\ &+ (2x)^p \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(p+k+1) F(\alpha-p-\frac{1}{2}-2k) x^{2k}}{k! \Gamma(2p+2k+2)}. \end{aligned}$$

Now, for $a = 1, b = 0, s = 1, c = 1, F = \zeta$, we first evaluate

$$\begin{aligned} \lim_{\alpha \rightarrow p+2m-1/2} \left(\frac{\sqrt{\pi}x^{\alpha-\frac{3}{2}}\Gamma(\frac{3}{4}-\frac{\alpha-p}{2})}{2^{\alpha+\frac{1}{2}}\Gamma(\frac{3}{4}+\frac{\alpha+p}{2})} \right. \\ \left. + 2^p \frac{(-1)^{m-1} \Gamma(p+m) \zeta(\alpha-p-2m+\frac{3}{2}) x^{2m+p-2}}{(m-1)! \Gamma(2p+2m)} \right). \end{aligned}$$

By bringing to the same denominator and applying the L'Hôpital rule, after the limiting process, we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{j_p(nx)}{n^{p+2m-1/2}} &= \frac{(-1)^{m-1} (\frac{x}{2})^{p+2m-\frac{3}{2}} \sqrt{\frac{\pi}{8x}}}{\Gamma(m)\Gamma(p+\frac{1}{2}+m)} \left(H_{m-1} + H_{p+m-\frac{1}{2}} - 2 \log \frac{x}{2} \right) \\ &+ (2x)^p \sum_{k=0}^{m-2} \frac{(-1)^k \Gamma(p+k+1) \zeta(2m-1-2k) x^{2k}}{k! \Gamma(2p+2k+2)} \\ &+ (-1)^{m-1} 2^{p+1} x^{p+2m-2} \sum_{n=1}^{\infty} \frac{(m+n)_p}{(2n)_{2p+2m}} \zeta(2n) \left(\frac{x}{2\pi} \right)^{2n}. \end{aligned}$$

Referring to (1.7), we evaluate

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{j_p(nx)}{n^{p+2m-1/2}} &= \frac{(-1)^{m-1} (\frac{x}{2})^{p+2m-\frac{3}{2}} \sqrt{\frac{\pi}{8x}}}{\Gamma(m)\Gamma(p+\frac{1}{2}+m)} \left(H_{m-1} + H_{p+m-\frac{1}{2}} - 2 \log \frac{x}{2} \right) \\ &+ (2x)^p \sum_{k=0}^{m-2} \frac{(-1)^k \Gamma(p+k+1) \zeta(2m-1-2k) x^{2k}}{k! \Gamma(2p+2k+2)} \\ &+ (-1)^{m-1} 4x^{p+2m-2} \sum_{k=1}^p C_k \sum_{j=0}^{2k-1} (-1)^j \prod_{i=1}^j (2k-i) \sum_{n=1}^{\infty} \frac{\zeta(2n) (\frac{x}{2\pi})^{2n}}{(2n)_{2m+j+1}}. \quad (2.11) \end{aligned}$$

We apply (1.4) to evaluate the sum of the series on the right side of (2.11).

Example 2.1 For $m = 3$ and $p = 2$, we apply (2.11) and (1.4) obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{j_2(nx)}{n^{15/2}} &= \frac{x^6 \left(\frac{3197}{1260} - \log x \right)}{15120} - \frac{x^4 \zeta(3)}{420} + \frac{x^2 \zeta(5)}{30} + \frac{4\pi^7 \zeta' \left(-7, \frac{x}{2\pi} \right)}{315x} \\ &\quad - \frac{4\pi^7 \zeta' \left(-7, 1 - \frac{x}{2\pi} \right)}{315x} - \frac{\pi^8 \zeta' \left(-8, \frac{x}{2\pi} \right)}{105x^2} - \frac{\pi^8 \zeta' \left(-8, 1 - \frac{x}{2\pi} \right)}{105x^2} \\ &\quad + \frac{2\pi^9 \zeta' \left(-9, \frac{x}{2\pi} \right)}{945x^3} - \frac{2\pi^9 \zeta' \left(-9, 1 - \frac{x}{2\pi} \right)}{945x^3}. \end{aligned}$$

References

- [1] M. Abramowitz, A. Stegun, *Handbook of mathematical functions*, Cambridge University Press, New York, 2010.
- [2] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag New York, 1976.
- [3] E. C. J. von Lommel, *Studien über die Besselschen Funktionen*, Leipzig: Teubner 1868.
- [4] S.D. Poisson, Mémoire sur la distribution de la chaleur dans les corps solides, *Journal de l'École Polytechnique* XII (cahier 19) (1823), 249–403.
- [5] A. P. Prudnikov, Y.A. Brychkov, O.I. Marichev, *Integrals and Series*, Vol. 2, *Special Functions*, Taylor & Francis, London, 1998.
- [6] S. B. Tričković, M. V. Vidanović, M.S. Stanković: *On the summation of series in terms of Bessel functions*, *Z. Anal. Anwen.* **25**, No. 3 (2006) 393–406.
- [7] S. B. Tričković, M. S. Stanković, M.V. Vidanović, *On the summation of trigonometric series*, *Int. Trans. Spec. Func.* **19**, No 6 (2008) 441–452.
- [8] S. B. Tričković, M. S. Stanković, *On the closed form of Clausen functions*, *Int. Trans. Spec. Func.* **34**, No 6 (2023) 469–477.