

# The Dance of Odd-Diffusive Particles: A Fourier Approach

Amelie Langer,<sup>1</sup> Abhinav Sharma,<sup>1,2</sup> Ralf Metzler,<sup>3,4</sup> and Erik Kalz<sup>3,\*</sup>

<sup>1</sup>University of Augsburg, Institute of Physics, D-86159 Augsburg, Germany

<sup>2</sup>Leibniz-Institute for Polymer Research, Institute Theory of Polymers, D-01069 Dresden, Germany

<sup>3</sup>University of Potsdam, Institute of Physics and Astronomy, D-14476 Potsdam, Germany

<sup>4</sup>Asia Pacific Centre for Theoretical Physics, KR-37673 Pohang, Republic of Korea

Odd-diffusive systems are characterized by transverse responses and exhibit unconventional behaviors in interacting systems. To address the dynamical interparticle rearrangements in a minimal system, we here exactly solve the problem of two hard disk-like interacting odd-diffusing particles. We calculate the probability density function (PDF) of the interacting particles in the Fourier-Laplace domain and find that oddness rotates all modes except the zeroth, resembling a “mutual rolling” of interacting odd particles. We show that only the first Fourier mode of the PDF, the polarization, enters the calculation of the force autocorrelation function (FACF) for generic systems with central-force interactions. An analysis of the polarization as a function of time reveals that the relative rotation angle between interacting particles overshoots before relaxation, thereby rationalizing the recently observed oscillating FACF in odd-diffusive systems.

## I. INTRODUCTION

The description of dissipative systems with an inherent broken time-reversal or parity symmetry has recently attracted considerable attention [1–5]. Relevant systems can be found in various domains of statistical physics, and include Brownian particles under the effect of Lorentz force [6], skyrmionic spin structures [7] and active chiral particles [1]. The transport coefficients of these systems show a characteristic transverse response to perturbations, which is encoded in antisymmetric off-diagonal elements in transport tensors. These characteristic elements behave odd under the transformation of the underlying (broken) symmetry, which serves as the namesake for these *odd* systems. Interestingly, a transverse response does not necessarily imply an anisotropic description of the system. In fact, in two spatial dimensions, odd transport coefficients represent the most general description of an isotropic physical system [8]. In this work, we specifically consider odd-diffusive systems [1, 2], which in two spatial dimensions are characterized by a diffusion tensor of the form

$$\mathbf{D} = D_0(\mathbf{1} + \kappa\epsilon), \quad (1)$$

where  $D_0$  is the bare diffusivity with physical dimensions  $[D_0] = \text{m}^2/\text{s}$ , and  $\kappa$  is the characteristic odd-diffusion parameter with  $[\kappa] = 1$ , encoding the transverse response.  $\mathbf{1}$  is the identity tensor and  $\epsilon$  the fully antisymmetric Levi-Civita symbol in two dimensions ( $\epsilon_{xy} = -\epsilon_{yx} = 1$  and  $\epsilon_{xx} = \epsilon_{yy} = 0$ ).

While the oddness parameter enters explicitly in the diffusion tensor in Eq. (1), the mean-squared displacement of a freely diffusing particle is determined only by the symmetric part of the diffusion tensor. However, in a system of interacting particles, oddness qualitatively alters the diffusive behavior by affecting the self-diffusion [2, 9]. The self-diffusion measures the dynamic

of a tagged particle in a crowded system and explicitly accounts for interactions of particles [10]. Independent of the microscopic details of the inter-particle interactions, the self-diffusion is usually reduced [11–21]. In odd-diffusive systems, however, it was recently shown that even purely repulsive interactions can enhance the self-diffusion [2]. Via expressing the self-diffusion as a time-integral of the (interaction) force autocorrelation function (FACF) [22], these findings could further be related to the unusual microscopic particle rearrangements in odd-diffusive systems [9].

For the overdamped equilibrium system under consideration here and consistent with the reduction of the self-diffusion [22], autocorrelation functions in general decay monotonically in time [23]. However, the enhancement of self-diffusion for odd systems can occur only if the FACF switches sign, i.e., becomes non-monotonic in time. This apparent contradiction could be resolved by recognizing that the time-evolution operator in an odd-diffusive system becomes non-Hermitian when  $\kappa \neq 0$  in Eq. (1), thereby breaking the monotonicity requirements on the FACF [9]. Surprisingly, in this work, it was further observed that the FACF even oscillates in time. While this is consistent with the observed enhancement of the self-diffusion in odd systems, a physical interpretation of this phenomenon remains elusive.

In the present work, we show that the non-monotonic FACF originates in the unusual dynamics of interacting odd particles. When a pair of odd-diffusive particles interacts, their motion resembles that of two mutually rotating particles [2]. This is despite the fact that the interaction potential is central-force, i.e., it acts along the vector that connects the centers of two particles. Our approach is based on the exact analytical derivation of the propagator of two interacting odd-diffusive particles. The joint probability density function (PDF) for the two particles separates into a center-of-mass PDF and a PDF of the relative coordinate, capturing the interplay of odd diffusion and interactions. We can express the relative PDF in a Fourier series to find that only certain modes

\* erik.kalz@uni-potsdam.de

enter the averages of observables, such as in the FACF. In particular, it is the polarization mode of the relative PDF, representing the positioning of the particles with respect to each other in time, which determines the full tensorial force autocorrelation behavior. By analyzing the polarization mode in time, we understand what originates the oscillating FACF. Interacting odd-diffusive particles mutually rotate further in time before they eventually relax. Figuratively they perform a dance, reminiscent of the classical step of the Viennese waltz.

The remainder of this work is organized as follows: In Section II we set the problem of two interacting odd-diffusive particles, which is then exactly solved in Section II A. The propagator can be put into the form of a Fourier series, of which the numerical analysis of the modes is presented in Section II B. In Section II C we show that only certain Fourier modes enter in averages of observables, in particular into the FACF. Finally, in Section III we summarize and give an extensive overview of systems, which can be subsumed under the terminology of odd diffusion. In Appendix A we present the detailed solution of the problem of interacting particles and Appendix B lists the relevant integral relations.

## II. INTERACTING ODD-DIFFUSIVE PARTICLES

We study the dynamics of two odd-diffusive particles at positions  $\mathbf{x}_1$  and  $\mathbf{x}_2$  in two spatial dimensions. The particles are assumed to interact with the potential energy  $U$ , which we assume to be of the hard-disk type

$$U(\mathbf{x}_1, \mathbf{x}_2) = U(r) = \begin{cases} \infty, & r \leq 1 \\ 0, & r > 1 \end{cases}, \quad (2)$$

where  $r = |\mathbf{x}_1 - \mathbf{x}_2|/d$  is the rescaled relative distance between the particles and  $d$  is the particle diameter.

The conditional joint PDF, i.e., the propagator for the particles to be found at positions  $(\mathbf{x}_1, \mathbf{x}_2)$  at time  $t$  given that they were at positions  $(\mathbf{x}_{1,0}, \mathbf{x}_{2,0})$  at time  $t_0$ ,  $P(t) = P(\mathbf{x}_1, \mathbf{x}_2, t | \mathbf{x}_{1,0}, \mathbf{x}_{2,0}, t_0)$ , evolves according to the time-evolution equation

$$\begin{aligned} \frac{\partial}{\partial t} P(t) = & \nabla_1 \cdot [\mathbf{D} \nabla_1 + \boldsymbol{\mu} \nabla_1 U(\mathbf{x}_1, \mathbf{x}_2)] P(t) \\ & + \nabla_2 \cdot [\mathbf{D} \nabla_2 + \boldsymbol{\mu} \nabla_2 U(\mathbf{x}_1, \mathbf{x}_2)] P(t), \end{aligned} \quad (3)$$

where the initial condition is given as  $P(t = t_0) = \delta(\mathbf{x}_1 - \mathbf{x}_{1,0}) \delta(\mathbf{x}_2 - \mathbf{x}_{2,0})$ . In Eq. (3),  $\nabla_1, \nabla_2$  are the partial differential operators for the positions of the particles, and  $\mathbf{D}$  is the odd-diffusion tensor of Eq. (1).  $\boldsymbol{\mu}$  is the mobility tensor, and we assume the fluctuation-dissipation relation (FDR) to hold

$$\mathbf{D} = k_B T \boldsymbol{\mu}, \quad (4)$$

where  $k_B$  is the Boltzmann constant and  $T$  the temperature of the solvent. Note that even though Eq. (3) looks

formally equivalent to a Fokker-Planck equation for the joint PDF  $P(t)$ , strictly spoken it is not, due to the anti-symmetric (odd-diffusive) elements in the diffusion tensor [24]. However, based on the assumption of the FDR Eq. (3) resembles equilibrium dynamics with a unique steady-state solution [9, 25].

### A. Analytical solution

We rescale space with the diameter of the particle  $d$ ,  $\mathbf{x}_i \rightarrow \mathbf{x}_i/d$ , and time by the natural time-scale of diffusing the radial distance of a particle diameter  $\tau_d = d^2/(2D_0)$ ,  $t \rightarrow \tau = t/\tau_d$ . Given the radial symmetry of the interaction potential  $U(r)$ , the time-evolution equation (3) can be written in terms of a center-of-mass coordinate  $\mathbf{x}_c = (\mathbf{x}_1 + \mathbf{x}_2)/2$  and a relative coordinate  $\mathbf{x} = \mathbf{x}_1 - \mathbf{x}_2$  as

$$\begin{aligned} \frac{\partial}{\partial \tau} P(\tau) = & \frac{1}{4} \nabla_{\mathbf{x}_c}^2 P(\tau) \\ & + \nabla_{\mathbf{x}} \cdot (\mathbf{1} + \kappa \boldsymbol{\epsilon}) [\nabla_{\mathbf{x}} + \beta \nabla_{\mathbf{x}} U(r)] P(\tau), \end{aligned} \quad (5)$$

where  $\beta = 1/k_B T$  and  $\nabla_{\mathbf{x}_c}, \nabla_{\mathbf{x}}$  are the partial differential operators corresponding to the center-of-mass and relative coordinates. Note that  $(\mathbf{1} + \kappa \boldsymbol{\epsilon}) = \mathbf{D}/D_0$  represents the dimensionless odd-diffusion tensor of Eq. (1). As Eq. (5) decouples the coordinates, the propagator can be written as  $P(\mathbf{x}_1, \mathbf{x}_2, \tau | \mathbf{x}_{2,0}, \mathbf{x}_{1,0}, \tau_0) = p_c(\mathbf{x}_c, \tau | \mathbf{x}_{c,0}, \tau_0) p(\mathbf{x}, \tau | \mathbf{x}_0, \tau_0)$  and the center-of-mass problem can be solved straight forwardly in the form

$$p_c(\mathbf{x}_c, \tau | \mathbf{x}_{c,0}) = \frac{1}{\pi \tau} \exp\left(-\frac{|\mathbf{x}_c - \mathbf{x}_{c,0}|^2}{\tau}\right), \quad (6)$$

where we have set  $\tau_0 = 0$  as the underlying stochastic process is time-translational invariant [26].

The relative PDF can be put into the form of a (Cartesian) multipole expansion [27–31] which in polar coordinates  $\mathbf{x} = (r, \varphi)$ ,  $\mathbf{x}_0 = (r_0, \varphi_0)$ , is given as

$$\begin{aligned} p(\mathbf{x}, \tau | \mathbf{x}_0) = & \Theta(r - 1) \left[ \varrho(r, \tau | r_0) + \boldsymbol{\sigma}(r, \tau | r_0) \cdot \mathbf{e}(\Delta\varphi) \right. \\ & \left. + \mathbf{Q}(r, \tau | r_0) : \left( \mathbf{e}(\Delta\varphi) \otimes \mathbf{e}(\Delta\varphi) - \frac{\mathbf{1}}{2} \right) + \dots \right], \end{aligned} \quad (7)$$

where  $\mathbf{e}(\Delta\varphi) = (\cos(\Delta\varphi), \sin(\Delta\varphi))^T$ ,  $\Delta\varphi = \varphi - \varphi_0$  is the angular difference.  $\mathbf{Q} : (\mathbf{e} \otimes \mathbf{e} - \mathbf{1}/2) = \sum_{\alpha, \beta=1}^2 Q_{\alpha\beta} (\mathbf{e}_\beta \mathbf{e}_\alpha - \delta_{\beta\alpha}/2)$  denotes the full contraction, where  $\mathbf{e} \otimes \mathbf{e}$  is the outer product. We again set  $\tau_0 = 0$  here. Note the multiplicative Heaviside function  $\Theta(r - 1)$  in Eq. (7), which is defined as  $\Theta(x) = 1$ , if  $x > 1$  and  $\Theta(x) = 0$  otherwise. This ensures the no-overlap condition of the hard-disk interaction potential in Eq. (2). The explicit Cartesian multipole expansion for the relative PDF in Eq. (7) constitutes the first terms of a Fourier

expansion for  $p(\mathbf{x}, \tau | \mathbf{x}_0)$  which reads

$$p(\mathbf{x}, \tau | \mathbf{x}_0) = \frac{\Theta(r-1)}{2\pi} \left[ a_0(r, \tau | r_0) + 2 \sum_{n=1}^{\infty} \begin{pmatrix} a_n(r, \tau | r_0) \\ b_n(r, \tau | r_0) \end{pmatrix} \cdot \mathbf{e}(n \Delta\varphi) \right]. \quad (8)$$

Here  $a_n$  and  $b_n$  are the Fourier coefficients of order  $n$ ,  $n \in \mathbb{N}_0$ . We introduce the notation  $p = \Theta(r-1) [p_0 + \sum_{n=1}^{\infty} p_n]$ , where  $p_0 = a_0/2\pi$  and  $p_n = [a_n \cos(n \Delta\varphi) + b_n \sin(n \Delta\varphi)]/\pi$ ,  $n \geq 1$  for our subsequent shorthand notation of the Fourier modes. The Cartesian modes in Eq. (7) are thus connected to the Fourier modes as

$$\varrho(r, \tau | r_0) = \frac{1}{2\pi} a_0(r, \tau | r_0), \quad (9)$$

---


$$a_n(r, \tau | r_0) = \frac{e^{-\frac{r_0^2 + r^2}{4\tau}}}{2\tau} I_n\left(\frac{r_0 r}{2\tau}\right) - \mathcal{L}^{-1} \left\{ k_n(r, s | r_0) \left[ s K'_n(\sqrt{s}) I'_n(\sqrt{s}) + (n\kappa)^2 K_n(\sqrt{s}) I_n(\sqrt{s}) - \delta(r_0 - 1) \frac{\sqrt{s} K'_n(\sqrt{s})}{2K_n(\sqrt{s})} \right] \right\}, \quad (12)$$

$$b_n(r, \tau | r_0) = -n\kappa \left( 1 - \frac{\delta(r_0 - 1)}{2} \right) \mathcal{L}^{-1} \{ k_n(r, s | r_0) \}, \quad (13)$$

where we used the abbreviation

$$k_n(r, s | r_0) = \frac{K_n(r\sqrt{s}) K_n(r_0\sqrt{s})}{(\sqrt{s} K'_n(\sqrt{s}))^2 + (n\kappa K_n(\sqrt{s}))^2}. \quad (14)$$

Here  $s$  denotes the (dimensionless) Laplace variable. The Laplace transform for a function  $f(\tau)$  is defined as  $\mathcal{L}\{f\}(s) = \int_0^\infty d\tau \exp(-s\tau) f(\tau)$ , using the rescaled time variable  $\tau = t/\tau_d$ . The prime denotes the derivative  $g'(a) = dg(x)/dx|_{x=a}$  and  $I_n(x)$ ,  $K_n(x)$  are the modified Bessel functions of the first kind and second kind, respectively [32]. As apparent from Eqs. (12) and (13), the inverse Laplace transformations remain unfeasible at the moment, and we rely on established numerical Laplace inversion methods.

We stress that the problem of solving for the full propagator of interacting particles in Eq. (3) can be formulated in the language of scattering theory [33], thus allowing to use well-developed tools from quantum field theory [34]. Odd diffusion here adds the perspective of non-Hermiticity with potentially new insights [9].

## B. Numerical results and Fourier modes

In Fig. 1 we compare the positional mode  $p_0 = \Theta(r-1)\varrho$ , the polarization mode  $p_1 = \Theta(r-1)[\boldsymbol{\sigma} \cdot \mathbf{e}]$ , and the nematic mode  $p_2 = \Theta(r-1)[\mathbf{Q} : (\mathbf{e} \otimes \mathbf{e} - \mathbf{1}/2)]$  for a normal ( $\kappa = 0$ ) and an odd-diffusive ( $\kappa = 1$ ) system of interacting particles. The modes are evaluated at fixed

which is the (scalar) mean positional PDF,

$$\boldsymbol{\sigma}(r, \tau | r_0) = \frac{1}{\pi} \begin{pmatrix} a_1(r, \tau | r_0) \\ b_1(r, \tau | r_0) \end{pmatrix}, \quad (10)$$

which is the (vectorial) polarization order PDF, and

$$\mathbf{Q}(r, \tau | r_0) = \frac{1}{\pi} \begin{pmatrix} a_2(r, \tau | r_0) & b_2(r, \tau | r_0) \\ b_2(r, \tau | r_0) & -a_2(r, \tau | r_0) \end{pmatrix}, \quad (11)$$

which is the (tensorial) nematic order PDF.

In Appendix A we provide the full solution to the relative problem, thereby following the original derivations in Refs. [9, 13], and show that the general Fourier coefficients  $a_n, b_n$  are given by

---

time  $\tau = 1$  and for  $(r_0, \varphi_0) = (1.01, 0)$ , i.e. particles are initially placed at a distance of 1.01 times their diameter along the (arbitrarily) chosen  $x$ -axis.

We observe that  $\kappa \neq 0$  does not affect the interacting particles' mean positional distribution  $p_0$ , but rotates higher order modes in comparison to  $\kappa = 0$ . We can understand this by observing from Eq. (13) that  $b_n \propto n\kappa$ , such that  $p_0$  is not affected by  $\kappa \neq 0$ —but for higher order modes  $b_n$  contributes for an odd-diffusive system. It is of interest to analyse the contribution of the higher-order modes to the time evolution of the relative PDF  $p$ . We define a measure of the significance of a mode of order  $n \geq 1$  as

$$\delta_n(\tau) = \frac{\int d\mathbf{x} |p_n(\mathbf{x}, \tau | \mathbf{x}_0)|}{\int d\mathbf{x} p_0(\mathbf{x}, \tau | \mathbf{x}_0)}. \quad (15)$$

We take the absolute value  $|p_n|$  in the definition of  $\delta_n(\tau)$ , as the full space integral for the polarization, nematic and all higher order modes vanishes due to the orthogonality of the harmonic functions,  $\int d\mathbf{x} p_n = 0$ ,  $n \geq 1$ . In contrast,  $\int d\mathbf{x} p_0 = 1$  (see Appendix B), which we only include in Eq. (15) to avoid numerical discretization errors.

Fig. 2 shows  $\delta_n(\tau)$  for  $n = 1, \dots, 5$  and for different values of the odd-diffusion parameter  $\kappa = 0, 1, 5$ . When considering  $\delta_n$  as a measure for the relevance of the  $n$ th order mode, we observe that the modes are ordered consecutively in their contribution to the relative PDF  $p$  and that considering higher order modes becomes important




FIG. 1. Plots of the first three Fourier modes  $p_0$ ,  $p_1$  and  $p_2$  of the relative PDF of interacting (odd-diffusive) particles  $p(r, \tau|r_0) = \Theta(r-1) \sum_{n=0}^{\infty} p_n(r, \tau|r_0)$  at fixed time  $\tau = 1$ . (a) - (c) show the modes for normal particles ( $\kappa = 0$ ) and (d) - (f) for odd-diffusive particles ( $\kappa = 1$ ). The initial positions are chosen as  $(r_0, \varphi_0) = (1.01, 0)$ , i.e. the particles are placed 1.01 times their diameter along an arbitrarily defined  $x$ -axis. The zeroth-order mode in (a) and (d) equals the mean positional PDF for the relative coordinate  $r$  (see Eq. (9)) which is unaffected by odd diffusion. The first order mode in (b) and (e), which corresponds to the (contracted) polarizational order PDF (see Eq. (10)), and the second order mode in (c) and (f), which corresponds to the (contracted) nematic order PDF (see Eq. (10)), as well as all higher order modes are affected by odd diffusion such that the modes are rotated in time compared to the  $\kappa = 0$  case, see also Fig. 4. Note that the gray circle in the middle of each plot is due to the excluded volume condition  $\Theta(r-1)$  of the hard-disk interaction of the particles.

for  $\tau \rightarrow 0$  as  $p(\tau) \rightarrow p(0) \propto \delta(\mathbf{x} - \mathbf{x}_0)$ , but higher order modes become less important for  $\tau \gg 0$ . As can be seen in Figs. 2(b), (c),  $\kappa > 0$  shifts the decay of  $\delta_n(\tau)$  to even shorter times.

In Fig. 3 we plot the full relative PDF  $p(\tau)$  at times  $\tau = 1, 5, 50$  for a normal ( $\kappa = 0$ ) and an odd-diffusive ( $\kappa = 1$ ) system. Again we choose the initial condition to be  $(r_0, \varphi_0) = (1.01, 0)$  and, based on the observation in Fig. 2, we truncated the Fourier series at  $n = 10$ . Comparing the normal and odd-diffusive relative PDF, we observe that the  $\kappa$ -induced rotation of every but the zeroth order Fourier mode persists into the full PDF. For  $\kappa = 0$ ,  $b_n = 0$  for all modes, and the Fourier representation of the relative PDF therefore only contains cosine terms. The relative PDF of a normal particle thus is symmetric around  $\Delta\varphi = 0, \pi$  for all times. This symmetry implies that particles encounter the space from both sides after a collision with equal likelihood. However, this does not hold for odd-diffusive particles.  $\kappa \neq 0$  introduces a handedness in the diffusive exploration of space. For interacting odd-diffusive particles, the additional rotational prob-

ability flux introduced via Eq. (1) results in a preferred direction after a collision depending on the sign of the odd-diffusion parameter  $\kappa$ . This effect was recently observed from Brownian dynamics simulations and termed as “mutual rolling” [2]. The interaction-induced symmetry breaking has far-reaching consequences for observable transport coefficients such as the self-diffusion coefficient. In odd-diffusive systems, even though resembling equilibrium overdamped dynamics, the self-diffusion can be enhanced by interactions instead of being reduced as for a normal system [2]. Even though, seemingly contradicting equilibrium statistical mechanics theorems [23, 35, 36], the interaction-enhanced self-diffusion could recently be rationalized by observing that the time-evolution in odd-diffusive systems (see Eq. (3)) becomes non-Hermitian for finite  $\kappa$  [9].




FIG. 2. Significance  $\delta_n(\tau)$  of a mode  $n$  as defined in Eq. (15), which measures the contribution of a Fourier mode of order  $n \geq 1$ ,  $p_n(\tau)$ , to the full relative PDF  $p(\tau) = \Theta(r-1) \sum_{n=0}^{\infty} p_n(\tau)$  as a function of time  $\tau$  for odd-diffusion parameter  $\kappa = 0, 1$  and  $5$  in (a), (b) and (c), respectively. We see that higher order modes are consecutively ordered in their contribution to  $p$  in time and become of comparable importance for  $\tau \rightarrow 0$  as  $p(\tau) \rightarrow p(0) \propto \delta(\mathbf{x} - \mathbf{x}_0)$ . Comparing (a) with (c) we observe that  $\kappa > 0$  shifts the characteristic decay to shorter times. Thus, for dynamics where one is not interested in the  $\tau \rightarrow 0$  limit, one can truncate the Taylor expansion at finite  $n$ .




FIG. 3. PDF  $p(r, \varphi, \tau | r_0, \varphi_0)$  for the relative coordinate of two hard interacting particles. (a) to (c) show normal particles ( $\kappa = 0$ ) and (d) to (f) show odd-diffusive particles ( $\kappa = 1$ ) at times  $\tau = 1, 5$  and  $50$ . The initial positions are chosen as  $(r_0, \varphi_0) = (1.01, 0)$ , i.e. the particles are placed 1.01 times their diameter along an arbitrarily defined  $x$ -axis. The Fourier expansion of  $p(\tau)$  in Eq. (7) is truncated at  $n = 10$  in the numerical evaluation for the plots, see also Fig. 2 for a rationale. (d) to (f) show the “mutual rolling effect”, i.e. that odd-diffusive particles rotate around each other after a collision [2], which is in contrast to a symmetric back-reflection for normal particles, visible in (a) - (c). Note the grey circle in the middle of each plot which is due to the excluded volume condition  $\Theta(r-1)$  of the hard-disk interaction of the particles.

### C. Relevance of the polarization mode for the force autocorrelation

The force autocorrelation tensor (FACT)  $\mathbf{C}_F(\tau)$  encodes the most detailed microscopic information in an interacting system. Via a Taylor-Green-Kubo relation [37–39] it encodes the particle-particle interaction effects in the self-diffusion. For a stationary system,  $\mathbf{C}_F(\tau)$  is defined as

$$\begin{aligned} \mathbf{C}_F(\tau) &= \langle \mathbf{F}(\vec{\mathbf{x}}) \otimes \mathbf{F}(\vec{\mathbf{x}}_0) \rangle \\ &= \int d\vec{\mathbf{x}} \int d\vec{\mathbf{x}}_0 \mathbf{F}(\vec{\mathbf{x}}) \otimes \mathbf{F}(\vec{\mathbf{x}}_0) P_N(\vec{\mathbf{x}}, \tau, \vec{\mathbf{x}}_0, \tau_0), \end{aligned} \quad (16)$$

where  $\vec{\mathbf{x}} = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$  and similarly  $\vec{\mathbf{x}}_0$  for a system of, in general,  $N$  particles.  $P_N$  is the  $N$ -particle joint PDF, which can be rewritten as  $P_N(\vec{\mathbf{x}}, \tau, \vec{\mathbf{x}}_0, \tau_0) = P_N(\vec{\mathbf{x}}, \tau | \vec{\mathbf{x}}_0) P_{\text{eq}}(\vec{\mathbf{x}}_0, \tau_0)$ , assuming the particles were in equilibrium at time  $\tau_0$ , which we take again to be 0. The force on the tagged particle (particle one) is  $\mathbf{F}(\vec{\mathbf{x}}) = -\nabla_1 U_N(\vec{\mathbf{x}})$ , and we assume a pairwise additive and radially symmetric potential  $U_N(\vec{\mathbf{x}}) = \sum_{i,j=1}^N U(r_{ij})/2$ , where  $r_{ij} = |\mathbf{x}_i - \mathbf{x}_j|, i \neq j$ . In the dilute limit, we can safely assume that only two-body correlations are important and thus ignore correlations between the untagged particles. The equilibrium PDF can be approximated as  $P_{\text{eq}}(\vec{\mathbf{x}}_0) = 1/(VZ_N) \prod_{i=2}^N \exp(-\beta U(r_{1i,0})/2)$ , where  $V \subset \mathbb{R}^2$  is the bounded space of diffusion and  $Z_N = \int d\vec{\mathbf{x}}_0 \prod_{i=2}^N \exp(-\beta U(r_{1i,0})/2)$  is the  $N$ -particle partition function. The conditional PDF, i.e., the  $N$ -particle propagator can be approximated as  $P_N(\vec{\mathbf{x}}, \tau | \vec{\mathbf{x}}_0) = (1/V) \prod_{i=2}^N p(\mathbf{x}_{1i}, \tau | \mathbf{x}_{1i,0})$ , where  $p(\mathbf{x}_{1i}, \tau | \mathbf{x}_{1i,0})$  is the PDF of the relative coordinate  $\mathbf{x}_{1i} = \mathbf{x}_1 - \mathbf{x}_i$ , similar to Eq. (7) but for a generic two-body interaction potential  $U$ . Following these approximations, all but one particle (particle two) can be integrated out in Eq. (16) and we denote as before,  $\mathbf{x} = \mathbf{x}_1 - \mathbf{x}_2$ . The central interaction force between the particles can be written as  $\mathbf{F}(\mathbf{x}) = F(r) \mathbf{e}(\varphi)$ , where  $\mathbf{e}(\varphi) = (\cos(\varphi), \sin(\varphi))^T$ , as before, and coincides with the radial unit vector. In the dilute limit, the FACT of Eq. (16) thus becomes

$$\begin{aligned} \mathbf{C}_F(\tau) &= \frac{N-1}{V} \int d\mathbf{x} \int d\mathbf{x}_0 F(r) F(r_0) \frac{e^{-\beta U(r_0)}}{Z_2} \\ &\quad \times p(\mathbf{x}, \tau | \mathbf{x}_0) \mathbf{e}(\varphi) \otimes \mathbf{e}(\varphi_0) \end{aligned} \quad (17)$$

$$\begin{aligned} &= \frac{N-1}{V} \pi \int dr \int dr_0 F(r) F(r_0) \frac{e^{-\beta U(r_0)}}{Z_2} \\ &\quad \times [a_1(r, \tau | r_0) \mathbf{1} - b_1(r, \tau | r_0) \boldsymbol{\epsilon}]. \end{aligned} \quad (18)$$

Here we used the orthogonality of the Fourier modes in the relative PDF  $p(\tau)$  (analogously to Eq. (7)) from Eq. (17) to Eq. (18) to find that only the polarization mode  $\boldsymbol{\sigma}(r, \tau | r_0) = (a_1(r, \tau | r_0), b_1(r, \tau | r_0))^T / \pi$  (analogously to Eq. (10)) contributes to the FACT. Note that relations analogous to Eqs. (7) and (10) also hold in a system with generic radially symmetric interaction po-

tential  $U(r)$ , specifically also for forces with transverse, odd components [7, 40–42].

We can specify Eq. (18) for the hard-disk interaction potential of Eq. (2) by observing that  $P_{\text{eq}}(\mathbf{x}_0) = \Theta(r_0 - 1)/V^2$  and that we can rewrite the singular interaction force via the trick  $\beta\Theta(r-1)\mathbf{F}(r) = \delta(r-1)\mathbf{e}(\varphi)$  [43], obeying the same generic form of a central force as before. We can perform the radial integral of Eq. (18) and find that the FACT of a hard-disk system in the dilute limit is given by  $\mathbf{C}_F(\tau) = \beta^{-2}\phi [a_1(1, \tau|1) \mathbf{1} - b_1(1, \tau|1) \boldsymbol{\epsilon}]$ , where  $\phi = (\pi d^2/4)(N/V)$  is the area fraction in dimensional form [9, 22]. Thus we understand that the behavior of the polarization mode  $\boldsymbol{\sigma}(\tau)$  in time governs the behavior of the FACT for interacting systems, in particular for a hard-disk interaction. Note specifically here, that the off-diagonal correlation ( $\propto b_1 \boldsymbol{\epsilon}$ ) is directly proportional to  $\kappa$ , and therefore might serve as a characteristic to odd diffusion [3, 9].

To further analyze the polarization mode in the hard system, we define

$$\gamma(\tau) = \arctan\left(\frac{b_1(1, \tau|1)}{a_1(1, \tau|1)}\right), \quad (19)$$

as the mean angle between the polarization vector  $\boldsymbol{\sigma}(1, \tau|1)$  as a function of time and its initial direction, which defines the  $x$ -axis in the system, see also inset in Fig. 4(a). We plot  $\gamma(\tau)$  for different values of  $\kappa$  as a function of time in Fig. 4(a) and use the  $\arctan 2(\cdot)$ -function for numerical evaluation to obtain angles within the interval  $(-\pi, \pi)$ . We observe that for  $\kappa = 0$  the initial direction of the polarization does not change with time and stays constant. Recalling the polarization mode in Fig. 1(b), the location of the extrema thus remain unchanged, meaning that the particles are symmetrically back-reflected from the center of the collision. For  $\kappa \neq 0$ , however,  $\gamma(\tau)$  changes in time (see again Fig. 1(e)). The mean-direction of the polarization develops an extremum  $\gamma_{\text{ext}}$  and relaxes to a constant, non-zero value  $\gamma_{\infty}$  for long times. For  $\kappa > 0.88$ ,  $\gamma_{\text{ext}} < -\pi/2$  and for  $\kappa > 1$  even  $\gamma_{\infty} < -\pi/2$ . The interval  $\kappa \in (0.88, 1)$  thereby aligns with the odd-diffusion interval, for which we reported an oscillating FACF earlier [9]. These sign-switches in the FACF coincide with what we observe now for the direction of  $\boldsymbol{\sigma}(\tau)$ , see also inset in Fig. 4(a):  $\gamma(\tau)$  crosses  $-\pi/2$  for some time (negative FACF) and eventually relaxes to angles larger than  $-\pi/2$  again (positive FACF).

We interpret this behavior as interacting odd-diffusive particles, which for  $|\kappa| > 0.88$  rotate more than  $\pi/2$  but relax to a steady state with a relative angle  $|\gamma_{\infty}| < \pi/2$  as long as  $|\kappa| < 1$ ; see Fig. 4(b) for a sketch. This behavior also rationalizes the oscillating FACF in the same interval, as  $\langle \mathbf{F}(\tau) \cdot \mathbf{F}(0) \rangle = \langle F(\tau) F(0) \cos(\pi/2) \rangle = 0$  here and the particles oscillate around that angle. However, a steady state angle  $|\gamma_{\infty}|$  which is smaller than the maximal angle  $|\gamma_{\text{ext}}|$  is a generic observation from Fig. 4(a). On average particles always rotate further in time, as they finally relax. Note that for  $\kappa \rightarrow -\kappa$ , we have that  $\gamma(\tau) \rightarrow -\gamma(\tau)$ , as  $a_n$  is even and  $b_n$  is an odd function

of  $\kappa$ . Thus,  $\kappa < 0$  results in the same phenomenology for the relaxation of  $\gamma(\tau)$  and thus for the FACF. Further, if  $|\kappa| \rightarrow \infty$ , we find that  $|\gamma_{\text{ext}}| \rightarrow |\gamma_{\infty}| \rightarrow \pi$ , which means, that at most odd particles exchange positions but do not rotate any further.

### III. DISCUSSION

We here presented the exact analytical solution and numerical evaluation of two hard disk-like interacting odd-diffusing particles in two spatial dimensions. Odd diffusion thereby is characterized by antisymmetric elements  $\propto \kappa$  in the diffusion tensor  $\mathbf{D} = D_0(\mathbf{1} + \kappa\epsilon)$ . Our analysis showed that the two-particle problem separates into a center-of-mass and a relative coordinate problem, of which the first can be solved straightforwardly, and the latter incorporates the nontrivial effects of interactions and odd diffusion. The relative PDF can be written as a Fourier series, and we observe that oddness rotates all apart from the zeroth order mode in time, which represents the unaffected positional distribution of the relative coordinate. The modes are consecutively ordered in their contribution to the relative PDF, and when summed up show a rotated form of the PDF. This effect is a characteristic of odd diffusion and has been termed as “mutual rolling” earlier [44] as the particles rotate around each other while interacting, in contrast to normal diffusive ( $\kappa = 0$ ) particles, which are symmetrically back-reflected when interacting.

The representation of the relative PDF in its Fourier modes becomes useful in the analysis of microscopic correlation functions, specifically the FACF. For any central interaction force, only the polarization mode of the relative PDF determines the correlation function. We conjecture here that a similar phenomenology holds for other (radially symmetric) observables of arbitrary tensorial order in an interacting system; the corresponding mode of the relative PDF might determine the expectation value and even the correlation function. We used the analytical access to the polarization mode, to understand the average configurations of the hard-disk-like interacting odd-diffusive particles. The maximal relative rotation angle overshoots the final relaxation on average, which quantitatively aligns with an oscillating FACF, recently reported for these systems [9].

Our work shines light on the novel way of particle interactions in odd-diffusive systems. Thereby it might serve as a reference case for interactions in the various odd-diffusive systems such as in equilibrium, e.g., Brownian particle under Lorentz force [6, 44–56], with a long-lasting history in statistical mechanics [57–59], or skyrmionic spin structures [60–62]. Here numerical studies recently reproduced the interaction-enhanced self-diffusion [7], which originates in the mutual rolling effect [2]. But there exist also non-equilibrium odd-diffusive systems, such as systems under shear [63, 64], or active chiral particles [1, 30, 42, 65–69]. In general,

for systems that break time-reversal symmetry, the relevance of off-diagonal correlation functions for transport properties has attracted considerable interest recently [1, 3, 5, 42, 67, 70, 71]. Odd diffusion further might serve as the unifying terminology for systems showing transverse responses such as systems with Magnus forces [72–74], Coriolis forces [75, 76], in complex (porous) environments [77–79], or to describe non-conservative force fields which are found in optical tweezer experiments [80–83]. Systems with an artificial transverse interaction component [41, 84] recently showed that odd interactions enhance the sampling of configurations in dense systems, originating again in the unique particle rearrangements in odd-diffusive systems. Transverse forces further have been the subject of interest in so-called linear-diffusive systems [85–89]. Odd diffusion is finally also relevant for strongly rotating [90, 91] or magnetized plasmas, as it can be found, e.g. in the realm of astrophysics, where antisymmetric transport is relevant to describe the movement of energetic particles moving through magnetized plasmas such as the interplanetary and interstellar medium [92–94].

### ACKNOWLEDGMENTS

*Acknowledgements.* The authors acknowledge the help of Timo J. Doerries in the creation of Fig. 4(b). A. S., R. M., and E. K. acknowledge support by the Deutsche Forschungsgemeinschaft (grants No. SH 1275/5-1, ME 1535/16-1 and SPP 2332 - 492009952).

#### Appendix A: Analytical solution of the relative problem

This Appendix closely follows Ref. [9] in its presentation of the analytical solution to the relative problem, which itself adapted the work of Hanna, Hess and Klein [13] to odd-diffusive systems.

The equation for the relative PDF  $p(\tau) = p(\mathbf{x}, \tau | \mathbf{x}_0)$ , where  $\mathbf{x} = \mathbf{x}_1 - \mathbf{x}_2$ , follows from Eq. (5) as

$$\frac{\partial}{\partial \tau} p(\tau) = \nabla_{\mathbf{x}} \cdot (\mathbf{1} + \kappa \epsilon) [\nabla_{\mathbf{x}} + \beta \nabla_{\mathbf{x}} U(r)] p(\tau), \quad (\text{A1})$$

and reads in (rescaled) polar coordinates  $\mathbf{x} = (r, \varphi)$  as

$$\begin{aligned} \frac{\partial}{\partial \tau} p(\tau) = & \frac{1}{r^2} \left( r \frac{\partial}{\partial r} r \frac{\partial}{\partial r} + r \frac{\partial}{\partial r} r \beta \frac{\partial U(r)}{\partial r} \right. \\ & \left. + \frac{\partial^2}{\partial \varphi^2} - \kappa r \beta \frac{\partial U(r)}{\partial r} \frac{\partial}{\partial \varphi} \right) p(\tau). \end{aligned} \quad (\text{A2})$$

Note that space is rescaled with the diameter of the particle  $d$ ,  $\mathbf{x} \rightarrow \mathbf{x}/d$ , and time is rescaled by the natural time-scale of diffusing the radial distance of a particle diameter  $\tau_d = d^2/(2D_0)$ ,  $t \rightarrow \tau = t/\tau_d$ . The initial condition  $p(\tau = 0) = \delta(\mathbf{x} - \mathbf{x}_0) \Theta(r - 1)$  in polar coordinates




FIG. 4. (a) Angle  $\gamma(\tau)$  between the polarization mode  $\sigma(r = 1, \tau|r_0 = 1)$ , Eq. (10), and its initial direction as function of time  $\tau$  for different odd-diffusion parameters  $\kappa$ , see Eq. (19) and the inset. The polarization mode constitutes the vectorial mode of the relative PDF  $p(\tau)$  for interacting particles and represents relative particle rearrangements. In a normal system ( $\kappa = 0$ ),  $\gamma(\tau) = 0$  and is constant in time, whereas it develops an extremum  $\gamma_{\text{ext}}$  different from its asymptotic steady state value  $\gamma_{\infty}$  for  $\kappa \neq 0$  in time. We observe that  $|\gamma_{\text{ext}}| < |\gamma_{\infty}|$ , indicating that particles on average rotate further in time than they finally relax to. Specifically for  $\kappa \in (0.88, 1.0)$ ,  $\gamma_{\text{ext}} < -\pi/2$ , but  $\gamma_{\infty} > -\pi/2$  which quantitatively aligns with the oscillating force autocorrelation function previously observed for this regime [9]. (b) The sketch qualitatively illustrates the scenario, that odd-diffusive particles with  $0.88 < \kappa < 1$  collide at  $\tau_0$  with each other along the (arbitrary)  $x$ -axis and rotate for  $\tau_1 > \tau_0$  to  $\gamma_{\text{ext}} < -\pi/2$ , and eventually relax at  $\tau_2 > \tau_1$  to  $\gamma_{\infty} > -\pi/2$ .

becomes

$$p(\tau = 0) = \frac{\delta(r - r_0)}{r_0} \delta(\varphi - \varphi_0) \Theta(r - 1). \quad (\text{A3})$$

The angular part of the initial condition can be expanded into a Fourier series  $\delta(\varphi - \varphi_0) = \sum_{n=-\infty}^{\infty} \exp(in(\varphi - \varphi_0))/2\pi$ , which, following Refs. [9, 13], we use as an ansatz to solve for  $p(\tau)$  as

$$p(\mathbf{x}, \tau|\mathbf{x}_0) = \frac{\Theta(r - 1)}{2\pi} \sum_{n=-\infty}^{\infty} R_n(r, \tau|r_0) e^{in(\varphi - \varphi_0)}. \quad (\text{A4})$$

Note that except for the radial functions  $R_n(r, \tau|r_0)$ , all other parts of the ansatz are time-independent.

Observing that for the hard-disk potential, see Eq. (2), we have that  $\exp(-\beta U(r)) = \Theta(r - 1)$ , we can replace the otherwise singular interaction force  $\partial U(r)/\partial r$  in Eq. (A2) via

$$-\beta\Theta(r - 1) \frac{\partial U(r)}{\partial r} = \frac{\partial}{\partial r} e^{-\beta U(r)} = \delta(r - 1). \quad (\text{A5})$$

Rewriting Eq. (A2) for the radial functions  $R_n(\tau) =$

$R_n(r, \tau|r_0)$ , we thus find

$$\frac{\partial}{\partial \tau} R_n(\tau) = \frac{1}{r^2} \left( r \frac{\partial}{\partial r} r \frac{\partial}{\partial r} - n^2 \right) R_n(\tau) \quad (\text{A6})$$

in the domain  $r \geq 1$  and for each order  $n \in \mathbb{Z}$ , and

$$\frac{\partial}{\partial r} R_n(\tau) = -\frac{in\kappa}{r} R_n(\tau) \quad (\text{A7})$$

to be satisfied additionally at  $r = 1$ . Eq. (A7) can be viewed as an extension of an ordinary von Neumann no-flux boundary condition which is recovered for  $\kappa = 0$ . This generalized condition can be found in the literature under the name of *oblique* boundary conditions, see for example [95]. Eq. (A6) is equipped with a second boundary condition, which can be found from the natural boundary condition on  $p(\tau)$ ,  $\lim_{r \rightarrow \infty} p(\tau) = 0$ , as  $\lim_{r \rightarrow \infty} R_n(\tau) = 0$  and the initial condition on  $R_n(\tau)$  translates from Eq. (A3) as  $R_n(\tau = 0) = \delta(r - r_0)/r_0$ . Note that Eq.(A6) together with the boundary condition (A7) also forms the basis of solving the problem by using the language of scattering theory [33], the only difference being the introduction of the the generalized

von-Neumann boundary condition for interacting odd-diffusive particles.

For the particular solution of Eq. (A6),  $R_n^{\text{part}}(\tau)$ , we make the ansatz

$$R_n^{\text{part}}(\tau) = \int_0^\infty du w_n(r, u|r_0) e^{-\tau u^2}, \quad (\text{A8})$$

which when inserted into Eq. (A6) shows that  $w_n(r, u|r_0)$  satisfy the Bessel equation [32] with solutions  $J_n(ur)$  and  $Y_n(ur)$  as the Bessel functions of first kind and second kind, respectively. Matching the particular solution with the initial condition  $R_n(\tau = 0) = \delta(r - r_0)/r_0$  and noting that the delta-distribution  $\delta(\cdot)$  can be expanded in Bessel functions of the first kind (see Eq. (B1)) we find that  $Y_n(ur)$  does not contribute to the particular solution.

After a Laplace transformation with (dimensionless) Laplace variable  $s$ , the homogeneous part of Eq. (A6) appears as the modified Bessel equation [32] with solutions  $I_n(r\sqrt{s})$  and  $K_n(r\sqrt{s})$ , as the modified Bessel functions of the first and second kind, respectively. The Laplace transformation for a function  $f(\tau)$  thereby is defined as  $\mathcal{L}\{f\}(s) = \int_0^\infty d\tau \exp(-s\tau)f(\tau)$ , noting that  $\tau = t/\tau_d$ . As the homogeneous solution has to satisfy the natural boundary condition on  $R_n(\tau)$ ,  $I_n(x)$  is no suitable solution as it diverges for  $x \rightarrow \infty$  for every  $n \in \mathbb{Z}$  [32, 9.7.1]. We conclude that the homogeneous part is solved by  $\mathcal{L}\{R_n^{\text{hom}}\}(s) = A_n(s|r_0) K_n(r\sqrt{s})$  for some amplitude  $A_n(s|r_0)$  to be determined by matching with the oblique boundary condition (A7).

The relative solution  $p(\tau)$  in the Laplace domain thus is given by

$$\mathcal{L}\{p\}(\mathbf{x}, s|\mathbf{x}_0) = \Theta(r-1) \sum_{n=-\infty}^{\infty} \frac{e^{in(\varphi-\varphi_0)}}{2\pi} \int_0^\infty du \frac{u J_n(ur_0)}{s+u^2} \left[ J_n(ur) - K_n(r\sqrt{s}) \frac{u J_n'(u) + in\kappa J_n(u)}{\sqrt{s} K_n'(\sqrt{s}) + in\kappa K_n(\sqrt{s})} \right]. \quad (\text{A9})$$

which was already found in Ref. [9]. There are two types of integrals appearing in Eq. (A9), which involve products of Bessel functions and which we list in Appendix B. The integrals  $\int_0^\infty du u J_n(ur_0) J_n(ub)/(s+u^2)$  can be evaluated using Eq. (B2), where  $b \in \{1, r\}$ . Depending on whether  $b = r > r_0$ ,  $r = r_0$  or  $r < r_0$ , Eq. (B2) gives different results but we only need to consider the latter two

for the cases  $b = 1 = r_0$  and  $1 < r_0$ . The second integral  $\int_0^\infty du u^2 J_n(ur_0) J_n'(u)/(s+u^2)$  involves the derivative of a Bessel function and therefore is more involved. We list the integral in Eq. (B5) and again need to differentiate whether  $r_0 = 1$  or  $r_0 > 1$ . Taking into account these different cases, we find for the Laplace transformed relative PDF

$$\mathcal{L}\{p\}(\mathbf{x}, s|\mathbf{x}_0) = \frac{\Theta(r-1)}{2\pi} \sum_{n=-\infty}^{\infty} e^{in(\varphi-\varphi_0)} \left[ \Theta(r_0-r) I_n(r\sqrt{s}) K_n(r_0\sqrt{s}) + \Theta(r-r_0) I_n(r_0\sqrt{s}) K_n(r\sqrt{s}) - \frac{K_n(r\sqrt{s}) K_n(r_0\sqrt{s})}{\sqrt{s} K_n'(\sqrt{s}) + in\kappa K_n(\sqrt{s})} \left( \sqrt{s} I_n'(\sqrt{s}) + in\kappa I_n(\sqrt{s}) - \frac{\delta(r_0-1)}{2K_n(\sqrt{s})} \right) \right]. \quad (\text{A10})$$

We can Laplace invert some parts of Eq. (A10) as

$$\begin{aligned} \mathcal{L}^{-1}\{I_n(r_0\sqrt{s}) K_n(r\sqrt{s})\} &= \mathcal{L}^{-1}\{I_n(r\sqrt{s}) K_n(r_0\sqrt{s})\} \\ &= \frac{\exp\left(-\frac{r_0^2+r^2}{4\tau}\right)}{2\tau} I_n\left(\frac{r_0 r}{2\tau}\right), \end{aligned} \quad (\text{A11})$$

[96, 13.96]. Together with decomposing  $e^{ix} = \cos x + i \sin x$  in Eq. (A10), this gives the Fourier expanded relative PDF as

$$\begin{aligned} p(\mathbf{x}, \tau|\mathbf{x}_0) &= \frac{\Theta(r-1)}{2\pi} \left[ a_0(r, \tau|r_0) \right. \\ &\left. + 2 \sum_{n=1}^{\infty} \begin{pmatrix} a_n(r, \tau|r_0) \\ b_n(r, \tau|r_0) \end{pmatrix} \cdot \begin{pmatrix} \cos(n(\varphi-\varphi_0)) \\ \sin(n(\varphi-\varphi_0)) \end{pmatrix} \right], \end{aligned} \quad (\text{A12})$$

where  $a_n$  and  $b_n$  are given in Eqs. (12) and (13) in the main text. Note that by using the symmetry in the order of the modified Bessel functions,  $I_{-n}(x) = I_n(x)$  and  $K_{-n}(x) = K_n(x)$  [32, 9.6.6], we can restrict the sum on positive modes only.

## Appendix B: Integral expressions

To evaluate the integrals in Eq. (A9), we use the following tabulated integrals, listed in this Appendix.

Following Arfken and Weber's *Mathematical Methods for Physicists* [97], we find that Bessel functions of (in-

teger) order  $n$ ,  $J_n$ , obey the integral relation

$$\int_0^\infty du u J_n(au) J_n(bu) = \frac{\delta(a-b)}{a}, \quad (\text{B1})$$

valid for  $n > -1/2$  and  $a, b$  some real-valued constants [97, 11.59].

To evaluate the remaining integrals of Bessel-functions, we draw on Gradshteyn and Ryzhik's *Table of Integrals, Series and Products* [98] and Abramowitz and Stegun's *Handbook of Mathematical Functions* [32]. The first relevant integral is

$$\int_0^\infty du \frac{u J_n(au) J_n(bu)}{u^2 + c^2} = \begin{cases} I_n(ac) K_n(bc), & 0 < a < b \\ I_n(ac) K_n(ac), & 0 < a = b \\ I_n(bc) K_n(ac), & 0 < b < a \end{cases}, \quad (\text{B2})$$

which is valid for  $n > -1$  and  $c > 0$  [98, 6.541, 6.535]. Here  $I_n, K_n$  are the modified Bessel functions of the first and second kind, and  $a, b, c$  some (real-valued) constants. Together with the symmetry relations for the order of Bessel functions,  $J_{-n}(x) = (-1)^n J_n(x)$ ,  $I_{-n}(x) = I_n(x)$  and  $K_{-n}(x) = K_n(x)$  [32, 9.1.5, 9.6.6], Eq. (B2) can be further extended to the relevant cases of  $n < 0$ .

The second relevant integral is  $\int_0^\infty du u^2 J_n(au) J'_n(bu)/(u^2 + c^2)$ , for some real-valued, positive constants  $a, b, c$ . Using  $dJ_n(x)/dx = (J_{n-1}(x) - J_{n+1}(x))/2$  [98, 8.471], the integral can be evaluated as

$$\int_0^\infty du \frac{u^2 J_{n_1}(au) J_{n_2}(bu)}{u^2 + c^2} = \begin{cases} (-1)^{\alpha_+} c I_{n_1}(ac) K_{n_2}(bc), & 0 < a < b \\ (-1)^{\alpha_-} c I_{n_2}(bc) K_{n_1}(ac), & 0 < b < a \end{cases}, \quad (\text{B3})$$

which is valid for  $n_2 > -1$  [98, 6.577]. The exponents are  $\alpha_\pm = (1 \pm (n_1 - n_2))/2 \in \mathbb{N}_0$ . We use Eq. (B3) for the case of  $n_1 = n \pm 1$  and  $n_2 = n$ . By relying on the symmetry relations for the orders of the Bessel functions, we can extend Eq. (B3) again to the negative cases of  $n_1 = -(n \pm 1)$  and  $n_2 = -n$ . Thus, together with [98, 8.468], we find

$$\int_0^\infty du \frac{u^2 J_n(au) J'_n(bu)}{u^2 + c^2} = \begin{cases} c K_n(bc) I'_n(ac), & 0 < a < b \\ \frac{n}{b} K_n(ac) I_n(bc), & 0 < b < a \end{cases}, \quad (\text{B4})$$

valid for all  $n \in \mathbb{Z}$ , but limited to  $b \neq a$ .

In order to generalize Eq. (B4) to the case of  $b = a > 0$ , which can not be found in Ref. [98], we generalize the relation, which was already derived in Ref. [9] for  $n = 1$  to an arbitrary  $n \in \mathbb{Z}$ . Therefore, by using the indefinite integral  $\int du J_n(u) J'_n(u) = J_n^2(u)/2 + \text{const}$ , we can partially integrate and find an integral of the form of Eq. (B2), which, using the Wronskian  $\mathcal{W}[I_n(x), K_n(x)] = I_n(x) K_{n+1}(x) + I_{n+1}(x) K_n(x) = 1/x$  [32, 9.1.15], gives

$$\int_0^\infty du \frac{u^2 J_n(au) J'_n(au)}{u^2 + c^2} = \begin{cases} c K_n(bc) I'_n(ac), & 0 < a < b \\ c K_n(ac) I'_n(ac) - \frac{1}{2a}, & 0 < a = b \\ \frac{n}{b} K_n(ac) I_n(bc), & 0 < b < a \end{cases}. \quad (\text{B5})$$

To prove the normalization of the zeroth order mode  $\int d\mathbf{x} p_0 = 1$ , Eq. (9) in the main text, we transform the integrals over  $I_0$  and  $K_0$  into higher order Bessel functions evaluated at the boundaries  $r = 1$  and  $r \rightarrow \infty$  by using the recursion relations [98, 8.486]

$$\left(\frac{1}{u} \frac{d}{du}\right)^m (u^n I_n(u)) = u^{n-m} I_{n-m}(u), \quad (\text{B6})$$

and

$$\left(\frac{1}{u} \frac{d}{du}\right)^m (u^n K_n(u)) = (-1)^m u^{n-m} K_{n-m}(u), \quad (\text{B7})$$

which specifically imply for  $m = 1$  and  $n = 1$  to  $u I_0(u) = d/du (u I_1(u))$  and  $u K_0(u) = -d/du (u K_1(u))$ . The integrals over  $I_0(u)$  and  $K_0(u)$  thus turn into evaluating  $u I_1(u)$  and  $u K_1(u)$  at the integration bounds and the upper bound vanishes by the asymptotic expansion of  $\lim_{u \rightarrow \infty} u K_n(u) \sim \lim_{u \rightarrow \infty} u \exp(-u) = 0$  [98, 8.451]. The remaining expressions from the lower bounds are evaluated by using the Wronskian  $\mathcal{W}[I_n(x), K_n(x)]$ , to be exactly 1.

- 
- [1] C. Hargus, J. M. Epstein, and K. K. Mandadapu, Odd diffusivity of chiral random motion, *Phys. Rev. Lett.* **127**, 178001 (2021).  
[2] E. Kalz, H. D. Vuijk, I. Abdoli, J.-U. Sommer, H. Löwen,

- and A. Sharma, Collisions enhance self-diffusion in odd-diffusive systems, *Phys. Rev. Lett.* **129**, 090601 (2022).  
[3] K. Yasuda, K. Ishimoto, A. Kobayashi, L.-S. Lin, I. Sou, Y. Hosaka, and S. Komura, Time-correlation functions

- for odd Langevin systems, *J. Chem. Phys.* **157**, 095101 (2022).
- [4] M. Fruchart, C. Scheibner, and V. Vitelli, Odd viscosity and odd elasticity, *Annu. Rev. Condens. Matter Phys.* **14**, 471 (2023).
- [5] C. Hargus, A. Deshpande, A. K. Omar, and K. K. Mandadapu, The flux hypothesis for odd transport phenomena, arXiv preprint arXiv:2405.08798 10.48550/arXiv.2405.08798 (2024).
- [6] H.-M. Chun, X. Durang, and J. D. Noh, Emergence of nonwhite noise in Langevin dynamics with magnetic Lorentz force, *Phys. Rev. E* **97**, 032117 (2018).
- [7] D. Schick, M. Weißenhofer, L. Rózsa, J. Rothörl, P. Virnau, and U. Nowak, Two levels of topology in skyrmion lattice dynamics, *Phys. Rev. Res.* **6**, 013097 (2024).
- [8] J. E. Avron, Odd viscosity, *J. Stat. Phys.* **92**, 543 (1998).
- [9] E. Kalz, H. D. Vuijk, J.-U. Sommer, R. Metzler, and A. Sharma, Oscillatory force autocorrelations in equilibrium odd-diffusive systems, *Phys. Rev. Lett.* **132**, 057102 (2024).
- [10] J. K. G. Dhont, *An Introduction to Dynamics of Colloids*, Vol. 2 (Elsevier, 1996).
- [11] G. K. Batchelor, Brownian diffusion of particles with hydrodynamic interaction, *J. Fluid Mech.* **74**, 1 (1976).
- [12] B. U. Felderhof, Diffusion of interacting Brownian particles, *J. Phys. A* **11**, 929 (1978).
- [13] S. Hanna, W. Hess, and R. Klein, Self-diffusion of spherical Brownian particles with hard-core interaction, **111**, 181 (1982).
- [14] J. R. Abney, B. A. Scalettar, and J. C. Owicki, Self diffusion of interacting membrane proteins, *Biophys. J.* **55**, 817 (1989).
- [15] K. Zahn, J. M. Méndez-Alcaraz, and G. Maret, Hydrodynamic interactions may enhance the self-diffusion of colloidal particles, *Phys. Rev. Lett.* **79**, 175 (1997).
- [16] S. D. Bembenek and G. Szamel, The role of attractive interactions in self-diffusion, *J. Phys. Chem. B* **104**, 10647 (2000).
- [17] R. B. Jones, Diffusion of tagged interacting spherically symmetric polymers, **97**, 113 (1979).
- [18] M. Medina-Noyola, Long-time self-diffusion in concentrated colloidal dispersions, *Phys. Rev. Lett.* **60**, 2705 (1988).
- [19] H. Löwen and G. Szamel, Long-time self-diffusion coefficient in colloidal suspensions: theory versus simulation, *J. Phys. Condens. Matter* **5**, 2295 (1993).
- [20] A. Imhof and J. K. G. Dhont, Long-time self-diffusion in binary colloidal hard-sphere dispersions, *Phys. Rev. E* **52**, 6344 (1995).
- [21] A. L. Thorneywork, R. E. Rozas, R. P. A. Dullens, and J. Horbach, Effect of hydrodynamic interactions on self-diffusion of quasi-two-dimensional colloidal hard spheres, *Phys. Rev. Lett.* **115**, 268301 (2015).
- [22] S. Hanna, W. Hess, and R. Klein, The velocity autocorrelation function of an overdamped Brownian system with hard-core interaction, *J. Phys. A* **14**, L493 (1981).
- [23] W. Feller, *An Introduction to Probability Theory and Its Applications*, 3rd ed. (John Wiley & Sons, New York, 1968).
- [24] B. Øksendal, *Stochastic Differential Equations*, 6th ed. (Springer Berlin, Heidelberg, 2003).
- [25] M. Doi and S. F. Edwards, *The Theory of Polymer Dynamics* (Clarendon Press, Oxford, 1988).
- [26] V. Balakrishnan, *Elements of Nonequilibrium Statistical Mechanics* (Springer International Publishing, 2021).
- [27] P.-G. de Gennes and J. Prost, *The Physics of Liquid Crystals*, 2nd ed. (Oxford University Press, 1995).
- [28] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (John Wiley & Sons, 2021).
- [29] M. E. Cates and J. Tailleur, When are active Brownian particles and run-and-tumble particles equivalent? Consequences for motility-induced phase separation, *Europhys. Lett.* **101**, 20010 (2013).
- [30] E. Kalz, A. Sharma, and R. Metzler, Field theory of active chiral hard disks: A first-principles approach to steric interactions, *J. Phys. A* 10.1088/1751-8121/ad5089 (2024).
- [31] M. te Vrugt and R. Wittkowski, Relations between angular and Cartesian orientational expansions, *AIP Adv.* **10**, 035106 (2020).
- [32] M. Abramowitz and I. A. Stegun, eds., *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables* (Dover, New York, 1965).
- [33] T. Franosch, F. Höfling, T. Bauer, and E. Frey, Persistent memory for a Brownian walker in a random array of obstacles, *Chem. Phys.* **375**, 540 (2010).
- [34] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, 4th ed. (Oxford University Press, 2021).
- [35] S. Leitmann and T. Franosch, Time-dependent fluctuations and superdiffusivity in the driven lattice Lorentz gas, *Phys. Rev. Lett.* **118**, 018001 (2017).
- [36] M. Caraglio and T. Franosch, Analytic solution of an active Brownian particle in a harmonic well, *Phys. Rev. Lett.* **129**, 158001 (2022).
- [37] G. I. Taylor, Diffusion by continuous movements, *Proc. Lond. Math. Soc.* **2**, 196 (1922).
- [38] M. S. Green, Markoff random processes and the statistical mechanics of time-dependent phenomena. II. Irreversible processes in fluids, *J. Chem. Phys.* **22**, 398 (1954).
- [39] R. Kubo, Statistical-mechanical theory of irreversible processes. I. General theory and simple applications to magnetic and conduction problems, *J. Phys. Soc. Jpn.* **12**, 570 (1957).
- [40] R. Shinde, J.-U. Sommer, H. Löwen, and A. Sharma, Strongly enhanced dynamics of a charged Rouse dimer by an external magnetic field, *PNAS Nexus* **1**, pgac119 (2022).
- [41] F. Ghimenti, L. Berthier, G. Szamel, and F. van Wijland, Sampling efficiency of transverse forces in dense liquids, *Phys. Rev. Lett.* **131**, 257101 (2023).
- [42] C. H. Batton and G. M. Rotskoff, Microscopic origin of tunable assembly forces in chiral active environments, *Soft Matter* 10.1039/D4SM00247D (2024).
- [43] For the hard-disk potential  $U(r) = 0$  for  $r > 1$  (rescaled space variable) and  $U(r) = \infty$  otherwise we observe that  $\exp(-\beta U(r)) = \Theta(r - 1)$ . We thus find on the one hand that  $\nabla \exp(-\beta U(r)) = -\beta \exp(-\beta U(r)) \nabla U(r) = \beta \Theta(r - 1) \mathbf{F}(r)$ , where  $\mathbf{F}(r) = -\nabla U(r)$  is the interaction force and on the other hand that  $\nabla \exp(-\beta U(r)) = \delta(r - 1) \mathbf{e}(\varphi)$ , where  $\mathbf{e}(\varphi) = (\cos(\varphi), \sin(\varphi))^T$  is the radial unit vector.
- [44] E. Kalz, *Diffusion under the Effect of Lorentz Force* (Springer Spektrum Wiesbaden, 2022).
- [45] D. S. Lemons and D. L. Kaufman, Brownian motion of a charged particle in a magnetic field, *IEEE Trans. Plasma Sci.* **27**, 1288 (1999).
- [46] R. Czopnik and P. Garbaczewski, Brownian motion in a

- magnetic field, *Phys. Rev. E* **63**, 021105 (2001).
- [47] H. Hayakawa, Langevin equation with Coulomb friction, *Physica D* **205**, 48 (2005).
- [48] T. P. Simões and R. E. Lagos, Kramers equation for a charged Brownian particle: The exact solution, **355**, 274 (2005).
- [49] J. I. Jiménez-Aquino and M. Romero-Bastida, Brownian motion of a charged particle in a magnetic field, *Revista Mexicana de Física E* **52**, 182 (2006).
- [50] L. J. Hou, Z. L. Mišković, A. Piel, and P. K. Shukla, Brownian dynamics of charged particles in a constant magnetic field, *Phys. Plasmas* **16**, 10.1063/1.3138746 (2009).
- [51] H. D. Vuijk, J. M. Brader, and A. Sharma, Anomalous fluxes in overdamped Brownian dynamics with Lorentz force, *J. Stat. Mech.: Theory Exp.* **2019** (6), 063203.
- [52] H.-M. Chun, L. P. Fischer, and U. Seifert, Effect of a magnetic field on the thermodynamic uncertainty relation, *Phys. Rev. E* **99**, 042128 (2019).
- [53] I. Abdoli, H. D. Vuijk, J.-U. Sommer, J. M. Brader, and A. Sharma, Nondiffusive fluxes in a Brownian system with Lorentz force, *Phys. Rev. E* **101**, 012120 (2020).
- [54] I. Abdoli, E. Kalz, H. D. Vuijk, R. Wittmann, J.-U. Sommer, J. M. Brader, and A. Sharma, Correlations in multithermostat Brownian systems with Lorentz force, *New J. Phys.* **22**, 093057 (2020).
- [55] J.-M. Park and H. Park, Thermodynamic uncertainty relation in the overdamped limit with a magnetic Lorentz force, *Phys. Rev. Res.* **3**, 043005 (2021).
- [56] I. Abdoli, R. Wittmann, J. M. Brader, J.-U. Sommer, H. Löwen, and A. Sharma, Tunable brownian magneto heat pump, *Sci. Rep.* **12**, 13405 (2022).
- [57] J. B. Taylor, Diffusion of plasma across a magnetic field, *Phys. Rev. Lett.* **6**, 262 (1961).
- [58] B. Kurşunoğlu, Brownian motion in a magnetic field, *Ann. Phys.* **17**, 259 (1962).
- [59] Karmeshu, Brownian motion of charged particles in a magnetic field, *Phys. Fluids* **17**, 1828 (1974).
- [60] C. Schütte, J. Iwasaki, A. Rosch, and N. Nagaosa, Inertia, diffusion, and dynamics of a driven skyrmion, *Phys. Rev. B* **90**, 174434 (2014).
- [61] R. E. Troncoso and Á. S. Núñez, Brownian motion of massive skyrmions in magnetic thin films, *Ann. Phys.* **351**, 850 (2014).
- [62] R. Gruber, M. A. Brems, J. Rothörl, T. Sparmann, M. Schmitt, I. Kononenko, F. Kammerbauer, M.-A. Syskaki, O. Farago, P. Virnau, and M. Kläui, 300-times-increased diffusive skyrmion dynamics and effective pinning reduction by periodic field excitation, *Adv. Mater.* **35**, 2208922 (2023).
- [63] C. Dieball and A. Godec, Coarse graining empirical densities and currents in continuous-space steady states, *Phys. Rev. Res.* **4**, 033243 (2022).
- [64] C. Dieball and A. Godec, Mathematical, thermodynamical, and experimental necessity for coarse graining empirical densities and currents in continuous space, *Phys. Rev. Lett.* **129**, 140601 (2022).
- [65] C. Reichhardt and C. J. O. Reichhardt, Active microrheology, Hall effect, and jamming in chiral fluids, *Phys. Rev. E* **100**, 012604 (2019).
- [66] P. L. Muzzeddu, H. D. Vuijk, H. Löwen, J.-U. Sommer, and A. Sharma, Active chiral molecules in activity gradients, *J. Chem. Phys.* **157**, 134902 (2022).
- [67] A. R. Poggioli and D. T. Limmer, Odd mobility of a passive tracer in a chiral active fluid, *Phys. Rev. Lett.* **130**, 158201 (2023).
- [68] C. W. Chan, D. Wu, K. Qiao, K. L. Fong, Z. Yang, Y. Han, and R. Zhang, Chiral active particles are sensitive reporters to environmental geometry, *Nat. Commun.* **15**, 1406 (2024).
- [69] F. Siebers, R. Bebon, A. Jayaram, and T. Speck, Collective Hall current in chiral active fluids: Coupling of phase and mass transport through traveling bands, *Proc. Natl. Acad. Sci. U.S.A.* **121**, e2320256121 (2024).
- [70] G. Pavliotis, Asymptotic analysis of the Green-Kubo formula, *IMA J. Appl. Math.* **75**, 951 (2010).
- [71] F. Vega Reyes, M. A. López-Castaño, and Á. Rodríguez-Rivas, Diffusive regimes in a two-dimensional chiral fluid, *Commun. Phys.* **5**, 1 (2022).
- [72] B. L. Brown, U. C. Täuber, and M. Pleimling, Effect of the Magnus force on skyrmion relaxation dynamics, *Phys. Rev. B* **97**, 020405 (2018).
- [73] C. J. O. Reichhardt and C. Reichhardt, Active rheology in odd-viscosity systems, *Europhys. Lett.* **137**, 66004 (2022).
- [74] X. Cao, D. Das, N. Windbacher, F. Ginot, M. Krüger, and C. Bechinger, Memory-induced Magnus effect, *Nat. Phys.* 10.1038/s41567-023-02213-1 (2023).
- [75] P. Welander, Note on the effect of rotation on diffusion processes, *Tellus* **18**, 63 (1966).
- [76] A. Brandenburg, A. Svedin, and G. M. Vasil, Turbulent diffusion with rotation or magnetic fields, *Mon. Not. R. Astron. Soc.* **395**, 1599 (2009).
- [77] D. L. Koch and J. F. Brady, The symmetry properties of the effective diffusivity tensor in anisotropic porous media, *Phys. Fluids* **30**, 642650 (1987).
- [78] J.-L. Auriault, C. Moyne, and H. P. Amaral Souto, On the asymmetry of the dispersion tensor in porous media, *Transport Porous Med.* **85**, 771 (2010).
- [79] S. Marbach, D. S. Dean, and L. Bocquet, Transport and dispersion across wiggling nanopores, *Nat. Phys.* **14**, 1108 (2018).
- [80] P. Wu, R. Huang, C. Tischer, A. Jonas, and E.-L. Florin, Direct measurement of the nonconservative force field generated by optical tweezers, *Phys. Rev. Lett.* **103**, 108101 (2009).
- [81] S. Sukhov and A. Dogariu, Non-conservative optical forces, *Rep. Prog. Phys.* **80**, 112001 (2017).
- [82] M. Mangeat, Y. Amarouchene, Y. Louyer, T. Guérin, and D. S. Dean, Role of nonconservative scattering forces and damping on Brownian particles in optical traps, *Phys. Rev. E* **99**, 052107 (2019).
- [83] G. Volpe, O. M. Maragò, H. Rubinsztein-Dunlop, G. Pesce, A. B. Stilgoe, *et al.*, Roadmap for optical tweezers, *J. Phys. Photonics* **5**, 022501 (2023).
- [84] F. Ghimenti, L. Berthier, G. Szamel, and F. van Wijland, Irreversible Boltzmann samplers in dense liquids: weak-coupling approximation and mode-coupling theory, *arXiv preprint arXiv:2404.14863* 10.48550/arXiv.2404.14863 (2024).
- [85] C. Kwon, P. Ao, and D. J. Thouless, Structure of stochastic dynamics near fixed points, *Proc. Natl. Acad. Sci. U.S.A.* **102**, 13029 (2005).
- [86] K. Turitsyn, M. Chertkov, V. Y. Chernyak, and A. Puliiafito, Statistics of entropy production in linearized stochastic systems, *Phys. Rev. Lett.* **98**, 180603 (2007).

- [87] C. Kwon, J. D. Noh, and H. Park, Nonequilibrium fluctuations for linear diffusion dynamics, *Phys. Rev. E* **83**, 061145 (2011).
- [88] J. D. Noh, C. Kwon, and H. Park, Multiple dynamic transitions in nonequilibrium work fluctuations, *Phys. Rev. Lett.* **111**, 130601 (2013).
- [89] J. du Buisson and H. Touchette, Dynamical large deviations of linear diffusions, *Phys. Rev. E* **107**, 054111 (2023).
- [90] H. Kählert, J. Carstensen, M. Bonitz, H. Löwen, F. Greiner, and A. Piel, Magnetizing a complex plasma without a magnetic field, *Phys. Rev. Lett.* **109**, 155003 (2012).
- [91] P. Hartmann, J. C. Reyes, E. G. Kostadinova, L. S. Matthews, T. W. Hyde, R. U. Masheyeva, K. N. Dzhumagulova, T. S. Ramazanov, T. Ott, H. Kählert, M. Bonitz, I. Korolov, and Z. Donkó, Self-diffusion in two-dimensional quasimagnetized rotating dusty plasmas, *Phys. Rev. E* **99**, 013203 (2019).
- [92] A. Shalchi, Applicability of the Taylor-Green-Kubo formula in particle diffusion theory, *Phys. Rev. E* **83**, 046402 (2011).
- [93] F. Effenberger, H. Fichtner, K. Scherer, S. Barra, J. Kleimann, and R. D. T. Strauss, A generalized diffusion tensor for fully anisotropic diffusion of energetic particles in the heliospheric magnetic field, *Astrophys. J.* **750**, 108 (2012).
- [94] A. Shalchi, Perpendicular transport of energetic particles in magnetic turbulence, *Space Sci. Rev.* **216**, 23 (2020).
- [95] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd ed. (Springer Berlin, Heidelberg, 2001).
- [96] F. Oberhettinger and L. Badii, *Tables of Laplace Transforms* (Springer Berlin, Heidelberg, 1973).
- [97] G. B. Arfken and H. J. Weber, *Mathematical Methods for Physicists*, 6th ed. (Academic Press, 2005).
- [98] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, 7th ed., edited by A. Jeffrey and D. Zwillinger (Academic Press, 2007).