An ∞ -Category of 2-Segal Spaces

Jonte Gödicke*

July 19, 2024

Abstract

Algebra objects in ∞ -categories of spans admit a description in terms of 2-Segal objects. We introduce a notion of span between 2-Segal objects and extend this correspondence to an equivalence of ∞ -categories. Additionally, for every ∞ -category with finite limits \mathcal{C} , we introduce a notion of a birelative 2-Segal object in \mathcal{C} and establish a similar equivalence with the ∞ -category of bimodule objects in spans. Examples of these concepts arise from algebraic and hermitian K-theory through the corresponding Waldhausen S_{\bullet} -construction. Apart from their categorical relevance, these concepts can be used to construct homotopy coherent representations of Hall algebras.

Contents

1	Introduction	2
	1.1 Outline	3
	1.2 Acknowledgments	3
2	2-Segal Conditions	4
	2.1 2-Segal Objects	5
	2.2 Birelative 2-Segal Objects	7
	2.3 Birelative Decomposition Spaces	12
3	Examples	14
	3.1 2-Segal Spans from Exact ∞ -Functors	15
	3.2 Relative 2-Segal Spans from Duality preserving ∞ -Functors	19
4	Modules in Span Categories	24
	4.1 Preliminaries	26
	4.2 Characterization of Bimodules	28
5	Module Morphisms in Span Categories	33
\mathbf{A}	Auxiliary Statements	42
	A.1 Proof of Proposition 4.11	42
	A.2 Proof of Proposition 4.16	46
em	*Universität Hamburg, Fachbereich Mathematik, Bundesstraße 55, 20146 Hamburg, Germ ail:jonte.goedicke@uni-hamburg.de or math@jonte-goedicke.com	ıany,

1 Introduction

A 2-Segal space, also known as a decomposition space [GCKT18], is a simplicial space $X_{\bullet} : \Delta^{op} \to S$ that satisfies a 2-dimensional analog of the famous Segal conditions introduced in [Seg68]. These concepts were initially introduced independently by Dyckerhoff and Kapranov [DK19] to describe the underlying structure of different Hall algebra constructions, and by Gálvez-Carrillo, Kock and Tonks to describe various coalgebra structures appearing in combinatorics [GCKT18]. Since then, the subject has undergone substantial developments, including the introduction of even higher Segal conditions [Pog17] to applications in different areas of mathematics like the theory of TFT's [DK18], Auslander-Reiten theory [DJW19] and Donaldson-Thomas theory [PS23].

The connection between 2-Segal spaces and Hall algebras arises from the algebraic interpretation of 2-Segal spaces. In [Ste21], it is shown that the space of 2-Segal spaces is equivalent to the space of algebra objects in the ∞ -category of spans of spaces. However, the appropriate notion of morphism between 2-Segal spaces, which under this equivalence corresponds to a morphism of algebras, has not been determined so far.

An initial proposal might be to define the corresponding notion of morphism between 2-Segal spaces as a morphism of simplicial spaces and thus to define the ∞ -category of 2-Segal spaces as a full subcategory of $\mathsf{Fun}(\Delta^{op}, \mathbb{S})$. Although this yields an interesting notion of morphism between 2-Segal spaces, it does not correspond to an algebra morphism. This fails for different reasons. On the one hand, a general morphism of simplicial spaces only corresponds to a lax algebra morphism [Wal16, Sect.4.2]. On the other hand, since the algebraic structure itself lives in an ∞ -category of spans an algebra morphism should also be formalized in this context. These considerations lead to the main result of this text:

Theorem 1.1. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence of ∞ -categories

 $2\text{-}\mathrm{Seg}^{\leftrightarrow}_{\Delta}(\mathcal{C}) \xrightarrow{\simeq} \mathsf{Alg}(\mathsf{Span}(\mathcal{C}^{\times}))$

between the subcategory of the ∞ -category Span(Fun(Δ^{op}, \mathbb{C})) of spans of simplicial objects with objects 2-Segal object and morphisms 2-Segal spans and the ∞ -category of algebra objects in Span(\mathbb{C}^{\times}).

A span between 2-Segal objects is called a 2-Segal span if the legs of the span satisfy conditions that were previously introduced under the names CULF and IKEO in [GCKT18]. There, the authors show that such morphisms induce algebra morphisms. The novel aspect of Theorem 1.1 is that every algebra morphism is in fact induced by spans of CULF and IKEO morphisms.

Analogous to the case of algebras, we introduce a multicolored version of the 2-Segal conditions on birelative simplicial spaces $X_{\bullet} : \Delta_{/[1]}^{op} \to S$. Using this, we extend the previous result to the case of bimodules:

Theorem 1.2. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence of ∞ -categories

$$\operatorname{BiSeg}_{\Delta}^{\leftrightarrow}(\operatorname{\mathcal{C}}) \xrightarrow{\simeq} \operatorname{BiMod}(\operatorname{Span}(\operatorname{\mathcal{C}}^{\times}))$$

between the subcategory of Span(Fun($\Delta_{/[1]}^{op}, \mathbb{C}$)) with objects birelative 2-Segal objects and with morphism birelative 2-Segal spans and the ∞ -category of bimodule objects in Span(\mathbb{C}^{\times}).

Similar conditions have been previously studied for left modules in the context of 1-categories in both [Wal16] and [You18], and for bimodules in terms of double Segal spaces in [Car20]. Our result generalizes these works into the context of ∞ -categories.

The prototypical example of a 2-Segal space is the Waldhausen S_{\bullet} -construction of an exact ∞ -category \mathcal{C} . It is thus reasonable to expect exact ∞ -functors $F : \mathcal{C} \to \mathcal{D}$ to induce algebra morphism between those. Indeed, we show that certain classes of exact ∞ -functors induce algebra morphisms between Waldhausen S_{\bullet} -constructions. Other examples arise from the relative Waldhausen S_{\bullet} -construction of an exact ∞ -functor F. This simplicial space fits into a sequence of simplicial spaces:

$$\mathcal{D}^{\simeq}_{\bullet} \xrightarrow{\iota_{\bullet}} S^{rel}_{\bullet}(F) \xrightarrow{\pi_{\bullet}} S_{\bullet}(\mathcal{C}) \tag{1}$$

which has first appeared in [Wal06] to construct the long exact sequence of relative K-theory. We show that the morphisms ι_{\bullet} and π_{\bullet} are indeed algebra morphisms. These morphisms underlie the construction of derived Hecke actions on cohomological Hall algebras [KV22].

For the construction of examples of modules, we consider 2-Segal spaces X_{\bullet} equipped with a twisted C_2 -action. Examples of these arise in hermitian K-theory as the Waldhausen S_{\bullet} -construction of an exact ∞ -category with duality. For every such 2-Segal space with a twisted C_2 action, the canonical morphism $Tw(X_{\bullet}) \to X_{\bullet} \times X_{\bullet}^{ev}$ is a relative 2-Segal object with C_2 -action. Taking homotopy fixed points yields:

Proposition 1.3. There exists an ∞ -functor from the ∞ -category of 2-Segal spaces with duality to the ∞ -category of relative 2-Segal spaces

$$\mathsf{Tw}^{\leftrightarrow}(-)^{\mathsf{C}_2}: 2\text{-}\mathrm{Seg}^{\leftrightarrow}_{\Delta}(\mathbb{S}) \to \mathrm{Rel2Seg}^{\leftrightarrow}_{\Delta}(\mathbb{S})$$

that maps a simplicial space with duality X_{\bullet} to the relative 2-Segal space $\mathsf{Tw}(X_{\bullet})^{\mathsf{C}_2} \to X_{\bullet}$.

In the example of the Waldhausen S_{\bullet} -construction of an exact ∞ -category with duality \mathcal{C} the simplicial space $\mathsf{Tw}(S_{\bullet}(\mathcal{C}))^{\mathsf{C}_2}$ is known as the \mathcal{R}_{\bullet} -construction [HS04] of \mathcal{C} . Consequently, the above Proposition expresses $\mathcal{R}_{\bullet}(\mathcal{C})$ as a module over $S_{\bullet}(\mathcal{C})$. This generalizes results of [You18] for proto-exact categories with duality into the ∞ -categorical context. Moreover, using the functoriality of this construction, we obtain examples of module morphisms from exact duality preserving functors between exact ∞ -categories. These results have applications to the construction of cohomological Hall algebra representations [DPS22] and orientifold Donaldson-Thomas-theory [You20].

1.1 Outline

We conclude this introduction with an outline of the structure of this paper. In Section 2, we define the ∞ -categories BiSeg $_{\Delta}^{\leftrightarrow}(S)$ and 2-Seg $_{\Delta}^{\leftrightarrow}(S)$. Therefore, we first recall in Subsection 2.1 the definition of 2-Segal objects and introduce the notion of 2-Segal spans. Then, in Subsection 2.2 we extend the definitions to the case of birelative simplicial objects and introduce the birelative 2-Segal conditions. Similarly to the 2-Segal case, we introduce a version of the decomposition space conditions in Subsection 2.3. We show that these are equivalent to the birelative 2-Segal conditions.

In Section 3, we discuss examples of the notions introduced in Section 2 that naturally appear in the context of algebraic and hermitian K-theory. To construct examples of modules, we construct an ∞ -functor from the ∞ -category of 2-Segal spaces with duality to the ∞ -category of relative 2-Segal objects. As a main application of this, we express the \mathcal{R}_{\bullet} -construction as a module over the S_{\bullet} -construction.

In Section 4 we prove the first half of our main theorem and characterize the space bimodule objects in span-categories in terms of birelative 2-Segal objects. To increase the level of readability, we have included some combinatorial lemmas into the Appendix A.

In Section 5 we finally prove our main theorem and extend the equivalence of spaces constructed in Section 4 to an equivalence of ∞ -categories. As a corollary, we obtain similar characterizations for the ∞ -category of algebra, left and right module objects.

1.2 Acknowledgments

I wish to thank my supervisor Tobias Dyckerhoff for his support and encouragement. I further thank Angus Rush for many discussions about spans and Joachim Kock for interesting disussions about the content of this text. The author is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC 2121 Quantum Universe" - 390833306 and the Collaborative Research Center - SFB 1624 Higher structures, moduli spaces and integrability" - 506632645.

2 2-Segal Conditions

Throughout this section, \mathcal{C} denotes an ∞ -category with finite limits. In this section, we introduce a variety of 2-*Segal type* conditions. These admit an interpretation in terms of homotopy coherent algebra in span categories. Before stating these definitions, let us motivate them for the example of the 2-Segal condition.

There exists an ∞ -category of spans Span(\mathcal{C}), whose objects are the objects of \mathcal{C} and whose 1-morphisms are spans of morphisms in \mathcal{C} . It further inherits a monoidal structure from the Cartesian product on \mathcal{C} . The datum of a coherent algebra object in Span(\mathcal{C}) consists

- an underlying object $X_1 \in \mathcal{C}$
- a multiplication span



• homotopy coherent associativity data

As the notations suggest this data organizes into a simplicial object X_{\bullet} . The higher simplices are encoded in the associativity data. Its lowest instance is an associativity 2-isomorphism

$$\alpha: X_2 \times_{X_1} X_2 \xrightarrow{\simeq} X_2 \times_{X_1} X_2$$

that makes the diagram:

2

$$\begin{array}{c} X_3 \\ (\partial_0,\partial_2) \\ X_2 \times X_1 \\ X_1 \times X_2 \\ X_1 \times X_1 \\ X_1 \times$$

commutes. One checks that the data of α is equivalent to an object X_3 and isomorphisms:

$$X_2 \times_{X_1} X_2 \xleftarrow{(\partial_0, \partial_2)} X_3 \xrightarrow{(\partial_3, \partial_1)} X_2 \times_{X_1} X_2 \tag{2}$$

s.t. the components ∂_i satisfy certain simplicial identities. Equation 2 is the lowest instance of the 2-Segal conditions (see Definition 2.1). Analogously the higher dimensional 2-Segal conditions encode the higher associativity of the algebra.

We now summarize the content of this section. In 2.1 we give the definitions of 2-Segal objects and 2-Segal spans. They assemble into an ∞ -category 2-Seg $_{\Delta}^{\leftrightarrow}(\mathbb{C})$ that is our candidate for the ∞ -category of 2-Segal objects. Afterwards we extend the discussion in 2.2 and introduce 2-Segal and Segal span conditions for the indexing categories $\Delta_{/[1]}^{op}$, Δ_{\geq}^{op} and Δ_{\leq}^{op} . We show in Section 4 that these admit an interpretation in terms of bi-, left, and right module objects in Span(\mathbb{C}) respectively. Finally, we provide in Subsection 2.3 a different characterization of these new 2-Segal conditions in terms of active-inert pullbacks. This generalizes the equivalence between 2-Segal and decomposition spaces [GCKT18] to this class of indexing categories.

2.1 2-Segal Objects

We frequently abuse notation and identify a simplicial object $X_{\bullet} : \Delta^{op} \to \mathbb{C}$ with its extension to $\operatorname{Fin}_{\geq}$ the category of all finite non empty linearly ordered sets. This will simplify the notation. Let us recall as a start the definition of a 2-Segal object.

Definition 2.1. [DK19, Def.2.3.1] Let $X_{\bullet} : \Delta^{op} \to \mathbb{C}$ be a simplicial object. X_{\bullet} is called 2-Segal, if for every $n \ge 3$ and $0 \le i < j \le n$ the map

$$X_n \to X_{\{0,1,\dots,i,j,j+1,\dots,n\}} \times_{X_{\{i,j\}}} X_{\{i,i+1,\dots,j\}}$$

is an equivalence. We denote by 2- Seg_{Δ}(\mathcal{C}) the full subcategory of Fun(Δ^{op} , \mathcal{C}) on 2-Segal objects.

The morphisms in the ∞ -category 2- Seg_{Δ}(\mathcal{C}) are morphisms of simplicial objects. Under the equivalence between 2-Segal objects and algebra objects in spans a general simplicial map between 2-Segal objects induces a lax-algebra morphism between the corresponding algebra objects [Wal16, Sect.4.2]. Let's sort out, what a strong algebra morphism looks like in terms of 2-Segal objects:

Definition 2.2. Let $\sigma : \Delta^1 \to \text{Span}(2\text{-}\text{Seg}_{\Delta}(\mathbb{C}))$ be a 1-morphism in the ∞ -category of spans of 2-Segal objects.¹ The morphism σ can be represented by a span



with X_{\bullet}, Y_{\bullet} and $M_{\bullet} : \Delta^{op} \to \mathbb{C}$ 2-Segal. The morphism σ is called a 2-Segal span, if for every $n \ge 0$ the induced diagrams



and

$$Y_n \underbrace{}_{\substack{i \neq 0 \\ i = 0}} Y_{\{i, i+1\}} \underbrace{}_{\substack{i = 0 \\ i = 0}} M_{\{i, i+1\}}$$

are pullback diagrams. We call s an active equifibered Δ^{op} -morphism and t a relative Segal Δ^{op} -morphism.

Notation 2.3. We introduce in Subsection 2.2 similar notions for other indexing categories than Δ . If the indexing category is clear from the context, we frequently abuse notation and drop the indexing category in the notation. E.g we will call an active equifibered Δ^{op} -morphism simply an active equifibered morphism when the indexing category is clear from the context.

Remark 2.4. The conditions imposed on the individual legs have previously been studied under the name CULF [GCKT18, Sect.4] and IKEO [GCKT18, Sect.8.5]. The authors further demonstrate that these types of morphisms induce algebra morphisms in spans. Our notation is motivated by the theory of algebraic patterns [BHS22].

¹See Construction 4.8

It is worth noting that in the definition of a 2-Segal span, we explicitly require the tip of the span to be 2-Segal as well. This is not an additional requirement. Indeed as demonstrated in [GCKT18, Lem.4.6], the source of an active equifibered morphism $f: X_{\bullet} \to Y_{\bullet}$ with target 2-Segal is itself 2-Segal. In the following, we interpret 2-Segal spans as the morphisms of a category and therefore have to study

their behavior under composition of spans:

Lemma 2.5. Let $\sigma : \Lambda_1^2 \to \text{Span}(2\text{-}\text{Seg}_{\Delta}(\mathcal{C}))$ be given by a composable pair of spans



s.t. each individual span is a 2-Segal span. Then also the composite span



is a 2-Segal span.

Proof. First, we show that s is active equifibered. For $n \neq 0$ consider the diagram:



We need to prove that the exterior diagram is a pullback diagram. By assumption, the lower square is a pullback diagram. Additionally, the upper square is a pullback square if and only if the outer rectangle in the diagram



is a pullback square. But this follows from the pasting law. On the other hand, d we show that t is relative Segal. We need to show that for every $n \ge 0$, the exterior rectangle in the diagram:



is a pullback. By assumption, the left square is a pullback square. Further, the right square is a pullback square if and only if the outer rectangle in the diagram:



is a pullback square. But this is again a consequence of the pasting law.

Definition 2.6. We define the ∞ -category 2-Seg $_{\Delta}^{\leftrightarrow}(\mathbb{C})$ of 2-Segal objects as the wide subcategory of Span(2-Seg $_{\Delta}(\mathbb{C})$) with morphisms given by 2-Segal spans.

2.2 Birelative 2-Segal Objects

In this subsection, we extend the framework of the previous paragraph to categories of functors with source $\Delta_{/[1]}$. We call such a functor $M_{\bullet}: \Delta_{/[1]}^{op} \to \mathcal{C}$ a birelative simplicial object. The corresponding birelative 2-Segal condition is a multi-coloured version of the 2-Segal condition from Definition 2.1:

Definition 2.7. Let $M_{\bullet} : \Delta_{[1]}^{op} \to \mathbb{C}$ be a birelative simplicial object. M_{\bullet} is called *birelative 2-Segal*, if for every $n \ge 3$, $f : [n] \to [1]$ and $0 \le i < j \le n$ the diagram



is Cartesian. We denote the full subcategory of $\operatorname{Fun}(\Delta_{/[1]}^{op}, \mathbb{C})$ generated by the birelative 2-Segal objects by Bi2Seg_{Δ}(\mathbb{C}).

As 2-Segal objects encode algebra objects, birelative 2-Segal objects encode bimodule objects. Let us also introduce a 2-Segal type condition for left and right modules. These already have been studied in the case of 1-categories in [Wal16] and [You18]. Denote by Δ_{\leq} (resp. Δ_{\geq}) the full subcategory of $\Delta_{/[1]}$ generated by those objects $f : [n] \rightarrow [1]$ that take the value 1 (resp. 0) at most ones. We call a functor with source Δ_{\leq} (resp. Δ_{\geq}) a left (resp. right) relative simplicial object. The corresponding 2-Segal condition reads as:

Definition 2.8. Let $M_{\bullet} : \Delta_{\leqslant}^{op} \to \mathbb{C}$ be a left relative simplicial object. M_{\bullet} is called *left relative 2-Segal*, if for every $n \ge 3$, $f : [n] \to [1] \in \Delta_{\leqslant}$ and $0 \le i < j \le n$ the diagram



is Cartesian. We denote the full subcategory of $\operatorname{Fun}(\Delta^{op}_{\leq}, \mathbb{C})$ generated by the left relative 2-Segal objects by $\operatorname{L2Seg}_{\Delta}(\mathbb{C})$. Similarly we define *right relative 2-Segal objects* and the ∞ -category of right relative 2-Segal objects $\operatorname{R2Seg}_{\Delta}(\mathbb{C})$.

Remark 2.9. The categories Δ_{\leq} and Δ_{\geq} defined above are equivalent to the categories, denoted with the same symbols, as defined in [Wal16, Def.3.2.5]. Further, our notions of left and right relative 2-Segal object coincide with the notion of a relative 2-Segal object in [Wal21, Def.3.5.1.].

To handle the cases of bi-, left and right relative 2-Segal conditions at once, we introduce the following notation:

Notation 2.10. For $\# \in \{/[1], \leq, \geq\}$ we call a functor $M_{\bullet} : \Delta^{op}_{\#} \to \mathbb{C}$ a #-relative simplicial object. Further, we call a #-relative simplicial object a #-relative 2-Segal object if it satisfies the corresponding 2-Segal conditions. We denote the ∞ -category of #-relative 2-Segal objects by #-2Seg_{Δ}(\mathbb{C}).

Example 2.11. Let $X_{\bullet} : \Delta^{op} \to \mathbb{C}$ be 2-Segal. We can associate to X_{\bullet} a birelative, left and right relative 2-Segal object

- (1) $X^b_{\bullet}: \Delta^{op}_{/[1]} \to \Delta^{op} \to \mathcal{C}$
- (2) $X^l_{\bullet} : \Delta^{op}_{\leq} \to \Delta^{op} \to \mathcal{C}$

$$(3) \ X^r_{\bullet}: \Delta^{op}_{\geqslant} \to \Delta^{op} \to \mathcal{C}$$

via precomposition with the corresponding forgetful functor. It follows from the 2-Segal conditions for X_{\bullet} , that these objects fulfill the respective (bi)relative Segal conditions. Under the equivalence of Corollary 5.2 and Theorem 5.1 these correspond to the regular bimodule (resp. left, right module) associated to the algebra object corresponding to X_{\bullet} .

We also introduce, in analogy to Definition 2.2, a notion of morphism between #-relative 2-Segal objects:

Definition 2.12. Let A_{\bullet}, B_{\bullet} and M_{\bullet} be #-relative 2-Segal. A #-relative 2-Segal span from A_{\bullet} to B_{\bullet} is given by a span



s.t. for every $n \ge 0$ and $f: [n] \to [1] \in \Delta_{\#}$ the diagrams



and



are pullback diagrams. We call s an active equifibered $\Delta^{op}_{\#}$ -morphism and t a relative Segal $\Delta^{op}_{\#}$ -morphism.

Analogously to Lemma 2.5 one can prove the following:

Lemma 2.13. Let $\sigma : \Lambda_1^2 \to \text{Span}(\#\text{-}2\text{Seg}_{\Delta}(\mathcal{C}))$ be given by a composable pair of spans



s.t. the individual spans are #-relative 2-Segal spans. Then also the composite span



is a #-relative 2-Segal span.

Proof. Similar to Lemma 2.5.

Definition 2.14. We define the ∞ -category Bi2Seg $_{\Delta}^{\leftrightarrow}(\mathbb{C})$ of birelative 2-Segal objects to be the wide subcategory of Span(Bi2Seg $_{\Delta}(\mathbb{C})$) with morphisms birelative 2-Segal spans.

Similarly we define the ∞ -category of left (resp. right) relative 2-Segal objects $L2Seg_{\Delta}^{\leftrightarrow}(\mathcal{C})$ (resp. $R2Seg_{\Delta}^{\leftrightarrow}(\mathcal{C})$)) as the wide subcategory of $Span(L2Seg_{\Delta}(\mathcal{C}))$ (resp. $Span(R2Seg_{\Delta}(\mathcal{C}))$) with morphisms given by left (resp. right) relative 2-Segal spans.

It has been shown in [Wal16, Prop.3.5.10] for 1-categories that left and right relative 2-Segal objects admit a different description in terms of morphisms of simplicial objects called relative 2-Segal objects. This description is useful in the study of examples. Let us therefore lift this description into the realm of ∞ -categories. We analyze the case of left relative 2-Segal objects. The case of right relative 2-Segal objects is analogous.

Definition 2.15. [You18, Def.2.2][Wal16, Prop.3.5.10] A morphism $\pi : X_{\bullet} \to Y_{\bullet}$ of simplicial objects is called *relative 2-Segal*, if

- 1 the source object X_{\bullet} is 1-Segal
- 2 the target object Y_{\bullet} is 2-Segal
- 3 for every $0 \leq i < j \leq n$ the following squares are Cartesian



We denote the full subcategory of $\operatorname{Fun}(\Delta^{0,p}, \mathbb{C})$ generated by relative 2-Segal morphisms by $\operatorname{Rel2Seg}_{\Delta}(\mathbb{C})$.

We also introduce a notion of morphism between relative 2-Segal objects:

Definition 2.16. Let $\pi_{\bullet}^i : X_{\bullet}^i \to Y_{\bullet}^i$ for $0 \le i \le 2$ be relative 2-Segal objects. A relative 2-Segal span from π_{\bullet}^0 to π_{\bullet}^2 is given by a span

$$\begin{array}{cccc} X^{0}_{\bullet} & \stackrel{s^{X}_{\bullet}}{\longleftarrow} & X^{1}_{\bullet} & \stackrel{t^{X}_{\bullet}}{\longrightarrow} & X^{2}_{\bullet} \\ \pi^{0}_{\bullet} & & & & \downarrow \pi^{1}_{\bullet} & & \downarrow \pi^{2}_{\bullet} \\ Y^{0}_{\bullet} & \stackrel{s^{Y}_{\bullet}}{\longleftarrow} & Y^{1}_{\bullet} & \stackrel{t^{Y}_{\bullet}}{\longrightarrow} & Y^{2}_{\bullet} \end{array}$$

of relative 2-Segal objects s.t.

(1) the span $Y^0_{\bullet} \xleftarrow{s^Y_{\bullet}} Y^1_{\bullet} \xrightarrow{t^Y_{\bullet}} Y^2_{\bullet}$ is 2-Segal

(2) For every $n \ge 1$ the square

$$\begin{array}{ccc} X_n^1 & \xrightarrow{s_n^X} & X_n^0 \\ \downarrow & & \downarrow \\ X_{\{0,n\}}^1 & \xrightarrow{s_1^X} & X_{\{0,n\}}^0 \end{array}$$

is Cartesian.

(3) For every $n \ge 1$ the diagram

$$\begin{array}{cccc} X_n^1 & & \underbrace{t_n^X} & & & X_n^2 \\ & & \downarrow & & \downarrow \\ Y_{\{0,1\}}^1 \times \cdots \times Y_{\{n-2,n-1\}}^1 \times X_{\{n-1,n\}}^1 & \longrightarrow Y_{\{0,1\}}^2 \times \cdots \times Y_{\{n-2,n-1\}}^2 \times X_{\{n-1,n\}}^2 \end{array}$$

is Cartesian.

We call the pair $(s^X_{\bullet}, s^Y_{\bullet})$ active equifibered and the pair $(t^X_{\bullet}, t^Y_{\bullet})$ relative Segal.

Definition 2.17. We define the the ∞ -category of relative 2-Segal objects as the wide subcategory Rel2Seg $_{\Delta}^{\leftrightarrow}(\mathbb{C})$ of the ∞ -category Span(Rel2Seg $_{\Delta}(\mathbb{C})$) with morphisms given by relative 2-Segal spans.

We now turn to the construction of an equivalence $\Theta_L^{\leftrightarrow} : L2Seg_{\Delta}^{\leftrightarrow}(\mathcal{C}) \xrightarrow{\simeq} Rel2Seg_{\Delta}^{\leftrightarrow}$. To do so, we construct an equivalence

 $\Theta_L : L2Seg_{\Delta}(\mathcal{C}) \xrightarrow{\simeq} Rel2Seg_{\Delta}(\mathcal{C})$

that induces $\Theta_L^{\leftrightarrow}$ on ∞ -categories of spans.

For the construction of Θ_L we use observation [Wal16, Rem.3.2.8]. First note that the category Δ_{\leq} contains two copies of the simplex category as full subcategories. We denote the corresponding fully faithful inclusion by $i_0^{\leq} : \Delta \to \Delta_{\leq}$ and $i_1^{\leq} : \Delta \to \Delta_{\leq}$. These are given on objects by

$$\begin{split} &i_0^\leqslant([n])(k) = 0 \quad \forall k \in [n] \\ &i_1^\leqslant([n])(k) = \begin{cases} 0 \quad \forall k \neq n+1 \\ 1 \quad k = n+1 \end{cases} \end{split}$$

Further, we observe that the morphisms $d_{n+1}: i_0^{\leq}([n]) \to i_1^{\leq}([n])$ assemble into a natural transformation $d_{\bullet+1}: i_0^{\leq}(-) \Rightarrow i_1^{\leq}(-)$. This datum is equivalent to the datum of a lax cocone



and therefore induces a unique map from the lax colimit. It follows from [GHN17, Thm.7.4] that the lax colimit is given by the total space of the cocartesian Grothendieck construction $\int_{\Delta^1} i d_{\Delta} \to \Delta$, where we identify the functor $i d_{\Delta} : \Delta \to \Delta$ with a morphism in Cat_{∞} . The morphism

$$\theta_L: \int_{\Delta^1} i d_\Delta \to \Delta_\leqslant$$

induced by Diagram 3 is given on the fiber over 0 by $i_0^{\leq} : \Delta \to \Delta_{\leq}$, on the fiber over 1 by $i_1^{\leq} : \Delta \to \Delta_{\leq}$ and on morphisms $f : ([n], 0) \to ([m], 1)$ by

$$i_0^{\leqslant}([n]) \xrightarrow{i_0^{\leqslant}(f)} i_0^{\leqslant}(m) \xrightarrow{d_{m+1}} i_1^{\leqslant}(m)$$

It follows from the explicit description that this functor is an equivalence.

Proposition 2.18. The functor $\Theta_L := \theta_L^* : \operatorname{Fun}(\Delta_{\leqslant}^{op}, \mathbb{C}) \to \operatorname{Fun}(\int_{\Delta^1} id_{\Delta}^{op}, \mathbb{C})$ restricts to an equivalence of ∞ -categories

$$\Theta_L : \mathrm{L2Seg}_{\Delta}(\mathcal{C}) \to \mathrm{Rel2Seg}_{\Delta}(\mathcal{C})$$

Proof. It follows from the universal property of the lax colimit that we have an equivalence of ∞ -categories

$$\mathsf{Fun}(\int_{\Delta^1} id^{op}_\Delta, \mathfrak{C}) \simeq \mathsf{Fun}(\Delta^1 \times \Delta^{op}, \mathfrak{C}) \simeq \mathsf{Fun}(\Delta^1, \mathsf{Fun}(\Delta^{op}, \mathfrak{C}))$$

Under this equivalence the functor Θ_L maps a left relative simplicial object $X_{\bullet} : \Delta_{\leqslant}^{op} \to \mathbb{C}$ to the natural transformation

$$(i_1^\leqslant)^*X_\bullet \xrightarrow{\partial_{n+1}} (i_0^\leqslant)^*X_\bullet$$

between simplicial objects. The claim follows from the observation that the functor Θ_L maps left relative 2-Segal objects to relative 2-Segal objects.

It follows from the discussion after Construction 4.8 that the ∞ -category of spans is functorial:

$$\mathsf{Span}(-): \mathsf{Cat}_{\infty}^{\mathsf{lex}} \to \mathsf{Cat}_{\infty}$$

We use this functor to conclude our comparison:

Corollary 2.19. The functor $\Theta_L^{\leftrightarrow} := \text{Span}(\Theta_L) : \text{Span}(\text{L2Seg}_{\Delta}(\mathcal{C})) \to \text{Span}(\text{Rel2Seg}_{\Delta}(\mathcal{C}))$ restricts to an equivalence of ∞ -categories

$$\Theta_L^{\leftrightarrow} : \mathrm{L2Seg}_{\Delta}^{\leftrightarrow}(\mathcal{C}) \to \mathrm{Rel2Seg}_{\Delta}^{\leftrightarrow}(\mathcal{C})$$

Proof. It suffices to show that $\Theta_L^{\leftrightarrow}$ maps left relative 2-Segal spans to relative 2-Segal spans. But this follows from unraveling the definitions.

Remark 2.20. Analogously to the proof of Corollary 2.19 one can construct an equivalence

$$\Theta_R^{\leftrightarrow} : \operatorname{R2Seg}_{\Delta}^{\leftrightarrow}(\mathcal{C}) \xrightarrow{\simeq} \operatorname{Rel2Seg}_{\Delta}^{\leftrightarrow}(\mathcal{C})$$

between the ∞ -category of right relative 2-Segal objects and relative 2-Segal objects.

We use Corollary 2.19 for the construction of examples of left relative 2-Segal objects and spans. This is particularly fruitful for the example of the hermitian Waldhausen construction that we discuss in Subsection 3.2.

Example 2.21. Let X_{\bullet} be 2-Segal. Denote by $P^{\lhd}(X)_{\bullet}$ the initial and by $P^{\rhd}(X)_{\bullet}$ the final path space [DK19, Section 6.2]. These come equipped with morphisms of simplicial sets $p_{\lhd} : P^{\lhd}(X)_{\bullet} \to X_{\bullet}$ and $p_{\rhd} : P^{\rhd}(X)_{\bullet} \to X_{\bullet}$. It follows from [DK19, Thm.6.3.2] that $P^{\lhd}(X)_{\bullet}$ is 1-Segal. The morphisms p_{\lhd} and p_{\triangleright} are examples of relative 2-Segal objects. These describe the regular left and right modules. Under the equivalence of Corollary 2.19 these get identified with the left and right relative 2-Segal objects from Example 2.11.

Example 2.22. Let X_{\bullet} be a 2-Segal object in \mathcal{C} . Recall that the edgewise subdivision functor $e : \Delta \to \Delta$ is defined on objects as $[n] \mapsto [n] * [n]^{op}$. Precomposition with e induces a functor

$$\mathsf{Tw}(-) := e^* : \mathsf{Fun}(\Delta^{op}, \mathfrak{C}) \to \mathsf{Fun}(\Delta^{op}, \mathfrak{C})$$

For a simplicial object X_{\bullet} , we call $\mathsf{Tw}(X)_{\bullet}$ the twisted arrow simplicial object. It follows from [BOO+20, Thm.2.9], that for X_{\bullet} 2-Segal the simplicial object $\mathsf{Tw}(X)_{\bullet}$ is 1-Segal. Moreover, it is easy to check that the morphism

$$\mathsf{Tw}(X)_{\bullet} \to X_{\bullet} \times X_{\bullet}^{op}$$

defines a relative 2-Segal object. This encodes the canonical bimodule action of X_{\bullet} on itself in terms of left modules.

2.3 Birelative Decomposition Spaces

An alternative but equivalent way to formalize the 2-Segal conditions are the decomposition space conditions of [GCKT18]. For the definition of these recall that the simplex category Δ admits a factorization system ($\Delta^{act}, \Delta^{int}$) generated by the active (depicted \rightarrow) and inert morphisms (depicted \rightarrow) [Lur17]. This factorization system has the special property that Δ admits active-inert-pushouts, i.e every cospan

$$\begin{array}{c}n \xrightarrow{f} m \\ \downarrow^{g} \\ \downarrow^{g} \\ l\end{array}$$

with f inert and g active admits an extension to a pushout diagram:

$$\begin{array}{ccc} n \longmapsto m \\ \downarrow & \downarrow \\ l \longmapsto k \end{array}$$

in Δ . A simplicial space $X_{\bullet} : \Delta^{op} \to S$ is then called a decomposition space if it maps active-inert pullbacks in Δ^{op} to pullback diagrams of spaces. The goal of this section is to extend the definition of a decomposition space to the category $\Delta_{/[1]}$. Therefore we first recall the definition of the active-inert factorization system on $\Delta_{/[1]}$

Notation 2.23. Given an object $f : [n] \to [1] \in \Delta_{/[1]}$, we will frequently denote it by f_n to indicate its dependence on the source. Given i < j in [n] we denote by $[i, j] := \{i, ..., j\} \subset [n]$ the subinterval from i to j and by $f_{[i,j]}$ the restriction of f_n to this subinterval.

Definition 2.24. A morphism $f: g_n^0 \to g_m^1$ in $\Delta_{/[1]}$ is called

- (1) active if the underlying morphism of linearly ordered sets $f : [n] \to [m]$ is endpoint preserving. We depict active morphisms by \twoheadrightarrow .
- (2) *inert* if the underlying morphism of linearly ordered sets $f : [n] \to [m]$ is a subinterval inclusion. We depict inert morphisms by \rightarrow

We denote the wide subcategory of $\Delta_{/[1]}$ spanned by the active (resp. inert) morphisms by $\Delta_{/[1]}^{act}$ (resp. $\Delta_{/[1]}^{int}$). Similarly, we call a morphism f in $\Delta_{/[1]}^{op}$ active (resp. inert) if it is active (reps. inert) as a morphism in $\Delta_{/[1]}$.

Proposition 2.25. Every morphism $f : g_n^0 \to g_m^1$ in $\Delta_{/[1]}$ admits an unique factorization $f = f'' \circ f'$ with f'' inert and f' active. In particular the subcategories $(\Delta_{/[1]}^{act}, \Delta_{/[1]}^{int})$ form a factorization system on $\Delta_{/[1]}$ [Lur06, Def.5.2.8.8].

Proof. This is a consequence of the dual of [Lur17, Prop.2.1.2.5] applied to the Cartesian fibration $\Delta_{/[1]} \rightarrow \Delta$. Alternatively one can see this more directly as follows. The underlying morphism of linearly ordered sets $f : [n] \rightarrow [m]$ admits an active-inert factorization given by the active map $f' : [n] \rightarrow [f(0), f(n)]$ and the inert map $f'' : [f(0), f(n)] \hookrightarrow [m]$. These morphisms admit unique extensions to morphisms in $\Delta_{/[1]}$.

To characterize the active-inert pullback diagrams in $\Delta_{/[1]}^{op}$, we need the following definition:

Definition 2.26. Let $g : [n] \to [1]$ and $f : [m] \to [1]$ be two objects in $\Delta_{/[1]}$. We will call g and f composable, if g(n) = f(0). The concatenation of g with f denoted $g \star f : [n + m] \to [1]$ is defined as the map:

$$(g * f)(k) = \begin{cases} g(k) & k \le n \\ f(k-n) & k \ge n \end{cases}$$

Similarly, we denote for two linearly ordered sets S, T by S * T the partially ordered set $S \coprod T/(s_{max} \sim t_{min})$, where s_{max} denotes the maximal element of S and t_{min} denotes the minimal element of T.

Given an active morphism $f^2 : h_k^0 \to h_l^1$ and an inert morphism $e^1 : g_n^0 \to h_k^0$ in $\Delta_{/[1]}$ we can consider an active-inert factorization of its composite $f^2 \circ e^1$. This factorization can be organized into a commutative square



It follows from the definition of active and inert morphisms, that this square is equivalent to the square

In analogy with [GCKT18, Sect.2.6], we call the squares of the form:

$$\begin{array}{c} g_n^0 \xrightarrow{f^1} & & & g_m^1 \\ & & & & \downarrow \\ & & & & \downarrow \\ h_a^0 \ast g_n^0 \ast h_b^0 \xrightarrow{}_{\mathsf{id} \ast f^1 \ast \mathsf{id}} & & & h_a^0 \ast g_m^1 \ast h_b^0 \end{array}$$

identity extension squares. These are precisely the active-inert pullbacks in $\Delta_{/[1]}^{op}$. **Proposition 2.27.** Let $\sigma : \Delta^1 \times \Delta^1 \to \Delta_{/[1]}$ be an identity extension square:

Then σ is a pushout square.

In analogy with [GCKT18], we introduce the following definition:

Definition 2.28. Let $X_{\bullet} : \Delta_{/[1]}^{op} \to \mathbb{C}$ be a birelative simplicial object. X_{\bullet} is called a *birelative decomposition space* if it sends every active-inert pullback in $\Delta_{/[1]}^{op}$ to a pullback square in \mathbb{C} , i.e

As in the case of 2-Segal spaces [GCKT18], we show that the birelative decomposition space condition is equivalent to the birelative 2-Segal condition:

Proposition 2.29. Let $X_{\bullet}: \Delta_{/[1]}^{op} \to \mathbb{C}$ be a birelative simplicial object. The following are equivalent:

- (1) X_{\bullet} is birelative 2-Segal
- (2) X_{\bullet} is a birelative decomposition space

Proof. Assume first that X_{\bullet} is a birelative decomposition space. Let $f : [n] \to [1]$ with $n \ge 3$ be an object of $\Delta_{/[1]}$ and $0 \le i < j \le n$. Consider the square



in $\Delta_{/[1]}$. Note that X_{\bullet} satisfies the birelative 2-Segal conditions if and only if it maps each such square to a pullback square. But this square is an identity extension square in $\Delta_{/[1]}$ and hence an active-inert pullback.

Conversely assume that X_{\bullet} is birelative 2-Segal. We need to show that the image under X_{\bullet} of every identity extension square



is a pullback diagram. We can extend every such to a rectangle:

$$\begin{array}{c} g^0_{\{0,n\}} = g^1_{\{0,m\}} & \longrightarrow & g^0_n & \stackrel{f^1}{\longrightarrow} & g^1_m \\ & & & \downarrow & & \downarrow \\ & & & \downarrow & & \downarrow \\ h^0_a \ast g^1_{\{0,m\}} \ast h^0_b & \longrightarrow & h^0_a \ast g^0_n \ast h^0_b & \stackrel{}{\longrightarrow} & h^0_a \ast g^1_m \ast h^0_b \end{array}$$

Since X_{\bullet} is birelative 2-Segal it maps the left square and the outer rectangle to pullback diagrams. The claim follows from the pasting law for pullbacks.

Remark 2.30. The active-inert factorization system on $\Delta_{[1]}$ restricts to an active inert factorization system on Δ_{\leq} and Δ_{\geq} . One can analogously to Definition 2.28 define a notion of left and right relative decomposition space and show that it is equivalent to the left and right relative 2-Segal conditions.

3 Examples

The fundamental example of a 2-Segal space is the Waldhausen S_{\bullet} -construction² of an exact ∞ -category, whose space of *n*-simplices $S_n(\mathcal{C})$ can be described as the space of length *n*-flags in \mathcal{C} . The 2-Segal condition then admits an interpretation as the third isomorphism theorem. It is therefore natural to

²See Definition 3.1

expect examples of active equifibered and relative Segal morphisms to arise from morphisms between exact ∞ -categories as well. Indeed every exact functor $F : \mathbb{C} \to \mathcal{D}$ induces a morphism of simplicial spaces $S_{\bullet}(F) : S_{\bullet}(\mathbb{C}) \to S_{\bullet}(\mathcal{D})$. We show that under the assumptions of Proposition 3.4 this morphism in fact gives rise to active equifibered and relative Segal morphisms. Moreover, for every exact functor $F : \mathbb{C} \to \mathcal{D}$ between exact ∞ -categories one can construct a 2-Segal space $S_{\bullet}^{rel}(F)$ called the relative S_{\bullet} -construction [Wal06]. This 2-Segal space fits into a sequence of simplicial spaces

$$\mathcal{D}_{\bullet} \xrightarrow{\iota_{\bullet}} S^{rel}_{\bullet}(F) \xrightarrow{\pi_{\bullet}} S_{\bullet}(\mathcal{C}) \tag{4}$$

that is important in algebraic K-theory since it induces the long exact sequence of relative K-theory [Wal06]. After recalling the definition of the Waldhausen and relative Waldhausen construction in the setting of exact ∞ -categories [Bar15], we show that the morphisms ι_{\bullet} and π_{\bullet} are examples of a relative Segal and an active equifibered morphism respectively.

These considerations induce on the one hand new examples of active equifibered and relative Segal morphisms. On the other hand, these constructions admit applications to the theory of Hall algebras. Indeed, every such morphism induces a (co)algebra morphism between any kind of Hall algebras. In particular, the morphism induced by the relative S_{\bullet} -construction is applicable for the construction of derived Hecke actions [KV22, Sect.5]. This is especially advantageous in the realm of ∞ -categories, where constructing homotopy coherent algebra morphisms can be challenging.

This discussion admits an analog for birelative simplicial spaces in the setting of hermitian K-theory. For exact 1-categories with exact duality $(\mathcal{C}, \mathsf{D}_{\mathcal{C}})$ it has been observed in [You18, Thm.3.6] that the hermitian \mathcal{R}_{\bullet} -construction of $(\mathcal{C}, \mathsf{D}_{\mathcal{C}})$ describes a relative 2-Segal space over $S_{\bullet}(\mathcal{C})$. The author applies this observation to construct representations of Hall algebras and to apply it in orientifold Donaldson-Thomas-theory [You20].

We extend in Section 3.2 their result to the setting of exact ∞ -categories with duality. Our proof is inspired by the construction of the real Waldhausen S_{\bullet} -construction as presented in [HSV19]. For an exact ∞ -category \mathcal{C} with duality functor $\mathsf{D}_{\mathcal{C}}$ the relative 2-Segal space from Example 2.22

$$\mathsf{Tw}(S_{\bullet}(\mathcal{C})) \to S_{\bullet}(\mathcal{C}) \times S_{\bullet}(\mathcal{C})^{rev}$$
(5)

admits an extension to a morphism of simplicial spaces with C_2 -action. For n = 1 we can explicitly describe the C_2 -action by the commutative diagram:



The hermitian \mathcal{R}_{\bullet} -construction then arises as the simplicial space of homotopy fixed points. Further the induced morphism exhibits $\mathcal{R}_{\bullet}(\mathcal{C}) \to S_{\bullet}(\mathcal{C})$ as a relative 2-Segal space. Using this method we finally construct examples of active equifibered and relative Segal maps between relative 2-Segal objects from exact duality preserving functors $F : \mathcal{C} \to \mathcal{D}$.

These results can be applied to the construction of ∞ -categorical Hall algebra representations and their morphisms. In particular, these constructions are essential for constructions of representations of categorified Hall algebra [PS23, DPS22].

3.1 2-Segal Spans from Exact ∞ -Functors

In this subsection we construct examples of active equifibered and relative Segal morphisms between Waldhausen S_{\bullet} -construction. For the reader's convenience, we first recall the construction of the Waldhausen S_{\bullet} -construction of an exact ∞ -categories. An exact ∞ -category consists of a triple ($\mathcal{C}, \mathcal{C}_{in}, \mathcal{C}^{eg}$) of an additive ∞ -category \mathcal{C} and two wide Waldhausen subcategories \mathcal{C}_{in} and \mathcal{C}^{eg} satisfying a list of compatibility conditions [Bar15, Def.3.1].We call the morphism in \mathcal{C}_{in} ingressive denoted \rightarrow and the morphism in \mathcal{C}^{eg} egressive denoted \rightarrow . For an exact ∞ -category \mathcal{C} , a bicartesian square



in \mathcal{C} is called an extension square. A functor $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \to (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ between exact ∞ -categories is called an exact ∞ -functor [Bar15, Def.4.1] if it preserves ingressive and egressive morphisms, zero objects and extension squares. We denote by Exact_{∞} the subcategory of $\mathsf{Fun}(\Lambda_2^2, \mathsf{Cat}_{\infty})$ spanned by exact ∞ -categories and exact functors.

Consider for every $n \ge 0$ the poset [n] as an ∞ -category and let $\mathsf{Ar}([n]) := \mathsf{Fun}([1], [n])$ be the associated ∞ -category of arrows. These assemble into a cosimplicial ∞ -category $\mathsf{Ar}([-]) : \Delta \to \mathsf{Cat}_{\infty}$.

Definition 3.1. Let $(\mathcal{C}, \mathcal{C}_{in}, \mathcal{C}^{eg})$ be an exact ∞ -category. We denote by $S_n(\mathcal{C}, \mathcal{C}_{in}, \mathcal{C}^{eg}) \subset \mathsf{Map}(\mathsf{Ar}([n]), \mathcal{C})$ the full subspace spanned by those functors $F : \mathsf{Ar}([n]) \to \mathcal{C}$ that satisfy the following conditions:

- (1) for every $0 \leq i \leq n$ we have F(i, i) = 0.
- (2) for every $0 \leq i \leq k \leq j \leq n$ the morphism $F(i,j) \rightarrow F(k,j)$ is ingressive.
- (3) for every $0 \le i \le j \le l \le n$ the morphism $F(i, j) \twoheadrightarrow F(i, l)$ is egressive
- (4) for every $0 \leq i \leq k \leq j \leq l \leq n$ the square

is bicartesian.

When *n* varies, the spaces $S_n(\mathbb{C}, \mathbb{C}_{in}, \mathbb{C}^{eg})$ assemble into a simplicial space called the Waldhausen S_{\bullet} -construction. When the classes of ingressive and egressive morphisms are clear from the context we abuse notation and denote the Waldhausen S_{\bullet} -construction by $S_{\bullet}(\mathbb{C})$. For completeness, we include a proof of the following statement. The original proof in the more general context of proto-exact ∞ -categories is given [DK19, Thm.7.3.3].

Proposition 3.2. [DK19, Thm.7.3.3] Let $(\mathcal{C}, \mathcal{C}_{in}, \mathcal{C}^{eg})$ be an exact ∞ -category. The simplicial space $S_{\bullet}(\mathcal{C})$ is 2-Segal.

Proof. We apply the path space criterion [DK19, Thm.6.3.2]. It therefore suffices to show that the two path spaces $(P^{\rhd}S)_{\bullet}$ and $(P^{\lhd}S)_{\bullet}$ are 1-Segal. It follows from [Lur17, Lem.1.2.2.4.] that for every $n \ge 0$ the projection maps

$$\operatorname{Map}(\Delta^{n-1}, \mathcal{C}^{\operatorname{eg}}) \xleftarrow{p} S_n(\mathcal{C}) \xrightarrow{p_h} \operatorname{Map}(\Delta^{n-1}, \mathcal{C}_{\operatorname{in}})$$

that are induced by restricting to the subposet

$$\{(0,n) < (1,n) < \ldots < (n-1,n)\} \hookrightarrow {\sf Ar}([n]) \longleftrightarrow \{(0,1) < (0,2) < \ldots < (0,n)\}$$

are equivalences. Through these equivalences the 1-Segal conditions for the two path spaces translate into the conditions that for every $n \ge 2$ the morphisms

$$\mathsf{Map}(\Delta^n, \mathcal{C}^{\mathsf{eg}}) \to \mathsf{Map}(\Delta^1 \amalg_{\Delta^0} \cdots \amalg_{\Delta^0} \Delta^1, \mathcal{C}^{\mathsf{eg}})$$

and

$$\mathsf{Map}(\Delta^n, \mathcal{C}_{\mathsf{in}}) \to \mathsf{Map}(\Delta^1 \amalg_{\Delta^0} \cdots \amalg_{\Delta^0} \Delta^1, \mathcal{C}_{\mathsf{in}})$$

are equivalences. But this condition is satisfied, since C^{eg} and C_{in} are ∞ -categories.

Let $(\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in})$ and $(\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be exact ∞ -categories and $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \to (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be an exact ∞ -functor. It follows from the construction of the Waldhausen construction that F induces a functor $S_{\bullet}(F) : S_{\bullet}(\mathcal{C}) \to S_{\bullet}(\mathcal{D})$ between the corresponding Waldhausen constructions. We analyze the conditions under which this morphism is active equifibered (resp. relative Segal). To do so, we need to introduce some definitions:

Definition 3.3. Let $(\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in})$ be an exact ∞ -category and let $\mathcal{C}_0 \subset \mathcal{C}$ be a full exact subcategory. We call \mathcal{C}_0

(1) extension closed if every exact-sequence $\sigma: \Delta^1 \times \Delta^1 \to \mathbb{C}$



in \mathfrak{C} with $C_1, C_3 \in \mathfrak{C}_0, \sigma$ factors through \mathfrak{C}_0 .

(2) closed under quotients (resp. subobjects) if for every egressive morphism $C_2 \twoheadrightarrow C_3$ (resp. ingressive morphism $C_1 \rightarrowtail C_2$) in \mathcal{C} with $C_2 \in \mathcal{C}_0$ also $C_3 \in \mathcal{C}_0$ (resp. $C_1 \in \mathcal{C}_0$)

We can then proof the following:

Proposition 3.4. Let $(\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in})$ and $(\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be exact ∞ -categories and $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \rightarrow (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be a fully faithful exact ∞ -functor. The induced morphism $S_{\bullet}(F)$ of simplicial spaces is

- (1) active equifibered if and only if the essential image of F is closed under quotients and subobjects.
- (2) relative Segal if and only if the essential image of F is extension closed.

Proof. To show (1), we need to show that for every $n \ge 1$ the morphism

$$\eta_1: \mathsf{Map}(\Delta^{n-1}, \mathcal{C}_{\mathsf{in}}) \to \mathsf{Map}(\Delta^{n-1}, \mathcal{D}) \times_{\mathcal{D}^{\simeq}} \mathcal{C}^{\simeq}$$

is an equivalence of spaces. It follows from the 2-out-of-3 property for fully faithful functors that η_1 is fully faithful. To prove the claim, it suffices to show that η_1 is essentially surjective. But this follows since $\mathcal{C} \subset \mathcal{D}$ is closed under subobjects.

The proof of (2) is similar. We need to show that for every $n \ge 1$ the functor

$$\eta_2: \mathsf{Map}(\Delta^{n-1}, \mathcal{C}_{\mathsf{in}}) \to \mathsf{Map}_{in}(\Delta^{n-1}, \mathcal{D}) \times_{\mathcal{D}^{\simeq} \times \cdots \times \mathcal{D}^{\simeq}} (\mathcal{C}^{\simeq} \times \cdots \times \mathcal{C}^{\simeq})$$

is an equivalence. It follows again from the 2-out-of-3 property for fully faithful functors that this functor is fully faithful. Moreover, since the essential image is closed under extensions, it is easy to see that the functor is also essentially surjective. \Box

For the construction of further active equifibered and relative Segal maps, we turn to the study of the Sequence 4. Let $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \to (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be an exact functor between exact ∞ -categories.

Following [DKSS21], we define the *relative* S_{\bullet} -construction of F as the simplicial space denoted $S_{\bullet}^{rel}(F)$ whose ∞ -category of *n*-simplices is defined as the pullback

Alternatively, we can also describe $S_n^{rel}(F)$ as the full subspace of the space of sections of the Grothendieck construction of the functor

$$\mathsf{Fun}(\Delta^{n-1}, \mathfrak{C}_{\mathsf{in}}) \xrightarrow{\mathsf{ev}_{n-1}} \mathfrak{C} \xrightarrow{F} \mathfrak{I}$$

with objects the ingressive sections. Since the full subcategory of $\operatorname{Fun}(\Delta^{op}, \mathbb{C})$ generated by 2-Segal objects is closed under limits, it follows that $S^{rel}_{\bullet}(F)$ is itself 2-Segal. The simplicial space $S^{rel}_{\bullet}(F)$ sits in a sequence of simplicial spaces

$$\mathcal{D}_{\bullet} \xrightarrow{\iota_{\bullet}} S_{\bullet}^{rel}(F) \xrightarrow{\pi_{\bullet}} S_{\bullet}(\mathcal{C})$$

where the maps $\iota_n : \mathcal{D} \to S_n(F)$ and $\pi_n : S_n(F) \to S_n(\mathcal{C})$ are given by

$$d \mapsto (0 \rightarrowtail \dots \rightarrowtail 0 \rightarrowtail b)$$

and

$$(a_0 \rightarrowtail \dots \rightarrowtail \dots \rightarrowtail a_{n-1} \rightarrowtail b) \mapsto (a_0 \rightarrowtail \dots \rightarrowtail a_{n-1})$$

respectively. These maps have the following properties:

Proposition 3.5. Let $F : \mathcal{C} \to \mathcal{D}$ be an exact functor between exact ∞ -categories. Then:

- (1) the morphism π_{\bullet} is active equifibered.
- (2) the morphism ι_{\bullet} is relative Segal.

Proof. Note that $S_n(F)$ can be written as the pullback



of the outer rectangle. To show the claim for π_{\bullet} , we need to show that the upper diagram is a pullback as well. But this follows from the pasting law for pullbacks. For (2) note that the pullback of spaces

$$\mathcal{D}^{\simeq} \times_{S_1^{rel}(F) \times_{S_0^{rel}(F)} \cdots \times_{S_0^{rel}(F)} S_1^{rel}(F)} S_n^{rel}(F)$$

can be identified with the full subspace of $S_n^{rel}(F) \simeq S_n(\mathcal{C}) \times_{S_n(\mathcal{D})} S_{n+1}(\mathcal{D})$ on those objects $(F_{\mathcal{C}}.G_{\mathcal{D}}, \alpha)$, s.t.

- (1) the restriction of $G_{\mathcal{D}}$ to the subposet $\{(0,n) < ... < (n-1,n)\}$ is constant
- (2) the restriction of $F_{\mathcal{C}}$ to the subposet $\{(0,1) < (1,2) < \dots < (n-1,n)\}$ is constant at 0

It then follows from an iterated Kan extension argument similar to [Lur17, Lem.1.2.2.4] that this subspace is equivalent to \mathcal{D}^{\simeq} .

As an application, we use the above results for the construction of the so-called derived Hecke actions from [KV22].

Proposition 3.6. Let \mathcal{C} be an ∞ -category with finite limits and $p: Y_{\bullet} \to X_{\bullet}$ be a relative 2-Segal object. Let $f: X_{\bullet} \to Z_{\bullet}$ be an active equifibered and $g: W_{\bullet} \to X_{\bullet}$ be relative Segal morphism. Then

- (1) the composite $f \circ p : Y_{\bullet} \to Z_{\bullet}$ is a relative 2-Segal object
- (2) the pullback $\pi: Y_{\bullet} \times_{X_{\bullet}} W_{\bullet} \to W_{\bullet}$ is a relative 2-Segal object

Proof. Since Y_{\bullet} is 1-Segal by definition, for the first case we only need to show that the composite $f \circ p$ is active equifibered. But this follows since active equifibered morphisms are closed under composition. For the second case note that it follows from the proof of Lemma 2.5 that the pullback of an active equifibered morphism along a relative Segal morphism is active equifibered. We therefore only need to show that $Y_{\bullet} \times_{X_{\bullet}} W_{\bullet}$ is 1-Segal. Consider the morphisms:

$$Y_{\bullet} \times_{X_{\bullet}} W_{\bullet} \to Y_{\bullet} \to *_{\bullet}$$

Since both of them are relative Segal their composite is so. In particular it follows that $Y_{\bullet} \times_{X_{\bullet}} W_{\bullet}$ is 1-Segal.

As a consequence we obtain:

Corollary 3.7. Let $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \to (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be an exact fully faithful functor between exact ∞ -categories whose essential image is extension closed. Then the simplicial space $S_{\bullet}(F)$ is 1-Segal

Proof. This follows from an application of Proposition 3.6 to the pullback diagram that defines $S_{\bullet}(F)$.

Example 3.8. Let $F : (\mathcal{C}, \mathcal{C}^{eg}, \mathcal{C}_{in}) \to (\mathcal{D}, \mathcal{D}^{eg}, \mathcal{D}_{in})$ be a fully faithful exact functor between exact ∞ -categories whose essential image is closed under extensions. Then the morphism

$$S_{\bullet}(F) \to S_{\bullet}(\mathcal{C})$$

defines a relative 2-Segal object by Corollary 3.7. Unraveling definitions this equips \mathcal{D} with the structure of an $S_{\bullet}(\mathcal{C})$ -module with action given by



This module structure is responsible for the so-called derived Heck actions as defined in [KV22, Sect.5.2]. Indeed repeating the analysis of [DK18, Sect.8] for the theory with transfer given by Borel-Moore homology and the above module structure recovers their derived Hecke actions. Further, it can be used to extend their derived Hecke action to the categorified Hall algebra of [PS23, DPS22].

3.2 Relative 2-Segal Spans from Duality preserving ∞ -Functors

In this subsection, we present a general construction of relative 2-Segal objects from 2-Segal spaces with duality. For this construction, we first have to recall some facts about ∞ -categories with duality. Since we only work with simplicial objects in S, we drop it from the notation.

Recall that the category Δ admits a canonical C₂-action. This action maps an object [n] to [n] and a morphism $f; [n] \rightarrow [m]$ to the composite morphism

$$[n] \simeq [n]^{op} \xrightarrow{f^{op}} [m]^{op} \simeq [m]$$

where the equivalence $[n] \simeq [n]^{op}$ is given by $i \mapsto n-i$. By functoriality of taking presheaves the C_2 -action on Δ induces one on $\mathcal{P}(\Delta)$ that maps a simplicial space X_{\bullet} to X_{\bullet}^{rev} . The restriction of this action to the full subcategory $\mathsf{Cat}_{\infty} \subset \mathcal{P}(\Delta)$ maps an ∞ -category to its opposite.

Definition 3.9. A simplicial space with duality is a homotopy fixed point with respect to the above C_2 -action on $\mathcal{P}(\Delta)$. Similarly, an ∞ -category with duality is a homotopy fixed point with respect to the C_2 -action on Cat_{∞} given by taking opposites.

More precisely an ∞ -category with duality is a section of the cocartesian fibration $\widetilde{Cat}_{\infty} \to BC_2$ encoding the C_2 -action on Cat_{∞} . We define the ∞ -category $Cat_{\infty}^{C_2}$ of ∞ -categories with duality and the ∞ -category of simplicial spaces with duality $\mathcal{P}(\Delta)^{C_2}$ as the respective ∞ -categories of homotopy fixed points.

Remark 3.10. The ∞ -categories Δ , $\mathcal{P}(\Delta)$ and Cat_{∞} also admit a trivial C_2 -action. We denote the respective category of homotopy fixed points with respect to this trivial action by $\Delta[\mathsf{C}_2]$, $\mathcal{P}(\Delta)[\mathsf{C}_2]$ and $\mathsf{Cat}_{\infty}[\mathsf{C}_2]$. In the following, we will call a homotopy fixed point with respect to this trivial action an object with C_2 -action.

Note that the non-trivial C_2 -action on $\mathcal{P}(\Delta)$ restricts to an C_2 -action on 2- Seg_{Δ} .

Definition 3.11. A 2-Segal space with duality is a homotopy fixed point with respect to the non-trivial C_2 -action on 2-Seg $_{\Delta}$.

We denote by 2- $\operatorname{Seg}_{\Delta}^{C_2}$ the ∞ -category of homotopy fixed points. By construction, this ∞ -category comes equipped with a forgetful functor to 2- $\operatorname{Seg}_{\Delta}(\mathbb{S})$. We call a morphism $f : X_{\bullet} \to Y_{\bullet}$ in 2- $\operatorname{Seg}_{\Delta}^{C_2}$ active equifibered (resp. relative Segal) if its image in 2- $\operatorname{Seg}_{\Delta}$ is active equifibered (resp. relative Segal). Similarly, we call a span



in 2-Seg^{C_2} a 2-Segal span with duality if s_{\bullet} is active equifibered and t_{\bullet} is relative Segal. Since the ∞ -category 2-Seg^{C_2} admits small limits, it makes sense to define:

Definition 3.12. We define the ∞ -category 2-Seg $_{\Delta}^{\leftrightarrow,C_2}$ of 2-Segal spaces with duality as the subcategory of Span(2-Seg $_{\Delta}^{C_2}$) with morphisms 2-Segal spans with duality.

Similarly, note that the trivial C_2 -action on $\mathcal{P}(\Delta)$ induces the trivial C_2 -action on Rel2Seg $_{\Delta}$.

Definition 3.13. A relative 2-Segal space with C_2 -action is a homotopy fixed point with respect to the trivial C_2 -action on Rel2Seg. We denote by Rel2Seg $_{\Delta}[C_2]$ the ∞ -category with homotopy fixed points.

As above the ∞ -category Rel2Seg[C₂] comes equipped with a forgetful functor to the ∞ -category Rel2Seg_{Δ}. We define active equifibered and relative Segal morphisms as for 2-Seg^{C₂}. The associated version of 2-Segal span is called a *relative 2-Segal span with* C₂-action.

Definition 3.14. We define the ∞ -category Rel2Seg $_{\Delta}^{\leftrightarrow}[C_2]$ of relative 2-Segal spaces with C_2 -action as the subcategory of Span(Rel2Seg $_{\Delta}[C_2]$) with morphisms relative 2-Segal spans with C_2 -action.

Our first goal in this section is the construction of a functor

$$\mathsf{Tw}^{\leftrightarrow}(\text{-})^{\mathsf{C}_2}: 2\text{-}\mathrm{Seg}_{\Delta}^{\leftrightarrow,\mathsf{C}_2} \to \mathrm{Rel2Seg}_{\Delta}^{\leftrightarrow}$$

that associates to every 2-Segal space with duality a relative 2-Segal space. Therefore, recall the definition of the edgewise subdivision functor $e : \Delta \to \Delta$ from Example 2.22. This functor becomes C₂-equivariant if we equip the source with the trivial and the target with the non-trivial C₂-action. By functoriality, this lifts to a C₂-equivariant functor

$$\mathsf{Tw} := e^* : \mathcal{P}(\Delta) \to \mathcal{P}(\Delta)$$

that induces a functor $\mathsf{Tw} : \mathcal{P}(\Delta)^{\mathsf{C}_2} \to \mathcal{P}(\Delta)[\mathsf{C}_2]$ on homotopy fixed points. The canonical inclusions $[n] \hookrightarrow [n] * [n]^{op}$ and $[n]^{op} \hookrightarrow [n] * [n]^{op}$ induce a natural transformation

$$\mathsf{Tw} \Rightarrow id \times (-)^{op} : \mathcal{P}(\Delta) \to \mathcal{P}(\Delta)$$

It follows from [HSV19, Lem.2.23] that the natural transformation is C_2 -equivariant and hence induces a natural transformation

$$\mathsf{Tw} \Rightarrow id \times (-)^{op} : \mathcal{P}(\Delta)^{\mathsf{C}_2} \to \mathcal{P}(\Delta)[\mathsf{C}_2]$$

on homotopy fixed points. In total, we can interpret this construction as a functor

$$\mathcal{P}(\Delta)^{\mathsf{C}_2} \xrightarrow{\mathsf{Tw}^{\rightarrow}(\operatorname{-})} \mathsf{Fun}(\Delta^1, \mathcal{P}(\Delta)[\mathsf{C}_2]) \xrightarrow{(-)^{\mathsf{C}_2}} \mathsf{Fun}(\Delta^1, \mathcal{P}(\Delta))$$

that associates to a simplicial space with duality X_{\bullet} the morphism

$$\mathsf{Tw}(X)_{\bullet}^{\mathsf{C}_2} \to (X_{\bullet} \times X_{\bullet}^{rev})^{\mathsf{C}_2}$$

Proposition 3.15. Let $X_{\bullet} \in \mathcal{P}(\Delta)^{\mathsf{C}_2}$ be a simplicial space with duality and let $X_{\bullet} \times X_{\bullet}^{rev}$ be the induced simplicial space with C_2 -action. There exists an equivalence of simplicial spaces

$$(X_{\bullet} \times X_{\bullet}^{rev})^{\mathsf{C}_2} \simeq X_{\bullet}$$

Proof. For any simplicial space X_{\bullet} , we can construct a C_2 -action on $X_{\bullet} \times X_{\bullet}$ via right Kan extension

$$* \xrightarrow{X_{\bullet}} \mathcal{P}(\Delta)$$

$$\downarrow \xrightarrow{X_{\bullet} \times X_{\bullet}} \mathcal{P}(\Delta)$$

$$\mathsf{BC}_{2}$$

By adjunction the projection onto the first factor $X_{\bullet} \times X_{\bullet}^{rev} \to X_{\bullet}$ induces a morphism of simplicial spaces with C_2 -action

$$\kappa: X_{\bullet} \times X_{\bullet}^{rev} \to X_{\bullet} \times X_{\bullet}$$

It follows from the construction that the functor underlying κ is given by $id \times \mathsf{D} : X_{\bullet} \times X_{\bullet}^{rev} \to X_{\bullet} \times X_{\bullet}$, where D denotes the duality on X_{\bullet} . Hence it is an equivalence. We therefore obtain an equivalence on homotopy fixed points. The claim follows from the transitivity of right Kan extensions

Proposition 3.16. The functor $\mathsf{Tw}^{\rightarrow}(-)$ constructed above restricts to a functor

$$\mathsf{Tw}^{\rightarrow}(\text{-}): 2\text{-}\mathrm{Seg}_{\Delta}^{\mathsf{C}_2} \to \mathrm{Rel}2\mathrm{Seg}_{\Delta}[\mathsf{C}_2]$$

Further, it preserves active equifibered and relative Segal morphisms.

Proof. It follows from Example 2.22 that the map $\mathsf{Tw}(X)_{\bullet} \to X_{\bullet} \times X_{\bullet}^{rev}$ is a relative 2-Segal object with C_2 -action. The second claim can be checked directly by looking at the corresponding pullback diagrams.

Proposition 3.17. The functor $(-)^{C_2}$: Fun $(\Delta^1, \mathcal{P}(\Delta)[C_2]) \rightarrow$ Fun $(\Delta^1, \mathcal{P}(\Delta))$ restricts to a functor

$$(-)^{\mathsf{C}_2} : \operatorname{Rel2Seg}_{\Delta}[\mathsf{C}_2] \to \operatorname{Rel2Seg}_{\Delta}$$

Further, it preserves active equifibered and relative Segal morphisms.

Proof. The claim follows from the fact that the functor $\lim_{BC_2}(-)$ preserves small limits.

Combining Proposition 3.17 and 3.16 we can obtain our first goal of this section

Proposition 3.18. The composite ∞ -functor

$$2\text{-}\mathrm{Seg}_{\Delta}^{\mathsf{C}_2} \xrightarrow{\mathsf{Tw}^{\rightarrow}(\text{-})} \mathrm{Rel2}\mathrm{Seg}_{\Delta}[\mathsf{C}_2] \xrightarrow{(-)^{\mathsf{C}_2}} \mathrm{Rel2}\mathrm{Seg}_{\Delta}$$

induces an ∞ -functor on the level of spans

$$\mathsf{Tw}^{\leftrightarrow}(\text{-})^{\mathsf{C}_2}: 2\text{-}\mathrm{Seg}_{\Delta}^{\leftrightarrow,\mathsf{C}_2} \to \mathrm{Rel2Seg}_{\Delta}^{\leftrightarrow}[\mathsf{C}_2] \to \mathrm{Rel2Seg}_{\Delta}^{\leftrightarrow}$$

We use this Proposition for the construction of examples of relative 2-Segal spaces and spans. In the previous section, we have shown that algebraic K-theory is a rich source of examples of active equifibered and relative Segal Δ^{op} -morphisms. The analog for relative 2-Segal spans is hermitian K-theory. The ideas behind hermitian K-theory originate from the fundamental work of Hesselholt and Madsen [HM15] on real algebraic K-theory. An ∞ -categorical formulation of these ideas is given in [HSV19]. The hermitian Waldhausen construction associates to every exact ∞ -category with duality a simplicial space with duality. For the construction of the hermitian Waldhausen construction, we follow the presentation from [HSV19, Sect.8.2]. To do so, we first recall some facts about exact ∞ -categories with duality and $S[C_2]$ -enriched ∞ -categories.

Recall from Subsection 3.1 that the ∞ -category Exact_{∞} of exact ∞ -categories is defined as the subcategory of $\mathsf{Exact}_{\infty} \subset \mathsf{Fun}(\Lambda_2^2,\mathsf{Cat}_{\infty})$ with objects exact ∞ -categories and morphisms exact functors. The ∞ -category Λ_2^2 carries a natural C₂-action. This action combines with the C₂-action on Cat_{∞} to an action on Fun($\Lambda_2^2,\mathsf{Cat}_{\infty}$) that is defined on objects as

$$(\mathfrak{C}_0 \to \mathfrak{C}_2 \leftarrow \mathfrak{C}_1) \mapsto (\mathfrak{C}_1^{op} \to \mathfrak{C}_2^{op} \leftarrow \mathfrak{C}_0^{op})$$

It is easy to see that this action restricts to an C_2 -action on the subcategory Exact_{∞} .

Definition 3.19. An *exact* ∞ -*category with duality* is a homotopy fixed point with respect to the above C_2 -action on Exact_{∞} . We call the ∞ -category of homotopy fixed points $\text{Exact}_{\infty}^{C_2}$ the ∞ -category of exact ∞ -categories with duality. A morphism in this ∞ -category is called an exact duality preserving functor.

Informally, an exact ∞ -category with duality \mathcal{C} is a Waldhausen ∞ -category s.t. the underlying ∞ category admits the structure of an ∞ -category with duality and the class of ingressive morphisms \mathcal{C}_{in} together with the opposites of the ingressive morphisms form the structure of an exact ∞ -category on \mathcal{C} . These serve as an input for the hermitian Waldhausen construction.

In [HSV19] the authors work with real exact ∞ -categories and equip the real Waldhausen S_{\bullet} -construction with the structures of a real simplicial space. Since the ∞ -category of exact ∞ -categories with duality forms a full subcategory of the ∞ -category of real exact ∞ -categories [HSV19, Rem.2.33], we can apply the construction of [HSV19, Sect 8.2] to our situation.

The key ingredient in the construction [HSV19, Sect.8.2] is enriched ∞ -category theory. In our situation, we are interested in ∞ -categories enriched over the ∞ -category $S[C_2]$ of spaces with C_2 -action. The amount of enriched ∞ -category theory necessary for this construction is described in [HSV19, App. A].³

Notation 3.20. In the following, we adopt the convention to denote $S[C_2]$ -enriched ∞ -categories by \underline{C} to distinguish them from their underlying ∞ -category C.

For an $S[C_2]$ -enriched ∞ -category \underline{C} the underlying ∞ -category is obtained by taking homotopy fixed points on Hom-spaces [HSV19, Sect.3.1]. It follows from [HSV19, Cor.2.8] that the ∞ -categories $Cat_{\infty}^{C_2}$ and $\mathcal{P}(\Delta)^{C_2}$ are Cartesian closed and are therefore enriched over $S[C_2]$. This enrichment restricts along the inclusion $\Delta^{C_2} \subset Cat_{\infty}^{C_2}$ to an enrichment of Δ . We denote the $S[C_2]$ -enriched ∞ -category Δ by $\underline{\Delta}$. Further, the ∞ -category $S[C_2]$ is naturally enriched over itself.

It follows from [HSV19, Prop.2.12] that there exists an equivalence $\underline{\mathcal{P}}(\Delta)^{\mathsf{C}_2} \simeq \underline{\mathsf{Fun}}_{\mathsf{S}[\mathsf{C}_2]}(\underline{\Delta}^{op}, \mathsf{S}[\mathsf{C}_2])$ of $\mathsf{S}[\mathsf{C}_2]$ -enriched ∞ -categories between the ∞ -category of simplicial spaces with duality and the ∞ -category of $\mathsf{S}[\mathsf{C}_2]$ -enriched functors. We will use this equivalence for the construction of the duality structures

³For a more general discussion of enriched ∞ -category theory see [GH15]

on the Waldhausen S_{\bullet} -construction.

The $S[C_2]$ -enriched ∞ -category <u>Exact</u>^{C_2}_{∞} [HSV19, Sect.7.1] is cotensored over $Cat_{\infty}^{C_2}$ [HSV19, Sect.8.2]. This cotensoring induces an $S[C_2]$ -enriched functor:

$$(-)^{-}: \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2} \times \underline{\mathsf{Cat}}_{\infty}^{\mathsf{C}_2, \mathsf{op}} \to \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2}$$

For an ∞ -category with duality \mathcal{I} and an exact ∞ -category with duality \mathcal{C} the underlying exact ∞ category of the cotensor \mathcal{C}^I is given by the functor category $\mathsf{Fun}(I, \mathcal{C})$. The exact structure is defined objectwise and the induced duality structure maps a functor $F: I \to \mathcal{C}$ to the composite functor

$$I \xrightarrow{\mathsf{D}_{\mathsf{l}}^{\mathsf{op}}} I^{op} \xrightarrow{F^{op}} \mathcal{C}^{op} \xrightarrow{\mathsf{D}_{\mathcal{C}}} \mathcal{C}$$

As the first step of our construction of the S_{\bullet} -construction, we define an $S[C_2]$ -enriched version of the ∞ -category of arrows. Analogously to the non-enriched case, we define it as the $S[C_2]$ -enriched functor

$$\underline{\mathsf{Ar}}(-):\underline{\Delta} \subset \underline{\mathsf{Cat}}_{\infty}^{\mathsf{C}_2} \xrightarrow{\underline{\mathsf{Hom}}([1],-)} \underline{\mathsf{Cat}}_{\infty}^{\mathsf{C}_2}$$

where $\underline{\text{Hom}}(-, -)$ denotes the $S[C_2]$ -enriched internal Hom-functor of $Cat_{\infty}^{C_2}$. For every [n] this induces a duality structure on the ∞ -category of Ar([n]). Note that for every $[n] \in \underline{\Delta}$ the induced duality structure on $C^{Ar([n])}$ restricts to a duality structure on $S_n(\mathcal{C})$. It follows that the $S[C_2]$ -enriched functor

$$(-)^{-}: \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2} \times \underline{\Delta}^{op} \to \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2}$$

restricts to an $S[C_2]$ -enriched functor

$$\underline{S}_{\bullet}(-): \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2} \times \underline{\Delta}^{op} \to \underline{\mathsf{Exact}}_{\infty}^{\mathsf{C}_2}$$

Such a functor is transpose to an $S[C_2]$ -enriched functor

$$\underline{S}^{\simeq}_{\bullet}(-): \underline{\mathsf{Exact}}^{\mathsf{C}_2}_{\infty} \to \mathsf{Fun}_{\mathscr{S}[\mathsf{C}_2]}(\underline{\Delta}^{op}, \underline{\mathsf{Exact}}^{\mathsf{C}_2}_{\infty}) \xrightarrow{(-)^{\simeq}} \mathsf{Fun}_{\mathscr{S}[\mathsf{C}_2]}(\underline{\Delta}^{op}, \underline{\mathscr{S}}[\mathsf{C}_2]) \simeq \underline{\mathcal{P}}(\Delta)^{\mathsf{C}_2}$$

Definition 3.21. Let \mathcal{C} be an exact ∞ -category with duality. The 2-Segal space with duality $\underline{S}^{\sim}_{\bullet}(\mathcal{C})$ is called the *Waldhausen S*_•-construction with duality.

The underlying simplicial space of \underline{S}_{\bullet} coincides with the Waldhausen S_{\bullet} -construction of an exact ∞ category as introduced in Definition 3.1. In particular, the Waldhausen S_{\bullet} -construction with duality is
an example of a 2-Segal space with duality.

We can now apply Proposition 3.18 to the Waldhausen S_{\bullet} -construction with duality.

Definition 3.22. [HS04, Sect.1.8] Let \mathcal{C} be an exact ∞ -category with duality. We call the simplicial object $\mathsf{Tw}(\underline{S}^{\simeq}_{\bullet}(\mathbb{C}))^{\mathsf{C}_2}$ the hermitian S_{\bullet} -construction and denote it by $\mathcal{R}_{\bullet}(\mathbb{C})$.

Remark 3.23. It is known by the fundamental work of Waldhausen [Wal06] that for every exact ∞ -category $\mathsf{Tw}(S_{\bullet}(\mathbb{C}))$ is equivalent to Quillens *Q*-construction $Q(\mathbb{C})$. In [CDH⁺20] the authors introduced a hermitian version of the *Q*-construction. We expect that for every exact ∞ -category with duality \mathbb{C} the equivalence between the $\mathsf{Tw}(S_{\bullet}(\mathbb{C}))$ and the *Q*-construction $Q(\mathbb{C})$ extends to an equivalence of simplicial spaces with C_2 -action.

The following result extends [You18, Thm.3.6] into the realm of ∞ -categories:

Corollary 3.24. Let \mathcal{C} be an exact ∞ -category with duality. The morphism

$$\mathcal{R}_{\bullet}(\mathcal{C}) \to S_{\bullet}(\mathcal{C})$$

is a relative 2-Segal object.

Proof. Apply Proposition 3.18 to $\underline{S}^{\sim}_{\bullet}(\mathcal{C})$.

We can further use Proposition 3.18 for the construction of relative 2-Segal spans. Let $F : \mathbb{C} \to \mathcal{D} \in \mathsf{Exact}_{\infty}^{\mathsf{C}_2}$ be a duality preserving exact functor. It follows from the construction of $\underline{S}_{\bullet}(-)$ that it induces a morphism

$$\underline{S}_{\bullet}(F): \underline{S}_{\bullet}(\mathcal{C}) \to \underline{S}_{\bullet}(\mathcal{D})$$

of simplicial spaces with duality.

Corollary 3.25. Let $F : \mathfrak{C} \to \mathfrak{D} \in \mathsf{Exact}_{\infty}^{\mathsf{C}_2}$ be a fully faithful duality preserving exact functor and consider the induced diagram

$$\begin{array}{c} \mathcal{R}_{\bullet}(\mathbb{C}) \xrightarrow{\mathcal{R}_{\bullet}(F)} \mathcal{R}_{\bullet}(\mathcal{D}) \\ & \downarrow \\ & \downarrow \\ S_{\bullet}(\mathbb{C}) \xrightarrow{S_{\bullet}(F)} S_{\bullet}(\mathcal{D}) \end{array}$$

Then the morphism $(\mathfrak{R}_{\bullet}(F), S_{\bullet}(F))$ is

- (1) active equifibered if F is closed under extensions and subobjects.
- (2) relative Segal if the essential image of F is closed under extensions.

Remark 3.26. The author is not aware of a hermitian analog of the relative S_{\bullet} -construction constructed in Subsection 3.1. An analog of the construction given in Subsection 3.1 for exact ∞ -categories can not be performed for exact ∞ -categories with duality. The problem is that the defining pullback diagram does not lift to a diagram of simplicial spaces with duality.

4 Modules in Span Categories

Let \mathcal{C} be an ∞ -category with finite limits. The goal of this section is to provide a characterization of bimodule objects in $\text{Span}_{\Delta}(\mathcal{C}^{\times})$ in terms of birelative 2-Segal objects in \mathcal{C} (see Definition 2.7). The proof we present here is a multicolored version of the proof provided in [Ste21, Sect.2], with many ideas drawn from there. For this, we use an explicit combinatorial description of the monoidal ∞ -category $\text{Span}_{\Delta}(\mathcal{C}^{\times})$ constructed in [DK19] that we recall in Subsection 4.1. After this preliminary subsection, we prove the claimed equivalence in Subsection 4.2.

Before we start, we sketch the strategy of the proof here. Let the functor $F : \Delta_{/[1]} \to \operatorname{Span}_{\Delta}(\mathbb{C}^{\times})$ represent a bimodule object. Unraveling definitions, we identify F in Corollary 4.11 with a functor $F : \operatorname{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{[1]} \to \mathbb{C}$ satisfying a list of conditions. In particular, F inverts a class of morphisms that we will denote by E. To deal with these conditions, we construct a localization functor \mathcal{L} : $\operatorname{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{[1]} \to \Delta_1^*$ along the morphisms E. The category Δ_1^* contains $\Delta_{/[1]}^{op}$ as a full subcategory and the remaining conditions translate under the restriction $\operatorname{Fun}(\Delta_1^*, \mathbb{C}) \to \operatorname{Fun}(\Delta_{/[1]}^{op}, \mathbb{C})$ to the birelative 2-Segal conditions (see Definition 2.7).

More explicitly we can describe this identification as follows. The category $\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}$ has objects $(f, \{i, j\})$ represented by diagrams of the form:



in Δ . Similarly a 1-morphism $(e^0, e^1) : (f_0, [i, j]) \to (f_1, [l, m])$ in $\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{[1]}$ can be represented by a commutative diagram in Δ of the form:



s.t. $e_0(i) \leq l \leq m \leq e_1(j)$. We denote by $X_{\bullet}: \Delta_{/[1]}^{op} \to \mathcal{C}$ the birelative simplicial object associated to F under the equivalence of Theorem 4.10. In terms of X_{\bullet} the value of F on (f, [i, j]) admits an interpretation as:

$$X_{g^{1}_{[(f(i),f(i+1)]}} \times \cdots \times X_{g^{1}_{[(f(j-1),f(j)]}}$$

Similarly, in terms of X_{\bullet} the value of F on the morphism (e^0, e^1) translates to the morphism:

 $X_{g^{1}_{[(f^{0}(i),f^{0}(i+1)]}} \times \dots \times X_{g^{1}_{[(f^{0}(j-1),f^{0}(j)]}} \to X_{h^{1}_{[(f^{1}(l),f^{1}(l+1)]}} \times \dots \times X_{h^{1}_{[(f^{1}(m-1),f^{1}(m)]}}$ position with the projection onto $X_{h^{1}}$ is given by:

whose composition with the projection onto $X_{h^1_{[(f^1(l), f^1(l+1)]}}$ is given by:

$$X_{g^{1}_{[(f^{0}(i),f^{0}(i+1)]}} \times \dots \times X_{g^{1}_{[(f^{0}(j-1),f^{0}(j)]}} \to X_{g^{1}_{[(f^{0}(i),f^{0}(i+1)]}} \xrightarrow{X_{e^{1}_{[f^{1}(l),f^{1}(l+1)]}}} X_{h^{1}_{[(f^{1}(l),f^{1}(l+1)]}} \to X_{h^{1$$

We unravel this in a specific example.

Example 4.1. Consider the objects $g : [3] \xrightarrow{0011} [1]$ and $id_{[1]} : [1] \to [1]$ in $\Delta_{/[1]}$. Denote the associated birelative simplicial object by X_{\bullet} . We denote by $f_{i,j} : [1] \to [3]$ the unique morphism with image $\{i, j\}$ for $i \leq j$. The morphism $f_{0,3}$ extends to an object $(f_{0,3} : \mathsf{id}_{[1]} \to g, \{0,1\})$ in $\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}$. Under the identification of the theorem the value of F on $(f_{0,3}, \{0,1\})$ identifies with

$$X_{g_{[f(0),f(1)]}} = X_g =: X_{(0,0,1,1)}$$

Next, consider the morphisms





in $\Delta_{/[1]}$. Under the identification with X_{\bullet} we can interpret the two morphisms $F(f_{0,3}, \{0, 1\}) \rightarrow F(\mathsf{id}_{[3]}, \{0, 3\})$ and $F(f_{0,3}, \{0, 1\}) \rightarrow F(\mathsf{id}_{[1]}, \{0, 1\})$ in \mathcal{C} as the span



This span describes the simultaneous action of the algebra objects $X_{(0,0)}$ and $X_{(1,1)}$ on the bimodule $X_{(0,1)}$.

4.1 Preliminaries

In this subsection, we introduce some notations and definitions essential for the combinatorics of the next section. Particularly, we recall an explicit construction of a monoidal structure on the ∞ -category of spans in the model of quasi-categories. We use this model in the following sections. First, we introduce some notation:

Notation 4.2. Let $f : [n] \to [1]$ be an object in $\Delta_{/[1]}$. We will say that f is supported at $S \subset \{0, 1\}$, if $\mathsf{Im}(f) = S$. We can uniquely represent such an object f by a sequence (0, 0, ..., 0, 1, ..., 1).

Definition 4.3. The interval category ∇ is the subcategory of Δ with objects given by those [n] with $n \ge 1$, and morphisms are maps that preserve maximal and minimal elements, also called active maps. The augmented interval category ∇^+ is the wide subcategory of Δ with morphisms being the active maps.

Definition 4.4. Let S be a linearly ordered set. An inner interstice of S is a pair $(n, n + 1) \in S \times S$, where n + 1 denotes the successor of n in S. We denote the set of inner interstices of S by $\mathbb{I}(S)$. This set comes equipped with a canonical linear order induced from S. This construction can be enhanced to a functor

 $\mathbb{I}: \nabla^{op}_+ \to \Delta_+$

that associates to a morphism $f:S\to T\in \nabla_+$ the morphism $\mathbb{I}(f):\mathbb{I}(T)\to\mathbb{I}(S)$

$$\mathbb{I}(f)(j, j+1) = (k, k+1) \quad ; f(k) \le j < j+1 \le f(k+1)$$

Definition 4.5. We define Δ^{\coprod} to be the category with objects $([n], i \leq j)$ consisting of a linearly ordered set [n] and a pair of objects $i \leq j \in [n]$. A morphism $f : ([n], i \leq j) \rightarrow ([m], k \leq l)$ is given by morphism f in Δ , s.t. $f(i) \leq k \leq l \leq f(j)$. An object $([n], i \leq j)$ represents a subinterval $\{i, ..., j\} \subset [n]$. We will frequently denote an object $([n], i \leq j)$ in Δ^{\coprod} by ([n], [i, j]) It is easy to check, that the forgetful functor $\pi_{\coprod} : \Delta^{\coprod} \rightarrow \Delta$ is a Cocoartesian fibration. For every morphism $f : [n] \rightarrow [m]$ a π_{\coprod} -cocartesian lift is given by the morphism $f : ([n], i \leq j) \rightarrow ([m], f(i) \leq f(j))$.

and

Construction 4.6. Given a morphism $f : [m] \to [n]$ over [1] we can uniquely decompose it as follows. We can decompose the source into a composite of $[m] := \{0,1\} * \{1,2\} * ... * \{m-1,m\}$. Restricting f to each individual interval yields a morphism $f_i := f|_{\{i-1,i\}} : \{i-1,i\} := [1_i] \to \{f(i-1) \leq f(i)\} := [n_i]$ in $\Delta_{/[1]}$ that preserves endpoints. We further denote $\{0 \leq f(0)\} := [n_l]$ and $\{f(m) \leq n\} := [n_r]$. Using this decomposition, we can uniquely reconstruct f as a morphism in $\Delta_{/[1]}$:

$$f = f_1 \ast \ldots \ast f_m : [1_1] \ast \ldots \ast [1_m] \rightarrow [n_1] \ast \ldots \ast [n_m] \hookrightarrow [n_l] \ast [n_1] \ast \ldots \ast [n_m] \ast [n_r]$$

We call this process decomposition.

We now proceed to recall a construction of the Cartesian monoidal structure and the ∞ -category of spans within the context of quasi-categories from [DK19, Chapt.10]. First we recall an explicit model for the Cartesian monoidal structure. In the following, we implicitly identify every 1-category with its nerve.

Construction 4.7. [Ste21, Constr.1.29] Let C be an ∞ -category with finite products. We define a simplicial set over Δ via the adjunction formula

$$\operatorname{Hom}_{\Delta}(K,\overline{\mathfrak{C}}^{\times}) \simeq \operatorname{Hom}_{\operatorname{Set}_{\Delta}}(K \times_{\Delta} \Delta^{\amalg}, \mathfrak{C})$$

This defines a Cartesian fibration $p: \bar{\mathbb{C}}^{\times} \to \Delta$. We define $\mathbb{C}^{\times} \subset \bar{\mathbb{C}}^{\times}$ to be the full subcategory on those objects $F: \Delta^{0} \times_{\Delta} \Delta^{\coprod} \to \mathbb{C}$, that display $F(\{i \leq j\})$ as a product of $F(\{k \leq k+1\})$ for $i \leq k < j$. The restricted functor $p: \mathbb{C}^{\times} \to \Delta$ is also a Cartesian fibration and exhibits the Cartesian monoidal structure on \mathbb{C} . A morphism $\tilde{\Phi}: \Delta^{1} \to \mathbb{C}^{\times}$ represented by a map $\Phi: \{[n] \to [m]\} \times_{\Delta} \Delta^{\coprod} \to \mathbb{C}$ is Cartesian if and only if Φ carries all π_{\coprod} -cocartesian edges of $\{[n] \xrightarrow{f} [m]\} \times_{\Delta} \Delta^{\coprod} \to \Delta$ to equivalences [Lur06, Cor.3.2.2.12]. This happens if and only if Φ maps all maps of the form $f: ([i, j] \subset [n]) \to ([f(i), f(j)] \subset [m])$ induced by a weakly monotone map $f: [n] \to [m]$ to equivalences.

Construction 4.8. [Ste21, Constr.1.33] Given $X_{\bullet} \in \mathsf{Set}_{\Delta}$. There exists an adjunction:

 $\mathsf{Tw}_X:(\mathsf{Set}_\Delta)_{/X}\longleftrightarrow(\mathsf{Set}_\Delta)_{/X}:\overline{\mathsf{Span}}_X$

by setting $\mathsf{Tw}_X(f: S \to X) = \mathsf{Tw}(S) \to \mathsf{Tw}(X) \to X$ and $\overline{\mathsf{Span}}_X(S \to X)$ to be the left vertical arrow of the pullback:

Let $p: S \to X$ a map of simplicial sets. An *n*-simplex in $\overline{\mathsf{Span}}_X(S)$ represented by a map $\phi: \mathsf{Tw}(\Delta^n) \to S$ is called a *Segal simplex* if, for every $\Delta^k \subset \Delta^n$ the composite diagram:

$$\{0,k\} * \mathsf{Tw}(\mathsf{Sp}(\Delta^k)) \subset \mathsf{Tw}(\Delta^k) \subset \mathsf{Tw}(\Delta^n) \xrightarrow{\phi} S$$

is a *p*-limit diagram. Here, $\mathsf{Sp}(\Delta^k)$ denotes the spine of Δ^k and we call the join $\{0, k\} * \mathsf{Tw}(\mathsf{Sp}(\Delta^k))$ the *Segal cone*. We denote by $\mathsf{Span}_X(S) \subset \overline{\mathsf{Span}}_X(S)$ the simplicial subset consisting of Segal simplices. In case $X \simeq *$ we adopt the notation $\mathsf{Span}_*(\mathcal{C}) := \mathsf{Span}(\mathcal{C})$.

It follows from [DK19, Thm.10.2.6] that for every ∞ -category \mathcal{C} the simplicial set Span(\mathcal{C}) is itself ∞ -category.

Proposition 4.9. [DK19, Prop.10.2.31] Let C be an ∞ -category with finite limits. Then the functor:

$$\operatorname{Span}_{\Delta}(\mathfrak{C}^{\times}) \to \Delta$$

is a Cartesian fibration providing a monoidal structure on $\mathsf{Span}(\mathfrak{C})$.

Note, that the construction of the ∞ -category $\text{Span}(\mathbb{C})$ is functorial. Indeed, every finite limit preserving functor $F : \mathbb{C} \to \mathcal{D}$ between ∞ -categories \mathbb{C} and \mathcal{D} induces a functor $\text{Span}(F) : \text{Span}(\mathbb{C}) \to \text{Span}(\mathcal{D})$. In particular, after passing to homotopy coherent nerves the construction Span(-) induces an ∞ -functor

$$\mathsf{Span}(-): \mathsf{Cat}^{\mathsf{lex}}_{\infty} \to \mathsf{Cat}_{\infty}$$

where $\mathsf{Cat}_{\infty}^{\mathsf{lex}}$ denotes the ∞ -category of ∞ -categories and finite product preserving functors.

4.2 Characterization of Bimodules

Let \mathcal{C} be an ∞ -category with finite limits and denote by \mathcal{C}^{\times} the associated Cartesian monoidal structure as constructed in Construction 4.7. Our main goal of this section is to prove the following:

Theorem 4.10. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence of spaces:

$$\mathsf{BiMod}(\mathsf{Span}_{\Lambda}(\mathfrak{C}^{\times}))^{\simeq} \simeq \mathsf{BiSeg}_{\Lambda}(\mathfrak{C})^{\simeq}$$

For better readability we have included some technical lemmas in the Appendix A. As a start, we unravel the datum of a cobimodule in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$. Such an object is given by a commutative triangle:



s.t F preserves inert morphisms and the adjoint morphism $\tilde{F} : \mathsf{Tw}(\Delta_{/[1]}) \to \mathbb{C}^{\times}$ maps every *n*-simplex $\Delta^n \to \Delta_{/[1]}$ to a Segal simplex. A precise discussion of these conditions is included in Appendix A.1. Here we just state the main result:

Proposition 4.11. A functor $F: \Theta_1 \to \mathbb{C}$ defines a bicomodule object, if and only if

- (1) F sends degenerate⁴ intervals to terminal objects.
- (2) F sends ever object $(\phi : g^0_{[n_0]} \to g^1_{[n_1]}, [i, j])$ together with its projection to subintervals to a product diagram⁵.
- (3) F sends morphisms every morphism

$$\sigma \simeq \begin{cases} [i,j] \subset g_{n_0}^0 \xrightarrow{g} g_{n_1}^1 \\ \downarrow^f & \bar{f} \\ \tilde{i}, \tilde{j}] \subset g_{m_0}^0 \xrightarrow{\tilde{g}} g_{m_1}^1 \end{cases}$$

s.t f restricts to an isomorphism $\{i, ..., j\} \rightarrow \{\tilde{i}, ..., \tilde{j}\}$ and \tilde{f} to an isomorphism $\{g(i), ..., g(j)\} \rightarrow \{\tilde{g}(\tilde{i}), ..., \tilde{g}(\tilde{j})\}$, to an equivalence. We denote by E the set of morphisms of the above form.

(4) F maps all Segal cone diagrams from Definition A.1 to limit diagrams.

Definition 4.12. We define $BiMod_{Sp}(\mathcal{C})$ as the full subcategory of $Fun(\Theta_1, \mathcal{C})$ on those functors that satisfy the conditions of Corollary 4.11 and denote by the full subcategory on $Fun^*(\Theta_1, \mathcal{C})$ those functors that map degenerate intervals to terminal objects.

Similarly, we denote by $\operatorname{BiMod}_{\pi}(\mathbb{C})$ the full subcategory of $\operatorname{Fun}(\Theta_1, \mathbb{C})$ on those functors that satisfy condition (1) and (2) of Corollary 4.11. Furthermore, we denote by Ω_1 the full subcategory of Θ_1 with objects those morphisms $([i, j] \subset g_{n_0}^0 \to g_{n_1}^1)$, s.t. the interval [i, j] is non-degenerate.

⁴An interval $[i, j] \subset [n]$ is called degenerate if i = j.

⁵See Construction 4.7

We first analyse condition (1):

Proposition 4.13. The restriction functor induces an equivalence of ∞ -categories

$$\operatorname{Fun}^*(\Theta_1, \mathcal{C}) \xrightarrow{\simeq} \operatorname{Fun}(\Omega_1, \mathcal{C})$$

Proof. The proof is the same as [Ste21, Cor.2.9].

To handle condition (3) of, we explicitly construct a localisation of Ω_1 along the class of morphisms E.

Definition 4.14. The category Δ_1^{\star} has objects consisting of pairs of a finite ordered set [k] and a [k]-indexed sequence of composable⁶ objects $(f_{n_0}, ..., f_{n_k})$ in $\Delta_{/[1]}$. By definition, these compose to a morphism

$$f := f_{n_0} * \dots * f_{n_k} : [n_0] * [n_1] * \dots * [n_k] \to [1]$$

A morphism $(g,\theta):((f_{n_0}^1,...f_{n_k}^1),[k])\to((f_{m_0}^2,...,f_{m_l}^2),[l])$ consists of a

- (1) a morphism $\theta : [l] \to [k]$
- (2) a commutative diagram

$$[m_0]*\ldots*[m_l] \xrightarrow{g} [n_0]*\ldots*[n_k]$$

$$\xrightarrow{f^2} [1] \swarrow f^1$$

s.t. for any
$$i \in [k]$$
 with $\theta^{-1}(i) = \{j_1, ..., j_p\}$, the restriction

$$g_i: [m_{j_1}] * \dots * [m_{j_p}] \to [n_0] * \dots * [n_k]$$

has image contained in $[n_i]$.

Construction 4.15. We define a functor $\mathcal{L} : \Omega_1 \to \Delta_1^*$ as follows:

- It maps an object $[i, j] \subset g_{n_0}^0 \xrightarrow{f} g_{n_1}^1$ to the object $(g_{[f(i), f(i+1)]}^1, ..., g_{[f(j-1), f(j)]}^1)$. The image is indexed by the inner interstices⁷ $\mathbb{I}([i, j])$ of the linearly ordered set $\{i, i+1, ..., j\}$
- A morphism in Ω_1 is given by

$$\sigma \simeq \left\{ \begin{array}{c} [i,j] \subset g_{n_0}^0 \xrightarrow{g} g_{n_1}^1 \\ \downarrow^f & \tilde{f} \\ [\tilde{i},\tilde{j}] \subset g_{m_0}^0 \xrightarrow{\tilde{g}} g_{m_1}^1 \end{array} \right\}$$

The functor \mathcal{L} maps σ to the map $\mathcal{L}(\sigma) = (h_{\sigma}, \mathbb{I}(f))$, whose second component is given by $\mathbb{I}(f) : \mathbb{I}([\tilde{i}, \tilde{j}]) \to \mathbb{I}([i, j])$. We define h_{σ} componentwise. Consider an inner interstices $\{p, p + 1\} \in \mathbb{I}([i, j])$. The restriction of h_{σ} to the component indexed by $\mathbb{I}(f)^{-1}(\{p, p + 1\})$ is given by:

$$[\tilde{g}(f(p)), \tilde{g}(f(p)+1)] * \dots * [\tilde{g}(f(p+1)-1), \tilde{g}(f(p+1))] \xrightarrow{\tilde{f}} [g(p), g(p+1)]$$

It follows from the commutativity of σ that this definition is well-defined.

 6 See Definition 2.26

 $^{^7\}mathrm{See}$ Definition 4.4

The technical proof of the following proposition is included in Appendix A.2:

Proposition 4.16. The functor $\mathcal{L} : \Omega_1 \to \Delta_1^*$ is an ∞ -categorical localisation at the morphisms E defined in Corollary 4.11 (3).

Definition 4.17. Let \mathcal{C} be an ∞ -category with finite limits. We define the category $\mathsf{Fun}^{bim}(\Delta_1^{\star}, \mathcal{C}) \subset \mathsf{Fun}(\Delta_1^{\star}, \mathcal{C})$ as the full subcategory with objects those functors, that

(I) send diagrams of the form



to equivalences

(II) send diagrams of the form



to equivalences

Proposition 4.18. Restriction along the functor $\mathcal{L} : \Omega_1 \to \Delta_1^*$ induces an equivalence of ∞ -categories

$$\mathsf{BiMod}_{Sp}(\mathfrak{C}) \simeq \mathsf{Fun}^{bim}(\Delta_1^{\star}, \mathfrak{C})$$

Proof. It follows from Proposition 4.16 that restriction along \mathcal{L} induces an equivalence

$$\operatorname{Fun}_{E^{-1}}(\Omega_1, \mathfrak{C}) \simeq \operatorname{Fun}(\Delta_1^{\star}, \mathfrak{C})$$

where we denote by $\operatorname{Fun}_{E^{-1}}(\Omega_1, \mathcal{C})$ the full subcategory of $\operatorname{Fun}(\Omega_1, \mathcal{C})$ on those functors, that map all morphisms in E to equivalences. To conclude, we only have to match the remaining conditions of Corollary 4.11. By construction, condition (2) on the left matches with condition (I) on the right. For condition (4) let σ be a 3-simplex

$$g_{n_0}^0 \xrightarrow{\phi_1} g_{n_1}^1 \xrightarrow{\phi_2} g_{n_2}^2 \xrightarrow{\phi_3} g_{n_3}^3$$

in Ω_1 and F an object of $\mathsf{BiMod}_{Sp}(\mathcal{C})$. Further let $[i, j] \subset [n_0]$ be a subinterval. The corresponding limit diagram for condition (4) reads as



It follows from the dual of [Lur06, Prop.4.4.2.2] that this diagram is a limit diagram, if and only if the diagram



is a pullback diagram. Combining this with the previous diagram, we get



It follows from the pasting property for pullbacks, that the diagram corresponding to any sub 2-simplex is a limit diagram. Iterating this we see, that condition (4) is satisfied if and only if it is satisfied on 2-simplices. Since condition (II) is precisely the image of condition (4) on 2-simplices under \mathcal{L} , we can conclude.

Using the above proposition, the problem has shifted to analyzing the conditions of Definition 4.17. Therefore, consider the full subcategory $i : \Delta_{/[1]}^{op} \to \Delta_1^*$ generated by objects $f : [n] \to [1] \in \Delta_{/[1]}$ with $n \ge 0$. Restriction and right Kan extension induce an adjunction

$$i^*: \operatorname{Fun}(\Delta_1^\star, \mathfrak{C}) \rightleftharpoons \operatorname{Fun}(\Delta_{/[1]}^{op}, \mathfrak{C}): i_*$$

We denote by $\operatorname{Fun}^{\times}(\Delta_{1}^{\star}, \mathbb{C}) \subset \operatorname{Fun}(\Delta_{1}^{\star}, \mathbb{C})$ the full subcategory on those functors that satisfy condition (I) of Definition 4.17.

Proposition 4.19. The functor $i_* : \operatorname{Fun}(\Delta_{/[1]}^{op}, \mathbb{C}) \to \operatorname{Fun}(\Delta_1^*, \mathbb{C})$ factors through the full subcategory $\operatorname{Fun}^{\times}(\Delta_1^*, \mathbb{C})$. Moreover, the induced functor

$$i_*: \operatorname{Fun}(\Delta^{op}_{/[1]}, \mathfrak{C}) \to \operatorname{Fun}(\Delta^{\star}_1, \mathfrak{C})$$

is an equivalence of ∞ -categories.

Proof. Let $F \in \mathsf{Fun}(\Delta_{[1]}^{op}, \mathbb{C})$ be a functor $(f_{m_1}, f_{m_2}, ..., f_{m_k})$ an object of Δ_1^{\star} . The value of i_*F on $(f_{m_1}, f_{m_2}, ..., f_{m_k})$ can be computed as the limit of F over the diagram indexed by the category $\mathcal{D} := (\Delta_{[1]}^{op})/(f_{m_1}, f_{m_2}, ..., f_{m_k})$. An object of the category \mathcal{D} consists of a choice $i \in \{1, ..., k\}$ together with a

morphism $h_l \to f_{m_i}$ in $\Delta_{/[1]}$. Note that a morphism from $(h_l \to f_{m_i})$ to $(e_p \to f_{m_j})$ in \mathcal{D} only exists if i = j. In this case, it is given by a commutative diagram



in $\Delta_{[1]}$. It follows, that for the value of $i_*F(f_{m_1},...,f_{m_k})$ is given by the limit



This proves the first claim. The second claim follows from [Lur06, Prop.4.3.2.15]

We need the following auxiliary lemma

Lemma 4.20. Let $F \in \operatorname{Fun}(\Delta_1^*, \mathbb{C})$ be a functor. Then F satisfies condition (II), if and only if it satisfies condition (II) where all but one of the g_2^i have source equal to [1]. We will call this condition (II').

Proof. It follows from the assumptions that (II) implies (II'). Let therefore F satisfy condition (II'). We consider the diagram in Δ_1^* displayed in Figure 1. Since F satisfies (II'), it follows that F maps



Figure 1: Proof of Lemma 4.20

the bottom right square and the exterior right rectangle of the diagram to pullback diagrams. By the pasting law, it follows that the functor F also sends the bottom left square to a pullback square. Hence, it follows from (II') that also the upper left square is a pullback square. Using the pasting law once again implies that the vertical left square is a pullback. Iterating this argument yields the claim.

After all these intermediate steps, we can finally prove:

Proposition 4.21. The restriction functor $i^* : \operatorname{Fun}(\Delta_1^*, \mathbb{C}) \to \operatorname{Fun}(\Delta_{/[1]}^{op}, \mathbb{C})$ descends to an equivalence

$$\operatorname{\mathsf{Fun}}^{\operatorname{bim}}(\Delta_1^{\star}, \mathfrak{C}) \simeq \operatorname{\mathsf{BiSeg}}_{\Delta}(\mathfrak{C})$$

where $\operatorname{BiSeg}_{\Delta}(\mathcal{C}) \subset \operatorname{Fun}(\Delta^{\operatorname{op}}_{/[1]}, \mathcal{C})$ denotes the full subcategory generated by birelative 2-Segal objects.

Proof. Let $F \in \mathsf{Fun}^{\times}(\Delta_1^{\star}, \mathfrak{C})$ be a functor and let

$$\begin{array}{c} g_{1} \longleftarrow & \ast_{0 \leqslant k < j-1} g_{1}|_{\{k,k+1\}} \ast g_{2} \ast (\star_{j \leqslant k < n} g_{1}|_{k,k+1}) \\ & \downarrow & \downarrow \\ (g_{1}|_{\{0,1\}}, ..., g_{1}|_{\{n-1,n\}}) \longleftarrow (g_{1}|_{\{0,1\}}, ..., g_{1}|_{\{j-2,j-1\}}, g_{2}, g_{1}|_{\{j,j+1\}}, ..., g_{1}|_{\{n-1,n\}}) \end{array}$$

be a diagram of type (II'). We can expand this diagram

$$\begin{array}{c} g_{1} < & & \\ & & & \\ & & & \\ & & & \\ & & & \\ (g_{1}|_{\{0,1\}}, \dots, g_{1}|_{\{n-1,n\}}) < & & \\ & & & \\ & & & \\ & & & \\ & & & \\ g_{1}|_{\{j-1,j\}} < & & \\ & & & \\ & & & \\ & & & \\ g_{2} \end{array} (g_{1}|_{\{j,1,1\}} * g_{2} * (*_{j \leq k < n}g_{1}|_{k,k+1}) \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ g_{1}|_{\{j-1,j\}} < & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ &$$

Since F satisfies condition (I) of Definition 4.17 it follows, that F maps the lower square to a pullback diagram. By the pasting lemma, the upper square is a pullback if and only if the outer rectangle is a pullback. But the latter can be identified with the opposite of the diagram



which is precisely one of the birelative 2-Segal conditions.

As a corollary, we obtain analogous results for left and right module objects in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$ Corollary 4.22. Let \mathbb{C} be an ∞ -category with finite limits. There exist equivalences of spaces

$$\mathsf{LMod}(\mathsf{Span}_{\Delta}(\mathfrak{C}^{\times}))^{\simeq} \simeq \mathsf{LSeg}_{\Delta}(\mathfrak{C})^{\simeq}$$

and

$$\mathsf{RMod}(\mathsf{Span}_{\Delta}(\mathcal{C}^{\times}))^{\simeq} \simeq \mathsf{RSeg}_{\Delta}(\mathcal{C})^{\simeq}$$

5 Module Morphisms in Span Categories

In the last section, we have constructed for every ∞ -category with finite limits \mathcal{C} an equivalence between the space of bimodule objects in Span(\mathcal{C}) and of birelative 2-Segal objects. The goal of this section is to extend this to an equivalence of ∞ -categories:

Theorem 5.1. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence of ∞ -categories

$$\operatorname{BiMod}(\operatorname{Span}_{\Delta}(\mathbb{C}^{\times})) \simeq \operatorname{Bi2Seg}_{\Delta}^{\leftrightarrow}(\mathbb{C})$$

between the ∞ -category of bimodule objects in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$ and the ∞ -category of birelative 2-Segal objects and birelative 2-Segal spans.

As a corollary of this we obtain a similar characterization for the ∞ -category of left (resp. right) module objects and algebra objects in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$.

Corollary 5.2. Let C be an ∞ -category with finite limits. There exists an equivalence

$$\mathsf{LMod}(\mathsf{Span}_{\Delta}(\mathfrak{C}^{\times})) \simeq \mathrm{L2Seg}_{\Delta}^{\leftrightarrow}(\mathfrak{C})$$

between the ∞ -category of left modules in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$ and the ∞ -category of left relative 2-Segal objects and left relative 2-Segal spans.

Corollary 5.3. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence

$$\mathsf{Alg}(\mathsf{Span}_{\Delta}(\mathcal{C}^{\times})) \simeq 2\operatorname{-Seg}_{\Delta}^{\leftrightarrow}(\mathcal{C})$$

between the ∞ -category of algebras in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$ and the ∞ -category of 2-Segal objects and 2-Segal spans.

Combining the above Corollary with Corollary 2.19, we obtain

Corollary 5.4. Let \mathcal{C} be an ∞ -category with finite limits. There exists an equivalence

$$\mathsf{LMod}(\mathsf{Span}_{\Lambda}(\mathcal{C}^{\times})) \simeq \mathrm{Rel2Seg}_{\Lambda}^{\leftrightarrow}(\mathcal{C})$$

between the ∞ -category of left modules in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$ and the ∞ -category of relative 2-Segal objects.

Before we prove these statements, let us explicitly describe the relation between algebra morphisms and Segal spans on the level of lowest dimensional simplices. Consider two algebra objects (X_1, μ^X) and (Y_1, μ^Y) in $\text{Span}(\mathbb{C}^{\times})$) with associated 2-Segal objects X_{\bullet} (resp. Y_{\bullet}). To construct an algebra morphism $F : (X_1, \mu^X) \to (Y_1, \mu^Y)$ between those we first have to provide a morphism on underlying objects, which is a span with tip denoted F_1 :



For this span to be part of an algebra morphism, we have to provide higher coherence data. At the lowest level this is given by an invertible 2-morphism $\alpha : F_1 \circ \mu^X \simeq \mu^Y \circ (F_1 \times F_1)$, i.e there has to exist an invertible morphism of spans α :



Since α is invertible, we can decompose α into two equivalences

$$(\partial_1^F, f_2) : F_2 \xrightarrow{\simeq} X_2 \times_{X_1} F_1$$

and

$$(g_2, \partial_2^F, \partial_0^F) : F_2 \xrightarrow{\simeq} Y_2 \times_{Y_1 \times Y_1} F_1$$

These conditions form the lowest dimensional instances of the active equifibered and relative Segal conditions respectively (see Definition 2.2). Similar calculations for higher coherence data on the algebra morphism construct the higher dimensional simplices of the simplicial object F_{\bullet} .

After these initial considerations, we now turn to the proof of Theorem 5.1. For this we will need the following lemma:

Lemma 5.5. Let \mathcal{C} be an ∞ -category with finite limits and consider a morphism:

$$F: \mathsf{Tw}(\Delta^n) \times (\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}) \to \mathfrak{C}$$

s.t. the associated morphism $\tilde{F} : \mathsf{Tw}(\Delta^n) \to \mathsf{Fun}((\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}), \mathfrak{C}) \text{ factors through } \mathsf{Bim}_{Sp}(\mathfrak{C})^{-8}$. Then F defines a morphism:

$$F:\Delta_{/[1]}\times\Delta^n\to\mathsf{Span}_\Delta(\mathcal{C})$$

if and only if for every 2-simplex $\Delta^2 \to \Delta^n \times \Delta_{/[1]}$ of the form

- $(1) \ (f_0:[n_0] \to [1], i) \xrightarrow{\phi_1} (f_1:[n_1] \to [1], i) \xrightarrow{\phi_2} (f_2:[n_2] \to [1], i)$
- $(2) \hspace{0.2cm} (f_{0}:[n_{0}] \rightarrow [1], i) \xrightarrow{i < j} (f_{0}:[n_{0}] \rightarrow [1], j) \xrightarrow{j < k} (f_{0}:[n_{0}] \rightarrow [1], k)$
- $(3) \ (f_0:[n_0] \to [1], i) \xrightarrow{\phi_1} (f_1:[n_1] \to [1], i) \xrightarrow{i < j} (f_2:[n_2] \to [1], j)$
- $(4) \ (f_0:[n_0] \to [1], i) \xrightarrow{i < j} (f_0:[n_0] \to [1], j) \xrightarrow{\phi_1} (f_1:[n_1] \to [1], j)$

the restriction $F|_{\Delta^2}$ is a Segal simplex.

Proof. The only if condition follows from the assumption. For the other direction, we first observe that for every morphism $\phi : f_{n_0} \to f_{n_1}$ in $\Delta_{/[1]}$, every i < j < k and every fixed subinterval $[l, m] \subset [n_0]$ the diagram



is a limit diagram. Indeed, since \tilde{F} factors through $\mathsf{Bim}_{\pi}(\mathfrak{C})$ the diagram is equivalent to a product of diagrams of the form



These are limit diagrams due to condition (II). Let $\sigma : \Delta^2 \to \Delta_{/[1]} \times \Delta^n$ be a 2-simplex represented by a composable pair of morphisms:

$$(f_0: [n_0] \to [1], i) \xrightarrow{\phi_1} (f_1: [n_1] \to [1], j) \xrightarrow{\phi_2} (f_2: [n_2] \to [1], k)$$

 $^{^{8}}$ See Definition 4.12



Figure 2: Image under F of the decomposition of the 2-simplex σ into a 4-simplex. The original 2-simplex is colored blue.

By Proposition A.2 need to check that for every interval $[l,m] \subset [n_0]$ the following square is a limit diagram:



To check this, we include this diagram into a fourfold composite of spans shown in Figure 2. The square is visualized by the blue part in the diagram. We denote the composite rectangle of two squares by the sum of their labels. It follows from (4), that the rectangles (IV), (IV) + (II), from (3) that the rectangles (III) + (VI), (VI) and from (1) that the square (V) are pullback squares. Moreover, as shown above, the square (I) is a pullback diagram. It follows from an iterated application of the pasting lemma that also the blue square is a pullback square.

Proof of Theorem 5.1. We prove the equivalence $\mathsf{CoBiMod}(\mathsf{Span}_{\Delta}(\mathbb{C}^{\times})) \simeq 2 - \mathsf{BiSeg}(\mathbb{C})^{op}$. The claimed result then follows from the equivalence $\mathsf{CoBiMod}(\mathsf{Span}_{\Delta}(\mathbb{C}^{\times})) \simeq \mathsf{BiMod}(\mathsf{Span}_{\Delta}(\mathbb{C}^{\times}))^{op}$. Unraveling definitions and using Proposition A.2 an *n*-simplex $\eta \in \mathsf{CoBiMod}(\mathsf{Span}_{\Delta}(\mathbb{C}^{\times}))_n$ is represented by a morphism

$$\eta: \mathsf{Tw}(\Delta^n) \times (\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}) \to \mathbb{C}$$

such that

(1) for every $0 \leq k \leq n$ the restriction $\eta_{\{k\}} : \{id_k\} \times (\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}) \to \mathbb{C}$ preserves inert morphisms, i.e. defines a cobimodule in $\mathsf{Span}_{\Delta}(\mathbb{C}^{\times})$.

(2) for every 2-simplex $\Delta^2 \to \Delta_{/[1]} \times \Delta^n$ of the form of Lemma A.3 depicted

$$(f_0: [n_0] \to [1], i) \xrightarrow{\phi_1} (f_1: [n_1] \to [1], j) \xrightarrow{\phi_2} (f_2: [n_2] \to [1], k)$$

with $0 \le i \le j \le k \le n$ and every subinterval $[p,q] \subset [n_0]$ the associated Segal cone diagram⁹ is a limit diagram:



. .

The datum of an *n*-simplex in $\mathsf{CoBiMod}(\mathsf{Span}_{\Delta}(\mathcal{C}^{\times}))$ corresponds via adjunction to a functor

 $\eta : \operatorname{\mathsf{Tw}}(\Delta^n) \to \operatorname{\mathsf{Fun}}(\operatorname{\mathsf{Tw}}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}, \mathfrak{C})$

We claim that η factors through the full subcategory

$$\mathsf{BiMod}_{Sp}(\mathfrak{C}) \subset \mathsf{Fun}(\mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\amalg}, \mathfrak{C}))$$

as defined in Definition 4.12. For the evaluation of η on objects of the form id_k , this follows from condition (1) and (2). Hence, let i < j be an object of $\mathsf{Tw}(\Delta^n)$. We need to check the conditions of Definition 4.12. All conditions except condition (3) of Definition 4.12 are satisfied by assumption. For this condition to hold it suffices to check that for every inert morphism $\phi : f_{n_0}^0 \to f_{n_1}^1 \in \Delta_{/[1]}$ and every subinterval $[k, l] \subset [n_0]$ the morphisms

(i) $\eta(\phi, [k, l], i < j) \to \eta(f_{n_1}^1, [\phi(k), \phi(l)], i < j)$

(ii)
$$\eta(\phi, [k, l], i < j) \to \eta(f_{n_0}^0, [k, l], i < j)$$

are equivalences. Therefore, consider the following 2-simplices:

- (i) $(f_{n_0}^0, i) \xrightarrow{\phi} (f_{n_1}^1, i) \xrightarrow{i < j} (f_{n_1}^1, j)$
- (ii) $(f_{n_0}^0, i) \xrightarrow{i < j} (f_{n_0}^0, j) \xrightarrow{\phi} (f_{n_1}^1, j)$

in $\Delta_{/[1]} \times_{\Delta} \Delta^n$. Since $\eta(-, -, i) \in \text{Bim}_{Sp}(\mathbb{C})$ for all $i \in [n]$, the pullback Diagrams 6 associated by η to these 2-simplices exhibit the morphisms in (i) and (ii) as pullbacks of equivalences. Hence they are itself equivalences.

Consequently, we can apply Lemma 5.5. It therefore, suffices to analyze condition (I) - (IV) of Lemma 5.5. Condition (I) and (II) imply that η descends to a morphism

$$\eta: \Delta^n \to \mathsf{Span}(\mathsf{Bim}_{\mathrm{Sp}}(\mathfrak{C}))$$

Under the equivalence $\mathsf{BiMod}_{Sp}(\mathcal{C}) \simeq \mathsf{Fun}^{\mathrm{bim}}(\Delta_1^{\star}, \mathcal{C})$ of Proposition 4.18 condition (III) reads as



 $^9 See$ Definition A.1

where the maps are the image under η of the diagram



in the category $\Delta_1^* \times \mathsf{Tw}(\Delta)$. Similarly condition (IV) is given by



Observe, that the above diagrams are products of individual square shaped diagrams. It therefore suffices to show that each individual square is a pullback diagram. For condition (*III*), we therefore need to check that for every $p \leq r \leq q - 1$ the diagram



is a pullback square. Similarly for condition (IV), we need to check that the diagram:



is a pullback diagram. Under the equivalence of Proposition 4.21, the map η identifies with a functor $\eta : \mathsf{Tw}(\Delta^n) \to \mathsf{Fun}(\Delta^{op}_{/[1]}, \mathbb{C})$. We denote the value of η on an object $i \leq j \in \mathsf{Tw}(\Delta^n)$ by $M^{i,j}_{\bullet}$, if i < j and by X^i_{\bullet} , if i = j.

Under this equivalence condition (III) translates into the condition that for every $f_l \in \Delta_{/[1]}^{op}$ and $0 \le i < j \le n$ the diagram



is a pullback diagram. Similarly, condition (IV) translates into the condition that for every $f_l \in \Delta_{/[1]}^{op}$ and $0 \leq i < j \leq n$ the diagram:



is pullback diagrams. By Lemma 5.6 and Lemma 5.7 the conditions suffice to hold for j = i + 1. But these are precisely the active equifibered and relative Segal conditions. This finishes the proof.

Lemma 5.6. Let $\tilde{\eta} : \Delta^n \to \text{Span}(\text{Fun}(\Delta_{/[1]}^{op}, \mathbb{C}))$ be a functor with adjoint $\eta : \text{Tw}(\Delta^n) \to \text{Fun}(\Delta_{/[1]}^{op}, \mathbb{C})$ and denote by $M^{i,j}_{\bullet}$ the value of η on the object $i \to j \in \text{Tw}(\Delta^n)$ if i < j and by X^i_{\bullet} , if i = j. Further let $f_l : [l] \to [1]$ be an object of $\Delta_{/[1]}$. Then the following are equivalent:

(1) the diagram



is a pullback diagram for all $0 \leq i < j \leq n$.

(2) the diagram



is a pullback diagram for all $0 \leq i < n$.

Proof. (1) implies (2) by assumption. To prove the converse we do induction on the difference p = j - i. The claim is true for p = 1 by assumption. Assume that the result holds for p - 1. Let $0 \le i < j \le n$ be objects of Δ^n with j - i = p. We denote j - 1 by k. Consider the diagram:



The upper left square is a pullback diagram by the induction assumption and the lower square is a pullback diagram by the definition of η . To show that the big horizontal rectangle is a pullback diagram, it therefore suffices to show that the big vertical rectangle is a pullback diagram. To show this, we factor the vertical rectangle as:



The lower square is a pullback square by the induction hypothesis and the upper square by the definition of η . It follows that the outer square is a pullback square. This finishes the proof.

Lemma 5.7. Let $\tilde{\eta} : \Delta^n \to \text{Span}(\text{Fun}(\Delta_{/[1]}^{op}, \mathbb{C}))$ be a functor with adjoint $\eta : \text{Tw}(\Delta^n) \to \text{Fun}(\Delta_{/[1]}^{op}, \mathbb{C})$ and denote by $M_{\bullet}^{i,j}$ the value of η on the object $i \to j \in \text{Tw}(\Delta^n)$ if i < j and by X_{\bullet}^i , if i = j. Further let $f_l : [l] \to [1]$ be an object of $\Delta_{/[1]}$. Then the following are equivalent: • the diagram



is a pullback diagram for all $0 \leq i < j \leq n$.

• the diagram



is a pullback diagram for all $0 \leq i < n$.

Proof. The proof is analogous to the one of Lemma 5.6. By assumption, (1) implies (2). For the converse, we do induction on the difference p = j - i. The case p = 1 follows by assumption. Assume therefore the result holds for p-1. Let $0 \le i < j \le n$ be objects of Δ^n and denote j-1 by k. Consider the diagram:

It follows from the inductive hypothesis that the bottom square is a pullback diagram. Further, the upper left square is a pullback by definition of η . Therefore, to show that the big vertical rectangle is a pullback diagram, it suffices to show that the big horizontal rectangle is a pullback. The horizontal rectangle factors as



The left square is a pullback square by the induction hypothesis and the right square is a pullback square by definition of η . Consequently, also the big horizontal square is a pullback diagram.

A Auxiliary Statements

The goal of this appendix is to present the technical proofs of Corollary 4.11 and 4.16. These proofs are included here to maintain the flow of the main argument presented in Section 4.

A.1 Proof of Proposition 4.11

The goal of this subsection is to prove Proposition 4.11. As a start, we unravel the datum of a cobimodule in $\text{Span}_{\Delta}(\mathbb{C}^{\times})$. Such an object is given by a commutative triangle:



s.t F preserves inert morphisms and the adjoint morphism $\tilde{F} : \mathsf{Tw}(\Delta_{/[1]}) \to \mathfrak{C}^{\times}$ maps every *n*-simplex $\Delta^n \to \Delta_{/[1]}$ to a Segal simplex¹⁰.

We analyze the second condition first. By the universal property of the Cartesian monoidal structure \tilde{F} is equivalent to a functor:

$$\overline{F}: \mathsf{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod} \to \mathcal{C}$$

Similarly an *n*-morphism $\eta: \Delta^n \to \mathsf{Fun}_{\Delta}(\Delta_{/[1]}, \mathsf{Span}_{\Delta}(\mathbb{C}^{\times}))$ is equivalent to a functor:

$$\bar{\eta}:\mathsf{Tw}(\Delta_{/[1]}\times\Delta^n)\times_{\Delta}\Delta^{\coprod}\simeq\mathsf{Tw}(\Delta^n)\times(\mathsf{Tw}(\Delta_{/[1]})\times_{\Delta}\Delta^{\coprod})\to\mathbb{C}$$

We will denote $\Theta_{n+1} := \mathsf{Tw}(\Delta_{/[1]} \times \Delta^n) \times_{\Delta} \Delta^{\coprod} \simeq \mathsf{Tw}(\Delta^n) \times \Theta_1$. The objects of this category are given by:

$$[i,j] \subset (g_p:[p] \to [1],k) \xrightarrow{f} (g_q:[q] \to [1],l)$$

where $0 \leq k \leq l \leq n, f : g_p \to g_q$ is a morphism in $\Delta_{/[n]}$ and [i, j] is a subinterval of [p].

Definition A.1. Let $\bar{\eta}$: $\mathsf{Tw}(\Delta^n \times \Delta_{/[1]}) \times_{\Delta} \Delta^{[]} \to \mathfrak{C}$ be a morphism of simplicial sets, $\sigma : \Delta^k \to \Delta^n \times \Delta_{/[1]}$ a k-simplex:

$$(g_0, i_0) \xrightarrow{\phi_1} \dots \xrightarrow{\phi_k} (g_k, i_k)$$

and $[i, j] \subset [n_0]$ a subinterval. The Segal cone of $\bar{\eta}$ associated to σ and [i, j] is the diagram:

$$\bar{\eta}(\psi_n, [i, j], i_0 < i_k) \xrightarrow{\bar{\eta}(\psi_n, [i, j], i_0 < i_k)} \bar{\eta}(\phi_k, [\psi_{k-1}(i), \psi_{k-1}(j)], i_{k-1} < i_k) \xrightarrow{\bar{\eta}(g_0, [i, j], i_0)} \bar{\eta}(g_1, [\psi(i), \psi(j)], i_1) \cdots \bar{\eta}(g_{k-1}, [\psi_{k-1}(i), \psi_{k-1}(j)], i_{k-1}) \xrightarrow{\bar{\eta}(g_k, [\psi_k(i), \psi_k(j)], i_k)} \bar{\eta}(g_k, [\psi_k(i), \psi_k(j)], i_k)$$

where we denote $\psi_i := \phi_i \circ \phi_{i-1} \circ \ldots \circ \phi_1$.

Proposition A.2. Let $\bar{\eta}$: $\mathsf{Tw}(\Delta^n \times \Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod} \to \mathbb{C}$ be a morphism of simplicial sets. Its adjoint η defines an n-morphism in $\mathsf{Fun}_{\Delta}(\Delta_{/[1]}, \mathsf{Span}_{\Delta}(\mathbb{C}^{\times}))$ if and only if for every k-simplex:



 $\underbrace{in \ \Delta_{/[1]} \times \Delta^n \text{ and every interval } \{i, j\} \subset [n_0] \text{ the associated Segal cone of } \bar{\eta} \text{ is a limit diagram in } \mathbb{C}.$ $\underbrace{in \ \Delta_{/[1]} \times \Delta^n \text{ and every interval } \{i, j\} \subset [n_0] \text{ the associated Segal cone of } \bar{\eta} \text{ is a limit diagram in } \mathbb{C}.$

Proof. By definition, we need to show that for every k-simplex $\Delta^k \to \Delta^n \times \Delta_{/[1]}$ given by

$$(g_0, i_0) \rightarrow (g_1, i_1) \rightarrow \dots \rightarrow (g_k, i_k)$$

the restriction of $\bar{\eta}$ to $\mathsf{Tw}(\Delta^k) \subset \mathsf{Tw}(\Delta_{/[1]} \times \Delta^n)$ is a Segal simplex in \mathfrak{C}^{\times} . According to [DK19, Lem.10.2.13] there exists a functor:

$$\tilde{H}_t: (\Delta^1 \times \mathsf{Tw}(\Delta^k)) \times_\Delta \Delta^{\coprod} \to \mathfrak{C}$$

adjoint to a homotopy:

$$H_t: (\Delta^1 \times \mathsf{Tw}(\Delta^k)) \to \mathfrak{C}^{\times}$$

such that $H_1 = \bar{\eta}$, the components of this homotopy are Cartesian morphisms and H_0 has image contained in $\mathcal{C}_{g_0}^{\times}$. Consequently, the restriction of $\bar{\eta}$ to the Segal cone¹¹ is a relative limit in \mathcal{C}^{\times} if and only if the restriction of H_0 to the Segal cone is a limit in $\mathcal{C}_{g_0}^{\times}$ [Lur06, Prop. 4.3.1.9].

We can check the latter for each interval $\{i, j\} \subset [n_0]$ individually. The restriction of \tilde{H}_0 to the Segal cone is represented in \mathcal{C} by the diagram

Since the components of the homotopy are Cartesian, it restricts to an equivalence between this diagram and the Segal cone diagram of $\bar{\eta}$ associated to σ and [i, j]. Therefore the simplex is Segal, if for all simplices σ and all subintervals [i, j], the Segal cone diagram of $\bar{\eta}$ associated to σ and [i, j] are limit diagrams.

In fact, it turns out to be sufficient to check this condition for 2-simplices:

Lemma A.3. Let \mathcal{C} be an ∞ -category with finite limits. For an n-simplex

$$F: \Delta^n \to \mathsf{Fun}_\Delta(\Delta_{/[1]}, \mathsf{Span}_\Delta(\mathfrak{C}^{\times}))$$

the following are equivalent:

- (1) F restricts to a functor $F: \Delta^n \to \operatorname{Fun}_{\Delta}(\Delta_{/[1]}, \operatorname{Span}_{\Delta}(\mathbb{C}^{\times}))$
- (2) for every non-degenerate 2-simplex $\Delta^2 \to \Delta_{/[1]} \times \Delta^n$ the restriction $F|_{\Delta^2}$ is a Segal simplex

Proof. This follows from the iterated application of [Lur06, Prop.4.2.3.8]

To study the first condition, recall that a morphism $\phi : \Delta^1 \to \Delta_{/[1]}$ is called inert if its image in Δ is inert. If we depict a morphism in $\Delta_{/[1]}$ by a commutative diagram



the morphism is inert when f is the inclusion of a subinterval. Let $\phi : \Delta^1 \to \Delta_{/[1]}$ be an inert morphism and F a bicomodule object in $\operatorname{Span}_{\Delta}(\mathbb{C}^{\times})$. By definition $F : \Delta_{/[1]} \to \operatorname{Span}_{\Delta}(\mathbb{C}^{\times})$ has to map inert morphisms to Cartesian morphisms. Unraveling the definition of the Cartesian fibration $\operatorname{Span}_{\Delta}(\mathbb{C}^{\times})$, it follows that the image of ϕ is Cartesian, if and only if the adjoint map $\widetilde{F \circ \phi} : \operatorname{Tw}(\Delta^1) \to \mathbb{C}^{\times}$ maps all morphisms in $\operatorname{Tw}(\Delta^1)$ to Cartesian morphisms. In other words for any morphism $\phi : g_{n_0}^0 \to g_{n_1}^1$ and any interval $[i, j] \subset [n_0]$

 $^{^{11}\}mathrm{See}$ Construction 4.8

- (1) the morphism $F(\phi, \{i, j\}) \to F(g_{n_0}^0, \{i, j\})$ induced by the source map $\phi \to g_{n_0}^0$ in $\mathsf{Tw}(\Delta_{/[1]})$ is an equivalence
- (2) the morphism $F(\phi, \{i, j\}) \to F(g_{n_1}^1, \{\phi(i), \phi(j)\})$ induced by the target map $\phi \to g_{n_1}^1$ in $\mathsf{Tw}(\Delta_{/[1]})$ is an equivalence

We derive some consequences of this

Lemma A.4. Suppose $F : \Delta_{/[1]} \to \operatorname{Span}_{\Delta}(\mathbb{C}^{\times})$ represents a cobimodule. Let $f : g_{n_0}^0 \to g_{n_1}^1$ be a morphism in $\Delta_{/[1]}$ viewed as an object in $\operatorname{Tw}(\Delta_{/[1]})$

(1) Denote for $[i,j] \subset [n_0]$ by $f|_{[i,j]}$ the restriction of f to $g^0_{[i,j]} \subset g^0_{n_0}$. Then the induced morphism

$$F(f|_{[i,j]}, [i,j]) \to F(f, [i,j])$$

is an equivalence.

(2) Let $\tilde{f}: g_{n_0}^0 \to g_{[l,k]}^1$ be a morphism with $[l, ..., k] \subset [n_1]$, s.t. the composite with the inert morphism $\phi: g_{[l,k]}^1 \hookrightarrow g_{n_1}^1$ is f. Then the induced morphism

$$F(f, [l, k]) \rightarrow F(\tilde{f}, [l, k])$$

is an equivalence.

Proof. The proof is analogous to [Ste21, Prop.2.2]

Lemma A.5. Suppose $F : \Delta_{/[1]} \to \text{Span}_{\Delta}(\mathbb{C}^{\times})$ represents a cobimodule. Let σ be a morphisms in $\text{Tw}(\Delta_{/[1]}) \times_{\Delta} \Delta^{\coprod}$ of the form:

$$\sigma \simeq \left\{ \begin{array}{c} [i,j] \subset g_{n_0}^0 \xrightarrow{g} g_{n_1}^1 \\ f & \uparrow \\ [\tilde{i},\tilde{j}] \subset g_{m_0}^0 \xrightarrow{g} g_{m_1}^1 \end{array} \right\}$$

s.t. f restricts to an isomorphism $[i, ..., j] \rightarrow [\tilde{i}, ..., \tilde{j}]$ and \tilde{f} restricts to an isomorphism $[g(i), ..., g(j)] \rightarrow [\tilde{g}(\tilde{i}), ..., \tilde{g}(\tilde{j})]$. Then F sends σ to an equivalence.

Proof. We decompose σ into the diagram

$$\begin{array}{c} g^0_{[i,j]} \xrightarrow{g_{[i,j]}} g^1_{n_1} \\ \downarrow & & \uparrow^{\mathrm{id}} \\ g^0_{n_0} \xrightarrow{g} g^1_{m_1} \\ f \downarrow & & \uparrow^{\bar{f}} \\ g^0_{m_0} \xrightarrow{g} g^1_{m_1} \end{array}$$

Note that the upper square gets mapped by F to an equivalence and that the composite of the left vertical maps is inert. So we can restrict by 2-out-of-3 to those σ , s.t the morphism f is inert. Such a

diagram σ can be further decomposed as:

$$\begin{array}{c} g^0_{[i,j]} \xrightarrow{g_{[i,j]}} g^1_{n_1} \\ \downarrow & \uparrow^{\bar{f}} \\ g^0_{[i,j]} \xrightarrow{\bar{g} \circ f} g^1_{m_1} \\ f \downarrow & \uparrow^{\mathrm{id}} \\ g^0_{m_0} \xrightarrow{\bar{q}} g^1_{m_1} \end{array}$$

Using the same argument as above applied to the lower square, we can reduce to the case that f = id. So let σ be a diagram of the form:

$$\begin{array}{ccc} g^0_{n_0} \xrightarrow{g_{[i,j]}} g^1_{n_1} \\ & & & \uparrow \\ \mathrm{id} & & & \uparrow \\ g^0_{n_0} \xrightarrow{g} g^1_{n_1} \end{array}$$

s.t \bar{f} induces an isomorphism $[\bar{g}(0), \bar{g}(n_0)] \rightarrow [g(0), g(n_0)]$. In particular, there exists a decomposition

$$\begin{array}{cccc} g_{n_0}^0 & \stackrel{g}{\longrightarrow} g_{n_1}^1 \\ & & & & \uparrow^{\bar{f}} \\ g_{n_0}^0 & \stackrel{\bar{g}}{\longrightarrow} g_{m_1}^1 \\ & & & \downarrow^{\psi} \\ g_{n_0}^0 & \stackrel{z}{\xrightarrow{\bar{g}}} g_{[\bar{g}(0),\bar{g}(n_0)]}^1 \end{array}$$

with ψ in ert. It follows by assumption that $f \circ \psi$ is itself in ert. Hence the claim follows from 2-out-of-3 again.

Putting these results together we have proven:

Proposition A.6. A functor $F: \Theta_1 \to \mathbb{C}$ defines a bicomodule object if and only if

- (1) F sends degenerate intervals to terminal objects.
- (2) F sends ever object $(\phi : g^0_{[n_0]} \to g^1_{[n_1]}, [i, j])$ together with its projection to subintervals to a product diagram¹².
- (3) F sends morphisms of the form

$$\sigma \simeq \left\{ \begin{array}{c} [i,j] \subset g_{n_0}^0 \xrightarrow{g} g_{n_1}^1 \\ \downarrow^f & \tilde{f} \\ [\tilde{i},\tilde{j}] \subset g_{m_0}^0 \xrightarrow{\tilde{g}} g_{m_1}^1 \end{array} \right\}$$

s.t. the morphism f restricts to an isomorphism $\{i, ..., j\} \rightarrow \{\tilde{i}, ..., \tilde{j}\}$ and the morphism \tilde{f} restricts to an isomorphism $\{g(i), ..., g(j)\} \rightarrow \{\tilde{g}(\tilde{i}), ..., \tilde{g}(\tilde{j})\}$ to equivalences. We denote by E the wide subcategory of Θ_1 with morphisms of the above form.

 $^{^{12}}$ See Construction 4.7

(4) F maps all Segal cone diagrams from Definition A.1 to limit diagrams.

A.2 Proof of Proposition 4.16

The goal of this subsection is to prove the following:

Proposition A.7. The functor $\mathcal{L} : \Omega_1 \to \Delta_1^*$ from Construction 4.15 is an ∞ -categorical localisation at the morphisms E defined in Corollary 4.11 (3).

To prove this we show that this functor satisfies the assumptions of [Wal21, Lem.3.1.1]. Therefore, we need to consider for any object $N = (e_{m_0}^0, ..., e_{m_{k-1}}^{k-1}) \in \Delta_1^{\star}$ the strict fiber of the functor \mathcal{L} at N. We denote this fiber by $\Omega_{1,N}$ and by $\Omega_{1,N}^E$ the subcategory of $\Omega_{1,N}$ with morphisms in E. In the first step, we construct an explicit initial object in $\Omega_{1,N}^E$

Construction A.8. For any $N = (e_{m_0}^0, ..., e_{m_{k-1}}^{k-1}) \in \Delta_1^*$ as above we define the object $I_N \in \Omega_{1,N}$ as

$$[0,k] \subset [k] \xrightarrow{f_N} [m_0] * \dots * [m_{k-1}]$$

$$e_k \xrightarrow{e_k} [1] \xleftarrow{e_{m_0}^0} * \dots * e_{m_{k-1}}^{k-1}$$

where the morphism f_N is defined by

$$f_N(i) = \begin{cases} 0 \in [m_i] & i \le k \\ m_{k-1} \in [m_{k-1}] & i = k \end{cases}$$

and e_k is the unique morphism that makes the triangle commute. In case that $e_{m_0}^0 * \ldots * e_{m_{k-1}}^{k-1}$ is supported at $\{0\}$ we denote by $I_N * id_{[1]}$ the object

$$[0,k] \subset [k] \xrightarrow{f_N} [m_0] * \dots * [m_{k-1}] * [1]$$

Similarly we denote by $\mathsf{id}_{[1]} * I_N$ the analogously defined object for the case that $e_{m_0}^0 * \ldots * e_{m_{k-1}}^{k-1}$ is supported at $\{1\}$.

Lemma A.9. For every N the category $\Omega_{1,N}^E$ has an initial object given by

- (1) $I_N * \operatorname{id}_{[1]} if e_{m_0}^0 * \dots * e_{m_{k-1}}^{k-1}$ is supported at 0.
- (2) I_N if $e_{m_0}^0 * ... * e_{m_{k-1}}^{k-1}$ is surjective.
- (3) $\operatorname{id}_{[1]} * I_N$ if $e_{m_0}^0 * \dots * e_{m_{k-1}}^{k-1}$ is supported at 0.

Proof. We will prove that the objects are initial. Let

$$[i,j] \subset [n] \xrightarrow{\alpha} [l] \xrightarrow{h} [1]$$

be an object in $\Omega_{1,N}^E$. We will construct a unique morphism Φ in $\Omega_{1,N}^E$ from the claimed initial object. In any of the above cases, we define the morphism $\phi : [k] \to [n]$ as the inert map that includes [k] as the interval $\{i, ..., j\}$. This is uniquely defined since every morphism in E has to induce an isomorphism between the chosen subintervals. We can decompose¹³ the object $[l] \xrightarrow{h} [1]$ into

$$[l_{left}] * \{\alpha(i), ..., \alpha(j)\} * [l_{right}] \xrightarrow{h_{left} * h * h_{right}} [1]$$

 $^{^{13}}$ See Construction 4.6

Since any morphism in E has to map $\{\alpha(i), ..., \alpha(j)\}$ isomorphically onto $[f_N(0), f_N(k)] = [n]$ the restriction of Φ to $\{\alpha(i), ..., \alpha(j)\}$ is uniquely determined. We show that our assumptions allow us to uniquely extend this arrow to the outer part. In case (2), we can uniquely extend. We only discuss case (1), since (3) is analogous. In case (1), we can uniquely extend Φ to $[l_{left}]$ by setting it constantly 0. An extension to $[l_{right}]$ is equivalent to a morphism to $\mathrm{id}_{[1]}$ in $\Delta_{/[1]}$. This morphism exists uniquely, since $\mathrm{id}_{[1]}$ is a final object of $\Delta_{/[1]}$.

We further need to show that the inclusion $\Omega_{1,N}^E \subset \Omega_{/N}^1$ is cofinal. Therefore we make the following preliminary considerations. By Quillen's theorem A [Lur06, Thm.4.1.3.1] this amounts to showing, that for every object

$$\{i, j\} \subset [n] \xrightarrow{\alpha} [l] \xrightarrow{h} [1]$$

whose image under \mathcal{L} is given by $(h_{l_{i+1}}, ..., h_{l_i})$, and every morphism

$$g: (h_{l_{i+1}}, ..., h_{l_j}) \to (e_{m_0}, ..., e_{m_{k-1}})$$

in Δ_1^{\star} the category $\Omega_{N,1}^E \times_{(\Omega_1)/N} ((\Omega_1)_{/N})_{g/}$ is contractible. We do so by constructing an explicit initial object. By definition, the morphism g amounts to a pair consisting of a morphism $\gamma : [k-1] \rightarrow \{i+1,...,j\}$ in Δ and a morphism $\bar{g} : [m_0] * ... * [m_{k-1}] \rightarrow [l_{i+1}] * ... * [l_j]$ in $\Delta_{/[1]}$. We denote by $[n_c] := \{p, ..., q\} \subset \{i, j\} \subset [n]$ the unique linearly ordered set s.t γ factors as

$$\gamma: [k-1] \to [n_c] \hookrightarrow [n_l] * [n_c] * [n_r]$$

The morphism \bar{g} has by construction image contained in $[l_c] := [l_{p+1}] * ... * [l_q]$. We introduce the following decompositions:

$$[l] = [l_l] * [l_c] * [l_r]$$
$$[m] = [m_l] * [m_c] * [m_r]$$

Lemma A.10. There exists a morphism in Ω_1

that extends to a morphism

$$\mu_{Z,N} := \left\{ \begin{array}{c} Z = [i,j] \subset d_{n_l} * d_{n_c} * d_{n_r} \xrightarrow{} h_{l_l} * h_{l_c} * h_{l_r} \\ \downarrow & \uparrow \\ Z_N = [0,k] \subset d_{n_l} * h_{\{0,l^1\}} * e_k * h_{\{0,l^2\}} * d_{n_r} \xrightarrow{} h_{l_l} * h_{l^1} * e_m * h_{l^2} * h_{l_r} \end{array} \right\}$$

covering g, s.t. $\mu_{Z,N}$ is an initial object in $\Omega^E_{N,1} \times_{(\Omega_1)/N} ((\Omega_1)/N)_{g/N}$

Proof. We define a morphism $\nu : \{p + 1, ..., q - 1\} \rightarrow [k]$ as $\mathbb{I}(\gamma)$ and extend it to a morphism $\nu : \{p, ..., q\} \rightarrow [1] * [k] * [1]$ by mapping endpoints to endpoints. We decompose $[l_c] = [l^1] * [l_c^m] * [l^2]$, where we denote by $[l_c^m]$ the subinterval of [l], that contains the image of \bar{g} . By construction, the map $\bar{g} : e_m \rightarrow h_{[l_c^m]}$ hits both endpoints. Therefore, we can extend it to a morphism

$$\bar{g}' := \mathsf{id}_{h_{l^1}} *g * id_{h_{l^2}} : h_{l^1} * e_m * h_{l^2} \to h_{l^c}$$

By construction \bar{g}' hits both endpoints and extends uniquely to a morphism over [1]. Similarly, we define

$$f_N: h_{\{0,l^1\}} * e_k * h_{\{0,l^2\}} \to h_{l^1} * e_m * h_{l^2}$$

to be f_N on e_k and to send endpoints to endpoints. This morphism also extends uniquely to a morphism over [1]. By decomposing the morphisms ν , h and $\bar{f}_N \circ \nu$, we obtain a not necessarily commuting diagram

$$\begin{array}{c} [p,q] \subset d_{[1_{p+1}]} * \dots * d_{[1_p]} & \longrightarrow & h_{l_{p+1}} * \dots * h_{l_q} \\ & \downarrow^{\nu} & & \bar{g'} \\ [0,k] \subset h_{\{0,l^1\}} * e_{k_{p+1}} * \dots * e_{k_q} * h_{\{0,l^2\}} & \longrightarrow & h_{l^1} * e_{m_{p+1}} * \dots * e_{m_q} * h_{l^2} \end{array}$$

This diagram commutes in $\Delta_{[1]}$ if and only if it commutes restricted to each individual $[1_r]$ with $r \in \{p + 1 \leq q\}$. Since all maps preserve endpoints, this is clear for r = p + 1 and r = q. In the other cases, it suffices to show, that \bar{g}' sends the endpoints of $[n_r]$ to the endpoints of $[m_r]$. This follows since \bar{g}' is induced by a morphism g in Δ_1^* .

Since all morphisms preserve endpoints, we can extend the diagram by taking star products with the morphisms $\mathrm{id}_{d_{n_l}}$, $\mathrm{id}_{d_{n_l}}$, $\mathrm{id}_{h_{l_l}}$, $\mathrm{id}_{h_{l_r}}$, $\alpha|_{d_{n_l}}: d_{n_l} \to d_{l_l}$ and $\alpha|_{d_{n_r}}: d_{n_r} \to d_{l_r}$

By construction, this diagram defines a morphism in Ω_1 covering g. We call this morphism $\mu_{Z,N}$. Let us show that $\mu_{Z,N}$ is initial. Suppose, we are given another morphism in Ω_1

covering g. We can decompose it as follows

It follows from [Ste21, Lem. A.3] that ρ is uniquely determined except at the endpoints by γ . Therefore, the restriction of the diagram Diagram 7 to d_{n_c} looks as follows:

$$\begin{array}{c} d_{n_c} \xrightarrow{h} h_{l_c} \\ & \downarrow^{\rho} & \downarrow^{\rho} \\ x_{a_c^1} * e_k * x_{a_c^2} \xrightarrow{\beta_c^1 * f_N * \beta_c^2} y_{b_c^1} * e_n * y_{b_c^2} \end{array}$$

Every morphism $Z_N \to X$ in E commuting with the above morphism $Z \to X$ and $\mu_{Z,M}$, must in

particular restrict to a commutative diagram



Moreover, since the bottom square is in E it must restrict to a commutative diagram



It follows that the component $h_{\{0,l^1\}} * e_k * h_{\{0,l^2\}} \to x_{a_c^1} * e_k * x_{a_c^2}$ is uniquely determined by the left hand triangle in Diagram 8. We can further decompose w, since it must restrict to \bar{g} on e_m , into

$$w = w^{1} * g * w^{2} : y_{b_{c}^{1}} * e_{m} * y_{b_{c}^{2}} \to h_{l^{1}} * h_{l_{c}^{m}} * h_{l^{2}}$$

Hence the component of the bottom morphism in the right triangle is uniquely determined and must be given by

$$w^1 * \mathsf{id}_{[m]} * w^2 : y_{b_c^1} * e_m * y_{b_c^2} \to h_{l^1} * e_m * h_{l^2}$$

We can now extend back to the full diagram



Note that the vertical arrows in the back square are given by identities when restricted to h_{l_l} , h_{l_r} , e_{m_l} and e_{m_r} . Hence the bottom square is uniquely determined by the morphisms $h_{l_l} \to x_{a_l}$, $h_{l_r} \to x_{a_r}$, $y_{b_l} \to e_{m_l}$ and $y_{b_r} \to e_{m_r}$. So there is a unique morphism $Z_N \to X$ with the desired properties.

Proof of Proposition 4.16. The result follows from combining Lemma A.10 and Lemma A.9 with [Wal21, Lem.3.1.1.]. \Box

References

- [Bar15] Clark Barwick. On exact ∞-categories and the theorem of the heart. Compositio Mathematica, 151(11):2160–2186, 2015.
- [BHS22] Shaul Barkan, Rune Haugseng, and Jan Steinebrunner. Envelopes for algebraic patterns. arXiv preprint arXiv:2208.07183, 2022.

- [BOO⁺20] Julia Bergner, Angélica Osorno, Viktoriya Ozornova, Martina Rovelli, and Claudia Scheimbauer. The edgewise subdivision criterion for 2-Segal objects. Proceedings of the American Mathematical Society, 148(1):71–82, 2020.
- [Car20] Louis Carlier. Incidence bicomodules, möbius inversion and a rota formula for infinity adjunctions. Algebraic & Geometric Topology, 20(1):169–213, 2020.
- [CDH⁺20] Baptiste Calmès, Emanuele Dotto, Yonatan Harpaz, Fabian Hebestreit, Markus Land, Kristian Moi, Denis Nardin, Thomas Nikolaus, and Wolfgang Steimle. Hermitian K-theory for stable ∞-categories ii: cobordism categories and additivity. arXiv preprint arXiv:2009.07224, 2020.
- [DJW19] Tobias Dyckerhoff, Gustavo Jasso, and Tashi Walde. Simplicial structures in higher Auslander–Reiten theory. *Advances in Mathematics 355*, 2019.
- [DK18] Tobias Dyckerhoff and Mikhail Kapranov. Triangulated surfaces in triangulated categories. Journal of the European Mathematical Society, 20(6):1473–1524, 2018.
- [DK19] Tobias Dyckerhoff and Mikhail Kapranov. *Higher Segal spaces*, volume 2244. Springer, 2019.
- [DKSS21] Tobias Dyckerhoff, Mikhail Kapranov, Vadim Schechtman, and Yan Soibelman. Spherical adjunctions of stable ∞-categories and the relative S-construction. arXiv preprint arXiv:2106.02873, 2021.
- [DPS22] Duiliu-Emanuel Diaconescu, Mauro Porta, and Francesco Sala. Cohomological Hall algebras and their representations via torsion pairs. *arXiv preprint arXiv:2207.08926*, 2022.
- [GCKT18] Imma Gálvez-Carrillo, Joachim Kock, and Andrew Tonks. Decomposition spaces, incidence algebras and möbius inversion i: basic theory. *Advances in Mathematics*, 331:952–1015, 2018.
- [GH15] David Gepner and Rune Haugseng. Enriched ∞ -categories via non-symmetric ∞ -operads. Advances in mathematics, 279:575–716, 2015.
- [GHN17] David Gepner, Rune Haugseng, and Thomas Nikolaus. Lax colimits and free fibrations in ∞-categories. *Documenta Mathematica 22 1225-1266*, 2017.
- [HM15] Lars Hesselholt and Ib Madsen. Real algebraic K-theory. Book project in progress, 2015.
- [HS04] Jens Hornbostel and Marco Schlichting. Localization in hermitian K-theory of rings. Journal of the London Mathematical Society 70.1: 77-124, 2004.
- [HSV19] Hadrian Heine, Markus Spitzweck, and Paul Verdugo. Real K-theory for Waldhausen infinity categories with genuine duality. *arXiv preprint arXiv:1911.11682*, 2019.
- [KV22] Mikhail Kapranov and Eric Vasserot. The cohomological Hall algebra of a surface and factorization cohomology. Journal of the European Mathematical Society 25.11: 4221-4289., 2022.
- [Lur06] Jacob Lurie. Higher Topos Theory. arXiv preprint math/0608040, 2006.
- [Lur17] Jacob Lurie. Higher Algebra, 2017.
- [Pog17] Thomas Poguntke. Higher Segal structures in algebraic K-theory. arXiv preprint arXiv:1709.06510, 2017.
- [PS23] Mauro Porta and Francesco Sala. Two-dimensional categorified Hall algebras. Journal of the European Mathematical Society 25.3: 1113-1205., 2023.

- [Seg68] Graeme Segal. Classifying spaces and spectral sequences. Publications Mathématiques de l'IHÉS, Volume 34, 1968.
- [Ste21] Walker H Stern. 2-Segal objects and algebras in spans. Journal of Homotopy and Related Structures, 16(2):297–361, 2021.
- [Wal06] Friedhelm Waldhausen. Algebraic K-theory of spaces. Algebraic and Geometric Topology: Proceedings of a Conference held at Rutgers University, New Brunswick, USA, Springer Berlin Heidelberg,, 2006.
- [Wal16] Tashi Walde. Hall monoidal categories and categorical modules. arXiv preprint arXiv:1611. 08241, 2016.
- [Wal21] Tashi Walde. 2-Segal spaces as invertible infinity-operads. Algebraic & Geometric Topology, 21(1):211–246, 2021.
- [You18] Matthew Young. Relative 2-Segal spaces. Algebraic & Geometric Topology, 18(2):975–1039, 2018.
- [You20] Matthew B Young. Representations of cohomological Hall algebras and Donaldson– Thomas theory with classical structure groups. *Communications in Mathematical Physics*, 380(1):273–322, 2020.