BALL CHARACTERIZATIONS IN PLANES AND SPACES OF CONSTANT CURVATURE, II THIS PDF-FILE IS NOT IDENTICAL WITH THE PRINTED PAPER.

J. JERÓNIMO-CASTRO*, E. MAKAI, JR.**

ABSTRACT. High proved the following theorem. If the intersections of any two congruent copies of a plane convex body are centrally symmetric, then this body is a circle. In our paper we extend the theorem of High to the sphere and the hyperbolic plane, and partly to spaces of constant curvature. We also investigate the dual question about the convex hull of the unions, rather than the intersections.

Let us have in H^2 proper closed convex subsets K, L with interior points, such that the numbers of the connected components of the boundaries of K and L are finite. We exactly describe all pairs of such subsets K, L, whose any congruent copies have an intersection with axial symmetry; there are nine cases. (The cases of S^2 and \mathbb{R}^2 were described in Part I, i.e., [5].)

Let us have in S^d , \mathbb{R}^d or H^d proper closed convex C^2_+ subsets K, L with interior points, such that all sufficiently small intersections of their congruent copies are symmetric w.r.t. a particular hyperplane. Then the boundary components of both K and L are congruent, and each of them is a sphere, a parasphere or a hypersphere.

Let us have a pair of convex bodies in S^d , \mathbb{R}^d or H^d , which have at any boundary points supporting spheres (for S^d of radius less than $\pi/2$). If the convex hull of the union of any congruent copies of these bodies is centrally symmetric, then our bodies are congruent balls (for S^d of radius less than $\pi/2$). An analogous statement holds for symmetry w.r.t. a particular hyperplane. For d = 2, suppose the existence of the above supporting circles (for S^2 of radius less than $\pi/2$), and, for S^2 , smoothness of K and L. If we suppose axial symmetry of all the above convex hulls, then our bodies are (incongruent) circles (for S^2 of radii less than $\pi/2$).

* Facultad de Ingeniería, Universidad Autónoma de Querétaro, Centro Universitario,

Cerro de las Campanas s/n C.P. 76010, Santiago de Querétaro, Qro. México, ME-XICO

ORCID ID: https://orcid.org/0000-0002-6601-0004

** Alfréd Rényi Mathematical Institute, Hungarian Research Network (HUN-REN),

¹⁹⁹¹ Mathematics Subject Classification. Mathematics Subject Classification 2020. 52A55.

Key words and phrases. spherical, Euclidean and hyperbolic planes and spaces, characterizations of ball/parasphere/hypersphere/half-space, convex bodies, proper closed convex sets with interior points, directly congruent copies, intersections, convex hulls of unions, central symmetry, symmetry w.r.t. a hyperplane, axial symmetry.

^{*}Research (partially) supported by CONACYT, SNI 38848

^{**}Research (partially) supported by Hungarian National Foundation for Scientific Research, grant nos. T046846, T043520, K68398, K81146, Research supported by ERC Advanced Grant "GeoScape", No. 882971.

H-1364 Budapest, Pf. 127, HUNGARY http://www.renyi.hu/~makai ORCID ID: https://orcid.org/0000-0002-1423-8613

E-mail address: * jeronimo@cimat.mx, jesusjero@hotmail.com ** makai.endre@renyi.hu

5. New results, Theorems 5–8

 S^d , \mathbb{R}^d , H^d , with $d \ge 2$, are the *d*-dimensional spherical, Euclidean and hyperbolic spaces, resp. Convexity of a set $K \subset H^d$ is defined as for $K \subset \mathbb{R}^d$. Convexity of $K \subset S^d$, with int $K \ne \emptyset$, is meant as follows: for any two non-antipodal points of K the shorter great circle arc connecting them belongs to K. Then for two antipodal points of K some great circle arc connecting them belongs to K. A convex body $Y \subset X$ is a compact convex set with nonempty interior. By a ball, or sphere in S^d , we mean one of radius at most $\pi/2$ (thus a ball is convex).

In the following theorem, the base line of a straight line is meant to be itself. We recall (from Part I, i.e., [5])

 $(*) \quad \begin{cases} X \text{ is } S^d, \ \mathbb{R}^d \text{ or } H^d, \text{ with } d \geq 2, \text{ and } K, L \subsetneqq X \text{ are closed convex sets} \\ \text{with interior points. Moreover, } \varphi, \psi : X \to X, \text{ sometimes with indices,} \\ \text{are orientation preserving congruences, with int} \left[(\varphi K) \cap (\psi L) \right] \neq \emptyset. \end{cases}$

Theorem 5. Assume (*) with d = 2 and let $X = H^2$. Then we have (3) \Longrightarrow (2) \Longrightarrow (1). If all connected components of the boundaries of both of K and L are hypercycles or straight lines, let their total number be finite. Then we have (3) \iff (2) \iff (1). Here:

- (1) For each φ, ψ we have that $(\varphi K) \cap (\psi L)$ admits some non-trivial congruence.
- (2) For each φ, ψ we have that $(\varphi K) \cap (\psi L)$ is axially symmetric.
- (3) We have either (A), or (B), or (C), or (D) or (E), where

(A): Any of K and L is a circle, a paracircle, a convex domain bounded by a hypercycle, or a half-plane. However, if one of K and L is a convex set bounded by a hypercycle or is a half-plane, then the other one is either a circle, or a congruent copy of the first one.

(B): One of K and L is a circle, and the other one is bounded either by two hypercycles, whose base lines coincide, or by a hypercycle, and its base line. (C): One of K and L is a circle, of radius r, say, and the other one is bounded by at least two hypercycles or straight lines (with all base lines different), whose mutual distances are at least 2r.

(D): One of K and L is a paracircle, and the other one is a parallel domain of some straight line, for some distance l > 0.

(E): K and L are congruent, and both are parallel domains of some straight lines, for some distance l > 0.

Remark 2. Observe that in Part I (i.e., [5]) Theorem 1 dealt with intersections of sufficiently small diameters and Theorems 3, 4, and this paper Theorem 5 dealt with all intersections. There is a question "in between" these: the case of all

compact intersections (observe that in Theorem 2 the intersections are compact). The case of central symmetry of all compact intersections, in [4] Theorem 3, in S^d , \mathbb{R}^d and H^d , under some regularity hypotheses (weaker than C_+^2), was clarified. There were found six cases. For central symmetry of all intersections [4] Theorem 2 gave that in S^d , \mathbb{R}^d and H^d , under the above mentioned regularity hypotheses, K and L were congruent balls, of radius at most $\pi/2$. This was just one of the above six cases.

Also here, for some non-trivial congruence, or axial symmetry the "in-between" question about all compact intersections, in S^2 , \mathbb{R}^2 and H^2 , can be posed, supposing the finiteness hypothesis in Theorem 5. Observe that for $X = S^2$ this is settled by Theorem 2. For $X = \mathbb{R}^2$ the answer is Theorem 1, (3). Namely, Theorem 3, (2) settles the case of all intersections (there are five cases), so these are examples also for all compact intersections. On the other hand, suppose that all compact intersections, with nonempty interior, are axially symmetric, or admit some non-trivial congruence. Then also all intersections, of sufficiently small diameter, are axially symmetric, or admit some non-trivial congruence. These are described in Theorem 1, (3) (there are six cases). There is just one case in Theorem 1, (3), not covered by Theorem 3, (2), namely the case of a half-plane and a parallel strip. However, these do not have a compact intersection with nonempty interior. Hence in this case the above properties "admitting some non-trivial congruence" and "axial symmetry" hold vacuously.

Problem 3. Can one decribe in H^2 the cases of admitting some non-trivial congruence, or axial symmetry of all compact intersections, at least when the numbers of connected components of bd K and of bd L are finite and when K and L are C_+^2 ? Here a larger number of cases can be expected. Namely, already our Theorem 5, about all intersections in H^2 , has five cases (and even Theorem 5, (A) comprises seven subcases). Moreover, [4], Theorem 3 about central symmetry of compact intersections in S^d , \mathbb{R}^d and H^d , when restricted to H^2 , has five cases – although three of these are contained in the cases in our Theorem 5. Clearly in all of the cases in both of these theorems, all compact intersections have some non-trivial symmetry.

Problem 4. Is the finiteness hypothesis in Theorem 5 necessary?

Remark 3. As will be seen from the proof of Theorem 5, namely in the proof of Lemma 5.10, rather than the finiteness hypothesis in Theorem 5, we may suppose only the following.

(1) One of K and L, e.g., the set K has a boundary component K_1 , such that some non-trivial arcs of S^1 , each with one endpoint at the infinite endpoints of K_1 , resp., and lying on the interior side of K_1 w.r.t. K, contain no infinite point of any other boundary component of K. Moreover, the other set L has two neighbourly boundary components L_1 and L_2 , with L_2 following L_1 in the positive sense, such that passing on bd L, meant in B^2 containing the model circle, in the positive sense from L_1 to L_2 , we pass on no other connected components of bd L, meant in H^2 . As follows from Theorem 5, namely in the proofs of Lemmas 5.7–5.10, rather than the respective finiteness hypotheses in Theorem 4 (in Part I, i.e., [4]) and Theorem 5, we may suppose the following.

(2) One of K and L has two boundary components with a common infinite point. (For Theorem 4 we have to consider that if all boundary components both of K and L are straight lines, then from Theorem 5, (3) only the case when both K and L are half-planes is possible. However their intersection is not always centrally symmetric.)

For the following Theorem 6 we will need the following weakening of the C^2 property.

 $(**) \quad \begin{cases} \text{Let for each } x \in \text{bd } K, \text{ and each } y \in \text{bd } L, \text{ there exist an } \varepsilon_1(x) > 0, \\ \text{and an } \varepsilon_1(y) > 0, \text{ such that } K \text{ and } L \text{ contain balls of radius } \varepsilon_1(x) \\ \text{and } \varepsilon_1(y), \text{ containing } x \text{ and } y \text{ in their boundaries, resp.} \end{cases}$

Moreover, we will need the following property, which together with (**) is a weak-ening of the C_+^2 property. (***)

Let for each $x \in \operatorname{bd} K$, and each $y \in \operatorname{bd} L$, there exist an $\varepsilon_2(x) > 0$ and an $\varepsilon_2(y) > 0$, such that the following holds. The set of points of K and L, lying at a distance at most $\varepsilon_2(x)$ and $\varepsilon_2(y)$ from x and from y, is contained in a ball B (for $X = S^d$, \mathbb{R}^d), or in a convex set B bounded by a hypersphere (for $X = H^d$). Moreover, bd B has sectional curvatures at least $\varepsilon_2(x)$ and $\varepsilon_2(y)$, and bd B contains x or y, resp.

Clearly (**) implies smoothness and (***) implies strict convexity, resp. Observe that both in (**) and (***) $\varepsilon_i(x) > 0$ and $\varepsilon_i(y) > 0$ can be decreased, and then (**) and (***) remain valid. ((**) and (***) are (A) and (B) in [4].)

An *inball of a convex body* $Y \subset X$ is a ball of maximal radius contained in Y.

Theorem 6. Assume (*). Let us assume C^2 for K and L (actually C^2 can be weakened to (**)). For $X = \mathbb{R}^d$ assume additionally that one of K and L has an extreme point. For $X = H^d$ assume C^2_+ for K and L (actually C^2_+ can be weakened to (**) and (***)). Then we have (1) \iff (2) \iff (3) \iff (4). Here

- (1) There exists some $\varepsilon = \varepsilon(K, L) > 0$, such that for each φ, ψ , for which diam $[(\varphi K) \cap (\psi L)] < \varepsilon$, we have that $(\varphi K) \cap (\psi L)$ is centrally symmetric.
- (2) For each φ, ψ and each x ∈ bd K and y ∈ bd L, there exists some ε = ε(K, L, x, y) > 0, such that the following holds. Suppose that φ, ψ and x, y satisfy the following hypotheses (A), (B) and (C).
 (A) diam [(φK) ∩ (ψL)] < ε.
 (B) (φK) ∩ (ψL) has a unique inball B₀.
 (C) [bd [(φK) ∩ (ψL)]] ∩ (bd B₀) = {φx, ψy}, where φx ∈ int (ψL) and ψy ∈ int (φK) are antipodal on bd B₀. Then we have that (φK) ∩ (ψL) is centrally symmetric.
- (3) The same as (2), but "centrally symmetric" in the conclusion replaced by "symmetric w.r.t. the orthogonal bisector hyperplane of $[\varphi x, \psi y]$ ".

(4) The connected components of the boundaries of both K and L are congruent spheres (for $X = S^d$ of radius at most $\pi/2$), or paraspheres, or congruent hyperspheres (for \mathbb{R}^d and H^d degeneration to hyperplanes being not admitted). For the case of congruent spheres or paraspheres we have that either K and L are congruent balls (for $X = S^d$ of radius at most $\pi/2$), or they are paraballs.

In Theorem 6, $(1) \Leftrightarrow (4)$ is the statement of [4], Theorem 1.

For Theorems 7 and 8, about (closed) convex hulls of unions (which is the question "dual" to that about the intersection), we use the hypothesis (****)

X is S^d , \mathbb{R}^d or H^d , with $d \ge 2$. Further, $K, L \subsetneq X$ are closed convex sets with interior points. For S^2 , both K and L lie in some open hemispheres. Moreover, $\varphi, \psi : X \to X$ are orientation preserving congruences. For $X = S^d$ the set $(\varphi K) \cup (\psi L)$ lies in an open hemisphere S of S^d .

The convex hull of a set $Y \subset H^d$ is defined as for \mathbb{R}^d . For $Y \subset S^d$, since we will use only sets $Y (= (\varphi K) \cup (\psi L))$ lying in an open half- S^d , namely S (cf. (****)), the convex hull of such a Y is the minimal convex set in S, containing Y. We may suppose that S is the open southern hemisphere, thus the collinear model exists for it. Thus, for any X, the image in the collinear model of the convex hull of such a $Y \subset X$ is the convex hull of the image in the collinear model of Y.

We say that a convex body Y in S^d , \mathbb{R}^d or H^d , has at its boundary point y a supporting sphere if the following holds. There exists a ball containing Y, for S^d of radius less than $\pi/2$, such that y belongs to the boundary of this ball. This boundary is called the *supporting sphere*. Observe that this implies compactness and strict convexity of Y, and for $X = S^d$ that Y is contained in an open hemisphere of S^d . Therefore, if K, L satisfy (****), then conv $[(\varphi K) \cup (\psi L)]$ is compact, hence closed. Therefore we will not need to take the closure of this set. Also, if Y is a convex body, satisfying this italicized hypothesis for all $y \in \operatorname{bd} Y$, then for S^d , \mathbb{R}^d and H^d , any existing sectional curvature of Y is greater than 0, 0 and 1, resp. For H^d this is a serious geometric restriction.

Let $D \subset X$ be a closed segment (to be specified later, in Theorems 7 and 8), with $D \subset S$ for S^d , and let $\varepsilon > 0$, with ε sufficiently small for S^d , cf. (*****) just below. Then we write $D(\varepsilon)$ for the (closed) convex hull of the union of two closed (d-1)-balls, of radius ε , lying in hyperplanes orthogonal to D at the endpoints of D, and of centres the two endpoints of D.

$$(****) \qquad \begin{cases} \text{For } S^d \text{ we choose } \varepsilon > 0 \text{ so that these two closed} \\ (d-1)\text{-balls lie in } S \text{ (or, equivalently, } D(\varepsilon) \subset S). \end{cases}$$

Observe that $D(\varepsilon)$ is centrally symmetric w.r.t. the midpoint of D, and also is symmetric w.r.t. the orthogonal bisector hyperplane of D.

Suppose that the midpoint of the segment D is the south pole o. Then the ball B(D), having D as a diametral chord, has as image in the collinear model a ball

about the centre 0 of the model. Further, the two bases of $D(\varepsilon)$ have as images in the collinear model two congruent closed (d-1)-balls. These lie in opposite tangent hyperplanes of the image of B(D) in the collinear model. Moreover, they have centres the images of the two endpoints of D in the collinear model. Moreover, the image of $D(\varepsilon)$ in the collinear model is a right cylinder over a (d-1)-ball, with axis of rotation (an axis of symmetry for d = 2) spanned by the image of D.

Theorem 7. Assume (****). Let both K and L have supporting spheres at any of their boundary points, for S^d of radii less than $\pi/2$. Then $(1) \iff (2) \iff (3) \iff (4) \iff (5) \iff (6) \iff (7)$. Here:

- (1) For each φ, ψ we have that conv $[(\varphi K) \cup (\psi L)]$ is centrally symmetric.
- (2) For each φ, ψ , provided that conv $[(\varphi K) \cup (\psi L)]$ has a unique diametral segment D, we have that conv $[(\varphi K) \cup (\psi L)]$ is symmetric w.r.t. the orthogonal bisector hyperplane of D.
- (3) Suppose that φ, ψ and $x \in \operatorname{bd} K$ and $y \in \operatorname{bd} L$ satisfy the following hypotheses (A) and (B).

(A) For S^d we have that diam $[\operatorname{conv} [(\varphi K) \cup (\psi L)]]$ is sufficiently close to π . For \mathbb{R}^d and H^d we have that this diameter is sufficiently large.

(B) conv $[(\varphi K) \cup (\psi L)]$ has $D := [\varphi x, \psi y]$ as its unique diametral segment. Then conv $[(\varphi K) \cup (\psi L)]$ is centrally symmetric.

- (4) The same as (3), but "centrally symmetric" changed to "symmetric w.r.t. the orthogonal bisector hyperplane of D".
- (5) Suppose that φ, ψ and x ∈ bd K and y ∈ bd L satisfy hypotheses (A) and (B) from (3). Then there exists an ε(K, L, x, y, φ, ψ) > 0 (for S^d an ε(K, L, x, y, φ, ψ, S) > 0, satisfying (*****) in place of ε there), such that the following holds. For 0 < ε < ε(K, L, x, y, φ, ψ) (for S^d for 0 < ε < ε(K, L, x, y, φ, ψ, S)) we have the following. For D := [φx, ψy] and for D(ε) defined before this theorem, [conv [(φK) ∪ (ψL)]] ∩ D(ε) is centrally symmetric.
- (6) The same as (5), but "centrally symmetric" changed to "symmetric w.r.t. the orthogonal bisector hyperplane of D".
- (7) K and L are congruent balls (for the case of S^d of radius less than $\pi/2$).

The equivalence of (1) and (7) in Theorem 7 slightly improves Theorem 4 of [4]. There we needed the additional hypothesis of smoothness of K and L for the same conclusion (but could add hypothesis (A) from Theorem 7, (3), to (1)).

Theorem 8. Assume (****) with d = 2. Also assume that both K and L have supporting circles at any of their boundary points, for S^2 of radii less than $\pi/2$. For S^2 also assume that K and L are C^1 .

Then $(1) \iff (2) \iff (3) \iff (4) \iff (5) \iff (6) \iff (7)$. Here:

- (1) For each φ, ψ we have that conv $[(\varphi K) \cup (\psi L)]$ is axially symmetric.
- (2) The same as (1), but "is axially symmetric" changed to "admits some nontrivial congruence".
- (3) Suppose that φ, ψ and $x \in \operatorname{bd} K$ and $y \in \operatorname{bd} L$ satisfy hypotheses (A) and (B) from (3) of Theorem 7. Then $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ is axially symmetric.

- (4) The same as (3), but "is axially symmetric" changed to "admits some nontrivial congruence".
- (5) Suppose that φ, ψ and x ∈ bd K and y ∈ bd L satisfy hypotheses (A) and
 (B) from (3) of Theorem 7. Then there exists an ε(K, L, x, y, φ, ψ) > 0 (for S² an ε(K, L, x, y, φ, ψ, S) > 0, satisfying (*****) in place of ε there), such that the following holds. For 0 < ε < ε(K, L, x, y, φ, ψ) (for S² for 0 < ε < ε(K, L, x, y, φ, ψ, S)) we have the following. For D := [φx, ψy] and for D(ε) defined before Theorem 7, [conv [(φK) ∪ (ψL)]] ∩ D(ε) is axially symmetric.
- (6) The same as (5), but "is axially symmetric" changed to "admits some nontrivial congruence".
- (7) K and L are (in general incongruent) circles (for the case of S^2 of radii less than $\pi/2$).

Remark 4. Probably smoothness for S^2 is not necessary for Theorem 8, (1) \iff (7). However, without the supporting circle hypothesis, Theorem 8, (1) \iff (7) does not remain true. Namely, besides the pairs K, L in Theorem 8, (7), also for the pairs K, L in Part I (i.e., [5]), Theorem 3, (2) (for \mathbb{R}^2), and in Theorem 5, (3), (A), (B), (D), (E) (for H^2), the set cl conv $[(\varphi K) \cup (\psi L)]$ admits some non-trivial congruences (actually, except for two parallel strips in \mathbb{R}^2 , axial symmetries). Still there are some evident examples, with closed convex hulls rather than convex hulls, satisfying (****), except for S^2 its last statement: for S^2 one of K and L is a halfsphere, and for \mathbb{R}^2 one of K and L is either a parallel strip, or a halfplane.

Problem 5. Are these the only pairs K, L in S^2 , \mathbb{R}^2 and H^2 , with this property? Observe that (5) and (6) of Theorem 7 are "localizations" of (3) and (4) of Theorem 7, resp., as well as (5) and (6) of Theorem 8 are "localizations" of (3) and (4) of Theorem 8, resp. Thus these are analogues of the "localizations" in Part I (i.e., [5]), Theorem 2, where (2), (6) and (7) are "localizations" of (1), (4) and (5), resp.

In the proofs of our Theorems we will use some ideas of [3].

6. Preliminaries

For 1. Introduction, and 3. Preliminaries we refer to Part I (i.e., [5]). Most of our notations are standard. However, we have to repete some notations from them.

X is S^d , \mathbb{R}^d or H^d , with $d \ge 2$. For $x, y \in X$ we write d(x, y) for their distance. In S^d , when saying *ball*, or *sphere*, we always mean one with radius at most $\pi/2$. A *paraball* is the closed convex set in H^d , bounded by a parasphere. For d = 2we say *paracircle*. For d = 2, and x_1, x_2 on the boundary of a closed convex set $K \subset X$ with interior points, "close" to each other, we write $\widehat{x_1x_2}$ for the (shorter, or unique) arc of bd K, which set K will be clear from the context. We will use the collinear and the conformal models of H^d , in the interior of the unit ball $B^d \subset \mathbb{R}^d$. We will speak about collinear and conformal models of S^d in \mathbb{R}^d . We mean by this the ones obtained by central projection (from the centre), or by stereographic projection (from the north pole), to the tangent hyperplane of S^d , at the south pole, in \mathbb{R}^{d+1} . These exist of course only on the open southern half-sphere, or on S^d minus the north pole, resp. Their images are \mathbb{R}^d . We call the *centre of the model* the south pole of S^d . The collinear and conformal models of \mathbb{R}^d are meant as itself, with *centre* the origin. Sometimes we will consider the (collinear or conformal) model circle of H^2 as the unit circle of the complex plane \mathbb{C} . Thus we will speak about its points 1, *i*, etc. *Smooth* will mean differentiable, which for closed convex sets with interior points is equivalent to C^1 . For a topological space Y we say that some property of a point $y \in Y$ holds *generically*, if it holds outside a nowhere dense closed subset.

A convex surface in X is the boundary of a proper closed convex subset of X with interior points. We will call a convex surface in \mathbb{R}^d at some of its points twice differentiable if the following holds. It is the graph, in a suitable rectangular coordinate system, of a function having a Taylor series expansion of second degree at this point, with an error term $o(\|\cdot\|^2\|)$. By [6], pp. 31-32 (in both editions), convex surfaces in \mathbb{R}^d are almost everywhere twice differentiable. This extends to S^d and H^d by using their collinear models.

Part I (i.e., [5]) contains Abstract, 1. Introduction, 2. New results: Theorems 1–4, 3. Preliminaries and 4. Proofs of Theorems 1–4. Part II contains Abstract (about Part II), 5. New results: Theorems 5–8, 6. Preliminaries and 7. Proofs of Theorems 5–8. The References in Part II contain only items referred to in Part II; more detailed references, in particular about hyperbolic geometry, cf. in Part I.

7. Proofs of Theorems 5–8

Before passing to the proof of Theorem 5, we introduce some terminology.

Suppose that Y is a connected real analytic manifold, and $f, g : Y \to \mathbb{R}$ are analytic functions. Then either f and g coincide, or else they cannot coincide on any subset of Y, whose closure contains some nonempty open subset of Y. This is the *principle of analytic continuation*. Otherwise said, in the second case, generically, for $y \in Y$ (i.e., except on a nowhere dense closed subset of Y), we have $f(y) \neq g(y)$.

Recall that a finite union of nowhere dense closed subsets is itself nowhere dense and closed. In **2** of the proof of Lemma 5.4, in **3–5** of the proof of Lemma 5.7, and in **3–6** of Lemma 5.8 we will have the following situation. On a connected real analytic manifold Y (in fact, on H^2 , or on the orientation preserving congruences of H^2) there are finitely many pairs of analytic functions, $f_1, g_1; \ldots; f_n, g_n: Y \to \mathbb{R}$, say, where f_i and g_i are different for all $1 \le i \le n$. Then generically, for $y \in Y$, we have that $f_i(y) \ne g_i(y)$ for all $1 \le i \le n$.

Before the proof of Theorem 5 we show a formula of trigonometrical type in H^2 . It is in a sense an analogue of the law of cosines for an angle of a triangle in H^2 . Namely, the law of cosines allows us, for two circles, or radii r, R, and distance of centres c, to determine the half central angle of the arc of the circle of radius r, lying in the circle of radius R. We will need an analogous formula, for a circle of radius r, and a hypercycle, with distance l from its base line, for the half central angle of the arc of the circle of radius r, lying in one of the sets bounded by the hypercycle. Here the distance c of the centre of the circle and the base line of the hypercycle is given. We consider c and l as signed distances. We admit degeneration to a straight line, i.e., we admit l = 0. This formula is surely known, but we could not locate a proof for it. Therefore we give its simple proof.

Lemma 5.1. Let $H \subset H^2$ be a hypercycle, with base line B, and with signed distance l from B. Let $H^- \subset H^2$ be the set of points at a signed distance at most lfrom B. Let $C \subset H^2$ be a circular line, with centre o, radius r, and with o having a signed distance c from B. Then $C \cap H \neq \emptyset$ if and only if $|c - l| \leq r$. If this inequality is satisfied, then for the half central angle ω of the arc $C \cap H^-$ of C we have

(5.1.1) $\sinh l = \cosh r \cdot \sinh c - \sinh r \cdot \cosh c \cdot \cos \omega.$

Proof. We use the conformal model. This shows that C and H have either two common points, or they are tangent to each other, or they are disjoint. (Their images are a circle, and a circular arc or a segment which cuts the model into two connected parts.) We suppose that B is a horizontal line containing the centre of the model, and the signed distance from B is positive in the open upper half-circle of the model.

Clearly we have $C \cap H \neq \emptyset$ if and only if $|c - l| \leq r$. From now on we suppose this inequality.

Let x be one of the points of $C \cap H$, and let y and z be the orthogonal projections of o and x to B. Thus d(x, z) = l. We let d := d(x, y). So we have to determine the angle $\omega = \angle xoy$ (for $o \in B$ the angle ω is defined by the evident limit procedure).

By the law of cosines for sides we have $\cosh d = \cosh r \cdot \cosh c - \sinh r \cdot \sinh c \cdot \cos \omega$. Now we calculate the angle $\alpha := \angle oyx$ (for $o \in B$ defined as a limit). Preliminarily let us suppose $l \neq 0$, that implies $d \neq 0$. Then by the law of sines we have $\sin^2 \alpha = \sin^2 \omega \cdot \sinh^2 r / \sinh^2 d$. Last, from the right triangle yxz we have $\sinh^2 d(x, z) = \sin^2(\pi/2 - \alpha) \cdot \sinh^2 d$. So, fixing r, c, and supposing $\cos \omega$ as given, we determine, by substitutions, successively, first $\cosh d$, then $\sin^2 \alpha$, then $\sinh^2 d(x, z)$. This last expression should equal $\sinh^2 l$. Solving this last equation for $\cos \omega$ (which is a quadratic equation), we obtain, by rearranging,

(5.1.2)
$$\begin{cases} \pm \sinh l = \cosh r \cdot \sinh c - \sinh r \cdot \cosh c \cdot \cos \omega \\ = \cosh r \cdot \cosh c \cdot (\tanh c - \tanh r \cdot \cos \omega). \end{cases}$$

The same equation holds also for l = 0. Namely then we have $\tanh c = \tanh r \cdot \cos \omega$, which is a trigonometric formula for a right triangle in H^2 .

We will show that (5.1.2) is satisfied with the plus sign, as stated in (5.1.1) in the lemma. Recall that $|c - l| \leq r$, or, equivalently, $C \cap H \neq \emptyset$, was assumed.

First suppose $c \ge 0$. We distinguish two cases.

- (1) We have $0 \le r \le c$.
- (2) We have $r > c \ge 0$.

In case (1) the entire circle C lies (not strictly) above B. By $C \cap H \neq \emptyset$ then H has some point at a non-negative signed distance from B. Thus $l \ge 0$. On the other hand, the expression in the middle of (5.1.2) lies in $[\sinh(c-r), \sinh(c+r)] \subset [0, \infty)$.

Hence both sides of (5.1.1) are non-negative, which means that in (5.1.2) the \pm sign is a + sign.

In case (2) $C \cap B$ consists of two points, and the line B cuts C to two nondegenerate arcs, both with these two points as endpoints. One of these arcs lies above B, the other one below B. For these two points we have l = 0, and from (5.1.2), the angle ω_0 corresponding to these points satisfies $\tanh c = \tanh r \cdot \cos \omega_0$. Moreover, the points of C corresponding to an angle $\omega \in [0, \omega_0)$, or $\omega \in (\omega_0, \pi/2]$, lie strictly below or above B, resp. Then l < 0, or l > 0, resp. On the other hand, in the last expression in (5.1.2), the factor $\tanh c - \tanh r \cdot \cos \omega$ vanishes for $\omega = \omega_0$, and for $\omega \in [0, \omega_0)$ or $\omega \in (\omega_0, \pi/2]$ it is negative or positive, resp. Hence once more, the left and right hand sides of (5.1.1) have the same signs, which means that in (5.1.2) the \pm sign is a + sign.

Last we extend the validity of (5.1.1) to c < 0. Let us apply (5.1.1) to $-c, -l, \pi - \omega$ rather than c, l, ω . Then the validity of (5.1.1) for these values implies its validity for c, l, ω .

Later, in the proof of Theorem 5, we will consider the case when $l \ge 0$; then, of course, c will vary in \mathbb{R} .

Remark 5. There is a well-known formalism between spherical and hyperbolical trigonometric relations. Namely, the hyperbolical ones can be obtained from the spherical ones by "choosing the radius of the ball as i". Applying this in the converse direction, (5.1.1) goes over to the following. Distance lines on S^2 with distance l are circles of radius $\pi/2 - l$, and similarly distance c from a line turns to distance $\pi/2 - c$ from the centre of this circle, while ω turns to $\pi - \omega$. Thus we obtain $\cos l = \cos r \cdot \cos c + \sin r \cdot \sin c \cdot \cos \omega$, i.e., the cosine law for sides.

In the proof of Theorem 5 we will allow that hypercycles degenerate to straight lines, i.e., hypercycle will mean either a proper hypercycle, or a straight line; i.e., the curvature is allowed to be 0 as well. The base line of a straight line is meant as itself.

Proof of Theorem 5. 1. The implication $(2) \Longrightarrow (1)$ is trivial.

Lemma 5.2. Assume (*) with d = 2 and $X = H^2$. Then the implication (3) \Longrightarrow (2) of Theorem 5 holds.

Proof. 1. We begin with case (A). A circle is axially symmetric w.r.t. any straight line passing through its centre. A paracircle is axially symmetric w.r.t. any straight line passing through its centre (its point at infinity). A half-plane, or a convex domain bounded by a hypercycle is axially symmetric w.r.t. any straight line, that intersects its boundary, or its base line orthogonally, resp. These imply, that if any of φK and ψL is either a circle or a paracircle, then their intersection is axially symmetric w.r.t. (any) straight line joining their centres. Suppose that one of φK and ψL is a circle, and the other one is a convex set bounded by a hypercycle, or is a half-plane. Then the straight line passing through the centre of the circle, and orthogonal to the base line of the hypercycle, or to the boundary of the half-plane, is an axis of symmetry of the intersection.

Last, let φK and ψL be congruent convex sets, each bounded by one hypercycle,

or let them be two half-planes. Consider the base lines of these hypercycles, or the boundaries of these half-planes, resp. There are four cases. These base lines

- (a) may coincide; or
- (b) may transversally intersect; or
- (c) may be parallel but distinct; or
- (d) may be ultraparallel.

In case (a) any straight line orthogonal to the common base line is an axis of symmetry. In case (b), $\varphi K \neq \psi L$, and bd (φK) and bd (ψL) intersect transversally at some point p (for this use the conformal model). Then (φK) \cap (ψL) has an inner angle at p, of measure less than π , and the bisector of this angle is an axis of symmetry of (φK) \cap (ψL). In case (c), if one of φK and ψL contains the other, the intersection is evidently axially symmetric. Otherwise, the symmetry axis of the base lines is an axis of symmetry of the intersection. In case (d), we consider the pair of points on the base lines, realizing the distance of these lines. The straight line connecting these points is orthogonal to both lines, and is an axis of symmetry of (φK) \cap (ψL).

2. We continue with case (B). Suppose that K is a circle, and L is bounded by two hypercycles, whose base lines coincide (one of them possibly degenerating to a straight line). Then the straight line passing through the centre of φK , and orthogonal to the above base line, is an axis of symmetry of $(\varphi K) \cap (\psi L)$.

3. We continue with case (C). Suppose that K is a circle of radius r, and the boundary hypercycle or straight line components of L have pairwise distances at least 2r. Then int (φK) can intersect at most one boundary component of ψL .

If int (φK) does not intersect any boundary component of ψL (and, by hypothesis, int $[(\varphi K) \cap (\psi L)] \neq \emptyset$), then $(\varphi K) \cap (\psi L) = \varphi K$ is a circle. Hence it is axially symmetric.

Suppose that int (φK) intersects exactly one boundary component ψL_1 of ψL . Then $(\varphi K) \cap (\psi L)$ is the same as the intersection of φK and of the closed convex set, bounded by ψL_1 , and containing ψL . This has an axis of symmetry, cf. case (A).

4. We continue with case (D). Let K be a paracircle, and L a parallel domain of some straight line, with some distance l > 0. Consider the common base line of the two hypercycles, bounding ψL . If the infinite point of the paracircle φK lies on this common base line, then this straight line is an axis of symmetry of $(\varphi K) \cap (\psi L)$. Suppose that the infinite point of φK does not lie on this common base line. Then there is a unique straight line that passes through the infinite point of φK , and is orthogonal to the common base line. Then this unique straight line is an axis of symmetry.

5. Last we turn to case (E). Consider the common base lines of the two hypercycles bounding φK , and of the two hypercycles bounding ψL . These two straight lines can coincide, or can transversally intersect, or can be parallel but distinct, or can be ultraparallel. In any case there is an axial symmetry interchanging these two straight lines. This axial symmetry interchanges the parallel domains of these straight lines, with distance l, as well. Hence it is an axial symmetry of the intersection of these parallel domains, i.e., of $(\varphi K) \cap (\psi L)$. Proof of Theorem 5, continuation. 2. Last we turn to the proof of $(1) \Longrightarrow (3)$. By Part I (i.e., [5]), Theorem 1, $(2) \Longrightarrow (3)$ we know that each boundary component of both K and L is either a cycle, or a straight line. Thus, for both K and L, we have the following possibilities: it is a circle, or a paracircle, or its boundary components are hypercycles and straight lines.

We make a case distinction. Either both $\operatorname{bd} K$ and $\operatorname{bd} L$ are connected, or one of them has several connected components.

Lemma 5.3. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. If both bd K and bd L are connected, then we have (A) of (3) of Theorem 5.

Proof. We have to investigate the cases when

(a) φK and ψL are one paracircle and one convex set bounded by a hypercycle or a straight line, or

(b) φK and ψL are two incongruent convex sets, both bounded by a hypercycle or a straight line,

and in both cases we have to find a contradiction.

Now it will be convenient to use the conformal model. In case (a), let the centre of the paracircle be one endpoint of the base line of the hypercycle, or one endpoint of the straight line. Then $\operatorname{bd}(\varphi K)$ and $\operatorname{bd}(\psi L)$ transversally intersect, at a single point, and this point p is the only non-smooth point of $(\varphi K) \cap (\psi L)$. In case (b), let the base lines of the two hypercycles, or the base line of the hypercycle and the straight line transversally intersect, resp. (two straight lines cannot occur). Then, also $\operatorname{bd}(\varphi K)$ and $\operatorname{bd}(\psi L)$ transversally intersect, at a single point, and this point p is the only non-smooth point of $(\varphi K) \cap (\psi L)$.

Both in case (a) and (b), any non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$ would have p as a fixed point. Moreover, the pair of the half-tangents at p would be preserved by this congruence. Thus, it would be an axial symmetry, w.r.t. the angle bisector of the inner angle of $(\varphi K) \cap (\psi L)$ at p. Thus, this axial symmetry should interchange the portions of the boundaries of φK and ψL , bounding $(\varphi K) \cap$ (ψL) . However, these portions of boundaries have different curvatures, which is a contradiction.

Proof of Theorem 5, continuation. 3. By Lemma 5.3 there remained the case when one of K and L has at least two boundary components. Observe that this rules out the cases when K, L are two circles, or two paracircles, or one circle and one paracircle. There remain the cases when one of K and L is bounded by several but finitely many hypercycles and straight lines, and the other one either is a circle, of some radius r, or is a paracircle, or also is bounded by finitely many (possibly one) hypercycles and straight lines. We will investigate these three cases separately, in Lemma 5.4, Lemma 5.5 and Lemmas 5.7–5.10, resp.

Suppose that one of K and L is bounded by hypercycles and straight lines. Then the boundary components φK_i of φK , or the boundary components ψL_i of ψL have a natural cyclic order, in the positive sense, on bd (φK), or bd (ψL), resp. We associate to φK , or to ψL a graph. Its vertices are the infinite points of the (φK_i)'s, or (ψL_i)'s, resp., and between two such points there is an edge, if they are the two infinite points of some φK_i , or ψL_i , resp. We say that this edge is φK_i , or ψL_i , resp. This graph either is a union of vertex-disjoint paths, or is a cycle. Here we admit a cycle of length 2, shortly a 2-cycle. Then the graph consists of two vertices, and two edges between these two vertices, which are two (φK_i) 's $((\psi L_i)$'s) with both infinite points in common.

If we have two edges in these graphs with a common vertex, and they are e.g., φK_1 and φK_2 , then by this notation we will mean that φK_2 follows φK_1 on bd (φK) in the positive sense. If φK_1 and φK_2 form a 2-cycle, then the notation is fixed some way. (A similar convention holds for ψL).

 $(5.1) \begin{cases} \text{Later in the proof of Theorem 5 we will allow that hypercycles} \\ \text{degenerate to straight lines, i.e., hypercycle will mean either a} \\ proper hypercycle, or a straight line; i.e., the curvature is allowed \\ to be 0 as well. The base line of a straight line is meant as itself. \end{cases}$

Lemma 5.4. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let K be a circle of radius r and centre o, and let the connected boundary components of L be at least two hypercycles or straight lines. Then (B) or (C) of (3) of Theorem 5 holds.

Proof. 1. We have to show that either L is bounded by two hypercycles with common base line, or the at least two boundary components of L have pairwise distances at least 2r. Let us suppose the contrary. That is, we have both that dist $(L_1, L_2) < 2r$, for some different boundary components L_1 and L_2 of L, and that L is not bounded by two hypercycles with a common base line. By dist $(\psi L_1, \psi L_2) < 2r$, we have, for some choice of φ , for φ_0 , say, that int $(\varphi_0 K)$ intersects both ψL_1 and ψL_2 . Then the same statement holds for all sufficiently small perturbations φ of φ_0 , hence also of the *centre* φ_{00} of $\varphi_0 K$.

(5.4.1) $\begin{cases}
\text{We are going to show that for an arbitrarily small, generic perturbation} \\
\varphi o \text{ of } \varphi_0 o \text{ we have that } (\varphi K) \cap (\psi L) \text{ admits no non-trivial congruence.}
\end{cases}$

Observe that (5.4.1) contradicts the hypotheses of this lemma, thus our above indirect assumption will lead to a contradiction.

2. We have that $\operatorname{int}(\varphi_0 K)$ intersects ψL_i for $i \in I(0, i)$ (where $|I(0, i)| \geq 2$ by $\{1, 2\} \subset I(0, i)$), and $\varphi_0 K$ touches ψL_i from outside (w.r.t. ψL) for $i \in I(0, t)$, and $\varphi_0 K$ is disjoint to ψL_i for $i \in I(0, d)$. For a sufficiently small perturbation φ of φ_0 we have for the analogously defined I(i), I(t) and I(d) that $\{1, 2\} \subset I(0, i) \subset I(i)$ and $I(0, d) \subset I(d)$, hence also

(5.4.2)
$$I(i) \cup I(t) \subset I(0, i) \cup I(0, t).$$

(However, generically $I(t) = \emptyset$.) Moreover, each of I(0, i), I(0, t), I(i) and I(t) is finite. (Observe that any compact set in H^2 , thus, e.g., a closed circle concentric with $\varphi_0 K$, of radius just a bit larger than r, intersects only finitely many (ψL_i) 's.)

Let φ be a small generic perturbation of φ_0 , thus φ_0 be a small generic perturbation of $\varphi_0 o$. Then $(\varphi K) \cap (\psi L)$ is a convex body. It is bounded, alternately, by finitely many, but at least two non-trivial arcs of $bd(\varphi K)$, and by same many non-trivial arcs of some (ψL_i) 's, for different (ψL_i) 's. (The case when some φK touches ψL_i at some point, from inside (w.r.t. ψL), gives that ψL_i does not cut off anything from φK , hence can be left out of consideration. The case when some φK touches ψL_i at some point, from outside (w.r.t. ψL), gives that $|(\varphi K) \cap (\psi L)| = 1$, contradicting $(int(\varphi K)) \cap (\psi L_1) \neq \emptyset$. Thus, by the conformal model, we may suppose that the neighbouring boundary arcs of $(\varphi K) \cap (\psi L)$ intersect transversally.) The curvatures of these arcs are greater than 1, or smaller than 1, resp., so each congruence admitted by $(\varphi K) \cap (\psi L)$ preserves both above types of arcs, separately. Thus the union of those boundary arcs of $(\varphi K) \cap (\psi L)$, which are arcs of bd (φK) , i.e., the set $[bd(\varphi K)] \cap (\psi L)$ is preserved by each congruence admitted by $(\varphi K) \cap (\psi L)$. Therefore also conv $[[bd(\varphi K)] \cap (\psi L)]$ is preserved by each congruence admitted by $(\varphi K) \cap (\psi L)$. Then conv [[bd $(\varphi K)] \cap (\psi L)$] is obtained from the circle φK , by cutting off disjoint circular segments, relatively open in φK , by finitely many, but at least two disjoint non-trivial chords. These have endpoints the points of intersection of the single (ψL_i) 's with $\operatorname{bd}(\varphi K)$, for all (ψL_i) 's with $i \in I(i)$.

(5.4.3) $\begin{cases} \text{We will attain that all these chords have different lengths} - \\ \text{moreover, this can be attained by a small, generic motion} \\ \text{of the centre } \varphi o \text{ of } \varphi K, \text{ from its original position } \varphi_0 o. \end{cases}$

By the law of sines, two such chords have equal length if and only if the corresponding half central angles, belonging to $(0, \pi) \subset [0, \pi]$ have equal sines. That is, the absolute values of the cosines of these angles coincide. We use formula (5.1.1) from Lemma 5.1. However, in Lemma 5.1 we dealt with the half central angle ω of the arc $C \cap H^-$ of C, while here we are interested in the central angle χ of the arc of bd (φK), "cut off" by ψL_i . Then $\chi = \pi - \omega$. We have several such equations like (5.1.1), corresponding to the (ψL_i)'s, for $i \in I(i)$, with respective fixed values rand l_i , and variable values c_i . We choose $l_i \geq 0$ for each i (this is the choice of the sign of the signed distance from the base line of ψL_i), and then c_i will vary in \mathbb{R} . For $l_i = 0$ we should have positive signed distance from ψL_i on the exterior side of the straight line ψL_i , w.r.t. ψL . We will use arbitrarily small generic perturbations φo of the centre $\varphi_0 o$ of our circle $\varphi_0 K$.

The half central angles of the arcs of $\operatorname{bd}(\varphi K)$, "cut off" by ψL_i , for $i \in I(i)$, are denoted by by χ_i . Then $\sum_{i \in I(i)} \chi_i < \pi$. Then for each distinct $i, j \in I(i)$ (recall from the paragraph following (5.4.1) that $|I(i)| \geq |I(0,i)| \geq 2$) we have $\chi_i + \chi_j \leq \sum_{i \in I(i)} \chi_i < \pi$. Hence for $|\cos \chi_i| = |\cos \chi_j|$ we have $\chi_i = \chi_j < \pi/2$. (Observe that $|I(0,i)| \geq 2$ was essential in this proof. For $I(i) = \emptyset$ and $I(t) = \{1\}$ we have the example when φK touches ψL_1 from outside w.r.t. ψL , hence it does not intersect any other ψL_i , and where the analogously defined χ_1 equals π , hence $\sum_{i \in I(i) \cup I(t)} \chi_i = \pi$.)

Hence, it suffices to exclude all pairwise equalities of finitely many expressions

for $\cos \chi = -\cos \omega$ – obtained from solving equations (5.1.1), for all $i \in I(0,i) \cup I(0,t)$ (which set contains I(i) by (5.4.2)) – namely those of the form on the right hand side of the following formula:

(5.4.4)
$$\cos \chi_i = (\sinh l_i - \cosh r \cdot \sinh c_i) / (\sinh r \cdot \cosh c_i) \,.$$

Observe that all these expressions are analytic in φo , since the c_i 's are analytic in φo , and r and the l_i 's are fixed. (In fact, in the collinear model the signed distance of a point (ξ, η) from a horizontal line of equation $\eta = \eta_0 \in (-1, 1)$ is the signed hyperbolic length of $[(\xi, \eta_0), (\xi, \eta)]$, which can be expressed by the cross-ratio formula, cf. [7].)

Moreover, none of these equalities, for distinct i, j in $I(0, i) \cup I(0, t)$, is an identity. Namely, let $\varphi' K$ touch a boundary component ψL_i of ψL from outside w.r.t. ψL . Then $(\varphi' K) \cap (\psi L)$ consists of one point. So, in particular, φK does not intersect any other boundary component ψL_j of ψL . Then in (5.4.4) $\cos \chi_i = -1$, however for $j \in [I(0,i) \cup I(0,t)] \setminus \{i\}$ the value for $\cos \chi_j$ from the right hand side of (5.4.4) is not in [-1,1]. (Else (5.1.1) and the second paragraph of the proof of Lemma 5.1 would give $l_j \in [c_j - r, c_j + r]$, so $|c_j - l_j| \leq r$, hence $[\operatorname{bd}(\varphi K)] \cap (\psi L_j) \neq \emptyset$, which is a contradiction.) Hence the *i*'th expression in the right hand side of (5.4.4), and any other *j*'th expression in the right hand side of (5.4.4), for $i, j \in I(0,i) \cup I(0,t)$, are not identical.

Therefore all our finitely many analytic equations, that the right hand sides of (5.4.4) are equal for distinct $i, j \in I(0,i) \cup I(0,t)$, are not identities. Hence each of them holds only for φo belonging to a nowhere dense closed subset. Therefore, supposing that φo belongs to some small closed neighbourhood A of $\varphi_{00}o$, except for φo belonging to a nowhere dense closed subset B of A, none of these equations hold. (Hence, by (5.4.2), also for any sufficiently small generic perturbation φ of φ_0 , and for distinct $i, j \in I(i)$, none of these equations holds.) That is, we have proved what was claimed in (5.4.3).

3. From now on we will suppose (5.4.3). There are two possibilities. Either (1) we can have at least three such chords, as in (5.4.3), or

(2) we always have exactly two such chords (then these correspond to the above L_1, L_2).

Any congruence admitted by $(\varphi K) \cap (\psi L)$ preserves φK , and conv $[[bd(\varphi K)] \cap (\psi L)]$, and also the above mentioned at least three, or exactly two disjoint chords, since their lengths are different. However, a non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$, preserving a single such chord (and that side of the chord, on which conv $[[bd(\varphi K)] \cap (\psi L)]$ lies), is an axial symmetry w.r.t. the orthogonal bisector straight line of this chord, which line contains φo . Since this is the unique non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$, any other one of these chords gives the same axial symmetry, hence also the same axis of symmetry of $(\varphi K) \cap (\psi L)$. However, there are no three disjoint circular segments, relatively open in φK , cut off by disjoint chords orthogonal to a single straight line, their common orthogonal bisector. Thus in case (1) we have a contradiction.

There remains case (2) when we can have only exactly two such chords. These

must correspond to the above considered hypercycles ψL_1 and ψL_2 .

Then φK does not even touch any other ψL_i from outside w.r.t. ψL , since then by a small motion of ψL we could attain that int (φK) intersects at least three (ψL_i) 's, which case was settled just above.

The above reasoning gives that, in this case, the orthogonal bisecting straight lines of the two chords coincide, furthermore, contain φo . However, the orthogonal bisecting straight lines of these two chords are orthogonal also to ψL_1 and ψL_2 , and hence also to the base lines of ψL_1 and ψL_2 . We have that these base lines are different, since their coincidence was excluded in the first paragraph of **1** of the proof of this lemma. Then they do not intersect, and thus are either parallel and distinct, or ultraparallel.

If they are parallel but distinct, then they admit no common orthogonal straight line, so we have a contradiction.

If they are ultraparallel, then they have exactly one common orthogonal straight line. This straight line depends on ψL only – let it be $l(\psi L)$. From above $\varphi o \in l(\psi L)$. Then in case (2) for $\varphi o \in A \setminus [B \cup (A \cap l(\psi L))]$ we have a contradiction.

Clearly $B \cup (A \cap l(\psi L))$ is a nowhere dense closed subset of A. Hence (5.4.1) holds for an arbitrarily small generic perturbation φo of $\varphi_0 o$, contrary to the hypotheses of this lemma. This shows that our indirect hypothesis in the first paragraph of **1** led to a contradiction, also in case (2).

Since both cases (1) and (2) led to a contradiction, therefore our indirect hypothesis was false. In other words, the statement of this lemma holds. \blacksquare

In the following lemmas, we will consider the (collinear or conformal) model circle of H^2 as embedded in the complex plane \mathbb{C} .

Lemma 5.5. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let K be a paracircle, and let the connected boundary components of L be at least two hypercycles or straight lines. Then (D) of (3) of Theorem 5 holds.

Proof. We consider the graph of ψL . It has at least two edges, by the hypothesis made after the proof of Lemma 5.3. Let ψL_1 be one of its edges. Let one endpoint of ψL_1 , say, 1, coincide with the centre of φK . Then $(\varphi K) \cap (\psi L)$ has the unique infinite point 1. Then each congruence admitted by $(\varphi K) \cap (\psi L)$ preserves this point 1, hence a congruence is either an axial symmetry w.r.t. an axis containing 1, or is a rotation about the infinite point 1. (Observe that by Part I (i.e., [5]), (2) it cannot be a nontrivial translation or a glide-reflection which is not a reflection, along a straight line having 1 as one of its infinite points.) However, the second case is impossible by Part I (i.e., [5]), (3).

In the first case let us consider the set $S = (\varphi K) \cap (\psi L_1) \subset \operatorname{bd} [(\varphi K) \cap (\psi L)]$, which is a closed semiinfinite arc of a hypercycle, with infinite point at 1, and finite endpoint on $\operatorname{bd}(\varphi K)$, and else lying in $\operatorname{int}(\varphi K)$. Then the image S' of Sby our axial symmetry is a hypercycle arc with the same above listed properties. Since $S' \subset \operatorname{bd} [(\varphi K) \cap (\psi L)] \subset [\operatorname{bd}(\varphi K)] \cup [\operatorname{bd}(\psi L)]$, therefore S', minus its finite endpoint, lies in $\operatorname{bd}(\psi L)$. Then taking closure also $S' \subset \operatorname{bd}(\psi L)$. Then $S' \subset \psi L_2$ for some boundary component ψL_2 of ψL , congruent to ψL_1 . Let ψL_2 follow ψL_1 at their common infinite point 1 in the positive sense. If also their other infinite points coincide, then ψL is a parallel domain of their coincident base lines, and then we have the statement of the lemma.

(5.5.1)

Therefore, for contradiction, let us suppose that the other endpoints of ψL_1 and ψL_2 are different, and we are going to show that this leads to a contradiction.

Since any three distinct points of the boundary circle of the model can be taken over by (the extension of) an orientation preserving congruence to any other three distinct points of the boundary circle of the model, of the same orientation, we may suppose that the other endpoints of ψL_1 and ψL_2 are, in the closure in \mathbb{C} of the model circle, very close to -1. Then all other boundary components of ψL are, in the closure in \mathbb{C} of the (conformal or collinear) model circle, close to -1.

We consider the conformal model. We fix the position of ψL_1 and ψL_2 , but we choose another φK so that it touches the boundary of the model circle very close (in \mathbb{C}) to 1, at a point in the open upper half-plane, and its interior intersects both ψL_1 and ψL_2 , and its image in the model is a circle of very small radius (in \mathbb{C}). Then φK is far (in \mathbb{C}) from all other boundary components of ψL . Hence, $(\varphi K) \cap (\psi L)$ is a convex arc-quadrangle, bounded by an arc of φK , an arc of ψL_2 , another arc of φK , and an arc of ψL_1 , in this cyclic order, in the positive sense. The endpoints of these arcs (the vertices of our arc-quadrangle) are the only non-smooth points of this arc-quadrangle, hence they are preserved by any congruence admitted by this arcquadrangle. This arc-quadrangle has two vertices on ψL_1 , and two vertices on ψL_2 , hence its vertices are the vertices of a strictly convex quadrangle (in the orientation inherited from bd $[(\varphi K) \cap (\psi L)]$). This arc-quadrangle is also compact, so by Part I (i.e., [5]), (2) and (3), it can admit only congruences which are rotations, or axial symmetries (according to whether they are orientation preserving, or orientation reversing).

Since a paracycle and a hypercycle (straight line) have different curvatures, neither of the diagonals can be an axis of symmetry, and there is no congruence which is a combinatorial 4-fold rotation. Therefore, our arc-quadrangle can admit only a rotation which is a combinatorial central symmetry, and by Part I (i.e., [5]), (4), this is also a geometrical central symmetry, with a unique centre. So, a non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$ is either a central symmetry, or an axial symmetry w.r.t. the common orthogonal bisector straight lines of the opposite arc-sides.

We begin with the case of central symmetry. Then the opposite arc-sides of $(\varphi K) \cap (\psi L)$ on the paracycle $\operatorname{bd}(\varphi K)$ are centrally symmetric images of each other. Then the paracycles containing these arc-sides have different infinite points (by the collinear model), hence are different. However, both are equal to $\operatorname{bd}(\varphi K)$, a contradiction.

We continue with the case of axial symmetry w.r.t. the common orthogonal bisector straight lines of the opposite arc-sides, lying on $\operatorname{bd}(\varphi K)$. However, a common orthogonal bisector straight line to the two opposite (thus disjoint) arcsides of $(\varphi K) \cap (\psi L)$, lying on $\operatorname{bd}(\varphi K)$, cannot exist. Namely, such orthogonal bisectors are different and parallel, so they cannot coincide.

We continue with the case of axial symmetry w.r.t. the common orthogonal

bisector straight lines of the opposite arc-sides, lying on ψL_1 and ψL_2 . However, a common orthogonal straight line to L_1 and L_2 is a common orthogonal straight line to their base lines as well. These base lines are parallel, and by (5.5.1) are different, hence admit no common orthogonal straight line, which is a contradiction. This shows that our indirect hypothesis (5.5.1) was false. This ends the proof.

Proof of Theorem 5, continuation. 4. Last, let both K and L be bounded by finitely many hypercycles and straight lines. Then, by the hypothesis made in the beginning of **3** of the proof of Theorem 5, either K, or L has at least two boundary components.

We will show that the graphs of φK and ψL must have only a few edges, and we will clarify the structure of these graphs, till we will obtain that we must have case (E) of (3) of our theorem.

We will make the following case distinction, for the graphs of φK and ψL . (5.2)

(1) Both graphs contain a pair of edges with at least one common end-point.

(2) One graph contains a pair of edges with at least one common endpoint,

but the other graph has only vertex-disjoint edges, and the number of these edges is at least two.

(3) One graph contains a pair of edges with at least one common endpoint, and the other graph has a single edge.(4) Both graphs have only vertex-disjoint edges, and at least one of the graphs contains at least two edges.

These cases are exhaustive, and mutually exclusive. They will be treated successively in the following Lemmas 5.7-5.10. Still we note that the finiteness hypothesis of Theorem 5 will not be needed in Lemmas 5.7–5.9, therefore it will not be included in their hypotheses.

5. In the proof of Lemmas 5.7–5.10 we will need "generic small perturbations of orientation preserving congruences of H^{2n} . For this to make sense, we have to give a topology on the set of orientation preserving congruences of H^2 . (Below φ_0, φ will be orientation preserving congruences of H^2 .) This will be the topology of uniform convergence on compacta in H^2 . This has as (open) neighbourhood base at some φ_0 the following:

 $\left\{\{\varphi \mid x \in K \Longrightarrow d(\varphi_0 x, \varphi x) < \varepsilon\} \mid K \subset H^2 \text{ is compact, and } \varepsilon > 0\right\}.$ (5.3)

(Since such a K is bounded in the metric of H^2 , rather than all such K, we may take only closed balls of some fixed centre and integer radii.) With this topology the maps $\varphi \mapsto \varphi x$, where $x \in H^2$, are continuous.

By [2], Theorem 2, the set of all congruences of H^2 (orientation preserving and reversing), with the topology defined by the same formula (5.3), is a topological group (i.e., the operations $(\varphi_1, \varphi_2) \mapsto \varphi_1 \varphi_2$ and $\varphi \mapsto \varphi^{-1}$ are continuous). This implies the same for its subgroup consisting of all orientation-preserving congruences of H^2 .

(5.4) Denote by Isom⁺(H^2) the set of orientation-preserving congruences of H^2 .

Now recall that in the conformal model, as a subspace of \mathbb{C} , the set $\text{Isom}^+(H^2)$ consists of the maps of the following form:

(5.5)
$$\begin{cases} z \mapsto f_{c,\alpha}(z) := c(z-\alpha)/(1-\overline{\alpha}z), \text{ where } |z| < 1, \\ \text{and } c \in \{z \in \mathbb{C} \mid |z| = 1\} = S^1, \text{ and } \alpha \in \text{int } B^2 \subset \mathbb{C} \end{cases}$$

(cf. [8]).

This is a real analytic manifold, with real analytic coordinates $c, a := \operatorname{Re} \alpha$ and $b := \operatorname{Im} \alpha$. The usual formulas for composition and inverse make this real analytic manifold a topological group, where the composition and the inverse are real analytic (a real analytic Lie group with these coordinates). Moreover, the action $(\varphi, z) \mapsto \varphi z$ is real analytic (in $c, a, b, \operatorname{Re} z, \operatorname{Im} z)$, and for given c and α it is complex analytic in z. Actually, rather than $z \in \operatorname{int} B^2$, we may take $z \in B^2$, and then the composition, inverse, and the action still are continuous (actually real analytic).

In the following Lemma 5.6 we will show that the above geometric definition of convergence on $\text{Isom}^+(H^2)$ is equivalent to a definition using the analytic functions from (5.5). This will be necessary in Lemma 5.8. This statement is surely known, but we could not locate a proof for it. Therefore we give its simple proof.

Lemma 5.6. A net $f_{c(\gamma),\alpha(\gamma)}(z)$ (defined in (5.5)) converges to $f_{c,\alpha}(z)$ in Isom⁺ (H²) in the topology of uniform convergence on compact in H², if and only if $(c(\gamma),\alpha(\gamma)) \to (c,\alpha)$ in $S^1 \times \operatorname{int} B^2$.

Proof. **1.** First we show that the map $S^1 \times \operatorname{int} B^2 \ni (c, \alpha) \mapsto f_{c,\alpha}(z) \in \operatorname{Isom}^+(H^2)$ (with $f_{c,\alpha}(z)$ from (5.5)), with $\operatorname{Isom}^+(H^2)$ endowed with the topology of uniform convergence on compact in H^2 , is continuous. We write $a := \operatorname{Re} \alpha$ and $b := \operatorname{Im} \alpha$.

Since $S^1 \times \operatorname{int} B^2$ is a metric space, therefore it suffices to consider only sequences $(c(n), \alpha(n)) \in S^1 \times \operatorname{int} B^2$, rather than general nets. Observe that $S^1 \times \operatorname{int} B^2 \ni (c(n), \alpha(n)), (c, \alpha)$ and $(c(n), \alpha(n)) \to (c, \alpha)$ imply $\alpha(n), \alpha \in rB^2$ for some $r \in (0, 1)$. Hence, letting $d_{\mathbb{C}}(\cdot, \cdot)$ being the distance in \mathbb{C} , we have $f_{c(n),\alpha(n)}(z) \to f_{c,\alpha}(z)$ in \mathbb{C} , i.e., $d_{\mathbb{C}}(f_{c(n),\alpha(n)}(z), f_{c,\alpha}(z)) \to 0$, pointwise for $z \in B^2$, but even uniformly on B^2 . Namely, $|(\partial/\partial c)f_{c,\alpha}(z)|, |(\partial/\partial a)f_{c,\alpha}(z)|, |(\partial/\partial b)f_{c,\alpha}(z)| \leq \operatorname{const}/(1-|\alpha|)^2 \leq \operatorname{const}/(1-r)^2$, uniformly for all $z \in B^2$. Therefore the same unifom convergence holds for any compact subset $C \subset \operatorname{int} B^2$. However, by Part I (i.e., [5]), (0.5), for distinct points of C, the distance $d(\cdot, \cdot)$ in H^2 , and the distance $d_{\mathbb{C}}(\cdot, \cdot)$ in the conformal model as a subspace of \mathbb{C} , have a quotient bounded from below and above. Hence (5.3) holds for $d(\cdot, \cdot)$, since from above it holds for the distance $d_{\mathbb{C}}(\cdot, \cdot)$. This shows that our map from $S^1 \times \operatorname{int} B^2$ to $\operatorname{Isom}^+(H^2)$, endowed with the topology of uniform convergence on compacta in H^2 , is continuous.

2. Second we show that also conversely, convergence of a net $f_{c(\gamma),\alpha(\gamma)}$ to $f_{c,\alpha}$, in the topology of uniform convergence on compact in H^2 , implies $(c(\gamma),\alpha(\gamma)) \rightarrow$

 (c, α) in $S^1 \times \operatorname{int} B^2$.

Let us choose the compact set $\{0\} \subset \operatorname{int} B^2$. Then

(5.6.1)
$$-c(\gamma)\alpha(\gamma) = f_{c(\gamma),\alpha(\gamma)}(0) \to f_{c,\alpha}(0) = -c\alpha.$$

Now observe that also $f_{c(\gamma),\alpha(\gamma)}^{-1} \to f_{c,\alpha}^{-1}$, by the topological group property of $\operatorname{Isom}^+(H^2)$. Let us choose the compact set $\{0, w\} \subset \operatorname{int} B^2$, where $w \neq 0$. Then

(5.6.2)
$$\alpha(\gamma) = f_{c(\gamma),\alpha(\gamma)}^{-1}(0) \to f_{c,\alpha}^{-1}(0) = \alpha,$$

(5.6.3)

$$(w+c(\gamma)\alpha(\gamma))/(c(\gamma)+\overline{\alpha(\gamma)}w)) = f_{c(\gamma),\alpha(\gamma)}^{-1}(w) \to f_{c,\alpha}^{-1}(w) = (w+c\alpha)/(c+\overline{\alpha}w).$$

Then, unless $\alpha = 0$, we have by (5.6.1) and (5.6.2), choosing γ sufficiently large, that

(5.6.4)
$$c(\gamma) = c(\gamma)\alpha(\gamma)/\alpha(\gamma) \to c\alpha/\alpha = c.$$

There remains the case $\alpha = 0$. In this case, taking in consideration |c| = 1 > |w| > 0, we can rewrite (5.6.3) as

(5.6.5)
$$(w + c(\gamma)\alpha(\gamma)) / (c(\gamma) + \overline{\alpha(\gamma)}w)) \to (w + c\alpha) / (c + \overline{\alpha}w) = w/c.$$

If $c(\gamma)$ had not converged to c, then by compactness of S^1 , for some subnet it would converge to some $c^* \in S^1 \setminus \{c\}$, and then the limit in (5.6.5) for this subnet would be $w/c^* \neq w/c$, a contradiction.

This ends the proof of the lemma. \blacksquare

Lemma 5.7. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let the connected boundary components of both K and L be hypercycles or straight lines. Moreover, suppose (5.2), case (1) from above. Then (E) of (3) of Theorem 5 holds.

Proof. 1. By (5.2), (1), each of the graphs of φK and ψL contains a path of length 2 or a 2-cycle. The corresponding boundary components of φK and ψL are denoted by $\varphi K_1, \varphi K_2$, and $\psi L_1, \psi L_2$, with φK_2 following φK_1 on bd (φK), and ψL_2 following ψL_1 on bd (ψL), according to the positive orientation. (If one of the graphs is a 2-cycle, then this does not work. Then we fix some notation, cf. below.)

We use the conformal model. Recall that any two or three distinct points on the boundary of the model can be taken by (the extension of) an orientation-preserving congruence to any other two or three distinct boundary points of the model, of the same orientation. Therefore we may suppose the following. The considered common vertex of φK_1 and φK_2 is $1 \in \mathbb{C}$. If their other vertices coincide, let them be -1. If their other vertices are different, let them be very close, in B^2 , to -1. Hence all other boundary components of φK (if any) are very close in B^2 , to -1, as well. Moreover, the considered common vertex of ψL_1 and ψL_2 is $i \in \mathbb{C}$. If their other vertices coincide, let them be -i. If their other vertices are different, let them be very close, in B^2 , to -i. Hence all other boundary components of ψL (if any) are very close in B^2 , to -i, as well.

Further, φK_2 follows φK_1 on bd (φK) at 1 in the positive sense, and similarly for $\psi L_2, \psi L_1$ at *i*. Then we have that the distance of 0 to $(\varphi K) \cap (\psi L)$ is small (possibly 0). Moreover, $(\varphi K) \cap (\psi L)$ is a convex arc-quadrangle, bounded by arcs of $\varphi K_1, \psi L_1, \varphi K_2, \psi L_2$, in this cyclic order, in the positive sense. In fact, all other boundary components, both of φK and of ψL , are, in B^2 , very close to the boundary of the model, hence cannot cut off parts of the above arc-quadrangle, which arcquadrangle is not close to the boundary of the model.

Thus $(\varphi K) \cap (\psi L)$ is a compact arc-quadrangle. Its vertices are its only nonsmooth points, hence they are preserved by any congruence admitted by this arcquadrangle. Analogously as in the proof of Lemma 5.5, by Part I (i.e., [5]), (2), (3) and (4) we have that the possible non-trivial congruences admitted by this arcquadrangle are the following:

(a) two fourfold rotations, and one central symmetry with a unique centre of symmetry, namely the midpoint of any of the two diagonals (which therefore coincide);(b) axial symmetries w.r.t. diagonals (which therefore bisect the angles at their endpoints);

(c) axial symmetries w.r.t. the common orthogonal bisector straight lines of two opposite arc-sides (which are also common orthogonal straight lines to the base-lines of the hypercycles containing these arc-sides), which simultaneosly interchange the remaining two opposite arc-sides (hence also the hypercycles containing them, and also their base-lines).

If we have a congruence which is a fourfold rotation, then we also have a congruence which is a central symmetry. Hence we need not exclude the case of a fourfold rotation, exclusion of a central symmetry will suffice.

Observe that a non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$ is a non-trivial congruence admitted by $(\varphi K_1) \cup (\varphi K_2) \cup (\psi L_1) \cup (\psi L_2)$ as well.

2. Next we investigate, for later use, two special cases, and we prove for them (E) of (3) of our theorem.

First suppose that both K and L are parallel domains of some straight lines. Thus the (positive) curvatures of K_1 and K_2 are equal, and also the (positive) curvatures of L_1 and L_2 are equal. We are going to show that also the curvatures of K_i and L_i coincide, i.e., that (E) of (3) of our theorem holds. Suppose, for contradiction, that, e.g., the (positive) curvature of K_i is less than the (positive) curvature of L_i .

Observe that K, or L, is symmetrical w.r.t. any straight line orthogonal to the common base line of the K_i 's, or of the L_i 's, resp. Hence any two congruent copies of K (and of L) are directly congruent if and only if they are indirectly congruent. (Thus K and L do not have any specific orientation.) Therefore we need not care about orientation preserving/reversing property of φ and ψ , it suffices to derive a contradiction for any φ and ψ . (Recall that in (*) we had φ, ψ as direct congruences.)

Now we choose a new pair φ, ψ (different from that in 1). This we make so that the infinite points of each of $\varphi K_1, \varphi K_2, \psi L_1, \psi L_2$ are ± 1 , and $\varphi K_1, \psi L_1$ are in the open lower half-plane, and $\varphi K_2, \psi L_2$ are in the open upper half-plane. Then we rotate ψL about the infinite point 1, counterclockwise, by a rotation of a small measure, obtaining a new $\psi' L$. Then, in the conformal model, the Euclidean tangents of the closures in B^2 of $\psi' L_1$ and $\psi' L_2$, at 1, are the same as those for ψL_1 and ψL_2 . Therefore, in the new position, $(\varphi K) \cap (\psi' L)$ is an arc-triangle, bounded by $\varphi K_2, \psi' L_2, \varphi K_1$, in this positive order. This has a unique infinite point, namely 1, and does not contain a paracircle with this infinite point. Hence by Part I (i.e., [5]), (2) and (3), a non-trivial congruence, admitted by it, is an axial symmetry w.r.t. a unique axis passing through 1.

Observe that this axial symmetry of $(\varphi K) \cap (\psi'L)$ is also an axial symmetry of $[(\varphi K) \cap (\psi'L)] \cap P = (\varphi K) \cap P$, for P a paracircle with centre 1, which in B^2 is a sufficiently small circle (satisfying $P \cap (\varphi K) \cap (\psi'L_2) = \emptyset$). We have symmetry of $(\varphi K) \cap P$, w.r.t. the common base line of φK_1 and φK_2 , i.e., the part of the real axis in the model. Even this is the unique axial symmetry of $(\varphi K) \cap P$, with axis of symmetry having 1 as an infinite point.

Then $(\varphi K) \cap (\psi' L)$ and $(\varphi K) \cap P$ coincide in some neighbourhood of $1 \in \mathbb{C}$. Hence also the unique axis of symmetry of $(\varphi K) \cap (\psi' L)$, having 1 as an infinite point, is the part of the real axis in the model. Then this axial symmetry should preserve the arc-side of $(\varphi K) \cap (\psi' L)$ on $\psi' L_2$. Therefore the transversal intersection of the real axis with $\psi' L_2$ in the conformal model should be an orthogonal intersection. However then in B^2 , the real axis should intersect the closure in B^2 of the hypercycle $\psi' L_2$, at 1, at a right angle as well, which is impossible. Hence, in the new position, $(\varphi K) \cap (\psi' L)$ admits no non-trivial congruence, which is a contradiction. Therefore our indirect hypotesis that, e.g., the curvature of φK_i was less than that of ψL_i , was false. This proves (E) of (3) of our theorem.

3. Next, as a second special case, we suppose that K and L are congruent, and both are bounded by two hypercycles with common base-line.

By a suitable notation, we have that the curvatures of K_1 and L_1 are equal, and also that the curvatures of K_2 and L_2 are equal. We are going to show that K_1 and K_2 , and then also L_1 and L_2 , each has the same curvature, i.e., that (E) of (3) of our theorem holds. Suppose, for contradiction, that, e.g., the (non-negative) curvature of K_1 is less than the (positive) curvature of K_2 .

Again, like in **2**, K, or L is symmetrical w.r.t. any straight line orthogonal to the common base line of the K_i 's, or of the L_i 's, resp. Therefore we derive a contradiction for any φ and ψ , independently of their orientation preserving/reversing property.

Now we choose a new pair φ, ψ (different from that in 1). Let us fix φK so that its points at infinity are ± 1 , and φK_1 lies in the closed lower half-plane, and φK_2 lies in the open upper half-plane (observe that possibly φK_1 is a straight line – but φK_2 is not). Let us obtain ψL by rotating φK about the infinite point 1 in positive sense a bit, with the image of φK_i being ψL_i . Then $(\varphi K) \cap (\psi L)$ is bounded by two arcs, one lying on φK_1 , the other one lying on ψL_2 . Then $(\varphi K) \cap (\psi L)$ has one non-smooth point, at $(\varphi K_1) \cap (\psi L_2)$. Hence a non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$ is an axial symmetry, w.r.t. the angle bisector of the angle of $(\varphi K) \cap (\psi L)$ at this point. However, this axial symmetry interchanges the two portions of the boundary, lying on φK_1 and ψL_2 . This contradicts our indirect hypothesis that the curvature of φK_1 is less than the curvature of φK_2 , i.e., the curvature of ψL_2 . This contradiction shows (E) of (3) of our theorem.

4. We turn to the general case. We begin with case (a), i.e., when $(\varphi K) \cap (\psi L)$ has a central symmetry. In this case we will show (E) of (3) of our theorem.

Then φK_1 and φK_2 have two common infinite points (images of each other by this symmetry), and the same statement holds for ψL_1 and ψL_2 , hence the graphs of both φK and ψL are 2-cycles. Clearly then the curvatures of φK_1 and φK_2 , as well as those of ψL_1 and ψL_2 , are equal, and are positive. That is, we have the case settled in **2**, and hence (E) of (3) of our theorem holds.

5. Each of the cases (b) and (c) in 1 describes two axial symmetries. Therefore, unless we have (E) of (3) of our theorem, we have to exclude simultaneously four possible axial symmetries. First observe that our question about $(\varphi K) \cap (\psi L)$, for arbitrary φ and ψ , is equivalent to the same question with ψ = identity. (5.7.1)

So let ψ = identity. (This hypothesis will be applied in the remainder of this proof.)

Then we are going to show that unless we have (E) of (3) of our theorem, we have the following. A generic arbitrary small perturbation φ' of φ (in the topology on Isom⁺(H^2), given by (5.3), cf. also (5.4)) simultaneously destroys all four possible axial symmetries from above.

Actually, in **6**, in case (α) (including the case of symmetry w.r.t. l_1), (E) of (3) of our theorem will be proved directly.

In the remaining three cases, i.e., symmetry w.r.t. l_2 , in case (β) , and symmetries w.r.t. l_3 and l_4 , we show the following. The *i*'th possible congruence (for $1 \le i \le 3$) admitted by $(\varphi'K) \cap (\psi L) = (\varphi'K) \cap L$ does not exist for φ' in a set of the form $U_i(\varphi) \setminus N_i(\varphi)$. Here $U_i(\varphi)$ is an open neighbourhood of φ , and $N_i(\varphi)$ is a relatively nowhere dense, relatively closed subset of $U_i(\varphi)$.

By elements of topology, this can be rewritten as follows. The *i*'th possible congruence $(1 \leq i \leq 3)$ admitted by $(\varphi'K) \cap L$ does not exist for φ' in a set of the form $U_i(\varphi) \setminus M_i$. Here $U_i(\varphi)$ is as above, and M_i is a nowhere dense, closed subset of $\text{Isom}^+(H^2)$. Then none of the investigated three axial symmetries hold on $\bigcap_{1 \leq i \leq 3} [U_i(\varphi) \setminus M_i] = [\bigcap_{1 \leq i \leq 3} U_i(\varphi)] \setminus [\bigcup_{1 \leq i \leq 3} M_i]$. Recall that a finite union of nowhere dense closed subsets is also nowhere dense and closed. Hence none of the investigated three axial symmetries hold on a set of the form $[\bigcap_{1 \leq i \leq 3} U_i(\varphi)] \setminus N(\varphi)$, where $N(\varphi)$ is a relatively nowhere dense, relatively closed subset of $\bigcap_{1 \leq i \leq 3} U_i(\varphi)$. Therefore, for a generic small perturbation φ' of φ , the set $(\varphi'K) \cap L$ will have none of the investigated three possible axial symmetries.

6. We turn to the case when $(\varphi K) \cap L$ has an axial symmetry w.r.t. a straight line l spanned by some diagonal. Then symmetry w.r.t. l interchanges $(\varphi K_1) \cup (\varphi K_2)$ and $L_1 \cup L_2$. Therefore these last two sets are also axially symmetric images of each other.

Hence the graphs of φK and L either

(α) are simultaneously 2-cycles, with edges $\varphi K_1, \varphi K_2$, and L_1, L_2 , or

(β) simultaneously contain paths of length 2, namely (φK_1)(φK_2) and L_1L_2 , resp.

Observe that if the symmetry w.r.t. l interchanges two arc-sides of our arcquadrangle, then it interchanges the entire hypercycles containing them, and also their base-lines. Hence in case (α) we have that bd (φK) = (φK_1) \cup (φK_2) and bd $L = L_1 \cup L_2$ are symmetric images of each other w.r.t. l. Hence also φK and Lare symmetric images of each other w.r.t. l. However then we have the situation discussed in **3**, and hence (E) of (3) of our theorem holds.

Recall that in **1** of this proof, φK and $\psi L = L$ were chosen as follows. The considered common infinite point of φK_1 and φK_2 was 1, the considered common infinite point of L_1 and L_2 was *i*, and the other end-points of φK_1 and φK_2 (possibly coinciding) were close to -1, and the other end-points of L_1 and L_2 (possibly coinciding) were close to -i.

By **1** of this proof $(\varphi K) \cap L$ is a compact arc-quadrangle, such that the distance of 0 to it is small (possibly is 0). It has two diagonals. One of these diagonals connects the intersection points $(\varphi K_1) \cap L_1$ and $(\varphi K_2) \cap L_2$, and spans the straight line l_1 . The other diagonal connects the intersection points $(\varphi K_2) \cap L_1$ and $(\varphi K_2) \cap L_1$ and $(\varphi K_1) \cap L_2$, and spans the straight line l_2 . (It will be convenient to identify a point x with the one-point set $\{x\}$.)

A. If l_1 is an axis of symmetry of $(\varphi K) \cap L$, then also the entire hypercycles φK_1 and L_1 , as well as φK_2 and L_2 are symmetric images of each other w.r.t. l_1 . Thus the common infinite point 1 of φK_1 and φK_2 has as symmetric image w.r.t. l_1 a common infinite point of L_1 and L_2 , close to -i in \mathbb{C} , hence different from their common infinite point *i*. Hence *L* is bounded by L_1 and L_2 . Similarly, φK is bounded by φK_1 and φK_2 . Then φK_1 and L_1 are symmetric images of each other w.r.t. l_1 . Since both φK and *L* are axially symmetric, therefore by $\mathbf{2} \ \varphi K_1$ and L_1 are also directly congruent. Thus we have case (α) , which above was shown to imply (E) of (3) of our theorem.

B. If l_2 is an axis of symmetry of $(\varphi K) \cap L$, then also the entire hypercycles φK_1 and L_2 , as well as φK_2 and L_1 , are symmetric images of each other w.r.t. l_2 . In case (α) again we have (E) of (3) of our theorem.

Therefore suppose case (β) . First we show that some generic small perturbation φ' has the property, that the analogously defined l'_2 (i.e., the straight line spanned by the vertices $(\varphi'K_2) \cap L_1$ and $(\varphi'K_1) \cap L_2$) is not an axis of symmetry of $(\varphi'K) \cap L$. More exactly, we will show that l'_2 is not an angle bisector of the angle of our perturbed arc-quadrangle at its vertex at $(\varphi'K_2) \cap L_1$.

The arc-sides of our perturbed arc-quadrangle on $\varphi' K_2$ and on L_1 (which by the conformal model intersect transversally at the vertex $(\varphi' K_2) \cap L_1$, hence the convex angle determined by them is unique) determine uniquely the angle-bisector a' of our perturbed arc-quadrangle at this vertex. (For φK_2 and/or L_1 straight line/s we have to consider also on which side of this/these straight lines our arc-quadrangle lies.)

Now we apply to φK a small non-zero translation along the base line of φK_2 , obtaining $\varphi' K$. This translation preserves φK_2 setwise – hence a' does not change by this translation. Moreover, it rotates φK_1 non-trivially about the common infinite point 1 of φK_2 and φK_1 . Then $(\varphi' K_1) \cap L_2 \not\subset a'$. Namely else $\emptyset \neq$ $(\varphi' K_1) \cap (L_2 \cap a') \subset (\varphi' K_1) \cap (\varphi K_1) = \emptyset$. (Since different rotated copies of φK_1 , about 1, are disjoint.) In other words, the perturbed diagonal and angle-bisector are different, hence we do not have axial symmetry of $(\varphi' K) \cap L$ w.r.t. the perturbed straight line l'_2 . Hence the investigated symmetry does not exist for the perturbed arc-quadrangle.

Let us choose a small connected open neighbourhood $U(\varphi)$ of φ . We may assume that the infinite points of $\varphi' K_1, \varphi' K_2, L_1, L_2$ (altogether six infinite points) are different, and cyclically follow each other in the same positive sense as those of $\varphi K_1, \varphi K_2, L_1, L_2$. As shown above, for some $\varphi' \in U(\varphi)$ we have that l'_2 does not bisect the investigated angle of our arc-quadrangle. Therefore the equality of the angles into which l'_2 divides our angle, for some $\varphi' \in U(\varphi)$, is expressed by an analytic equation for φ' , which is not an identity. Hence, by the principle of analytic continuation, this equality is satisfied only for φ' in a relatively nowhere dense relatively closed subset of $U(\varphi)$. Hence, all such (φ') 's, with $(\varphi'K) \cap L$ satisfying the symmetry property in **B**, in case (β) , form a nowhere dense closed set in $U(\varphi)$.

7. There remained the case, when $(\varphi K) \cap L$ has an axial symmetry w.r.t. a common orthogonal bisector straight line of two opposite edges. If these edges lie on φK_1 and φK_2 , then let this line be l_3 . Hence l_3 is a common orthogonal straight line to the base-lines of φK_1 and φK_2 . Moreover, this axial symmetry interchanges the arc-sides on L_1 and L_2 , hence also the base-lines of L_1 and L_2 (case (c) in 1).

Observe that a common orthogonal straight line to the parallel base-lines of φK_1 and φK_2 exists only if these base-lines coincide. In this case the graph of φK is a 2-cycle, and φK is bounded just by φK_1 and φK_2 . Let the other common infinite point of φK_1 and φK_2 be -1 (and let $\psi L = L$ be as described in the third paragraph of **1** of the proof of this lemma). The axis l_3 of our symmetry is a straight line orthogonal to this common base line, i.e., to the real axis.

On the other hand, l_3 is the unique axis of symmetry interchanging the parallel base-lines of L_1 and L_2 , hence contains *i*. (Since our axial symmetry w.r.t. l_3 , admitted by $(\varphi K) \cap L$, yields an axial symmetry, admitted by each of φK_1 , φK_2 , and $L_1 \cup L_2$.)

Thus l_3 is the imaginary axis. Hence the common base-line of φK_1 and φK_2 , and the unique axis of symmetry of $L_1 \cup L_2$, interchanging L_1 and L_2 , are orthogonal. However, all such φ 's form a closed set in our topology (given by (5.3)), which is moreover nowhere dense. Namely, a small rotation of φK , about the point of intersection of the above two lines, destroying this orthogonality, is possible.

For the case of axial symmetry, w.r.t. the common orthogonal bisector straight line l_4 to the opposite edges on L_1 and L_2 , we proceed analogously. (Observe that the roles of φ and ψ can be changed: we may fix φ as identity, and then ψ will vary.)

8. Hence, by 5 of this proof, we obtain the following. In 6, case (α) was settled (which included the case of symmetry w.r.t. l_1). In the remaining cases we had the following. Case (β), with symmetry w.r.t. l_2 , was investigated in 6. The cases of symmetries w.r.t. l_3 and l_4 were investigated in 7. In all these remaining three cases we obtained the following. For a generic small perturbation φ' of φ , we had that $(\varphi'K) \cap L$ had none of the remaining three axial symmetries. Summing up: we have proved (E) of (3) of our theorem. \blacksquare

Lemma 5.8. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let the connected boundary components both of K and L be hypercycles or straight lines. Moreover, suppose (5.2), case (2). Then this leads to a contradiction.

Proof. 1. Let, e.g., the graph of φK consist of vertex disjoint edges, whose number is at least 2. Let us choose two vertex-disjoint edges of this graph, φK_1 , φK_2 , say. Further, let the graph of ψL contain a path of length 2 or a 2-cycle, consisting of ψL_1 and ψL_2 , where ψL_2 follows ψL_1 in the positive orientation on bd (ψL) (if we have a 2-cycle, then their numeration is done in some way). We are going to show that this case cannot occur.

We fix φK and thus φK_1 and φK_2 , and will choose ψL in the following way. The set φK lies in the closed convex set bounded by φK_1 and φK_2 . Then we have relatively open arcs I_1 and I_2 of the boundary of the (conformal) model, bounded by the infinite points of φK_1 and φK_2 , resp., and lying outside the closure in the closed unit circle B^2 of the closed convex set bounded by φK_1 and φK_2 . We choose the (considered) common infinite point of ψL_1 and ψL_2 close to the midpoint 1 of I_1 , with ψL_2 following ψL_1 there in the positive sense, and the other infinite points of ψL_1 and ψL_2 (possibly coinciding) close to the midpoint of I_2 . We suppose that φK_1 separates 1 and φK_2 .

Then $(\varphi K) \cap (\psi L)$ is contained in a compact arc-quadrangle Q, bounded by arcs lying on $\varphi K_1, \psi L_2, \varphi K_2, \psi L_1$, in this order, say.

Observe that all boundary components of ψL , other than ψL_1 and ψL_2 , are in the model very close to the boundary of the model, hence cannot cut off parts of this arc-quadrangle Q, which arc-quadrangle is not close to this boundary. So these boundary components have no arcs on bd $[(\varphi K) \cap (\psi L)]$. So we need not deal with these boundary components.

However, there may exist several boundary components φK_i of φK , with $i \neq 1, 2$, which cut off parts of this arc-quadrangle, hence have disjoint non-trivial closed arcs on bd $[(\varphi K) \cap (\psi L)]$. However, there can be only finitely many of these, since each compact set in H^2 meets only finitely many (φK_i) 's.

Since we investigate case (5.2), (2), we have that the (φK_i) 's have no common endpoints. Of course, ψL_1 and ψL_2 have at least one common endpoint. However, by construction, neither ψL_1 and any φK_i (including φK_1 and φK_2), nor ψL_2 and any φK_i (including φK_1 and φK_2), have any common endpoint.

2. We are going to show that

(5.8.1) $\begin{cases} \text{any non-trivial congruence admitted by } (\varphi K) \cap (\psi L) \text{ is a congruence} \\ \text{admitted by } Q \text{ as well; moreover, it is a congruence mapping each} \\ \text{arc-side of } Q \text{ either to itself, or to the opposite arc-side of } Q. \end{cases}$

We have that $\operatorname{bd} [(\varphi K) \cap (\psi L)]$ consists of arcs, following each other, in the positive sense, lying on $\varphi K_1, \psi L_2, \varphi K_{i(1)}, \psi L_2, \varphi K_{i(2)}, ..., \varphi K_{i(j)}, \psi L_2, \varphi K_2, \psi L_1, \varphi K_{i(j+1)}, \psi L_1, \varphi K_{i(j+2)}, ..., \varphi K_{i(k)}, \psi L_1$, in this order, say. From all of these arcs only those lying on ψL_1 and ψL_2 lie on different hypercycles, which have at least one infinite end-point in common.

Let us introduce a symmetric relation \mathcal{R} on the arc-sides of $(\varphi K) \cap (\psi L)$. For two such arc-sides S_1, S_2 we write $S_1\mathcal{R}S_2$, if the hypercycles spanned by these sides are different, and have at least one common end-point. Clearly any nontrivial congruence χ of $(\varphi K) \cap (\psi L)$ preserves this relation \mathcal{R} (i.e., $S_1\mathcal{R}S_2 \iff$ $\chi(S_1)\mathcal{R}\chi(S_2)$), hence also the set of arc-sides $\mathcal{S} := \{S_1 \mid \exists S_2 \text{ such that } S_1\mathcal{R}S_2\}$. Observe that the relation \mathcal{R} induces a complete bipartite graph on the vertex set \mathcal{S} , with classes \mathcal{L}_i , for i = 1, 2, where \mathcal{L}_i is the set of arc-sides of $(\varphi K) \cap (\psi L)$, lying on ψL_i , for i = 1, 2.

Therefore, each non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$ preserves the two-element set $\{\mathcal{L}_1, \mathcal{L}_2\}$. Of course, also the cyclic order of the arc-sides of $(\varphi K) \cap (\psi L)$ is preserved, up to inversion. Let the first end-point of the first arc-side and last endpoint of the last arc-side in \mathcal{L}_i (i.e., lying on ψL_i), in the positive order, be $v_{i,1}$ and $v_{i,2}$. Then the set $\{\{v_{1,1}, v_{1,2}\}, \{v_{2,1}, v_{2,2}\}\}$ is preserved by each non-trivial congruence admitted by $(\varphi K) \cap (\psi L)$, as well. So (5.8.1) is proved.

By (5.8.1) we need to discuss only the congruences of the arc-quadrangle Q, more exactly only those of them, which satisfy the property in the second half of (5.8.1). Therefore, like in the proof of Lemma 5.7, last paragraph of **1**, the possible non-trivial congruences admitted by Q, to be investigated and to be excluded, are central symmetry, and axial symmetries w.r.t. common orthogonal side-bisector straight lines of opposite sides.

Below, if there is an axial symmetry interchanging ψL_1 and ψL_2 (which is unique, if it exists, and is determined by ψL), then its axis of symmetry will be denoted by $a(\psi L)$. Like in the proof of Lemma 5.7, **3**, we set $\psi = identity$.

3. We begin with the case of central symmetry of Q. The central symmetry interchanges the arc-sides lying on φK_1 and φK_2 , hence also φK_1 and φK_2 , hence also the infinite points of φK_1 , and the infinite points of φK_2 . Thus its centre must be the intersection of the straight lines connecting the interchanged endpoints. (This determines the interchanged pairs of infinite points uniquely: these pairs separate each other on the boundary of the model.) We denote this centre of symmetry, which is determined by φK , by $c(\varphi K)$. Also, by central symmetry, the graph of $\psi L = L$ must be a 2-cycle, with L_1 and L_2 having the same curvatures. Then $c(\varphi K)$ lies on the common base line of L_1 and L_2 , which is a(L). (Else the distances of $c(\varphi K)$ from L_1 and L_2 would be different.) Hence $c(\varphi K)$ lies on a(L). Clearly the set of such φ 's is a nowhere dense closed set.

4. We continue with the case of axial symmetry of Q w.r.t. the common orthogonal bisector of the arc-sides lying on L_1 and L_2 . Such a common orthogonal straight line is orthogonal to the base lines of L_1 and L_2 as well, hence it exists only if the base lines of L_1 and L_2 coincide (they cannot be parallel but different), i.e., the graph of L is a 2-cycle. Then this symmetry interchanges φK_1 and φK_2 , hence its axis is the axis of the unique axial symmetry interchanging φK_1 and φK_2 . We denote this axis of symmetry, which is determined by φK , by $a'(\varphi K)$. Then reflection w.r.t. $a'(\varphi K)$ preserves the common base line of L_1 and L_2 . Therefore this common base line either coincides with the axis of reflection, or intersects it orthogonally. The first case is impossible, since L_1 and L_2 , and hence also their common base line is not contained in the closed convex set bounded by φK_1 and φK_2 . Hence $a'(\varphi K)$ intersects orthogonally the common base line of L_1 and L_2 , which is a(L). That is, $a'(\varphi K)$ intersects orthogonally a(L). Clearly the set of such φ 's is a nowhere dense closed set.

5. We continue with the case of axial symmetry of Q w.r.t. the common orthogonal bisector straight line of the arc-sides lying on φK_1 and φK_2 . This axis is the unique straight line orthogonal to φK_1 and φK_2 (and hence also to their base lines). We denote this axis of symmetry, which is determined by φK , by $a''(\varphi K)$. Hence, $a''(\varphi K)$ coincides with a(L). Clearly the set of such φ 's is a nowhere dense closed set.

6. Considering all three possible cases in **3**, **4**, **5**, we apply **3** of the proof of Lemma 5.7. Thus we obtain that for a generic small perturbation of φ , we have that $(\varphi K) \cap L$ admits none of the possible three non-trivial congruences. This is a contradiction, which proves the lemma.

Lemma 5.9. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let the connected boundary components both of K and L be hypercycles or straight lines. Moreover, suppose (5.2), case (3). Then this leads to a contradiction.

Proof. Let, e.g., the graph of φK contain a single edge φK_1 , i.e., this is the unique boundary component of φK , and let the graph of ψL contain a path of length 2 or a 2-cycle. We are going to show that this is impossible.

Let the graph of ψL contain two edges ψL_1 and ψL_2 with a common vertex at 1. Let the other endpoints of ψL_1 and ψL_2 be close to -1. We consider the conformal model. We consider ψL_1 and ψL_2 as fixed, and φK as being in a small Euclidean neighbourhood of 1. Then φK does not intersect any other ψL_i , so we may leave them out of consideration. Moreover, let φK_1 intersect both ψL_1 and ψL_2 . Then φK lies on that side of φK_1 , as 1.

Then $(\varphi K) \cap (\psi L)$ is an arc-triangle, with one infinite vertex at 1. Hence any of its non-trivial congruences is an axial symmetry, w.r.t. an axis passing through 1 (it cannot be a non-trivial rotation about the infinite point 1, by Part I (i.e., [5]), (3)). Moreover, ψL_1 and ψL_2 , as well as their base lines, are images of each other by this axial symmetry. Then this axis of symmetry, interchanging these base lines, is uniquely determined by ψL (else $(\varphi K) \cap (\psi L)$ would admit a non-trivial rotation about the infinite point 1, which is excluded by (3)). This axis intersects the arcside of $(\varphi K) \cap (\psi L)$ on φK_1 orthogonally. However, a small generic rotation of φK , and thus also of φK_1 , about the intersection point of the above axis of symmetry and φK_1 , destroys orthogonality of this intersection, hence destroys also this unique congruence.

In the proof of the next lemma, in case (5.2), (4), we will need boundaries of closed convex sets K, L with interior points, with finitely many (but at least one) hypercycle or straight line boundary components in H^2 , and also their boundaries in the (conformal or collinear) model circle together with its boundary S^1 . The second boundaries of closed convex sets with interior points are homeomorphic to S^1 , and contain the first ones. In case (5.2), (4), with the finiteness hypothesis, their set theoretical difference (second one minus first one) is the union of at least one but finitely many disjoint closed proper circular subarcs of S^1 (in case (5.2), (4)).

Lemma 5.10. Assume (*) with d = 2 and $X = H^2$. Suppose (1) of Theorem 5. Let the connected boundary components both of K and L be finitely many hypercycles or straight lines. Moreover, suppose (5.2), case (4). Then this leads to a contradiction.

Proof. By hypothesis, both graphs are finite. By (5.2), (4), both graphs contain only vertex-disjoint edges, and, e.g., the graph of L has at least two edges. We are going to show that this is impossible.

Let ψL_1 and ψL_2 denote two neighbourly boundary components of ψL , with ψL_2 following ψL_1 in the positive orientation. (That is, passing on the boundary of ψL , meant in one of the model circles together with its boundary circle, from ψL_1 to ψL_2 , in the positive sense, there are no other connected components of bd (ψL) between them. Here we use finiteness of the set of the boundary components of L. Cf. also Remark 3.) Then, denoting by ψl_1 the second infinite point of ψL_1 , and by ψl_2 the first infinite point of ψL_2 (both taken in the positive orientation), the counterclockwise open arc (ψl_1)(ψl_2) of S^1 contains no infinite point of any boundary component of ψL . Then the base lines of ψL_1 and ψL_2 are symmetric images of each other w.r.t. a unique axis $a(\psi L)$, determined by ψL . The infinite points of $a(\psi L)$ are $p(\psi L)$ and $q(\psi L)$, where $p(\psi L)$ lies in the counterclockwise open arc (ψl_1)(ψl_2) $\subset S^1$. We suppose that $a(\psi L)$ lies on the real axis, and then $p(\psi L) = 1$ and $q(\psi L) = -1$.

Let φK_1 be a boundary component of φK . Let its infinite end-points be $\varphi k'_1$ and $\varphi k''_1$, following each other in this order in the positive sense, on the boundary of φK , meant in the model together with its boundary circle. Let us begin with the position, when $\varphi k'_1 = \psi l_2$ and $\varphi k''_1 = \psi l_1$, and φK lies on the same side of φK_1 , as 1.

Now let us translate φK , and thus also φK_1 , along the real axis a bit, so that $\varphi k'_1$ and $\varphi k''_1$ move on S^1 in the direction from 1 to -1, both on the respective open arcs of S^1 from 1 to -1, in which they are contained. For the new congruent copy of K we will not apply a new notation, but will preserve the old notation φK . We want to determine the intersection $(\varphi K) \cap (\psi L)$.

Let the boundary components of φK , or of ψL , be, in the positive sense, $\varphi K_1, \varphi K_2, ..., \varphi K_n$, or $\psi L_1, ..., \psi L_m$, resp. By (5.2), (4), using the collinear model, we see that any of φK and ψL can be obtained from a convex polygon, with all vertices at infinity (not counted to the polygon), whose number of vertices is even, in the following way. We have that φK or ψL contains the polygon, and bd (φK) or bd (ψL) is obtained from the boundary of the polygon the following way. We replace each second side of the polygon by a hypercycle with this base line (outwards of the polygon), and remove each other second side of the polygon. (Including the case when this convex polygon is a 2-gon, i.e., a segment.)

Then we may suppose that all boundary components of the new φK , except φK_1 (if any), lie strictly on the other side of the straight line $(\psi l_1)(\psi l_2)$, as the new φK_1 . (Here we use finiteness of the set of the boundary components of K. Cf.

also Remark 3.) All these are boundary components of $(\varphi K) \cap (\psi L)$ as well. There is still one boundary component of $(\varphi K) \cap (\psi L)$. This begins at ψl_2 , then passes on ψL_2 , then passes on φK_1 , then on some $\psi L_{i(1)}$, then once more on φK_1 , then on some $\psi L_{i(2)}$, ..., then on some $\psi L_{i(k)}$, then once more on φK_1 , then on ψL_1 , and ends at ψl_1 , with all these arc-sides being nondegenerate. (There can be no common arc of φK_1 and any ψL_i . For ψL_1 and ψL_2 this is evident. For L_i , with $i \geq 3$, this could occur only if we had $\varphi K_1 = \psi L_i$, and then ψL_i would intersect both ψL_1 and ψL_2 , a contradiction.) Then

(5.10.1)
$$\begin{cases} \text{any non-trivial congruence admitted by } (\varphi K) \cap (\psi L) \text{ preserves} \\ \text{this unique non-smooth boundary component } C \text{ of } (\varphi K) \cap (\psi L). \end{cases}$$

The extension to B^2 of a congruence admitted by C, in particular, preserves the pair of its infinite points. That is, either each of them is mapped to itself, or they are exchanged by the congruence. If each of them is mapped to itself by the congruence, then the semi-infinite hypercycle-arc side of C lying on ψL_2 is also mapped to itself by the congruence. Then by induction one sees that each hypercycle-arc side of Cis mapped to itself. Thus all three vertices of some triangle (one infinite and two finite vertices of the first two arc-sides of C) are preserved, hence the congruence is trivial.

If the pair of infinite points of C is exchanged by the congruence, then similarly one sees that all the hypercycle-arc sides of C are mapped to all the hypercyclearc sides of C, but their order is reversed. The congruence preserves also the property on which side of C is $(\varphi K) \cap (\psi L)$ contained. Hence the congruence is orientation-reversing, and then by Part I (i.e., [5]), (5) it is an axial symmetry. Then it exchanges the hypercycle-arc sides of C lying on ψL_1 and ψL_2 , hence it exchanges ψL_1 and ψL_2 as well. Therefore its axis is $a(\psi L)$, which lies on the ξ -axis, cf. the second paragraph of this proof. Moreover, it exchanges the first and last arcs of φK_1 on C (these may coincide). Hence this congruence maps φK_1 onto itself. Therefore its axis $a(\psi L)$, which intersects φK_1 transversally, is orthogonal to φK_1 .

However, a small rotation of φK , and thus also of φK_1 , about the intersection point of the axis of symmetry $a(\psi L)$ and φK_1 , preserves the combinatorial structure of $(\varphi K) \cap (\psi L)$ (only possibly the set of indices $\{i(1), ..., i(k)\}$ will change, but this does not invalidate the above considerations). Simultaneously, this small rotation destroys orthogonality of the intersection of $a(\psi L)$ and φK_1 , hence destroys also this unique non-trivial congruence.

Proof of Theorem 5, continuation. 6. Now the earlier parts of the proof of Theorem 5, and Lemmas 5.1-5.10 end the proof of Theorem 5.

Proof of Theorem 6. 1. We have $(4) \Longrightarrow (1)$ by [4], Theorem 1. We have evidently $(1) \Longrightarrow (2)$. There remains to prove $(4) \Longrightarrow (3)$ and $(2) \Longrightarrow (4)$ and $(3) \Longrightarrow (4)$. These we do in the next three lemmas.

Lemma 6.1. In Theorem 6 we have $(4) \Longrightarrow (3)$.

Proof. We copy the proof of [4], Lemma 1.3, which says just $(4) \implies (1)$ in our Theorem 6.

1. Let K and L be congruent balls (in S^d , \mathbb{R}^d or H^d).

For $\varphi K = \psi L$ we do not have Theorem 6, (2), (C), hence the implication (A) \land (B) \land (C) \Longrightarrow (3) is satisfied. Further let $\varphi K \neq \psi L$.

We denote by o the midpoint of the segment S with endpoints the centres of φK and ψL , and by H the orthogonal bisector hyperplane of S. Then by $\operatorname{int} [(\varphi K) \cap (\psi L)] \neq \emptyset$ the length of S is less than the diameter of K, hence $o \in \operatorname{int} [(\varphi K) \cap (\psi L)]$. Then φK and ψL are symmetric images of each other w.r.t. H. Hence both $(\varphi K) \cap (\psi L)$ and S are symmetric w.r.t. H. We claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from $\operatorname{bd} [(\varphi K) \cap (\psi L)]$ is attained only for z = o.

We suppose that the image of H in the conformal model lies on the $x_1 \ldots x_{d-1}$ coordinate hyperplane, the image of S lies on the x_d -axis, whence the image of o is 0. We say that φK lies lower than ψL . We may suppose that $z \in (\varphi K) \cap (\psi L)$ lies on or above H, and $z \notin bd(\varphi K)$. Then the minimum distance of z and the part of $bd(\varphi K)$ above H is realized by a geodesic segment, from z to some point of $bd(\varphi K)$ above H, which orthogonally intersects $bd(\varphi K)$. Then this geodesic segment lies on a radius of φK . Fixing this radius and varying z on it, the maximum distance is attained only for $z \in H$. Consider a ball φK_0 , concentric with φK , such that $o \in bd(\varphi K_0)$.

Then on every radius of φK , the signed distances of the intersection points of this radius with $\operatorname{bd}(\varphi K)$ and $\operatorname{bd}(\varphi_1 K)$ are constant. By the conformal model, $(\varphi K_0) \setminus \{o\}$ lies strictly below H. Therefore the maximum distance of $z \in (\varphi K) \cap (\psi L)$, lying on or above H, to the part of $\operatorname{bd}(\varphi K)$ above H, is attained only for z = o.

This implies our claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from bd $[(\varphi K) \cap (\psi L)]$ is attained only for z = o. It also follows that the farthest points of bd $[(\varphi K) \cap (\psi L)]$ from o are the intersections of the line, spanned by S, with bd $[(\varphi K) \cap (\psi L)]$.

Therefore $(\varphi K) \cap (\psi L)$ has a unique inball, with centre o. So Theorem 6, (2), (B) is satisfied. Moreover, Theorem 6, (2), (C) is also satisfied, with $\{\varphi x, \psi y\}$ being the intersection of bd $[(\varphi K) \cap (\psi L)]$ with the straight line, spanned by S. Last, Theorem 6, (3) is also satisfied, with the orthogonal bisector hyperplane of $S = [\varphi x, \psi y]$ being H.

2. Now let us consider any compact intersection of two paraballs φK and ψL , with nonempty interior (in H^d). Then, like in [4], Lemma 1.3, their infinite points, φk and ψl , say, are different. Let the other points of bd (φK) and bd (ψL) on the line (φk)(ψl) be $\varphi k'$ and $\psi l'$. Like in [4], Lemma 1.3, the order of these points on this line is $\varphi k, \psi l', \varphi k', \psi l$, and these points are different. We denote by o the midpoint of $S := [\varphi k', \psi l']$ (then $o \in int [(\varphi K) \cap (\psi L)])$, and by H the orthogonal bisector hyperplane of S. Then φK and ψL are symmetric images of each other w.r.t. H. Hence both (φK) $\cap (\psi L)$ and S are symmetric w.r.t. H. We claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from bd $[(\varphi K) \cap (\psi L)]$ is attained only for z = o.

We suppose that the image of H in the conformal model lies on the $x_1 \ldots x_{d-1}$ coordinate hyperplane, and the image of φk or ψl is $(0, \ldots, 0, -1)$ or $(0, \ldots, 0, 1)$,

resp., whence the image of o is 0. We say that φK lies lower than ψL . Like in 1, we may suppose that $z \in (\varphi K) \cap (\psi L)$ lies on or above H, and $z \notin bd(\varphi K)$. Then the minimum distance of z and the part of $bd(\varphi K)$ above H is realized by a geodesic segment, from z to some point of $bd(\varphi K)$ above H, which orthogonally intersects $bd(\varphi K)$. Then this geodesic segment lies on a straight line with infinite point φk . Fixing this line and varying z on it, the maximum distance is attained only for $z \in H$. Let us consider another paracycle $\varphi_1 K$, having φk as its infinite point, and with $o \in bd(\varphi_1 K)$ (this is unique).

Then on every straight line with φk as an infinite point, the signed distances of the intersection points of this line with $\operatorname{bd}(\varphi K)$ and $\operatorname{bd}(\varphi_1 K)$ are constant. (The intersection point with $\operatorname{bd}(\varphi_1 K)$ lies between φk and the intersection point with $\operatorname{bd}(\varphi K)$.) However, by the conformal model, $(\varphi_1 K) \setminus \{o\}$ lies strictly below H. Therefore the maximum distance of $z \in (\varphi K) \cap (\psi L)$, lying on or above H, to the part of $\operatorname{bd}(\varphi K)$ above H, is attained only for z = o.

This implies our claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from $\operatorname{bd} [(\varphi K) \cap (\psi L)]$ is attained only for z = o. It also follows that the farthest points of $\operatorname{bd} [(\varphi K) \cap (\psi L)]$ from o are $\varphi k'$ and $\psi l'$.

Therefore $(\varphi K) \cap (\psi L)$ has a unique inball, with centre o. So Theorem 6, (2), (B) is satisfied. Moreover, Theorem 6, (2), (C) is also satisfied, with $\{\varphi x, \psi y\}$ (where x = k' and y = l') being the intersection of $\operatorname{bd} [(\varphi K) \cap (\psi L)]$ with the straight line $(\varphi k)(\psi l)$. Last, Theorem 6, (3) is also satisfied, with the orthogonal bisector hyperplane of $S = [\varphi k', \psi l'] = [\varphi x, \psi y]$ being H.

3. There remains the case when the connected components of both K and L are congruent hyperspheres, with common positive distance, λ , say, from their respective base hyperplanes (in H^d). In [4], proof of Lemma 1.3 it was shown that different hypersphere boundary components of φK , or of ψL , have a distance at least 2λ , and supposing diam $[(\varphi K) \cap (\psi L)] < 2\lambda$, for some boundary components φK_i of φK and ψL_j of ψL , we have $(\varphi K) \cap (\psi L) = (\varphi K_i^*) \cap (\psi L_j^*)$. Here φK_i^* , or ψL_j^* , is the closed convex set bounded by φK_i , or by ψL_j , resp. (Later we will use only compactness of $(\varphi K_i^*) \cap (\psi L_j^*)$.) By [4], Lemma 1.1, the parallel domain of the base hyperplane of K_i , or of L_j , with distance λ , lies in K, or in L, resp.

Suppose that the base hyperplanes of φK_i and ψL_j have a common finite point. Then $\emptyset \neq \text{int} [(\varphi K_i^*) \cap (\psi L_j^*)]$. Moreover, $(\varphi K_i^*) \cap (\psi L_j^*)$ has some infinite point, hence diam $[(\varphi K) \cap (\psi L)] = \infty$, a contradiction.

Suppose that the base hyperplanes of φK_i and ψL_j have a (unique) common infinite point p. Then in the conformal model, as a subspace of \mathbb{R}^d , the stereoangles of φK_i^* and ψL_j^* at p are greater than 1/4 times the total stereoangle, while the conformal model ball, as a subspace of \mathbb{R}^d , has at p a stereoangle 1/2 times the total stereoangle. Hence $\varphi K_i^* \cap \psi L_j^* \neq \emptyset$, and it has p as an infinite point, again implying diam $[(\varphi K) \cap (\psi L)] = \infty$, a contradiction.

Hence the base hyperplanes of φK_i and ψL_j have no common finite or infinite point. Let S be the segment realizing the distance of these two base hyperplanes, o be its midpoint, and H be its orthogonal bisector hyperplane. We say that the base hyperplane of φK_i lies below that of ψL_j . Then φK_i lies above its base hyperplane, and ψL_j lies below its base hyperplane. Namely, in the remaining three cases, $(\varphi K_i^*) \cap (\psi L_j^*)$ would have infinite points, and then diam $[(\varphi K) \cap (\psi L)] = \infty$, a contradiction. Then φK and ψL are symmetric images of each other w.r.t. H. Hence both $(\varphi K) \cap (\psi L)$ and S are symmetric w.r.t. H. Also, using (*), int $[(\varphi K) \cap (\psi L)] \neq \emptyset$ is symmetric w.r.t. o, hence $o \in int[(\varphi K) \cap (\psi L)]$. We claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from bd $[(\varphi K) \cap (\psi L)]$ is attained only for z = o.

We suppose that the image of H in the conformal model lies on the $x_1 \ldots x_{d-1}$ hyperplane, and the image of S lies on the x_d -axis, whence the image of o is 0. Like in 1 and 2, we may suppose that $z \in (\varphi K) \cap (\psi L)$ lies on or above H, and $z \notin bd(\varphi K_i)$. Then the minimum distance of z and the part of φK_i above His realized by a geodesic segment, from z to some point of φK_i above H, which orthogonally intersect φK_i . Thus this geodesic segment lies on a straight line orthogonal to the base hyperplane of φK_i . Fixing this line and varying z on it, the maximum distance is attained only for $z \in H$. Let us consider another hypersphere $\varphi \tilde{K}_i$, with the same base hyperplane as φK_i , such that $o \in \varphi \tilde{K}_i$ (this is unique).

Then on every straight line, orthogonal to our base hyperplane, the signed distances of the intersection points of this line with φK_i and $\varphi \tilde{K}_i$ are constant. However, by the conformal model, $(\varphi \tilde{K}_i) \setminus \{o\}$ lies strictly below H. Therefore the maximum distance of $z \in (\varphi K) \cap (\psi L)$, lying on or above H, to the part of φK_i above H, is attained only for z = o.

This implies our claim that for $z \in (\varphi K) \cap (\psi L)$, the maximum distance from bd $[(\varphi K) \cap (\psi L)]$ is attained only for z = o. It also follows that the farthest points of bd $[\varphi K) \cap (\psi L)]$ from o are the two points of intersection of bd $[(\varphi K) \cap (\psi L)]$ and the line spanned by S.

Therefore $(\varphi K) \cap (\psi L)$ has a unique inball, with centre *o*. So Theorem 6, (2), (B) is satisfied. Moreover, Theorem 6, (2), (C) is also satisfied, with $\{\varphi x, \psi y\}$ being the intersection of bd $[\varphi K) \cap (\psi L)$ with the straight line spanned by *S*. Last, Theorem 6, (3) is also satisfied, with the orthogonal bisector hyperplane of $[\varphi x, \psi y]$ being *H*.

We note that in [4], as a part of the proof of Lemma 1.6, uniqueness of the inball of $(\varphi K) \cap (\psi L)$, for sufficiently small diameter, was proved for K, L being balls in S^d and \mathbb{R}^d , and for K, L having congruent hypersphere boundary components (with hyperplanes excluded) in H^d . In the special case of our Lemma 6.1, the proof in this paper proves a bit more, is shorter, and all three cases **1-3** are treated analogously.

Lemma 6.2. In Theorem 6 we have $(2) \Longrightarrow (4)$.

Proof. Observe that [4], proof of Theorem 1, (1) \implies (2) (i.e., our Theorem 6, (1) \implies (4)) did not use central symmetry of all intersections (φK) \cap (ψL) of "small" diameter, but only those which satisfied (B) and (C) of Theorem 6, (2). (Moreover, in [4], Theorem 1 there occured $\varepsilon = \varepsilon(K, L)$, while in our Theorem 6, (2) we have $\varepsilon = \varepsilon(K, L, x, y)$. This difference will be discussed later.)

In fact, in [4], Lemma 1.6 (2), and in its proof (just below (1.6.1)) the centre of symmetry c of $(\varphi K) \cap (\psi L)$ was identified as the point O such that for some

 $\varepsilon_1 > 0$ we had that $B(O, \varepsilon_1)$ was the unique inball of $(\varphi K) \cap (\psi L)$. This is just our Theorem 6, (2) (B). (Observe that in case of [4], Lemma 1.6 (1) we had that $X = S^d$ and K, L were halfspheres, and then Theorem 6 (4) held.)

To show Theorem 6, (2) (C), we have to recall the construction of $B(O, \varepsilon_1)$ from [4]. In [4], (A) (which is (**) in this paper), for $x \in \operatorname{bd} K$ and $y \in \operatorname{bd} L$, there was asserted the existence of two balls, with the properties stated there. However, if the radii of these balls were chosen less than $\varepsilon_1(x)$, or $\varepsilon_1(y)$, resp. (cf. [4], (1.5.7)), then these balls, except their points x or y, lied actually in int K, or int L, resp. We supposed that the radii of these new balls were equal to some $\varepsilon_1 \in (0, \min\{\varepsilon_1(x), \varepsilon_1(y)\}/2).$

In the first paragraph of the proof of [4] Lemma 1.5, it was proved that either $X = S^d$ and K, L were halfspheres, and then Theorem 6 (4) held, or, e.g., K had an exposed point. Suppose this second case. Then in [4] Lemma 1.7 (2) it was proved that both K and L were strictly convex. ([4] Lemma 1.7 (1) asserted the first case from the beginning of this paragraph.) In [4] (1.5.4), we chose a point $O \in X$ ("origin"), and chose φ_0 and ψ_0 so that $\varphi_0 x = O$ and $\psi_0 y = O$, and $\varphi_0 K$ and $\psi_0 L$ touched each other from outside, at O (recall (**) $\Longrightarrow C^1$).

Then in [4] (1.5.6) and (1.5.7) we moved $\varphi_0 K$ and $\psi_0 L$, and with them the new balls as rigidly attached to them, toward each other. The centres of the new balls moved along the common normal line of $\varphi_0 K$ and $\psi_0 L$ at O, through the distance ε_1 , while we allowed any simultaneous, independent rotations of $\varphi_0 K$ and $\psi_0 L$ about their original common normal. The new positions of $\varphi_0 K$ and $\psi_0 L$ were denoted by φK and ψL . Then the centres of the new balls moved to their common new position O. From above, originally these new balls, except for their points $\varphi_0 x$ and $\psi_0 y$, resp., were contained in int $(\varphi_0 K)$ and int $(\psi_0 L)$, resp. Therefore in their common new position these new balls lied in $[int (\varphi K)] \cap [int (\psi L)] = int [(\varphi K) \cap (\psi L)]$, except for two points of their common normal (hence were antipodal), and were the centres of the new balls before the translation and rotations.

The later steps of the proof only used central symmetry in the above described positions. Moreover, observe that in [4], proof of Theorem 1, (1) \implies (2) in (1) there occured the hypothesis $\varepsilon = \varepsilon(K, L)$, stronger than in our Theorem 6, (2). However, actually in the proof only one $x \in \text{bd } K$ and one $y \in \text{bd } L$ were used at a time, to prove equality of all sectional curvatures of bd K at x with all sectional curvatures of bd L at y (and their positivity for $X = \mathbb{R}^d$ and $X = H^d$). Hence we may admit that ε depends also on x and y, as in our Theorem 6, (2).

In fact, recall that φK and ψL were not uniquely determined, but they could be independently rotated about the original common normal line. (This line contained O. Suppose that the image of O in one of the models is 0. Then the images of these rotations in the model are Euclidean rotations about the image of this line in the model.) Therefore any two-dimensional normal section of bd (φK) could become the centrally symmetric image of any two-dimensional normal section of bd (ψL), with the central symmetry interchanging φx and ψy . Hence the centre of symmetry was the midpoint of [$\varphi x, \psi y$]. (For S^d another centre of symmetry was the point of S^d antipodal to this midpoint. However, the central symmetries w.r.t. these two centres as maps coincide, so we may leave the other centre of symmetry out of consideration.) From above the curvature of any two-dimensional normal section of $\operatorname{bd}(\varphi K)$ at φx , and the curvature of any two-dimensional normal section of $\operatorname{bd}(\psi L)$ at ψy coincide, as claimed above. What was written above, readily implies also the local rotational symmetry of both $\operatorname{bd} K$ and $\operatorname{bd} L$, at any of their points x and y (w.r.t. the normal of $\operatorname{bd} K$ at x, and that of $\operatorname{bd} L$ at y – which exist by (**)).

In [4], Lemma 1.5 (1) constantness and equality (and for \mathbb{R}^d and H^d also positivity) of all sectional curvatures of bd K and bd L at all their points x and y were proved. In [4], Lemma 1.5 (2) the local rotational symmetry of both bd K and bd L, at any of their points x and y was proved. From these the global statement of [4], Theorem 1, (2), i.e., the global statement of (4) of our theorem, was deduced in [4], Lemmas 1.8 and 1.9. This ends the proof of (2) \Longrightarrow (4) in Theorem 6.

Lemma 6.3. In Theorem 6 we have $(3) \Longrightarrow (4)$.

Proof. We repeat the proof of Lemma 6.2, except that we replace the first sentence in italics, in its last but three paragraph, by the following. Therefore any twodimensional normal section of $\operatorname{bd}(\varphi K)$ could become the symmetric image of any two-dimensional normal section of $\operatorname{bd}(\psi L)$, via the symmetry w.r.t. the orthogonal bisector hyperplane of $[\varphi x, \psi y]$.

Proof of Theorem 6, continuation. 2. Now the proof of Theorem 6 follows from the earlier parts of the proof of Theorem 6, and from Lemmas 6.1-6.3.

The proof of Theorem 7 will use several ideas from the proof of Theorem 4 in [4].

Proof of Theorem 7. 1. Suppose (7) of Theorem 7. By (****), for S^d the centres of φK and ψL are not antipodal. Hence the midpoint of the (shorter) segment joining these centres is a centre of symmetry of

(7.1)
$$M := \operatorname{conv} \left[(\varphi K) \cup (\psi L) \right].$$

Hence Theorem 7, (7) \implies (1) holds. Moreover, if we suppose (7) of Theorem 7, then either $\varphi K = \psi L$ and then (2) of Theorem 7 holds vacuously, or else there is a unique diametral segment D of conv $[(\varphi K) \cup (\psi L)]$, and the orthogonal bisector hyperplane of D is the orthogonal bisector hyperplane of the segment with endpoints the centres of K and L, and M is symmetrical w.r.t. it. Hence Theorem 7, (7) \implies (2) holds.

The implications Theorem 7, $(1) \Longrightarrow (3)$ and $(2) \Longrightarrow (4)$ are evident.

There remains to show Theorem 7, $(3) \Longrightarrow (5)$, $(4) \Longrightarrow (6)$, $(5) \Longrightarrow (7)$ and $(6) \Longrightarrow (7)$.

Lemma 7.1. Assume (****). Assume the hypotheses of Theorem 7. Then in Theorem 7 we have $(3) \Longrightarrow (5)$ and $(4) \Longrightarrow (6)$.

Proof. Suppose (3) or (4) of Theorem 7, resp. As in part **2** of the proof of Theorem 4 in [4] (this part does not use smoothness of K and L), we have the following. For any $x \in \text{bd } K$ and $y \in \text{bd } L$, there exists a ball $B(o, R) \subset X$ (for S^d we may

suppose even that B(o, R) lies in the open southern hemisphere, which implies that B(o, R) does not contain any antipodal pair of points of S^d), with centre o and radius R (less than $\pi/2$ but arbitrarily close to $\pi/2$ for S^d , and arbitrarily large for \mathbb{R}^d and H^d), and there exist φ and ψ with the following properties.

(1) $\varphi x, \psi y$ are antipodal points of bd B(o, R) (thus $\varphi x \neq \psi y$; but, in the case of S^d , they are not antipodal in S^d),

(2) $\varphi K, \psi L \subset B(o, R)$; even φK and ψL are contained in balls of radius some r < R, contained in B(o, R), and tangent to B(o, R) at φx and ψy , resp.,

(3) by (2), $(\operatorname{bd} B(o, R)) \cap [\operatorname{bd} (\varphi K)] = \{\varphi x\}$, and $(\operatorname{bd} B(o, R)) \cap [\operatorname{bd} (\psi L)] = \{\psi y\}$. Then, in the collinear model, any point of $\operatorname{bd} B(o, R)$, except φx and ψy , can be separated by a hyperplane, intersecting int B(o, R), from both of φK and ψL , hence also from M. Therefore $(\operatorname{bd} B(o, R)) \cap (\operatorname{bd} M) = \{\varphi x, \psi y\}$.

(4) B(o, R) is the unique ball of minimal radius containing M.

Then (4) implies that any congruence admitted by M leaves B(o, R) invariant. This holds in particular for central symmetry, and for symmetry w.r.t. the orthogonal bisector hyperplane of the unique diametral segment $D := [\varphi x, \psi y]$ of M. Hence for central symmetry the centre of symmetry is o, and both of these symmetries interchange φx with ψy . (For S^d another centre of symmetry is -o, but the central symmetries w.r.t. o and -o, as set maps, are identical, so we may leave -o out of consideration.)

By its definition, $D(\varepsilon)$ is also centrally symmetric w.r.t. o, and also is symmetric w.r.t. the orthogonal bisector hyperplane of D. Therefore in case (3) of Theorem 7, we have that $M \cap D(\varepsilon)$ is also centrally symmetric (w.r.t. o). Similarly, in case (4) of Theorem 7, we have that $M \cap D(\varepsilon)$ is also symmetric w.r.t. the orthogonal bisector hyperplane of D. Hence (5) or (6) of Theorem 7 holds, resp., for any $\varepsilon(K, L, x, y, \varphi, \psi, S) \in (0, \pi/2)$ for S^d , and for any $\varepsilon(K, L, x, y, \varphi, \psi) \in (0, \infty)$ for \mathbb{R}^d and H^d .

So there remains to prove $(5) \Longrightarrow (7)$ and $(6) \Longrightarrow (7)$ of Theorem 7.

Lemma 7.2. Assume (****). Assume the hypotheses of Theorem 7. Then both of (5) and (6) of Theorem 7 imply that both of K and L are C^1 .

Proof. It suffices to give the proof for K, the other case is analogous.

Suppose that $x \in \text{bd } K$ is not a smooth point of bd K. Recall that convex bodies are almost everywhere smooth (cf. [6]). (This takes over from \mathbb{R}^d to S^d and H^d via the collinear models.) Then let $y \in \text{bd } L$ be a smooth point of bd L. Further, let obe the midpoint of D (from Theorem 7, (3), (B)).

Now we consider the collinear model, so that o becomes the centre 0 of the model. Then the image of $B(o, R) \subset X$ from the proof of Lemma 7.1 is a Euclidean ball in the collinear model, with centre 0. The images of points, sets in this collinear model, as a subset of \mathbb{R}^d , will be denoted by ', like $(\varphi x)'$, (B(o, R))', etc. The radius of (B(o, R))' is denoted by R', and $D(\varepsilon)'$ is in this model a right cylinder, of height 2R', and has bases (d-1)-balls, of radius ε' . Further, diam' (\cdot) denotes the diameter in the Euclidean sense, in the collinear model, as a subset of \mathbb{R}^d . Moreover, we suppose that (1)-(4) from the proof of Lemma 7.1 hold.

Let $(\varphi x)'$ be the north pole and $(\psi y)'$ the south pole of (B(o, R))'. Then the horizontal supporting planes of (B(o, R))' at its poles are supporting hyperplanes

of M' (cf. (7.1)) and $M' \cap D(\varepsilon)'$ as well. At $(\psi y)'$ this is the unique supporting hyperplane of $(\psi L)'$, but at $(\varphi x)'$ there are also other supporting planes of $(\varphi K)'$, even there are ones which are not horizontal, but are arbitrarily close to horizontal. Let H' be one such supporting plane.

We have diam' $[(\varphi K)']$, diam' $[(\psi L)'] < \text{diam'} [(B(o, R))']$. Therefore the highest point of $(\psi L)'$ lies strictly below the height of the north pole. By a suitable, almost horizontal choice of the supporting plane H' we can attain that the ball segment cut off by H' from (B(o, R))', lying "above" H', lies entirely above the highest point of $(\psi L)'$. Then both $(\varphi K)'$ and $(\psi L)'$ lie "below" H', hence also M' and $M' \cap D(\varepsilon)'$ lie "below" H', and then H' is a supporting hyperplane of $M' \cap D(\varepsilon)'$ at $(\varphi x)'$.

In case of central symmetry of $M' \cap D(\varepsilon)'$ w.r.t. o' = 0 (for Theorem 7, (5)) then -H' is a supporting hyperplane of $M' \cap D(\varepsilon)'$ at $(\psi y)' = -(\varphi x)'$. In case of symmetry of $M' \cap D(\varepsilon)'$ w.r.t. the orthogonal bisector hyperplane of D, we have that the symmetrical image \tilde{H}' of H' w.r.t. this orthogonally bisecting hyperplane (for Theorem 7, (6)) is a supporting hyperplane of $M' \cap D(\varepsilon)'$ at $(\psi y)' = -(\varphi x)'$. Since being a supporting plane is a local property, therefore then -H', or \tilde{H}' is a supporting hyperplane of M' at $(\psi y)'$ as well, resp. Then -H', or \tilde{H}' is also a supporting hyperplane of $(\psi L)' \subset M'$ at $(\psi y)'$, resp., which is not horizontal, a contradiction to smoothness of bd L at y.

In Lemma 7.3 smoothness of K and L follows from Lemma 7.2. However, also for the proof of Theorem 8 we will use Lemma 7.3, where smoothness of K and L will follow from the hypotheses of Theorem 8, for S^2 , and from Lemma 8.2 for \mathbb{R}^2 and H^2 . On the other hand, strict convexity of K and L follows both from the hypotheses of Theorem 7 and Theorem 8.

Lemma 7.3. Assume (****). Assume the hypotheses of Theorem 7. Let K and L be C^1 , and let $x \in \text{bd } K$ and $y \in \text{bd } L$. Then, in the situation described in (1) to (4) of the proof of Lemma 7.1, and with M from (7.1), we have the following. In some open neighbourhood of φx , we have that $\text{bd}(\varphi K)$ and bd M coincide. Also, in some open neighbourhood of ψy , we have that $\text{bd}(\psi L)$ and bd M coincide.

Proof. It suffices to give the proof for $\operatorname{bd}(\varphi K)$, the case of $\operatorname{bd}(\psi L)$ is analogous.

It suffices to show that in some neighbourhood of $(\varphi x)'$ we have that $\operatorname{bd} [(\varphi K)']$ and $\operatorname{bd} M'$ coincide.

We use the collinear model, and the coordinate system from the proof of Lemma 7.2 on it. We use ' as in the proof of Lemma 7.2 (e.g., o' = 0).

Let $(\varphi x)' = R'e_d$; then $(\psi y)' = -R'e_d$. Denoting by h(M', u) the support function of M', etc., for $u \in S^{d-1}$, we have

(7.3.1)
$$\begin{cases} h(M', e_d) = h((\varphi K)', e_d) = R' \text{ and} \\ h((\psi L)', e_d) \le \operatorname{diam}'[(\psi L)'] - R' =: R' - c' < R'. \end{cases}$$

Since $M', (\varphi K)', (\psi L)' \subset (B(o, R))' = B(0, R')$, therefore for their support functions we have for $u_1, u_2 \in S^{d-1}$ that $|h(M', u_1) - h(M', u_2)| \leq R' \cdot ||u_1 - u_2||$,

etc.

Let $u \in S^d$, with $R' \cdot ||u - e_d|| < c'/2$. That is, u can vary in the open spherical cap $C' \subset S^{d-1}$, of centre e_d and spherical radius $2 \arcsin [c'/(4R')]$. Then $h((\varphi K)', u) > h((\varphi K)', e_d) - c'/2 = R' - c'/2$, and $h((\psi L)', u) < h((\psi L)', e_d) + c'/2 < R' - c'/2$. This implies that

(7.3.2)
$$\begin{cases} u \in C' \Longrightarrow h(M', u) = h\left(\operatorname{conv}\left[(\varphi K)' \cup (\psi L)'\right], u\right) \\ = \max\{h\left((\varphi K)', u\right), h\left((\psi L)', u\right)\} = h\left((\varphi K)', u\right). \end{cases}$$

By the supporting sphere hypothesis, K and L, hence also $(\varphi K)'$ and $(\psi L)'$ are strictly convex, and by the hypothesis of this lemma they are C^1 . Then their spherical maps (i.e., $(\varphi x^*)' \in \operatorname{bd}[(\varphi K)']$ is mapped to the outer unit normal of $(\varphi K)'$ at $(\varphi x^*)'$, and similarly for $(\psi L)'$, are homeomorphisms $\operatorname{bd}[(\varphi K)'] \to S^{d-1}$, and $\operatorname{bd}[(\psi L)'] \to S^{d-1}$, resp.

It will be convenient to identify a singleton set $\{a\}$ with a. The support function of a convex body $A \subset \mathbb{R}^d$ can be extended from S^{d-1} to \mathbb{R}^d , in a positively homogeneous way (at 0 it is defined as 0). Now recall the following (cf. [1], [9]). (1) Suppose that for the supporting hyperplane H(A, u) of A, with outer unit nor-

(1) Suppose that for the supporting hyperplane H(A, u) of A, with outer unit normal $u \in S^{d-1}$, the intersection $A \cap H(A, u)$ is a singleton. Then the gradient of the extended support function of A at u exists, and equals this singleton point. (2) Conversely, if this gradient exists at $u \in S^{d-1}$, then $A \cap H(A, u)$ is a singleton.

Applying (1) to the strictly convex body $(\varphi K)'$, we get that the gradient of the extended support function of $(\varphi K)'$ exists at any $u \in S^{d-1}$, and its value is the singleton $(\varphi K)' \cap H((\varphi K)', u)$. However, by (7.3.2), on the open subset $C' \subset S^{d-1}$ the extended support function of $(\varphi K)'$ equals the extended support function of M'. Hence the gradient of this second function also exists on C', and its value is the same singleton $(\varphi K)' \cap H((\varphi K)', u)$. Then applying (2) to the convex body M', we get that for $u \in C'$ the set $M' \cap H(M', u)$ is a singleton. Once more applying (1), but now for the convex body M', we get that this singleton equals the gradient of M' at $u \in C'$. Thus $(\varphi K)' \cap H((\varphi K)', u)$ and $M' \cap H(M', u)$ are the (existing) gradients of the same function, at the same point $u \in C'$, hence they are equal.

Summing up: for $u \in C'$ the singleton support sets of $(\varphi K)'$ and M' at u coincide. The union of all these singleton sets, for all $u \in C'$, are the inverse spherical images of the open subset $C' \subset S^{d-1}$, for $(\varphi K)'$ and M'. But the spherical map and the inverse spherical map for $(\varphi K)'$ are homeomorphisms, hence the inverse spherical image I' of $C' \subset S^{d-1}$ is open in bd $[(\varphi K)']$. By the invariance of domain theorem for bd (M') we have that I' is open in bd (M') as well.

Now we prove local coincidence of $\operatorname{bd} [(\varphi K)']$ and $\operatorname{bd} (M')$ at $(\varphi x)'$. In fact, let $B'((\varphi x)', \delta')$ be a ball of centre $(\varphi x)'$ and radius δ' (meant in \mathbb{R}^2), contained in int B(0,1). Let $\delta' > 0$ be so small, that $I' \supset [\operatorname{int} B'((\varphi x)', \delta')] \cap \operatorname{bd} [(\varphi K)']$ and $I' \supset [\operatorname{int} B'((\varphi x)', \delta')] \cap \operatorname{bd} (M')$. Then

(7.3.3)
$$\begin{cases} z' \in [\operatorname{int} B'((\varphi x)', \delta')] \cap \operatorname{bd} [(\varphi K)'] \Longrightarrow z' \in I' \subset \operatorname{bd} (M').\\ \operatorname{Similarly}, z' \in [\operatorname{int} B'((\varphi x)', \delta')] \cap \operatorname{bd} (M') \Longrightarrow z' \in \operatorname{bd} [(\varphi K)'] \end{cases}$$

The next lemma will be used also in the proof of Theorem 8. Observe that, under the hypotheses of Lemma 7.2, both K and L are C^1 . In the proof of Theorem 8, the C^1 property of K and L is a hypothesis of Theorem 8 for S^2 , while for \mathbb{R}^2 and H^2 , under the hypotheses of Lemma 8.2, it will follow from Lemma 8.2.

Lemma 7.4. Assume (****). Assume the hypotheses of Theorem 7. Let K and L be C^1 , and let $x \in \operatorname{bd} K$ and $y \in \operatorname{bd} L$. Then, in the situation described in (1) to (4) of the proof of Lemma 7.1, in some neighbourhood of φx , and in some neighbourhood of ψy , we have that $\operatorname{bd} M$ and $\operatorname{bd} [M \cap D(\varepsilon)]$ coincide.

Proof. We use the collinear model, and the coordinate system from the proof of Lemma 7.2 on it. We use ' as in the proof of Lemma 7.2.

By the supporting sphere hypothesis, both K and L are strictly convex. Moreover, by the hypothesis of this lemma, they are also C^1 . Then $(\varphi K)'$ and $(\psi L)'$ are also strictly convex and C^1 .

For our lemma it suffices to show that

(7.4.1)
$$\begin{cases} \text{ in some neighbourhood of } (\varphi x)', \text{ and of } (\psi y)', \text{ we} \\ \text{ have that } \mathrm{bd} (M') \text{ and } \mathrm{bd} [[M \cap D(\varepsilon)]'] \text{ coincide.} \end{cases}$$

Let $[D_h(\varepsilon)]'$ be the closed parallel strip, whose boundary hyperplanes are spanned by the two (horizontal) bases of the cylinder $[D(\varepsilon)]'$. Let $[D_v(\varepsilon)]'$ be the closed bothway infinite cylinder, with directrices spanned by the (vertical) directrices of the cylinder $[D(\varepsilon)]'$.

We assert that

(7.4.2)
$$\begin{cases} \text{the neighbourhood of } (\varphi x)', \text{ or of } (\psi y)', \text{ whose existence} \\ \text{is claimed in (7.4.1), can be chosen as } [\text{int } D_v(\varepsilon)]'. \end{cases}$$

First observe that $\{\varphi x, \psi y\}' \subset \operatorname{bd} M'$ and $\{\varphi x, \psi y\}' \subset \operatorname{bd} [M \cap D(\varepsilon)]'$. Therefore it remains to show that $[(\operatorname{bd} M) \setminus \{\varphi x, \psi y\}]'$ and $[[\operatorname{bd} [M \cap D(\varepsilon)]] \setminus \{\varphi x, \psi y\}]'$ coincide in some neighbourhood of $(\varphi x)'$, or of $(\psi y)'$, resp., where this neighbourhood is $[\operatorname{int} D_v(\varepsilon)]'$. That is, we have to prove

(7.4.3)
$$\begin{cases} [(\operatorname{bd} M) \setminus \{\varphi x, \psi y\}]' \cap [\operatorname{int} D_v(\varepsilon)]' = \\ [[\operatorname{bd} [M \cap D(\varepsilon)]] \setminus \{\varphi x, \psi y\}]' \cap [\operatorname{int} D_v(\varepsilon)]'. \end{cases}$$

We are going to prove that both sides of (7.4.3) contain the other one.

Recall the formulas $\operatorname{bd}(A \cup B) \subset (\operatorname{bd} A) \cup (\operatorname{bd} B)$ and $\operatorname{bd}(A \cap B) \subset (\operatorname{bd} A) \cup (\operatorname{bd} B)$. Then by the proof of Lemma 7.1, (2) and (3) we have

(7.4.4)
$$\begin{cases} [(\operatorname{bd} M) \setminus \{(\varphi x), (\psi y)\}]', [[\operatorname{bd} [M \cap D(\varepsilon)]] \setminus \{\varphi x, \psi y\}]' \\ \subset [\operatorname{int} B(o, R)]' \subset [\operatorname{int} D_h(\varepsilon)]', \end{cases}$$

which implies

$$\begin{array}{l} (7.4.5) \\ \left\{ \begin{array}{l} [(\mathrm{bd}\,M) \setminus \{\varphi x, \psi y\}]' = [(\mathrm{bd}\,M) \setminus \{\varphi x, \psi y\}]' \cap [\mathrm{int}\,D_h(\varepsilon)]' \text{ and} \\ \left[\left[\mathrm{bd}\,[M \cap D(\varepsilon)]\right] \setminus \{\varphi x, \psi y\}\right]' = \left[\left[\mathrm{bd}\,[M \cap D(\varepsilon)]\right] \setminus \{\varphi x, \psi y\}\right]' \cap [\mathrm{int}\,D_h(\varepsilon)]'. \end{array} \right. \end{array}$$

Intersecting the right hand side sets in (7.4.5) with $[\operatorname{int} D_v(\varepsilon)]'$, we get

(7.4.6)
$$\begin{cases} [(\operatorname{bd} M) \setminus \{\varphi x, \psi y\}]' \cap [\operatorname{int} D(\varepsilon)]' \text{ and} \\ [[\operatorname{bd} [M \cap D(\varepsilon)]] \setminus \{\varphi x, \psi y\}]' \cap [\operatorname{int} D(\varepsilon)]', \text{ resp.}, \end{cases}$$

and we have to show that both of these sets contains the other one.

We have
$$(7.4.7)$$

$$\begin{cases} (\operatorname{IAA}) \setminus \{\varphi x, \psi y\}]' \cap [\operatorname{int} D(\varepsilon)]' = \left[\left[\operatorname{bd} \left[[M \cap D(\varepsilon)] \cup [M \setminus D(\varepsilon)] \right] \right] \setminus \left\{ \varphi x, \psi y\} \right]' \cap [\operatorname{int} D(\varepsilon)]' \subset \left\{ \left[\left[\operatorname{bd} [M \cap D(\varepsilon)] \right] \setminus \{\varphi x, \psi y\} \right]' \cap [\operatorname{int} D(\varepsilon)]' \right\} \cup \left\{ \left[\operatorname{cl} \left[X \setminus D(\varepsilon) \right] \right]' \cap [\operatorname{int} D(\varepsilon)]' \right\} = \left[\left[\operatorname{bd} \left[M \cap D(\varepsilon) \right] \right] \setminus \{\varphi x, \psi y\} \right]' \cap [\operatorname{int} D(\varepsilon)]', \end{cases}$$

i.e., the first set in (7.4.6) is contained in the second set in (7.4.6).

On the other hand,

(7.4.8)
$$\begin{cases} \left[\left[\operatorname{bd} \left[M \cap D(\varepsilon) \right] \right] \setminus \{\varphi x, \psi y\} \right]' \cap \left[\operatorname{int} D(\varepsilon) \right]' \subset \left[\left[\left(\operatorname{bd} M \right) \cup \left[\operatorname{bd} D(\varepsilon) \right] \right] \\ \setminus \{\varphi x, \psi y\} \right]' \cap \left[\operatorname{int} D(\varepsilon) \right]' \subset \left\{ \left[\left(\operatorname{bd} M \right) \setminus \{\varphi x, \psi y\} \right]' \cap \left[\operatorname{int} D(\varepsilon) \right]' \right\} \cup \\ \left\{ \left[\operatorname{bd} D(\varepsilon) \right]' \cap \left[\operatorname{int} D(\varepsilon) \right]' \right\} = \left[\left(\operatorname{bd} M \right) \setminus \{\varphi x, \psi y\} \right]' \cap \left[\operatorname{int} D(\varepsilon) \right]', \end{cases}$$

i.e., the second set in (7.4.6) is contained in the first set in (7.4.6). This ends the proof that the two sets in (7.4.6) coincide. Then the two sets in (7.4.3) also coincide, and (7.4.1) is proved, and hence the statement of the lemma is proved.

Lemma 7.5. Assume (****). Assume the hypotheses of Theorem 7. Let K and L be C^1 . Then both of (5) and (6) of Theorem (6) imply that both of bd K and bd L are at each of their boundary points twice differentiable (cf. 6. Preliminaries). Moreover, in the situation described in (1) to (4) in the proof of Lemma 7.1, for any 2-plane P containing $D = [\varphi x, \psi y]$, the curvatures of $(\varphi K) \cap P$ at φx and that of $(\psi L) \cap P$ at ψy coincide.

Proof. As stated in 6. Preliminaries, convex surfaces are almost everywhere twice differentiable, in the sense as given there. If, say, bd K is not twice differentiable at its point x, then we will choose $y \in \text{bd } L$ so that bd L is twice differentiable at y.

Now let us consider the situation described in (1) to (4) in the proof of Lemma 7.1. Then, by Lemma 7.3, in some neighbourhood of φx we have that $\operatorname{bd}(\varphi K)$ and $\operatorname{bd} M$ coincide. Then $\operatorname{bd} M$ is not twice differentiable at its point φx . Similarly, in some neighbourhood of ψy we have that $\operatorname{bd}(\psi L)$ and $\operatorname{bd} M$ coincide. Then $\operatorname{bd} M$ is twice differentiable at its point φx .

However, the central symmetry of $M \cap D(\varepsilon)$ about the centre o (in case of (5) of Theorem 7), or its symmetry w.r.t. the orthogonal bisector hyperplane of D (in

case of (6) of Theorem 7), interchanges φx and ψy , as well as some of their open neighbourhoods in bd M. Then bd M cannot be not twice differentiable at φx , but twice differentiable at ψy . This contradiction proves the first statement of the lemma.

The second statement of the lemma follows analogously: two congruent, twice differentiable curves have at their points φx and ψy , corresponding to each other by the congruence, equal curvatures.

The following lemma is an analogue of [4], Lemma 1.8.

Lemma 7.6. Assume (****). Assume the hypotheses of Theorem 7. Then both (5) and (6) of Theorem 7 imply that both bd K and bd L are, at each of their boundary points, locally, relatively open subsets of spherical surfaces, of some fixed sectional curvature $\kappa > 0$ for S^d and \mathbb{R}^d , and $\kappa > 1$ for H^d . For $X = S^d$ the radius of these spherical surfaces is less than $\pi/2$.

Proof. Again consider the situation described in (1) to (4) of the proof of Lemma 7.1. Like in the proof of Lemma 6.2, observe that φK and ψL are not uniquely determined, but we may apply any rotations about the line spanned by D to any of them, independently of each other. Thus varying φ and ψ in this way, by Lemma 7.3, the central symmetry of $M \cap D(\varepsilon)$ about the centre o (in case of Theorem 7, (5)), or its symmetry w.r.t. the orthogonal bisector hyperplane of D (in case of Theorem 7, (6)), may interchange locally a 2-dimensional normal section of M, hence of φK at φx , with any 2-dimensional normal section of M, hence of ψL at ψy . This immediately implies that both bd (φK) and bd (ψL) are locally rotationally symmetric about the line spanned by D. By Lemma 7.5, the curvatures of the 2-dimensional normal sections of φK at φx , and of ψL at ψy , exist and coincide.

Therefore, all these (congruent) 2-dimensional normal sections have curvatures at φx and ψy , resp., and all these curvatures are equal to some number $\kappa \geq 0$. Then the supporting sphere hypothesis implies that $\kappa > 0$ for S^d and \mathbb{R}^d , and $\kappa > 1$ for H^d .

We still have to show that these 2-dimensional normal sections are circular arcs. We claim that the 2-dimensional normal sections N of $[\operatorname{bd}(\varphi K)] \cap D(\varepsilon)$ containing φx (or of $[\operatorname{bd}(\psi L)] \cap D(\varepsilon)$ containing ψy) are 2-dimensional normal sections at any of their other points φx^* (or ψy^*), lying in the same open halfspace bounded by the orthogonal bisector hyperplane of D, as x (or y). This means just that the normal of $\operatorname{bd}(\varphi K)$ at φx^* lies in the 2-plane spanned by N, which is evident for d = 2. However, for $d \geq 3$, local rotational symmetry of $\operatorname{bd}(\varphi K)$ at φx , which necessarily has as rotation axis the normal of $\operatorname{bd}(\varphi K)$ at φx^* . Thus the uniquely determined normal of $\operatorname{bd}(\varphi K)$ at φx^* lies in any of these hyperplanes, so it lies in their intersection, which is just the 2-plane spanned by this rotation axis and φx^* . Thus this intersection is spanned by N, proving our claim.

Therefore N has at each of its points φx^* , lying in the same open halfspace bounded by the orthogonal bisector hyperplane of D, as x, the constant curvature κ . Thus N locally is a cycle. The already proved inequalities for κ in this Lemma imply that N locally is a circular arc. Therefore bd (φK) is locally a spherical surface, of sectional curvature κ . For $X = S^d$, by the supporting sphere hypothesis, this spherical surface has a radius less $\pi/2$.

Proof of Theorem 7, continuation. 2. We apply Lemma 1.9 of [4]. Its hypothesis, for the case of the existence of the supporting spheres at any boundary point, is just the statement of our Lemma 7.6. (Additionally there is as hypothesis (1) of Theorem 1 or (1) of Theorem 4 of [4]. However, these are actually not used in the proof of Lemma 1.9 of [4]. Observe still that the C_{α} 's in the proof of Lemma 1.9 of [4] are correctly unions of equivalence classes there.) Its conclusion implies that the connected components both of K and L are congruent spheres (for $X = S^d$ of radius at most $\pi/2$), paraspheres or hyperspheres, but in case of congruent spheres K and L are congruent balls. However, the inequalities for κ in our Lemma 7.6 exclude paraspheres and hyperspheres. So K and L are congruent balls, for $X = S^d$ of radius less than $\pi/2$, by Lemma 7.6. This ends the proof of Theorem 7.

Proof of Theorem 8. 1. We have the evident implications $(7) \Longrightarrow (1)$, and $(1) \Longrightarrow (2) \Longrightarrow (4)$, and $(1) \Longrightarrow (3) \Longrightarrow (4)$, and $(5) \Longrightarrow (6)$. Therefore we have to prove only the implications $(3) \Longrightarrow (5)$, and $(4) \Longrightarrow (6)$, and $(6) \Longrightarrow (7)$.

We will use the notation M from (7.1).

Lemma 8.1. Assume (****) with d = 2. Assume the hypotheses of Theorem 8. Then in Theorem 8 we have $(3) \Longrightarrow (5)$, and $(4) \Longrightarrow (6)$.

Proof. We begin with copying the proof of Lemma 7.1. We may suppose that there is a circle B(o, R), satisfying (1)-(4) from the proof of Lemma 7.1, hence also (B) from Theorem 7 (3), and also satisfying (A) from Theorem 7 (3). Hence the hypotheses of Theorem 8, (3) and (4) are satisfied.

Observe that any congruence admitted by M is a congruence admitted by its unique diametral segment D as well. Therefore the non-trivial congruences admitted by M form a subset of {central symmetry w.r.t. o, axial symmetry w.r.t. the straight line spanned by D, axial symmetry w.r.t. the orthogonal bisector line of D}. Observe that $D(\varepsilon)$ admits all these three non-trivial congruences.

Hence, in case of Theorem 8 (3), axial symmetry of M has as axis one of the above two axes. Moreover, $D(\varepsilon)$ has the same axis of symmetry (in case of both axes). Therefore $M \cap D(\varepsilon)$ also has the same axis of symmetry, i.e., (5) is satisfied.

In case of Theorem 8 (4), in an analogous way, any non-trivial congruence admitted by M is a non-trivial congruence admitted by $D(\varepsilon)$ as well, hence also $M \cap D(\varepsilon)$ admits this non-trivial congruence, i.e., (6) is satisfied.

Proof of Theorem 8, continuation. 2. Observe that if for K and L we have supporting circles, of radius some r (with $r < \pi/2$ for S^2), then we have supporting circles of any larger radius (also less than $\pi/2$ for S^2) as well. Now we define R as $(r + \pi/2)/2 < \pi/2$ for S^2 , and as 2r for \mathbb{R}^2 and H^2 . If for S^2 we have that $r < \pi/2$ is arbitrarily close to π , or for \mathbb{R}^2 and H^2 we have that r is arbitrarily large, then the same statements hold for R rather than r. Occasionally we will use some other R > r. Then we suppose (1)-(4) in the proof of Lemma 7.1, meant with such an Ras written above. In the proof of the implication (6) \Longrightarrow (7) of this theorem we will use only sets M with diameter 2R (which for S^2 can be arbitrarily close to π , and for \mathbb{R}^2 and H^2 can be arbitrarily large), and having unique diametral segments, as will be clear from the proof. Hence the hypothesis Theorem 7, (3) (A) will be satisfied. Also, by (3) of the proof of Lemma 7.1, the hypothesis Theorem 7, (3) (B) will be satisfied.

Lemma 8.2. Assume (****) with d = 2. Assume the hypotheses of Theorem 8. Then for $X = \mathbb{R}^2$ and $X = H^2$, (6) of Theorem 8 implies that K and L are C^1 .

Proof. 1. Let us suppose that, e.g., K is not smooth, and for $x \in \operatorname{bd} K$ it has an inner angle $\alpha \in (0, \pi)$. Since $\operatorname{bd} L$ is almost everywhere smooth, we choose $y \in \operatorname{bd} L$ as a smooth point of of $\operatorname{bd} L$. Now let us consider, for this x and this y, the situation described in (1) to (4) in the proof of Lemma 7.1. We suppose that the centre o of B(o, R) is mapped to the point 0 in the collinear model. Further we suppose, as in the proof of Lemma 7.3, that $(\varphi x)' = R'e_2$ and $(\psi y)' = -R'e_2$. We will investigate $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ in a sufficiently small neighbourhood of φx . Since $D(\varepsilon) \cap B(o, R)$ is a fixed neighbourhood of $D = [\varphi x, \psi y]$ relative to B(o, R), therefore for any fixed ε a sufficiently small neighbourhood of φx relative to B(o, R), is contained in $D(\varepsilon)$. We are going to ensure that $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ is not even locally axially symmetric at φx w.r.t. the straight line spanned by D. This will then imply that $[\operatorname{conv} [(\varphi K) \cup (\psi L)]] \cap D(\varepsilon)$ is not axially symmetric w.r.t. this line.

2. Since each congruence admitted by conv $[(\varphi K) \cup (\psi L)]$ preserves the endpoints of its unique diametral segment D (whose midpoint is o), therefore the only possible non-trivial congruences admitted by conv $[(\varphi K) \cup (\psi L)]$ are central symmetry w.r.t. o and axial symmetry w.r.t. the orthogonal bisector of D (these interchange φx and ψy), and axial symmetry w.r.t. the line spanned by D. We will consider φx and ψy as the north and south pole of B(o, R) (thus the images in the collinear model of φx and ψy lie on the positive and negative vertical coordinate axis, resp.).

3. We are going to show that we can avoid axial symmetry of conv $[(\varphi K) \cup (\psi L)]$ w.r.t. the line spanned by D, by suitably choosing φ and ψ . Even, this line will not be locally, at φx , an axis of symmetry of conv $[(\varphi K) \cup (\psi L)]$. For this aim we will have to choose a sufficiently large R, greater than the fixed radius r, in dependence of K and L. In the situation described in (1) to (4) in the proof of Lemma 7.1, we have that φK and ψL are contained in balls $B_{\varphi x}$ and $B_{\psi y}$, say, of radius r, both contained in B(o, R), and containing φx and ψy , resp. Then the angle subtended by ψL at φx is at most the angle subtended by $B_{\psi y}$ at φx , which is $\beta := 2 \arcsin[r/(2R-r)]$ or $\beta := 2 \arcsin[\sinh r/\sinh(2R-r)]$ (by the law of sines), for \mathbb{R}^2 or H^2 , resp. We choose R > r so large, that $\beta < \alpha$ should be satisfied. Even actually both legs of the angle subtended by ψL at φx enclose an angle at most $\beta/2 < \alpha/2 < \pi/2$ with the unit vector $-e_2$ (meant in the collinear model).

Our aim is to show that φK can be rotated through a small angle about φx , while preserving ψL , so that conv $[(\varphi K) \cup (\psi L)]$ becomes at φx not even locally axially symmetric w.r.t. the straight line spanned by D.

We will consider the angular domains subtended by φK and ψL at φx . These will be considered as subsets of the lower semicircle of S^1 , which we identify as usual with the angular interval $[-\pi, 0]$. Let these angular intervals for φK and ψL be $[u_1, u_2]$ and $[v_1, v_2]$, resp. Here $-\pi \leq u_1 < u_2 \leq 0$, but $-\pi = u_1$ and $u_2 = 0$ cannot hold simultaneously, by $\alpha \in (0, \pi)$. Further,

(8.2.1)
$$-\pi < -\pi/2 - \beta/2 \le v_1 < -\pi/2 < v_2 \le -\pi/2 + \beta/2 < 0.$$

Here the first and sixth inequalities hold by $\beta < \alpha < \pi$. The second and fifth inequalities hold since both legs of the angle, subtended by ψL at φx , enclose an angle at most $\beta/2$ with $-e_2$ (in the collinear model). The third and fourth inequalities hold by smoothness of bd (ψL) at ψy , which implies $D \cap \operatorname{int} (\psi L) \neq \emptyset$. Moreover, $\pi > \angle (u_1, u_2) = \alpha > \beta \ge \angle (v_1, v_2) > 0$. Then the collinear model shows that the angular domain subtended at φx by conv $[(\varphi K) \cup (\psi L)]$ is

(8.2.2)
$$A := [\min\{u_1, v_1\}, \max\{u_2, v_2\}].$$

(The subtended angular domain clearly contains A. Conversely, there are straight lines in X through φx , of direction vectors corresponding to $\min\{u_1, v_1\}, \max\{u_2, v_2\}$, such that the following holds. The images of these lines in the collinear model have the images both of φK and ψL in the collinear model in the lower side of these image non-vertical lines.) Since $-\pi \leq u_1 < u_2 \leq 0$, but $-\pi = u_1$ and $u_2 = 0$ cannot hold simultaneously, and $-\pi < v_1 < -\pi/2 < v_2 < 0$, therefore the angle of the angular domain A lies in $(0, \pi)$. That is, φx is not a smooth point of conv $[(\varphi K) \cup (\psi L)]$.

Then local symmetry, at φx , of conv $[(\varphi K) \cup (\psi L)]$, w.r.t. the straight line spanned by D, would imply symmetry of the angular domain A w.r.t. the same line (*i.e.*, as a subset of $[-\pi, 0]$, w.r.t. $-\pi/2$). We will investigate the possibilities about the relative position of the angular intervals $[u_1, u_2]$ and $[v_1, v_2]$, and in each case we will find a small rotation of φK about φx , destroying the symmetry of Aw.r.t. the straight line spanned by D.

We may assume that $u_2 < 0$, since else min $\{u_1, v_1\} > -\pi$, and max $\{u_2, v_2\} = 0$, and then A is not symmetric w.r.t. the straight line spanned by D. Analogously we may assume that $u_1 > -\pi$. So in the following we assume $-\pi < u_1 < u_2 < 0$.

By $\angle(u_1, u_2) = \alpha > \beta \ge \angle(v_1, v_2)$ we have either $u_2 > v_2$, or $u_1 < v_1$. These two cases are symmetrical, so it will suffice to deal with the case $u_2 > v_2$. Then we distinguish the following cases:

(A) $u_1 < v_1$,

(B) $v_1 \le u_1$.

4. In Case (A) we have $A = [u_1, u_2] \subset (\pi, 0)$. Then we will show that one can rotate φK about φx through any sufficiently small angle (thus preserving the equality $A = [u_1, u_2]$), and axial symmetry w.r.t. the straight line spanned by D occurs only for one position, namely when the straight line spanned by D is the angle-bisector of the angular domain A. So generically A will not be axially symmetric w.r.t. the straight line spanned by D. By this small rotation we preserve the inclusion $\varphi K \subset (\operatorname{int} B(o, R)) \cup \{\varphi x\}$.

We consider the chords of $B_{\varphi x}$ from φx , in the directions of u_1 and u_2 , and consider the intersection C of $B_{\varphi x}$ with the angular domain with vertex at φx and legs of directions u_1, u_2 . Then $\varphi K \subset C$ (by the collinear model). Now let us choose some $r^* \in (r, R)$, and let $B'_{\varphi x}$ be a circle of radius r^* , containing φx in its

44

boundary, and let $B'_{\varphi x} \subset (\operatorname{int} B(o, R)) \cup \{\varphi x\}$. Then all boundary points of C, lying on bd $B_{\varphi x}$, except φx , originally lied in $\operatorname{int} B'_{\varphi x}$, and had a distance from bd $B'_{\varphi x}$ at least some positive number. So for sufficiently small rotations about φx they still will remain in $\operatorname{int} B'_{\varphi x}$. Therefore the rotated copy of C, being the convex hull of the rotated above mentioned boundary points of C and of φx , lies in $(\operatorname{int} B'_{\varphi x}) \cup \{\varphi x\}$. Hence we have the situation as described in (1) to (4) in the proof of Lemma 7.1, however with r replaced by r^* . Generically the straight line spanned by D is not the angle-bisector of the angular domain A. Hence generically we do not have at φx even a local symmetry of $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ w.r.t. the straight line spanned by D, so also we do not have its global symmetry w.r.t. the same line. This ends the proof of Case (A).

5. In case (B) we have $A = [v_1, u_2]$. Here $v_1 \ge -\pi/2 - \beta/2$, and $u_2 = u_1 + \alpha > v_1 + \beta \ge -\pi/2 + \beta/2$, hence $(v_1 + u_2)/2 > -\pi/2$. Therefore the angular domain A, as a subset of $[-\pi, 0]$, is not symmetric w.r.t. $-\pi/2$, hence this angular domain is not symmetric w.r.t. the straight line spanned by D (cf. 3). Hence also conv $[(\varphi K) \cup (\psi L)]$ is not symmetric w.r.t. this line. This ends the proof of Case (B).

6. As proved in 4 and 5, there are φ and ψ , such that $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ is not axially symmetric w.r.t. the straight line spanned by D. Therefore, by 2, the only possible non-trivial congruences admitted by this set are central symmetry w.r.t. the midpoint o of D, and axial symmetry w.r.t. the orthogonal bisector of D. Both of these symmetries interchange φx and ψy .

In **3** we have seen that φx is not a smooth point of $\operatorname{conv} [(\varphi K) \cup (\psi L)]$. On the other hand, by the choice of y in **1** we have that $\psi y \in \operatorname{bd} (\psi L)$ is a smooth point of ψL . Thus the inner angle of ψL at ψy is π , which is at most the inner angle of $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ at ψy , which is at most π . Hence ψy is a smooth point of $\operatorname{conv} [(\varphi K) \cup (\psi L)]$.

However, as above shown, the only possible non-trivial congruences admitted by $\operatorname{conv} [(\varphi K) \cup (\psi L)]$ should interchange φx and ψy . This is a contradiction, proving Lemma 8.2.

Proof of Theorem 8, continuation. 3. We use the sign ' as introduced in the proof of Lemma 7.2, for the images in the collinear model, as a subset of \mathbb{R}^2 . We use, e.g., (B(o, R))' and $D(\varepsilon)'$ and diam'(\cdot) as in the proof of Lemma 7.2. As in the proof of Lemma 8.2, 1, we suppose o' = 0 and $(\varphi x)' = R'e_2$ and $(\psi y)' = -R'e_2$.

For any $x \in \operatorname{bd} K$ we have by (2) in the proof of Lemma 7.1 that φK is included not only in the circle B(o, R) of radius R, but even in a circle $B_{\varphi x}$ ($\subset B(o, R)$), of radius r (< R), Moreover, $B_{\varphi x}$ also contains φx in its boundary, and is tangent at φx to B(o, R). The analogous statement holds also for ψL .

Recall that both K and L are C^1 , for S^2 by hypothesis of Theorem 8, and for \mathbb{R}^2 and H^2 by Lemma 8.2. They are also strictly convex, by the supporting sphere hypothesis of Theorem 8. Hence the spherical maps $\operatorname{bd}(\varphi K)' \to S^1$ and $\operatorname{bd}(\psi L)' \to S^1$ are homeomorphisms.

Combining Lemmas 7.3 and 7.4 (for d = 2), we get that

J. JERÓNIMO-CASTRO, E. MAKAI, JR.

(8.1) $\begin{cases} \text{ in the situation described in (1) to (4) of the proof of Lemma 7.1,} \\ \text{ in a neighbourhood of } \varphi x, \text{ or of } \psi y, \text{ we have that bd } [M \cap D(\varepsilon)] \text{ coincides with bd } (\varphi K), \text{ or with bd } (\psi L), \text{ resp.} \end{cases}$

By Theorem 8, (6), $M \cap D(\varepsilon)$ admits some non-trivial congruence. Such a congruence also preserves the unique diametral segment D of $M \cap D(\varepsilon)$, hence also its midpoint o. Thus

(8.2) $\begin{cases} a \text{ non-trivial congruence admitted by } M \cap D(\varepsilon) \text{ is either a} \\ central symmetry w.r.t. o, or an axial symmetry w.r.t. the straight line spanned by D or w.r.t. the orthogonal bisector line of D. \end{cases}$

We say that $x \in \operatorname{bd} K$ is a point of local symmetry of $\operatorname{bd} K$ if the following holds. The point x has some open arc neighbourhood relative to $\operatorname{bd} K$, which is symmetric w.r.t. the normal of $\operatorname{bd} K$ at x. Analogously we define points of local symmetry of $\operatorname{bd} L$.

We make a case distinction. Either

(1) each $x \in \text{bd } K$, and each $y \in \text{bd } L$ is a point of local symmetry of bd K, and of bd L, resp., or

(2) some $x_0 \in \text{bd } K$, or some $y_0 \in \text{bd } L$, is not a point of local symmetry of bd K, or of bd L, resp.

In the second statement of Lemma 8.3 compactness of K and L will not be required.

Lemma 8.3. Assume (****) with d = 2. Assume the hypotheses of Theorem 8, and (6) of Theorem 8. Let K and L be C^1 . Then case (1) in **3** of the proof of Theorem 8 implies (7) of Theorem 8. More exactly, suppose that K, L are C^1 , and satisfy (*) – except possibly int $[(\varphi K) \cap (\psi L)] \neq \emptyset$. Then case (1) in **3** of the proof of Theorem 8 is equivalent to that each connected component of the boundaries of K and L is a cycle or a straight line.

Proof. **1.** If each connected component of the boundaries of K and L is a cycle or a straight line, then case (1) in **3** of the proof of Theorem 8 holds evidently.

2. Conversely, suppose case (1) in **3** of the proof of Theorem 8. Then consider any connected component K_i of bd K, with K satisfying (****). We will show that it is a cycle, or a straight line. (The proof of this statement for L is analogous. Actually we need this statement for K only in the compact case, when bd K is connected.)

We claim that K_i is symmetric w.r.t. the normal of K_i , at any $x \in K_i$. In fact, suppose the contrary. Then let $\widehat{x_1x_2} \neq K_i$ be a maximal open counterclockwise arc of K_i , symmetric w.r.t. its normal at x.

Possibly $x_1 = x_2$. However, this can occur only for K compact. Then $K_i = \operatorname{bd} K$ equals the closure of $\widehat{x_1x_2}$, hence it is symmetric w.r.t. the normal of K_i at x, as asserted. Hence further we may assume $x_1 \neq x_2$.

The normals of K_i at x_1 and x_2 are the one-sided normals of the closure of the

46

open arc $\widehat{x_1x_2} \subset K_i$ at x_1 and x_2 . Therefore these normals are symmetric images of each other w.r.t. the normal of K_i at x. Observe that both x_1, x_2 are points of local symmetry of K_i . Hence they are midpoints of sufficiently short counterclockwise open arcs $\widehat{x_{1,1}x_{1,2}}$ and $\widehat{x_{2,1}x_{2,2}}$ of K_i , of equal lengths, and symmetric w.r.t. the normals of K_i at x_1 and x_2 , resp. Then $(\widehat{x_{1,1}x_{1,2}}) \cup (\widehat{x_{1}x_{2,2}}) \cup (\widehat{x_{2,1}x_{2,2}})$ is an open arc of K_i , of midpoint x, symmetric w.r.t. the normal of K_i at x, strictly containing the maximal such open arc. This is a contradiction, proving symmetry of K_i w.r.t. its normal at any of its points $x \in K_i$.

We have that K_i is twice differentiable at some of its points x_0 . Let $x \in K_i$ be arbitrary. Then let x^* be the midpoint of (one of) the $\operatorname{arc}(s)$ $\widehat{x_0x} \subset K_i$. Then the image of x_0 w.r.t. the normal of K_i at x^* is x. Hence K_i is twice differentiable at x, and the curvatures of K_i at x_0 and x coincide. Hence K_i is of constant curvature. That is, it is either a cycle, or a straight line.

Then by the supporting circle hypothesis $\operatorname{bd} K = K_i$ is a circle, so K is a circle.

Lemma 8.4. Assume (****) with d = 2. Assume the hypotheses of Theorem 8, and (6) of Theorem 8. Let K and L be C^1 . Then case (2) in **3** of the proof of Theorem 8 leads to a contradiction.

Proof. Suppose, e.g., that some point $x_0 \in \operatorname{bd} K$ is not a point of local symmetry of bd K. (The proof for L is analogous.) Then consider for this x_0 and any $y \in \operatorname{bd} L$ the situation described in (1) to (4) in the proof of Lemma 7.1. By (8.1) bd (φK) and bd $[M \cap D(\varepsilon)]$ at φx_0 locally coincide. Therefore φx_0 is not a point of local symmetry of bd $[M \cap D(\varepsilon)]$ either.

Then conv $[M \cap D(\varepsilon)]$ is not symmetric w.r.t. the line spanned by D. Hence by (8.2) it is symmetric either w.r.t. o, or w.r.t. the orthogonal bisector of D. In both cases the symmetric image of φx_0 is ψy . Moreover, in both cases, the symmetric image of a short open arc of bd $[M \cap D(\varepsilon)]$, with midpoint φx_0 , is a short open arc of bd $[M \cap D(\varepsilon)]$, with midpoint ψy . Then, using (8.1), if bd (φK) at φx_0 is not twice differentiable, then by this symmetry bd (ψL) at any $\psi y \in bd(\psi L)$ is not twice differentiable either. This is a contradiction. Therefore $bd(\varphi K)$ at φx_0 is twice differentiable, and, by this symmetry, $bd(\psi L)$ is everywhere twice differentiable. Moreover, by this symmetry, the curvature of $bd(\varphi K)$ at φx_0 coincides with the curvature of $bd(\psi L)$ at any $\psi y \in bd(\psi L)$. Therefore ψL has constant curvature. Then, by the supporting circle hypothesis, L is a circle.

Then the fixed $\varphi x_0 \in \operatorname{bd} [M \cap D(\varepsilon)]$ and any $\psi y \in \operatorname{bd} [M \cap D(\varepsilon)]$ can become images of each other either w.r.t. o, or w.r.t. the orthogonal bisector of D. Moreover, sufficiently short open arcs of $\operatorname{bd} [M \cap D(\varepsilon)]$, of midpoints φx_0 and ψy , become then images of each other w.r.t. one of these symmetries. Then, on one hand, $\varphi x_0 \in \operatorname{bd} [M \cap D(\varepsilon)]$ is not a point of local symmetry of $M \cap D(\varepsilon)$. On the other hand, ψL and $M \cap D(\varepsilon)$ coincide in a neighbourhood of ψy . Hence, ψy is a point of local symmetry of $M \cap D(\varepsilon)$. This contradicts the fact that φx_0 and ψy , as well as the above short open arcs of $\operatorname{bd} [M \cap D(\varepsilon)]$, can be symmetric images of each other by such a symmetry of $M \cap D(\varepsilon)$. Thus we have obtained a contradiction.

Proof of Theorem 8, continuation. 4. Now the proof of Theorem 8 follows

from the earlier parts of the proof of Theorem 8, and from Lemmas 8.1-8.4. We only have to take into consideration that the supporting circle hypothesis excludes the paracycle and the hypercycle boundary components of K and L, so the connected boundary components of K and L are circles. Then, by the last paragraph of the proof of Lemma 1.7, K and L are circles.

Acknowledgements. The authors express their gratitude to I. Bárány, for carrying the problem, and bringing the two authors together; to B. Csikós for pointing out that Lemma 8.3 is valid for $K_i \subset X$ being only a C^1 1-manifold, which is also a closed subset. Namely if it can be translated in itself (this follows from symmetry w.r.t. the normal at any of its points) then it is a cycle or a straight line. W

References

- [1] T. Bonnesen, W. Fenchel, *Theorie der konvexen Körper*, Berichtigter Reprint (Theory of convex bodies, corrected reprint, in German), Springer, Berlin-New York, 1974.
- [2] J. Flachsmeyer, On the convergence of motions, In: Gen. Top. Rel. Modern Anal. Alg. V (Proc. 5th Prague Top. Symp., 1981), Sigma Ser. Pure Math. 3, Heldermann, Berlin, 1983, 183-188.
- R. High, Characterization of a disc, Solution to problem 1360 (posed by P. R. Scott), Math. Magazine 64 (1991), 353-354.
- [4] J. Jerónimo-Castro, E. Makai, Jr., Ball characterizations in spaces of constant curvature, Studia Sci. Math. Hungar. 55 (2018), 421-478.
- [5] J. Jerónimo-Castro, E. Makai, Jr., Ball characterizations in planes and spaces of constant curvature, I, submitted.
- [6] R. Schneider, Convex bodies: the Brunn-Minkowski theory; Convex bodies: the Brunn-Minkowski theory, Second expanded edition, Encyclopedia of Math. and its Appls., Vol. 44; 151, Cambridge Univ. Press, Cambridge, 1993; 2014.
- [7] Beltrami-Klein model, Wikipedia.
- [8] Möbius transformation, Ch. Subgroups of the Möbius group, Wikipedia.
- [9] Support function, Wikipedia.