Multiplicity of the ground state of the interaction system of a massless Bose field

Toshimitsu Takaesu ¹

Faculty of Science and Technology, Gunma University, Gunma, Japan

[Abstract] An interaction system of a massless Bose field is investigated. We suppose some conditions for the commutation relation of the interaction and annihilation operators. It is proven that if the ground state exists, its multiplicity is finite. Application to the Wigner-Weisskopf model is also considered.

Key words: Quantum fields, Fock spaces, Ground states, Spectral analysis.

1 Introduction and Main Result

In this paper, we consider an interaction system of a massless Bose field. Let \mathcal{K} be a Hilbert space over \mathbb{C} . The state space for the system is given by

$$\mathcal{H} = \mathcal{K} \otimes \mathcal{F}_{\mathbf{h}},\tag{1}$$

where \mathcal{F}_b denotes the boson Fock space over $L^2(\mathbb{R}^d)$. The free Hamiltonian is given by

$$H_0 = K \otimes \mathbb{1} + \mathbb{1} \otimes H_b, \tag{2}$$

where K is an operator on \mathcal{K} and $H_b = d\Gamma(\omega)$ is the second quantization of $\omega = \omega(\mathbf{k})$, $\mathbf{k} \in \mathbb{R}^d$. We suppose the following condition.

(H.1) *K* is self-adjoint and bounded from below.

$$(\mathrm{H.2})\ \omega \in C(\mathbb{R}), \inf_{\mathbf{k} \in \mathbb{R}} \omega(\mathbf{k}) = 0, \lim_{|\mathbf{k}| \to \infty} \omega(\mathbf{k}) = \infty.$$

The total Hamiltonian is defined by

$$H = H_0 + H_{\rm I}. \tag{3}$$

Suppose the condition below.

- (H.3) $H_{\rm I}$ is symmetric.
- (H.4) H is self-adjoint and bounded from below.

¹E-mail: t-takaesu@hotmail.co.jp

The annihilation operator is denoted by a(f) and the creation operator by $a^{\dagger}(f)$. It holds that $a(\alpha f + \beta g) = \alpha^* a(f) + \beta^* a(g)$, $\alpha, \beta \in \mathbb{C}$, and $a(f) = (a^{\dagger}(f))^*$ where X^* denotes the adjoint of operator X. The creation operators and annihilation operators satisfy the canonical commutation relations

$$[a(f), a^{\dagger}(g)] = \langle f, g \rangle, \tag{4}$$

$$[a(f), a(g)] = [a^{\dagger}(f), a^{\dagger}(g)] = 0.$$
 (5)

The main interest in this paper is to investigate the multiplicity of the ground state of H. The ground state of a system in quantum field theory is an important mathematical subject and has been studied by many researchers. One of the most difficult aspect of the analysis is that we have to investigate embedded eigenvalues. For the multiplicity of the ground states, an evaluation of the upper bound of an abstract system of a massless Bose field is obtained in [12]. The result in [12] can be applied to many important models, but for sufficiently small values of coupling constants. In this paper, we consider an evaluation of the multiplicity of the ground state, and it is independent of coupling constants.

Suppose the conditions below.

- (H.5) *K* has a compact resolvent.
- (H.6) There exist $L_0 \ge 0$ and $R_0 \ge 0$ such that for all $\Psi \in \mathcal{D}(H)$,

$$||H_0\Psi|| \le L_0||H\Psi|| + R_0||\Psi||.$$

We consider some conditions for the commutation relation of the interaction and annihilation operators. The weak commutator of X and Y is defined by

$$[X,Y]^0 \langle \Phi, \Psi \rangle = \langle X^* \Phi, Y \Psi \rangle - \langle Y^* \Phi, X \Psi \rangle.$$

Assume the following.

(H.7) There exists $G_I : \mathbb{R}^d \to \mathcal{L}(\mathcal{H})$ such that for all $\Phi, \Psi \in \mathcal{D}(H)$,

$$[H_{\mathrm{I}}, \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Psi \rangle = \int_{\mathbb{R}^{d}} f(\mathbf{k})^{*} \langle \Phi, G_{\mathrm{I}}(\mathbf{k}) \Psi \rangle d\mathbf{k}.$$

In addition, there exist $\zeta_l \ge 0$ and $\tau_l \ge 0$, l = 0, 1, such that for all $\Psi \in \mathcal{D}(H)$,

$$\int_{\mathbb{R}^d} \frac{1}{\omega(\mathbf{k})^{2l}} \|G_{\mathrm{I}}(\mathbf{k})\Psi\|^2 d\mathbf{k} \leq \zeta_l \|H\Psi\|^2 + \tau_l \|\Psi\|^2.$$

(H.8) It holds that $\omega \in C^1(\mathbb{R}^d \setminus I_0)$ where $I_0 = \{\mathbf{a}_l\}_{l=1}^{\infty}$, $\mathbf{a}_l \in \mathbb{R}^d$, $l \in \mathbb{N}$, and $\sup_{\mathbf{k} \in \mathbb{R} \setminus I_0} |\partial_{k_j} \omega(\mathbf{k})| < \infty$, $j = 1, \ldots, d$.

(H.9) There exist $\mu_0 \ge 0$ and $\nu_0 \ge 0$ such that for all $\Psi \in \mathcal{D}(H)$,

$$\int_{\mathbb{R}^d} \frac{\left|\partial_{k_j} \omega(\mathbf{k})\right|^2}{\omega(\mathbf{k})^4} \|G_{\mathrm{I}}(\mathbf{k})\Psi\|^2 d\mathbf{k} \leq \mu_0 \|H\Psi\|^2 + \nu_0 \|\Psi\|^2.$$

(H.10) $G_{\rm I}(\mathbf{k})$ is strongly differentiable for all $\mathbf{k} \in \mathbb{R}^d \setminus \{J_0\}$, where $J_0 = \{\mathbf{b}_l\}_{l=1}^{\infty}$, $\mathbf{b}_l \in \mathbb{R}^d$, $l \in \mathbb{N}$. There exist $\mu_j \geq 0$ and $\nu_j \geq 0$, $j = 1, \ldots d$, such that for all $\Psi \in \mathcal{D}(H)$,

$$\int_{\mathbb{R}^d} \frac{1}{\omega(\mathbf{k})^2} \|\partial_{k_j} G_{\mathbf{I}}(\mathbf{k}) \Psi\|^2 d\mathbf{k} \leq \mu_j \|H\Psi\|^2 + \nu_j \|\Psi\|^2.$$

The following is the main result.

Theorem 1.1 Suppose (H.1)-(H.10). Then if H has a ground state, its multiplicity is finite.

The point of Theorem 1.1 is that it is independent of coupling constants, as seen in the application to a concrete example in Section 3. The outline of the main theorem is as follows. First, we prepare for the pull-through formula. Second, we obtain the boson derivative bound [8]. Last, according to the argument of approximation of localized momentum [7], we show that if the multiplicity of the ground state is infinite, then it yields a contradiction. The strategy of the proof of the main theorem has been applied to a model of quantum electrodynamics (QED) in [15].

We briefly review other references to the multiplicity of the ground states. The uniqueness of the ground states in the massive case for the generalized spin-boson model [2] and the non-relativistic QED model [5] are investigated. An abstract Peron-Frobenius theory and its applications are considered in [14]. By the renormalization group analysis, the uniqueness of the ground state is proven in [9]. In addition, it is known that for the Hamiltonian which is represented by functional integration, the uniqueness of the ground state can be shown by probabilistic methods [13].

This paper is organized as follows. In section 2, we prove the main theorem. In section 3, the application of the main theorem is investigated.

2 Proof of Main Result

2.1 Preliminary

In the following, basic terms and properties of Fock pace are explained. For the detail, refer to e.g., [1]. The Fock vacuum is defined by $\Omega_0 = \{1,0,0...\} \in \mathcal{F}_b$. The finite particle subspace on a subspace \mathcal{M} is defined by

$$\mathfrak{F}_{\mathrm{b,fin}}(\mathfrak{M}) = \mathrm{L.h.}\left\{\Omega_0, \ a^{\dagger}(f_1) \dots a^{\dagger}(f_n)\Omega_0 \ \middle| \ f_j \in \mathfrak{M}, j = 1, \dots, n, n \in \mathbb{N}\right\}.$$

Let

$$\mathfrak{D}_{b} = \mathfrak{F}_{b, fin} \left(\mathfrak{D}(\boldsymbol{\omega}) \right).$$

It is seen that H_b is essentially self-adjoint on \mathcal{D}_b .

It holds that for all $f \in \mathcal{D}(\omega^{-1/2})$ and $\Psi \in \mathcal{D}(H_{\mathsf{h}}^{1/2})$,

$$||a(f)\Psi|| \le ||\frac{f}{\sqrt{\omega}}|| ||H_b^{1/2}\Psi||,$$
 (6)

$$||a^{\dagger}(f)\Psi|| \le ||\frac{f}{\sqrt{\omega}}|| ||H_{b}^{1/2}\Psi|| + ||f|||\Psi||.$$
 (7)

For all $f \in \mathcal{D}(\omega)$ and $\Psi \in \mathcal{D}_b$, it follows that

$$[H_{\mathsf{b}}, a(f)]\Psi = -a(\omega f)\Psi. \tag{8}$$

By (8), it is proven that for all $f \in \mathcal{D}(\omega) \cap \mathcal{D}(\omega^{-1/2})$ and $\Psi \in \mathcal{D}(H_h^{1/2})$,

$$[H_{b}, a(f)]^{0} \langle \Phi, \Psi \rangle = -\langle \Phi, a(\omega f) \Psi \rangle. \tag{9}$$

By (4) and (5), it is also shown that for all $f, h \in \mathcal{D}(\omega^{-1/2})$ and $\Psi \in \mathcal{D}(H_b^{1/2})$,

$$[a^{\dagger}(h), a(f)]^{0} \langle \Phi, \Psi \rangle = -\langle f, h \rangle \langle \Phi, \Psi \rangle, \tag{10}$$

$$[a(h), a(f)]^{0}\langle \Phi, \Psi \rangle = 0. \tag{11}$$

2.2 Pull-Through Formula and Derivative Bound

From this subsection onwards, we assume the existence of the ground state of H. Let Ω be the normalized ground state of H:

$$H\Omega = E_0(H)\Omega, \quad ||\Omega|| = 1,$$

where $E_0(H) = \inf \sigma(H)$. The kernel of an annihilation operator is defined by

$$(a(\mathbf{k})\Psi)^{(n)}(\mathbf{k}_1,\ldots,\mathbf{k}_n)=\sqrt{n+1}\Psi^{(n+1)}(\mathbf{k},\mathbf{k}_1,\ldots,\mathbf{k}_n).$$

Let $f \in \mathcal{D}(\omega^{-1/2})$. Then it holds that for all $\Phi \in \mathcal{F}_b$ and $\Psi \in \mathcal{D}(H_b^{1/2})$,

$$\langle \Phi, a(f)\Psi \rangle = \int_{\mathbb{R}^d} f(\mathbf{k})^* \langle \Phi, a(\mathbf{k})\Psi \rangle d\mathbf{k}.$$
 (12)

Let $N_b = d\Gamma(1)$. It holds that

$$\int_{\mathbb{R}^d} \langle a(\mathbf{k}) \Psi, a(\mathbf{k}) \Psi \rangle d\mathbf{k} = \|N_b^{1/2} \Psi\|^2, \quad \Psi \in \mathcal{D}(N_b^{1/2}).$$
 (13)

It also holds that

$$\int_{\mathbb{D}_d} \boldsymbol{\omega}(\mathbf{k}) \langle a(\mathbf{k}) \Psi, a(\mathbf{k}) \Psi \rangle d\mathbf{k} = \| H_b^{1/2} \Psi \|^2, \quad \Psi \in \mathcal{D}(H_b^{1/2}). \tag{14}$$

Proposition 2.1 (Pull-Through Formula) Suppose (H.1)-(H.4) and (H.7). Then,

$$1 \otimes a(\mathbf{k})\Omega = (H - E_0(H) + \omega(\mathbf{k}))^{-1}G_1(\mathbf{k})\Omega.$$

(Proof) Let $f \in \mathcal{D}(\omega^{-1/2}) \cap \mathcal{D}(\omega)$ and $\Phi \in \mathcal{D}(H)$. From (9), we have

$$[H, \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Omega \rangle = -\langle \Phi, \mathbb{1} \otimes a(\omega f) \Omega \rangle + [H_{\mathbf{I}}, \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Omega \rangle. \tag{15}$$

Since $H\Omega = E_0(H)\Omega$, we also see that

$$[H, \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Omega \rangle = \langle H\Phi, \mathbb{1} \otimes a(f)\Omega \rangle - E_{0}(H) \langle \mathbb{1} \otimes a^{\dagger}(f)\Phi, \Omega \rangle$$
$$= \langle (H - E_{0}(H))\Phi, \mathbb{1} \otimes a(f)\Omega \rangle. \tag{16}$$

By (15) and (16), we have

$$\langle (H - E_0(H))\Phi, \mathbb{1} \otimes a(f)\Omega \rangle + \langle \Phi, \mathbb{1} \otimes a(\omega f)\Omega \rangle = [H_I, \mathbb{1} \otimes a(f)]^0(\Phi, \Omega).$$

By (12) and (H.7),

$$\int_{\mathbb{R}^d} f(\mathbf{k})^* \langle (H - E_0(H) + \omega(\mathbf{k})) \Phi, \mathbb{1} \otimes a(\mathbf{k}) \Omega \rangle d\mathbf{k} = \int_{\mathbb{R}^d} f(\mathbf{k})^* \langle \Phi, G_{\mathbf{I}}(\mathbf{k}) \Omega \rangle d\mathbf{k}.$$

Let $\Phi = (H - E_0(H) + \omega(\mathbf{k})^{-1}\Xi, \Xi \in \mathcal{H}$. Then we have

$$\int_{\mathbb{R}^d} f(\mathbf{k})^* \langle \Xi, \mathbb{1} \otimes a(\mathbf{k}) \Omega \rangle d\mathbf{k} = \int_{\mathbb{R}^d} f(\mathbf{k})^* \langle \Xi, (H - E_0(H) + \omega(\mathbf{k}))^{-1} G_{\mathrm{I}}(\mathbf{k}) \Omega \rangle d\mathbf{k}. \quad (17)$$

Since $\mathcal{D}(\boldsymbol{\omega}^{-1/2}) \cap \mathcal{D}(\boldsymbol{\omega})$ is dense in $L^2(\mathbb{R}^d)$, (17) yields that

$$\langle \Xi, \mathbb{1} \otimes a(\mathbf{k}) \Omega \rangle = \langle \Xi, (H - E_0(H) + \omega(\mathbf{k}))^{-1} G_{\mathbf{I}}(\mathbf{k}) \Omega \rangle. \tag{18}$$

Since (18) holds for all $\Xi \in \mathcal{H}$, we have

$$1 \otimes a(\mathbf{k})\Omega = (H - E_0(H) + \omega(\mathbf{k}))^{-1}G_1(\mathbf{k})\Omega.$$

Thus, the assertion follows. \Box

Lemma 2.2 (Boson Number Bound) Suppose (H.1)-(H.4) and (H.7). Then,

$$\int_{\mathbb{D}^d} \| \mathbf{1} \otimes a(\mathbf{k}) \Omega \|^2 d\mathbf{k} \le c_1,$$

where $c_1 = E_0(H)^2 \zeta_1 + \tau_1$.

(**Proof**) By Proposition 2.1 and (H.5),

$$\begin{split} \int_{\mathbb{R}^d} \| \mathbf{1} \otimes a(\mathbf{k}) \Omega \|^2 d\mathbf{k} &\leq = \int_{\mathbb{R}^d} \left\| (H - E_0(H) + \omega(\mathbf{k}))^{-1} G_{\mathbf{I}}(\mathbf{k}) \Omega \right\|^2 d\mathbf{k} \\ &\leq \int_{\mathbb{R}^d} \frac{1}{\omega(\mathbf{k})^2} \| G_{\mathbf{I}}(\mathbf{k}) \Omega \|^2 d\mathbf{k} \\ &\leq \zeta_1 \| H \Omega \|^2 + \tau_1 \| \Omega \|^2. \end{split}$$

Since $H\Omega = E_0(H)\Omega$ and $||\Omega|| = 1$, the proof is obtained. \square

Using Proposition 2.1, we prove the following.

Proposition 2.3 (Boson Derivative Bound) Suppose (H.1)-(H.4) and (H.7)-(H.10). Then,

$$\int_{\mathbb{R}^d} \|\partial_{k_j} (\mathbb{1} \otimes a(\mathbf{k})) \Omega\|^2 d\mathbf{k} \leq \xi_j,$$

where $\xi_j = 2((\mu_0 + \mu_j)E_0(H)^2 + v_0 + v_j).$

(**Proof**) By Proposition 2.1, (H.8) and the strong differentiability of the resolvent, we have for all $\mathbf{k} \in \mathbb{R}^d \setminus (I_0 \cup J_0)$,

$$\begin{split} \partial_{k_j}(\mathbb{1} \otimes a(\mathbf{k}))\Omega &= \partial_{k_j} \left((H - E_0(H) + \omega(\mathbf{k}))^{-1} G_{\mathbf{I}}(\mathbf{k}) \Omega \right) \\ &= - (H - E_0(H) + \omega(\mathbf{k}))^{-2} \left(\partial_{k_j} \omega(\mathbf{k}) \right) G_{\mathbf{I}}(\mathbf{k}) \Omega \\ &+ (H - E_0(H) + \omega(\mathbf{k}))^{-1} \partial_{k_j} G_{\mathbf{I}}(\mathbf{k}) \Omega. \end{split}$$

By (H.9) and (H.10),

$$\int_{\mathbb{R}^{d}} \|\partial_{k_{j}}(\mathbb{1} \otimes a(\mathbf{k}))\Omega\|^{2} d\mathbf{k}$$

$$\leq 2 \left(\int_{\mathbb{R}^{d}} \frac{|\partial_{k_{j}}\omega(\mathbf{k})|^{2}}{\omega(\mathbf{k})^{4}} \|G_{\mathbf{I}}(\mathbf{k})\Omega\|^{2} d\mathbf{k} + \int_{\mathbb{R}^{d}} \frac{1}{\omega(\mathbf{k})^{2}} \|\partial_{k_{j}}G_{\mathbf{I}}(\mathbf{k})\Omega\|^{2} d\mathbf{k} \right)$$

$$\leq 2 \left(\mu_{0} \|H\Omega\|^{2} + \nu_{0} \|\Omega\|^{2} + \mu_{j} \|H\Omega\|^{2} + \nu_{j} \|\Omega\|^{2} \right)$$

$$\leq 2 \left((\mu_{0} + \mu_{j})E_{0}(H)^{2} + \nu_{0} + \nu_{j} \right).$$

Thus the assertion follows. \Box

Let *X* be an operator in $L^2(\mathbb{R}^d)$. We set

$$\Gamma(X) = \bigoplus_{n=0}^{\infty} (\otimes^n X).$$

Lemma 2.4 Let J be a self-adjoint operator on $L^2(\mathbb{R}^d)$. Suppose that J is non-negative and $||J|| \le 1$. Then, it follows that for all $\Psi \in \mathcal{D}(N_b)$,

$$\|(\mathbb{1} - \Gamma(J))\Psi\|^2 \le (\Psi, d\Gamma(1 - J)\Psi). \tag{19}$$

(**Proof**) Since $||J|| \le 1$, it holds that $\mathcal{D}(d\Gamma(J)) = \mathcal{D}(N_b)$. Let $\Psi = {\Psi^{(n)}}_{n=0}^{\infty} \in \mathcal{D}(N_b)$. Let $0 \le \lambda_i \le 1$, $j = 1, \dots, n$, $n \in \mathbb{N}$. Then we see that

$$(1 - \lambda_1 \dots \lambda_n)^2 \le (1 - \lambda_1 \dots \lambda_n) \le \sum_{j=1}^n (1 - \lambda_j).$$
(20)

By (20) and the spectral decomposition theorem, we have for each n-particle components,

$$\|(\mathbb{1} - \Gamma(J))\Psi^{(n)}\|^2 = (\Psi^{(n)}, (\mathbb{1} - \Gamma(J))^2\Psi^{(n)}) \le (\Psi^{(n)}, d\Gamma(1 - J)\Psi^{(n)}). \tag{21}$$

From (21), the lemma follows. \Box

We consider a localization estimate of momentum, which is investigated in [6], [7] (see also [11];Section 4.1.3). Let $\chi \in C_0^{\infty}(\mathbb{R}^d)$. We suppose that $0 \le \chi \le 1$ and $\chi(\mathbf{k}) = 1$ for $|\mathbf{k}| \le 1$. Let $\chi_R(\mathbf{k}) = \chi(\frac{\mathbf{k}}{R})$.

Proposition 2.5 Suppose (H.1)-(H.4) and (H.7)-(H.10). Then,

$$\|1\!\!1 \otimes (1\!\!1 - \Gamma(\chi_R(-i\nabla_{\mathbf{k}})))\Omega\|^2 \le c_1^{1/2} \left(\frac{d+1}{R^4}c_1 + \frac{d+1}{R^2}\sum_{j=1}^d \xi_j\right)^{1/2}.$$

(**Proof**) Let $\hat{\mathbf{k}} = -i\nabla_{\mathbf{k}}$. From Lemma 2.2 and 2.4, we have

$$\|\mathbf{1} \otimes (\mathbf{1} - \Gamma(\boldsymbol{\chi}_{R}(\hat{\mathbf{k}}))\Omega)\|^{2}$$

$$\leq (\Omega, \mathbf{1} \otimes d\Gamma(1 - \boldsymbol{\chi}_{R}(\hat{\mathbf{k}}))\Omega)$$

$$\leq \int_{\mathbb{R}^{d}} (\mathbf{1} \otimes a(\mathbf{k})\Omega, \mathbf{1} \otimes (1 - \boldsymbol{\chi}_{R}(\hat{\mathbf{k}}))\mathbf{1} \otimes a(\mathbf{k})\Omega) d\mathbf{k}$$

$$\leq \left(\int_{\mathbb{R}^{d}} \|\mathbf{1} \otimes a(\mathbf{k})\Omega\|^{2} d\mathbf{k}\right)^{1/2} \left(\int_{\mathbb{R}^{d}} \|(1 - \boldsymbol{\chi}_{R}(\hat{\mathbf{k}}))\mathbf{1} \otimes a(\mathbf{k})\Omega\|^{2} d\mathbf{k}\right)^{1/2}$$

$$= c_{1}^{1/2} \left(\int_{\mathbb{R}^{d}} \|(1 - \boldsymbol{\chi}_{R}(\hat{\mathbf{k}}))\mathbf{1} \otimes a(\mathbf{k})\Omega\|^{2} d\mathbf{k}\right)^{1/2}.$$
(22)

Then, we see that

$$\int_{\mathbb{R}^{d}} \|(1 - \chi_{R}(\hat{\mathbf{k}})) \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$= \int_{\mathbb{R}^{d}} \|\frac{(1 - \chi_{R}(\hat{\mathbf{k}}))}{1 + \hat{\mathbf{k}}^{2}} (1 + \hat{\mathbf{k}}^{2}) \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$\leq (d+1) \int_{\mathbb{R}^{d}} \|\frac{(1 - \chi_{R}(\hat{\mathbf{k}}))}{1 + \hat{\mathbf{k}}^{2}} \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$\leq (d+1) \int_{\mathbb{R}^{d}} \|\frac{(1 - \chi_{R}(\hat{\mathbf{k}}))}{1 + \hat{\mathbf{k}}^{2}} \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$\leq (d+1) \int_{\mathbb{R}^{d}} \|\frac{(1 - \chi_{R}(\hat{\mathbf{k}}))}{1 + \hat{\mathbf{k}}^{2}} \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$+ (d+1) \sum_{j=1}^{d} \int_{\mathbb{R}^{d}} \left\| \frac{(1 - \chi_{R}(\hat{\mathbf{k}}))\hat{k}_{j}}{1 + \hat{\mathbf{k}}^{2}} \partial_{k_{j}} (\mathbb{1} \otimes a(\mathbf{k})) \Omega \right\|^{2} d\mathbf{k}.$$
 (24)

We see that

$$\sup_{\mathbf{k} \in \mathbb{R}^d} \left| \frac{(1 - \chi_R(\mathbf{k}))}{1 + \mathbf{k}^2} \right| \le \frac{1}{R^2},\tag{25}$$

and,

$$\sup_{\mathbf{k}\in\mathbb{R}^d} \left| \frac{(1-\chi_R(\mathbf{k}))k_j}{1+\mathbf{k}^2} \right| \le \frac{1}{R}, \quad j=1,\dots d.$$
 (26)

By the spectral decomposition theorem, we apply (25) and (26) to (24), and then,

$$\int_{\mathbb{R}^{d}} \|(1 - \chi_{R}(\hat{\mathbf{k}})) \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$\leq \frac{d+1}{R^{4}} \int_{\mathbb{R}^{d}} \|\mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k} + \frac{d+1}{R^{2}} \sum_{j=1}^{d} \int_{\mathbb{R}^{d}} \|\partial_{k_{j}} \mathbf{1} \otimes a(\mathbf{k}) \Omega\|^{2} d\mathbf{k}$$

$$\leq \frac{d+1}{R^{4}} c_{1} + \frac{d+1}{R^{2}} \sum_{j=1}^{d} \xi_{j}.$$
(27)

Here we used Lemma 2.2 and Proposition 2.3 in the last line. From (22) and (27), the lemma follows. \Box

From Proposition 2.5, the next corollary follows.

Corollary 2.6 Suppose (H.1)-(H.4) and (H.7)-(H.10). Then,

$$\|1\!\!1 \otimes (1\!\!1 - \Gamma(\chi_R(-i\nabla_\mathbf{k})))\Omega\| \le \frac{c_2}{\sqrt{R}}, \quad R \ge 1,$$
(28)

where
$$c_2 = (d+1)^{1/4} (c_1^{1/2} + c_1^{1/4} (\sum_{j=1}^d \xi_j)^{1/4}).$$

(Proof of Theorem 1.1)

Suppose that dim ker $(H - E_0(H)) = \infty$. Then there exists a sequence of the normalized

ground states $\{\Omega_j\}_{j=1}^{\infty}$ of H. Here note that $\{\Omega_j\}_{j=1}^{\infty}$ is an orthonormal system of H, and hence, w- $\lim_{i\to\infty}\Omega_j=0$. Let $P_{\rm b}$ be the spectral projection of $N_{\rm b}$. It is seen that

$$\begin{split} 1\!\!1\otimes 1\!\!1 &= 1\!\!1\otimes P_{\mathrm{b}}([0,n]) + 1\!\!1\otimes P_{\mathrm{b}}([n+1,\infty)) \\ &= 1\!\!1\otimes P_{\mathrm{b}}([0,n])\Gamma(\chi_R(-i\nabla_{\mathbf{k}})) \\ &+ 1\!\!1\otimes \left(P_{\mathrm{b}}([0,n])(1\!\!1 - \Gamma(\chi_R(-i\nabla_{\mathbf{k}})))\right) + 1\!\!1\left(\otimes P_{\mathrm{b}}([n+1,\infty))\right). \end{split}$$

Then we have

$$\left\| \mathbb{1} \otimes \left(P_{b}([0,n]) \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}})) \right) \Omega_{j} \right\|$$

$$\geq 1 - \left\| \mathbb{1} \otimes \left(P_{b}([0,n](\mathbb{1} - \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}}))) \right) \Omega_{j} \right\| - \left\| \mathbb{1} \otimes P_{b}([n+1,\infty)) \Omega_{j} \right\|$$

$$\geq 1 - \left\| \mathbb{1} \otimes \left(\mathbb{1} - \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}})) \right) \Omega_{j} \right\| - \left\| \mathbb{1} \otimes P_{b}([n+1,\infty)) \Omega_{j} \right\|. \tag{29}$$

From Corollary 2.6, we have

$$\left\| \mathbb{1} \otimes \left(\mathbb{1} - \Gamma(\chi_R(-i\nabla_{\mathbf{k}})) \right) \Omega_j \right\| \le \frac{c_2}{\sqrt{R}}. \tag{30}$$

By Lemma 2.2 and (13), we have

$$\| \mathbb{1} \otimes P_{b}([n+1,\infty))\Omega_{j} \| \leq \frac{1}{\sqrt{n+1}} \| \mathbb{1} \otimes N_{b}^{1/2}\Omega_{j} \| \leq \frac{\sqrt{c_{1}}}{\sqrt{n+1}}.$$
 (31)

Applying (30) and (31) to (29),

$$\left\| \mathbb{1} \otimes \left(P_{\mathbf{b}}([0,n]) \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}})) \right) \Omega_{j} \right\| \ge 1 - \frac{c_{2}}{\sqrt{R}} - \frac{\sqrt{c_{1}}}{\sqrt{n+1}}.$$
 (32)

The left-hand side of (32) is estimated as

$$\left\| \mathbb{1} \otimes \left(P_{b}([0,n]) \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}})) \right) \Omega_{j} \right\|^{2}$$

$$\leq \left\| (H_{0} + i) \Omega_{j} \right\| \left\| (H_{0} - i)^{-1} \left(\mathbb{1} \otimes (P_{b}([0,n]) \Gamma(\chi_{R}(-i\nabla_{\mathbf{k}}))) \right) \Omega_{j} \right\|.$$
(33)

By (H.6),

$$||(H_{0}+i)\Omega_{j}|| \leq ||H_{0}\Omega_{j}|| + ||\Omega_{j}|| \leq L_{0} ||H\Omega_{j}|| + R_{0} ||\Omega_{j}|| + ||\Omega_{j}|| \leq L_{0}E_{0}(H) + R_{0} + 1.$$
(34)

Applying (33) and (34) to (32), and taking R > 0 and $n \in \mathbb{N}$ such that $1 - \frac{c_2}{\sqrt{R}} - \frac{\sqrt{c_1}}{\sqrt{n+1}} > 0$, we have

$$\left\| (H_0 - i)^{-1} \left(1 \otimes (P_b([0, n]) \Gamma(\chi_R(-i\nabla_{\mathbf{k}}))) \right) \Omega_j \right\| \ge \frac{1}{c_3} \left(1 - \frac{c_2}{\sqrt{R}} - \frac{\sqrt{c_1}}{\sqrt{n+1}} \right)^2.$$
 (35)

where $c_3 = L_0 E_0(H) + R_0 + 1$. It is seen that

$$(H_{0}-i)^{-1} \left(\mathbb{1} \otimes (P_{b}([0,n])\Gamma(\chi_{R}(-i\nabla_{\mathbf{k}}))) \right)$$

$$= \left((H_{0}-i)^{-1} ((K-i) \otimes (H_{b}-i)) \right)$$

$$\times \left((K-i)^{-1} \otimes (P_{b}([0,n])(H_{b}-i)^{-1}\Gamma(\chi_{R}(-i\nabla_{\mathbf{k}}))) \right). \tag{36}$$

Here, we see that $(H_0-i)^{-1}((K-i)\otimes(H_b-i))$ is bounded. We also see that $(K-i)^{-1}\otimes(P_b([0,n])(H_b-i)^{-1}\Gamma(\chi_R(-i\nabla_{\mathbf{k}})))$ is compact. Hence, by (36) we see that $(H_0-i)^{-1}(\mathbb{1}\otimes(P_b([0,n])\Gamma(\chi_R(-i\nabla_{\mathbf{k}}))))$ is compact. Since w- $\lim_{j\to\infty}\Omega_j=0$, it holds that s- $\lim_{j\to\infty}(H_0-i)^{-1}(\mathbb{1}\otimes(P_b([0,n])\Gamma(\chi_R(-i\nabla_{\mathbf{k}}))))\Omega_j=0$. On the other hand, (35) yields that $\lim_{j\to\infty}\left\|(H_0+1)^{-1}(\mathbb{1}\otimes(P_b([0,n])\Gamma(\chi_R(-i\nabla_{\mathbf{k}}))))\Omega_j\right\|>0$, and this is a contradiction. Hence dim ker $(H-E_0(H))<\infty$. \square

3 Applications

3.1 Wigner-Weisskopf Model

The Wigner-Weisskopf model Model describes a two-level atom interacting with a scalar Bose field. It is a concrete example of the models investigated in [2] and [7]. For the properties of the ground state, refer to e.g., [3], [4], [10]. The total Hamiltonian acting in $\mathbb{C}^2 \otimes \mathcal{F}_b$ is defined by

$$H_{\alpha} = \zeta c^* c \otimes \mathbb{1} + \mathbb{1} \otimes H_{\mathsf{b}} + H_{\mathsf{I}}(\alpha), \tag{37}$$

where

$$H_{\rm I}(\alpha) = \alpha \left(c^* \otimes a(\rho_{\rm I}) + c \otimes a(\rho_{\rm I})^* \right). \tag{38}$$

Here $\varsigma \in \mathbb{R} \setminus \{0\}$ is a constant, $c = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, $H_b = d\Gamma(\omega)$ with $\omega(\mathbf{k}) = |\mathbf{k}|$, $\mathbf{k} \in \mathbb{R}^d$, and $\rho_I \in L^2(\mathbb{R}^d)$. It is seen that c and c^* has the following fermionic property;

$$\{c, c^*\} = 1$$
,
 $\{c, c\} = \{c^*, c^*\} = 0$.

We see that $||c|| \le 1$ and $||c^*|| \le 1$. From the definition of H_{α} , it is seen that (H.1)-(H.3) are satisfied. For the self-adjointness of H_{α} , we suppose the condition below.

(A.1)
$$\int_{\mathbb{R}^d} \frac{|\rho_{\rm I}(\mathbf{k})|^2}{\omega(\mathbf{k})} d\mathbf{k} < \infty$$
.

By (6) and (7), it holds that for all $\Psi \in \mathcal{D}(\mathbb{1} \otimes H_b^{1/2})$,

$$||H_{\mathrm{I}}(\alpha)\Psi|| = |\alpha| ||(c^* \otimes a(\rho_{\mathrm{I}}) + c \otimes a(\rho_{\mathrm{I}})^*)\Psi||$$

$$\leq |\alpha| (||\mathbb{1} \otimes a(\rho_{\mathrm{I}})\Psi|| + ||\mathbb{1} \otimes a(\rho_{\mathrm{I}})^*\Psi||)$$

$$\leq 2|\alpha| ||\frac{\rho_{\mathrm{I}}}{\sqrt{\omega}}|| ||\mathbb{1} \otimes H_{\mathrm{b}}^{1/2}\Psi|| + |\alpha||\rho_{\mathrm{I}}|| ||\Psi||.$$
(39)

By the spectral decomposition theorem, we see that for all $\varepsilon > 0$,

$$\|\mathbb{1} \otimes H_{b}^{1/2} \Psi\| \leq \varepsilon \|\mathbb{1} \otimes H_{b} \Psi\| + \frac{1}{2\varepsilon} \|\Psi\|$$

$$\leq \varepsilon \|H_{0} \Psi\| + \frac{1}{2\varepsilon} \|\Psi\|. \tag{40}$$

Here we used $c^*c \ge 1$ in the last line. From (39) and (40), we have

$$||H_{\mathbf{I}}(\alpha)\Psi|| \le c_{\mathbf{I}}(\alpha, \varepsilon)||H_{\mathbf{0}}\Psi|| + d_{\mathbf{I}}(\alpha, \varepsilon)||\Psi||, \tag{41}$$

where $c_{\rm I}(\alpha,\varepsilon)=2\varepsilon|\alpha|\|\frac{\rho_{\rm I}}{\sqrt{\omega}}\|$ and $d_{\rm I}(\alpha,\varepsilon)=\frac{|\alpha|}{\varepsilon}\|\frac{\rho_{\rm I}}{\sqrt{\omega}}\|+|\alpha|\|\rho_{\rm I}\|$. Taking $\varepsilon>0$ such as $\varepsilon<\frac{1}{2\alpha\|\frac{\rho_{\rm I}}{\sqrt{\omega}}\|}$, then H_{α} is self-adjoint for all $\alpha\in\mathbb{R}$ by the Kato-Rellich theorem. Hence (H.4) is satisfied. Since ςc^*c is a matrix, (H.5) follows. From (41), it is seen that for all $\Psi\in\mathcal{D}(H_0)$,

$$||H_0\Psi|| \le ||H_\alpha\Psi|| + ||H_I(\alpha)\Psi||$$

$$< ||H_\alpha\Psi|| + c_I(\alpha,\varepsilon)||H_0\Psi|| + d_I(\alpha,\varepsilon)||\Psi||.$$

Let $\varepsilon > 0$ such as $1 - c_{\rm I}(\alpha, \varepsilon) > 0$, then we have

$$||H_0\Psi|| \le L_{\mathrm{I}}(\varepsilon,\alpha)||H_\alpha\Psi|| + R_{\mathrm{I}}(\varepsilon,\alpha)||\Psi||, \tag{42}$$

where $L_{\rm I}(\varepsilon,\alpha) = \frac{1}{1-c_{\rm I}(\alpha,\varepsilon)}$ and $R_{\rm I}(\varepsilon,\alpha) = \frac{d_{\rm I}(\alpha,\varepsilon)}{1-c_{\rm I}(\alpha,\varepsilon)}$. From (42), (H.6) is satisfied. For the existence of the ground state, we introduce a stronger condition than (A.1).

(A.2)
$$\int_{\mathbb{R}^d} \frac{|\rho_{\rm I}(\mathbf{k})|^2}{\omega(\mathbf{k})^2} d\mathbf{k} < \infty$$
.

Then, it is proven that H_{α} has the ground state for all values of $\alpha \in \mathbb{R}$, ([7]; Theorem 1). Next, let us consider the multiplicity of the ground state of H_{α} . We assume the conditions below, which include a stronger condition than (A.2).

(B.1)
$$\int_{\mathbb{R}^d} \frac{|\rho_{\rm I}(\mathbf{k})|^2}{\omega(\mathbf{k})^4} d\mathbf{k} < \infty$$
.

(B.2)
$$\rho_{\mathrm{I}} \in C^{1}(\mathbb{R}^{d})$$
 and $\int_{\mathbb{R}^{d}} \frac{|\partial_{k_{j}} \rho_{\mathrm{I}}(\mathbf{k})|^{2}}{\omega(\mathbf{k})^{2}} d\mathbf{k} < \infty, j = 1, \dots, d.$

Let us check the conditions (H.7)-(H.10). From (10) and (11), we have

$$[H_{\mathbf{I}}(\alpha), \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Psi \rangle$$

$$= \alpha \left([c^{*} \otimes a(\rho_{\mathbf{I}}), \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Psi \rangle + [c \otimes a(\rho_{\mathbf{I}})^{*}, \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Psi \rangle \right)$$

$$= -\alpha \langle f, \rho_{\mathbf{I}} \rangle \langle \Phi, c \otimes \mathbb{1} \Psi \rangle. \tag{43}$$

From (43), we have

$$[H_{\mathbf{I}}(\alpha), \mathbb{1} \otimes a(f)]^{0} \langle \Phi, \Psi \rangle = \int_{\mathbb{D}^{d}} f(\mathbf{k})^{*} \langle \Phi, Q_{\alpha}(\mathbf{k}) \Psi \rangle d\mathbf{k}, \tag{44}$$

where $Q_{\alpha}(\mathbf{k}) = -\alpha \rho_{\rm I}(\mathbf{k})c \otimes \mathbb{1}$. Then, it holds that

$$\int_{\mathbb{R}^d} \frac{1}{\omega(\mathbf{k})^{2l}} \|Q_{\alpha}(\mathbf{k})\Psi\|^2 d\mathbf{k} \le \alpha^2 \left(\int_{\mathbb{R}^d} \frac{|\rho_{\mathbf{I}}(\mathbf{k})|^2}{\omega(\mathbf{k})^{2l}} d\mathbf{k} \right) \|\Psi\|, \quad l = 0, 1, \tag{45}$$

and hence (H.7) holds. Since $\omega(\mathbf{k}) = |\mathbf{k}|$, we see that $\sup_{\mathbb{R}^d \setminus \{\mathbf{0}\}} |\partial k_j \omega(\mathbf{k})| = \sup_{\mathbb{R}^d \setminus \{\mathbf{0}\}} \frac{|k_j|}{\omega(\mathbf{k})} \le 1$, and (H.8) is satisfied. It is seen that

$$\int_{\mathbb{R}^d} \frac{|\partial k_j \omega(\mathbf{k})|}{\omega(\mathbf{k})^4} \|Q_{\alpha}(\mathbf{k})\Psi\|^2 d\mathbf{k} \le \alpha^2 \left(\int_{\mathbb{R}^d} \frac{|\rho_{\mathbf{I}}(\mathbf{k})|^2}{\omega(\mathbf{k})^4} d\mathbf{k} \right) \|\Psi\|, \tag{46}$$

and

$$\int_{\mathbb{R}^d} \frac{1}{\omega(\mathbf{k})^2} \|\partial_{k_j} Q_{\alpha}(\mathbf{k}) \Psi\|^2 d\mathbf{k} \le \alpha^2 \left(\int_{\mathbb{R}^d} \frac{|\partial_{k_j} \rho_{\mathrm{I}}(\mathbf{k})|^2}{\omega(\mathbf{k})^4} d\mathbf{k} \right) \|\Psi\|. \tag{47}$$

Hence (H.9) and (H.10) are satisfied. Thus the following theorem holds.

Theorem 3.1 *Suppose* (B.1) and (B.2). Then, dim ker $(H_{\alpha} - E_0(H_{\alpha})) < \infty$ for all values of $\alpha \in \mathbb{R}$.

3.2 Application to Other Models

The generalized spin-boson model [2] includes various models. Among them, we consider a lattice spin system interacting with phonon ([2];Example 1.3). Let Λ be a finite set of the d-dimensional square lattice \mathbb{Z}^d . Let $\mathbf{S} = (S^{(n)})_{n=1}^N$ be a N component spin. Here $S^{(n)}$ acts on \mathbb{C}^s with $s \in \mathbb{N}$. Let $\mathcal{K}_{\Lambda} = \bigotimes_{i \in \Lambda} \mathcal{K}_i$ with $\mathcal{K}_i = \mathbb{C}^s$. Let \mathbf{S}_i be the spin at site $i \in \Lambda$. The Hamiltonian acting in $\mathcal{K}_{\Lambda} \otimes \mathcal{F}_b$, is given by

$$H_{\Lambda}(\beta) = \left(-\sum_{(i,j)\subset\Lambda} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j\right) \otimes \mathbb{1} + \mathbb{1} \otimes H_{b} + \beta \sum_{j\in\Lambda} \sum_{n=1}^{N} S_j^{(n)} \otimes \left(a(\boldsymbol{\rho}_j^{(n)}) + a^{\dagger}(\boldsymbol{\rho}_j^{(n)})\right)$$

where $J_{ij} \in \mathbb{R}$, $i, j \in \Lambda$, and $\rho_j^{(n)} \in L^2(\mathbb{R}^d)$, $j \in \Lambda$, j = 1, ..., N. Without going into details, Theorem 1.1 can be applied to $H_{\Lambda}(\beta)$ in the same way.

Acknowledgements

This work is supported by JSPS KAKENHI 20K03625.

Conflict of Interest

The author states that there is no conflict of interest.

References

- [1] A. Arai, Analysis on Fock spaces and mathematical theory of quantum fields: An introduction to mathematical analysis of quantum fields, World Scientific Publishing, 2018.
- [2] A. Arai and M. Hirokawa, On the existence and uniqueness of ground states of a generalized spin-boson model, *J. Funct. Anal.*, **151**, 455-503, (1997).
- [3] A. Arai and M. Hirokawa, Ground states of a general class of quantum field Hamiltonians, *Rev. Math. Phys.*, **12**, 1085-1135, (2000).
- [4] A. Arai and M. Hirokawa, Stability of ground states in sectors and its application to the Wigner-Weisskopf Model, *Rev. Math. Phys.*, **13**, 513-528, (2001).
- [5] V. Bach, J. Fröhlich and I. M. Sigal, Quantum electrodynamics of confined nonrelativistic particles, *Adv. Math.*, **137**, 299-395, (1998).
- [6] J. Dereziński and C. Gérard, Asymptotic completeness in quantum field theory: Massive Pauli-Fierz Hamiltonian, *Rev. Math. Phys.*, **11**, 383-450, (1999).
- [7] C. Gérard, On the existence of ground states for massless Pauli-Fierz hamiltonians, *Ann. H. Poincaré*, **1**, 443-459, (2000).
- [8] M. Griesemer, E. Lieb and M. Loss, Ground states in non-relativistic quantum electrodynamics, *Invent. Math.*, **145**, 557-595, (2001).
- [9] D. Hasler and I. Herbst, Uniqueness of the ground state in Feshbach renormalization analysis, *Lett. Math. Phys.*, **100**, 171-180, (2012).
- [10] M. Hirokawa, Remarks on the ground state energy of the spin-boson model. An Application of the Wigner-Weisskopf model, *Rev. Math. Phys.*, **13**, 221-251, (2001).
- [11] F. Hiroshima, Ground states of quantum field models: Perturbation of embedded eigenvalues, Springer, (2019).
- [12] F. Hiroshima, Multiplicity of ground states in quantum field models; Applications of asymptotic fields, *J. Funct. Anal.*, **224**, 431-470, (2005).
- [13] F. Hiroshima and J. Lörinczi, Feynman-Kac-Type theorems and Gibbs measures on path space: Applications in rigorous quantum field Theory, De Gruyter, (2020).

- [14] T. Miyao, Nondegeneracy of ground states in the nonrelativistic quantum field theory, *J. Operator Theory*, **64**, 207-241, (2010).
- [15] T. Takaesu, Ground states of quantum electrodynamics with cutoffs, *J. Mah. Phys.*, **59**, 072301, (2018).