A Construction of Quantum Stabilizer Codes from Classical Codes and Butson Hadamard Matrices

Bülent Saraç, Damla Acar

Abstract

In this paper, we give a constructive proof to show that if there exist a classical linear code $\mathscr{C} \subseteq \mathbb{F}_q^n$ of dimension k and a classical linear code $\mathscr{D} \subseteq \mathbb{F}_{q^k}^m$ of dimension s, where q is a power of a prime number p, then there exists an $[nm, ks, \delta]_q$ quantum stabilizer code with δ determined by \mathscr{C} and \mathscr{D} by identifying the stabilizer group of the code. In the construction, we use a particular type of Butson Hadamard matrices equivalent to multiple Kronecker products of the Fourier matrix of order p. We also consider the same construction of a quantum code for a general normalized Butson Hadamard matrix and search for a condition for the quantum code to be a stabilizer code.

1 Introduction and Preliminaries

Quantum error-correcting codes have experienced rapid growth since their inception by Shor in [10]. One of the most important classes of these codes is quantum stabilizer codes, first introduced by Gottesman [4] and Calderbank et al. [3]. Stabilizer codes have been extensively studied due to their relatively simple encoding algorithms and their structural advantages, which allow for connections to classical error-correcting codes for analysis.

In this paper, we propose a method for constructing quantum error-correcting codes that generalize Shor's code and the quantum codes discussed in [11]. Our approach encompasses both binary and non-binary schemes. We use quantum digits (qudits for short) over \mathbb{F}_q as a unit of quantum information, where \mathbb{F}_q denotes the finite field with q elements. The states of qudits are simply vectors in \mathbb{C}^q . This vector space is equipped with the standard inner product, with respect to which there is an orthonormal basis, whose elements are denoted $|x\rangle$, where x runs through the elements of \mathbb{F}_q . Now the state of a system of n qudits are represented by vectors in $(\mathbb{C}^q)^{\otimes n}$. The set $\{|x_1\rangle \otimes \cdots \otimes |x_n\rangle : x_1, \ldots, x_n \in \mathbb{F}_q\}$ forms an orthonormal basis for the Hilbert space $(\mathbb{C}^q)^{\otimes n}$. For the sake of simplicity, we denote the basis vector $|x_1\rangle \otimes \cdots \otimes |x_n\rangle$ as $|x_1 \dots x_n\rangle$. Thus every vector in $(\mathbb{C}^q)^{\otimes n}$ is a linear combination of the vectors $|\mathbf{x}\rangle$, where $\mathbf{x} \in \mathbb{F}_q^n$.

A quantum code of length n is a nonzero subspace of $(\mathbb{C}^q)^{\otimes n}$. We denote a q-ary quantum code which encodes k qudits of information into n-qudits as $[n, k]_q$.

In order to provide information about the types of errors a quantum code can detect or correct, we first define the unitary operators on \mathbb{C}^q by

$$X(a) |x\rangle = |x+a\rangle, \ Z(b) |x\rangle = \omega^{\operatorname{tr}(bx)} |x\rangle$$

where $a, b, x \in \mathbb{F}_q$, tr denotes the trace function from \mathbb{F}_q to its prime field \mathbb{F}_p , and ω is a primitive *p*-th root of unity. Note that the operators X(a) and Z(b), for nonzero *a* and *b*, coincide with the usual bit flip and phase flip errors, respectively, on (binary) quantum bits (or qubits). Let $\mathcal{E} = \{X(a)Z(b) \mid a, b \in \mathbb{F}_q\}$. Then \mathcal{E} forms a basis for the vector space of linear operators on \mathbb{C}^q . Further, we observe that the following properties are satisfied, which make \mathcal{E} a nice error basis on \mathbb{C}^q (see [8]):

- (NE1) \mathcal{E} contains the identity operator.
- (NE2) $X(a)Z(b)X(a')Z(b') = \omega^{\operatorname{tr}(ba')}X(a+a')Z(b+b')$ for every $a, a', b, b' \in \mathbb{F}_q$. Therefore the product of two elements of \mathcal{E} is a scalar multiple of another element of \mathcal{E} .
- (NE3) If M, N are distinct elements of \mathcal{E} then $\operatorname{Tr}(M^{\dagger}N) = 0$.

Remark 1. Let p = 2. If we take $\omega = -1$ for the binary case it results in omission of complex phases. Thus we use ω as a primitive 4-th root of unity; more precisely, the imaginary unit *i*.

One can extend \mathcal{E} to a suitable error basis on n qudits, as follows. For $\mathbf{a} = (a_1, \ldots, a_n) \in \mathbb{F}_q^n$, we write $X(\mathbf{a}) = X(a_1) \otimes \cdots \otimes X(a_n)$ and $Z(\mathbf{a}) = Z(a_1) \otimes \cdots \otimes Z(a_n)$ for the tensor products of n error operators. Then $\mathcal{E}_n = \{X(\mathbf{a})Z(\mathbf{b}) \mid \mathbf{a}, \mathbf{b} \in \mathbb{F}_q^n\}$ forms a nice error basis on $(\mathbb{C}^q)^{\otimes n}$ (see [7]).

Binary stabilizer codes were first introduced by Gottesman in [4]. They have proved to form an important class of quantum codes because they provide an algebraic approach for error correction. Later, the stabilizer formation was extended to non-binary quantum error-correcting codes in [1] and investigated in detail in [6].

Let $\mathcal{P}_n = \{\omega^c X(\mathbf{a}) Z(\mathbf{b}) : \mathbf{a}, \mathbf{b} \in \mathbb{F}_{q^k}, c \in \mathbb{F}_p\}$. Then \mathcal{P}_n is a finite group of order pq^{2n} for p > 2. Note that when p = 2, we take $\omega = i$ and $c \in \{0, 1, 2, 3\}$; hence $|\mathcal{P}_n| = 4q^{2n}$. \mathcal{P}_n is called the error group associated with \mathcal{E}_n (see [6]).

Given a quantum code Q of length n, one may think of the subgroup

$$\operatorname{Stab}(Q) = \{ E \in \mathcal{P}_n : E\mathbf{v} = \mathbf{v} \text{ for all } \mathbf{v} \in Q \}$$

of \mathcal{P}_n , called the *stabilizer group* of Q. Note that $\operatorname{Stab}(Q)$ is an abelian subgroup of \mathcal{P}_n with $\operatorname{Stab}(Q) \cap \mathcal{Z}(\mathcal{P}_n) = \{I\}$. Conversely, given an abelian subgroup \mathcal{S} of \mathcal{P}_n with $\mathcal{S} \cap \mathcal{Z}(\mathcal{P}_n) = \{I\}$, one can define the quantum code

$$\operatorname{Fix}(\mathcal{S}) = \{ \mathbf{v} \in (\mathbb{C}^q)^{\otimes n} : E\mathbf{v} = \mathbf{v} \text{ for all } E \in \mathcal{S} \},\$$

the joint eigenspace of S associated with the eigenvalue 1. Observe that if Q is a quantum code, then $Q \subseteq \operatorname{Fix}(\operatorname{Stab}(Q))$. If the equality holds, then we call Q a stabilizer code. Equivalently, a quantum code Q is a stabilizer code if and only if there exists an abelian subgroup S of \mathcal{P}_n with $S \cap \mathcal{Z}(\mathcal{P}_n) = \{I\}$ such that $Q = \operatorname{Fix}(S)$. If Q is a q-ary stabilizer code of length n, then $S = \operatorname{Stab}(Q)$ is a finite abelian subgroup of \mathcal{P}_n of order $q^n/\dim(Q)$. It follows that if S has a minimal generating set (of r elements, say), then $|S| = q^r$ since S is abelian and every non-central element of \mathcal{P}_n has order q, which gives that Q is an $[n, n - r]_q$ -code.

For a subgroup S of \mathcal{P}_n , we write $C_{\mathcal{P}_n}(S)$ to denote the centralizer of S in \mathcal{P}_n , namely $C_{\mathcal{P}_n}(S) = \{E \in \mathcal{P}_n : ES = SE$ for every $S \in S\}$. Also, the subgroup of \mathcal{P}_n generated by S and $\mathcal{Z}(\mathcal{P}_n)$ is $\mathcal{SZ}(\mathcal{P}_n)$. The following proposition, proved in [7], shows that the error detection capability of a stabilizer code is closely related to the subgroups $C_{\mathcal{P}_n}(S)$ and $\mathcal{SZ}(\mathcal{P}_n)$.

Proposition 1.1. Suppose that $S \leq \mathcal{P}_n$ is the stabilizer group of a stabilizer code Q with dim Q > 1. An error E in \mathcal{P}_n is detectable by the quantum code Q if and only if either E is an element of $SZ(\mathcal{P}_n)$ or E does not belong to the centralizer $C_{\mathcal{P}_n}(S)$.

The weight of an error operator $E = \omega^c E_1 \otimes \cdots \otimes E_n \in \mathcal{P}_n$ is defined by $\operatorname{wt}(E) = \#\{i \mid E_i \neq I\}$, that is the number of tensor factors that are nonidentity. We say that a quantum code Q has minimal distance d if d is a positive integer such that every error in \mathcal{P}_n of weight less than d can be detected by Qwhereas there is an error in \mathcal{P}_n of weight d which cannot be detected by Q. Thus, it follows easily from Proposition 1.1 that the minimum distance of a stabilizer code Q with the stabilizer group S is given by

$$d(Q) = \begin{cases} \min\{\operatorname{wt}(E) \mid E \in C_{\mathcal{P}_n}(\mathcal{S}) \setminus \mathcal{S}\}, & \text{if } S \subsetneq C_{\mathcal{P}_n}(\mathcal{S}) \\ \min\{\operatorname{wt}(E) \mid E \in \mathcal{S} \setminus \{I\}\}, & \text{if } \mathcal{S} = C_{\mathcal{P}_n}(\mathcal{S}) \end{cases}$$

It is known that a code of minimum distance d can correct all errors of weight at most |(d-1)/2|.

With any error of type $\omega^c X(\mathbf{a})Z(\mathbf{b})$, we associate the vector $(\mathbf{a} \mid \mathbf{b})$ in \mathbb{F}_q^{2n} , where $\mathbf{a}, \mathbf{b} \in \mathbb{F}_q^n$, and define a map ψ by

$$\psi: \mathcal{P}_n \to \mathbb{F}_q^{2n}, \ \omega^c X(\mathbf{a}) Z(\mathbf{b}) \longmapsto (\mathbf{a} \mid \mathbf{b}).$$
 (1.1)

Note that ψ is a group homomorphism from the multiplicative group \mathcal{P}_n onto the additive group \mathbb{F}_q^{2n} , through which some multiplicative properties in \mathcal{P}_n translates into additive properties in \mathbb{F}_q^{2n} . For instance, if $E = \omega^c X(\mathbf{a})Z(\mathbf{b})$ and $E' = \omega^{c'}X(\mathbf{a}')Z(\mathbf{b}')$ are the elements of \mathcal{P}_n , then the commutator $[E, E'] = E^{-1}E'^{-1}EE'$ is equal to $\omega^{\mathrm{tr}(\mathbf{b}.\mathbf{a}'-\mathbf{b}'.\mathbf{a})}I$, and hence the errors E and E' commute if and only if $\mathrm{tr}(\mathbf{b}.\mathbf{a}'-\mathbf{b}'.\mathbf{a}) = 0$. This motivates the definition of a function $\langle \cdot, , \cdot \rangle_s : \mathbb{F}_q^{2n} \times \mathbb{F}_q^{2n} \to \mathbb{F}_p$ by

$$\langle (\mathbf{a} \mid \mathbf{b}), (\mathbf{a'} \mid \mathbf{b'}) \rangle_s := \operatorname{tr}(\mathbf{b}.\mathbf{a'} - \mathbf{b'}.\mathbf{a}),$$

which is an inner product on \mathbb{F}_q^{2n} as an \mathbb{F}_p -space, called the *trace symplec*tic inner product. It follows that the errors $E = \omega^c X(\mathbf{a})Z(\mathbf{b})$ and $E' = \omega^{c'}X(\mathbf{a'})Z(\mathbf{b'})$ in \mathcal{P}_n commute if and only if the vectors $(\mathbf{a} \mid \mathbf{b})$ and $(\mathbf{a'} \mid \mathbf{b'})$ in \mathbb{F}_q^{2n} are orthogonal according to the trace symplectic inner product. We sometimes express the orthonality of the vectors $(\mathbf{a} \mid \mathbf{b})$ and $(\mathbf{a'} \mid \mathbf{b'})$ by writing $(\mathbf{a} \mid \mathbf{b}) \perp_s (\mathbf{a'} \mid \mathbf{b'})$.

Let \overline{S} be the image of a subgroup S of \mathcal{P}_n under ψ . Then \overline{S} is an \mathbb{F}_p -subspace of \mathbb{F}_q^{2n} . The symplectic dual of \overline{S} (denoted \overline{S}^{\perp_s}) consists of the elements of \mathbb{F}_q^{2n} that are orthogonal to every element in \overline{S} . Note that ψ maps $C_{\mathcal{P}_n}(S)$ onto \overline{S}^{\perp_s} . In view of Proposition 1.1, we define the *symplectic weight* of a vector ($\mathbf{a} \mid \mathbf{b}$) in \mathbb{F}_q^{2n} (denoted swt($\mathbf{a} \mid \mathbf{b}$)) to be the weight of the error $X(\mathbf{a})Z(\mathbf{b})$ in \mathcal{P}_n . Equivalently, swt($\mathbf{a} \mid \mathbf{b}$) is the number of the *i*'s such that $(a_i, b_i) \neq (0, 0)$ for $\mathbf{a} = (a_1, \ldots, a_n)$ and $\mathbf{b} = (b_1, \ldots, b_n)$.

A Butson-Hadamard (BH) matrix H of order n is an $n \times n$ matrix whose entries are primitive roots of unity such that $HH^{\dagger} = nI_n$, where H^{\dagger} is the complex conjugate transpose of H and I_n is the identity matrix of order n. Thus the rows of a Butson Hadamard matrix are pairwise orthogonal with respect to the Hermitian inner product. If the entries of H are k-th roots of unity then we say that H is a BH(n, k) matrix. Butson Hadamard matrices were introduced by Butson in [2] and has been studied in various aspects. Two Butson Hadamard matrices are said to be equivalent if they are of the same type, say BH(n, k), and one can be obtained from the other by permuting rows (or columns) or multiplying rows (or columns) by a k-th root of unity. It is readily checked that every BH matrix is equivalent to a BH matrix whose first row and first column consist of 1's, which is called normalized. One of the simplest examples of a normalized BH matrix is the Fourier matrix $[\zeta_n^{(i-1)(j-1)}]_{i,j=1}^n \in BH(n, n)$, where ζ_n is a primitive n-th rooth of unity. Moreover, one can construct a BH matrix $[\zeta_n^{\mathbf{xy}^T}]_{\mathbf{x},\mathbf{y}\in\mathbb{F}_p^n} \in BH(p^n, p)$, which is, indeed, the n-fold Kronecker product of the Fourier matrix of order p.

The organization of our paper is as follows. In the second section, we give our general method for constructing a quantum code using p-ary and q-ary classical codes and a $BH(q^k, p)$ matrix, where p is a prime and q is a power of p. In the third section, we look for a BH matrix for which our construction gives a stabilizer quantum code. In particular, we show that if the BH matrix used in the construction has a particular form such that it is equivalent to a Kronecker product of the Fourier matrix of order p, then the resulting quantum code is a stabilizer code. In the next section, we search for a converse, and consider a certain q-ary classical code in the construction. In particular, we prove that if the resulting quantum code is a stabilizer code, then the BH matrix used in the construction must be equivalent to a Kronecker product of the Fourier matrix of order p. In Section 5, we give an algorithm of error correction for quantum stabilizer codes obtained by our construction as stabilizer codes. The last section is given as appendix devoted to provide a proof of the fact that the BH matrices considered in Section 3 and Theorem 4.2 are equivalent to a Knocker product of Fourier matrices.

2 The General Construction

Let $\mathscr{C} \subseteq \mathbb{F}_q^n$ be a *q*-ary $[n, k, d_1]$ classical linear code, where $1 \leq k < r$. We shall use the codewords of \mathscr{C} to create quantum states in $(\mathbb{C}^q)^{\otimes n}$, as follows: Let $\{f_{\lambda} : \lambda \in \mathbb{F}_{q^k}\}$ be a set of functions from \mathscr{C} into \mathbb{F}_p and let

$$\phi_{\lambda} := \frac{1}{\sqrt{q^k}} \sum_{\mathbf{c} \in \mathscr{C}} \omega^{f_{\lambda}(\mathbf{c})} | \mathbf{c} \rangle \tag{2.1}$$

for all $\lambda \in \mathbb{F}_{q^k}$, where $\omega = e^{2\pi i/p}$. Note that the ϕ_{λ} 's form an orthonormal basis for a subspace of $(\mathbb{C}^q)^{\otimes n}$ if and only if the $q^k \times q^k$ matrix

$$H = [\omega^{f_{\lambda}(\mathbf{c})}]_{\lambda \in \mathbb{F}_{q^k}, \mathbf{c} \in \mathscr{C}}, \qquad (2.2)$$

with rows indexed by the elements of \mathbb{F}_{q^k} and columns indexed by the elements of \mathscr{C} , both written in a fixed order, forms a $BH(q^k, p)$ matrix.

Let $\mathscr{D} \subseteq \mathbb{F}_{a^k}^m$ be a classical linear code of dimension s and define

$$\Phi_{\Lambda} := \phi_{\lambda_1} \otimes \ldots \otimes \phi_{\lambda_m} \in (\mathbb{C}^q)^{\otimes nm}$$
(2.3)

for all $\Lambda = (\lambda_1, \ldots, \lambda_m) \in \mathscr{D}$. We form a quantum code, denoted $Q_H(\mathscr{C}, \mathscr{D})$, of length nm to be the subspace of $(\mathbb{C}^q)^{\otimes nm}$ spanned by Φ_{Λ} for all $\Lambda \in \mathscr{D}$, i.e.,

$$Q_H(\mathscr{C},\mathscr{D}) := \operatorname{span}\{\Phi_\Lambda : \Lambda \in \mathscr{D}\}.$$
(2.4)

Note that $Q_H(\mathscr{C}, \mathscr{D})$ is a q^{ks} -dimensional subspace of $(\mathbb{C}^q)^{\otimes nm}$, namely, $Q_H(\mathscr{C}, \mathscr{D})$ is an $[\![nm, ks, \delta]\!]_q$ quantum code where δ is the minimum distance of $Q_H(\mathscr{C}, \mathscr{D})$. On the other hand, there exists a one-to-one correspondance $\nu : \mathbb{F}_q^{ks} \to \mathscr{D}$, and so one can set the logical state $|\mathbf{a}\rangle_L$ as $\Phi_{\nu(\mathbf{a})}$ for each $\mathbf{a} \in \mathbb{F}_q^{ks}$. Then $Q_H(\mathscr{C}, \mathscr{D})$ is a quantum code of length nm that encodes ks logical q-states.

Example 1. If p = q = 2, $\mathscr{C} = \mathscr{D} = \{000, 111\} \subseteq \mathbb{F}_2^3$, and

$$H = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

in (2.2), then the construction of the quantum code $Q_H(\mathscr{C}, \mathscr{D})$ as in (2.4) coincides with the well-known Shor's 9-qubit code.

Example 2. Let p = q = 3, $\mathscr{C} = \mathscr{D} = \{000, 111, 222\}$, and

$$H = \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}$$

in (2.2), where $w = e^{2\pi i/3}$. Then the quantum code obtained as in (2.4) is the same as the nine-quirt error correcting code considered in Sect. V of [9].

More generally, if $\mathscr{C} = \mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_q\} \subseteq \mathbb{F}_q^m$, the one-dimensional linear code over \mathbb{F}_{q^k} spanned by $(1, 1, \ldots, 1) \in \mathbb{F}_q^m$, and H is any BH(q, p)

matrix, then the quantum code $Q_H(\mathscr{C}, \mathscr{D})$ in (2.4) is among the $[m^2, 1, m]_q$ quantum error-correcting codes studied in [5]. In view of the next section, we see that if the matrix H is chosen to be a normalized Butson-Hadamard matrix of Fourier type, then $Q_H(\mathscr{C}, \mathscr{D})$ turns out to be a stabilizer code.

Remark 2. Let \mathscr{D} be a q^k -ary linear code of length m. If there exists a positive integer i with $1 \leq i \leq m$ such that for every codeword Λ in \mathscr{D} , the i-th coordinate of Λ is zero, then we can project \mathscr{D} onto a q^k -ary linear code \mathscr{D}' of length m-1by deleting the i-th coordinate of each codeword of \mathscr{D} , where the minimum distance remains unaltered, and use \mathscr{D}' instead of \mathscr{D} in the construction of $Q_H(\mathscr{C}, \mathscr{D})$. Thus, throughout this note, we shall assume that the code \mathscr{D} in the above construction, is non-trivial (i.e., neither 0 nor $\mathbb{F}_{q^k}^m$) and that for every integer $1 \leq i \leq m$, there exists a codeword in \mathscr{D} whose i-th coordinate is equal to 1.

Proposition 2.1. Let \mathscr{C} be a q-ary linear code of dimension k and length n, and let \mathscr{D} be a q^k -ary linear code of dimension s and length m (where 0 < s < m). Let H and H' be $BH(q^k, p)$ matrices. If $Q_H(\mathscr{C}, \mathscr{D}) = Q_{H'}(\mathscr{C}, \mathscr{D})$, then H and H' are row-equivalent BH matrices. The converse also holds when $\mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_{q^k}\} \subseteq \mathbb{F}_{q^k}^m$ is the one-dimensional linear code over \mathbb{F}_{q^k} .

Proof. Assume that $Q_H(\mathscr{C}, \mathscr{D}) = Q_{H'}(\mathscr{C}, \mathscr{D})$. Given $\lambda \in \mathbb{F}_{q^k}$, we use the notations ϕ_{λ} or ϕ'_{λ} accordingly the coefficients of the quantum states in (2.1) come from H or H'. Similarly, we write Φ'_{Λ} to mean the tensor products of states of the form ϕ'_{λ} in (2.3).

Since the sets $\{\phi_{\lambda} : \lambda \in \mathbb{F}_{q^k}\}$ and $\{\phi'_{\lambda} : \lambda \in \mathbb{F}_{q^k}\}$ are orthonormal, they are independent; hence we have

$$\operatorname{span}\{\phi_{\lambda}:\lambda\in\mathbb{F}_{q^{k}}\}=\operatorname{span}\{|\mathbf{c}\rangle:\mathbf{c}\in\mathscr{C}\}=\operatorname{span}\{\phi_{\lambda}':\lambda\in\mathbb{F}_{q^{k}}\}.$$

It follows that there exist $b_{\lambda\mu} \in \mathbb{C}$ $(\lambda, \mu \in \mathbb{F}_{q^k})$ such that

$$\phi_{\lambda} = \sum_{\mu \in \mathbb{F}_{q^k}} b_{\lambda\mu} \phi'_{\mu} \tag{2.5}$$

for all $\lambda \in \mathbb{F}_{q^k}$. On the other hand, since $Q_H(\mathscr{C}, \mathscr{D}) = Q_{H'}(\mathscr{C}, \mathscr{D})$, there exist $a_{\Lambda M} \in \mathbb{C}$ $(\Lambda, M \in \mathscr{D})$ such that $\Phi_{\Lambda} = \sum_{M \in \mathscr{D}} a_{\Lambda M} \Phi'_M$ for all $\Lambda \in \mathscr{D}$. Rewriting Φ_{Λ} for $\Lambda = (\lambda_1, \ldots, \lambda_m) \in \mathscr{D}$ as

$$\Phi_{\Lambda} = \left(\sum_{\mu \in \mathbb{F}_{q^k}} b_{\lambda_1 \mu} \phi'_{\mu}\right) \otimes \cdots \otimes \left(\sum_{\mu \in \mathbb{F}_{q^k}} b_{\lambda_1 \mu} \phi'_{\mu}\right)$$
$$= \sum_{(\mu_1, \dots, \mu_m) \in \mathbb{F}_{q^k}^m} \left(\prod_{i=1}^m b_{\lambda_i \mu_i}\right) (\phi'_{\mu_1} \otimes \cdots \otimes \phi'_{\mu_m}),$$

we see that for all $\Lambda = (\lambda_1, \ldots, \lambda_m) \in \mathscr{D}, \prod_{i=1}^m b_{\lambda_i \mu_i} = a_{\Lambda M}$ if $M = (\mu_1, \ldots, \mu_m) \in \mathscr{D}$ and $\prod_{i=1}^m b_{\lambda_i \mu_i} = 0$ if $M = (\mu_1, \ldots, \mu_m) \notin \mathscr{D}$.

Considering the equation (2.5), we see that there exists a function σ : $\mathbb{F}_{q^k} \to \mathbb{F}_{q^k}$ such that $b_{\lambda\sigma(\lambda)} \neq 0$. Since we assume that \mathscr{D} is non-trivial, at least one of the standard basis element of $\mathbb{F}_{q^k}^m$ does not belong to \mathscr{D} . Without loss of generality, we assume that $(1, 0, \ldots, 0) \notin \mathscr{D}$. Let $\lambda \in \mathbb{F}_{q^k}$. By our assumption on \mathscr{D} (see Remark 2), there exists $(\lambda_1, \ldots, \lambda_m) \in \mathscr{D}$ with $\lambda_1 = \lambda$. Then we must have $(\sigma(\lambda_1), \ldots, \sigma(\lambda_m)) \in \mathscr{D}$. Let $\alpha \in \mathbb{F}_{q^k} \setminus \{0\}$. Then $(\alpha + \sigma(\lambda_1), \sigma(\lambda_2), \ldots, \sigma(\lambda_m)) \notin \mathscr{D}$. This gives that $b_{\lambda\nu}.b_{\lambda_2\sigma(\lambda_2)}\ldots b_{\lambda_m\sigma(\lambda_m)} = 0$, where $\nu = \alpha + \sigma(\lambda)$; hence $b_{\lambda\nu} = 0$. It follows that $b_{\lambda\mu} = 0$ for all $\mu \in \mathbb{F}_{q^k}$ with $\mu \neq \sigma(\lambda)$. Therefore, $\phi_{\lambda} = b_{\lambda\sigma(\lambda)}\phi'_{\sigma(\lambda)}$ for all $\lambda \in \mathbb{F}_{q^k}$, and so σ is a permutation on \mathbb{F}_{q^k} . It is now straightforward to check that $b_{\lambda\sigma(\lambda)}$ is a power of ω for all $\lambda \in \mathbb{F}_{q^k}$. This completes the proof of the first assertion since we also have $H = [b_{\lambda\mu}]H'$. Now the second assertion follows since in case $\mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_q\} \subseteq \mathbb{F}_q^m$, the row-equivalence of H and H' implies that Φ_{Λ} is a constant multiple of Φ'_{Λ} for every $\Lambda \in \mathscr{D}$.

3 A Quantum Stabilizer Code

In this section, we use our general construction of quantum codes described in the preceding section to produce quantum stabilizer codes by choosing a particular set $\{f_{\lambda} : \lambda \in \mathbb{F}_{q^k}\}$ of functions from \mathscr{C} into \mathbb{F}_p in the formation of the ϕ_{λ} 's in (2.1). We start with two lemmas.

Lemma 3.1. Let $\mathscr{C} \subseteq \mathbb{F}_q^n$ be a linear code over \mathbb{F}_q and let $\mathbf{u} \in \mathbb{F}_q^n$. Then $\operatorname{tr}_{q/p}(\mathbf{u}.\mathbf{c}) = 0$ for all $\mathbf{c} \in \mathscr{C}$ if and only if $\mathbf{u} \in \mathscr{C}^{\perp}$.

Proof. It is enough to prove the "only if" part of the statement. So, suppose that $\operatorname{tr}_{q/p}(\mathbf{u}.\mathbf{c}) = 0$ for all $\mathbf{c} \in \mathscr{C}$. Choose an arbitrary nonzero element $\lambda \in \mathbb{F}_q$. Then $\mathscr{C} = \{\lambda \mathbf{c} : \mathbf{c} \in \mathscr{C}\}$, and so $\operatorname{tr}_{q/p}(\lambda(\mathbf{u}.\mathbf{c})) = 0$ for all $\mathbf{c} \in \mathscr{C}$. Since $\lambda \in \mathbb{F}_q$ is arbitrary, this implies that $\mathbf{u}.\mathbf{c} = 0$ for all $\mathbf{c} \in \mathscr{C}$; hence $\mathbf{u} \in \mathscr{C}^{\perp}$.

Lemma 3.2. Let $\mathscr{C} \subseteq \mathbb{F}_q^n$ be a linear code over \mathbb{F}_q and let $\phi = \sum_{\mathbf{c} \in \mathscr{C}} \omega^{f(\mathbf{c})} | \mathbf{c} \rangle \in (\mathbb{C}^q)^{\otimes n}$, where $f : \mathscr{C} \to \mathbb{F}_p$ is a function. Let $\mathbf{u} \in \mathbb{F}_q^n$. Then $Z(\mathbf{u})$ stabilizes ϕ if and only if $\mathbf{u} \in \mathscr{C}^{\perp}$.

Proof. This is clear by Lemma 3.1 since $Z(\mathbf{u})\phi = \sum_{\mathbf{c}\in\mathscr{C}} \omega^{f(c)+\operatorname{tr}_{q/p}(\mathbf{u}.\mathbf{c})} |\mathbf{c}\rangle$.

Let $\mathscr{C} \subseteq \mathbb{F}_q^n$ be a classical linear code over \mathbb{F}_q of dimension k, where $q = p^r$ and $1 \leq k < n$, and $\mathscr{D} \subseteq \mathbb{F}_{q^k}^m$ be a classical linear code over \mathbb{F}_{q^k} of dimension s with $1 \leq s < m$. Note that \mathscr{C} is also a vector space over \mathbb{F}_p by restriction of scalars. Let $\mathcal{L}(\mathscr{C}, \mathbb{F}_p)$ be the \mathbb{F}_p -dual of \mathscr{C} . That is, $\mathcal{L}(\mathscr{C}, \mathbb{F}_p)$ is the set of all \mathbb{F}_p -linear transformations from \mathscr{C} to \mathbb{F}_p . Then $\mathcal{L}(\mathscr{C}, \mathbb{F}_p)$ is a vector space over \mathbb{F}_p of dimension rk; in other words, $\mathcal{L}(\mathscr{C}, \mathbb{F}_p)$ contains q^k linear transformations of \mathbb{F}_p -spaces. Also, there exists an \mathbb{F}_p -space isomorphism $\kappa : \mathbb{F}_{q^k} \to \mathcal{L}(\mathscr{C}, \mathbb{F}_p)$. We set $f_{\lambda} = \kappa(\lambda)$ for each $\lambda \in \mathbb{F}_{q^k}$, form the matrix $H = [\omega^{f_{\lambda}(\mathbf{c})}]_{\lambda \in \mathbb{F}_{q^k}, \mathbf{c} \in \mathscr{C}}$ which is necessarily a BH matrix, and define $Q_H(\mathscr{C}, \mathscr{D})$ as in (2.4). As shown in Corollary A.6, H is equivalent to the rk-fold Kronecker product of the Fourier matrix of order p. Note that the above setting of f_{λ} 's yields that $f_{\lambda_1} + f_{\lambda_2} = f_{\lambda_1+\lambda_2}$ and $f_{c\lambda} = cf_{\lambda}$ for all $\lambda, \lambda_1, \lambda_2 \in \mathbb{F}_{q^k}$ and $c \in \mathbb{F}_p$. Given any $\Lambda = (\lambda_1, \ldots, \lambda_m) \in \mathscr{D}$, define a mapping $F_{\Lambda} : \mathscr{C}^m \to \mathbb{F}_p$ by $F_{\Lambda}(\mathbf{c}_1, \ldots, \mathbf{c}_m) = f_{\lambda_1}(\mathbf{c}_1) + \cdots + f_{\lambda_m}(\mathbf{c}_m)$ for all $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \mathscr{C}^m$. Then F_{Λ} is a linear transformation of \mathbb{F}_p -spaces. Moreover, by definition of f_{λ} 's above, the set $\{F_{\Lambda} : \Lambda \in \mathscr{D}\}$ is a vector space over \mathbb{F}_p in a natural way since $F_{\Lambda_1} + F_{\Lambda_2} = F_{\Lambda_1+\Lambda_2}$ and $h.F_{\Lambda} = F_{h\Lambda}$ for all $h \in \mathbb{F}_p$ and $\Lambda, \Lambda_1, \Lambda_2 \in \mathscr{D}$.

For $\mathbf{x} \in \mathbb{F}_q^n$, define the mapping $\rho_{\mathbf{x}} : \mathscr{C} \to \mathbb{F}_p$ by $\rho_{\mathbf{x}}(\mathbf{c}) = \operatorname{tr}_{q/p}(\mathbf{c}.\mathbf{x})$. Then $\rho_{\mathbf{x}}$ is a linear transformation of \mathbb{F}_p -spaces, and given $\mathbf{x}, \mathbf{y} \in \mathbb{F}_q^n$, $\rho_{\mathbf{x}} = \rho_{\mathbf{y}}$ if and only if $\mathbf{x} - \mathbf{y} \in \mathscr{C}^{\perp}$ by Lemma 3.1. Thus we sometimes write $\rho_{\overline{\mathbf{x}}}$ for $\rho_{\mathbf{x}}$, where $\overline{\mathbf{x}}$ denotes the image of \mathbf{x} under the canonical projection $\mathbb{F}_q^n \to \mathbb{F}_q^n / \mathscr{C}^{\perp}$. Note that for each $\lambda \in \mathbb{F}_q$, there corresponds \mathbf{x}_{λ} such that $f_{\lambda} = \rho_{\overline{\mathbf{x}}_{\lambda}}$. This correspondence yields an \mathbb{F}_p -space isomorphism $\Theta : \mathbb{F}_{q^k} \to \mathbb{F}_q^n / \mathscr{C}^{\perp}$, where $\Theta(\lambda) = \overline{\mathbf{x}}_{\lambda}$ for which $f_{\lambda} = \rho_{\overline{\mathbf{x}}_{\lambda}}$. Define

$$\mathscr{D}^{\Theta} := \{ (\mathbf{x}_{\lambda_1}, \dots, \mathbf{x}_{\lambda_m}) : (\lambda_1, \dots, \lambda_m) \in \mathscr{D} \text{ and } \mathbf{x}_{\lambda_i} \in \Theta(\lambda_i) \text{ for all } 1 \le i \le m \}.$$

That is,

$$\mathscr{D}^{\Theta} = \bigcup_{(\lambda_1, \dots, \lambda_m) \in \mathscr{D}} \Theta(\lambda_1) \times \dots \times \Theta(\lambda_m).$$

Then, clearly, \mathscr{D}^{Θ} is an additive code over \mathbb{F}_q . Note that all the vectors in $(\mathscr{C}^{\perp})^{(m)}$ are elements of \mathscr{D}^{Θ} corresponding to the zero codeword in \mathscr{D} . Thus, $(\mathscr{C}^{\perp})^{(m)} \subseteq \mathscr{D}^{\Theta}$.

Now we are ready to state and prove the main theorem of this section.

Theorem 3.3. With the above notation, $Q_H(\mathscr{C}, \mathscr{D})$ is an $[nm, ks, \delta]_q$ quantum stabilizer code and $\delta = \min\{d(\mathscr{C}), \ell\}$, where $\ell = \min\{\operatorname{wt}(\mathbf{X}) : \mathbf{X} \in \mathscr{D}^{\Theta} \setminus (\mathscr{C}^{\perp})^{(m)}\}$. Moreover, the stabilizer group \mathcal{S} of $Q_H(\mathscr{C}, \mathscr{D})$ consists of the errors of the form $X(\mathbf{c}_1, \ldots, \mathbf{c}_m)Z(\mathbf{d}_1, \ldots, \mathbf{d}_m)$, where $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)$ and $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$, and the centralizer $C_{\mathcal{P}_n}(\mathcal{S})$ of \mathcal{S} consists of the errors of the form $\omega^c X(\mathbf{u}_1, \ldots, \mathbf{u}_m)Z(\mathbf{v}_1, \ldots, \mathbf{v}_m)$, where $c \in \mathbb{F}_p$, $\mathbf{u}_1, \ldots, \mathbf{u}_m \in \mathscr{C}$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_m) \in \mathscr{D}^{\Theta}$.

Proof. If $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$, then $Z(\mathbf{d}_1, \ldots, \mathbf{d}_m) \in \mathscr{C}^{\perp}$ by Lemma 3.2. On the other hand, given $\mathbf{e} \in \mathscr{C}$ and $\lambda \in \mathbb{F}_{q^k}$, we have

$$X(\mathbf{e})\phi_{\lambda} = \sum_{\mathbf{c}\in\mathscr{C}} \omega^{f_{\lambda}(\mathbf{c})} |\mathbf{c} + \mathbf{e}\rangle = \sum_{\mathbf{c}\in\mathscr{C}} \omega^{f_{\lambda}(\mathbf{c}-\mathbf{e})} |\mathbf{c}\rangle = \omega^{-f_{\lambda}(e)}\phi_{\lambda},$$

and so

$$X(\mathbf{c}_1,\ldots,\mathbf{c}_m)\Phi_{\Lambda}=\omega^{F_{\Lambda}(\mathbf{c}_1,\ldots,\mathbf{c}_m)}\Phi_{\Lambda}$$

for all $\mathbf{c}_1, \ldots, \mathbf{c}_m \in \mathscr{C}$ and $\Lambda \in \mathscr{D}$. Thus, given codewords $\mathbf{c}_1, \ldots, \mathbf{c}_m$ of \mathscr{C} , $X(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \mathscr{S}$ if and only if $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)$. Therefore, \mathscr{S} contains all the errors of the form $X(\mathbf{c}_1, \ldots, \mathbf{c}_m)Z(\mathbf{d}_1, \ldots, \mathbf{d}_m)$ for which $\mathbf{c}_1, \ldots, \mathbf{c}_m \in \mathscr{C}$ with $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)$ and $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$. Clearly, the number of the errors $Z(\mathbf{d}_1, \ldots, \mathbf{d}_m)$, where $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$ is equal to $|\mathscr{C}^{\perp}| = q^{(n-k)m}$. We shall show that the number of errors $X(\mathbf{c}_1, \ldots, \mathbf{c}_m)$, where $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_{\Lambda})$ is equal to $q^{k(m-s)}$.

Let $\{F_{\Lambda_1}, \ldots, F_{\Lambda_{rks}}\}$ be an \mathbb{F}_p -basis for $\{F_{\Lambda} : \Lambda \in \mathscr{D}\}$. Each F_{Λ_i} can be represented by a $1 \times rkm$ matrix, say R_i with respect to a fixed ordered basis of \mathscr{C}^m . Then the $rks \times rks$ matrix

$$\mathcal{R} = \begin{pmatrix} R_1 \\ \vdots \\ R_{rks} \end{pmatrix}$$

represents the \mathbb{F}_p -linear transformation $F : \mathscr{C}^m \to \mathbb{F}_p^{rks}$ defined by $F(x) = (F_{\Lambda_1}(x), \ldots, F_{\Lambda_{rks}}(x))$ with respect to the same ordered basis of \mathscr{C}^m . Since

$$c_1R_1 + \dots + c_{rks}R_{rks} = 0$$
 if and only if $c_1F_{\Lambda_1} + \dots + c_{rks}F_{\Lambda_{rks}} = 0$

for any $c_1, \ldots, c_{rks} \in \mathbb{F}_p$, we see that the set $\{R_1, \ldots, R_{rks}\}$ is linearly independent over \mathbb{F}_p . Thus \mathcal{R} has rank rks, or equivalently, has nullity rk(m-s). Since $\ker(F) = \bigcap_{i=1}^{rks} \ker(F_{\Lambda_i}), \dim_{\mathbb{F}_p} \left(\bigcap_{\Lambda \in \mathscr{D}} \ker(F_{\Lambda})\right) = \dim_{\mathbb{F}_p} \left(\bigcap_{i=1}^{rks} \ker(F_{\Lambda_i})\right) = rk(m-s)$. Then the number of errors $X(\mathbf{c}_1, \ldots, \mathbf{c}_m)$, where $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_{\Lambda})$ is equal to $p^{rk(m-s)} = q^{k(m-s)}$.

It follows that $|\mathcal{S}| \geq q^{(n-k)m}q^{k(m-s)} = q^{nm-ks}$. On the other hand, since $Q_H(\mathscr{C}, \mathscr{D}) \subseteq \operatorname{Fix}(\mathcal{S})$, we have

$$q^{ks} = \dim_{\mathbb{F}_q}(Q_H(\mathscr{C}, \mathscr{D})) \le \dim_{\mathbb{F}_q}(\operatorname{Fix}(\mathcal{S})) = \frac{q^{nm}}{|\mathcal{S}|}$$

and so $|\mathcal{S}| \leq q^{nm-ks}$. This gives that $|\mathcal{S}| = q^{nm-ks}$, and hence \mathcal{S} consists of the errors of the form $X(\mathbf{c}_1, \ldots, \mathbf{c}_m) Z(\mathbf{d}_1, \ldots, \mathbf{d}_m)$ for which $\mathbf{c}_1, \ldots, \mathbf{c}_m \in \mathscr{C}$ with $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)$ and $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$. This shows, in particular, that $Q_H(\mathscr{C}, \mathscr{D})$ is a stabilizer code.

Finally, we shall show that $\delta = \min\{d(\mathscr{C}), \ell\}$. To see this, we first need to determine $\overline{\mathcal{S}}^{\perp_s}$. We claim that $\overline{\mathcal{S}}^{\perp_s}$ consists of $(\mathbf{u}_1, \ldots, \mathbf{u}_m \mid \mathbf{v}_1, \ldots, \mathbf{v}_m)$ for which $\mathbf{u}_1, \ldots, \mathbf{u}_m \in \mathscr{C}$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_m) \in \mathscr{D}^{\Theta}$. By above, we see that $\overline{\mathcal{S}}$ consists of the sequences $(\mathbf{c}_1, \ldots, \mathbf{c}_m \mid \mathbf{d}_1, \ldots, \mathbf{d}_m)$ such that $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)$ and $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$. Let $(\mathbf{U} \mid \mathbf{V}) = (\mathbf{u}_1, \ldots, \mathbf{u}_m \mid \mathbf{v}_1, \ldots, \mathbf{v}_m)$ be such that $\mathbf{u}_1, \ldots, \mathbf{u}_m \in \mathscr{C}$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_m) \in \mathscr{D}^{\Theta}$. Let $(\mathbf{C} \mid \mathbf{D}) = (\mathbf{c}_1, \ldots, \mathbf{c}_m \mid \mathbf{d}_1, \ldots, \mathbf{d}_m) \in \overline{\mathcal{S}}$. By the choice of $\mathbf{v}_1, \ldots, \mathbf{v}_m$, there exists $\Lambda = (\lambda_1, \ldots, \lambda_m) \in \mathscr{D}$ such that $\mathbf{v}_i \in \Theta(\lambda_i)$ for each $1 \leq i \leq m$. In other words, $f_{\lambda_i} = \rho_{\mathbf{v}_i}$ for each $1 \leq i \leq m$. This gives that

$$\operatorname{tr}_{q/p}(\mathbf{C}.\mathbf{V}) = \sum_{i=1}^{m} \operatorname{tr}_{q/p}(\mathbf{c}_{i}.\mathbf{v}_{i}) = \sum_{i=1}^{m} f_{\lambda_{i}}(\mathbf{c}_{i}) = F_{\Lambda}(\mathbf{c}_{1},\ldots,\mathbf{c}_{m}) = 0.$$
(3.1)

It follows that $\langle (\mathbf{C} \mid \mathbf{D}), (\mathbf{U} \mid \mathbf{V}) \rangle_s = \operatorname{tr}_{q/p}(\mathbf{d}.\mathbf{u} - \mathbf{v}.\mathbf{c}) = 0$; hence $(\mathbf{U} \mid \mathbf{V}) \in \overline{\mathcal{S}}^{\perp_s}$. Now let $(\mathbf{U} \mid \mathbf{V}) = (\mathbf{u}_1, \dots, \mathbf{u}_m \mid \mathbf{v}_1, \dots, \mathbf{v}_m) \in \overline{\mathcal{S}}^{\perp}$. Note that given any $\mathbf{d} \in \overline{\mathcal{S}}^{\perp_s}$. \mathscr{C}^{\perp} , $(0, \ldots, 0 \mid \mathbf{d}, 0, \ldots, 0)$, $(0, \ldots, 0 \mid 0, \mathbf{d}, 0, \ldots, 0)$, \ldots , $(0, \ldots, 0 \mid 0, \ldots, 0, \mathbf{d})$ all lie in $\overline{\mathcal{S}}$. Thus we have $\operatorname{tr}_{q/p}(\mathbf{u}_i.\mathbf{d}) = 0$ for all $\mathbf{d} \in \mathscr{C}^{\perp}$ and $1 \leq i \leq m$. It follows, from Lemma 3.1, that $\mathbf{u}_i \in \mathscr{C}$ for all $1 \leq i \leq m$. Now we shall show that $\mathbf{v} \in \mathscr{D}^{\Theta}$. To see this, it is enough to show that \mathscr{D}^{Θ} is equal to the additive code

$$\mathcal{V} := \{ (\mathbf{z}_1, \dots, \mathbf{z}_m) : \sum_{i=1}^m \operatorname{tr}_{q/p}(\mathbf{z}_i \cdot \mathbf{c}_i) = 0 \text{ for all } (\mathbf{c}_1, \dots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda) \}$$

$$(3.2)$$

over \mathbb{F}_q . By (3.1), we see that $\mathscr{D}^{\Theta} \subseteq \mathcal{V}$. To see the reverse inclusion, define $R_{\mathbf{y}_1,\ldots,\mathbf{y}_m} : \mathbb{F}_q^{nm} \to \mathbb{F}_p$ by $R_{\mathbf{y}_1,\ldots,\mathbf{y}_m}(\mathbf{z}_1,\ldots,\mathbf{z}_m) = \operatorname{tr}_{q/p}(\sum_{i=1}^m \mathbf{y}_i.\mathbf{z}_i)$ for all $\mathbf{y}_1,\ldots,\mathbf{y}_m,\mathbf{z}_1,\ldots,\mathbf{z}_m \in \mathbb{F}_q^n$. Note that $\mathcal{V} = \bigcap \{\ker(R_{\mathbf{c}_1,\ldots,\mathbf{c}_m}) : (\mathbf{c}_1,\ldots,\mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)\}$. Moreover, $\{R_{\mathbf{c}_1,\ldots,\mathbf{c}_m} : (\mathbf{c}_1,\ldots,\mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)\}$ is an \mathbb{F}_p -space, in a natural way, and the correspondence $(\mathbf{c}_1,\ldots,\mathbf{c}_m) \mapsto R_{\mathbf{c}_1,\ldots,\mathbf{c}_m}$ is an \mathbb{F}_p -space isomorphism. Thus, by similar arguments as used above, one can see that the \mathbb{F}_p -dimension of \mathcal{V} is equal to

$$\dim_{\mathbb{F}_p}(\mathbb{F}_q^{nm}) - \dim_{\mathbb{F}_p}\left(\bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda)\right) = rnm - rkm + rks,$$

and so $|\mathcal{V}| = q^{(n-k)m}q^{ks}$. One can also see that $|\mathscr{D}^{\Theta}| = q^{(n-k)m}.q^{ks}$. Since we already have $\mathscr{D}^{\Theta} \subseteq \mathcal{V}$, we must have the equality $\mathscr{D}^{\Theta} = \mathcal{V}$. Therefore, $\overline{\mathcal{S}}^{\perp_s}$ consists of $(\mathbf{u}_1, \ldots, \mathbf{u}_m \mid \mathbf{v}_1, \ldots, \mathbf{v}_m)$ for which $\mathbf{u}_1, \ldots, \mathbf{u}_m \in \mathscr{C}$ and $(\mathbf{v}_1, \ldots, \mathbf{v}_m) \in \mathscr{D}^{\Theta}$.

Let $\mathbf{c} \in \mathscr{C}$ and suppose that $(\mathbf{c}, \mathbf{0}, \dots, \mathbf{0}) \in \ker(F_{\Lambda})$ for all $\Lambda \in \mathscr{D}$. By our assumption on \mathscr{D} (see Remark 2) the first coordinates of the elements of \mathscr{D} form up \mathbb{F}_{q^k} . It follows that $c \in \bigcap_{\lambda \in \mathbb{F}_{q^k}} \ker(f_{\lambda}) = 0$. This gives that for every nonzero $\mathbf{c} \in \mathscr{C}$, the element $(\mathbf{c}, \mathbf{0}, \dots, \mathbf{0} \mid \mathbf{0}, \dots, \mathbf{0})$ of $(\mathbb{F}_q^n)^{2m}$ lies in $\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}$. Therefore, $\operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}) \leq d(\mathscr{C})$. On the other hand, for any nonzero $\Lambda = (\lambda_1, \dots, \lambda_m) \in \mathscr{D}$, an element $\mathbf{D} = (\mathbf{x}_{\lambda_1}, \dots, \mathbf{x}_{\lambda_m}) \in \mathscr{D}^{\Theta}$ cannot belong to $(\mathscr{C}^{\perp})^{(m)}$; hence the element $(\mathbf{0}, \dots, \mathbf{0} \mid \mathbf{D})$ of $(\mathbb{F}_q^n)^{2m}$ belongs to $\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}$. Therefore, we also have $\operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}) \leq \ell$. Consequently, $\operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}) \leq \min\{d(\mathscr{C}), \ell\}$.

Now suppose that $\operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}) < \min\{d(\mathscr{C}), \ell\}$. Let $(\mathbf{C} \mid \mathbf{D}) \in \overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}$ such that $\operatorname{swt}(\mathbf{C} \mid \mathbf{D}) = \operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}})$. Since $\operatorname{swt}(\mathbf{C} \mid \mathbf{D}) < d(\mathscr{C})$ and $\mathbf{C} \in \mathscr{C}^{(m)}$, we must have $\mathbf{C} = \mathbf{0}$. But since $\mathbf{D} \notin (\mathscr{C}^{\perp})^{(m)}$, we get $\ell \leq \operatorname{swt}(\mathbf{C} \mid \mathbf{D}) < \ell$, a contradiction. Therefore, $\operatorname{swt}(\overline{\mathcal{S}}^{\perp_s} \setminus \overline{\mathcal{S}}) = \min\{d(\mathscr{C}), \ell\}$.

Corollary 3.4. Let the situation be as in Theorem 3.3. If $d(\mathscr{C}) \leq d(\mathscr{D})$, then $\delta = d(\mathscr{C})$.

Proof. It is not difficult to see that the number ℓ in Theorem 3.3 is at least $d(\mathcal{D})$. Now the result follows since $\delta = \min\{d(\mathcal{D}), \ell\}$.

Corollary 3.5. Let the situation be as in Theorem 3.3. Suppose that $\mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_{q^k}\} \subset \mathbb{F}_{q^k}^m$. Then $\delta = \min\{d(\mathscr{C}), m\}$ and the stabilizer group

of $Q_H(\mathscr{C}, \mathscr{D})$ consists of the errors $X(\mathbf{c}_1, \dots, \mathbf{c}_m) Z(\mathbf{d}_1 \dots, \mathbf{d}_m)$, where $\mathbf{c}_1, \dots, \mathbf{c}_m \in \mathscr{C}$ with $\sum_{i=1}^m \mathbf{c}_i = 0$ and $\mathbf{d}_1, \dots, \mathbf{d}_m \in \mathscr{C}^{\perp}$.

Proof. It is known from the proof of Theorem 3.3 that \mathscr{D}^{Θ} is equal to the code

$$\mathcal{V} := \{ (\mathbf{z}_1, \dots, \mathbf{z}_m) : \sum_{i=1}^m \operatorname{tr}_{q/p}(\mathbf{z}_i \cdot \mathbf{c}_i) = 0 \text{ for all } (\mathbf{c}_1, \dots, \mathbf{c}_m) \in \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda) \}$$

over \mathbb{F}_q . Since $\mathscr{D} = \{(\lambda, \dots, \lambda) : \lambda \in \mathbb{F}_{q^k}\} \subset \mathbb{F}_{q^k}^m, \bigcap_{\Lambda \in \mathscr{D}} \ker(F_\Lambda) = \{(c_1, \dots, c_m) \in \mathscr{C}^{(m)} : c_1 + \dots + c_m \in \bigcap_{\lambda \in \mathbb{F}_{q^k}} f_\lambda = 0\}$ such that f_λ 's are all linear transformations from \mathscr{C} to \mathbb{F}_p . So $\mathscr{D}^{\Theta} = \{(c_1, \dots, c_m) \in \mathscr{C}^{(m)} : c_1 + \dots + c_m = 0\}$. Therefore the stabilizer group of $Q_H(\mathscr{C}, \mathscr{D})$ consists of the errors $X(\mathbf{c}_1, \dots, \mathbf{c}_m)Z(\mathbf{d}_1 \dots \mathbf{d}_m)$, where $\mathbf{c}_1, \dots, \mathbf{c}_m \in \mathscr{C}$ with $\sum_{i=1}^m \mathbf{c}_i = 0$ and $\mathbf{d}_1, \dots, \mathbf{d}_m \in \mathscr{C}^{\perp}$. For a suitable λ , x_λ could be $(10 \dots 0)$. Then wt(\mathbf{X}) = m, where \mathbf{X} is the m-fold tensor product of x_λ . The result follows since $\ell = m$.

Proposition 3.6. Let the situation be as in Theorem 3.3, where $\mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_{q^k}\} \subset \mathbb{F}_{q^k}^m$. Then

$$Q_H(\mathscr{C},\mathscr{D}) = span \left\{ \sum_{\substack{(\mathbf{c}_1, \dots, \mathbf{c}_m) \in \mathscr{C}^{(m)} \\ \mathbf{c}_1 + \dots + \mathbf{c}_m = c}} |\mathbf{c}_1 \dots \mathbf{c}_m \rangle : \mathbf{c} \in \mathscr{C} \right\}.$$

Proof. Let

$$\mathcal{A} = \left\{ \sum_{\substack{(\mathbf{c}_1, \dots, \mathbf{c}_m) \in \mathscr{C}^{(m)} \\ \mathbf{c}_1 + \dots + \mathbf{c}_m = c}} |\mathbf{c}_1 \dots \mathbf{c}_m \rangle : \mathbf{c} \in \mathscr{C} \right\}.$$

Since for any $\Lambda = (\lambda, \ldots, \lambda) \in \mathscr{D}$,

$$\Phi_{\Lambda} = \phi_{\lambda}^{\otimes m} = \frac{1}{\sqrt{q^{km}}} \sum_{(\mathbf{c}_{1}, \dots, \mathbf{c}_{m}) \in \mathscr{C}^{(m)}} \omega^{f_{\lambda}(\sum_{i=1}^{m} \mathbf{c}_{i})} |\mathbf{c}_{1} \dots \mathbf{c}_{m}\rangle$$
$$= \frac{1}{\sqrt{q^{km}}} \sum_{(\mathbf{c}_{1}, \dots, \mathbf{c}_{m}) \in \mathscr{C}^{(m)}} \omega^{f_{\lambda}(\mathbf{c})} \sum_{\mathbf{c}_{1} + \dots + \mathbf{c}_{m} = \mathbf{c}} |\mathbf{c}_{1} \dots \mathbf{c}_{m}\rangle,$$

we have $Q_H(\mathscr{C}, \mathscr{D}) \subseteq \operatorname{span} \mathcal{A}$. Since $\dim(Q_H(\mathscr{C}, \mathscr{D})) = q^k = \dim(\operatorname{span} \mathcal{A})$, we have the equality $Q_H(\mathscr{C}, \mathscr{D}) = \operatorname{span} \mathcal{A}$.

4 In Search of a Converse

Lemma 4.1. Let \mathscr{C} be a non-empty subset of \mathbb{F}_q^n and let $\phi_i = \sum_{\mathbf{c} \in \mathscr{C}} \omega^{\alpha_i(\mathbf{c})} | \mathbf{c} \rangle$, $\psi_i = \sum_{\mathbf{c} \in \mathscr{C}} \omega^{\beta_i(\mathbf{c})} | \mathbf{c} \rangle$ be elements of $(\mathbb{C}^q)^{\otimes n}$ for every $1 \leq i \leq m$, where $\omega = e^{2\pi i/p}$ and the α_i and β_i are functions from \mathscr{C} into \mathbb{F}_p . Suppose that

$$\phi_1 \otimes \cdots \otimes \phi_m = \psi_1 \otimes \cdots \otimes \psi_m$$

Then there exist $b_r \in \mathbb{F}_p$ $(1 \le r \le m)$ such that $\psi_r = \omega^{b_r} \phi_r$ for all $1 \le r \le m$.

Proof. By assumption, we have the equality

$$\sum_{(\mathbf{c}_1,\ldots,\mathbf{c}_m)\in\mathscr{C}^m}\omega^{\sum_{i=1}^m\alpha_i(\mathbf{c}_i)}|\mathbf{c}_1\ldots\mathbf{c}_m\rangle=\sum_{(\mathbf{c}_1,\ldots,\mathbf{c}_m)\in\mathscr{C}^m}\omega^{\sum_{i=1}^m\beta_i(\mathbf{c}_i)}|\mathbf{c}_1\ldots\mathbf{c}_m\rangle,$$

where \mathscr{C}^m denotes the *m*-fold Cartesian product of \mathscr{C} . Hence $\sum_{i=1}^m \alpha_i(\mathbf{c}_i) = \sum_{i=1}^m \beta_i(\mathbf{c}_i)$ for every $(\mathbf{c}_1, \ldots, \mathbf{c}_m) \in \mathscr{C}^m$. Fix a $\mathbf{c}_0 \in \mathscr{C}$. Then

$$\alpha_r(\mathbf{c}) + \sum_{\substack{i=1\\i\neq r}}^m \alpha_i(\mathbf{c}_0) = \beta_r(\mathbf{c}) + \sum_{\substack{i=1\\i\neq r}}^m \beta_i(\mathbf{c}_0)$$

for all $\mathbf{c} \in \mathscr{C}$ and $1 \leq r \leq m$. Let

$$b_r = \sum_{\substack{i=1\\i\neq r}}^m \left(\alpha_i(\mathbf{c}_0) - \beta_i(\mathbf{c}_0) \right).$$

Then $\alpha_r(\mathbf{c}) + b_r = \beta_r(\mathbf{c})$ for all $\mathbf{c} \in \mathscr{C}$ and $1 \leq r \leq m$. It, therefore, follows that $\psi_r = \omega^{b_r} \phi_r$ for all $1 \leq r \leq m$.

Theorem 4.2. Let H be a normalized $BH(q^k, p)$ matrix, where p is a prime number and $q = p^r$ for some positive integer r. Let $\mathscr{C} \subseteq \mathbb{F}_q^n$ be a classical linear code over \mathbb{F}_q of dimension k, where $1 \leq k < n$, and $\mathscr{D} = \{(\lambda, \ldots, \lambda) : \lambda \in \mathbb{F}_{q^k}\} \subset \mathbb{F}_{q^k}^m$ be the one-dimensional linear code over \mathbb{F}_{q^k} . If the quantum code $Q_H(\mathscr{C}, \mathscr{D})$ is a stabilizer code, then H is equivalent to the rk-fold Kronecker product of the Fourier matrix of order p.

Proof. Let $\mathscr{C} = {\mathbf{c}_1 = 0, \mathbf{c}_2, \dots, \mathbf{c}_{q^k}}$. We can write $H = [\omega^{f_i(\mathbf{c}_j)}]_{1 \le i,j \le q^k}$, where $\omega = e^{2\pi i/p}$ and $f_i : \mathscr{C} \to \mathbb{F}_p$ is a function for each $1 \le i \le q^k$. Now, $Q_H(\mathscr{C}, \mathscr{D})$ is the linear span of ${\phi_1^{\otimes m}, \dots, \phi_{q^k}^{\otimes m}}$ over \mathbb{F}_q , where

$$\phi_i = \frac{1}{\sqrt{q^k}} \sum_{j=1}^{q^k} \omega^{f_i(\mathbf{c}_j)} | \mathbf{c}_j \rangle \in (\mathbb{C}^q)^{\otimes n}$$

Suppose that $Q = Q_H(\mathscr{C}, \mathscr{D})$ is a stabilizer code and let $\mathcal{S} = \operatorname{Stab}(Q)$. By Lemma 3.2, $(\mathbf{0}, \ldots, \mathbf{0} \mid \mathbf{d}_1, \ldots, \mathbf{d}_m) \in \overline{\mathcal{S}}$ for all $\mathbf{d}_1, \ldots, \mathbf{d}_m \in \mathscr{C}^{\perp}$. Let $\mathbf{s} = (\mathbf{u}_1, \ldots, \mathbf{u}_m \mid \mathbf{v}_1, \ldots, \mathbf{v}_m) \in \overline{\mathcal{S}}$. Then \mathbf{s} is sympletically orthogoanl to the elements of $\overline{\mathcal{S}}$ of the forms $(\mathbf{0}, \ldots, \mathbf{0} \mid \mathbf{d}, \mathbf{0}, \ldots, \mathbf{0}), \ldots, (\mathbf{0}, \ldots, \mathbf{0} \mid \mathbf{0}, \ldots, \mathbf{0}, \mathbf{d})$ for all $\mathbf{d} \in \mathscr{C}^{\perp}$. Thus, $\operatorname{tr}_{q/p}(\mathbf{u}_i.d) = 0$ for all $1 \leq i \leq m$ and $\mathbf{d} \in \mathscr{C}^{\perp}$. Therefore, $\mathbf{u}_i \in \mathscr{C}$ for all $1 \leq i \leq m$ by Lemma 3.1. Note that there exists $h \in \mathbb{F}_p$ such that $\omega^h X(\mathbf{u}_1, \ldots, \mathbf{u}_m) Z(\mathbf{v}_1, \ldots, \mathbf{v}_m) \phi_j^{\otimes m} = \phi_j^{\otimes m}$ for all $1 \leq i \leq m$ and $\mathbf{u} \in \mathscr{C}^{\perp}$. This gives that $\omega^h X(\mathbf{u}_1) Z(\mathbf{v}_1) \phi_j \otimes \cdots \otimes X(\mathbf{u}_m) Z(\mathbf{v}_m) \phi_j = \phi_j \otimes \cdots \otimes \phi_j$, and so there exists $h_{ij} \in \mathbb{F}_p$ $(1 \leq i \leq m, 1 \leq j \leq q^k)$ such that $X(\mathbf{u}_i) Z(\mathbf{v}_i) \phi_j = \omega^{h_{ij}} \phi_j$ for all $1 \leq i \leq m$ and $1 \leq j \leq q^k$ by Lemma 4.1. Since $X(\mathbf{u}_i)Z(\mathbf{v}_i)\phi_j = \sum_{\mathbf{c}\in\mathscr{C}} \omega^{f_j(\mathbf{c}-\mathbf{u}_i)} \omega^{\operatorname{tr}_{q/p}(\mathbf{v}_i,\mathbf{c})} |\mathbf{c}\rangle$, we have

$$f_j(\mathbf{c} - \mathbf{u}_i) + \operatorname{tr}_{q/p}(\mathbf{v}_i.\mathbf{c}) = f_j(\mathbf{c}) + h_{ij}$$
(4.1)

for all $\mathbf{c} \in \mathscr{C}$, $1 \leq i \leq m$, and $1 \leq j \leq q^k$. Since H is assumed to be normalized, $f_1(\mathbf{c}) = 0$ for all $\mathbf{c} \in \mathscr{C}$ and $f_i(0) = 0$ for all $1 \leq i \leq q^k$. In particular, (4.1) yields $\operatorname{tr}_{q/p}(\mathbf{v}_i.\mathbf{c}) = h_{i1}$ for all $\mathbf{c} \in \mathscr{C}$ and $1 \leq i \leq m$. Substituting $\mathbf{c} = 0$ and j = 1 in (4.1), we obtain $h_{i1} = 0$ for all $1 \leq i \leq m$. Hence $\mathbf{v}_i \in \mathscr{C}^{\perp}$ for all $1 \leq i \leq m$ by Lemma 3.1. Since $\sum_{i=1}^m h_{ij} = -h$, this also shows that h = 0. Note that (4.1) turns into $f_j(\mathbf{c} - \mathbf{u}_i) = f_j(\mathbf{c}) + h_{ij}$, where $h_{ij} = f_j(-\mathbf{u}_i) = -f_j(\mathbf{u}_i)$ for all $1 \leq i \leq m$ and $1 \leq j \leq q^k$, and $\sum_{i=1}^m f_j(\mathbf{u}_i) = 0$ for all $1 \leq j \leq q^k$. It follows that

$$f_j(\mathbf{c} - \mathbf{u}_i) = f_j(\mathbf{c}) - f_j(\mathbf{u}_i) \tag{4.2}$$

for all $\mathbf{c} \in \mathscr{C}$, $1 \leq i \leq m$, and $1 \leq j \leq q^k$. Replacing \mathbf{c} by $\mathbf{c} + \mathbf{u}_i$ in (4.2), we obtain that $f_j(\mathbf{c} + \mathbf{u}_i) = f_j(\mathbf{c}) + f_j(\mathbf{u}_i)$ for all $\mathbf{c} \in \mathscr{C}$, $1 \leq i \leq m$, and $1 \leq j \leq q^k$. In particular, we have $f_j(\sum_{i=1}^m \mathbf{u}_i) = \sum_{i=1}^m f_j(\mathbf{v}_i) = 0$ for all $1 \leq j \leq q^k$. Since H, whose rank is q^k , has its first column consisting of 1's, this is possible only when $\sum_{i=1}^m \mathbf{u}_i = 0$. It follows that \overline{S} is contained in

$$\mathcal{A} := \{ (\mathbf{u}_1, \dots, \mathbf{u}_m \mid \mathbf{v}_1, \dots, \mathbf{v}_m) : \mathbf{v}_i \in \mathscr{C}^{\perp}, \ \mathbf{u}_i \in \mathscr{C}, \ \forall 1 \le i \le m \text{ with } \sum_{i=1}^m \mathbf{u}_i = 0 \}.$$

Rewriting \mathcal{A} as

$$\{(\mathbf{u}_1,\ldots\mathbf{u}_{m-1},-\sum_{i=1}^{m-1}\mathbf{u}_i\mid\mathbf{v}_1,\ldots,\mathbf{v}_m):\mathbf{u}_1,\ldots\mathbf{u}_m\in\mathscr{C},\ \mathbf{v}_1,\ldots,\mathbf{v}_m\in\mathscr{C}^{\perp}\},\$$

we have $|\overline{\mathcal{S}}| = q^{nm-k} = (q^k)^{m-1} (q^{n-k})^m = |\mathcal{A}|$; hence $\overline{\mathcal{S}} = \mathcal{A}$. Now (4.2) gives that $f_j(\mathbf{c} - \mathbf{c}') = f_j(\mathbf{c}) - f_j(\mathbf{c}')$ for all $\mathbf{c}, \mathbf{c}' \in \mathscr{C}$ and $1 \leq j \leq q^k$, proving that $f_j : \mathscr{C} \to \mathbb{F}_p$ is a linear transformation of \mathbb{F}_p -spaces.

A Appendix: Equivalence of Fourier-type BH Matrices

Throughout this brief note, R will denote a finite Frobenius ring and M a finite R-bimodule. We also denote the set of all non-degenerate bilinear forms on M by BLF(M).

Let $B: M \times M \longrightarrow R$ be a non-degenerate bilinear form on M. Associated to B are there right R-module homomorphisms, for all $x \in M$, defined by

$$B(x): \qquad M \longrightarrow R$$

$$y \longmapsto B(x,y)$$

that is B(x)(y) = B(x, y) for all $x, y \in M$. It follows that $B(x) \in M^* = \text{Hom}(M_R, R_R)$ for every $x \in M$. Note the following properties:

- $B(x_1 + x_2) = B(x_1) + B(x_2)$ for all $x_1, x_2 \in M$.
- B(rx) = rB(x) in the left *R*-module M^* for all $r \in R$ and $x \in M$.
- B(x) = B(x') if and only if x = x' for all $x, x' \in M$.
- $M^* = \{B(x) : x \in M\}.$

Let $\operatorname{Aut}(_RM)$ (respectively, $\operatorname{Aut}(M_R)$) denote the group of left (respectively, right) R-module automorphisms of M, equipped with the usual composition of maps. Given $\gamma \in \operatorname{Aut}(_RM)$ and $\eta \in \operatorname{Aut}(M_R)$, one can define two mappings assosiated with B as follows:



Observe that both B' and B'' are non-degenerate bilinear forms. Thus both groups $\operatorname{Aut}(_RM)$ and $\operatorname{Aut}(M_R)$ act on $\operatorname{BLF}(M)$. We write $B' = B \cdot \gamma$ and $B'' = B \cdot \eta$.

Theorem A.1. Let B be any non-degenerate bilinear form on M. Then we have

$$BLF(M) = \{B \cdot \gamma : \gamma \in Aut(_RM)\}\$$
$$= \{B \cdot \eta : \eta \in Aut(M_R)\}.$$

Proof. The arguments above the theorem show that $\{B \cdot \gamma : \gamma \in \operatorname{Aut}(_RM)\}$ lies in $\operatorname{BLF}(M)$.

Conversely, let B' be any other non-degenerate bilinear form on M. We shall show that $B' = B \cdot \gamma$ for a suitable $\gamma \in \operatorname{Aut}(_R M)$.

Let $M = \{0, x_1, ..., x_n\}$. Since

$$M^* = \{B(0), B(x_1), \dots, B(x_n)\} = \{B'(0), B'(x_1), \dots, B'(x_n)\},\$$

there exists $\sigma \in S_n$ such that $B'(x_i) = B(x_{\sigma(i)})$. Now define $\gamma : M \to M$ by $\gamma(0) = 0$ and $\gamma(x_i) = x_{\sigma(i)}$ for each $i = 1, \ldots, n$. Notice that

$$B'(x,y) = B(\gamma(x),y)$$

for all $x, y \in M$. Thus we complete the proof by showing that $\gamma \in Aut(_RM)$. Let $x_i + x_j = x_k$. Since

$$B(x_{\sigma(k)}) = B'(x_k)$$

= $B'(x_i + x_j)$
= $B'(x_i) + B'(x_j)$
= $B(x_{\sigma(i)}) + B(x_{\sigma(j)})$
= $B(x_{\sigma(i)} + x_{\sigma(j)})$

we have $x_{\sigma(i)} + x_{\sigma(j)} = x_{\sigma(k)}$. It follows that

$$\gamma(x_i + x_j) = \gamma(x_k) = x_{\sigma(k)} = x_{\sigma(i)} + x_{\sigma(j)} = \gamma(x_i) + \gamma(x_j),$$

i.e. γ is additive. On the other hand if $r \in R$ and $rx_i = x_j$, then

$$B(x_{\sigma(j)}) = B'(x_j)$$

= B'(rx_i)
= rB'(x_i)
= rB(x_{\sigma(i)})
= B(rx_{\sigma(i)})

hence $x_{\sigma(j)} = r x_{\sigma(i)}$, which yields

$$\gamma(rx_i) = \gamma(x_j) = x_{\sigma(j)} = rx_{\sigma(i)} = r\gamma(x_i),$$

as desired. This completes the proof of the first equality. By symmetric arguments, one easily prove the other equality. $\hfill \Box$

Proposition A.2. Let χ be a generating character of R and let B, B' be two non-degenerate bilinear forms of the R-bimodule $M = \{x_0 = 0, x_1, \ldots, x_n\}$. Then the matrices

$$H = [\chi(B(x_i, x_j))]_{0 \le i, j \le n}$$

and

$$H' = [\chi(B'(x_i, x_j))]_{0 \le i,j \le n}$$

are equivalent (by row permutation).

Proof. There exists an $\sigma \in S_n$ such that $B'(x_i, x_j) = B(x_{\sigma(i), x_j})$ for all $0 \leq i, j \leq n$ by the proof of Theorem A.1. Thus $H' = [\chi(B'(x_i, x_j))] = [B(x_{\sigma(i)}, x_j)]$ is H with rows permuted by σ .

Lemma A.3. Let χ and χ' be two generating characters of the ring R. Then there exists a unit element $a \in R$ such that $\chi' = \chi^a$.

Proof. Since χ is a generating character then there exists an element r of R such that $\chi' = \chi r$. Similarly, there exists an element r' of R such that $\chi = \chi' r'$. Therefore $\chi(1 - rr') = 0$ and $\chi \neq 0$. Then $\chi' = \chi r = \chi^r$ and r is a unit element of R.

Proposition A.4. Let χ and χ' be two generating characters of the ring R and let B be a non-degenerate bilinear form on the R-bimodule $M = \{x_0 = 0, x_1, \ldots, x_n\}$. Then the matrices

$$H = [\chi(B(x_i, x_j))]_{0 \le i,j \le n}$$

and

$$H' = [\chi'(B(x_i, x_j))]_{0 \le i,j \le n}$$

are equivalent (by row permutation).

Proof. By above lemma, there exists a unit element $a \in R$ such that $\chi' = \chi^a$. Since a is unit, we have aM = M. It follows that there exists $\sigma \in S_n$ such that $ax_i = x_{\sigma(i)}$ for all i = 1, ..., n. Now

$$H' = [\chi'(B(x_i, x_j))] = [\chi^a(B(x_i, x_j))] = [\chi(aB(x_i, x_j))] = [\chi(B(ax_i, x_j))] = [\chi(B(x_{\sigma(i)}, x_j))]$$

is clearly the matrix $[\chi(B(x_i, x_i))]$ with rows permuted by σ .

Proposition A.5. For the *R*-bimodule $M = \{x_0 = 0, x_1, \dots, x_n\}$, the matrices of the form

$$[\chi(B(x_i, x_j))]_{0 \le i,j \le n},$$

where χ is a generating character of R and $B: M \times M \to R$ is a non-degenerate bilinear form on M, are all equivalent.

Proof. For every non-degenerate bilinear form B' on M distinct from B according to Proposition A.2, $H' = [\chi(B'(x_i, x_j))]_{0 \le i,j \le n}$ is equivalent to H. Similarly for every χ' distinct from χ according to Proposition A.4, $H'' = [\chi'(B(x_i, x_j))]_{0 \le i,j \le n}$ is equivalent to H. This completes the proof. \Box

Corollary A.6. Let $H = [\omega^{f_i(\mathbf{c}_j)}]_{1 \leq i,j \leq q^k}$, where $\omega = e^{2\pi i/p}$ and $f_i : \mathscr{C} \to \mathbb{F}_p$ is a linear transformation for each $1 \leq i \leq q^k$. Then H is equivalent to a Kronecker product of Fourier matrices.

Proof. Since $f_i \in \mathcal{L}(\mathscr{C}, \mathbb{F}_p)$ for $1 \leq i, j \leq q^k$ and \mathscr{C} is isomorphic to \mathbb{F}_{q^k} then we use the composition of \mathbb{F}_p -space isomorphisms $\kappa : \mathscr{C} \to \mathbb{F}_{q^k} \to \mathcal{L}(\mathscr{C}, \mathbb{F}_p)$. Let $B : \mathscr{C} \times \mathscr{C} \to \mathbb{F}_p$ be a transformation defined as $B(c, c') = f_i(c')$ such that $\kappa(c) = f_i$. Then B is a non-degenerate bilinear form. Also $\chi : \mathbb{F}_p \to \mathbb{C}^*, \chi(a) = \omega^a$ is a generating character for \mathbb{F}_p . If we combine these we say

$$H = [\omega^{f_i(\mathbf{c}_j)}]_{1 \le i,j \le q^k} = [\chi(B(c_i, c_j))]_{1 \le i,j \le q^k}.$$

On the other hand $B' : \mathbb{F}_{q^k} \times \mathbb{F}_{q^k} \to \mathbb{F}_{q^k}, B(i,j) = ij$ is a non-degenerate bilinear form and $\chi' : \mathbb{F}_{q^k} \to \mathcal{C}^*, \chi'(a) = \omega^a$ is a generating character. Then the Fourier matrix $F_{ij} = [\omega^{ij}] = [\chi'(B'(i,j))]_{1 \leq i,j \leq q^k}$. The desired result is obtained by Proposition A.5.

References

- A. Ashikhmin and E. Knill. Nonbinary quantum stabilizer codes. *Ieee Transactions on Information Theory*, 47(7):3065–3072, 2001. Ashikhmin, A Knill, E AMS Spring Eastern Section Meeting Apr 28-29, 2001 Hoboken, nj Ams 1557-9654.
- [2] AT Butson. Generalized hadamard matrices. Proceedings of the American Mathematical Society, 13(6):894–898, 1962.
- [3] A Robert Calderbank and Peter W Shor. Good quantum error-correcting codes exist. *Physical Review A*, 54(2):1098, 1996.
- [4] Daniel Gottesman. Stabilizer codes and quantum error correction. California Institute of Technology, 1997.
- [5] W. F. Ke, K. F. Lai, and R. B. Zhang. Quantum codes from hadamard matrices. *Linear & Multilinear Algebra*, 58(7):847–854, 2010.
- [6] A. Ketkar, A. Klappenecker, S. Kumar, and P. K. Sarvepalli. Nonbinary stabilizer codes over finite fields. *Ieee Transactions on Information The*ory, 52(11):4892–4914, 2006. Ketkar, Avanti Klappenecker, Andreas Kumar, Santosh Sarvepalli, Pradeep Kiran Sarvepalli, Pradeep/0000-0001-8047-6946 1557-9654.
- [7] Avanti Ketkar, Andreas Klappenecker, Santosh Kumar, and Pradeep Kiran Sarvepalli. Nonbinary stabilizer codes over finite fields. *IEEE transactions* on information theory, 52(11):4892–4914, 2006.
- [8] Emanuel Knill. Non-binary unitary error bases and quantum codes. arXiv preprint quant-ph/9608048, 1996.
- [9] R. Majumdar, S. Basu, S. Ghosh, and S. Sur-Kolay. Quantum errorcorrecting code for ternary logic. *Physical Review A*, 97(5), 2018.
- [10] Peter W Shor. Scheme for reducing decoherence in quantum computer memory. *Physical review A*, 52(4):R2493, 1995.
- [11] K.F. Lai W.F. Ke and R.B. Zhang. Quantum codes from hadamard matrices. *Linear and Multilinear Algebra*, 58(7):847–854, 2010.