

BIHARMONIC FUNCTIONS AND BI-EIGENFUNCTIONS ON SOME MODEL SPACES

YE-LIN OU

ABSTRACT

In this paper, we first give a convenient formula for bi-Laplacian on a sphere and the complete description of its eigenvalues, buckling eigenvalues, and their corresponding eigenfunctions. We then show that the radial (or rotationally symmetric) solutions for biharmonic equation on the model space $(\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r) g^{S^{m-1}})$ can be given by an integral formula. We also prove that the model space always admits proper biharmonic functions as the products of any eigenfunctions of the factor sphere with certain radial functions. Many explicit examples of proper biharmonic functions on space forms are given. Finally, we give a complete classification of proper biharmonic functions with positive Laplacian on the punctured Euclidean space.

1. BI-LAPLACIAN, BI-EIGENFUNCTIONS AND BUCKLING EIGENFUNCTIONS ON A SPHERE

In this section, we derive a formula for the bi-Laplacian on functions on a sphere using the Euclidean bi-Laplacian and give a complete description of all bi-eigenvalues, buckling eigenvalues and the corresponding bi-eigenfunctions and buckling eigenfunctions on the sphere S^m .

Our sign convention for the Laplace operator acting on functions on a Riemannian manifold (M, g) is $\Delta = \operatorname{div}(\nabla f)$, and the bi-Laplace operator is denoted by $\Delta^2 f = \Delta(\Delta f)$.

1.1 A formula for bi-Laplacian on functions on the sphere S^m

Date: 7/21/2024.

2020 Mathematics Subject Classification. 58E20, 53C12.

Key words and phrases. Bi-Laplacian, bi-eigenvalues, biharmonic functions, bi-eigenfunctions, spherical harmonics, model space.

Let $\mathbf{i} : S^m \rightarrow \mathbb{R}^{m+1}$ denote the standard isometric embedding of the unit sphere into Euclidean space. It is well known (see e.g., [9], Lemma 9.6) that the Euclidean Laplacian and the Laplacian on a sphere are related by

$$(1) \quad \Delta_{S^m}(F \circ \mathbf{i}) = \left(\Delta_{\mathbb{R}^{m+1}} F - \frac{\partial^2 F}{\partial r^2} - m \frac{\partial F}{\partial r} \right) \circ \mathbf{i}.$$

where $\frac{\partial}{\partial r}$ denotes the directional derivative in the radial direction.

Using the fact that the radial projection $\pi : \mathbb{R}^{m+1} \rightarrow S^m, \pi(x) = \frac{x}{|x|}$ is a harmonic morphism (a horizontally weakly conformal harmonic map) we derive the following convenient formula to compute the bi-Laplacian on a sphere using the data and the operators on the ambient Euclidean space.

Theorem 1.1. *For any function $f : S^m \supseteq U \rightarrow \mathbb{R}$, we have*

$$(2) \quad \Delta_{S^m}^2 f = \left(\Delta_{\mathbb{R}^{m+1}}^2 f \left(\frac{x}{|x|} \right) \right)_{S^m} + 2(m-3) \Delta_{S^m} f, \text{ or}$$

$$(3) \quad \Delta_{S^m}^2 f = \left(\Delta_{\mathbb{R}^{m+1}}^2 f \left(\frac{x}{|x|} \right) + 2(m-3) \Delta_{\mathbb{R}^{m+1}} f \left(\frac{x}{|x|} \right) \right)_{S^m}.$$

Proof. It is well known (see e.g. [2]) that the radial projection $\pi : \mathbb{R}^{m+1} \rightarrow S^m$ with $\pi(x) = \frac{x}{|x|}$ is a horizontally homothetic harmonic morphism with dilation $\lambda = \frac{1}{|x|}$. It follows that for any function $f : S^m \supseteq U \rightarrow \mathbb{R}$, we have

$$(4) \quad \Delta_{\mathbb{R}^{m+1}}(f \circ \pi) = \lambda^2 (\Delta_{S^m} f) \circ \pi, \quad \forall x \in \pi^{-1}(U) \subseteq \mathbb{R}^{m+1} \setminus \{0\}, \text{ i.e.,}$$

$$\Delta_{\mathbb{R}^{m+1}} f \left(\frac{x}{|x|} \right) = \frac{1}{|x|^2} (\Delta_{S^m} f) \left(\frac{x}{|x|} \right), \quad \forall x \in \pi^{-1}(U) \subseteq \mathbb{R}^{m+1} \setminus \{0\}, \text{ or}$$

$$(5) \quad (\Delta_{S^m} f) \circ \pi = |x|^2 \Delta_{\mathbb{R}^{m+1}} \left(f \left(\frac{x}{|x|} \right) \right), \quad \forall x \in \pi^{-1}(U) \subseteq \mathbb{R}^{m+1} \setminus \{0\}.$$

Evaluating the above equation on S^m we have

$$(6) \quad \Delta_{S^m} f = \left(\Delta_{\mathbb{R}^{m+1}} f \left(\frac{x}{|x|} \right) \right)_{S^m}.$$

Applying $\Delta_{\mathbb{R}^{m+1}}$ to both sides of (4), using the product rule for Laplace operator, and a straightforward computation yields

$$\begin{aligned} \Delta_{\mathbb{R}^{m+1}}^2(f \circ \pi) &= \Delta_{\mathbb{R}^{m+1}}(\Delta_{\mathbb{R}^{m+1}}(f \circ \pi)) = \Delta_{\mathbb{R}^{m+1}}(\lambda^2 (\Delta_{S^m} f) \circ \pi) \\ &= \lambda^4 (\Delta_{S^m}^2 f) \circ \pi + 2 \langle \nabla \lambda^2, \nabla((\Delta_{S^m} f) \circ \pi) \rangle + (\Delta_{\mathbb{R}^{m+1}} \lambda^2) (\Delta_{S^m} f) \circ \pi \\ (7) \quad &= \lambda^4 (\Delta_{S^m}^2 f) \circ \pi + 2 (\Delta_{S^m} f)_\alpha \circ \pi \langle \nabla \lambda^2, \nabla \pi^\alpha \rangle + (\Delta_{\mathbb{R}^{m+1}} \lambda^2) (\Delta_{S^m} f) \circ \pi. \end{aligned}$$

Using the fact that π is horizontally homothetic and hence $\langle \nabla \lambda^2, \nabla \pi^\alpha \rangle = 0, \forall \alpha = 1, 2, \dots, m$, together with $\lambda^2 = \frac{1}{|x|^2}$, $\Delta_{\mathbb{R}^{m+1}} \lambda^2 = -\frac{2(m-3)}{|x|^4}$, we have

$$\begin{aligned} \Delta_{\mathbb{R}^{m+1}}^2(f \circ \pi) &= \frac{1}{|x|^4}(\Delta_{S^m}^2 f) \circ \pi - \frac{2(m-3)}{|x|^4}(\Delta_{S^m} f) \circ \pi, \text{ or} \\ (8) \quad |x|^4 \Delta_{\mathbb{R}^{m+1}}^2 f \left(\frac{x}{|x|} \right) &= (\Delta_{S^m}^2 f) \left(\frac{x}{|x|} \right) - 2(m-3)(\Delta_{S^m} f) \left(\frac{x}{|x|} \right). \end{aligned}$$

Evaluate both sides of (8) on $x \in S^m$ (i.e., $|x| = 1$) gives (2). Using (2) and (6) we have

$$\left(\Delta_{\mathbb{R}^{m+1}}^2 f \left(\frac{x}{|x|} \right) \right)_{S^m} = \Delta_{S^m}^2 f - 2(m-3) \left(\Delta_{\mathbb{R}^{m+1}} f \left(\frac{x}{|x|} \right) \right)_{S^m},$$

from which we obtain (3). □

As immediate consequences, we have

Corollary 1.2. (i) For any function $f : S^3 \supseteq U \rightarrow \mathbb{R}$, we have

$$(9) \quad \Delta_{S^3}^2 f = \left(\Delta_{\mathbb{R}^4}^2 f \left(\frac{x}{|x|} \right) \right)_{S^3}.$$

(ii) A function $f : S^2 \supseteq U \rightarrow \mathbb{R}$ is a buckling eigenfunction of eigenvalue $\mu = 2$, i.e., $\Delta_{S^2}^2 f = -2\Delta_{S^2} f$ if and only if $\left(\Delta_{\mathbb{R}^3}^2 f \left(\frac{x}{|x|} \right) \right)_{S^2} = 0$.

Example 1. The restriction of $f(x) = \ln(1 - x_3)$, $x \in \mathbb{R}^3$ is a proper biharmonic function on S^2 . In fact, in this case, we have $(f \circ \pi)(x) = f\left(\frac{x}{|x|}\right) = \ln\left(1 - \frac{x_3}{|x|}\right)$. A straightforward computation yields

$$(10) \quad \Delta_{\mathbb{R}^3} \left(f\left(\frac{x}{|x|}\right) \right) = -\frac{1}{|x|^2}, \quad \Delta_{\mathbb{R}^3}^2 \left(f\left(\frac{x}{|x|}\right) \right) = -\frac{2}{|x|^4}.$$

Substituting these into (6) and (3) we have, respectively

$$(11) \quad \Delta_{S^2} f = \left(\Delta_{\mathbb{R}^3} \left(f\left(\frac{x}{|x|}\right) \right) \right)_{|x|=1} = 1 \neq 0,$$

$$(12) \quad \Delta_{S^2}^2 f = \left(\Delta_{\mathbb{R}^3}^2 f \left(\frac{x}{|x|} \right) - 2\Delta_{\mathbb{R}^3} f \left(\frac{x}{|x|} \right) \right)_{S^2} \equiv 0.$$

Thus, $f(x) = \ln(1 - x_3)$, $x \in S^2$ is a proper biharmonic function on the sphere $S^2 \setminus \{N\}$.

Example 2. The restriction of $f(x) = (ax_1 + bx_2 + cx_3)/\sqrt{1 - x_4^2}$ to $S^3 \setminus \{N, S\}$ is a family of proper biharmonic function on 3-sphere.

In fact, $f(\frac{x}{|x|}) = (ax_1 + bx_2 + cx_3)/\sqrt{\sum_{i=1}^3 x_i^2}$, and a straightforward computation gives

$$(13) \quad \Delta_{\mathbb{R}^4} \left(f\left(\frac{x}{|x|}\right) \right) = \Delta_{\mathbb{R}^4} \left(\frac{ax_1 + bx_2 + cx_3}{\sqrt{x_1^2 + x_2^2 + x_3^2}} \right) = \frac{-2(ax_1 + bx_2 + cx_3)}{(x_1^2 + x_2^2 + x_3^2)^{3/2}},$$

$$\Delta_{\mathbb{R}^4}^2 \left(f\left(\frac{x}{|x|}\right) \right) = \Delta_{\mathbb{R}^4} \left(\frac{-2(ax_1 + bx_2 + cx_3)}{(x_1^2 + x_2^2 + x_3^2)^{3/2}} \right) = 0.$$

It follows from (9) that $\Delta_{S^3}^2 f = \left(\Delta_{\mathbb{R}^4}^2 f\left(\frac{x}{|x|}\right) \right)_{S^3} = 0$.

Corollary 1.3. *$f = F|_{S^m}$ is the restriction of a homogeneous polynomial F of degree k on \mathbb{R}^{m+1} , then we have*

$$(14) \quad \Delta_{S^m}^2 f = k(k+2)(k-1+m)(k+m-3)f(x) + 2(m-3)\Delta_{S^m} f$$

$$+ \left(\Delta_{\mathbb{R}^{m+1}}^2 F(x) - 2k(k+m-3)\Delta_{\mathbb{R}^{m+1}} F(x) \right)_{S^m}, \text{ or}$$

$$(15) \quad \Delta_{S^m}^2 f = k^2(k-1+m)^2 f$$

$$+ \left(\Delta_{\mathbb{R}^{m+1}}^2 F(x) - 2[k^2 + (k-1)(m-3)]\Delta_{\mathbb{R}^{m+1}} F(x) \right)_{S^m}.$$

Proof. It is easily checked (see also [1], Lemma 4.4) that if $f : S^m \rightarrow \mathbb{R}$ with $f = F|_{S^m}$ is a restriction of a homogeneous polynomial F of degree k , then we have

$$(16) \quad \nabla |x|^\alpha = \alpha |x|^{\alpha-2} x,$$

$$(17) \quad \Delta_{\mathbb{R}^{m+1}} |x|^\alpha = \alpha(\alpha-1+m)|x|^{\alpha-2},$$

$$(18) \quad \langle x, \nabla F(x) \rangle = kF(x).$$

Since $f = F|_{S^m}$ is a restriction of a homogeneous polynomial F of degree k , we have $f(\frac{x}{|x|}) = F(\frac{x}{|x|}) = |x|^{-k}F(x)$. A straightforward computation using (16)-(18) gives

$$(19) \quad \Delta_{\mathbb{R}^{m+1}} f \left(\frac{x}{|x|} \right) = \Delta_{\mathbb{R}^{m+1}} (|x|^{-k}F(x))$$

$$= (\Delta_{\mathbb{R}^{m+1}} |x|^{-k}) F(x) + |x|^{-k} \Delta_{\mathbb{R}^{m+1}} F(x) + 2\langle \nabla |x|^{-k}, \nabla F(x) \rangle$$

$$= -k(k-1+m)|x|^{-k-2} F(x) + |x|^{-k} \Delta_{\mathbb{R}^{m+1}} F(x).$$

Evaluate both sides of the above on S^m we have

$$(20) \quad (\Delta_{S^m} F|_{S^m})(x) = \left(\Delta_{\mathbb{R}^{m+1}} |x|^{-k} F|_{S^m}(x) \right)_{|x|=1}$$

$$= -k(k-1+m) F|_{S^m}(x) + (\Delta_{\mathbb{R}^{m+1}} F(x))|_{S^m}.$$

Applying $\Delta_{\mathbb{R}^{m+1}}$ to both sides of (19) and a further calculation using (16)-(18) yields

$$\begin{aligned}
 (21) \quad \Delta_{\mathbb{R}^{m+1}}^2 f \left(\frac{x}{|x|} \right) &= \Delta_{\mathbb{R}^{m+1}} [-k(k-1+m) |x|^{-k-2} F(x) + |x|^{-k} \Delta_{\mathbb{R}^{m+1}} F(x)] \\
 &= -k(k-1+m) \Delta_{\mathbb{R}^{m+1}} [|x|^{-k-2} F(x)] + \Delta_{\mathbb{R}^{m+1}} [|x|^{-k} \Delta_{\mathbb{R}^{m+1}} F(x)] \\
 &= k(k+2)(k-1+m)(k+m-3) |x|^{-k-4} F(x) - 2k(k+m-3) |x|^{-k-2} \Delta_{\mathbb{R}^{m+1}} F(x) \\
 &\quad + |x|^{-k} \Delta_{\mathbb{R}^{m+1}}^2 F(x).
 \end{aligned}$$

Evaluating both sides of (21) on S^m and substituting the result into (2) gives (14). Finally, (15) is obtained by substituting (20) into (14). \square

Remark 1. Note that Formulas (6) and (20) were proved in [14] in a different way.

1.2 Bi-eigenvalues, buckling eigenvalues, and their eigenfunctions on the sphere S^m

Definition 1.4. A function $f \in C^\infty(M)$ is called an eigenfunction of the Laplace operator with eigenvalue λ if it solves the PDE

$$(22) \quad \Delta f = -\lambda f.$$

The set of all eigenvalues (called the spectrum) of Δ is denoted by $\text{Spec}_\Delta(M)$. The eigenspace corresponding to the eigenvalue λ_i is denoted by

$$V(\lambda_i) = \{f \in C^\infty(M) \mid \Delta f = -\lambda_i f\}.$$

A function $f \in C^\infty(M)$ is called an eigenfunction of the bi-Laplace operator with eigenvalue μ if it solves the PDE

$$(23) \quad \Delta^2 f = \mu f.$$

For convenience, eigenfunctions and eigenvalues of the bi-Laplace operator are also called **bi-eigenfunctions** and **bi-eigenvalue** of the Laplacian respectively.

The set of all bi-eigenvalues of Δ is denoted by $\text{Spec}_{\Delta^2}(M)$. The bi-eigenspace corresponding to the bi-eigenvalue μ_i is denoted by

$$V^2(\mu_i) = \{f \in C^\infty(M) \mid \Delta^2 f = \mu_i f\}.$$

A function $f \in C^\infty(M)$ is called a **buckling eigenfunction** with a **buckling eigenvalue** ν if it solves the PDE

$$(24) \quad \Delta^2 f = -\nu \Delta f.$$

The set of all buckling-eigenvalues is denoted by $\text{Spec}_{\Delta^b}(M)$. The buckling-eigenspace corresponding to the buckling-eigenvalue ν_i is denoted by

$$V^b(\nu_i) = \{f \in C^\infty(M) \mid \Delta^2 f = -\nu_i \Delta f\}.$$

It is well known (see e.g., [17]) that all of the above three types of eigenvalues are countable so we can list them as

$$\begin{aligned} 0 = \lambda_0 < \lambda_1 < \lambda_2 \leq \lambda_3 \leq \cdots + \infty & \quad (\text{Laplacian eigenvalues}) \\ 0 = \mu_0 < \mu_1 < \mu_2 \leq \mu_3 \leq \cdots + \infty & \quad (\text{bi - eigenvalues}) \\ 0 = \nu_0 < \nu_1 < \nu_2 \leq \nu_3 \leq \cdots + \infty & \quad (\text{Buckling eigenvalues}) \end{aligned}$$

Remark 2. One can easily check the following

(i) The square of any eigenvalue of the Laplacian Δ is an eigenvalue of the bi-Laplacian Δ^2 , and any eigenfunction is automatically a bi-eigenfunction, i.e.,

$$(25) \quad \text{Spec}_{\Delta^2}(M) \supset \{\lambda^2 \mid \forall \lambda \in \text{Spec}_\Delta(M)\}, \text{ and } V^2(\lambda^2) \supset V(\lambda).$$

(ii) The set of all buckling eigenvalues on (M, g) is the same as the set of all eigenvalues of the Laplacian Δ , and a buckling eigenfunction is precisely a function whose Laplacian is an eigenfunction of Δ , i.e.,

$$(26) \quad \text{Spec}_{\Delta^b}(M) = \text{Spec}_\Delta(M), \text{ and } V^b(\lambda) = \{f \in C^\infty(M) \mid \Delta f \in V(\lambda)\}.$$

For eigenvalues and the corresponding eigenspaces of the Laplacian on a sphere, we have the following well-known results.

Proposition 1.5. (see e.g., [14] Proposition 22.2.) *The restriction to S^m of any harmonic homogeneous polynomial of degree k on \mathbb{R}^{m+1} is an eigenfunction of the Laplacian Δ_{S^m} with eigenvalue $\lambda_k = k(m+k-1)$. Conversely, any eigenfunction of Δ_{S^m} on the sphere S^m is a restriction of a homogeneous polynomial of degree k .*

Now we are ready to prove the following theorem which gives complete descriptions of bi-eigenvalues and buckling eigenvalues and the corresponding eigenspaces of the sphere.

Theorem 1.6. (I) All eigenvalues of bi-Laplacian on S^m come from a square of an eigenvalue of the Laplacian, and bi-eigenfunction on S^m are precisely the restrictions of harmonic homogenous polynomial in \mathbb{R}^{m+1} . This can be described as

$$(27) \quad \text{Spec}_{\Delta^2}(S^m) = \{\lambda_k^2 \mid \lambda_k = k(m+k-1), \forall k \in \mathbb{N}\}, \text{ and}$$

$$(28) \quad V^2(\lambda_k^2) = \mathcal{H}^k,$$

where \mathcal{H}^k denotes the set of harmonic homogeneous polynomials of degree k . In particular, the first nonzero eigenvalue of the bi-Laplacian on S^m is $\mu_1 = m^2$.

(II) The set of all buckling eigenvalues on S^m is the same as the set of all eigenvalues of the Laplacian Δ on S^m , and buckling eigenfunctions on S^m are precisely the restrictions of polynomial of the form $h_k + c|x|^2$. This can be described as

$$\text{Spec}_{\Delta^b}(S^m) = \text{Spec}_{\Delta}(S^m) = \{\lambda_k = k(m+k-1) \mid \forall k \in \mathbb{N}\}, \text{ and}$$

$$V^b(\lambda_k) = \mathcal{H}^k \oplus \{c|x|^2\}.$$

In particular, the first nonzero buckling eigenvalue of S^m is $\mu_1 = m$.

Proof. Recall that for a closed manifold (M, g) , $C^\infty(M)$ is an inner product space with the inner product defined by

$$(29) \quad \langle \alpha, \beta \rangle = \int_M \alpha \beta dv_g$$

and the Laplace operator $\Delta : C^\infty(M) \rightarrow C^\infty(M)$ is a self-adjoint operator, so we have

$$(30) \quad \langle \Delta \alpha, \beta \rangle = \langle \alpha, \Delta \beta \rangle, \quad \forall \alpha, \beta \in C^\infty(M).$$

Note also that the space $L^2(S^m)$ of all real-valued square-integrable functions on S^m is a Hilbert space with the inner product also defined by (29).

Let λ_k be an eigenvalue of the Laplacian on S^m , i.e., $\Delta_{S^m} f = -\lambda f$ for some function on S^m . Then, as in Remark 2, λ_k^2 is a bi-eigenvalue. Conversely, if $\Delta_{S^m}^2 f = \mu f$ for a constant μ , we are to prove that $\mu = \lambda_k^2$ for one and only one $k \in \mathbb{N}$. It is well known (see e.g., [1], Theorem 5.12) that $L^2(S^m) = \bigoplus_{i=0}^{\infty} \mathcal{H}^i$ as the direct sum of spherical harmonic polynomials of homogeneous degree k . For any smooth function f on S^m , we have $f = \sum_{i=0}^{\infty} h_i$ in $L^2(S^m)$. For $k \in \mathbb{N}$, we

have

$$\begin{aligned}
(31) \quad \|h_k\|_{L^2(S^m)} &= \langle f, h_k \rangle_{L^2(S^m)} = \frac{1}{\lambda_k^2} \langle f, \Delta_{S^m}^2 h_k \rangle_{L^2(S^m)} = \frac{1}{\lambda_k^2} \langle \Delta_{S^m}^2 f, h_k \rangle_{L^2(S^m)} \\
&= \frac{1}{\lambda_k^2} \langle \mu f, h_k \rangle_{L^2(S^m)} = \frac{\mu}{\lambda_k^2} \langle f, h_k \rangle_{L^2(S^m)} = \\
&= \frac{\mu}{\lambda_k^2} \langle f, h_k \rangle_{L^2(S^m)} = \frac{\mu}{\lambda_k^2} \|h_k\|_{L^2(S^m)}.
\end{aligned}$$

It follows that

$$(32) \quad \left(1 - \frac{\mu}{\lambda_k^2}\right) \|h_k\|_{L^2(S^m)} = 0.$$

Case (i): if $\mu \neq \lambda_k^2$ for any $k \geq 1$, then $h_k = 0, \forall k \geq 1$ in $L^2(S^m)$ and hence $f = h_0$ is a constant function. In this case, we have $\Delta_{S^m}^2 f = 0 = \mu f$ and $\mu = 0$.

Case (ii): if $\mu = \lambda_k^2$ for some $k \in \mathbb{N}$, i.e., $\Delta_{S^m}^2 f = \mu f = \lambda_k^2 f$, we will show that f is orthogonal to \mathcal{H}^l for any $l \geq 1$ and $l \neq k$ in $L^2(S^m)$. In fact, for any $l \geq 1$ and any $h_l \in \mathcal{H}^l$ we have

$$\begin{aligned}
(33) \quad \langle h_l, f \rangle_{L^2(S^m)} &= \frac{1}{\lambda_l^2} \langle \Delta_{S^m}^2 h_l, f \rangle_{L^2(S^m)} = \frac{1}{\lambda_l^2} \langle h_l, \Delta_{S^m}^2 f \rangle_{L^2(S^m)} \\
&= \frac{1}{\lambda_l^2} \langle h_l, \lambda_k^2 f \rangle_{L^2(S^m)} = \frac{\lambda_k^2}{\lambda_l^2} \langle h_l, f \rangle_{L^2(S^m)}.
\end{aligned}$$

From this and the fact that $\frac{\lambda_k^2}{\lambda_l^2} \neq 1$ we conclude that f is orthogonal to \mathcal{H}^l for any $l \geq 1$ and any $l \neq k$ in $L^2(S^m)$, so we conclude that f must be of the form $f = h_0 + h_k \in \mathcal{H}^0 \oplus \mathcal{H}^k$. In this case, $\Delta_{S^m}^2 f = \lambda_k^2 f$ if and only if $h_0 = 0$. Thus, any bi-eigenfunction corresponding to the bi-eigenvalue λ_k^2 is a restriction of a harmonic homogeneous polynomial of degree k . This completes the proof of Statement (I).

For Statement (II), by Remark 2, we only need to prove that the buckling eigenspace corresponding to the buckling eigenvalue λ_k is $V^b(\lambda_k) = \mathcal{H}^k \oplus \{c|x|^2 \mid c \in \mathbb{R}\}$. In fact, if $f \in C^\infty(S^m)$ is a buckling eigenfunction corresponding to eigenvalue λ_k such that $\Delta_{S^m}^2 f = -\lambda_k \Delta_{S^m} f$, then, rewriting it as $\Delta_{S^m}(\Delta_{S^m} f) = -\lambda_k(\Delta_{S^m} f)$, we see that $\Delta_{S^m} f$ is an eigenfunction of Δ_{S^m} with eigenvalue λ_k . By Proposition 1.5, we have

$$(34) \quad \Delta_{S^m} f = q_k|_{S^m} f \text{ for some } q_k \in \mathcal{H}^k.$$

Using the unique representation of f in $L^2(S^m) = \bigoplus_{i=0}^{\infty} \mathcal{H}^i$ we have $f = \sum_{i=0}^{\infty} h_i$. We compute the i th components of f for $i \geq 1$ to have

$$(35) \quad \begin{aligned} \langle f, h_i \rangle_{L^2(S^m)} &= -\frac{1}{\lambda_i} \langle f, \Delta_{S^m} h_i \rangle_{L^2(S^m)} = -\frac{1}{\lambda_i} \langle \Delta_{S^m} f, h_i \rangle_{L^2(S^m)} \\ &= -\frac{1}{\lambda_i} \langle q_k, h_i \rangle_{L^2(S^m)} = 0, \quad \forall i \geq 1, \text{ and } i \neq k, \end{aligned}$$

where the 3rd “=” was obtained by using (34) and the last “=” holds because $(q_k, h_i) \in \mathcal{H}^k \oplus \mathcal{H}^i$ and hence $\langle q_k, h_i \rangle_{L^2(S^m)} = 0, \forall i \geq 1$, and $i \neq k$. It follows from (35) that $h_i = 0$ for all $i \neq 0, k$ and hence $f = \sum_{i=0}^{\infty} h_i = (h_k + h_0)|_{S^m}$ or $f = (h_k + c|x|^2)|_{S^m}$ for $h_0 = c$, which completes the proof of the theorem. \square

Corollary 1.7. *The restriction of any polynomial of homogeneous degree 2 to S^m is a buckling eigenfunction (i.e., solution of $\Delta_{S^m}^2 f = -\mu \Delta_{S^m} f$) on sphere with $\mu = 2(m+1)$.*

Proof. It is well known (see e.g., [1]) any homogeneous polynomial of degree 2 can be written as $p_2(x) = h_2(x) + |x|^2 h_0$. From this and Statement (II) of Theorem 1.6 we obtain the corollary. \square

Remark 3. Although the buckling eigenvalue $\mu = 2(m+1)$ is the second eigenvalue of the Laplacian Δ_{S^m} , the buckling eigenspace $V^b(\mu)$ corresponds to $\mu = 2(m+1)$ has one dimension bigger than the eigenspace $V(\mu)$ of the Laplace operator. The buckling eigenspace includes all homogeneous polynomials of degree 2 whilst the latter consists of only all **harmonic** homogeneous polynomials of degree 2.

For any integer $k \geq 2$, the k -Laplacian is define to be $\Delta^k f = \Delta(\Delta^{k-1})$. We define μ to be an eigenvalue of the k -Laplacian if there is a nonconstant function f such that $\Delta^k f = (-1)^k \mu f$. Then, a similar proof gives the following description of all eigenvalues of the k -Laplacian on the sphere.

Corollary 1.8. *The spectrum of the k -Laplacian on the sphere S^m is*

$$\text{Spec}_{\Delta^k}(S^m) = \{\lambda_i^k \mid \lambda_i = i(m+i-1), \forall i \in \mathbb{N}\}.$$

In particular, the first nonzero eigenvalue of Δ^k on S^m is m^k .

Many recent work (see e.g., [5, 15, 4, 17, 4, 16, 18]) study bi-eigenvalue and buckling bi-eigenvalue problems on a domain of a Riemannian manifold with

boundary conditions like

$$\textit{The Clamped plate problem} : \begin{cases} \Delta^2 f = \mu f & \text{on } M \\ f|_{\partial M} = \frac{\partial f}{\partial n}|_{\partial M} = 0, \end{cases}$$

$$\textit{The buckling eigenvalue problem} : \begin{cases} \Delta^2 f = -\nu \Delta f & \text{on } M \\ f|_{\partial M} = \frac{\partial f}{\partial n}|_{\partial M} = 0. \end{cases}$$

2. RADIAL BIHARMONIC FUNCTIONS ON MODEL SPACE $M_\sigma^m(o)$

It is well known that a model space refers to the Riemannian manifold given by the warped product

$$M_\sigma^m(o) = \left((0, +\infty) \times S^{m-1}, dr^2 + \sigma^2(r) g^{S^{m-1}} \right),$$

where $(S^{m-1}, g^{S^{m-1}})$ is the standard $(m-1)$ -dimensional unit sphere, and $\sigma : (0, \infty) \rightarrow (0, \infty)$ is a smooth function.

This model space plays a very important role in mathematics and physics, and it includes all three space forms \mathbb{R}^m , S^m , and H^m as special cases corresponding to $\sigma(r) = r$, $\sin r$, and $\sinh r$ respectively.

In this section we will show that the biharmonic equation for radial (or rotationally symmetric) solutions on the model space $M_\sigma^m(o) = (\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$ is completely solvable. We give an integral formula for the solutions which are used to produce families of infinitely many proper biharmonic functions on space forms.

Lemma 2.1. *The Laplacian and the bi-Laplacian operators on the model space $M_\sigma^m(o) = (\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$ can be described as*

$$(36) \quad \Delta_\sigma f = f_{rr} + (m-1) \frac{\sigma'}{\sigma} f_r + \frac{1}{\sigma^2} \Delta_{S^{m-1}} f$$

$$(37) \quad \Delta_\sigma^2 f = (\Delta_\sigma f)_{rr} + (m-1) \frac{\sigma'}{\sigma} (\Delta_\sigma f)_r + \frac{1}{\sigma^2} \Delta_{S^{m-1}} (\Delta_\sigma f).$$

In particular, for a radial function $f(r, \theta) = f(r)$, we have

$$(38) \quad \Delta_\sigma f = f'' + (m-1) \frac{\sigma'}{\sigma} f'$$

$$(39) \quad \Delta_\sigma^2 f = (\Delta_\sigma f)'' + (m-1) \frac{\sigma'}{\sigma} (\Delta_\sigma f)'$$

Proof. Let h denote the standard metric on the sphere S^{m-1} and $g = dr^2 + \sigma^2(r)h$ be the warped product metric. $\bar{\Gamma}_{ab}^c$ (respectively Γ_{ij}^k) the connection coefficients

of g and h respectively. We use a local coordinate system $\{r, u^i\}$ on $M_\sigma^m(o)$ where $\{u^i\}$ is a local coordinate system on S^{m-1} . We will also use the following indices and their ranges: $a, b, c, d = 0, 1, 2, \dots, m-1$ $i, j, k, l = 1, 2, \dots, m-1$ for tensors on $(M_\sigma^m(o), g = dr^2 + \sigma^2(r)h)$ and (S^{m-1}, h) respectively. Then, it is easily checked that

$$g_{00} = g^{00} = 1, \quad g_{0i} = g^{0i} = 0, \quad g_{ij} = \sigma^2(r)h_{ij}, \quad g^{ij} = \sigma^{-2}h^{ij}.$$

By a straightforward computation we have, for a function $f = f(x_0, x_1, \dots, x_{m-1})$ with $(x_1, \dots, x_{m-1}) = (u_1, \dots, u_{m-1}) \in S^{m-1}$,

$$\begin{aligned} (40) \quad \Delta_\sigma f &= g^{ab}f_{ab} - g^{ab}\bar{\Gamma}_{ab}^c f_c = g^{00}f_{00} + g^{ij}f_{ij} - g^{00}\bar{\Gamma}_{00}^c f_c - g^{ij}\bar{\Gamma}_{ij}^c f_c \\ &= g^{00}f_{00} + g^{ij}f_{ij} - g^{00}\bar{\Gamma}_{00}^0 f_0 - g^{00}\bar{\Gamma}_{00}^k f_k - g^{ij}\bar{\Gamma}_{ij}^0 f_0 - g^{ij}\bar{\Gamma}_{ij}^k f_k \\ &= f_{rr} + g^{ij}f_{ij} - \bar{\Gamma}_{00}^k f_k - g^{ij}\bar{\Gamma}_{ij}^0 f_0 - g^{ij}\bar{\Gamma}_{ij}^k f_k \\ &= f_{rr} + (m-1)\frac{\sigma'}{\sigma}f_r + \frac{1}{\sigma^2}\Delta_{S^{m-1}}f \end{aligned}$$

where in obtaining the last equality we have used the identities of the connection coefficients of the warped product metrics

$$\bar{\Gamma}_{ij}^0 = -\sigma\sigma' h_{ij}, \quad \bar{\Gamma}_{ij}^k = \Gamma_{ij}^k.$$

$$\Delta_\sigma^2 f = (\Delta_\sigma f)_{rr} + (m-1)\frac{\sigma'}{\sigma}(\Delta_\sigma f)_r + \frac{1}{\sigma^2}\Delta_{S^{m-1}}(\Delta_\sigma f).$$

For radial function $f(r, u) = f(r)$, we have $\Delta_{S^{m-1}}f = 0$ and hence the formulas (38) and (39) follow. \square

As a straightforward application we obtain the following complete description of all radial biharmonic functions on the model space $(M_\sigma^m(o), g = dr^2 + \sigma^2(r)g^{S^{m-1}})$.

Theorem 2.2. *A radial function $f(r)$ on the model space $(\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$ is biharmonic if and only if*

$$\begin{aligned} (41) \quad f(r) &= c_4 + c_3 \int \frac{1}{\sigma^{m-1}} dr \\ &+ c_2 \left[\int \frac{1}{\sigma^{m-1}} dr \int \sigma^{m-1} dr - \int \left(\int \frac{1}{\sigma^{m-1}} dr \right) \sigma^{m-1} dr \right] \\ &+ c_1 \left[\int \frac{1}{\sigma^{m-1}} dr \int \left(\int \frac{1}{\sigma^{m-1}} dr \right) \sigma^{m-1} dr - \int \left(\int \frac{1}{\sigma^{m-1}} dr \right)^2 \sigma^{m-1} dr \right]. \end{aligned}$$

Proof. By (39), a radial function $f(r, u) = f(r)$ is biharmonic on $M_\sigma^m(o)$ if and only if $y = \Delta_\sigma f$ is a solution of the second order ordinary homogeneous differential equation

$$(42) \quad y'' + (m-1)\frac{\sigma'}{\sigma}y' = 0.$$

It is easy to check that the general solutions of (42) are

$$(43) \quad y = c_1y_1 + c_2y_2, \quad \text{with } y_1 = \int \frac{1}{\sigma^{m-1}}dr, \quad y_2 = -1.$$

Thus, the biharmonicity of $f(r, u) = f(r)$ leads to the inhomogeneous 2nd order ODE

$$(44) \quad \Delta f = c_1y_1 + c_2y_2, \quad \text{with } y_1 = \int \frac{1}{\sigma^{m-1}}dr, \quad y_2 = -1, \quad \text{or}$$

$$(45) \quad f'' + (m-1)\frac{\sigma'}{\sigma}f' = c_1y_1 + c_2y_2, \quad \text{with } y_1 = \int \frac{1}{\sigma^{m-1}}dr, \quad y_2 = -1.$$

Note that the homogeneous part of this equation is exactly (42) whose general solution is known to be (43). By the theory of the general solutions of linear ordinary differential equations, we conclude that the general solutions of (44) can be written as

$$(46) \quad f = Ay_1 + By_2 + y_{p_1} + y_{p_2},$$

where y_{p_1} and y_{p_2} are a particular solution of

$$(47) \quad y'' + (m-1)\frac{\sigma'}{\sigma}y' = c_1 \int \frac{1}{\sigma^{m-1}}dr, \quad \text{and}$$

$$(48) \quad y'' + (m-1)\frac{\sigma'}{\sigma}y' = c_2,$$

respectively. By using the method of variation of parameters (see e.g., [19], §4.6) we find that

$$(49) \quad y_{p_1} = c_1 \left(y_1 \int y_1 \sigma^{m-1} dr - \int y_1^2 \sigma^{m-1} dr \right)$$

$$(50) \quad y_{p_2} = c_2 \left(y_1 \int \sigma^{m-1} dr - \int y_1 \sigma^{m-1} dr \right).$$

$$(51)$$

Therefore, a radial function $f(r)$ is biharmonic function on the model space $(M_\sigma^m(o) = (\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}}))$ if and only if

$$(52) \quad f(r) = c_3 y_1 + c_4 + c_1 \left(y_1 \int y_1 \sigma^{m-1} dr - \int y_1^2 \sigma^{m-1} dr \right)$$

$$(53) \quad + c_2 \left(y_1 \int \sigma^{m-1} dr - \int y_1 \sigma^{m-1} dr \right),$$

where $y_1 = \int \frac{1}{\sigma^{m-1}} dr$. From this we obtain the integral formula for biharmonic functions on the model space. \square

Applying Theorem 2.2 with $\sigma(r) = r, \sin r, \sinh r$ respectively we obtain the following descriptions of radial biharmonic functions on space forms.

Proposition 2.3. (i) On the Euclidean space $(\mathbb{R}^m \setminus \{0\}, dr^2 + r^2 g^{S^{m-1}})$, a radial functions $f(r)$ is a biharmonic if and only if

$$(54) \quad f(r) = \begin{cases} c_4 + c_3 \ln r + c_2 r^2 + c_1 r^2 \ln r, & m = 2, \\ c_4 + c_3 r^{-2} + c_2 r^2 + c_1 \ln r, & m = 4, \text{ and} \\ c_4 + c_3 r^{2-m} + c_2 r^2 + c_1 r^{4-m}, & m \neq 2, 4. \end{cases}$$

(ii) On the sphere $(S^m, dr^2 + \sin^2 r g^{S^{m-1}})$, a radial function $f(r)$ is a biharmonic if and only if

$$(55)$$

$$\begin{aligned} f(r) &= c_4 + c_3 \int \frac{1}{\sin^{m-1} r} dr \\ &+ c_2 \left[\int \frac{1}{\sin^{m-1} r} dr \int \sin^{m-1} r dr - \int \left(\int \frac{1}{\sin^{m-1} r} dr \right) \sin^{m-1} r dr \right] \\ &+ c_1 \left[\int \frac{1}{\sin^{m-1} r} dr \int \left(\int \frac{1}{\sin^{m-1} r} dr \right) \sin^{m-1} r dr - \int \left(\int \frac{1}{\sin^{m-1} r} dr \right)^2 \sin^{m-1} r dr \right]. \end{aligned}$$

(iii) On the hyperbolic space $(H^m, dr^2 + \sinh^2 r g^{S^{m-1}})$, a radial function $f(r)$ is a biharmonic if and only if

$$(56)$$

$$\begin{aligned} f(r) &= c_4 + c_3 \int \frac{1}{\sinh^{m-1} r} dr \\ &+ c_2 \left[\int \frac{1}{\sinh^{m-1} r} dr \int \sinh^{m-1} r dr - \int \left(\int \frac{1}{\sinh^{m-1} r} dr \right) \sinh^{m-1} r dr \right] \\ &+ c_1 \left[\int \frac{1}{\sinh^{m-1} r} dr \int \left(\int \frac{1}{\sinh^{m-1} r} dr \right) \sinh^{m-1} r dr - \int \left(\int \frac{1}{\sinh^{m-1} r} dr \right)^2 \sinh^{m-1} r dr \right]. \end{aligned}$$

Note that for $m = 3$ we obtain the following examples which was obtained in [3]

Example 3. (I) A radial function $f(r)$ on 3-sphere $(S^3, dr^2 + \sin^2 r g^{S^2})$ is biharmonic if and only if

$$(57) \quad f(r) = a_1 + a_2 \cot r + a_3 r + a_4 r \cot r,$$

where a_i are arbitrary constants.

(II) A radial function $f(r)$ on hyperbolic space $(H^3, dr^2 + \sinh^2 r g^{S^2})$ is biharmonic if and only if

$$(58) \quad f(r) = a_1 + a_2 \coth r + a_3 r + a_4 r \coth r,$$

where a_i are arbitrary constants.

In Fact, using Proposition 2.3 with $m = 3$ we know that a radial function $f(r)$ on $(S^3, dr^2 + \sin^2 r g^{S^2})$ is biharmonic if and only if

$$(59) \quad \begin{aligned} f(r) = & c_4 + c_3 \int \frac{1}{\sin^2 r} dr \\ & c_2 \left[\int \frac{1}{\sin^2 r} dr \int \sin^2 r dr - \int \left(\int \frac{1}{\sin^2 r} dr \right) \sin^2 r dr \right] \\ & + c_1 \left[\int \frac{1}{\sin^2 r} dr \int \left(\int \frac{1}{\sin^2 r} dr \right) \sin^2 r dr - \int \left(\int \frac{1}{\sin^2 r} dr \right)^2 \sin^2 r dr \right]. \end{aligned}$$

A straightforward computation gives

$$(60) \quad f(r) = c_4 - c_3 \cot r - \frac{c_2}{2} r \cot r - \frac{c_1}{2} (r + \cot r)$$

$$(61) \quad = c_4 - (c_3 + \frac{c_1}{2}) \cot r - \frac{c_2}{2} r \cot r - \frac{c_1}{2} r,$$

from which we obtained Statement (I). A similar proof gives Statement (II)

Remark 4. (i) Note that in the family (57) $a_1 + a_2 \cot r$ are harmonic and $a_3 r + a_4 r \cot r$ are proper biharmonic on S^3 . More precisely, $a_4 r \cot r$ is a family of quasi-harmonic functions (i.e., $\Delta^{S^3}(a_4 r \cot r) = \text{constant} \neq 0$) and the function $a_3 r$ is proper biharmonic. Similarly, in the family (58) $a_1 + a_2 \coth r$ are harmonic and $a_3 r + a_4 r \coth r$ are proper biharmonic on H^3 . Furthermore, $a_4 r \coth r$ is quasi-harmonic and $a_3 \mathbb{R}$ is proper biharmonic.

(ii) An interesting result proved in [3] states that the distance function $d(r) = r$ is proper biharmonic in a Riemannian manifold (M^m, g) if and only if $m = 1$, or $m = 3$ and the manifold is a space form. Our example above and Proposition

2.3 (i) confirm that the function $u(r) = r$ in 3-dimensional space form is indeed proper biharmonic.

Corollary 2.4. (A) A radial function $f(r)$ on the sphere $(S^2, dr^2 + \sin^2 r d\theta^2)$ is biharmonic if and only if

$$(62) \quad f(r) = a_1 + a_2 \ln \tan \frac{r}{2} + a_3 \ln \sin r \\ + a_4 \left[\ln \sin r \ln \tan \frac{r}{2} - 2 \int \cot r \ln \tan \frac{r}{2} dr \right],$$

where a_i are arbitrary constants. In particular, $f(r) = \ln \sin r$ is a proper biharmonic function.

(B) A radial function $f(r)$ on the hyperbolic space $(H^2, dr^2 + \sinh^2 r d\theta^2)$ is biharmonic if and only if

$$(63) \quad f(r) = a_1 + a_2 \ln \tanh \frac{r}{2} + a_3 \ln \sinh r \\ + a_4 \left[\ln \sinh r \ln \tanh \frac{r}{2} - 2 \int \coth r \ln \tanh \frac{r}{2} dr \right],$$

where a_i are arbitrary constants. In particular, $f(r) = \ln \sinh r$ is a proper biharmonic function.

Proof. Applying formula (55) with $m = 2$ we have

$$(64) \quad f(r) = c_4 + c_3 \int \frac{1}{\sin r} dr \\ + c_2 \left[\int \frac{1}{\sin r} dr \int \sin r dr - \int \left(\int \frac{1}{\sin r} dr \right) \sin r dr \right] \\ + c_1 \left[\int \frac{1}{\sin r} dr \int \left(\int \frac{1}{\sin r} dr \right) \sin r dr - \int \left(\int \frac{1}{\sin r} dr \right)^2 \sin r dr \right].$$

A straightforward calculation yields

$$(65) \quad f(r) = (c_4 + c_2 \ln 2) + (c_3 - c_1 \ln 2) \ln \tan \frac{r}{2} - c_2 \ln \sin r \\ + c_1 \left[\ln \sin r \ln \tan \frac{r}{2} - 2 \int \cot r \ln \tan \frac{r}{2} dr \right].$$

Relabeling the arbitrary constants we obtain (62). A similar proof gives Statement (B). \square

Radial biharmonic functions on space forms with conformally flat models: One can easily check that by performing a coordinate change $r = 2 \tan^{-1} t$ the warped product model for a sphere $(S^m, dr^2 + \sin^2 r g^{S^{m-1}})$ can be

written as the conformally flat model $\left(S^m, \frac{4(dt^2+t^2g^{S^{m-1}})}{(1+t^2)^2}\right)$. Similarly, the hyperbolic space $(H^m, dr^2 + \sinh^2 r g^{S^{m-1}})$ as can be written as $\left(H^m, \frac{4(dt^2+t^2g^{S^{m-1}})}{(1-t^2)^2}\right)$.

Using these models and the transformations $r = 2 \tan^{-1} t$ and $r = 2 \tanh^{-1} t$ respectively, Proposition 2.3 reads

Proposition 2.5. (i) A radial function $u : \left(S^m, \frac{4(dt^2+t^2g_{S^{m-1}})}{(1+t^2)^2}\right) \rightarrow \mathbb{R}$, $u = u(t)$ on the sphere is biharmonic if and only if

$$(66) \quad u(t) = c_1 + c_2 u_2 + c_3 \left[u_2 \int \frac{t^{m-1}}{(1+t^2)^m} dt - \int \frac{u_2 t^{m-1}}{(1+t^2)^m} dt \right] \\ + c_4 \left[u_2 \int \frac{u_2 t^{m-1}}{(1+t^2)^m} dr - \int \frac{u_2^2 t^{m-1}}{(1+t^2)^m} dt \right],$$

where $u_2 = \int \frac{(1+t^2)^{m-2}}{t^{m-1}} dt$.

(ii) A radial function $u : \left(H^m, \frac{4(dt^2+t^2g_{S^{m-1}})}{(1-t^2)^2}\right) \rightarrow \mathbb{R}$, $u = u(t)$ on the hyperbolic space is biharmonic if and only if

$$(67) \quad u(t) = c_1 + c_2 u_2 + c_3 \left[u_2 \int \frac{t^{m-1}}{(1-t^2)^m} dr - \int \frac{u_2 t^{m-1}}{(1-t^2)^m} dt \right] \\ + c_4 \left[u_2 \int \frac{u_2 t^{m-1}}{(1-t^2)^m} dt - \int \frac{u_2^2 t^{m-1}}{(1-t^2)^m} dr \right],$$

where $u_2 = \int \frac{(1-t^2)^{m-2}}{t^{m-1}} dr$.

A straightforward checking using (66) and (67) with $m = 2$ respectively we recover the following radial biharmonic functions on S^2 and H^2 which was obtained in [7].

Corollary 2.6. (see also [7]) (i) A radial function $u : \left(S^2, \frac{4(dt^2+t^2d\theta^2)}{(1+t^2)^2}\right) \rightarrow \mathbb{R}$ with $u = u(t)$ is biharmonic if and only if

$$(68) \quad u(t) = c_1 + c_2 \ln t + c_3 \ln(1+t^2) + c_4 \left[\ln t \ln(1+t^2) - 2 \int \frac{\ln(1+t^2)}{t} dt \right].$$

(ii) A radial function $u : \left(H^2, \frac{4(dt^2+t^2d\theta^2)}{(1-t^2)^2}\right) \rightarrow \mathbb{R}$ with $u = u(t)$ is biharmonic if and only if

$$(69) \quad u(t) = c_1 + c_2 \ln t + c_3 \ln(1-t^2) + c_4 \left[\ln t \ln(1-t^2) - 2 \int \frac{\ln(1-t^2)}{t} dt \right].$$

Example 4. The function $u(t) = \arctan t$ is a proper biharmonic function on $\left(S^3, \frac{4(dt^2+t^2g^{S^2}}{(1+t^2)^2}\right) \rightarrow \mathbb{R}$. Note that in this model, one can easily described the

function by using the Cartesian coordinates from \mathbb{R}^4 in which S^3 lies. In fact, this function can be viewed as $\mathbb{R}^4 \supset S^3 \rightarrow \mathbb{R}$ and $u(x) = \arctan \sqrt{\frac{1+x_4}{1-x_4}}$ for any $(x_1, x_2, x_3, x_4) \in S^3 \setminus \{N\}$.

Note that radial quasi-harmonic functions on Poincaré n-ball $B_\alpha^n = (D^n, (1 - |x|^2)^{2\alpha} g_0)$ were studied in [11] and [12], where the authors proved that there exist bounded quasi-harmonic functions on B_α^n if and only if $\alpha \in (-1, \frac{1}{n-2})$.

3. BIHARMONIC FUNCTIONS ON MODEL SPACES VIA SPHERICAL HARMONIC FUNCTIONS

In their study of the generalized Almansi property the authors in [8] explored the possibility of constructing biharmonic functions on the model space $M_\sigma^m(o) = (\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$ via the products of a radial function and a local harmonic function. It turns out that the only model space that has this property is the Euclidean space.

In this section, we will prove that the model space $M_\sigma^m(o)$ always admits proper biharmonic functions $f(r, \theta) = u(r)v_k(\theta)$ as a product of any eigenfunction on the factor sphere S^{m-1} and certain radial functions.

This provides an effective way to construct infinitely many proper biharmonic functions on the model space including the three space forms.

Theorem 3.1. *For any eigenfunction $v_k : S^{m-1} \rightarrow \mathbb{R}$ there exist radial functions $u(r)$ such that the product $p(r, \theta) = u(r)v_k(\theta)$ is a biharmonic functions on the model space $M_\sigma^m(o) = (\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$. In particular, there are infinitely many proper biharmonic functions on space forms given by the product of any eigenfunctions with some radial functions.*

Proof. Using polar coordinates $(r, \theta) \in \mathbb{R}^+ \times S^{m-1} = \mathbb{R}^m \setminus \{0\}$ and (36), we have the Laplacian on the function $f(r, \theta) = u(r)v_k(\theta)$ on $M_\sigma^m(o)$ as

$$(70) \quad \Delta f(r, \theta) = [u'' + (m-1)\frac{\sigma'}{\sigma}u']v_k(\theta) + \frac{u}{\sigma^2}\Delta^{S^{m-1}}(v_k(\theta)).$$

Since $v_k(\theta)$ is a λ_k -eigenfunction on S^{m-1} (i.e., a restriction of a harmonic homogeneous polynomial of degree k in \mathbb{R}^m to S^{m-1}), we have

$$(71) \quad \Delta_{S^{m-1}}v_k(\theta) = -k(m+k-2)v_k(\theta)$$

and hence (70) becomes

$$(72) \quad \Delta f(r, \theta) = \left[u'' + (m-1) \frac{\sigma'}{\sigma} u' - \frac{k(m+k-2)}{\sigma^2} u \right] v_k(\theta).$$

Thus, a function $f(r, \theta) = u(r)v_k(\theta)$ is harmonic on the model space $M_\sigma^m(o)$ if and only if the radial function $u(r)$ solves the 2nd order linear homogeneous equation

$$(73) \quad u'' + (m-1) \frac{\sigma'}{\sigma} u' - \frac{k(m+k-2)}{\sigma^2} u = 0.$$

By the well known existence theory of linear differential equations (see e.g., [19], Theorem 4.14), we have the general solution of (73) given by $u = c_1 u_1 + c_2 u_2$.

This means that $f(r, \theta) = u(r)v_k(\theta)$ is harmonic if and only if $f(r, \theta) = (c_1 u_1(r) + c_2 u_2(r))v_k(\theta)$.

On the other hand, recall that $p(r, \theta) = u(r)v_k(\theta)$ is biharmonic on the model space if and only if $\Delta p(r, \theta)$ is harmonic on the model space. Looking for biharmonic function $p(r, \theta) = u(r)v_k(\theta)$ with $\Delta p(r, \theta) = (c_1 u_1 + c_2 u_2)v_k(\theta)$ and using (72) we are lead to the following non-homogeneous 2nd order differential equation

$$(74) \quad u'' + (m-1) \frac{\sigma'}{\sigma} u' - \frac{k(m+k-2)}{\sigma^2} u = c_1 u_1 + c_2 u_2.$$

By the method of variation of parameters (see, e.g., [19] §4.6) we have the general solution of (74) given by

$$(75) \quad u(r) = c_1 u_1(r) + c_2 u_2(r) + c_3 u_{p_1}(r) + c_4(r) u_{p_2}(r),$$

where

$$(76) \quad u_{p_1}(r) = u_1 \int \frac{-u_1 u_2}{u_1 u_2' - u_1' u_2} dr + u_2 \int \frac{u_1^2}{u_1 u_2' - u_1' u_2} dr, \text{ and}$$

$$(77) \quad u_{p_2}(r) = u_1 \int \frac{-u_2^2}{u_1 u_2' - u_1' u_2} dr + u_2 \int \frac{u_1 u_2}{u_1 u_2' - u_1' u_2} dr.$$

Summarizing the above, we conclude that for any eigenfunction $v_k(\theta)$ on S^{m-1} the product $p(r, \theta) = u(r)v_k(\theta)$ with $u(r)$ given by (75) is a biharmonic function on the model space $(\mathbb{R}^+ \times S^{m-1}, dr^2 + \sigma^2(r)g^{S^{m-1}})$. This completes the proof of the theorem. □

Theorem 3.1 (Formula (75)) provides an effective method to construct biharmonic functions on model space including the three space forms.

Corollary 3.2. *On the punctured Euclidean plane $(\mathbb{R}^2 \setminus \{0\}, dr^2 + r^2 d\theta^2)$, each function in the family*

$$(78) \quad p(r, \theta) = (c_3 + c_4 r^4) v_2(\theta)$$

is a proper biharmonic function. In particular,

$p(r, \theta) = a \cos 2\theta + b \sin 2\theta = \frac{a(x^2 - y^2) + b(xy)}{x^2 + y^2}$ is a family of bounded proper biharmonic functions on the punctured Euclidean plane.

Proof. Using Theorem 3.1 with $m = 2, k = 2, \sigma(r) = r$ Equation (73) becomes

$$(79) \quad u'' + \frac{1}{r} u' - \frac{4}{r^2} u = 0$$

which has the fundamental set of solutions $u_1(r) = r^2, u_2(r) = r^{-2}$. Using formulas (76) and (77) we have

$$u_{p_1}(r) = \frac{r^4}{12}, \quad u_{p_2}(r) = -\frac{1}{4}.$$

Thus, we obtain the family (78) of proper biharmonic functions. Recalling that $v_2(\theta)$ is the restriction of a harmonic homogeneous polynomial of degree 2, so we have

$$(80) \quad v_2(\theta) = [a(x^2 - y^2) + bxy]_{|S^1} = \frac{a(x^2 - y^2) + b(xy)}{x^2 + y^2} = a \cos 2\theta + b \sin 2\theta,$$

which is clearly a bounded function on the puncture plane $\mathbb{R}^2 \setminus \{0\}$. \square

Remark 5. (i) The interesting problem of classifying all bounded biharmonic functions on a punctured Euclidean space was studied in [10], where the authors obtained the following nice classification.

Theorem ([10]): The linear space $BH^2(\mathbb{R}^m \setminus \{0\})$ of bounded biharmonic functions on $\mathbb{R}^m \setminus \{0\}$ is

$$(81) \quad BH^2(\mathbb{R}^m \setminus \{0\}) = \begin{cases} \text{span}\{1, \cos 2\theta, \sin 2\theta\}, & \text{for } m = 2; \\ \text{span}\{1, \cos r \cos \theta, \sin r \sin \theta, \cos r\} & \text{for } m = 3; \\ \text{Span}\{1\}, & \text{for } m \geq 4. \end{cases}$$

(ii) Our Corollary 3.2 does confirm that the functions $\cos 2\theta, \sin 2\theta$ are indeed bounded proper biharmonic on the punctured plane.

Example 5. On the sphere $(S^2 \setminus \{\pm N\}, dr^2 + \sin^2 r d\theta^2)$, each function in the family

$$(82) \quad p(r, \theta) = \left[c_3 \left(-\sin^2 \frac{r}{2} \left(\cot \frac{r}{2} + \tan \frac{r}{2} \right) + 2 \tan \frac{r}{2} \ln \sin \frac{r}{2} \right) \right. \\ \left. + c_4 \left(\sin^2 \frac{r}{2} \left(\cot \frac{r}{2} + \tan \frac{r}{2} \right) + 2 \cot \frac{r}{2} \ln \cos \frac{r}{2} \right) \right] v_1(\theta)$$

is a proper biharmonic function. In particular,

$p(r, \theta) = (2 \tan \frac{r}{2} \ln \sin \frac{r}{2} + 2 \cot \frac{r}{2} \ln \cos \frac{r}{2})v_1(\theta)$ is a proper biharmonic function on 2-sphere.

In fact, using Theorem 3.1 with $m = 2, k = 1, \sigma(r) = \sin r$, Equation (73) becomes

$$(83) \quad u'' + \cot r u' - \csc^2 r u = 0$$

which has the general solution $u(r) = c_1 u_1(r) + c_2 u_2(r)$ for $u_1 = \cot \frac{r}{2}, u_2 = \tan \frac{r}{2}$. Using formulas (76) and (77) we have

$$\begin{aligned} u_{p_1}(r) &= \frac{1}{2} \cot \frac{r}{2} - \sin^2 \frac{r}{2} (\cot \frac{r}{2} + \tan \frac{r}{2}) + 2 \tan \frac{r}{2} \ln \sin \frac{r}{2}, \\ u_{p_2}(r) &= -\frac{1}{2} \tan \frac{r}{2} + \sin^2 \frac{r}{2} (\cot \frac{r}{2} + \tan \frac{r}{2}) + 2 \cot \frac{r}{2} \ln \cos \frac{r}{2}. \end{aligned}$$

Noting that the first term in both $u_{p_1}(r)$ and $u_{p_2}(r)$ above are harmonic, by discarding them from the families we obtain the claimed family of proper biharmonic functions. The particular one is obtained by choosing $c_3 = c_4 = 1$.

Similarly, by using Theorem 3.1 with $m = 2, k = 1, \sigma(r) = \sinh r$ we have

Example 6. On the hyperbolic plane $(H^2, dr^2 + \sinh^2 r d\theta^2)$, each function in the family

$$(84) \quad \begin{aligned} p(r, \theta) &= [c_3(-\sinh^2 \frac{r}{2} (\coth \frac{r}{2} - \tanh \frac{r}{2}) + 2 \tanh \frac{r}{2} \ln \sinh \frac{r}{2}) \\ &+ c_4(-\sinh^2 \frac{r}{2} (\coth \frac{r}{2} - \tanh \frac{r}{2}) + 2 \coth \frac{r}{2} \ln \cosh \frac{r}{2})] v_1(\theta) \end{aligned}$$

is a proper biharmonic function. In particular,

$p(r, \theta) = (2 \tanh \frac{r}{2} \ln \sinh \frac{r}{2} - 2 \coth \frac{r}{2} \ln \cosh \frac{r}{2})v_1(\theta)$ is a proper biharmonic function on the hyperbolic plane H^2 .

Example 7. On the sphere $(S^2 \setminus \{\pm N\}, dr^2 + \sin^2 r d\theta^2)$, each function in the family

$$(85) \quad \begin{aligned} p(r, \theta) &= [c_3(1 + \cos^2 \frac{r}{2} - \sin^2 \frac{r}{2} \tan^2 \frac{r}{2} + 4 \tan^2 \frac{r}{2} \ln \sin \frac{r}{2}) \\ &+ c_4(\cos^2 \frac{r}{2} - \sin^2 \frac{r}{2} \tan^2 \frac{r}{2} + 2 \cot^2 \frac{r}{2} \ln \cos \frac{r}{2})] v_2(\theta) \end{aligned}$$

is a proper biharmonic function. In particular,

$p(r, \theta) = (1 + 4 \tan^2 \frac{r}{2} \ln \sin \frac{r}{2} - 2 \cot^2 \frac{r}{2} \ln \cos \frac{r}{2})v_2(\theta)$ is a proper biharmonic function on 2-sphere.

In fact, with $m = k = 2, \sigma(r) = \sin r$ Equation (73) becomes

$$(86) \quad u'' + \cot r u' - 4 \csc^2 r u = 0$$

which has a set of fundamental solutions $u_1 = \cot^2 \frac{r}{2}$, $u_2 = \tan^2 \frac{r}{2}$. Using formulas (76) and (77) and discarding the harmonic parts we have

$$\begin{aligned} u_{p_1}(r) &= 1 + \cos^2 \frac{r}{2} - \sin^2 \frac{r}{2} \tan^2 \frac{r}{2} + 4 \tan^2 \frac{r}{2} \ln \sin \frac{r}{2}, \\ u_{p_2}(r) &= \cos^2 \frac{r}{2} - \sin^2 \frac{r}{2} \tan^2 \frac{r}{2} + 2 \cot^2 \frac{r}{2} \ln \cos \frac{r}{2}, \end{aligned}$$

which gives the claimed family of proper biharmonic functions on S^2 . The particular one is obtained by choosing $c_3 = -c_4 = 1$.

4. BIHARMONIC FUNCTIONS ON PUNCTURED EUCLIDEAN SPACES

For harmonic functions on a Euclidean space we have the following well known Liouville type theorems.

Liouville Theorem: Any bounded harmonic function on \mathbb{R}^m is constant.

Generalization 1: Any positive harmonic function on \mathbb{R}^m is constant.

Generalization 2 (see e.g., [1], Corollaries 3.3, 3.14): Any positive harmonic function on $u : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}$ takes the form

$$(87) \quad u(x) = \begin{cases} \text{constant}, & m = 2 \\ a + b|x|^{-(m-2)}, & \text{for } a, b > 0, \quad m > 2. \end{cases}$$

For biharmonic functions on a Euclidean space we have

Liouville Theorem ([6]): Any bounded biharmonic function on \mathbb{R}^m is constant.

Fact: There are positive biharmonic function on \mathbb{R}^m which is NOT constant, e.g., $u(x) = 1 + |x|^2$.

For bounded biharmonic functions on the punctured Euclidean space we have **Theorem** ([10]): The linear space $BH^2(\mathbb{R}^m \setminus \{0\})$ of bounded biharmonic functions on $\mathbb{R}^m \setminus \{0\}$ is

$$BH^2(\mathbb{R}^m \setminus \{0\}) = \begin{cases} \text{span}\{1, \cos 2\theta, \sin 2\theta\}, & \text{for } m = 2; \\ \text{span}\{1, \cos r \cos \theta, \sin r \sin \theta, \cos r\} & \text{for } m = 3; \\ \text{Span}\{1\}, & \text{for } m \geq 4. \end{cases}$$

In particular, any bounded biharmonic function on $\mathbb{R}^m \setminus \{0\}$ with $m \geq 4$ is constant.

Regarding biharmonic functions on $u : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}$ we also have

Almansi Theorem (1899): Any biharmonic function u on $\mathbb{R}^m \setminus \{0\}$ (or more generally, a star-shaped region) takes the form

$$(88) \quad u = h_1 + |x|^2 h_2,$$

where h_1, h_2 are two harmonic functions.

A little more detailed description is the following

Lemma A ([13], **Lemma 2.2**): Any biharmonic function u on $\mathbb{R}^m \setminus \{0\}$ takes the form

$$(89) \quad u = \begin{cases} h_1 + |x|^2 h_2 + (ax_1 + bx_2) \ln |x|, & \text{for } m = 2; \\ h_1 + |x|^2 h_2 + a \ln |x|, & \text{for } m = 4; \\ h_1 + |x|^2 h_2, & \text{for } m \neq 2, 4. \end{cases}$$

where a, b are arbitrary constants and h_1, h_2 are two arbitrary harmonic functions.

Note that both formulas for biharmonic functions given in (88) and (89) depend on two arbitrary harmonic functions. To the author's knowledge, no further classification has been found in the literature.

Here, we give a complete classification of biharmonic functions with condition $\Delta u > 0$ on the punctured Euclidean space.

Theorem 4.1. *A function $u : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}$ is biharmonic with $\Delta u > 0$ if and only if*

$$(90) \quad u(x) = \begin{cases} c_1 + c_2 \ln |x| + a|x|^2, & \text{for } m = 2; \\ c_1 + c_2|x|^{-2} + a|x|^2 + b \ln |x|, & \text{for } m = 4; \\ c_1 + c_2|x|^{-(m-2)} + a|x|^2 + b|x|^{-(m-4)}, & \text{for } m \neq 2, 4, \end{cases}$$

where c_i are arbitrary constants, for $m = 2, 3, 4$, $a, b \geq 0$, $a^2 + b^2 \neq 0$, and for $m \neq 2, 3, 4$, the constants $a \geq 0$, $b \leq 0$, $a^2 + b^2 \neq 0$.

Proof. Note that $u : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}$ is biharmonic with $\Delta u > 0$ means that Δu is a positive harmonic function on $\mathbb{R}^m \setminus \{0\}$. So, by (87), we have

$$(91) \quad \Delta u = \begin{cases} C, & m = 2 \\ a + b|x|^{-(m-2)}, & \text{for } a, b > 0, \quad m > 2. \end{cases}$$

Using polar coordinates $x = (r, \theta) \in \mathbb{R}^+ \times S^{m-1} \equiv \mathbb{R}^m \setminus \{0\}$ with $|x| = r, \theta = \frac{x}{|x|} \in S^{m-1}$. Then it is known (see e.g., [14], § 22.3) that for any smooth function $u : \mathbb{R}^m \setminus \{0\} \rightarrow \mathbb{R}$ is a limit of a sequence $f_n = \sum_{i=1}^{t_n} a_{ni}(r)b_{ni}(\theta)$, where $b_{ni}(\theta) \in C^\infty(S^{m-1})$. Using the well-known fact that $C^\infty(S^{m-1}) \subset L^2(S^{m-1}) = \bigoplus_{i=0}^\infty \mathcal{H}^i$ and the bases $\{v_{k1}, v_{k2}, \dots, v_{kn_k}\}$ for \mathcal{H}^k , we can express f_n as

$$(92) \quad f_n = \sum_{k=0}^{\infty} \sum_{i=1}^{n_k} u_{ki}^n(r)v_{ki}(\theta).$$

Since u and f_n are closed in C^∞ topology for $n \geq N$, we conclude that u is biharmonic function with $\Delta u > 0$, then so is f_n a biharmonic function with $\Delta f_n > 0$ for $n \geq N$.

Using (38) with $\sigma = r$ and the fact that $\Delta^{S^{m-1}}(u_{ki}(r)v_{ki}(\theta)) = -k(m+k-2)u_{ki}v_{ki}$, we have

$$(93) \quad \Delta f_n = \sum_{k=0}^{\infty} \sum_{i=1}^{n_k} [(u_{ki}^n)'' + (m-1)r^{-1}(u_{ki}^n)' - k(m+k-2)r^{-2}u_{ki}^n] v_{ki}.$$

It follows that for any n , f_n is a solution of (91) if and only if

$$(94) \quad (u_{ki}^n)'' + (m-1)r^{-1}(u_{ki}^n)' - k(m+k-2)r^{-2}u_{ki}^n = 0, \quad \forall k > 0, i, \text{ and}$$

$$(95) \quad \begin{aligned} & (u_{01}^n)'' + (m-1)r^{-1}(u_{01}^n)' - k(m+k-2)r^{-2}u_{01}^n \\ &= \begin{cases} C, & m = 2 \\ a + b|x|^{-(m-2)}, & \text{for } a, b > 0, \quad m > 2. \end{cases} \end{aligned}$$

One can easily check that the general solution of Equation (94) is $u_{ki}^n(r) = c_1 r^k + c_2 r^{-(m+k-2)}$ which means $u_{ki}^n(r)v_{ki}$ is a harmonic function for all $k > 0, i$. By solving Equation (95) with condition $\Delta u_{01}^n > 0$ we have the general solutions

$$(96) \quad v_{01}^n(x) = \begin{cases} c_1 + c_2 \ln|x| + a|x|^2, & \text{for } m = 2, \\ c_1 + c_2|x|^{-2} + a|x|^2 + b \ln|x|, & \text{for } m = 4, \\ c_1 + c_2|x|^{-(m-2)} + a|x|^2 + b|x|^{-(m-4)}, & \text{for } m \neq 2, 4, \end{cases}$$

where for $m = 2$, the constant $a > 0$, for $m = 3, 4$, the constants $a, b \geq 0, a^2 + b^2 \neq 0$, and for $m \neq 2, 3, 4$, the constants $a \geq 0, b \leq 0, a^2 + b^2 \neq 0$.

Summarizing the above we obtain that $f_n = \sum_{k=0}^{\infty} \sum_{i=1}^{n_k} u_{ki}^n(r)v_{ki}^n(\theta)$ is a biharmonic function on $\mathbb{R}^m \setminus \{0\}$ with $\Delta f_n > 0$ if and only if $f_n = v_{01}^n$ given by (96), which does not depends on n . Since $u(r, \theta) = \lim_{n \rightarrow \infty} f_n = v_{01}^n$, we obtain the theorem. □

Remark 6. (i) We can check that by opposite sign choices for the constants a, b in Theorem 4.1, we obtain all biharmonic functions on the punctured Euclidean space with negative Laplacian, i.e., $\Delta u < 0$.

(ii) Note also that there are many biharmonic functions on $\mathbb{R}^m \setminus \{0\}$ whose Laplacian do not have a fixed sign. For example, $u = r^2 \ln r$ is biharmonic on $\mathbb{R}^2 \setminus \{0\}$ with $\Delta u = 4(1 + \ln r)$ which can be any real number.

REFERENCES

- [1] S. Axler, P. Bourdon, and W. Ramey, *Harmonic Function Theory*, Springer, New York, NY, 1992, GMT Volume 137.
- [2] P. Baird and J. C. Wood, *Harmonic morphisms between Riemannian manifolds*, London Math. Soc. Monogr. (N.S.) No. 29, Oxford Univ. Press (2003).
- [3] R. Caddeo, *Riemannian manifolds on which the distance function is biharmonic*. Rend. Sem. Mat. Univ. Politec. Torino 40, 93-101 (1982).
- [4] D. G. Chen, Q. M. Cheng, Q. L. Wang, and C. Y. Xia, *On eigenvalues of a system of elliptic equations and biharmonic operator*. J. Math. Anal. Appl. 387, 1146-1159 (2012).
- [5] Q. M. Cheng and H.C. Yang, *Inequalities for eigenvalues of a clamped plate problem*, Trans. Amer. Math. Soc. 358 (2006) 2625-2635.
- [6] R. R. Huilgol *On Liouville's theorem for biharmonic functions*, SIAM J. Applied Math., 20 (1) (1971), pp. 37-39.
- [7] H. Lee, *Biharmonic functions on S^2 and H^2* , MS thesis, Texas A & M University-Commerce, May 2024.
- [8] S. Montaldo and A. Ratto, *A Note on the Almansi Property*, Mediterr. J. Math. (2017) 14:200, DOI 10.1007/s00009-017-1003-x 1660-5446/17/050001-13
- [9] Y. -L. Ou and B. -Y. Chen, *Biharmonic submanifolds and biharmonic maps in Riemannian geometry*, World Scientific, Hackensack, N. J., 2020.
- [10] L. Sario and C. Wang, *Generators of the Space of Bounded Biharmonic Functions*, Math. Z., 127 (8) (1972), 273-280.
- [11] L. Sario and C. Wang, *Radial quasiharmonic functions*, Pacific J. Math., 46 (1973), 515-522.
- [12] L. Sario and C. Wang, *Quasiharmonic functions on the Poincaré N -ball*, Rend. Mat. 6 (1973)1-4.
- [13] S. R. Sario, M. Nakai, C. Wang, and L. O. Chung, *Classification Theory of Riemannian Manifolds: Harmonic, Quasiharmonic and Biharmonic Functions*, Lecture Notes in Math 605, Springer-Verlag, Berlin Heidelberg New York, 1977.
- [14] M. A. Shubin, *Pseudodifferential Operators and Spectral Theory*, 2nd edition. Springer, 2001.
- [15] Q. Wang and C. -Y. Xia, *Universal bounds for eigenvalues of the biharmonic operator on Riemannian manifolds*, J. Funct. Analysis, 245 (2007) 334-352.
- [16] Q. Wang and C. Xia, *Sharp Lower Bounds for the First Eigenvalues of the Bi-Laplace Operator*, preprint 2021, arXiv:1802.05502v5.
- [17] C. -Y. Xia, *Eigenvalues on Riemannian manifolds*, IMPA Mathematical Publications, 29th Brazilian Mathematics Colloquium, Rio de Janeiro, 2013.
- [18] L. Zhang and Y. Zhao, *The lower bounds of the first eigenvalues for the biharmonic operator on manifolds*, J. Inequalities and Applications, (5) (2016),
- [19] D. G Zill and M. R. Cullen, *Differential Equations with boundary-value problems*, 7th edition, Brooks/Cole CENGAGE Learning, USA, 2009.

DEPARTMENT OF MATHEMATICS,
TEXAS A & M UNIVERSITY-COMMERCE,
COMMERCE, TX 75429, U S A.
E-MAIL: YELIN.OU@TAMUC.EDU