

Formations of Finite Groups in Polynomial Time: the \mathfrak{F} -Hypercenter

Viachaslau I. Murashka
{mvimath@yandex.ru}

Faculty of Mathematics and Technologies of Programming, Francisk Skorina Gomel State
University, Sovetskaya 104, Gomel, 246028, Belarus

Abstract

For a wide family of formations \mathfrak{F} (which includes Baer-local formations) it is proved that the \mathfrak{F} -hypercenter of a permutation finite group can be computed in polynomial time. In particular, the algorithms for computing the \mathfrak{F} -hypercenter for the following classes of groups are suggested: hereditary local formations with the Shemetkov property, rank formations, formations of all quasinilpotent, Sylow tower, p -nilpotent, supersoluble, w -supersoluble and SC -groups. For some of these formations algorithms for the computation of the intersection of all maximal \mathfrak{F} -subgroups are suggested.

Keywords. Finite group; \mathfrak{F} -hypercenter; Baer-local formation; permutation group computation; polynomial time algorithm.

AMS(2010). 20D10, 20B40.

Introduction

All groups considered here are finite. Recall that a class of groups is a collection \mathfrak{X} of groups with the property that if $G \in \mathfrak{X}$ and if $H \simeq G$, then $H \in \mathfrak{X}$. The theory of classes of groups is well developed nowadays (for example, see [6, 8, 10, 25]) and has various applications (for example, in the theory of formal languages [5], in the solution of Yang-Baxter equation [3] and etc.) The main its tasks are to construct classes of groups and to recognize is a given group belongs to a given class or not. With a class of groups \mathfrak{X} one can associate the canonical subgroups such as the \mathfrak{X} -residual, the \mathfrak{X} -radical, a \mathfrak{X} -projector, a \mathfrak{X} -injector, the \mathfrak{X} -hypercenter and etc. Even if a group does not belong to \mathfrak{X} , then these subgroups encode some of its properties associated with \mathfrak{X} .

The algorithms for computing the \mathfrak{X} -residual and \mathfrak{X} -projectors (of a soluble group) were presented in [9, 11]. The algorithms for computing the \mathfrak{X} -radical and \mathfrak{X} -injectors (of a soluble group) were presented in [11]. Nevertheless the algorithms for the computation of the \mathfrak{X} -hypercenter which plays an important role in the theory of formations were not suggested before.

A group can be represented in different ways. In this paper we will consider only permutation groups because the computational theory of permutation groups is well developed (see [22]). We leave the reader to decide how our algorithms can be applied in the case of non-permutation groups. The aim of this paper is to find effective algorithms (which runs in polynomial time for permutation groups) for the computation of the \mathfrak{F} -hypercenter for a wide family of formations \mathfrak{F} of not necessary soluble groups.

1 The Main Result

Recall that the hypercenter of a group is just the final member of its upper central series. The concept of the hypercenter was widely studied and generalized by different mathematicians

(Baer [2], Huppert [12], Shemetkov [23], Skiba [26] and many other). Some of their results are presented in [10, Chapter 1]. In the final form the notion of hypercenter appeared in [25]. Let \mathfrak{X} be a class of groups. A chief factor H/K of G is called \mathfrak{X} -central in G provided that the semidirect product $(H/K) \rtimes (G/C_G(H/K))$ of H/K with $G/C_G(H/K)$ corresponding to the action by conjugation of G on H/K belongs to \mathfrak{X} (see [25, p. 127–128]). The \mathfrak{X} -hypercenter $Z_{\mathfrak{X}}(G)$ is the greatest normal subgroup of G all whose G -composition factors are \mathfrak{X} -central.

From Barnes-Kegel Theorem [8, IV, Proposition 1.5] it follows that if \mathfrak{F} is a formation and $G \in \mathfrak{F}$, then $G = Z_{\mathfrak{F}}(G)$. The converse of this statement is false as we can see on the example of the class of all abelian groups. Shemetkov [24] asked to described all formations $\mathfrak{F} = (G \mid G = Z_{\mathfrak{F}}(G))$ (such formations are called Z -saturated [16]). The solution of this problem was started in [4, 16]. The method for calculating the \mathfrak{F} -hypercenter of a given group may be a useful tool in the solution of this problem.

In [18] the example of a permutation group of degree n was given such that it has a quotient with no faithful representations of degree less than $2^{n/4}$. That is why the given definition of the \mathfrak{X} -central chief factor is not good from the computational point of view. Therefore we will consider a more general definition:

Definition 1 ([17, Definition 2]). Let f be a function which assigns 0 or 1 to every group G and its chief factor H/K such that

- (1) $f(H/K, G) = f(M/N, G)$ whenever H/K and M/N are G -isomorphic chief factors of G ;
- (2) $f(H/K, G) = f((H/N)/(K/N), G/N)$ for every $N \trianglelefteq G$ with $N \leq K$.

Such functions f will be called chief factor functions. Denote by $\mathcal{C}(f)$ the class of groups

$$(G \mid G \simeq 1 \text{ or } f(H/K, G) = 1 \text{ for every chief factor } H/K \text{ of the group } G).$$

In particular if $f(H/K, G) = 1$ iff H/K is \mathfrak{X} -central, then f is a chief factor function. So Z -saturated formations (and hence local and Baer-local formations) are the particular cases of this construction. With each chief factor function we can associate the following subgroup:

Definition 2. Denote by $Z(G, f)$ the greatest normal subgroup of G such that $f(H/K, G) = 1$ for every chief factor H/K of G below it.

The subgroup $Z(G, f)$ becomes the \mathfrak{X} -hypercenter of G with the right choice of f . In particular, if f checks if H/K is central (resp. cyclic) in G , then $Z(G, f)$ is the (resp. supersoluble) hypercenter.

Example 1. Let f_1 and f_2 check if H/K is abelian and if $G/C_G(H/K)$ is soluble respectively. Then $\mathcal{C}(f_1) = \mathcal{C}(f_2) = \mathfrak{S}$ is the class of all soluble groups. Note that $Z(G, f_1)$ is the soluble radical of G and $Z(G, f_2)$ is the soluble hypercenter of G . If we consider the semidirect product of the alternating group A_5 of degree 5 and its faithful simple module over \mathbb{F}_5 (it exists by [8, B, Theorem 10.3]), then we will see that these subgroups are different.

The main result of the paper is

Theorem 1. *Assume that $f(H/T, G)$ can be computed in polynomial time (in n) for every group $G \leq S_n$ and its chief factor H/T . Then the subgroup $Z(G/K, f)$ is well defined and can be computed in polynomial time (in n) for every $K \trianglelefteq G \leq S_n$.*

In section 4 we will discuss how this theorem can be applied to various formations.

2 Preliminaries

All unexplained notations and terminologies are standard. The reader is referred to [6, 8, 10] if necessary. Recall that $Z(G)$ denotes the center of G ; $O_p(G)$ is the greatest normal p -subgroup of G ; $O^\pi(G)$ is the smallest normal subgroup of G of π -index; $\Omega_1(G)$ denotes the subgroup that is generated by elements of order p for a p -group G ; S_n denotes the symmetric group of degree n ; \mathfrak{G}_π and \mathfrak{S} denote the classes of all π -groups and soluble groups respectively.

Recall that a *formation* is a class of groups \mathfrak{F} which is closed under taking epimorphic images (i.e. from $G \in \mathfrak{F}$ and $N \trianglelefteq G$ it follows that $G/N \in \mathfrak{F}$) and subdirect products (i.e. from $G/N_1 \in \mathfrak{F}$ and $G/N_2 \in \mathfrak{F}$ it follows that $G/(N_1 \cap N_2) \in \mathfrak{F}$). If \mathfrak{F} is a non-empty formation, then in every group G exists the \mathfrak{F} -residual, i.e. the smallest normal subgroup of G with $G/G^\mathfrak{F} \in \mathfrak{F}$.

We use standard computational conventions of abstract finite groups equipped with polynomial-time procedures to compute products and inverses of elements (see [22, Chapter 2]). For both input and output, groups are specified by generators. We will consider only $G = \langle S \rangle \leq S_n$ with $|S| \leq n^2$. If necessary, Sims' algorithm [22, Parts 4.1 and 4.2] can be used to arrange that $|S| \leq n^2$. Quotient groups are specified by generators of a group and its normal subgroup. We need the following well known basic tools in our proofs (see, for example [13] or [22]). Note that some of them are obtained mod CFSG.

Theorem 2. *Given $A, B \trianglelefteq G \leq S_n$ with $A \leq B$, in polynomial time one can solve the following problems:*

1. Find $C_{G/A}(B/A)$.
2. Find G' and hence $G^\mathfrak{S}$.
3. Find $|G|$.
4. Find $Z(G/A)$, $O_p(G/A)$ and $O^\pi(G/A)$.
5. Find a chief series for G containing A and B .
6. Given $H = \langle S_1 \rangle, K = \langle S_2 \rangle \leq G$ find $\langle H, K \rangle = \langle S_1, S_2 \rangle$ and $[H, K] = \langle \{[s_1, s_2] \mid s_1 \in S_1, s_2 \in S_2\} \rangle^{(H, K)}$.
7. Check if G/A is simple.

The following lemma plays an important role in our proves.

Lemma 1 ([1]). *Given $G \leq S_n$ every chain of subgroups of G has at most $2n - 3$ members for $n \geq 2$.*

3 Proof of Theorem 1

The proof of Theorem 1 is based on the following 4 lemmas.

Lemma 2. *The following statements hold:*

- (1) $Z(G, f)$ is well defined for any group G .
- (2) Let $Z/K = Z(G/K, f)$. Then Z is the greatest normal subgroup of G such that it contains K and $f(H/T, G) = 1$ for every chief factor H/T of G with $K \leq T \leq H \leq Z$.

Proof. (1) Let M and N be normal subgroups of a group G . Then from [8, A, The Isomorphism Theorems(2)] it follows that every chief factor of G below MN is G -isomorphic to a chief factor of G below either M or N . So if $f(H/K, G) = 1$ for every chief factor H/K of G below M and N , then $f(H/K, G) = 1$ for every chief factor H/K of G below MN by (1) of Definition 1. It means that the subgroup $Z(G, f)$ is well defined.

(2) Let Z be the greatest normal subgroup of G such that it contains K and $f(H/T, G) = 1$ for every chief factor H/T of G with $K \leq T \leq H \leq Z$. Now $f((H/K)/(T/K), G/K) = f(H/T, G) = 1$ for every chief factor H/T of G such that $(H/K)/(T/K)$ is a chief factor of G/K below Z/K by (2) of Definition 1. Hence $Z/K \leq Z(G/K, f)$. Assume that $f((M/K)/(Z/K), G/K) = 1$ for some chief factor $(M/K)/(Z/K)$ of G/K . Then $f(M/Z, G) = 1$ for some chief factor M/Z of G by (2) of Definition 1, a contradiction with the definition of Z . Thus $Z/K = Z(G/K, f)$. \square

Lemma 3. *Let H/T be a chief factor of G , $K \trianglelefteq G$, $K \leq T$, $Z_1/K = Z(G/K, f) \cap T/K$ and $Z_2/K = Z(G/K, f) \cap H/K$. Then $Z_2/Z_1 \not\cong 1$ if and only if $f(H/T, G) = 1$ and $H/Z_1 = T/Z_1 \times Z/Z_1$ for some $Z \trianglelefteq G$. In this case $Z_2 = Z$.*

Proof. Note that $Z_1/K = Z_2/K \cap T/K = (Z_2 \cap T)/K$. Hence $Z_1 = Z_2 \cap T$. Now

$$Z_2/Z_1 = Z_2/(Z_2 \cap T) \simeq Z_2T/T \leq H/T.$$

Hence Z_2/Z_1 is G -isomorphic to either 1 or H/T .

Assume that $Z_2/Z_1 \not\cong 1$. It means that $1 = f(Z_2/Z_1, G) = f(H/T, G)$ and Z_2/Z_1 is a minimal normal subgroup of H/Z_1 not contained in T/Z_1 . Hence $H/Z_1 = T/Z_1 \times Z_2/Z_1$.

Assume now that $f(H/T, G) = 1$ and $H/Z_1 = T/Z_1 \times Z/Z_1$ for some $Z \trianglelefteq G$. Note that $f(A/B, G) = 1$ for every chief factor A/B of G between Z_1 and K . From $H/T = ZT/T \simeq Z/(Z \cap T) = Z/Z_1$ it follows that $f(Z/Z_1, G) = 1$. It means that

$$Z_1/K < Z/K \leq Z(G/K, f) \cap H/K = Z_2/K.$$

Therefore $Z_2/Z_1 \not\cong 1$ and $Z = Z_2$. \square

Lemma 4. *In the notations of Lemma 3 if H/T is non-abelian and Z_1 is given, then we can compute Z_2 in polynomial time.*

Proof. Assume that H/T is non-abelian. If $f(H/T, G) = 0$, then $Z_1 = Z_2$ by Lemma 3. So we can assume that $f(H/T, G) = 1$. Let $C/Z_1 = C_{H/Z_1}(T/Z_1)^\ominus$. We claim that $Z_2/Z_1 \not\cong 1$ iff $|C/Z_1| = |H/T|$. In this case if $Z_2/Z_1 \not\cong 1$, then $Z_2/Z_1 = C/Z_1$.

Suppose that $Z_2/Z_1 \not\cong 1$. From Lemma 3 it follows that $H/Z_1 = T/Z_1 \times Z/Z_1$ for some $Z \trianglelefteq G$. Then $Z/Z_1 \leq C_{H/Z_1}(T/Z_1)$. Hence

$$C_{H/Z_1}(T/Z_1) = TZ/Z_1 \cap C_{H/Z_1}(T/Z_1) = (C_{H/Z_1}(T/Z_1) \cap T/Z_1)(Z/Z_1) = Z(T/Z_1) \times (Z/Z_1).$$

Recall that $H/T \simeq Z/Z_1$ is a direct product of simple non-abelian groups. It means that $C_{H/Z_1}(T/Z_1)^\ominus = Z/Z_1 = Z_2/Z_1$ by Lemma 3. Thus $|C/Z_1| = |H/T|$.

Suppose that $|C/Z_1| = |H/T|$. Note that C/Z_1 is non-abelian (because it is the soluble residual) and $C/Z_1 \cap T/Z_1 \leq Z(T/Z_1)$ is abelian. Therefore $C/Z_1 \not\leq T/Z_1$. From

$$C/Z_1 \text{ char } C_{H/Z_1}(T/Z_1) = C_{G/Z_1}(T/Z_1) \cap H/Z_1 \trianglelefteq G/Z_1$$

it follows that $C/Z_1 \trianglelefteq G$. It means that $H = TC$. Note that $H/T = CT/T \simeq C/(C \cap T)$. From $|C/Z_1| = |H/T|$ it follows that $|C \cap T| = |Z_1|$. Note that $Z_1 \leq C \cap T$. Thus $C \cap T = Z_1$. It means that $H/Z_1 = T/Z_1 \times C/Z_1$. Thus $C/Z_1 = Z_2/Z_1$ by Lemma 3.

By the statement of Theorem 1 we can compute $f(H/T, G)$ in polynomial time. Now $C/Z_1 = C_{H/Z_1}(T/Z_1)$ can be computed in polynomial time by 1 of Theorem 2. Note that $(C/Z_1)^\mathfrak{S} = C^\mathfrak{S}Z_1/Z_1$. Now we can compute $C^\mathfrak{S}$ by 2 of Theorem 2. According to 3 of Theorem 2 we can check if $|H/T| = |C/Z_1|$ in polynomial time. \square

Lemma 5. *In the notations of Lemma 3 if H/T is abelian and Z_1 is given, then we can compute Z_2 in polynomial time.*

Proof. Assume that H/T is abelian. If $f(H/T, G) = 0$, then $Z_1 = Z_2$ by Lemma 3. So we can assume that $f(H/T, G) = 1$.

Note that H/T is a p -group for some prime p and every minimal normal p -subgroup of a group is centralized by its the greatest normal p -subgroup [8, A, Lemma 13.6]. Also note that every minimal normal p -subgroup is generated by elements of order p . Hence all minimal normal p -subgroups of H/Z_1 are subgroups of $P/Z_1 = \Omega_1(Z(\mathcal{O}_p(H/Z_1)))$.

If $P \leq T$, then $H/Z_1 \neq T/Z_1 \times Z/Z_1$ for all $Z \trianglelefteq G$. Thus $Z_2 = Z_1$ by Lemma 3. Assume now that $P \not\leq T$. Note that P/Z_1 is the direct product of G -indecomposable subgroups $P_1/Z_1, \dots, P_m/Z_1$ by the Krull-Remak-Schmidt Theorem. Note that $H/Z_1 = T/Z_1 \times Z/Z_1$ for some $Z \trianglelefteq G$ iff $P/Z_1 = (P \cap T)/Z_1 \times Z/Z_1$ for some G -simple subgroup Z/Z_1 of P/Z_1 . Let $(P \cap T)/Z_1$ be the direct product of G -indecomposable subgroups $T_1/Z_1, \dots, T_l/Z_1$. If the required Z exists, then P/Z_1 is the direct product of G -indecomposable subgroups $T_1/Z_1, \dots, T_l/Z_1$ and Z/Z_1 . Hence $l = m - 1$. Moreover if P/Z_1 is the direct product of G -indecomposable subgroups $Q_1/Z_1, \dots, Q_m/Z_1$, then these subgroups can be numbered in such way that P/Z_1 is the direct product of $T_1/Z_1, \dots, T_{m-1}/Z_1$ and Q_m/Z_1 by the Krull-Remak-Schmidt Theorem. Note that in these case Q_m/Z_1 is G -simple and not contained in T/Z_1 . Therefore the required subgroup Z exists only in case when any direct decomposition of P/Z_1 into the product of G -indecomposable subgroups has G -simple subgroup not contained in T/Z_1 .

Note that P/Z_1 can be viewed as a G -module over \mathbb{F}_p and G -indecomposable and G -simple subgroups of $P/Z_1 \leq G/Z_1$ are in one to one correspondence of indecomposable and simple submodules of a G -module P/Z_1 .

From 4 of Theorem 2 it follows that we can compute $Z(\mathcal{O}_p(H/Z_1))$ in polynomial time. Since $Z(\mathcal{O}_p(H/Z_1))$ is abelian, we can find P/Z_1 in polynomial time by taking the right powers of generators of $Z(\mathcal{O}_p(H/Z_1))$. According to [22, p. 155] for each generator g of G in polynomial time we can find the linear transformation which this element induces on P/Z_1 . Let R be the algebra generated by the above mentioned transformations. Now by [7, Theorem 5] we can decompose P/Z_1 into the sum $P_1 \oplus \dots \oplus P_k$ of indecomposable submodules in polynomial time. For each P_i we can check either it is simple or not by [20, Corollary 5.4] and for each simple P_i we can find the subgroup P_i/Z_1 of P/Z_1 to which it corresponds. If $P_i/Z_1 \not\leq T/Z_1$, then P_i/Z_1 is the required subgroup. \square

3.1 Proof of the Theorem

We can compute a chief series $K = G_0 \trianglelefteq G_1 \trianglelefteq \dots \trianglelefteq G_m = G$ of G in polynomial time by 5 of Theorem 2. Let $Z_i = G_i \cap Z(G/K, f)$. We see that $Z_{i-1} = Z_i \cap G_{i-1}$ and $Z_0 = K$. Now using Lemmas 4 and 5 if we know Z_{i-1} then we can compute Z_i in polynomial time. Note that $m \leq 2n$ by Lemma 1. Hence we can compute $Z(G/K, f)$ in polynomial time (see Algorithm 1).

Algorithm 1: GFHYPERCENTER(G, K, f)

Result: $Z(G/K, f)$.

Data: A normal subgroup K of a group G and a chief factor function f .

$Z \leftarrow K$;

Compute a chief series $K = G_0 \trianglelefteq G_1 \trianglelefteq \cdots \trianglelefteq G_m = G$ of G ;

for $i \in \{1, \dots, m\}$ **do**

if $f(G_i/G_{i-1}, G) = 1$ **then**

if $|\pi(G_i/G_{i-1})| > 1$ **then**

$C/Z \leftarrow C_{G_i/Z}(G_{i-1}/Z)$;

$S/Z \leftarrow \langle C^{\mathfrak{S}}, Z \rangle / Z$;

if $|S/Z| = |G_i/G_{i-1}|$ **then**

$Z \leftarrow S$;

end

else

$p \leftarrow \pi(G_i/G_{i-1})[1]$;

$P/Z \leftarrow \Omega_1(Z(O_p(G_i/Z)))$;

 For each generator g of G find the linear transformation which this element induces on P/Z ;

 Let R be the algebra generated by above mentioned transformations.

 Decompose the R -module P/Z into the sum $P_1 \oplus \cdots \oplus P_k$ of indecomposable submodules;

for $j \in \{1, \dots, k\}$ **do**

if P_j is a simple module **then**

 Find the corresponding to P_j subgroup P_j/Z ;

if $\langle P_j/Z, G_{i-1}/Z \rangle = G_i/Z$ **then**

$Z \leftarrow P_j$;

 Leave the current **for** cycle;

end

end

end

end

end

end

return Z/K

4 Applications

Recall [26] that $\text{Int}_{\mathfrak{F}}(G)$ denotes the intersection of all \mathfrak{F} -maximal subgroups of G . L.A. Shemetkov posed the following question on Gomel Algebraic seminar in 1995: «For what non-empty (normally) hereditary local (Baer-local) formations \mathfrak{F} do the equality $\text{Int}_{\mathfrak{F}}(G) = Z_{\mathfrak{F}}(G)$ hold for every group G ?» The solution to this question in case when \mathfrak{F} is a hereditary saturated formation was obtained in [26]. For some class of Baer-local formation the answer to this question was given in [15]. Nevertheless this problem is still open. We will show that Theorem 1 can be used to compute $\text{Int}_{\mathfrak{F}}(G)$ for some formations \mathfrak{F} .

4.1 Baer local and local formation

The notion of \mathfrak{F} -hypercenter plays an important role in the study for Baer-local and local formations. Let f be a function which assigns to every simple group J a possibly empty formation $f(J)$. Now extend the domain of f . If G is the direct product of simple groups isomorphic to J , then let $f(G) = f(J)$. If J is a cyclic group of order p , then let $f(p) = f(J)$. Such functions f are called Baer functions [8, IV, Definitions 4.9]. A formation \mathfrak{F} is called Baer-local (or composition, see [10, p. 4] and [23]) if for some Baer function f holds:

$$\mathfrak{F} = (G \mid G/C_G(H/K) \in f(H/K) \text{ for every chief factor } H/K \text{ of } G).$$

Theorem 3. *Let \mathfrak{F} be a Baer-local formation defined by f . Assume that $(G/K)^{f(J)}$ can be computed in polynomial time for every $K \trianglelefteq G \leq S_n$ and a simple group J . Then $Z_{\mathfrak{F}}(G/K)$ can be computed in polynomial time for every $K \trianglelefteq G \leq S_n$ and a simple group J .*

Proof. Here we assume that if $f(J) = \emptyset$, then the computation of $(G/K)^{f(J)}$ returns “is not defined”. Let $f(H/K, G) = 1$ iff $G/C_G(H/K) \in f(H/K)$. This condition is equivalent to $G^{f(H/K)}$ is defined and $[G^{f(H/K)}, H] \subseteq K$. Hence it can be checked in polynomial time by 3 and 6 of Theorem 2 and the statement of the theorem. By analogy with the proof of [17, Theorem 7] one can show that f is a chief factor function. Then $\mathfrak{F} = \mathcal{C}(f)$. Hence we can compute $G^{\mathfrak{F}}$ in polynomial time by [17, Theorem 1].

Let H/K be a chief factor of G and $T \simeq (H/K) \rtimes G/C_G(H/K)$. Note that T is a primitive group of type 1 or 3. Then $T/C_T(H/K) \simeq G/C_G(H/K)$ by [8, A, Theorem 15.2(1, 3)]. So if $T \in \mathfrak{F}$, then $G/C_G(H/K) \in f(H/K) \cap \mathfrak{F}$. From the other hand if $G/C_G(H/K) \in f(H/K) \cap \mathfrak{F}$, then $T \in \mathfrak{F}$. Note that $G/C_G(H/K) \in (f(H/K) \cap \mathfrak{F})$ iff $[G^{\mathfrak{F}}G^{f(H/K)}, H] \subseteq K$. Let $f_1(H/K, G) = 1$ iff $G^{f(H/K)}$ is defined and $[G^{\mathfrak{F}}G^{f(H/K)}, H] \subseteq K$. Note that we can check this condition in polynomial time by 3 and 6 of Theorem 2, f_1 is a chief factor function and $Z(G, f_1) = Z_{\mathfrak{F}}(G)$ for any group G . Thus $Z_{\mathfrak{F}}(G/K)$ can be computed in polynomial time for any $K \trianglelefteq G \leq S_n$ by Theorem 1. \square

A Baer-local formation defined by Baer function f is called local if $f(J) = \bigcap_{p \in \pi(J)} f(p)$ for every simple group J . In this case to define \mathfrak{F} we need only to know the values of f on primes.

Corollary 3.1. *Let \mathfrak{F} be a local formation locally defined by f . Assume that $(G/K)^{f(p)}$ can be computed in polynomial time for every $K \trianglelefteq G \leq S_n$ and a prime p . Then $Z_{\mathfrak{F}}(G/K)$ can be computed in polynomial time.*

According to [17, the proof of Corollary 1] for the class \mathfrak{U} of supersoluble groups, the class $w\mathfrak{U}$ of widely supersoluble groups [27], the class \mathfrak{NA} of groups G such that all Sylow subgroups of $G/F(G)$ are abelian [27], the class $sm\mathfrak{U}$ of groups with submodular Sylow subgroups [29, 30], the class of strongly supersoluble groups $s\mathfrak{U}$ [29] and the class $sh\mathfrak{U}$ of groups all of whose Schmidt subgroups are supersoluble [14] we can compute the described in the statement of Corollary 3.1 $f(p)$ -residuals in polynomial time.

Corollary 3.2. *Let $\mathfrak{F} \in \{\mathfrak{U}, w\mathfrak{U}, s\mathfrak{U}, sm\mathfrak{U}, \mathfrak{NA}, sh\mathfrak{U}\}$. Then $Z_{\mathfrak{F}}(G/K)$ can be computed in a polynomial time for every $K \trianglelefteq G \leq S_n$.*

Recall [28] that a group is called c -supersoluble in the terminology of Vedernikov (SC -group in the terminology of Robinson [19]) if every its chief factor is a simple group. It was proved the the class \mathfrak{U}_c of all c -supersoluble groups is a Baer-local formation [28].

Theorem 4. *In polynomial time one can compute $\text{Int}_{\mathfrak{U}_c}(G/K)$ and $Z_{\mathfrak{U}_c}(G/K)$ for $K \trianglelefteq G \leq S_n$.*

Proof. Let f_1 checks if H/K is simple. Then $f_1(H/K, G)$ can be computed in polynomial time by 7 of Theorem 2. Therefore $\mathfrak{U}_c = \mathcal{C}(f_1)$. It is clear that $Z(G, f_1)$ is the greatest normal subgroup of G all whose G -composition factors are simple. Therefore $Z(G, f_1)H \in \mathfrak{U}_c$ for every \mathfrak{U}_c -subgroup H of G . Hence $Z(G, f_1) = \text{Int}_{\mathfrak{U}_c}(G)$.

From the other hand let $G \simeq \text{Aut}(P\Omega_8^+(p^f))$ for an odd p and $P\Omega_8^+(p^f) \simeq I \trianglelefteq G$. Note that $I \leq \text{Int}_{\mathfrak{U}_c}(G)$. Now $G/C_G(I) \notin \mathfrak{U}_c$ because it contains a normal section isomorphic to S_4 . Note that an abelian chief factor is \mathfrak{U}_c -central iff it is simple. If H/K is a non-abelian factor, then $T = (H/K) \rtimes (G/C_G(H/K))$ has two minimal normal subgroups whose quotients are isomorphic to $G/C_G(H/K)$ by [8, A, Theorem 15.2(3)]. Now $T \in \mathfrak{U}_c$ iff $G/C_G(H/K) \in \mathfrak{U}_c$. The last is equivalent to $[G^{\mathfrak{U}_c}, H] \subseteq K$. According to [17, Theorem 1] we can compute $G^{\mathfrak{U}_c}$ in polynomial time. Therefore we can check if H/K is \mathfrak{U}_c -central in G in polynomial time and hence compute $Z_{\mathfrak{U}_c}(G/K)$. \square

Recall that a group is called quasinilpotent if every its element induces an inner automorphism on every its chief factor. The class of all quasinilpotent groups is denoted by \mathfrak{N}^* .

Theorem 5. *In polynomial time one can compute $\text{Int}_{\mathfrak{N}^*}(G/K) = Z_{\mathfrak{N}^*}(G/K)$ for $K \trianglelefteq G \leq S_n$.*

Proof. According to [15, Corollary 1] $\text{Int}_{\mathfrak{N}^*}(G) = Z_{\mathfrak{N}^*}(G)$ holds for every group G . From [15, Remark 2] $Z_{\mathfrak{N}^*}(G)$ is the greatest normal subgroup of G such that every element of G induces an inner automorphism on every chief factor of G below $Z_{\mathfrak{N}^*}(G)$, i.e. $Z_{\mathfrak{N}^*}(G) = Z(G, f)$ where f checks if every element of G induces an inner automorphism on H/K that is $G = HC_G(H/K)$ or $G/K = (H/K)C_{G/K}(H/K)$. This condition can be checked in polynomial time by Theorem 2. It is straightforward to check that f is a chief factor function. Thus in polynomial time one can compute $\text{Int}_{\mathfrak{N}^*}(G/K) = Z_{\mathfrak{N}^*}(G/K)$ for $K \trianglelefteq G \leq S_n$ by Theorem 1. \square

4.2 Formations with the Shemetkov property

Recall [21] (see also [6, Chapter 6]) that a formation \mathfrak{F} has the Shemetkov property (resp. in \mathfrak{S}) if every (resp. soluble) \mathfrak{F} -critical group is either a Schmidt group or a cyclic group of prime order. Note that if a hereditary local formation \mathfrak{F} has the Shemetkov property (and contains all nilpotent groups) then it can be locally defined by f where $f(p) = \mathfrak{S}_{g(p)}$ and g assigns to a prime p a set of primes $g(p)$ with $p \in g(p)$. The converse is not true. Nevertheless a hereditary local formation \mathfrak{F} of soluble groups has the Shemetkov property in \mathfrak{S} and contains all nilpotent groups iff \mathfrak{F} is locally defined by f where $f(p) = \mathfrak{S}_{g(p)}$ and g assigns to a prime p a set of primes $g(p)$ with $p \in g(p)$. With the help of the next result one can compute the \mathfrak{F} -hypercenter and the intersection of all maximal \mathfrak{F} -subgroups of a (resp. soluble) group G where \mathfrak{F} is a hereditary formation with the Shemetkov property (resp. in \mathfrak{S}).

Theorem 6. *Let g be a function which assigns to a prime p a set of primes $g(p)$ with $p \in g(p)$ and $h(p) = \mathfrak{S}_{g(p)}$. Assume that $g(p) \cap \pi(G)$ can be computed in a polynomial time for every $p \in \pi(G)$. Let \mathfrak{F} be a local formation defined by h . Then $\text{Int}_{\mathfrak{F}}(G/K)$ and $Z_{\mathfrak{F}}(G/K)$ can be computed in a polynomial time for every $K \trianglelefteq G \leq S_n$.*

Proof. Let prove that $H/K \leq \text{Int}_{\mathfrak{F}}(G/K)$ iff $G/C_G(H/K) \in h(p)$ for every $p \in \pi(H/K)$.

Assume that $G/C_G(H/K) \in h(p)$ for every $p \in \pi(H/K)$. Let M/K be an \mathfrak{F} -maximal subgroup of G/K . Let $T = MH$. Note that $T/C_T(U/V) \in h(p)$ for all $p \in \pi(U/V)$ and every chief factor U/V of T/K below H/K . From $T/H \in \mathfrak{F}$ it follows that $T/C_T(U/V) \in h(p)$

for all $p \in \pi(U/V)$ and every chief factor U/V of T/K above H/K . Thus $T/K \in \mathfrak{F}$. Hence $T/K = M/K$. Therefore $H/K \leq \text{Int}_{\mathfrak{F}}(G/K)$.

Assume that $H/K \leq \text{Int}_{\mathfrak{F}}(G/K)$. It means that $\pi(H/K) \subseteq g(p)$ for all $p \in \pi(H/K)$. Let M/K be an \mathfrak{F} -maximal subgroup of G/K and $K = H_0 \trianglelefteq H_1 \trianglelefteq \cdots \trianglelefteq H_m = H$ be a part of chief series of M/K . Then $(M/K)/C_{M/K}(H_i/H_{i-1}) \in h(p)$ for all $p \in \pi(H/K)$. Let $C/K = \bigcap_{i=1}^m C_{M/K}(H_i/H_{i-1})$. Now $(C/K)/C_{M/K}(H/K)$ is a $\pi(H/K)$ -group by [A, Corollary 12.4(a)]. Since $h(p)$ is a formation, $(M/K)/(C/K) \in h(p)$ for all $p \in \pi(H/K)$. Thus $(M/K)/C_{M/K}(H/K) \in \mathfrak{G}_{g(p)} = h(p)$ for all $p \in \pi(H/K)$. Since $g(p) \neq \emptyset$ for all prime p , we see that every element of G/K belongs to some its \mathfrak{F} -maximal subgroup. From $(M/K)/C_{M/K}(H/K) \simeq MC_{G/K}(H/K)/C_{G/K}(H/K)$ it follows that $G/C_G(H/K)$ is a $\mathfrak{G}_{g(p)}$ -group for every $p \in \pi(H/K)$, i.e. $G/C_G(H/K) \in h(p)$ for every $p \in \pi(H/K)$.

From [26, Theorem C(e)] it follows that if $I \leq \text{Int}_{\mathfrak{F}}(G)$, then $\text{Int}_{\mathfrak{F}}(G/I) = \text{Int}_{\mathfrak{F}}(G)/I$. Therefore $\text{Int}_{\mathfrak{F}}(G/K) = \text{Z}(G/K, \mathfrak{f})$ where \mathfrak{f} checks if $G/C_G(H/K) \in h(p)$ for every $p \in \pi(H/K)$. The last condition is equivalent to $[\text{O}^{g(p)}, H] \subseteq K$ and can be checked in polynomial time by Theorem 2. Thus the statement of theorem follows from Theorem 1 and Corollary 3.1. \square

Let φ be some linear ordering on \mathbb{P} . Recall [8, IV, Examples 3.4(g)] that a group G is called a Sylow tower group of type φ if it has normal Hall $\{p_1, \dots, p_t\}$ -subgroups for all $1 \leq t \leq k$ where $\pi(G) = \{p_1, \dots, p_k\}$ and $p_i >_{\varphi} p_j$ for $i < j$. Let $g(p) = \{q \mid q \leq_{\varphi} p\}$. Note that the class of all Sylow tower group of type φ can be locally defined by $h(p) = \mathfrak{G}_{g(p)}$.

Corollary 6.1. *Let φ be a linear ordering on \mathbb{P} such that we can check in polynomial time if $p \leq_{\varphi} q$ for any primes $p, q \leq n$. If \mathfrak{F} is the class of all Sylow tower group of type φ , then $\text{Int}_{\mathfrak{F}}(G/K)$ and $\text{Z}_{\mathfrak{F}}(G/K)$ can be computed in a polynomial time for every $K \trianglelefteq G \leq S_n$.*

Note that the class of all p -nilpotent groups can be locally defined by h where $h(p) = \mathfrak{G}_p$ and $h(q) = \mathfrak{G}$ for $q \neq p$.

Corollary 6.2. *Let p be a prime. If \mathfrak{F} is the class of all p -nilpotent groups, then $\text{Int}_{\mathfrak{F}}(G/K)$ and $\text{Z}_{\mathfrak{F}}(G/K)$ can be computed in a polynomial time for every $K \trianglelefteq G \leq S_n$.*

4.3 Rank formations

If H/K is a chief factor of G , then $H/K = H_1/K \times \cdots \times H_k/K$ where H_i are isomorphic simple groups. The number k is called the *rank* of H/K in G .

Definition 3 ([8, Chapter VII, Definition 2.3]). A *rank function* R is a map which associates with each prime p a set $R(p)$ of natural numbers. With each rank function R we associate a class of soluble groups

$$\mathfrak{F}(R) = (G \in \mathfrak{S} \mid G \simeq 1 \text{ or for all } p \in \mathbb{P} \text{ each } p\text{-chief factor of } G \text{ has rank in } R(p)).$$

This class is a formation of soluble groups. If $R(p) = \{1, 2\}$ for all $p \in \mathbb{P}$, then formation $\mathfrak{F}(R)$ is non-local by [8, Chapter VII, Theorem 2.18] and hence is non-Baer-local. If $R(p) = \{1\}$ for all $p \in \mathbb{P}$, then $\mathfrak{F}(R) = \mathfrak{A}$.

Theorem 7. *Let R be a rank function. Assume that one can test if $n \in R(p)$ in polynomial (in n) time for every $p \in \mathbb{P}$. Then $\text{Z}_{\mathfrak{F}(R)}(G/K)$ can be computed in polynomial time for every $K \trianglelefteq G \leq S_n$.*

Proof. From 3 of Theorem 2 we can compute the rank of H/K in polynomial time. Hence we can check if $G/K \in \mathfrak{F}(\mathcal{R})$ for any $K \trianglelefteq G \leq S_n$ in polynomial time by 3 and 5 of Theorem 2 and the statement of the theorem. Note that a chief factor H/K is $\mathfrak{F}(R)$ -central iff H/K is a p -group for some prime p , the rank of H/K is in $R(p)$ and $G/C_G(H/K) \in \mathfrak{F}(R)$. Note that $C_G(H/K)/K = C_{G/K}(H/K)$. Hence we can check if H/K is $\mathfrak{F}(R)$ -central in polynomial time by Theorem 2. Thus the statement of Theorem 7 follows from Theorem 1. \square

Acknowledgments

This work is supported by BRFFR $\Phi 23PH\Phi$ -237.
I am grateful to A. F. Vasil'ev for helpful discussions.

References

- [1] L. BABAI, *On the length of subgroup chains in the symmetric group*, Comm. Algebra, 14 (1986), pp. 1729–1736.
- [2] R. BAER, *Supersoluble Immersion*, Canad J. Math., 11 (1959), pp. 353–369.
- [3] A. BALLESTER-BOLINCHES, R. ESTEBAN-ROMERO, P. JIMÉNEZ-SERAL, AND V. PÉREZ-CALABUIG, *On Yang-Baxter groups*, Quaest. Math., 46 (2023), pp. 1273–1281.
- [4] A. BALLESTER-BOLINCHES AND M. D. PEREZ-RAMOS, *On a question of L. A. Shemetkov*, Comm. Algebra, 27 (1999), pp. 5615–5618.
- [5] A. BALLESTER-BOLINCHES, J.-E. PIN, AND X. SOLER-ESCRIVÁ, *Formations of finite monoids and formal languages: Eilenberg's variety theorem revisited*, Forum Math., 26 (2014), pp. 1737–1761.
- [6] A. BALLESTER-BOLLINCHES AND L. M. EZQUERRO, *Classes of Finite Groups*, vol. 584 of Math. Appl., Springer Netherlands, 2006.
- [7] A. L. CHISTOV, G. IVANYOS, AND M. KARPINSKI, *Polynomial time algorithms for modules over finite dimensional algebras*, in International Symposium on Symbolic and Algebraic Computation, 1997, pp. 68–74.
- [8] K. DOERK AND T. O. HAWKES, *Finite Soluble Groups*, vol. 4 of De Gruyter Exp. Math., De Gruyter, Berlin, New York, 1992.
- [9] B. EICK AND C. R. B. WRIGHT, *Computing Subgroups by Exhibition in Finite Solvable Groups*, J. Symb. Comput., 33 (2002), pp. 129–143.
- [10] W. GUO, *Structure Theory for Canonical Classes of Finite Groups*, Springer-Verlag, Berlin, Heidelberg, 2015.
- [11] B. HÖFLING, *Computing Projectors, Injectors, Residuals and Radicals of Finite Soluble Groups*, J. Symb. Comput., 32 (2001), pp. 499–511.
- [12] B. HUPPERT, *Zur Theorie der Formationen*, Arch. Math., 19 (1969), pp. 561–574.
- [13] W. M. KANTOR AND E. M. LUKS, *Computing in quotient groups*, in Proceedings of the Twenty-Second Annual ACM Symposium on Theory of Computing, Baltimore, May 14–16, New York, 1990, ACM, pp. 524–534.

- [14] V. S. MONAKHOV, *Finite groups with a given set of Schmidt subgroups*, Math. Notes, 58 (1995), pp. 1183–1186.
- [15] V. I. MURASHKA, *On the \mathfrak{F} -hypercenter and the intersection of \mathfrak{F} -maximal subgroups of a finite group*, J. Group Theory, 21 (2018), pp. 463–473.
- [16] ———, *On Questions Posed by Shemetkov, Ballester-Bolínches, and Perez-Ramos in Finite Group Theory*, Math. Notes., 112 (2022), pp. 932–939.
- [17] ———, *Formations of finite groups in polynomial time: \mathfrak{F} -residuals and \mathfrak{F} -subnormality*, J. Symb. Comput., 122 (2024), p. 102271.
- [18] P. M. NEUMANN, *Some algorithms for computing with finite permutation groups*, in Proc. of Groups-St. Andrews 1985, E. Robertson and C. Campbell, eds., no. 121 in London Math. Soc. Lect. Note, Cambridge U. Press, 1987, pp. 59–92.
- [19] D. J. S. ROBINSON, *The structure of finite groups in which permutability is a transitive relation*, J. Austral. Math. Soc., 70 (2001), pp. 143–160.
- [20] L. RÓNYAI, *Computing the structure of finite algebras*, J. Symb. Comput., 9 (1990), pp. 355–373.
- [21] V. N. SEMENCHUCK AND A. F. VASIL'EV, *Studies of normaland subgroup structure of finite groups*, Nauka i tehnika, 1984, ch. Characterization of local formations \mathfrak{F} by properties of minimal non- \mathfrak{F} -groups, pp. 175–181.
- [22] Á. SERESS, *Permutation Group Algorithms*, Cambridge University Press, Cambridge, 2003.
- [23] L. A. SHEMETKOV, *Graduated formations of groups*, Mathematics of the USSR-Sbornik, 23 (1974), pp. 593–611.
- [24] ———, *Frattnini extensions of finite groups and formations*, Comm. Algebra, 25 (1997), pp. 955–964.
- [25] L. A. SHEMETKOV AND A. N. SKIBA, *Formations of algebraic systems*, Nauka, Moscow, 1989. In Russian.
- [26] A. N. SKIBA, *On the \mathfrak{F} -hypercentre and the intersection of all \mathfrak{F} -maximal subgroups of a finite group*, J. Pure Appl. Algebra, 216 (2012), pp. 789–799.
- [27] A. F. VASIL'EV, T. I. VASIL'EVA, AND V. N. TYUTYANOV, *On the finite groups of supersoluble type*, Sib. Math. J., 51 (2010), pp. 1004–1012.
- [28] A. F. VASIL'EV AND T. I. VASILYEVA, *On finite groups whose principal factors are simple groups*, Russian Math. (Iz. VUZ), 41 (1997), pp. 8–12.
- [29] V. A. VASILYEV, *Finite groups with submodular sylow subgroups*, Sib. Math. J., 56 (2015), pp. 1019–1027.
- [30] I. ZIMMERMANN, *Submodular subgroups in finite groups*, Math. Z., 202 (1989), pp. 545–557.