

Large deviations of Dyson Brownian motion on the circle and multiradial SLE_{0+}

Osama Abuzaid ^{*}, Vivian Olsiewski Healey [†], and Eveliina Peltola [‡]

Abstract

The main motivation of the present work is to investigate the asymptotic behavior as $\kappa \rightarrow 0+$ of multiradial Schramm-Loewner evolution, SLE_κ . We show that this process with the common parameterization satisfies a finite-time large deviation principle (LDP) in the Hausdorff metric with non-negative rate function, the multiradial Loewner energy. We also characterize the large-time behavior of curves with finite energy and zero energy (whose driving functions correspond to the trigonometric Calogero-Moser system).

The first half of this article is of independent interest regardless of SLE theory. It is devoted to proving a finite-time LDP for Dyson Brownian motion on the circle for a fixed number n of particles as the coupling parameter $\beta = 8/\kappa$ tends to ∞ . To our knowledge, in the literature large deviations of Dyson Brownian motion has only been considered for fixed β and as n tends to ∞ . While the non-Lipschitz drift precludes the application of the Freidlin-Wentzell theorem, we show that the rate function has the same form as in Freidlin-Wentzell theory for diffusions with uniformly Lipschitz drift.

In the second half of this article, we turn to proving an LDP for multiradial SLE_κ . Here, the main technical difficulty is that the SLE_κ curves have a common target point, preventing the usual configurational, or global, approach. Instead, we make careful use of the contraction principle from the LDP for Dyson Brownian motion (proven in the first part of the article), combined with topological results in Loewner theory: we show that finite-energy multiradial Loewner hulls are always disjoint unions of simple curves, except possibly at their common endpoint. A key to this is obtained from a derivative estimate for the radial Loewner map in terms of the energy of its driving function.

^{*}Department of Mathematics and Systems Analysis, Aalto University, Finland. osama.abuzaid@aalto.fi

[†]Department of Mathematics, Texas State University, US. healey@txstate.edu

[‡]Department of Mathematics and Systems Analysis, Aalto University, Finland; and Institute for Applied Mathematics, University of Bonn, Germany. eveliina.peltola@aalto.fi

Contents

1	Introduction	3
1.1	Large deviations of Dyson Brownian motion on the circle	5
1.2	Large deviations of multiradial SLE_{0+}	7
1.3	Finite-energy systems	9
2	LDP for Dyson Brownian motion on the circle	11
2.1	Dyson Brownian motion and n -radial Bessel process	11
2.2	Construction of n -radial Bessel process	12
2.3	Multiradial Dirichlet energy	14
2.4	LDP for Dyson Brownian motion on the circle	19
3	LDP for multiradial SLE_{0+}	22
3.1	Multiradial Loewner equation and multiradial SLE_{κ}	22
3.1.1	Multiradial SLE_{κ} , for $\kappa \in (0, 4]$	24
3.1.2	Time changes	25
3.2	Derivative estimate for finite-energy Loewner chains for $n = 1$	26
3.3	Finite-energy hulls are simple radial multichords	29
3.4	Proof of the LDP for multiradial SLE_{0+}	34
4	Large-time behavior of finite-energy systems	37
4.1	Zero-energy systems: existence, uniqueness, and asymptotics	37
4.2	Finite-energy systems	40

1 Introduction

Schramm-Loewner evolution (SLE_κ) is a natural model of a random interface arising from two-dimensional conformal geometry. SLE_κ curves have two equivalent characterizations: they can be defined in purely geometric and probabilistic terms (as curves satisfying conformal invariance and the domain Markov property), or they can be defined in terms of a one-parameter family of slit domains arising from the solutions to the Loewner equation with driving function $\sqrt{\kappa} B$, where B is a standard Brownian motion [Sch00]. These two perspectives are often referred to as the “configurational” (or “global”) and “dynamical” (or “local”) interpretations of SLE_κ , respectively. Their interplay allows for a rich theory that employs tools from diverse disciplines, including conformal geometry [LSW03, Wan19a], stochastic analysis [RS05, Dub07, MS16b], interacting particle systems [Car03, ABKM20], Teichmüller theory [Wan19b, Bis19], and algebraic geometry [PW24].

The roughness of SLE_κ curves depends on a parameter $\kappa \geq 0$. In particular, for different values of κ , variants of SLE_κ curves describe scaling limits of interfaces in a variety of statistical physics models (e.g., [LSW04, Smi06, Sch06, SS09]). The close relationship with discrete statistical physics models also allows discrete intuition and enumerative analysis to inform conjectures about SLE_κ itself, as applied to the theory of multiple SLEs in [KL07, BPW21, HL21]. Interestingly, SLE_κ curves are also very closely related to conformal field theory [BB03, BB04, FW03, FK04, CDR06, KS07, Dub15, Pel19], the Gaussian free field [Dub09, KM13, MS16a, She16], and random matrix theory [Car03, CLM23].

Natural variants of SLE_κ can be constructed from the so-called chordal SLE_κ by change of measure. For instance, multiple-curve chordal SLE_κ (where each curve connects two distinct boundary points) has been investigated in many works, including [BBK05, Dub07, KL07, Law09b, KP16, PW19, BPW21]. When $\kappa \in (0, 4]$, it is the measure absolutely continuous with respect to the product measure on n independent SLE_κ curves with Radon-Nikodym derivative

$$\mathbb{1}\{\gamma^j \cap \gamma^k = \emptyset \text{ for all } j \neq k\} \exp\left(\frac{c}{2} \sum_{i=2}^n \mu^{\text{loop}}[L_i(\gamma)]\right), \quad (1.1)$$

where $\mu^{\text{loop}}[L_i(\gamma)]$ is the Brownian loop measure of loops that intersect at least i of the curves $\gamma = (\gamma^1, \dots, \gamma^n)$, and c is a parameter known as the *central charge*,

$$c = \frac{(6 - \kappa)(3\kappa - 8)}{2\kappa}.$$

(See, e.g., [KL07, Law09b, PW19] for this definition, and [LSW04] for the construction of Brownian loop measure.) However, these measures are mutually singular when $\kappa = 0$.

Recently, a large deviation principle (LDP) for multichordal SLE_κ as $\kappa \rightarrow 0+$ was established in [PW24]. In that result, the convergence takes place in the Hausdorff metric, and the (good) rate function is termed the *multichordal Loewner energy*. The results in [PW24] have far-reaching applications. The authors show that there is a unique arrangement of curves that minimize the multichordal Loewner energy for given boundary data, and the union of these curves is the real locus of a real rational function, thus providing an alternate proof of the Shapiro conjecture in real enumerative geometry [Sot00, EG02]. The authors

also show that the Loewner potential (which differs from the Loewner energy by a function of the boundary data) has a simple expression in terms of zeta-regularized determinants of Laplacians (similar to the loop case in [Wan19b]) and is the semiclassical $c \rightarrow -\infty$ limit of certain CFT correlation functions (see [Dub15, KKP19, Pel19, ABKM20]).

In the present work, we investigate the asymptotic behavior as $\kappa \rightarrow 0+$ of multiradial SLE_κ (a multiple SLE in the disk where all curves have the origin as their common target point). We show that multiradial SLE_κ satisfies a finite-time LDP in the Hausdorff metric with good rate function that we call the *multiradial Loewner energy* (see Theorem 1.8). Notably, in contrast to the existing literature, establishing an LDP is technically much more difficult in the present multiradial case, for instance because the curves have a common target point. Indeed, we shall proceed by quite different methods¹ than in [PW24].

A key to our approach is that we work with *parameterized* curves — this difference in perspective is a result of the different way that multiple SLE_κ is constructed in the radial (in contrast to the chordal) setting. In the radial setting, the common target point causes essential difficulties for a “configurational” approach to defining multiple SLE_κ , since the Brownian loop measure in (1.1) blows up when curves intersect. This difficulty was addressed in the construction of multiradial SLE_κ in [HL21], whose main result is the construction of multiradial SLE_κ for $\kappa \leq 4$ as the solution to the multiradial Loewner equation for driving functions that evolve according to Dyson Brownian motion on the circle² with a particular repulsive strength. (See Definition 1.6 and Section 3.1 for details.)

The connection between multiradial SLE_κ and Dyson Brownian motion was first described by Cardy in the physics literature [Car03]. Loewner evolution driven by Dyson Brownian motion has gained recent interest in [Kat16, KK21, CM22, CLM23, FWY24]. An investigation of chordal Loewner evolution driven by a branching particle system (varying n) evolving according to Dyson Brownian motion for $\beta = \infty$ appears in [HM23]. However, asymptotic results linking SLE and Dyson Brownian motion have thus far focused on the setting where the number of curves tends to infinity (cf. [dMS16, HK18, HS21]).

The description of multiradial SLE_κ in terms of the corresponding driving functions provides the key tool in deriving the multiradial Loewner energy (Definition 1.5). Accordingly, much of the present work is devoted to proving an LDP for Dyson Brownian motion on the circle, which is of independent interest (see Theorem 1.2). Therefore, we have organized this article in such a way that, after the introduction of the main concepts and results, Sections 2 & 4 only address Dyson Brownian motion independently of Loewner theory (thus suitable for readers in a general probability audience), while Section 3 contains our main results in Loewner theory (assuming some familiarity with basic techniques in stochastic analysis and complex geometry). We will recall concepts from LDP theory along the way.

¹A finite-time parameterized single-curve LDP for chordal SLE_{0+} recently appeared in [Gus23] and is extended to infinite time in [AP24].

²We use “*Dyson Brownian motion on the circle*” to refer to the evolution of points on the circle, while the “*radial Bessel process*” refers to the evolution of the arguments of the same process. See Definition 2.1.

1.1 Large deviations of Dyson Brownian motion on the circle

Fix an integer $n \geq 1$. Let $(\mathbb{R}/2\pi\mathbb{Z})^n$ be the torus with periodic boundary conditions, and let \mathcal{X}_n denote the subset of elements admitting representatives $\boldsymbol{\theta} = (\theta^1, \dots, \theta^n)$ satisfying

$$\theta^1 < \theta^2 < \dots < \theta^n < \theta^1 + 2\pi. \quad (1.2)$$

Throughout, we use the convention that $\theta^{n+j} = \theta^j + 2\pi$ for all j . Let $C([0, \infty), \mathcal{X}_n)$ denote the space of continuous functions $\boldsymbol{\theta}_t = (\theta_t^1, \dots, \theta_t^n)$ from $[0, \infty)$ to \mathcal{X}_n . Consider the unique strong solution $\mathbf{U}_t^\kappa := (U_t^1, \dots, U_t^n)$ in $C([0, \infty), \mathcal{X}_n)$ to the system of SDEs³

$$dU_t^j = 2 \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + \sqrt{\kappa} dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}, \quad (1.3)$$

up to the collision time

$$\tau_{\text{coll}} := \inf \left\{ t \geq 0 : \min_{1 \leq j < k \leq n} |e^{iU_t^j} - e^{iU_t^k}| = 0 \right\}, \quad (1.4)$$

where W_t^1, \dots, W_t^n are independent Brownian motions. We shall see that the process $(e^{iU_t^1}, \dots, e^{iU_t^n})$ is nothing but a variant of Dyson Brownian motion on the circle (Section 2.1). In particular, the existence of a unique strong solution to (1.3) follows from the analogous result for Dyson Brownian motion [AGZ10]. While using the parameter κ in the context of Dyson Brownian motion is non-standard, our choice of \mathbf{U}^κ is strongly motivated by its connection to SLE_κ . Setting $\kappa = 0$ in (1.3) motivates the next definition.

Definition 1.1. The *multiradial Dirichlet energy*⁴ $J: C([0, \infty), \mathcal{X}_n) \rightarrow [0, +\infty]$ is the limit

$$J(\boldsymbol{\theta}) := \lim_{T \rightarrow \infty} J_T(\boldsymbol{\theta}) \in [0, +\infty], \quad \boldsymbol{\theta} \in C([0, \infty), \mathcal{X}_n),$$

where for each $T > 0$, the (*truncated*) *multiradial Dirichlet energy* of $\boldsymbol{\theta}$ is

$$J_T(\boldsymbol{\theta}) := \begin{cases} \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi^j(\boldsymbol{\theta}_s) \right|^2 ds, & \text{if } \boldsymbol{\theta} \text{ is absolutely continuous on } [0, T], \\ \infty, & \text{otherwise,} \end{cases}$$

$$\text{where } \phi^j(\boldsymbol{\theta}) := \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{\theta^j - \theta^k}{2}\right). \quad (1.5)$$

Analogously, for each $a > 0$, the *multiradial Dirichlet energy with parameter a* is defined in terms of J_T^a by replacing in (1.5) the function $\phi^j(\boldsymbol{\theta})$ by $\phi_a^j(\boldsymbol{\theta}) := \frac{a}{4} \phi^j(\boldsymbol{\theta}) = \frac{a}{4} \sum_{k \neq j} \cot\left(\frac{\theta^j - \theta^k}{2}\right)$.

The original study of Dyson Brownian motion dates back to [Dys62], one of the founding articles of random matrix theory. For fixed $n \in \mathbb{N}$ and $\beta \geq 1$, Dyson Brownian motion may be defined as the unique strong solution (X_t^1, \dots, X_t^n) in the Weyl chamber of type A_{n-1} ,

$$\{(x^1, \dots, x^n) \in \mathbb{R}^n \mid x^1 < x^2 < \dots < x^n\},$$

³Notice that Equation (1.3) is Equation (2.2) from Definition 2.1 with $\alpha = 4/\kappa$.

⁴This is the usual Dirichlet energy when $n = 1$ (see Equation (2.16)).

to the SDEs

$$dX_t^j = \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \frac{dt}{X_t^j - X_t^k} + \sqrt{\frac{2}{\beta}} dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}. \quad (1.6)$$

In the present work, we consider the analogue of this process on the circle, where the radial Bessel-type equation (1.3) plays the role of Equation (1.6). In particular, after a suitable time change (see Section 2.1), we see that the relationship between β and κ is

$$\beta = \frac{8}{\kappa}, \quad (1.7)$$

which matches the prediction of Cardy from the physics literature [Car03].

For particular values of β , Dyson Brownian motion describes the evolution of the ordered eigenvalues of symmetric, Hermitian, and symplectic matrix Brownian motions (corresponding to the self-dual Gaussian ensembles GOE, GUE, and GSE, for $\beta = 1, 2, 4$, respectively — see [AGZ10, Chapter 4]). For $\beta = 2$, Dyson Brownian motion has the same law as n independent Brownian motions conditioned on nonintersection [KT03]. Furthermore, for general $\beta \in (0, +\infty]$ one can construct ensembles of Jacobi matrices whose eigenvalues correspond to (1.6), see [DE02, GK20]. It would be particularly interesting to investigate the fluctuations near the large deviation limit of the Dyson Brownian motion (1.6) in the sense of Theorem 1.2 and its relation with the $\beta = \infty$ process considered in [GK20]. Lastly, let us remark that a new geometric construction of Dyson Brownian motion for general $\beta \in (0, +\infty]$ has recently appeared in [HIM23] — interestingly, this construction relies on tools from Riemannian geometry and mean curvature flow.

In Section 2, we prove a large deviation principle for \mathbf{U}^κ satisfying (1.3) as $\kappa \rightarrow 0+$, with good⁵ rate function being the multiradial Dirichlet energy J_T (Definition 1.1). Our result also implies an LDP for Dyson Brownian motion on the circle. To state the result, for fixed $T \in (0, \infty)$ and $\boldsymbol{\theta}_0 \in \mathcal{X}_n$ we denote by $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$ the space of continuous functions $\boldsymbol{\theta}$ from $[0, T]$ to \mathcal{X}_n started at $\boldsymbol{\theta}_0$. We endow $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$ with the metric

$$d_{[0, T]}(\boldsymbol{\theta}, \boldsymbol{\omega}) := \sup_{t \in [0, T]} |\boldsymbol{\theta}_t - \boldsymbol{\omega}_t| = \sup_{t \in [0, T]} \left((\theta_t^1 - \omega_t^1)^2 + \dots + (\theta_t^n - \omega_t^n)^2 \right)^{1/2}. \quad (1.8)$$

Then, $(C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n), d_{[0, T]})$ is a Polish space (as a separable complete metric space).

Theorem 1.2 (LDP for Dyson Brownian motion on the circle). *Fix $T \in (0, \infty)$. Let \mathbf{U}^κ be the unique strong solution to (1.3), started at $\mathbf{U}_0^\kappa = \boldsymbol{\theta}_0 \in \mathcal{X}_n$. The family $(\mathbf{P}^\kappa)_{\kappa > 0}$ of laws induced by \mathbf{U}^κ satisfies the following LDP in $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$ with good rate function J_T :*

For any closed subset F and open subset O of $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$, we have

$$\overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^\kappa[\mathbf{U}^\kappa \in F] \leq - \inf_{\boldsymbol{\theta} \in F} J_T(\boldsymbol{\theta}), \quad (1.9)$$

$$\underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^\kappa[\mathbf{U}^\kappa \in O] \geq - \inf_{\boldsymbol{\theta} \in O} J_T(\boldsymbol{\theta}). \quad (1.10)$$

⁵A good rate function is a rate function for which all level sets are compact.

In fact, in Section 2.4 we prove a more general result, Theorem 2.12, involving the multiradial Dirichlet energy with parameter a . Theorem 1.2 is the special case of this with $a = 4$. Allowing a general parameter $a > 0$ is useful, e.g., in applications to SLE variants. The proof of the LDP is a careful application of Varadhan’s lemma (Lemma D) relying on properties of the multiradial Dirichlet energy derived in Section 2.3, to transport the well-known LDP of Brownian motion from Schilder’s theorem (Theorem C). Note that Schilder’s theorem also gives the basic case of $n = 1$ of Theorem 1.2.

Remark 1.3. Theorem 1.2 is stated for the radial Bessel process \mathbf{U}^κ , but it is equivalent to an LDP for Dyson Brownian motion on the circle as $\beta = \frac{8}{\kappa} \rightarrow \infty$, by considering $\exp(i\mathbf{U}_t^\kappa)$ and applying the contraction principle (Theorem E) to the continuous function⁶ $-i \log(\cdot)$.

Large deviation results for Dyson Brownian motion (for fixed β) as $n \rightarrow \infty$ have been considered, e.g. in [GZ02, GZ02], and are closely connected to random matrix theory. In contrast, our Theorem 1.2 holds for fixed n as $\beta \rightarrow \infty$, thereby filling a gap in the literature.

More generally, large deviation theory for stochastic differential equations (or SPDEs) is a field of independent interest, and Theorem 1.2 fits very naturally in this context. The multiradial Dirichlet energy J_T is exactly the rate function that would be predicted by applying Freidlin-Wentzell theory to the diffusion (1.3) (see [FW84], originally published in Russian in 1979, and [DZ10] for a survey). However, since the drift in (1.3) is not uniformly Lipschitz continuous, the Freidlin-Wentzell theorem does not apply directly. Our Theorem 1.2 can thus be considered as an extension of the Freidlin-Wentzell theorem to a diffusion with non-Lipschitz drift.

1.2 Large deviations of multiradial SLE₀₊

We will mainly work on Loewner flows in the following setup. The *multiradial Loewner equation* with the common parameterization is the solution to the boundary value problem

$$\partial_t g_t(z) = g_t(z) \sum_{j=1}^n \frac{z_t^j + g_t(z)}{z_t^j - g_t(z)}, \quad g_0(z) = z, \quad z \in \overline{\mathbb{D}}, \quad t \geq 0, \quad (1.11)$$

where $z_t^1, \dots, z_t^n \in \partial\mathbb{D}$ are cyclically ordered points on the unit circle, called the *driving functions*. It is most common to study (1.11) for driving functions that are non-intersecting and continuous in time, in which case the mappings g_t that satisfy (1.11) generate a locally growing family of compact subsets K_t of $\overline{\mathbb{D}}$ (hulls⁷) satisfying $g_t(\mathbb{D} \setminus K_t) = \mathbb{D}$. In fact, each $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ is the unique conformal mapping that satisfies $g_t(0) = 0$ and $g_t'(0) > 0$. Throughout, we refer to this map as the *uniformizing map normalized at the origin*.

The parameterization in (1.11) guarantees that $g_t'(0) = e^{nt}$. If each hull K_t is a union of n disjoint connected components (e.g., as in Figure 3.1), the “common parameterization” implies that, roughly, each component is locally growing at the same rate (see [HL21] for more details). See also Equation (3.1) for a more general case involving weights.

It will be convenient to use the angle coordinates $z_t^j = \exp(i\theta_t^j)$, where the driving function becomes $\boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in C([0, \infty), \mathcal{X}_n)$. For each $t \geq 0$, the *Loewner transform*

⁶Throughout, we use the principal branch of the logarithm, so that angles are taken to lie in $[0, 2\pi)$.

⁷A hull is a compact set $K \subset \overline{\mathbb{D}}$ such that $\mathbb{D} \setminus K$ is simply connected, $0 \in \mathbb{D} \setminus K$, and $\overline{K} \cap \overline{\mathbb{D}} = K$.

$\mathcal{L}_t: C([0, t], \mathcal{X}_n) \rightarrow \mathcal{C}$ sends driving functions to hulls,

$$K_t = \mathcal{L}_t(\boldsymbol{\theta}) := \{z \in \overline{\mathbb{D}} : \tau_z \leq t\} \subset \mathcal{C}, \quad (1.12)$$

where \mathcal{C} is the set of non-empty compact subsets of $\overline{\mathbb{D}}$, and τ_z is the swallowing time of z ,

$$\tau_z := \sup \left\{ t \geq 0 : \inf_{s \in [0, t]} \min_{1 \leq j \leq n} |g_s(z) - e^{i\theta_s^j}| > 0 \right\} \in [0, +\infty].$$

We endow \mathcal{C} with the Hausdorff metric $d_{\mathcal{C}}: \mathcal{C} \times \mathcal{C} \rightarrow [0, \infty)$ defined by

$$d_{\mathcal{C}}(K, K') := \inf \{ \delta > 0 : K \subset \mathcal{B}_{K'}(\delta) \text{ and } K' \subset \mathcal{B}_K(\delta) \}, \quad (1.13)$$

where $\mathcal{B}_K(\delta) := \bigcup_{x \in K} \mathcal{B}_x(\delta)$. Then, $(\mathcal{C}, d_{\mathcal{C}})$ is a compact metric space.

Definition 1.4. Fix distinct points $x^1, \dots, x^n \in \partial\mathbb{D}$. We call an n -tuple $\boldsymbol{\gamma} = (\gamma^1, \dots, \gamma^n)$ such that $\gamma^1, \dots, \gamma^n$ are curves⁸ in $\overline{\mathbb{D}}$ with $\gamma^j(0) = x^j$, and $\lim_{t \rightarrow \infty} \gamma_t^j = 0$ for each j , a *radial multichord* in $(\mathbb{D}; x^1, \dots, x^n)$. We naturally identify $\boldsymbol{\gamma}$ with the union $\bigcup_j \gamma^j \in \mathcal{C}$.

Definition 1.5. For each $T \in (0, \infty)$, we define the (*truncated*) *multiradial Loewner energy* of a radial multichord to be the Dirichlet energy of its driving function in the common parameterization (Definition 1.1). More generally, we define the energy functional $I_T: \mathcal{C} \rightarrow [0, +\infty]$ on the metric space $(\mathcal{C}, d_{\mathcal{C}})$ by

$$I_T(K) := \inf_{\boldsymbol{\theta} \in \mathcal{L}_T^{-1}(K)} J_T(\boldsymbol{\theta}), \quad T > 0, \quad K \in \mathcal{C}. \quad (1.14)$$

Note that $I_T(K) = \infty$ if there is no driving function that generates K in time T in the common parameterization. We can also define the energy in a general domain D by conformal equivalence using a conformal mapping $\varphi: D \rightarrow \mathbb{D}$ with $\varphi(0) = 0$ and $\varphi'(0) > 0$:

$$I_T(\tilde{K}; D) := I_T(K), \quad \text{where } \tilde{K} \subset \overline{D} \text{ and } \varphi(\tilde{K}) = K \subset \overline{\mathbb{D}}.$$

We next give the definition of n -radial SLE_{κ} that we will use for the remainder of this work. Remark 3.5 and the discussion following it in Section 3.1 offer additional justification for this definition and comparison to [HL21].

Definition 1.6. Fix $\boldsymbol{\theta}_0 \in \mathcal{X}_n$ and $\mathbf{z}_0 = (e^{i\theta_0^1}, \dots, e^{i\theta_0^n})$. For each parameter $0 < \kappa \leq 4$, n -radial SLE_{κ} with the common parameterization started from \mathbf{z}_0 is the random radial multichord $\boldsymbol{\gamma}_t$ for which the uniformizing conformal mappings $g_t: \mathbb{D} \setminus \boldsymbol{\gamma}_t \rightarrow \mathbb{D}$ satisfy Equation (1.11) with driving functions $z_t^j = e^{iU_t^j}$ for $j = 1, \dots, n$, where $\mathbf{U}_t^{\kappa} = (U_t^1, \dots, U_t^n)$ is the unique strong solution in $C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$ to the SDEs (1.3) started at $\mathbf{U}_0^{\kappa} = \boldsymbol{\theta}_0$. Note that we have $\tau_{\text{coll}} = +\infty$ almost surely in this case.

Remark 1.7. A key aspect of Definition 1.6 is the specific weight “2” in front of the drift term in (1.3). Other values of this weight give rise to other variants of SLE, including so-called *locally independent* SLE_{κ} when the drift is instead multiplied by “1”, see [HL21]. The reason to define multiradial SLE_{κ} using (1.3) (rather than another drift) is that this is the drift strength that appears when considering large time T truncations of the chordal Radon-Nikodym derivative (1.1) and then taking $T \rightarrow \infty$. See also Section 3.1, where we discuss Loewner evolutions with various weight functions and Remark 3.6 concerning SLE with spiral.

⁸Note that the definition of a radial multichord allows the curves to intersect.

Our main result is the following finite-time LDP for multiradial SLE_{0+} .

Theorem 1.8 (LDP for multiradial SLE). *Fix $T \in (0, \infty)$. The initial segments $\gamma_{[0,T]}^\kappa \in \mathcal{C}$ of multiradial SLE_κ curves satisfy the following LDP in \mathcal{C} with good rate function I_T :*

For any Hausdorff-closed subset F and Hausdorff-open subset O of \mathcal{C} , we have

$$\overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{P}^\kappa [\gamma_{[0,T]}^\kappa \in F] \leq - \inf_{K \in F} I_T(K), \quad (1.15)$$

$$\underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{P}^\kappa [\gamma_{[0,T]}^\kappa \in O] \geq - \inf_{K \in O} I_T(K). \quad (1.16)$$

We prove Theorem 1.8 in Section 3.4. To this end, the idea is to make careful use of the contraction principle and derive Theorem 1.8 from the LDP for Dyson Brownian motion (Theorem 1.2). The usage of the contraction principle will be enabled by topological results concerning Loewner theory and finite-energy hulls. These results enable us to essentially disregard the discontinuities of the Loewner transform. We show that finite-energy multiradial Loewner hulls are always disjoint unions of simple curves (see Theorem 1.9 below). The strategy is to first derive a derivative estimate for the single-chord radial Loewner map in terms of the energy of its driving function (see Theorem 3.9), and then to use complex analysis techniques to pull this result to the case of several curves.

Theorem 1.9. *Consider a multiradial Loewner chain with the common parameterization for which the uniformizing conformal mappings $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ satisfy Equation (1.11) with driving functions $z_t^j = e^{i\theta_t^j}$ for $j = 1, \dots, n$, where $\boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$. If $J_T(\boldsymbol{\theta}) < \infty$, then the hull $K_T = \mathcal{L}_T(\boldsymbol{\theta})$ consists of n pairwise disjoint simple curves.*

We prove Theorem 1.9 in Section 3.3. The key inputs are the derivative estimate in the case of $n = 1$ (Theorem 3.9 in Section 3.2), which is a weighted, radial generalization of a result appearing in [FS17], and a sort of generalized conformal restriction property (see Proposition 3.12), which we will utilize to pull the $n = 1$ result to general $n \geq 2$.

1.3 Finite-energy systems

Finally, in Section 4 we analyze the interacting particle system corresponding to finite-energy driving functions. In Theorem 1.10, we characterize the large-time behavior of the driving functions of finite-energy curves: these functions eventually approach an equally-spaced configuration (1.17). For zero-energy systems, the limit is static, see Equation (1.18).

When considering zero-energy systems in the context of Hamiltonian dynamics, the associated particle system is called the trigonometric Calogero-Moser system; its study dates back to the original articles [Cal71, Mos75]. Connections between chordal SLE and other Calogero-Moser systems have appeared recently in [ABKM20]. The existence and uniqueness (up to rotation) of a stable equilibrium for the trigonometric Calogero-Moser system has been considered, e.g., in [Mul11], though our proofs were developed independently. Instead of leveraging the connection to Hamiltonian dynamics, our approach depends on explicit analysis of the deterministic PDE obtained by setting $\kappa = 0$ in (1.3). The existence and uniqueness of the zero-energy flow for each starting point $\boldsymbol{\theta}_0$ is stated in Proposition 4.2 in Section 4.1.

Theorem 1.10 (Asymptotic configuration of finite-energy systems). *Fix an integer $n \geq 2$. Consider a function $\theta \in C_{\theta_0}([0, \infty), \mathcal{X}_n)$. If $J(\theta) < \infty$, then we have*

$$\lim_{t \rightarrow \infty} (\theta_t^{j+1} - \theta_t^j) = \frac{2\pi}{n}, \quad \text{for all } j \in \{1, \dots, n\}. \quad (1.17)$$

Furthermore, if $J(\theta) = 0$, then there exists $\zeta \in [0, 2\pi)$ such that

$$\lim_{t \rightarrow \infty} \theta_t = (\zeta, \zeta + \frac{2\pi}{n}, \dots, \zeta + \frac{(n-1)2\pi}{n}), \quad (1.18)$$

and the convergence is exponentially fast with exponential rate n .

We prove Theorem 1.10 in Section 4.2, where we also discuss the rate of convergence for finite-energy systems (Remark 4.9 & Proposition 4.10).

To understand what this result means for the zero-energy curves, let the angle $\zeta \in [0, 2\pi)$ be fixed, and let θ^ζ denote the constant configuration

$$\theta_t^\zeta \equiv \theta^\zeta := (\zeta, \zeta + \frac{2\pi}{n}, \dots, \zeta + \frac{(n-1)2\pi}{n}), \quad \text{for all } t \geq 0.$$

By symmetry, we see that the constant driving functions $\exp(i\theta_t^\zeta)$ generate the “pizza pie” configuration of curves: the union of straight lines in $\overline{\mathbb{D}}$ from the points $\exp(i(\zeta + 2\pi j/n))$, $j = 1, \dots, n$, to the origin. Theorem 1.10 implies that for large enough times, the zero-energy driving functions approach this configuration. Thus, we expect that the union of curves $g_T(\gamma_{[T, \infty)})$ approaches the pizza pie configuration, though we do not prove this in the present article. Understanding the asymptotic configurations of general finite-energy systems in detail will be key for establishing an infinite-time LDP for radial SLE processes.

Acknowledgments

- Part of this project was initiated while V.O.H. and E.P. participated in a program hosted by the Mathematical Sciences Research Institute (MSRI) in Berkeley, California, in Spring 2022, supported by the National Science Foundation (Grant No. DMS-1928930).
- Part of this project was performed while E.P. was visiting the Institute for Pure and Applied Mathematics (IPAM), which is supported by the National Science Foundation (Grant No. DMS-1925919).
- O.A. is supported by the Academy of Finland grant number 340461 “Conformal invariance in planar random geometry.”
- V.O.H. is partially supported by an AMS Simons Research Enhancement Grant for PUI Faculty.
- This material is part of a project that has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (101042460): ERC Starting grant “Interplay of structures in conformal and universal random geometry” (ISCoURaGe) and from the Academy of Finland grant number 340461 “Conformal invariance in planar random geometry.” E.P. is also supported by the Academy of Finland Centre of Excellence Programme grant number

346315 “Finnish centre of excellence in Randomness and Structures (FiRST)” and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy EXC-2047/1-390685813, as well as the DFG collaborative research centre “The mathematics of emerging effects” CRC-1060/211504053.

- V.O.H. would like to thank Nestor Guillen for useful conversations about PDEs.

2 LDP for Dyson Brownian motion on the circle

The main result of this section is Theorem 1.2, a finite-time LDP for the n -radial Bessel process (equivalently, for Dyson Brownian motion on the circle). We will justify the upper and lower bounds (1.9, 1.10) separately, by applying Varadhan’s lemma (Lemma D) somewhat similarly as in [PW24, Proof of Theorem 5.11], relying on Schilder’s theorem for Brownian motion (Theorem C) as key input. We however first need to control the difference of the rate function J_T to the usual Dirichlet energy E_T appearing in Schilder’s theorem — see in particular Definition 2.5 and Lemmas 2.6 & 2.8. Moreover, because the change of measure from independent Brownian motions to the n -radial Bessel process contains a factor that is not uniformly bounded, we need a specific tail estimate (Lemma 2.7).

Before addressing the proof of the main Theorem 1.2, we compare our setup to the literature in Sections 2.1 and 2.2 — indeed, the setup in Theorem 1.2 with the SDEs (1.3) is equivalent to that for the more commonly considered Dyson Brownian motion and n -radial Bessel process (see Proposition 2.2). We then gather known facts from [HL21, Section 5] in Section 2.2, and proceed to the main results in Sections 2.3–2.4.

2.1 Dyson Brownian motion and n -radial Bessel process

Definition 2.1 ([HL21]). The n -radial Bessel process on $\frac{1}{2}\mathcal{X}_n := \{\boldsymbol{\theta} \mid 2\boldsymbol{\theta} \in \mathcal{X}_n\}$ with parameter $\alpha \in \mathbb{R}$ is the process $\boldsymbol{\Theta}_t^\alpha = (\Theta_t^1, \dots, \Theta_t^n)$ satisfying

$$d\Theta_t^j = \alpha \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot(\Theta_t^j - \Theta_t^k) dt + dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}, \quad (2.1)$$

where W_t^1, \dots, W_t^n are independent Brownian motions. *Dyson Brownian motion on the circle* is the process $e^{2i\boldsymbol{\Theta}_t^\alpha} = (e^{2i\Theta_t^1}, \dots, e^{2i\Theta_t^n})$. Note that $\mathbf{U}_t^\kappa := 2\boldsymbol{\Theta}_{\kappa t/4}^\alpha \in \mathcal{X}_n$ satisfies

$$dU_t^j = \frac{\alpha\kappa}{2} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + \sqrt{\kappa} dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}. \quad (2.2)$$

In particular, the SDE (1.3) appearing in our main Theorem 1.2 is (2.2) with $\alpha = 4/\kappa$. More generally, if $\alpha = a/\kappa$, for $a > 0$, as in Theorem 2.12, then the equivalent SDE is

$$dU_t^j = \frac{a}{2} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + \sqrt{\kappa} dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}. \quad (2.3)$$

The half-angle convention in the definition of the n -radial Bessel process is also convenient for direct comparison with Dyson Brownian motion (X_t^1, \dots, X_t^n) on the real line, which satisfies the SDEs (1.6). For random matrix theory applications, this process is more commonly written using the time change $\tilde{X}_t^j = X_{t/n}^j$ so that (1.6) is equivalent to

$$d\tilde{X}_t^j = \frac{1}{n} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \frac{1}{\tilde{X}_t^j - \tilde{X}_t^k} dt + \sqrt{\frac{1}{n}} \sqrt{\frac{2}{\beta}} dW_t^j, \quad \text{for all } j \in \{1, \dots, n\}.$$

Using $\alpha = 4/\kappa$ in Definition 2.1 we find the relationship $\beta = 8/\kappa$ between β and κ [Car03].

Dyson Brownian motion on the circle is also referred to in the literature as the *Dyson circular ensemble* [FWY24]. Although Definition 2.1 holds for any α , we will restrict our attention to $\alpha \geq 1$, which corresponds to $\kappa \leq 4$ in Equation (1.7). Comparison to the usual Bessel process shows that $\alpha = 1$ corresponds to the phase transition for recurrence and transience. The existence of a unique strong solution to (2.1) (actually for any $\alpha \geq \frac{1}{2}$) follows from the analogous result for Dyson Brownian motion [AGZ10] (see also [Law24]).

2.2 Construction of n -radial Bessel process

We now recall the construction of n -radial Bessel process from [HL21]. We include a few of the computations that justify the definition, since we use a different parameterization and other slightly different notation. We begin by defining the non-negative functions

$$F(\boldsymbol{\theta}) := \prod_{1 \leq j < k \leq n} \left| \sin \left(\frac{\theta^k - \theta^j}{2} \right) \right|, \quad (2.4)$$

$$\psi^j(\boldsymbol{\theta}) := \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \csc^2 \left(\frac{\theta^j - \theta^k}{2} \right), \quad (2.5)$$

$$\psi(\boldsymbol{\theta}) := \sum_{j=1}^n \psi^j(\boldsymbol{\theta}) = 2 \sum_{1 \leq j < k \leq n} \csc^2 \left(\frac{\theta^j - \theta^k}{2} \right), \quad \boldsymbol{\theta} \in \mathcal{X}_n. \quad (2.6)$$

The next two results come from [HL21], though they are stated there with different notation and a different parameterization convention⁹. Since these differences make cross-referencing cumbersome, we include short proof sketches here.

Suppose that $\mathbf{B}_t = (B_t^1, \dots, B_t^n)$ is an n -dimensional standard Brownian motion in \mathbb{R}^n defined on the filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{P})$, where \mathcal{F}_\bullet is its natural right-continuous completed filtration. Fix $\boldsymbol{\theta}_0 \in \mathcal{X}_n$ and define $\mathbf{U}_t^\kappa = (U_t^1, \dots, U_t^n) \in \mathcal{X}_n$ by

$$\mathbf{U}_t^\kappa = \boldsymbol{\theta}_0 + \sqrt{\kappa} \mathbf{B}_t, \quad \mathbf{U}_0^\kappa = \boldsymbol{\theta}_0, \quad (2.7)$$

stopped at the collision time

$$\tau_{\text{coll}} := \inf \left\{ t \geq 0 : \min_{1 \leq j < k \leq n} |e^{iU_t^j} - e^{iU_t^k}| = 0 \right\} = \inf \{ t \geq 0 : \mathbf{U}_t^\kappa \notin \mathcal{X}_n \}.$$

⁹Later, we will set $\alpha = a/\kappa$ (with the case $\alpha = 4/\kappa$ playing a central role), but for now we use the general parameter α in the interest of recording the version of the formulas and computations that corresponds to this choice of parameterization, which is more standard in the literature.

Proposition A ([HL21]). Fix $\kappa > 0$ and $\alpha \in \mathbb{R}$. Write $F_\alpha(\boldsymbol{\theta}) := (F(\boldsymbol{\theta}))^\alpha$. Define

$$M_t^{\kappa,\alpha} := F_\alpha(\mathbf{U}_t^\kappa) \exp\left(-\frac{1}{2} \int_0^t \frac{\Delta F_\alpha(\mathbf{U}_s^\kappa)}{F_\alpha(\mathbf{U}_s^\kappa)} ds\right), \quad t < \tau_{\text{coll}}. \quad (2.8)$$

Then, the process $M_t^{\kappa,\alpha}$ is a local martingale on $(\Omega, \mathcal{F}_t, \mathbb{P})$, satisfying

$$\frac{dM_t^{\kappa,\alpha}}{M_t^{\kappa,\alpha}} = \alpha\sqrt{\kappa} \sum_{1 \leq j < k \leq n} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dB_t^j. \quad (2.9)$$

Also, if $\alpha \geq \frac{1}{2}$, then $M_t^{\kappa,\alpha}$ is a martingale and $\tau_{\text{coll}} = \infty$ almost surely.

Proof sketch. Differentiating F_α with respect to θ^j and using the notations ϕ^j and ψ^j from (1.5, 2.5), we see that

$$\frac{\partial_j F_\alpha(\mathbf{U}_t^\kappa)}{F_\alpha(\mathbf{U}_t^\kappa)} = \frac{\alpha\sqrt{\kappa}}{2} \phi^j(\mathbf{U}_t^\kappa), \quad (2.10)$$

$$\frac{\partial_j^2 F_\alpha(\mathbf{U}_t^\kappa)}{F_\alpha(\mathbf{U}_t^\kappa)} = \frac{\alpha^2 \kappa}{4} (\phi^j(\mathbf{U}_t^\kappa))^2 - \frac{\alpha\kappa}{4} \psi^j(\mathbf{U}_t^\kappa), \quad j = 1, \dots, n. \quad (2.11)$$

This gives the claim by Itô's formula. \square

Corollary B ([HL21]). Fix $\kappa > 0$ and $\alpha \in \mathbb{R}$. Let $\mathbb{P}^{\kappa,\alpha}$ be the probability measure that is absolutely continuous with respect to \mathbb{P} with Radon-Nikodym derivative

$$\frac{d\mathbb{P}_t^{\kappa,\alpha}}{d\mathbb{P}_t} = \frac{M_t^{\kappa,\alpha}}{M_0^{\kappa,\alpha}}, \quad t < \tau_{\text{coll}}.$$

Then, the process $\mathbf{U}_t^\kappa = (U_t^1, \dots, U_t^n) \in C_{\theta_0}([0, \infty), \mathcal{X}_n)$ satisfies

$$dU_t^j = \frac{\alpha\kappa}{2} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + \sqrt{\kappa} dW_t^j, \quad t < \tau_{\text{coll}}, \quad (2.12)$$

where W_t^1, \dots, W_t^n are independent Brownian motions with respect to $\mathbb{P}^{\kappa,\alpha}$.

Proof sketch. By the Girsanov theorem, Equation (2.9) implies that the B_t^j satisfy under \mathbb{P}

$$dB_t^j = \frac{\alpha\sqrt{\kappa}}{2} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + dW_t^j,$$

and Equation (2.12) then follows from the definition (2.7) of \mathbf{U}^κ . \square

Let $\mathbb{P}^{\kappa,a}$ be the measure absolutely continuous with respect to \mathbb{P} with Radon-Nikodym derivative obtained from the (local) martingale $M_t^{\kappa,\alpha}$ of Proposition A with $\alpha = a/\kappa$:

$$\frac{d\mathbb{P}_t^{\kappa,a}}{d\mathbb{P}_t} = \frac{M_t^{\kappa,a/\kappa}}{M_0^{\kappa,a/\kappa}}, \quad t < \tau_{\text{coll}}.$$

Combining this change of measure with Corollary B, we can formulate the existence and uniqueness of the process \mathbf{U}^κ satisfying the SDEs (1.3), central to the present work.

Proposition 2.2. *Suppose that $\mathbf{B}_t = (B_t^1, \dots, B_t^n)$ is an n -dimensional standard Brownian motion in \mathbb{R}^n defined on the filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{P})$, where \mathcal{F}_\bullet is its natural right-continuous completed filtration. Fix $\boldsymbol{\theta}_0 \in \mathcal{X}_n$ and define $\mathbf{U}_t^\kappa = (U_t^1, \dots, U_t^n) \in \mathcal{X}_n$ by*

$$\mathbf{U}_t^\kappa = \boldsymbol{\theta}_0 + \sqrt{\kappa} \mathbf{B}_t, \quad \mathbf{U}_0^\kappa = \boldsymbol{\theta}_0.$$

Then, \mathbf{U}_t^κ is the unique strong solution to the system of SDEs (1.3) in the measure $\mathbb{P}^{\kappa,4}$.

Proof. Setting $\alpha = 4/\kappa$ (so $a = 4$), we see that $\mathbb{P}_t^{\kappa,4} = \mathbb{P}_t^{\kappa,\alpha}$ from Corollary B up to the collision time (1.4). In particular, by (2.12) the processes U_t^j satisfy SDEs (1.3) where W_t^1, \dots, W_t^n are independent Brownian motions with respect to the measure $\mathbb{P}^{\kappa,4}$. \square

Remark 2.3. For each additional parameter $\rho \in \mathbb{R}$, one can similarly define an n -radial Bessel process *with spiraling rate* ρ as the measure $\mathbb{P}^{\kappa,\alpha,\rho}$ absolutely continuous with respect to \mathbb{P} with Radon-Nikodym derivative obtained from the (local) martingale

$$M_t^{\kappa,\alpha,\rho} := F_{\alpha,\rho}(\mathbf{U}_t^\kappa) \exp\left(-\frac{1}{2} \int_0^t \frac{\Delta F_{\alpha,\rho}(\mathbf{U}_s^\kappa)}{F_{\alpha,\rho}(\mathbf{U}_s^\kappa)} ds\right), \quad t < \tau_{\text{coll}}, \quad (2.13)$$

where

$$F_{\alpha,\rho}(\boldsymbol{\theta}) = F_\alpha(\boldsymbol{\theta}) \exp\left(\frac{\rho}{\kappa} \sum_{j=1}^n \theta^j\right),$$

and the process $\mathbf{U}_t^\kappa = (U_t^1, \dots, U_t^n) \in C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$ satisfies

$$dU_t^j = \frac{\alpha\kappa}{2} \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot\left(\frac{U_t^j - U_t^k}{2}\right) dt + \rho dt + \sqrt{\kappa} dW_t^j, \quad t < \tau_{\text{coll}}, \quad (2.14)$$

where W_t^1, \dots, W_t^n are independent Brownian motions with respect to $\mathbb{P}^{\kappa,\alpha,\rho}$. With the parameter choice $\alpha = 4/\kappa$, this process generates spiraling multiradial SLE_κ curves (see Remark 3.6 and [MS17, WW24]). As a consequence of our Theorem 2.12, the n -radial Bessel process with spiraling rate ρ also satisfies an LDP with the good rate function

$$J_T^{a,\rho}(\boldsymbol{\theta}) := \begin{cases} \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - (2\phi_a^j(\boldsymbol{\theta}_s) + \rho) \right|^2 ds, & \text{if } \boldsymbol{\theta} \text{ is absolutely continuous on } [0, T], \\ \infty, & \text{otherwise.} \end{cases}$$

This follows, for example, by applying the contraction principle (Theorem E) to the continuous map sending the function $t \mapsto \boldsymbol{\theta}_t$ to the function $t \mapsto \boldsymbol{\theta}_t + \rho t$.

2.3 Multiradial Dirichlet energy

From Proposition 2.2, we learn that the Dyson-type process \mathbf{U}_t^κ solving (1.3), equivalent to the n -radial Bessel process by (2.2), and to the Dyson Brownian motion on the circle via $\exp(i\mathbf{U}_t^\kappa) = \exp(2i \boldsymbol{\Theta}_{\kappa t/4}^\alpha)$, is a Girsanov transform of n -dimensional standard Brownian motion \mathbf{B} . From Schilder's classical theorem, one readily obtains an LDP for n -dimensional Brownian motion \mathbf{B} , whose components are independent (Theorem C).

Denote by $C_0([0, T], \mathbb{R}^n)$ the space of continuous functions $\boldsymbol{\theta}: [0, T] \rightarrow \mathbb{R}^n$ started at $\boldsymbol{\theta}_0 = \mathbf{0}$, equipped with the supremum norm $\|\boldsymbol{\theta}\|_{[0, T]} := \sup_{t \in [0, T]} |\boldsymbol{\theta}_t|$. The rate function is the n -dimensional Dirichlet energy

$$E_T(\boldsymbol{\theta}) := \sum_{j=1}^n E_T(\theta^j), \quad \boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in C_0([0, T], \mathbb{R}^n), \quad (2.15)$$

where $E_T(\theta)$ is the Dirichlet energy of $\theta \in C_0([0, T], \mathbb{R})$:

$$E_T(\theta) := \begin{cases} \frac{1}{2} \int_0^T \left| \frac{d}{dt} \theta_t \right|^2 dt, & \text{if } \theta \text{ is absolutely continuous on } [0, T], \\ \infty, & \text{otherwise.} \end{cases} \quad (2.16)$$

Remark 2.4. The (Cameron-Martin) space of absolutely continuous functions on $[0, T]$ with square-integrable derivative coincides with the Sobolev space $W_0^{1,2}([0, T], \mathbb{R}^n)$ that has the norm

$$\|\boldsymbol{\theta}\|_{1,2;[0, T]} := \left(\sum_{j=1}^n \int_0^T |\theta_t^j|^2 dt + \sum_{j=1}^n \int_0^T \left| \frac{d}{dt} \theta_t^j \right|^2 dt \right)^{1/2},$$

thanks to the ACL characterization of Sobolev spaces [AIM09, Lemma A.5.2] (note that as such, this fails for $T = \infty$). We will thus identify all these spaces:

$$H_0^1([0, T], \mathbb{R}^n) = W_0^{1,2}([0, T], \mathbb{R}^n) = \{\boldsymbol{\theta} \in C_0([0, T], \mathbb{R}^n) \mid E_T(\boldsymbol{\theta}) < \infty\}$$

Let us also note that if $\boldsymbol{\theta} \in H_0^1([0, T], \mathbb{R}^n)$, then $\boldsymbol{\theta}$ is $\frac{1}{2}$ -Hölder continuous by Morrey's inequality (but may have arbitrarily large Hölder norm), cf. [Eva10, Theorem 4, page 280].

Theorem C (Direct consequence of Schilder's theorem; see, e.g., [DZ10], Chapter 5.2). *Fix $T \in (0, \infty)$. The process $(\sqrt{\kappa} \mathbf{B}_t)_{t \in [0, T]}$ satisfies the following LDP in $C_0([0, T], \mathbb{R}^n)$, with good rate function E_T :*

For any closed subset F and open subset O of $C_0([0, T], \mathbb{R}^n)$, we have

$$\begin{aligned} \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{P}[\sqrt{\kappa} \mathbf{B}_{[0, T]} \in F] &\leq - \inf_{\boldsymbol{\theta} \in F} E_T(\boldsymbol{\theta}), \\ \underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{P}[\sqrt{\kappa} \mathbf{B}_{[0, T]} \in O] &\geq - \inf_{\boldsymbol{\theta} \in O} E_T(\boldsymbol{\theta}). \end{aligned}$$

A convenient tool for proving an LDP when a family of probability measures is absolutely continuous with respect to another family for which an LDP is already known is provided by the classical Varadhan's lemma. We will use it in combination with Theorem C.

Lemma D (Varadhan's lemma; see, e.g., [DZ10], Lemmas 4.3.4 and 4.3.6). *Suppose that the probability measures $(\mathbb{P}^\kappa)_{\kappa > 0}$ satisfy an LDP in a topological space X with good rate function E . Let $\Phi: X \rightarrow \mathbb{R}$ be a function bounded from above. Then, the following hold.*

1. *If Φ is upper semicontinuous, then for any closed subset F of X ,*

$$\overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}^\kappa \left[\exp \left(\frac{1}{\kappa} \Phi(X) \right) \mathbb{1}\{X \in F\} \right] \leq - \inf_{x \in F} (E(x) - \Phi(x)).$$

2. If Φ is lower semicontinuous, then for any open subset O of X ,

$$\liminf_{\kappa \rightarrow 0^+} \kappa \log \mathbb{E}^\kappa \left[\exp \left(\frac{1}{\kappa} \Phi(X) \right) \mathbb{1}\{X \in O\} \right] \geq - \inf_{x \in O} (E(x) - \Phi(x)).$$

In order to apply Varadhan's lemma in the measure $\mathbf{P}^{\kappa,a}$, we first find a suitable function $\Phi_t^{\kappa,a}$ so that the martingale $M_t^{\kappa,a/\kappa}$ from Proposition A with $\alpha = a/\kappa$ takes the form

$$\frac{M_t^{\kappa,a/\kappa}}{M_0^{\kappa,a/\kappa}} = \exp \left(\frac{1}{\kappa} \Phi_t^{\kappa,a}(\mathbf{B}) \right), \quad t \geq 0. \quad (2.17)$$

Using the notations $\phi_a^j(\boldsymbol{\theta}) := \frac{a}{4} \phi^j(\boldsymbol{\theta})$ and $\psi_a(\boldsymbol{\theta}) := \frac{a}{4} \psi(\boldsymbol{\theta})$ analogous to (1.5, 2.6), we have

$$\frac{\Delta F_{a/\kappa}(\mathbf{U}_t^\kappa)}{F_{a/\kappa}(\mathbf{U}_t^\kappa)} = -\psi_a(\mathbf{U}_t^\kappa) + \frac{4}{\kappa} \sum_{j=1}^n (\phi_a^j(\mathbf{U}_t^\kappa))^2, \quad t \geq 0.$$

Therefore, using (2.4, 2.8, 2.10), we see that Equation (2.17) holds with

$$\begin{aligned} \Phi_t^{\kappa,a}(\mathbf{B}) &= a \log F(\mathbf{U}_t^\kappa) - a \log F(\mathbf{U}_0^\kappa) + \frac{1}{2} \int_0^t \left(\kappa \psi_a(\mathbf{U}_s^\kappa) - 4 \sum_{j=1}^n (\phi_a^j(\mathbf{U}_s^\kappa))^2 \right) ds \\ &=: \Phi_t^{\kappa,a}(\mathbf{U}^\kappa), \end{aligned}$$

Definition 2.5. For any $\kappa \geq 0$, $a > 0$, and $T \in (0, \infty)$ the *radial interaction functional* is

$$\begin{aligned} \Phi_T^{\kappa,a} &: C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n) \rightarrow \mathbb{R}, \\ \Phi_T^{\kappa,a}(\boldsymbol{\theta}) &:= G_a(\boldsymbol{\theta}_0) - G_a(\boldsymbol{\theta}_T) + \frac{1}{2} \int_0^T \left(\kappa \psi_a(\boldsymbol{\theta}_s) - \sum_{j=1}^n (2\phi_a^j(\boldsymbol{\theta}_s))^2 \right) ds, \end{aligned} \quad (2.18)$$

where

$$G_a(\boldsymbol{\theta}) := -a \log F(\boldsymbol{\theta}) = -a \log \prod_{1 \leq j < k \leq n} \left| \sin \left(\frac{\theta^k - \theta^j}{2} \right) \right| \geq 0. \quad (2.19)$$

In the next lemma, we gather useful properties of the functional $\Phi_T^{\kappa,a}$.

Lemma 2.6. Fix $T \in (0, \infty)$ and $\boldsymbol{\theta}_0 \in \mathcal{X}_n$. Equation (2.18) defines a continuous functional with respect to the metric (1.8), and for each $\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$, we have

$$\Phi_T^{\kappa,a}(\boldsymbol{\theta}) = \Phi_T^{0,a}(\boldsymbol{\theta}) + \frac{\kappa}{2} \int_0^T \psi_a(\boldsymbol{\theta}_s) ds \xrightarrow{\kappa \rightarrow 0^+} \Phi_T^{0,a}(\boldsymbol{\theta}), \quad (2.20)$$

and this limit is monotonically decreasing. Furthermore, $\Phi_T^{0,a}$ is bounded from above as

$$\Phi_T^{0,a}(\boldsymbol{\theta}) \leq G_a(\boldsymbol{\theta}_0), \quad \boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n). \quad (2.21)$$

Lastly, the following identity holds:

$$\Phi_t^{\kappa,a}(\boldsymbol{\theta}) = a \log \frac{F(\boldsymbol{\theta}_t)}{F(\boldsymbol{\theta}_0)} + a\kappa \frac{n(n^2 - 1)}{24} t + \frac{2(\kappa - a)}{a} \int_0^t \sum_{j=1}^n (\phi_a^j(\boldsymbol{\theta}_s))^2 ds. \quad (2.22)$$

Proof. The identities (2.20) follow by noting that $\log F(\boldsymbol{\theta}) \leq 0$, as $0 \leq F(\boldsymbol{\theta}) \leq 1$ by (2.4). As ψ_a is non-negative, we see that this limit is monotonically decreasing. The continuity and the bound (2.21) are clear. Lastly, using [HL21, Lemma 5.1],

$$\psi(\boldsymbol{\theta}) = \sum_{j=1}^n \left(\sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot \left(\frac{\theta^j - \theta^k}{2} \right) \right)^2 + \frac{n(n^2 - 1)}{3} = \sum_{j=1}^n (\phi^j(\boldsymbol{\theta}))^2 + \frac{n(n^2 - 1)}{3}$$

(for the functions $\psi(\boldsymbol{\theta})$ in (2.6) and ϕ^j in (1.5)), we obtain (2.22). \square

The following technical tail estimate is needed in the proof of Theorem 1.2.

Lemma 2.7. *Fix $\kappa \in (0, 4]$ and $a \geq \kappa$. Fix $\epsilon > 0$ and consider the stopping time*

$$\tau_\epsilon = \tau_\epsilon(\boldsymbol{\theta}) = \inf \left\{ t \geq 0 \mid \min_{1 \leq j < k \leq n} |\theta_t^j - \theta_t^k| < \epsilon \right\}. \quad (2.23)$$

For each initial configuration $\boldsymbol{\theta}_0 \in \mathcal{X}_n$, there exist constants $R = R(\epsilon, \boldsymbol{\theta}_0, a) \in (0, \infty)$ and $C = C(n, T, a) \in (0, \infty)$ independent of κ such that $\lim_{\epsilon \rightarrow 0} R(\epsilon, \boldsymbol{\theta}_0, a) = +\infty$ and

$$\mathbf{P}^{\kappa, a}[\tau_\epsilon \leq T] \leq C e^{-R/\kappa}. \quad (2.24)$$

Proof. Recall (cf. Proposition 2.2) that $\mathbf{P}^{\kappa, a}$ is the probability measure absolutely continuous with respect to \mathbb{P} with Radon-Nikodym derivative (2.17), given by the martingale (rather than simply a local martingale) $M_t^{\kappa, a/\kappa}$ from Proposition A. Therefore, since $\tau_\epsilon \wedge T$ is a stopping time bounded by T , by the optional stopping theorem (OST), we have

$$\begin{aligned} \mathbb{E}_T \left[\frac{M_T^{\kappa, a/\kappa}}{M_0^{\kappa, a/\kappa}} \mathbb{1}\{\tau_\epsilon \leq T\} \right] &= \mathbb{E}_T \left[\mathbb{E}_T \left[\frac{M_T^{\kappa, a/\kappa}}{M_0^{\kappa, a/\kappa}} \mathbb{1}\{\tau_\epsilon \leq T\} \mid \mathcal{F}_{\tau_\epsilon \wedge T} \right] \right] \\ &= \mathbb{E}_T \left[\frac{M_{\tau_\epsilon \wedge T}^{\kappa, a/\kappa}}{M_0^{\kappa, a/\kappa}} \mathbb{1}\{\tau_\epsilon \leq T\} \right] = \mathbb{E}_T \left[\frac{M_{\tau_\epsilon}^{\kappa, a/\kappa}}{M_0^{\kappa, a/\kappa}} \mathbb{1}\{\tau_\epsilon \leq T\} \right]. \end{aligned} \quad (2.25)$$

Thus, we obtain

$$\begin{aligned} \mathbf{P}^{\kappa, a}[\tau_\epsilon \leq T] &= \mathbb{E}_T \left[\exp \left(\frac{1}{\kappa} \Phi_T^{\kappa, a}(\mathbf{U}^\kappa) \right) \mathbb{1}\{\tau_\epsilon \leq T\} \right] \\ &= \mathbb{E}_T \left[\exp \left(\frac{1}{\kappa} \Phi_{\tau_\epsilon}^{\kappa, a}(\mathbf{U}^\kappa) \right) \mathbb{1}\{\tau_\epsilon \leq T\} \right] \quad [\text{by OST, (2.25)}] \\ &\leq \mathbb{E}_T \left[\exp \left(\frac{a}{\kappa} \log \frac{F(\mathbf{U}_{\tau_\epsilon}^\kappa)}{F(\boldsymbol{\theta}_0)} + \frac{an(n^2 - 1)}{24} \tau_\epsilon \right) \mathbb{1}\{\tau_\epsilon \leq T\} \right] \quad [\text{by (2.22), as } \kappa \leq a] \\ &\leq e^{\frac{an(n^2 - 1)}{24} T} F(\boldsymbol{\theta}_0)^{-a/\kappa} \left(\frac{\epsilon}{2} \right)^{a/\kappa}. \quad [\text{by (2.4)}] \end{aligned}$$

Hence, (2.24) holds with $R(\epsilon, \boldsymbol{\theta}_0, a) = -a \log \left(\frac{\epsilon}{2F(\boldsymbol{\theta}_0)} \right)$ and $C(n, T, a) = e^{\frac{an(n^2 - 1)}{24} T}$. \square

We next express the multiradial Dirichlet energy J_T^a with parameter a (cf. Def. 1.1),

$$J_T^a(\boldsymbol{\theta}) := \begin{cases} \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi_a^j(\boldsymbol{\theta}_s) \right|^2 ds, & \text{if } \boldsymbol{\theta} \text{ is absolutely continuous on } [0, T], \\ \infty, & \text{otherwise.} \end{cases}$$

in terms of the functional $\Phi_T^{0,a}$ (cf. Definition 2.5) and the sum of independent Dirichlet energies (2.15) appearing in Schilder's theorem, denoted E_T .

Lemma 2.8. *Fix $T \in (0, \infty)$, $a > 0$, and $\boldsymbol{\theta}_0 \in \mathcal{X}_n$. For any $\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$, we have*

$$J_T^a(\boldsymbol{\theta}) = E_T(\boldsymbol{\theta}) - \Phi_T^{0,a}(\boldsymbol{\theta}), \quad \text{where} \quad E_T(\boldsymbol{\theta}) := \sum_{j=1}^n E_T(\theta^j). \quad (2.26)$$

Proof. On the one hand, if $\boldsymbol{\theta}$ is not absolutely continuous, then $J_T^a(\boldsymbol{\theta}) = \infty$, since the last term on the righthand side of (2.18) with $\kappa = 0$ is negative. On the other hand, if $\boldsymbol{\theta}$ is absolutely continuous, then by (2.4), we have

$$\begin{aligned} \log \frac{F(\boldsymbol{\theta}_t)}{F(\boldsymbol{\theta}_0)} &= \sum_{1 \leq j < k \leq n} \log \left| \sin \left(\frac{\theta_t^k - \theta_t^j}{2} \right) \right| - \sum_{1 \leq j < k \leq n} \log \left| \sin \left(\frac{\theta_0^k - \theta_0^j}{2} \right) \right| \\ &= \frac{1}{2} \int_0^t \sum_{j=1}^n \left(\frac{d}{ds} \theta_s^j \right) \sum_{\substack{1 \leq k \leq n \\ k \neq j}} \cot \left(\frac{\theta_s^k - \theta_s^j}{2} \right) ds = \frac{1}{2} \int_0^t \sum_{j=1}^n \left(\frac{d}{ds} \theta_s^j \right) \phi^j(\boldsymbol{\theta}_s) ds, \end{aligned}$$

which together with Definition 2.5 and (2.19) implies that

$$\begin{aligned} E_T(\boldsymbol{\theta}) - \Phi_T^{0,a}(\boldsymbol{\theta}) &= \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j \right|^2 ds - \frac{1}{2} \int_0^T \sum_{j=1}^n \left(4\phi_a^j(\boldsymbol{\theta}_s) \left(\frac{d}{ds} \theta_s^j \right) - (2\phi_a^j(\boldsymbol{\theta}_s))^2 \right) ds \\ &= \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi_a^j(\boldsymbol{\theta}_s) \right|^2 ds = J_T^a(\boldsymbol{\theta}). \end{aligned}$$

□

Corollary 2.9. *We have $J_T^a(\boldsymbol{\theta}) < \infty$ if and only if $E_T(\theta^j) < \infty$ for all $j \in \{1, \dots, n\}$ and*

$$\tau_{\text{coll}} = \tau_{\text{coll}}(\boldsymbol{\theta}) := \inf \left\{ t \geq 0 : \min_{1 \leq j < k \leq n} |e^{i\theta_t^j} - e^{i\theta_t^k}| = 0 \right\} > T. \quad (2.27)$$

Proof. If $\tau_{\text{coll}} > T$, then we see from Definition 2.5 that $\Phi_T^{0,a}(\boldsymbol{\theta}) > -\infty$. If furthermore $E_T(\theta^j) < \infty$ for all j , then we see from (2.26) from Lemma 2.8 that $J_T^a(\boldsymbol{\theta}) < \infty$. This proves the converse implication. To prove (the contraposition of) the forward implication, note that $\Phi_T^{0,a}$ is bounded from above by (2.21), so $E_T(\theta^j) = \infty$ readily implies that $J_T^a(\boldsymbol{\theta}) = \infty$, while if $\tau_{\text{coll}} \leq T$, then monotonicity of the energy in time gives

$$\begin{aligned} J_T^a(\boldsymbol{\theta}) &\geq J_{\tau_{\text{coll}}}^a(\boldsymbol{\theta}) = \overline{\lim}_{t \rightarrow \tau_{\text{coll}}^-} J_t^a(\boldsymbol{\theta}) \\ &= \overline{\lim}_{t \rightarrow \tau_{\text{coll}}^-} (E_t(\boldsymbol{\theta}) - \Phi_t^{0,a}(\boldsymbol{\theta})) \quad [\text{by (2.26) from Lemma 2.8}] \\ &\geq \overline{\lim}_{t \rightarrow \tau_{\text{coll}}^-} (G_a(\boldsymbol{\theta}_t) - G_a(\boldsymbol{\theta}_0)) \quad [\text{by (2.18), as } E_t(\boldsymbol{\theta}) \geq 0] \\ &= \infty. \quad [\text{by (2.19)}] \end{aligned}$$

This concludes the proof. □

We also have a similar (unidirectional) result for infinite time, to be used in Section 4.

Corollary 2.10. *If $J^a(\boldsymbol{\theta}) := \lim_{T \rightarrow \infty} J_T^a(\boldsymbol{\theta}) < \infty$, then $E(\boldsymbol{\theta}) := \lim_{T \rightarrow \infty} E_T(\boldsymbol{\theta}) < \infty$.*

Proof. Using Lemma 2.8 and the definition (2.18) of $\Phi_T^{0,a}$, we see that

$$\begin{aligned} E(\boldsymbol{\theta}) &= \lim_{T \rightarrow \infty} (J_T^a(\boldsymbol{\theta}) + \Phi_T^{0,a}) && \text{[by (2.26) from Lemma 2.8]} \\ &\leq \lim_{T \rightarrow \infty} J_T^a(\boldsymbol{\theta}) + G_a(\boldsymbol{\theta}_0) = J^a(\boldsymbol{\theta}) + G_a(\boldsymbol{\theta}_0) < \infty. && \text{[by (2.18), as } J_T^a(\boldsymbol{\theta}) \geq 0 \text{]} \end{aligned}$$

□

2.4 LDP for Dyson Brownian motion on the circle

We are now prepared to prove the LDP for the radial Bessel process (Theorem 1.2).

Recall (e.g., from [DZ10]) that for a topological space X , a *rate function* I is a lower semicontinuous mapping $I: X \rightarrow [0, \infty]$ (i.e., for all $c \in [0, \infty)$, the level set $I^{-1}[0, c]$ is a closed subset of X). We note that in Theorems 1.2 and 1.8, the space X is a metric space ($C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$, or \mathcal{C} , respectively), so it is sufficient to check the lower semicontinuity property on sequences. Recall also that a *good rate function* is a rate function for which all level sets are compact subsets of X , which implies lower semicontinuity.

Lemma 2.11. *The multiradial Dirichlet energy J_T^a in Definition 1.1 is a good rate function.*

Proof. Fix $c \geq 0$, and let $(\boldsymbol{\theta}_{(k)})_{k \in \mathbb{N}}$ be a sequence in $(J_T^a)^{-1}[0, c]$. We have

$$\begin{aligned} E_T(\boldsymbol{\theta}_{(k)}) &\leq E_T(\boldsymbol{\theta}_{(k)}) - \Phi_T^{0,a}(\boldsymbol{\theta}_{(k)}) + G_a(\boldsymbol{\theta}_0) && \text{[by (2.21)]} \\ &= J_T^a(\boldsymbol{\theta}_{(k)}) + G_a(\boldsymbol{\theta}_0) && \text{[by (2.26) from Lemma 2.8]} \\ &\leq c + G_a(\boldsymbol{\theta}_0). \end{aligned}$$

Since E_T is a good rate function (by Schilder's theorem), we can pass to a subsequence, also denoted by $(\boldsymbol{\theta}_{(k)})_{k \in \mathbb{N}}$, which converges to some element

$$\boldsymbol{\theta} \in (E_T)^{-1}[0, c + G_a(\boldsymbol{\theta}_0)].$$

By continuity of $\Phi_T^{0,a}$ from Lemma 2.6 and lower semicontinuity of E_T (by Schilder's theorem), we obtain using Lemma 2.8 the estimate

$$J_T^a(\boldsymbol{\theta}) = E_T(\boldsymbol{\theta}_{(k)}) - \Phi_T^{0,a}(\boldsymbol{\theta}) \leq \liminf_{k \rightarrow \infty} (E_T(\boldsymbol{\theta}_{(k)}) - \Phi_T^{0,a}(\boldsymbol{\theta}_{(k)})) = \liminf_{k \rightarrow \infty} J_T^a(\boldsymbol{\theta}_{(k)}) \leq c,$$

yielding $\boldsymbol{\theta} \in (J_T^a)^{-1}[0, c]$. This shows that $(J_T^a)^{-1}[0, c]$ is compact, so J_T^a is good. □

We now conclude with the proof of the first main result of the present work (Theorem 1.2). In fact, we shall prove a slightly more general result, Theorem 2.12 — Theorem 1.2 then follows as the special case where $a = 4$.

Theorem 2.12. *Fix $T \in (0, \infty)$ and $a > 0$. For $\kappa < 2a$, let \mathbf{U}^κ be the unique strong solution to (2.3) with $\alpha = a/\kappa$, started at $\mathbf{U}_0^\kappa = \boldsymbol{\theta}_0 \in \mathcal{X}_n$. The family $(\mathbf{P}^{\kappa,a})_{\kappa > 0}$ of laws induced by \mathbf{U}^κ satisfies the following LDP in $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$ with good rate function J_T^a :*

For any closed subset F and open subset O of $C_{\theta_0}([0, T], \mathcal{X}_n)$, we have

$$\overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^{\kappa, a}[\mathbf{U}^\kappa \in F] \leq - \inf_{\theta \in F} J_T^a(\theta), \quad (2.28)$$

$$\underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^{\kappa, a}[\mathbf{U}^\kappa \in O] \geq - \inf_{\theta \in O} J_T^a(\theta). \quad (2.29)$$

Proof of Theorem 1.2. Theorem 1.2 follows by setting $a = 4$ in Theorem 2.12. \square

Proof of Theorem 2.12. We already know that J_T^a is a good rate function by Lemma 2.11. We will use Schilder's theorem (Theorem C) combined with Varadhan's lemma (Lemma D), to prove the lower & upper bounds (2.28, 2.29). The upper bound (2.28) is the harder one.

Recall that \mathbb{P} is the probability measure for n independent Brownian motions, satisfying an LDP from Schilder's theorem, with good rate function E_T . Also, $\mathbf{P}^{\kappa, a}$ is the probability measure that is absolutely continuous with respect to \mathbb{P} with Radon-Nikodym derivative given by (2.17), where $\Phi_T^{\kappa, a}$ is the functional from Definition 2.5:

$$\frac{d\mathbf{P}_T^{\kappa, a}}{d\mathbb{P}_T} = \exp\left(\frac{1}{\kappa} \Phi_T^{\kappa, a}(\mathbf{U}^\kappa)\right). \quad (2.30)$$

Let \mathbb{E}_T be the expected value with respect to \mathbb{P}_T . Using the Radon-Nikodym derivative (2.30), for any Borel set $H \subseteq C_{\theta_0}([0, T], \mathcal{X}_n)$, we have

$$\kappa \log \mathbf{P}_T^{\kappa, a}[\mathbf{U}_{[0, T]}^\kappa \in H] = \kappa \log \mathbb{E}_T \left[\exp\left(\frac{1}{\kappa} \Phi_T^{\kappa, a}(\mathbf{U}^\kappa)\right) \mathbb{1}\{\mathbf{U}_{[0, T]}^\kappa \in H\} \right] \quad (2.31)$$

Lower bound. Fix an open set $O \subset C_{\theta_0}([0, T], \mathcal{X}_n)$. Without loss of generality, we may assume that

$$M_O := \inf_{\theta \in O} (E_T(\theta) - \Phi_T^{0, a}(\theta)) < \infty.$$

Fix $\varepsilon > 0$ and $\theta^{(\varepsilon)} \in O$ such that $E_T(\theta^{(\varepsilon)}) - \Phi_T^{0, a}(\theta^{(\varepsilon)}) \leq M_O + \varepsilon$, which in particular implies that $\Phi_T^{0, a}(\theta^{(\varepsilon)}) > -\infty$. As $\Phi_T^{0, a}$ is continuous by Lemma 2.6, we can pick an open neighborhood $O^{(\varepsilon)} \subset O$ of $\theta^{(\varepsilon)} \in C_{\theta_0}([0, T], \mathcal{X}_n)$ such that $\Phi_T^{0, a} \geq \Phi_T^{0, a}(\theta^{(\varepsilon)}) - 1$ on $O^{(\varepsilon)}$.

Now, let Φ be the lower semicontinuous function equaling $\Phi_T^{0, a}$ on $O^{(\varepsilon)}$ and $\Phi_T^{0, a}(\theta^{(\varepsilon)}) - 2$ otherwise. Since $\Phi_T^{0, a} \leq \Phi_T^{\kappa, a}$, by applying Item 2 of Varadhan's lemma (Lemma D) to the set $O^{(\varepsilon)}$ and the function Φ , combined with Schilder's theorem (Theorem C), we obtain

$$\begin{aligned} & \underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}_T \left[\exp\left(\frac{1}{\kappa} \Phi_T^{\kappa, a}(\mathbf{U}^\kappa)\right) \mathbb{1}\{\mathbf{U}_{[0, T]}^\kappa \in O\} \right] \\ & \geq \underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}_T \left[\exp\left(\frac{1}{\kappa} \Phi_T^{0, a}(\mathbf{U}^\kappa)\right) \mathbb{1}\{\mathbf{U}_{[0, T]}^\kappa \in O^{(\varepsilon)}\} \right] \\ & \geq - \inf_{\theta \in O^{(\varepsilon)}} (E_T(\theta) - \Phi(\theta)) \geq -(E_T(\theta^{(\varepsilon)}) - \Phi_T^{0, a}(\theta^{(\varepsilon)})) \\ & \geq -M_O - \varepsilon \xrightarrow{\varepsilon \searrow 0} -M_O. \end{aligned}$$

Upper bound. Fix a closed set $F \subset C_{\theta_0}([0, T], \mathcal{X}_n)$. We separate the proof into two parts. First, assume that for the function $\psi_a = \frac{a}{4}\psi$ defined via (2.6), we have

$$\Psi_a(F) := \sup_{\theta \in F} \int_0^T \psi_a(\theta_s) ds < \infty, \quad (2.32)$$

which implies by Lemma 2.6 that

$$\Phi_T^{\kappa,a}(\boldsymbol{\theta}) = \Phi_T^{0,a}(\boldsymbol{\theta}) + \frac{\kappa}{2} \int_0^T \psi_a(\boldsymbol{\theta}_s) \, ds \leq \Phi_T^{0,a}(\boldsymbol{\theta}) + \frac{\kappa}{2} \Psi_a(F), \quad \text{for all } \boldsymbol{\theta} \in F.$$

Fix $\varepsilon, M > 0$. Note that $\Phi(\boldsymbol{\theta}; M, \varepsilon) := \max\{\Phi_T^{0,a}(\boldsymbol{\theta}) + \frac{\varepsilon}{2} \Psi_a(F), -M\}$ is a continuous function by Lemma 2.6, and $\Phi_T^{\kappa,a}(\boldsymbol{\theta}) \leq \Phi(\boldsymbol{\theta}; M, \varepsilon)$ for all $\boldsymbol{\theta} \in F$ and for all $\kappa \in [0, \varepsilon]$. Applying Item 1 of Varadhan's lemma (Lemma D) to the set F and the continuous function $\Phi(\boldsymbol{\theta}; M, \varepsilon)$, combined with Schilder's theorem (Theorem C), we obtain

$$\begin{aligned} & \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}_T \left[\exp \left(\frac{1}{\kappa} \Phi_T^{\kappa,a}(\mathbf{U}^\kappa) \right) \mathbb{1}\{\mathbf{U}_{[0,T]}^\kappa \in F\} \right] \\ & \leq \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}_T \left[\exp \left(\frac{1}{\kappa} \Phi(\mathbf{U}^\kappa; M, \varepsilon) \right) \mathbb{1}\{\mathbf{U}_{[0,T]}^\kappa \in F\} \right] \\ & \leq - \inf_{\boldsymbol{\theta} \in F} (E_T(\boldsymbol{\theta}) - \Phi(\boldsymbol{\theta}; M, \varepsilon)) \xrightarrow[\varepsilon \searrow 0]{M \nearrow \infty} - \inf_{\boldsymbol{\theta} \in F} (E_T(\boldsymbol{\theta}) - \Phi_T^{0,a}(\boldsymbol{\theta})). \end{aligned}$$

Next, if (2.32) does not hold, we fix $\epsilon > 0$ and consider the stopping time τ_ϵ (2.23) from Lemma 2.7. Note that the set $F(\epsilon) := F \cap \{\tau_\epsilon > T\}$ is closed, and (2.32) holds on $F(\epsilon)$:

$$\Psi_a(F(\epsilon)) := \sup_{\boldsymbol{\theta} \in F(\epsilon)} \int_0^T \psi_a(\boldsymbol{\theta}_s) \, ds < \infty.$$

Therefore, we see that

$$\begin{aligned} & \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbb{E}_T \left[\exp \left(\frac{1}{\kappa} \Phi_T^{\kappa,a}(\mathbf{U}^\kappa) \right) \mathbb{1}\{\mathbf{U}_{[0,T]}^\kappa \in F\} \right] \\ & \leq \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \left(\mathbf{P}^{\kappa,a}[\mathbf{U}^\kappa \in F(\epsilon)] + \mathbf{P}^{\kappa,a}[\tau_\epsilon \leq T] \right) \end{aligned}$$

is bounded from above by the maximum of the two terms

$$\begin{aligned} \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^{\kappa,a}[\mathbf{U}^\kappa \in F(\epsilon)] & \leq - \inf_{\boldsymbol{\theta} \in F(\epsilon)} (E_T(\boldsymbol{\theta}) - \Phi_T^{0,a}(\boldsymbol{\theta})) \leq - \inf_{\boldsymbol{\theta} \in F} (E_T(\boldsymbol{\theta}) - \Phi_T^{0,a}(\boldsymbol{\theta})), \\ \overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^{\kappa,a}[\tau_\epsilon \leq T] & \leq -R(\epsilon, \boldsymbol{\theta}_0, a) \xrightarrow[\epsilon \searrow 0]{} -\infty, \end{aligned}$$

where to bound the first term we used the first part of the proof and the fact that $F(\epsilon) \subseteq F$, and noted that the second term is exponentially small thanks to (2.24) in Lemma 2.7.

Conclusion. By Lemma 2.8, the asserted inequalities (2.28, 2.29) follow from the above bounds together with (2.31). \square

Remark 2.13. It follows from the goodness of the multiradial Dirichlet energy (Lemma 2.11) that it attains its minimum on $C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$. Moreover, from Theorem 1.2 we see that the minimum equals zero: taking $F = O = C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$, (2.28, 2.29) together imply that

$$\min_{\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)} J_T^a(\boldsymbol{\theta}) = \inf_{\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)} J_T^a(\boldsymbol{\theta}) = \lim_{\kappa \rightarrow 0+} \kappa \log \underbrace{\mathbf{P}^{\kappa,a}[\mathbf{U}^\kappa \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)]}_{=1} = 0.$$

3 LDP for multiradial SLE_{0+}

The goal of this section is to prove the main result, Theorem 1.8, which is a finite-time LDP for the n -radial SLE_κ process as $\kappa \rightarrow 0+$. Recall that in Definition 1.6, n -radial SLE is defined as the Loewner chain whose (n -dimensional) driving process is $(e^{i\Theta_t^1}, \dots, e^{i\Theta_t^n})$, where Θ is the n -radial Bessel process (cf. Corollary 3.4). As Theorem 1.2 gives an LDP for the n -radial Bessel process, it would be convenient to just apply the Loewner transform and use the contraction principle (recalled in Theorem E) to deduce an LDP for multiradial SLE. Unfortunately, the standard contraction principle cannot be applied directly, since the Loewner transform (1.12) is not continuous for the Hausdorff metric, but only in the so-called Carathéodory sense. As the latter topology is not very useful for addressing geometric properties of hulls, we need to address the discontinuities of the Loewner transform under the Hausdorff metric (1.13). From the chordal case [PW24], we know that discontinuities of the Loewner transform (for the Hausdorff metric) occur at hulls with non-empty interiors. In contrast, we show that finite-energy hulls are simple radial multichords (Sections 3.2 & 3.3 — see in particular Theorem 1.9). This allows us to sidestep the discontinuities: we can apply the contraction principle on a smaller space where the Loewner transform is continuous, and then extend the LDP to the full space using Proposition 3.13 and Lemma 3.15.

Proving that finite-energy hulls are simple radial multichords (Theorem 1.9) is the main work of this section. In the chordal case with $n = 1$, an analogous result has been verified by two methods. On the one hand, following the methodology of Lind, Marshall, and Rohde [LMR10], Wang used quasiconformal maps to argue that each finite-energy ($n = 1$) hull is a quasi-arc [Wan19a], which was later generalized to the case of n -multichords in [PW24]. However, for radial multichords with $n \geq 2$, quasiconformal maps do not seem to present the most natural geometric setup, so we do not follow this approach here. (See [AP24] for an elaboration of this approach in the case of one radial chord.)

On the other hand, motivated by rough path theory, in [FS17] Friz & Shekhar derived a strong derivative estimate for the Loewner uniformizing map near the tip for drivers with finite Dirichlet energy (2.16). This can be used via standard arguments to imply that the Loewner hulls thus obtained are in fact simple curves. In the present work, we employ the strategy used by Friz & Shekhar combined with a generalized version of the restriction property [LSW03] (see Proposition 3.12). As a by-product, we obtain a radial version of the main theorem of [FS17], but generalized to allow weight functions λ — see Theorem 3.9.

3.1 Multiradial Loewner equation and multiradial SLE_κ

We will now consider a more general (weighted) version of the Loewner equation (Equation (3.1) below, which generalizes (1.11)). This allows us to consider Loewner flow with a more general time-dependent parameterization and enables us to reparameterize radial multichords when necessary. We restrict our attention to “nice” weights as follows.

Definition 3.1. A *weight function* is a càdlàg (i.e., right-continuous with left limits) and locally integrable function $\lambda: [0, \infty) \rightarrow (0, \infty)$.

Notice that if λ is a weight function, then the map $t \mapsto \int_0^t \lambda_s ds$ is strictly increasing

(hence, it can be used to define a time change), and λ is bounded on compact time intervals.

Multiradial Loewner equation. For any weight function λ , we define the *multiradial Loewner equation with weight λ* as the boundary value problem

$$\partial_t \mathfrak{g}_t(z) = \lambda_t \mathfrak{g}_t(z) \sum_{j=1}^n \frac{w_t^j + \mathfrak{g}_t(z)}{w_t^j - \mathfrak{g}_t(z)}, \quad \mathfrak{g}_0(z) = z, \quad z \in \overline{\mathbb{D}}, \quad t \geq 0, \quad (3.1)$$

where the driving functions $w_t^1, \dots, w_t^n \in \partial\mathbb{D}$ are non-intersecting and continuous in time. The solution $\mathfrak{g}_t = \mathfrak{g}_t^\lambda$ to (3.1) is called the *Loewner chain with λ -common parameterization*. Then, $\mathfrak{g}_t: \mathbb{D} \setminus K_t^\mathfrak{g} \rightarrow \mathbb{D}$ is the uniformizing map normalized at the origin, and in the parameterization in (3.1), we have

$$\log \mathfrak{g}'_t(0) = n \int_0^t \lambda_s ds.$$

Note also that the map \mathfrak{h}_t related to \mathfrak{g}_t via $\mathfrak{g}_t(e^{iu}) = \exp(i\mathfrak{h}_t(u))$, and with $w_t^j = \exp(i\theta_t^j)$ for $j = 1, \dots, n$, satisfies

$$\partial_t \mathfrak{h}_t(u) = \lambda_t \sum_{j=1}^n \cot \left(\frac{\mathfrak{h}_t(u) - \theta_t^j}{2} \right). \quad (3.2)$$

We say that the generated hulls $K_t^\mathfrak{g}$ have the *λ -common parameterization*.

An even more general version of (3.1) could be obtained by weighting each term in the sum by a different weight λ_t^j , which would allow the components of the generated hull to be parameterized at different rates, but this is not needed for the present work.

Definition 3.2. Fix $T \in (0, \infty)$. Let $\theta \in C([0, T], \mathcal{X}_n)$, let λ be a weight function, and let \mathfrak{g} and $K_t^\mathfrak{g}$ be as in (3.1). We say that θ *generates a radial multichord $\gamma_{[0, T]}$* in \mathbb{D} with the λ -common parameterization if $t \mapsto \gamma_t^j$ is a continuous map from $[0, T]$ to $\overline{\mathbb{D}}$ such that $\gamma_0^j \in \partial\mathbb{D}$ for each $1 \leq j \leq n$, the image $\gamma_{[0, t]}$ generates $K_t^\mathfrak{g}$ for all $t \in [0, T]$, and the concatenations of γ^j with any simple curves from γ_T^j to the origin form a radial multichord (as in Definition 1.4). We call the radial multichord $\gamma_{[0, T]}$ *simple* if its each component γ^j is injective, $\gamma_{(0, T]} \subset \mathbb{D}$, and furthermore $\gamma_{[0, T]}^j \cap \gamma_{[0, T]}^k$ for all $j \neq k$.

When $n = 1$, we call $\gamma_{[0, T]}$ a (simple) radial chord in \mathbb{D} with the λ -parameterization. In this case, γ is also often referred to as the *Loewner trace* in the literature.

Remark 3.3. In general, the geometry of Loewner hulls depends on both the weight function and the driving function; reparameterizing a hull allows one to focus on whichever is more convenient. For example, the well-known phase transition for ($n = 1$) chordal SLE $_\kappa$ [RS05] from almost surely simple ($\kappa \leq 4$) to self-touching to space-filling ($\kappa \geq 8$) can be understood by performing a time change so that the driving function is standard Brownian motion B_t and analyzing the resulting weight function. Indeed, the curves generated by the weighted chordal Loewner equation

$$\partial_t g_t(z) = \frac{\alpha}{g_t(z) - B_t}, \quad g_0(z) = z, \quad z \in \overline{\mathbb{H}}, \quad t \geq 0,$$

are almost surely simple if $\alpha \geq 1/2$ and space-filling if $\alpha \leq 1/4$. In this case, the time change allows for comparison between the weight function $\alpha = 2/\kappa$ and the parameter of the usual Bessel process on the real line. In this context, Item 2 in Theorem 3.9 is rather surprising: there, we show that driving functions with finite energy generate simple radial multichords for *any* weight function that is uniformly bounded away from zero.

3.1.1 Multiradial SLE $_{\kappa}$, for $\kappa \in (0, 4]$

The next corollary clarifies the relationship between the measures \mathbf{P}^{κ} discussed in Section 2 and n -radial SLE $_{\kappa}$ processes (Definition 1.6).

Corollary 3.4. *Suppose that $\mathbf{B}_t = (B_t^1, \dots, B_t^n)$ is an n -dimensional standard Brownian motion in \mathbb{R}^n defined on the filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{P})$, where \mathcal{F}_{\bullet} is its natural right-continuous completed filtration. Fix $\boldsymbol{\theta}_0 \in \mathcal{X}_n$ and define $\mathbf{U}_t^{\kappa} = (U_t^1, \dots, U_t^n) \in \mathcal{X}_n$ by*

$$\mathbf{U}_t^{\kappa} = \boldsymbol{\theta}_0 + \sqrt{\kappa} \mathbf{B}_t, \quad \mathbf{U}_0^{\kappa} = \boldsymbol{\theta}_0, \quad \text{for } 0 < \kappa \leq 4.$$

Let $z_t^j := e^{iU_t^j}$ for $j = 1, \dots, n$. Then, in the measure \mathbf{P}^{κ} appearing in Proposition 2.2, the process (z_t^1, \dots, z_t^n) comprises the driving functions for n -radial SLE $_{\kappa}$ started from \mathbf{z}_0 .

Proof. This is an immediate consequence of Proposition 2.2 and Definition 1.6. □

Remark 3.5. In [HL21] the authors consider (for $0 < \kappa \leq 4$) a sequence of measures $\mu_{t,T}^{\kappa}$ that are absolutely continuous with respect to \mathbb{P}_t , with Radon-Nikodym derivative that is a large time T truncation of the chordal Radon-Nikodym derivative (1.1). For each fixed t , as $T \rightarrow \infty$, the measures $\mu_{t,T}^{\kappa}$ converge in the finite variation distance to \mathbf{P}_t^{κ} . Combining this convergence result with Corollary 3.4 justifies the definition of multiradial SLE $_{\kappa}$ that we use in this work (Definition 1.6).

Since the present work relies on the construction of n -radial SLE $_{\kappa}$ in [HL21], it is worthwhile to briefly address our differing choice of parameterization. In [HL21], for each $\kappa \in (0, 4]$, the authors describe n -radial SLE $_{\kappa}$ as the Loewner chain generated by the multiradial Loewner equation (3.1) with weight $\lambda \equiv 4/\kappa$ and driving functions $w_t^j = e^{2i\Theta_t^j}$, $j = 1, \dots, n$, where $\Theta_t^{\alpha} = (\Theta_t^1, \dots, \Theta_t^n)$ is the n -radial Bessel process with parameter $\alpha = 4/\kappa$ from Definition 2.1¹⁰. However, the dependence of the weight function on κ (i.e., using the $\lambda \equiv 4/\kappa$ -common parameterization) poses complications as $\kappa \rightarrow 0+$, so this setup is not amenable to large deviations analysis. Thus, it will be more convenient for us to consider the corresponding process up to the time change $t \mapsto \frac{\kappa}{4} t$, so that the curves have the 1-common parameterization. In this setup, we consider the uniformizing conformal mappings $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ normalized at the origin and satisfying the multiradial Loewner equation (3.1) with $\lambda \equiv 1$ (i.e., Equation (1.11)). In particular, we note that the Loewner hulls generated by (1.11) with the 1-common parameterization are the same as those generated by (3.1) with the $\lambda = 4/\kappa$ -common parameterization, if

$$g_t(z) = \mathfrak{g}_{\kappa t/4}(z) \quad \text{and} \quad z_t^j = w_{\kappa t/4}^j.$$

We shall address more general time changes in the next Section 3.1.2.

¹⁰See [HL21, Theorem 3.12 and the discussion following Corollary 3.13].

Remark 3.6. For each $a > 0$, $\kappa \leq \min\{a, 4\}$, and an additional parameter $\rho \in \mathbb{R}$, one can similarly define multiradial SLE_κ with spiraling rate ρ (and with the common parameterization) as the random radial multichord γ_t for which the uniformizing conformal mappings $g_t: \mathbb{D} \setminus \gamma_t \rightarrow \mathbb{D}$ satisfy Equation (1.11) with driving functions $z_t^j = e^{iU_t^j}$ for $j = 1, \dots, n$, where $\mathbf{U}_t^k = (U_t^1, \dots, U_t^n)$ is the strong solution in $C_{\theta_0}([0, \infty), \mathcal{X}_n)$ to the SDEs (2.14) with $\alpha = a/\kappa$. The results of the present work apply to derive an LDP for this process as well (i.e., a version of Theorem 1.8), with the good rate function obtained from Remark 2.3 similarly as in (1.14). Indeed, note that by Corollary 2.9 the finite-energy condition $J_T^a(\boldsymbol{\theta}) < \infty$ is independent of a while, by applying the inequality $|A + B|^2 \leq 2(|A|^2 + |B|^2)$ below, for any $\rho, \rho' \in \mathbb{R}$ we have

$$\begin{aligned} J_T^{a,\rho}(\boldsymbol{\theta}) &= \frac{1}{2} \int_0^T \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - (2\phi_a^j(\boldsymbol{\theta}_s) + \rho) \right|^2 ds \\ &\leq \frac{1}{2} \int_0^T \sum_{j=1}^n \left(2 \left| \frac{d}{ds} \theta_s^j - (2\phi_a^j(\boldsymbol{\theta}_s) + \rho') \right|^2 + 2|\rho' - \rho|^2 \right) ds \\ &= 2J_T^{a,\rho'}(\boldsymbol{\theta}) + nt|\rho' - \rho|^2. \end{aligned}$$

From this, we conclude that the multiradial Dirichlet energy $J_T^{a,\rho}$ with spiral defined in Remark 2.3 is finite if and only if the multiradial Dirichlet energy $J_T = J_T^{4,0}$ without spiral in Definition 1.1 is finite. Thus, Theorem 1.9 also holds with the assumption $J_T(\boldsymbol{\theta}) < \infty$ replaced by the assumption $J_T^{a,\rho}(\boldsymbol{\theta}) < \infty$. Using this fact, one can check that also the proof of Theorem 1.8 applies verbatim to the spiraling case (up to footnote 12 on page 36).

Remark 3.7. Multiradial SLE_κ curves (allowing a spiraling rate ρ), are expected to satisfy the so-called re-sampling property: for each curve γ^j in $\boldsymbol{\gamma} = (\gamma^1, \dots, \gamma^n)$, conditionally on the other curves $\{\gamma^k, k \neq j\}$, the law of γ^j is that of the chordal SLE_κ in its natural connected component. To prove this property, one should first show that the n -radial SLE_κ is supported on radial multichords (as in Definition 1.4), continuous at the origin, and elsewhere pairwise disjoint (cf. [Law13]). This follows from [MS17] by using a coupling of SLE_κ curves as flow lines of the Gaussian free field. See also [WW24], where variants of this process are considered. We will not need these properties in the present work.

3.1.2 Time changes

Remark 3.3 describes the application of a particular time change to a Loewner chain with $n = 1$. More generally, we see that (3.1) is related to (1.11) by the following time change. Let λ_t and \mathfrak{g}_t as in (3.1), and define

$$\sigma(t) := \int_0^t \lambda_s ds, \quad \tau(t) := \sigma^{-1}(t), \quad \text{and} \quad g_t := \mathfrak{g}_{\tau(t)}. \quad (3.3)$$

Then, we have $\frac{d}{dt} \tau(t) = 1/\lambda_{\tau(t)}$, so the chain rule shows that g_t satisfies (1.11) with $z_t^j = w_{\tau(t)}^j$. Consequently, this time change allows us to conveniently move between the 1-common parameterization and the λ -common parameterization as needed.

Most importantly, this allows us to reparameterize radial multichords: the property of having finite truncated multiradial Dirichlet energy is preserved under a large class of time changes, as the next lemma states.

Lemma 3.8. Fix $T \in (0, \infty)$. Let $\sigma: [0, T] \rightarrow [0, \sigma(T)]$ be strictly increasing and differentiable, with $\sigma(0) = 0$, and suppose that $\dot{\sigma}(t) := \frac{d}{dt}\sigma(t)$ is uniformly bounded away from zero and infinity, i.e.,

$$\|\dot{\sigma}\|_{[0, T]} := \sup_{t \in [0, T]} |\dot{\sigma}(t)| \in (0, \infty) \quad \text{and} \quad \left\| \frac{1}{\dot{\sigma}} \right\|_{[0, T]} := \sup_{t \in [0, T]} \frac{1}{|\dot{\sigma}(t)|} \in (0, \infty).$$

For $\theta \in C([0, T], \mathcal{X}_n)$, set $\hat{\theta}_t := \theta_{\sigma(t)}$, and $\hat{T} := \sigma^{-1}(T)$. Then, we have

$$J_T(\theta) < \infty \quad \text{if and only if} \quad J_{\hat{T}}(\hat{\theta}) < \infty.$$

In particular, if the time change σ is defined by (3.3) for λ bounded away from zero and infinity, then the conclusion of Lemma 3.8 is that on finite time intervals, finite-energy drivers for 1-common and λ -common parameterizations coincide up to time change.

Proof. Since σ is strictly increasing and differentiable, we have $\dot{\sigma}(t) > 0$ for all t , so we can estimate the Dirichlet energy of each $\hat{\theta}^j$ by

$$\left\| \frac{1}{\dot{\sigma}} \right\|_{[0, T]} E_T(\theta^j) \leq E_{\hat{T}}(\hat{\theta}^j) \leq \|\dot{\sigma}\|_{[0, T]} E_T(\theta^j).$$

Moreover, we have $\tau_{\text{coll}}(\theta) > T$ if and only if $\tau_{\text{coll}}(\hat{\theta}) > \hat{T}$, and Corollary 2.9 thus implies that $J_T(\theta) < \infty$ is equivalent to $J_{\hat{T}}(\hat{\theta}) < \infty$. \square

3.2 Derivative estimate for finite-energy Loewner chains for $n = 1$

In this section, we consider solutions to the (single) radial Loewner equation (3.2) (with $n = 1$) with some weight function $\lambda: [0, T] \rightarrow (0, \infty)$. A well-known condition for the property that the driving function $\theta \in C([0, T], \mathbb{R})$ generates a radial chord $\gamma_{[0, T]}$ in \mathbb{D} is an estimate for the derivative of the inverse map $\mathfrak{f}_t := \mathfrak{h}_t^{-1}$ near the driving function θ_t (locally) uniformly in time. More precisely, to verify the existence of the Loewner trace γ , it suffices to show the existence of the radial limit at its tip (see, e.g., [RS05, Theorem 4.1] or [Kem17, Theorem 6.4]):

$$\gamma_t := \lim_{y \rightarrow 0^+} \exp(i \mathfrak{f}_t(\theta_t + iy)), \quad \text{uniformly for all } t \in [0, T]. \quad (3.4)$$

It is not hard to check (see, e.g., [FS17, Appendix] or [RS05, Theorem 3.6]) that the limit (3.4) exists uniformly in time if there exists a constant $a \in (0, 1)$ such that

$$|\mathfrak{f}'_t(\theta_t + iy)| \lesssim y^{a-1}, \quad \text{for all } y > 0 \text{ and } t \in [0, T]. \quad (3.5)$$

When θ has finite energy, the derivative estimate (3.5) holds (in a very strong form), and hence, the Loewner trace (3.4) exists and is continuous in time.

A chordal version of the next result appeared in [FS17, Theorem 2(i)] without any weight function. Theorem 3.9 includes a general weight function and thanks to its radial setup should be useful in applications to various planar growth processes.

Theorem 3.9. Fix $n = 1$ and $T \in (0, \infty)$. Let $\lambda: [0, T] \rightarrow (0, \infty)$ be a weight function that is uniformly bounded away from zero, i.e.,

$$\left\| \frac{1}{\lambda} \right\|_{[0, T]} := \sup_{t \in [0, T]} \frac{1}{|\lambda_t|} < \infty.$$

Let $\theta \in H_0^1([0, T], \mathbb{R})$ (i.e., absolutely continuous such that $\theta_0 = 0$ and $E_T(\theta) < \infty$).

1. Then, we have

$$|\mathfrak{f}'_t(\theta_t + iy)| \leq \exp\left(\frac{1}{2} \|\frac{1}{\lambda}\|_{[0,T]} E_T(\theta)\right), \quad \text{for all } y > 0, \quad (3.6)$$

where $\mathfrak{f}_t := \mathfrak{h}_t^{-1}$ is the inverse of the Loewner map \mathfrak{h}_t satisfying (3.2) with $n = 1$.

2. Moreover, θ generates a simple radial chord $\gamma_{[0,T]}$ in \mathbb{D} with the λ -parameterization.

The proof of Item 1 uses a computation similar to that in [FS17, Proof of Theorem 4]. (Though such computations have been used in earlier works, including [Lin05, MR05, Law09a, LMR10, LJV11].) The proof of Item 2 relies on the bound (3.5) implied by Item 1 together with an argument that the resulting curve is indeed *simple*, which differs from prior arguments used in the chordal case (that in the literature rely on the specific form of the chordal Loewner equation, or scale-invariance which is absent in the radial case). Alternatively, one could estimate the quasiconformal distortion to show that radial finite-energy hulls are quasislits as in [MR05, LMR10, AP24].

Proof. Fix $t \geq 0$ and write $\varrho_s := \theta_t - \theta_{t-s}$. Then, the (mirror) backward Loewner flow

$$\mathfrak{p}_s(z) := \mathfrak{h}_{t-s}(\mathfrak{f}_t(z + \theta_t)) - \theta_t, \quad 0 \leq s \leq t,$$

satisfies $\mathfrak{p}_t(z) = \mathfrak{f}_t(z + \theta_t) - \theta_t$ and the backward Loewner equation

$$\partial_s \mathfrak{p}_s(z) = -\ell_s \cot\left(\frac{\mathfrak{p}_s(z) + \varrho_s}{2}\right), \quad \mathfrak{p}_0(z) = z, \quad \ell_s := \lambda_{t-s}, \quad 0 \leq s \leq t.$$

Writing

$$\mathfrak{p}_s(z) + \varrho_s = X_s + iY_s \quad \text{and} \quad N_s := \cos(X_s) - \cosh(Y_s),$$

we find for the inverse Loewner map the equation

$$\partial_s \log |\mathfrak{f}'_s(z + \theta_t)| = \partial_s \log |p'_s(z)| = \ell_s \frac{(1 - \cos(X_s) \cosh(Y_s))}{N_s^2}, \quad 0 \leq s \leq t.$$

Writing also $G_s := \varrho_s - X_s$, we obtain

$$\partial_s X_s = \partial_s \varrho_s - \partial_s G_s, \quad \partial_s Y_s = \ell_s \frac{\sinh(Y_s)}{N_s}, \quad \partial_s G_s = \ell_s \frac{\sin(X_s)}{N_s}.$$

Now, a straightforward computation shows that

$$\partial_s \log |\mathfrak{f}'_s(z + \theta_t)| = -\ell_s \frac{\sinh^2(Y_s)}{N_s^2} + \ell_s \frac{\cosh(Y_s)}{N_s} = -\frac{\sinh(Y_s)}{N_s} (\partial_s Y_s) + \frac{\partial_s \sinh(Y_s)}{\sinh(Y_s)}.$$

To write this in a more useful form, let us compute

$$\begin{aligned} \frac{\partial_s N_s}{N_s} &= \frac{\sinh(Y_s)}{N_s} (\partial_s Y_s) + \frac{\sin(X_s)}{N_s} (\partial_s X_s) \\ &= \frac{\sinh(Y_s)}{N_s} (\partial_s Y_s) + \frac{1}{\ell_s} (\partial_s G_s) (\partial_s \varrho_s - \partial_s G_s). \end{aligned}$$

Putting the above computations together, we see that

$$\log |\mathfrak{f}'_s(z + \theta_t)| = \log \left(\frac{\sinh(Y_t)}{\sinh(Y_0)} \right) - \log \left(\frac{N_t}{N_0} \right) + \int_0^t \left((\partial_s G_s)(\partial_s \varrho_s) - (\partial_s G_s)^2 \right) \frac{ds}{\ell_s}. \quad (3.7)$$

To evaluate (3.6), take $z = iy$, with $y > 0$, so that $X_0 = 0$ and $Y_0 = y$. Then, since Y_s and $\partial_s Y_s$ are positive, we see that

$$\begin{aligned} \log \left(\frac{\sinh(Y_t)}{\sinh(Y_0)} \right) - \log \left(\frac{N_t}{N_0} \right) &\leq \log \left(\frac{\sinh(Y_t)}{\sinh(Y_0)} \right) - \log \left(\frac{\cosh(Y_t) - 1}{\cosh(Y_0) - 1} \right) \\ &= \int_0^t \left(\frac{\cosh(Y_s)}{\sinh(Y_s)} - \frac{\sinh(Y_s)}{\cosh(Y_s) - 1} \right) (\partial_s Y_s) ds \\ &= - \int_0^t \frac{\partial_s Y_s}{\sinh(Y_s)} ds \leq 0. \end{aligned}$$

Finally, noting that $\frac{1}{4}(\partial_s \varrho_s)^2 \geq (\partial_s G_s)(\partial_s \varrho_s) - (\partial_s G_s)^2$, we obtain from (3.7) the sought estimate (3.6):

$$\log |\mathfrak{f}'_t(iy + \theta_t)| \leq \frac{1}{4} \int_0^t (\partial_s \varrho_s)^2 \frac{ds}{\ell_s} = \frac{1}{4} \int_0^t (\partial_s \theta_s)^2 \frac{ds}{\lambda_s} \leq \frac{1}{2} \|\frac{1}{\lambda}\|_{[0,T]} E_T(\theta).$$

This proves Item 1. To prove Item 2, note first that the estimate (3.6) already implies that, for every $t \in [0, T]$ and $0 < y < y' \leq y_0$, we have

$$\begin{aligned} |\mathfrak{f}_t(\theta_t + iy) - \mathfrak{f}_t(\theta_t + iy')| &\leq \int_y^{y'} |\mathfrak{f}'_t(\theta_t + iu)| du \\ &\leq y_0 \exp \left(\frac{1}{2} \|\frac{1}{\lambda}\|_{[0,T]} E_T(\theta) \right) \xrightarrow{y_0 \rightarrow 0} 0. \end{aligned}$$

This shows that the radial limit (3.4) exists uniformly in time and in particular is continuous in time. By arguments similar to [RS05, Theorem 4.1], this then implies that θ generates a radial chord $\gamma_{[0,T]}$ in \mathbb{D} . It remains to show that γ is simple. Observe that if γ is not simple, then there exists a time $\tau \in [0, T]$ such that one of the following holds:

- (i) γ intersects the boundary at some point $\gamma(\tau) = x \in \partial\mathbb{D} \setminus \{1\}$ at time $\tau = \tau_x$; or
- (ii) at time τ , the curve γ intersects its own past, so $\gamma(\tau) = \gamma(\tau')$ for some $0 \leq \tau' < \tau$.

If scenario (ii) occurs, then for any intermediate time $s \in (\tau', \tau)$, the part $t \mapsto \mathfrak{g}_s(\gamma_{s+t}) =: \tilde{\gamma}_t$ of the curve after time s hits $\partial\mathbb{D} \setminus \{e^{i\theta_s}\}$ at time $t = \tau - s$. By additivity of the Dirichlet energy (2.16), the energy of the driving function $\tilde{\theta}$ of $\tilde{\gamma}$ satisfies $E_{T-s}(\tilde{\theta}) \leq E_T(\theta)$, so scenario (ii) reduces to scenario (i). It thus remains to show that scenario (i) cannot occur. Thanks to Lemma 3.8, by making a time change we may assume without loss of generality that $\lambda_t \equiv 1$. Suppose, towards a contradiction, that scenario (i) occurs for some $\gamma(\tau) = x \in \partial\mathbb{D} \setminus \{1\}$. Consider the time-evolution (3.2) (with $n = 1$) of $\xi_t := \mathfrak{h}_t(-i \log x) \in (0, 2\pi)$:

$$\frac{d}{dt} \xi_t = \cot \left(\frac{\xi_t - \theta_t}{2} \right) = \cot \left(\frac{\omega_t}{2} \right), \quad \text{where} \quad \omega_t := \xi_t - \theta_t$$

satisfies $\omega_0 = \xi_0 \in (0, 2\pi)$. At the hitting time τ_x to x , we have $\omega_{\tau_x} \in \{0, 2\pi\}$, and

$$\infty > 2E_T(\theta) \geq 2E_{\tau_x}(\theta) \geq 2 \overline{\lim}_{t \rightarrow \tau_x^-} E_t(\theta)$$

$$\begin{aligned}
&= \overline{\lim}_{t \rightarrow \tau_x^-} \int_0^t \left(\frac{d}{ds} \omega_s - \frac{d}{ds} \zeta_s \right)^2 ds \\
&= \overline{\lim}_{t \rightarrow \tau_x^-} \int_0^t \left(\left(\frac{d}{ds} \omega_s \right)^2 - 2 \left(\frac{d}{ds} \omega_s \right) \cot \left(\frac{\omega_s}{2} \right) + \cot^2 \left(\frac{\omega_s}{2} \right) \right) ds \\
&\geq -2 \underline{\lim}_{t \rightarrow \tau_x^-} \int_0^t \left(\frac{d}{ds} \omega_s \right) \cot \left(\frac{\omega_s}{2} \right) ds \\
&= -4 \underline{\lim}_{t \rightarrow \tau_x^-} \int_0^t \frac{d}{ds} \left(\log \sin \left(\frac{\omega_s}{2} \right) \right) ds \\
&= -4 \underline{\lim}_{t \rightarrow \tau_x^-} \log \left(\frac{\sin \left(\frac{\omega_t}{2} \right)}{\sin \left(\frac{\omega_0}{2} \right)} \right) = \infty.
\end{aligned}$$

This contradiction shows that scenario (i) cannot occur, and finishes the proof. \square

Remark 3.10. As a consequence of the proof of Theorem 3.9, we get the following form for the derivative of the inverse Loewner chain (compare with [FS17, Proposition 1]):

$$\log |f'_s(z + \theta_t)| = \log \left(\frac{\sinh(Y_t)}{\sinh(Y_0)} \right) - \log \left(\frac{N_t}{N_0} \right) + \int_0^t \left((\partial_s G_s)(\partial_s \varrho_s) - (\partial_s G_s)^2 \right) \frac{1}{\ell_s} ds.$$

By a closer investigation of the above computation, it should also be possible to extend other results in [FS17] (for example, [FS17, Theorem 4] in the context of Itô-Föllmer type integrals). Such generalizations would be, however, beyond the applications that we have in mind in the present work, so we shall not attempt to do this.

3.3 Finite-energy hulls are simple radial multichords

The purpose of this section is to prove Theorem 1.9. The proof comprises a few steps. We first show that for each $t \in [0, T]$, the hull K_t consists of n disjoint sets which only touch the boundary $\partial\mathbb{D}$ at the starting points z_0^1, \dots, z_0^n (Proposition 3.11). We then derive a generalized version of the restriction property (Proposition 3.12, cf. [LSW03]), which enables us to pass from the case of one radial curve to the case of several curves. We combine these results with Theorem 3.9 to finish the proof of Theorem 1.9 in the end.

Proposition 3.11. *Consider a multiradial Loewner chain with the 1-common parameterization for which the uniformizing conformal mappings $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ satisfy Equation (1.11) with driving functions $z_t^j = e^{i\theta_t^j}$ for $j = 1, \dots, n$, where $\boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in C_{\theta_0}([0, T], \mathcal{X}_n)$. If $J_T(\boldsymbol{\theta}) < \infty$, then we have*

$$K_t = \bigsqcup_{j=1}^n K_t^j, \quad \text{for each } t \in [0, T],$$

where K_t^j are pairwise disjoint connected hulls such that $K_t^j \cap \partial\mathbb{D} = \{e^{i\theta_t^j}\}$, for all j .

Proof. We will first prove that $K_t \cap \partial\mathbb{D} = \{e^{i\theta_0^1}, \dots, e^{i\theta_0^n}\}$ for all $t \in [0, T]$. As the first step, we show that none of the boundary points $x \in \partial\mathbb{D} \setminus \{e^{i\theta_0^1}, \dots, e^{i\theta_0^n}\}$ can be swallowed when the energy is finite. Consider the swallowing times

$$\tau_x := \min_{1 \leq j \leq n} \tau_x^j \quad \text{where} \quad \tau_x^j := \sup \left\{ t \geq 0 : \inf_{s \in [0, t]} |g_s(x) - e^{i\theta_s^j}| > 0 \right\}.$$




Figure 3.1: Illustration of a finite-energy hull and paths separating its components, as in the proof of Proposition 3.11.

Towards a contradiction, suppose that $\tau_x \leq T$. On the one hand, Corollary 2.9 shows that $T < \tau_{\text{coll}}$. On the other hand, if $\tau_x^j = \tau_x^i$ for some $i \neq j$, then

$$|e^{i\theta_s^j} - e^{i\theta_s^i}| \leq |e^{i\theta_s^j} - g_s(x)| + |g_s(x) - e^{i\theta_s^i}| \xrightarrow{s \rightarrow \tau_x^j} 0,$$

which shows that $\tau_{\text{coll}} \leq \tau_x^j$. Hence, we may without loss of generality assume that $\{j_0\} := \arg \min_j \tau_x^j = \{1\}$, so that $\tau_x = \tau_x^{j_0} = \tau_x^1 \leq T$. Consider the time-evolution

$$g_t(e^{iu}) = \exp(ih_t(u)) \quad \text{and} \quad \xi_t := h_t(-i \log x) \in (0, 2\pi), \quad t < \tau_x,$$

and denote

$$\omega_t^j := \xi_t - \theta_t^j, \quad j = 1, \dots, n, \quad \text{so that} \quad \omega_{\tau_x}^1 \in \{0, 2\pi\}.$$

From (3.2) (with $\lambda_t \equiv 1$) we see that

$$\frac{d}{dt} \xi_t = \sum_{j=1}^n \cot\left(\frac{\omega_t^j}{2}\right), \quad t < \tau_x.$$

We will now estimate the multiradial Dirichlet energy of $\boldsymbol{\theta}$ under the assumption that $\tau_x = \tau_x^1 \leq T$, which will lead to a contradiction with the finiteness of the energy:

$$\begin{aligned} \infty &> 2J_T(\boldsymbol{\theta}) \geq 2J_{\tau_x}(\boldsymbol{\theta}) = \int_0^{\tau_x} \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi^j(\boldsymbol{\theta}_s) \right|^2 ds \geq \int_0^{\tau_x} \left| \frac{d}{ds} \theta_s^1 - 2\phi^1(\boldsymbol{\theta}_s) \right|^2 ds \\ &= \int_0^{\tau_x} |V_s - Z_s|^2 ds, \end{aligned} \tag{3.8}$$

$$\text{where} \quad V_s := \frac{d}{ds} \omega_s^1 - \cot\left(\frac{\omega_s^1}{2}\right),$$

$$Z_s := \sum_{j=2}^n \left(\cot\left(\frac{\omega_s^j}{2}\right) + 2 \cot\left(\frac{\omega_s^1 - \omega_s^j}{2}\right) \right).$$

We will show that the righthand side of (3.8) is infinite, which gives a contradiction. First, as $\tau_{\text{coll}} > \tau_x$, there exists a constant $R \in (0, \infty)$ such that $|Z_s| \leq R$ for all $s \leq \tau_x$, so

$$\int_0^{\tau_x} |Z_s|^2 ds \leq R\tau_x.$$

Second, the same computation as in the end of the proof of Theorem 3.9 shows that

$$\lim_{t \rightarrow \tau_x^-} \int_0^t |V_s|^2 ds \geq -4 \lim_{t \rightarrow \tau_x^-} \log \left(\frac{\sin(\frac{\omega_t^1}{2})}{\sin(\frac{\omega_0^1}{2})} \right) = +\infty,$$

since $\omega_{\tau_x}^1 \in \{0, 2\pi\}$. We conclude that

$$\begin{aligned} \infty > (3.8) &= \int_0^{\tau_x} |V_s - Z_s|^2 ds \\ &\geq \underbrace{\lim_{t \rightarrow \tau_x^-} \int_0^t |V_s|^2 ds}_{= +\infty} - \underbrace{\int_0^{\tau_x} |Z_s|^2 ds}_{\leq R\tau_x \in [0, +\infty)} - \underbrace{2 \int_0^{\tau_x} |V_s - Z_s| |Z_s| ds}_{\leq 2\sqrt{R\tau_x} \left(\int_0^{\tau_x} |V_s - Z_s|^2 ds \right)^{1/2}} = \infty, \end{aligned}$$

since the third term is finite by the Cauchy-Schwarz inequality and (3.8):

$$0 \leq 2 \int_0^{\tau_x} |V_s - Z_s| |Z_s| ds \leq \left| 2\sqrt{R\tau_x} \left(\int_0^{\tau_x} |V_s - Z_s|^2 ds \right)^{1/2} \right| \leq 2\sqrt{R\tau_x} \sqrt{2J_{\tau_x}(\boldsymbol{\theta})} < \infty.$$

This gives the sought contradiction — so we conclude that for all $x \in \partial\mathbb{D} \setminus \{e^{i\theta_0^1}, \dots, e^{i\theta_0^n}\}$, we have $\tau_x > T$. We have thus shown that $K_t \cap \partial\mathbb{D} = \{e^{i\theta_0^1}, \dots, e^{i\theta_0^n}\}$ for all $t \in [0, T]$.

To finish, we will prove that $K_t = \sqcup_{j=1}^n K_t^j$ is a union of pairwise disjoint connected hulls K_t^j . Indeed, pick n boundary points $e^{iu^1}, e^{iu^2}, \dots, e^{iu^n} \in \partial\mathbb{D}$ such that $u^j \in (\theta_0^j, \theta_0^{j+1})$ for each j , with the convention that $\theta_0^{n+1} = \theta_0^1 + 2\pi$. Pick n simple paths $\eta^1, \eta^2, \dots, \eta^n$ in $\overline{\mathbb{D}} \setminus K_t$ such that each η^j connects the origin to e^{iu^j} in \mathbb{D} and $\eta^1, \eta^2, \dots, \eta^n$ only intersect at the origin — see Figure 3.1 for an illustration. Then, $\cup_j \eta^j$ separates K_t into n disjoint connected hulls K_t^j satisfying $K_t^j \cap \partial\mathbb{D} = \{e^{i\theta_0^j}\}$, for $j = 1, \dots, n$. \square

We will now prove a generalized version of the restriction property (Proposition 3.12). In the 1-common parameterization, the uniformizing Loewner maps $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ satisfy (1.11) with $z_t^j = e^{i\theta_t^j}$, and the map h_t related to g_t via $g_t(e^{iu}) = \exp(ih_t(u))$ satisfies

$$\partial_t h_t(u) = \sum_{j=1}^n \cot \left(\frac{h_t(u) - \theta_t^j}{2} \right). \quad (3.9)$$

See Figure 3.2 for an illustration of the setup of Proposition 3.12.

Proposition 3.12. *Consider a multiradial Loewner chain with the 1-common parameterization for which the uniformizing conformal mappings $g_t: \mathbb{D} \setminus K_t \rightarrow \mathbb{D}$ satisfy Equation (1.11) with driving functions $z_t^j = e^{i\theta_t^j}$ for $j = 1, \dots, n$, where $\boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in C_{\theta_0}([0, T], \mathcal{X}_n)$. Suppose that $J_T(\boldsymbol{\theta}) < \infty$.*

Fix $k \in \{1, \dots, n\}$ and consider the partition $K_t = R_t^k \sqcup K_t^k$, where K_t^k is the connected component of K_t containing $e^{i\theta_0^k}$ as in Proposition 3.11, and $R_t^k = K_t \setminus K_t^k$ its complement. Define the following quantities, as shown in Figure 3.2:




Figure 3.2: Illustration of the setup of Proposition 3.12.

- Write $\tilde{R}_t^k := g_t(R_T^k \setminus R_t^k)$ for $t \in [0, T]$.
- Let $\varphi_t^k: \mathbb{D} \setminus \tilde{R}_t^k \rightarrow \mathbb{D}$ denote the uniformizing map normalized at the origin.
- Write $\hat{K}_t^k = \varphi_0^k(K_t^k)$.
- Let $\hat{g}_t^k: \mathbb{D} \setminus \hat{K}_t^k \rightarrow \mathbb{D}$ denote the uniformizing map normalized at the origin.
- Define $\hat{h}_t^k \in [0, 2\pi)$ by $\hat{g}_t^k(e^{iu}) = \exp(i\hat{h}_t^k(u))$ for $u \in \mathbb{H}$ such that $e^{iu} \in \mathbb{D} \setminus \hat{K}_t^k$.
- Define $\chi_t^k \in [0, 2\pi)$ by $\varphi_t^k(e^{iu}) = \exp(i\chi_t^k(u))$ for $u \in \mathbb{H}$ such that $e^{iu} \in \mathbb{D} \setminus \tilde{R}_t^k$.

Then, we have

$$\partial_t \hat{h}_t^k(u) = \lambda_t^k \cot\left(\frac{\hat{h}_t^k(u) - \hat{\theta}_t^k}{2}\right), \quad t \in [0, \hat{\tau}_u), \quad (3.10)$$

where $t \mapsto \lambda_t^k := ((\chi_t^k)'(\theta_t^k))^2$ is a continuous weight function, $t \mapsto \hat{\theta}_t^k := \chi_t^k(\theta_t^k)$, is the driving function, and

$$\hat{\tau}_u := \sup\left\{t \geq 0 : \inf_{s \in [0, t]} |\hat{g}_s^k(e^{iu}) - e^{i\hat{\theta}_s^k}| > 0\right\}.$$

Furthermore, we have $E_T(\lambda^k) < \infty$ and $E_T(\hat{\theta}^k) < \infty$.

Proof. See Figure 3.2 for the setup. Note that $\hat{g}_t^k = \varphi_t^k \circ g_t \circ (\varphi_0^k)^{-1}$, since all uniformizing maps are normalized at the origin. Similarly, $\hat{h}_t^k = \chi_t^k \circ h_t \circ (\chi_0^k)^{-1}$.

We see that the hulls $(\hat{K}_t^k)_{t \geq 0}$ are locally growing, since the hulls $(K_t^k)_{t \geq 0}$ are locally growing and contained in the domain of φ_0^k , which is a homeomorphism that extends

continuously to the boundary. This implies that \hat{g}_t^k satisfies the weighted single-curve radial Loewner equation for some weight λ_t^k , with driving function $e^{i\hat{\theta}_t^k} = \varphi_t^k(e^{i\theta_t^k})$. Consequently, \hat{h}_t^k satisfies an equation of the form (3.10) with $\hat{\theta}_t^k = \chi_t^k(\theta_t)$, again with some weight λ_t^k .

In order to find λ_t^k , we compute the time derivative of $\hat{h}_t(u) = \chi_t^k \circ h_t \circ (\chi_0^k)^{-1}(u)$ using the chain rule, substituting into (3.9), and then setting the result equal to the righthand side of (3.10), which shows that

$$\lambda_t^k \cot\left(\frac{\chi_t^k(v) - \chi_t^k(\theta_t^k)}{2}\right) = (\partial_t \chi_t^k)(v) + (\chi_t^k)'(v) \sum_{j=1}^n \cot\left(\frac{v - \theta_t^j}{2}\right), \quad (3.11)$$

where $v = (h_t \circ (\chi_0^k)^{-1})(u)$ for $e^{iu} \in \mathbb{D} \setminus \hat{K}_t^k$. The above equation (3.11) holds whenever $e^{iv} \in \mathbb{D} \setminus \tilde{R}_t^k$. We will solve for λ_t^k and evaluate the limit as $v \rightarrow \theta^* := \theta_t^k$. For this purpose, we define the notation θ^* to be clear that this value is fixed, even as we consider χ_s^k for $s \in [t - \epsilon, t + \epsilon]$. However, due to the singularity at θ^* , we first take care to check that all relevant maps are jointly continuous in a neighborhood of (t, θ^*) .

The conformal mapping χ_t^k is well-defined and extends continuously to the boundary in an \mathbb{H} -neighborhood around θ^* , so by Schwarz reflection, χ_t^k extends conformally to a neighborhood of θ^* in the complex plane. Moreover, the conformal maps χ_s^k are continuously differentiable in s , and we can find $\epsilon > 0$ and \mathbb{H} -neighborhood \mathcal{O} around θ^* such that the map $(v, s) \mapsto \partial_s \chi_s^k(v)$ exists and is jointly continuous on $\bar{\mathcal{O}} \times [t - \epsilon, t + \epsilon]$. Again extending by Schwarz reflection, there exists a \mathbb{C} -neighborhood $\mathcal{U} \ni \theta^*$ such that for $s \in [t - \epsilon, t + \epsilon]$, each $\partial_s \chi_s^k$ is conformal on \mathcal{U} , the map $(v, s) \mapsto \partial_s \chi_s^k(v)$ is jointly continuous on $\bar{\mathcal{U}} \times [t - \epsilon, t + \epsilon]$. Without loss of generality, we may assume that \mathcal{U} is simply connected with rectifiable boundary. Finally, for each $m \in \mathbb{N}$, the map¹¹ $(z, t) \mapsto \partial_t \chi_t^{(m)}$ is jointly continuous on $\bar{\mathcal{U}} \times [t - \epsilon, t + \epsilon]$, which we can see by applying the Cauchy differentiation formula,

$$\frac{\chi_{t+s}^{(m)}(z) - \chi_t^{(m)}(z)}{s} = \frac{m!}{2\pi i} \int_{\partial \mathcal{U}} \frac{\chi_{t+s}(w) - \chi_t(w)}{s} \frac{dw}{(w-z)^m}, \quad \text{for all } z \in \mathcal{U},$$

and the dominated convergence theorem, which yield

$$\partial_t \chi_t^{(m)}(z) = \frac{m!}{2\pi i} \int_{\partial \mathcal{U}} \frac{\partial_t \chi_t(w)}{(w-z)^m} dw = \left(\frac{d}{dz}\right)^m \partial_t \chi_t(z).$$

Next, we Laurent expand both sides of (3.11) around the singularity θ^* , to obtain

$$\frac{2\lambda_t^k}{\chi_t^k(v) - \chi_t^k(\theta_t^k)} + O(1) = (\partial_t \chi_t^k)(v) + (\chi_t^k)'(v) \left(\frac{2}{v - \theta_t^k} + O(1) \right), \quad v \rightarrow \theta^*.$$

We can solve for λ_t^k by multiplying both sides by $\frac{1}{2}(\chi_t^k(v) - \chi_t^k(\theta_t^k))$ and taking the limit as $v \rightarrow \theta^*$ (which is justified by the continuity checks above):

$$\lambda_t^k = \lim_{v \rightarrow \theta_t^k} \left((\chi_t^k)'(v) \frac{\chi_t^k(v) - \chi_t^k(\theta_t^k)}{v - \theta_t^k} \right) = ((\chi_t^k)'(\theta_t^k))^2 \in (0, \infty).$$

Let us also note that $(\chi_t^k(\theta^*))' \neq 0$, since (by Schwarz reflection, as above) χ_t^k is conformal in a neighborhood of θ^* . Finally, the joint continuity of $(s, v) \mapsto (\chi_s^k)'(v)$ implies that λ_t^k is

¹¹Here, we use the superscript “(m)” to denote the m :th complex derivative.

continuous and therefore locally integrable. We thus conclude that $\lambda_t^k = ((\chi_t^k)'(\theta_t^k))^2$ is a continuous weight function. This proves the asserted equality (3.10).

Finally, we check that $E_T(\hat{\theta}_t^k) < \infty$ and $E_T(\lambda_t^k) < \infty$: differentiating, we have

$$\begin{aligned}\frac{d}{dt}\hat{\theta}_t^k &= \frac{d}{dt}\chi_t^k(\theta_t^k) = \partial_t\chi_t^k(\theta_t^k) + ((\chi_t^k)'(\theta_t^k))\frac{d}{dt}\theta_t^k, \\ \frac{d}{dt}\lambda_t^k &= \frac{d}{dt}((\chi_t^k)'(\theta_t^k))^2 = 2(\partial_t(\chi_t^k)'(\theta_t^k) + ((\chi_t^k)''(\theta_t^k))\frac{d}{dt}\theta_t^k).\end{aligned}$$

By joint continuity, all of $\partial_t\chi_t^k(\theta_t^k)$, and $(\chi_t^k)'(\theta_t^k)$, and $\partial_t(\chi_t^k)'(\theta_t^k)$, and $(\chi_t^k)''(\theta_t^k)$ are uniformly bounded on the compact interval $[0, T]$, while by Corollary 2.9, the function θ_t^k has finite Dirichlet energy and so $t \mapsto \frac{d}{dt}\theta_t^k$ is integrable on $[0, T]$ (as θ^k is absolutely continuous). Thus, we see that $E_T(\hat{\theta}_t^k) < \infty$ and $E_T(\lambda_t^k) < \infty$, proving the last claim. \square

Proof of Theorem 1.9. The case of $n = 1$ is covered by Theorem 3.9, so we will consider the case where $n > 1$. By Proposition 3.11, the hull $K_t = \bigsqcup_{j=1}^n K_t^j$ is a disjoint union of n connected components K_t^j containing the starting points θ_0^j , with $1 \leq j \leq n$. It thus suffices to show that each such connected component is generated by a simple curve.

Fix $k \in \{1, \dots, n\}$. By conjugating by a suitable rotation, we may assume without loss of generality that $\theta_0^k = 0$. With notation from Proposition 3.12, the map \hat{h}_t^k satisfies the (single) radial Loewner equation (3.2) (with $n = 1$) parametrized by the continuous weight function λ^k and with driving function $\hat{\theta}^k$, which has finite Dirichlet energy $E_T(\hat{\theta}^k) < \infty$. Thus, by Theorem 3.9 we know that \hat{K}^k is a simple radial chord, so $K_T^k = (\varphi_0^k)^{-1}(\hat{K}_T^k)$ is also a simple curve (as a conformal image of such). As the choice of the index $k \in \{1, \dots, n\}$ was arbitrary, we conclude that every connected component of K_T is a simple curve. \square

3.4 Proof of the LDP for multiradial SLE $_{0+}$

In this section, we prove the main result, Theorem 1.8. Let us begin by recalling that the Loewner transform \mathcal{L}_t , defined in (1.12), sends driving functions to the hulls generated by the multiradial Loewner equation (1.11) with 1-common parameterization. Hence, it would be natural to apply the contraction principle, (Theorem E) to deduce the LDP for multiradial SLE $_\kappa$ from the LDP for Dyson Brownian motion (Theorem 1.2). However, as the Loewner transform is not continuous, we cannot do this directly. Instead, we first restrict \mathcal{L}_t into a subset with full measure where it is continuous, and use Proposition 3.13 below, which will allow us to derive the large deviations result for multiradial SLE $_{0+}$.

Theorem E (Contraction principle, [DZ10, Theorem 4.2.1]). *Let X and Y be Hausdorff topological spaces, and let $f: X \rightarrow Y$ be a continuous map. Suppose that the family $(\mathbf{P}^\kappa)_{\kappa>0}$ of probability measures satisfies an LDP in X with good rate function $I: X \rightarrow [0, +\infty]$, that is, for any closed subset F_0 and open subset O_0 of X , we have*

$$\overline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^\kappa[F_0] \leq - \inf_{x \in F_0} I(x) \quad \text{and} \quad \underline{\lim}_{\kappa \rightarrow 0+} \kappa \log \mathbf{P}^\kappa[O_0] \geq - \inf_{x \in O_0} I(x),$$

and the level set $I^{-1}[0, c]$ is a compact subset of X , for all $c \in [0, \infty)$. Define

$$J(y) := \inf_{x \in f^{-1}\{y\}} I(x), \quad y \in Y.$$

Then, the family $(\mathbb{P}^\kappa)_{\kappa>0} := (\mathbb{P}^\kappa \circ f^{-1})_{\kappa>0}$ of pushforward probability measures satisfies an LDP in Y with good rate function J : for any closed subset F and open subset O of Y ,

$$\overline{\lim}_{\kappa \rightarrow 0^+} \kappa \log \mathbb{P}^\kappa[F] \leq - \inf_{y \in F} J(y) \quad \text{and} \quad \underline{\lim}_{\kappa \rightarrow 0^+} \kappa \log \mathbb{P}^\kappa[O] \geq - \inf_{y \in O} J(y),$$

and the level set $J^{-1}[0, c]$ is a compact subset of Y , for all $c \in [0, \infty)$.

Proposition 3.13 (Restricted LDP). *Let X be a Hausdorff topological space, $(\mathbb{P}^\kappa)_{\kappa>0}$ a family of probability measures on X , and $I: X \rightarrow [0, +\infty]$ a rate function. Suppose $A \subset X$ is a measurable subset such that $I^{-1}[0, \infty) \subset A$ and $\mathbb{P}^\kappa[A] = 1$ for every $\kappa > 0$. Then, the family $(\mathbb{P}|_A^\kappa)_{\kappa>0}$ of restricted measures satisfies an LDP in A with rate function $I|_A$ if and only if the family $(\mathbb{P}^\kappa)_{\kappa>0}$ satisfies an LDP in X with rate function I . Specifically, for every closed $F \subset X$ and open $O \subset X$, the following equivalences hold:*

$$\begin{aligned} \overline{\lim}_{\kappa \rightarrow 0} \kappa \log \mathbb{P}^\kappa[F] \geq - \inf_{x \in F} I(x) &\iff \overline{\lim}_{\kappa \rightarrow 0} \kappa \log \mathbb{P}|_A^\kappa[F \cap A] \geq - \inf_{x \in F \cap A} I|_A(x), \\ \underline{\lim}_{\kappa \rightarrow 0} \kappa \log \mathbb{P}^\kappa[O] \leq - \inf_{x \in O} I(x) &\iff \underline{\lim}_{\kappa \rightarrow 0} \kappa \log \mathbb{P}|_A^\kappa[O \cap A] \geq - \inf_{x \in O \cap A} I|_A(x). \end{aligned}$$

Furthermore, I is a good rate function if and only if $I|_A$ is a good rate function.

Proof. Since $\mathbb{P}^\kappa[A] = 1$ and $I^{-1}[0, \infty) \subset A$, for any measurable set $B \subset X$, we have

$$\mathbb{P}^\kappa[B] = \mathbb{P}^\kappa[B \cap A] = \mathbb{P}|_A^\kappa[B \cap A] \quad \text{and} \quad \inf_{x \in B} I(x) = \inf_{x \in B \cap A} I(x) = \inf_{x \in B \cap A} I|_A(x).$$

The equivalences follow. The last claim follows from the assumption $I^{-1}[0, \infty) \subset A$. \square

Remark 3.14. In Proposition 3.13 it is enough to assume that $A \subset X$ is some (not necessarily measurable) subset such that $I^{-1}[0, \infty) \subset A$ and for every $\kappa > 0$ there is some measurable set $A^\kappa \subset A$ such that $\mathbb{P}^\kappa[A^\kappa] = 1$. We equip A with the subspace topology induced from X , and define the restricted measures by $\mathbb{P}|_A^\kappa[E] := \mathbb{P}^\kappa[E \cap A^\kappa]$.

Recall that a *hull* is a compact set $K \subset \overline{\mathbb{D}}$ such that $\mathbb{D} \setminus K$ is simply connected, $0 \in \mathbb{D} \setminus K$, and the closure $\overline{K} \cap \overline{\mathbb{D}} = K$ in \mathbb{C} . For each hull K , we denote by $g_K: \mathbb{D} \setminus K \rightarrow \mathbb{D}$ the uniformizing map normalized at the origin, i.e., satisfying $g_K(0) = 0$ and $g'_K(0) > 0$. We call $\log g'_K(0)$ the *capacity* of K , so that the complement of K has conformal radius $1/g'_K(0) = e^{-\log g'_K(0)}$. For each fixed $T \in (0, \infty)$, we denote

$$\mathcal{K}_T = \{\text{hulls } K \subset \overline{\mathbb{D}} \text{ of capacity } nT\}. \quad \mathcal{K} := \bigcup_{T \geq 0} \mathcal{K}_T. \quad (3.12)$$

We endow the space \mathcal{K} of hulls with the coarsest (Carathéodory) topology for which a sequence $(K_{(k)})_{k \in \mathbb{N}}$ in \mathcal{K} converges to $K \in \mathcal{K}$ if and only if the associated functions $g_{K_{(k)}}^{-1}$ converge to g_K^{-1} uniformly on compact subsets of \mathbb{D} . By [Dur83, Theorem 3.1], this is equivalent to the *Carathéodory kernel convergence* of the complementary domains $D_{(k)} := \mathbb{D} \setminus K_{(k)}$ to $D := \mathbb{D} \setminus K$ with respect to the origin: for any subsequence $(D_{(k_j)})_{j \in \mathbb{N}}$ we have $D = \bigcup_{j \geq 1} (\bigcap_{i \geq j} D_{(k_i)})_0$, denoting by V_0 the connected component of a set $V \subset \overline{\mathbb{D}}$ containing the origin. Because we require that $\overline{K} \cap \overline{\mathbb{D}} = K$ for any hull K , we see that for two hulls $K \neq \tilde{K}$, we have $\mathbb{D} \setminus K \neq \mathbb{D} \setminus \tilde{K}$, which shows that the Carathéodory topology on the set \mathcal{K} has the Hausdorff (T2) property. (This is required in the contraction principle.)

Although $\mathcal{K} \subset \mathcal{C}$ is contained in the set of compact subsets of $\overline{\mathbb{D}}$, the Carathéodory and Hausdorff (1.13) topologies on \mathcal{K} are not comparable. However, we can characterize their difference in the following useful manner (via a radial analogue of [PW24, Lemma 2.3]).

Lemma 3.15. *Suppose that a sequence $(K_{(k)})_{k \in \mathbb{N}}$ in \mathcal{K} converges to $K \in \mathcal{K}$ in the Carathéodory sense and to $\tilde{K} \in \mathcal{C}$ in the Hausdorff metric. Then $\mathbb{D} \setminus K = (\mathbb{D} \setminus \tilde{K})_0$. In particular, we have $\mathbb{D} \cap K = \mathbb{D} \cap \tilde{K}$ if and only if $\mathbb{D} \setminus \tilde{K}$ is connected.*

Proof. This follows by the same proof as [PW24, Lemma 2.3]. \square

The Loewner transform $\mathcal{L}_T: C([0, T], \mathcal{X}_n) \rightarrow \mathcal{C}$ sends driving functions to hulls,

$$\mathcal{L}_T(\boldsymbol{\theta}) := \{z \in \overline{\mathbb{D}} \mid \tau_z \leq T\} \subset \mathcal{K}_T \subset \mathcal{C}.$$

It is well-known that \mathcal{L}_T is continuous in the Carathéodory sense (see [MS16b, Proposition 6.1] for a proof for general Loewner chains). While \mathcal{L}_T is not continuous in the Hausdorff metric, its discontinuities occur outside of the set of simple curves (cf. [PW24, Lemma 2.4]).

Proof of Theorem 1.8. We can write the Loewner transform as a composition $\mathcal{L}_T = \iota \circ \mathcal{L}'_T$, where $\mathcal{L}'_T: C([0, T], \mathcal{X}_n) \rightarrow \mathcal{K}_T$ is the Loewner transform to the set (3.12) of hulls of capacity nT , and $\iota: \mathcal{K} \hookrightarrow \mathcal{C}$ is the inclusion of the hulls to the compact subsets of $\overline{\mathbb{D}}$. Now, the map \mathcal{L}'_T is continuous in the Carathéodory sense (e.g., by [MS16b, Proposition 6.1]), so Theorem 1.2 and the contraction principle (Theorem E) together imply that the initial segments $\gamma_{[0, T]}^\kappa$ of multiradial SLE $_\kappa$ curves with laws $(\mathbb{P}^\kappa)_{\kappa > 0}$ satisfy an LDP in \mathcal{K}_T (in the Carathéodory sense) with good rate function $I'_T: \mathcal{K}_T \rightarrow [0, +\infty]$ defined similarly to (1.14),

$$I'_T(K) := \inf_{\boldsymbol{\theta} \in (\mathcal{L}'_T)^{-1}(K)} J_T(\boldsymbol{\theta}).$$

Next, denote by $\mathcal{A} \subset \mathcal{K}_T$ the set of simple radial multichords with total capacity nT which are generated by a driving function in the 1-common parameterization (as in Definition 3.2). For $\kappa \leq 4$,¹² we have $\mathbb{P}^\kappa[\mathcal{A}] = 1$, while by Theorem 1.9, the set \mathcal{A} contains all finite-energy hulls. Thus, we deduce from Proposition 3.13¹³ that the family $(\mathbb{P}^\kappa)_{\kappa > 0}$ satisfies an LDP in \mathcal{A} in the Carathéodory sense and with good rate function $I'_T|_{\mathcal{A}}$.

Now, we claim that the restricted map $\iota|_{\mathcal{A}}: \mathcal{A} \hookrightarrow \mathcal{C}$ is continuous, when the former space carries the Carathéodory topology and the latter the Hausdorff metric. Indeed, suppose that a sequence $(\boldsymbol{\eta}_{(k)})_{k \in \mathbb{N}}$ of simple radial multichords in \mathcal{A} converges to $\boldsymbol{\eta} \in \mathcal{A}$ in the Carathéodory sense. By compactness of \mathcal{C} , passing to a subsequence, $\boldsymbol{\eta}_{(k)}$ also converge in the Hausdorff metric to some $\tilde{\boldsymbol{\eta}} \in \mathcal{C}$. Then Lemma 3.15 implies that $\mathbb{D} \cap \boldsymbol{\eta} = \mathbb{D} \cap \tilde{\boldsymbol{\eta}}$ (since otherwise, $\boldsymbol{\eta}$ would have non-empty interior). Furthermore, since $\boldsymbol{\eta}$ is a hull and $\tilde{\boldsymbol{\eta}}$ is compact, this implies that $\boldsymbol{\eta} = \overline{\boldsymbol{\eta} \cap \mathbb{D}} = \overline{\mathbb{D} \cap \tilde{\boldsymbol{\eta}}} \subseteq \tilde{\boldsymbol{\eta}}$. Now, if $x \in (\partial\mathbb{D}) \setminus \boldsymbol{\eta}$, then the sets $\boldsymbol{\eta}_{(k)}$ avoid x for large enough k , so $x \notin \tilde{\boldsymbol{\eta}}$. It follows that $\boldsymbol{\eta}$ and $\tilde{\boldsymbol{\eta}}$ agree on the interior of the disk and also on the boundary, so $\boldsymbol{\eta} = \tilde{\boldsymbol{\eta}}$. This shows that $\iota|_{\mathcal{A}}: \mathcal{A} \hookrightarrow \mathcal{C}$ is continuous. Therefore, we can apply the contraction principle (Theorem E) again to deduce

¹²In the context of Remark 3.6 concerning other SLE variants, this is replaced by $\kappa \leq 4 \wedge a$.

¹³Note that \mathcal{A} contains a \mathbb{P}^κ -measurable subset of full measure, because the solution of the Loewner equation is measurable with respect to the driving process (whose law is \mathbb{P}^κ). See also Remark 3.14.

that the pushforward measures $(\mathbb{P}^\kappa \circ (\iota|_{\mathcal{A}}))_{\kappa>0}$ satisfy an LDP in $\iota(\mathcal{A}) \subset \mathcal{C}$ in the Hausdorff metric with good rate function $(I'_T \circ \iota^{-1})|_{\iota(\mathcal{A})} = (I_T)|_{\iota(\mathcal{A})}$. From this, we conclude again using Proposition 3.13 that the initial segments $\gamma_{[0,T]}^\kappa$ of multiradial SLE $_\kappa$ curves with laws $(\mathbb{P}^\kappa)_{\kappa>0}$ indeed satisfy the LDP (1.15, 1.16) in \mathcal{C} with good rate function I_T . \square

4 Large-time behavior of finite-energy systems

By the definition of multiradial Loewner energy, finite-energy curves are exactly those whose driving functions have finite multiradial Dirichlet energy. Intuitively, as $t \rightarrow \infty$ we expect the interacting particle system that describes these driving functions to approach an equally-spaced configuration. This section makes these ideas precise. Given an initial configuration, Proposition 4.2 in Section 4.1 concerns the existence and uniqueness of the zero-energy flow for all time, and Proposition 4.5 concerns its convergence to the static equally-spaced configuration. We consider the convergence of finite-energy systems in Section 4.2 (see in particular Proposition 4.7 and Remark 4.9). Theorem 1.10 follows by collecting the results of this section:

Proof of Theorem 1.10. This is the content of Propositions 4.5 & 4.7 below. \square

4.1 Zero-energy systems: existence, uniqueness, and asymptotics

To begin, we will show that infinite-time zero-energy flows do exist. Clearly, the n -dimensional Dirichlet energy E_T appearing in (2.15) is non-negative and attains the minimum $E_T(\boldsymbol{\theta}_0) = 0$ at the constant function $\boldsymbol{\theta} \equiv \boldsymbol{\theta}_0$. Although the sign of the functional Φ_T^0 is not clear from its formula (2.18), Proposition 4.2 below shows that also J_T attains the minimum zero. We investigate the minimizers in more detail in Proposition 4.5.

Lemma 4.1. *Fix $\boldsymbol{\theta} \in \mathcal{X}_n$ and, using the convention that $\theta^{n+1} = \theta^1 + 2\pi$ as in (1.2), let $j \in \arg \min_{1 \leq k \leq n} (\theta^{k+1} - \theta^k)$. Then, the functions (1.5) satisfy*

$$\phi^{j+1}(\boldsymbol{\theta}) - \phi^j(\boldsymbol{\theta}) \geq \pi - n \left(\frac{\theta^{j+1} - \theta^j}{2} \right) \geq 0. \quad (4.1)$$

Proof. To streamline the notation, we make the change of variables $\omega^k := \theta^k/2$. Rewriting the lefthand side of (4.1) in terms of ω^k using (1.5) yields

$$\phi^{j+1}(\boldsymbol{\theta}) - \phi^j(\boldsymbol{\theta}) = \sum_{k=1}^{n-1} \left(\cot(\omega^{j+1} - \omega^{j+1-k}) - \cot(\omega^j - \omega^{j-k}) \right) \quad (4.2)$$

A key observation is that the sum is telescoping when the ω^j are equally-spaced on $[0, \pi)$, an idea that will be used again in the proof of Proposition 4.5. Note that

$$\cot(u) - \cot(v) \geq v - u, \quad 0 < u \leq v < \pi, \quad (4.3)$$

since $\frac{d}{du} \cot(u) = -\csc^2(u) \leq -1$. We can use this to bound terms in (4.2) as

$$\cot(\omega^{j+1} - \omega^k) - \cot(\omega^j - \omega^{k-1}) \geq (\omega^j - \omega^{k-1}) - (\omega^{j+1} - \omega^k)$$

$$= -(\omega^{j+1} - \omega^j) + (\omega^k - \omega^{k-1}).$$

Substituting these bounds into (4.2), then adding and subtracting $(\omega^{j+1} - \omega^j)$, we obtain

$$\phi^{j+1}(\boldsymbol{\theta}) - \phi^j(\boldsymbol{\theta}) \geq \left(\sum_{k=1}^n (\omega^k - \omega^{k-1}) \right) - n(\omega^{j+1} - \omega^j) = \pi - n(\omega^{j+1} - \omega^j),$$

which gives the asserted inequality (4.1) (after substituting back $\omega^k = \theta^k/2$). \square

Proposition 4.2. *The system of differential equations on $C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$ given by*

$$\frac{d}{dt}\theta_t^j = 2\phi^j(\boldsymbol{\theta}_t) \quad \text{for all } t \geq 0 \text{ and } j \in \{1, \dots, n\} \quad (4.4)$$

has a unique solution for each initial configuration $\boldsymbol{\theta}_0 \in \mathcal{X}_n$.

Proof. Using the convention that $\theta^{n+1} = \theta^1 + 2\pi$ as in (1.2), write

$$\Delta_{\boldsymbol{\theta}} := \min_{1 \leq j \leq n} |\theta^{j+1} - \theta^j| \in \left[0, \frac{2\pi}{n}\right], \quad \boldsymbol{\theta} = (\theta^1, \dots, \theta^n) \in \mathcal{X}_n, \quad (4.5)$$

and consider the space

$$\mathcal{X}_n^\varepsilon = \{\boldsymbol{\theta} \in \mathcal{X}_n \mid \Delta_{\boldsymbol{\theta}} > \varepsilon\}, \quad \varepsilon > 0.$$

For each fixed $\varepsilon > 0$, each map $\boldsymbol{\theta} \mapsto 2\phi^j(\boldsymbol{\theta})$ appearing on the righthand side of the ODE (4.4) is Lipschitz on $\mathcal{X}_n^\varepsilon$ by (1.5), so for $\boldsymbol{\theta}_0 \in \mathcal{X}_n^\varepsilon$ a unique solution exists in $\mathcal{X}_n^\varepsilon$ up until the time when the boundary $\partial\mathcal{X}_n^\varepsilon$ is hit. (Indeed, the Picard-Lindelöf theorem implies that the unique solution exists on a time interval whose length depends only on the Lipschitz constant, which in turn only depends on ε .) Thus, it suffices to show that any solution to (4.4) with initial condition in $\mathcal{X}_n^\varepsilon$ stays in $\mathcal{X}_n^\varepsilon$ without hitting the boundary. To establish this, we show that the map $t \mapsto \Delta(t) := \Delta_{\boldsymbol{\theta}_t}$ is non-decreasing. As Δ is differentiable almost everywhere, it suffices to show that $\frac{d}{dt}\Delta(t) \geq 0$ for each $t > 0$ where it exists.

To this end, note that the ODE (4.4) and Lemma 4.1 together imply that

$$\frac{d}{dt}(\theta_t^{j+1} - \theta_t^j) = 2(\phi^{j+1}(\boldsymbol{\theta}_t) - \phi^j(\boldsymbol{\theta}_t)) \geq 2\pi - n\Delta(t) \geq 0, \quad (4.6)$$

for any index $j \in A_t := \arg \min_{1 \leq k \leq n} (\theta_t^{k+1} - \theta_t^k)$. From this, we deduce that

$$\frac{d}{dt}\Delta(t) = \min_{j \in A_t} \frac{d}{dt}(\theta_t^{j+1} - \theta_t^j) \geq 0. \quad (4.7)$$

This concludes the proof (see also Remark 4.3). \square

Remark 4.3. Equations (4.6, 4.7) imply that if $\Delta(t) < 2\pi/n$, then $\frac{d}{dt}\Delta(t) > 0$, while if $\Delta(t) = 2\pi/n$, then $\Delta(t)$ stays constant after time t , since $\frac{d}{dt}\Delta(t) = 0$.

Remark 4.4. Remark 2.13 could be used to give an alternative, indirect proof of Proposition 4.2. Indeed, for every $T \geq 0$, there exists at least one minimizer $\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, T], \mathcal{X}_n)$ with $J_T(\boldsymbol{\theta}) = 0$. From Definition 1.1, we see that such minimizers must satisfy Equation (4.5). Since $T \geq 0$ was arbitrary, we can extend the solution for all times, $\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$.

The next result shows that, for any initial configuration, the zero-energy particle system eventually approaches a static equally-spaced configuration, exponentially fast.

Proposition 4.5. *Suppose that $J(\boldsymbol{\theta}) = 0$. Then, there exists $\zeta \in \mathbb{R}$ such that*

$$\lim_{t \rightarrow \infty} \boldsymbol{\theta}_t = \left(\zeta, \zeta + \frac{2\pi}{n}, \dots, \zeta + \frac{(n-1)2\pi}{n} \right), \quad (4.8)$$

and the convergence is exponentially fast with exponential rate n .

Proof. We will first show that

$$d(\boldsymbol{\theta}_t) := \max_{1 \leq j \leq n} \left| \frac{2\pi}{n} - (\theta_t^{j+1} - \theta_t^j) \right| \xrightarrow{t \rightarrow \infty} 0 \quad (4.9)$$

and this convergence happens exponentially fast at rate n . In fact, for this it is actually sufficient to show that the *smallest* gap approaches $2\pi/n$ as $t \rightarrow \infty$. Indeed, consider

$$y_t := \frac{2\pi}{n} - \min_{1 \leq j \leq n} (\theta_t^{j+1} - \theta_t^j) = \frac{2\pi}{n} - \Delta(t) \geq 0, \quad (4.10)$$

$$Y_t := \max_{1 \leq j \leq n} (\theta_t^{j+1} - \theta_t^j) - \frac{2\pi}{n} \geq 0, \quad (4.11)$$

where $\Delta(t) := \min_j |\theta_t^{j+1} - \theta_t^j|$. Since the gaps $(\theta_t^{j+1} - \theta_t^j)$ sum up to 2π , we see that $Y_t \leq (n-1)y_t$, and thus,

$$d(\boldsymbol{\theta}_t) = \max\{y_t, Y_t\} \leq (n-1)y_t. \quad (4.12)$$

From Definition 1.1 and the assumption $J(\boldsymbol{\theta}) = 0$, we know that $\boldsymbol{\theta}$ satisfies the ODE (4.4) from Proposition 4.2. From (4.7), we then deduce that, for almost every $t \geq 0$, there exists an index j such that

$$\frac{d}{dt} y_t = -2(\phi^{j+1}(\boldsymbol{\theta}_t) - \phi^j(\boldsymbol{\theta}_t)) \leq -ny_t \quad \implies \quad y_t \leq y_0 e^{-nt}.$$

It follows that

$$\left| \frac{2\pi}{n} - (\theta_t^{j+1} - \theta_t^j) \right| \leq (n-1)y_t \leq (n-1)y_0 e^{-nt}, \quad j \in \{1, \dots, n\}. \quad (4.13)$$

As $t \rightarrow \infty$, the righthand side approaches zero exponentially fast with rate n , yielding (4.9).

It remains to prove the convergence of $\boldsymbol{\theta}_t$ to the static equally-spaced configuration (4.8). From (4.13), we have

$$\frac{2\pi}{n} - (n-1)e^{-nt} \leq \theta_t^{j+1} - \theta_t^j \leq \frac{2\pi}{n} + (n-1)e^{-nt}, \quad j \in \{1, \dots, n\}. \quad (4.14)$$

Fix t_0 such that $(n-1)e^{-nt_0} < \frac{2\pi}{n^2}$. Then $(\theta_t^j - \theta_t^{j-k}) \in (0, \pi)$ and $(\theta_t^{j+k} - \theta_t^j) \in (0, \pi)$ for all $k \in \{1, 2, \dots, \lfloor n/2 \rfloor\}$ and $t \geq t_0$. Applying (4.14) k times and using the identity $\frac{d}{du} \cot(u) = -\csc^2(u) \leq -1$, we see that

$$\cot\left(\frac{k\pi}{n} + \frac{k(n-1)}{2}e^{-nt}\right) \leq \cot\left(\frac{\theta_t^j - \theta_t^{j-k}}{2}\right) \leq \cot\left(\frac{k\pi}{n} - \frac{k(n-1)}{2}e^{-nt}\right), \quad (4.15)$$

and similarly for $\cot\left(\frac{\theta_t^{j+k}-\theta_t^j}{2}\right)$. We thus obtain

$$\begin{aligned}
\left|\frac{d}{dt}\theta_t^j\right| &\leq 2 \sum_{k=1}^{\lfloor n/2 \rfloor - 1} \left| \cot\left(\frac{\theta_t^j - \theta_t^{j-k}}{2}\right) - \cot\left(\frac{\theta_t^{j+k} - \theta_t^j}{2}\right) \right| \\
&\leq 2 \sum_{k=1}^{\lfloor n/2 \rfloor - 1} \left(\cot\left(\frac{k\pi}{n} - \frac{k(n-1)}{2}e^{-nt}\right) - \cot\left(\frac{k\pi}{n} + \frac{k(n-1)}{2}e^{-nt}\right) \right) \quad [\text{by (4.15)}] \\
&\leq 2(n-1)e^{-nt} \sum_{k=1}^{\lfloor n/2 \rfloor - 1} k \quad [\text{by (4.3)}] \\
&\leq \frac{(n-1)^2(n-3)}{4} e^{-nt} \xrightarrow{t \rightarrow \infty} 0,
\end{aligned}$$

when n is odd, and similarly,

$$\begin{aligned}
\left|\frac{d}{dt}\theta_t^j\right| &\leq 2 \underbrace{\left| \cot\left(\frac{\theta_t^j - \theta_t^{j-n/2}}{2}\right) \right|}_{\leq \frac{n(n-1)}{4} e^{-nt}} + 2 \underbrace{\sum_{k=1}^{\lfloor n/2 \rfloor - 1} \left| \cot\left(\frac{\theta_t^j - \theta_t^{j-k}}{2}\right) - \cot\left(\frac{\theta_t^{j+k} - \theta_t^j}{2}\right) \right|}_{\leq \frac{(n-1)^2(n-3)}{4} e^{-nt}} \\
&\xrightarrow{t \rightarrow \infty} 0,
\end{aligned}$$

when n is even. In particular, we see that there exists a constant $c \in (0, \infty)$ such that

$$\int_{t_0}^{\infty} \left|\frac{d}{dt}\theta_t^j\right| dt \leq cn^3 e^{-nt_0} < \infty,$$

which shows that, first of all, $\lim_{t \rightarrow \infty} \theta_t^j$ exists and is given by (4.8) for some $\zeta \in \mathbb{R}$, and second of all, the convergence happens with exponential rate n . \square

Remark 4.6. If $J_T^{a,\rho}(\theta) = 0$, we get instead

$$\frac{d}{dt}y_t = -2\left((\phi_a^{j+1}(\theta_t) + \rho) - (\phi_a^j(\theta_t) + \rho)\right) = -\frac{a}{4}\left(2(\phi^{j+1}(\theta_t) - \phi^j(\theta_t))\right) \leq -\frac{a}{4}ny_t.$$

Hence, the proofs of Propositions 4.2 and 4.5 give that the differential equation

$$\frac{d}{dt}\theta_t^j = 2\phi_a^j(\theta_j) + \rho$$

has a unique solution for each initial configuration $\theta_0 \in \mathcal{X}_n$, and it satisfies

$$\lim_{t \rightarrow \infty} (\theta_t - \rho t) = \left(\zeta, \zeta + \frac{2\pi}{n}, \dots, \zeta + \frac{(n-1)2\pi}{n}\right),$$

for some $\zeta \in \mathbb{R}$, where the convergence is exponentially fast with exponential rate $\frac{a}{4}n$.

4.2 Finite-energy systems

We will now show that any function with finite multiradial Dirichlet energy converges to an equally-spaced system in the long run. However, if the multiradial energy is non-zero, it is possible that the convergence rate is very slow and that the system continues slow rotation for all time. (Compare to Proposition 4.5 for zero-energy systems, and see Remark 4.8.)

Proposition 4.7. Consider a function $\boldsymbol{\theta} \in C_{\theta_0}([0, \infty), \mathcal{X}_n)$. If $J(\boldsymbol{\theta}) < \infty$, then we have

$$\lim_{t \rightarrow \infty} (\theta_t^{j+1} - \theta_t^j) = \frac{2\pi}{n}, \quad \text{for all } j \in \{1, \dots, n\}. \quad (4.16)$$

Remark 4.8. Proposition 4.7 states that for finite-energy systems, the points $e^{i\theta_t^1}, \dots, e^{i\theta_t^n}$ eventually approach equal spacing around the circle — but, in contrast to Proposition 4.5, it is not true that a system with finite energy necessarily converges to a *static* equally-spaced configuration. For instance, let us consider the system defined by n equally-spaced copies of a single driver θ^1 :

$$\boldsymbol{\theta}_t = (\theta_t^1, \theta_t^1 + \frac{2\pi}{n}, \dots, \theta_t^1 + \frac{(n-1)2\pi}{n}), \quad t \geq 0,$$

so that $\phi^j(\boldsymbol{\theta}_t) = 0$ for every j and t . If $\theta^1 \in C_{\theta_0^1}([0, \infty), \mathcal{X}_1)$ has finite Dirichlet energy $E(\theta^1) < \infty$, then $\boldsymbol{\theta}$ has finite multiradial Dirichlet energy:

$$J(\boldsymbol{\theta}) = \frac{1}{2} \int_0^\infty \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi^j(\boldsymbol{\theta}_s) \right|^2 ds = \frac{1}{2} \int_0^\infty \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j \right|^2 ds = n E(\theta^1) < \infty.$$

However, this system may slowly spiral, for example if $\theta_t^1 = \log(t+1)$.

Proof of Proposition 4.7. As in the proof of Proposition 4.5, now under the assumption that $J(\boldsymbol{\theta}) < \infty$, we first prove (4.16), which will follow by proving the convergence (4.9) (in this case with unspecified rate). By the observation in Equation (4.12), it actually suffices to show that the quantity y_t (4.10) approaches zero as $t \rightarrow \infty$. To this end, we will first show that $J(\boldsymbol{\theta}) < \infty$ implies

$$D(\delta) := \{t \in [0, \infty) : y_t \geq \delta\} \quad \text{has finite Lebesgue measure for any } \delta > 0. \quad (4.17)$$

Thereafter, we will show that if $J(\boldsymbol{\theta}) < \infty$, then $D(\delta)$ is a bounded set for every $\delta > 0$. This is equivalent with $\overline{\lim}_{t \rightarrow \infty} y_t \leq \delta$ for every $\delta > 0$. As y is non-negative, by taking $\delta \rightarrow 0$ we may then conclude that $\lim_{t \rightarrow \infty} y_t = 0$.

Fix $\delta > 0$. On the one hand, Lemma 4.1 implies that if $y_t \geq \delta$, then $2 \max_{1 \leq j \leq n} |\phi^j(\boldsymbol{\theta}_t)| \geq \frac{n}{2} \delta$. On the other hand, the triangle inequality yields

$$2 \max_{1 \leq j \leq n} |\phi^j(\boldsymbol{\theta}_t)| \leq \max_{1 \leq j \leq n} \left| \frac{d}{dt} \theta_t^j - 2\phi^j(\boldsymbol{\theta}_t) \right| + \max_{1 \leq j \leq n} \left| \frac{d}{dt} \theta_t^j \right|. \quad (4.18)$$

Hence, if $y_t \geq \delta$, then at least one term on the righthand side of (4.18) is greater than or equal to $\frac{n}{4} \delta$. This allows us to bound the Lebesgue measure $\nu(D(\delta))$ of the set $D(\delta)$ as

$$\nu(D(\delta)) \leq \nu(S) + \nu(R), \quad (4.19)$$

$$\text{where } S := \left\{ t \in [0, \infty) : \max_{1 \leq j \leq n} \left| \frac{d}{dt} \theta_t^j - 2\phi^j(\boldsymbol{\theta}_t) \right| \geq \frac{n}{4} \delta \right\},$$

$$R := \left\{ t \in [0, \infty) : \max_{1 \leq j \leq n} \left| \frac{d}{dt} \theta_t^j \right| \geq \frac{n}{4} \delta \right\}.$$

To bound the righthand side of (4.19), we note that each term on the righthand side of (4.18) is square-integrable:

$$\int_0^\infty \left(\max_{1 \leq j \leq n} \left| \frac{d}{ds} \theta_s^j - 2\phi^j(\boldsymbol{\theta}_s) \right| \right)^2 ds \leq \int_0^\infty \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j - 2\phi^j(\boldsymbol{\theta}_s) \right|^2 ds = 2J(\boldsymbol{\theta}),$$

$$\int_0^\infty \left(\max_{1 \leq j \leq n} \left| \frac{d}{ds} \theta_s^j \right| \right)^2 ds \leq \int_0^\infty \sum_{j=1}^n \left| \frac{d}{ds} \theta_s^j \right|^2 ds = 2E(\boldsymbol{\theta}),$$

so that

$$\nu(D(\delta)) \leq \frac{32}{n^2 \delta^2} (J(\boldsymbol{\theta}) + E(\boldsymbol{\theta})) < \infty,$$

since $J(\boldsymbol{\theta}) < \infty$ by assumption and $E(\boldsymbol{\theta}) < \infty$ by Corollary 2.10. This verifies (4.17).

Next, suppose that $D(\delta)$ is unbounded. Then, there exists a sequence $(t_{(k)})_{k \in \mathbb{N}}$ such that $t_{(k)} \rightarrow \infty$ as $k \rightarrow \infty$ and $y_{t_{(k)}} \geq \delta$ for all k . Since $\nu(D(\delta/2)) < \infty$, we may assume (passing to a subsequence if necessary) that on each interval $(t_{(k)}, t_{(k+1)})$, the function y_t exits $D(\delta/2)$. Set

$$s_{(k)} := \max\{t \leq t_{(k)} : y_t = \delta/2\}, \quad k \in \mathbb{N}.$$

Since the set $D(\delta/2)$ has finite Lebesgue measure, the length of the intervals $(s_{(k)}, t_{(k)})$ approaches zero as $k \rightarrow \infty$, so for any $\epsilon > 0$ we can find an index k_ϵ such that

$$|t_{(k_\epsilon)} - s_{(k_\epsilon)}| < \epsilon.$$

By construction, for every $k \in \mathbb{N}$ and $i \in A_{t_{(k)}} := \arg \min_{1 \leq k \leq n} (\theta_{t_{(k)}}^{k+1} - \theta_{t_{(k)}}^k)$ we also have

$$\begin{aligned} \frac{\delta}{2} &\leq y_{t_{(k)}} - y_{s_{(k)}} = \min_{1 \leq j \leq n} (\theta_{s_{(k)}}^{j+1} - \theta_{s_{(k)}}^j) - \min_{1 \leq j \leq n} (\theta_{t_{(k)}}^{j+1} - \theta_{t_{(k)}}^j) \\ &\leq |\theta_{s_{(k)}}^{i+1} - \theta_{t_{(k)}}^{i+1}| + |\theta_{t_{(k)}}^i - \theta_{s_{(k)}}^i|. \end{aligned}$$

Hence, we see that there exists an index j such that $|\theta_{t_{(k_\epsilon)}}^j - \theta_{s_{(k_\epsilon)}}^j| \geq \delta/4$. We thus obtain (using also the Cauchy–Schwarz inequality)

$$\infty > E(\boldsymbol{\theta}) \geq \frac{1}{2} \int_{s_{(k_\epsilon)}}^{t_{(k_\epsilon)}} \left| \frac{d}{du} \theta_u^j \right|^2 du \geq \frac{|\theta_{t_{(k_\epsilon)}}^j - \theta_{s_{(k_\epsilon)}}^j|^2}{2|t_{(k_\epsilon)} - s_{(k_\epsilon)}|} \geq \frac{\delta^2}{32\epsilon} \xrightarrow{\epsilon \rightarrow 0} \infty,$$

which is a contradiction. This shows that $D(\delta)$ is bounded for every $\delta > 0$ and, in particular, that (4.16) holds. \square

Remark 4.9. In contrast with Proposition 4.5, finite-energy systems do not necessarily enjoy exponential rate of convergence to the equally-spaced configuration (4.16). In fact, it is even possible to construct systems of arbitrarily small energies with polynomial convergence rates. To demonstrate this, let us consider the case of two drivers, $n = 2$. Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a continuous L^2 -function, and suppose θ^1, θ^2 satisfy the differential equations

$$\begin{aligned} \frac{d}{dt} \theta_t^1 &= 2 \cot \left(\frac{\theta_t^1 - \theta_t^2}{2} \right) - f(t), \\ \frac{d}{dt} \theta_t^2 &= 2 \cot \left(\frac{\theta_t^2 - \theta_t^1}{2} \right), \quad t \geq 0, \end{aligned}$$

with initial configuration $\boldsymbol{\theta}_0 = (\theta_0^1, \theta_0^2) = (0, \pi)$. Then, $\boldsymbol{\theta} = (\theta^1, \theta^2) \in C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$ has multiradial Diriclet energy

$$J_T(\boldsymbol{\theta}) = \frac{1}{2} \int_0^T f(s)^2 ds,$$

and $u_t := (\theta_t^2 - \theta_t^1) - \pi$ satisfies the differential equation

$$\frac{d}{dt}u_t = 4 \cot\left(\frac{u_t + \pi}{2}\right) + f(t), \quad \text{with initial configuration } u_0 = 0.$$

Now, note that $d(\boldsymbol{\theta}_t) \geq |u_t|$. Next, consider a function $v: [0, \infty) \rightarrow \mathbb{R}$ satisfying the differential equation

$$\frac{d}{dt}v_t = -4v_t + f(t), \quad \text{with initial configuration } v_0 = 0.$$

Since $x \mapsto \cot(\frac{x+\pi}{2})$ is 1-Lipschitz on $[0, \frac{\pi}{2}]$, we have $u_t \geq v_t$ for all times before v_t exits the interval $[0, \frac{\pi}{2}]$. Choosing $f(t) = \frac{\varepsilon}{t+1}$ for $\varepsilon \in (0, \frac{\pi}{2})$ gives rise to the function

$$v_t = \varepsilon e^{-4t} \int_0^t \frac{e^{4s}}{s+1} ds \geq \varepsilon \frac{1 - e^{-4t}}{4(t+1)},$$

which never exits $[0, \frac{\pi}{2}]$, and thus, we find that

$$d(\boldsymbol{\theta}_t) \geq u_t \geq v_t \geq \varepsilon \frac{1 - e^{-4t}}{4(t+1)} = O(t^{-1}).$$

This gives a polynomial lower bound for the convergence rate to the equally-spaced configuration (4.16) for the system $\boldsymbol{\theta} = (\theta^1, \theta^2)$ having energy

$$J(\boldsymbol{\theta}) = \int_0^\infty \frac{\varepsilon^2}{2(t+1)^2} dt = \frac{\varepsilon^2}{2},$$

which can be made arbitrarily small by taking $\varepsilon \rightarrow 0$.

Finally, we treat the convergence rate for systems with (locally) finite energy. For zero-energy systems, it recovers the exponential rate of convergence of (4.9) from the proof of Proposition 4.5.

Proposition 4.10. *Consider a function $\boldsymbol{\theta} \in C_{\boldsymbol{\theta}_0}([0, \infty), \mathcal{X}_n)$. If $J_T(\boldsymbol{\theta}) < \infty$ for every $T \geq 0$, then we have*

$$d(\boldsymbol{\theta}_t) \leq (n-1)e^{-nt} \left(2\sqrt{2} \int_0^t e^{ns} \sqrt{\partial_s J_s(\boldsymbol{\theta})} ds + d(\boldsymbol{\theta}_0) \right), \quad t \geq 0, \quad (4.20)$$

$$\text{where } d(\boldsymbol{\theta}_t) := \max_{1 \leq j \leq n} \left| \frac{2\pi}{n} - (\theta_t^{j+1} - \theta_t^j) \right|.$$

Proof. We will show the slightly stronger claim for y_t (4.10) that

$$y_t \leq e^{-nt} \left(2\sqrt{2} \int_0^t e^{ns} \sqrt{\partial_s J_s(\boldsymbol{\theta})} ds + y_0 \right), \quad t \geq 0. \quad (4.21)$$

The asserted bound (4.20) then follows from (4.12).

Similarly as in (4.7), we deduce that for almost all times t , we have

$$\frac{d}{dt}y_t = \left(\frac{d}{dt}\theta_t^j - 2\phi^j(\boldsymbol{\theta}_t) \right) - \left(\frac{d}{dt}\theta_t^{j+1} - 2\phi^{j+1}(\boldsymbol{\theta}_t) \right) - 2(\phi^{j+1}(\boldsymbol{\theta}_t) - \phi^j(\boldsymbol{\theta}_t)) \quad (4.22)$$

for some $j \in A_t := \arg \min_{1 \leq k \leq n} (\theta_t^{k+1} - \theta_t^k)$. It follows from Lemma 4.1 that

$$2(\phi^{j+1}(\boldsymbol{\theta}_t) - \phi^j(\boldsymbol{\theta}_t)) \geq ny_t,$$

and

$$|\frac{d}{dt}\theta_t^k - 2\phi^k(\boldsymbol{\theta}_t)| \leq \left(\sum_{i=1}^n |\frac{d}{dt}\theta_t^i - 2\phi^i(\boldsymbol{\theta}_t)|^2 \right)^{1/2} = \sqrt{2}\sqrt{\partial_t J_t(\boldsymbol{\theta})}, \quad k \in \{1, \dots, n\}.$$

Plugging these back to (4.22) yields

$$\frac{d}{dt}y_t \leq -ny_t + 2\sqrt{2}\sqrt{\partial_t J_t(\boldsymbol{\theta})},$$

which implies (4.21) and concludes the proof. \square

References

- [ABKM20] Tom Alberts, Sung-Soo Byun, Nam-Gyu Kang, and Nikolai Makarov. Pole dynamics and an integral of motion for multiple SLE(0). Preprint in arXiv:2011.05714, 2020.
- [AGZ10] Greg W. Anderson, Alice Guionnet, and Ofer Zeitouni. *An Introduction to Random Matrices*. Number 118 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2010.
- [AIM09] Kari Astala, Tadeusz Iwaniec, and Gaven Martin. *Elliptic partial differential equations and quasiconformal mappings in the plane*, volume 48 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2009.
- [AP24] Osama Abuzaid and Eveliina Peltola. Large deviations of radial SLE₀₊. In preparation, 2024.
- [BB03] Michel Bauer and Denis Bernard. Conformal field theories of stochastic Loewner evolutions. *Comm. Math. Phys.*, 239(3):493–521, 2003.
- [BB04] Michel Bauer and Denis Bernard. CFTs of SLEs: the radial case. *Phys. Lett. B*, 583(3-4):324–330, 2004.
- [BBK05] Michel Bauer, Denis Bernard, and Kalle Kytölä. Multiple Schramm-Loewner evolutions and statistical mechanics martingales. *J. Stat. Phys.*, 120(5-6):1125–1163, 2005.
- [Bis19] Christopher J. Bishop. Weil-Petersson curves, conformal energies, β -numbers, and minimal surfaces. Preprint in <https://www.math.stonybrook.edu/~bishop/papers/wpbeta.pdf>, 2019.
- [BPW21] Vincent Beffara, Eveliina Peltola, and Hao Wu. On the uniqueness of global multiple SLEs. *Ann. Probab.*, 49(1):400–434, 2021.
- [Cal71] Francesco Calogero. Solution of the one-dimensional n -body problems with quadratic and/or inversely quadratic pair potentials. *J. Math. Phys.*, 12:419–436, 1971.
- [Car03] John L. Cardy. Stochastic Loewner evolution and Dyson’s circular ensembles. *J. Phys. A*, 36(24):L379–L386, 2003.

- [CDR06] John L. Cardy, Benjamin Doyon, and Valentina G. Riva. Identification of the stress-energy tensor through conformal restriction in SLE and related processes. *Comm. Math. Phys.*, 268(3):687–716, 2006.
- [CLM23] Andrew Campbell, Kyle Luh, and Vlad Margarint. Rate of convergence in multiple SLE using random matrix theory. Preprint in arXiv:2301.04722, 2023.
- [CM22] Chen and Vlad Margarint. Perturbations of multiple Schramm-Loewner evolution with two non-colliding Dyson Brownian motions. *Stochastic Process. Appl.*, 151:553–570, 2022.
- [DE02] Ioana Dumitriu and Alan Edelman. Matrix models for beta ensembles. *J. Math. Phys.*, 43(11):5830–5847, 2002.
- [dMS16] Andrea del Monaco and Sebastian Schleißinger. Multiple SLE and the complex Burgers equation. *Math. Nachr.*, 289(16):2007–2018, 2016.
- [Dub07] Julien Dubédat. Commutation relations for SLE. *Comm. Pure Appl. Math.*, 60(12):1792–1847, 2007.
- [Dub09] Julien Dubédat. SLE and the free field: partition functions and couplings. *J. Amer. Math. Soc.*, 22(4):995–1054, 2009.
- [Dub15] Julien Dubédat. SLE and Virasoro representations: localization. *Comm. Math. Phys.*, 336(2):695–760, 2015.
- [Dur83] Peter L. Duren. *Univalent Functions*. Grundlehren der mathematischen Wissenschaften. Springer-Verlag, New York, 1983.
- [Dys62] Freeman J. Dyson. A Brownian-motion model for the eigenvalues of a random matrix. *J. Math. Phys.*, 3:1191–1198, 1962.
- [DZ10] Amir Dembo and Ofer Zeitouni. *Large deviations techniques and applications*, volume 38 of *Stochastic Modelling and Applied Probability*. Springer-Verlag, Berlin, 2010.
- [EG02] Alexandre Eremenko and Andrei Gabrielov. Rational functions with real critical points and the B. and M. Shapiro conjecture in real enumerative geometry. *Ann. of Math.*, 155(1):105–129, 2002.
- [Eva10] Lawrence C. Evans. *Partial Differential Equations*. Graduate Studies in Mathematics. American Mathematical Society, 2 edition, 2010.
- [FK04] Roland Friedrich and Jussi Kalkkinen. On conformal field theory and stochastic Loewner evolution. *Nucl. Phys. B*, 687(3):279–302, 2004.
- [FS17] Peter K. Friz and Atul Shekhar. On the existence of SLE trace: finite energy drivers and non-constant κ . *Probab. Theory Related Fields*, 169(1-2):353–376, 2017.
- [FW84] Mark I. Freidlin and Alexander D. Wentzell. *Random Perturbations of Dynamical Systems*. Springer-Verlag, New York, 1984.
- [FW03] Roland Friedrich and Wendelin Werner. Conformal restriction, highest weight representations and SLE. *Comm. Math. Phys.*, 243(1):105–122, 2003.
- [FWY24] Yu Feng, Hao Wu, and Lu Yang. Multiple Ising interfaces in annulus and $2N$ -sided radial SLE. *Int. Math. Res. Not. (IMRN)*, 2024(6):5326–5372,

2024.

- [GK20] Vadim Gorin and Victor Kleptsyn. Universal objects of the infinite beta random matrix theory. Preprint in arXiv:2009.02006, 2020.
- [Gus23] Vladislav Guskov. A large deviation principle for the Schramm-Loewner evolution in the uniform topology. *Ann. Fenn. Math.*, 48(1):389–410, 2023.
- [GZ02] Alice Guionnet and Ofer Zeitouni. Large deviations asymptotics for spherical integrals. *J. Funct. Anal.*, 188(2):461–515, 2002.
- [HIM23] Ching-Peng Huang, Dominik Inauen, and Govind Menon. Motion by mean curvature and Dyson Brownian motion. *Electron. Commun. Probab.*, 28(34):1–10, 2023.
- [HK18] Ikkei Hotta and Makoto Katori. Hydrodynamic limit of multiple SLE. *J. Stat. Phys.*, 171:166–188, 2018.
- [HL21] Vivian O. Healey and Gregory F. Lawler. N-sided radial Schramm-Loewner evolution. *Probab. Theory Related Fields*, 181(1-3):451–488, 2021.
- [HM23] Vivian O. Healey and Govind Menon. Scaling limits of branching Loewner evolutions and the Dyson superprocess. Preprint in arXiv:2310.02858, 2023.
- [HS21] Ikkei Hotta and Sebastian Schleißinger. Limits of radial multiple SLE and a Burgers-Loewner differential equation. *J. Theoret. Probab.*, 34(2):755–783, 2021.
- [Kat16] Makoto Katori. *Bessel Processes, Schramm-Loewner Evolution, and the Dyson Model*, volume 11 of *SpringerBriefs in Mathematical Physics*. Springer, 2016.
- [Kem17] Antti Kemppainen. *Schramm-Loewner evolution*, volume 24 of *SpringerBriefs in Mathematical Physics*. Springer International Publishing, 2017.
- [KK21] Makoto Katori and Shinji Koshida. Three phases of multiple SLE driven by non-colliding Dyson’s Brownian motions. *J. Phys. A*, 54(32, Paper No. 325002):1–19, 2021.
- [KKP19] Alex Karrila, Kalle Kytölä, and Eveliina Peltola. Conformal blocks, q -combinatorics, and quantum group symmetry. *Ann. Inst. Henri Poincaré D*, 6(3):449–487, 2019.
- [KL07] Michael J. Kozdron and Gregory F. Lawler. The configurational measure on mutually avoiding SLE paths. *Fields Inst. Commun.*, 50:199–224, 2007.
- [KM13] Nam-Gyu Kang and Nikolai G. Makarov. Gaussian free field and conformal field theory. *Asterisque*, 353, 2013.
- [KP16] Kalle Kytölä and Eveliina Peltola. Pure partition functions of multiple SLEs. *Comm. Math. Phys.*, 346(1):237–292, 2016.
- [KS07] Maxim Kontsevich and Yuri Suhov. On Malliavin measures, SLE, and CFT. *P. Steklov I. Math.*, 258(1):100–146, 2007.
- [KT03] Makoto Katori and Tanemura. Noncolliding Brownian motions and Harish-Chandra formula. *Electron. Comm. Probab.*, 8:112–121, 2003.
- [Law09a] Gregory F. Lawler. Multifractal analysis of the reverse flow for the Schramm-Loewner evolution. In *Progress in Probability*, volume 61 of *Fractal Geometry*

and *Stochastics IV*, pages 73–107. Birkhäuser Basel, 2009.

- [Law09b] Gregory F. Lawler. Partition functions, loop measure, and versions of SLE. *J. Stat. Phys.*, 134(5-6):813–837, 2009.
- [Law13] Gregory F. Lawler. Continuity of radial and two-sided radial SLE at the terminal point. In *In the tradition of Ahlfors-Bers*, volume 590 of *Contemp. Math.*, pages 101–124. Amer. Math. Soc., Providence, RI, 2013.
- [Law24] Gregory F. Lawler. Schramm-Loewner evolution. Draft book, available at <http://www.math.uchicago.edu/~lawler/bookmaster.pdf>, 2024.
- [Lin05] Joan R. Lind. A sharp condition for the Loewner equation to generate slits. *Ann. Acad. Sci. Fenn. Math.*, 30:143–158, 2005.
- [LJV11] Gregory F. Lawler and Fredrik Johansson-Viklund. Optimal Hölder exponent for the SLE path. *Duke Math. J.*, 159(3):351–383, 2011.
- [LMR10] Joan Lind, Donald E. Marshall, and Steffen Rohde. Collisions and spirals of Loewner traces. *Duke Math. J.*, 154(3):527–573, 2010.
- [LSW03] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Conformal restriction: the chordal case. *J. Amer. Math. Soc.*, 16(4):917–955, 2003.
- [LSW04] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.*, 32(1B):939–995, 2004.
- [Mos75] Jürgen Moser. Three integrable Hamiltonian systems connected with isospectral deformations. *Adv. Math.*, 16(2):197–220, 1975.
- [MR05] Donald E. Marshall and Steffen Rohde. The Loewner differential equation and slit mappings. *J. Amer. Math. Soc.*, 18(4):763–778, 2005.
- [MS16a] Jason Miller and Scott Sheffield. Imaginary geometry I: interacting SLEs. *Probab. Theory Related Fields*, 164(3-4):553–705, 2016.
- [MS16b] Jason Miller and Scott Sheffield. Quantum Loewner evolution. *Duke Math. J.*, 165(17):3241–3378, 2016.
- [MS17] Jason Miller and Scott Sheffield. Imaginary geometry IV: interior rays, whole-plane reversibility, and space-filling trees. *Probab. Theory Related Fields*, 169(3-4):729–869, 2017.
- [Mul11] Gregory Muller. 2D locus configurations and the trigonometric Calogero-Moser system. *J. Nonlinear Math. Phys.*, 18(3):475–482, 2011.
- [Pel19] Eveliina Peltola. Towards a conformal field theory for Schramm-Loewner evolutions. *J. Math. Phys.*, 60(10):103305, 2019. Special issue (Proc. ICMP, Montreal, July 2018).
- [PW19] Eveliina Peltola and Hao Wu. Global and local multiple SLEs for $\kappa \leq 4$ and connection probabilities for level lines of GFF. *Comm. Math. Phys.*, 366(2):469–536, 2019.
- [PW24] Eveliina Peltola and Yilin Wang. Large deviations of multichordal SLE_{0+} , real rational functions, and zeta-regularized determinants of Laplacians. *J. Eur. Math. Soc.*, 26(2):469–535, 2024.

- [RS05] Steffen Rohde and Oded Schramm. Basic properties of SLE. *Ann. of Math.*, 161(2):883–924, 2005.
- [Sch00] Oded Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118(1):221–288, 2000.
- [Sch06] Oded Schramm. Conformally invariant scaling limits, an overview and a collection of problems. In *Proceedings of the ICM 2006, Madrid, Spain*, volume 1, pages 513–543. European Mathematical Society, 2006.
- [She16] Scott Sheffield. Conformal weldings of random surfaces: SLE and the quantum gravity zipper. *Ann. Probab.*, 44(5):3474–3545, 2016.
- [Smi06] Stanislav Smirnov. Towards conformal invariance of 2D lattice models. In *Proceedings of the ICM 2006, Madrid, Spain*, volume 2, pages 1421–1451. European Mathematical Society, 2006.
- [Sot00] Frank Sottile. Real Schubert calculus: polynomial systems and a conjecture of Shapiro and Shapiro. *Experiment. Math.*, 9(2):161–182, 2000.
- [SS09] Oded Schramm and Scott Sheffield. Contour lines of the two-dimensional discrete Gaussian free field. *Acta Math.*, 202(1):21–137, 2009.
- [Wan19a] Yilin Wang. The energy of a deterministic Loewner chain: reversibility and interpretation via SLE_{0+} . *J. Eur. Math. Soc.*, 21(7):1915–1941, 2019.
- [Wan19b] Yilin Wang. Equivalent descriptions of the Loewner energy. *Invent. Math.*, 218:573–621, 2019.
- [WW24] Yilin Wang and Hao Wu. Commutation relations for two-sided radial SLE. Preprint in arXiv:2405.07082, 2024.