

A BGK-type model for multi-component gas mixtures undergoing a bimolecular chemical reaction

G. Martalò¹, A. J. Soares², R. Travaglini^{3,1}

¹Department of Mathematical, Physical and Computer Sciences, University of Parma, Parco Area delle Scienze 53/A, 43124, Parma, Italy

²Centre of Mathematics of the University of Minho, Campus de Gualtar, 4710-057 Braga, Portugal

³ Istituto Nazionale di Alta Matematica “Francesco Severi”, Piazzale Aldo Moro 5, 00185, Roma, Italy

Abstract

Abstract. We propose a new kinetic BGK-type model for a mixture of four monatomic gases, undergoing a bimolecular and reversible chemical reaction. The elastic and reactive interactions are described separately by distinct relaxation terms and the mechanical operator is the sum of binary BGK contributions, one for each pair of interacting species. In this way, our model separately incorporates the effects of mechanical processes and chemical reactions. Additionally, it retains the effects of inter-species interactions which are proper of the mixture. The dependence of Maxwellian attractors on the main macroscopic fields is explicitly expressed by assuming that the exchange rates for momentum and energy of mechanical and chemical operators coincide with the ones of the corresponding Boltzmann terms. Under suitable hypotheses, the entropy dissipation is proven by means of an H-functional prescribing the relaxation of the distribution functions to the equilibrium.

Keywords: *Kinetic theory; BGK equations; Reactive mixtures; Exchange rates*

AMS subject classification 2020: 82C40; 76P05; 80A30; 35Q20

1. Introduction

Kinetic theory provides a natural framework to describe the evolution of rarefied gas mixtures. The classical description in the kinetic setting is given by Boltzmann equations that describe the evolution of the distribution functions. This formulation, though, results in being very complicated to deal with, due to the presence of integral collision operators with quadratic non-linearities. Indeed, the Boltzmann equation is well known in literature [19, 20, 25], but the investigation of its mathematical properties [14, 23], as well as the construction of efficient numerical schemes [16, 21, 33, 34], continues to motivate challenging problems that are still in progress.

To overcome the difficulties associated with the integral collision term, some alternative approaches have been proposed. The most famous one is based on the pioneering paper by Bhatnagar, Gross, and Krook [2]. In this work, the authors suggested replacing the

Boltzmann collision operator with a linear non-integral term, prescribing the relaxation towards an equilibrium Maxwellian attractor. Being the description formulated for a single gas, consistent generalizations to gas mixtures were highly desirable, especially concerning their application to physical processes. The first consistent model has been built up in [1], where the authors assume that, in the evolution equation of each component, the collision phenomenon is governed by a unique relaxation operator, which collects the effects of the interactions among all the constituents.

It is essential to point out that the extension of the BGK model to a gas mixture is not unique. In the BGK attractors, indeed, several free parameters intervene, and they can be suitably chosen to reproduce some basic properties of the original Boltzmann operators. For this reason, in the same spirit, different formulations can be found in the literature for inert and reactive mixtures of monatomic and polyatomic gases [9, 10, 11, 12, 13, 18, 24, 27, 28].

Among the others, we refer to [15], where the proposed model of BGK-type for inert mixtures mimics the same structure of the Boltzmann one, i.e. the collision operator is the sum of binary terms, each one accounting for the exchanges, occurring via mechanical interactions, of any pair of components. This formulation has the great advantage of distinguishing the different collision types and, consequently, allows the investigation of some multiple-scale regimes [5, 25], which are often present in plasma physics. For example, as described in [22], in the presence of heavy and light components in the mixture, momentum and energy exchanges proceed at different scales, according to the mass ratio. A typical example is given by a mixture of ions and electrons, as detailed in [31].

From this perspective, recently a class of mixed Boltzmann-BGK models has been presented [3, 4], where the accuracy of the Boltzmann operator is required to describe the dominant phenomenon, while the slow process is modeled by more manageable BGK terms.

In this manuscript, we extend the work [15] and propose a new model by considering a mixture of four monatomic gases undergoing elastic collisions and a bimolecular reversible chemical reaction [7]. As done for the inert mixture, the collision operator for each involved species is given by a sum of BGK terms, each term being distinguished by its specific set of parameters. Assuming that the exchange rates for momentum and energy of BGK mechanical and chemical operators coincide with the corresponding exchange rates of each Boltzmann integral operator, we can ensure the consistency of the model. The extension of the model to a reactive mixture represents a non-trivial problem because the rearrangement of masses and redistribution of energy among the reactive species lead to cumbersome computations.


Assuming the Maxwell molecule intermolecular potential, it is possible to explicitly calculate the production terms in the mechanical contributions for both Boltzmann and BGK formulations. This results in an exact relationship between the parameters of local attractors and the macroscopic fields of each species. Regarding chemical contributions, a reasonable input distribution function is required to compute the production term in the Boltzmann setting. To this aim, we follow the approach proposed in [29], where the distribution function is approximated by a proper perturbation of the Maxwellian function in the constituent reference frame. In other words, it results in a linearization with respect to species velocities and temperatures and corresponds to a not-so-far-away deviation from the mechanical equilibrium. Other choices are admissible, like the Grad-type approximation leading to 13-moment equations in the hydrodynamic limit [6].

After this introduction, the paper is organized as follows. In Section 2, after recalling the classical Boltzmann equation for a reactive mixture of monatomic gases, we propose a new BGK-type model, by introducing a kinetic operator as the sum of relaxation terms.

In Section 3, we explicitly compute the auxiliary parameters in the Maxwellian attractors of BGK terms. This can be done by assuming that the exchange rates for Boltzmann and BGK operators are the same for any type of interaction. An entropy estimate is proven in Section 4. In particular, we show the existence of an H -functional prescribing the relaxation of the distribution functions towards the mechanical and chemical equilibrium, under the hypotheses of species velocity equalization and the mass action law. Lastly, some concluding remarks are given in Section 5.

2. Kinetic equations

We consider a reactive mixture of four monatomic rarefied gases G_i , $i = 1, 2, 3, 4$, undergoing elastic mechanical collisions and a bimolecular reversible chemical reaction of type



Each i -component of the mixture is endowed with its own molecular mass m_i and internal energy E_i , such that $\Delta E = E_3 + E_4 - E_1 - E_2 > 0$, meaning that the forward reaction is endothermic.

Let $f_i = f_i(\mathbf{x}, \mathbf{v}, t)$ be the distribution function of the particles of the i -th component in the phase space, depending on space variable $\mathbf{x} \in \mathbb{R}^3$, microscopic velocity $\mathbf{v} \in \mathbb{R}^3$ and time $t \in \mathbb{R}_+$. We use the notation $\mathbf{f} = [f_1, f_2, f_3, f_4]$ for the vector collecting all the distribution functions. The time-space evolution of each function is governed by a proper kinetic equation

$$\frac{\partial f_i}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f_i = \mathcal{Q}_i[\mathbf{f}], \quad i = 1, 2, 3, 4, \quad (2)$$

where the operator \mathcal{Q}_i , at the right-hand side, takes into account both mechanical and chemical interactions.

For each component of the mixture, we define the main macroscopic fields as suitable moments of the distribution function. In particular, the number density n_i , mean velocity \mathbf{u}_i and temperature T_i for the i -th component are computed as

$$n_i = \int_{\mathbb{R}^3} f_i(\mathbf{v}) d\mathbf{v}, \quad \mathbf{u}_i = \frac{1}{n_i} \int_{\mathbb{R}^3} \mathbf{v} f_i(\mathbf{v}) d\mathbf{v}, \quad T_i = \frac{m_i}{3n_i} \int_{\mathbb{R}^3} |\mathbf{v} - \mathbf{u}_i|^2 f_i(\mathbf{v}) d\mathbf{v}. \quad (3)$$

Mixture observable quantities are defined as a combination of the corresponding species ones, as

$$\begin{aligned} n &= \sum_{i=1}^4 n_i, \quad \rho = \sum_{i=1}^4 \rho_i = \sum_{i=1}^4 m_i n_i, \quad \mathbf{u} = \frac{1}{\rho} \sum_{i=1}^4 \rho_i \mathbf{u}_i, \\ T &= \frac{1}{n} \left[\sum_{i=1}^4 n_i T_i + \frac{1}{3} \sum_{i=1}^4 m_i n_i (u_i^2 - u^2) \right]. \end{aligned} \quad (4)$$

2.1. Boltzmann formulation

The more relevant equation in Kinetic Theory is the one based on the Boltzmann formulation. The collision operator is split into a sum of two contributions

$$\mathcal{Q}_i^{Bol} = \tilde{\mathcal{Q}}_i^{Bol} + \hat{\mathcal{Q}}_i^{Bol}, \quad (5)$$

where $\tilde{\mathcal{Q}}_i^{Bol}$ takes into account the mechanical encounters, whereas $\hat{\mathcal{Q}}_i^{Bol}$ considers the effects of the chemical reaction. As usual in Boltzmann formulation for mixtures [19, 20, 25],

the mechanical contribution is decomposed in a sum of binary terms, each one describing the elastic interactions of the pair of i and j components, namely

$$\tilde{\mathcal{Q}}_i^{Bol} = \sum_{j=1}^4 \tilde{\mathcal{Q}}_{ij}^{Bol}. \quad (6)$$

Both the mechanical and chemical terms are non-linear operators of integral type. More precisely, the mechanical terms are given by

$$\tilde{\mathcal{Q}}_{ij}^{Bol} = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} g \tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}') [f_i(\mathbf{v}') f_j(\mathbf{w}') - f_i(\mathbf{v}) f_j(\mathbf{w})] d\mathbf{w} d\hat{\Omega}', \quad i, j = 1, 2, 3, 4, \quad (7)$$

where $\mathbf{g} = \mathbf{v} - \mathbf{w} = g\hat{\Omega}$ denotes the relative velocity of the colliding pair, with g being its modulus and $\hat{\Omega}$ the corresponding unit vector. Pre- and post-collision velocities are related by the conservation laws of mass, momentum, and total energy during the collisions, that is

$$\begin{aligned} m_i + m_j &= m_i + m_j \\ m_i \mathbf{v} + m_j \mathbf{w} &= m_i \mathbf{v}' + m_j \mathbf{w}' \\ \frac{1}{2} m_i v^2 + E_i + \frac{1}{2} m_j w^2 + E_j &= \frac{1}{2} m_i (v')^2 + E_i + \frac{1}{2} m_j (w')^2 + E_j. \end{aligned} \quad (8)$$

The quantity $\tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}')$ in (7) denotes the elastic cross-section [19], and takes into account the inter-molecular potential, and it is assumed to satisfy proper symmetry conditions

$$\tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}') = \tilde{\sigma}_{ij}(g, -\hat{\Omega} \cdot \hat{\Omega}') = \tilde{\sigma}_{ji}(g, \hat{\Omega} \cdot \hat{\Omega}'). \quad (9)$$

In this paper, we assume that the mixture particle interact according to the Maxwell molecule model [30], and therefore

$$\tilde{\lambda}_{ij}^{(0)} = \int_{\mathbb{S}^2} g \tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}') d\hat{\Omega}' = \text{constant}. \quad (10)$$

Concerning chemical interactions, they can be described in an analogous way, by means of a Boltzmann-like operator. In particular, the chemical contribution to the kinetic equation of the i -th component is given by [32]

$$\hat{\mathcal{Q}}_i^{Bol} = \iint_{\mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{ij}^{hk}) g \hat{\sigma}_{ijhk}(g, \hat{\Omega} \cdot \hat{\Omega}') \left[\left(\frac{\mu_{ij}}{\mu_{hk}} \right)^3 f_h(\mathbf{v}') f_k(\mathbf{w}') - f_i(\mathbf{v}) f_j(\mathbf{w}) \right] d\mathbf{w} d\hat{\Omega}', \quad (11)$$

where H is the Heaviside function and $\mu_{ij} = m_i m_j / (m_i + m_j)$ denotes the reduced mass of the colliding pair. The quantity $\delta_{ij}^{hk} = 2\Delta E / \mu_{ij}$ provides a proper threshold for the modulus of relative velocity to be overcome in order to guarantee the occurrence of the exothermic reaction.

As above, pre- and post-interaction velocities fulfill the conservation

$$\begin{aligned} m_i + m_j &= m_h + m_k \\ m_i \mathbf{v} + m_j \mathbf{w} &= m_h \mathbf{v}' + m_k \mathbf{w}' \\ \frac{1}{2} m_i v^2 + E_i + \frac{1}{2} m_j w^2 + E_j &= \frac{1}{2} m_h (v')^2 + E_h + \frac{1}{2} m_k (w')^2 + E_k, \end{aligned} \quad (12)$$

Moreover, the quantity $\hat{\sigma}_{ijhk}(g, \hat{\Omega} \cdot \hat{\Omega}')$ denotes the reactive cross-section. Here, we assume a chemical potential of Maxwell type also for chemically reacting interactions,

$$\hat{\lambda}_{ijhk}^{(0)} = \int_{\mathbb{S}^2} g \hat{\sigma}_{ijhk}(g, \hat{\Omega} \cdot \hat{\Omega}') d\hat{\Omega}' = \text{constant}. \quad (13)$$

and symmetry relations similar to those in (9) hold for reactive interactions.

As well known, collision equilibria of the reactive mixture entail both mechanical and chemical equilibrium. They are defined by distribution functions of Maxwell type, for which collision operators (7) and (11) simultaneously vanish, that is

$$f_i^M(\mathbf{v}) = n_i \mathcal{M}(\mathbf{v}; \mathbf{u}, T) = n_i \left(\frac{m_i}{2\pi T} \right)^{\frac{3}{2}} \exp \left[-\frac{m_i}{2T} (\mathbf{v} - \mathbf{u})^2 \right], \quad (14)$$

where, in addition, species number densities fulfill the mass action law

$$\frac{n_1 n_2}{n_3 n_4} = \left(\frac{m_1 m_2}{m_3 m_4} \right)^{\frac{3}{2}} \exp \left(\frac{\Delta E}{T} \right). \quad (15)$$

Mechanically, the equilibrium states are described by equating species velocities and temperatures, so that all constituents share a common velocity and a common temperature, which are those of the mixture. The mass action law (15) gives a characterization of such states from a chemical point of view and represents the chemical equilibrium condition of the chemical reaction (1).

In addition, in the space homogeneous conditions, it is possible to prove the global stability of such equilibria by showing that the function

$$\mathcal{H}[\mathbf{f}] = \sum_{i=1}^4 \int_{\mathbb{R}^3} f_i(\mathbf{v}) \log \left(\frac{f_i(\mathbf{v})}{m_i^3} \right) d\mathbf{v} \quad (16)$$

is a Lyapunov functional for the kinetic system (2) with the corresponding collisional operator Q_i^{Bol} , thus providing, consequently, an entropy estimation.

2.2. A BGK model

Due to the complexity of the integral form of the Boltzmann operators, some alternative formulations are highly desirable. The most common approach is the so called BGK model, which replaces the Boltzmann terms by simpler relaxation operators. These operators are defined in terms of suitable relaxation attractors involving free parameters that must be specified to ensure the mathematical and physical consistency of the model.

Despite their simplicity, these operators, being non-integral and linear with respect to the distribution functions, effectively reproduce the main results of the Boltzmann description, including the conservation laws of mass, momentum, and total energy, correct equilibria, and entropy inequality.

It is important to note that the BGK model for a single gas can be extended to mixtures in several ways. In particular, the usual conservation laws provide few constraints when setting the free auxiliary parameters in the relaxation attractors, and additional conditions are required.

Among the models proposed in the literature, we consider, in this paper, the one proposed in [15] for inert mixtures. Here, we extend this model to a reactive mixture in the presence of the chemical reaction given in (1). The aim is to replicate the same structure of the Boltzmann collision operator, as a sum of many terms, one for each type of interaction. In this way, the resulting model is able to capture a broader range of mixture effects. Accordingly, the BGK operator can be decomposed as

$$Q_i^{BGK} = \tilde{Q}_i^{BGK} + \hat{Q}_i^{BGK} = \sum_{j=1}^4 \tilde{Q}_{ij}^{BGK} + \hat{Q}_i^{BGK}. \quad (17)$$

with mechanical terms \tilde{Q}_{ij}^{BGK} and chemical terms \hat{Q}_i^{BGK} , given by

$$\tilde{Q}_{ij}^{BGK} = \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i), \quad (18)$$

and

$$\hat{Q}_i^{BGK} = \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i), \quad (19)$$

respectively, where $\tilde{\nu}_{ij}$ represents the frequency of mechanical collisions between i -th and j -th components, and $\hat{\nu}_{ij}^{hk}$ the frequency of reactive collisions obeying the chemical law (1). Moreover,

$$\mathcal{M}_{ij} = \tilde{n}_{ij} \mathbf{M}_i(\mathbf{v}; \tilde{\mathbf{u}}_{ij}, \tilde{T}_{ij}) \quad \text{and} \quad \mathcal{M}_i = \hat{n}_i \mathbf{M}_i(\mathbf{v}; \hat{\mathbf{u}}_i, \hat{T}_i)$$

are attractors depending on some fictitious parameters to be determined, namely \tilde{n}_{ij} , $\tilde{\mathbf{u}}_{ij}$, \tilde{T}_{ij} , and \hat{n}_i , $\hat{\mathbf{u}}_i$, \hat{T}_i .

3. Computation of auxiliary parameters

In order to determine the auxiliary parameters in the Maxwellian attractors, we follow the procedure proposed in [15]. Hence, we assume that the exchange rates of momentum and energy for Boltzmann and BGK formulations coincide for any type of interaction.

For this purpose, we define

$$\begin{aligned} \tilde{P}_{ij}^{(k)} &= \int_{\mathbb{R}^3} \varphi_i^{(k)}(\mathbf{v}) \tilde{Q}_{ij}^{Bol}(\mathbf{v}) d\mathbf{v} \quad , \quad \hat{P}_i^{(k)} = \int_{\mathbb{R}^3} \varphi_i^{(k)}(\mathbf{v}) \hat{Q}_i^{Bol}(\mathbf{v}) d\mathbf{v} \quad , \\ \tilde{\Pi}_{ij}^{(k)} &= \int_{\mathbb{R}^3} \varphi_i^{(k)}(\mathbf{v}) \tilde{Q}_{ij}^{BGK}(\mathbf{v}) d\mathbf{v} \quad , \quad \hat{\Pi}_i^{(k)} = \int_{\mathbb{R}^3} \varphi_i^{(k)}(\mathbf{v}) \hat{Q}_i^{BGK}(\mathbf{v}) d\mathbf{v} \quad , \end{aligned} \quad (20)$$

where

$$\varphi_i^{(0)}(\mathbf{v}) = m_i \quad , \quad \varphi_i^{(1)}(\mathbf{v}) = m_i \mathbf{v} \quad , \quad \varphi_i^{(2)}(\mathbf{v}) = \frac{1}{2} m_i v^2 + E_i \quad . \quad (21)$$

3.1. Parameters in mechanical terms

As anticipated above, we assume that

$$\tilde{\Pi}_{ij}^{(k)} = \tilde{P}_{ij}^{(k)} \quad , \quad i, j = 1, 2, 3, 4 \quad , \quad k = 0, 1, 2 \quad . \quad (22)$$

To compute the production terms coming from Boltzmann formulation, we consider the following weak form of Boltzmann operators

$$\tilde{P}_{ij}^{(k)} = \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} g \tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}') \left(\varphi_i^{(k)}(\mathbf{v}') - \varphi_i^{(k)}(\mathbf{v}) \right) f_i(\mathbf{v}) f_j(\mathbf{w}) d\mathbf{v} d\mathbf{w} d\hat{\Omega}' \quad , \quad (23)$$

and, under the hypothesis of Maxwell molecule potential, we can compute explicitly

$$\begin{aligned} \tilde{P}_{ij}^{(0)} &= 0 \\ \tilde{P}_{ij}^{(1)} &= \tilde{\lambda}_{ij}^{(0)} \mu_{ij} n_i n_j (\mathbf{u}_j - \mathbf{u}_i) \\ \tilde{P}_{ij}^{(2)} &= \tilde{\lambda}_{ij}^{(0)} \alpha_{ij} \alpha_{ji} n_i n_j [(m_i \mathbf{u}_i + m_j \mathbf{u}_j) \cdot (\mathbf{u}_j - \mathbf{u}_i) + 3(T_j - T_i)] \quad , \end{aligned} \quad (24)$$

where we notice that

$$\int_{\mathbb{S}^2} \hat{\Omega}' \tilde{\sigma}_{ij}(g, \hat{\Omega} \cdot \hat{\Omega}') d\hat{\Omega}' = \mathbf{0} \quad (25)$$

by parity arguments, and we made use of the relation between pre- and post-collision velocities

$$\begin{aligned}\mathbf{v}' &= \alpha_{ij}\mathbf{v} + \alpha_{ji}\mathbf{w} + \alpha_{ji}g'\widehat{\Omega}' \\ \mathbf{w}' &= \alpha_{ij}\mathbf{v} + \alpha_{ji}\mathbf{w} - \alpha_{ij}g'\widehat{\Omega}',\end{aligned}\tag{26}$$

$\alpha_{ij} = m_i/(m_i + m_j)$ denoting the mass ratio.

For what concerns the production terms from the BGK model, we can easily obtain that

$$\begin{aligned}\widetilde{\Pi}_{ij}^{(0)} &= \widetilde{\nu}_{ij}m_i(\widetilde{n}_{ij} - n_i) \\ \widetilde{\Pi}_{ij}^{(1)} &= \widetilde{\nu}_{ij}m_i(\widetilde{n}_{ij}\widetilde{\mathbf{u}}_{ij} - n_i\mathbf{u}_i) \\ \widetilde{\Pi}_{ij}^{(2)} &= \widetilde{\nu}_{ij}\left[\frac{1}{2}m_i(\widetilde{n}_{ij}\widetilde{\mathbf{u}}_{ij}^2 - n_i\mathbf{u}_i^2) + \frac{3}{2}(\widetilde{n}_{ij}\widetilde{T}_{ij} - n_iT_i) + E_i(\widetilde{n}_{ij} - n_i)\right],\end{aligned}\tag{27}$$

and hence, by assumption (22), we get

$$\begin{aligned}\widetilde{n}_{ij} &= n_i \\ \widetilde{\mathbf{u}}_{ij} &= (1 - a_{ij})\mathbf{u}_i + a_{ij}\mathbf{u}_j \\ \widetilde{T}_{ij} &= (1 - b_{ij})T_i + b_{ij}T_j + \gamma_{ij}|\mathbf{u}_i - \mathbf{u}_j|^2\end{aligned}\tag{28}$$

with

$$a_{ij} = \frac{\widetilde{\lambda}_{ij}^{(0)}}{\widetilde{\nu}_{ij}}\alpha_{ji}n_j, \quad b_{ij} = 2a_{ij}\alpha_{ij}, \quad \gamma_{ij} = \frac{1}{3}m_ia_{ij}(2\alpha_{ji} - a_{ij}).\tag{29}$$

The positivity of auxiliary temperatures is guaranteed under the condition $\widetilde{\nu}_{ij} \geq \frac{1}{2}\widetilde{\lambda}_{ij}^{(0)}n_j$.

Further details can be found in [15].

3.2. Parameters in chemical terms

As done for mechanical contributions, we impose that the exchange rates of mass, momentum, and total energy due to chemical interactions are the same for Boltzmann and BGK operators, i.e.

$$\widehat{\Pi}_i^{(k)} = \widehat{P}_i^{(k)}, \quad i = 1, 2, 3, 4, \quad k = 0, 1, 2.\tag{30}$$

Production terms in the BGK model can be computed explicitly, while the corresponding contributions in the Boltzmann formulation are approximated by considering a quasi-equilibrium configuration [29]. We consider

$$f_i(\mathbf{v}) \simeq n_i \left(\frac{m_i}{2\pi T}\right)^{3/2} \exp\left[-\frac{m_i}{2T}|\mathbf{v} - \mathbf{u}|^2\right] \mathcal{E}_i(\mathbf{v}), \quad i = 1, 2, 3, 4,\tag{31}$$

where

$$\mathcal{E}_i(\mathbf{v}) = \mathcal{A}_i + \mathcal{B}_i(\mathbf{v} - \mathbf{u}) \cdot (\mathbf{u}_i - \mathbf{u}) + \mathcal{C}_i|\mathbf{v} - \mathbf{u}|^2\tag{32}$$

and

$$\mathcal{A}_i = 1 - \frac{3}{2}\frac{T_i - T}{T}, \quad \mathcal{B}_i = \frac{m_i}{2T}, \quad \mathcal{C}_i = \frac{m_i}{2T}\frac{T_i - T}{T}.\tag{33}$$

We get

$$\begin{aligned}\widehat{n}_i &= n_i + \frac{\widehat{P}_i^{(0)}}{m_i\widehat{\nu}_{ij}^{hk}} \\ \widehat{\mathbf{u}}_i &= \frac{1}{\widehat{n}_i} \left(n_i\mathbf{u}_i + \frac{\widehat{P}_i^{(1)}}{m_i\widehat{\nu}_{ij}^{hk}} \right) \\ \widehat{T}_i &= \frac{1}{\widehat{n}_i} \left\{ n_iT_i + \frac{2}{3} \left[\frac{\widehat{P}_i^{(2)}}{\widehat{\nu}_{ij}^{hk}} - \frac{1}{2}m_i(\widehat{n}_i\widehat{\mathbf{u}}_i^2 - n_i\mathbf{u}_i^2) - E_i(\widehat{n}_i - n_i) \right] \right\},\end{aligned}\tag{34}$$

where production terms $\widehat{P}_i^{(0)}$, $\widehat{P}_i^{(1)}$, $\widehat{P}_i^{(2)}$ are computed in detail in Appendix A.

We can notice that

$$\frac{\widehat{P}_1^{(0)}}{m_1} = \frac{\widehat{P}_2^{(0)}}{m_2} = -\frac{\widehat{P}_3^{(0)}}{m_3} = -\frac{\widehat{P}_4^{(0)}}{m_4} \quad (35)$$

and hence the partial and total conservation of mass follows; moreover, also global momentum and energy preservation are correctly reproduced, i.e. it holds

$$\begin{aligned} \sum_{i=1}^4 \widehat{P}_i^{(1)} &= \mathbf{0} \\ \sum_{i=1} \widehat{P}_i^{(2)} + \lambda_i \frac{\widehat{P}_i^{(0)}}{m_i} \Delta E &= 0, \text{ for any fixed } i = 1, 2, 3, 4, \end{aligned} \quad (36)$$

where $\lambda_1 = \lambda_2 = 1 = -\lambda_3 = -\lambda_4$ are the stoichiometric coefficients.

Since $\mathcal{M}_i = \widehat{n}_i \mathbf{M}_i(\mathbf{v}; \widehat{\mathbf{u}}_i, \widehat{T}_i)$ is a distribution function for each species $i = 1, 2, 3, 4$, the Maxwellian must be positive and measurable, and hence we must require that

$$\widehat{n}_i > 0 \quad \text{and} \quad \widehat{T}_i > 0. \quad (37)$$

The condition (37) is fulfilled at least in the quasi-equilibrium configuration, since the production terms \widehat{P}_i deduced from Boltzmann operators are sufficiently small.

4. Entropy properties

For what concerns the entropy growth, regrettably, proving rigorously an \mathcal{H} -theorem for the model proposed above seems unfeasible. An entropy estimate can be established only under specific assumptions, more precisely, assuming the equalization of fictitious species velocities and temperatures in the chemical BGK contributions, i.e.

$$\widehat{\mathbf{u}}_i = \widehat{\mathbf{u}}, \quad \widehat{T}_i = \widehat{T}. \quad (38)$$

Moreover, we assume that auxiliary densities fulfill the following algebraic constraint accounting for the mass action law (15),

$$\frac{\widehat{n}_1 \widehat{n}_2}{\widehat{n}_3 \widehat{n}_4} = \left(\frac{m_1 m_2}{m_3 m_4} \right)^{\frac{3}{2}} \exp \left(\frac{\Delta E}{\widehat{T}} \right). \quad (39)$$

Note that such assumptions (38) and (39) are distinctive hypotheses when proving an \mathcal{H} -theorem for a BGK model for reactive mixtures, as can be seen, for instance, in the model proposed in [26] and then reformulated in [8], or in the papers [17] and [18]. In our work, thus, the general shape of attractors allows to reproduce the correct conservation laws, whereas we need the extra assumptions (38) and (39) to prove an \mathcal{H} -theorem. More in detail, it is possible to prove that the \mathcal{H} -functional for chemically reactive mixtures, already introduced in (16) for the Boltzmann system, that is

$$\mathcal{H}[f_i] = \sum_{i=1}^4 \int_{\mathbb{R}^3} f_i(\mathbf{v}) \log \left(\frac{f_i(\mathbf{v})}{m_i^3} \right) d\mathbf{v}, \quad (40)$$

is a Lyapunov functional for our BGK model proposed in Subsection 2.2.

The following main result holds.

Theorem 4.1 (*\mathcal{H} -theorem*) *Let us assume conditions (38) and (39) on the auxiliary parameters of chemical terms. Under space homogeneous conditions, for all measurable distribution functions $f_i \geq 0$, $i = 1, 2, 3, 4$, we have that*

$$\frac{d\mathcal{H}}{dt} \leq 0, \quad \text{for all } t \geq 0. \quad (41)$$

Proof. Under space homogeneous conditions, one has

$$\begin{aligned} \frac{d\mathcal{H}}{dt} &= \sum_{i=1}^4 \int_{\mathbb{R}^3} \frac{\partial f_i}{\partial t} \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \sum_{i=1}^4 \int_{\mathbb{R}^3} f_i \frac{m_i^3}{f_i} \frac{1}{m_i^3} \frac{\partial f_i}{\partial t} d\mathbf{v} \\ &= \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} \frac{d}{dt} \int_{\mathbb{R}^3} (\mathcal{M}_{ij} - f_i) d\mathbf{v} \\ &\quad + \sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \frac{d}{dt} \sum_{i=1}^4 \hat{\nu}_{ij}^{hk} \int_{\mathbb{R}^3} (\mathcal{M}_i - f_i) d\mathbf{v}, \end{aligned} \quad (42)$$

where, taking into account the assumptions on the auxiliary parameters, the chemical attractor is given by

$$\mathcal{M}_i = \hat{n}_i \mathbf{M}_i(\mathbf{v}; \hat{\mathbf{u}}, \hat{T}) = \hat{n}_i \left(\frac{m_i}{2\pi\hat{T}} \right)^{\frac{3}{2}} \exp \left[-\frac{m_i}{2\hat{T}} (\mathbf{v} - \hat{\mathbf{u}})^2 \right]. \quad (43)$$

Concerning the attractors, we observe that

$$\int_{\mathbb{R}^3} (\mathcal{M}_{ij} - f_i) d\mathbf{v} = \tilde{n}_{ij} - n_i = n_i - n_i = 0 \quad (44)$$

and

$$\sum_{i=1}^4 \hat{\nu}_{ij}^{hk} \int_{\mathbb{R}^3} (\mathcal{M}_i - f_i) d\mathbf{v} = \sum_{i=1}^4 \hat{\nu}_{ij}^{hk} (\hat{n}_i - n_i) = \sum_{i=1}^4 \frac{\hat{P}_i^{(0)}}{m_i} = 0. \quad (45)$$

Therefore, coming back to (42), we obtain

$$\begin{aligned} \frac{d\mathcal{H}}{dt} &= \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} \\ &= \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v} \\ &\quad + \sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{\mathcal{M}_i}{m_i^3} \right) d\mathbf{v} \end{aligned} \quad (46)$$

Now, we treat separately the three contributions on the right-hand side of expression (46).

The first contribution can be dealt with as done in [15] and we show that

$$\sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} \leq 0. \quad (47)$$

More precisely, first, we rewrite

$$\begin{aligned} & \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} \\ &= \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log(f_i) d\mathbf{v} - 3 \sum_{i=1}^4 \sum_{j=1}^4 \log(m_i) \tilde{\nu}_{ij} \int_{\mathbb{R}^3} (\mathcal{M}_{ij} - f_i) d\mathbf{v}. \end{aligned}$$

Here, the second term vanishes, again due to (44), while the first one can be estimated by means of the well-known inequality

$$(y - x) \log x \leq y(\log y - 1) - x(\log x - 1). \quad (48)$$

One has

$$\begin{aligned} & \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log(f_i) d\mathbf{v} \\ & \leq \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} \left[\int_{\mathbb{R}^3} \mathcal{M}_{ij} (\log \mathcal{M}_{ij} - 1) d\mathbf{v} - \int_{\mathbb{R}^3} f_i (\log f_i - 1) d\mathbf{v} \right] \\ & = \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} \left[\int_{\mathbb{R}^3} \mathcal{M}_{ij} \log \mathcal{M}_{ij} d\mathbf{v} - \int_{\mathbb{R}^3} f_i \log f_i d\mathbf{v} + \int_{\mathbb{R}^3} (f_i - \mathcal{M}_{ij}) d\mathbf{v} \right]. \quad (49) \end{aligned}$$

As above, the last term vanishes, since f_i and \mathcal{M}_{ij} share the same number density, due to (44). Using classical results for a single component gas, the second contribution can be estimated as follows [15],

$$\int_{\mathbb{R}^3} f_i \log f_i d\mathbf{v} \geq \int_{\mathbb{R}^3} \mathcal{M}_{ii} \log \mathcal{M}_{ii} d\mathbf{v}. \quad (50)$$

We can compute explicitly each component in the first term of (49), obtaining

$$\int_{\mathbb{R}^3} \mathcal{M}_{ij} \log \mathcal{M}_{ij} d\mathbf{v} = n_i \left[\log(n_i) + \frac{3}{2} \log \left(\frac{m_i}{2\pi \tilde{T}_{ij}} \right) - \frac{3}{2} \right]. \quad (51)$$

By summing up,

$$\begin{aligned} & \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log(f_i) d\mathbf{v} \\ & \leq \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} \left[\int_{\mathbb{R}^3} \mathcal{M}_{ij} \log \mathcal{M}_{ij} d\mathbf{v} - \int_{\mathbb{R}^3} \mathcal{M}_{ii} \log \mathcal{M}_{ii} d\mathbf{v} \right] \\ & = -\frac{3}{2} \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} n_i \log \left(\frac{\tilde{T}_{ij}}{T_i} \right). \quad (52) \end{aligned}$$

From conditions (28) characterizing the auxiliary parameters in mechanical terms, we get

$$\tilde{T}_{ij} \geq (1 - b_{ij})T_i + b_{ij}T_j \quad (53)$$

and by recalling Jensen's inequality

$$\log [(1 - b)x + by] \geq (1 - b) \log x + b \log y, \quad (54)$$

we obtain

$$\log \left(\frac{\tilde{T}_{ij}}{T_i} \right) \geq b_{ij} \log \left(\frac{T_j}{T_i} \right). \quad (55)$$

Consequently, for the first contribution in (46), we get

$$\begin{aligned} & \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} \\ & \leq -\frac{3}{2} \sum_{i=1}^4 \sum_{j=1}^4 \tilde{\nu}_{ij} n_i b_{ij} \log \left(\frac{T_j}{T_i} \right) \\ & = -\frac{3}{2} \sum_{i=1}^4 \sum_{\substack{j=1 \\ j>i}}^4 \tilde{\nu}_{ij} n_i b_{ij} \left[\log \left(\frac{T_j}{T_i} \right) + \log \left(\frac{T_i}{T_j} \right) \right] = 0, \end{aligned} \quad (56)$$

since $\tilde{\nu}_{ij} n_i b_{ij} = \tilde{\nu}_{ji} n_j b_{ji}$.

Coming back to (46), and considering the second contribution, it is easy to prove that

$$\int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v} \leq 0. \quad (57)$$

More precisely, we have

$$\int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v} = \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} \mathcal{M}_i \left(1 - \frac{f_i}{\mathcal{M}_i} \right) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v} \leq 0, \quad (58)$$

by means of usual convexity arguments of the function $(1-x) \log x$.

Finally, considering the third contribution in (46), and we show that it vanishes, that is

$$\sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{\mathcal{M}_i}{m_i^3} \right) d\mathbf{v} = 0. \quad (59)$$

In fact, we have

$$\begin{aligned} & \sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{\mathcal{M}_i}{m_i^3} \right) d\mathbf{v} \\ & = \sum_{i=1}^4 \hat{\nu}_{ij}^{hk} \left[\log(\hat{n}_i) - \frac{3}{2} \log(m_i) - \frac{3}{2} \log(2\pi\hat{T}) \right] \int_{\mathbb{R}^3} (\mathcal{M}_i - f_i) d\mathbf{v} \\ & \quad - \sum_{i=1}^4 \hat{\nu}_{ij}^{hk} \frac{m_i}{2\hat{T}} \int_{\mathbb{R}^3} |\mathbf{v} - \hat{\mathbf{u}}|^2 (\mathcal{M}_i - f_i) d\mathbf{v}. \end{aligned} \quad (60)$$

Concerning the first contribution on the right-hand side of (60), we notice that

$$\begin{aligned}
& \sum_{i=1}^4 \widehat{\nu}_{ij}^{hk} \left[\log(\widehat{n}_i) - \frac{3}{2} \log(m_i) - \frac{3}{2} \log(2\pi\widehat{T}) \right] \int_{\mathbb{R}^3} (\mathcal{M}_i - f_i) d\mathbf{v} \\
&= \sum_{i=1}^4 \widehat{\nu}_{ij}^{hk} \left[\log(\widehat{n}_i) - \frac{3}{2} \log(m_i) - \frac{3}{2} \log(2\pi\widehat{T}) \right] (\widehat{n}_i - n_i) \\
&= \sum_{i=1}^4 \frac{\widehat{P}_i^{(0)}}{m_i} \left[\log(\widehat{n}_i) - \frac{3}{2} \log(m_i) - \frac{3}{2} \log(2\pi\widehat{T}) \right] \\
&= \frac{\widehat{P}_1^{(0)}}{m_1} \sum_{i=1}^4 \lambda_i \left[\log(\widehat{n}_i) - \frac{3}{2} \log(m_i) - \frac{3}{2} \log(2\pi\widehat{T}) \right] \\
&= \frac{\widehat{P}_1^{(0)}}{m_1} \log \left[\frac{\widehat{n}_1 \widehat{n}_2}{\widehat{n}_3 \widehat{n}_4} \left(\frac{m_3 m_4}{m_1 m_2} \right)^{3/2} \right].
\end{aligned} \tag{61}$$

For what concerns the second sum in (60), by expanding the squared term and bearing in mind the conservation laws of total momentum and energy, it reduces to

$$\begin{aligned}
& - \sum_{i=1}^4 \widehat{\nu}_{ij}^{hk} \frac{m_i}{2\widehat{T}} \int_{\mathbb{R}^3} |\mathbf{v} - \widehat{\mathbf{u}}|^2 (\mathcal{M}_i - f_i) d\mathbf{v} \\
&= \sum_{i=1}^4 \widehat{\nu}_{ij}^{hk} \frac{E_i}{\widehat{T}} (\widehat{n}_i - n_i) = \frac{\widehat{P}_1^{(0)}}{m_1} \sum_{i=1}^4 \lambda_i \frac{E_i}{\widehat{T}} \\
&= - \frac{\widehat{P}_1^{(0)}}{m_1} \frac{\Delta E}{\widehat{T}}
\end{aligned} \tag{62}$$

Putting together the results obtained in (61) and (62), we can rewrite (60) as

$$\begin{aligned}
& \sum_{i=1}^4 \int_{\mathbb{R}^3} \widehat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{\mathcal{M}_i}{m_i^3} \right) d\mathbf{v} = \frac{\widehat{P}_1^{(0)}}{m_1} \left\{ \log \left[\frac{\widehat{n}_1 \widehat{n}_2}{\widehat{n}_3 \widehat{n}_4} \left(\frac{m_3 m_4}{m_1 m_2} \right)^{3/2} \right] - \frac{\Delta E}{\widehat{T}} \right\} \\
&= 0,
\end{aligned} \tag{63}$$

where assumption (39) associated to the mass action law is used to justify the vanishing of the above integral term. This completes the proof. \square

As a consequence of Theorem 4.1, we can prove that the equilibrium states for our BGK model are defined by the Maxwellian distribution functions given by (14), (15), that assure the equilibrium of the reactive mixture.

Theorem 4.2 *Let assume conditions (38) and (39) on the auxiliary parameters of chemical terms. Under space homogeneous conditions, for all measurable distribution functions $f_i \geq 0$, $i = 1, 2, 3, 4$, we have that*

$$\frac{d\mathcal{H}}{dt}(t) = 0 \quad \text{iff} \quad f_i = f_i^M, \quad \text{for } i = 1, 2, 3, 4. \tag{64}$$

Proof. In Theorem 4.1, we have proven that

$$\frac{d\mathcal{H}}{dt} = \sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \widetilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} + \sum_{i=1}^4 \int_{\mathbb{R}^3} \widehat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v}, \tag{65}$$

where

$$\sum_{i=1}^4 \sum_{j=1}^4 \int_{\mathbb{R}^3} \tilde{\nu}_{ij} (\mathcal{M}_{ij} - f_i) \log \left(\frac{f_i}{m_i^3} \right) d\mathbf{v} \leq 0 \quad (66)$$

and

$$\sum_{i=1}^4 \int_{\mathbb{R}^3} \hat{\nu}_{ij}^{hk} (\mathcal{M}_i - f_i) \log \left(\frac{f_i}{\mathcal{M}_i} \right) d\mathbf{v} \leq 0 \quad (67)$$

since conditions (47), (57) and (59) hold.

Therefore, the time derivative of the \mathcal{H} -function vanishes if and only if relations (66) and (67) hold as equalities. However, as discussed in [15], the equality in (66) implies that the auxiliary velocities $\tilde{\mathbf{u}}_i$ and temperatures \tilde{T}_i share a common value $\tilde{\mathbf{u}}$ and \tilde{T} , respectively, and hence $f_i = n_i \mathbf{M}_i(\mathbf{v}; \tilde{\mathbf{u}}, \tilde{T})$, for $i = 1, 2, 3, 4$. Thus, all constituents of the mixture share a common velocity $\tilde{\mathbf{u}}$ and a common temperature \tilde{T} , assuring mechanical equilibrium of the mixture.

Now, plugging in the above distributions f_i in (67), the equality implies that also $n_i \mathbf{M}_i(\mathbf{v}; \tilde{\mathbf{u}}, \tilde{T}) = \mathcal{M}_i$. Under condition (38), this gives $\mathbf{M}_i(\mathbf{v}; \tilde{\mathbf{u}}, \tilde{T}) = \mathbf{M}_i(\mathbf{v}; \hat{\mathbf{u}}, \hat{T})$ and therefore $\tilde{\mathbf{u}} = \hat{\mathbf{u}}$ and $\tilde{T} = \hat{T}$. Now, condition (39) implies that distributions f_i also verify the chemical equilibrium condition and therefore $f_i = f_i^M$, for $i = 1, 2, 3, 4$. This completes the proof. \square

5. Conclusions

We have proposed a BGK-type kinetic model for a reactive mixture of four monatomic gases undergoing a bimolecular reversible chemical reaction. Although the classical Boltzmann description is able to take into account the intermolecular potential of the microscopic collisional, a BGK formulation is highly desirable, being more manageable from both the analytical and the numerical point of view.

The model proposed in this work extends the description of paper [15] for an inert mixture, where the collision operator is the sum of BGK terms, one for each pair of interacting species. We have assumed that the exchange rates of momentum and energy coincide for Boltzmann and BGK formulations, both in mechanical terms, as done in [15], and in chemical contributions. We point out that the Boltzmann production terms for the chemical operator cannot be computed explicitly, unless a proper approximation of the distribution function is assumed. In this paper, we consider the perturbation of the Maxwellian distribution used in [29], corresponding to a quasi-equilibrium configuration.

The new model proposed here has the great advantage of modeling separately the effects of a chemical reaction and the ones of the mechanical interactions. Moreover, as observed in [5], the mechanical collision operator splits in a sum of binary terms, which takes into account the individual role of any pair of interacting gas species and, thus retains the mixture effects in the collisional dynamics.

Therefore, the BGK equations of this model mimic the structure of the Boltzmann formulation and hence provide the possibility to derive macroscopic evolution equations for the main fields in different hydrodynamic chemical regimes, according to the dominant collisional phenomenon.

As usual in the framework of the reactive mixtures, two distinctive features emerge. At first, the number density of each constituent is not preserved, since particles can change their nature when they interact at the chemical level. This implies that auxiliary number densities appearing in the Maxwellian attractors for chemical contributions result in non-linear functions of masses, species concentration, global auxiliary temperature, and energy

threshold, whose shape derives from model assumptions. The second feature regards the characterization of equilibria from a chemical point of view, which allows us to reproduce correctly the mass action law of chemistry.

Due to these characteristics, some technical difficulties may arise when proving the consistency properties of the model, such as the positivity of number densities and temperatures, as well as the entropy estimate. As regards the latter issue, a proper H-theorem has been proved by assuming in the BGK terms for chemical interactions the equalization of the auxiliary velocities and temperatures and an additional algebraic constraint relating auxiliary densities to global temperature.

Our future projects will involve further development of the ideas in this paper to extend a recent mixed Boltzmann-BGK model to reactive gaseous mixtures. Moreover, it would be interesting to test the proposed model in space-dependent problems, such as the analysis of the one-dimensional shock wave structure.

Acknowledgments

The work of authors G.M. and R.T. was performed in the frame of activities supported by Italian National Group of Mathematical Physics (GNFM-INdAM) and by the University of Parma (Italy). R.T. is a post-doc fellow supported by the National Institute of Advanced Mathematics (INdAM), Italy.

The research work of R.T. and A.J.S. was carried out in the frame of activities sponsored by the Cost Action CA18232, by the Portuguese Projects UIDB/00013/2020 (<https://doi.org/10.54499/UIDB/00013/2020>) of CMAT-UM (<https://doi.org/10.54499/UIDP/00013/2020>), and by the Portuguese national funds (OE), through the Project FCT/MCTES PTDC/03091/2022, “Mathematical Modelling of Multi-scale Control Systems: applications to human diseases – CoSysM3” (<https://doi.org/10.54499/2022.03091.PTDC>).

The work of R.T. was carried out in the frame of activities sponsored by University of Parma through the action Bando di Ateneo 2022 per la ricerca co-funded by MUR-Italian Ministry of Universities and Research - D.M. 737/2021 - PNR - PNRR - NextGenerationEU, Project “Collective and self-organised dynamics: kinetic and network approaches”.

The research work of A.J.S. was carried out in the frame of activities sponsored by Pessoa Project Ref.2021.09255.CBM.

G.M. also thanks the support by the Italian PRIN 2022 PNRR Research Project “Mathematical Modelling for a Sustainable Circular Economy in Ecosystems” (P2022PSMT7) funded by the European Union – NextGenerationEU and by MUR-Italian Ministry of Universities and Research.

A. Appendix – computation of integrals

In this appendix, we include supplementary material, showing the details of the computations involved in the determination of the production terms of mass, momentum, and energy. The computations are quite lengthy and somewhat intricate. We choose to provide more details, focusing solely on the production terms of one constituent. For the other constituents, the computations are rather similar.

More specifically, we calculate the source terms $\widehat{P}_1^{(k)}$, $k = 0, 1, 2$, for the first constituent. By using relations (12), such terms can be written as

$$\widehat{P}_1^{(k)} = \frac{(m_1 m_2)^{3/2}}{(2\pi T)^3} \left[\left(\frac{m_1 m_2}{m_3 m_4} \right)^{3/2} \exp\left(\frac{\Delta E}{T}\right) n_3 n_4 \mathcal{R}_{1+}^k - n_1 n_2 \mathcal{R}_{1-}^{(k)} \right], \quad (68)$$

where

$$\begin{aligned} \mathcal{R}_{1+}^{(k)} &= \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \varphi_1^{(k)}(\mathbf{v}) \\ &\quad \times \mathcal{E}_3(\mathbf{v}') \mathcal{E}_4(\mathbf{w}') \exp\left[-\frac{m_1}{2T} |\mathbf{v} - \mathbf{u}|^2 - \frac{m_2}{2T} |\mathbf{w} - \mathbf{u}|^2\right] d\mathbf{v} d\mathbf{w} d\widehat{\Omega}' \\ \mathcal{R}_{1-}^{(k)} &= \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \varphi_1^{(k)}(\mathbf{v}) \\ &\quad \times \mathcal{E}_1(\mathbf{v}) \mathcal{E}_2(\mathbf{w}) \exp\left[-\frac{m_1}{2T} |\mathbf{v} - \mathbf{u}|^2 - \frac{m_2}{2T} |\mathbf{w} - \mathbf{u}|^2\right] d\mathbf{v} d\mathbf{w} d\widehat{\Omega}'. \end{aligned} \quad (69)$$

Similar expressions for other constituents can be recovered through equivalent forms of the weak formulation of the collision operator [32].

A.1. Some preliminaries

It is convenient to introduce the following change of variable with unit jacobian,

$$(\mathbf{v}, \mathbf{w}) \longrightarrow (\mathbf{v}^*, \mathbf{w}^*) \longrightarrow (\mathbf{G}^*, \mathbf{g}^*), \quad (70)$$

where

$$\mathbf{v}^* = \mathbf{v} - \mathbf{u}, \quad \mathbf{w}^* = \mathbf{w} - \mathbf{u}, \quad \mathbf{G}^* = \alpha_{12} \mathbf{v}^* + \alpha_{21} \mathbf{w}^*, \quad \mathbf{g}^* = \mathbf{v}^* - \mathbf{w}^* = \mathbf{g}. \quad (71)$$

Therefore, we can rewrite

$$-\frac{m_1}{2T} |\mathbf{v} - \mathbf{u}|^2 - \frac{m_2}{2T} |\mathbf{w} - \mathbf{u}|^2 = -\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2, \quad (72)$$

and some of the following integrals intervene

$$\begin{aligned} I_{\alpha\beta\gamma}^\ell &= \int_{\mathbb{R}^3} (G_1^*)^{2\alpha} (G_2^*)^{2\beta} (G_3^*)^{2\gamma} (G^*)^{2\ell} \exp\left(-\frac{m_1 + m_2}{2T} (G^*)^2\right) d\mathbf{G}^* \\ Y_{\alpha\beta\gamma}^\ell &= \int_{\mathbb{R}^3} H(g^2 - \delta_{12}^{34}) g_1^{2\alpha} g_2^{2\beta} g_3^{2\gamma} g^{2\ell} \exp\left(-\frac{\mu_{12}}{2T} g^2\right) d\mathbf{g} \end{aligned} \quad (73)$$

when computing $\mathcal{R}_{1\pm}^{(k)}$. Both types of integrals can be computed explicitly in terms of complete and incomplete Gamma function

$$\Gamma(\alpha) = \int_0^{+\infty} \tau^{\alpha-1} e^{-\tau} d\tau, \quad \Gamma(\alpha, \beta) = \int_\beta^{+\infty} \tau^{\alpha-1} e^{-\tau} d\tau \quad (74)$$

A.2. Production of mass

We have to compute $\mathcal{R}_{1\pm}^{(0)}$, by using the test function $\varphi_1^{(0)}(\mathbf{v}) = m_1$. From (32), we have

$$\mathcal{E}_3(\mathbf{v}') \mathcal{E}_4(\mathbf{w}') = [\mathcal{A}_3 + \mathcal{B}_3 \mathbf{v}'^* \cdot \mathbf{u}_3^* + \mathcal{C}_3 (v'^*)^2] [\mathcal{A}_4 + \mathcal{B}_4 \mathbf{w}'^* \cdot \mathbf{u}_4^* + \mathcal{C}_4 (w'^*)^2], \quad (75)$$

where

$$\mathbf{v}'^* = \mathbf{G}^* + \alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \widehat{\Omega}', \quad \mathbf{w}'^* = \mathbf{G}^* - \alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \widehat{\Omega}', \quad (76)$$

and hence the term $\mathcal{R}_{1+}^{(0)}$ can be rewritten as sum of several integrals in terms of \mathbf{G}^* and \mathbf{g} ; by parity arguments, some of these terms vanish and, consequently, $\mathcal{R}_{1+}^{(0)}$ reduces to

$$\mathcal{R}_{1+}^{(0)} = \mathcal{J}_{1+}^{(0)} + \mathcal{J}_{2+}^{(0)} + \mathcal{J}_{3+}^{(0)} + \mathcal{J}_{4+}^{(0)} + \mathcal{J}_{5+}^{(0)}, \quad (77)$$

where

$$\begin{aligned} J_{1+}^{(0)} &= m_1 \mathcal{A}_3 \mathcal{A}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\ &\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\ &= m_1 \mathcal{A}_3 \mathcal{A}_4 \widehat{\lambda}_{1234}^{(0)} I_{000}^0 Y_{000}^0 \end{aligned} \quad (78)$$

$$\begin{aligned} J_{2+}^{(0)} &= m_1 \mathcal{A}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\ &\quad \times \left[(G^*)^2 - 2\alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\ &\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\ &= m_1 \mathcal{A}_3 \mathcal{C}_4 \widehat{\lambda}_{1234}^{(0)} \left[I_{000}^1 Y_{000}^0 + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right] \end{aligned} \quad (79)$$

$$\begin{aligned} J_{3+}^{(0)} &= m_1 \mathcal{A}_4 \mathcal{C}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\ &\quad \times \left[(G^*)^2 + 2\alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\ &\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\ &= m_1 \mathcal{A}_4 \mathcal{C}_3 \widehat{\lambda}_{1234}^{(0)} \left[I_{000}^1 Y_{000}^0 + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right] \end{aligned} \quad (80)$$

$$\begin{aligned}
J_{4+}^{(0)} &= m_1 \mathcal{B}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\left(\mathbf{G}^* + \alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \widehat{\Omega}' \right) \cdot \mathbf{u}_3^* \right] \\
&\quad \times \left[\left(\mathbf{G}^* - \alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \widehat{\Omega}' \right) \cdot \mathbf{u}_4^* \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\sum_{i=1}^3 (G_i^*)^2 u_{3i}^* u_{4i}^* - \alpha_{34} \alpha_{43} \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \sum_{i,j=1}^3 \widehat{\Omega}'_i \widehat{\Omega}'_j u_{3i}^* u_{4j}^* \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_3 \mathcal{B}_4 \left[\widehat{\lambda}_{1234}^{(0)} Y_{000}^0 (I_{100}^0 u_{31}^* u_{41}^* + I_{010}^0 u_{32}^* u_{42}^* + I_{001}^0 u_{33}^* u_{43}^*) \right. \\
&\quad \left. - \alpha_{34} \alpha_{43} \frac{\mu_{12}}{\mu_{34}} I_{000}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \widehat{\Lambda}_{1234}^{(2)} : (\mathbf{u}_3^* \otimes \mathbf{u}_4^*) \right] \\
&= m_1 \mathcal{B}_3 \mathcal{B}_4 \left[\widehat{\lambda}_{1234}^{(0)} I_{100}^0 Y_{000}^0 \mathbf{u}_3^* \cdot \mathbf{u}_4^* - \alpha_{12} \alpha_{21} I_{000}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \widehat{\Lambda}_{1234}^{(2)} : (\mathbf{u}_3^* \otimes \mathbf{u}_4^*) \right]
\end{aligned} \tag{81}$$

with

$$I_{100}^0 = I_{010}^0 = I_{001}^0 \tag{82}$$

and

$$\widehat{\Lambda}_{1234}^{(2)} = \int_{\mathbb{S}^2} (\widehat{\Omega}' \otimes \widehat{\Omega}') g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') d\widehat{\Omega}' = \text{constant} . \tag{83}$$

$$\begin{aligned}
J_{5+}^{(0)} &= m_1 \mathcal{C}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 + 2\alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times \left[(G^*)^2 - 2\alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{C}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 - 4\alpha_{34} \alpha_{43} \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \sum_{ij=1}^3 G_i^* G_j^* \widehat{\Omega}'_i \widehat{\Omega}'_j \right. \\
&\quad \quad \left. + \alpha_{34}^2 \alpha_{43}^2 \left(\frac{\mu_{12}}{\mu_{34}} \right)^2 (g^2 - \delta_{12}^{34})^2 + (\alpha_{34}^2 + \alpha_{43}^2) \frac{\mu_{12}}{\mu_{34}} (G^*)^2 (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{C}_3 \mathcal{C}_4 \left\{ \widehat{\lambda}_{1234}^{(0)} I_{000}^2 Y_{000}^0 - 4\alpha_{12} \alpha_{21} I_{100}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \text{tr}(\widehat{\Lambda}_{1234}^{(0)}) \right. \\
&\quad \left. + \widehat{\lambda}_{1234}^{(0)} \alpha_{12}^2 \alpha_{21}^2 I_{000}^0 \left[Y_{000}^2 - 2\delta_{12}^{34} Y_{000}^1 + (\delta_{12}^{34})^2 Y_{000}^0 \right] \right. \\
&\quad \left. + \widehat{\lambda}_{1234}^{(0)} (\alpha_{34}^2 + \alpha_{43}^2) \frac{\mu_{12}}{\mu_{34}} I_{000}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right\}
\end{aligned} \tag{84}$$

Analogously, for the loss term, we have that

$$\mathcal{E}_1(\mathbf{v}) \mathcal{E}_2(\mathbf{w}) = [\mathcal{A}_1 + \mathcal{B}_1 \mathbf{v}^* \cdot \mathbf{u}_1^* + \mathcal{C}_1 (v^*)^2] [\mathcal{A}_2 + \mathcal{B}_2 \mathbf{w}^* \cdot \mathbf{u}_2^* + \mathcal{C}_2 (w^*)^2], \tag{85}$$

and $\mathcal{R}_{1-}^{(0)}$ reduces to

$$\mathcal{R}_{1-}^{(0)} = \mathcal{J}_{1-}^{(0)} + \mathcal{J}_{2-}^{(0)} + \mathcal{J}_{3-}^{(0)} + \mathcal{J}_{4-}^{(0)} + \mathcal{J}_{5-}^{(0)}, \tag{86}$$

where

$$\begin{aligned}
J_{1-}^{(0)} &= m_1 \mathcal{A}_1 \mathcal{A}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_1 \mathcal{A}_2 \widehat{\lambda}_{1234}^{(0)} I_{000}^0 Y_{000}^0
\end{aligned} \tag{87}$$

$$\begin{aligned}
J_{2-}^{(0)} &= m_1 \mathcal{A}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 - 2\alpha_{12} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{12}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_1 \mathcal{C}_2 \widehat{\lambda}_{1234}^{(0)} (I_{000}^1 Y_{000}^0 + \alpha_{12}^2 I_{000}^0 Y_{000}^1)
\end{aligned} \tag{88}$$

$$\begin{aligned}
J_{3-}^{(0)} &= m_1 \mathcal{A}_2 \mathcal{C}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_2 \mathcal{C}_1 \widehat{\lambda}_{1234}^{(0)} (I_{000}^1 Y_{000}^0 + \alpha_{21}^2 I_{000}^0 Y_{000}^1)
\end{aligned} \tag{89}$$

$$\begin{aligned}
J_{4-}^{(0)} &= m_1 \mathcal{B}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') [(\mathbf{G}^* + \alpha_{21} \mathbf{g}) \cdot \mathbf{u}_1^*] \\
&\quad \times [(\mathbf{G}^* - \alpha_{12} \mathbf{g}) \cdot \mathbf{u}_2^*] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\sum_{i=1}^3 (G_i^*)^2 u_{1i}^* u_{2i}^* - \alpha_{12} \alpha_{21} \sum_{i=1}^3 g_i^2 u_{1i}^* u_{2i}^* \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_1 \mathcal{B}_2 \widehat{\lambda}_{1234}^{(0)} (I_{100}^0 Y_{000}^0 - \alpha_{12} \alpha_{21} I_{000}^0 Y_{000}^1) \mathbf{u}_1^* \cdot \mathbf{u}_2^*
\end{aligned} \tag{90}$$

with

$$Y_{100}^0 = Y_{010}^0 = Y_{001}^0. \tag{91}$$

$$\begin{aligned}
J_{5-}^{(0)} &= m_1 \mathcal{C}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \\
&\quad \times [(G^*)^2 - 2\alpha_{12} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{12}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{C}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 - 4\alpha_{12} \alpha_{21} \sum_{i=1}^3 (G_i^*)^2 g_i^2 + \alpha_{12}^2 \alpha_{21}^2 g^4 + (\alpha_{12}^2 + \alpha_{21}^2) (G^*)^2 g^2 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{C}_1 \mathcal{C}_2 \widehat{\lambda}_{1234}^{(0)} [I_{000}^2 Y_{000}^0 - 12\alpha_{12} \alpha_{21} I_{100}^0 Y_{100}^0 + \alpha_{12}^2 \alpha_{21}^2 I_{000}^0 Y_{000}^2 + (\alpha_{12}^2 + \alpha_{21}^2) I_{000}^1 Y_{000}^1]
\end{aligned} \tag{92}$$

A.3. Production of momentum

We have to compute $\mathcal{R}_{1\pm}^{(1)}$, by using the test function $\varphi_1^{(0)}(\mathbf{v}) = m_1 \mathbf{v}^*$. More precisely, we have that $\mathcal{R}_{1+}^{(1)}$ is a vector and its first component $(\mathcal{R}_{1+}^{(1)})_1$ reads as

$$(\mathcal{R}_{1+}^{(1)})_1 = (\mathcal{J}_{1+}^{(1)})_1 + (\mathcal{J}_{2+}^{(1)})_1 + (\mathcal{J}_{3+}^{(1)})_1 + (\mathcal{J}_{4+}^{(1)})_1, \tag{93}$$

where, from (75),

$$\begin{aligned}
(\mathcal{J}_{1+}^{(1)})_1 &= m_1 \mathcal{A}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\left(\mathbf{G}^* - \alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \widehat{\Omega}' \right) \cdot \mathbf{u}_4^* \right] (G_1^* + \alpha_{21} g_1) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') (G_1^*)^2 u_{41}^* \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_3 \mathcal{B}_4 \widehat{\lambda}_{1234}^{(0)} I_{100}^0 Y_{000}^0 u_{41}^*
\end{aligned} \tag{94}$$

$$\begin{aligned}
\left(J_{2+}^{(1)}\right)_1 &= m_1 \mathcal{A}_4 \mathcal{B}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\left(\mathbf{G}^* + \alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \widehat{\Omega}' \right) \cdot \mathbf{u}_3^* \right] (G_1^* + \alpha_{21} g_1) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_4 \mathcal{B}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') (G_1^*)^2 u_{31}^* \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_4 \mathcal{B}_3 \widehat{\lambda}_{1234}^{(0)} I_{100}^0 Y_{000}^0 u_{31}^*
\end{aligned} \tag{95}$$

$$\begin{aligned}
\left(J_{3+}^{(1)}\right)_1 &= m_1 \mathcal{B}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\left(\mathbf{G}^* + \alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \widehat{\Omega}' \right) \cdot \mathbf{u}_3^* \right] \\
&\quad \times \left[(G^*)^2 - 2\alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}} (g^2 - \delta_{12}^{34}) \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times (G_1^* + \alpha_{21} g_1) \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G_1^*)^2 (G^*)^2 u_{31}^* + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (G_1^*)^2 (g^2 - \delta_{12}^{34}) u_{31}^* \right. \\
&\quad \left. - 2\alpha_{12} \alpha_{21} (G_1^*)^2 (g^2 - \delta_{12}^{34}) ((\widehat{\Omega}' \otimes \widehat{\Omega}') \cdot \mathbf{u}_3^*)_1 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_3 \mathcal{C}_4 \left\{ \widehat{\lambda}_{1234}^{(0)} \left[I_{100}^1 Y_{000}^0 + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{100}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right] u_{31}^* \right. \\
&\quad \left. - 2\alpha_{12} \alpha_{21} I_{100}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) (\widehat{\Lambda}_{1234}^{(2)} \cdot \mathbf{u}_3^*)_1 \right\}
\end{aligned} \tag{96}$$

$$\begin{aligned}
\left(J_{4+}^{(1)}\right)_1 &= m_1 \mathcal{B}_4 \mathcal{C}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[\left(\mathbf{G}^* - \alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \widehat{\Omega}' \right) \cdot \mathbf{u}_4^* \right] \\
&\quad \times \left[(G^*)^2 + 2\alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times (G_1^* + \alpha_{21} g_1) \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_4 \mathcal{C}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G_1^*)^2 (G^*)^2 u_{41}^* + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (G_1^*)^2 (g^2 - \delta_{12}^{34}) u_{41}^* \right. \\
&\quad \left. - 2\alpha_{12} \alpha_{21} (G_1^*)^2 (g^2 - \delta_{12}^{34}) (\widehat{\Omega}' \otimes \widehat{\Omega}') \cdot \mathbf{u}_4^* \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_4 \mathcal{C}_3 \left\{ \widehat{\lambda}_{1234}^{(0)} \left[I_{100}^1 Y_{000}^0 + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{100}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right] u_{41}^* \right. \\
&\quad \left. - 2\alpha_{12} \alpha_{21} I_{100}^0 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) (\widehat{\Lambda}_{1234}^{(2)} \cdot \mathbf{u}_4^*) \right\}
\end{aligned} \tag{97}$$

We can proceed analogously for the other components of the vector $\mathcal{R}_{1+}^{(1)}$. For what concerns the loss term $\mathcal{R}_{1-}^{(1)}$, the first component is given by

$$\left(\mathcal{R}_{1-}^{(1)}\right)_1 = \left(\mathcal{J}_{1-}^{(1)}\right)_1 + \left(\mathcal{J}_{2-}^{(1)}\right)_1 + \left(\mathcal{J}_{3-}^{(1)}\right)_1 + \left(\mathcal{J}_{4-}^{(1)}\right)_1, \tag{98}$$

where, from (85),

$$\begin{aligned}
\left(J_{1-}^{(1)}\right)_1 &= m_1 \mathcal{A}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(\mathbf{G}^* - \alpha_{12} \mathbf{g}') \cdot \mathbf{u}_2^*] (G_1^* + \alpha_{21} g_1) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') [(G_1^*)^2 u_{21}^* - \alpha_{12} \alpha_{21} g_1^2 u_{21}^*] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_1 \mathcal{B}_2 \widehat{\lambda}_{1234}^{(0)} (I_{100}^0 Y_{000}^0 - \alpha_{12} \alpha_{21} I_{000}^0 Y_{100}^0) u_{41}^*
\end{aligned} \tag{99}$$

$$\begin{aligned}
(J_{2-}^{(1)})_1 &= m_1 \mathcal{A}_2 \mathcal{B}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(\mathbf{G}^* + \alpha_{21} \mathbf{g}') \cdot \mathbf{u}_2^*] (G_1^* + \alpha_{21} g_1) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_2 \mathcal{B}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') [(G_1^*)^2 u_{21}^* + \alpha_{21}^2 g_1^2 u_{21}] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{A}_2 \mathcal{B}_1 \widehat{\lambda}_{1234}^{(0)} (I_{100}^0 Y_{000}^0 + \alpha_{21}^2 I_{000}^0 Y_{100}^0) u_{41}^*
\end{aligned} \tag{100}$$

$$\begin{aligned}
(J_{3-}^{(1)})_1 &= m_1 \mathcal{B}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(\mathbf{G}^* + \alpha_{21} \mathbf{g}) \cdot \mathbf{u}_1^*] [(G^*)^2 - 2\alpha_{12} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{12}^2 g^2] \\
&\quad \times (G_1^* + \alpha_{21} g_1) \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G_1^*)^2 (G^*)^2 u_{11}^* - 4\alpha_{12} \alpha_{21} (G_1^*)^2 g_1^2 u_{11}^* + \alpha_{12}^2 (G_1^*)^2 g^2 u_{11}^* \\
&\quad + \alpha_{21}^2 (G^*)^2 g_1^2 u_{11}^* + \alpha_{12}^2 \alpha_{21}^2 g_1^2 g^2 u_{11}^*] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_1 \mathcal{C}_2 \widehat{\lambda}_{1234}^{(0)} (I_{100}^1 Y_{000}^0 - 4\alpha_{12} \alpha_{21} I_{100}^0 Y_{100}^0 + \alpha_{12}^2 I_{100}^0 Y_{000}^1 \\
&\quad + \alpha_{21}^2 I_{000}^1 Y_{100}^0 + \alpha_{12}^2 \alpha_{21}^2 I_{000}^0 Y_{100}^1) u_{11}^*
\end{aligned} \tag{101}$$

$$\begin{aligned}
(J_{4-}^{(1)})_1 &= m_1 \mathcal{B}_2 \mathcal{C}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(\mathbf{G}^* - \alpha_{12} \mathbf{g}) \cdot \mathbf{u}_1^*] [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \\
&\quad \times (G_1^* + \alpha_{21} g_1) \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_2 \mathcal{C}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G_1^*)^2 (G^*)^2 u_{11}^* + 2\alpha_{21} (\alpha_{21} - \alpha_{12}) (G_1^*)^2 g_1^2 u_{11}^* + \alpha_{21}^2 (G_1^*)^2 g^2 u_{11}^* \\
&\quad - \alpha_{12} \alpha_{21} (G^*)^2 g_1^2 u_{11}^* - \alpha_{12} \alpha_{21}^3 g_1^2 g^2 u_{11}^*] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= m_1 \mathcal{B}_2 \mathcal{C}_1 \widehat{\lambda}_{1234}^{(0)} [I_{100}^1 Y_{000}^0 + 2\alpha_{21} (\alpha_{21} - \alpha_{12}) I_{100}^0 Y_{100}^0 + \alpha_{21}^2 I_{100}^0 Y_{000}^1 \\
&\quad - \alpha_{12} \alpha_{21} I_{000}^1 Y_{100}^0 + \alpha_{12} \alpha_{21}^3 I_{000}^0 Y_{100}^1] u_{11}^*
\end{aligned} \tag{102}$$

Similar expressions can be deduced for the other constituents.

A.4. Production of energy

We use now the test function $\varphi_1^{(2)} = \frac{1}{2} m_1 (v^*)^2$; the gain term can be written as

$$\mathcal{R}_{1+}^{(2)} = \mathcal{J}_{1+}^{(2)} + \mathcal{J}_{2+}^{(2)} + \mathcal{J}_{3+}^{(2)} + \mathcal{J}_{4+}^{(2)} + \mathcal{J}_{5+}^{(2)}, \tag{103}$$

where

$$\begin{aligned}
J_{1+}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_3 \mathcal{A}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \quad (104) \\
&= \frac{1}{2} m_1 \mathcal{A}_3 \mathcal{A}_4 \widehat{\lambda}_{1234}^{(0)} (I_{000}^1 Y_{000}^0 + \alpha_{21}^2 I_{000}^0 Y_{000}^1)
\end{aligned}$$

$$\begin{aligned}
J_{2+}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 - 2\alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 + \alpha_{21}^2 (G^*)^2 g^2 + \alpha_{34}^2 (G^*)^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) + \alpha_{21}^2 \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) g^2 \right] \quad (105) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_3 \mathcal{C}_4 \widehat{\lambda}_{1234}^{(0)} \left[I_{000}^2 Y_{000}^0 + \alpha_{21}^2 I_{000}^1 Y_{000}^1 + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right. \\
&\quad \left. + \alpha_{21}^2 \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^0 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) \right]
\end{aligned}$$

$$\begin{aligned}
J_{3+}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_4 \mathcal{C}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 + 2\alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_4 \mathcal{C}_3 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 + \alpha_{21}^2 (G^*)^2 g^2 + \alpha_{43}^2 (G^*)^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) + \alpha_{21}^2 \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 + \delta_{12}^{34}) g^2 \right] \quad (106) \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_4 \mathcal{C}_3 \widehat{\lambda}_{1234}^{(0)} \left[I_{000}^2 Y_{000}^0 + \alpha_{21}^2 I_{000}^1 Y_{000}^1 + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right. \\
&\quad \left. + \alpha_{21}^2 \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^0 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) \right]
\end{aligned}$$

$$\begin{aligned}
J_{4+}^{(2)} &= \frac{1}{2} m_1 \mathcal{B}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(\mathbf{G}^* + \alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \widehat{\Omega}') \cdot \mathbf{u}_3^* \right] \left[(\mathbf{G}^* - \alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34})} \widehat{\Omega}') \cdot \mathbf{u}_4^* \right] \\
&\quad \times \left[(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2 \right] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{B}_3 \mathcal{B}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \left[(G^*)^2 \sum_{i=1}^3 (G_i^*)^2 u_{3i}^* u_{4i}^* + \alpha_{21}^2 g^2 \sum_{i=1}^3 (G_i^*)^2 u_{3i}^* u_{4i}^* - \alpha_{34} \alpha_{43} (G^*)^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right. \\
&\quad \left. \times \sum_{i,j=1}^3 \widehat{\Omega}'_i \widehat{\Omega}'_j u_{3i}^* u_{4j}^* - \alpha_{34} \alpha_{43} \alpha_{21}^2 \frac{\mu_{12}}{\mu_{34}} g^2 (g^2 - \delta_{12}^{34}) \sum_{i,j=1}^3 \widehat{\Omega}'_i \widehat{\Omega}'_j u_{3i}^* u_{4j}^* \right] \quad (107) \\
&= \frac{1}{2} m_1 \mathcal{B}_3 \mathcal{B}_4 \left[\widehat{\lambda}_{1234}^{(0)} (I_{100}^1 u_{31}^* u_{41}^* + I_{010}^1 u_{32}^* u_{42}^* + I_{001}^1 u_{33}^* u_{43}^*) Y_{000}^0 \right. \\
&\quad + \widehat{\lambda}_{1234}^{(0)} \alpha_{21}^2 (I_{100}^0 u_{31}^* u_{41}^* + I_{010}^0 u_{32}^* u_{42}^* + I_{001}^0 u_{33}^* u_{43}^*) Y_{000}^1 \\
&\quad - \alpha_{12} \alpha_{21} I_{000}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \widehat{\Lambda}_{1234}^{(2)} : (\mathbf{u}_3^* \otimes \mathbf{u}_4^*) \\
&\quad \left. - \alpha_{12} \alpha_{21}^3 I_{000}^0 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) \widehat{\Lambda}_{1234}^{(2)} : (\mathbf{u}_3^* \otimes \mathbf{u}_4^*) \right] \\
&= \frac{1}{2} m_1 \mathcal{B}_3 \mathcal{B}_4 \left\{ \widehat{\lambda}_{1234}^{(0)} (I_{100}^1 Y_{000}^0 + \alpha_{21}^2 I_{100}^0 Y_{000}^1) \mathbf{u}_3^* \cdot \mathbf{u}_4^* - \right. \\
&\quad \left. \alpha_{12} \alpha_{21} [I_{000}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) + \alpha_{21}^2 I_{000}^0 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1)] \widehat{\Lambda}_{1234}^{(2)} : (\mathbf{u}_3^* \otimes \mathbf{u}_4^*) \right\}
\end{aligned}$$

$$\begin{aligned}
J_{5+}^{(2)} &= \frac{1}{2} m_1 \mathcal{C}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 + 2\alpha_{43} \sqrt{\frac{\mu_{12}}{\mu_{34}}(g^2 - \delta_{12}^{34})} \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times \left[(G^*)^2 - 2\alpha_{34} \sqrt{\frac{\mu_{12}}{\mu_{34}}(g^2 - \delta_{12}^{34})} \mathbf{G}^* \cdot \widehat{\Omega}' + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (g^2 - \delta_{12}^{34}) \right] \\
&\quad \times \left[(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2 \right] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{C}_3 \mathcal{C}_4 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^6 + \alpha_{21}^2 (G^*)^4 g^2 + \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (G^*)^4 (g^2 - \delta_{12}^{34}) + \alpha_{21}^2 \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} (G^*)^2 g^2 (g^2 - \delta_{12}^{34}) \right. \\
&\quad - 4\alpha_{34} \alpha_{43} \frac{\mu_{12}}{\mu_{34}} (G^*)^2 (g^2 - \delta_{12}^{34}) \sum_{i,j=1}^3 G_i^* G_j^* \widehat{\Omega}'_i \widehat{\Omega}'_j \\
&\quad - 4\alpha_{21}^2 \alpha_{34} \alpha_{43} \frac{\mu_{12}}{\mu_{34}} g^2 (g^2 - \delta_{12}^{34}) \sum_{i,j=1}^3 G_i^* G_j^* \widehat{\Omega}'_i \widehat{\Omega}'_j \\
&\quad + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (G^*)^4 (g^2 - \delta_{12}^{34}) + \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} (G^*)^2 g^2 (g^2 - \delta_{12}^{34}) \\
&\quad \left. + \alpha_{34}^2 \alpha_{43}^2 \frac{\mu_{12}^2}{\mu_{34}^2} (G^*)^2 (g^2 - \delta_{12}^{34})^2 + \alpha_{21}^2 \alpha_{34}^2 \alpha_{43}^2 \frac{\mu_{12}^2}{\mu_{34}^2} g^2 (g^2 - \delta_{12}^{34})^2 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{C}_3 \mathcal{C}_4 \left[\widehat{\lambda}_{1234}^{(0)} I_{000}^3 Y_{000}^0 + \widehat{\lambda}_{1234}^{(0)} \alpha_{21}^2 I_{000}^2 Y_{000}^1 + \widehat{\lambda}_{1234}^{(0)} \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^2 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \right. \\
&\quad + \widehat{\lambda}_{1234}^{(0)} \alpha_{21}^2 \alpha_{34}^2 \frac{\mu_{12}}{\mu_{34}} I_{000}^1 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) - 4\alpha_{12} \alpha_{21} I_{100}^1 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \text{tr}(\widehat{\Lambda}_{1234}^{(0)}) \\
&\quad - 4\alpha_{12} \alpha_{21}^3 I_{100}^0 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) \text{tr}(\widehat{\Lambda}_{1234}^{(0)}) + \widehat{\lambda}_{1234}^{(0)} \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} I_{100}^2 (Y_{000}^1 - \delta_{12}^{34} Y_{000}^0) \\
&\quad + \widehat{\lambda}_{1234}^{(0)} \alpha_{43}^2 \frac{\mu_{12}}{\mu_{34}} I_{100}^1 (Y_{000}^2 - \delta_{12}^{34} Y_{000}^1) + \widehat{\lambda}_{1234}^{(0)} \alpha_{12}^2 \alpha_{21}^2 I_{100}^1 (Y_{000}^2 - 2\delta_{12}^{34} Y_{000}^1 + \delta_{12}^{34} Y_{000}^0) \\
&\quad \left. + \widehat{\lambda}_{1234}^{(0)} \alpha_{12}^2 \alpha_{21}^4 I_{100}^0 (Y_{000}^3 - 2\delta_{12}^{34} Y_{000}^2 + \delta_{12}^{34} Y_{000}^1) \right] \tag{108}
\end{aligned}$$

Analogously, we can rewrite

$$\mathcal{R}_{1-}^{(2)} = \mathcal{J}_{1-}^{(2)} + \mathcal{J}_{2-}^{(2)} + \mathcal{J}_{3-}^{(2)} + \mathcal{J}_{4-}^{(2)} + \mathcal{J}_{5-}^{(2)}, \tag{109}$$

where

$$\begin{aligned}
J_{1-}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_1 \mathcal{A}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2 \right] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* dg d\widehat{\Omega}' \tag{110} \\
&= \frac{1}{2} m_1 \mathcal{A}_1 \mathcal{A}_2 \widehat{\lambda}_{1234}^{(0)} (I_{000}^1 Y_{000}^0 + \alpha_{21}^2 I_{000}^0 Y_{000}^1)
\end{aligned}$$

$$\begin{aligned}
J_{2-}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 - 2\alpha_{12} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{12}^2 g^2] [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 + \alpha_{21}^2 (G^*)^2 g^2 - 4\alpha_{12} \alpha_{21} \sum_{i=1}^3 (G_i^*)^2 g_i^2 + \alpha_{12}^2 (G^*)^2 g^2 + \alpha_{12}^2 \alpha_{21}^2 g^4 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_1 \mathcal{C}_2 \widehat{\lambda}_{1234}^{(0)} [I_{000}^2 Y_{000}^0 + \alpha_{21}^2 I_{000}^1 Y_{000}^1 - 12\alpha_{12} \alpha_{21} I_{100}^0 Y_{100}^0 + \alpha_{12}^2 I_{000}^1 Y_{000}^1 \\
&\quad + \alpha_{12}^2 \alpha_{21}^2 I_{000}^0 Y_{000}^2]
\end{aligned} \tag{111}$$

$$\begin{aligned}
J_{3-}^{(2)} &= \frac{1}{2} m_1 \mathcal{A}_2 \mathcal{C}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2]^2 \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_2 \mathcal{C}_1 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^4 + \alpha_{21}^2 (G^*)^2 g^2 + 4\alpha_{21}^2 \sum_{i=1}^3 (G_i^*)^2 g_i^2 + \alpha_{21}^2 (G^*)^2 g^2 + \alpha_{21}^4 g^4 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{A}_2 \mathcal{C}_1 \widehat{\lambda}_{1234}^{(0)} [I_{000}^2 Y_{000}^0 + 2\alpha_{21}^2 I_{000}^1 Y_{000}^1 + 12\alpha_{21}^2 I_{100}^0 Y_{100}^0 + \alpha_{21}^4 I_{000}^0 Y_{000}^2]
\end{aligned} \tag{112}$$

$$\begin{aligned}
J_{4-}^{(2)} &= \frac{1}{2} m_1 \mathcal{B}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(\mathbf{G}^* + \alpha_{21} \mathbf{g}) \cdot \mathbf{u}_1^*] [(\mathbf{G}^* - \alpha_{12} \mathbf{g}) \cdot \mathbf{u}_2^*] \\
&\quad \times [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2] \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{B}_1 \mathcal{B}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^2 \sum_{i=1}^3 (G_i^*)^2 u_{3i}^* u_{4i}^* + \alpha_{21}^2 g^2 \sum_{i=1}^3 (G_i^*)^2 u_{3i}^* u_{4i}^* - 2\alpha_{12} \alpha_{21} \sum_{i=1}^3 (G_i^*)^2 g_i^2 u_{1i}^* u_{2i}^* \right. \\
&\quad \left. + 2\alpha_{21}^2 \sum_{i=1}^3 (G_i^*)^2 g_i^2 u_{1i}^* u_{2i}^* - \alpha_{12} \alpha_{21} (G^*)^2 \sum_{i=1}^3 g_i^2 u_{1i}^* u_{2i}^* - \alpha_{12} \alpha_{21}^3 g^2 \sum_{i=1}^3 g_i^2 u_{1i}^* u_{2i}^* \right] \\
&= \frac{1}{2} m_1 \mathcal{B}_1 \mathcal{B}_2 \widehat{\lambda}_{1234}^{(0)} \{ I_{100}^1 Y_{000}^0 + \alpha_{21}^2 I_{100}^0 Y_{000}^1 + 2\alpha_{21} (\alpha_{21} - \alpha_{12}) I_{100}^0 Y_{100}^0 \\
&\quad - \alpha_{12} \alpha_{21} I_{000}^1 Y_{100}^0 - \alpha_{12} \alpha_{21}^3 I_{000}^0 Y_{100}^1 \} \mathbf{u}_1^* \cdot \mathbf{u}_2^*
\end{aligned} \tag{113}$$

$$\begin{aligned}
J_{5-}^{(2)} &= \frac{1}{2} m_1 \mathcal{C}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times [(G^*)^2 - 2\alpha_{12} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{12}^2 g^2] [(G^*)^2 + 2\alpha_{21} \mathbf{G}^* \cdot \mathbf{g} + \alpha_{21}^2 g^2]^2 \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{C}_1 \mathcal{C}_2 \int_{\mathbb{R}^3 \times \mathbb{R}^3 \times \mathbb{S}^2} H(g^2 - \delta_{12}^{34}) g \widehat{\sigma}_{1234}(g, \widehat{\Omega} \cdot \widehat{\Omega}') \\
&\quad \times \left[(G^*)^6 + (\alpha_{12}^2 + 2\alpha_{21}^2) (G^*)^4 g^2 + \alpha_{21}^2 (2\alpha_{12}^2 + \alpha_{21}^2) (G^*)^2 g^4 + \alpha_{12}^2 \alpha_{21}^2 g^6 \right. \\
&\quad \left. + 4\alpha_{21} (\alpha_{21} - 2\alpha_{12}) (G^*)^2 \sum_{i=1}^3 (G_i^*)^2 g_i^2 + 4\alpha_{12} \alpha_{21}^2 (\alpha_{12} - 2\alpha_{21}) g^2 \sum_{i=1}^3 (G_i^*)^2 g_i^2 \right] \\
&\quad \times \exp \left[-\frac{m_1 + m_2}{2T} (G^*)^2 - \frac{\mu_{12}}{2T} g^2 \right] d\mathbf{G}^* d\mathbf{g} d\widehat{\Omega}' \\
&= \frac{1}{2} m_1 \mathcal{C}_1 \mathcal{C}_2 \widehat{\lambda}_{1234}^{(0)} \left[I_{000}^3 Y_{000}^0 + (\alpha_{12}^2 + 2\alpha_{21}^2) I_{000}^2 Y_{000}^1 + \alpha_{21}^2 (2\alpha_{12}^2 + \alpha_{21}^2) I_{000}^1 Y_{000}^2 \right. \\
&\quad \left. + \alpha_{12}^2 \alpha_{21}^2 I_{000}^0 Y_{000}^3 + 12\alpha_{21} (\alpha_{21} - 2\alpha_{12}) I_{100}^1 Y_{100}^0 + 12\alpha_{12} \alpha_{21}^2 (\alpha_{12} - 2\alpha_{21}) I_{100}^0 Y_{100}^1 \right] \tag{114}
\end{aligned}$$

Declarations

Competing interests: The authors declare no competing interests directly or indirectly related to the work submitted for publication.

Conflict-of-interest statement: The authors have no conflicts of interest to declare.

Data availability statement: The authors declare that no datasets were generated or analyzed during the current study.

References

- [1] P. Andries, K. Aoki, B. Perthame, A consistent BGK-type model for gas mixtures, *J. Stat. Phys.*, **106**, 993–1018, 2002.
- [2] P.L. Bhatnagar, E.P. Gross, K. Krook, A model for collision processes in gases, *Phys. Rev.*, **94**, 511–524, 1954.
- [3] M. Bisi, W. Boscheri, G. Dimarco, M. Groppi, G. Martalò, A new mixed Boltzmann-BGK model for mixtures solved with an IMEX finite volume scheme on unstructured meshes, *Appl. Math. Comput.*, **433**, 127416, 2022.
- [4] M. Bisi, M. Groppi, E. Lucchin, G. Martalò, A mixed Boltzmann-BGK model for inert gas mixtures, *Kinet. Relat. Models* early access, 2023.
- [5] M. Bisi, M. Groppi, G. Martalò, Macroscopic equations for inert gas mixtures in different hydrodynamic regimes, *J. Phys. A: Math. Theor.*, **54**(8), 085201, 2021.
- [6] M. Bisi, M. Groppi, G. Spiga, Grad's distribution functions in the kinetic equations for a chemical reaction, *Contin. Mech. Thermodyn.*, **14**, 207–222, 2002.
- [7] M. Bisi, M. Groppi, G. Spiga, Kinetic modelling of bimolecular chemical reactions, in *Kinetic Methods for Nonconservative and Reacting Systems*, G. Toscani ed., Quaderni di Matematica, **16**, 1–143, Aracne Editrice, Roma, 2005.
- [8] M. Bisi, M. Groppi, G. Spiga, Kinetic problems in rarefied gas mixtures, *AIP Conf. Proc.*, **1084**, 487–92, 2008.

- [9] M. Bisi, M. Groppi, G. Spiga, Kinetic Bhatnagar–Gross–Krook model for fast reactive mixtures and its hydrodynamic limit, *Phys. Rev. E*, **81**, 036327 1–9, 2010.
- [10] M. Bisi, R. Monaco, A.J. Soares, A BGK model for reactive mixtures of polyatomic gases with continuous internal energy, *J. Phys. A - Math. Theor.*, **51**, 125501 1–29, 2018.
- [11] M. Bisi, R. Travaglini, A BGK model for mixtures of monoatomic and polyatomic gases with discrete internal energy, *Phys. A: Stat. Mech. Appl.*, **547**, 124441, 2020.
- [12] M. Bisi, R. Travaglini, A kinetic BGK relaxation model for a reacting mixture of polyatomic gases, in *Recent Advances in Kinetic Equations and Applications*, F. Salvarani ed., Springer INdAM Series, **48**, Springer International Publishing, 2021.
- [13] M. Bisi, R. Travaglini, BGK model for a mixture with two reversible reactions, in *INdAM Workshop The Legacy of Carlo Cercignani: from Kinetic Theory to Turbulence Modeling*, Springer Nature Singapore, 59–72, 2021.
- [14] M. Briant, E. S. Daus, The Boltzmann equation for a multi-species mixture close to global equilibrium, *Arch. Ration. Mech. Anal.*, **222**, 1367–1443, 2016.
- [15] A.V. Bobylev, M. Bisi, M. Groppi, G. Spiga, I.F. Potapenko, A general consistent BGK model for gas mixtures, *Kinet. Relat. Models*, **11**, 1377–1393, 2018.
- [16] A. Bondesan, L. Boudin, B. Grec, A numerical scheme for a kinetic model for mixtures in the diffusive limit using the moment method, *Numer. Meth. Part. D. E.*, **35**(3), 1184–1205, 2019.
- [17] S. Brull, V. Pavan, J. Schneider, Derivation of a BGK model for mixtures, *European Journal Of Mechanics B/Fluids*, **33**, 74–86, 2012.
- [18] S. Brull, J. Schneider, Derivation of a BGK model for reacting gas mixtures, *Commun. Math. Sci.*, **12**, 1199–1223, 2014.
- [19] C. Cercignani, *The Boltzmann Equation and its Applications*, Springer, New York, 1988.
- [20] S. Chapman, T.G. Cowling, *The Mathematical Theory of Non-Uniform Gases*, Cambridge University Press, Cambridge, 1970.
- [21] A. Crestetto, C. Klingenberg, M. Pirner, Kinetic/fluid micro-macro numerical scheme for a two component gas mixture, *Multiscale Model. Simul.*, **18**(2), 970–998, 2020.
- [22] V. S. Galkin, N. K. Makashev, Kinetic derivation of the gas-dynamic equation for multicomponent mixtures of light and heavy particles, *Fluid Dyn.*, **29**(1), 140–155, 1994.
- [23] I. M. Gamba, M. Pavić-Cölić, On existence and uniqueness to homogeneous Boltzmann flows of monatomic gas mixtures, *Arch. Ration. Mech. Anal.*, **235**(1), 723–781, 2020.
- [24] V. Garzó, A. Santos, J.J. Brey, A kinetic model for a multicomponent gas, *Phys. Fluids*, **1**(2), 380–383, 1989.
- [25] V. Giovangigli, *Multicomponent Flow Modeling. Series on Modeling and Simulation in Science, Engineering and Technology*, Birkhäuser, Boston, 1999.

- [26] M. Groppi, G. Spiga, Grad's distribution functions in the kinetic equations for a chemical reaction, *Phys. Fluids*, **16**, 4273–4284, 2004.
- [27] J.R. Haack, C.D. Hauck, M.S. Murillo, A conservative, entropic multispecies BGK model, *J. Stat. Phys.*, **168**, 826–856, 2017
- [28] C. Klingenberg, M. Pirner, G. Puppo, A consistent kinetic model for a two–component mixture with an application to plasma, *Kinet. Relat. Models*, **10**, 445–465, 2017.
- [29] G.M. Kremer, M. Pandolfi Bianchi, A.J. Soares, A relaxation kinetic model for transport phenomena in a reactive flow, *Phys. Fluids*, **18**, 037104 1–15, 2006.
- [30] J.C. Maxwell, XIII. The Bakerian Lecture. – On the viscosity or internal friction of air and other gases, *Philos. Trans. Royal Soc*, **156**, 249–268, 1866.
- [31] M. Pirner, *Kinetic modelling of gas mixtures*, Würzburg University Press, Würzburg, 2018.
- [32] A. Rossani, G. Spiga, A note on the kinetic theory of chemically reacting gases, *Physica A*, **272**(3-4), 563–573, 1999.
- [33] L. Wu, J. Zhang, , J. M. Reese, , Y. Zhang, A fast spectral method for the Boltzmann equation for monatomic gas mixtures, *J. Comput. Phys.*, **298**, 602–621, 2015.
- [34] Y. Zhang, , L. Zhu, R. Wang, Z. Guo, Discrete unified gas kinetic scheme for all Knudsen number flows. III. Binary gas mixtures of Maxwell molecules, *Phys. Rev. E*, **97**(5), 053306, 2018.