

ON THE REDUCTION OF POWERS OF SELF-ADJOINT OPERATORS

SALIMA KEBLI AND MOHAMMED HICHEM MORTAD*

ABSTRACT. Let $T \in B(H)$ be such that T^n is self-adjoint for some $n \in \mathbb{N}$ with $n \geq 3$. The paper's primary aim is to establish the conditions that lead to the self-adjointness of T . We pay particular attention to the case where $T^3 = 0$ and how it implies T is complex symmetric.

1. INTRODUCTION

First, we assume that readers are familiar with definitions and concepts of bounded linear operators. A suitable reference for this work is [14]. We will recall specific definitions, however.

In this manuscript, we denote a complex Hilbert space as H , which can be infinite-dimensional. The algebra of bounded linear operators from H into itself is denoted as $B(H)$. Let $T \in B(H)$. Call T self-adjoint when $T^* = T$, where T^* is the usual adjoint, which is the conjugate transpose of T if T is a matrix. We say that T is normal if $TT^* = T^*T$. Say that $T \in B(H)$ is complex symmetric if $T = CT^*C$ for some conjugation $C \in B(H)$, where a conjugation is a conjugate-linear operator $C \in B(H)$ that is both involutive ($C^2 = I$) and isometric (see, e.g., [8]).

Call T a square root of S , where $S \in B(H)$, provided $T^2 = S$.

The operator T is called positive, denoted as $T \geq 0$, if $\langle Tx, x \rangle \geq 0$ for all $x \in H$ (in the context of matrices, this means positive semi-definite). It is well-known that each positive operator has a unique positive square root. Since T^*T is positive, it has a unique positive square root, which we denote by $|T|$, as is customary.

Remember that any operator T in $B(H)$ can be expressed as $T = A + iB$, where A and B are self-adjoint. This decomposition is called

2020 *Mathematics Subject Classification*. Primary 47A62. Secondary 47B15, 47B44, 47B65, 47A05.

Key words and phrases. Self-adjoint operators; Normal operators; Real and Imaginary parts; Positive Operators; Nilpotent operators; Powers of operators; Complex symmetric operators.

* Corresponding author.

the Cartesian decomposition of T . As per convention, we represent A as $\operatorname{Re} T$ and B as $\operatorname{Im} T$, where $\operatorname{Re} T = (T+T^*)/2$ and $\operatorname{Im} T = (T-T^*)/2i$.

It is evident that T is self-adjoint if and only if $B = 0$. Also, $T = 0$ if and only if $A = B = 0$. Furthermore, T is normal if and only if $AB = BA$.

That a normal operator with a real spectrum is self-adjoint is typically proven using the spectral theorem. However, a new simple proof based on the Cartesian decomposition can be found in [15].

Now, consider the equation $T^2 = S$, where $S, T \in B(H)$. It is of some interest to know when T belongs to the same class of S . For example, if S is self-adjoint (respectively normal), when is T self-adjoint (respectively normal)? Just when $H = \mathbb{C}^2$, taking $S = 0$, which is obviously self-adjoint, shows that T can be anything, e.g., $T = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$,

which is not even normal (in fact, not quite anything as such a matrix is necessarily complex symmetric, as in Theorem 5 in [8]). On the other hand, the equation $T^2 = I$, still in a bi-dimensional space, can have an infinitude of self-adjoint solutions. See, e.g., Question 6.2.1 in [17]. Also, the shift operator on ℓ^2 does not have any square root. See Problem 151 in [10]. In [13], the authors provided complete descriptions of the set of square roots of certain classical operators, such as the Volterra and the Cesàro operators, among others.

Some authors have sought for conditions on T that place it in the same class as that of S . For example, C. R. Putnam showed in [20] that if $T \in B(H)$ satisfies $T^2 = S$, where $S \geq 0$ and $\operatorname{Re} T \geq 0$, then T is necessarily the unique positive square root of S . He also obtained in [19] that if T^2 is self-adjoint and $\operatorname{Re}\langle Tx, x \rangle \neq 0$ for all $x \in H$, then T is self-adjoint. Some other results can be consulted in [6], [12], [16], [19], and [21]. Special cases involving possibly unbounded operators are discussed in [4], [7] and [18].

Regarding the class of normal operators, it is also known that if T^2 is normal and $\operatorname{Re}(T) \geq 0$, then T is normal. This result has appeared in at least two papers in the literature ([5] and [19]). We may also add [7], which discusses both a specific and a more general version; specific because $T^2 = 0$, and more general as it deals with $T^n = 0$, where $n \geq 3$. We digress to say that this result already appeared in [12] (Lemma 3) but the authors of [7] were not aware of it, and besides, their proof is different, simpler, and also dealt with unbounded operators.

There are further results that provide conditions for the operator T to be in the class of, for example, T^n . This type of results differs from what we are after in this manuscript, as we wanted some results that

make sense even for matrices. For example, some of these conditions involve non-normal operators such as hyponormal or quasinormal ones, but these results are not applicable when $\dim H$ is finite because, for example, hyponormality is equivalent to normality in that case. Other results involving unbounded symmetric and self-adjoint operators are also straightforward when $T \in B(H)$, as these concepts coincide in such a case.

In the end, this paper aims to continue the investigation, focusing on the more general and challenging case of higher powers while mainly restricting ourselves to self-adjoint operators.

2. THE CASE OF THE SELF-ADJOINTNESS OF T^3

It is plain that if $T \in B(H)$ is self-adjoint, then so are all its powers T^n . Now, if T^2 is self-adjoint, then T^{2p} is self-adjoint for all $p \in \mathbb{N}$, but, e.g., T^3 need not be self-adjoint. Similarly, if T^3 is self-adjoint, so are T^{3q} for any $q \in \mathbb{N}$ and T^2 could be non-self-adjoint. Therefore, the following simple observation might be helpful.

Proposition 2.1. *Let $T \in B(H)$ be such that T^2 and T^3 are self-adjoint. Then T^n is self-adjoint for any $n \geq 4$, with $n \in \mathbb{N}$.*

Proof. Let $n \geq 4$, where $n \in \mathbb{N}$. Write $n = 2p + 3q$ for some $p, q \in \mathbb{N}$. Since commuting self-adjoint operators are self-adjoint, as are powers of self-adjoint operators, we have

$$T^n = T^{2p+3q} = T^{2p}T^{3q} = T^{2p}T^{3q} = (T^2)^p(T^3)^q,$$

which shows that T^n is self-adjoint. □

Remark. The preceding result for the class of normal operators first appeared in [11], then in [1]. The proof presented here for self-adjoint operators may also be adapted to the normal ones by remembering that the product of two commuting normal operators is normal.

Remark. It is worth noting that the previous result and remark also apply to unbounded operators without any changes or additions.

From now on, we will be interested in forms of the converse of the previous result. In the introduction, we have already mentioned that if T^2 is normal while $\operatorname{Re} T \geq 0$, then T is normal. This result is not valid anymore if we replace "normal" with "self-adjoint" (witness $T = iI$). Now, we inquire whether T remains self-adjoint if only T^3 is self-adjoint while also assuming $\operatorname{Re} T \geq 0$. The answer is again negative.

Example 2.2. For instance, consider

$$T = \begin{pmatrix} 0 & i \\ i & 1 \end{pmatrix},$$

which is defined on \mathbb{C}^2 . Then $T^3 = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, which is self-adjoint,

but T itself is not. Observe, in the end, that $\operatorname{Re} T = \begin{pmatrix} 0 & 0 \\ 0 & 1/2 \end{pmatrix}$, which is clearly positive semi-definite.

In the above example, $T^2 = \begin{pmatrix} -1 & i \\ i & 0 \end{pmatrix}$ indicating that T^2 is not self-adjoint. Therefore, we are inclined to propose the following result:

Theorem 2.3. *Let $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. If T^3 and T^2 are self-adjoint with $A \geq 0$ (or $A \leq 0$), then T is self-adjoint.*

We present two proofs: the first is based on Cartesian decomposition, while the second is somewhat simple but relies on a result by M. R. Embry.

First proof of Theorem 2.3. It is easy to see that

$$T^2 = A^2 - B^2 + i(AB + BA)$$

and

$$T^3 = A^3 - B^2A - AB^2 - BAB + i(A^2B - B^3 + ABA + BA^2).$$

Since $A^3 - B^2A - AB^2 - BAB$ and $A^2B - B^3 + ABA + BA^2$ are self-adjoint, they form the real and imaginary parts of T^3 respectively.

Given that T^3 is self-adjoint, its imaginary part must vanish, leading to $B^3 = A^2B + ABA + BA^2$. Furthermore, T^2 is self-adjoint; we must have $AB + BA = 0$, or $AB = -BA$. Thus, $A^2B = BA^2$. If $A \geq 0$, then $AB = BA$; and if $A \leq 0$, we obtain $-AB = -BA$. In either case, $AB = BA$. Consequently, $AB = 0$. Returning to the imaginary part of T^3 , we see that $B^3 = 0$, from which we derive $B = 0$. Thus, T is self-adjoint. \square

Second proof of Theorem 2.3. Since T^2 is self-adjoint, the equation $T^2T = TT^2$ gives $T^2T^* = T^*T^2$. So

$$T^*TTT^* = T^*T^*TT = TT^*T^*T (= T^4)$$

(such a T is commonly known as binormal). Thus, $|T||T^*| = |T^*||T|$. When $\operatorname{Re} T \geq 0$, Theorem 2 in [5] implies the normality of T . By

replacing T with $-T$, we observe that the condition $\operatorname{Re} T \leq 0$ also ensures the normality of T .

We now prove that T is self-adjoint. Since T is normal, it suffices to show that its spectrum is a subset of \mathbb{R} . Let $\lambda \in \sigma(T)$. Since both T^2 and T^3 are self-adjoint, it follows by the spectral mapping theorem that λ^2 and λ^3 are real, which means that λ must also be real. Therefore, we conclude that T is self-adjoint. \square

Remark. Notice that the self-adjointness of T^2 and T^3 does not even yield the normality of $T \in B(H)$. For instance, any nilpotent non-zero 2×2 matrix is a counterexample.

Remark. Example 2.2 already supplies a non-normal matrix T such that T^2 is not normal either, yet T^3 is self-adjoint and $\operatorname{Re} T \geq 0$. This example is somehow discouraging as we also have $\operatorname{Re} T^2 \leq 0$. Thus, it seems hard to obtain the normality of a T if T^m is normal for some $n \geq 3$ and if $\operatorname{Re} T^m \geq 0$ (or ≤ 0) for a certain (or all) m .

A similar reasoning yields the following result:

Theorem 2.4. *Let $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. If T^3 and T^2 are self-adjoint with $B \geq 0$ (or $B \leq 0$), then T is self-adjoint.*

Thus, we have an intriguing way of characterizing the self-adjointness of T through its real and imaginary parts, which conclusively eliminates the possibility of finding superior results.

Corollary 2.5. *Let $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. Assume that T^3 is self-adjoint. If $A \geq 0$ or $A \leq 0$ or $B \geq 0$ or $B \leq 0$, then*

$$T^2 \text{ is self-adjoint} \iff T \text{ is self-adjoint.}$$

Next, we treat the case of positive operators, which, in light of Theorem 2.3, has now become straightforward to deal with.

Corollary 2.6. *Let $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. If T^3 and T^2 are positive with $B \geq 0$ (or $B \leq 0$), then T is positive.*

Proof. Since positivity is stronger than self-adjointness, by Theorem 2.3, we can conclude that T is self-adjoint. Let λ be an a priori complex number in $\sigma(T)$. Since $T^2, T^3 \geq 0$, we can infer that $\lambda^2, \lambda^3 \geq 0$. Since the condition $\lambda^2 \geq 0$ implies the realness of λ , $\lambda^3 \geq 0$ yields $\lambda \geq 0$, which means that T is positive, marking the end of the proof. \square

Remark. If we assume that $\operatorname{Re} T \geq 0$, then T is positive if T^2 is positive, that is, without needing to add the positivity of T^3 (a similar

conclusion is reached if we suppose $\operatorname{Re} T \leq 0$ in lieu). We already alluded that this result appeared in [20]. To show that the above corollary is not covered by the result of [20], let $T = -I$. Then $T^2 \geq 0$ and $\operatorname{Im} T = 0$, yet T is not positive, hence the need for assuming $T^3 \geq 0$.

If the positivity of the real or imaginary part of T is dropped, then T remains self-adjoint but under a specific assumption. For instance, if $T \in B(H)$ is invertible and T^2 and T^3 are self-adjoint, then T is also self-adjoint. This was demonstrated in [4] for two relatively prime numbers (which need not be just 2 and 3) and for an unbounded self-adjoint T . The subsequent result represents a slight improvement of this finding.

Proposition 2.7. *Let $T \in B(H)$ be such that T^2 and T^3 are self-adjoint. If $\ker T = \ker T^2$ or $\ker T^* = \ker T^2$, then T is self-adjoint.*

Proof. Since T^2 and T^3 are self-adjoint, we have $T^2 T^* = T^* T^2$ and $T^3 T^* = T^* T^3$. Thus,

$$T^3 T^* = T^* T^3 = T^* T^2 T = T^2 T^* T,$$

that is, $T^2(TT^* - T^*T) = 0$. Since $\ker T = \ker T^2$, we even have $T(TT^* - T^*T) = 0$, or equivalently, $TTT^* = TT^*T$. This says that T^* is quasinormal. But a hyponormal operator, which is a weaker notion than quasinormality, is self-adjoint as soon as it has a real spectrum (see [22]). Arguing as in the second proof of Theorem 2.3, we can establish the self-adjointness of T^* or T .

In case $\ker T^* = \ker T^2$, we have

$$T^2(TT^* - T^*T) = 0 \implies T^*(TT^* - T^*T) = 0,$$

which means that T is quasinormal. Thus, and as above, T is self-adjoint. \square

What other generalizations of Theorem 2.3 are possible? One could conjecture the following:

Conjecture 2.8. *If the real part of $T \in B(H)$ is positive and both T^3 and T^4 are self-adjoint, then T is self-adjoint.*

Before attempting to address this conjecture, however, readers should first consider the following example:

Example 2.9. Let

$$T = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

which is not self-adjoint. Then T^2 is not self-adjoint because

$$T^2 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

However,

$$T^3 = T^4 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

are self-adjoint (and idempotent as well). Observe in the end that the self-adjoint $\operatorname{Re}T$ is not positive (or negative) semi-definite, as it has eigenvalues of opposite signs, namely $\frac{\sqrt{2}}{2}$ and $-\frac{\sqrt{2}}{2}$.

Therefore, this example does not answer the conjecture above, but it could be useful for other purposes. Still related to the above conjecture, we have the following observation:

Proposition 2.10. *Let $T \in B(H)$ be such that T^3 and T^4 are self-adjoint. If $\operatorname{Re}T \geq 0$ and $\operatorname{Re}(T^2) \geq 0$, then T is self-adjoint.*

Proof. Since T^3 is self-adjoint, so is T^6 . Thus, by Corollary 2.5 and the self-adjointness of T^4 , T^2 too is self-adjoint as $\operatorname{Re}(T^2) \geq 0$. Since $\operatorname{Re}T \geq 0$, the self-adjointness of both T^2 and T^3 yields that of T using the same corollary. \square

Remark. The conclusion of the previous result stays unchanged when $\operatorname{Im}T$ (or $\operatorname{Im}T^2$) substitutes $\operatorname{Re}T$ (or $\operatorname{Re}T^2$) and when " ≥ 0 " is exchanged with " ≤ 0 ".

3. THE GENERAL CASE OF AN ODD POWER

Theorem 3.1. *Let $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. Let $n \in \mathbb{N}$ be such that $n \geq 2$. Suppose T^{2n+1} and T^2 are self-adjoint with $A \geq 0$. Then T is self-adjoint.*

Proof. Since T^2 is self-adjoint, we have $AB + BA = 0$. As shown in the first proof of Theorem 2.3, we can deduce that $BA = AB = 0$. To find the imaginary part of T^{2n+1} , which is nil, there is no need to expand T^{2n+1} using the Binomial theorem. It is evident that each expression of the form $A^p B^q$, for any $p, q \in \mathbb{N}$, is zero due to the condition $AB = BA = 0$. Thus, we are left with $B^{2n+1} = 0$, which then leads to $B = 0$, making T self-adjoint, as wished. \square

Remark. It is primordial to have an odd power in the preceding theorem. Indeed, if $T = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$. Then $T^2 = -I$ and $T^4 = I$ are self-adjoint, $\operatorname{Re} T = 0$, and yet T is not self-adjoint. What went wrong is the fact that $T^3 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$ is not self-adjoint.

Remark. Here, too, and as in Corollary 2.5, if T^{2n+1} is self-adjoint and $\operatorname{Re} T \geq 0$, then T is self-adjoint if and only if T^2 is self-adjoint.

4. THE CASE OF NILPOTENT OPERATORS

If $T \in B(H)$ is such that $T^3 = 0$, then there is, a priori, no reason why we should have $T^2 = 0$, unless $\dim H = 2$. The following result provides a way of reducing the index of nilpotence for an operator on an infinite-dimensional space.

Theorem 4.1. *Let $T \in B(H)$ be such that $T^3 = 0$, where $\dim H \geq 3$. If $\operatorname{Re}(T^2) \geq 0$ or $\operatorname{Im}(T^2) \geq 0$, then $T^2 = 0$.*

Remark. Readers will notice from the proof of the previous result that we will obtain the same outcome if we substitute " ≥ 0 " with " ≤ 0 ".

The above result could be shown using various methods. We have chosen one based on the Cartesian decomposition. First, we require the following auxiliary result.

Lemma 4.2. *Let $R, S \in B(H)$ be self-adjoint, with one being positive. Assume $S^2 = R^2$ and $SR = -RS$. Then it follows that $R = S = 0$.*

Proof. Let us assume $R \geq 0$ without loss of generality. Since $SR = -RS$, we can deduce that $SR^2 = R^2S$, and therefore $SR = RS$. Thus, $SR = 0$, which implies $SR^2 = S^2R = 0$. As a result, $R^3 = S^3 = 0$, and consequently, $R = S = 0$, as desired. \square

Remark. There are self-adjoint non-zero matrices R and S that satisfy $S^2 = R^2$ and $SR = -RS$. Indeed, consider two of the Pauli spin matrices on \mathbb{C}^2 , namely:

$$R = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ and } S = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$$

(more commonly denoted by σ_x and σ_y). It is clear that $RS = -SR$ and $R^2 = S^2$. However, neither matrices R and S are positive semi-definite. More generally, if S and R are two self-adjoint operators and one of them is positive, then $SR = \lambda RS \neq 0$, where $\lambda \in \mathbb{C}$, implies that $\lambda = 1$ only. This result is originally from [2], and a different proof can be found in [3].

Proof of Theorem 4.1. Write $T = A + iB$, where $A, B \in B(H)$ are self-adjoint. We have

$$T^3 = A^3 - B^2A - AB^2 - BAB + i(A^2B - B^3 + ABA + BA^2).$$

Since $A^3 - B^2A - AB^2 - BAB$ and $A^2B - B^3 + ABA + BA^2$ are self-adjoint, and $T^3 = 0$, it ensues that

$$A^3 - B^2A - AB^2 - BAB = 0 \text{ and } A^2B - B^3 + ABA + BA^2 = 0.$$

These last two equations may be rewritten in different forms, e.g.,

$$\begin{aligned} (A^2 - B^2)A &= (AB + BA)B, & (B^2 - A^2)B &= (AB + BA)A, \\ A(A^2 - B^2) &= B(AB + BA), & B(B^2 - A^2) &= A(AB + BA). \end{aligned}$$

Therefore,

$$(AB + BA)A^2 = (B^2 - A^2)BA \text{ and } (A^2 - B^2)AB = (AB + BA)B^2,$$

and so

$$(AB + BA)(A^2 - B^2) = -(A^2 - B^2)(AB + BA).$$

On the other hand,

$$(A^2 - B^2)A^2 = (AB + BA)BA \text{ and } (B^2 - A^2)B^2 = (AB + BA)AB,$$

thereby

$$(A^2 - B^2)^2 = (AB + BA)^2.$$

Since $A^2 - B^2$ and $AB + BA$ are both self-adjoint and, $A^2 - B^2 \geq 0$ (or $AB + BA \geq 0$), Lemma 4.2 implies that $A^2 - B^2 = AB + BA = 0$. Since $T^2 = A^2 - B^2 + i(AB + BA)$, it is seen that $T^2 = 0$, as required. \square

Corollary 4.3. *Let $T \in B(H)$ be such that $T^3 = 0$, where $\dim H \geq 3$. If $\operatorname{Re}(T^2) \geq 0$ or $\operatorname{Im}(T^2) \geq 0$, then T is complex symmetric.*

Proof. Based on Theorem 4.1, we have $T^2 = 0$, and according to Theorem 5 in [8], it follows that T is complex symmetric. \square

Remark. The preceding corollary is interesting in the sense that the authors of [9] showed in their Theorem 2 that for every finite $n \geq 3$ and for every H with $\dim H \geq n$, there is an algebraic operator T on H , that is, $p(T) = 0$ for some polynomial p , of degree n , which is not a complex symmetric operator.

We conclude with the following simple observation.

Proposition 4.4. *Let $T \in B(H)$ be such that $T^n = 0$ for some $n \geq 2$, where $n \in \mathbb{N}$. Then T^{n-1} is self-adjoint if and only if $T^{n-1} = 0$.*

Proof. We only show that the self-adjointness of T^{n-1} implies $T^{n-1} = 0$. Since $T^n = 0$, it follows that $(T^{n-1})^2 = T^{2n-2} = T^n T^{n-2} = 0$, which yields $T^{n-1} = 0$, as wished. \square

REFERENCES

1. S.A. Alzuraiqi, A.B. Patel. On n -normal operators, *General Math. Notes*, **1** (2010) 61-73.
2. J. A. Brooke, P. Busch, D. B. Pearson. Commutativity up to a factor of bounded operators in complex Hilbert space, *R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.*, **458/2017** (2002) 109-118.
3. Ch. Chellali, M. H. Mortad. Commutativity up to a factor for bounded and unbounded operators, *J. Math. Anal. Appl.*, **419/1** (2014), 114-122.
4. S. Dehimi, M. H. Mortad. Unbounded operators having self-adjoint, subnormal or hyponormal powers, *Math. Nachr.*, **296/9** (2023) 3915-3928.
5. M. R. Embry. Conditions implying normality in Hilbert space, *Pacific J. Math.*, **18** (1966) 457-460.
6. M. R. Embry. n th roots of operators, *Proc. Amer. Math. Soc.*, **19** (1968) 63-68.
7. N. Frid, M. H. Mortad, S. Dehimi. When nilpotence implies the zeroness of linear operators, *Khayyam J. Math.*, **8/2** (2022) 163-173.
8. S. R. Garcia. Aluthge transforms of complex symmetric operators, *Integral Equations Operator Theory*, **60/3** (2008) 357-367.
9. S. R. Garcia, W. R. Wogen. Some new classes of complex symmetric operators, *Trans. Amer. Math. Soc.*, **362/11** (2010) 6065-6077.
10. P. R. Halmos. A Hilbert space problem book, *Springer*, 1982 (2nd edition).
11. A. A. Jibril. On n -power normal operators, *Arab. J. Sci. Eng. Sect. A Sci.*, **33/2** (2008) 247-251.
12. S. Kurepa. On n -th roots of normal operators, *Math. Z.*, **78** (1962) 285-292.
13. J. Mashreghi, M. Ptak, W. T. Ross. The square roots of some classical operators, *Studia Math.*, **269/1** (2023) 83-106.
14. M. H. Mortad. An operator theory problem book, *World Scientific Publishing Co.*, (2018).
15. M. H. Mortad. On the invertibility of the sum of operators, *Anal. Math.*, **46/1** (2020) 133-145.
16. M. H. Mortad. On the existence of normal square and n th roots of operators, *J. Anal.*, **28/3** (2020) 695-703.
17. M. H. Mortad. Counterexamples in operator theory. *Birkhäuser/Springer*, Cham (2022).
18. M. H. Mortad, Certain properties involving the unbounded operators $p(T)$, TT^* , and T^*T ; and some applications to powers and n th roots of unbounded operators, *J. Math. Anal. Appl.*, **525/2** (2023). Paper No. 127159, 26 pp.
19. C. R. Putnam. On square roots of normal operators, *Proc. Amer. Math. Soc.*, **8** (1957) 768-769.
20. C. R. Putnam. On square roots and logarithms of self-adjoint operators, *Proc. Glasgow Math. Assoc.*, **4** (1958) 1-2.
21. H. Radjavi, P. Rosenthal. On roots of normal operators, *J. Math. Anal. Appl.*, **34** (1971) 653-664.
22. J. G. Stampfli. Hyponormal operators and spectral density, *Trans. Amer. Math. Soc.*, **117** (1965) 469-476.

(BOTH AUTHORS) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ORAN 1,
AHMED BEN BELLA, B.P. 1524, EL MENOVAR, ORAN 31000, ALGERIA.

Email address: skebli@yahoo.com, kebli.salima@univ-oran1.dz

Email address: mhmortad@gmail.com, mortad.hichem@univ-oran1.dz.