

SOME CONDITIONAL FKG INEQUALITIES FOR MARKOV CHAINS AND RANDOM PROCESSES

ALEXANDRE LEGRAND

ABSTRACT. This paper is interested in proving correlation inequalities of the FKG-type for a broad range of random processes. The two pivotal tools which yield these correlation inequalities are that many discrete-time Markov chains, even under a very general conditioning, satisfy the FKG inequality, and that such inequalities are preserved through weak convergences. In particular, we prove that Lévy processes, Bessel processes and several conditioned Brownian processes have positive correlations.

The *FKG inequality* is a correlation inequality named after Fortuine, Ginibre and Kasteleyn [8], who proved a sufficient condition for its validity in the Ising and other lattice models (see also [10]). It has seen many applications and generalizations in statistical mechanics and general probability theory, see e.g. [2, 6, 11, 15, 21, 23, 24] and references therein, and it is a recurrent tool in proofs of stochastic domination results, as well as computations of lower bounds on first moments or event probabilities. In particular, Barbato [1] proved that the Brownian motion, as well as solutions of stochastic differential equations with Lipschitz-continuous drift coefficients, satisfy the FKG inequality; and this result has already seen several applications, the reader can refer to [5, 9, 22, 25] for some examples.

The present paper aims to contribute to this literature in a few directions. First, we extend the FKG inequality in [1] from the Brownian motion to general Lévy processes, which we achieve by presenting an alternate proof. Then, we show an FKG inequality for a class of conditional distributions of Markov chain trajectories. Not only this allows us to prove that many other general processes in continuous time, such as Bessel processes and conditioned Brownian processes, satisfy the FKG inequality too; this result also relates to the matter of correlation inequalities for conditional distributions —a topic which has notably aroused interest in the domains of spin systems (see [19] and references therein) as well as percolation [3, 4].

1. MAIN RESULTS

1.1. The FKG inequality. Consider $(E, \mathcal{E}, \preceq)$ a measurable set endowed with a measurable partial order (i.e. $\{(u, v) \in E^2; u \preceq v\} \in \mathcal{E}^{\otimes 2}$). A function $f : E \rightarrow \mathbb{R}$ is said to be *increasing* (or non-decreasing) on E if $u \preceq v$ implies $f(u) \leq f(v)$ for all $u, v \in E$. This paper will mostly be interested in the following cases, for some $n \in \mathbb{N}$ or $T > 0$:

- (i) $E = \mathbb{R}^n$, $\mathcal{E} = \text{Bor}(\mathbb{R}^n)$ and for $u, v \in E$, $u \preceq v$ if and only if $u_i \leq v_i$ for all $1 \leq i \leq n$,
- (ii) $E = \mathcal{C}([0, T], \mathbb{R})$, $\mathcal{E} = \mathcal{E}_{\mathcal{C}}$ is the σ -algebra induced by the uniform convergence topology, and for $u, v \in E$, $u \preceq v$ if and only if $u(t) \leq v(t)$ for all $t \in [0, T]$.

2020 Mathematics Subject Classification. Primary: 60J10, 60E15 ; Secondary: 60G15, 60G51.

Key words and phrases. FKG inequality, conditional positive correlations, Markov chains.

(iii) $E = \mathcal{D}([0, T], \mathbb{R})$ is the set of right-continuous with left-hand limits (rcll) functions on $[0, T]$, $\mathcal{E} = \mathcal{E}_{\mathcal{D}}$ is the σ -algebra induced by the Skorokhod metric, and for $u, v \in E$, $u \preceq v$ if and only if $u(t) \leq v(t)$ for all $t \in [0, T]$.

Definition 1.1. Let \mathbf{P} a probability measure on $(E, \mathcal{E}, \preceq)$, and X a random variable with law \mathbf{P} : then X is *associated* (or *positively correlated*) if for all increasing functions $f, g \in L^2(\mathbf{P})$, one has

$$(1.1) \quad \mathbf{E}[f(X)g(X)] \geq \mathbf{E}[f(X)]\mathbf{E}[g(X)].$$

We shall abusively write that \mathbf{P} is associated whenever $X \sim \mathbf{P}$ is.

When X is associated, we may also say that it “satisfies the FKG inequality”. Our first result is an extension of [1, Theorem 4] from the Brownian motion to all Lévy processes.

Theorem 1.2. *Let X be a one-dimensional Lévy process. Then it is associated.*

The proof of this theorem relies on elementary arguments on random walks and weak convergence of measures, however the author is not aware of a previous reference stating this property. Moreover, even though the proof displayed here is simpler than that of [1] (which relies on non-trivial measure theory and the introduction of an auxiliary order relation), it seems that some of their arguments could be extended to continuous random processes with independent, non-stationary increments.

1.2. Conditional association of Markov chains. The second (and most central) result of this paper is that some Markov chains distributions, after conditioning on a quite generic event, are associated. Then, we combine this statement with convergence arguments below to deduce that several processes in continuous time are associated. Let us introduce the following definitions.

Definition 1.3. We say that an event $A \in \mathcal{E}$ is *max/min-stable* if for $u, v \in A$, one has $u \vee v \in A$ and $u \wedge v \in A$, where $u \vee v$ (resp. $u \wedge v$) denotes the smallest common upper bound (resp. largest lower bound) of u and v in E .

For the sake of simplicity, in this paper we always assume that $u \vee v$, $u \wedge v$ are well-defined for all $u, v \in E$ —which is the case if $E = \mathbb{R}^d$, $E = \mathcal{C}([0, T], \mathbb{R})$ or $E = \mathcal{D}([0, T], \mathbb{R})$.

Definition 1.4. Let $\mathbf{p} := (p_{x,y})_{x,y \in \mathcal{X}}$ a probability transition kernel on some countable subset $\mathcal{X} \subset \mathbb{R}$ (i.e. $p_{x,y} \geq 0$ and $\sum_y p_{x,y} = 1$). For $\mathcal{X}_1 \subset \mathcal{X}$, we say that \mathbf{p} has \mathcal{X}_1 -*unfavourable crossings* (abbreviated \mathcal{X}_1 -u.c.) if for all $u_1, v_1 \in \mathcal{X}_1$, $u_2, v_2 \in \mathcal{X}$ one has,

$$(1.2) \quad p_{u_1, u_2} \times p_{v_1, v_2} \leq p_{u_1 \vee v_1, u_2 \vee v_2} \times p_{u_1 \wedge v_1, u_2 \wedge v_2}.$$

For \mathbf{P} a (purely atomic) probability measure on \mathbb{R} , we say that \mathbf{P} has \mathcal{X}_1 -u.c. if the transition kernel of the random walk with increment distribution \mathbf{P} has \mathcal{X}_1 -u.c..

We now provide some examples of Markov chains on \mathbb{Z} which have (or not) unfavourable crossings. If \mathbf{P} is an atomic measure on some measurable space E , we write abusively $\mathbf{P}(u) := \mathbf{P}(\{u\})$ for $u \in E$.

Lemma 1.5. (i) *Discrete Laplace: let $\beta > 0$ and $\mathbf{P}(z) := C e^{-\beta|z|} \mathbf{1}_{z \in \mathbb{Z}}$, where $C = C(\beta)$ is a normalizing constant. Then \mathbf{P} has \mathbb{Z} -u.c. (i.e. the transition kernel of the random walk with increment distribution \mathbf{P} has \mathbb{Z} -u.c.).*

(ii) *Unitary steps: assume $p_{i,i+1} + p_{i,i-1} = 1$ for all $i \in \mathbb{Z}$. Then \mathbf{p} has both $(2\mathbb{Z})$ -u.c. and $(2\mathbb{Z} + 1)$ -u.c..*

Example 1.6. (i) Let $\alpha > 1$ and $\mathbf{P}(z) := C(1 + |z|)^{-\alpha} \mathbf{1}_{z \in \mathbb{Z}}$, where $C = C(\alpha)$ is a normalizing constant. Then \mathbf{P} does not have \mathbb{Z} -u.c..

(ii) Let $\varepsilon \in [0, 1/3)$, and $\mathbf{P} := \frac{1-\varepsilon}{2}(\delta_{-1} + \delta_{+1}) + \varepsilon\delta_0$. Then \mathbf{P} does not have \mathbb{Z} -u.c..

The proofs of Lemma 1.5 and Example 1.6 are very straightforward and are left as an exercise to the reader (nonetheless let us mention that a proof of Lemma 1.5.(i) can be found in [18, (B.5)]).

We may now state our central result. For a general random variable X on some space (E, \mathcal{E}) , we denote its law with \mathbf{P}_X . Moreover recall that for any $A \in \mathcal{E}$ such that $\mathbf{P}_X(A) > 0$, the law of X conditioned to A is defined with $\mathbf{P}_X(B|A) := \mathbf{P}_X(B \cap A)/\mathbf{P}_X(A)$, $B \in \mathcal{E}$. We have the following.

Theorem 1.7. *Let $X = (X_1, \dots, X_n)$ be the first n steps of a Markov chain on some countable set $\mathcal{X} \subset \mathbb{R}$ started from some $x_0 \in \mathcal{X}$. Assume \mathcal{X} is locally finite, and let \mathbf{p} denote the transition kernel of X . Let $A \subset \mathcal{X}^n$, and assume the following:*

(H1) *for all $k \in \{0, \dots, n-1\}$, \mathbf{p} has $\text{Supp}(\mathbf{P}_{X_k})$ -u.c.,*

(H2) *$\mathbf{P}_X(A) > 0$ and A is max/min-stable,*

then, $\mathbf{P}_X(\cdot|A)$ is associated.

Example 1.8. Let X be a random walk on \mathbb{Z} started from some $x_0 \in \mathbb{Z}$ and with increment distribution \mathbf{P} . If \mathbf{P} has \mathbb{Z} -unfavourable crossings, or if $\mathbf{P}(\{-1, +1\}) = 1$ (so \mathbf{P} has both $(2\mathbb{Z})$ - and $(2\mathbb{Z} + 1)$ -unfavourable crossings by Lemma 1.5.(ii)), then (H1) holds. In particular, let us mention that the specific case of the random walk on \mathbb{Z} with discrete Laplace increments has already been studied in [18, Proposition B.1].

Remark 1.1. One can clearly take $A := \mathcal{X}^n$ in the theorem, hence \mathbf{P}_X is associated as soon as \mathbf{p} satisfies (H1).

Remark 1.2. One could be tempted to replace the assumption (H1) with “ \mathbf{p} has \mathcal{X} -u.c.”, however the latter is a strictly stronger assumption since it excludes some periodic Markov chains (e.g. compare Lemma 1.5.(ii) with Example 1.6.(ii) for $\varepsilon = 0$).

When combined with some weak convergence results, Theorem 1.7 allows us to show that several continuous processes are positively correlated.

Corollary 1.9. *Let $E = \mathcal{C}([0, T], \mathbb{R})$ for some $T > 0$. Then, the distributions of the following Brownian processes $(B_t)_{t \in [0, T]}$ in $(\mathcal{C}([0, T], \mathbb{R}), \mathcal{E}_{\mathcal{C}}, \preceq)$, started from $B_0 = 0$, are associated:*

(1) *the Brownian bridge, $B_T = x$, $x \in \mathbb{R}$,*

(2) *the Brownian excursion, $B_T = 0$, $B_t \geq 0$ for all $t \in [0, T]$,*

(3) *the Brownian meander, $B_t \geq 0$ for all $t \in [0, T]$,*

(4) *the Brownian motion conditioned to remain in some time-dependent interval, $B_t \in [a(t), b(t)]$ for all $t \in [0, T]$, where $a, b \in \mathcal{D}([0, T], [-\infty, +\infty])$ satisfy $\mathbf{P}_\omega(\forall t \in [0, T], \omega_t \in [a(t), b(t)]) > 0$, for \mathbf{P}_ω the canonical Wiener measure on $\mathcal{C}([0, T], \mathbb{R})$.*

Corollary 1.10. *Let $E = \mathcal{C}([0, T], \mathbb{R})$. Then the Bessel process $(X_t)_{t \in [0, T]}$ of index $\nu \in (-1, +\infty)$, i.e. with generator $\frac{1}{2} \frac{d^2}{dx^2} + \frac{2\nu+1}{2x} \frac{d}{dx}$, and started from some $x_0 \geq 0$, is associated.*

Let us clarify that these corollaries of Theorem 1.7 are not meant to be exhaustive, but illustrative; the reader may think of many similar applications other than Corollaries 1.9, 1.10.

1.3. Comments.

Association for SDE solutions. As mentioned above, association (i.e. the FKG inequality) has already been proven to hold for some class of random diffusion processes in [1]. Not only this paper presents an alternate proof in the Brownian case, our results can also be seen as an extension of those of [1] in the case of SDE solutions. More precisely, the later paper only considers SDE solutions with a Lipschitz drift term, which rules out solutions of SDEs with barriers (of any kind), such as Bessel and some conditioned Brownian processes. We conjecture that many other solutions of SDEs (with barriers) also satisfy the FKG inequality: noticeably, if a random process is associated, applying to it an increasing change of variable preserves the FKG inequality (see Lemma 2.4 below, and [1, Lemma 8]). We leave these very general considerations for further work.

Another direction in which we expect our results can be pushed is the case of SDEs driven by a Lévy process, see e.g. [16]. Again, we deduce from Lemma 2.4 below and Theorem 1.2 that any increasing function of a Lévy process is associated, but we do not delve deeper in this direction in the present paper.

A final way in which our results can be extended is that of SDE solution approximation. If there exists a sequence of Markov chains which are associated and converge weakly to an SDE solution, then it is associated (see Lemma 2.1 below). Therefore, it would be interesting to prove that some usual approximation schemes from the SDE literature are associated, implying the FKG inequality for very general SDE solutions.

Conditional association. The matter of proving association (or other correlation inequalities) for conditional distributions has been investigated in various settings, see e.g. [4, 3, 19] or even [22, Lemma 2.3]. However, to the author’s knowledge, the vast majority of existing results make strong assumptions on the conditioning set $A \subset E$. In Theorem 1.7, the assumption (H2) on the set A seems to be minimal, whereas the crucial assumption (H1) only involves the transition kernel of the Markov chain *without conditioning*.

The assumption (H1) is closely related to a correlation property called the “FKG condition” (see (3.2) below), which is central to the proof of Theorem 1.7. In general, the FKG condition implies association (see notably [8, 24]), but they are not equivalent: counter-examples in various settings can be found e.g. in [7, 19], and in this paper we provide another one with random walk distributions in Section 4. As a matter of fact, one can see the association of some conditional distributions as an intermediary property between the FKG condition and association without conditioning —see in particular [19], and the notions of “downward FKG” and “downward conditional association” therein. Therefore, since we prove in Lemma 3.3 below that the assumption (H1) actually implies the FKG condition, it is not surprising that it yields conditional association for generic sets A .

We believe that there are (at least) two interesting directions in which this matter could be developed, in the case of trajectories of general Markov chains on a totally ordered state space $\mathcal{X} \subset \mathbb{R}$. First, it would be interesting to determine if the assumption (H1) is necessary and sufficient for the FKG condition to hold (we prove in Section 4 below that this is the case for random walks). Second, when (H1) and/or the FKG condition fails, it would be interesting to determine if some conditional association inequalities still hold, at the cost of a stronger assumption than (H2) on the conditioning set, similarly to [3] or [19].

Organization of the paper. In Section 2 we prove that general random walks are associated, which yields Theorem 1.2 with a weak convergence argument. In Section 3 we prove Theorem 1.7, and we apply it to some well-chosen conditioned Markov chains in order to deduce Corollaries 1.9

and 1.10. Finally, in Section 4 we further discuss the FKG condition for random walks, showing in that case that it is equivalent to (H1).

2. ASSOCIATION AND WEAK CONVERGENCE

In this section, we state the main lemma that allows us to push the FKG inequality through weak convergences. In particular this yields Theorem 1.2, which also shows that the standard Brownian motion is positively correlated with an alternate, simpler proof to that of [1].

Theorem 1.2 is a consequence of the following, quite elementary statement.

Lemma 2.1. *Let $(E, \mathcal{E}, \preceq)$ a metric space endowed with its Borel σ -algebra and a measurable partial order. Let (\mathbf{P}_n) , $n \geq 1$ a sequence of probability measures on (E, \mathcal{E}) , which converges to some probability measure \mathbf{P}_∞ . Assume that \mathbf{P}_n is associated for all $n \geq 1$; then \mathbf{P}_∞ is associated.*

In order to prove the lemma, we recall the following classical property.

Lemma 2.2. *Let $(E, \mathcal{E}, \preceq, \mathbf{P})$ a metric space endowed with its Borel σ -algebra, a measurable partial order and a probability measure. The inequality (1.1) holds for all increasing functions $f, g \in L^2(\mathbf{P})$ if and only if it holds for all Lipschitz-continuous, non-negative, bounded and increasing functions.*

This follows from a standard integral approximation result, which we do not detail in this paper.

Proof of Lemma 2.1. Let f, g be bounded, increasing and continuous functions: then, under the assumptions of Lemma 2.1, f, g and \mathbf{P}_n satisfy (1.1) for any $n \geq 1$. Taking the limit $n \rightarrow +\infty$, so does \mathbf{P}_∞ . Applying Lemma 2.2, this finishes the proof. \square

We now turn to the proof of Theorem 1.2. It is a consequence of the following claim.

Proposition 2.3. *Let \mathbf{P} be any probability measure on \mathbb{R} , and let $X = (X_0, X_1, \dots, X_n)$ a random walk with independent increments with law \mathbf{P} , and started from some (random) initial value $X_0 \sim \mathbf{P}_{X_0}$. Then \mathbf{P}_X is associated.*

Before proving this proposition, let us state some fundamental results.

Lemma 2.4. *Let $(E_1, \mathcal{E}_1, \preceq_1, \mathbf{P}_1)$ be a partially ordered probability space. Let $(E_2, \mathcal{E}_2, \preceq_2)$ be a partially ordered measurable space. Let $f : E_1 \rightarrow E_2$ a measurable, increasing function, and let \mathbf{P}_2 be the push-forward measure of \mathbf{P}_1 by f . If \mathbf{P}_1 is associated, then so is \mathbf{P}_2 .*

The proof of this lemma is straightforward, see e.g. [1, Proposition 2] for the details. Moreover, we recollect the following classical result.

Theorem 2.5. (i) *Let \mathbf{P} be a probability measure on $(\mathbb{R}, \text{Bor}(\mathbb{R}), \leq)$. Then \mathbf{P} is associated.*

(ii) *Let \mathbf{P} be a product probability measure on $(\mathbb{R}^n, \text{Bor}(\mathbb{R}^n), \preceq)$, $n \geq 1$. Then \mathbf{P} is associated.*

Proof. The first statement follows from the fact that \leq is a total order on \mathbb{R} , and from a direct computation of $\mathbf{E}[(f(X) - f(X'))(g(X) - g(X'))] \geq 0$ where X' is an independent copy of $X \sim \mathbf{P}$ and f, g are increasing (we leave the details to the reader). The second statement can then be deduced from [1, Theorem 3] or [24, Theorem 3]. \square

Proof of Proposition 2.3. Define,

$$(2.1) \quad \begin{aligned} & \mathbb{R}^{n+1} \mapsto \mathbb{R}^{n+1} \\ \varphi : & (z_i)_{0 \leq i \leq n} \rightarrow \left(\sum_{i=0}^k z_i \right)_{0 \leq k \leq n}, \end{aligned}$$

and notice that it is increasing, i.e. for $u, v \in \mathbb{R}^{n+1}$, $u \preceq v$ implies $\varphi(u) \preceq \varphi(v)$. Let $X = (X_0, \dots, X_n)$ a random walk started from X_0 , and let $U := \varphi^{-1}(X)$. In particular $\mathbf{P}_U = \mathbf{P}_{X_0} \otimes \mathbf{P}^{\otimes n}$ is a product measure, so it is associated by Theorem 2.5. Since \mathbf{P}_X is the push-forward of \mathbf{P}_U by φ , by Lemma 2.4 it is associated as well, finishing the proof. \square

Proof of Theorem 1.2. For the sake of presentation, we first detail the proof in the specific case of the Brownian motion, then we extend the argument to general Lévy processes. For $n \geq 1$, let $Y^n = (Y_0, \dots, Y_n)$ be the first steps of a random walk, and assume that its increments are centered with variance 1. For $T > 0$, define

$$(2.2) \quad \widehat{X}_t^n := n^{-1/2} Y_{\lfloor nt/T \rfloor}, \quad t \in [0, T],$$

which is a random process in $\mathcal{D}([0, T], \mathbb{R})$. Notice that \widehat{X}^n is an increasing function of Y^n , so by Lemma 2.4 and Proposition 2.3 it is associated. Moreover by Donsker's theorem, \widehat{X}^n converges weakly to the Brownian motion as $n \rightarrow +\infty$ (provided that X_0^n converges weakly to some real random variable B_0). Using Lemma 2.1, we deduce that the Brownian motion is associated.

For a general Lévy process X , the fact that it is infinitely divisible implies that there exists a sequence of random walks $(S_k^n)_{0 \leq k \leq m_n}$, $n, m_n \in \mathbb{N}$, whose rcll interpolations $\widehat{X}_t^n := S_{\lfloor m_n t/T \rfloor}^n$, $t \in [0, T]$ converge weakly to X in $\mathcal{D}([0, T], \mathbb{R})$ as $n \rightarrow +\infty$, see e.g. [14, Theorem 23.14]. Clearly the process \widehat{X}^n can be written as an increasing function of a random walk, hence $\mathbf{P}_{\widehat{X}^n}$ is associated. By Lemma 2.1, so is \mathbf{P}_X , which finishes the proof. \square

3. CONDITIONAL ASSOCIATION FOR MARKOV CHAINS

In this section we prove Theorem 1.7, and then Corollaries 1.9, 1.10. For the sake of simplicity, in this section we only consider Markov chains and processes started from a deterministic initial value a.s.. Then, our results may be extended to general initial values with the following lemma, which we include for the sake of completeness.

Lemma 3.1. *Let I a generic set and $E \subset \mathbb{R}^I$ be a partially ordered measurable space stable by translation (e.g. \mathbb{R}^n , $\mathcal{C}([0, T], \mathbb{R})$ or $\mathcal{D}([0, T], \mathbb{R})$). Let Y be a random process in E . Let X_0 be a real random variable, and assume that for almost every X_0 , $\mathbf{P}_Y(\cdot | X_0)$ is associated. Then, \mathbf{P}_{Y+X_0} is associated.*

Notice that Y and X_0 are not supposed to be independent, so this lemma is much more general than the argument used in Proposition 2.3 (in particular it applies to mixtures of Markov chains).

Proof. This is straightforward: let $f, g : E \rightarrow \mathbb{R}$ be increasing functions (one can also assume they are bounded, recall Lemma 2.2), then one has

$$(3.1) \quad \begin{aligned} \mathbf{E}[f(Y + X_0)g(Y + X_0)] &= \mathbf{E}[\mathbf{E}[f(Y + X_0)g(Y + X_0)|X_0]] \\ &\geq \mathbf{E}[\mathbf{E}[f(Y + X_0)|X_0]\mathbf{E}[g(Y + X_0)|X_0]] \\ &\geq \mathbf{E}[\mathbf{E}[f(Y + X_0)|X_0]]\mathbf{E}[\mathbf{E}[g(Y + X_0)|X_0]] = \mathbf{E}[f(Y + X_0)]\mathbf{E}[g(Y + X_0)], \end{aligned}$$

where the first inequality comes from the fact that, for almost every X_0 , Y is associated and the functions $y \mapsto f(y + X_0)$, $y \mapsto g(y + X_0)$ are increasing on E ; and the second inequality comes from the observation that $x \mapsto \mathbf{E}[f(Y + X_0)|X_0 = x]$, $x \mapsto \mathbf{E}[g(Y + X_0)|X_0 = x]$ are increasing functions on \mathbb{R} , and since X_0 is a real random variable, by Theorem 2.5.(i) it is associated. This finishes the proof of the lemma. \square

3.1. Proof of Theorem 1.7. Let us first introduce the following definition.

Definition 3.2. Let $(E, \mathcal{E}, \preceq, \mathbf{P})$ be a partially ordered probability space, and assume \mathbf{P} is a purely atomic measure. We say that \mathbf{P} satisfies the *FKG condition* if,

$$(3.2) \quad \mathbf{P}(u \vee v)\mathbf{P}(u \wedge v) \geq \mathbf{P}(u)\mathbf{P}(v), \quad \text{for } \mathbf{P}\text{-a.e. } u, v \in E.$$

Then, the proof of our main result is achieved in three short steps, which are contained in the following lemmata. Recall assumptions (H1) and (H2) from Theorem 1.7.

Lemma 3.3. *Let $n \geq 1$, and let $X = (X_1, \dots, X_n)$ be the first n steps of a Markov chain on some countable subset $\mathcal{X} \subset \mathbb{R}$, started from $X_0 = x_0 \in \mathcal{X}$ and with transition kernel \mathbf{p} . If \mathbf{p} satisfies (H1), then for all $1 \leq k \leq n$, $\mathbf{P}_{(X_1, \dots, X_k)}$ satisfies the FKG condition on \mathbb{R}^k .*

Proof. For $(u_j)_{1 \leq j \leq n}$ and $1 \leq k \leq n$, write $u^k := (u_1, \dots, u_k)$, and similarly $X^k := (X_1, \dots, X_k)$. Since $\mathbf{p} = (p_{x,y})_{x,y \in \mathcal{X}}$ satisfies (H1), this implies that for \mathbf{P}_{X^k} -a.e. $u^k, v^k \in \mathcal{X}^k$, one has

$$(3.3) \quad \begin{aligned} \mathbf{P}_{X^k}(u^k \vee v^k)\mathbf{P}_{X^k}(u^k \wedge v^k) &= \prod_{j=0}^{k-1} p_{u_j \vee v_j, u_{j+1} \vee v_{j+1}} \times \prod_{j=0}^{k-1} p_{u_j \wedge v_j, u_{j+1} \wedge v_{j+1}} \\ &\geq \prod_{j=0}^{k-1} p_{u_j, u_{j+1}} \times \prod_{j=0}^{k-1} p_{v_j, v_{j+1}} = \mathbf{P}_{X^k}(u^k)\mathbf{P}_{X^k}(v^k), \end{aligned}$$

where $u_0 = v_0 := x_0$, which finishes the proof. \square

Lemma 3.4. *Let $(E, \mathcal{E}, \preceq, \mathbf{P})$ be a partially ordered probability space, and assume that \mathbf{P} is purely atomic and satisfies the FKG condition. Let $A \in \mathcal{E}$ which satisfies (H2). Then, $\mathbf{P}(\cdot|A)$ satisfies the FKG condition.*

Proof. This is straightforward. For $u, v \in A$, one has $u \wedge v, u \vee v \in A$ by assumption. Hence,

$$\mathbf{P}(u \vee v|A)\mathbf{P}(u \wedge v|A) = \frac{1}{\mathbf{P}(A)^2}\mathbf{P}(u \vee v)\mathbf{P}(u \wedge v) \geq \frac{1}{\mathbf{P}(A)^2}\mathbf{P}(u)\mathbf{P}(v) = \mathbf{P}(u|A)\mathbf{P}(v|A),$$

for \mathbf{P} -a.e. $u, v \in A$, which finishes the proof. \square

Lemma 3.5. *Let \mathbf{P} a probability measure on a countable, locally finite subset of $(\mathbb{R}^n, \text{Bor}(\mathbb{R}^n), \preceq)$ which satisfies the FKG condition. Then it is associated.*

Proof. This is essentially the content of [24]. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}$ a non-negative, bounded and increasing function, and define the probability measure \mathbf{Q} on \mathbb{R}^n with

$$\frac{d\mathbf{Q}}{d\mathbf{P}}(u) = \frac{g(u)}{\mathbf{E}[g(u)]}, \quad \text{for } \mathbf{P}\text{-a.e. } u \in \mathbb{R}^n,$$

where $\mathbf{E} = \mathbf{E}_{\mathbf{P}}$ denotes the expectation with respect to \mathbf{P} (we assume that $\mathbf{E}[g(u)] > 0$, otherwise $g \equiv 0$ \mathbf{P} -a.e. and (1.1) holds for all f). Since \mathbf{P} satisfies (3.2), one clearly has for \mathbf{P} -a.e. $u, v \in \mathbb{R}^n$,

$$\mathbf{Q}(u \vee v)\mathbf{P}(u \wedge v) = g(u \vee v)\mathbf{P}(u \vee v)\mathbf{P}(u \wedge v) \geq g(u)\mathbf{P}(u)\mathbf{P}(v) = \mathbf{Q}(u)\mathbf{P}(v).$$

Therefore, it follows¹ from [24, Theorem 3] that for $f : \mathbb{R}^n \rightarrow \mathbb{R}$ a bounded increasing function, one has $\mathbf{E}_{\mathbf{Q}}[f] \geq \mathbf{E}_{\mathbf{P}}[f]$, which is exactly (1.1). Recalling Lemma 2.2, the inequality (1.1) also holds for all increasing $f, g \in L^2(\mathbf{P})$, finishing the proof. \square

¹Indeed, since $\text{Supp}(\mathbf{P})$ is locally finite, there exists a σ -finite counting measure μ on some set $A \subset \mathbb{R}$ such that $\text{Supp}(\mathbf{P}) \subset A^n$. Letting $\omega := \mu^{\otimes n}$ in [24], this yields the result (details are left to the reader).

With these lemmata at hand, the proof of Theorem 1.7 is immediate.

Proof of Theorem 1.7. Under the assumptions (H1, H2), Lemmata 3.3, 3.4 imply that the measure $\mathbf{P}_X(\cdot|A)$ satisfies the FKG condition. By Lemma 3.5, it is associated, completing the proof. \square

3.2. Proof of the corollaries. Let $T > 0$, $X^n = (X_1, \dots, X_n)$ a random vector in \mathbb{R}^n , and $X_0 := x_0 \in \mathbb{R}$. Similarly to (2.2), one can define the rcl (resp. continuous) interpolation of X in $\mathcal{D}([0, T], \mathbb{R})$ (resp. $\mathcal{C}([0, T], \mathbb{R})$) with, for $t \in [0, T]$,

$$(3.4) \quad \begin{aligned} \widehat{X}_t^n &:= \sum_{j=0}^n X_j \mathbf{1}_{[Tj/n, T(j+1)/n)}(t), \\ \text{resp. } \widetilde{X}_t^n &:= \sum_{j=0}^n [X_j + (\frac{nt}{T} - j)(X_{j+1} - X_j)] \mathbf{1}_{[Tj/n, T(j+1)/n)}(t). \end{aligned}$$

Both are increasing functions of X^n ; hence by Lemma 2.4, if the vector X^n is associated, then the processes \widehat{X}^n and \widetilde{X}^n are associated as well. Moreover By Lemma 2.1, if the sequence of processes \widehat{X}^n (or \widetilde{X}^n) converges weakly, then the limiting process is associated as well. Therefore, each corollary of Theorem 1.7 will be proven by determining a sequence of Markov chains X^n with transition kernel \mathbf{p}^n , and sets A_n in $\mathcal{E}_{\mathcal{D}}$ (or $\mathcal{E}_{\mathcal{C}}$) such that,

- \mathbf{p}^n satisfies (H1) for $n \geq 1$,
- A_n satisfies (H2) for $n \geq 1$,
- the sequence of processes \widehat{X}^n (or \widetilde{X}^n) conditioned on A_n , $n \geq 1$, converges weakly to the desired process.

Proof of Corollary 1.9. We prove the corollary simultaneously in the four cases. Recall Lemma 1.5.(i) and let Y^n be the first n steps of a random walk with discrete Laplace² increment for some $\beta > 0$. We may take $\beta > 0$ such that the Laplace increments have unit variance, and we let $X^n := n^{-1/2}Y^n$. Then for all $n \geq 1$, the transition kernel \mathbf{p} of the Markov chain X^n satisfies (H1).

We then define the sets $A_{j,n}$, $n \geq 1$ differently for each case $j \in \{1, 2, 3, 4\}$. Let,

$$(3.5) \quad A_j = A_{j,n} := \begin{cases} \{x \in \mathcal{D}([0, T], \mathbb{R}) \mid x_T = 0\} & \text{for } j = 1, \\ \{x \in \mathcal{D}([0, T], \mathbb{R}) \mid x_T = 0; \forall t, x_t \geq 0\} & \text{for } j = 2, \\ \{x \in \mathcal{D}([0, T], \mathbb{R}) \mid \forall t, x_t \geq 0\} & \text{for } j = 3, \\ \{x \in \mathcal{D}([0, T], \mathbb{R}) \mid \forall t, x_t \in [a(t), b(t)]\} & \text{for } j = 4, \end{cases}$$

Then, one clearly has in the four cases that $\mathbf{P}_{\widehat{X}^n}(A_j) > 0$ (also $\mathbf{P}_{\widetilde{X}^n}(A_j) > 0$), and that A is max/min-stable, so (H2) holds.

By Theorem 1.7, we deduce in each case $j \in \{1, 2, 3, 4\}$ that $\mathbf{P}_{\widehat{X}^n}(\cdot|A_j)$ (similarly $\mathbf{P}_{\widetilde{X}^n}(\cdot|A_j)$) is associated for all $n \in \mathbb{N}$. Therefore, it only remains to confirm in each case that the conditioned process converges weakly to the desired limit. Those results are classical: for the Brownian bridge $j = 1$, this is the content of [20]. For the Brownian excursion $j = 2$, this can be found in [13]. For the Brownian meander $j = 3$, we refer to [12]. Finally in the case $j = 4$, by assumption the event A_4 has positive probability for the Wiener measure, therefore the weak convergence can be deduced directly from Donsker's theorem (details are left to the reader). This concludes the proof of the corollary. \square

²Of course, one could also consider a simple random walk (recall Lemma 1.5.(ii)), provided that they restrict themselves to $n \in 2\mathbb{N}$ in the cases $j \in \{1, 2\}$ below.

Proof of Corollary 1.10. For $\nu \in \mathbb{R}$, we may define a Markov chain $(Y_k)_{k \geq 0}$ on \mathbb{N} with transition probabilities that satisfy,

$$p_{i,i+1} = 1 - p_{i,i-1} := \frac{1}{2} \left(1 + \frac{2\nu + 1}{2i} + o(i^{-1}) \right), \quad \text{if } i \geq 1,$$

$$p_{0,1} = 1,$$

and $p_{i,j} = 0$ otherwise. Since Y only makes unitary steps, by Lemma 1.5.(ii) its transition kernel satisfies (H1). For $n \geq 1$, letting $X^n := n^{-1/2}(Y_1, \dots, Y_n)$ and \tilde{X}^n its piecewise linear interpolation, one deduces from Theorem 1.7, Remark 1.1 and Lemma 2.4 that $\mathbf{P}_{\tilde{X}^n}$ is associated. Moreover when $\nu > -1$, the convergence of the process \tilde{X}^n to the Bessel process of index ν is proven in [17, Theorem 5.1]. By Lemma 2.1, this concludes the proof of the corollary. \square

4. SOME RANDOM WALK COUNTER-EXAMPLES TO THE FKG CONDITION

Recall the assumption (H1), and that it implies that X^k satisfies the FKG condition for all $k \leq n$ by Lemma 3.3. We now prove that reciprocal holds for (lattice) random walks.

Proposition 4.1. *Let $X^n = (X_1, \dots, X_n)$ the n first steps of a random walk on \mathbb{Z} started from some $x_0 \in \mathbb{Z}$, with increment probability measure denoted \mathbf{P} . Assume that (H1) does not hold, then there exists $m \in \mathbb{N}$ such that X^m does not satisfy the FKG condition (3.2).*

Proof. Recall the notation $u^k := (u_1, \dots, u_k)$. If (H1) fails, there exists $k \in \mathbb{N}$, $u_k \leq v_k$ in $\text{Supp}(X_k)$ and $u_{k+1} \geq v_{k+1}$ in \mathbb{Z} such that,

$$\mathbf{P}(u_{k+1} - u_k) \mathbf{P}(v_{k+1} - v_k) > \mathbf{P}(u_{k+1} - v_k) \mathbf{P}(v_{k+1} - u_k).$$

By assumption there exists two paths $u^k = (u_1, \dots, u_k)$ and $v^k = (v_1, \dots, v_k)$ such that $\mathbf{P}_{X^k}(u^k) > 0$, $\mathbf{P}_{X^k}(v^k) > 0$. For $j \geq k$, let us define by induction,

$$u_{j+1} := \begin{cases} u_j + (u_{k+1} - u_k) & \text{if } j - k \text{ is even,} \\ u_j + (v_{k+1} - v_k) & \text{if } j - k \text{ is odd,} \end{cases} \quad \text{and } v_{j+1} := \begin{cases} v_j + (v_{k+1} - v_k) & \text{if } j - k \text{ is even,} \\ v_j + (u_{k+1} - u_k) & \text{if } j - k \text{ is odd.} \end{cases}$$

Then, one has $u_j \leq v_j$ if $j - k$ is even, and $v_j \leq u_j$ if $j - k$ is odd. Therefore, one observes that

$$\frac{\mathbf{P}_{X^m}(u^m \wedge v^m) \mathbf{P}_{X^m}(u^m \vee v^m)}{\mathbf{P}_{X^m}(u^m) \mathbf{P}_{X^m}(v^m)} \leq C \left(\frac{\mathbf{P}(u_{k+1} - v_k) \mathbf{P}(v_{k+1} - u_k)}{\mathbf{P}(u_{k+1} - u_k) \mathbf{P}(v_{k+1} - v_k)} \right)^m,$$

for all $m \geq k$ and some $C > 0$. Taking m large enough, this concludes the proof. \square

Recalling Example 1.6, this provides a few examples of random walk distributions that do not satisfy the FKG condition. This contrasts with the fact that all random walks are associated, recall Proposition 2.3.

REFERENCES

- [1] D. Barbato. FKG inequality for brownian motion and stochastic differential equations. *Electronic Communications in Probability [electronic only]*, 10:7–16, 2005.
- [2] V. Belitsky, P. A. Ferrari, N. Konno, and T. M. Liggett. A strong correlation inequality for contact processes and oriented percolation. *Stochastic Processes and their Applications*, 67(2):213–225, 1997.
- [3] J. van den Berg, O. Häggström, and J. Kahn. Some conditional correlation inequalities for percolation and related processes. *Random Structures & Algorithms*, 29(4):417–435, 2006.
- [4] J. van den Berg and J. Kahn. A correlation inequality for connection events in percolation. *The Annals of Probability*, 29(1):123 – 126, 2001.

- [5] Y. Bröcker and C. Mukherjee. Localization of the Gaussian multiplicative chaos in the Wiener space and the stochastic heat equation in strong disorder. *Ann. Appl. Probab.*, 29(6):3745–3785, 2019.
- [6] L. Caffarelli. Monotonicity properties of optimal transportation and the FKG and related inequalities. *Communications in Mathematical Physics*, 214:547–563, 2000.
- [7] P. C. Fishburn, P. G. Doyle, and L. A. Shepp. The match set of a random permutation has the FKG property. *The Annals of Probability*, 16(3):1194 – 1214, 1988.
- [8] C. M. Fortuin, P. W. Kasteleyn, and J. Ginibre. Correlation inequalities on some partially ordered sets. *Comm. Math. Phys.*, 22:89–103, 1971.
- [9] S. Ganguly and A. Hammond. The geometry of near ground states in Gaussian polymer models. *Electron. J. Probab.*, 28:Paper No. 60, 80, 2023.
- [10] T. E. Harris. A lower bound for the critical probability in a certain percolation process. *Proc. Cambridge Philos. Soc.*, 56:13–20, 1960.
- [11] R. Holley. Remarks on the FKG inequalities. *Communications in Mathematical Physics*, 36(3):227 – 231, 1974.
- [12] D. L. Iglehart. Functional central limit theorems for random walks conditioned to stay positive. *Ann. Probability*, 2:608–619, 1974.
- [13] W. D. Kaigh. An invariance principle for random walk conditioned by a late return to zero. *Ann. Probability*, 4(1):115–121, 1976.
- [14] O. Kallenberg. *Foundations of modern probability*, volume 99 of *Probability Theory and Stochastic Modelling*. Springer, Cham, third edition, [2021] ©2021.
- [15] T. Kamae, U. Krengel, and G. L. O’Brien. Stochastic inequalities on partially ordered spaces. *Ann. Probability*, 5(6):899–912, 1977.
- [16] H. Kunita. Stochastic differential equations based on lévy processes and stochastic flows of diffeomorphisms. In M. M. Rao, editor, *Real and Stochastic Analysis: New Perspectives*, pages 305–373, Boston, MA, 2004. Birkhäuser Boston.
- [17] J. Lamperti. A new class of probability limit theorems. *J. Math. Mech.*, 11:749–772, 1962.
- [18] A. Legrand and N. Pétrelis. Surface transition in the collapsed phase of a self-interacting walk adsorbed along a hard wall. *Ann. Probab.*, 50(4):1538–1588, 2022.
- [19] T. Liggett. Conditional association and spin systems. *ALEA - Latin American journal of probability and mathematical statistics*, 1:1–19, 2006.
- [20] T. M. Liggett. An invariance principle for conditioned sums of independent random variables. *J. Math. Mech.*, 18:559–570, 1968.
- [21] T. M. Liggett. *Interacting particle systems*. Classics in Mathematics. Springer-Verlag, Berlin, 2005. Reprint of the 1985 original.
- [22] B. Mallein and P. Mił oś. Brownian motion and random walk above quenched random wall. *Ann. Inst. Henri Poincaré Probab. Stat.*, 54(4):1877–1916, 2018.
- [23] I. Mezić. FKG inequalities in cellular automata and coupled map lattices. *Physica D: Nonlinear Phenomena*, 103(1):491–504, 1997. Lattice Dynamics.
- [24] C. J. Preston. A generalization of the fkg inequalities. *Communications in Mathematical Physics*, 36:233–241, 1974.
- [25] E. Schertzer, R. Sun, and J. M. Swart. Stochastic flows in the Brownian web and net. *Mem. Amer. Math. Soc.*, 227(1065):vi+160, 2014.

UNIVERSITÉ CLAUDE BERNARD LYON 1, INSTITUT CAMILLE JORDAN, UMR 5208. 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE

Email address: legrand@math.univ-lyon1.fr