## Deriving friction force using fractional calculus

Georgii Koniukov

University of Montpellier, 34090 Montpellier, France

## Abstract

The friction force is derived using fractional calculus by considering the non-uniform flow of time in dissipative processes. The approach incorporates inhomogeneous velocity without unphysical approximations, resulting in a Lagrangian where the order of fractional derivatives measures time intervals. The fractional term in the Lagrangian provides correct Euler-Lagrange, and ultimately, the Hamilton equations, and vanishes during energy change measurements, like a ghost.

## CONTENTS

I.	Introduction	2
II.	The Riemann-Liouville and Caputo fractional calculus	3
III.	Conceptual basis for fractional friction forces	4
IV.	Lagrangian	6
V.	Hamilton's principle and derivation of the friction force	7
VI.	Hamilton equations	9
VII.	Energy change	10
VIII.	Summary and Concluding Remarks on the Physical Significance	11
	References	12

#### I. INTRODUCTION

Since the mature formulation of the action principle by Euler, Hamilton, and Lagrange, it has been established that the equations of motion for dissipative linear dynamical systems with constant coefficients cannot be derived from a variational principle. In 1873, Lord Rayleigh introduced the Rayleigh dissipation function to handle the effects of velocityproportional frictional forces in Lagrangian mechanics [1]. In this situation, two scalar functions are required to define the equation of motion for a single system, which is not a very natural approach. Some limitations were rigorously demonstrated by Bauer in 1931 [2], who proved the impossibility of obtaining a dissipation term proportional to the first-order time derivative from such a principle. Over the years, various methods have been developed to address this issue, including the Bateman approach with auxiliary coordinates [4] and time-dependent Lagrangians [3]. Unfortunately, these methods often result in non-physical Lagrangians that fail to provide meaningful relations for the system's momentum and Hamiltonian. In 1975 Dekker [5, 6] introduces an innovative approach utilizing a Lagrangian that produces two first-order equations as complex conjugates, which is logical given the frequent appearance of complex numbers in the context of dissipation. These equations can be combined to yield a real, second-order equation of motion. However, it is important to note that this method is specifically applicable to a damped oscillator.

In 1996-1997, Riewe discovered a gap in Bauer's proof, which originally assumed that all derivatives were of integer order. Riewe incorporated Riemann-Liouville fractional time derivatives into the Lagrangian [7], successfully deriving equations of motion that included non-conservative forces such as friction. While innovative due to the redefinition of the action principle, this work relied on non-physical approximations and assumptions, such as inappropriately changing the limits of integration in fractional derivatives, applying an unphysical limit where the observation time interval is artificially set to zero, and selecting the fractional derivative order as  $\frac{1}{2}$ .

In 2014, Lazo and Krumreich refined Riewe's approach by replacing the Riemann-Liouville derivative with Caputo's fractional derivative and utilizing the semigroup property of fractional integral operators [8], which made the calculations more straightforward. However, the physical interpretation of fractional derivatives remained unclear. Additionally, issues such as the inappropriate modification of integration limits in fractional derivatives, the unphysical time limit, and the selection of the fractional derivative order as  $\frac{1}{2}$  persisted.

Despite these advances, a unified and physically meaningful method to incorporate nonconservative forces within the framework of Lagrangian mechanics is still lacking. In this paper, we propose a novel approach that leverages the properties of fractional derivatives to derive nonconservative forces directly from a Lagrangian. Our method addresses the limitations of previous approaches by avoiding non-physical time limits and approximations, thereby providing a more natural and consistent description of dissipative systems. This new framework not only preserves the elegance of the variational principle but also enhances our understanding of the underlying physics in systems with friction and other nonconservative forces.

The structure of the paper is as follows: Section II introduces the necessary mathematical background on fractional calculus, focusing on the Riemann-Liouville and Caputo derivatives. Section III discusses the conceptual basis for modeling friction forces using fractional calculus. Section IV introduces the Lagrangian for systems with fractional dissipative forces. Section V presents the application of Hamilton's principle and derives the corresponding Euler-Lagrange equations. Section VI derives the Hamilton equations for the system. Section VII examines the changes in energy within the system. Finally, Section VIII discusses the physical implications of our findings and outlines potential directions for future research.

### II. THE RIEMANN-LIOUVILLE AND CAPUTO FRACTIONAL CALCULUS

All formal mathematical definitions in this section have been adopted from the paper by Lazo and Krumreich [8].

Let  $t_0, t_1, t_1 - t_0 \in \mathbb{R}_+$  and  $\rho(t) \in C^{[\alpha]+2}(\mathbb{R})$ , the notation [.] denotes the ceiling function, which returns the smallest integer greater than or equal to the given value.

**Definition II.1** (Integral operators).

$$_{t_0}I_t^{\alpha}\rho(t) = \frac{1}{\Gamma(\alpha)}\int_{t_0}^t \frac{\rho(\tau)}{(t-\tau)^{1-\alpha}}d\tau \qquad (left)$$
(1)

and

$${}_{t}I^{\alpha}_{t_{1}}\rho(t) = \frac{1}{\Gamma(\alpha)} \int_{t}^{t_{1}} \frac{\rho(\tau)}{(\tau-t)^{1-\alpha}} d\tau \qquad (right)$$

$$\tag{2}$$

**Definition II.2** (Riemann-Liouville fractional derivatives).

$${}_{t_0}D_t^{\alpha}\rho(t) = D_t^{[\alpha]+1} {}_{t_0}I_t^{1-\{\alpha\}}\rho(t) = \frac{d^{[\alpha]+1}}{dt^{[\alpha]+1}} \frac{1}{\Gamma(1-\{\alpha\})} \int_{t_0}^t \frac{\rho(\tau)}{(t-\tau)^{\{\alpha\}}} d\tau \quad (left)$$
(3)

and

$${}_{t}D_{t_{1}}^{\alpha}\rho(t) = (-1)^{[\alpha]+1}D_{t}^{[\alpha]+1} {}_{t}I_{t_{1}}^{1-\{\alpha\}}\rho(t) = \frac{d^{[\alpha]+1}}{dt^{[\alpha]+1}}\frac{(-1)^{[\alpha]+1}}{\Gamma\left(1-\{\alpha\}\right)}\int_{t}^{t_{1}}\frac{\rho(\tau)}{(\tau-t)^{\{\alpha\}}}d\tau \quad (right) \quad (4)$$

**Definition II.3** (Caputo fractional derivatives).

$${}_{t_0}^C D_t^{\alpha} \rho(t) = {}_{t_0} I_t^{1-\{\alpha\}} D_t^{[\alpha]+1} \rho(t) = \frac{1}{\Gamma(1-\{\alpha\})} \int_{t_0}^t \frac{\rho^{([\alpha]+1)}(\tau)}{(t-\tau)^{\{\alpha\}}} d\tau \qquad (left)$$
(5)

and

$${}_{t}^{C}D_{t_{1}}^{\alpha}\rho(t) = (-1)^{[\alpha]+1} {}_{t}I_{t_{1}}^{1-\{\alpha\}}D_{t}^{[\alpha]+1}\rho(t) = \frac{(-1)^{[\alpha]+1}}{\Gamma(1-\{\alpha\})} \int_{t}^{t_{1}} \frac{\rho^{([\alpha]+1)}(\tau)}{(\tau-t)^{\{\alpha\}}} d\tau \qquad (right)$$
(6)

**Remark II.3.1** (Relations between Riemann-Liouville and Caputo fractional derivatives).

$${}_{t_0}^C D_t^{\alpha} \rho(t) = {}_{t_0} D_t^{\alpha} \rho(t) - \sum_{k=0}^{[\alpha]} \frac{\rho^{(k)}(t_0)}{\Gamma(k-\alpha+1)} (t-t_0)^{k-\alpha}$$
(7)

and

$${}_{t}^{C}D_{t_{1}}^{\alpha}\rho(t) = {}_{t}D_{t_{1}}^{\alpha}\rho(t) - \sum_{k=0}^{[\alpha]} \frac{\rho^{(k)}(t_{1})}{\Gamma(k-\alpha+1)}(t_{1}-t)^{k-\alpha}$$
(8)

## **III. CONCEPTUAL BASIS FOR FRACTIONAL FRICTION FORCES**

According to the Noether's theorem, each symmetry in a physical system corresponds to a conservation law. In particular, the conservation of energy is a direct result of time invariance, meaning that the laws of physics are unchanged over time. This relationship is well-documented in classical mechanics and field theory, where the invariance of the Lagrangian with respect to time translations leads to energy conservation. Noether's seminal paper, "Invariante Variationsprobleme" [9] laid the groundwork for these fundamental concepts, establishing the deep connection between symmetries and conservation laws in physics. However, when considering dissipative systems where energy is not conserved, time invariance must be broken. This can be achieved by introducing time-dependent elements or making the system explicitly time-inhomogeneous.

Here by time inhomogeneity, we refer to introducing non-linearity in the time dependence of the system's parameters. By breaking time invariance, these models should describe the transfer and dissipation of energy within a system, providing a more comprehensive understanding of real-world physical processes where ideal conservation laws do not strictly apply.

Let's consider an evolution from time  $t_0$  until  $t_1$  in one-dimesional case. When time  $\tau$  is linear one may try to find the position q(t) at any moment t between  $t_0$  and  $t_1$  by integrating velocity v:

$$q(t) = \int_{t_0}^t v(\tau) d\tau + q(t_0)$$
(9)

Or

$$q(t) = q(t_1) - \int_{t}^{t_1} v(\tau) d\tau$$
(10)

For dissipative systems one may consider non-linearity of time  $(t - \tau)^{\alpha}$  (where t shows dependence on the translation of time). For the first try we will consider only  $0 < \alpha < 1$ . One may postulate as well inhomogeneous velocity  $\rho(\tau)$  to find the displacement  $\Delta q(t)$  by integrating such velocity  $\rho(\tau)$ :

$$q(t) \propto \int_{t_0}^t \rho(\tau) d(t-\tau)^{\alpha} + q(t_0)$$
 (11)

However, since the Hamilton principle is evaluated in terms  $d\tau$ , not  $d(t - \tau)^{\alpha}$ :

$$q(t) \propto \int_{t_0}^t \rho(\tau) d(t-\tau)^{\alpha} + q(t_0) = -\alpha \int_{t_0}^t \frac{\rho(\tau)}{(t-\tau)^{1-\alpha}} d\tau + q(t_0)$$
(12)

This reminds of Riemann-Liouville fractional integrals II.1. Then to have consistency with

their mathematical properties we will choose the constant  $\frac{1}{\Gamma(\alpha)}$  instead of  $-\alpha$ :

$$q(t) - q(t_0) \equiv {}_{t_0} I_t^{\alpha} \rho(t) = \frac{1}{\Gamma(\alpha)} \int_{t_0}^t \frac{\rho(\tau)}{(t-\tau)^{1-\alpha}} \mathrm{d}\tau$$
(13)

Since  $\rho(\tau)$  is inhomogeneous velocity, there could be an analogy with 10 for the  $\tau > t$ , and one might introduce as well ( $0 < \beta < 1$  for the first try):

$$q(t_1) - q(t) \equiv {}_t I_{t_1}^{\beta} \rho(t) = \frac{1}{\Gamma(\beta)} \int_t^{t_1} \frac{\rho(\tau)}{(\tau - t)^{1 - \beta}} d\tau$$
(14)

And, thus,

$$q(t) = {}_{t_0}I_t^{\alpha}\rho(t) + q(t_0) = q(t_1) - {}_{t}I_{t_1}^{\beta}\rho(t)$$
(15)

Let's remark on the consequences that should be true  $\forall t \in [t_0; t_1]$  VIII:

$$q(t_1) - q(t_0) = {}_{t_0} I_t^{\alpha} \rho(t) + {}_t I_{t_1}^{\beta} \rho(t)$$
(16)

The following sections are dedicated to the search for the values of  $\alpha$  and  $\beta$  V.

## IV. LAGRANGIAN

Since  $\rho(t)$  represents velocity, one may consider a Lagrangian where  $\rho^2(t)$  serves as the dissipative component of the kinetic energy (representing energy loss) and  $\frac{\gamma}{2}$  is an aspect ratio. We will adopt a quadratic form similar to Rayleigh [1]. The negative sign preceding the "kinetic" term resembles the ghost term introduced by Lee and Wick [11]. However, in the work by Riewe [7], this term is complex. All these points emphasize that this term is likely non-measurable.

$$L(t) = \frac{m\dot{q}^2(t)}{2} - \frac{\gamma\rho^2(t)}{2} - U(q)$$
(17)

There is no energy loss at the start of observation, therefore  $\rho(t_0) = 0$ . It is needed now to express  $\rho(t)$  in terms of q(t) to write the equations of motion later. As we defined in 13 (but with  $q(t_0) = 0$  as our frame of reference):

$$q(t) = \frac{1}{\Gamma(\alpha)} \int_{t_0}^t \frac{\rho(\tau)}{(t-\tau)^{1-\alpha}} \mathrm{d}\tau$$
(18)

Let's consider:

$${}_{t_0}^C D_t^{\alpha} q(t) = {}_{t_0}^C D_t^{\alpha} {}_{t_0} I_t^{\alpha} \rho(t) = {}_{t_0} I_t^{1-\{\alpha\}} D_t^{[\alpha]+1} {}_{t_0} I_t^{\alpha} \rho(t)$$
(19)

Since  $0 < \alpha < 1$ :

$${}_{t_0}I_t^{1-\{\alpha\}}D_t^{[\alpha]+1} {}_{t_0}I_t^{\alpha}\rho(t) = {}_{t_0}I_t^{1-\alpha}\frac{d}{dt} {}_{t_0}I_t^{\alpha}\rho(t)$$
(20)

According to 3:

$${}_{t_0}I_t^{1-\alpha}\frac{d}{dt} {}_{t_0}I_t^{\alpha}\rho(t) = {}_{t_0}I_t^{1-\alpha} {}_{t_0}D_t^{1-\alpha}\rho(t)$$
(21)

According to II.3.1 and absence of dissipative kinetic energy at the beginning of observation  $(\rho(t_0) = 0)$  we can change Riemann's fractional derivative of  $\rho(t)$  with Caputo's:

$${}_{t_0}I_t^{1-\alpha} {}_{t_0}D_t^{1-\alpha}\rho(t) = {}_{t_0}I_t^{1-\alpha} {}_{t_0}^C D_t^{1-\alpha}\rho(t) = {}_{t_0}I_t^{1-\alpha} {}_{t_0}I_t^{1-\{1-\alpha\}} D_t^{[1-\alpha]+1}\rho(t) = {}_{t_0}I_t^{1-\alpha} {}_{t_0}I_t^{\alpha} \frac{d}{dt}\rho(t)$$

$$(22)$$

According to the properties of Riemann-Liouville integrals [10] and the fact that  $\rho(t_0) = 0$ :

$${}_{t_0}I_t^{1-\alpha} {}_{t_0}I_t^{\alpha}\frac{d}{dt}\rho(t) = {}_{t_0}I_t^1\frac{d}{dt}\rho(t) = \rho(t)$$
(23)

Thus,

$$L(t) = \frac{m\dot{q}^{2}(t)}{2} - \frac{\gamma \left(\frac{C}{t_{0}}D_{t}^{\alpha}q(t)\right)^{2}}{2} - U(q)$$
(24)

#### V. HAMILTON'S PRINCIPLE AND DERIVATION OF THE FRICTION FORCE

In this chapter, we will derive the Euler-Lagrange equation for Hamilton's principle where Lagrangian 24 depends on the Caputo Dearivatve [12]. The action S is:

$$S = \int_{t_0}^{t_1} L\left(t, q(t), \dot{q}(t), \ _{t_0}^C D_t^{\alpha} q(t)\right) \mathrm{d}t$$
(25)

We will consider a case where we have initial condition  $q(t_0) = 0$ .

Let q(t) be the extremal function for the action 25. Then if we do the variation  $q \mapsto q + \varepsilon \eta$   $(\eta(t_0) = \eta(t_1) = 0; \ \dot{\eta}(t_0) = \dot{\eta}(t_1) = 0)$ . The condition for the extremum is obtained when the first Gâteaux variation is zero:

$$\delta S = \lim_{\epsilon \to 0} \frac{S[q + \epsilon \eta] - S[q]}{\epsilon} = \int_{t_0}^{t_1} \left[ \eta \frac{\partial L}{\partial q} + \dot{\eta} \frac{\partial L}{\partial \dot{q}} + {}^{C}_{t_0} D^{\alpha}_t \eta \cdot \frac{\partial L}{\partial {}^{C}_{t_0} D^{\alpha}_t q} \right] \mathrm{d}t = 0.$$
(26)

Let's integrate by parts term by term  $(\eta(t_0) = \eta(t_1) = 0)$ :

$$\int_{t_0}^{t_1} \dot{\eta} \frac{\partial L}{\partial \dot{q}} dt = \eta(t_1) \frac{\partial L}{\partial \dot{q}} \Big|_{t=t_1} - \int_{t_0}^{t_1} \eta \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} dt$$
(27)

The term with fractional derivative:

$$\int_{t_0}^{t_1} \frac{\partial L}{\partial t_0^C D_t^{\alpha} q} \cdot \int_{t_0}^C D_t^{\alpha} \eta \, \mathrm{d}t = \int_{t_0}^{t_1} \frac{\partial L}{\partial t_0^C D_t^{\alpha} q} \cdot \left[ \int_{t_0}^t \frac{\dot{\eta}(\tau) \mathrm{d}\tau}{(t-\tau)^{\alpha}} \right] \mathrm{d}t$$
(28)

We can change the order of integrals according to the integration domain [13] and then integrate by parts:



FIG. 1. Integration domain

$$= \int_{t_0}^{t_1} \dot{\eta}(\tau) \cdot \left[ \int_{\tau}^{t_1} \frac{\frac{\partial L}{\partial t_0 D_t^{\alpha} q} dt}{(t-\tau)^{\alpha}} \right] d\tau = \eta(\tau) \left[ \int_{\tau}^{t_1} \frac{\frac{\partial L}{\partial t_0 D_t^{\alpha} q} dt}{(t-\tau)^{\alpha}} \right] \Big|_{t_0}^{t_1} - \int_{t_0}^{t_1} \eta(\tau) \cdot \frac{d}{d\tau} \left[ \int_{\tau}^{t_1} \frac{\frac{\partial L}{\partial t_0 D_t^{\alpha} q} dt}{(t-\tau)^{\alpha}} \right] d\tau$$
(29)
Since  $\eta(t_0) = 0$ ,  $\int_{\tau}^{t_1} \frac{\frac{\partial L}{\partial t_0 D_t^{\alpha} q} dt}{(t-\tau)^{\alpha}} \Big|_{\tau=t_1} = 0$  and 3:

$$= \int_{t_0}^{t_1} \eta(\tau) \cdot {}_{\tau} D^{\alpha}_{t_1} \frac{\partial L}{\partial {}_{t_0}^C D^{\alpha}_t q} \mathrm{d}\tau$$
(30)

Thus,

$$0 = \delta S = \int_{t_0}^{t_1} \left[ \eta \frac{\partial L}{\partial q} - \eta \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} + \eta \cdot {}_t D^{\alpha}_{t_1} \frac{\partial L}{\partial {}_{t_0}^C D^{\alpha}_t q} \right] dt$$
(31)

And the Euler-Lagrange equation:

$$\frac{\partial L}{\partial q} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}} + {}_{t} D^{\alpha}_{t_1} \frac{\partial L}{\partial {}^{C}_{t_0} D^{\alpha}_{t} q} = 0$$
(32)

Let's consider the last term repeating the whole algorithm from IV:

$$\gamma {}_{t}D^{\alpha}_{t_{1}} {}^{C}_{t_{0}}D^{\alpha}_{t}q = \gamma {}_{t}D^{\alpha}_{t_{1}}\rho$$

$$(33)$$

Let's calculate (remembering that Caputo derivative of a constant  $q(t_1)$  is 0):

$${}_{t}^{C}D_{t_{1}}^{\beta}q(t) = {}_{t}^{C}D_{t_{1}}^{\beta}\left(q(t_{1}) - {}_{t}I_{t_{1}}^{\beta}\rho(t)\right) = -{}_{t}^{C}D_{t_{1}}^{\beta} {}_{t}I_{t_{1}}^{\beta}\rho(t)$$
(34)

Then by using II.3.1 (since  $_{t_1}I_{t_1}^{\beta}\rho = 0$ ):

$$= -{}_{t}D^{\beta}_{t_{1}} {}_{t}I^{\beta}_{t_{1}}\rho(t) = -\left(-\frac{d}{dt} {}_{t}I^{1-\beta}_{t_{1}} {}_{t}I^{\beta}_{t_{1}}\rho(t)\right)$$
(35)

Again we use the semigroup property of Riemann-Liouville integrals [10]:

$$= \frac{d}{dt} {}_{t}I^{1}_{t_{1}}\rho(t) = \frac{d}{dt}\frac{1}{\Gamma(1)}\int_{t}^{t_{1}}\frac{\rho(\tau)}{(\tau-t)^{0}}d\tau = \frac{d}{dt}\int_{t}^{t_{1}}\rho(\tau)d\tau = -\rho(t)$$
(36)

Thus, the equation 33 can be rewritten:

$$\gamma {}_{t}D^{\alpha}_{t_{1}}\rho = -\gamma {}_{t}D^{\alpha}_{t_{1}} {}_{t}^{C}D^{\beta}_{t_{1}}q(t) = -\gamma \left(-\frac{d}{dt} {}_{t}I^{1-\alpha}_{t_{1}}\left(-{}_{t}I^{1-\beta}_{t_{1}}\frac{d}{dt}\right)\right)q(t) = -\gamma \frac{d}{dt} {}_{t}I^{2-\alpha-\beta}_{t_{1}}\dot{q}(t)$$
(37)

Now, it is possible to see the important restriction on  $\alpha$  and  $\beta$ . Since the friction force should be proportional  $\dot{q}(t) \Rightarrow \alpha + \beta = 1$ :

$$= -\gamma \frac{d}{dt} {}_{t} I^{1}_{t_{1}} \dot{q}(t) = \gamma \dot{q}(t)$$
(38)

And the Euler-Lagrange equation:

$$m\ddot{q} + \gamma\dot{q} + \frac{\partial U(q)}{\partial q} = 0 \tag{39}$$

As a remark: adding  ${}_{t_0}^C D_t^{\alpha} q \cdot {}_t^C D_{t_1}^{1-\alpha}$  term to the Lagrangian does not change the equation of motion like adding  $\frac{df}{dt}$  where f is arbitrary.

## VI. HAMILTON EQUATIONS

To obtain a physically meaningful Hamiltonian and corresponding equations of motion, it is essential to perform a Legendre transformation. In this context, we will treat the fractional derivative of q as an independent projection, as it is not influenced by the velocity at the present time t, but rather by the velocities at previous times. Remembering 24:

$$L(q, q^{(\alpha)}, \dot{q}) = \frac{m\dot{q}^2}{2} - \frac{\gamma \left(q^{(\alpha)}\right)^2}{2} - U(q)$$
(40)

In order to make H not dependent on  $\dot{q}, q^{(\alpha)}$  we will define canonical  $p \equiv \frac{\partial L}{\partial \dot{q}}, p_{\alpha} \equiv \frac{\partial L}{\partial q^{(\alpha)}}$ :

$$H(p, p_{\alpha}, q) \equiv p\dot{q} + p_{\alpha}q^{(\alpha)} - L(q, q^{(\alpha)}, \dot{q})$$
(41)

$$dH = \dot{q}dp + q^{(\alpha)}dp_{\alpha} - \frac{\partial L}{\partial q}dq$$
(42)

Meanwhile:

$$dH = \frac{\partial H}{\partial p}dp + \frac{\partial H}{\partial p_{\alpha}}dp_{\alpha} + \frac{\partial H}{\partial q}dq$$
(43)

Thus, according to 32:

$$\frac{\partial H}{\partial p} = \dot{q}$$

$$\frac{\partial H}{\partial p_{\alpha}} = q^{(\alpha)}$$
(44)

$$\begin{pmatrix} \frac{\partial H}{\partial q} = -\frac{d}{dt}p + {}_{t}D^{\alpha}_{t_{1}}p_{\alpha} \\
H(p, p_{\alpha}, q) = \frac{p^{2}}{2m} - \frac{p^{2}_{\alpha}}{2\gamma} + U(q)$$
(45)

And, actually, it is now evident that inhomogeneous velocity 13 is a canonical momenta  $p_{\alpha}$ , and it, probably, should be  $\alpha$ -dependent VIII:

$$p_{\alpha}^2 = (\gamma \rho)^2 \tag{46}$$

### VII. ENERGY CHANGE

With the inclusion of the new dissipative term in the Lagrangian, we will observe that, in this case, the energy and the Hamiltonian do not coincide.

$$\frac{dH}{dt} = \frac{\partial H}{\partial p}\frac{dp}{dt} + \frac{\partial H}{\partial p_{\alpha}}\frac{dp_{\alpha}}{dt} + \frac{\partial H}{\partial q}\frac{dq}{dt} = \frac{\partial H}{\partial p}\frac{dp}{dt} + \frac{\partial H}{\partial p_{\alpha}}\frac{dp_{\alpha}}{dt} + \left(-\frac{dp}{dt} + {}_{t}D_{t_{1}}^{\alpha}p_{\alpha}\right)\frac{dq}{dt} = \frac{C}{t_{0}}D_{t}^{\alpha}q\cdot\dot{p_{\alpha}} + \dot{q}\cdot{}_{t}D_{t_{1}}^{\alpha}p_{\alpha} \quad (47)$$

Here we can define the anticommutator:

$$\frac{dH}{dt} = \{q, p_{\alpha}\} \equiv {}^{C}_{t_{0}}D^{\alpha}_{t}q \cdot \dot{p_{\alpha}} + \dot{q} \cdot {}_{t}D^{\alpha}_{t_{1}}p_{\alpha} = -\frac{p_{\alpha}}{\gamma}\dot{p_{\alpha}} + \dot{q}(-\gamma\dot{q})$$
(48)

On the other side we can introduce the energy E:

$$H(p, p_{\alpha}, q) = \frac{p^2}{2m} - \frac{p_{\alpha}^2}{2\gamma} + U(q) \equiv E - \frac{p_{\alpha}^2}{2\gamma}$$
(49)

$$\frac{dH}{dt} = \frac{dE}{dt} - \frac{p_{\alpha}}{\gamma} \dot{p_{\alpha}}$$
(50)

By comparing 48 and 50 we can obtain the energy dissipation law:

$$\frac{dE}{dt} = -\gamma \dot{q}^2 \tag{51}$$

# VIII. SUMMARY AND CONCLUDING REMARKS ON THE PHYSICAL SIG-NIFICANCE

- The concept of fractional derivatives lies in the non-uniform flow of time for dissipative processes, potentially offering a new understanding of the flow of time in physical processes. Philosophically, this can be understood as the different perceptions of time flow during energy expenditure/active engagement versus periods of inactivity.
- The use of Caputo fractional derivatives, both left and right, carries the same physical meaning; the difference lies in whether we determine the coordinate by knowing the future or the past. To define the Caputo fractional derivative, we integrate the velocity. The same applies to the usage of Riemann-Liouville derivatives, but we integrate the coordinate to define them.
- The dissipative term in the Lagrangian resembles "ghost term" [11] because it gives the correct Euler-Lagrange and Hamilton equations of motion, but disappears when we measure energy change.
- When calculating  $\frac{dH}{dt}$ , natural commutation relations between canonical generalized coordinates arise, caused by integration by parts of the terms with fractional derivatives in the action principle.
- The condition that the sum of  $\alpha + \beta = 1$ , probably, means that the order of the fractional derivative is a measure of the time intervals of the past and future, respectively.



FIG. 2. Derivative's fractional order as a measure of time interval

- Since we did not consider  $\alpha = \alpha(t)$  in the derivation, this, probably, means that for each time t nature recalculates the Hamilton principle with new fixed  $\alpha$ .
- It might be worth considering the Lagrangian as a function of the fractional derivative order  $L(0, \alpha, 1)$ , with boundary values at zero and one, representing kinetic and

potential energy, respectively.

- It is natural to continue the study of this method in quantum field cases and to rigorously define all concepts in the language of functional spaces and possibly interpret Planck's constant *i*ħ as a complex viscosity/damping coefficient, since it does not change the energy as the real viscosity coefficient γ, but stays with the first-order time derivative.
- Furthermore, we hope that the insights gained from this study could enhance our understanding of the Navier-Stokes equations, which govern the motion of fluid substances. These equations, fundamental to fluid dynamics, describe how the velocity field of a fluid evolves over time under the influence of various forces, including viscosity [14].
- J. W. Strutt, Some General Theorems relating to Vibrations, Proceedings of the London Mathematical Society, vol. s1-4, no. 1, pp. 357-368, 1871. doi: 10.1112/plms/s1-4.1.357.
- [2] P. S. Bauer, *Dissipative Dynamical Systems: I*, Proceedings of the National Academy of Sciences of the United States of America, vol. 17, no. 5, pp. 311-314, May 1931. doi: 10.1073/pnas.17.5.311. PMID: 16577367; PMCID: PMC1076051.
- K. W. H. Stevens, The Wave Mechanical Damped Harmonic Oscillator, Proceedings of the Physical Society, vol. 72, p. 1027, 1958.
- [4] H. Bateman, On Dissipative Systems and Related Variational Principles, Physical Review, vol. 38, pp. 815, August 1931.
- [5] H. Dekker, Classical and Quantum Mechanics of the Damped Harmonic Oscillator, Physics Reports (Review Section of Physics Letters), vol. 80, no. 1, pp. 1-112, 1981.
- [6] H. Dekker, On the Quantization of Dissipative Systems in the Lagrange-Hamilton Formalism, Zeitschrift f
  ür Physik B, vol. 21, pp. 295-300, 1975.
- [7] F. Riewe, Nonconservative Lagrangian and Hamiltonian mechanics, Physical Review E, vol. 53, no. 2, pp. 1890-1899, February 1996.
- [8] M. J. Lazo and C. E. Krumreich, The action principle for dissipative systems, Journal of Mathematical Physics, vol. 55, 122902, 2014. doi: 10.1063/1.4903991.

- [9] Emmy Noether, Invariante Variationsprobleme, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse
- [10] S. G. Samko, A. A. Kilbas, and O. I. Marichev, Fractional Integrals and Derivatives—Theory and Applications, Gordon and Breach, Linghorne, PA, 1993, Chapter 1, p. 34.
- [11] T. D. Lee and G. C. Wick, Negative metric and the unitarity of the S-matrix, Nuclear Physics
   B, vol. 9, no. 2, pp. 209-243, Feb. 1969. doi: 10.1016/0550-3213(69)90098-4.
- [12] O. P. Agrawal, Generalized Euler Lagrange Equations and Transversality Conditions for FVPs in terms of the Caputo Derivative, Journal of Vibration and Control, vol. 13, no. 9-10, pp. 1217-1237, September 2007.
- [13] E. R. Love and L. C. Young, On Fractional Integration by Parts, Proceedings of the London Mathematical Society, [Received 26 April, 1937.—Read 20 May, 1937].
- [14] G. G. Stokes, On the Theories of the Internal Friction of Fluids in Motion, and of the Equilibrium and Motion of Elastic Solids, Mathematical and Physical Papers by George Gabriel Stokes, Cambridge University Press, pp. 75-129, 1880.