Well-conditioned Primal-Dual Interior-point Method for Accurate Low-rank Semidefinite Programming

Hong-Ming Chiu University of Illinois Urbana-Champaign hmchiu2@illinois.edu Richard Y. Zhang University of Illinois Urbana-Champaign ryz@illinois.edu

Abstract

We describe how the low-rank structure in an SDP can be exploited to reduce the periteration cost of a convex primal-dual interior-point method down to $O(n^3)$ time and $O(n^2)$ memory, even at very high accuracies. A traditional difficulty is the dense Newton subproblem at each iteration, which becomes progressively ill-conditioned as progress is made towards the solution. Preconditioners have previously been proposed to improve conditioning, but these can be expensive to set up, and become ineffective as the preconditioner itself becomes increasingly ill-conditioned at high accuracies. Instead, we present a *well-conditioned reformulation* of the Newton subproblem that is cheap to set up, and whose condition number is guaranteed to remain bounded over all iterations of the interior-point method. In theory, applying an inner iterative method to the reformulation reduces the per-iteration cost of the outer interior-point method to $O(n^3)$ time and $O(n^2)$ memory. We also present a *well-conditioned preconditioner* that theoretically increases the outer per-iteration cost to $O(n^3r^3)$ time and $O(n^2r^2)$ memory, where r is an upper-bound on the solution rank, but in practice greatly improves the convergence of the inner iterations.

1 Introduction

Given problem data $\mathcal{A}: \mathcal{S}^n \to \mathbb{R}^m$ and $b \in \mathbb{R}^m$ and $C \in \mathcal{S}^n$, we seek to solve the standard-form semidefinite program (SDP)

$$X^{\star} = \arg\min_{X \in \mathcal{S}^n} \{ \langle C, X \rangle : \mathcal{A}(X) = b, \quad X \succeq 0 \},$$
(1a)

to high accuracy, under the assumption of a unique low-rank solution

$$X^{\star}$$
 is unique, $r^{\star} \stackrel{\text{def}}{=} \operatorname{rank}(X^{\star}) \ll n.$ (1b)

Here, S^n denotes the set of $n \times n$ real symmetric matrices with inner product $\langle A, B \rangle = \text{tr}(AB)$, and we write $A \succeq B$ to mean that A - B is positive semidefinite. Without loss of generality, we assume that the constraints in \mathcal{A} are linearly independent, meaning that $\mathcal{A}^T(y) = 0$ holds if and only if y = 0.

Currently, the low-rank SDP (1) is most commonly approached using the nonconvex low-rank factorization of Burer and Monteiro [5]. The basic idea is to factor $X = UU^T$ into an $n \times r$ low-rank factor U, in which $r \geq r^*$, and then to locally optimize over U. The dramatic reduction in the number of variables, from $O(n^2)$ to as few as O(n), allows the approach to achieve state-of-the-art results across many applications [3, 28, 12]. Unfortunately, the nonconvexity of the factorization

can make the approach surprisingly ineffective elsewhere. As a motivating example, consider the ℓ_1 linear regression problem with a trace regularizer [9, 13, 21],

$$\min_{X \succeq 0} \|\mathcal{A}(X) - b\|_{\ell_1} + \lambda \cdot \operatorname{tr}(X), \tag{2}$$

which can be recognized as an instance of (1) by rewriting

$$\min_{v,w\in\mathbb{R}^m,X\in\mathcal{S}^n}\left\{\mathbf{1}^T v + \mathbf{1}^T w + \lambda \cdot \operatorname{tr}(X): \begin{array}{c} v - w + \mathcal{A}(X) = b, \\ v \ge 0, w \ge 0, X \succeq 0. \end{array}\right\}$$
(3)

The Burer-Monteiro approach can be applied, by substituting $X = UU^T$ into either the original formulation (2) or the SDP reformulation (3). But in both cases, local optimization can struggle to converge, because the nonconvexity of the factorization exacerbates the problem's underlying nonsmoothness. To our best knowledge, no algorithm can consistently solve a generic instance of (2) to high accuracy, particularly when the number of equations scales as $m = O(n^2)$ with respect to the matrix order n.




Figure 1: Reconstruction error $||X - X^*||_F$ v.s. runtime for solving an instance of (2) using our proposed method (our method); spectrally preconditioned interior-point method by Zhang and Lavaei [39] (ZL17); nonlinear programming solvers fmincon [23] and knitro [6]; and first-order subgradient method method (GD). (See Section 7.2 for details.)

In this paper, we revisit classical primal-dual interior-point methods, because they maintain the convexity of the SDP (1), and therefore enjoy simple and rapid global convergence to ϵ accuracy in $O(\sqrt{n}\log(1/\epsilon))$ iterations, irrespective of any underlying nonsmoothness. Our contribution is to use the low-rank structure to reduce the per-iteration cost of the interior-point method to $O(n^3)$ time and $O(n^2)$ memory. As we explain below, the main difficulty is to be able to maintain these figures even at high accuracies with very small values of μ .

1.1 Newton subproblem

A perennial difficulty for interior-point methods is the expensive Newton subproblem that must be solved at each iteration. In its simplest form, an interior-point method works by applying Newton's method (with fixed or adaptive stepsize) to the logarithmic penalized SDP and its dual

$$\begin{aligned} X(\mu) &= \arg\min_{X\in\mathcal{S}^n} \{ \langle C, X \rangle - \mu \log \det(X) : \mathcal{A}(X) = b \}, \\ y(\mu), S(\mu) &= \arg\max_{S\in\mathcal{S}^n, y\in\mathbb{R}^m} \{ \langle b, y \rangle + \mu \log \det(S) : \mathcal{A}^T(y) + S = C \}, \end{aligned}$$

while progressively decreasing the duality gap parameter $\mu > 0$ after each iteration, noting that (1) is recovered in the limit $\mu = 0^+$. At each iteration, Newton's method computes its search direction $(\Delta X, \Delta y, \Delta S)$ by solving the least-squares problem

$$\min_{\Delta X \in \mathcal{S}^n} \left\{ \left\langle \tilde{C}, \Delta X \right\rangle + \frac{1}{2} \| W^{-1/2}(\Delta X) W^{-1/2} \|_F^2 : \mathcal{A}(\Delta X) = \tilde{b} \right\},\tag{4a}$$

$$\max_{\Delta S \in \mathcal{S}^n, \Delta y \in \mathbb{R}^m} \left\{ \left\langle \tilde{b}, \Delta y \right\rangle - \frac{1}{2} \| W^{1/2}(\Delta S) W^{1/2} \|_F^2 : \mathcal{A}^T(\Delta y) + \Delta S = \tilde{C} \right\},\tag{4b}$$

in which the scaling point $W \in S_{++}^n$ and the residuals $\tilde{b} \in \mathbb{R}^m$, $\tilde{C} \in S^n$ are iteration- and algorithmdependent. In this paper, we will focus on the primal-dual Nesterov–Todd scaling [25, 26], which is computed from the current iterates X, S as follows

$$W = X^{1/2} (X^{1/2} S X^{1/2})^{-1/2} X^{1/2} = [S^{1/2} (S^{1/2} X S^{1/2})^{-1/2} S^{1/2}]^{-1},$$
(4c)

and is used in popular solvers like SeDuMi [30], SDTP3 [34], and MOSEK [24].

Among all existing interior-point methods, a common critical issue is that the scaling point $W \in S_{++}^n$ becomes *dense* and *ill-conditioned* with decreasing values of μ . While (4) can be solved directly, the density of W renders its Karush–Kuhn–Tucker (KKT) equations dense and costly to solve, even when the data C and \mathcal{A} are themselves highly sparse. The $\Theta(n^3m + m^2n^2 + m^3)$ time and $\Theta(n^2m + m^2)$ memory it takes to form and factor these dense equations limit m and n to moderate values.

Alternatively, (4) can be solved iteratively, noting that if the data C and \mathcal{A} are suitably sparse, then each matrix-vector product with (4) can be evaluated in just $O(n^3)$ time. Numerically exact methods iterate until reaching the numerical floor, hence yielding a search direction that is indistinguishable from the true solution of (4) under finite precision [35], but the number of iterations needed for numerical exactness can grow as $O(1/\mu)$. Inexact methods truncate after a much smaller number of iterations [32], but the inaccurate search direction that result can slow the convergence of the outer interior-point method. In the end, all existing interior-point methods face a steep costaccuracy tradeoff: one can pay as much $O(n^6)$ time per-iteration to achieve very high accuracy of $\mu \approx 10^{-12}$, or one can accept coarse accuracy of $\mu \approx 10^{-2}$ to reduce costs to as low as $O(n^3)$ time per-iteration.

1.2 Prior work: Spectral preconditioning

Toh and Kojima [33] first made the critical observation that the scaling point $W \in S_{++}^n$ becomes ill-conditioned as $\operatorname{cond}(W) = \Theta(1/\mu)$ only because its eigenvalues split into two clusters,

$$W = \underbrace{Q\Lambda Q^T}_{\text{top } r \text{ eigenvalues}} + \underbrace{Q_{\perp}\Lambda_{\perp}Q_{\perp}^T}_{\text{bottom } n-r \text{ eigenvalues}}$$

with $\Lambda = \Theta(1/\sqrt{\mu})$ and $\Lambda_{\perp} = \Theta(\sqrt{\mu})$, but that each cluster remained individually well-conditioned, with cond(Λ) = $\Theta(1)$ and cond(Λ_{\perp}) = $\Theta(1)$, under strict complementarity and nondegeneracy assumptions. Based on this insight, they formulated a spectral preconditioner to counteract the illconditioning between the two clusters, and proved that the resulting preconditioned method solves (4) to ϵ accuracy in $O(\log(1/\epsilon))$ iterations for all $0 < \mu \leq 1$. While extremely effective, the cost of setting up the preconditioner is unfortunately very high, and this made the overall solution time comparable to a direct solution of (4).

Inspired by the above, Zhang and Lavaei [39] proposed a spectral preconditioner that is much cheaper to set up. Their key idea is to rewrite the scaling point as the low-rank perturbation of a well-conditioned matrix, and then to approximate the well-conditioned part by the identity matrix

$$W = \underbrace{Q(\Lambda - \tau)Q^{T}}_{\text{low-rank}} + \tau \cdot \underbrace{(QQ^{T} + \tau^{-1}Q_{\perp}\Lambda_{\perp}Q_{\perp}^{T})}_{\text{well-conditioned}} \approx UU^{T} + \tau I$$

where $\tau \approx \|\Lambda_{\perp}\|$. Substituting $W \approx UU^T + \tau I$ into (4) reveals a low-rank perturbed problem that admits a closed-form solution via the Sherman–Morrison–Woodbury formula, which they proposed to use as a preconditioner. They proved that the preconditioner costs $O(n^3r^3)$ time and $O(n^2r^2)$ to set up, assuming, for $U, V \in \mathbb{R}^{n \times r}$ and $y \in \mathbb{R}^m$, that the two matrix-vector products

$$(U,V) \mapsto \mathcal{A}(UV^T + VU^T) = \left[\left\langle A_i, UV^T + VU^T \right\rangle \right]_{i=1}^m$$
$$(U,y) \mapsto \left[\mathcal{A}^T(y) \right] U = \sum_{i=1}^m y_i A_i U$$

can both be performed in $O(n^2r)$ time and $O(n^2)$ storage. Under the same strict complementarity and nondegeneracy assumptions as Toh and Kojima [33], they proved that the resulting preconditioned method solves (4) to ϵ accuracy in $O(\log(1/\epsilon))$ iterations for all $0 < \mu \leq 1$. Running the iterations to the numerical floor yields a numerically exact search direction in $O(n^3r^3)$ time and $O(n^2r^2)$ memory. This same preconditioning idea was also used in the solver Loraine [17].

While fast, a critical weakness of the Zhang–Lavaei preconditioner is that it abruptly stops working once the outer interior-point method reaches higher accuracies. This is an inherent limitation; to precondition a matrix with a diverging condition number $\Theta(1/\mu^2)$, the preconditioner's condition number must also diverge as $\Theta(1/\mu^2)$. As μ approaches zero, it becomes impossible to solve the preconditioner accurately enough for it to remain effective. This issue is further exacerbated by the use of the Sherman–Morrison–Woodbury formula, which is itself notorious for numerical instability. In practice, the Zhang–Lavaei preconditioner limits the outer interior-point method to medium accuracies of $\mu \approx 10^{-6}$.

1.3 This work: Well-conditioned reformulation and preconditioner

In this paper, we derive a well-conditioned reformulation of the Newton subproblem (4) that costs just $O(n^3)$ time and $O(n^2)$ memory to set up. In principle, as the reformulation is already wellconditioned by construction, it can be solved using any iterative method to ϵ accuracy in $O(\log(1/\epsilon))$ iterations. Running these iterations to the numerical floor yields a numerically exact solution in $O(n^3)$ time and $O(n^2)$ memory, even as the scaling point W itself becomes arbitrarily ill-conditioned with $\mu \to 0^+$. In practice, the convergence rate can be substantially improved by the use of a wellconditioned preconditioner. We prove, under the same fast matrix-vector product assumption as Zhang and Lavaei, that the resulting preconditioned iterations also compute a numerically exact solution in $O(n^3r^3)$ time and $O(n^2r^2)$ memory. But unlike Zhang and Lavaei, our well-conditioned preconditioner to work even as the outer interior-point method reaches extremely high accuracies like $\mu \approx 10^{-12}$. Our main theoretical result is a rigorous guarantee that the reformulation and the preconditioner both remain well-conditioned for all $0 < \mu \leq 1$. The result hinges on two critical assumptions, that can be interpreted as more precise versions of the *strict complementarity* and *nondegeneracy* assumptions adopted by prior work. First, we assume that primal-dual iterates X, S used to compute the scaling W in (4) satisfy the following

$$\left\|\frac{1}{\mu}X^{1/2}SX^{1/2} - I\right\| < 1, \qquad \|X - X^*\| = O(\mu), \qquad \|S - S^*\| = O(1).$$
(A1)

Indeed, (A1) is exactly equivalent to strict complementarity $X^* + S^* \succ 0$ if the iterates are kept perfectly centered $\frac{1}{\mu}X^{1/2}SX^{1/2} = I$ [20]. Therefore, (A1) is a stronger version of strict complementarity that additionally requires the solver to keep its iterates X, S sufficiently centered. In this paper, we omit the rigorous details on how this is actually achieved, and instead verify experimentally that the popular interior-point method SeDuMi [30] satisfies the assumption. Our experiments suggest that the assumption is mild and holds generically; we defer a rigorous justification to future work.

Second, we assume that the linear operator \mathcal{A} is injective with respect to the tangent space of the manifold of rank- r^* positive semidefinite matrices, evaluated at the unique primal solution $X^* = U^*U^{*T}$:

$$\mathcal{A}(U^{\star}V^{T} + VU^{\star T}) = 0 \quad \Longleftrightarrow \quad U^{\star}V^{T} + VU^{\star T} = 0.$$
(A2)

This is a stronger version of the dual nondegeneracy condition of Alizadeh, Haeberly, and Overton [1], which under (A1), is equivalent to uniqueness of the primal solution X^* . It is stronger because it is possible for a rank- r^* primal solution to be unique with just $m = \frac{1}{2}r^*(r^* + 1)$ constraints, but (A2) can hold only when the number of constraints m is no less than the dimension of the tangent space

$$m \ge \frac{1}{2}r^{\star}(r^{\star}+1) + r^{\star}(n-r^{\star}) = nr^{\star} - \frac{1}{2}r^{\star}(r^{\star}-1).$$

Fortunately, many rigorous proofs of primal uniqueness, such as for matrix sensing [27, 10], matrix completion [8], phase retrieval [11], actually work by establishing (A2).

1.4 Related work

Our well-conditioned reformulation is inspired by Toh [31]. Indeed, the relationship between our reformulation and the Zhang–Lavaei preconditioner [39] can be viewed as analogous to Toh's reformulation with respect to the Toh–Kojima preconditioner [33]. Compared to prior work, we are the first to elucidate a precise, experimentally-verified, numerical prediction of the condition number, based on quantifying the same assumptions of strict complementarity and nondegeneracy used in prior work. In elucidating this dependence, we find that primal nondegeneracy is not actually needed for well-conditioning to hold. However, conditioning is significantly improved by dual nondegeneracy, which roughly corresponds to a steeper and more well-defined unique primal solution X^* that is much better than nearby points.

We mention that first-order methods have also been derived to exploit the low-rank structure of the SDP (1) without giving up the convexity [7, 37, 36]. These methods gain their efficiency by keeping all their iterates low-rank, and by performing convex updates on the low-rank iterates $X = UU^T$ while maintaining them in low-rank factored form $X_+ = U_+U_+^T$. While convex first-order methods cannot become stuck at a spurious local minimum, they still require poly $(1/\epsilon)$ iterations to converge to an ϵ -accurate solution of (1). In contrast, our proposed method achieves the same in just $O(\log(1/\epsilon))$ iterations, for an exponential factor improvement. Finally, for low-rank SDPs with small treewidth sparsity, chordal conversion methods [14, 18] compute an ϵ accurate solution in guaranteed $O(n^{1.5} \log(1/\epsilon))$ time and O(n) memory [40, 38]. Where the property holds, chordal conversion methods achieve state-of-the-art speed and reliability. Unfortunately, many real-world problems do not enjoy small treewidth sparsity [22]. For low-rank SDPs that are sparse but non-chordal, our proposed method can solve them in as little as $O(n^{3.5} \log(1/\epsilon))$ time and $O(n^2)$ memory.

Notations

Write $\mathbb{R}^{n \times n} \supseteq S^n \supseteq S^n_+ \supseteq S^n_{++}$ respectively as the sets of $n \times n$ real matrices, real symmetric matrices, positive semidefinite matrices, and positive definite matrices. Write vec : $\mathbb{R}^{n \times n} \to \mathbb{R}^{n^2}$ as the usual column-stacking vectorization, and $A \otimes B$ as the usual Kronecker product satisfying the identity $(B \otimes A) \operatorname{vec}(X) = \operatorname{vec}(AXB^T)$. Write $\|\cdot\|$ and $\|\cdot\|_F$ as spectral norm and Frobenius norm, respectively.

Let Ψ_n denote the $n^2 \times n(n+1)/2$ basis for the set of vectorized real symmetric matrices $\operatorname{vec}(\mathcal{S}^n) \subseteq \operatorname{vec}(\mathbb{R}^{n \times n})$. Concretely, we have

$$\Psi_n^T \operatorname{vec}(X) = (X_{1,1}, \sqrt{2}X_{2,1}, \dots, \sqrt{2}X_{n,1}, X_{2,2}, \sqrt{2}X_{3,2}, \dots, \sqrt{2}X_{n,2}, \dots, X_{n,n})$$

and $\Psi_n \Psi_n^T \operatorname{vec}(X) = \operatorname{vec}(X)$ for all $X \in S^n$. Like the identity matrix, we will frequently suppress the subscript *n* if the dimensions can be inferred from context. Using the basis matrix, we define the symmetric vectorization of X,

$$\operatorname{vec}_{\mathcal{S}} : \operatorname{vec}(\mathcal{S}^n) \to \mathbb{R}^{n(n+1)/2}, \qquad \operatorname{vec}_{\mathcal{S}}(X) \stackrel{\text{def}}{=} \Psi_n^T \operatorname{vec}(X),$$

and note that by $\operatorname{vec}(X) = \Psi_n \operatorname{vec}_{\mathcal{S}}(X)$, it follows that

$$\langle X, S \rangle = \langle \operatorname{vec}_{\mathcal{S}}(X), \operatorname{vec}_{\mathcal{S}}(S) \rangle = \langle \operatorname{vec}(X), \operatorname{vec}(S) \rangle$$
 for all $X, S \in \mathcal{S}^n$

Given $A, B \in \mathbb{R}^{n \times r}$, we define their symmetric Kronecker product as

$$A \otimes_{\mathcal{S}} B = B \otimes_{\mathcal{S}} A \stackrel{\text{def}}{=} \frac{1}{2} \Psi_n^T (A \otimes B + B \otimes A) \Psi_r$$

so that $(B \otimes_{\mathcal{S}} A) \operatorname{vec}_{\mathcal{S}}(X) = \operatorname{vec}_{\mathcal{S}}(AXB^T)$ holds for all $X \in \mathcal{S}^r$. (See also [1, 33, 31] for earlier references on the symmetric vectorization and Kronecker product.)

2 Preliminaries

2.1 MINRES

Given a system of equations Ax = b with a symmetric A and right-hand side b, we define the corresponding minimum residual (MINRES) iterations with respect to preconditioner P, and initial point x_0 implicitly in terms of its Krylov optimality condition.

Definition 2.1 (Minimum residual). Given $A \in S^n$ and $P \in S^n_{++}$ and $b, x_0 \in \mathbb{R}^n$, we define MINRES_k $(A, b, P, x_0) \stackrel{\text{def}}{=} P^{1/2} y_k$ where

$$y_k = \arg\min_{y} \{ \|\tilde{A}y - \tilde{b}\| : y \in y_0 + \operatorname{span}\{\tilde{b}, \tilde{A}\tilde{b}, \dots, \tilde{A}^{k-1}\tilde{b}\} \}$$

and $\tilde{A} = P^{-1/2}AP^{-1/2}$ and $\tilde{b} = P^{-1/2}b$ and $y_0 = P^{-1/2}x_0$.

The precise implementation details of MINRES are complicated, and can be found in the standard reference [16, Algorithm 4]. We only note that the method uses O(n) memory over all iterations, and that each iteration costs O(n) time plus the the matrix-vector product $p \mapsto Ap$ and the preconditioner linear solve $r \mapsto P^{-1}r$. In exact arithmetic, MINRES terminates with the exact solution in *n* iterations. With round-off noise, however, exact termination does not occur. Instead, the behavior of the iterates are better predicted by the following bound.

Proposition 2.2 (Symmetric indefinite). Given $A \in S^n$ and $P \in S^n_{++}$ and $b, x_0 \in \mathbb{R}^n$, let $x_k = MINRES_k(A, b, P, x_0)$. Then, we have $||Ax_k - b|| \le \epsilon$ in at most

$$k \le \left\lceil \frac{1}{2} \kappa \log \left(\frac{2 \|Ax^0 - b\|}{\epsilon} \right) \right\rceil \text{ iterations}$$

where $\kappa = \text{cond}(P^{-1/2}AP^{-1/2}).$

Our main contribution is to propose a well-conditioned reformulation Ax = b for the Newton subproblem (4), whose condition number $\kappa = \text{cond}(A)$ remains bounded over all iterations of the interior-point method. Therefore, applying Proposition 2.2 shows that it takes $O(\log(1/\epsilon))$ iterations to solve this reformulation to ϵ accuracy.

2.2 PCG

Given a system of equations Ax = b with a symmetric A and right-hand side b, we define the corresponding preconditioned conjugate gradient (PCG) iterations with respect to preconditioner P, and initial point x_0 explicitly in terms of its iterations.

Definition 2.3 (Preconditioned conjugate gradients). Given $A, P \in S^n$ and $b, x_0 \in \mathbb{R}^n$, we define $PCG_k(A, b, P, x_0) \stackrel{\text{def}}{=} x_k$ where for $j \in \{0, 1, 2, ..., k\}$

$$\begin{aligned} x_{j+1} &= x_j + \alpha_j p_j, \quad P z_{j+1} = r_{j+1} = r_j - \alpha_j A p_j, \quad p_{j+1} = z_{j+1} + \beta_j p_j, \\ \alpha_j &= \langle r_i, z_j \rangle / \langle p_j, A p_j \rangle, \qquad \beta_j = \langle r_{j+1}, z_j \rangle / \langle r_j, z_j \rangle, \end{aligned}$$

and $Pz_0 = r_0 = b - Ax_0$ and $p_0 = z_0$.

We can verify from Definition 2.3 that the method uses O(n) memory over all iterations, and that each iteration costs O(n) time plus the the matrix-vector product $p \mapsto Ap$ and the preconditioner linear solve $r \mapsto P^{-1}r$. The following is a classical iteration bound for when PCG is used to solve a symmetric positive definite system of equations with a positive definite preconditioner.

Proposition 2.4 (Symmetric positive definite). Given $A, P \in S_{++}^n$ and $b, x_0 \in \mathbb{R}^n$, let $x_k = \text{PCG}_k(A, b, P, x_0)$. Then, both $||Ax_k - b|| \leq \epsilon$ holds in at most

$$k \le \left\lceil \frac{1}{2} \sqrt{\kappa} \log \left(\frac{2\sqrt{\kappa} \|Ax_0 - b\|}{\epsilon} \right) \right\rceil \text{ iterations}$$

where $\kappa = \text{cond}(P^{-1/2}AP^{-1/2}).$

In the late 1990s and early 2000s, it was pointed out in several parallel works [19, 15, 29] that PCG can also be used to solve *indefinite* systems, with the help of an indefinite preconditioner. The proof of Proposition 2.5 follows immediately by substituting into Definition 2.3.

Proposition 2.5 (Indefinite preconditioning). Given $A, P \in S_{++}^n$ and $C \in S_{++}^r$ and $B \in \mathbb{R}^{n \times r}$ and $x_0, f \in \mathbb{R}^n$ and $g \in \mathbb{R}^r$, let

$$\begin{bmatrix} u_k \\ v_k \end{bmatrix} = \operatorname{PCG}_k \left(\begin{bmatrix} A & B \\ B^T & -C \end{bmatrix}, \begin{bmatrix} f \\ g \end{bmatrix}, \begin{bmatrix} P & B \\ B^T & -C \end{bmatrix}, \begin{bmatrix} x_0 \\ C^{-1}(B^T x_0 - g) \end{bmatrix} \right),$$
(5a)

$$x_k = \text{PCG}_k(A + BC^{-1}B^T, f + BC^{-1}g, P + BC^{-1}B^T, x_0).$$
(5b)

In exact arithmetic, we have $u_k = x_k$ and $v_k = C^{-1}(B^T x_k - g)$ for all $k \ge 0$.

Therefore, the indefinite preconditioned PCG (5a) is guaranteed to converge because it is mathematically equivalent to PCG on the underlying positive definite Schur complement system (5b). Nevertheless, (5a) is more preferable when the matrix C is close to singular, because the two indefinite matrices in (5a) can remain well-conditioned even as $\operatorname{cond}(C) \to \infty$, but their two Schur complements in (5b) must become increasingly ill-conditioned. Once their condition numbers approach $\approx 10^{12}$, the preconditioning effect will abruptly fail in double-precision arithmetic. Also, even if $u_k = x_k$ if computed to high accuracy, (5b) would still require recovering $v_k = C^{-1}(B^T x_k - g)$ by inverting the near-singular matrix C.

3 Proposed method and summary of results

Given scaling point $W \in S_{++}^n$ and residuals $\tilde{b} \in \mathbb{R}^m$, $\tilde{C} \in S^n$, our goal is to compute $\Delta X \in S^n$ and $\Delta y \in \mathbb{R}^m$ in closed-form via the Karush–Kuhn–Tucker equations

$$\left\| \begin{bmatrix} -(W \otimes_{\mathcal{S}} W)^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \end{bmatrix} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(\Delta X) \\ \Delta y \end{bmatrix} - \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(\tilde{C}) \\ \tilde{b} \end{bmatrix} \right\| \leq \epsilon \qquad (\operatorname{Newt} \epsilon)$$

where $\mathbf{A} = [\operatorname{vec}_{\mathcal{S}}(A_1), \dots, \operatorname{vec}_{\mathcal{S}}(A_m)]^T$, and then recover $\Delta S = \tilde{C} - \mathcal{A}^T(\Delta y)$. We focus exclusively on achieving a small enough ϵ as to be considered numerically exact.

As mentioned in the introduction, the main difficulty is that the scaling point W becomes progressively ill-conditioned as the accuracy parameter is taken $\mu \to 0^+$. The following is a more precise statement of (A1).

Assumption 1. There exist absolute constants $L \ge 0$ and $0 \le \delta < 1$ such that $W = X^{1/2} (X^{1/2} S X^{1/2})^{-1/2} X^{1/2}$ where $X, S \in \mathcal{S}_{++}^n$ satisfy

$$\left\|\frac{1}{\mu}X^{1/2}SX^{1/2} - I\right\| \le \delta, \qquad \|X - X^{\star}\| \le L \cdot \mu, \qquad \|S - S^{\star}\| \le L,$$

with respect to $X^{\star}, S^{\star} \in \mathcal{S}^n_+$ satisfying $X^{\star}S^{\star} = 0$ and $\chi_1^{-1} \cdot I \preceq X^{\star} + S^{\star} \preceq I$.

Under Assumption 1, the eigenvalues of W split into a size- r^* cluster that diverges as $\Theta(1/\sqrt{\mu})$, and a size- $(n - r^*)$ cluster that converges to zero as $\Theta(\sqrt{\mu})$. Therefore, its condition number scales as $\operatorname{cond}(W) = \Theta(1/\mu)$, and this causes the entire system to become ill-conditioned with a condition number like $\Theta(1/\mu^2)$.

Lemma 3.1. Under Assumption 1, let $r = \operatorname{rank}(X^*)$ and $0 < \mu \leq 1$. The eigenvalues of W satisfy

$$C_1/\sqrt{\mu} \geq \lambda_1(W) \geq \lambda_r(W) \geq 1/(C_2\sqrt{\mu}), C_1\sqrt{\mu} \geq \lambda_{r+1}(W) \geq \lambda_n(W) \geq \sqrt{\mu}/C_1,$$

where $C_1 = \frac{1+L}{1-\delta}$, and $C_2 = 4\chi_1 + \frac{2L^2\chi_1}{1-\delta}$.

Proof. Write $w_i \equiv \lambda_i(W)$ and $x_i^* \equiv \lambda_i(X^*)$ and $s_i^* \equiv \lambda_i(S^*)$, and let # denote the matrix geometric mean operator of Ando [2]. Substituting the matrix arithmetic-mean geometric-mean [2, Corollary 2.1] inequalities $\sqrt{\mu}W = X \# (\mu S^{-1}) \preceq \frac{1}{2}(X + \mu S^{-1})$ and $\sqrt{\mu}W^{-1} = S \# (\mu X^{-1}) \preceq \frac{1}{2}(S + \mu X^{-1})$ into $\|\frac{1}{\mu}X^{1/2}SX^{1/2} - I\| \leq \delta$ yields

$$\frac{1}{1+\delta} \cdot X \preceq \sqrt{\mu} W \preceq \frac{1}{1-\delta} \cdot X, \quad \frac{1}{1+\delta} \cdot S \preceq \sqrt{\mu} W^{-1} \preceq \frac{1}{1-\delta} \cdot S,$$

which we combine with $||X - X^*|| \le L\mu$ and $||S - S^*|| \le L$ to obtain

$$\sqrt{\mu}w_{1} \leq \frac{x_{1}^{\star} + L\mu}{1 - \delta}, \quad \sqrt{\mu}w_{n}^{-1} \leq \frac{s_{1}^{\star} + L}{1 - \delta}, \quad \sqrt{\mu}w_{r+1} \leq \frac{0 + L\mu}{1 - \delta},$$
$$\sqrt{\mu}w_{r} \geq \frac{x_{r}^{\star} - L\mu}{1 + \delta}, \quad \sqrt{\mu}w_{r}^{-1} \leq \frac{0 + L}{1 - \delta}.$$

The first row yields $\sqrt{\mu}w_1 \leq C_1$ for $0 < \mu \leq 1$, and $\sqrt{\mu}w_n^{-1} \leq C_1$ and $w_{r+1} \leq C_1\sqrt{\mu}$. We can select a *fixed* point $\hat{\mu} > 0$ and combine the second row to yield

$$\sqrt{\mu}w_r \ge \max\left\{\frac{x_r^{\star} - L\mu}{1 + \delta}, \frac{1 - \delta}{L}\mu\right\} \ge \min\left\{\frac{x_r^{\star} - L\hat{\mu}}{1 + \delta}, \frac{1 - \delta}{L}\hat{\mu}\right\},\$$

which is valid for all $\mu > 0$. In particular, if we choose $\hat{\mu}$ such that $x_r^{\star} - L\hat{\mu} = \frac{1}{2}x_r^{\star}$, then $\frac{x_r^{\star} - L\mu}{1+\delta} \ge \frac{1}{4}x_r^{\star} = \frac{1}{4\chi_1} \ge \frac{1}{C_2}$ and $\frac{1-\delta}{L}\hat{\mu} = \frac{1-\delta}{L}(\frac{x_r^{\star}}{2L}) \ge \frac{1-\delta}{2L^2\chi_1} \ge \frac{1}{C_2}$.

To derive a reformulation of (Newt- ϵ) that remains well-conditioned even as $\mu \to 0^+$, our approach is to construct a *low-rank plus well-conditioned* decomposition of the scaling matrix

$$W \otimes_{\mathcal{S}} W = \underbrace{\mathbf{Q} \boldsymbol{\Sigma}^{-1} \mathbf{Q}^T}_{\text{low-rank}} + \underbrace{\boldsymbol{\tau}^2 \cdot \mathbf{E}}_{\text{well-conditioned}}.$$

First, we choose a rank parameter $r \ge \operatorname{rank}(X^*)$, and partition the orthonormal eigendecomposition of the scaling matrix W into two parts

$$W = \underbrace{Q\Lambda Q^{T}}_{\text{top } r \text{ eigenvalues}} + \underbrace{Q_{\perp}\Lambda_{\perp}Q_{\perp}^{T}}_{\text{bottom } n-r \text{ eigenvalues}}$$
(6a)

Then, we choose a threshold parameter $\tau = \frac{1}{2}\lambda_{r+1}(W)$ and define

$$\mathbf{E} = E \otimes_{\mathcal{S}} E, \qquad E = QQ^T + \tau^{-1} \cdot Q_{\perp} \Lambda_{\perp} Q_{\perp}^T, \tag{6b}$$

$$\mathbf{Q} = [Q \otimes_{\mathcal{S}} Q, \quad \sqrt{2\Psi_n^T} (Q \otimes Q_\perp)], \tag{6c}$$

$$\Sigma = \operatorname{diag}(\Lambda \otimes_{\mathcal{S}} \Lambda - \tau^2 I, \quad (\Lambda - \tau I) \otimes \Lambda_{\perp})^{-1}.$$
(6d)

The statement below characterizes this decomposition.

Proposition 3.2 (Low-rank plus well-conditioned decomposition). Given $W \in S_{++}^n$, choose $1 \leq r \leq n$ and $0 < \tau < \lambda_r(W)$, and define $\mathbf{E}, \mathbf{Q}, \mathbf{\Sigma}$ as in (6). Then, $W \otimes_{\mathcal{S}} W = \mathbf{Q} \mathbf{\Sigma}^{-1} \mathbf{Q}^T + \tau^2 \cdot \mathbf{E}$ holds with $\mathbf{\Sigma} \succ 0$, and $\mathbf{E} \succeq 0$, and

$$\operatorname{colsp}(\mathbf{Q}) = \{\operatorname{vec}_{\mathcal{S}}(VQ^T) : V \in \mathbb{R}^{n \times r}\}, \qquad \mathbf{Q}^T \mathbf{Q} = I_d$$

where $d = nr - \frac{1}{2}r(r-1)$. Moreover, under Assumption 1, if $r \ge \operatorname{rank}(X^*)$ and $\tau = \frac{1}{2}\lambda_{r+1}(W)$, then

$$\begin{array}{rcl} 4 & \geq & \lambda_{\max}(\mathbf{E}) & \geq & \lambda_{\min}(\mathbf{E}) & \geq & 1/C_1^4 \\ C_1^4 & \geq & \tau^2 \cdot \lambda_{\max}(\mathbf{\Sigma}) & \geq & \tau^2 \cdot \lambda_{\min}(\mathbf{\Sigma}) & \geq & \mu^2/(4C_1^4) \end{array}$$

for all $0 < \mu \leq 1$, where C_1, C_2 are as defined in Lemma 3.1. If additionally $r = \operatorname{rank}(X^*)$, then $\mu \cdot C_1^3 C_2 \geq \tau^2 \cdot \lambda_{\max}(\Sigma)$ for all $0 < \mu \leq 1$.

Proof. The result follows from straightforward linear algebra and by substituting Lemma 3.1. For completeness, we provide a proof in Appendix A. \Box

We propose using an iterative Krylov subspace method to solve the following

$$\left\| \begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2 \cdot \mathbf{\Sigma} \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} - \begin{bmatrix} b + \tau^2 \mathbf{A}\mathbf{E}c \\ \tau^2 \cdot \mathbf{Q}^Tc \end{bmatrix} \right\| \le \epsilon,$$
(7a)

and then recover a solution to $(Newt-\epsilon)$ via the following

$$\Delta y = \tau^{-2} \cdot u, \qquad \operatorname{vec}_{\mathcal{S}}(\Delta X) = \mathbf{E}(\mathbf{A}^T u - \tau^2 c) + \mathbf{Q}v.$$
(7b)

Our main result is a theoretical guarantee that our specific choice of $\mathbf{E}, \mathbf{Q}, \mathbf{\Sigma}$ in (6) results in a bounded condition number in (7a), even as $\mu \to 0^+$. Notably, the well-conditioning holds for any rank parameter $r \ge \operatorname{rank}(X^*)$; this is important, because the exact value of $\operatorname{rank}(X^*)$ is often not precisely known in practice. The following is a more precise version of (A2) stated in a scale-invariant form.

Assumption 2 (Tangent space injectivity). Factor $X^* = U^*U^{*T}$ with $U^* \in \mathbb{R}^{n \times r}$. Then, there exists a condition number $1 \le \chi_2 < \infty$ such that

$$\|(\mathbf{A}\mathbf{A}^T)^{-1/2}\mathbf{A} \operatorname{vec}_{\mathcal{S}}(U^*V^T + VU^*)\| \ge \chi_2^{-1} \cdot \|U^*V^T + VU^*\|_F \quad \text{for all } V \in \mathbb{R}^{n \times r}.$$

Theorem 3.3 (Well-conditioning). Let Assumption 1 and Assumption 2 hold with parameters $L, \delta, \chi_1, \chi_2$. Given $W \in S^n_{++}$, select rank $(X^*) \leq r \leq n$ and $\tau = \frac{1}{2}\lambda_{r+1}(W)$, and define $\mathbf{E}, \mathbf{Q}, \boldsymbol{\Sigma}$ as in (6). For \mathbf{G} satisfying $\gamma_{\max} \mathbf{A} \mathbf{A}^T \succeq \mathbf{G} \succeq \gamma_{\min} \mathbf{A} \mathbf{A}^T$ with $\gamma_{\max} \geq \gamma_{\min} > 0$, we have

$$\operatorname{cond}\left(\begin{bmatrix}\mathbf{G} & \mathbf{A}\mathbf{Q}\\ \mathbf{Q}^{T}\mathbf{A}^{T} & -\tau^{2}\cdot\boldsymbol{\Sigma}\end{bmatrix}\right) = O(\operatorname{cond}(\mathbf{A}\mathbf{A}^{T})\cdot\gamma_{\max}^{4}\cdot\gamma_{\min}^{-8}\cdot L^{12}\cdot(1-\delta)^{-11}\cdot\chi_{1}^{2}\cdot\chi_{2}^{6})$$

for all $0 < \mu \leq 1$, with no dependence on $1/\mu$.

The most obvious way to use this result is to solve the well-conditioned indefinite system in (7a) using MINRES, and then use (7b) to recover a solution to (Newt- ϵ). In theory, this immediately allows us to complete an interior-point method iteration in $O(n^3)$ time and $O(n^2)$ memory.

Corollary 3.4. Given $W \in S_{++}^n$, $\mathbf{A} \in \mathbb{R}^{m \times \frac{1}{2}n(n+1)}$, $\tilde{b} \in \mathbb{R}^m$, $\tilde{C} \in S^n$, suppose that Assumption 1 and Assumption 2 hold with parameters $L, \delta, \chi_1, \chi_2$. Select rank $(X^*) \leq r \leq n$ and $\tau = \frac{1}{2}\lambda_{r+1}(W)$, and define $\mathbf{E}, \mathbf{Q}, \Sigma$ as in (6). Let

$$\begin{bmatrix} u_k \\ v_k \end{bmatrix} = \text{MINRES}_k \left(\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2 \cdot \mathbf{\Sigma} \end{bmatrix}, \begin{bmatrix} \tilde{b} + \tau^2 \mathbf{A}\mathbf{E} \operatorname{vec}_{\mathcal{S}}(\tilde{C}) \\ \tau^2 \cdot \mathbf{Q}^T \operatorname{vec}_{\mathcal{S}}(\tilde{C}) \end{bmatrix}, I, 0 \right)$$

and recover ΔX_k and Δy_k from (7b). Then, the residual condition (Newt- ϵ) is satisfied in at most

$$k = O\left(\operatorname{cond}(\mathbf{A}\mathbf{A}^T) \cdot L^{20} \cdot (1-\delta)^{-19} \cdot \chi_1^2 \cdot \chi_2^6 \cdot \log\left(\frac{L^4}{\mu \cdot \epsilon \cdot (1-\delta)^4}\right)\right) \text{ iterations}$$

for all $0 < \mu \leq 1$, with a logarithmic dependence on $1/\mu$. Moreover, suppose that $X \mapsto \mathbf{A} \operatorname{vec}_{\mathcal{S}}(X)$ and $y \mapsto \mathbf{A}^T y$ can be evaluated in at most $O(n^3)$ time and $O(n^2)$ storage for all $X \in S^n$ and $y \in \mathbb{R}^m$. Then, this algorithm can be implemented in $O(n^3k)$ time and $O(n^2)$ storage.

In practice, while the condition number of the augmented matrix in (7a) does remain bounded as $\mu \to 0^+$, it can grow to be very large, and MINRES can still require too many iterations to be effective. We propose an indefinite preconditioning algorithm. Below, recall that $PCG_k(A, b, P, x_0)$ denotes the k-th iterate generated by PCG when solving Ax = b using preconditioner P and starting from an initial x_0 .

Assumption 3 (Fast low-rank matrix-vector product). For $U, V \in \mathbb{R}^{n \times r}$ and $y \in \mathbb{R}^m$, the two matrixvector products $(U, V) \mapsto \mathcal{A}(UV^T + VU^T)$ and $y \mapsto [\mathcal{A}^T(y)]U$ can both be performed in at most $O(n^2r)$ time and $O(n^2)$ storage.

Corollary 3.5. Given $W \in S_{++}^n$, $\mathbf{A} \in \mathbb{R}^{m \times \frac{1}{2}n(n+1)}$, $\tilde{b} \in \mathbb{R}^m$, $\tilde{C} \in S^n$, suppose that Assumption 1 and Assumption 2 hold with parameters $L, \delta, \chi_1, \chi_2$. Select rank $(X^*) \leq r \leq n$ and $\tau = \frac{1}{2}\lambda_{r+1}(W)$, and define $\mathbf{E}, \mathbf{Q}, \mathbf{\Sigma}$ as in (6). Let $v_0 = -\mathbf{\Sigma}^{-1}\mathbf{Q}^T \tilde{c}$ where $\tilde{c} = \operatorname{vec}_{\mathcal{S}}(\tilde{C})$, and let

$$\begin{bmatrix} u_k \\ v_k \end{bmatrix} = \operatorname{PCG}_k \left(\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2 \cdot \boldsymbol{\Sigma} \end{bmatrix}, \begin{bmatrix} \tilde{b} + \tau^2 \mathbf{A}\mathbf{E}\tilde{c} \\ \tau^2 \cdot \mathbf{Q}^T\tilde{c} \end{bmatrix}, \begin{bmatrix} \beta I & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2 \cdot \boldsymbol{\Sigma} \end{bmatrix}, \begin{bmatrix} 0 \\ v_0 \end{bmatrix} \right)$$

where β is chosen to satisfy $\lambda_{\min}(\mathbf{AEA}^T) \leq \beta \leq \lambda_{\max}(\mathbf{AEA}^T)$, and recover ΔX_k and Δy_k from (7b). Then, the residual condition (Newt- ϵ) is satisfied in at most

$$k = \left\lceil \frac{\kappa_E \cdot \kappa_{\mathbf{A}}}{2} \log \left(\frac{\kappa_E \cdot \kappa_{\mathbf{A}} \cdot \|\tilde{b} + \mathbf{A} \operatorname{vec}_{\mathcal{S}}(W\tilde{C}W)\|}{2\epsilon} \right) \right\rceil \text{ iterations}$$

where $\kappa_E = \text{cond}(E) = \sqrt{\text{cond}(\mathbf{E})}$ and $\kappa_{\mathbf{A}} = \sqrt{\text{cond}(\mathbf{A}\mathbf{A}^T)}$, with no dependence on $1/\mu$. Under Assumption 3, this algorithm can be implemented with overall cost of $O(n^3r^3k)$ time and $O(n^2r^2)$ storage.

As we explained in the discussion around Proposition 2.5, the significance of the indefinite preconditioned PCG in Corollary 3.5 is that both the indefinite matrix and the indefinite preconditioner are well-conditioned for all $0 < \mu \leq 1$ via Theorem 3.3. Therefore, the preconditioner will continue to work even with very small values of μ .

In practice, the iteration count in Corollary 3.5 closely matches experimental observations. The rank parameter $r \ge r^*$ controls the tradeoff between the cost of the preconditioner and the reduction in iteration count. A larger value of r leads to a montonously smaller cond(E) and faster convergence, but also a cubically higher cost.

4 Proof of Well-Conditioning

Our proof of Theorem 3.3 is based on the following.

Lemma 4.1. Suppose that $\gamma_{\max} \mathbf{A} \mathbf{A}^T \succeq \mathbf{G} \succeq \gamma_{\min} \mathbf{A} \mathbf{A}^T$. Then,

$$\operatorname{cond}\left(\begin{bmatrix}\mathbf{G} & \mathbf{A}\mathbf{Q}\\ \mathbf{Q}^{T}\mathbf{A} & -\tau^{2}\boldsymbol{\Sigma}\end{bmatrix}\right) = O(\gamma_{\max} \cdot \gamma_{\min}^{-5} \cdot \operatorname{cond}(\mathbf{A}\mathbf{A}^{T}) \cdot \operatorname{cond}(\mathbf{C}))$$

where $\mathbf{C} = \tau^2 \mathbf{\Sigma} + \mathbf{Q}^T \mathbf{A}^T \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}$.

Proof. Block diagonal preconditioning with $\mathbf{P} = (\mathbf{A}\mathbf{A}^T)^{-1/2}$ yields

$$\mathbf{M} = \begin{bmatrix} \mathbf{G} & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} = \begin{bmatrix} \mathbf{P}^{-1} & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} \mathbf{P}\mathbf{G}\mathbf{P} & \mathbf{P}\mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T\mathbf{P} & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} \mathbf{P}^{-1} & 0 \\ 0 & I \end{bmatrix}.$$

Let $\tilde{\mathbf{G}} = \mathbf{P}\mathbf{G}\mathbf{P}$ and $\tilde{\mathbf{A}} = \mathbf{P}\mathbf{A}$. We perform a block-triangular decomposition

$$\tilde{\mathbf{M}} = \begin{bmatrix} \tilde{\mathbf{G}} & \tilde{\mathbf{A}}\mathbf{Q} \\ \mathbf{Q}^T \tilde{\mathbf{A}}^T & -\tau^2 \boldsymbol{\Sigma} \end{bmatrix} = \begin{bmatrix} I & 0 \\ \mathbf{Q}^T \tilde{\mathbf{A}}^T \tilde{\mathbf{G}}^{-1} & I \end{bmatrix} \begin{bmatrix} \tilde{\mathbf{G}} & 0 \\ 0 & -\mathbf{C} \end{bmatrix} \begin{bmatrix} I & \tilde{\mathbf{G}}^{-1} \tilde{\mathbf{A}} \mathbf{Q} \\ 0 & I \end{bmatrix},$$

where $\mathbf{C} = \tau^2 \mathbf{\Sigma} + \mathbf{Q}^T \tilde{\mathbf{A}}^T \tilde{\mathbf{G}} \tilde{\mathbf{A}} \mathbf{Q} = \tau^2 \mathbf{\Sigma} + \mathbf{Q}^T \mathbf{A}^T \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}$. Substituting $\|\tilde{\mathbf{G}}\| \leq \gamma_{\max}$ and $\|\tilde{\mathbf{G}}^{-1}\| \leq \gamma_{\min}^{-1}$ yields $\operatorname{cond}(\tilde{\mathbf{M}}) \leq (1 + \gamma_{\min}^{-1})^4 (\gamma_{\max} + \|\mathbf{C}\|) (\gamma_{\min}^{-1} + \|\mathbf{C}^{-1}\|)$. The desired estimate follows from $\operatorname{cond}(\mathbf{M}) \leq (1 + \|\mathbf{A}\mathbf{A}^T\|)(1 + \|(\mathbf{A}\mathbf{A}^T)^{-1}\|) \operatorname{cond}(\mathbf{M})$.

Given that **G** is already assumed to be well-conditioned, Lemma 4.1 says that the augmented matrix is well-conditioned if and only if the Schur complement **C** is well-conditioned. Proposition 3.2 assures us that $\|\mathbf{C}\|$ is always bounded as $\mu \to 0^+$, so the difficulty of the proof is to show that $\|\mathbf{C}^{-1}\|$ also remains bounded.

In general, we do not know the true rank $r^* = \operatorname{rank}(X^*)$ of the solution. If we choose $r > r^*$, then both terms $\tau^2 \Sigma$ and $\mathbf{Q}^T \mathbf{A}^T \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}$ will become singular in the limit $\mu = 0^+$, but their sum \mathbf{C} will nevertheless remain non-singular. To understand why this occurs, we need to partition the columns of Q into the dominant $r^* = \operatorname{rank}(X^*)$ eigenvalues and the $r - r^*$ excess eigenvalues, as in

$$W = \underbrace{Q_1 \Lambda_1 Q_1^T}_{\text{top } r^* \text{ eigenvalues}} + \underbrace{Q_2 \Lambda_2 Q_2^T}_{\text{next } r - r^* \text{ eigenvalues}} + \underbrace{Q_\perp \Lambda_\perp Q_\perp^T}_{\text{bottom } n - r \text{ eigenvalues}}$$
(8)

We emphasize that the partitioning $Q = [Q_1, Q_2]$ is done purely for the sake of analysis. Our key insight is that the matrix **Q** inherents a similar partitioning.

Lemma 4.2 (Participation of **Q** and **\Sigma**). Given $W \in S_{++}^n$ in (8), choose $0 < \tau < \lambda_r(W)$, let $Q = [Q_1, Q_2]$ and $\Lambda = \operatorname{diag}(\Lambda_1, \Lambda_2)$. Define

$$\mathbf{Q} = [Q \otimes_{\mathcal{S}} Q, \ \sqrt{2} \Psi_n^T (Q \otimes Q_\perp)], \quad \mathbf{\Sigma} = \operatorname{diag}(\Lambda \otimes_{\mathcal{S}} \Lambda - \tau^2 I, \ (\Lambda - \tau I) \otimes \Lambda_\perp)^{-1}.$$

Then, there exists permutation matrix Π so that $\mathbf{Q}\Pi = [\mathbf{Q}_1, \mathbf{Q}_2]$ where

$$\mathbf{Q}_1 = [Q_1 \otimes_{\mathcal{S}} Q_1, \ \sqrt{2} \Psi_n^T (Q_1 \otimes [Q_2, Q_\perp])], \quad \mathbf{Q}_2 = [Q_2 \otimes_{\mathcal{S}} Q_2, \ \sqrt{2} \Psi_n^T (Q_2 \otimes Q_\perp)],$$

and $\Pi^T \Sigma \Pi = \text{diag}(\Sigma_1, \Sigma_2)$ where

$$\begin{split} \boldsymbol{\Sigma}_1 &= \operatorname{diag}(\Lambda_1 \otimes_{\mathcal{S}} \Lambda_1 - \tau^2 I, \quad \Lambda_1 \otimes \operatorname{diag}(\Lambda_2, \Lambda_\perp) - \tau I \otimes \operatorname{diag}(I, \Lambda_\perp))^{-1}, \\ \boldsymbol{\Sigma}_2 &= \operatorname{diag}(\Lambda_2 \otimes_{\mathcal{S}} \Lambda_2 - \tau^2 I, \quad (\Lambda_2 - \tau I) \otimes \Lambda_\perp)^{-1}. \end{split}$$

Proof. Let $[Q_1, Q_2, Q_{\perp}] = I_n$ without loss of generality. For any $B \in S^r$ and $N \in \mathbb{R}^{n \times (n-r)}$, we verify that

$$\mathbf{Q}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B)\\\sqrt{2}\operatorname{vec}(N)\end{bmatrix} = \operatorname{vec}_{\mathcal{S}}\left(\begin{bmatrix}B & N^{T}\\N & 0\end{bmatrix}\right) = \operatorname{vec}_{\mathcal{S}}\left(\begin{bmatrix}B_{11} & B_{21}^{T} & N_{1}^{T}\\B_{21} & B_{22} & N_{2}^{T}\\N_{1} & N_{2} & 0\end{bmatrix}\right)$$
$$= \mathbf{Q}_{1}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B_{11})\\\sqrt{2}\operatorname{vec}\left(\begin{bmatrix}B_{21}\\N_{1}\end{bmatrix}\right)\end{bmatrix} + \mathbf{Q}_{2}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B_{22})\\\operatorname{vec}(N_{2})\end{bmatrix} = [\mathbf{Q}_{1}, \mathbf{Q}_{2}]\Pi^{T}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B)\\\sqrt{2}\operatorname{vec}(N)\end{bmatrix}$$

where we have partitioned B and N appropriately. Next, we verify that

$$\begin{aligned} \mathbf{Q} \mathbf{\Sigma}^{-1} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(B) \\ \sqrt{2} \operatorname{vec}(N) \end{bmatrix} &= \mathbf{Q} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(\Lambda B \Lambda - \tau^{2}I) \\ \sqrt{2} \operatorname{vec}[\Lambda_{\perp} N(\Lambda - \tau I)] \end{bmatrix} \\ &= \mathbf{Q}_{1} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(\Lambda_{1} B_{11} \Lambda_{1} - \tau B_{11}) \\ \sqrt{2} \operatorname{vec} \left(\begin{bmatrix} \Lambda_{2} B_{21} \Lambda_{1} \\ \Lambda_{\perp} N_{1} \Lambda_{1} \end{bmatrix} - \tau \begin{bmatrix} B_{21} \\ \Lambda_{\perp} N_{1} \end{bmatrix} \right) \end{bmatrix} + \mathbf{Q}_{2} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(\Lambda_{2} B_{22} \Lambda_{2} - \tau B_{22}) \\ \operatorname{vec}[\Lambda_{\perp} N(\Lambda_{2} - \tau I)] \end{bmatrix} \\ &= \mathbf{Q}_{1} \mathbf{\Sigma}_{1}^{-1} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(B_{11}) \\ \sqrt{2} \operatorname{vec} \left(\begin{bmatrix} B_{21} \\ N_{1} \end{bmatrix} \right) \end{bmatrix} + \mathbf{Q}_{2} \mathbf{\Sigma}_{2}^{-1} \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(B_{22}) \\ \operatorname{vec}(N_{2}) \end{bmatrix}. \end{aligned}$$

Applying Lemma 4.2 allows us to further partition the Schur complement C into blocks:

$$\Pi^{T}\mathbf{C}\Pi = \begin{bmatrix} \tau^{2}\boldsymbol{\Sigma}_{1} \\ \tau^{2}\boldsymbol{\Sigma}_{2} \end{bmatrix} + \begin{bmatrix} \mathbf{Q}_{1}^{T} \\ \mathbf{Q}_{2}^{T} \end{bmatrix} \mathbf{A}^{T}\mathbf{G}^{-1}\mathbf{A}\begin{bmatrix} \mathbf{Q}_{1} & \mathbf{Q}_{2} \end{bmatrix}.$$

In the limit $\mu = 0^+$, the following two lemmas assert that the two diagonal blocks $\tau^2 \Sigma_2$ and $\mathbf{Q}_1^T \mathbf{A}^T \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}_1$ will remain nonsingular. This is our key insight for why **C** will also remain non-singular.

Lemma 4.3 (Eigenvalue bounds on Σ_2). Given $W \in S_{++}^n$, choose $\tau = \frac{1}{2}\lambda_{r+1}(W)$, and define Σ_2 as in Lemma 4.2. Under Assumption 1, $\tau^2 \lambda_{\min}(\Sigma_2) \ge 1/(4C_1^4)$ where C_1 is defined in Lemma 3.1.

Proof. Write $w_i \equiv \lambda_i(W)$. Indeed, $\tau^{-2}\lambda_{\min}^{-1}(\Sigma_2) \leq 4w_{r^*+1}^2/w_n^2$ since $\lambda_{\min}^{-1}(\Sigma) \leq \max\{\lambda_{\max}(\Lambda_2 \otimes \Lambda_{\perp}), \lambda_{\max}(\Lambda_2 \otimes_{\mathcal{S}} \Lambda_2)\} \leq w_{r^*+1}^2$. Substituting Lemma 3.1 yields the desired bound.

Lemma 4.4 (Tangent space injectivity). Given $U \in \mathbb{R}^{n \times r}$, let $Q = \operatorname{orth}(U)$ and let Q_{\perp} to be its orthogonal complement, and define $\mathbf{Q} = [\mathbf{Q}_B, \mathbf{Q}_N]$ where $\mathbf{Q}_B = Q \otimes_S Q$ and $\mathbf{Q}_N = \sqrt{2} \Psi_n^T (Q \otimes Q_{\perp})$. Then, we have $\lambda_{\min}^{1/2}(\mathbf{Q}^T \mathbf{A}^T \mathbf{A} \mathbf{Q}) = \eta_{\mathbf{A}}(U)$ where

$$\eta_{\mathbf{A}}(U) \stackrel{\text{def}}{=} \min_{V \in \mathbb{R}^{n \times r}} \{ \| \mathbf{A} \operatorname{vec}_{\mathcal{S}}(UV^T + VU^T) \| : \| UV^T + VU^T \|_F = 1 \}.$$

Proof. Let rank (U) = k. The characterization of $\mathbf{Q} = [\mathbf{Q}_B, \mathbf{Q}_N]$ where $\mathbf{Q}_B = Q \otimes_{\mathcal{S}} Q$ and $\mathbf{Q}_N = \sqrt{2} \Psi_n^T (Q \otimes Q_\perp)$ in Proposition 3.2 yields

$$\lambda_{\min}^{1/2}(\mathbf{Q}^T \mathbf{A}^T \mathbf{A} \mathbf{Q}) = \min_{\|h\|=1} \|\mathbf{A} \mathbf{Q} h\| \stackrel{\text{(a)}}{=} \min_{\|\mathbf{Q} h\|=1} \|\mathbf{A} \mathbf{Q} h\| \stackrel{\text{(b)}}{=} \eta_{\mathbf{A}}(U).$$

Step (a) follows from $\mathbf{Q}^T \mathbf{Q} = I_d$ with $d = nk - \frac{1}{2}k(k-1)$. Step (b) is by substituting $\operatorname{colsp}(\mathbf{Q}) = \{\operatorname{vec}_{\mathcal{S}}(QV^T) : V \in \mathbb{R}^{n \times k}\} = \{\operatorname{vec}_{\mathcal{S}}(U\overline{V}^T) : \overline{V} \in \mathbb{R}^{n \times r}\}.$

In order to be able to accommodate small but nonzero values of $\mu > 0$, we will need to be able to perturb Lemma 4.4.

Lemma 4.5 (Injectivity perturbation). Let $Q, \hat{Q} \in \mathbb{R}^{n \times r}$ have orthonormal columns. Then, $|\eta_{\mathbf{A}}(Q) - \eta_{\mathbf{A}}(\hat{Q})| \leq 10 \|\mathbf{A}\| \| (I - \hat{Q}\hat{Q}^T)Q \|$.

We will defer the proof of Lemma 4.5 in order to prove our main result. Below, we factor $X^* = Q^* \Lambda^* Q^{*T}$. We will need the following claim.

Claim 4.6. Under Assumption 1, $\|(I - Q^*Q^{*T})Q_1\|^2 \le \mu \cdot \frac{C_2L}{\sqrt{1-\delta}}$. Proof. Lemma 3.1 says $W \succeq \lambda_{r^*}(W)Q_1Q_1^T \succeq (C_2\sqrt{\mu})^{-1}Q_1Q_1^T$ and therefore $\frac{1}{C_2\sqrt{\mu}}\|Q_1^TQ_{\perp}^*\|^2 \le \langle W, Q_{\perp}^*Q_{\perp}^{*T} \rangle = \langle (X^{1/2}SX^{1/2})^{-1/2}, X^{1/2}Q_{\perp}^*Q_{\perp}^{*T}X^{1/2} \rangle$ $\le \|(X^{1/2}SX^{1/2})^{-1/2}\| \cdot \langle X, Q_{\perp}^*Q_{\perp}^{*T} \rangle$

 $=\lambda_{\min}^{-1/2}(X^{1/2}SX^{1/2})\cdot\left\langle X-X^{\star},Q_{\perp}^{\star}Q_{\perp}^{\star T}\right\rangle\leq\mu\cdot\frac{L}{\sqrt{(1-\delta)\mu}}.$

Proof of Theorem 3.3. Write $\mathbf{K}_{ij} \stackrel{\text{def}}{=} \mathbf{Q}_i^T \mathbf{A}^T \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}_j$ for $i, j \in \{1, 2\}$. Under Assumption 1, substituting Claim 4.6 into Lemma 4.5 with $\mathbf{B} = (\mathbf{A}\mathbf{A}^T)^{-1/2}\mathbf{A}$ and taking $\eta_{\mathbf{B}}(U^*) = 1/\chi_2$ from Assumption 2 yields

$$\lambda_{\min}^{1/2}(\mathbf{Q}_1^T \mathbf{B}^T \mathbf{B} \mathbf{Q}_1) = \eta_{\mathbf{B}}(Q_1) \ge \eta_{\mathbf{B}}(Q^*) - 5 \|\mathbf{B}\| \| (I - Q^* Q^{*T}) Q_1 \|$$
$$\ge \frac{1}{\chi_2} - \mu \cdot \frac{5L}{C_2 \sqrt{1 - \delta}}.$$

In particular, if $0 < \mu \leq \mu_0 \stackrel{\text{def}}{=} \frac{C_2 \sqrt{1-\delta}}{6L}$, then $\lambda_{\min}^{1/2}(\mathbf{Q}_1^T \mathbf{B}^T \mathbf{B} \mathbf{Q}_1) \geq \frac{1}{6\chi_2}$ and

$$\mathbf{K}_{11} = \mathbf{Q}_{1}^{T} \mathbf{A}^{T} \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}_{1} \succeq \frac{1}{\gamma_{\max}} \mathbf{Q}_{1}^{T} \mathbf{A}^{T} (\mathbf{A} \mathbf{A}^{T})^{-1} \mathbf{A} \mathbf{Q}_{1}$$
$$= \frac{1}{\gamma_{\max}} \mathbf{Q}_{1}^{T} \mathbf{B}^{T} \mathbf{B} \mathbf{Q}_{1} \succeq \frac{1}{36\chi_{2}^{2} \gamma_{\max}}.$$
(9)

This implies $\mathbf{C} \succ 0$ via the steps below:

$$\Pi^{T} \mathbf{C} \Pi = \begin{bmatrix} \tau^{2} \boldsymbol{\Sigma}_{1} \\ \tau^{2} \boldsymbol{\Sigma}_{2} \end{bmatrix} + \begin{bmatrix} \mathbf{K}_{11} & \mathbf{K}_{21}^{T} \\ \mathbf{K}_{21} & \mathbf{K}_{22} \end{bmatrix}$$
$$\succeq \begin{bmatrix} 0 & 0 \\ 0 & \tau^{2} \boldsymbol{\Sigma}_{2} \end{bmatrix} + \begin{bmatrix} I \\ \mathbf{K}_{21} \mathbf{K}_{11}^{-1} & I \end{bmatrix} \begin{bmatrix} \mathbf{K}_{11} \\ 0 \end{bmatrix} \begin{bmatrix} I & \mathbf{K}_{21} \mathbf{K}_{11}^{-1} \\ I \end{bmatrix}$$
$$= \begin{bmatrix} I \\ \mathbf{K}_{21} \mathbf{K}_{11}^{-1} & I \end{bmatrix} \begin{bmatrix} \mathbf{K}_{11} \\ \tau^{2} \boldsymbol{\Sigma}_{2} \end{bmatrix} \begin{bmatrix} I & \mathbf{K}_{21} \mathbf{K}_{11}^{-1} \\ I \end{bmatrix}.$$

Indeed, substituting $\|\mathbf{K}_{11}^{-1}\| \leq 36\chi_2^2\gamma_{\max}$ from (9) and $\|(\tau^2\Sigma_2)^{-1}\| \leq 4C_1^4$ from Lemma 4.3 and $C_1 = O(L \cdot (1-\delta)^{-1})$ yields

$$\begin{aligned} \|\mathbf{C}^{-1}\| &\leq (1 + \|\mathbf{K}_{21}\mathbf{K}_{11}^{-1}\|)^2 (\|\mathbf{K}_{11}^{-1}\| + \|(\tau^2 \boldsymbol{\Sigma}_2)^{-1}\|) \\ &\leq (1 + 36\chi_2^2 \gamma_{\max} \cdot \gamma_{\min}^{-1})^2 \left(36\chi_2^2 \cdot \gamma_{\max} + 4C_1^4\right) \\ &= O(\gamma_{\max}^3 \cdot \gamma_{\min}^{-2} \cdot \chi_2^6 \cdot L^4 \cdot (1 - \delta)^{-4}). \end{aligned}$$

The second line uses the hypothesis $\mathbf{G} \succeq \gamma_{\min} \mathbf{A} \mathbf{A}^T \succ 0$ to bound

$$\|\mathbf{K}_{21}\| \le \|\mathbf{A}^T \mathbf{G}^{-1} \mathbf{A}\| \le \gamma_{\min}^{-1} \|\mathbf{A}^T (\mathbf{A} \mathbf{A}^T)^{-1} \mathbf{A}\| = \gamma_{\min}^{-1}$$

Otherwise, if $\mu_0 < \mu \leq 1$, then substituting $\|(\tau^2 \Sigma)^{-1}\| \leq 4C_1^4/\mu^2$ from Proposition 3.2 and $C_2 = O(\chi_1 \cdot L^2 \cdot (1-\delta)^{-1})$ yields

$$\|\mathbf{C}^{-1}\| \le \|(\tau^2 \mathbf{\Sigma})^{-1}\| \le 4C_1^4/\mu_0^2 = 4C_1^4 \cdot \frac{36L^2C_2^2}{1-\delta} = O(L^8 \cdot \chi_1^2 \cdot (1-\delta)^{-7}).$$

Finally, for all $0 < \mu \leq 1$,

$$\|\mathbf{C}\| \le \|\tau^{2} \mathbf{\Sigma}\| + \|\mathbf{Q}^{T} \mathbf{A}^{T} \mathbf{G}^{-1} \mathbf{A} \mathbf{Q}\| \le C_{1}^{4} + \gamma_{\min}^{-1} = O(\gamma_{\min}^{-1} \cdot L^{4} \cdot (1-\delta)^{-4}).$$

Substituting $\|\mathbf{C}\|$ and $\|\mathbf{C}^{-1}\|$ into Lemma 4.1 with the following simplification

$$\xi(1+\gamma_{\min}^{-1})^4(\gamma_{\max}+\|\mathbf{C}\|)(\gamma_{\min}^{-1}+\|\mathbf{C}^{-1}\|) = O(\gamma_{\max}\cdot\gamma_{\min}^{-5}\cdot\operatorname{cond}(\mathbf{A}\mathbf{A}^T)\cdot\operatorname{cond}(\mathbf{C}))$$

yields the desired bound.

We now turn to the proof of Lemma 4.5. We first need to prove a technical lemma on the distance between orthogonal matrices. Below, Orth(n) denotes the set of $n \times n$ orthonormal matrices.

Lemma 4.7. Let $Q, \hat{Q} \in \mathbb{R}^{n \times r}$ and $Q_{\perp}, \hat{Q}_{\perp} \in \mathbb{R}^{n \times (n-r)}$ satisfy $[Q, Q_{\perp}] \in \text{Orth}(n)$ and $[\hat{Q}, \hat{Q}_{\perp}] \in \text{Orth}(n)$. Then,

$$\max\left\{\min_{R\in\operatorname{Orth}(r)}\|Q-\hat{Q}R\|,\min_{R_{\perp}\in\operatorname{Orth}(n-r)}\|Q_{\perp}-\hat{Q}_{\perp}R_{\perp}\|\right\}\leq\sqrt{2}\|Q^{T}\hat{Q}_{\perp}\|.$$

Proof. Let $\sigma_{\max}(\hat{Q}_{\perp}^T Q) = \sin \theta$. Define $\operatorname{polar}(M) = \arg \max_{R \in \operatorname{Orth}(n)} \langle M, R \rangle = UV^T$ where $M = U\Sigma V^T$ is the usual SVD. The choice of $R = \operatorname{polar}(\hat{Q}^T Q)$ yields

$$\|Q - \hat{Q}R\|^{2} = \|\hat{Q} - QR^{T}\|^{2} \le \|I - \hat{Q}^{T}QR^{T}\|^{2} + \|\hat{Q}_{\perp}^{T}QR^{T}\|^{2}$$

$$\stackrel{(a)}{=} [1 - \sigma_{\min}(\hat{Q}^{T}Q)]^{2} + \sigma_{\max}^{2}(\hat{Q}_{\perp}^{T}Q)$$

$$\stackrel{(b)}{=} (1 - \cos\theta)^{2} + \sin^{2}\theta \le 2\sin^{2}(\theta) \text{ for } \theta \in [0, \pi/2].$$

Step (a) is because our choice of R renders $\hat{Q}^T Q R^T = R Q^T \hat{Q} \succeq 0$. Step (b) is because of $\sigma_{\min}(\hat{Q}^T Q) = \cos \theta$, which follows from $\lambda_{\min}(Q^T \hat{Q} \hat{Q}^T Q) = \lambda_{\min}(Q^T [I - \hat{Q}_{\perp} \hat{Q}_{\perp}^T] Q) = 1 - \lambda_{\max}(Q^T \hat{Q}_{\perp} \hat{Q}_{\perp}^T Q)$. The proof for $\|Q_{\perp} - \hat{Q}_{\perp} R_{\perp}\|^2 \leq 2 \sin^2(\theta)$ follows identical steps with the choice of $R_{\perp} = \operatorname{polar}(\hat{Q}_{\perp}^T Q_{\perp})$.

We will now use Lemma 4.7 to prove Lemma 4.5.

Proof of Lemma 4.5. Given that $\eta(Q) = \eta(QR)$ for $R \in \operatorname{Orth}(r)$, we assume without loss of generality that $\|Q - \hat{Q}\| \leq \sqrt{2} \|Q^T \hat{Q}_{\perp}\|$ and $\|Q_{\perp} - \hat{Q}_{\perp}\| \leq \sqrt{2} \|Q^T \hat{Q}_{\perp}\|$ as in Lemma 4.7. Applying Weyl's inequality for singular values yields

$$|\eta(Q) - \eta(\hat{Q})| = |\sigma_d(\mathbf{A}\mathbf{Q}) - \sigma_d(\mathbf{A}\hat{\mathbf{Q}})| \le \|\mathbf{A}\mathbf{Q} - \mathbf{A}\hat{\mathbf{Q}}\| \le \|\mathbf{A}\|\|\mathbf{Q} - \hat{\mathbf{Q}}\|$$

Next, observe that

$$\begin{aligned} \left(\mathbf{Q} - \hat{\mathbf{Q}}\right) \begin{bmatrix} \operatorname{vec}_{\mathcal{S}}(B) \\ \frac{1}{\sqrt{2}} \operatorname{vec}(N) \end{bmatrix} &= \operatorname{vec}_{\mathcal{S}} \left(\begin{bmatrix} Q & Q_{\perp} \end{bmatrix} \begin{bmatrix} B \\ N \end{bmatrix} Q^{T} - \begin{bmatrix} \hat{Q} & \hat{Q}_{\perp} \end{bmatrix} \begin{bmatrix} B \\ N \end{bmatrix} \hat{Q}^{T} \right) \\ &= \operatorname{vec}_{\mathcal{S}} \left(\begin{bmatrix} Q & Q_{\perp} \end{bmatrix} \begin{bmatrix} B \\ N \end{bmatrix} (Q - \hat{Q})^{T} \right) \\ &+ \operatorname{vec}_{\mathcal{S}} \left(\left(\begin{bmatrix} Q & Q_{\perp} \end{bmatrix} - \begin{bmatrix} \hat{Q} & \hat{Q}_{\perp} \end{bmatrix}\right) \begin{bmatrix} B \\ N \end{bmatrix} \hat{Q}^{T} \right), \end{aligned}$$

and therefore

$$\begin{aligned} \frac{1}{\sqrt{2}} \|\mathbf{Q} - \hat{\mathbf{Q}}\| &\leq \|Q - \hat{Q}\| + \|[Q, Q_{\perp}] - [\hat{Q}, \hat{Q}_{\perp}]\| \\ &\leq \|Q - \hat{Q}\| + \sqrt{\|Q - \hat{Q}\|^2 + \|Q_{\perp} - \hat{Q}_{\perp}\|^2} \leq (1 + \sqrt{2}) \cdot \|Q^T \hat{Q}_{\perp}\|. \end{aligned}$$

Finally, we note that $(1 + \sqrt{2}) \cdot 2 \approx 4.828 < 5$.

5 Solution via MINRES

Let us estimate the cost of solving (7) using MINRES, as in Corollary 3.4. In the initial setup, we spend $O(n^3)$ time and $O(n^2)$ memory to explicitly compute the diagonal matrix Σ , and to implicitly set up \mathbf{E}, \mathbf{Q} via their constituent matrices $E \in S^n$, $Q \in \mathbb{R}^{n \times r}$, and $Q_{\perp} \in \mathbb{R}^{n \times (n-r)}$. Afterwards, each matrix-vector product with $\mathbf{E}, \mathbf{Q}, \mathbf{Q}^T$ can be evaluated in $O(n^3)$ time and $O(n^2)$ memory, because $\mathbf{E} \operatorname{vec}_{\mathcal{S}}(X) = \operatorname{vec}_{\mathcal{S}}(EXE)$ and

$$\mathbf{Q}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B)\\\operatorname{vec}(N)\end{bmatrix} = \operatorname{vec}_{\mathcal{S}}\left(\begin{bmatrix} Q & Q_{\perp}\end{bmatrix}\begin{bmatrix} B\\\sqrt{2}N\end{bmatrix}Q^{T}\right), \quad \mathbf{Q}^{T}\operatorname{vec}_{\mathcal{S}}(X) = \begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(Q^{T}XQ)\\\sqrt{2}\operatorname{vec}(Q_{\perp}^{T}XQ)\end{bmatrix}.$$

Therefore, the per-iteration cost of MINRES, which is dominated by a single matrix-vector product with the augmented matrix in (7a), is $O(n^3)$ time and $O(n^2)$ memory, plus a single matrix-vector product with **A** and **A**^T. This is also the cost of evaluating the recovery equation in (7b).

We now turn to the number of iterations. Proposition 2.2 says that MINRES converges to ϵ residual in $O(\kappa \log(1/\epsilon))$ iterations, where κ is the condition estimate from Theorem 3.3, but we need the following lemma to ensure that small residual in (7a) would actually recover through (7b) a small-residual solution $\Delta X, \Delta y$ to our original problem (Newt- ϵ).

Lemma 5.1 (Propagation of small residuals). Given $\mathbf{D} = \mathbf{Q} \mathbf{\Sigma}^{-1} \mathbf{Q}^T + \tau^2 \cdot \mathbf{E}$ where $\mathbf{Q}^T \mathbf{Q} = I$ and $\mathbf{E} \succ 0$ and $\mathbf{\Sigma} \succ 0$. Let

$$\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} - \begin{bmatrix} b + \tau^2 \cdot \mathbf{A}\mathbf{E}c \\ \tau^2 \cdot \mathbf{Q}^Tc \end{bmatrix} = \begin{bmatrix} p \\ d \end{bmatrix}.$$
 (10)

Then, $y = \tau^{-2} \cdot u$ and $x = \mathbf{E}(\mathbf{A}^T u - \tau^2 c) + \mathbf{Q}v$ satisfy the following

$$\begin{bmatrix} -\mathbf{D}^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} c \\ b \end{bmatrix} = \begin{bmatrix} \tau^{-2} \mathbf{E}^{-1} \mathbf{Q} \mathbf{C}^{-1} d \\ p \end{bmatrix}$$
(11)

where $\mathbf{C} = \tau^2 \mathbf{\Sigma} + \mathbf{Q}^T \mathbf{E}^{-1} \mathbf{Q}$.

Proof. Define (x^*, u^*, v^*) as the exact solution to the following system of equations

$$\begin{bmatrix} -\mathbf{E}^{-1} & \mathbf{A}^T & \mathbf{E}^{-1}\mathbf{Q} \\ \mathbf{A} & 0 & 0 \\ \mathbf{Q}^T\mathbf{E}^{-1} & 0 & -\mathbf{C} \end{bmatrix} \begin{bmatrix} x^* \\ u^* \\ v^* \end{bmatrix} = \begin{bmatrix} \tau^2 c \\ b \\ 0 \end{bmatrix}.$$

Our key observation is that the matrix has two possible block factorizations

$$\begin{bmatrix} I & & \\ -\mathbf{A}\mathbf{E} & I & 0 \\ -\mathbf{Q}^T & 0 & I \end{bmatrix} \begin{bmatrix} -\mathbf{E}^{-1} & & & \\ & \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ & & \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} I & -\mathbf{E}\mathbf{A}^T & -\mathbf{Q} \\ & I & 0 \\ & & 0 & I \end{bmatrix}$$
(12)

$$= \begin{bmatrix} I & 0 & -\mathbf{E}^{-1}\mathbf{Q}\mathbf{C}^{-1} \\ 0 & I & 0 \\ & & I \end{bmatrix} \begin{bmatrix} -\tau^{2}\mathbf{D}^{-1} & \mathbf{A}^{T} \\ \mathbf{A} & 0 \\ & & -\mathbf{C} \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & I \\ -\mathbf{C}^{-1}\mathbf{Q}^{T}\mathbf{E}^{-1} & I \end{bmatrix},$$
(13)

where $\tau^2 \mathbf{D}^{-1} = \mathbf{E}^{-1} - \mathbf{E}^{-1} \mathbf{Q} \mathbf{C}^{-1} \mathbf{Q}^T \mathbf{E}^{-1}$ is via the Sherman–Morrison–Woodbury identity. We verify from (12) and (13) respectively that

$$\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} u^* \\ v^* \end{bmatrix} = \begin{bmatrix} b+\tau^2\cdot\mathbf{A}\mathbf{E}c \\ \tau^2\cdot\mathbf{Q}^Tc \end{bmatrix}, \quad \begin{bmatrix} -\tau^2\mathbf{D}^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \end{bmatrix} \begin{bmatrix} x^* \\ u^* \end{bmatrix} = \begin{bmatrix} \tau^2c \\ b \end{bmatrix},$$

and $x^{\star} = \mathbf{E}\mathbf{A}^T u^{\star} + \mathbf{Q}v^{\star}$. Now, for a given (x, u, v), the associated errors are

$$\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} u - u^* \\ v - v^* \end{bmatrix} = \begin{bmatrix} p \\ d \end{bmatrix}, \qquad x - x^* = \begin{bmatrix} \mathbf{E}\mathbf{A}^T & \mathbf{Q} \end{bmatrix} \begin{bmatrix} u - u^* \\ v - v^* \end{bmatrix},$$

and these can be rearranged as

$$\begin{bmatrix} -\mathbf{E}^{-1} & & \\ & \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ & & \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} I & -\mathbf{E}\mathbf{A}^T & -\mathbf{Q} \\ & I & 0 \\ & 0 & I \end{bmatrix} \begin{bmatrix} x - x^* \\ u - u^* \\ v - v^* \end{bmatrix} = \begin{bmatrix} 0 \\ p \\ d \end{bmatrix}.$$

It follows from our first block-triangular factorization (12) that

$$\begin{bmatrix} -\mathbf{E}^{-1} & \mathbf{A}^T & \mathbf{E}^{-1}\mathbf{Q} \\ \mathbf{A} & 0 & 0 \\ \mathbf{Q}^T\mathbf{E}^{-1} & 0 & -\mathbf{C} \end{bmatrix} \begin{bmatrix} x - x^* \\ u - u^* \\ v - v^* \end{bmatrix} = \begin{bmatrix} I \\ -\mathbf{A}\mathbf{E} & I & 0 \\ -\mathbf{Q}^T & 0 & I \end{bmatrix} \begin{bmatrix} 0 \\ p \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ p \\ d \end{bmatrix}.$$

It follows from our second block-triangular factorization (13) that

$$\begin{bmatrix} -\tau^2 \mathbf{D}^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \\ & & -\mathbf{C} \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & I \\ -\mathbf{C}^{-1} \mathbf{Q}^T \mathbf{E}^{-1} & I \end{bmatrix} \begin{bmatrix} x - x^* \\ u - u^* \\ v - v^* \end{bmatrix} = \begin{bmatrix} \mathbf{E}^{-1} \mathbf{Q} \mathbf{C}^{-1} d \\ p \\ d \end{bmatrix}.$$

We obtain (11) by isolating the first two block-rows, rescaling the first block-row by τ^{-2} , and then substituting (x^*, u^*) as an exact solution.

It follows that the residuals between (7) and (Newt- ϵ) are related as

$$\left\| \begin{bmatrix} -\mathbf{D}^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} b \\ c \end{bmatrix} \right\| \le \frac{16C_1^4C_2^2}{\mu} \cdot \left\| \begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} - \begin{bmatrix} b + \tau^2 \cdot \mathbf{A}\mathbf{E}c \\ \tau^2 \cdot \mathbf{Q}^Tc \end{bmatrix} \right\|$$

where we used $\|\tau^{-2}\mathbf{E}^{-1}\mathbf{Q}\mathbf{C}^{-1}\| \leq \tau^{-2}\|\mathbf{E}^{-1}\|\|\mathbf{E}\|$, and substituted $\tau = \frac{1}{2}\lambda_{r+1}(W) \geq \sqrt{\mu}/C_1$ where $C_1 = \frac{1+L}{1-\delta}$ from Lemma 3.1, and $\|\mathbf{E}\| \leq 4$ and $\|\mathbf{E}^{-1}\| \leq C_1^4$ from Proposition 3.2. We obtain Corollary 3.4 by substituting Theorem 3.3 into Proposition 2.2, setting $\gamma_{\max} = \|\mathbf{E}\| = 4$ and $\gamma_{\min}^{-1} = \|\mathbf{E}^{-1}\| = C_1^4$ and then performing enough iterations to achieve a residual of $\frac{\mu\epsilon}{16C_1^4}$.

6 Solution via indefinite PCG

Let us estimate the cost of solving (7) using indefinite PCG, as in Corollary 3.5. After setting up $\mathbf{E}, \mathbf{Q}, \mathbf{Q}^T$ in the same way as using MINRES, we spend $O(d \cdot n^2 r) = O(n^3 r^2)$ time and $O(d^2) = O(n^2 r^2)$ memory to explicitly form the $d \times d$ Schur complement of the preconditioner

$$\mathbf{C} = \tau^2 \mathbf{\Sigma} + \beta^{-1} \mathbf{Q}^T \mathbf{A}^T \mathbf{A} \mathbf{Q},$$

where $d = nr - \frac{1}{2}r(r-1)$. This can be done using d matrix-vector products with **AQ** and **Q**^T**A**^T, which under Assumption 3, can each be evaluated in at $O(n^2r)$ time and $O(n^2)$ memory:

$$\mathbf{AQ}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(B)\\\operatorname{vec}(N)\end{bmatrix} = \mathcal{A}(QU^T + UQ^T), \qquad \mathbf{Q}^T\mathbf{A}^T y = \begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(Q^T)\\\sqrt{2}\operatorname{vec}(Q_{\perp}^T V)\end{bmatrix}$$

where $U = QB + \sqrt{2}Q_{\perp}N$ and $V = \mathcal{A}^T(y)Q$. Once **C** is formed, the indefinite preconditioner can be applied in its block triangular form

$$\begin{bmatrix} \beta I & \mathbf{A} \mathbf{Q} \\ \mathbf{Q}^T \mathbf{A}^T & -\tau^2 \boldsymbol{\Sigma} \end{bmatrix}^{-1} = \begin{bmatrix} I & -\mathbf{A} \mathbf{Q} \\ 0 & I \end{bmatrix} \begin{bmatrix} \beta I & 0 \\ 0 & -\mathbf{C}^{-1} \end{bmatrix} \begin{bmatrix} I & 0 \\ -\mathbf{Q}^T \mathbf{A}^T & I \end{bmatrix}$$

as a linear solve with \mathbf{C} , and a matrix-vector products with each of $\mathbf{A}\mathbf{Q}$ and $\mathbf{Q}^T\mathbf{A}^T$, for $O(d^3) = O(n^3r^3)$ time and $O(d^2) = O(n^2r^2)$ memory. Moreover, note that under Assumption 3, each matrix-vector product with \mathbf{A} and \mathbf{A}^T costs $O(n^3)$ time and $O(n^2)$ memory. Therefore, the per-iteration cost of PCG, which is dominated by a single matrix-vector product with the augmented matrix in (7a) and a single application of the indefinite preconditioner, is $O(n^3r^3)$ time and $O(n^2r^2)$ memory.

We now turn to the number of iterations. Proposition 2.5 says that indefinite PCG will generate identical iterates u_k to regular PCG applied to the following problem

$$\underbrace{\mathbf{A}(\mathbf{E} + \tau^{-2}\mathbf{Q}\boldsymbol{\Sigma}^{-1}\mathbf{Q}^T)\mathbf{A}^T}_{\mathbf{H}}u_k = \underbrace{b + \tau^2\mathbf{A}(\mathbf{E} + \tau^{-2}\mathbf{Q}\boldsymbol{\Sigma}^{-1}\mathbf{Q}^T)c}_{g}$$

with preconditioner $\tilde{\mathbf{H}} = \beta I + \tau^{-2} \mathbf{A} \mathbf{Q} \boldsymbol{\Sigma}^{-1} \mathbf{Q}^T \mathbf{A}^T$ and iterates v_k that exactly satisfy

$$\mathbf{Q}^T \mathbf{A}^T u_k - \tau^2 \cdot \mathbf{\Sigma} v_k = \tau^2 \cdot \mathbf{Q}^T c.$$

Plugging these iterates back into (7a) yields

$$\begin{bmatrix} \mathbf{A}\mathbf{E}\mathbf{A}^T & \mathbf{A}\mathbf{Q} \\ \mathbf{Q}^T\mathbf{A}^T & -\tau^2\boldsymbol{\Sigma} \end{bmatrix} \begin{bmatrix} u_k \\ v_k \end{bmatrix} - \begin{bmatrix} b+\tau^2 \cdot \mathbf{A}\mathbf{E}c \\ \tau^2 \cdot \mathbf{Q}^Tc \end{bmatrix} = \begin{bmatrix} \mathbf{H}u_k - g \\ 0 \end{bmatrix},$$

and plugging into Lemma 5.1 and Proposition 2.4 shows that the $\Delta X, \Delta y$ recovered from (7b) will satisfy

$$\left\| \begin{bmatrix} -\mathbf{D}^{-1} & \mathbf{A}^T \\ \mathbf{A} & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} - \begin{bmatrix} b \\ c \end{bmatrix} \right\| = \left\| \mathbf{H}u_k - g \right\| \le 2\sqrt{\kappa} \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^k \|g\|$$

where $\kappa = \operatorname{cond}(\tilde{\mathbf{H}}^{-1/2}\mathbf{H}\tilde{\mathbf{H}}^{-1/2})$. We use the following argument to estimate κ . Claim 6.1. Let $\mathbf{E} \succ 0$ satisfy $\lambda_{\min}(\mathbf{E}) \leq \beta \leq \lambda_{\max}(\mathbf{E})$. Then

$$\operatorname{cond}((\beta I + \mathbf{U}\mathbf{U}^T)^{-1/2}(\mathbf{E} + \mathbf{U}\mathbf{U}^T)(\beta I + \mathbf{U}\mathbf{U}^T)^{-1/2}) \le \operatorname{cond}(\mathbf{E})$$

Proof. We have $\mathbf{E} + \mathbf{U}\mathbf{U}^T \preceq \lambda_{\max}(\mathbf{E})I + UU^T \preceq (\lambda_{\max}(\mathbf{E})/\beta)(\beta I + UU^T)$ because $\lambda_{\max}(\mathbf{E})/\beta \ge 1$. We have $\mathbf{E} + \mathbf{U}\mathbf{U}^T \succeq \lambda_{\min}(\mathbf{E})I + UU^T \succeq (\lambda_{\min}(\mathbf{E})/\beta)(\beta I + UU^T)$ because $\lambda_{\min}(\mathbf{E})/\beta \le 1$.

Therefore, given that $\lambda_{\min}(\mathbf{AEA}^T) \leq \beta \leq \lambda_{\max}(\mathbf{AEA}^T)$ holds by hypothesis, it takes at most k iterations to achieve $\|\mathbf{H}u_k - g\| \leq \epsilon$, where

$$k = \left\lceil \frac{1}{2} \sqrt{\operatorname{cond}(\mathbf{A} \mathbf{E} \mathbf{A}^T)} \log \left(\frac{\operatorname{cond}(\mathbf{A} \mathbf{E} \mathbf{A}^T) \cdot \|g\|}{2\epsilon} \right) \right\rceil$$

Finally, we bound $\operatorname{cond}(\mathbf{AEA}^T) \leq \operatorname{cond}(\mathbf{E}) \cdot \operatorname{cond}(\mathbf{AA}^T)$ and $\operatorname{cond}(\mathbf{E}) \leq \operatorname{cond}(E)^2$.

7 Experiments

The impetus of this paper is to solve the *robust* low-rank matrix recovery problem [9, 21, 13], which seeks to recover a low-rank ground truth matrix X^* , given a linear measurement operator $\mathcal{A} : \mathbb{R}^{n \times n} \to \mathbb{R}^m$ and linear measurements $b \approx \mathcal{A}(X^*)$. Specific to the robust variant is the assumption that some of the measurements are grossly corrupted, meaning that $\varepsilon = b - \mathcal{A}(X^*)$ contains mostly zero elements, except for a small number of *outliers*. The problem admits a wellknown convex relaxation

$$\min_{X \in \mathbb{R}^{n \times n}} \quad \|\mathcal{A}(X) - b\|_{\ell_1} + \lambda \|X\|_* \tag{14}$$

where $||b||_{\ell_1} \equiv \sum_i |b_i|$ denotes the usual vector ℓ_1 norm, and the nuclear norm regularizer $||X||_* \equiv \sum_i \sigma_i(X)$ is used to encourage lower rank in X.

Despite its ostensibly familiar form, the nonsmooth nature of (14) makes the problem surprisingly difficult to solve. The classical way of reformulating (14) into an SDP and applying a generalpurpose interior-point method incurs at least $\Omega((n+m)^3)$ time and $\Omega((n+m)^2)$ memory. This is particularly concerning for problem instances with $m = O(n^2)$ measurements, such as the robust PCA problem [4], as the per-iteration cost would scale at least $\Omega(n^6)$ time.

Unfortunately, as we explained in the introduction, the Burer–Monteiro approach is not able to solve (14) satisfactorily. For the symmetric, positive semidefinite variant of (14) that we introduced earlier as (2), the Burer–Monteiro approach would either solve the unconstrained nonsmooth problem

$$\min_{U \in \mathbb{R}^{n \times r}} f(U) = \|\mathcal{A}(UU^T) - b\|_{\ell_1} + \lambda \|U\|_F^2,$$
(15)

or the constrained twice-differentiable problem

$$\min_{U \in \mathbb{R}^{n \times r}, v, w \in \mathbb{R}^m} \quad \mathbf{1}^T v + \mathbf{1}^T w + \lambda \|U\|_F^2 \quad \text{s.t.} \quad \begin{array}{l} v - w + \mathcal{A}(UU^T) = b, \\ v \ge 0, \quad w \ge 0, \end{array}$$
(16)

in which $r \geq \operatorname{rank}(X^*)$ is a search rank parameter. The machine learning area prefers to locally optimize the nonsmooth objective in (15) using the subgradient method $U_{\text{new}} = U - \alpha \cdot \partial f(U)$ with step-size $\alpha \in (0, 1]$, which is commonly called GD in this context. The operations research area prefers applying a nonlinear programming (NLP) solver like fmincon [23] and knitro [6] to (16). But in either case, convergence can both be very slow, due to the underlying nonsmoothness of the problem, and also very unreliable, due to sensitivity to the initial point. In the end, both approaches can fail by getting stuck at a *spurious* local minimum.

Instead, we present experimental evidence to show that our proposed approach is the most reliable way of solving instances of (14) to high accuracies. In the remainder of this section, we consider the symmetric, positive semidefinite instance (2) with $\lambda = 1$ and \mathcal{A} chosen according to the "matrix completion" model, meaning that $\mathcal{A}(X) = P_m \operatorname{vec}_{\mathcal{S}}(X)$ where P_m is the matrix that randomly permutes and then subsamples m out of n(n+1)/2 elements in $\operatorname{vec}_{\mathcal{S}}(X)$. In Section 7.1, we validate Assumptions 1, 2 and 3 for the SDP reformulation (3) with this particular choice of \mathcal{A} , as well as the bounded condition number predicted in Theorem 3.3, and the cubic time complexity predicted in Corollaries 3.4 and 3.5. In Section 7.2, we compare the practical performance of our method against GD, fmincon, and knitro, and also the prior approach of Zhang and Lavaei [39] denoted as ZL17. In Section 7.3, we confirm that our method consistently reaches accuracies of $\mu \approx 10^{-12}$, whereas ZL17 is fundamentally limited to accuracies of $\mu \approx 10^{-6}$.

All of our experiments are performed on an Apple laptop in MATLAB R2021a, running a silicon M1 pro chip with 10-core CPU, 16-core GPU, and 32GB of RAM. In each case, the ground truth is chosen as $X^* = GG^T$ in which the elements of $G \in \mathbb{R}^{n \times r^*}$ are sampled $\operatorname{vec}(G) \sim \mathcal{N}(0, I_{nr^*})$. A small

number m_o of Gaussian outliers ε are added to the measurements, meaning that $b = \mathcal{A}(X^*) + \varepsilon$ where $\varepsilon_i = 0$ except m_o randomly sampled elements with $\epsilon_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 10^4)$. We implement our method by modifying the source code of SeDuMi [30] to use (7) to solve (Newt- ϵ) while keeping all other parts unchanged. Specifically, we decompose the scaling point W at each iteration as in (6) with the rank parameter $r = \arg \max_{i \leq \hat{r}} \lambda_i(W) / \lambda_{i+1}(W)$ for some fixed \hat{r} , and then solve (7) using PCG as in Corollary 3.5 with $\beta = 3$. We choose to modify SeDuMi in order to be consistent with ZL17.

7.1 Validation of our theoretical results

We start by validating Assumptions 1, 2 and 3 for the SDP reformulation (3). It is obvious that Assumption 3 holds for our choice of measurement operator $\mathcal{A}(X) = P_m \operatorname{vec}_{\mathcal{S}}(X)$ because for any $U, V \in \mathbb{R}^{n \times r}$ and $Y \in \mathcal{S}^n, y \in \mathbb{R}^m$ such that $\operatorname{vec}_{\mathcal{S}}(Y) = P_m^T y$, both $\mathcal{A}(UV^T + VU^T) =$ $P_m \operatorname{vec}_{\mathcal{S}}(UV^T + VU^T)$ and $\mathcal{A}^T(y)U = YU$ can be performed in $O(n^2r)$ time and $O(n^2)$ storage. In Figure 7.1, using a small-scaled robust matrix completion problem on a rank-2, size 50 × 50 ground truth matrix, with 1275 constraints and 25 outliers, we empirically verify Assumption 1 holds with $\delta = 0.9$ and L = 10, and Assumption 2 holds with $\chi_2 = 2.7$. For this result, we run **our method** and extract the primal variable X, dual variable S, and scaling point W at each iteration. For each X, S and W, the corresponding μ is chosen such that $\|\frac{1}{\mu}X^{1/2}SX^{1/2} - I\| = 0.1$, and the corresponding matrix **Q** is constructed as in (6) with the rank parameter $r = \arg \max_{i \leq 5} \lambda_i(W)/\lambda_{i+1}(W)$.




Figure 2: Verifying Assumption 1 and Assumption 2 hold for robust matrix completion problem (3). Left. Verifying Assumption 1, $\|\frac{1}{\mu}X^{1/2}SX^{1/2} - I\| \leq \delta$, $\|X - X^*\| \leq L\mu$ and $\|S - S^*\| \leq L$, holds for (3) with $\delta = 0.9$ and L = 10. Right. Verifying Assumption 2, $\min_x \|(\mathbf{A}\mathbf{A}^T)^{-1/2}\mathbf{A}\mathbf{Q}x\|/\|\mathbf{Q}x\| \geq \chi_2^{-1}$, holds for (3) with $\chi_2 = 2.7$.

As the three assumptions hold for (3), Theorem 3.3 provides an upper bound on the condition number of the corresponding augmented system (7), which is independent of μ . In contrast, the condition number of KKT (Newt- ϵ) explodes as $\mu \to 0$. In the left-hand-side of Figure 7.1, we plot the condition number of the augmented system (7) and KKT equations (Newt- ϵ) with respect to different values of μ , as well as the upper bound predicted in Theorem 3.3. Here, we use the same choice of W and μ in Figure 7.1. For each pair of W and μ , the corresponding augmented system is constructed as in (7) using the rank parameter $r = \arg \max_{i \leq 5} \lambda_i(W)/\lambda_{i+1}(W)$, and the corresponding KKT equations is constructed as in (Newt- ϵ). We see that the condition number of the augmented system is upper bounded by the bound predicted in Theorem 3.3. On the other hand, the condition number of the KKT equations explodes as $\mu \to 0$ (we note that it get stuck at 10^{25} due to finite precision).

Corollaries 3.4 and 3.5 show that the system can be efficiently solved using either MINRES or PCG. In practice, PCG is much more preferable as the preconditioner rapidly speeds up the convergence compared to MINRES. In the right-hand-side of Figure 7.1, we plot the relative residual of MINRES and PCG for solving the augmented system (7) obtained using different values of rank parameter $r = \{2, 5, 10, 15, 20\}$. For this result, we run **our method** with a fixed rank parameter r and extract the augmented system at the last iteration. We then solve the system using both MINRES (as in Corollary 3.4) and PCG (as in Corollary 3.5) with the same initial point. As predicted in Corollaries 3.4 and 3.5, PCG converges significantly faster than MINRES.

Finally, we verify Corollary 3.5 that our method achieves cubic time complexity for robust matrix completion problems when rank $(X^*) \ll n$, with no dependence on the number of constraints m. In this experiment, we consider two cases, m = O(n) and $m = O(n^2)$, to verify the time complexity of our method is indeed invariant to m. In Figure 7.1, we plot the average setup time for the augmented system (7) and the average per PCG iteration time against the size of the ground truth n. In particular, the setup time includes the time to decompose $W = Q\Lambda Q^T + Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$ and construct $E = QQ^T + \tau^{-1}Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$ in order to implicitly setup \mathbf{Q} , \mathbf{E} and explicitly construct Σ as in (6), and explicitly form the Schur complement $\mathbf{C} = \tau^2 \Sigma + \beta^{-1} \mathbf{Q}^T \mathbf{A}^T \mathbf{A} \mathbf{Q}$ via O(nr) matrix-vector product with $\mathbf{A}\mathbf{Q}$ and $\mathbf{Q}^T \mathbf{A}^T$ and compute its Cholesky factor. In order to accurately measure the cubic time complexity, we fix $m_o = 0$ and set rank $(X^*) = 2$ so that rank $(X^*) \ll n$ for sufficiently large n. We set the number of constraints to be m = 20n for the case of m = O(n), and m = n(n+1)/2for the case of $m = O(n^2)$. As shown in Figure 7.1, the setup time and per PCG time exhibit cubic time complexity for both cases.

7.2 Comparison against other approaches

We compare the practical performance of our method against fmincon, knitro, GD, and also the prior approach of Zhang and Lavaei [39] denoted as ZL17. We start by demonstrating the efficiency and efficacy of our method by solving a robust matrix completion problem on a rank-2, size 200×200




Figure 3: Verifying Theorem 3.3, Corollary 3.4 and Corollary 3.5 on robust matrix completion problem (3). Left. Verifying the condition number upper bound predicted in Theorem 3.3 for the augmented system (7). The upper bound is independent of μ . In contrast, the condition number of the KKT equations (Newt- ϵ) explodes as $\mu \to 0$. Right. Convergence of PCG and MINRES for solving augmented systems (7) constructed using different values of rank parameter r. As predicted in Corollaries 3.4 and 3.5, PCG converges significantly faster then MINRES.




Figure 4: Verifying cubic time complexity of our method for solving rank-2 robust matrix completion problems (3) with m = O(n) and $m = O(n^2)$ number of constraints. In both cases, our method exhibits cubic time complexity. Left. Average setup time and per PCG iteration time for solving (3) with m = O(n) constraints. Right. Average setup time and per PCG iteration time for solving (3) with $m = O(n^2)$ constraints.

ground truth matrix X^* , with m = 20100 constraints and $m_o = 100$ outliers. In Figure 1, we plot the reconstruction error $||X - X^*||_F$ against runtime for our method, ZL17, fmincon, knitro and GD. Here, the maximum rank parameter \hat{r} for our method, and the search rank r for fmincon, knitro and GD are all set to be r = 15. We initialized fmincon, knitro and GD at the same initial point that is drawn from standard Gaussian. The step-size of GD is set be $\alpha = 10^{-6}$. From Figure 1, we see that our method is the only one that is able to achieve reconstruction error below 10^{-8} . Though ZL17 achieves reconstruction error 10^{-6} at the fastest rate, it unfortunately becomes illconditioned and therefore is unable to make any progress. Comparing to nonlinear programming solvers, both fmincon and knitro eventually reach the same reconstruction error similar to ZL17 but at a significantly slower rate than our method. Finally, GD converges extremely slowly for this problem.

We also compare the reconstruction error and runtime for solving several instances of rank-2 robust matrix completion problem to demonstrate that **our method** can consistently achieve the smallest reconstruction error under reasonable runtime. The results are shown in Table 1. Here, the maximum rank parameter \hat{r} for **our method**, and the search rank r for fmincon, knitro and GD are all set to be min $\{n/10, 15\}$. We run fmincon and knitro for 1000 iterations, and GD for 300,000 iterations, starting at the same initial point that is drawn from unit Gaussian. For these three methods, we report their best reconstruction error and the time they achieve their best reconstruction error in Table 1. We note that we tune and use the best possible step-size for GD in each case.

7.3 Comparison of accuracy vs ZL17

We confirm that our method consistently reaches accuracies of $\mu \approx 10^{-12}$, whereas ZL17 is fundamentally limited to accuracies of $\mu \approx 10^{-6}$ due to ill-conditioning. In Table 2, we show the number of cumulative PCG iterations that our method and ZL17 perform before achieving accuracies of $\mu \approx 10^{-2}, 10^{-4}, 10^{-6}, 10^{-8}, 10^{-10}$ and 10^{-12} . The number of cumulative PCG iterations is ∞ if the corresponding accuracy is not achievable by the algorithm. For this result, the simulation settings

n	m	m_o	our method		ZL17		fmincon		knitro		GD	
			Error	Time	Error	Time	Error	Time	Error	Time	Error	Time
50 50 50	800 1000 1200	6 12 24	2.8e-12 2.5e-12 1.6e-12	1.3s 2.4s 3s	2.1e-09 8.0e-06 1.1e-05	1.1s 0.8s 1.0s	3.7e-07 1.6e-08 5.2e-10	8s 15s 125s	4.2e-08 1.1e-09 5.1e-10	1s 1.5s 2.4s	1.4e-02 1.3e-02 2.0e-02	53s 62s 70s
100 100 100	3000 4000 5000	12 24 48	2.2e-12 8.7e-12 5.2e-12	6.2s 7.3s 11s	9.7e-06 7.9e-07 1.1e-05	2.0s 2.2s 2.6s	1.2e-06 5.9e-07 1.3e-06	384s 231s 301s	1.3e-09 3.2e-09 7.3e-09	103s 111s 339s	3.5e-02 5.4e-02 9.5e-02	56s 69s 97s
200	12000 16000 20000	50	2.9e-12 3.3e-12 4.0e-12	20s 53s 52s	1.6e-06 2.0e-05 1.0e-05	8.0s 14s 15s	1.2e-02 4.6e-02 9.1e-07	846s 920s 436s	1.7e-03 1.3e-03 1.3e-03	$\begin{array}{c} 475s \\ 442s \\ 390s \end{array}$	1.3e-01 1.6e-01 2.4e-01	258s 195s 198s

Table 1: Reconstruction error (Error) and runtime (Time) for solving a size $n \times n$ robust matrix completion problem (3) with m constraints and m_o outliers. Our method consistently achieves reconstruction error around 10^{-12} in all cases.

for our method and ZL17 is exactly the same as those in Table 1, and μ is selected using the same procedure in Figure 7.1. As shown in Table 2, we see that our method is able to achieve the same accuracies μ compared to ZL17 using fewer or similar number of cumulative PCG iterations. Most importantly, our method consistently achieves accuracies $\mu \approx 10^{-12}$ under reasonable cumulative PCG iteration while ZL17 usually fails at $\mu \approx 10^{-6}$.

Table 2: The number of cumulative PCG iterations required for our method and ZL17 to achieve accuracies of $\mu \approx 10^{-2}, 10^{-4}, 10^{-6}, 10^{-8}, 10^{-10}$ and 10^{-12} when solving a size $n \times n$ robust matrix completion problem (3) with m constraints and m_o outliers. The number of iterations is ∞ if the corresponding accuracy is not achievable. Our method consistently achieves $\mu \approx 10^{-12}$ under reasonable number of PCG iterations while ZL17 fail to achieve accuracies below 10^{-6} in all cases.

n	m	m_o	our method						ZL17						
			1e-2	1e-4	1e-6	1e-8	1e-10	1e-12	1e-2	1e-4	1e-6	1e-8	1e-10	1e-12	
50	800	6	452	614	718	878	957	1038	290	614	878	∞	∞	∞	
50	1000	12	285	559	901	1250	1537	1826	213	559	833	∞	∞	∞	
50	1200	24	331	722	1048	1257	1362	2210	331	644	1048	∞	∞	∞	
100	3000	12	327	491	864	996	1128	1328	360	760	∞	∞	∞	∞	
100	4000	24	403	655	905	1338	1681	2027	319	854	1775	∞	∞	∞	
100	5000	48	390	563	899	1081	1198	1549	584	1111	2339	∞	∞	∞	
200	10000	25	307	598	764	840	995	1511	386	491	804	∞	∞	∞	
200	15000	50	404	655	1244	1934	2399	3182	404	890	1553	∞	∞	∞	
200	20000	100	453	549	922	1679	2193	3105	453	643	1015	∞	∞	∞	

References

- [1] F. ALIZADEH, J.-P. A. HAEBERLY, AND M. L. OVERTON, Complementarity and nondegeneracy in semidefinite programming, Mathematical Programming, 77 (1997), pp. 111–128.
- [2] T. ANDO, Concavity of certain maps on positive definite matrices and applications to hadamard products, Linear Algebra and its Applications, 26 (1979), pp. 203–241.
- [3] N. BOUMAL, Nonconvex phase synchronization, SIAM Journal on Optimization, 26 (2016), pp. 2355–2377.
- [4] T. BOUWMANS AND E. H. ZAHZAH, Robust pca via principal component pursuit: A review for a comparative evaluation in video surveillance, Computer Vision and Image Understanding, 122 (2014), pp. 22–34.
- [5] S. BURER AND R. D. MONTEIRO, A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization, Math. Program., 95 (2003), pp. 329–357.
- [6] R. H. BYRD, J. NOCEDAL, AND R. A. WALTZ, K nitro: An integrated package for nonlinear optimization, Large-scale nonlinear optimization, (2006), pp. 35–59.
- [7] J.-F. CAI, E. J. CANDÈS, AND Z. SHEN, A singular value thresholding algorithm for matrix completion, SIAM Journal on Optimization, 20 (2010), pp. 1956–1982.
- [8] E. CANDÈS AND B. RECHT, *Exact matrix completion via convex optimization.*, Foundations of Computational Mathematics, 9 (2009).
- [9] E. J. CANDÈS, X. LI, Y. MA, AND J. WRIGHT, Robust principal component analysis?, Journal of the ACM (JACM), 58 (2011), pp. 1–37.
- [10] E. J. CANDES AND Y. PLAN, Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements, IEEE Transactions on Information Theory, 57 (2011), pp. 2342–2359.
- [11] E. J. CANDES, T. STROHMER, AND V. VORONINSKI, Phaselift: Exact and stable signal recovery from magnitude measurements via convex programming, Communications on Pure and Applied Mathematics, 66 (2013), pp. 1241–1274.
- [12] H.-M. CHIU AND R. Y. ZHANG, Tight certification of adversarially trained neural networks via nonconvex low-rank semidefinite relaxations, in International Conference on Machine Learning, PMLR, 2023, pp. 5631–5660.
- [13] L. DING, L. JIANG, Y. CHEN, Q. QU, AND Z. ZHU, Rank overspecified robust matrix recovery: Subgradient method and exact recovery, Advances in Neural Information Processing Systems, 34 (2021), pp. 26767–26778.
- [14] M. FUKUDA, M. KOJIMA, K. MUROTA, AND K. NAKATA, Exploiting sparsity in semidefinite programming via matrix completion I: General framework, SIAM J. Optim., 11 (2001), pp. 647– 674.
- [15] N. I. GOULD, Iterative methods for ill-conditioned linear systems from optimization, Springer, 2000.

- [16] A. GREENBAUM, Iterative methods for solving linear systems, SIAM, 1997.
- [17] S. HABIBI, M. KOČVARA, AND M. STINGL, Loraine-an interior-point solver for low-rank semidefinite programming, Optimization Methods and Software, (2023), pp. 1–31.
- [18] S. KIM, M. KOJIMA, M. MEVISSEN, AND M. YAMASHITA, Exploiting sparsity in linear and nonlinear matrix inequalities via positive semidefinite matrix completion, Math. Program., 129 (2011), pp. 33–68.
- [19] L. LUKŠAN AND J. VLČEK, Indefinitely preconditioned inexact newton method for large sparse equality constrained non-linear programming problems, Numerical linear algebra with applications, 5 (1998), pp. 219–247.
- [20] Z.-Q. LUO, J. F. STURM, AND S. ZHANG, Superlinear convergence of a symmetric primaldual path following algorithm for semidefinite programming, SIAM Journal on Optimization, 8 (1998), pp. 59–81.
- [21] J. MA AND S. FATTAHI, Global convergence of sub-gradient method for robust matrix recovery: Small initialization, noisy measurements, and over-parameterization, Journal of Machine Learning Research, 24 (2023), pp. 1–84.
- [22] S. MANIU, P. SENELLART, AND S. JOG, An experimental study of the treewidth of realworld graph data, in 22nd International Conference on Database Theory (ICDT 2019), Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2019.
- [23] Matlab optimization toolbox, 2021. The MathWorks, Natick, MA, USA.
- [24] MOSEK APS, The MOSEK optimization toolbox for MATLAB manual. Version 7.1 (Revision 28)., 2015.
- [25] Y. E. NESTEROV AND M. J. TODD, Self-scaled barriers and interior-point methods for convex programming, Mathematics of Operations research, 22 (1997), pp. 1–42.
- [26] —, Primal-dual interior-point methods for self-scaled cones, SIAM Journal on optimization, 8 (1998), pp. 324–364.
- [27] B. RECHT, M. FAZEL, AND P. A. PARRILO, Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization, SIAM review, 52 (2010), pp. 471–501.
- [28] D. M. ROSEN, L. CARLONE, A. S. BANDEIRA, AND J. J. LEONARD, Se-sync: A certifiably correct algorithm for synchronization over the special euclidean group, The International Journal of Robotics Research, 38 (2019), pp. 95–125.
- [29] M. ROZLOZNÍK AND V. SIMONCINI, Krylov subspace methods for saddle point problems with indefinite preconditioning, SIAM journal on matrix analysis and applications, 24 (2002), pp. 368– 391.
- [30] J. F. STURM, Using SeDuMi 1.02, a MATLAB toolbox for optimization over symmetric cones, Optimization methods and software, 11 (1999), pp. 625–653.
- [31] K.-C. TOH, Solving large scale semidefinite programs via an iterative solver on the augmented systems, SIAM Journal on Optimization, 14 (2004), pp. 670–698.

- [32] —, An inexact primal-dual path following algorithm for convex quadratic sdp, Mathematical programming, 112 (2008), pp. 221–254.
- [33] K.-C. TOH AND M. KOJIMA, Solving some large scale semidefinite programs via the conjugate residual method, SIAM Journal on Optimization, 12 (2002), pp. 669–691.
- [34] K.-C. TOH, M. J. TODD, AND R. H. TÜTÜNCÜ, Sdpt3-a matlab software package for semidefinite programming, version 1.3, Optimization methods and software, 11 (1999), pp. 545–581.
- [35] L. VANDENBERGHE AND S. BOYD, A primal-dual potential reduction method for problems involving matrix inequalities, Mathematical programming, 69 (1995), pp. 205–236.
- [36] H. YANG, L. LIANG, L. CARLONE, AND K.-C. TOH, An inexact projected gradient method with rounding and lifting by nonlinear programming for solving rank-one semidefinite relaxation of polynomial optimization, Mathematical Programming, 201 (2023), pp. 409–472.
- [37] A. YURTSEVER, J. A. TROPP, O. FERCOQ, M. UDELL, AND V. CEVHER, Scalable semidefinite programming, SIAM Journal on Mathematics of Data Science, 3 (2021), pp. 171–200.
- [38] R. Y. ZHANG, Parameterized complexity of chordal conversion for sparse semidefinite programs with small treewidth, arXiv preprint arXiv:2306.15288, (2023).
- [39] R. Y. ZHANG AND J. LAVAEI, Modified interior-point method for large-and-sparse low-rank semidefinite programs, in IEEE 56th Conference on Decision and Control, 2017.
- [40] —, Sparse semidefinite programs with guaranteed near-linear time complexity via dualized clique tree conversion, Mathematical programming, 188 (2021), pp. 351–393.

A Proof of Proposition 3.2

We break the proof into three parts. Below, recall that we have defined Ψ_n as the basis matrix for $\operatorname{vec}(\mathcal{S}^n)$ on $\operatorname{vec}(\mathbb{R}^{n\times n}) = \mathbb{R}^{n^2}$, in order to define $\operatorname{vec}_{\mathcal{S}}(X) \stackrel{\text{def}}{=} \Psi_n^T \operatorname{vec}(X) = \frac{1}{2} \Psi_n^T \operatorname{vec}(X + X^T)$ for $X \in \mathbb{R}^{n \times n}$, and $A \otimes_{\mathcal{S}} B = B \otimes_{\mathcal{S}} A \stackrel{\text{def}}{=} \frac{1}{2} \Psi_n^T (A \otimes B + B \otimes A) \Psi_r$ for $A, B \in \mathbb{R}^{n \times r}$.

Lemma A.1 (Correctness). Let $W = Q\Lambda Q^T + Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$ where $[Q, Q_{\perp}]$ is orthonormal, and pick $0 < \tau < \lambda_{\min}(\Lambda)$. Define $E = QQ^T + \tau^{-1} \cdot Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$, $\mathbf{Q}_B = Q \otimes_{\mathcal{S}} Q$, $\mathbf{Q}_N = \sqrt{2}\Psi_n^T(Q \otimes Q_{\perp})$, $\mathbf{\Sigma}_B^{-1} = \Lambda \otimes_{\mathcal{S}} \Lambda - \tau^2 I$, and $\mathbf{\Sigma}_N^{-1} = (\Lambda - \tau I) \otimes \Lambda_{\perp}$. Then

$$W \otimes_{\mathcal{S}} W = [\mathbf{Q}_B, \mathbf{Q}_N] \begin{bmatrix} \mathbf{\Sigma}_B & \\ & \mathbf{\Sigma}_N \end{bmatrix}^{-1} [\mathbf{Q}_B, \mathbf{Q}_N]^T + \tau^2 \cdot E \otimes_{\mathcal{S}} E$$

Proof. Write $W_X = \tau \cdot Q(\Lambda - \tau I)Q^T$ and $W_S = \tau^{-1}Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$. For arbitrary $Y \in S^n$, it follows from $W = \tau E + \tau^{-1}W_X$ that

$$\Psi_n^T(W \otimes W)\Psi_n \operatorname{vec}_{\mathcal{S}}(Y) = \operatorname{vec}_{\mathcal{S}}((\tau E + \tau^{-1}W_X)Y(\tau E + \tau^{-1}W_X))$$
$$= \operatorname{vec}_{\mathcal{S}}\left[\tau^2 \cdot EYE + 2 \cdot EYW_X + \tau^{-2} \cdot W_XYW_X\right].$$

Now, substituting $E = QQ^T + W_S$ yields our desired claim:

$$= \operatorname{vec}_{\mathcal{S}} \left[\tau^{2} \cdot EYE + 2 \cdot W_{S}YW_{X} + \left(2QQ^{T} + \tau^{-2}W_{X} \right) YW_{X} \right]$$
$$= \tau^{2} \cdot \underbrace{\operatorname{vec}_{\mathcal{S}}(EYE)}_{\mathbf{E}y} + \underbrace{\operatorname{vec}_{\mathcal{S}}(2W_{S}YW_{X})}_{\mathbf{Q}_{N}\boldsymbol{\Sigma}_{N}^{-1}\mathbf{Q}_{N}^{T}y} + \underbrace{\operatorname{vec}_{\mathcal{S}} \left[\left(2QQ^{T} + \tau^{-2}W_{X} \right) YW_{X} \right]}_{\mathbf{Q}_{B}\boldsymbol{\Sigma}_{B}^{-1}\mathbf{Q}_{B}^{T}y}.$$

Indeed, we verify that $\operatorname{vec}(2W_SYW_X) = (\sqrt{2}Q \otimes Q_{\perp})[(\Lambda - \tau I) \otimes \Lambda_{\perp}](\sqrt{2}Q \otimes Q_{\perp})^T \operatorname{vec}(Y) = \mathbf{Q}_N[(\Lambda - \tau I) \otimes \Lambda_{\perp}]\mathbf{Q}_N^T \operatorname{vec}(Y)$ and $\operatorname{vec}_{\mathcal{S}}\left[\left(2QQ^T + \tau^{-2}W_X\right)YW_X\right] = \mathbf{Q}_B[(\Lambda + \tau I) \otimes_{\mathcal{S}}(\Lambda - \tau I)]\mathbf{Q}_B^T \operatorname{vec}_{\mathcal{S}}(Y)$, where we used $(A + B) \otimes_{\mathcal{S}} (A - B) = A \otimes_{\mathcal{S}} A - B \otimes_{\mathcal{S}} B$ because $A \otimes_{\mathcal{S}} B = B \otimes_{\mathcal{S}} A$.

Lemma A.2 (Column span of **Q**). Let $[Q, Q_{\perp}]$ be orthonormal with $Q \in \mathbb{R}^{n \times r}$. Define **Q** = $[\mathbf{Q}_B, \mathbf{Q}_N]$ where $\mathbf{Q}_B = Q \otimes_S Q$ and $\mathbf{Q}_N = \sqrt{2} \Psi_n^T (Q \otimes Q_{\perp})$. Then, $\operatorname{colsp}(\mathbf{Q}) = \{\operatorname{vec}_S(VQ^T) : V \in \mathbb{R}^{n \times r}\}$ and $\mathbf{Q}^T \mathbf{Q} = I_d$ where $d = nr - \frac{1}{2}r(r-1)$.

Proof. Without loss of generality, let $[Q, Q_{\perp}] = I_n$. We observe that

$$\mathbf{Q}\begin{bmatrix}\operatorname{vec}_{\mathcal{S}}(H_1)\\\operatorname{vec}(H_2)\end{bmatrix} = \operatorname{vec}_{\mathcal{S}}\left(\begin{bmatrix}H_1 & \frac{1}{\sqrt{2}}H_2^T\\\frac{1}{\sqrt{2}}H_2 & 0\end{bmatrix}\right) = \operatorname{vec}_{\mathcal{S}}\left(\begin{bmatrix}H_1\\\sqrt{2}H_2\end{bmatrix}Q^T\right)$$

holds for any arbitrary (possibly nonsymmetric) $H_1 \in \mathbb{R}^{r \times r}$ and $H_2 \in \mathbb{R}^{(n-r) \times r}$, and that $\|\mathbf{Q}h\| = \|h\|$ for all $h \in \mathbb{R}^d$, so $\mathbf{Q}^T \mathbf{Q} = I_d$.

Lemma A.3 (Eigenvalue bounds on **E** and **\Sigma**). Given $W \in S_{++}^n$, write $w_i \equiv \lambda_i(W)$ for $i \in \{1, 2, ..., n\}$. Choose $r \geq 1$ and $\tau = \frac{1}{2}w_{r+1}$, and define $\mathbf{E} = E \otimes_{\mathcal{S}} E$ where $E = QQ^T + \tau^{-1} \cdot Q_{\perp}\Lambda_{\perp}Q_{\perp}^T$, and $\mathbf{\Sigma} = \operatorname{diag}(\mathbf{\Sigma}_B, \mathbf{\Sigma}_N)$ where $\mathbf{\Sigma}_B^{-1} = \Lambda \otimes_{\mathcal{S}} \Lambda - \tau^2 I$, and $\mathbf{\Sigma}_N^{-1} = (\Lambda - \tau I) \otimes \Lambda_{\perp}$. Then,

$$4 \geq \lambda_{\max}(\mathbf{E}) \geq \lambda_{\min}(\mathbf{E}) \geq w_n^2/w_{r+1}^2, w_{r+1}^2/(w_n w_r) \geq \tau^2 \cdot \lambda_{\max}(\mathbf{\Sigma}) \geq \tau^2 \cdot \lambda_{\min}(\mathbf{\Sigma}) \geq w_n^2/(4w_1^2).$$

Proof. We have $\lambda_{\max}(\mathbf{E}) = \lambda_{\max}^2(E)$ and $\lambda_{\max}(E) = \max\{1, \lambda_{\max}(\Lambda_{\perp})/\tau\} = \max\{1, 2w_{r+1}/w_{r+1}\} \leq 2$. Moreover, $\lambda_{\min}(\mathbf{E}) = \lambda_{\min}^2(E)$ and $\lambda_{\min}(E) = \min\{1, \lambda_{\min}(\Lambda_{\perp})/\tau\} = \min\{1, 2w_n/w_{r+1}\} \geq w_n/w_{r+1}$. We have $\tau^{-2}\lambda_{\max}^{-1}(\mathbf{\Sigma}) \geq (\frac{1}{2}w_rw_n)/(\frac{1}{2}w_{r+1})^2 \geq w_rw_n/w_{r+1}^2$, because $\lambda_{\max}^{-1}(\mathbf{\Sigma}) = \min\{\lambda_{\max}^{-1}(\mathbf{\Sigma}_B), \lambda_{\max}^{-1}(\mathbf{\Sigma}_N)\}$ and $\lambda_{\max}^{-1}(\mathbf{\Sigma}_B) = \lambda_{\min}(\Lambda \otimes_S \Lambda - \tau^2 I) \geq w_r^2 - \frac{1}{4}w_{r+1}^2 \geq \frac{3}{4}w_r^2$, while $\lambda_{\max}^{-1}(\mathbf{\Sigma}_N) = \lambda_{\min}[(\Lambda - \tau I) \otimes \Lambda_{\perp}] \geq (w_r - \frac{1}{2}w_{r+1})w_n \geq \frac{1}{2}w_rw_n$. Finally, we have $\tau^{-2}\lambda_{\min}^{-1}(\mathbf{\Sigma}) \leq w_1^2/(\frac{1}{2}w_{r+1})^2 \leq 4w_1^2/w_{r+1}^2$, because $\lambda_{\min}^{-1}(\mathbf{\Sigma}) = \max\{\lambda_{\min}^{-1}(\mathbf{\Sigma}_B), \lambda_{\min}^{-1}(\mathbf{\Sigma}_N)\} \leq w_1^2$.