

# Two Infinite Families of Solutions for Singular Superlinear Equations on Exterior Domains

Abdeali Diwan, Joseph Iaia

University of North Texas

July 22, 2024

## Abstract

In this paper, we study radial solutions of  $\Delta u + K(|x|)f(u) = 0$  in the exterior of the ball of radius  $R > 0$  in  $\mathbb{R}^N$  with  $N > 2$  where  $f$  grows superlinearly at infinity and is singular at 0 with  $f \sim \frac{1}{|u|^{q-1}u}$  where  $0 < q < 1$ . We also assume  $K(r) \sim |r|^{-\alpha}$  for large  $r$  and establish the existence of two infinite families of solutions when  $N + q(N - 2) < \alpha < 2(N - 1)$ .

## 1 Introduction

In this paper, we are interested in radial solutions of

$$\Delta u + K(|x|)f(u) = 0 \text{ on } \mathbb{R}^N \setminus B_R, \quad (1)$$

$$u|_{\partial B_R} = 0, \quad \lim_{|x| \rightarrow \infty} u(x) = 0 \quad (2)$$

when  $N > 2$  and where  $B_R$  is the ball of radius  $R > 0$  centered at the origin. Assuming  $u(x) = u(|x|) = u(r)$  this becomes

$$u'' + \frac{N-1}{r}u' + K(r)f(u) = 0 \text{ for } r > R, \quad (3)$$

$$u(R) = 0, \quad \lim_{r \rightarrow \infty} u(r) = 0. \quad (4)$$

We make the following assumptions on  $f$ . We assume  $f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$  is odd, locally Lipschitz, and

$$f(u) = |u|^{p-1}u + g(u) \text{ for } p > 1 \text{ where } \lim_{|u| \rightarrow \infty} \frac{g(u)}{|u|^p} = 0, \quad (\text{H1})$$

$$f(u) = \frac{1}{|u|^{q-1}u} + g_1(u) \text{ for } 0 < q < 1 \text{ and } g_1 \text{ is locally Lipschitz with } g_1(0) = 0. \quad (\text{H2})$$

We also define  $F(u) = \int_0^u f(t) dt$ . Since  $f$  is integrable at 0 it follows that  $F$  is continuous and  $F(0) = 0$ . In addition, since  $f$  is odd then  $F$  is even and since  $f(u) > 0$  for  $u > 0$  it follows that  $F(u) > 0$  for  $u \neq 0$ . We also assume  $K > 0$  and  $K'$  are continuous on  $[R, \infty)$  and there are positive constants  $K_0, K_1$ , and  $\alpha$  such that

$$K_0 r^{-\alpha} \leq K \leq K_1 r^{-\alpha}, \quad 2(N-1) > -\frac{rK'}{K}, \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{rK'}{K} = -\alpha \text{ with } N+q(N-2) < \alpha < 2(N-1). \quad (\text{H3})$$

**Main Theorem:** Under the assumptions (H1)-(H3) there exist two infinite families of sign-changing solutions to (3)-(4) on  $[R, \infty)$ .

Numerous papers have proved the existence of *positive* solutions of these equations with various nonlinearities  $f(u)$  and for various functions  $K(|x|) \sim |x|^{-\alpha}$  with  $\alpha > 0$ . See for example [4]-[5]. We will prove the existence of two infinite families of *sign-changing* solutions.

## 2 Preliminaries

Let  $R > 0$  and define  $R_1 = R^{2-N}$ . We begin our analysis by first making the change of variables  $u(r) = v(r^{2-N}) = v(t)$  to (3)-(4) and obtaining

$$v''(t) + h(t)f(v(t)) = 0 \text{ for } 0 < t < R_1, \quad (5)$$

$$v(0) = 0, v(R_1) = 0 \quad (6)$$

where

$$h(t) = \frac{t^{\frac{2(N-1)}{2-N}} K(t^{\frac{1}{2-N}})}{(N-2)^2}. \quad (7)$$

It follows from (H3) that  $h' < 0$  and  $\lim_{t \rightarrow 0^+} \frac{th'}{h} = -\tilde{\alpha}$  where  $\tilde{\alpha} = \frac{2(N-1)-\alpha}{N-2}$ . In addition, (H3) implies there exist positive constants  $h_0$  and  $h_1$  such that

$$h_0 t^{-\tilde{\alpha}} \leq h \leq h_1 t^{-\tilde{\alpha}} \text{ on } (0, R_1]. \quad (8)$$

Also from (H3) it follows that

$$0 < \tilde{\alpha} < 1 - q < 1. \quad (9)$$

We now attempt to solve the initial value problem

$$v'' + h(t)f(v) = 0 \text{ for } 0 < t < R_1, \quad (10)$$

$$v(0) = 0, \quad v'(0) = a > 0 \quad (11)$$

and then try to find values of  $a$  so that  $v(R_1) = 0$ .

### 3 Existence of Solutions

#### 3.1. Integral Equation Formulation

First, we prove the existence of a solution of equations (10)-(11) on  $[0, \epsilon]$  for some  $\epsilon > 0$ . In order to do this, we reformulate (10)-(11) as an appropriate integral equation. Assuming  $v$  is a solution of (10)-(11), then integrating (10) on  $(0, t)$  and using (11) gives

$$v' + \int_0^t h(s)f(v) ds = a. \quad (12)$$

Integrating again on  $(0, t)$  gives

$$v + \int_0^t \int_0^s h(x)f(v) dx ds = at. \quad (13)$$

To ensure the integral in (12)-(13) is defined, we observe it follows from (11) that for sufficiently small  $t > 0$

$$\frac{at}{2} < v < at.$$

Furthermore, from conditions (H1)-(H2), it follows that there exists a constant  $f_1 > 0$  such that  $0 < f(v) \leq f_1(v^{-q} + v^p)$  for  $v > 0$ . Also, it follows from this and (8) that for  $v > 0$

$$0 < h(t)f(v(t)) \leq f_1 h_1 (t^{-\tilde{\alpha}} v^{-q} + t^{-\tilde{\alpha}} v^p) \leq f_1 h_1 \left( \left( \frac{a}{2} \right)^{-q} t^{-\tilde{\alpha}-q} + a^p t^{-\tilde{\alpha}+p} \right). \quad (14)$$

It follows from (9) and (H1) that  $1 - \tilde{\alpha} - q > 0$  and  $1 - \tilde{\alpha} + p > 0$ . Thus it follows from (14) that  $h(t)f(v)$  is integrable near  $t = 0$ , and thus the integrals in (12) and (13) are defined and are continuous functions.

#### 3.2. Contraction Mapping Principle

Using (H2), we see that (13) is equivalent to

$$v + \int_0^t \int_0^s h(x) \left( \frac{1}{v^q} + g_1(v) \right) dx ds = at. \quad (15)$$

Substituting  $v = tw$  into (15) gives

$$w = a - \frac{1}{t} \int_0^t \int_0^s h(x) \left( \frac{1}{x^q w^q} + g_1(xw) \right) dx ds. \quad (16)$$

We also define  $w(0) = a$ .

Next let  $\epsilon > 0$ ,  $a > 0$ , and

$$X = \{w \in C[0, \epsilon] \mid w(0) = a \text{ and } \|w - a\| \leq \frac{a}{2}\}$$

where  $C[0, \epsilon]$  is the set of continuous functions on  $[0, \epsilon]$  and  $\|\cdot\|$  is the supremum norm, i.e.

$$\|w\| = \sup_{[0, \epsilon]} |w(x)|.$$

It is well-known that  $(X, \|\cdot\|)$  is a Banach space [1]. We now define  $T : X \rightarrow C[0, \epsilon]$

$$Tw = a - \frac{1}{t} \int_0^t \int_0^s h(x) \left( \frac{1}{x^q w^q} + g_1(xw) \right) dx ds.$$

It can be shown that  $T : X \rightarrow X$  and  $T$  is a contraction for  $\epsilon$  sufficiently small, say for  $\epsilon = \epsilon_0$ . These details are carried out in [2].

The contraction mapping principle then ensures the existence of a unique  $w$  in  $X$  such that  $Tw = w$ , by choosing any function  $w_1$  in  $X$  and then defining

$$w_{n+1} = T(w_n).$$

Using the fact that  $T$  is a contraction it follows that  $\{w_n\}$  forms a Cauchy sequence in  $X$  and hence converges to some  $w \in X$  with

$$w = \lim_{n \rightarrow \infty} w_{n+1} = \lim_{n \rightarrow \infty} T(w_n) = Tw.$$

Thus  $w$  is a fixed point of  $T$  on  $[0, \epsilon_0]$ .

Finally letting  $v = tw$  then we see that  $v$  is a solution of (10)-(11) on  $[0, \epsilon_0]$ .

We henceforth denote this solution as  $v_a$ .

### 3.3. Existence and Uniqueness of Solutions on $[0, R_1]$

We now show that the solution  $v_a$  is defined on all of  $[0, R_1]$ . Let

$$E = \frac{1}{2}v_a'^2 + hF(v_a) \quad (17)$$

and notice it follows from (10) that

$$E' = h'F(v_a) \leq 0. \quad (18)$$

Hence it follows that  $E$  is nonincreasing and so integrating this on  $[0, t]$  gives

$$\frac{1}{2}v_a'^2 + hF(v_a) \leq \frac{1}{2}a^2. \quad (19)$$

Since  $F(v_a) \geq 0$  it follows from (19) that  $v_a$  and  $v_a'$  are defined and bounded on the largest interval of existence  $[0, T] \subset [0, R_1]$  where they are defined. Since this upper bound is independent of  $T$  it follows then that  $v_a$  and  $v_a'$  are defined and continuous on all of  $[0, R_1]$ . In addition, it follows that  $v_a$  and  $v_a'$  vary continuously with  $a > 0$ .

## 4 Behavior of Solutions for Small Values of $a$

**Lemma 4.1:** Assume (H1)-(H3) and suppose  $v_a$  satisfies (10)-(11). If  $a > 0$  is sufficiently small, then  $v_a$  has a first zero,  $z_{a,1}$ , with  $0 < z_{a,1} < R_1$  and  $\lim_{a \rightarrow 0^+} z_{a,1} = 0$ . In addition,  $\lim_{a \rightarrow 0^+} |v_a'(z_{a,1})| = 0$ .

**Proof:** Let us suppose by way of contradiction that  $|v_a| > 0$  on  $(0, R_1]$ . Since  $v_a(0) = 0$  and  $v_a'(0) = a > 0$  it follows that  $v_a$  is positive and since  $f(v_a) > 0$  when  $v_a > 0$  it follows from (10) that  $v_a'' < 0$ . Integrating this twice on  $(0, t)$  gives

$$0 < v_a < at \quad (20)$$

and therefore

$$\frac{1}{v_a^q} > \frac{1}{a^q t^q}. \quad (21)$$

It follows from (H1) and (H2) that  $f \rightarrow \infty$  as  $v \rightarrow \infty$  and since  $g_1$  is continuous it follows that  $g_1(v_a) \geq -C$  for some constant  $C > 0$ . Using this and (21) in (12) gives:

$$v_a' + \int_0^t \left( \frac{h_0 x^{-\tilde{\alpha}-q}}{a^q} - C x^{-\tilde{\alpha}} \right) dx \leq v_a' + \int_0^t h(s) \left( \frac{1}{v_a^q} + g_1(v_a) \right) ds = a.$$

Carrying out the integration on the left-hand side and using (9) it follows then that

$$v_a' + \frac{h_0 t^{1-\tilde{\alpha}-q}}{a^q(1-\tilde{\alpha}-q)} - \frac{C t^{1-\tilde{\alpha}}}{1-\tilde{\alpha}} \leq a.$$

Integrating again on  $(0, t)$  and rewriting gives

$$v_a \leq at - \frac{h_0 t^{2-\tilde{\alpha}-q}}{a^q(1-\tilde{\alpha}-q)(2-\tilde{\alpha}-q)} + \frac{C t^{2-\tilde{\alpha}}}{(1-\tilde{\alpha})(2-\tilde{\alpha})}. \quad (22)$$

Notice for fixed positive  $t$  that if we choose  $a$  sufficiently small then the right hand-side of (22) becomes negative contradicting the assumption that  $v_a > 0$ . Thus  $v_a$  has a first zero,  $z_{a,1}$ , for  $a > 0$  sufficiently small. Then evaluating (22) at  $t = z_{a,1}$  and rewriting gives:

$$z_{a,1}^{1-\tilde{\alpha}-q} \leq \frac{a^{q+1} + Ca^q z_{a,1}^{1-\tilde{\alpha}}}{h_0} \leq \frac{a^{q+1} + Ca^q R_1^{1-\tilde{\alpha}}}{h_0} \rightarrow 0 \text{ as } a \rightarrow 0^+. \quad (23)$$

In addition, it follows from (19) that  $v_a'(z_{a,1}) \leq a^2$  and hence  $\lim_{a \rightarrow 0^+} |v_a'(z_{a,1})| = 0$ .

This completes the proof of the lemma.  $\square$

In a similar way we can show  $v_a$  has any finite number of zeros if  $a > 0$  is sufficiently small and  $\lim_{a \rightarrow 0^+} |v_a'(z_a)| = 0$  at each zero,  $z_a$ , of  $v_a$ .

## 5 Behavior of Solutions for Large Values of $a$

**Lemma 5.1:** Assume (H1)-(H3) and suppose  $v_a$  satisfies (10)-(11). Then  $\lim_{a \rightarrow \infty} \max_{[0, R_1]} v_a = \infty$ .

**Proof:** Recall  $F(u) = \int_0^u f(t) dt$  and suppose now by way of contradiction that  $v_a \leq C_0$  where  $C_0$  is independent of  $a$ . Using (20) and (H1)-(H2) then it follows that there are constants  $C_1, C_2$  and  $C_3$  so that

$$F(v_a) \leq C_1 v_a^{1-q} + C_2 v_a^{p+1} = v_a^{1-q} (C_1 + C_2 v_a^{p-q}) \leq a^{1-q} t^{1-q} (C_1 + C_2 C_1^{p-q}) \leq C_3 a^{1-q} t^{1-q}. \quad (24)$$

Next we see that it follows from (17)-(18) that

$$\frac{1}{2} v_a'^2 + hF(v_a) + \int_0^t (-h'(s))F(v_a) ds = \frac{1}{2} a^2. \quad (25)$$

Using (8) and (24) we see

$$hF(v_a) + \int_0^t (-h'(s))F(v_a) ds \leq C_3 h_1 a^{1-q} t^{-\tilde{\alpha}} t^{1-q} + \int_0^t 2\tilde{\alpha} C_3 h_1 a^{1-q} s^{-\tilde{\alpha}-1} s^{1-q} ds = C_4 a^{1-q} t^{1-\tilde{\alpha}-q}$$

where  $C_4 = h_1(C_3 + 2\tilde{\alpha})$ . Using this in (25) we obtain

$$|v_a'| \geq a^{\frac{1-q}{2}} \sqrt{a^{1+q} - C_4 R_1^{1-\tilde{\alpha}-q}}.$$

Thus we see for sufficiently large  $a$  that the right-hand side of the above is positive and so  $v_a$  is increasing for  $a$  sufficiently large. Then integrating on  $(0, t)$  gives

$$C_0 \geq v_a \geq a^{\frac{1-q}{2}} \left( \sqrt{a^{1+q} - C_4 R_1^{1-\tilde{\alpha}-q}} \right) t.$$

Thus for fixed  $t > 0$  and  $a$  sufficiently large we see that the right-hand side of the above gets larger than  $C_0$  and so we obtain a contradiction. Thus the lemma holds.  $\square$

**Lemma 5.2:** Assume (H1)-(H3) and suppose  $v_a$  satisfies (10)-(11). Then  $v_a$  has a local maximum,  $M_a$ , if  $a > 0$  is sufficiently large. In addition,  $M_a \rightarrow 0$  and  $|v_a(M_a)| \rightarrow \infty$  as  $a \rightarrow \infty$ .

**Proof:** Assume by way of contradiction that  $v_a$  is increasing on  $[0, R_1]$  for all  $a > 0$ . Then it follows from Lemma 5.1 and for any  $t_0 > 0$  that  $\min_{[t_0, R_1]} v_a(t) = v_a(t_0) \rightarrow \infty$  as  $a \rightarrow \infty$ . Then we see

$$\min_{[t_0, R_1]} \frac{h(t)f(v_a(t))}{v_a(t)} \geq h(R_1) \min_{[t_0, R_1]} \frac{f(v_a(t))}{v_a(t)} = C_a \rightarrow \infty \text{ as } a \rightarrow \infty.$$

Rewriting (10) we obtain

$$v_a'' + \left( \frac{h(t)f(v_a)}{v_a} \right) v_a = 0. \quad (26)$$

Next, let  $t_0 > 0$  and consider

$$y'' + C_a y = 0 \quad (27)$$

with  $y(t_0) = v_a(t_0) > 0$ ,  $y'(t_0) = v_a'(t_0) > 0$ . Now multiply (26) by  $y$ , (27) by  $v_a$ , and subtract to obtain

$$(yv_a' - y'v_a)' + \left( \frac{h(t)f(v_a)}{v_a} - C_a \right) yv_a = 0. \quad (28)$$

We know that  $y$  is a linear combination of  $\sin(\sqrt{C_a}t)$  and  $\cos(\sqrt{C_a}t)$  and as is well-known, any interval of length  $\frac{\pi}{\sqrt{C_a}}$  has a zero of  $y$ . Next we choose  $a$  sufficiently large so that  $\left( \frac{h(t)f(v_a)}{v_a} - C_a \right) > 0$  on  $[0, R_1]$ . Then by assumption we have  $v_a' > 0$  and  $v_a > 0$  on  $(t_0, t_0 + \frac{\pi}{\sqrt{C_a}})$  and so integrating (28) on  $(t_0, t)$  gives  $yv_a' - y'v_a < 0$  on  $(t_0, t)$ . It follows then that  $(\frac{y}{v_a})' < 0$  and so we integrate on  $(t_0, t)$  to get  $v_a < y$  on  $(t_0, t_0 + \frac{\pi}{\sqrt{C_a}})$ . Also, as mentioned above  $y$  has a smallest zero,  $Z$ , on  $(t_0, t_0 + \frac{\pi}{\sqrt{C_a}})$  and so we see  $v_a(Z) < y(Z) = 0$ . Now since  $v_a(t_0) > 0$ ,  $v_a'(t_0) > 0$ , and  $v_a(Z) < 0$  it follows that  $v_a$  must have a local maximum on  $(t_0, Z)$  which contradicts our assumption. Thus there is an  $M_a \in (0, R_1)$  at which  $v_a$  has a local maximum if  $a$  is sufficiently large.

It follows from Lemma 5.1 that  $v_a(M_a) \rightarrow \infty$  as  $a \rightarrow \infty$ .

Next integrating (10) on  $(t, M_a)$

$$v_a' = \int_t^{M_a} hf(v_a) ds \geq c_0 \int_t^{M_a} hv_a^p ds \geq c_0 v_a^p \int_t^{M_a} h ds.$$

Dividing by  $v_a^p$  and integrating again gives

$$\frac{v_a^{1-p}(M_a)}{p-1} \geq \frac{v_a^{1-p}(M_a) - v^{1-p}(t)}{p-1} \geq c_0 \int_t^{M_a} \int_s^{M_a} h \, dx \, ds.$$

Now as mentioned above, the left-hand side goes to zero as  $a \rightarrow \infty$  and so this is only possible in the above inequality if  $M_a \rightarrow 0$  as  $a \rightarrow \infty$ .  $\square$

**Lemma 5.3:** Assume (H1)-(H3) and suppose  $v_a$  satisfies (10)-(11). Then  $v_a$  has a zero  $z_a$  if  $a > 0$  is sufficiently large. Also  $z_a \rightarrow 0$  and  $|v'_a(z_a)| \rightarrow \infty$  as  $a \rightarrow \infty$ .

**Proof:** Assume by way of contradiction that  $v_a(t) > 0$  on  $(M_a, R_1)$ . Now let  $E_1 = \frac{1}{2} \frac{v_a'^2}{h} + F(v_a)$ . It then follows that  $E_1' = -\frac{v_a'^2 h'}{h^2} \geq 0$  and so

$$\frac{1}{2} \frac{v_a'^2}{h} + F(v_a) \geq F(v_a(M_a)) \text{ on } (M_a, R_1).$$

Rewriting, integrating on  $(M_a, t)$ , and changing variables gives

$$\int_{v_a(t)}^{M_a} \frac{1}{\sqrt{2}\sqrt{F(v_a(M_a)) - F(s)}} \, ds = \int_{M_a}^t \frac{-v'_a(s)}{\sqrt{2}\sqrt{F(v_a(M_a)) - F(v_a)}} \, ds \geq \int_{M_a}^t \sqrt{h} \, ds.$$

Assuming  $v_a(t) > 0$  on  $(0, R_1)$  it follows then that

$$\int_0^{M_a} \frac{1}{\sqrt{2}\sqrt{F(v_a(M_a)) - F(s)}} \, ds \geq \int_{M_a}^{R_1} \sqrt{h} \, ds.$$

After another change of variables we obtain

$$\frac{v_a(M_a)}{\sqrt{2}\sqrt{F(v_a(M_a))}} \int_0^1 \frac{1}{\sqrt{1 - \frac{F(v_a(M_a)s)}{F(v_a(M_a))}}} \, ds \geq \int_{M_a}^{R_1} \sqrt{h} \, ds. \quad (29)$$

However, it follows that since  $f$  is superlinear then  $\frac{v_a(M_a)}{\sqrt{F(v_a(M_a))}} \rightarrow 0$  as  $a \rightarrow \infty$  and by the dominated convergence theorem

$$\int_0^1 \frac{1}{\sqrt{1 - \frac{F(v_a(M_a)s)}{F(v_a(M_a))}}} \, ds \rightarrow \int_0^1 \frac{1}{\sqrt{1 - s^{p+1}}} \, ds$$

hence the left-hand side of (29) goes to 0 as  $a \rightarrow \infty$  and thus this forces  $\int_0^{R_1} \sqrt{h} \, ds = 0$  which is false.

Thus it must be the case that  $v_a$  has a zero,  $z_a$ , on  $(M_a, R_1)$ . Further, evaluating the above integral at  $t = z_a$  gives

$$\int_0^{M_a} \frac{1}{\sqrt{2}\sqrt{F(v_a(M_a)) - F(s)}} \, ds \geq \int_{M_a}^{z_a} \sqrt{h} \, ds.$$

Since the left-hand side goes to 0 as  $a \rightarrow \infty$  it follows then that  $z_a - M_a \rightarrow 0$  as  $a \rightarrow \infty$  and since  $M_a \rightarrow 0$  by Lemma 5.2 it follows that  $z_a \rightarrow 0$  as  $a \rightarrow \infty$ .



It follows from this then that if  $M_a < z_a$  then  $F(v_a(M_a) = E_1(M_a) \leq E_1(z_a) = \frac{1}{2} \frac{v_a'^2(z_a)}{h(z_a)} \leq \frac{1}{2} \frac{v_a'^2(z_a)}{h(R_1)}$  and since the left-hand side goes to infinity by Lemma 5.2 it follows then that  $|v_a'(z_a)| \rightarrow \infty$  as  $a \rightarrow \infty$ .  $\square$

In a similar way we can show  $v_a$  has any finite number of zeros if  $a > 0$  is sufficiently large and  $\lim_{a \rightarrow \infty} |v_a'(z_a)| = \infty$  at each zero  $z_a$  and  $z_a \rightarrow 0$  as  $a \rightarrow \infty$ .

## 6 Proof of the Main Theorem

**Lemma 6.1:** Assume (H1)-(H3) and suppose  $v_a$  satisfies (10)-(11). If  $z_a > 0$  and  $v_a(z_a) = 0$  then  $v_a'(z_a) \neq 0$ . In addition,  $v_a$  has a finite number of zeros on  $[0, R_1]$  for each  $a > 0$ .

**Proof:** Recall from above that  $E_1 = \frac{1}{2} \frac{v_a'^2}{h} + F(v_a)$  and that  $E_1' = -\frac{v_a'^2 h'}{h^2} \geq 0$ . Hence integrating on  $(0, t)$  we see that  $\frac{1}{2} \frac{v_a'^2}{h} + F(v_a) > 0$  for  $t > 0$ . It follows from this then that if  $v_a(z_a) = 0$  and  $z_a > 0$  then  $E_1(z_a) = \frac{1}{2} \frac{v_a'^2(z_a)}{h(z_a)} > 0$  and thus  $v_a'(z_a) \neq 0$ .

It follows from this that  $v_a$  cannot have an infinite number of zeros for if so then there would exist  $z_k$  such that  $0 < z_k < z_{k+1} \leq R_1$  such that  $v_a(z_k) = 0$  and there would also exist local extrema,  $M_k$ , with  $z_k < M_k < z_{k+1} \leq R_1$  and  $v_a'(M_k) = 0$ . But then the  $z_k$  are increasing and bounded above so there is a  $z^* \leq R_1$  such that  $z_k \rightarrow z^* \leq R_1$ . Also  $M_k \rightarrow z^*$  and it follows from continuity that  $v_a(z^*) = v_a'(z^*) = 0$  contradicting the first part of the lemma.  $\square$

**Proof of the Main Theorem** Let  $n$  be a nonnegative integer. Let

$$S_n = \{a | v_a > 0 \text{ has exactly } n \text{ zeros on } (0, R_1)\}.$$

It follows from Lemma 6.1 that there is smallest  $n_0 \geq 0$  such that  $S_{n_0}$  is nonempty.

Then let

$$a_{n_0} = \sup S_{n_0}.$$

It can then be show that  $v_{a_{n_0}}$  has exactly  $n_0$  zeros on  $(0, R_1)$  and  $v_{a_{n_0}}(R_1) = 0$ . See [2] for details.

Then by continuity with respect to  $a$  it follows that if  $a > a_{n_0}$  and  $a$  is close to  $a_{n_0}$  that  $v_a$  will have exactly  $n_0 + 1$  zeros on  $(0, R_1)$  and thus  $S_{n_0+1}$  is nonempty. Then letting  $a_{n_0+1} = \sup S_{n_0+1}$  we can similarly show  $v_{a_{n_0}}$  has exactly  $n_0 + 1$  zeros on  $(0, R_1)$  and  $v_{a_{n_0+1}}(R_1) = 0$ . Continuing in this way we can show that there are an infinite number of solutions.

In a similar way we can let

$$a_{n_0}^* = \inf S_{n_0}$$

and proceed to get a second infinite family of solutions.

Finally we set  $u_{a_n}(r) = v_{a_n}(r^{2-N})$  and  $u_{a_n^*}(r) = v_{a_n^*}(r^{2-N})$  thus we obtain two infinite families of solutions  $u_{a_n}$  and  $u_{a_n^*}$ . This completes the proof of the Main Theorem..  $\square$

## References

- [1] L. Evans, *Partial Differential Equations*, 2nd ed, Graduate Studies in Mathematics, Vol 19, AMS, 2010.
- [2] J. Iaia, Existence of Two Infinite Families of Solutions for Singular Superlinear Equations on Exterior Domains, *Electronic Journal of Differential Equations*, Vol. 2024 (2024), No. 06, pp. 1-14.
- [3] J. Iaia, Existence and Nonexistence of Sign-changing Solutions for Singular Superlinear Equations on Exterior Domains, *Communications on Pure and Applied Analysis*, 22(11) 3218-3232, 2023.
- [4] E.K. Lee, R. Shivaji, B. Son, "Positive radial solutions to classes of singular problems on the exterior of a ball," *Journal of Mathematical Analysis and Applications*, 434, No. 2, 1597-1611, 2016.
- [5] E. Lee, L. Sankar, R. Shivaji, "Positive solutions for infinite semipositone problems on exterior domains," *Differential and Integral Equations*, Volume 24, Number 9/10, 861-875, 2011.

University of North Texas

P.O. Box 311430

Denton, Texas 76203

abdealidiwan@my.unt.edu

iaia@unt.edu