# arXiv:2407.14511v1 [math.GR] 18 Jun 2024

# Counting subgroups of the groups $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ : a survey

László Tóth Department of Mathematics University of Pécs Ifjúság útja 6, 7624 Pécs, Hungary E-mail: ltoth@gamma.ttk.pte.hu

New Frontiers in Number Theory and Applications, Trends in Mathematics, Birkhäuser, Cham, 2024, Guàrdia, J., Minculete, N., Savin, D., Vela, M., Zekhnini, A. (eds), pp. 385–409 https://doi.org/10.1007/978-3-031-51959-8\_18

### Abstract

We present a survey of exact and asymptotic formulas on the number of cyclic subgroups and total number of subgroups of the groups  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ , where  $k \ge 2$  and  $n_1, \ldots, n_k$ are arbitrary positive integers.

2010 Mathematics Subject Classification: 20K01, 20K27, 05A15, 11A25, 11N37 Key Words and Phrases: finite abelian group, subgroup, number of subgroups, number of cyclic subgroups, Goursat's lemma, asymptotic formula

•

# Contents

T	Introduction	2
2	Number of cyclic subgroups	3
3	Number of cyclic subgroups in the case $n_1 = \cdots = n_k$	6
4	Number of all subgroups in the case $k = 2$	7
5	Number of all subgroups in the case $k = 3$	13
6	Number of all subgroups in the case $k = 4$	15
7	Final remarks	<b>21</b>

### 1 Introduction

Throughout this chapter we use the following notation:  $\mathbb{N} := \{1, 2, \ldots\}, \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ ; the prime power factorization of  $n \in \mathbb{N}$  is  $n = \prod_p p^{\nu_p(n)}$ , where the product is over the primes p and all but a finite number of the exponents  $\nu_p(n)$  are zero;  $\mathbb{Z}_n$  denotes the additive group of residue classes modulo n (the cyclic group of order  $n \in \mathbb{N}$ );  $(n_1, \ldots, n_k)$  and  $gcd(n_1, \ldots, n_k)$ denote the greatest common divisor of  $n_1, \ldots, n_k \in \mathbb{N}$ ,  $[n_1, \ldots, n_k]$  and  $lcm(n_1, \ldots, n_k)$  denote the least common multiple of  $n_1, \ldots, n_k \in \mathbb{N}$ ;  $id_t(n) = n^t, \sigma_t(n) = \sum_{d|n} d^t$   $(n \in \mathbb{N}, t \in \mathbb{C})$ ;  $\sigma_0(n) = \tau(n)$  and  $\sigma_1(n) = \sigma(n)$  denote the number and the sum of (positive) divisors of  $n \in \mathbb{N}$ ;  $\varphi$  is Euler's arithmetic function;  $\mu$  is the Möbius function; f \* g is the convolution of the functions  $f, g: \mathbb{N} \to \mathbb{C}$ ;  $\zeta$  is the Riemann zeta function; the sums  $\sum_p$  and products  $\prod_p$  and  $\times_p$  are taken over the primes p.

Let G be a finite abelian group of order n. Let s(G) and c(G) denote the total number of subgroups of G and the number of its cyclic subgroups, respectively. Also, let  $G = \times_p G_p$  be the primary decomposition of G, where  $|G_p| = p^{\nu_p(n)}$  (p prime). Then

$$s(G) = \prod_{p} s(G_{p}), \qquad c(G) = \prod_{p} c(G_{p}).$$
(1)

Therefore, the problem of counting the subgroups of G reduces to p-groups. Formulas for the total number of subgroups of a given type of an abelian p-group were established by Bhowmik [2], Delsarte [11], Miller [24, 25], Shokuev [34], Yeh [51], and others. Also see the papers by Stehling [35] and Suzuki [39], and the monographs by Butler [7] and Schmidt [32]. One of these formulas is in terms of the gaussian coefficients  $\begin{bmatrix} r \\ k \end{bmatrix}_p = \prod_{i=1}^k \frac{p^{r-k+i}-1}{p^i-1}$ . Namely, let  $G_{(p)}$  be a p-group of rank r and type  $\lambda = (\lambda_1, \ldots, \lambda_r)$ , that is,

$$G_{(p)} \simeq \mathbb{Z}_{p^{\lambda_1}} \times \dots \times \mathbb{Z}_{p^{\lambda_r}},$$
(2)

with  $\lambda_1 \ge \ldots \ge \lambda_r \ge 1$ , where  $\lambda$  is a partition of  $|\lambda| = \lambda_1 + \cdots + \lambda_r$ . Then the number  $s_{\mu}(G_{(p)})$  of subgroups of type  $\mu$  (with  $\mu \le \lambda$ , meaning that  $\mu_i \le \lambda_i$  for  $1 \le i \le r$ ) of  $G_{(p)}$  is

$$s_{\mu}(G_{(p)}) = \prod_{j=1}^{\lambda_1} p^{\mu'_{j+1}(\lambda'_j - \mu'_j)} \begin{bmatrix} \lambda'_j - \mu'_{j+1} \\ \mu'_j - \mu'_{j+1} \end{bmatrix}_p,$$

where  $\lambda'$  and  $\mu'$  are the conjugates (according to the Ferrers diagrams) of  $\lambda$  and  $\mu$ , respectively. Hence  $s_{\mu}(G_{(p)})$  is a polynomial in p, with integer coefficients, depending only on  $\lambda$  and  $\mu$  (it is a sum of Hall polynomials).

It follows that the total number of subgroups of  $G_{(p)}$  is

$$s(G_{(p)}) = \sum_{0 \leq k \leq |\lambda|} \sum_{\substack{\mu \leq \lambda \\ |\mu| = k}} s_{\mu}(G_{(p)}),$$

and as a consequence, for abelian p-groups the total number of subgroups and the number of subgroups of a given order are polynomials in p with integer coefficients. In fact, it can be shown

that all coefficients of these polynomials are non-negative. However, it is a difficult task to make explicit these polynomials, even in the case of p-groups of small rank.

There are more simple formulas for the number of cyclic subgroups of p-groups. Consider the p-group  $G_{(p)}$  given by (2). The number of cyclic subgroups of order  $p^{\nu}$  of  $G_{(p)}$  is

$$c_{p^{\nu}}(G_{(p)}) = \frac{p^{j_{\nu}} - 1}{p - 1} p^{\lambda_{j_{\nu}+1} + \dots + \lambda_r + \lambda_{r+1} + (\nu - 1)(j_{\nu} - 1)},$$

where  $\lambda_{j_{\nu}} \ge \nu > \lambda_{j_{\nu}+1}$  and  $\lambda_{r+1} = 0$ . See Miller [25], Yeh [51].

If G is a finite abelian group, then one can define the generalized counting function

$$\sigma_t(G) = \sum_{H \leqslant G} |H|^t, \tag{3}$$

where the sum is over all subgroups H of G and  $t \in \mathbb{C}$ . If t = 0, then  $\sigma_0(G) = s(G)$ , the number of all subgroups of G. Ramaré [31] investigated the function  $\sigma_t(G)$  in the case when Gis a *p*-group of type (2), proved an involved but fully explicit formula for  $\sigma_t(G)$ , where G is an arbitrary abelian *p*-group, and deduced the rationality of a related generating series.

Instead of p-groups we prefer to consider the group  $G := \mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ , where  $n_1, \ldots, n_k$ are arbitrary positive integers. Let  $s(n_1, \ldots, n_k)$  and  $c(n_1, \ldots, n_k)$  stand for the total number of subgroups of G and the number of its cyclic subgroups, respectively. According to (1), the functions  $s(n_1, \ldots, n_k)$  and  $c(n_1, \ldots, n_k)$  are multiplicative functions of k variables. Therefore,  $s(n, \ldots, n)$  and  $c(n, \ldots, n)$  are multiplicative in n, as functions of a single variable. See the survey of the author [47] on general properties of multiplicative arithmetic functions of several variables. As well known, in the case k = 1 one has  $s(n) = c(n) = \tau(n)$  for every  $n \in \mathbb{N}$ ,  $\tau(n)$ denoting the number of divisors of n. Also note that  $\sigma_t(\mathbb{Z}_n) = \sum_{d|n} d^t = \sigma_t(n)$ .

Here we present a survey concerning the total number of subgroups (of a given order) and the number of cyclic subgroups (of a given order) of the group  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ , where  $k \ge 2$ . Some new results are also included. We consider, in particular, the cases k = 2, k = 3 and k = 4, and present exact and asymptotic formulas on the functions  $s(n_1, \ldots, n_k)$  and  $c(n_1, \ldots, n_k)$ , where  $n_1, \ldots, n_k \in \mathbb{N}$ . We also discuss the representation of the subgroups in the cases k = 2 and k = 3, and offer some new results for  $\sigma_t(G)$  in the case when  $G = \mathbb{Z}_m \times \mathbb{Z}_n$   $(m, n \in \mathbb{N})$ . We point out that there are known asymptotic formulas for the sums  $\sum_{n_1,\ldots,n_k \leq x} c(n_1,\ldots,n_k)$  with  $k \ge 2$ , see Section 2, and for  $\sum_{n_1,n_2 \leq x} s(n_1,n_2)$ , see Section 4. However, no asymptotic formula for  $\sum_{n_1,\ldots,n_k \leq x} s(n_1,\ldots,n_k)$  is known the literature in the case  $k \ge 3$ .

The new results are given together with their proofs. These are Propositions 1, 2, Theorems 1, 2, 3, 4 and Corollaries 1, 2. To be precise, Proposition 2 and the results of Section 6 are included in the arXiv preprint by the author [48], but have not been published elsewhere.

### 2 Number of cyclic subgroups

For  $c(n_1, \ldots, n_k)$ , the number of cyclic subgroups of the group  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$  we have the following compact formula, valid for every  $k \ge 2$  and every  $n_1, \ldots, n_k \in \mathbb{N}$ :

$$c(n_1,\ldots,n_k) = \sum_{d_1|n_1,\ldots,d_k|n_k} \frac{\varphi(d_1)\cdots\varphi(d_k)}{\varphi(\operatorname{lcm}(d_1,\ldots,d_k))},$$
(4)

where  $\varphi$  is Euler's function. Formula (4) was derived by the author in [44] using the orbit counting lemma (Burnside's lemma) and in [45] by some simple number-theoretic arguments.

Consequently, the number of all cyclic subgroups of an arbitrary finite abelian group G can be described by formula (4) in a compact form. Namely, use the primary decomposition of G, mentioned above, or consider the invariant factor decomposition of G given by

$$G \simeq \mathbb{Z}_{n_1} \times \dots \times \mathbb{Z}_{n_r},\tag{5}$$

with  $n_1, \ldots, n_r \in \mathbb{N} \setminus \{1\}, n_j \mid n_{j+1} \ (1 \leq j \leq r-1).$ 

Note that if |G| > 1, then the number r in (5) is uniquely determined, it represents the minimal number of generators of the group G, and is called the rank of G. In the case of abelian p-groups this notion of rank recovers the rank mentioned above.

As discussed in the Introduction,  $c(n_1, \ldots, n_k)$  is a multiplicative function of k variables. This property is a direct consequence of convolutional formula (4).

Let  $c_{\delta}(n_1, \ldots, n_k)$  denote the number of cyclic subgroups of order  $\delta$  of the group  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ . For every  $n_1, \ldots, n_k \in \mathbb{N}$  and  $\delta \mid \text{lcm}(n_1, \ldots, n_k)$  we have, see Tóth [45, Th. 2],

$$c_{\delta}(n_1, \dots, n_k) = \frac{1}{\varphi(\delta)} \sum_{e|\delta} \gcd(e, n_1) \cdots \gcd(e, n_k) \mu(\delta/e)$$
$$= \frac{1}{\varphi(\delta)} \sum_{\substack{d_1|n_1, \dots, d_k|n_k\\ \operatorname{lcm}(d_1, \dots, d_r) = \delta}} \varphi(d_1) \cdots \varphi(d_k).$$

In the case k = 2, identity (4) reduces to

$$c(n_1, n_2) = \sum_{d_1|n_1, d_2|n_2} \varphi(\gcd(d_1, d_2)),$$
(6)

which can also be written as

$$c(n_1, n_2) = \sum_{d \mid \text{gcd}(n_1, n_2)} (\mu * \varphi)(d) \tau(n_1/d) \tau(n_2/d).$$

Note that for every prime p and  $1 \leq a \leq b$  we have

$$c(p^{a}, p^{b}) = 2(1 + p + p^{2} + \dots + p^{a-1}) + (b - a + 1)p^{a},$$

and (6) leads to the Dirichlet series representation

$$\sum_{n_1,n_2=1}^{\infty} \frac{c(n_1,n_2)}{n_1^{s_1} n_2^{s_2}} = \frac{\zeta^2(s_1)\zeta^2(s_2)\zeta(s_1+s_2-1)}{\zeta^2(s_1+s_2)}.$$

By using a variant of the hyperbola method, Nowak and Tóth [26, 27] proved the asymptotic formula

$$\sum_{n_1, n_2 \leqslant x} c(n_1, n_2) = x^2 \sum_{j=0}^3 B_j (\log x)^j + O\left(x^{\frac{1117}{701} + \varepsilon}\right),\tag{7}$$

where  $B_3 = 12/\pi^4$ ,  $B_2$ ,  $B_1$ ,  $B_0$  are explicit constants, and  $1117/701 \doteq 1.593437$ . They also proved a similar formula with the same error term for the sum  $\sum_{n_1,n_2 \leq x,(n_1,n_2)>1} c(n_1,n_2)$  concerning the number of cyclic subgroups of the groups  $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2}$  having rank two, and pointed out that this error can be replaced by  $O\left(x^{\frac{3-\theta}{2-\theta}+\varepsilon}\right)$ , where  $\theta$  is the exponent in Dirichlet's divisor problem. The *O*-term stated above was obtained from the sharpest result known at the time due to Huxley, namely from  $\theta = 131/416$ . It was pointed out by Tóth and Zhai [49] that asymptotic formula (7) holds for the slight better exponent  $4427/2779 \doteq 1.593019$  by using the exponent  $\theta = 517/1648$  obtained by Bourgain and Watt [4]<sup>1</sup>. Note that the limit of this approach is  $11/7 \doteq 1.571428$  with  $\theta = 1/4$ .

Tóth and Zhai [49] proved, by using a two dimensional Perron formula and the complex integration method that the error term of (7) can be refined into  $O\left(x^{3/2}(\log x)^{13/2}\right)$ . Concerning the case k = 3, the same authors [50] proved, again by using a multidimensional Perron formula and the complex integration method that

$$\sum_{n_1, n_2, n_3 \leqslant x} c(n_1, n_2, n_3) = x^3 \sum_{j=0}^7 C_j (\log x)^j + O\left(x^{8/3 + \varepsilon}\right)$$

where  $C_j$   $(0 \le j \le 7)$  are some explicit constants.

As a consequence of certain more general results, it was obtained by Essouabri, Salinas Zavala and Tóth [12, Cor. 5], based on identity (4), by deep analytic methods that for every  $k \ge 2$ ,

$$\sum_{n_1,\dots,n_k \leqslant x} c(n_1,\dots,n_k) = x^k Q(\log x) + O\left(x^{k-\beta}\right),\tag{8}$$

where Q(t) is a polynomial in t of degree  $2^k - 1$  and  $\beta$  is a positive real number. More exactly, by considering a large class of mutivariable multiplicative functions  $f : \mathbb{N}^k \to \mathbb{R}_+$ , it was proved in [12] the existence of the meromorphic continuation and some related properties of the associated multiple Dirichlet series

$$\sum_{n_1,\dots,n_k=1}^{\infty} \frac{f(n_1,\dots,n_k)}{n_1^{s_1}\cdots n_k^{s_k}},$$

and then formula (8) was deduced by combining those results and La Bretèche's multivariable Tauberian theorems [5, Ths. 1, 2].

In paper [12] a result similar to (8), but with respect to the Hölder norm condition  $(n_1^s + \cdots + n_k^s)^{1/s} \leq x$ , where  $s \geq 1$  is a fixed real number, was also deduced. Sui and Liu [36] obtained an asymptotic formula for the hyperbolic summation  $\sum_{n_1n_2 \leq x} c(n_1, n_2)$ , and studied the order of magnitude of the corresponding error term. Kiuchi and Saad Eddin [20] deduced an asymptotic formula for the weighted sum  $\sum_{n_1n_2 \leq x} c(n_1, n_2) \log \frac{x}{n_1n_2}$ .

<sup>&</sup>lt;sup>1</sup>This paper has been withdrawn by the second author in July 2023.

# 3 Number of cyclic subgroups in the case $n_1 = \cdots = n_k$

Let  $f_k(n)$  denote the number of subgroups of index n of the group  $\mathbb{Z}^k$  (sublattices of the lattice  $\mathbb{Z}^k$ ). It is known in the literature that for every  $k, n \in \mathbb{N}$  one has

$$f_k(n) = \sum_{d_1 d_2 \cdots d_k = n} d_1^0 d_2^1 d_3^2 \cdots d_k^{k-1},$$
$$\sum_{n=1}^{\infty} \frac{f_k(n)}{n^s} = \zeta(s)\zeta(s-1)\cdots\zeta(s-k+1).$$

These identities (having applications in crystallography) lead to the asymptotic formula

$$\sum_{n \leqslant x} f_k(n) = \frac{\zeta(2)\zeta(3)\cdots\zeta(k)}{k} x^k + O\left(x^{k-1}(\log x)^{2/3}\right),\tag{9}$$

valid for every  $k \ge 2$ . See Grady [14, Lemma 1] (case k = 2), Gruber [15], Knopfmacher [21, Corollary (2.13)], Zou [52]. Note that the error term in (9) comes from the error due to Walfisz concerning the function  $\sigma(n) = \sum_{d|n} d$ , namely

$$\sum_{n \le x} \sigma(n) = \frac{\pi^2}{12} x^2 + O\left(x(\log x)^{2/3}\right).$$
(10)

Now let  $c_k(n)$  denote the number of cyclic subgroups of the group  $\mathbb{Z}_n^k = \mathbb{Z}_n \times \cdots \times \mathbb{Z}_n$  (k times). Here  $c_k(n)$  is multiplicative in n, and we have, see Haukkanen and Sivaramakrishnan [18, Th. 4]),

$$c_k(n) = \sum_{d|n} \frac{\varphi_k(d)}{\varphi(d)},\tag{11}$$

where  $\varphi_k(n) = n^k \prod_{p|n} (1 - 1/p^k)$  is the Jordan function of order k and  $\varphi(n) = \varphi_1(n)$  is Euler's function. Note that (11) also follows from identity (4). Also see Schulte [33].

In the case k = 2 one has  $c_2(n) = \sum_{d|n} \psi(d)$ , where  $\psi(n) = n \prod_{p|n} (1 + 1/p)$  is the Dedekind function. It was mentioned in our arXiv preprint [26] that

$$\sum_{n \leqslant x} c_2(n) = \frac{5}{4} x^2 + O\left(x(\log x)^{5/3}\right),$$

the error term coming from the result of Walfisz for the function  $\psi$  (the same error as in (10)).

Now we prove the following result, which is similar to (9).

**Proposition 1.** For every fixed  $k \ge 3$ ,

$$\sum_{n \leq x} c_k(n) = \frac{\zeta(k)}{k} H(k) x^k + O\left(x^{k-1} \log x\right),$$

where

$$H(k) = \prod_{p} \left( 1 + \frac{1}{p^2} + \frac{1}{p^3} + \dots + \frac{1}{p^k} \right).$$

*Proof.* We use standard arguments. The Dirichlet series of  $c_k(n)$  is, by (11),

$$\sum_{n=1}^{\infty} \frac{c_k(n)}{n^s} = \zeta(s) \prod_p \sum_{\nu=0}^{\infty} \frac{\varphi_k(p^{\nu})}{\varphi(p^{\nu})p^{\nu s}}$$
$$= \zeta(s)\zeta(s-k+1)H(s),$$

where

$$H(s) = \sum_{n=1}^{\infty} \frac{h(n)}{n^s} = \prod_p \left( 1 + \frac{1}{p^{s-k+2}} + \frac{1}{p^{s-k+3}} + \dots + \frac{1}{p^s} \right)$$

is absolutely convergent for  $\Re s > k - 1$ . Hence

$$c_k(n) = \sum_{de=n} h(d)\sigma_{k-1}(e),$$

with  $\sigma_{k-1}(n) = \sum_{d|n} d^{k-1}$ . We have

$$\sum_{n \leqslant x} c_k(n) = \sum_{d \leqslant x} h(d) \sum_{e \leqslant x/d} \sigma_{k-1}(e) = \sum_{d \leqslant x} h(d) \left( \frac{\zeta(k)}{k} \left( \frac{x}{d} \right)^k + O\left( \left( \frac{x}{d} \right)^{k-1} \right) \right)$$
$$= \frac{\zeta(k)}{k} x^k \sum_{d=1}^\infty \frac{h(d)}{d^k} + O\left( x^k \sum_{d>x} \frac{|h(d)|}{d^k} \right) + O\left( x^{k-1} \sum_{d \leqslant x} \frac{|h(d)|}{d^{k-1}} \right).$$

The sum of the series in the main term is H(k). Also,  $h(p) = \frac{p^{k-1}-1}{p-1} \leq 2p^{k-2}$  for any prime p and  $h(p^{\nu}) = 0$  for any prime power  $p^{\nu}$  with  $\nu \geq 2$ . So,  $h(n) \ll 2^{\omega(n)} n^{k-2}$ , where  $\omega(n) = \sum_{p|n} 1$ . This gives

$$\sum_{d>x} \frac{|h(d)|}{d^k} \ll \sum_{d>x} \frac{2^{\omega(d)}}{d^2} \ll \frac{\log x}{x}.$$

Furthermore,

$$\sum_{d \leqslant x} \frac{|h(d)|}{d^{k-1}} \leqslant \prod_{p \leqslant x} \sum_{\nu=0}^{\infty} \frac{|h(p^{\nu})|}{p^{\nu(k-1)}} = \prod_{p \leqslant x} \left( 1 + \frac{|h(p)|}{p^{k-1}} \right)$$
$$= \prod_{p \leqslant x} \left( 1 + \frac{1}{p} + \frac{1}{p^2} + \dots + \frac{1}{p^{k-1}} \right) \ll \log x$$

by Mertens' theorem.

# 4 Number of all subgroups in the case k = 2

Consider the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , where  $m, n \in \mathbb{N}$ . Note that  $\mathbb{Z}_m \times \mathbb{Z}_n$  is isomorphic to  $\mathbb{Z}_{(m,n)} \times \mathbb{Z}_{[m,n]}$ , and it has rank two, assuming that (m, n) > 1.

Goursat's lemma for groups can be stated as follows:

i) Let G and H be arbitrary groups. Then there is a bijection between the set of all subgroups of  $G \times H$  and the set of all 5-tuples  $(A, B, C, D, \Psi)$ , where  $B \leq A \leq G$ ,  $D \leq C \leq H$  and  $\Psi : A/B \to C/D$  is an isomorphism (here  $\leq$  denotes subgroup and  $\leq$  denotes normal subgroup). More precisely, the subgroup corresponding to  $(A, B, C, D, \Psi)$  is

$$K = \{ (g,h) \in A \times C : \Psi(gB) = hD \}.$$

ii) Assume that G and H are finite groups and assume that the subgroup K of  $G \times H$  corresponds to the 5-tuple  $(A_K, B_K, C_K, D_K, \Psi_K)$  under the above bijection. Then one has  $|A_K| \cdot |D_K| = |K| = |B_K| \cdot |C_K|$ .

For the history of Goursat's lemma, its proof and discussion see the references given in Tóth [46]. Also see the paper by Bauer et. al. [3].

For every  $m, n \in \mathbb{N}$  let

$$J_{m,n} := \left\{ (a, b, c, d, \ell) \in \mathbb{N}^5 : a \mid m, b \mid a, c \mid n, d \mid c, \frac{a}{b} = \frac{c}{d}, \\ \ell \leqslant \frac{a}{b}, \operatorname{gcd}\left(\ell, \frac{a}{b}\right) = 1 \right\},$$

$$(12)$$

and for  $(a, b, c, d, \ell) \in J_{m,n}$  define

$$K_{a,b,c,d,\ell} := \left\{ \left( i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d} \right) : 0 \le i \le a - 1, 0 \le j \le d - 1 \right\}.$$

$$\tag{13}$$

Using Goursat's lemma, the author [46, Th. 3.1] gave the following representation of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ .

Let  $m, n \in \mathbb{N}$ .

i) The map  $(a, b, c, d, \ell) \mapsto K_{a,b,c,d,\ell}$  is a bijection between the set  $J_{m,n}$  and the set of subgroups of  $(\mathbb{Z}_m \times \mathbb{Z}_n, +)$ .

ii) The invariant factor decomposition of the subgroup  $K_{a,b,c,d,\ell}$  is

$$K_{a,b,c,d,\ell} \simeq \mathbb{Z}_{\gcd(b,d)} \times \mathbb{Z}_{\operatorname{lcm}(a,c)},\tag{14}$$

where  $gcd(b, d) \mid lcm(a, c)$ .

iii) The order of the subgroup  $K_{a,b,c,d,\ell}$  is ad and its exponent is lcm(a,c).

iv) The subgroup  $K_{a,b,c,d,\ell}$  is cyclic if and only if gcd(b,d) = 1.

The next figure represents the subgroup  $K_{6,2,18,6,1}$  of  $\mathbb{Z}_{12} \times \mathbb{Z}_{18}$ . It has order 36 and is isomorphic to  $\mathbb{Z}_2 \times \mathbb{Z}_{18}$ .

According to the above properties, the total number s(m, n) of subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$  can be obtained by counting the elements of the set  $J_{m,n}$ . We have the following simple compact formulas, see the author [46, Th. 4.1]. For every  $m, n \in \mathbb{N}$ , s(m, n) is given by

$$s(m,n) = \sum_{i|m,j|n} \gcd(i,j) \tag{15}$$

$$= \sum_{d|\gcd(m,n)} \varphi(d)\tau\left(\frac{m}{d}\right)\tau\left(\frac{n}{d}\right).$$
(16)

17					٠						•	
16			٠			•			٠			
15	٠						٠	•	•		•	
14	•	•	•	•	•	•	•	•	•	•	٠	•
13		•	٠				•	•	٠			•
12	٠		•			•	٠	•	•		•	
11	•	•	•	•	٠	•	•	•	•	•	٠	•
10	•	•	٠	•	•	•	•	•	٠	•	•	•
9	٠	•		•	•	•	٠		•	•	•	•
8	•	•	•	•	٠	•	•	•	•	•	٠	•
7	•	•	٠	•	•	•	•	•	٠	•	•	•
6	٠	•	•	•	•	•	•	•	•	•	•	•
5	•	•	•	•	٠	•	·	·	•	•	٠	•
4	·	•	٠	•	•	·	·	·	٠	•	•	•
3	٠	•	•	·	·	•	•	·	•	·	•	·
2	·	•	•	•	•	•	·	•	•	•	•	·
1	·	•	٠	•	•	•	·	•	٠	•	•	·
0	•	•	•	•	•	•	•	•	•	•	•	•
	0	1	2	3	4	5	6	7	8	9	10	11

Representation and properties of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ , including identities (15) and (16) were also obtained by Hampejs, Holighaus, Tóth and Wiesmeyr [16, Th. 4] by tooking a different approach.

It is a direct consequence of (15) or (16) that for every prime p and  $1 \leq a \leq b$ ,

$$s(p^{a}, p^{b}) = \frac{(b-a+1)p^{a+2} - (b-a-1)p^{a+1} - (a+b+3)p + (a+b+1)}{(p-1)^{2}}.$$
 (17)

Formula (17) was also deduced, applying Goursat's lemma for groups by Călugăreanu [9, Sect. 4] and Petrillo [29, Prop. 2], using the concept of the fundamental group lattice by Tărnăuceanu [41, Prop. 2.9], [42, Th. 3.3], and from a general recurrence relation by Ramaré [31, Eq. (10)].

We also remark that (15) is a special case of an identity representing the total number of subgroups of a class of groups formed as cyclic extensions of cyclic groups, deduced by Calhoun [8] and having a laborious proof.

Let  $s_{\delta}(m, n)$  denote the number of subgroups of order  $\delta$  of  $\mathbb{Z}_m \times \mathbb{Z}_n$ . For every  $m, n, \delta \in \mathbb{N}$  such that  $\delta \mid mn$ ,

$$s_{\delta}(m,n) = \sum_{\substack{i \mid \gcd(m,\delta) \\ j \mid \gcd(n,\delta) \\ \delta \mid ij}} \varphi(ij/\delta))$$
$$= \sum_{d \mid \gcd(m,n,\delta)} \varphi(d) N(m/d, n/d, \delta/d),$$

where N(a, b, c) is the number of solutions  $(x, y, z, t) \in \mathbb{N}^4$  of the system of equations xy = a, zt = b, xz = c, see Tóth [46, Th. 4.3], Hampejs, Holighaus, Tóth and Wiesmeyr [16, Th. 4].

For the number of subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$  with a given isomorphism type  $\mathbb{Z}_A \times \mathbb{Z}_B$  we have the following formula, see the author [46, Th. 4.5].

Let  $m, n \in \mathbb{N}$  and let  $A, B \in \mathbb{N}$  such that  $A \mid B, AB \mid mn$ . Let  $A \mid \text{gcd}(m, n)$ . Then the number  $N_{A,B}(m, n)$  of subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ , which are isomorphic to  $\mathbb{Z}_A \times \mathbb{Z}_B$  is given by

$$N_{A,B}(m,n) = \sum_{\substack{i|m,j|n\\AB|ij\\lcm(i,j)=B}} \varphi\left(\frac{ij}{AB}\right),$$

where  $\varphi$  is Euler's function. If  $A \nmid \operatorname{gcd}(m, n)$ , then  $N_{A,B}(m, n) = 0$ .

The following result is concerning the quotient group  $(\mathbb{Z}_m \times \mathbb{Z}_n)/K_{a,b,c,d,\ell}$ .

**Proposition 2.** Let  $m, n \in \mathbb{N}$ . For every subgroup  $K_{a,b,c,d,\ell}$  defied by (13), we have the invariant factor decomposition

$$(\mathbb{Z}_m \times \mathbb{Z}_n)/K_{a,b,c,d,\ell} \simeq \mathbb{Z}_{\gcd\left(\frac{m}{a},\frac{n}{c}\right)} \times \mathbb{Z}_{\operatorname{lcm}\left(\frac{m}{b},\frac{n}{d}\right)},$$

where  $\operatorname{gcd}\left(\frac{m}{a}, \frac{n}{c}\right) \mid \operatorname{lcm}\left(\frac{m}{b}, \frac{n}{d}\right).$ 

*Proof.* Let  $K := K_{a,b,c,d,\ell}$ . The group  $\mathbb{Z}_m \times \mathbb{Z}_n$  is generated by (0,1) and (1,0). Therefore, the quotient group  $\mathbb{Z}_m \times \mathbb{Z}_n/K$  is generated by (0,1) + K and (1,0) + K.

First we show that the order of (0,1) + K is  $\frac{n}{d}$ . Indeed, this follows from the following properties:

•  $t(0,1) \in K$  if and only if there is  $(i,j) \in \{0,1,\ldots,a-1\} \times \{0,1,\ldots,d-1\}$  such that  $(0,t) = (i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d})$ , where the last condition is equivalent, in turn, to i = 0 and  $t \equiv j\frac{n}{d} \pmod{n}$ ;

• the least such  $t \in \mathbb{N}$  is  $\frac{n}{d}$ .

Next we show that the order of (1,0) + K is  $\frac{m}{h}$ . Indeed, observe that

•  $u(1,0) \in K$  if and only if there is  $(i,j) \in \{0,1,\ldots,a-1\} \times \{0,1,\ldots,d-1\}$  such that  $(u,0) = (i\frac{m}{a}, i\ell\frac{n}{c} + j\frac{n}{d});$ 

• therefore,  $u \equiv i\frac{m}{a} \pmod{m}$  and  $i\ell\frac{n}{c} + j\frac{n}{d} \equiv 0 \pmod{n}$ , that is  $\frac{n}{c}(i\ell + j\frac{c}{d}) \equiv 0 \pmod{n}$ ,  $i\ell + j\frac{c}{d} \equiv 0 \pmod{c}$ , the latter is a linear congruence in j and it turns out that for a fixed i it has solution in j if  $gcd\left(\frac{c}{d},c\right) = \frac{c}{d} \mid i\ell$ , that is  $\frac{c}{d} \mid i$ , since  $gcd\left(\ell,\frac{c}{d}\right) = 1$  by (12);

• the least such  $i \in \mathbb{N}$  is  $i = \frac{c}{d}$ ;

• finally, the least such  $u \in \mathbb{N}$  is  $u = \frac{c}{d} \cdot \frac{m}{a} = \frac{a}{b} \cdot \frac{m}{a} = \frac{m}{b}$ , representing the order of (1,0) + K. We deduce that the exponent of  $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$  is  $\operatorname{lcm}(\frac{m}{b}, \frac{n}{d})$ .

Now,  $\mathbb{Z}_m \times \mathbb{Z}_n \simeq \mathbb{Z}_{\text{gcd}(m,n)} \times \mathbb{Z}_{\text{lcm}(m,n)}$  is a group of rank  $\leq 2$ . Therefore, the quotient group  $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$  has also rank  $\leq 2$ . That is,  $(\mathbb{Z}_m \times \mathbb{Z}_n)/K \simeq \mathbb{Z}_v \times \mathbb{Z}_w$  for unique v and w, where  $v \mid w$  and  $vw = \frac{mn}{ad}$ , since |K| = ad. Here the exponent of  $\mathbb{Z}_v \times \mathbb{Z}_w$  is w, and we obtain that the

exponent of  $(\mathbb{Z}_m \times \mathbb{Z}_n)/K$  is  $\operatorname{lcm}(\frac{m}{h}, \frac{n}{d}) = w$ . Hence

r

$$v = \frac{mn}{adw} = \frac{mn}{ad \operatorname{lcm}(\frac{m}{b}, \frac{n}{d})} = \frac{mn}{ad \operatorname{lcm}(\frac{mn}{bn}, \frac{mn}{dm})} = \frac{mn}{ad \frac{mn}{\gcd(bn, dm)}}$$
$$= \frac{\gcd(bn, dm)}{ad} = \gcd\left(\frac{n}{d} \cdot \frac{b}{a}, \frac{m}{a}\right) = \gcd\left(\frac{n}{d} \cdot \frac{d}{c}, \frac{m}{a}\right) = \gcd\left(\frac{m}{a}, \frac{n}{c}\right).$$
mpletes the proof.

This completes the proof.

Based on formulas (15) and (16), Nowak and Tóth [26, 27] proved the following results. For every  $z, w \in \mathbb{C}$  with  $\Re z > 1, \Re w > 1$ ,

$$\sum_{n,n=1}^{\infty} \frac{s(m,n)}{m^z n^w} = \frac{\zeta^2(z)\zeta^2(w)\zeta(z+w-1)}{\zeta(z+w)},$$

and

$$\sum_{n,n \le x} s(m,n) = x^2 \sum_{r=0}^{3} A_r (\log x)^r + O\left(x^{\frac{1117}{701} + \varepsilon}\right),$$
(18)

for every fixed  $\varepsilon > 0$ , where  $A_3 = 2/\pi^2$ ,  $A_2, A_1, A_0$  are explicit constants, and 1117/701  $\doteq$ 1.593437. Nowak and Tóth [26, 27] also proved a similar formula with the same error term for the sum  $\sum_{n_1,n_2 \leq x, (n_1,n_2)>1} s(n_1,n_2)$  concerning the number of all subgroups of the rank two groups  $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2}$ . The error term of (18) is the same as of (7), and can be replaced by  $O\left(x^{\frac{3-\theta}{2-\theta}+\varepsilon}\right)$ , where  $\theta$  is the exponent in Dirichlet's divisor problem. Tóth and Zhai [49] proved, by using a two dimensional Perron formula and the complex integration method that the error term of (18) can be refined into  $O(x^{3/2}(\log x)^{13/2})$ .

Sui and Liu [36] deduced an asymptotic formula for the hyperbolic summation  $\sum_{mn \leq x} s(m, n)$ , and studied the order of magnitude of the corresponding error term. Kiuchi and Saad Eddin [20] deduced an asymptotic formula for the weighted sum  $\sum_{mn \leq x} s(m, n) \log \frac{x}{mn}$ .

Sui and Liu [37] proved by a method which is elementary in essence, but using some preliminary results of analytic nature, that

$$\sum_{\substack{m,n\in\mathbb{N}\\n^2+n^2\leqslant x}} s(m,n) = xP(\log x) + O\left(x^{17/22+\varepsilon}\right),$$

where P(t) is a polynomial in t of degree three. By using a two dimensional Perron formula and the complex integration method, the same authors [38] proved that for every fixed  $k \ge 2$ ,

$$\sum_{m,n \leq x} s(m^k, n^k) = C_k x^{k+1} + O\left(x^{k+1/2+\varepsilon}\right),$$

where  $C_k$  is an absolute constant.

It was mentioned in our arXiv preprint [26] that

$$\sum_{n \le x} s(n,n) = \frac{5\pi^2}{24} x^2 + O\left(x(\log x)^{8/3}\right),\tag{19}$$

the error term coming from the result of Walfisz for the Dedekind function  $\psi(n) = n \prod_{p|n} (1 + 1/p)$ .

Tărnăuceanu and Tóth [43, Prop. 3.2] proved that the sum of exponents of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$  is  $\sigma(m)\sigma(n)$  for every  $m, n \in \mathbb{N}$ . Therefore, the arithmetic mean of exponents of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$  is  $\sigma(m)\sigma(n)/s(m,n)$ . Now consider the case m = n, and let AE(n)stand for the arithmetic mean of exponents of the subgroups of  $\mathbb{Z}_n \times \mathbb{Z}_n$ . We have

$$AE(n) = \frac{\sigma(n)^2}{s(n,n)},$$

and, see same authors [43, Prop. 3.3],

$$\sum_{n \leqslant x} AE(n) = \frac{C}{2}x^2 + O\left(x\log^3 x\right),$$

where

$$C := \prod_{p} \left( 1 - \frac{1}{p} \right) \sum_{\nu=0}^{\infty} \frac{(p^{\nu+1} - 1)^2}{p^{2\nu}(p^{\nu+2} + p^{\nu+1} - (2\nu+3)p + 2\nu + 1)}$$

A generalization of the function s(m, n) is  $\sigma_t(m, n)$ , defined as the sum of t-th powers of the orders of subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ , that is,

$$\sigma_t(m,n) = \sum_{H \leqslant \mathbb{Z}_m \times \mathbb{Z}_n} |H|^t,$$

which is multiplicative. If t = 0, then  $\sigma_0(m, n) = s(m, n)$ . Here we prove the following related results.

**Theorem 1.** Let  $m, n \in \mathbb{N}$  and  $t \in \mathbb{C}$ . Then

$$\sigma_t(m,n) = \sum_{i|m,j|n} \operatorname{lcm}(i,j)^t (\operatorname{id}_t * \varphi)(\operatorname{gcd}(i,j))$$
(20)

$$= \sum_{d|\gcd(m,n)} \varphi(d) d^t \sigma_t\left(\frac{m}{d}\right) \sigma_t\left(\frac{n}{d}\right).$$
(21)

*Proof.* We use the representation of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ , given by (13). We obtain

$$\sigma_t(m,n) = \sum_{\substack{(a,b,c,d,\ell) \in J_{m.n}}} |K_{a,b,c,d,\ell}|^t = \sum_{\substack{a|m \ c|n \ a/b = c/d = e \\ b|a \ d|c}} \sum_{\substack{\ell \le e \\ (\ell,e) = 1}} (ad)^t$$
(22)

Let m = ax, a = by, n = cz, c = dt. Rearranging the terms of (22),

$$\sigma_t(m,n) = \sum_{bxe=m} \sum_{dze=n} \varphi(e)(bde)^t = \sum_{\substack{ix=m\\jz=n}} i^t \sum_{\substack{be=i\\de=j}} (j/e)^t \varphi(e)$$
(23)

$$=\sum_{\substack{i|m\\j|n}} (ij)^t \sum_{\substack{e|\gcd(i,j)\\j|n}} \frac{\varphi(e)}{e^t} = \sum_{\substack{i|m\\j|n}} \frac{(ij)^t}{\gcd(i,j)^t} (\mathrm{id}_t \ast \varphi)(\gcd(i,j)),$$

using that for every  $n \in \mathbb{N}$ ,

$$\sum_{e|n} \frac{\varphi(e)}{e^t} = \frac{1}{n^t} \sum_{e|n} e^t \ \varphi(n/e) = \frac{1}{n^t} (\mathrm{id}_t * \varphi)(n),$$

and finishing the proof of (20). Now write (23) as follows:

$$\sigma_t(m,n) = \sum_{\substack{ek=m\\e\ell=n}} \varphi(e)e^t \sum_{\substack{bx=k\\dz=\ell}} (bd)^t = \sum_{\substack{ek=m\\e\ell=n}} \varphi(e)e^t \sum_{\substack{bk} b^t} \sum_{\substack{d|\ell}} d^t$$
$$= \sum_{\substack{e|\gcd(m,n)}} \varphi(e)e^t \sigma_t\left(\frac{m}{e}\right) \sigma_t\left(\frac{n}{e}\right),$$

giving (21).

Note that in the case t = 0, (20) and (21) recover identities (15) and (16), respectively.

**Theorem 2.** We have the Dirichlet series representation

$$\sum_{m,n=1}^{\infty} \frac{\sigma_t(m,n)}{m^z n^w} = \frac{\zeta(z)\zeta(z-t)\zeta(w)\zeta(w-t)\zeta(z+w-t-1)}{\zeta(z+w-t)}.$$
(24)

and for t = 1,

$$\sum_{m,n \leqslant x} \sigma_1(m,n) = \frac{\pi^6}{864\zeta(3)} x^4 + O\left(x^3 (\log x)^{5/3}\right).$$
(25)

*Proof.* Here (24) is a direct consequence of identity (21). If t = 1, then using (21) we have

$$\sum_{m,n\leqslant x}\sigma_1(m,n) = \sum_{ek\leqslant x, e\ell\leqslant x}\varphi(e)e\sigma(k)\sigma(\ell) = \sum_{e\leqslant x}\varphi(e)e\left(\sum_{k\leqslant x/e}\sigma(k)\right)^2,$$

and (25) follows applying formula (10), with Walfisz' error term, and using standard arguments.  $\Box$ 

# 5 Number of all subgroups in the case k = 3

Now consider the group  $\Gamma := \mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r$ , where  $m, n, r \in \mathbb{N}$ . By using only simple group-theoretic and number-theoretic arguments, Hampejs and Tóth [17] proved the following result concerning the representation of the subgroups of  $\Gamma$ .

(i) Choose  $a, b, c \in \mathbb{N}$  such that  $a \mid m, b \mid n, c \mid r$ .

(ii) Compute  $A := \operatorname{gcd}(a, n/b), B := \operatorname{gcd}(b, r/c), C := \operatorname{gcd}(a, r/c).$ 

(iii) Compute

$$X := \frac{ABC}{\gcd(a(r/c), ABC)}.$$

(iv) Let s := at/A, where  $0 \le t \le A - 1$ . (v) Let

$$v := \frac{bX}{B \operatorname{gcd}(t, X)} w$$
, where  $0 \le w \le B \operatorname{gcd}(t, X)/X - 1$ .

(vi) Find a solution  $u_0$  of the linear congruence

$$(r/c)u \equiv rvs/(bc) \pmod{a}$$
.

(vii) Let  $u := u_0 + az/C$ , where  $0 \le z \le C - 1$ . (viii) Consider

$$U_{a,b,c,t,w,z} := \langle (a,0,0), (s,b,0), (u,v,c) \rangle$$

 $=\{(ia+js+ku,jb+kv,kc): 0\leqslant i\leqslant n/a-1, 0\leqslant j\leqslant n/b-1, 0\leqslant k\leqslant n/c-1\}.$ 

Then  $U_{a,b,c,t,w,z}$  is a subgroup of order mnr/(abc) of  $\Gamma$ . Moreover, there is a bijection between the set of sixtuples (a, b, c, t, w, z) satisfying conditions (i)-(viii) and the set of subgroups of  $\Gamma$ .

Let  $P(n) = \sum_{k=1}^{n} \gcd(k, n) = \sum_{d|n} d\varphi(n/d)$  be the gcd-sum function. Note that the function P is multiplicative and  $P(p^{\nu}) = (\nu + 1)p^{\nu} - \nu p^{\nu-1}$  for every prime power  $p^{\nu}$  ( $\nu \in \mathbb{N}$ ).

For every  $m, n, r \in \mathbb{N}$  the total number of subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r$  is given by

$$s(m,n,r) = \sum_{a|m,b|n,c|r} \frac{ABC}{X^2} P(X),$$
(26)

where A, B, C are defined above. The number of subgroups of order  $\delta$  ( $\delta \mid mnr$ ) is given by (26) with the additional condition that the summation is subject to  $abc = mnr/\delta$ . If one of m, n, r is 1, then formula (26) reduces to (15).

It follows that for every prime p and every  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{N}$ ,  $s(p^{\lambda_1}, p^{\lambda_2}, p^{\lambda_3})$  is a polynomial in p with integer coefficients. Using different arguments, namely the formulas in terms of the gaussian coefficients mentioned in the Introduction, it was shown by Tărnăuceanu and Tóth [43, Cor. 2.2] that the corresponding explicit formula is, with  $\lambda_1 \ge \lambda_2 \ge \lambda_3 \ge 1$ ,

$$s(p^{\lambda_1}, p^{\lambda_2}, p^{\lambda_3}) = \frac{F(p)}{(p^2 - 1)^2(p - 1)}$$

where

$$F(p) = (\lambda_3 + 1)(\lambda_1 - \lambda_2 + 1)p^{\lambda_2 + \lambda_3 + 5} + 2(\lambda_3 + 1)p^{\lambda_2 + \lambda_3 + 4} - 2(\lambda_3 + 1)(\lambda_1 - \lambda_2)p^{\lambda_2 + \lambda_3 + 3} - 2(\lambda_3 + 1)p^{\lambda_2 + \lambda_3 + 2} + (\lambda_3 + 1)(\lambda_1 - \lambda_2 - 1)p^{\lambda_2 + \lambda_3 + 1} - (\lambda_1 + \lambda_2 - \lambda_3 + 3)p^{2\lambda_3 + 4} - 2p^{2\lambda_3 + 3} + (\lambda_1 + \lambda_2 - \lambda_3 - 1)p^{2\lambda_3 + 2} + (\lambda_1 + \lambda_2 + \lambda_3 + 5)p^2 + 2p - (\lambda_1 + \lambda_2 + \lambda_3 + 1).$$

This formula was also obtained by Oh [28, Cor. 2.2], using different arguments.

The final result of this paragraph is an asymptotic formula for the number s(n, n, n) of subgroups of  $\mathbb{Z}_n^3$ . Define the multiplicative function h by

$$s(n,n,n) = \sum_{d|n} d^2 \tau(d) h(n/d) \quad (n \in \mathbb{N})$$

and let  $H(z) = \sum_{n=1}^{\infty} h(n) n^{-z}$  be the Dirichlet series of h.

We have, see Hampejs and Tóth [17, Th. 2.3] that for every  $\varepsilon > 0$ ,

$$\sum_{n \le x} s(n, n, n) = \frac{x^3}{3} \left( H(3)(\log x + 2\gamma - 1) + H'(3) \right) + O\left(x^{2+\theta+\varepsilon}\right),$$
(27)

where H'(z) is the derivative of H(z),  $\gamma$  is Euler's constant and  $\theta$  is the exponent in Dirichlet's divisor problem.

# 6 Number of all subgroups in the case k = 4

Now consider the group  $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$ , and let N(m, n, r, s) and N(m, n, r, s; k) denote the total number of its subgroups and the number of its subgroups of order k, respectively.

We state and prove the following results, already given in the arXiv paper by the author [48]. Namely, we deduce direct formulas for N(m, n, r, s) and N(m, n, r, s; k), where  $m, n, r, s, k \in \mathbb{N}$ ,  $k \mid mnrs$ , see Theorems 3 and 4. As direct consequences, in Corollaries 1 and 2 we derive formulas for the number of subgroups of *p*-groups of rank four. We do not obtain the explicit form of the corresponding polynomials, but those can be computed in special cases. We present some tables for such polynomials, computed using Corollaries 1 and 2. For the proofs we use the representation of the subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$ , Goursat's lemma (see Section 4) and Proposition 2. A table of values for N(n) := N(n, n, n, n) is also included.

**Theorem 3.** For every  $m, n, r, s \in \mathbb{N}$  we have

$$N(m, n, r, s) = \sum \varphi(x_3)\varphi(y_3)\varphi(z_3)\varphi(t_3)F(u, v), \qquad (28)$$

where  $\varphi$  is Euler's function, F(u, v) represents the number of automorphisms of the group  $\mathbb{Z}_u \times \mathbb{Z}_v$ , and the sum is over all

 $(x_1, x_2, x_3, x_4, x_5, y_1, y_2, y_3, y_4, y_5, z_1, z_2, z_3, z_4, z_5, t_1, t_2, t_3, t_4, t_5) \in \mathbb{N}^{20}$ 

satisfying the following 10 conditions:

(1)  $x_1x_2x_3 = m$ , (2)  $x_3x_4x_5 = n$ , (3)  $y_1y_2y_3 = \gcd(x_1, x_4)$ ,

- (4)  $y_3y_4y_5 = x_3 \operatorname{lcm}(x_1, x_4),$
- $(4) \ g_3g_4g_5 = x_3 \operatorname{rem}(x_1, x_4)$
- (5)  $z_1 z_2 z_3 = r$ ,
- (6)  $z_3 z_4 z_5 = s$ ,

(7)  $t_1 t_2 t_3 = \gcd(z_1, z_4),$ (8)  $t_3 t_4 t_5 = z_3 \operatorname{lcm}(z_1, z_4),$ (9)  $\gcd(y_2, y_5) = \gcd(t_2, t_5) =: u,$ (10)  $y_3 \operatorname{lcm}(y_2, y_5) = t_3 \operatorname{lcm}(t_2, t_5) =: v.$ 

*Proof.* We apply Goursat's lemma for the groups  $G = \mathbb{Z}_m \times \mathbb{Z}_n$  and  $H = \mathbb{Z}_r \times \mathbb{Z}_s$ . Let

$$A = K_{a,b,c,d,\ell} \leqslant \mathbb{Z}_m \times \mathbb{Z}_n,$$

where  $a \mid m, b \mid a, c \mid n, d \mid c, \frac{a}{b} = \frac{c}{d} = e, 1 \leq \ell \leq e, \gcd(\ell, e) = 1$ . Here, by (14),

$$A \simeq \mathbb{Z}_{m_1} \times \mathbb{Z}_{n_1},$$

where  $m_1 = \gcd(b, d), n_1 = \operatorname{lcm}(a, c)$ . Also, let

$$B \simeq K_{a_1, b_1, c_1, d_1, \ell_1} \leqslant A,$$

where  $a_1 \mid m_1, b_1 \mid a_1, c_1 \mid n_1, d_1 \mid c_1, \frac{a_1}{b_1} = \frac{c_1}{d_1} = e_1, 1 \leq \ell_1 \leq e_1, \gcd(\ell_1, e_1) = 1$ . We have, by using Proposition 2,

$$A/B \simeq \mathbb{Z}_u \times \mathbb{Z}_v,\tag{29}$$

where  $u = \gcd(m_1/a_1, n_1/c_1), v = \operatorname{lcm}(m_1/b_1, n_1/d_1).$ 

In a similar way, let

$$C = K_{\alpha,\beta,\gamma,\delta,\lambda} \leqslant \mathbb{Z}_r \times \mathbb{Z}_s,$$

where  $\alpha \mid r, \beta \mid \alpha, \gamma \mid s, \delta \mid \gamma, \frac{\alpha}{\beta} = \frac{\gamma}{\delta} = \epsilon, 1 \leq \lambda \leq \epsilon, \gcd(\lambda, \epsilon) = 1$ . Here,

 $C \simeq \mathbb{Z}_{r_1} \times \mathbb{Z}_{s_1},$ 

where  $r_1 = \gcd(\beta, \delta), s_1 = \operatorname{lcm}(\alpha, \gamma)$ . Furthermore, let

$$D \simeq K_{\alpha_1,\beta_1,\gamma_1,\delta_1,\lambda_1} \leqslant C,$$

with  $\alpha_1 \mid r_1, \beta_1 \mid \alpha_1, \gamma_1 \mid s_1, \delta_1 \mid \gamma_1, \frac{\alpha_1}{\beta_1} = \frac{\gamma_1}{\delta_1} = \epsilon_1, 1 \leq \lambda_1 \leq \epsilon_1, \gcd(\lambda_1, \epsilon_1) = 1$ . Here,

$$C/D \simeq \mathbb{Z}_{\eta} \times \mathbb{Z}_{\theta},\tag{30}$$

where  $\eta = \gcd(r_1/\alpha_1, s_1/\gamma_1), \ \theta = \operatorname{lcm}(r_1/\beta_1, s_1/\delta_1).$ 

According to (29) and (30), the quotient groups A/B and C/D are isomorphic. if and only if  $u = \eta$  and  $v = \theta$ . In this case there are F(u, v) isomorphisms  $A/B \simeq C/D$ .

It follows that

$$N(m, n, r, s) = \sum F(u, v),$$

where the sum is over all

$$(a, b, c, d, \ell, a_1, b_1, c_1, d_1, \ell_1, \alpha, \beta, \gamma, \delta, \lambda, \alpha_1, \beta_1, \gamma_1, \delta_1, \lambda_1) \in \mathbb{N}^{20}$$

satisfying the above properties.

More exactly, let m = ax, a = by, n = cz, c = dt, where y = t = e. Hence m = bxe, n = dze. Also,  $m_1 = b_1 x_1 e_1, n_1 = d_1 z_1 e_1$ , where  $m_1 = \gcd(b, d), n_1 = \operatorname{lcm}(a, c) = e \operatorname{lcm}(b, d)$ . Similarly, let  $r = \alpha X, \alpha = \beta Y, s = \gamma Z, \gamma = \delta T$ , where  $Y = T = \epsilon$ . Hence  $r = \beta X \epsilon, s = \delta Z \epsilon$ . Also,  $r_1 = \beta_1 X_1 \epsilon_1, s_1 = \delta_1 Z_1 \epsilon_1$ , where  $r_1 = \gcd(\beta, \delta), r_1 = \operatorname{lcm}(\alpha, \gamma) = \epsilon \operatorname{lcm}(\beta, \delta)$ .

We deduce that

$$N(m, n, r, s) = \sum F(u, v),$$

where the sum is over all

 $(b, x, e, d, z, \ell, b_1, x_1, e_1, d_1, z_1, \ell_1, \beta, X, \epsilon, \delta, Z, \lambda, \beta_1, X_1, \epsilon_1, \delta_1, Z_1, \lambda_1) \in \mathbb{N}^{24}$ 

satisfying the following conditions:

(1a) bxe = m, (2a) dze = n, (3a)  $b_1x_1e_1 = \gcd(b, d)$ , (4b)  $1 \le \ell \le e, \gcd(\ell, e) = 1$ , (5a)  $\beta X \epsilon = r$ , (6a)  $\delta Z \epsilon = s$ , (7a)  $\beta_1 X_1 \epsilon_1 = \gcd(\beta, \delta)$ , (8a)  $\delta_1 Z_1 \epsilon_1 = \epsilon \operatorname{lcm}(\beta, \delta)$ , (8b)  $1 \le \ell_1 \le e_1, \gcd(\ell_1, e_1) = 1$ , (9a)  $\gcd(x_1, z_1) = \gcd(X_1, Z_1) = u$ , (10a)  $e_1 \operatorname{lcm}(x_1, z_1) = \epsilon_1 \operatorname{lcm}(X_1, Z_1) = v$ , (10b)  $1 \le \lambda \le \epsilon, \gcd(\lambda, \epsilon) = 1$ , (10c)  $1 \le \lambda_1 \le \epsilon_1, \gcd(\lambda_1, \epsilon_1) = 1$ .

Note that, according to (4b), (8b), (10b) and (10c), the variables  $\ell$ ,  $\ell_1$ ,  $\lambda$  and  $\lambda_1$  take  $\varphi(e)$ ,  $\varphi(e_1)$ ,  $\varphi(\epsilon)$ , respectively  $\varphi(\epsilon_1)$  distinct values, independently from the other variables. Now formula (28) follows by introducing the notation

 $\begin{array}{l} x_1 := b, \, x_2 := x, \, x_3 := e, \, x_4 := d, \, x_5 := z, \, y_1 := b_1, \, y_2 := x_1, \, y_3 := e_1, \, y_4 := d_1, \, y_5 := z_1 \\ z_1 := \beta, \, z_2 := X, \, z_3 := \epsilon, \, z_4 := \delta, \, z_5 := Z, \, t_1 := \beta_1, \, t_2 := X_1, \, t_3 := \epsilon_1, \, t_4 := \delta_1, \, t_5 := Z_1. \end{array}$ 

We remark that the number F(m, n) of automorphisms of the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , used above, can be computed as follows. For every  $m, n \in \mathbb{N}$ ,

$$F(m,n) = \prod_{p} F(p^{\nu_{p}(m)}, p^{\nu_{p}(n)}),$$

hence  $(m, n) \mapsto F(m, n)$  is a multiplicative arithmetic function of two variables. For prime powers  $p^a, p^b$  one has, cf. [19, Th. 4.1], [13, Cor. 3],

$$F(p^{a}, p^{b}) = \begin{cases} p^{2a}\varphi(p^{a})\varphi(p^{b}), & \text{if } 0 \leq a < b; \\ p^{a}\varphi_{2}(p^{a})\varphi(p^{a}), & \text{if } 0 \leq a = b; \end{cases}$$

where  $\varphi_2$  is the Jordan function of order 2 given by  $\varphi_2(n) = n^2 \prod_{p|n} (1 - \frac{1}{p^2})$ . We deduce that  $F(1, p^b) = \varphi(p^b)$  for  $b \in \mathbb{N}_0$  and

$$F(p^{a}, p^{b}) = \begin{cases} p^{3a+b} \left(1 - \frac{1}{p}\right)^{2}, & \text{if } 1 \leq a < b; \\ p^{4a} \left(1 - \frac{1}{p}\right) \left(1 - \frac{1}{p^{2}}\right), & \text{if } 1 \leq a = b. \end{cases}$$

**Theorem 4.** For every  $k, m, n, r, s \in \mathbb{N}$  such that  $k \mid mnrs$ , the number N(m, n, r, s; k) is given by the sum (28) with the additional constraint

 $(11) \quad x_1 x_3 x_4 t_1 t_3 t_4 = k.$ 

*Proof.* According to the second part of Goursat's lemma, see Section 4, we have k = |A||D| = |B||C|. Here |A| = ad = bde,  $|B| = a_1d_1 = b_1d_1e_1$ . Similarly,  $|C| = \alpha\delta = \beta\delta\epsilon$ ,  $|D| = \alpha_1\delta_1 = \beta_1\delta_1\epsilon_1$ . Hence condition k = |A||D| = |B||C| is equivalent to  $bde\beta_1\delta_1\epsilon_1 = k$  and to  $x_1x_3x_4t_1t_3t_4 = k$  with the changing of notation of above. Note that relation k = |B||C| is a consequence of the other constraints and needs not be considered.

**Corollary 1.** For every  $a, b, c, d \in \mathbb{N}_0$  we have

$$N(p^{a}, p^{b}, p^{c}, p^{d}) = \sum \varphi(p^{x_{3}})\varphi(p^{y_{3}})\varphi(p^{z_{3}})\varphi(p^{t_{3}})F(p^{u}, p^{v}),$$
(31)

where the sum is over all

 $(x_1, x_2, x_3, x_4, x_5, y_1, y_2, y_3, y_4, y_5, z_1, z_2, z_3, z_4, z_5, t_1, t_2, t_3, t_4, t_5) \in \mathbb{N}_0^{20}$ 

satisfying the following 10 conditions:

(i)  $x_1 + x_2 + x_3 = a$ , (ii)  $x_3 + x_4 + x_5 = b$ , (iii)  $y_1 + y_2 + y_3 = \min(x_1, x_4)$ , (iv)  $y_3 + y_4 + y_5 = x_3 + \max(x_1, x_4)$ , (v)  $z_1 + z_2 + z_3 = c$ , (vi)  $z_3 + z_4 + z_5 = d$ , (vii)  $t_1 + t_2 + t_3 = \min(z_1, z_4)$ , (viii)  $t_3 + t_4 + t_5 = z_3 + \max(z_1, z_4)$ , (ix)  $\min(y_2, y_5) = \min(t_2, t_5) =: u$ , (x)  $y_3 + \max(y_2, y_5) = t_3 + \max(t_2, t_5) =: v$ .

*Proof.* Apply Theorem 3 in the case  $(m, n, r, s) = (p^a, p^b, p^c, p^d)$ .

**Corollary 2.** For every  $a, b, c, d, k \in \mathbb{N}_0$  with  $k \leq a + b + c + d$  the number  $N(p^a, p^b, p^c, p^d; p^k)$  is given by the sum (31) with the additional constraint

(xi)  $x_1 + x_3 + x_4 + t_1 + t_3 + t_4 = k$ .

*Proof.* Apply Theorem 4 in the case 
$$(m, n, r, s) = (p^a, p^b, p^c, p^d), k := p^k$$
.

[]

Consider the polynomials  $N(p^a, p^b, p^c, p^d)$  and  $N(p^a, p^b, p^c, p^d; p^k)$ , where p is prime,  $1 \leq a \leq b \leq c \leq d, 0 \leq k \leq a+b+c+d =: n$ . It is known that  $N(p^a, p^b, p^c, p^d; p^k) = N(p^a, p^b, p^c, p^d; p^{n-k})$  for every  $0 \leq k \leq n$ . The following unimodality result is also known:  $N(p^a, p^b, p^c, p^d; p^k) - N(p^a, p^b, p^c, p^d; p^{k-1})$  has non-negative coefficients for every  $1 \leq k \leq \lfloor n/2 \rfloor$ . More generally, similar properties hold for every abelian p-group. See Butler [6], Takegahara [40].

In what follows we give the polynomials representing the values of  $N(p^a, p^b, p^c, p^d)$  and  $N_{p^k}(p^a, p^b, p^c, p^d)$ , where  $1 \leq a \leq b \leq c \leq d \leq 2$ , also a = b = c = d = 3, with  $0 \leq k \leq \lfloor n/2 \rfloor$ , computed using Corollaries 1 and 2. Some more tables are given in [48].

N(p, p, p, p)	$5 + 3p + 4p^2 + 3p^3 + p^4$
N(p, p, p, p; 1)	1
N(p, p, p, p; p)	$1 + p + p^2 + p^3$
$N(p, p, p, p; p^2)$	$1 + p + 2p^2 + p^3 + p^4$

$N(p, p, p, p^2)$	$6 + 4p + 6p^2 + 6p^3 + 2p^4$
$N(p, p, p, p^2; 1)$	1
$N(p, p, p, p^2; p)$	$1 + p + p^2 + p^3$
$N(p, p, p, p^2; p^2)$	$1 + p + 2p^2 + 2p^3 + p^4$

$N(p, p, p^2, p^2)$	$7 + 5p + 8p^2 + 9p^3 + 6p^4 + p^5$
$N(p, p, p^2, p^2; 1)$	1
$N(p, p, p^2, p^2; p)$	$1 + p + p^2 + p^3$
$N(p, p, p^2, p^2; p^2)$	$1 + p + 2p^2 + 2p^3 + 2p^4$
$N(p,p,p^2,p^2;p^3)$	$1 + p + 2p^2 + 3p^3 + 2p^4 + p^5$

$N(p, p^2, p^2, p^2)$	$8 + 6p + 10p^2 + 12p^3 + 10p^4 + 6p^5 + 2p^6$
$N(p, p^2, p^2, p^2; 1)$	1
$N(p, p^2, p^2, p^2; p)$	$1 + p + p^2 + p^3$
$N(p, p^2, p^2, p^2; p^2)$	$1 + p + 2p^2 + 2p^3 + 2p^4 + p^5$
$N(p, p^2, p^2, p^2; p^3)$	$1 + p + 2p^2 + 3p^3 + 3p^4 + 2p^5 + p^6$

$N(p^2, p^2, p^2, p^2)$	$9 + 7p + 12p^2 + 15p^3 + 14p^4 + 11p^5 + 9p^6 + 3p^7 + p^8$
$N(p^2, p^2, p^2, p^2; 1)$	1
$N(p^2, p^2, p^2, p^2; p)$	$1 + p + p^2 + p^3$
$N(p^2, p^2, p^2, p^2; p^2)$	$1 + p + 2p^2 + 2p^3 + 2p^4 + p^5 + p^6$
$N(p^2, p^2, p^2, p^2; p^3)$	$1 + p + 2p^2 + 3p^3 + 3p^4 + 3p^5 + 2p^6 + p^7$
$N(p^2, p^2, p^2, p^2; p^4)$	$1 + p + 2p^2 + 3p^3 + 4p^4 + 3p^5 + 3p^6 + p^7 + p^8$

$N(p^3, p^3, p^3, p^3, p^3)$	$13 + 11p + 20p^2 + 27p^3 + 30p^4 + 31p^5 + 37p^6 + 30p^7$
	$+26p^8 + 18p^9 + 9p^{10} + 3p^{11} + p^{12}$
$N(p^3, p^3, p^3, p^3; 1)$	1
$N(p^3, p^3, p^3, p^3; p)$	$1 + p + p^2 + p^3$
$N(p^3, p^3, p^3, p^3; p^2)$	$1 + p + 2p^2 + 2p^3 + 2p^4 + p^5 + p^6$
$N(p^3, p^3, p^3, p^3; p^3)$	$1 + p + 2p^2 + 3p^3 + 3p^4 + 3p^5 + 3p^6 + 2p^7 + p^8 + p^9$
$N(p^3, p^3, p^3, p^3; p^4)$	$1 + p + 2p^2 + 3p^3 + 4p^4 + 4p^5 + 5p^6 + 4p^7 + 4p^8 + 2p^9$
	$+p^{10}$
$N(p^3, p^3, p^3, p^3; p^5)$	$1 + p + 2p^2 + 3p^3 + 4p^4 + 5p^5 + 6p^6 + 6p^7 + 5p^8 + 4p^9$
	$+2p^{10}+p^{11}$
$N(p^3, p^3, p^3, p^3; p^6)$	$1 + p + 2p^2 + 3p^3 + 4p^4 + 5p^5 + 7p^6 + 6p^7 + 6p^8 + 4p^9$
	$+3p^{10} + p^{11} + p^{12}$

In our preprint [48] we have formulated the following conjectures, confirmed by the above examples. For every  $1 \leq a \leq b \leq c \leq d$ , the degree of the polynomial  $N(p^a, p^b, p^c, p^d)$  is 2a+b+c. For every  $1 \leq m$  the degree of the polynomial  $N(p^m, p^m, p^m, p^m)$  is 4m and its leading coefficient is 1.

These conjectures have been answered in the affirmative by Chew, Chin and Lim [10]. They used some different arguments, namely by constructing a recurrence relation for the number of subgroups of rank four *p*-groups, they considered the number of subgroups contained in the maximal subgroups. They also deduced an explicit formula for  $N(p^a, p^b, p^c, p^d)$ , having main term  $(d - c + 1)(b - a + 1)p^{2a+b+c}$ , where  $1 \le a \le b \le c \le d$ .

Admasu and Sehgal [1] used a different approach and found an explicit formula for the polynomials  $N(p^m, p^m, p^m, p^m)$  with  $m \in \mathbb{N}$ , also proving the second conjecture of above.

The values of N(n) := N(n, n, n, n) for  $1 \le n \le 30$  are given by the next table.

n	N(n)	n	N(n)	n	N(n)
1	1	11	19156	21	774224
2	67	12	420396	22	1283452
3	212	13	35872	23	318532
4	1983	14	244684	24	9187868
5	1 1 2 0	15	237440	25	810969
6	14204	16	821335	26	2403424
7	3652	17	99472	27	2222704
8	43339	18	1610211	28	7241916
9	24033	19	152404	29	783904
10	75040	20	2220960	30	15908480

# 7 Final remarks

Identity (4) by the author is representing the number  $c(n_1, \ldots, n_k)$  of cyclic subgroups of the group  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$  with arbitrary  $k \ge 2$ . We note that it has also been cited, resp. applied by Kumar [22, 23], Plavnik et. al. [30]. It is not known any similar compact identity for the total number  $s(n_1, \ldots, n_k)$  of subgroups. However, Ramaré [31, Eq. (20)] provided an explicit but involved formula to compute  $s(n_1, \ldots, n_k)$  and also the corresponding generalized function (3), by using the primary decomposition of  $\mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_k}$ .

We mention two other related known results. Let  $G_{(p)}$  be an arbitrary *p*-group of type  $\lambda = (\lambda_1, \ldots, \lambda_r)$ , that is  $G_{(p)} \simeq \mathbb{Z}_{p^{\lambda_1}} \times \cdots \times \mathbb{Z}_{p^{\lambda_r}}$ , where  $\lambda_1 \ge \ldots \ge \lambda_r \ge 1$ . Tărnăuceanu [42, Prop. 3.2] proved that the number of maximal subgroups of  $G_{(p)}$  is  $1 + p + \cdots + p^{k-1}$ . Chew, Chin and Lim [10, Prop. 5.1] showed, by using a general recurrence relation for the number of subgroups of rank four, and extending their method to higher rank, that the number of subgroups of  $G_{(p)}$  is  $1 + \lambda_1 + \cdots + \lambda_k \pmod{p}$ .

As remarked in the Introduction, an open problem is to find an asymptotic formula for the sum  $\sum_{n_1,\ldots,n_k \leq s} s(n_1,\ldots,n_k)$  with  $k \geq 3$ . See Section 4 for known results on  $\sum_{m,n \leq x} s(m,n)$ . It would be also interesting to study asymptotic properties of the one variable function  $s(n,\ldots,n)$ , and to find an asymptotic formula for the sum  $\sum_{n \leq x} s(n,\ldots,n)$  with  $k \geq 4$ . See the known formulas (19) and (27) corresponding to the cases k = 2 and k = 3, respectively.

### References

- F. K. Admasu and A. Sehgal, Counting subgroups of fixed order in finite abelian groups, J. Discrete Math. Sci. Cryptogr. 24 (2021), no. 1, 263–276.
- [2] G. Bhowmik, Evaluation of divisor functions of matrices, Acta Arith. 74 (1996), 155–159.
- [3] K. Bauer, D. Sen, P. Zvengrowski, A generalized Goursat lemma, Tatra Mt. Math. Publ. 64 (2015), 1–19.
- [4] J. Bourgain and N. Watt, Mean square of zeta function, circle problem and divisor problem revisited, Preprint, 2017, https://doi.org/10.48550/arXiv.1709.04340. This paper has been withdrawn by the second author in July 2023.
- [5] R. de La Bretèche, Estimation de sommes multiples de fonctions arithmétiques, Compositio Math. 128 (2001), 261–298.
- [6] L. M. Butler, A unimodality result in the enumeration of subgroups of a finite abelian group, Proc. Amer. Math. Soc. 101 (1987), 771–775.
- [7] L. M. Butler, Subgroup Lattices and Symmetric Functions, Mem. Amer. Math. Soc., vol. 112, no. 539, 1994.
- [8] W. C. Calhoun, Counting the subgroups of some finite groups, Amer. Math. Monthly 94 (1987), 54–59.

- [9] G. Călugăreanu, The total number of subgroups of a finite abelian group, Sci. Math. Japan.
   60 (2004), 157–167.
- [10] C. Y. Chew, A. Y. M. Chin, and C. S. Lim, The number of subgroups of finite abelian p-groups of rank 4 and higher, *Comm. Algebra* 48 (2020), no. 4, 1538–1547.
- [11] S. Delsarte, Fonctions de Möbius sur les groupes abeliens finis, Annals of Math. 49 (1948), 600–609.
- [12] D. Essouabri, C. Salinas Zavala, and L. Tóth, Mean values of multivariable multiplicative functions and applications to the average number of cyclic subgroups and multivariable averages associated with the LCM function, J. Number Theory 236 (2022), 404–442.
- [13] M. Golasiński and D. L. Gonçalves, On automorphisms of finite abelian p-groups, Math. Slovaca 58 (2008), 405–412.
- [14] M. J. Grady, A group theoretic approach to a famous partition formula, Amer. Math. Monthly 112 (2005), 645–651.
- [15] B. Gruber, Alternative formulae for the number of sublattices, Acta Cryst. (1997) A53, 807–808.
- [16] M. Hampejs, N. Holighaus, L. Tóth, and C. Wiesmeyr, Representing and counting the subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , J. Numbers (2014), Article ID 491428.
- [17] M. Hampejs and L. Tóth, On the subgroups of finite Abelian groups of rank three, Annales Univ. Sci. Budapest., Sect. Comp. 39 (2013), 111–124.
- [18] P. Haukkanen and R. Sivaramakrishnan, On certain trigonometric sums in several variables, *Collect. Math.* 45 (1994), 245–261.
- [19] Ch. J. Hillar and D. L. Rhea, Automorphisms of finite abelian groups, Amer. Math. Monthly 114 (2007), 917–923.
- [20] I. Kiuchi and S. Saad Eddin, On the weighted average number of subgroups of  $\mathbb{Z}_m \times \mathbb{Z}_n$  with  $mn \leq x$ , Int. J. Number Theory 18 (2022), 2005–2013.
- [21] J. Knopfmacher, Solomon's zeta function and enumeration of lattices over orders, Analysis 5 (1985), 29–42.
- [22] P. Kumar, Maximal cyclic subgroups of a finite abelian p-group of rank two, Quasigroups Related Systems 28 (2020), 237–242.
- [23] P. Kumar, A note on Shen's conjecture on groups with given same-order type, Math. Notes 111 (2022), 899–902.
- [24] G. A. Miller, On the subgroups of an abelian group, Ann. of Math. (2) 6 (1904), 1–6.

- [25] G. A. Miller, Number of the subgroups of any given abelian group, Proc. Nat. Acad. Sci. U.S.A. 25 (1939), 258–262.
- [26] W. G. Nowak and L. Tóth, On the average number of subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , Preprint, 2013, arXiV version of [27], https://doi.org/10.48550/arXiv.1307.1414.
- [27] W. G. Nowak and L. Tóth, On the average number of subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , Int. J. Number Theory 10 (2014), 363–374.
- [28] J.-M. Oh, An explicit formula for the number of subgroups of a finite abelian p-group up to rank 3, Commun. Korean Math. Soc. 28 (2013), 649–667.
- [29] J. Petrillo, Counting subgroups in a direct product of finite cyclic groups, College Math. J. 42 (2011), 215–222.
- [30] J. Plavnik, A. Schopieray, Z. Yu, and Q. Zhang, Modular tensor categories, subcategories, and Galois orbits, *Transform. Groups*, accepted for publication (2023), https://doi.org/10.1007/s00031-022-09787-9
- [31] O. Ramaré, Rationality of the zeta function of the subgroups of abelian p-groups, Publ. Math. Debrecen 90 (2017), 91–105.
- [32] R. Schmidt, Subgroup Lattices of Groups, de Gruyter Expositions in Mathematics 14, de Gruyter, Berlin, 1994.
- [33] J. Schulte, Über die Jordansche Verallgemeinerung der Eulerschen Funktion, Results Math. 36 (1999), 354–364.
- [34] V. N. Shokuev, An expression for the number of subgroups of a given order of a finite p-group, Math. Notes 12 (1972), 774–778 (1973).
- [35] T. Stehling, On computing the number of subgroups of a finite abelian group, Combinatorica 12 (1992), 475–479.
- [36] Y. Sui and D. Liu, On the error term concerning the number of subgroups of the groups  $\mathbb{Z}_m \times \mathbb{Z}_n$  with  $mn \leq x, J$ . Number Theory **216** (2020), 264–279.
- [37] Y. Sui and D. Liu, Error term concerning number of subgroups of group  $\mathbb{Z}_m \times \mathbb{Z}_n$  with  $m^2 + n^2 \leq x$ , Front. Math. China 17 (2022), 987–999.
- [38] Y. Sui and D. Liu, On the average number of subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n$  with k-th power sequence, Ramanujan J. 58 (2022), 1333–1351.
- [39] M. Suzuki, On the lattice of subgroups of finite groups, Trans. Amer. Math. Soc., 70 (1951), 345–371.
- [40] Y. Takegahara, On Butler's unimodality result, Combinatorica 18 (1998), 437–439.

- [41] M. Tărnăuceanu, A new method of proving some classical theorems of abelian groups, Southeast Asian Bull. Math. 31 (2007), 1191–1203.
- [42] M. Tărnăuceanu, An arithmetic method of counting the subgroups of a finite abelian group, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) 53(101) (2010), 373–386.
- [43] M. Tărnăuceanu and L. Tóth, On the number of subgroups of a given exponent in a finite abelian group, Publ. Inst. Math. (Beograd) (N.S.) 101(115) (2017), 121–133.
- [44] L. Tóth, Menon's identity and arithmetical sums representing functions of several variables, Rend. Sem. Mat. Univ. Politec. Torino 69 (2011), 97–110.
- [45] L. Tóth, On the number of cyclic subgroups of a finite Abelian group, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) 55(103) (2012), 423–428.
- [46] L. Tóth, Subgroups of finite Abelian groups having rank two via Goursat's lemma, Tatra Mt. Math. Publ. 59 (2014), 93–103.
- [47] L. Tóth, Multiplicative arithmetic functions of several variables: a survey, in *Mathematics Without Boundaries, Surveys in Pure Mathematics*. Th. M. Rassias, P. Pardalos (Eds.), Springer, 2014, 483–514.
- [48] L. Tóth, The number of subgroups of the group  $\mathbb{Z}_m \times \mathbb{Z}_n \times \mathbb{Z}_r \times \mathbb{Z}_s$ , Preprint, 2016, https://arxiv.org/abs/1611.03302
- [49] L. Tóth and W. Zhai, On the error term concerning the number of subgroups of the groups  $\mathbb{Z}_m \times \mathbb{Z}_n$  with  $m, n \leq x$ , Acta Arith. 183 (2018), no. 3, 285–299.
- [50] L. Tóth and W. Zhai, On the average number of cyclic subgroups of the groups  $\mathbb{Z}_{n_1} \times \mathbb{Z}_{n_2} \times \mathbb{Z}_{n_3}$  with  $n_1, n_2, n_3 \leq x$ , Res. Number Theory 6 (2020), no. 1, Art. 12, 33 pp.
- [51] Y. Yeh, On prime power abelian groups, Bull. Amer. Math. Soc. 54 (1948), 323–327.
- [52] Y. M. Zou, Gaussian binomials and the number of sublattices, Acta Cryst. 62 (2006), 409–410.