

INDEPENDENT GUE PROCESSES OF PERFECT MATCHINGS ON RAIL-YARD GRAPHS

ZHONGYANG LI

ABSTRACT. We study perfect matchings on the rail-yard graphs in which the right boundary condition is given by the empty partition and the left boundary can be divided into finitely many alternating line segments where all the vertices along each line segment are either removed or remained. When the edge weights satisfy certain conditions, we show that the distributions of the locations of certain types of dimers near the right boundary converge to the spectra of independent GUE minor processes. The proof is based on new quantitative analysis of a formula to compute Schur functions at general points discovered in [28].

1. INTRODUCTION

A Gaussian Unitary ensemble (GUE) is an $\infty \times \infty$ random Hermitian matrix $[H_{i,j}]_{i,j \in \mathbb{N}}$ whose entries are given by independent random variables such that $H_{i,i} \sim \mathcal{N}(0, 1)$ and for $i < j$, $H_{i,j} \sim \mathcal{N}(0, \frac{1}{2}) + \mathbf{i}\mathcal{N}(0, \frac{1}{2})$ is a standard complex Gaussian random variable, whose real part and imaginary part are independent.

A perfect matching, or a dimer cover of a graph is a subset of edges satisfying the constraint that each vertex is incident to exactly one edge. Perfect matchings are natural models in statistical mechanics, for example, dimer configurations on the hexagonal lattice model the molecule structure of graphite. See [20, 11] for more details.

Perfect matchings on 2-dimensional Euclidean lattice have been studied extensively. Among a number of interesting topics including phase transition ([22]), limit shape ([8, 21, 6, 25, 23, 28, 31, 26, 33, 29, 5, 3]), conformal invariance ([18, 19, 9, 24, 10, 2]), the relations between distributions of random perfect matchings and spectra of GUE minor processes have attracted significant interest among probabilists for a long time. Here is an incomplete list. Since the work of Johansson and Nordenstam [17], it is known that the uniform random domino tilings of large size Aztec diamonds behave, near the point of tangency of the inscribed arctic circle with the edge of the Aztec diamond, like the spectra of the principal minors of a GUE-matrix. This was also done in the work of Okounkov and Reshetikhin [39] for plane partitions. In the case of the uniform perfect matching on a hexagon lattice, the result was proved in [13, 37]. The convergence of locations of certain types of dimers near the tangent points to the spectra of GUE process in the scaling limit when the graph is a square-hexagon lattice with non-uniform weights was shown in [6]. It was then proved in [12] whenever a boundary of the domain has three adjacent straight

segments inclined under 120 degrees to each other, the asymptotics of uniform random lozenge tilings near the middle segment is described by the GUE.

Multiple GUE minor processes arising from the same statistical mechanical model have gained significant attention recently, see for example [1, 36].

Rail-yard graphs are a general class of graphs introduced in [7] on which the random dimer coverings form Schur processes. Limit shapes and height fluctuations of dimer coverings on rail-yard graphs were studied in [32, 33, 30]. The main result we proved in this paper, roughly speaking, is as follows

Theorem 1.1. *When the edge weights satisfy certain conditions, the distributions of the locations of certain types of dimers near the right boundary of a rail-yard graph converge to the spectra of independent GUE minor processes in the scaling limit.*

See Propositions 6.1 and 5.2 for precise statements.

The major differences of this paper and previous works lie in the following

- Previous works focus on GUE and dimer coverings on either hexagonal lattice or square grid; while our graph is the general rail-yard graph.
- Previous works focus on uniform dimer coverings and study the distribution of certain types of dimers near one tangent points; when multiple tangent points are indeed discussed (see for example [1]), these tangent points are formed due to the structure of the graph; while multiple tangent points in our paper are formed only because of the edge weights. Moreover, while [1] considers 2 GUE processes, our results applies to n GUE processes for arbitrary finite n .
- We develop new techniques of asymptotic analysis of a formula to compute Schur functions at general points discovered in [28]. We also introduce new difference operators acting on schur polynomials, which, instead of acting on all the variables in a symmetric way, acts on part of the variables in a symmetric way, which give us the marginal distribution of certain parts of the random partition.

The organization of the paper is as follows. In Section 2, we review the definitions of weighted rail-yard graphs, dimer coverings and known results to compute the partition function of dimer coverings and Schur generating functions on such graphs. We also introduce piecewise boundary conditions and the corresponding edge weights. In Section 3, we express the Schur generating function with given piecewise boundary condition as the product of Schur generating functions with boundary conditions given by one piece of the original boundary condition, when the edge weights satisfy certain conditions. We also construct subgraphs of rail yard graphs on which the Schur generating function of dimer coverings is given by factors of the original Schur generating function. The proof is based on new analysis of Schur generating functions inspired by sampling algorithms in [7]. In Section 4, we introduce new difference operators, and prove that its action on the Schur generating function gives the marginal distribution of certain parts of the random partitions corresponding to random dimer coverings on rail yard graphs. In Section 5, we prove that different parts of random partitions are asymptotically independent. In Section 6, we prove that near the right boundary, the distribution of each part of the random partition

converges to the spectra of GUE matrices when the size of the graph goes to infinity. In Section A, we include a combinatorial formula to compute Schur functions proved in [28].

2. BACKGROUND

In this section, we review the definitions and properties of the Gaussian Unitary Ensemble (GUE), weighted rail-yard graphs, dimer coverings and known results to compute the partition function of dimer coverings and Schur generating functions on such graphs. We then introduce the piecewise boundary condition and corresponding edge weights.

2.1. Spectra of GUE matrices. A matrix of the GUE of size k is diagonalizable with real eigenvalues $\epsilon_1 \geq \epsilon_2 \geq \dots \geq \epsilon_k$, whose distribution $\mathbb{P}_{\text{GUE}_k}$ on \mathbb{R}^k has a density with respect to the Lebesgue measure on \mathbb{R}^k proportional to:

$$\prod_{1 \leq i < j \leq k} (\epsilon_i - \epsilon_j)^2 \exp\left(-\sum_{i=1}^k \epsilon_i^2\right),$$

See [35, Theorem 3.3.1]. Here is the main theorem we prove in this section.

The following lemma represents the Schur function as a matrix integral over the unitary group, the so-called Harish–Chandra–Itzykson–Zuber integral:

Lemma 2.1 ([15, 16]). *Let $\lambda \in \mathbb{Y}_N$ be a partition, and let B be an $N \times N$ diagonal matrix given by*

$$B = \text{diag}[\lambda_1 + N - 1, \dots, \lambda_j + N - j, \dots, \lambda_N + N - N]$$

Let $(a_1, a_2, \dots, a_N) \in \mathbb{C}^N$, and let A be an $N \times N$ diagonal matrix given by

$$A = \text{diag}[a_1, \dots, a_N].$$

Then,

$$(2.1) \quad \frac{s_\lambda(e^{a_1}, \dots, e^{a_N})}{s_\lambda(1, \dots, 1)} = \prod_{1 \leq i < j \leq N} \frac{a_i - a_j}{e^{a_i} - e^{a_j}} \int_{U(N)} e^{\text{Tr}(U^* A U B)} dU,$$

where dU is the Haar probability measure on the unitary group $U(N)$.

Recall the following lemma giving a necessary and sufficient condition for the distribution of eigenvalues of $k \times k$ GUE random matrix.

Lemma 2.2. *Let $(q_1, \dots, q_k) \in \mathbb{R}^k$ be a random vector with distribution \mathbb{P} and $Q = \text{diag}[q_1, \dots, q_k]$ the $k \times k$ diagonal matrix obtained by putting the q_i on the diagonal.*

Then \mathbb{P} is $\mathbb{P}_{\text{GUE}_k}$ if and only if for any matrix P ,

$$\mathbb{E} \int_{U(k)} \exp[\text{Tr}(P U Q U^*)] dU = \exp\left(\frac{1}{2} \text{Tr} P^2\right).$$

It is in fact enough to check the case when P is diagonal, with real coefficients.

2.2. Weighted rail-yard graphs. Let $l, r \in \mathbb{Z}$ such that $l \leq r$. Let

$$[l..r] := [l, r] \cap \mathbb{Z},$$

i.e., $[l..r]$ is the set of integers between l and r . For a positive integer m , we use the notation $[m]$ to denote the set of all the integers from 1 to m , i.e.

$$[m] := \{1, 2, \dots, m\}.$$

Consider two binary sequences indexed by integers in $[l..r]$

- the LR sequence $\underline{a} = \{a_l, a_{l+1}, \dots, a_r\} \in \{L, R\}^{[l..r]}$;
- the sign sequence $\underline{b} = (b_l, b_{l+1}, \dots, b_r) \in \{+, -\}^{[l..r]}$.

The rail-yard graph $RYG(l, r, \underline{a}, \underline{b})$ with respect to integers l and r , the LR sequence \underline{a} and the sign sequence \underline{b} , is the bipartite graph with vertex set $[2l - 1..2r + 1] \times \{\mathbb{Z} + \frac{1}{2}\}$. A vertex is called even (resp. odd) if its abscissa is an even (resp. odd) integer. Each even vertex $(2m, y)$, $m \in [l..r]$ is incident to 3 edges, two horizontal edges joining it to the odd vertices $(2m - 1, y)$ and $(2m + 1, y)$ and one diagonal edge joining it to

- the odd vertex $(2m - 1, y + 1)$ if $(a_m, b_m) = (L, +)$;
- the odd vertex $(2m - 1, y - 1)$ if $(a_m, b_m) = (L, -)$;
- the odd vertex $(2m + 1, y + 1)$ if $(a_m, b_m) = (R, +)$;
- the odd vertex $(2m + 1, y - 1)$ if $(a_m, b_m) = (R, -)$.

See Figure 2.2 for an example of a rail-yard graph.

The left boundary (resp. right boundary) of $RYG(l, r, \underline{a}, \underline{b})$ consists of all odd vertices with abscissa $2l - 1$ (resp. $2r + 1$). Vertices which do not belong to the boundaries are called inner. A face of $RYG(l, r, \underline{a}, \underline{b})$ is called an inner face if it contains only inner vertices.

We assign edge weights to a rail yard graph $RYG(l, r, \underline{a}, \underline{b})$ as follows:

- all the horizontal edges have weight 1; and
- each diagonal edge adjacent to a vertex with abscissa $2i$ has weight x_i .

2.3. Dimer coverings and pure dimer coverings.

Definition 2.3. A dimer covering is a subset of edges of $RYG(l, r, \underline{a}, \underline{b})$ such that

- (1) each inner vertex of $RYG(l, r, \underline{a}, \underline{b})$ is incident to exactly one edge in the subset;
- (2) each left boundary vertex or right boundary vertex is incident to at most one edge in the subset;
- (3) only a finite number of diagonal edges are present in the subset.

See Figure 2.2 for an example of a dimer covering on a rail yard graph.

2.4. Partitions. Let N be a non-negative integer. A length- N partition is a non-increasing sequence $\lambda = (\lambda_i)_{i \geq 0}^N$ of non-negative integers. Let \mathbb{Y}_N be the set of all the length- N partitions. The size of a partition $\lambda \in \mathbb{Y}_N$ is defined by

$$|\lambda| = \sum_{i=1}^N \lambda_i.$$




FIGURE 2.1. A rail yard graph with LR sequence $\underline{a} = \{L, R, R, L\}$ and sign sequence $\underline{b} = \{+, +, -, -\}$. Odd vertices are represented by red points, and even vertices are represented by blue points. Dark lines represent a pure dimer covering. Assume that above the horizontal line $y = 4$, only horizontal edges with an odd vertex on the left are present in the dimer configuration; and below the horizontal line $y = -4$, only horizontal edges with an even vertex on the left are present in the dimer configuration.

We denote the length of a partition by $l(\lambda)$. In particular, if $\lambda \in \mathbb{Y}_N$, then $l(\lambda) = N$. Two partitions $\lambda \in \mathbb{Y}_N$ and $\mu \in \mathbb{Y}_{N-1}$ are called interlaced, and written by $\lambda \succ \mu$ or $\mu \prec \lambda$ if

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \lambda_3 \geq \dots \geq \mu_{N-1} \geq \lambda_N.$$

A length- ∞ partition is a non-increasing sequence $\lambda = (\lambda_i)_{i \geq 0}^\infty$ of non-negative integers which vanishes eventually. Each length- N partition can be naturally extended to an length- ∞ partition by adding infinitely many 0's at the end of the sequence.

When representing partitions by Young diagrams, $\mu \prec \lambda$ means λ/μ is a horizontal strip. The conjugate partition λ' of λ is a partition whose Young diagram $Y_{\lambda'}$ is the image of the Young diagram Y_λ of λ by the reflection along the main diagonal. More precisely

$$\lambda'_i := |\{j \geq 0 : \lambda_j \geq i\}|, \quad \forall i \geq 1.$$

The skew Schur functions are defined in Section I.5 of [34].

Definition 2.4. Let λ, μ be partitions of finite length. Define the skew Schur functions as follows

$$s_{\lambda/\mu} = \det \left(h_{\lambda_i - \mu_j - i + j} \right)_{i,j=1}^{l(\lambda)}$$

Here for each $r \geq 0$, h_r is the r th complete symmetric function defined by the sum of all monomials of total degree r in the variables x_1, x_2, \dots . More precisely,

$$h_r = \sum_{1 \leq i_1 \leq i_2 \leq \dots \leq i_r} x_{i_1} x_{i_2} \cdots x_{i_r}$$

If $r < 0$, $h_r = 0$.

Define the Schur function as follows

$$s_\lambda = s_{\lambda/\emptyset}.$$

For a dimer covering M of $RYG(l, r, \underline{a}, \underline{b})$, we associate a particle-hole configuration to each odd vertex of $RYG(l, r, \underline{a}, \underline{b})$ as follows: let $m \in [l..(r+1)]$ and $k \in \mathbb{Z}$: if the odd endpoint $(2m-1, k + \frac{1}{2})$ is incident to a present edge in M on its right (resp. left), then associate a hole (resp. particle) to the odd endpoint $(2m-1, k + \frac{1}{2})$. When M is a pure dimer covering, it is not hard to check that there exists $N > 0$, such that when $y > N$, only holes exist and when $y < -N$, only particles exist.

We associate a partition $\lambda^{(M,m)}$ to the column indexed by m of particle-hole configurations, which corresponds to a pure dimer covering M adjacent to odd vertices with abscissa $(2m-1)$ as follows. Assume

$$\lambda^{(M,m)} = (\lambda_1^{(M,m)}, \lambda_2^{(M,m)}, \dots),$$

Then for $i \geq 1$, $\lambda_i^{(M,m)}$ is the total number of holes in M along the vertical line $x = 2m-1$ below the i th highest particles. Let $l(\lambda^{(M,m)})$ be the total number of nonzero parts in the partition $\lambda^{(M,m)}$.

We define the charge $c^{(M,m)}$ on column $(2m-1)$ for the configuration M as follows:

$$(2.2) \quad c^{(M,m)} = \begin{aligned} & \text{number of particles on column } (2m-1) \text{ in the upper half plane} \\ & - \text{number of holes on column } (2m-1) \text{ in the lower half plane} \end{aligned}$$

Recall the following lemma proved in [30].

Lemma 2.5. *Let M be a dimer covering of $RYG(l, r, \underline{a}, \underline{b})$. Then for each $m \in [l..r]$, the central charge $c^{(M,m)}$ is a constant independent of m .*

Proof. See Lemma 3.2 of [30]. □

The weight of a dimer covering M of $RYG(l, r, \underline{a}, \underline{b})$ is defined as follows

$$w(M) := \prod_{i=l}^r x_i^{d_i(M)},$$




FIGURE 2.2. Particle-hole configuration corresponding to the dimer covering in Figure 2.2. Particles are represented by black dots, while holes are represented by circles. Each particle/hole represents the configuration of an odd vertex (represented by red dots) in the same location of Figure 2.2. The corresponding sequence of partitions (from the left to the right) is given by $\emptyset \prec (2, 0, \dots) \prec' (3, 1, 1, \dots) \succ' (2, 0, \dots) \succ \emptyset$.

where $d_i(M)$ is the total number of present diagonal edges of M incident to an even vertex with abscissa $2i$.

Let $\lambda^{(l)}, \lambda^{(r+1)}$ be two partitions. The partition function $Z_{\lambda^{(l)}, \lambda^{(r+1)}}(G, \underline{x})$ of dimer coverings on $RYG(l, r, \underline{a}, \underline{b})$ whose configurations on the left (resp. right) boundary correspond to partition $\lambda^{(l)}$ (resp. $\lambda^{(r+1)}$) is the sum of weights of all such dimer coverings on the graph. Given the left and right boundary conditions $\lambda^{(l)}$ and $\lambda^{(r+1)}$, respectively, the probability of a dimer covering M is then defined by

$$(2.3) \quad \Pr(M | \lambda^{(l)}, \lambda^{(r+1)}) := \frac{w(M)}{Z_{\lambda^{(l)}, \lambda^{(r+1)}}(G, \underline{x})}.$$

Note that pure dimer coverings have left and right boundary conditions given by

$$(2.4) \quad \lambda^{(l)} = \lambda^{(r+1)} = \emptyset;$$

respectively.

The bosonic Fock space \mathcal{B} is the infinite dimensional Hilbert space spanned by the orthonormal basis vectors $|\lambda\rangle$, where λ runs over all the partitions. Let $\langle \lambda |$ denote the dual basis vector. Let x be a formal or a complex variable. Introduce the operators $\Gamma_{L+}(x)$,

$\Gamma_{L-}(x), \Gamma_{R+}(x), \Gamma_{R-}(x)$ from \mathcal{B} to \mathcal{B} as follows

$$\begin{aligned}\Gamma_{L+}(x)|\lambda\rangle &= \sum_{\mu \prec \lambda} x^{|\lambda|-|\mu|} |\mu\rangle; & \Gamma_{R+}(x)|\lambda\rangle &= \sum_{\mu' \prec \lambda'} x^{|\lambda|-|\mu'|} |\mu'\rangle; \\ \Gamma_{L-}(x)|\lambda\rangle &= \sum_{\mu \succ \lambda} x^{|\mu|-|\lambda|} |\mu\rangle; & \Gamma_{R-}(x)|\lambda\rangle &= \sum_{\mu' \succ \lambda'} x^{|\mu'|-|\lambda|} |\mu'\rangle;\end{aligned}$$

Such operators were first studied in [38] for random partitions.

Lemma 2.6. *Let $a_1, a_2 \in \{L, R\}$. We have the following commutation relations for the operators $\Gamma_{a_1, \pm}, \Gamma_{a_2, \pm}$.*

$$\Gamma_{a_1, +}(x_1)\Gamma_{a_2, -}(x_2) = \begin{cases} \frac{\Gamma_{a_2, -}(x_2)\Gamma_{a_1, +}(x_1)}{1-x_1x_2} & \text{if } a_1 = a_2 \\ (1+x_1x_2)\Gamma_{a_2, -}(x_2)\Gamma_{a_1, +}(x_1) & \text{if } a_1 \neq a_2. \end{cases}$$

Moreover,

$$\Gamma_{a_1, b}(x_1)\Gamma_{a_2, b}(x_2) = \Gamma_{a_2, b}(x_2)\Gamma_{a_1, b}(x_1);$$

for all $a_1, a_2 \in \{L, R\}$ and $b \in \{+, -\}$.

Proof. See Proposition 7 of [7]; see also [40, 4]. □

Given the definitions of the operators $\Gamma_{a, b}(x)$ with $a \in \{L, R\}$, $b \in \{+, -\}$, it is straightforward to check the following lemma.

Lemma 2.7. *The partition function of dimer coverings on a rail yard graph $G = RYG(l, r, \underline{a}, \underline{b})$ with left and right boundary conditions given by $\lambda^{(l)}, \lambda^{(r+1)}$, respectively, is*

$$(2.5) \quad Z_{\lambda^{(l)}, \lambda^{(r+1)}}(G; \mathbf{x}) = \langle \lambda^{(l)} | \Gamma_{a_l b_l}(x_l) \Gamma_{a_{l+1} b_{l+1}}(x_{l+1}) \cdots \Gamma_{a_r b_r}(x_r) | \lambda^{(r+1)} \rangle;$$

where

$$(2.6) \quad \mathbf{x} := (x_l, \dots, x_r)$$

denote the edge weights.

Corollary 2.8. (1) *Assume for all $i \in [l, r]$,*

$$(2.7) \quad (a_i, b_i) \neq (R, -).$$

Let g_l be the number of nonzero entries in $\lambda^{(l)}$, then

$$(2.8) \quad g_l \leq |i \in [l..r] : a_i = L, b_i = -|.$$

The partition function of dimer coverings on a rail yard graph $G = RYG(l, r, \underline{a}, \underline{b})$ with left and right boundary conditions given by $\lambda^{(l)}, \emptyset$, respectively, can be computed as follows:

$$(2.9) \quad Z_{\lambda^{(l)}, \emptyset}(G; \mathbf{x}) = s_{\lambda^{(l)}} \left(\mathbf{x}^{(L, -)} \right) \prod_{l \leq i < j \leq r; b_i = +, b_j = -} z_{i, j}$$

where

$$(2.10) \quad z_{ij} = \begin{cases} 1 + x_i x_j & \text{if } a_i \neq a_j \\ \frac{1}{1 - x_i x_j} & \text{if } a_i = a_j \end{cases}$$

and

$$\mathbf{x}^{(L,-)} := \{x_i : (a_i, b_i) = (L, -)\}.$$

(2) Assume for all $i \in [l, r]$,

$$(2.11) \quad (a_i, b_i) \neq (L, -).$$

Let g'_l be the number of nonzero entries in $\lambda^{(l)}$, then

$$g'_l \leq |i \in [l..r] : a_i = L, b_i = -|.$$

The partition function of dimer coverings on a rail yard graph $G = RYG(l, r, \underline{a}, \underline{b})$ with left and right boundary conditions given by $\lambda^{(l)}, \emptyset$, respectively, can be computed as follows:

$$(2.12) \quad Z_{\lambda^{(l)}, \emptyset}(G; \mathbf{x}) = s_{[\lambda^{(l)}]'}(\mathbf{x}^{(R,-)}) \prod_{l \leq i < j \leq r; b_i = +, b_j = -} z_{i,j}$$

where

$$\mathbf{x}^{(R,-)} := \{x_i : (a_i, b_i) = (R, -)\}.$$

Proof. See Corollary 2.5 of [32]. □

Assumption 2.9. For any $i, j \in [l..r]$, $i < j$, $a_i = a_j$ and $b_i = +$, $b_j = -$ we have

$$x_i x_j < 1.$$

2.5. Schur generating functions.

Definition 2.10. Let $\mathbf{X} = (x_1, \dots, x_N) \in \mathbb{C}^N$. Let ρ be a probability measure on \mathbb{Y}_N . The Schur generating function $\mathcal{S}_{\rho, \mathbf{X}}(u_1, \dots, u_N)$ with respect to parameters \mathbf{X} is the symmetric Laurent series in (u_1, \dots, u_N) given by

$$\mathcal{S}_{\rho, \mathbf{X}}(u_1, \dots, u_N) = \sum_{\lambda \in \mathbb{Y}} \rho(\lambda) \frac{s_{\lambda}(u_1, \dots, u_N)}{s_{\lambda}(x_1, \dots, x_N)}.$$

Lemma 2.11. Let $t \in [l..r]$. Assume (2.7) holds. Let

$$\begin{aligned} \mathbf{x}^{(L,-,>t)} &= \{x_i : i \in [t+1..r], a_i = L, b_i = -\}. \\ \mathbf{x}^{(L,-,\leq t)} &= \{x_i : i \in [l..t], a_i = L, b_i = -\} \end{aligned}$$

Let

$$\mathbf{u}^{(L,-,>t)} = \{u_i : i \in [t+1..r], a_i = L, b_i = -\}.$$

Let ρ^t be the probability measure for the partitions corresponding to dimer configurations adjacent to the column of odd vertices labeled by $2t-1$ in $RYG(l, r, \underline{a}, \underline{b})$, conditional on the left and right boundary condition $\lambda^{(l)}$ and \emptyset , respectively.

For $i \in [l..r]$, let

$$w_i = \begin{cases} u_i & \text{if } i \in [t+1..r], a_i = L, b_i = - \\ x_i & \text{otherwise} \end{cases}$$

and

$$\xi_{ij} = \begin{cases} 1 + w_i w_j & \text{if } a_i \neq a_j \\ \frac{1}{1 - w_i w_j} & \text{if } a_i = a_j \end{cases}.$$

Then the generating Schur function $\mathcal{S}_{\rho^t, \mathbf{x}^{(L, -, > t)}}$ is given by:

$$\mathcal{S}_{\rho^t, \mathbf{x}^{(L, -, > t)}}(\mathbf{u}^{(L, -, > t)}) = \frac{s_{\lambda^{(l)}}(\mathbf{u}^{(L, -, > t)}, \mathbf{x}^{(L, -, \leq t)})}{s_{\lambda^{(l)}}(\mathbf{x}^{(L, -)})} \prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}}.$$

Proof. See Lemma 2.8 of [32]. □

2.6. Piecewise Boundary Conditions.

Assumption 2.12. Let $\epsilon > 0$ be a small positive parameter.

- (1) Let $l^{(\epsilon)} < r^{(\epsilon)}$ be the integers representing the left and right boundary of the rail-yard graph depending on ϵ such that

$$\lim_{\epsilon \rightarrow 0} \epsilon l^{(\epsilon)} = l^{(0)} < r^{(0)} = \lim_{\epsilon \rightarrow 0} \epsilon r^{(\epsilon)}$$

so that the scaling limit of the sequence of rail-yard graphs $\{\epsilon RYG(l^{(\epsilon)}, r^{(\epsilon)}, \underline{a}^{(\epsilon)}, \underline{b}^{(\epsilon)})\}_{\epsilon > 0}$, as $\epsilon \rightarrow 0$, has left boundary given by $x = 2l^{(0)}$ and right boundary given by $x = 2r^{(0)}$.

- (2) Assume for each $\epsilon > 0$, the weight of each diagonal edge incident to an even vertex with abscissa $2i$, for $i \in [l^{(\epsilon)}..r^{(\epsilon)}]$ is $x_{i,\epsilon}$.

Suppose that Assumption 2.12 holds. For each $\epsilon > 0$, define

$$N_{L,-}^{(\epsilon)} := \left| \left\{ j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j^{(\epsilon)} = L, b_j^{(\epsilon)} = - \right\} \right|.$$

Let π be the bijection from $[N_{L,-}^{(\epsilon)}]$ to $\{j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j^{(\epsilon)} = L, b_j^{(\epsilon)} = -\}$, such that

- for any $i, j \in [N_{L,-}^{(\epsilon)}]$ and $i < j$, $\pi(i) < \pi(j)$.

Let

$$(2.13) \quad N^{(\epsilon)} := r^{(\epsilon)} - l^{(\epsilon)} + 1;$$

for all $\epsilon > 0$; i.e. $N^{(\epsilon)}$ is the total number of even columns in the rail-yard graph.

Let $\Sigma_{L,-,\epsilon}$ be the group consisting of all the bijections from $\{j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j^{(\epsilon)} = L, b_j^{(\epsilon)} = -\}$ to itself.

Let $\Sigma_{L,-,\epsilon}^X$ be a subgroup of $\Sigma_{L,-,\epsilon}$ preserving the values of the diagonal edge weights $x_{i,\epsilon}$'s, i.e.

$$\Sigma_{L,-,\epsilon}^X := \{\sigma \in \Sigma_{L,-,\epsilon} : x_{\sigma(i),\epsilon} = x_{i,\epsilon}, \forall i \in [l^{(\epsilon)}..r^{(\epsilon)}], \text{ s.t. } a_i^{(\epsilon)} = L, b_i^{(\epsilon)} = -\}.$$

Note that $\pi\Sigma_{L,-,\epsilon}$, defined by

$$(2.14) \quad \pi\Sigma_{L,-,\epsilon} := \{\pi\xi : \xi \in \Sigma_{L,-,\epsilon}\},$$

consists of all the bijections from $[N_{L,-}^{(\epsilon)}]$ to $\{j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j^{(\epsilon)} = L, b_j^{(\epsilon)} = -\}$. Let $\sigma_0 \in \pi\Sigma_{L,-,\epsilon}$ such that

$$(2.15) \quad x_{\sigma_0(1),\epsilon} \geq x_{\sigma_0(2),\epsilon} \geq \dots \geq x_{\sigma_0(N_{L,-}^{(\epsilon)})\epsilon}.$$

Assumption 2.13. Suppose Assumption 2.12 holds. Let $s \in [N_{L,-}^{(\epsilon)}]$ such that s is independent of ϵ as $\epsilon \rightarrow 0$. Assume there exist positive integers $K_1^{(\epsilon)}, K_2^{(\epsilon)}, \dots, K_s^{(\epsilon)}$, such that

- (1) let $n > 0$ be the total number of distinct elements in $\{x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon)})\epsilon}\}$; then n is finite, fixed and independent of ϵ .
- (2) $\sum_{t=1}^s K_t^{(\epsilon)} = N_{L,-}^{(\epsilon)}$;
- (3) Assume the partition corresponding to the left boundary condition is $\lambda^{(l,\epsilon)}$ is given by

$$\lambda^{(l,\epsilon)} = \left(\lambda_1, \dots, \lambda_{N_{L,-}^{(\epsilon)}} \right).$$

such that

$$(2.16) \quad \mu_1^{(\epsilon)} > \dots > \mu_s^{(\epsilon)}$$

are all the distinct elements in $\left\{ \lambda_1, \lambda_2, \dots, \lambda_{N_{L,-}^{(\epsilon)}} \right\}$.

(4)

$$\begin{aligned} \lambda_1 &= \lambda_2 = \dots = \lambda_{K_s^{(\epsilon)}} = \mu_1^{(\epsilon)}; \\ \lambda_{K_s^{(\epsilon)}+1} &= \lambda_{K_s^{(\epsilon)}+2} = \dots = \lambda_{K_s^{(\epsilon)}+K_{s-1}^{(\epsilon)}} = \mu_2^{(\epsilon)}; \\ &\dots \\ \lambda_{1+\sum_{t=2}^s K_t^{(\epsilon)}} &= \lambda_{2+\sum_{t=2}^s K_t^{(\epsilon)}} = \dots = \lambda_{\sum_{t=1}^s K_t^{(\epsilon)}} = \mu_s^{(\epsilon)}; \end{aligned}$$

(5) For $i \in [n]$, let

$$(2.17) \quad J_i = \{t \in [s] : \exists q \in [N_{L,-}^{(\epsilon)}], \text{ s.t. } x_{\sigma_0(q)} \text{ is the } i\text{th largest elements in } \{x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon)})\epsilon}\}, \text{ and } \lambda_q = \mu_t^{(\epsilon)}\}$$

such that

- (a) For any $i_1, i_2 \in [n]$ with $i_1 < i_2$, if $l \in J_{i_1}$, and $t \in J_{i_2}$, then $l < t$.
- (b) For any p, q satisfying $1 \leq p \leq s$ and $1 \leq q \leq s$, and $q > p$

$$C_1 N_{L,-}^{(\epsilon)} \leq \mu_p^{(\epsilon)} - \mu_q^{(\epsilon)} \leq C_2 N_{L,-}^{(\epsilon)}$$

where C_1, C_2 are sufficiently large constants independent of ϵ as $\epsilon \rightarrow 0$.

Assumption 2.14. Suppose that Assumption 2.12 holds. For $i \in [l^{(\epsilon)}..r^{(\epsilon)}]$, suppose the weights x_i change with respect to ϵ , and denote it by $x_{i,\epsilon}$. Let σ_0 be defined as in (2.15). More precisely, we order elements in the set

$$(2.18) \quad \{x_{i,\epsilon}\}_{i \in [l^{(\epsilon)}, r^{(\epsilon)}], a_{i,p}=L, b_{i,p}=-}$$

by (2.15). Assume

$$x_{\sigma_0(1),\epsilon} = x_{\sigma_0(1)} > 0$$

i.e., the maximal weight in (2.18) is a positive constant independent of ϵ .

Suppose that Assumption 2.13 holds. Moreover, assume that

$$\liminf_{\epsilon \rightarrow 0} \epsilon \log \left(\min_{x_{i,\epsilon} > x_{j,\epsilon}} \frac{x_{i,\epsilon}}{x_{j,\epsilon}} \right) \geq \alpha > 0,$$

where α is a sufficiently large positive constant independent of ϵ .

For $N_{L,-}^{(\epsilon)} \geq 1$, let $\lambda^{(l)} \in \mathbb{Y}_{N_{L,-}^{(\epsilon)}}$ be the left boundary partition satisfying Assumption 2.13. Let

$$\Omega = \left(\Omega_1 < \Omega_2 < \dots < \Omega_{N_{L,-}^{(\epsilon)}} \right) = \left(\lambda_{N_{L,-}^{(\epsilon)}}^{(l)} - N_{L,-}^{(\epsilon)}, \lambda_{N_{L,-}^{(\epsilon)}-1}^{(l)} - N_{L,-}^{(\epsilon)} + 1, \dots, \lambda_1^{(l)} - 1 \right)$$

Indeed, $\Omega_1, \dots, \Omega_{N_{L,-}^{(\epsilon)}}$ are the locations of the $N_{L,-}^{(\epsilon)}$ topmost particles on the left boundary of the rail-yard graph, assuming that the vertical coordinate of the $N_{L,-}^{(\epsilon)}$ highest particles is $-N_{L,-}^{(\epsilon)}$ when the left boundary condition is given by the empty partition. Under Assumption 2.13, we may assume

$$(2.19) \quad \Omega = (A_1, A_1 + 1, \dots, B_1 - 1, B_1, \\ A_2, A_2 + 1, \dots, B_2 - 1, B_2, \dots, A_s, A_s + 1, \dots, B_s - 1, B_s).$$

where

$$B_i - A_i + 1 = K_i.$$

and

$$\sum_{i=1}^s (B_i - A_i + 1) = N_{L,-}^{(\epsilon)}.$$

Suppose as $\epsilon \rightarrow 0$,

$$(2.20) \quad A_i = a_i N_{L,-}^{(\epsilon)} + o(N_{L,-}^{(\epsilon)}), \quad B_i = b_i N_{L,-}^{(\epsilon)} + o(N_{L,-}^{(\epsilon)}), \quad \text{for } i \in [s],$$

and $a_1 < b_1 < \dots < a_s < b_s$ are fixed parameters independent of ϵ and satisfy $\sum_{i=1}^s (b_i - a_i) = 1$. Under Assumption 2.13, it is straightforward to check that for $i \in [s]$

$$\begin{aligned} b_i &= \lim_{N \rightarrow \infty} \frac{\mu_{s-i+1} + \sum_{t=1}^i K_t}{N_{L,-}^{(\epsilon)}} - 1 \\ a_i &= \lim_{N \rightarrow \infty} \frac{\mu_{s-i+1} + \sum_{t=1}^{i-1} K_t}{N_{L,-}^{(\epsilon)}} - 1 \end{aligned}$$

Assumption 2.15. For each $i \in [n]$, write

$$\begin{aligned} N_{L,-,i}^{(\epsilon)} &= |\{j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j = L, b_j = -\}, \\ & \quad x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(2),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon)})}, \epsilon\}| \end{aligned}$$

Suppose that for each $i \in [n]$,

$$\lim_{\epsilon \rightarrow 0} \frac{N_{L,-,i}^{(\epsilon)}}{N_{L,-}^{(\epsilon)}} = \theta_i \in (0, 1); \quad \text{and} \quad \lim_{\epsilon \rightarrow 0} \frac{N_{L,-}^{(\epsilon)}}{N^{(\epsilon)}} = \rho \in (0, 1];$$

where $N^{(\epsilon)}$ is defined by (2.13); and $\sum_{i \in [n]} \theta_i = 1$.

3. POSITIONS OF V-EDGES IN A ROW AND EIGENVALUES OF GUE RANDOM MATRIX

In this section, we prove the following results.

- (1) Under Assumptions 2.13 and 2.14 regarding edge weights, as $\epsilon \rightarrow 0$, we approximately express the Schur generating function with respect to the distribution of partitions near the right boundary into the product of Schur generating functions, each of which depend on one of the distinct weights in

$$\{x_{j,\epsilon} : j \in [l^{(\epsilon)}..r^{(\epsilon)}] : a_j = L, b_j = -\}$$

Let \mathcal{S} denote the collection of these Schur generating functions as factors of the original one.

- (2) We construct subgraphs of the rail-yard graphs such that the distribution of the partitions corresponding to dimer configurations near its right boundary has a Schur generating function as a factor in \mathcal{S} .

3.1. Product of Schur generating functions. For each $\sigma \in \pi\Sigma_{L,-,\epsilon}$ as defined by (2.14) and $j \in [N_{L,-}^{(\epsilon)}]$, let

$$(3.1) \quad \eta_j^\sigma(N_{L,-}^{(\epsilon)}) = |\{k \in [N_{L,-}^{(\epsilon)}] : k > j, x_{\sigma(k),\epsilon} \neq x_{\sigma(j),\epsilon}\}|.$$

For $1 \leq i \leq n$, let

$$(3.2) \quad \Phi^{(i,\sigma)}(N_{L,-}^{(\epsilon)}) = \{\lambda_j^{(i)} + \eta_j^\sigma : x_{\sigma(j)} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1)}, x_{\sigma_0(2)}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon)})}, \epsilon\}\}$$

and let $\phi^{(i,\sigma)}(N_{L,-}^{(\epsilon)})$ (sometimes we just write $\phi^{(i,\sigma)}$) be the partition with length $|\{1 \leq j \leq N : x_j = x_i\}|$ obtained by decreasingly ordering all the elements in $\Phi^{(i,\sigma)}(N_{L,-}^{(\epsilon)})$.

Proposition 3.1. *Suppose Assumptions 2.14, 2.13 and 2.15 hold, and let α be given as in Assumption 2.14. For each given $\{a_i, b_i\}_{i=1}^n$, when α is sufficiently large, for any $\sigma \notin \bar{\sigma}_0$ we have*

$$(3.3) \quad \left| \frac{\left(\prod_{i=1}^n x_{i,\epsilon}^{|\phi^{(i,\sigma_0)}(N_{L,-}^{(\epsilon)})|} \right) \left(\prod_{i=1}^n s_{\phi^{(i,\sigma_0)}(N_{L,-}^{(\epsilon)})}(1, \dots, 1) \right)}{\left(\prod_{i=1}^n x_{i,\epsilon}^{|\phi^{(i,\sigma)}(N_{L,-}^{(\epsilon)})|} \right) \left(\prod_{i=1}^n s_{\phi^{(i,\sigma)}(N_{L,-}^{(\epsilon)})}(1, \dots, 1) \right)} \right| \\ \times \left| \frac{\left(\prod_{i < j, x_{\sigma_0(i),\epsilon} \neq x_{\sigma_0(j),\epsilon}} \frac{1}{x_{\sigma_0(i),\epsilon} - x_{\sigma_0(j),\epsilon}} \right)}{\left(\prod_{i < j, x_{\sigma(i),\epsilon} \neq x_{\sigma(j),\epsilon}} \frac{1}{x_{\sigma(i),\epsilon} - x_{\sigma(j),\epsilon}} \right)} \right| \geq e^{C\epsilon^{-2}}$$

where $C > 0$ is a constant independent of ϵ and σ , and increases as α increases. Indeed, we have

$$\lim_{\alpha \rightarrow \infty} C = \infty.$$

Proof. Same arguments as the proof of Proposition 4.5 in [28]. \square

Recall that the Schur generating function was defined in Definition 2.10. By Lemma 2.11, we obtain

$$(3.4) \quad \mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}, \mathbf{x}^{(L,-,\leq t)}) = \frac{s_{\lambda^{(t)}}(\mathbf{u}^{(L,-,>t)}, \mathbf{x}^{(L,-,\leq t)})}{s_{\lambda^{(t)}}(\mathbf{x}^{(L,-)})} \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -} \prod_{a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}}.$$

When the edge weights satisfy Assumption 2.14 we have

$$\prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -} \prod_{a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}} = \left(1 + O(e^{-C\epsilon^{-1}})\right) \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_{j,\epsilon} = x_{\sigma_0(1),\epsilon}} \prod_{a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}}.$$

By Proposition A.1 Corollary A.2 and Proposition 3.1.

$$(3.5) \quad \frac{s_{\lambda^{(t)}}(\mathbf{u}^{(L,-,>t)}, \mathbf{x}^{(L,-,\leq t)})}{s_{\lambda^{(t)}}(\mathbf{x}^{(L,-)})} \\ = \left(1 + e^{-C\epsilon^{-2}}\right) \prod_{i=1}^n \frac{s_{\phi^{(i,\sigma_0)}}(\underline{u}^{(i)}, 1, \dots, 1)}{s_{\phi^{(i,\sigma_0)}}(1, \dots, 1)} \prod_{1 \leq i < j \leq N_{L,-}^{(\epsilon)}, x_{\sigma_0(i),\epsilon} \neq x_{\sigma_0(j),\epsilon}} \frac{x_{\sigma_0(i),\epsilon} - x_{\sigma_0(j),\epsilon}}{w_{\sigma_0(i),\epsilon} - w_{\sigma_0(j),\epsilon}},$$

where

$$\underline{u}^{(i)} = \left\{ \frac{u_j}{x_{j,\epsilon}} : x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon),\epsilon}),\epsilon}\}, \right. \\ \left. a_j = L, b_j = -, t+1 \leq j \leq r \right\}$$

Under Assumption 2.14 we have

$$\begin{aligned}
 (3.6) \quad & \prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}} \frac{x_{\sigma_0(i), \epsilon} - x_{\sigma_0(j), \epsilon}}{w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon}} \\
 &= \left(1 + e^{-C\epsilon^{-1}}\right) \prod_{i \in [N_{L, -}^{(\epsilon)}]: \sigma_0(i) \in [t+1..r]} \left(\frac{x_{\sigma_0(i), \epsilon}}{u_{\sigma_0(i)}}\right)^{|j \in [N_{L, -}^{(\epsilon)}]: j > i; x_{\sigma_0(i)} \neq x_{\sigma_0(j)}|}
 \end{aligned}$$

Recall the following lemma proved in [27].

Lemma 3.2. (1) Assume that $x_1, \dots, x_N \in \mathbb{C}$ such that

$$(3.7) \quad |\{i \in [N] : x_i = 0\}| = b \in [N]$$

Let $\lambda \in \mathbb{Y}_N$ such that there are exactly $a \in [N]$ components of λ taking value 0. If $a < b$ Then

$$(3.8) \quad s_\lambda(x_1, \dots, x_N) = 0.$$

(2) Let N be a positive integer. Let $b \in [N]$. Assume that

$$(u_1, \dots, u_{N-b}, 0^b) \in \mathbb{C}^N$$

where

$$0^b = (0, \dots, 0) \in \mathbb{R}^b$$

Let $\tilde{\lambda} \in \mathbb{Y}_N$ such that there are exactly $a \in [N]$ components of $\tilde{\lambda}$ taking value 0. If $a \geq b$, let

$$(3.9) \quad \lambda := (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_{N-b}) \in \mathbb{GT}_{N-b}$$

Then

$$s_{\tilde{\lambda}}(u_1, \dots, u_{N-b}, 0^b) = s_\lambda(u_1, \dots, u_{N-b}).$$

Proof. See Lemmas 3.1 and 3.2 in [27]. □

For $i \in [n]$, let

$$\Lambda^{(i, \sigma)} := \{\lambda_j^{(l)} : x_{\sigma(j), \epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1), \epsilon}, x_{\sigma_0(2), \epsilon}, \dots, x_{\sigma_0(N_{L, -}^{(\epsilon)})}, \epsilon\}\},$$

and let $\lambda^{(i, \sigma)}$ be the partition formed by elements in $\Lambda^{(i, \sigma)}$.

Let

$$\begin{aligned} \mathbf{u}^{(L,-,>t,i)} &= \left\{ u_j : x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon),\epsilon}),\epsilon}\}, \right. \\ &\quad \left. a_j = L, b_j = -, t+1 \leq j \leq r \right\} \\ \mathbf{x}^{(L,-,\leq t,i)} &= \left\{ x_{j,\epsilon} : x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon),\epsilon}),\epsilon}\}, \right. \\ &\quad \left. a_j = L, b_j = -, l \leq j \leq t \right\} \\ \mathbf{x}^{(L,-,i)} &= \left\{ x_{j,\epsilon} : x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(1),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon),\epsilon}),\epsilon}\}, \right. \\ &\quad \left. a_j = L, b_j = -, l \leq j \leq r \right\} \end{aligned}$$

By (3.5),(3.6), we obtain

$$\begin{aligned} \frac{s_{\lambda^{(l)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\lambda^{(l)}}(\mathbf{x}^{(L,-)})} &= \left(1 + e^{-C\epsilon^{-1}}\right) \prod_{i=1}^n \frac{s_{\phi^{(i,\sigma_0)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\phi^{(i,\sigma_0)}}(\mathbf{x}^{(L,-,i)})} \\ &\quad \times \prod_{j \in [N_{L,-}^{(\epsilon)}] : \sigma_0(j) \in [t+1..r]} \left(\frac{x_{\sigma_0(j),\epsilon}}{u_{\sigma_0(j)}} \right)^{\eta_j^{\sigma_0(N_{L,-}^{(\epsilon)})}}, \\ &= \left(1 + e^{-C\epsilon^{-1}}\right) \prod_{i=1}^n \frac{s_{\lambda^{(i,\sigma_0)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\lambda^{(i,\sigma_0)}}(\mathbf{x}^{(L,-,i)})} \end{aligned}$$

Then we can compute the Schur generating function as follows

(3.10)

$$\begin{aligned} &\mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}) \\ &= (1 + e^{-C\epsilon^{-1}}) \prod_{i=1}^n \frac{s_{\lambda^{(i,\sigma_0)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\lambda^{(i,\sigma_0)}}(\mathbf{x}^{(L,-,i)})} \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_{j,\epsilon} = x_{\sigma_0(1),\epsilon}} \prod_{i,j} \frac{\xi_{ij}}{z_{ij}}. \end{aligned}$$

Lemma 3.3.

$$\begin{aligned} (3.11) \quad &\frac{s_{\lambda^{(1,\sigma)}}(\mathbf{u}^{(L,-,>t,1)}, \mathbf{x}^{(L,-,\leq t,1)})}{s_{\lambda^{(1,\sigma)}}(\mathbf{x}^{(L,-,1)})} \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_{j,\epsilon} = x_{\sigma_0(1),\epsilon}} \prod_{i,j} \frac{\xi_{ij}}{z_{ij}} \\ &= \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda^{(1,\sigma)}}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,1}^{(>t,\epsilon)}})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,1}^{(>t,\epsilon)}})} \end{aligned}$$

For each $i \in [2..n]$,

$$(3.12) \quad \frac{s_{\lambda^{(i,\sigma)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\lambda^{(i,\sigma)}}(\mathbf{x}^{(L,-,i)})} = \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda^{(i,\sigma)}}^t(\mu) \frac{s_{\mu}(\underline{u}^{(i)}, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,i}^{(>t,\epsilon)}})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,i}^{(>t,\epsilon)}})}$$

where

(1)

$$N_{L,-}^{(>t,\epsilon)} = |\{j \in [t+1..r^{(\epsilon)}] : a_j = L, b_j = -\}|$$

$$N_{L,-,i}^{(>t,\epsilon)} = |\{j \in [t+1..r^{(\epsilon)}] : a_j = L, b_j = -, x_{j,\epsilon} \text{ is the } i\text{th greatest element in } \{x_{\sigma(1),\epsilon}, x_{\sigma(2),\epsilon}, \dots, x_{\sigma(N_{L,-}^{(\epsilon)})}, \epsilon\}\}|$$

(2) $\tilde{\rho}_{\lambda^{(1,\sigma)}}^t$ is the probability measure for the random partition corresponding to dimer configurations incident to odd vertices with abscissa $2t-1$ on a rail-yard graph $\text{RYG}(l, r, \underline{a}, \underline{b})$, left boundary condition given by

$$\tilde{\lambda}^{(1,\sigma)} = (\lambda^{(1,\sigma)}, 0, \dots, 0), \text{ s.t. } l(\tilde{\lambda}^{(1,\sigma)}) = N_{L,-}^{(\epsilon)}$$

and right boundary condition given by the empty partition; and the weight $\tilde{x}_{j,\epsilon}^{(1)}$ of diagonal edges incident to the column of even vertices with abscissa $2j$ given by

(a) $x_{j,\epsilon}$, if one of the following two conditions holds

(i) $(a_j, b_j) = (L, -)$, and $x_{j,\epsilon} = x_{\sigma_0(1),\epsilon}$; or

(ii) $(a_j, b_j) \neq (L, -)$

(b) 0, otherwise.

(3) If $i \in [2..n]$, $\tilde{\rho}_{\lambda^{(i,\sigma)}}^t$ is the probability measure for the random partition corresponding to dimer configurations incident to the column of odd vertices with abscissa $2t-1$ labeled by $(L, -)$, counting from the right on a rail-yard graph $\text{RYG}(l, r, (L, L, L, \dots, L), \underline{b})$, left boundary condition given by

$$\tilde{\lambda}^{(i,\sigma)} = (\lambda^{(i,\sigma)}, 0, \dots, 0), \text{ s.t. } l(\tilde{\lambda}^{(i,\sigma)}) = N_{L,-}^{(\epsilon)}$$

and right boundary condition given by the empty condition; and the weight $\tilde{x}_{j,\epsilon}^{(i)}$ of diagonal edges incident to the column of even vertices with abscissa $2j$ given by

(a) $x_{j,\epsilon}$, if $a_j = L, b_j = -$; and $x_{j,\epsilon}$ is the i th greatest element in $\{x_{\sigma_0(1),\epsilon}, x_{\sigma_0(2),\epsilon}, \dots, x_{\sigma_0(N_{L,-}^{(\epsilon)})}, \epsilon\}$

(b) 0, otherwise.

Proof. By Lemma 3.2,

$$\frac{s_{\lambda^{(i,\sigma)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)})}{s_{\lambda^{(i,\sigma)}}(\mathbf{x}^{(L,-,i)})} = \frac{s_{\lambda^{(i,\sigma)}}(\underline{u}^{(i)}, 1, \dots, 1)}{s_{\lambda^{(i,\sigma)}}(1, \dots, 1)} = \frac{s_{\tilde{\lambda}^{(i,\sigma)}}(\underline{u}^{(i)}, 1, \dots, 1, 0, \dots, 0)}{s_{\tilde{\lambda}^{(i,\sigma)}}(1, \dots, 1, 0, \dots, 0)}$$

By Lemma 2.7 we obtain

$$\tilde{\rho}_{\lambda^{(1,\sigma)}}^t(\mu) = \frac{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \cdots \Gamma_{a_t b_t}(\tilde{x}_{t,\epsilon}^{(1)}) | \mu \rangle \langle \mu | \Gamma_{a_{t+1} b_{t+1}}(\tilde{x}_{t+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle}{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \Gamma_{a_{l+1} b_{l+1}}(\tilde{x}_{l+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle}$$

By Lemma 2.6 and Corollary 2.8(1), we have

$$\begin{aligned} & \langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \Gamma_{a_{l+1} b_{l+1}}(\tilde{x}_{l+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle \\ &= s_{\tilde{\lambda}^{(1,\sigma)}} \left(\mathbf{x}^{(L,-,1)}, 0, \dots, 0 \right) \prod_{l \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j} \\ &= s_{\lambda^{(1,\sigma)}} \left(\mathbf{x}^{(L,-,1)} \right) \prod_{l \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j} \end{aligned}$$

and

$$\begin{aligned} & \langle \mu | \Gamma_{a_{t+1} b_{t+1}}(\tilde{x}_{t+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle \\ &= s_{\mu} \left(\mathbf{x}^{(L,-,>t,1)}, 0, \dots, 0 \right) \prod_{t+1 \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j}. \end{aligned}$$

Hence we have

$$\begin{aligned} & \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda^{(1,\sigma)}}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})} \\ &= \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \frac{s_{\mu}(\mathbf{u}^{(L,-,>t,1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})}{s_{\mu}(\mathbf{x}^{(L,-,>t,1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})} \\ & \times \frac{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \cdots \Gamma_{a_t b_t}(\tilde{x}_{t,\epsilon}^{(1)}) | \mu \rangle s_{\mu}(\mathbf{x}^{(L,-,>t,1)}, 0, \dots, 0) \prod_{t+1 \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j}}{s_{\lambda^{(1,\sigma)}}(\mathbf{x}^{(L,-,1)}) \prod_{l \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j}} \\ &= \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \frac{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \cdots \Gamma_{a_t b_t}(\tilde{x}_{t,\epsilon}^{(1)}) | \mu \rangle s_{\mu}(\mathbf{u}^{(L,-,>t,1)}, 0, \dots, 0) \prod_{t+1 \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j}}{s_{\lambda^{(1,\sigma)}}(\mathbf{x}^{(L,-,1)}) \prod_{l \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} z_{i,j}} \\ &= \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \frac{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \cdots \Gamma_{a_t b_t}(\tilde{x}_{t,\epsilon}^{(1)}) | \mu \rangle \langle \mu | \Gamma_{a_{t+1} b_{t+1}}(\tilde{u}_{t+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{u}_{r,\epsilon}^{(1)}) | \emptyset \rangle}{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \Gamma_{a_{l+1} b_{l+1}}(\tilde{x}_{l+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle} \\ & \prod_{l \leq i < j \leq r; b_i=+, b_j=-, x_{j,\epsilon}=x_{\sigma_0(1),\epsilon}} \frac{z_{i,j}}{\xi_{i,j}} \end{aligned}$$

where for $i \in [t + 1..r^{(\epsilon)}]$

$$\tilde{u}_{i,\epsilon}^{(1)} = \begin{cases} u_{i,\epsilon}; & \text{if } x_{i,\epsilon} = x_{\sigma_0(1),\epsilon} \\ 0; & \text{otherwise.} \end{cases}$$

Then we have

$$\begin{aligned} & \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda^{(1,\sigma)}}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,1}^{(>t,\epsilon)}})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,1}^{(>t,\epsilon)}})} \\ &= \frac{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \cdots \Gamma_{a_t b_t}(\tilde{x}_{t,\epsilon}^{(1)}) \Gamma_{a_{t+1} b_{t+1}}(\tilde{u}_{t+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{u}_{r,\epsilon}^{(1)}) | \emptyset \rangle}{\langle \tilde{\lambda}^{(1,\sigma)} | \Gamma_{a_l b_l}(\tilde{x}_{l,\epsilon}^{(1)}) \Gamma_{a_{l+1} b_{l+1}}(\tilde{x}_{l+1,\epsilon}^{(1)}) \cdots \Gamma_{a_r b_r}(\tilde{x}_{r,\epsilon}^{(1)}) | \emptyset \rangle} \prod_{l \leq i < j \leq r; b_i = +, b_j = -, x_{j,\epsilon} = x_{\sigma_0(1),\epsilon}} \frac{z_{i,j}}{\xi_{i,j}} \end{aligned}$$

Then (3.11) follows.

The identity (3.12) can be proved similarly. \square

By (3.10), (3.11) and (3.12), we have

$$(3.13) \quad \begin{aligned} & \mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}) \\ &= (1 + e^{-C\epsilon^{-1}}) \prod_{i=1}^n \left(\sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda^{(i,\sigma_0)}}^t(\mu) \frac{s_{\mu}(\underline{u}^{(i)}, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,i}^{(>t,\epsilon)}})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)} - N_{L,-,i}^{(>t,\epsilon)}})} \right) \end{aligned}$$

By Corollaries 2.8(1), A.2 and (3.3), we have

$$(3.14) \quad \begin{aligned} & Z_{\lambda^{(l)}, \emptyset}(G; \mathbf{x}) = (1 + e^{-C\epsilon^{-1}}) \left(s_{\lambda^{(1,\sigma_0)}}(\mathbf{x}^{(L,-,1)}) \prod_{l \leq i < j \leq r; b_i = +, b_j = -, x_{j,\epsilon} = x_{\sigma_0(1)}} z_{i,j} \right) \\ & \quad \times \prod_{i=2}^n s_{\lambda^{(i,\sigma_0)}}(\mathbf{x}^{(L,-,i)}) \end{aligned}$$

In Lemma 3.3, we constructed rail-yard graphs $RYG(l, r, \underline{a}, \underline{b})$ such that partition functions of perfect matchings on these subgraphs are

$$(3.15) \quad s_{\lambda^{(1,\sigma)}}(\mathbf{x}^{(L,-,1)}) \prod_{l \leq i < j \leq r; b_i = +, b_j = -, x_{j,\epsilon} = x_{\sigma_0(1)}} z_{i,j}$$

and

$$(3.16) \quad s_{\lambda^{(i,\sigma)}}(\mathbf{x}^{(L,-,i)}); \quad \forall i \in [2..n]$$

respectively.

Now we discuss (3.15); which is equal to

$$(3.17) \quad s_{\tilde{\lambda}^{(1,\sigma)}}(\mathbf{x}^{(L,-,1)}, 0^{N_{L,-}^{(\epsilon)} - N_{L,-,i}^{(\epsilon)}}) \prod_{l \leq i < j \leq r; b_i = +, b_j = -, x_{j,\epsilon} = x_{\sigma_0(1)}} z_{i,j}$$

Consider the weighted rail-yard graph given as in Lemma 3.3(2). Then (3.15), or equivalently (3.17), gives the partition function of dimer configurations on such a graph with left boundary condition given by $\tilde{\lambda}^{(1,\sigma)}$ and right boundary condition given by the empty partition. Let $\mathcal{M}_{1,\sigma}$ be the set consisting of all the dimer configurations on such a graph with given boundary conditions. For each $M_{1,\sigma} \in \mathcal{M}_{1,\sigma}$, let

$$\tilde{\lambda}^{(1,\sigma)} (= \nu^{(M_{1,\sigma},l)}, \nu^{(M_{1,\sigma},l-1)}, \dots, \nu^{(M_{1,\sigma},r-1)}) (= \emptyset)$$

be the sequence of partitions corresponding to $M_{1,\sigma}$. For each $t \in [l..r]$, recall that $l_+(\nu^{M_{1,\sigma},t})$ is the total number of nonzero parts in $\nu^{(M_{1,\sigma},t)}$.

Assume $2 \leq i \leq n$, then (3.16) is equal to

$$(3.18) \quad s_{\tilde{\lambda}^{(i,\sigma)}} \left(\mathbf{x}^{(L,-,1)}, 0^{N_{L,-}^{(\epsilon)} - N_{L,-,i}^{(\epsilon)}} \right)$$

Consider the weighted rail-yard graph given as in Lemma 3.3(3). Then (3.16), or equivalently (3.18), gives the partition function of dimer configurations on such a graph with left boundary condition given by $\tilde{\lambda}^{(i,\sigma)}$ and right boundary condition given by the empty partition. Let $\mathcal{M}_{i,\sigma}$ be the set consisting of all the dimer configurations on such a graph with given boundary conditions. For each $M \in \mathcal{M}_{i,\sigma}$, let

$$\tilde{\lambda}^{(i,\sigma)} (= \nu^{(M_{i,\sigma},l)}, \nu^{(M_{i,\sigma},l-1)}, \dots, \nu^{(M_{i,\sigma},r-1)}) (= \emptyset)$$

be the sequence of partitions corresponding to $M_{i,\sigma}$.

Then it is straightforward to check the following lemma

Lemma 3.4. *Assume $2 \leq i \leq n$, we have*

$$\left| \nu_1^{(M_{i,\sigma},t)} \right| \leq \tilde{\lambda}_1^{(i,\sigma)} \leq \lambda_1^{(l)}$$

Moreover, we have

Lemma 3.5. *Assume (2.7) holds.*

$$l_+(\nu^{(M_{1,\sigma},t)}) \leq N_{L,-,1}^{(>t,\epsilon)}$$

Proof. Consider dimer configurations on the graph $RYG(t, r, \underline{a}, \underline{b})$ with left boundary condition given by $\nu^{(M_{1,\sigma},t)}$, right boundary condition given by the empty partition, and edge weights satisfies Lemma 3.3(2). First of all, if the edge weights satisfy $\tilde{x}_{j,\epsilon}^{(1)} = 0$ for some j , then we must have

$$\nu^{(M_{1,\sigma},j)} = \nu^{(M_{1,\sigma},j+1)}, \quad \forall M \in \mathcal{M}_{1,\sigma}$$

We may reorder the operators Γ_{a_i, b_i} for $i \in [t..r] \setminus \{j : \tilde{x}_{j,\epsilon}^{(1)} = 0\}$, to make sure that all the operators with $b_i = -$ are before all the operators with $b_i = +$. Since (2.7) holds, whenever $b_i = -$ we have $a_i = L$. By Algorithms ?? and ??, we obtain a sequence of partitions

$$\mu^{(M,t)} \succ \tilde{\nu}^{(M,t+1)} \succ \dots \succ \tilde{\nu}^{(M,t+N_{L,-,1}^{(>t,\epsilon)})} = \emptyset \subseteq \dots \subseteq \emptyset$$

Therefore the total number of nonzero parts of $\nu^{(M_1, \sigma, t)}$ is at most $N_{L, -, 1}^{(>t, \epsilon)}$, and the lemma follows. \square

Lemma 3.6. *Suppose that (2.7) and Assumption 2.9 hold. Assume c is a constant satisfying*

$$(3.19) \quad c > \left| \frac{\log 2}{\log[\max_{i \in [l..r], b_i = +, a_i = L} x_i x_{\sigma_0(1)}]} \right|$$

Then for any $\delta > 0$,

$$(3.20) \quad \mathbb{P} \left(\max_{t \in [l..r], M \in \mathcal{M}_{1, \sigma}} \nu_1^{(M, t)} \geq c\delta[N^{(\epsilon)}]^2 + N^{(\epsilon)} + \lambda_1^{(1, \sigma)} \right) < N^{(\epsilon)} \left(\frac{1}{2} \right)^{\delta N^{(\epsilon)}}$$

Proof. Let $M \in \mathcal{M}_{1, \sigma}$ be arbitrary. For each $j \in [l..r - 1]$, the case

$$\nu_1^{(M, j+1)} > \nu_1^{(M, j)}$$

can occur only when either $\nu^{(M, j)} \prec \nu^{(M, j+1)}$ or $\nu^{(M, j)} \prec' \nu^{(M, j+1)}$. Moreover, if $\nu^{(M, j)} \prec' \nu^{(M, j+1)}$, we have

$$(3.21) \quad \nu_1^{(M, j+1)} \leq \nu_1^{(M, j)} + 1.$$

Consider all the $M \in \mathcal{M}_{1, \sigma}$ such that there exists $t \in [l..r]$ such that

$$(3.22) \quad \nu_1^{(M, t)} \geq c\delta[N^{(\epsilon)}]^2 + N^{(\epsilon)} + \lambda_1^{(1, \sigma)}$$

and let $\mathcal{Z}_{1, +}$ be the partition function (weighted sum) of all such M 's satisfying (3.22). Let \mathcal{Z}_1 be the partition function of all dimer configurations in $\mathcal{M}_{1, \sigma}$. Note that \mathcal{Z}_1 is given by (3.15). Then

$$\mathbb{P} \left(\max_{t \in [l..r], M \in \mathcal{M}_{1, \sigma}} \nu_1^{(M, t)} \geq c\delta[N^{(\epsilon)}]^2 + N^{(\epsilon)} + \lambda_1^{(1, \sigma)} \right) = \frac{\mathcal{Z}_{1, +}}{\mathcal{Z}_1}$$

Note that when $a_i = L$, $b_i = +$, $a_j = L$, $b_j = -$ and $x_{j, \epsilon} = x_{\sigma_0(1)}$, we have

$$z_{i, j} = \frac{1}{1 - x_i x_{\sigma_0(1)}} = \sum_{k=0}^{\infty} [x_i x_{\sigma_0(1)}]^k$$

Let A_1 be the contribution of $\nu_1^{(M, t)} - \lambda_1^{(1, \sigma)}$ from \prec' and let A_2 be the contribution of $\nu_1^{(M, t)} - \lambda_1^{(1, \sigma)}$ from \prec . By (3.21) we see that

$$A_1 \leq N^{(\epsilon)} - N_{L, -, \epsilon}^{(\epsilon)};$$

When (3.22) holds, we have

$$A_2 \geq c\delta[N^{(\epsilon)}]^2.$$

Since \mathcal{Z}_1 can be computed by (3.15), $\mathcal{Z}_{1, +}$ corresponds to monomials in the expansion of $\prod_{a_i = a_j = L, b_i = +, b_j = -, x_{j, \epsilon} = x_{\sigma_0(1)}} z_{i, j}$, in which the total degree of $x_i x_{\sigma_0(1)}$ for all $i \in [l..r]$, $a_i = L$, $b_i = +$ is at least $c\delta[N^{(\epsilon)}]^2$. Then there exists an i with $a_i = L$ and $b_i = +$ and the degree of $x_i x_{\sigma_0(1)}$ is at least $c\delta N^{(\epsilon)}$.

Therefore we have

$$(3.23) \quad \frac{\mathcal{Z}_{1,+}}{\mathcal{Z}_1} \leq N^{(\epsilon)} \left[\max_{i \in [l..r], b_i=+, a_i=L} x_i x_{\sigma_0(1)} \right]^{c\delta N^{(\epsilon)}}$$

Then (3.20) follows from (3.19) and (3.23). \square

4. MARGINAL DISTRIBUTION

In this section, we introduce new difference operators, and prove that its action on the Schur generating function gives the marginal distribution of certain parts of the random partitions corresponding to random dimer coverings on rail yard graphs.

Definition 4.1. For $1 \leq i \leq n$, let

$$V_i(\mathbf{u}^{(L,-,>t,i)}) := \prod_{u_j, u_r \in \mathbf{u}^{(L,-,>t,i)}; j < r} (u_j - u_r)$$

$$V_{i,*}(\mathbf{u}^{(L,-,>t,i)}) := \left[\prod_{u_j \notin \mathbf{u}^{(L,-,>t,i)}, u_r \in \mathbf{u}^{(L,-,>t,i)}; j < r} (u_j - u_r) \right] \left[\prod_{u_j \notin \mathbf{u}^{(L,-,>t,i)}, u_r \in \mathbf{u}^{(L,-,>t,i)}; j > r} (u_r - u_j) \right].$$

Introduce the family $\mathcal{D}_{i,k}$ of differential operators acting on symmetric functions f with variables $\mathbf{u}^{(L,-,>t,i)}$ as follows:

$$(4.1) \quad \mathcal{D}_{i,k} f = \frac{1}{V_i V_{i,*}} \left(\sum_{u_j \in \mathbf{u}^{(L,-,<t,i)}} \left(u_j \frac{\partial}{\partial u_j} \right)^k \right) (V_i V_{i,*} \cdot f).$$

Let

$$\bar{\mathcal{D}}_{i,k} f = \frac{1}{V_i} \left(\sum_{u_j \in \mathbf{u}^{(L,-,<t,i)}} \left(u_j \frac{\partial}{\partial u_j} \right)^k \right) (V_i \cdot f)$$

Define

$$V(\mathbf{u}^{(L,-,>t)}) = \prod_{u_j, u_r \in \mathbf{u}^{(L,-,>t)}; j < r} (u_j - u_r)$$

and

$$(4.2) \quad \mathcal{D}_k f = \frac{1}{V} \left(\sum_{u_j \in \mathbf{u}^{(L,-,>t)}} \left(u_j \frac{\partial}{\partial u_j} \right)^k \right) (V \cdot f).$$

It is straightforward to check the following lemma

Lemma 4.2.

$$\mathcal{D}_k = \sum_{i=1}^n \mathcal{D}_{i,k}.$$

By the Schur branching formula (3.4), Proposition A.1 and Corollary A.2, we have for each $1 \leq h \leq n$

$$\begin{aligned} \mathcal{D}_{h,k} \mathcal{S}_{\rho^t, \mathbf{x}^{(L, -, > t)}}(\mathbf{u}^{(L, -, > t)}) &= \mathcal{D}_{h,k} \left[\frac{s_{\lambda^{(t)}}(\mathbf{u}^{(L, -, > t)}, \mathbf{x}^{(L, -, \leq t)})}{s_{\lambda^{(t)}}(\mathbf{x}^{(L, -)})} \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}} \right] \\ &= g(\bar{\sigma}_0) + \sum_{\bar{\sigma} \in \left[\Sigma_{N_{L,-}^{(\epsilon)}} / \Sigma_{N_{L,-}^X} \right]^r, \bar{\sigma} \neq \bar{\sigma}_0} g(\bar{\sigma}) \end{aligned}$$

where for each $\bar{\sigma} \in \left[\Sigma_{N_{L,-}^{(\epsilon)}} / \Sigma_{N_{L,-}^X} \right]^r$, we have

$$\begin{aligned} g(\bar{\sigma}) &:= \frac{1}{s_{\lambda^{(t)}}(\mathbf{x}^{(L, -)})} \mathcal{D}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -} \frac{\xi_{ij}}{z_{ij}} \right) \right. \\ &\quad \left. \left[\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}} \frac{1}{w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon}} \right] \left((-1)^{f(\bar{\sigma})} \left(\prod_{i=1}^n s_{\phi^{(i, \sigma)}}(\mathbf{u}^{(L, -, > t, i)}, \mathbf{x}^{(L, -, \leq t, i)}) \right) \right) \right] \end{aligned}$$

and

$$(-1)^{f(\bar{\sigma})} = \frac{\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma(i), \epsilon} \neq x_{\sigma(j), \epsilon}} (w_{\sigma(i), \epsilon} - w_{\sigma(j), \epsilon})}{\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}} (w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon})}.$$

Then we have

$$\begin{aligned} g(\bar{\sigma}) &= \frac{(-1)^{f(\bar{\sigma})} \left(\prod_{i=1..n, i \neq h} s_{\phi^{(i, \sigma)}}(\mathbf{u}^{(L, -, > t, i)}, \mathbf{x}^{(L, -, \leq t, i)}) \right)}{s_{\lambda^{(t)}}(\mathbf{x}^{(L, -)})} \left(\prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -, x_j \notin \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \\ &\quad \times \left[\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}, x_{\sigma_0(i), \epsilon} \notin \mathbf{x}^{(L, -, > t, h)}, x_{\sigma_0(j), \epsilon} \notin \mathbf{x}^{(L, -, > t, h)}} \frac{1}{w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon}} \right] A(\bar{\sigma}) \end{aligned}$$

where

$$\begin{aligned} A(\bar{\sigma}) &:= \mathcal{D}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \left(s_{\phi^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right) \right. \\ &\quad \left. \left[\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}, [x_{\sigma_0(i)} \in \mathbf{x}^{(L, -, > t, h)} \text{ or } x_{\sigma_0(i)} \in \mathbf{x}^{(L, -, > t, h)}]} \frac{1}{w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon}} \right] \right] \end{aligned}$$

Let

$$\underline{u}_j := \frac{u_j}{x_j}; \quad \forall u \in \mathbf{u}^{(L, -)}$$

then \underline{u}_j is in a neighborhood of 1, and we can write

$$\mathcal{D}_{i,k} = \frac{1}{V_i V_{i,*}} \left(\sum_{\underline{u}_j \in \mathbf{u}^{(L,-,<t,i)}} \left(\underline{u}_j \frac{\partial}{\partial \underline{u}_j} \right)^k \right) V_i V_{i,*}$$

Assume $i < j$, $a_i = a_j = L$, $b_i = b_j = -$, $x_{\sigma_0(i),\epsilon} \neq x_{\sigma_0(j),\epsilon}$, [$\sigma_0(i) = d$ or $\sigma_0(j) = d$], $x_d \in \mathbf{x}^{(L,-,>t,h)}$. Under Assumption 2.14 we have

$$(4.3) \quad \left(\frac{\partial}{\partial \underline{u}_d} \right)^g \frac{1}{w_{\sigma_0(i),\epsilon} - w_{\sigma_0(j),\epsilon}} = \left(-\frac{x_{d,\epsilon}}{w_{\sigma_0(i),\epsilon}} \right)^g \frac{1}{w_{\sigma_0(i),\epsilon}} (1 + O(e^{-cN}))$$

Note that under Assumption 2.14, (4.3) is exponentially small when $\sigma_0(j) = d$ compared to when $\sigma_0(i) = d$.

Assume $u_j \notin \mathbf{u}^{(L,-,>t,h)}$, $u_d \in \mathbf{u}^{(L,-,>t,i)}$; $j \neq d$, then

$$(4.4) \quad \frac{1}{u_j - u_d} \left(\underline{u}_d \frac{\partial}{\partial \underline{u}_d} \right)^g (u_j - u_d) = -\frac{x_d}{u_j - u_d}$$

Under Assumption 2.14, (4.4) is exponentially small when $u_j > u_d$ compared to when $u_j < u_d$.

Moreover, assume $i \in [l..t]$, $b_i = +$, $j \in [t+1..r]$, $a_d = L$, $b_d = -$, $x_d \in \mathbf{x}^{(L,-,>t,h)}$, we have

$$\left(\frac{\partial}{\partial \underline{u}_d} \right)^g \frac{\xi_{id}}{z_{id}} =$$

- $\frac{1}{z_{id}} \left(\frac{x_{i,\epsilon} x_{d,\epsilon}}{1 - x_{i,\epsilon} u_{d,\epsilon}} \right)^g \frac{1}{1 - x_{i,\epsilon} u_{d,\epsilon}}$, if $b_i = L$;
- $\frac{x_{i,\epsilon} x_{d,\epsilon}}{z_{id}}$, if $b_i = R$ and $g = 1$;
- 0, if $b_i = R$ and $g > 1$.

In each case above, we have

$$\left(\frac{\partial}{\partial \underline{u}_d} \right)^g \frac{\xi_{id}}{z_{id}} = O(e^{-cN^{(\epsilon)}}); \text{ If } x_{d,\epsilon} \neq x_{\xi_0(1),\epsilon}$$

Therefore we have

$$\begin{aligned}
 A(\bar{\sigma}) &= \left(1 + O(e^{-cN^{(\epsilon)}})\right) \mathcal{D}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \left(s_{\phi^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right) \right. \\
 &\quad \left. \left[\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i)}, \epsilon \neq x_{\sigma_0(j)}, \epsilon, [x_{\sigma_0(i)} \in \mathbf{x}^{(L, -, > t, h)} \text{ or } x_{\sigma_0(i)} \in \mathbf{x}^{(L, -, > t, h)}]} \frac{1}{w_{\sigma_0(i), \epsilon}} \right] \right] \\
 &= \left(1 + O(e^{-cN^{(\epsilon)}})\right) \mathcal{D}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \left(s_{\lambda^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right) \right] \\
 &= \frac{\left(1 + O(e^{-cN^{(\epsilon)}})\right)}{V_{h,*}} \bar{\mathcal{D}}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \right. \\
 &\quad \left. \left(s_{\lambda^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right) V_{h,*} \right] \\
 &= \frac{\left(1 + O(e^{-cN^{(\epsilon)}})\right)}{Q_h} \bar{\mathcal{D}}_{h,k} \left[\left(\prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L, -, > t, h)}} \frac{\xi_{ij}}{z_{ij}} \right) \right. \\
 &\quad \left. \left(s_{\lambda^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right) Q_h \right]
 \end{aligned}$$

where

$$Q_h = \left[\prod_{x_j \in \mathbf{x}^{(L, -, > t, h)}} \underline{u}_j \right]^{\sum_{i=h+1}^n N_{L, -, i}^{(> t, \epsilon)}}$$

Moreover, if $h > 1$,

$$A(\bar{\sigma}) = \frac{\left(1 + O(e^{-cN^{(\epsilon)}})\right)}{Q_h} \bar{\mathcal{D}}_{h,k} \left(Q_h s_{\lambda^{(h, \sigma)}}(\mathbf{u}^{(L, -, > t, h)}, \mathbf{x}^{(L, -, \leq t, h)}) \right)$$

When $h = 1$, we construct a rail-yard graph as in Lemma 3.3(2) with left boundary condition given by

$$\tilde{\lambda}^{(1, \sigma)} = (\lambda^{(1, \sigma)}, 0, \dots, 0), \text{ s.t. } l(\tilde{\lambda}^{(1, \sigma)}) = N_{L, -}^{(\epsilon)}$$

Let $\tilde{\rho}_{\lambda^{(1, \sigma)}}^t$ be the probability measure for the random partition corresponding to dimer configurations incident to odd vertices with abscissa $2t - 1$.

When $h > 1$, we construct a rail-yard graph as in Lemma 3.3(3) with left boundary condition given by

$$\tilde{\lambda}^{(h, \sigma)} = (\lambda^{(h, \sigma)}, 0, \dots, 0), \text{ s.t. } l(\tilde{\lambda}^{(h, \sigma)}) = N_{L, -}^{(\epsilon)}$$

Let $\tilde{\rho}_{\lambda^{(h, \sigma_0)}}^t$ be the probability measure for the random partition corresponding to dimer configurations incident to the column of odd vertices with abscissa $2t - 1$.

Following the same arguments as in Lemma 3.3 with σ_0 replaced by σ , we obtain for $h = 1$

$$\begin{aligned}
A(\bar{\sigma}) &= (1 + O(e^{-cN^{(\epsilon)}})) s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)}) \\
&\times \mathcal{D}_{1,k} \left[\left(\prod_{i \in [l..t], b_i = +} \prod_{j \in [t+1..r], a_j = L, b_j = -, x_j \in \mathbf{x}^{(L,-,>t,1)}} \frac{\xi_{ij}}{z_{ij}} \right) \frac{s_{\lambda(1,\sigma)}(\mathbf{u}^{(L,-,>t,1)}, \mathbf{x}^{(L,-,\leq t,1)})}{s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)})} \right] \\
&= (1 + O(e^{-cN^{(\epsilon)}})) s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)}) \mathcal{D}_{1,k} \left[\sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda(1,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})} \right] \\
&= (1 + O(e^{-cN^{(\epsilon)}})) s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)}) \mathcal{D}_{1,k} \left[\sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda(1,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})} \right] \\
&= \frac{(1 + O(e^{-cN^{(\epsilon)}}))}{Q_h} s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)}) \bar{\mathcal{D}}_{1,k} \left[Q_h \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon)}}} \tilde{\rho}_{\lambda(1,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)}, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})}{s_{\mu}(1, \dots, 1, 0^{N_{L,-}^{(>t,\epsilon)}} - N_{L,-,1}^{(>t,\epsilon)})} \right]
\end{aligned}$$

By Lemma 3.2, we obtain when $h = 1$

$$A(\bar{\sigma}) = \frac{(1 + O(e^{-cN^{(\epsilon)}}))}{Q_1} s_{\lambda(1,\sigma)}(\mathbf{x}^{(L,-,1)}) \bar{\mathcal{D}}_{1,k} \left[Q_1 \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon,1)}}} \tilde{\rho}_{\lambda(1,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)})}{s_{\mu}(1, \dots, 1)} \right]$$

Similarly by Lemmas 3.2 and 3.3, we obtain that for $h > 1$

$$A(\bar{\sigma}) = \frac{(1 + O(e^{-cN^{(\epsilon)}}))}{Q_h} s_{\lambda(h,\sigma)}(\mathbf{x}^{(L,-,h)}) \bar{\mathcal{D}}_{h,k} \left[Q_h \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon,h)}}} \tilde{\rho}_{\lambda(h,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(1)})}{s_{\mu}(1, \dots, 1)} \right]$$

Note that for all $1 \leq h \leq n$, we have

$$\begin{aligned}
&\left| \bar{\mathcal{D}}_{h,k} \left[Q_h \sum_{\mu \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon,h)}}} \tilde{\rho}_{\lambda(h,\sigma)}^t(\mu) \frac{s_{\mu}(\underline{u}^{(h)})}{s_{\mu}(1, \dots, 1)} \right] \right|_{\underline{u}^{(h)} = (1, \dots, 1)} \\
&= \mathbb{E} \tilde{\rho}_{\lambda(h,\sigma)}^t \left[\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,j)} - i \right)^k \right]
\end{aligned}$$

Hence we obtain when $1 \leq h \leq n$,

$$\begin{aligned}
 g(\bar{\sigma})|_{\underline{u}^{(h)}=(1,\dots,1)} &= \frac{(-1)^{f(\bar{\sigma})} \left(\prod_{i=[1..n], i \neq h} \mathcal{S}_{\phi^{(t,\sigma)}}(\mathbf{u}^{(L,-,>t,i)}, \mathbf{x}^{(L,-,\leq t,i)}) \right)}{Q_h \mathcal{S}_{\lambda^{(t)}}(\mathbf{x}^{(L,-)})} \\
 &\times \left(\prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -, x_j \notin \mathbf{x}^{(L,-,>t,h)}} \frac{\xi_{ij}}{z_{ij}} \right) \\
 &\times \left[\prod_{i < j, a_i = a_j = L, b_i = b_j = -, x_{\sigma_0(i), \epsilon} \neq x_{\sigma_0(j), \epsilon}, x_{\sigma_0(i), \epsilon} \notin \mathbf{x}^{(L,-,>t,h)}, x_{\sigma_0(j), \epsilon} \notin \mathbf{x}^{(L,-,>t,h)}} \frac{1}{w_{\sigma_0(i), \epsilon} - w_{\sigma_0(j), \epsilon}} \right] \\
 &\times (1 + O(e^{-cN^{(\epsilon)}})) \mathcal{S}_{\lambda^{(h,\sigma)}}(\mathbf{x}^{(L,-,h)}) \mathbb{E}_{\bar{\rho}_{\lambda^{(h,\sigma)}}^t} \left[\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,j)} - i \right)^k \right]
 \end{aligned}$$

By Lemmas 3.6 and 3.4, we have

$$\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,j)} - i \right)^k = O(|N^{(\epsilon)}|^{2k+1}).$$

Together with Proposition 3.1, one sees that for any $\bar{\sigma} \neq \bar{\sigma}_0$, we have $|g(\bar{\sigma})|$ is exponentially small compared to $|g(\bar{\sigma}_0)|$. Hence we have

$$\begin{aligned}
 \mathcal{D}_{h,k} \mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}) \Big|_{\underline{u}=(1,\dots,1)} &= (1 + O(e^{-cN^{(\epsilon)}})) g(\bar{\sigma}_0) \\
 &= (1 + O(e^{-cN^{(\epsilon)}})) \mathbb{E}_{\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t} \left[\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,j)} - i \right)^k \right]
 \end{aligned}$$

Similarly, one can show that

$$\mathcal{D}_{h,k}^l \mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}) \Big|_{\underline{u}=(1,\dots,1)} = (1 + O(e^{-cN^{(\epsilon)}})) \mathbb{E}_{\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t} \left[\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,h)} - i \right)^k \right]^l$$

and for h_1, h_2, \dots, h_g distinct with $g < n$, we have

$$\begin{aligned}
 (4.5) \quad &\left[\prod_{j=1}^g \mathcal{D}_{h_j, k_j}^{l_j} \right] \mathcal{S}_{\rho^t, \mathbf{x}^{(L,-,>t)}}(\mathbf{u}^{(L,-,>t)}) \Big|_{\underline{u}=(1,\dots,1)} \\
 &= (1 + O(e^{-cN^{(\epsilon)}})) \prod_{j=1}^g \mathbb{E}_{\bar{\rho}_{\lambda^{(h_j, \sigma_0)}}^t} \left[\sum_{i=1}^{N_{L,-}^{(>t,\epsilon,h_j)}} \left(\mu_i + \sum_{j=h}^n N_{L,-}^{(>t,\epsilon,h_j)} - i \right)^{k_j} \right]^{l_j}
 \end{aligned}$$

5. INDEPENDENCE

In this section, we prove that different parts of random partitions are asymptotically independent.

It is straightforward to check the following

$$\mathcal{D}_k s_\lambda(u_1, \dots, u_N) = \left[\sum_{i=1}^N (\lambda_i + N - i)^k \right] s_\lambda(u_1, \dots, u_N)$$

Recall the Schur generating function defined as in 2.10. We have

$$\left[[\mathcal{D}_k]^l \mathcal{S}_{\rho, \mathbf{x}}(u_1, \dots, u_N) \right]_{(u_1, \dots, u_N) = \mathbf{x}} = \mathbb{E}_\rho \left[\sum_{i=1}^N (\lambda_i + N - i)^k \right]^l$$

Applying the formula to $\mathcal{S}_{\rho^t, \mathbf{x}^{(L, -, > t)}}(\mathbf{u}^{(L, -, > t)})$, where ρ^t is the distribution of partitions corresponding to the dimer configurations adjacent to the odd vertices in column $(2t - 1)$, we obtain

$$(5.1) \quad \mathbb{E}_{\rho^t} \prod_{j=1}^r \left[\sum_{i=1}^{N_{L, -}^{(> t, \epsilon)}} [\lambda_i + N_{L, -}^{(> t, \epsilon)} - i]^{k_j} \right]^{l_j} = \left(\prod_{j=1}^r [\mathcal{D}_{k_j}]^{l_j} \right) \mathcal{S}_{\rho^t, \mathbf{x}^{(L, -, > t)}}(\mathbf{u}^{(L, -, > t)}) \Big|_{\mathbf{u}^{(L, -, > t)} = \mathbf{x}^{(L, -, > t)}}$$

where $N_{L, -}^{(> t, \epsilon)}$ is given as in Lemma 3.3(1). Recall that n is the total number of distinct elements in $\mathbf{x}^{L, -}$. For $1 \leq s \leq n$, let

$$I_s^{(> t)} = \sum_{d=1}^s N_{L, -, d}^{(> t, \epsilon)}$$

and define

$$A_{s, k}^{(> t)} = \sum_{i=I_{s-1}^{(> t)} + 1}^{I_s^{(> t)}} [\lambda_i + N_{L, -}^{(> t, \epsilon)} - i]^k$$

Then we obtain

$$(5.2) \quad \mathbb{E}_{\rho^t} \prod_{j=1}^r \left[\sum_{i=1}^{N_{L, -}^{(> t, \epsilon)}} [\lambda_i + N_{L, -}^{(> t, \epsilon)} - i]^{k_j} \right]^{l_j} = \mathbb{E}_{\rho^t} \prod_{j=1}^r \left[\sum_{s=1}^n A_{s, k_j}^{(> t)} \right]^{l_j}$$

It is straightforward to verify the following lemma:

Proposition 5.1. *Consider perfect matchings on a rail-yard graph $RYG(l, r, \underline{a}, \underline{b})$ satisfying Assumptions 2.12, 2.13, 2.14. Let*

$$\left(\lambda_1, \dots, \lambda_{N_{L, -}^{(> t, \epsilon)}} \right)$$

be the random partition corresponding to dimer configurations adjacent to a column of odd vertices with abscissa $2t - 1$.

Conditional on an event with probability at least $1 - e^{-cN^\epsilon}$.

(1) The random partition

$$((\lambda_i)_{i=1}^{I_1^{(>t)}}, 0, \dots, 0) \in \mathbb{Y}_{L,-}^{N^{(>t),\epsilon}}$$

has the distribution $\tilde{\rho}_{\lambda(1,\sigma_0)}^t$, where $\tilde{\rho}_{\lambda(1,\sigma_0)}^t$ is given as in Lemma 3.3(2).

(2) The random partition

$$((\lambda_i)_{i=I_{s-1}^{(>t)}+1}^{I_s^{(>t)}}, 0, \dots, 0) \in \mathbb{Y}_{L,-}^{N^{(>t),\epsilon}}$$

has distribution $\tilde{\rho}_{\lambda(s,\sigma_0)}^t$, where $\tilde{\rho}_{\lambda(s,\sigma_0)}^t$ is given as in Lemma 3.3(3).

Proof. The lemma follows from Corollary A.2 and Proposition 3.1 and Lemma 3.3. \square

Proposition 5.2. *The random partitions*

$$(\lambda_i)_{i=1}^{I_1^{(>t)}}, (\lambda_i)_{i=I_1^{(>t)}+1}^{I_2^{(>t)}}, \dots, (\lambda_i)_{i=I_{n-1}^{(>t)}+1}^{I_n^{(>t)}}$$

are asymptotically independent in the sense that if

$$\prod_{j=1}^r \left[\sum_{s=1}^n A_{s,k_j}^{(>t)} \right]^{l_j} = \sum_i \prod_{s=1}^n f_i(A_{s,k_1}^{(>t)}, A_{s,k_2}^{(>t)}, \dots, A_{s,k_n}^{(>t)})$$

Then

$$\mathbb{E} \prod_{j=1}^r \left[\sum_{s=1}^n A_{s,k_j}^{(>t)} \right]^{l_j} = \left(1 + O(e^{-cN^\epsilon}) \right) \left[\sum_i \prod_{s=1}^n \mathbb{E} f_i(A_{s,k_1}^{(>t)}, A_{s,k_2}^{(>t)}, \dots, A_{s,k_n}^{(>t)}) \right]$$

Proof. The lemma follows from (5.1), (5.2), Lemma (4.2), (4.5) and Lemma 5.1. \square

6. CONVERGENCE TO THE SPECTRA OF GUE MATRICES

The main aim of this section is to prove the following proposition:

Proposition 6.1. *Consider perfect matchings on a rail-yard graph $RYG(l, r, \underline{a}, \underline{b})$ satisfying Assumptions 2.12, 2.13, 2.14. Let*

$$\left(\lambda_1, \dots, \lambda_{N_{L,-}^{(>t),\epsilon}} \right)$$

be the random partition corresponding to dimer configurations adjacent to a column of odd vertices with abscissa $2t - 1$.

Assume t is fixed and is finite. Assume $1 \leq h \leq n$. For random partition $(\lambda_i)_{i=I_{h-1}^{(>t)}+1}^{I_h^{(>t)}}$, and $1 \leq i \leq N_{L,-}^{(\epsilon,h)}$, define

$$b_{N_{L,-}^{(\epsilon,h)},i} = \frac{\lambda_{i+I_{h-1}^{(>t)}} + N_{L,-}^{(\epsilon,h)} - i}{\sqrt{N_{L,-}^{(\epsilon,h)}}} - \sqrt{N_{L,-}^{(\epsilon,h)}} A_h$$

where A_h, B_h are given by (6.6) (6.7) (6.8), (6.9). Then the distribution of $\left\{ b_{N_{L,-}^{(\epsilon,1)},i} \right\}_{i=1}^{N_{L,-}^{(\epsilon,h)}}$ converges to $\text{GUE}_{N_{L,-}^{(\epsilon,h)}}$

Given Proposition 5.1, it suffices to prove the corresponding result for random partitions with distributions $\tilde{\rho}_{\lambda^{(h,\sigma_0)}}^t$.

It is straightforward to verify the following lemma:

Lemma 6.2. *Let $\{J_i\}_{i \in [n]}$ be defined as in Assumption 2.13. Under Assumptions 2.13 2.15, assume that*

$$J_i = \begin{cases} \{d_i, d_i + 1, \dots, d_{i+1} - 1\} & \text{if } 1 \leq i \leq n - 1 \\ \{d_n, d_n + 1, \dots, s\} & \text{if } i = n \end{cases}$$

where d_1, \dots, d_n are positive integers satisfying

$$1 = d_1 < d_2 < \dots < d_n \leq s$$

Let

$$d_{n+1} := s + 1.$$

For $j \in [s]$, let a_j, b_j be given by (2.20). If $i \in [n]$, for $0 \leq k \leq d_{n+1} - d_n - 1$, let

$$(6.1) \quad \beta_{i,k} = \sum_{j=1}^{s-d_i-k} \frac{a_{j+1} - b_j}{\theta_i} + \sum_{r=d_i}^{d_i+k-1} \frac{b_r - a_r}{\theta_i}$$

and let

$$(6.2) \quad \gamma_{i,k} = \beta_{i,k} + \frac{b_{s-d_i-k+1} - a_{s-d_i-k+1}}{\theta_i} = \frac{b_{s-d_i-k+1} - a_1}{\theta_i}$$

Then for $i \in [n]$ the counting measures of $\lambda^{(i,\sigma_0)}$ converge weakly to a limit measure \mathbf{m}_i as $\epsilon \rightarrow 0$. Moreover, if $i \in [n]$, for $0 \leq k \leq d_{i+1} - d_i - 1$, the limit counting measure \mathbf{m}_i is a probability measure on $[\beta_{i,1}, \gamma_{i,d_{i+1}-d_i-1}]$ with density given by

$$\frac{d\mathbf{m}_i}{dy} = \begin{cases} 1, & \text{if } \beta_{i,k} < y < \gamma_{i,k}; \\ 0, & \text{if } \gamma_{i,k} \leq y \leq \beta_{i,k+1}. \end{cases}$$

As we have seen in Lemma 6.2, under Assumption 2.13, as $\epsilon \rightarrow 0$, the counting measures of $\phi^{(i,\sigma_0)}(N_{L,-}^{(\epsilon)})$ converges weakly to a limit measure \mathbf{m}_i .

Let

$$P_k = \text{diag}[v_1, \dots, v_k], \quad Q_k = \text{diag}[b_{k1}, \dots, b_{kk}].$$

By Lemma 2.1, we have for any k :

$$(6.3) \quad \int_{U(k)} \exp[\text{Tr}(P_k U Q_k U^*)] dU = \left[\prod_{1 \leq i < j \leq k} \frac{e^{v_i} - e^{v_j}}{v_i - v_j} \right] \frac{s_{\lambda^k(N)}(e^{v_1}, \dots, e^{v_k})}{s_{\lambda^k(N)}(1, \dots, 1)}$$

For $1 \leq h \leq n$, let $\tilde{\rho}_{\lambda^{(h,\sigma_0)}}^t$ be given as in Lemma 3.3; and let $\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t$ be the restriction of $\tilde{\rho}_{\lambda^{(h,\sigma_0)}}^t$ to the first $N_{L,-}^{(>t,\epsilon,h)} = k$ parts. Then

$$(6.4) \quad \begin{aligned} & \sum_{\lambda \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon,h)}}} \bar{\rho}_{\lambda^{(h,\sigma_0)}}^t \int_{U(k)} \exp\left[\frac{1}{\sqrt{N}} \text{Tr}(P_k U Q_k U^*)\right] dU \\ &= \prod_{1 \leq i < j \leq k} \left[\sqrt{N} \frac{e^{v_i/\sqrt{N}} - e^{v_j/\sqrt{N}}}{v_i - v_j} \right] \times \sum_{\lambda \in \mathbb{Y}_{N_{L,-}^{(>t,\epsilon,h)}}} \bar{\rho}_{\lambda^{(h,\sigma_0)}}^t \frac{s_{\lambda}(e^{v_1/\sqrt{N}}, \dots, e^{v_k/\sqrt{N}})}{s_{\lambda}(1, \dots, 1)} \\ &= \prod_{1 \leq i < j \leq k} \left[\sqrt{N} \frac{e^{v_i/\sqrt{N}} - e^{v_j/\sqrt{N}}}{v_i - v_j} \right] \times \mathcal{S}_{\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t}(e^{v_1/\sqrt{N}}, \dots, e^{v_k/\sqrt{N}}) \end{aligned}$$

note that the denominator of the last fraction being exactly $\mathcal{S}_{\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t}(e^{v_1/\sqrt{N}}, \dots, e^{v_k/\sqrt{N}})$ by Definition 2.10.

We then use Lemma 3.3 to re-express the Schur generating function:

$$\mathcal{S}_{\bar{\rho}_{\lambda^{(1,\sigma_0)}}^t}(\mathbf{u}^{(L,-,>t,1)}) = \frac{s_{\lambda^{(1,\sigma)}}(\underline{u}^{(L,-,>t,1)}, \mathbf{1}^{N_{L,-}^{(\leq t,1)}})}{s_{\lambda^{(1,\sigma)}}(\mathbf{1}^{N_{L,-}^{(\leq t,1)}})} \prod_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_j, \epsilon = x_{\sigma_0(1)}, \epsilon} \prod_{z_{ij}} \frac{\xi_{ij}}{z_{ij}},$$

and for $2 \leq h \leq k$

$$\mathcal{S}_{\bar{\rho}_{\lambda^{(h,\sigma_0)}}^t}(\mathbf{u}^{(L,-,>t,h)}) = \frac{s_{\lambda^{(h,\sigma_0)}}(\underline{u}^{(L,-,>t,h)}, \mathbf{1}^{N_{L,-}^{(\leq t,h)}})}{s_{\lambda^{(h,\sigma_0)}}(\mathbf{1}^{N_{L,-}^{(\leq t,h)}})},$$

This ratio of Schur functions appears also when using again Lemma 2.1 to rewrite the matrix integral over $U(N_{L,-}^{(h,\sigma_0)})$ this time. More precisely, let

$$R_{N_{L,-}^{(\epsilon,h)}} = \text{diag} [v_1, \dots, v_k, 0, \dots, 0],$$

$$Q_{N_{L,-}^{(\epsilon,h)}} = \text{diag} \left[\lambda_1^{(h,\sigma_0)} + N_{L,-}^{(\epsilon,h)} - 1, \lambda_2^{(h,\sigma_0)} + N_{L,-}^{(\epsilon,h)} - 2, \dots, \lambda_{N_{L,-}^{(\epsilon,h)}}^{(h,\sigma_0)} \right].$$

For $1 \leq i \leq k$, let $\underline{u}_i = e^{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}}$. Then,

$$(6.5) \quad \frac{s_{\lambda^{(h,\sigma_0)}}(1, \dots, 1, \underline{u}^{(L,-, > t, h)})}{s_{\lambda^{(h,\sigma_0)}}(1, \dots, 1)} = \int_{U(N_{L,-}^{(\epsilon,h)})} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} \left(R_{N_{L,-}^{(\epsilon,h)}} U Q_{N_{L,-}^{(\epsilon,h)}} U^* \right) \right] dU \times$$

$$\left[\prod_{1 \leq i < j \leq k} \sqrt{N_{L,-}^{(\epsilon,h)}} \frac{e^{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} - e^{\frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,h)}}}}}{v_i - v_j} \right]^{-1} \left[\prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq N_{L,-}^{(\epsilon,h)} - k}} \frac{e^{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} - 1}{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} \right]^{-1}$$

Plugging (6.5) into (6.4), one gets if $h \geq 2$:

$$\log \sum_{\lambda^{(\epsilon,h)} \in \mathbb{Y}_{N_{L,-}^{(\epsilon,h)}}} \bar{\rho}_{N_{L,-}^{(\epsilon,h)}}^t \int_{U(k)} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} (P_k U Q_k U^*) \right] dU$$

$$= \log \int_{U(N_{L,-}^{(h,\sigma_0)})} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(h,\sigma_0)}}} \text{Tr} \left(R_{N_{L,-}^{(h,\sigma_0)}} U Q_{N_{L,-}^{(h,\sigma_0)}} U^* \right) \right] dU - \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq N_{L,-}^{(h,\sigma_0)} - k}} \log \left[\frac{e^{\frac{v_i}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}} - 1}{\frac{v_i}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}} \right]$$

$$- \sum_{1 \leq i < j \leq k} \log \left[\sqrt{N_{L,-}^{(h,\sigma_0)}} \frac{e^{\frac{v_i}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}} - e^{\frac{v_j}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}}}{v_i - v_j} \right].$$

If $h = 1$,

$$\begin{aligned}
 & \log \sum_{\lambda^{(1,\sigma_0)} \in \mathbb{Y}_{N_{L,-}^{(1,\sigma_0)}}} \bar{\rho}_{N_{L,-}^{(1,\sigma_0)}}^t \int_{U^{(k)}} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} (P_k U Q_k U^*) \right] dU \\
 &= \log \int_{U(N_{L,-}^{(\epsilon,h)})} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} \left(R_{N_{L,-}^{(\epsilon,h)}} U Q_{N_{L,-}^{(\epsilon,h)}} U^* \right) \right] dU - \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq N_{L,-}^{(\epsilon,h)} - k}} \log \left[\frac{e^{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} - 1}{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} \right] \\
 &- \sum_{1 \leq i < j \leq k} \log \left[\frac{\sqrt{N_{L,-}^{(\epsilon,h)}} e^{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} - e^{\frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,h)}}}}}{v_i - v_j} \right] + \sum_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_j, \epsilon = x_{\sigma_0(1)}, \epsilon} \sum \log \frac{\xi_{ij}}{z_{ij}}
 \end{aligned}$$

Expand the sums using the two expansions $\log \frac{e^u - e^v}{u - v} = \frac{u+v}{2} + \frac{(u-v)^2}{24} + O(|u|^3 + |v|^3)$, and $\log \left(\frac{1+ye^v}{1+y} \right) = \frac{y}{1+y} v + \frac{y}{2(1+y)^2} v^2 + O(|v|^3)$ as u and v tend to 0, to get that:

$$\begin{aligned}
 & \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq N_{L,-}^{(\epsilon,h)} - k}} \log \left[\frac{e^{v_i/\sqrt{N_{L,-}^{(\epsilon,h)}}} - 1}{\frac{v_i}{\sqrt{N_{L,-}^{(\epsilon,h)}}}} \right] = (N_{L,-}^{(\epsilon,h)} - k) \sum_{1 \leq i \leq k} \left[\frac{v_i}{2\sqrt{N_{L,-}^{(\epsilon,h)}}} + \frac{v_i^2}{24N_{L,-}^{(\epsilon,h)}} \right] + O([N_{L,-}^{(\epsilon,h)}]^{-\frac{1}{2}}), \\
 & \sum_{1 \leq i < j \leq k} \log \left[\frac{\sqrt{N_{L,-}^{(h,\sigma_0)}} e^{\frac{v_i}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}} - e^{\frac{v_j}{\sqrt{N_{L,-}^{(h,\sigma_0)}}}}}{v_i - v_j} \right] = O([N_{L,-}^{(\epsilon,h)}]^{-\frac{1}{2}}).
 \end{aligned}$$

Moreover,

$$\begin{aligned}
 & \sum_{i \in [l..t], b_i = +; j \in [t+1..r], a_j = L, b_j = -; x_j, \epsilon = x_{\sigma_0(1)}, \epsilon} \sum \log \frac{\xi_{ij}}{z_{ij}} \\
 &= \sum_{i \in [l..t], a_i = L; b_i = +; j=1}^k \log \frac{1 - x_i x_{\sigma_0(1)}}{1 - x_i x_{\sigma_0(1)} e^{\frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,1)}}}}} + \sum_{i \in [l..t], a_i = L; b_i = +; j=1}^k \log \frac{1 + x_i x_{\sigma_0(1)} e^{\frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,1)}}}}}{1 + x_i x_{\sigma_0(1)}} \\
 &= \sum_{i \in [l..t], a_i = L; b_i = +; j=1}^k \left[\frac{x_i x_{\sigma_0(1)}}{1 - x_i x_{\sigma_0(1)}} \frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,1)}}} + \frac{x_i x_{\sigma_0(1)}}{2(1 - x_i x_{\sigma_0(1)})^2} \frac{v_j^2}{N_{L,-}^{(\epsilon,1)}} + O([N_{L,-}^{(\epsilon,1)}]^{-\frac{3}{2}}) \right] \\
 &+ \sum_{i \in [l..t], a_i = L; b_i = +; j=1}^k \left[\frac{x_i x_{\sigma_0(1)}}{1 + x_i x_{\sigma_0(1)}} \frac{v_j}{\sqrt{N_{L,-}^{(\epsilon,1)}}} + \frac{x_i x_{\sigma_0(1)}}{2(1 + x_i x_{\sigma_0(1)})^2} \frac{v_j^2}{N_{L,-}^{(\epsilon,1)}} + O([N_{L,-}^{(\epsilon,1)}]^{-\frac{3}{2}}) \right]
 \end{aligned}$$

Then we use Lemma 6.3 below to get the asymptotic behavior of the integral over $U(N_{L,-}^{(\epsilon,h)})$, to obtain that

$$\begin{aligned} & \log \int_{U(N_{L,-}^{(\epsilon,h)})} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} \left(R_{N_{L,-}^{(\epsilon,h)}} U Q_{N_{L,-}^{(\epsilon,h)}} U^* \right) \right] dU \\ &= \psi_{1,h} \left(\sum_{i=1}^k v_i \right) + \frac{\psi_{2,h} - \psi_{1,h}^2}{2} \left(\sum_{i=1}^k v_i^2 \right) + o(1), \end{aligned}$$

where $\psi_{1,h}$ and $\psi_{2,h}$ are respectively the first and second moments of the limiting measure \mathbf{m}_h (see Lemma 6.2), as $N_{L,-}^{(\epsilon,h)}$ goes to infinity. Bringing all the pieces together, and defining

$$(6.6) \quad A_1 = \psi_{1,1} - \frac{1}{2} + \sum_{i \in [l..t], a_i=L; b_i=+} \frac{x_i x_{\sigma_0(1)}}{1 - x_i x_{\sigma_0(1)}} + \sum_{i \in [l..t], a_i=L; b_i=+} \frac{x_i x_{\sigma_0(1)}}{1 + x_i x_{\sigma_0(1)}},$$

(6.7)

$$B_1 = \psi_{2,1} - \psi_{1,1}^2 - \frac{1}{12} + \sum_{i \in [l..t], a_i=L; b_i=+} \frac{x_i x_{\sigma_0(1)}}{[1 + x_i x_{\sigma_0(1)}]^2} + \sum_{i \in [l..t], a_i=L; b_i=+} \frac{x_i x_{\sigma_0(1)}}{[1 - x_i x_{\sigma_0(1)}]^2},$$

and for $h \geq 2$, define

$$(6.8) \quad A_h = \psi_{1,h} - \frac{1}{2},$$

$$(6.9) \quad B_h = \psi_{2,h} - \psi_{1,h}^2,$$

one finally obtain that

$$\int_{U(k)} \exp \left[\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}}} \text{Tr} (P_k U Q_k U^*) \right] dU = \exp \left(\sqrt{N_{L,-}^{(\epsilon,h)}} A_h \sum_i v_i + \frac{1}{2} B_h \sum_i v_i^2 + o(1) \right).$$

Therefore, according to Lemma 2.2, the distribution of the diagonal coefficients of

$$\frac{1}{\sqrt{N_{L,-}^{(\epsilon,h)}} B_h} (Q_k - \sqrt{N_{L,-}^{(\epsilon,h)}} A_h \text{Id}_k)$$

which are exactly the $\tilde{b}_{kl}^{(N)}$, converges to $\mathbb{P}_{\text{GUE}_k}$.

Lemma 6.3. *Let $k \in \mathbb{N}$ be fixed. For any N , let $\omega(N) \in \mathbb{GT}_N^+$ and*

$$\begin{aligned} q_i^{(N)} &= \omega_i(N) + N - i, \quad Q_N = \text{diag} [q_1^{(N)}, \dots, q_N^{(N)}], \\ P_N &= \text{diag} [v_1, \dots, v_k, 0, \dots, 0]. \end{aligned}$$

Assume that

- $(\omega(N))_{N \in \mathbb{N}}$ is a regular sequence of signatures,
- as $N \rightarrow \infty$, the counting measures $\mathbf{m}_{\omega(N)}$ converge weakly to \mathbf{m}_ω ,

Then as $N \rightarrow \infty$, we have the following asymptotic expansion

$$(6.10) \quad \log \int_{U(N)} \exp \left[\frac{1}{\sqrt{N}} \text{Tr} (P_N U Q_N U^*) \right] dU = K_1 p_1(v_1, \dots, v_k) N^{1/2} + \frac{K_2}{2!} p_2(v_1, \dots, v_k) + o(1)$$

where K_d is $(d-1)!$ multiplying the d th free cumulant of the measure \mathbf{m}_ω (in particular we have $K_1 = \psi_1, K_2 = \psi_2 - \psi_1^2$) and the p_d is the d th power sum:

$$p_d(v_1, \dots, v_k) = \sum_{i=1}^k v_i^d.$$

Proof. See [37, Corollary 6]; see also [14]. □

APPENDIX A. A FORMULA TO COMPUTE SCHUR FUNCTIONS

Let $\lambda(N) \in \mathbb{Y}_N$. Let Σ_N be the permutation group of N elements and let $\sigma \in \Sigma_N$. Let

$$X = (x_1, \dots, x_N).$$

Assume that there exists a positive integer $n \in [N]$ such that x_1, \dots, x_n are pairwise distinct and $\{x_1, \dots, x_n\} = \{x_1, \dots, x_N\}$. For $j \in [N]$, let

$$(A.1) \quad \eta_j^\sigma(N) = |\{k : k > j, x_{\sigma(k)} \neq x_{\sigma(j)}\}|.$$

For $1 \leq i \leq n$, let

$$(A.2) \quad \Phi^{(i,\sigma)}(N) = \{\lambda_j(N) + \eta_j^\sigma(N) : x_{\sigma(j)} = x_i\}$$

and let $\phi^{(i,\sigma)}(N)$ be the partition with length $|\{1 \leq j \leq N : x_j = x_i\}|$ obtained by decreasingly ordering all the elements in $\Phi^{(i,\sigma)}(N)$. Let Σ_N^X be the subgroup Σ_N that preserves the value of X ; more precisely

$$\Sigma_N^X = \{\sigma \in \Sigma_N : x_{\sigma(i)} = x_i, \text{ for } i \in [N]\}.$$

Let $[\Sigma/\Sigma_N^X]^r$ be the collection of all the right cosets of Σ_N^X in Σ_N . More precisely,

$$[\Sigma/\Sigma_N^X]^r = \{\Sigma_N^X \sigma : \sigma \in \Sigma_N\},$$

where for each $\sigma \in \Sigma_N$

$$\Sigma_N^X \sigma = \{\xi \sigma : \xi \in \Sigma_N^X\}$$

and $\xi \sigma \in \Sigma_N$ is defined by

$$\xi \sigma(k) = \xi(\sigma(k)), \text{ for } k \in [N].$$

Let $k \in [N]$. Let

$$(A.3) \quad w_i = \begin{cases} u_i & \text{if } 1 \leq i \leq k \\ x_i & \text{if } k+1 \leq i \leq N \end{cases}$$

Proposition A.1. Let $\{w_i\}_{i \in [N]}$ be given by (A.3). Define

$$\underline{u}^{(i)} = \left\{ \begin{array}{l} u_j : x_j = x_i \\ x_i \end{array} \right\}$$

Then we have the following formula

$$(A.4) \quad \begin{aligned} & s_\lambda(w_1, \dots, w_N) \\ &= \sum_{\bar{\sigma} \in [\Sigma_N / \Sigma_N^X]^r} \left(\prod_{i=1}^n x_i^{|\phi^{(i, \sigma)}(N)|} \right) \left(\prod_{i=1}^n s_{\phi^{(i, \sigma)}(N)}(\underline{u}^{(i)}, 1, \dots, 1) \right) \\ & \quad \times \left(\prod_{i < j, x_{\sigma(i)} \neq x_{\sigma(j)}} \frac{1}{w_{\sigma(i)} - w_{\sigma(j)}} \right) \end{aligned}$$

where $\sigma \in \bar{\sigma} \cap \Sigma_N$ is a representative.

Proof. See Proposition 4.1 of [32]. □

Corollary A.2.

$$(A.5) \quad \begin{aligned} & s_\lambda(x_1, \dots, x_N) \\ &= \sum_{\bar{\sigma} \in [\Sigma_N / \Sigma_N^X]^r} \left(\prod_{i=1}^n x_i^{|\phi^{(i, \sigma)}(N)|} \right) \left(\prod_{i=1}^n s_{\phi^{(i, \sigma)}(N)}(1, \dots, 1) \right) \left(\prod_{i < j, x_{\sigma(i)} \neq x_{\sigma(j)}} \frac{1}{x_{\sigma(i)} - x_{\sigma(j)}} \right) \end{aligned}$$

where $\sigma \in \bar{\sigma} \cap \Sigma_N$ is a representative.

Proof. The corollary follows from Proposition A.1 by letting $u_i = x_i$ for all $i \in [k]$. □

Acknowledgements. ZL acknowledges support from National Science Foundation DMS 1608896 and Simons Foundation grant 638143.

REFERENCES

- [1] M. Adler and P. van Moerbeke. Coupled GUE minor processes and domino tilings. *Int. Math. Res. Not. IMRN*, 21:10987–11044, 2015.
- [2] M. Basok and D. Chelkak. Double dimers, conformal loop ensembles and isomonodromic deformations. *J. Eur. Math. Soc.*, 23:2787–2832, 2019.
- [3] T. Berggren and A. Borodin. Geometry of the doubly periodic aztec dimer model. 2023.
- [4] D. Betea, C. Boutillier, J. Bouttier, G. Chapuy, S. Corteel, and M. Vuletic. Perfect sampling algorithm for schur processes. *Markov Processes and Related Fields, proceedings of Inhomogenous Random Systems 2015*, 3:381–418, 2018.
- [5] A. Borodin and M. Duits. Biased 2×2 periodic aztec diamond and an elliptic curve. *Probab. Theory Related Fields*, 187:259–315, 2023.
- [6] C. Boutillier and Z. Li. Limit shape and height fluctuations of random perfect matchings on square-hexagon lattices. *Annales de Institut Fourier*. [arXiv:1709.09801](https://arxiv.org/abs/1709.09801).
- [7] Cédric Boutillier, Jérémie Bouttier, Guillaume Chapuy, Sylvie Corteel, and Sanjay Ramassamy. Dimers on rail yard graphs. *Ann. Inst. Henri Poincaré D, Comb. Phys. Interact. (AIHPD)*, 4(4):479–539, 2017.

- [8] H. Cohn, R. Kenyon, and J. Propp. A variational principle for domino tilings. *J. Amer. Math. Soc.*, 14:297–346, 2001.
- [9] J. Dubedat. Dimers and families of cauchy-riemann operators i. *Journal of the American Mathematical Society*, 28:1063–1167, 2015.
- [10] J. Dubedat. Double dimers, conformal loop ensembles and isomonodromic deformations. *J. Eur. Math. Soc.*, 21:1–54, 2019.
- [11] V. Gorin. *Lectures on random lozenge tilings*. Cambridge University Press, 2021.
- [12] V. Gorin and A. Aggarwal. *Probability Theory and Related Fields*, 184:1139–1166, 2022.
- [13] V. Gorin and G. Panova. Asymptotics of symmetric polynomials with applications to statistical mechanics and representation theory. *Ann. Probab.*, 43:3052–3132, 2015.
- [14] I.P. Goulden, M. Guay-Paquet, and J. Novak. Monotone Hurwitz numbers and the HCIZ integral. *Annales mathématiques Blaise Pascal*, 21:71–89, 2014.
- [15] Harish-Chandra. Differential operators on a semisimple Lie algebra. *Amer. J. Math.*, 79:87–120, 1957.
- [16] C. Itzykson and J. B. Zuber. The planar approximation. II. *J. Math. Phys.*, 21:411–421, 1980.
- [17] K. Johansson and E. Nordenstam. Eigenvalues of gue minors. *Electron. J. Probab.*, 11:1342–1371, 2006.
- [18] R. Kenyon. Conformal invariance of domino tiling. *Ann. Probab.*, 28:759–795, 2000.
- [19] R. Kenyon. Dominos and the Gaussian free field. *Ann. Probab.*, 29:1128–1137, 2001.
- [20] R. Kenyon. The dimer model. In *Exact methods in low-dimensional statistical physics and quantum computing*, pages 341–361. Oxford University Press, 2010.
- [21] R. Kenyon and A. Okounkov. Limit shapes and the complex Burgers equation. *Acta Mathematica*, 199:263–302, 2007.
- [22] R. Kenyon, A. Okounkov, and S. Sheffield. Dimers and amoebae. *Ann. Math.*, 163:1019–1056, 2006.
- [23] R. Kenyon and I. Prause. Gradient variational problems in \mathbb{R}^2 . 2020. [arXiv:2006.01291](https://arxiv.org/abs/2006.01291).
- [24] Z. Li. Conformal invariance of dimer heights on isoradial double graphs. *Ann. Inst. H. Poincaré Sect. D.*, 4:273–307, 2017.
- [25] Z. Li. Asymptotics of schur functions on almost staircase partitions. *Electronic Communications in Probability*, 25:13pp, 2020.
- [26] Z. Li. Asymptotics of schur functions on almost staircase partitions. *Electronic Communications in Probability*, 25:13pp, 2020.
- [27] Z. Li. Limit shape of perfect matchings on contracting bipartite graphs. *International Mathematics Research Notices*, 2021. [arXiv:2011.10161](https://arxiv.org/abs/2011.10161).
- [28] Z. Li. Schur function at generic points and limit shape of perfect matchings on contracting square hexagon lattices with piecewise boundary conditions. *International Mathematics Research Notices*, 2021. [arXiv:1807.06175](https://arxiv.org/abs/1807.06175).
- [29] Z. Li. Asymptotics of dimer coverings on free boundary rail-yard graphs. 2023. <https://arxiv.org/abs/2304.00650>.
- [30] Z. Li. Asymptotics of dimer coverings on free boundary rail-yard graphs. 2023. <https://arxiv.org/pdf/2304.00650>.
- [31] Z. Li. Limit shape of perfect matchings on rail-yard graphs. *International Mathematics Research Notices*, 2023:12231–12297, 2023.
- [32] Z. Li. Limit shape of perfect matchings on rail-yard graphs. *International Mathematics Research Notices*, 2023. <https://arxiv.org/pdf/2110.11351>.
- [33] Z. Li and M. Vuletić. Asymptotics of pure dimer coverings on rail-yardgraphs. 2021. [arXiv:2110.11393](https://arxiv.org/abs/2110.11393).
- [34] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford University Press, 1998.
- [35] M.L. Mehta. *Random Matrices*. Academic Press, 2004.
- [36] H. Narayanan, S. Sheffield, and T. Tao. Sums of gue matrices and concentration of hives from correlation decay of eigengaps. 2023. <https://arxiv.org/abs/2306.11514>.
- [37] J. Novak. Lozenge tilings and Hurwitz numbers. *J. Stat. Phys.*, 161:509–517, 2015.
- [38] A. Okounkov. Infinite wedge and random partitions. *Selecta Math.*, 7:57–81, 2001.

- [39] Andrei Okounkov and Nicolai Reshetikhin. The birth of a random matrix. *Mosc. Math. J.*, 6(3):553–566, 2006.
- [40] B. Young. Generating functions for coloured 3d young diagrams and the Donaldson-Thomas invariants of orbifolds. *Duke Math J.*, 152:115–153, 2010.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT 06269-3009,
USA

Email address: `zhongyang.li@uconn.edu`

URL: <https://mathzhongyangli.wordpress.com>