

UNIFORM DISPERSION IN GROWTH MODELS ON HOMOGENEOUS TREES

VALDIVINO V. JUNIOR, FÁBIO P. MACHADO, AND ALEJANDRO ROLDÁN-CORREA

ABSTRACT. We consider the dynamics of a population spatially structured in colonies that are vulnerable to catastrophic events occurring at random times, which randomly reduce their population size and compel survivors to disperse to neighboring areas. The dispersion behavior of survivors is critically significant for the survival of the entire species. In this paper, we consider an uniform dispersion scheme, where all possible survivor groupings are equally probable. The aim of the survivors is to establish new colonies, with individuals who settle in empty sites potentially initiating a new colony by themselves. However, all other individuals succumb to the catastrophe. We consider the number of dispersal options for surviving individuals in the aftermath of a catastrophe to be a fixed value d within the neighborhood. In this context, we conceptualize the evolution of population dynamics occurring over a homogeneous tree. We investigate the conditions necessary for these populations to survive, presenting pertinent bounds for survival probability, the number of colonized vertices, the extent of dispersion within the population, and the mean time to extinction for the entire population.

1. INTRODUCTION

Some biological populations frequently face catastrophic events, such as epidemics and natural disasters, which can lead to the extinction of the species. Dispersal of individuals during catastrophes serves as a strategy to enhance a species' chances of survival. This strategy increases genetic diversity within separated populations, reduces intraspecific competition for resources, and helps individuals avoid predation or infections. Dispersal also allows for the colonization of new habitats less affected by the catastrophe, ultimately increasing the chance of long-term survival for at least a portion of the population. For further information on dispersal in the biological context, please refer to Ronce [8].

In [2, 3, 4, 5, 6, 9], models have been introduced to examine various dispersal strategies in populations experiencing different types of catastrophes. These models aim to assess the influence of these strategies on population viability, contrasting them with scenarios where no dispersal occurs. Their analysis seeks to determine the best strategy (dispersion or no dispersion) by evaluating survival probability and population extinction times when a specific strategy is employed. In particular, in [3, 4, 5, 6], the authors consider that

Date: July 23, 2024.

2010 Mathematics Subject Classification. 60J80, 60J85, 92D25.

Key words and phrases. Branching processes, Catastrophes, Population dynamics.

Research supported by FAPESP (2023/13453-5 and 2022/08948-2) and Universidad de Antioquia.

when a colony of individuals suffers a catastrophe, the size of this population is reduced according to some probability law (binomial or geometric), and the surviving individuals disperse to neighboring sites to establish new colonies. New colonies can only form on empty sites. Among the individuals that go to the same empty site, only one survives and the others die. Three dispersion schemes were considered, when each colony has d neighboring sites:

- *Optimal Dispersion:* When r individuals survive a catastrophe, there are exactly $\min\{r, d\}$ successful attempts to establish new colonies.
- *Independent Dispersion:* Each survivor picks a neighboring site at random and tries to create a new colony there.
- *Uniform Dispersion:* For every r survivors, consider all sets of numbers $r_1, \dots, r_d \in \mathbb{N}$ (*occupancy set of numbers*) that are a solution to $r_1 + r_2 + \dots + r_d = r$. For each of these sets exactly one of the r_i individuals (if $r_i > 0$, and 0 otherwise) will succeed in colonizing the neighboring vertex i , $i = 1, \dots, d$. Observe that r_i may be 0. Here we consider that all possible survivor groupings are equally probable. So, the probability of having exactly $y \leq \min\{r, d\}$ successful attempts when the number of survivors is r , is

$$\frac{\binom{r-1}{y-1}}{\binom{d+r-1}{r}} \binom{d}{y}.$$

Taking everything into account, the population growth of each colony follows a birth and death process denoted as \mathcal{N} . Each colony is associated with an independent exponential random time, with a mean of 1, indicating the occurrence time of a catastrophe. When a catastrophe occurs, it results in a reduction in the colony's size, following a specific probability distribution, \mathcal{L} . Although the paper is presented in a general context, several results detail specific cases. In particular, we examine the scenario where

- \mathcal{N} follows a Poisson process with rate λ , denoted by $\mathcal{P}(\lambda)$, and
- \mathcal{L} follows a binomial distribution whose parameters are the colony's size and p , denoted by $\text{Bin}(p)$.

Next, consider the number of dispersal options for surviving individuals in the aftermath of a catastrophe to be a fixed value d within the neighborhood. In this context, we conceptualize the evolution of population dynamics occurring over \mathbb{T}^d , a homogeneous tree where every vertex has $d + 1$ nearest neighbors, and over \mathbb{T}_+^d , a tree whose only difference from \mathbb{T}^d is that its origin has degree of d .

Individuals remaining after the catastrophe are distributed among the nearest neighbor vertices, according to *uniform dispersion*. Individuals that go to a vertex already occupied by a colony die. Among the individuals that go to the same (empty) vertex to create a new colony there, only one succeeds, the others die. Therefore, when a catastrophe occurs in a colony, that colony is replaced by 0, 1, \dots or d colonies, each colony is started by a single individual. We assume that initially all vertices of \mathbb{G} are empty, except one called the

origin, which has a colony with a single individual. We denote this model by $C(\mathbb{G}; \mathcal{N}, \mathcal{L})$ where \mathbb{G} is either \mathbb{T}^d or \mathbb{T}_+^d .

This paper is divided into four sections. In Section 2, we introduce results on phase transition, survival probability, the number of colonies created in the model $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. Besides that, we study the reach on the graph and the mean extinction time of the model $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. In Section 3, we prove the results presented in Sections 2. Finally, in Section 4, we contrast *uniform dispersion* with *independent dispersion*, by comparing our model with the model introduced by Machado et al. [5].

2. MAIN RESULTS

In order to establish the main results on $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, we begin by introducing some definitions and notations.

Let N_t be the number of individuals in a colony at time t (after its creation) and T be the time in which the catastrophe occurs in that colony. We denote by N the number of individuals that survive the catastrophe before dispersing. In particular, for $C(\mathbb{G}; \mathcal{P}(\lambda), \text{Bin}(p))$, the distribution of N is given by

$$\mathbb{P}[N = n] = \begin{cases} \frac{1-p}{\lambda p + 1}, & n = 0, \\ \left(\frac{\lambda p}{\lambda p + 1}\right)^n \frac{\lambda + 1}{\lambda(\lambda p + 1)}, & n \geq 1, \end{cases} \quad (2.1)$$

and the probability generating function is

$$\mathbb{E}[s^N] = \frac{1 - p(1-s)}{1 + \lambda p(1-s)}. \quad (2.2)$$

For more details, see Junior *et al* [2, Proof of Lemma 4.3].

Observe that model $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ is a continuous-time stochastic process with state space $\mathbb{N}_0^{\mathcal{V}(\mathbb{T}^d)}$, where $\mathcal{V}(\mathbb{T}^d)$ is the set of vertices of \mathbb{T}^d . The evolution of this process in time is denoted by η_t . For a vertex $x \in \mathcal{V}(\mathbb{T}^d)$, $\eta_t(x)$ is the number of individuals at time t at vertex x . We consider $|\eta_t| := \sum_{x \in \mathcal{V}(\mathbb{T}^d)} \eta_t(x)$, the total number of individuals present in \mathbb{T}^d at time t . Analogously for $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$.

2.1. Phase Transition.

Definition 2.1. We say that the process $(C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ or $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L}))$ *survives* if at every instant of time there is at least one alive colony. For V_d , the survival event,

$$V_d = \{|\eta_t| > 0, \text{ for all } t \geq 0\}.$$

In the following result, we establish sufficient conditions for the probability of survival to be greater than or equal to zero.

Theorem 2.2. Consider $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and let V_d be the survival event. Then $\mathbb{P}(V_d) = 0$ if

$$\mathbb{E} \left[\frac{N}{N+d} \right] \leq \frac{1}{d+1}$$

and $\mathbb{P}(V_d) > 0$ if

$$\mathbb{E} \left[\frac{N}{N+d} \right] > \frac{1}{d}.$$

Corollary 2.3. Consider $C(\mathbb{T}^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and let V_d be the survival event. Then $\mathbb{P}(V_d) = 0$ if

$$\Phi \left(\frac{\lambda p}{\lambda p + 1}, 1, d+1 \right) \geq \frac{(\lambda p + 1)[p(\lambda d + d + 1) - 1]}{pd(d+1)(\lambda + 1)}$$

and $\mathbb{P}(V_d) > 0$ if

$$\Phi \left(\frac{\lambda p}{\lambda p + 1}, 1, d+1 \right) < \frac{(\lambda p + 1)[p(\lambda d + d - \lambda) - 1]}{pd^2(\lambda + 1)},$$

where $\Phi(z, s, a)$ is the Lerch Transcendent Function given by

$$\Phi(z, s, a) = \sum_{j=0}^{\infty} \frac{z^j}{(a+j)^s}, \quad |z| < 1. \quad (2.3)$$

In particular, the Lerch Transcendent Function satisfies

$$\Phi(z, 1, a) = \frac{1}{z^a} \left[\ln \left(\frac{1}{1-z} \right) - \sum_{j=1}^{a-1} \frac{z^j}{j} \right]. \quad (2.4)$$

for every natural number a .

Example 2.4. Consider $C(\mathbb{T}^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and let $p_c(d, \lambda)$ defined by

$$p_c(d, \lambda) = \inf\{p \geq 0; \mathbb{P}(V_d) > 0\}.$$

Then, for $\lambda = 10$ and $d = 30$, by using Corollary 2.3, we have that

$$0.0962 \leq p_c(30, 10) \leq 0.0996.$$

With independent dispersion scheme, using [5, Theorem 3.2] for the analogous parameter, we have

$$0.0936 \leq p_c(30, 10) \leq 0.0969.$$

Due to the intersection of the ranges obtained from the rigorous results currently available, it is not possible to make a definitive comparison between the models with uniform dispersion and independent dispersion.

2.2. Probability of Survival.

Theorem 2.5. Consider $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and let V_d be the survival event. Then

$$\sum_{r=1}^{d+1} \left[(1 - \rho^r) \binom{d+1}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right] \leq \mathbb{P}(V_d) \leq 1 - \psi$$

where ψ and ρ are, respectively, the smallest non-negative solutions of

$$\sum_{y=1}^{d+1} \left[s^y \binom{d+1}{y} \sum_{n=y}^{\infty} \frac{\binom{n-1}{y-1}}{\binom{n+d}{d}} \mathbb{P}(N=n) \right] = s - \mathbb{P}(N=0) \text{ and}$$

$$\sum_{y=0}^d \left[s^y \binom{d}{y} \sum_{n=y}^{\infty} \frac{\binom{n}{y}}{\binom{n+d}{d}} \mathbb{P}(N=n) \right] = s.$$

Example 2.6. Consider $C(\mathbb{T}^{10}; \mathcal{P}(5), \text{Bin}(\frac{3}{5}))$. Then,

$$\mathbb{P}(N=n) = \begin{cases} \frac{1}{10}, & n=0; \\ \frac{3}{10} \left(\frac{3}{4}\right)^n, & n \geq 1. \end{cases}$$

Applying Theorem 2.5 we have $\psi = 0.141484$ and $\rho = 0.162176$. So,

$$\frac{3}{10} \sum_{r=1}^{11} \left[(1 - 0.162176^r) \binom{11}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+10}{10}} \left(\frac{3}{4}\right)^n \right] \leq \mathbb{P}(V_{10}) \leq 1 - 0.141484$$

$$0.85153 \leq \mathbb{P}(V_{10}) \leq 0.858516.$$

Theorem 2.7. Consider $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and let V_d be the survival event. We have that

$$\lim_{d \rightarrow \infty} \mathbb{P}(V_d) = 1 - \nu$$

where ν is the smallest non-negative solution of $\mathbb{E}(s^N) = s$.

Corollary 2.8. Consider $C(\mathbb{T}^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and let V_d be the survival event. Then

$$\lim_{d \rightarrow \infty} \mathbb{P}(V_d) = \max \left\{ 0, \frac{p(\lambda+1) - 1}{\lambda p} \right\}.$$

2.3. The reach of the process. We define M_d as the reach of the process $(C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ or $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L}))$, which, in simpler terms, represents the distance from the origin to the farthest vertex where a colony is formed. Observe M_d is infinite if and only if the process survives.

To study properties of M_d we need the definitions of a few technical quantities.

Definition 2.9. Consider $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ We define the quantities

$$\alpha := d \mathbb{E} \left[\frac{N}{N+d} \right]$$

$$\beta := (d+1) \mathbb{E} \left[\frac{N}{N+d} \right] = \alpha + \mathbb{E} \left[\frac{N}{N+d} \right]$$

$$D := \max \left\{ 2; \frac{\beta}{\beta - \mathbb{P}(N \neq 0)} \right\}$$

$$B := d(d-1) \left[\mathbb{E} \left(\frac{N(N-1)}{(N+d-1)(N+d-2)} \right) \right]$$

Lemma 2.10. For $C(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$, the quantities of the Definition 2.9 are given by

$$\begin{aligned}\alpha &= \frac{dp(\lambda+1)}{(\lambda p+1)^2} \left[\lambda p + 1 - d\Phi \left(\frac{\lambda p}{\lambda p+1}, 1, d+1 \right) \right] \\ \beta &= \frac{(d+1)p(\lambda+1)}{(\lambda p+1)^2} \left[\lambda p + 1 - d\Phi \left(\frac{\lambda p}{\lambda p+1}, 1, d+1 \right) \right] \\ D &= \max \left\{ 2; \frac{\beta(\lambda p+1)}{\beta+p(\beta\lambda-\lambda-1)} \right\} \\ B &= \frac{d\lambda p^2(d-1)(\lambda+1)}{(\lambda p+1)^3} \left[\frac{d^2+d(\lambda p-2)+2}{d} - \frac{(d-1)(d+2\lambda p)}{(\lambda p+1)} \Phi \left(\frac{\lambda p}{\lambda p+1}, 1, d+1 \right) \right]\end{aligned}$$

Theorem 2.11. Consider M_d , the reach of $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. Assume that

$$\mathbb{E} \left[\frac{N}{N+d} \right] < \frac{1}{d+1}.$$

We have that

$$\frac{[1 + D(1 - \beta)][1 - \beta^{n+1}]}{1 + D(1 - \beta) - \beta^{n+1}} \leq \mathbb{P}(M_d \leq n) \leq \frac{[1 + \frac{\alpha(1-\alpha)}{B}](1 - \alpha^{n+1})}{1 + \frac{\alpha(1-\alpha)}{B} - \alpha^{n+1}}$$

where α, β, B and D are given in Definition 2.9.

Theorem 2.12. Let M_d be the reach of $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. When $d \rightarrow \infty$ we have that

$$M_d \xrightarrow{\mathcal{D}} M,$$

where $\mathbb{P}(M \leq m) = g_{m+1}(0)$, $g(s) = \mathbb{E}(s^N)$ and $g_{m+1}(s) = g(g(\cdots g(s)) \cdots)$ ^{$m+1$ times}.

Example 2.13. Let M_d be the reach of $C(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$. We have that $M_d \xrightarrow{\mathcal{D}} M$ where

$$\mathbb{P}(M \leq n) = \frac{1 - [(\lambda+1)p]^{n+1}}{1 - \frac{\lambda p}{1-p} [(\lambda+1)p]^{n+1}} \text{ and } \mathbb{E}(M) = \frac{(1-p-\lambda p)}{\lambda p} \sum_{n=0}^{\infty} \frac{[(\lambda+1)p]^{n+1}}{\frac{1-p}{\lambda p} - [(\lambda+1)p]^{n+1}}.$$

2.4. Number of colonies.

Theorem 2.14. Consider $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and let I_d be the number of colonies created during the process. If

$$\mathbb{E} \left[\frac{N}{N+d} \right] < \frac{1}{d+1}$$

then

$$\frac{1 + \mathbb{E} \left[\frac{N}{N+d} \right]}{1 - d\mathbb{E} \left[\frac{N}{N+d} \right]} \leq \mathbb{E}(I_d) \leq \frac{1}{1 - (d+1)\mathbb{E} \left[\frac{N}{N+d} \right]}$$

In addition, if $\mathbb{E}(N) < 1$ (the subcritical case),

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \frac{1}{1 - \mathbb{E}(N)}.$$

Corollary 2.15. Consider $C(\mathbb{T}^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and let I_d be the number of colonies created during the process. If

$$\Phi\left(\frac{\lambda p}{\lambda p + 1}, 1, d + 1\right) > \frac{(\lambda p + 1)[p(\lambda d + d + 1) - 1]}{pd(d + 1)(\lambda + 1)},$$

then

$$\frac{d + \alpha}{d(1 - \alpha)} \leq \mathbb{E}(I_d) \leq \frac{d}{d - (d + 1)\alpha},$$

where α is given in Lemma 2.10. Besides, if $(\lambda + 1)p < 1$ then

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \frac{1}{1 - (\lambda + 1)p}.$$

Example 2.16. Let I_d be the total number of colonies created in $C(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$ with $\lambda = 9$ and $p = \frac{99}{1000}$. If $d = 800$ we have that

$$74.5761 \leq \mathbb{E}(I_{800}) \leq 82.0181.$$

Besides, $\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = 100$. In the model with scheme independent dispersion, [5, Theorem 3.23], we have

$$81.1729 \leq \mathbb{E}(I_{800}) \leq 90.0900.$$

As in example 2.4, due to the intersection of the ranges obtained from the rigorous results currently available, it is not possible to make a definitive comparison.

2.5. Extinction time.

Definition 2.17. Let be η_t the process $C(\mathbb{G}; \mathcal{N}, \mathcal{L})$. We define the extinction time of the process $C(\mathbb{G}; \mathcal{N}, \mathcal{L})$ by

$$\tau := \inf\{t > 0 : |\eta_t| = 0\}.$$

Theorem 2.18. Consider $C(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and let $\tau_u(d)$ be the extinction time of the process. Assume that

$$\mathbb{E}\left[\frac{N}{N + d}\right] < \frac{1}{d + 1}.$$

Then

$$\int_0^1 \frac{1 - s}{G_L(s) - s} ds \leq \mathbb{E}(\tau_u(d)) \leq \int_0^1 \frac{1 - s}{G_U(s) - s} ds$$

where

$$G_L(s) = \sum_{y=0}^d s^y \left[\sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d}{y} \binom{n}{y}}{\binom{n+d}{d}} \right] \right]$$

and

$$G_U(s) = \mathbb{P}(N = 0) + \sum_{y=1}^{d+1} \left[\binom{d+1}{y} s^y \sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{n-1}{y-1}}{\binom{n+d}{d}} \right] \right].$$

3. PROOFS

To prove our results we define auxiliary processes on the graphs \mathbb{T}^d and \mathbb{T}_+^d , whose understanding will provide bounds for the processes defined in Section 2.

In the first two auxiliary processes, denoted by $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, every time a catastrophe occurs in a colony, according to uniform dispersion, the survival individuals disperse over neighboring vertices that are further from the origin than the vertex where that colony was placed. In other words, individuals do not disperse to sites that have already been colonized. We refer to this process as *Self Avoiding*.

In the last two auxiliary processes, represented by $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, surviving individuals disperse to any of the $d + 1$ neighboring vertices according to the uniform dispersion. However, those who move backward towards the origin die, as this direction is deemed inhospitable or infertile. We refer to this process as *Move Forward or Die*. In both processes, every new colony starts with only one individual.

The main idea behind the proofs is the identification of an underlying branching process related to the models. After doing that we can apply results of the theory of Branching Processes, including the less known and new results presented in [5].

3.1. The Self Avoiding Model.

Proposition 3.1. *Consider $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and let V_d be the survival event. We have that $\mathbb{P}(V_d) > 0$ if and only if*

$$\mathbb{E} \left[\frac{N}{N + d - 1} \right] > \frac{1}{d}.$$

Proof of Proposition 3.1

Next, observe that the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ behaves as a homogeneous branching process. Every vertex $x \in \mathbb{T}^d$ which is colonized produces Y new colonies (whose distribution depends only on \mathcal{N} and \mathcal{L}) on the d neighbor vertices which are located further from the origin than x is. By numbering those d vertices, we can represent the random variable Y as

$$Y = \sum_{i=1}^d \mathcal{I}_i$$

where \mathcal{I}_i is the indicator variable of event the vertex i receives at least one individual.

Thus,

$$\mathbb{E}(Y) = \sum_{i=1}^d \mathbb{P}(\mathcal{I}_i = 1) = d \sum_{n \geq 1} \mathbb{P}(\mathcal{I}_1 = 1 | N = n) \mathbb{P}(N = n).$$

Now, note that

$$\mathbb{P}(\mathcal{I}_1 = 1 | N = n) = \frac{n}{n + d - 1},$$

and so

$$\mathbb{E}(Y) = d \sum_{n \geq 1} \left[\frac{n}{n + d - 1} \mathbb{P}(N = n) \right] = d \mathbb{E} \left(\frac{N}{N + d - 1} \right). \quad (3.1)$$

From the theory of homogeneous branching processes we see that $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ (and also $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$) survives if and only if $\mathbb{E}(Y) > 1$.

Proposition 3.2. *Consider the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. Let V_d be the survival event and I_d the number of colonies created during the process. Then*

$$\mathbb{P}(V_d) = \sum_{r=1}^{d+1} \left[(1 - \psi^r) \binom{d+1}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right]$$

where ψ , the extinction probability for the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, is the smallest non-negative solution of

$$\sum_{y=1}^d \left[s^y \binom{d}{y} \sum_{n=y}^{\infty} \frac{\binom{n-1}{y-1}}{\binom{n+d-1}{d-1}} \mathbb{P}(N = n) \right] = s - \mathbb{P}(N = 0)$$

On the subcritical regime, which means

$$\mathbb{E} \left[\frac{N}{N+d-1} \right] < \frac{1}{d}$$

it holds that

$$\mathbb{E}(I_d) = 1 + \frac{(d+1)\mathbb{E}\left(\frac{N}{N+d}\right)}{1 - d\mathbb{E}\left(\frac{N}{N+d-1}\right)}.$$

Proof of Proposition 3.2

When a colony placed at the origin collapses, Y_R new colonies are formed by the survivors at its neighboring vertices. If a colony located outside the origin collapses, Y new colonies are created on neighboring vertices. Observe that the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ behaves as a non-homogeneous branching process $\{Z_n\}_{n \geq 0}$ with

$$Z_0 = 1, Z_{n+l} = \sum_{i=1}^{Z_n} X_{n,i}, n \geq 0,$$

where $X_{1,1}$ and $X_{n,i}, n > 1$ are distributed as Y_R and Y , respectively. So, we have that

$$\mathbb{P}(V_d) = \sum_{r=0}^{d+1} \mathbb{P}(V_d | Y_R = r) \mathbb{P}(Y_R = r).$$

Given that $Y_R = r$ one have r independent $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ processes living on r independent rooted trees. So, we have that $\mathbb{P}(V_d^C | Y_R = r) = \psi^r, r = 0, 1, 2, \dots, d+1$, where ψ is the smallest non-negative solution of $E(s^Y) = s$.

By using the Total Probability Theorem,

$$\begin{aligned} \mathbb{P}(Y_R = y) &= \sum_{n=r}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d+1}{y} \binom{n-1}{y-1}}{\binom{n+d}{d}} \right] \text{ for } y = 0, 1, 2, \dots, d+1. \\ \mathbb{P}(Y = y) &= \sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d}{y} \binom{n-1}{y-1}}{\binom{n+d-1}{d-1}} \right] \text{ for } y = 0, 1, 2, \dots, d. \end{aligned} \quad (3.2)$$

The probability generating function of Y is

$$G_U(s) = \mathbb{E}(s^Y) = \mathbb{P}(N = 0) + \sum_{y=1}^d \left[\binom{d}{y} s^y \sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{n-1}{y-1}}{\binom{n+d-1}{d-1}} \right] \right] \quad (3.3)$$

Thus,

$$\mathbb{P}(V_d) = \sum_{r=1}^{d+1} \left[(1 - \psi^r) \binom{d+1}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right].$$

As for the second part of the proposition, note that

$$\mathbb{E}(I_d) = \sum_{r=0}^{d+1} \mathbb{E}(I_d | Y_R = r) \mathbb{P}(Y_R = r)$$

Besides, from (3.1) we have

$$\mathbb{E}(I_d | Y_R = r) = r\mu + 1 \text{ where } \mu = \left[1 - d\mathbb{E} \left(\frac{N}{N+d-1} \right) \right]^{-1},$$

see Stirzaker [10, Exercise 2b, p. 280].

Proposition 3.3. *Consider the process $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. Let V_d be the survival event and I_d the number of colonies created during the process. Then*

$$\lim_{d \rightarrow \infty} \mathbb{P}(V_d) = 1 - \nu \quad (3.4)$$

where ν is the smallest non-negative solution of $\mathbb{E}(s^N) = s$. Besides that, if $\mathbb{E}(N) < 1$ (the subcritical case) then

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \frac{1}{1 - \mathbb{E}(N)}. \quad (3.5)$$

Proof of Proposition 3.3. In order to prove (3.4) one has to apply [5, Proposition 4.2], observing that $Y \xrightarrow{\mathcal{D}} N$ and $Y_R \xrightarrow{\mathcal{D}} N$, when $d \rightarrow \infty$. Moreover to prove (3.5) observe that

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \lim_{d \rightarrow \infty} \sum_{r=0}^{d+1} \mathbb{E}(I_d | Y_R = r) \mathbb{P}(Y_R = r).$$

As $Y \xrightarrow{\mathcal{D}} N$ and $Y_R \xrightarrow{\mathcal{D}} N$ when $d \rightarrow \infty$ then

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d | Y_R = r) = \lim_{d \rightarrow \infty} r \frac{1}{1 - \mathbb{E}(Y)} + 1 = \frac{r}{1 - \mathbb{E}(N)} + 1$$

and the result follows from the Dominated Convergence Theorem [11, Theorem 9.1 p. 26]. \square

Proposition 3.4. *Let M_d be the reach of the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, that is, the distance from the origin to the farthest vertex where a colony is formed. Assuming*

$$\mathbb{E} \left[\frac{N}{N + d - 1} \right] < \frac{1}{d}$$

Then

$$\frac{[1 + D(1 - \mu_u)](1 - B_u^{n+1})}{1 + D(1 - \mu_u - \mu_u^{n+1})} \leq \mathbf{P}(M_d \leq n) \leq \frac{\left[1 + \frac{\mu_u(1 - \mu_u)}{B_u}\right](1 - \mu_u^{n+1})}{1 + \frac{\mu_u(1 - \mu_u)}{B_u} - \mu_u^{n+1}}$$

where

$$\begin{aligned} \mu_u &= d \mathbb{E} \left[\frac{N}{N + d - 1} \right] \\ D &= \max \left\{ 2; \frac{\mu_u}{\mu_u - \mathbb{P}(N \neq 0)} \right\} \\ B_u &= d(d - 1) \left[\mathbb{E} \left(\frac{N(N - 1)}{(N + d - 1)(N + d - 2)} \right) \right] \end{aligned}$$

Moreover,

$$M_d \xrightarrow{\mathcal{D}} M,$$

where $\mathbb{P}(M \leq m) = g_{m+1}(0)$, being $g(s) = \mathbb{E}(s^N)$ and $g_{m+1}(s) = g(g(\cdots g(s)) \cdots)$ ^{$m+1$ times}.

Proof of Proposition 3.4

Every vertex $x \in \mathbb{T}^d$ which is colonized produces Y new colonies (whose distribution depends only on \mathcal{N} and \mathcal{L}) on the d neighbor vertices which located are further from the origin than x is. By numbering those d vertices, we can represent the random variable Y as

$$Y = \sum_{i=1}^d \mathcal{I}_i$$

where \mathcal{I}_i is the indicator variable of event the vertex i receives at least one individual.

From (3.1) we obtained $\mathbb{E}(Y)$. Now

$$Y^2 = \left(\sum_{i=1}^d \mathcal{I}_i \right)^2 = \sum_{i=1}^d \mathcal{I}_i^2 + 2 \sum_{1 \leq i < j \leq d} \mathcal{I}_i \mathcal{I}_j$$

$$\mathbb{E}(Y^2) = d\mathbb{E}(\mathcal{I}_1^2) + d(d-1)\mathbb{E}(\mathcal{I}_1 \mathcal{I}_2)$$

On the other hand

$$\mathbb{E}(\mathcal{I}_1^2) = \mathbb{P}(\mathcal{I}_1 = 1) = \sum_n [\mathbb{P}(\mathcal{I}_1 = 1 | N = n) \mathbb{P}(N = n)] = \sum_n \left(\frac{n}{n+d-1} \right) \mathbb{P}(N = n)$$

$$= \mathbb{E} \left(\frac{N}{N+d-1} \right)$$

and

$$\mathbb{E}(\mathcal{I}_1 \mathcal{I}_2) = \mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1) = \sum_n [\mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1 | N = n) \mathbb{P}(N = n)]$$

where

$$\mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1 | N = n) = \frac{n(n-1)}{(n+d-1)(n+d-2)}$$

Then

$$\mathbb{E}(Y^2) = d\mathbb{E} \left(\frac{N}{N+d-1} \right) + d(d-1)\mathbb{E} \left(\frac{N(N-1)}{(N+d-1)(N+d-2)} \right)$$

$$\mathbb{E}(Y(Y-1)) = d(d-1)\mathbb{E} \left(\frac{N(N-1)}{(N+d-1)(N+d-2)} \right)$$

Then the result follows from [1, Theorem 1 p. 331] with $\mu_u = \mathbb{E}(Y)$ and $B_u = \mathbb{E}(Y(Y-1))$.

The convergence $M_d \xrightarrow{\mathcal{D}} M$ follows from the fact that $Y \xrightarrow{\mathcal{D}} N$ when $d \rightarrow \infty$ and from [5, Proposition 4.1].

Proposition 3.5. *Let $\tau_u(d)$ be the extinction time of the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. If*

$$\mathbb{E} \left[\frac{N}{N+d-1} \right] < \frac{1}{d}$$

then

$$\mathbb{E}[\tau_u(d)] = \int_0^1 \frac{1-y}{G_U(y) - y} dy.$$

where $G_U(s)$ is given by (3.3).

Proof. Let Z_t be the number of colonies at time t in the model $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. Observe that Z_t is a continuous-time branching process with $Z_0 = 1$. Each particle (colony) in Z_t survives an exponential time of rate 1 and right before death produces $Y \leq d$ particles (colonies are created right after a catastrophe) with distribution given by (3.2) and probability generating function given by (3.3). If $G'_U(1) \leq 1$, then $\mathbb{P}[\tau_u(d) < \infty] = 1$ and the result following from Narayan [7]. □

3.2. Move Forward or Die Model.

Proposition 3.6. *Consider $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and let V_d be the survival event. We have that $\mathbb{P}(V_d) > 0$ if and only if*

$$\mathbb{E} \left[\frac{N}{N+d} \right] > \frac{1}{d}.$$

Proof of Proposition 3.6

Firstly, note that for fixed \mathcal{N} and \mathcal{L} , both processes $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ either both have positive probability of survival or neither do

Next, observe that the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ behaves as a homogeneous branching process. Every vertex $x \in \mathbb{T}^d$ which is colonized produces Y new colonies (whose distribution depends only on \mathcal{N} and \mathcal{L}) on the d neighbor vertices which are located further from the origin than x is. By numbering those d vertices, we can represent the random variable Y as

$$Y = \sum_{i=1}^d \mathcal{I}_i$$

where \mathcal{I}_i is the indicator variable of event the vertex i receives at least one individual. Thus,

$$\mathbb{E}(Y) = \sum_{i=1}^d \mathbb{P}(\mathcal{I}_i = 1) = d \sum_{n \geq 1} \mathbb{P}(\mathcal{I}_1 = 1 | N = n) \mathbb{P}(N = n).$$

Now, note that

$$\mathbb{P}(\mathcal{I}_1 = 1 | N = n) = \frac{n}{n+d},$$

and so

$$\mathbb{E}(Y) = d \sum_{n \geq 1} \left[\frac{n}{n+d} \mathbb{P}(N = n) \right] = d \mathbb{E} \left(\frac{N}{N+d} \right). \quad (3.6)$$

From the theory of homogeneous branching processes we see that $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ (and also $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$) survives if and only if $\mathbb{E}(Y) > 1$.

Proposition 3.7. *Consider the process $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. Let V_d be the survival event and I_d the number of colonies created during the process. Then*

$$\mathbb{P}(V_d) = \sum_{r=1}^{d+1} \left[(1 - \rho^r) \binom{d+1}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right]$$

where ρ , the extinction probability for the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, is the smallest non-negative solution of

$$\sum_{y=0}^d \left[s^y \binom{d}{y} \sum_{n=y}^{\infty} \frac{\binom{n}{y}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right] = s.$$

On the subcritical regime, which means

$$\mathbb{E} \left[\frac{N}{N+d} \right] < \frac{1}{d}$$

it holds that

$$\mathbb{E}(I_d) = \frac{1 + \mathbb{E} \left(\frac{N}{N+d} \right)}{1 - d \mathbb{E} \left(\frac{N}{N+d} \right)}.$$

Proof of Proposition 3.7

When a colony placed at the origin collapses, Y_R new colonies are formed by the survivors at its neighboring vertices. If a colony located outside the origin collapses, Y new colonies are created on neighboring vertices. Observe that the process $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ behaves as a non-homogeneous branching process $\{Z_n\}_{n \geq 0}$ with

$$Z_0 = 1, Z_{n+l} = \sum_{i=1}^{Z_n} X_{n,i}, n \geq 0,$$

where $X_{1,1}$ and $X_{n,i}, n > 1$ are distributed as Y_R and Y , respectively. So, we have that

$$\mathbb{P}(V_d) = \sum_{r=0}^{d+1} \mathbb{P}(V_d | Y_R = r) \mathbb{P}(Y_R = r).$$

Given that $Y_R = r$ one have r independent $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ processes living on r independent rooted trees. So, we have that $\mathbb{P}(V_d^C | Y_R = r) = \rho^r, r = 0, 1, 2, \dots, d+1$, where ρ is the smallest non-negative solution of $E(s^Y) = s$.

By using the Total Probability Theorem,

$$\begin{aligned} \mathbb{P}(Y_R = y) &= \sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d+1}{y} \binom{n-1}{y-1}}{\binom{n+d}{d}} \right] \text{ for } y = 0, 1, 2, \dots, d+1. \\ \mathbb{P}(Y = y) &= \sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d}{y} \binom{n}{y}}{\binom{n+d}{d}} \right] \text{ for } y = 0, 1, 2, \dots, d. \end{aligned} \quad (3.7)$$

The probability generating function of Y is

$$G_L(s) = \mathbb{E}(s^Y) = \sum_{y=0}^d s^y \left[\sum_{n=y}^{\infty} \left[\mathbb{P}(N = n) \frac{\binom{d}{y} \binom{n}{y}}{\binom{n+d}{d}} \right] \right]. \quad (3.8)$$

Thus,

$$\mathbb{P}(V_d) = \sum_{r=1}^{d+1} \left[(1 - \rho^r) \binom{d+1}{r} \sum_{n=r}^{\infty} \frac{\binom{n-1}{r-1}}{\binom{n+d}{d}} \mathbb{P}(N = n) \right]$$

As for the second part of the proposition, note that

$$\mathbb{E}(I_d) = \sum_{r=0}^{d+1} \mathbb{E}(I_d | Y_R = r) \mathbb{P}(Y_R = r)$$

Besides, from (3.6) we have

$$\mathbb{E}(I | Y_R = r) = r\mu + 1 \text{ onde } \mu = \left[1 - d\mathbb{E} \left(\frac{N}{N+d} \right) \right]^{-1},$$

see Stirzaker [10, Exercise 2b, p. 280].

Proposition 3.8. *Consider the process $L(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. Let V_d be the survival event and I_d the number of colonies created during the process. Then*

$$\lim_{d \rightarrow \infty} \mathbb{P}(V_d) = 1 - \nu \quad (3.9)$$

where ν is the smallest non-negative solution of $\mathbb{E}(s^N) = s$. Besides that, if $\mathbb{E}(N) < 1$ (the subcritical case) then

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \frac{1}{1 - \mathbb{E}(N)}. \quad (3.10)$$

Proof of Proposition 3.8. In order to prove (3.9) one has to apply [5, Proposition 4.2], observing that $Y \xrightarrow{\mathcal{D}} N$ and $Y_R \xrightarrow{\mathcal{D}} N$, when $d \rightarrow \infty$. Moreover to prove (3.10) observe that

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d) = \lim_{d \rightarrow \infty} \sum_{r=0}^{d+1} \mathbb{E}(I_d | Y_R = r) \mathbb{P}(Y_R = r).$$

As $Y \xrightarrow{\mathcal{D}} N$ and $Y_R \xrightarrow{\mathcal{D}} N$ when $d \rightarrow \infty$ then

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_d | Y_R = r) = \lim_{d \rightarrow \infty} r \frac{1}{1 - \mathbb{E}(Y)} + 1 = \frac{r}{1 - \mathbb{E}(N)} + 1$$

and the result follows from the Dominated Convergence Theorem [11, Theorem 9.1 p. 26]. \square

Proposition 3.9. *Let M_d be the reach of the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, that is, the distance from the origin to the farthest vertex where a colony is formed. Assuming*

$$\mathbb{E} \left[\frac{N}{N+d} \right] < \frac{1}{d}$$

Then

$$\frac{[1 + D(1 - \mu_l)](1 - B_l^{n+1})}{1 + D(1 - \mu_l - \mu_l^{n+1})} \leq \mathbf{P}(M_d \leq n) \leq \frac{\left[1 + \frac{\mu_l(1 - \mu_l)}{B_l} \right] (1 - \mu_l^{n+1})}{1 + \frac{\mu_L(1 - \mu_l)}{B_l} - \mu_l^{n+1}}$$

where

$$\begin{aligned}\mu_l &= d\mathbb{E}\left[\frac{N}{N+d}\right] \\ D &= \max\left\{2; \frac{\mu_l}{\mu_l - \mathbb{P}(N \neq 0)}\right\} \\ B_l &= d(d-1)\left[\mathbb{E}\left(\frac{N(N-1)}{(N+d-1)(N+d)}\right)\right]\end{aligned}$$

Moreover,

$$M_d \xrightarrow{\mathcal{D}} M,$$

where $\mathbb{P}(M \leq m) = g_{m+1}(0)$, being $g(s) = \mathbb{E}(s^N)$ and $g_{m+1}(s) = g(g(\cdots g(s)) \cdots)$ ^{$m+1$ times}.

Proof of Proposition 3.9

Every vertex $x \in \mathbb{T}^d$ which is colonized produces Y new colonies (whose distribution depends only on \mathcal{N} and \mathcal{L}) on the d neighbor vertices which located are further from the origin than x is. By numbering those d vertices, we can represent the random variable Y as

$$Y = \sum_{i=1}^d \mathcal{I}_i$$

where \mathcal{I}_i is the indicator variable of event the vertex i receives at least one individual.

From (3.6) we obtained $\mathbb{E}(Y)$. Now

$$\begin{aligned}Y^2 &= \left(\sum_{i=1}^d \mathcal{I}_i\right)^2 = \sum_{i=1}^d \mathcal{I}_i^2 + 2 \sum_{1 \leq i < j \leq d} \mathcal{I}_i \mathcal{I}_j \\ \mathbb{E}(Y^2) &= d\mathbb{E}(\mathcal{I}_1^2) + d(d-1)\mathbb{E}(\mathcal{I}_1 \mathcal{I}_2)\end{aligned}$$

On the other hand

$$\begin{aligned}\mathbb{E}(\mathcal{I}_1^2) &= \mathbb{P}(\mathcal{I}_1 = 1) = \sum_n [\mathbb{P}(\mathcal{I}_1 = 1 | N = n) \mathbb{P}(N = n)] = \sum_n \left(\frac{n}{n+d}\right) \mathbb{P}(N = n) \\ &= \mathbb{E}\left(\frac{N}{N+d}\right)\end{aligned}$$

and

$$\mathbb{E}(\mathcal{I}_1 \mathcal{I}_2) = \mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1) = \sum_n [\mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1 | N = n) \mathbb{P}(N = n)]$$

where

$$\mathbb{P}(\mathcal{I}_1 = 1; \mathcal{I}_2 = 1 | N = n) = \frac{n(n-1)}{(n+d-1)(n+d)}$$

Then

$$\begin{aligned}\mathbb{E}(Y^2) &= d\mathbb{E}\left(\frac{N}{N+d}\right) + d(d-1)\mathbb{E}\left(\frac{N(N-1)}{(N+d-1)(N+d)}\right) \\ \mathbb{E}(Y(Y-1)) &= d(d-1)\mathbb{E}\left(\frac{N(N-1)}{(N+d-1)(N+d)}\right).\end{aligned}$$

Then the result follows from [1, Theorem 1 p. 331] with $\mu_l = \mathbb{E}(Y)$ and $B_l = \mathbb{E}(Y(Y-1))$.

The convergence $M_d \xrightarrow{D} M$ follows from the fact that $Y \xrightarrow{D} N$ when $d \rightarrow \infty$ and from [5, Proposition 4.1].

Proposition 3.10. *Let $\tau_u(d)$ be the extinction time of the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. If*

$$\mathbb{E}\left[\frac{N}{N+d}\right] < \frac{1}{d}$$

then

$$\mathbb{E}[\tau_u(d)] = \int_0^1 \frac{1-y}{G_L(y)-y} dy.$$

where $G_L(s)$ is given by (3.8).

Proof. Analogous to the proof of Proposition 3.5. □

3.3. Proofs of the Main Results. First we define a coupling between the processes $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ in such a way that the latter is stochastically dominated by the former. Every colony in $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ is associated to a colony in $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. As a consequence, if the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ dies out, the same happens to $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$.

At every catastrophe time at a vertex x in the model $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$, a non-empty group of individuals that tries to colonize the neighbor vertex to x which is closer to the origin than x will create there a new colony provided that that vertex is empty. In the model $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ the same non-empty group of individuals that tries to colonize the same vertex, immediately dies.

Next we define a coupling between the processes $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$ in such a way that the former is stochastically dominated by the latter. Every colony in $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ can be associated to a colony in $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$. Thus, if the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$ dies out, the same happens to $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$.

At every catastrophe time at a vertex x in the model $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$, we associate the neighbor vertex to x which is closer to the origin than x to the extra vertex on the model $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$. In the model $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$, a non-empty group of individuals that tries to colonize the neighbor vertex to x which is closer to the origin than x will create there a new colony provided that that vertex is empty. In the model $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$ the same non-empty group of individuals that tries to colonize the extra vertex, founds a new colony there.

Proof of Theorem 2.2 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.1 and 3.6.

Proof of Corollary 2.3 The result follows from Theorem 2.2 and Equation (2.1).

Proof of Theorem 2.5 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.2 and 3.7.

Proof of Theorem 2.7 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.3 and 3.8.

Proof of Corollary 2.8 The result follows from Theorem 2.7 and Equation (2.2).

Proof of Theorem 2.11 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.4 and 3.9.

Proof of Theorem 2.12 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.4 and 3.9.

Proof of Theorem 2.14 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.2 and 3.7.

Proof of Theorem 2.18 The result follows from the fact that the process $C(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ stochastically dominates the process $L(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and by its turn, is stochastically dominated by the process $U(\mathbb{T}_+^{d+1}; \mathcal{N}, \mathcal{L})$, together with Propositions 3.5 and 3.10.

4. COMPARISON WITH MODEL OF INDEPENDENT DISPERSION

In this section, we compare the Self Avoiding Models with independent dispersion and uniform dispersion. The model with uniform dispersion is $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ considered in Section 3.1 and the model with independent dispersion is considered in [5, Section 4.1] and we denote it here by $I(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$. The following results indicate that independent dispersion is a better strategy than uniform dispersion in order to extend population survival.

Theorem 4.1. *Consider the processes $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and $I(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ with survival probability $P(V_u)$ and $P(V_i)$, respectively. If $P(V_i) = 0$ then $P(V_u) = 0$.*

Proof. Observe that for a fixed \mathbb{T}_+^d , \mathcal{N} and \mathcal{L} , both processes, $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and $I(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, behave as a homogeneous branching processes. Every vertex $x \in \mathbb{T}_+^d$ which is colonized produces a random number of new colonies (whose distribution depends only on \mathcal{N} , \mathcal{L} and of dispersion scheme) on the d neighbor vertices which located are further from the origin than x is. We denote by Y_u those number in the process $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and Y_i in the process $I(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$. By numbering those d vertices, we can represent the random variables Y_u and Y_i as

$$Y_u = \sum_{i=1}^d \mathcal{I}_i, \quad \text{and} \quad Y_i = \sum_{i=1}^d \mathcal{U}_i,$$

where \mathcal{I}_i and \mathcal{U}_i are the indicator variables of event the vertex i receives at least one individual in the processes $U(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$ and $I(\mathbb{T}_+^d; \mathcal{N}, \mathcal{L})$, respectively.

Observe that

$$\begin{aligned} \mathbb{E}(Y_u | N = n) &= \sum_{i=1}^d \mathbb{E}(\mathcal{U}_i | N = n) = \sum_{i=1}^d \mathbb{P}(\mathcal{U}_i = 1 | N = n) = d\mathbb{P}(\mathcal{U}_1 = 1 | N = n) \\ &= \frac{dn}{n + d - 1}. \end{aligned}$$

$$\begin{aligned} \mathbb{E}(Y_i | N = n) &= \sum_{i=1}^d \mathbb{E}(\mathcal{I}_i | N = n) = \sum_{i=1}^d \mathbb{P}(\mathcal{I}_i = 1 | N = n) = d\mathbb{P}(\mathcal{I}_1 = 1 | N = n) \\ &= d \left(1 - \left(\frac{d-1}{d} \right)^n \right). \end{aligned}$$

Note that for all $n \geq 0$ and $d \geq 1$ integers,

$$1 - \left(\frac{d-1}{d} \right)^n \geq \frac{n}{n + d - 1}.$$

Thus,

$$\mathbb{E}(Y_i) = \mathbb{E}[\mathbb{E}(Y_i | N)] \geq \mathbb{E}[\mathbb{E}(Y_u | N)] = \mathbb{E}(Y_u). \quad (4.1)$$

The result is obtained through the comparison of the mean numbers of offspring in the branching processes. \square

Theorem 4.2. *Let $I_u(d)$ and $I_i(d)$ be the numbers of colonies created during the processes $U(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$ and $I(\mathbb{T}^d; \mathcal{N}, \mathcal{L})$, respectively.*

(1) *Suppose that*

$$\mathbb{E} \left[\frac{N}{N + d - 1} \right] < \frac{1}{d}.$$

Then

$$\mathbb{E}(I_u(d)) = 1 + \frac{(d+1)\mathbb{E}\left(\frac{N}{N+d}\right)}{1 - d\mathbb{E}\left(\frac{N}{N+d-1}\right)}.$$

(2) Let $G(s)$ be the probability generating function of N . Suppose that

$$G\left(\frac{d-1}{d}\right) < \frac{d-1}{d}$$

then

$$\mathbb{E}(I_i(d)) = 1 + \frac{(d+1)\left(1 - G\left(\frac{d}{d+1}\right)\right)}{1 - d\left(1 - G\left(\frac{d-1}{d}\right)\right)}.$$

(3) For every $d \geq 2$, it holds that $\mathbb{E}(I_u(d)) \leq \mathbb{E}(I_i(d))$. Besides, if $\mathbb{E}(N) < 1$,

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_i(d)) = \lim_{d \rightarrow \infty} \mathbb{E}(I_u(d)) = \frac{1}{1 - \mathbb{E}(N)}.$$

Proof. Items (1) and (2) are Propositions 3.2 and [5, Proposition 4.4], respectively. Item (3) follows from (4.1), Proposition 3.3 and [5, Proposition 4.5]. \square

Corollary 4.3. Consider $U(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and $I(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$.

(1) Suppose that

$$\Phi\left(\frac{\lambda p}{\lambda p + 1}, 1, d + 1\right) > \frac{(\lambda p + 1)[p(\lambda d + d + 1) - 1]}{pd(d + 1)(\lambda + 1)}.$$

Then

$$\mathbb{E}(I_u(d)) = 1 + \frac{(d+1)(\lambda+1)\left(\lambda p + 1 - d\Phi\left(\frac{\lambda p}{\lambda p + 1}, 1, d\right)\right)}{\lambda(\lambda p + 1)\left(1 - \frac{d(\lambda+1)p}{(\lambda p + 1)^2}\left[\lambda p + 1 - (d-1)\Phi\left(\frac{\lambda p}{\lambda p + 1}, 1, d\right)\right]\right)},$$

(2) Suppose that

$$p < \frac{d+1}{d + \lambda(d-1)}$$

Then

$$\mathbb{E}(I_i(d)) = 1 + \frac{(d+1)(\lambda+1)p(d + \lambda p)}{(d + \lambda p + 1)(d(1 - p(\lambda + 1)) + \lambda p)}$$

(3) If $(\lambda + 1)p < 1$,

$$\lim_{d \rightarrow \infty} \mathbb{E}(I_i(d)) = \lim_{d \rightarrow \infty} \mathbb{E}(I_u(d)) = \frac{1}{1 - (\lambda + 1)p}.$$

Proof. Follows from Theorem 4.2 and equations (2.1) and (2.2). \square

Theorem 4.4. Let $\tau_u(d)$ and $\tau_i(d)$ be the extinction times of the processes $U(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and $I(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$.

(1) Suppose that

$$\Phi\left(\frac{\lambda p}{\lambda p + 1}, 1, d + 1\right) > \frac{(\lambda p + 1)[p(\lambda d + d + 1) - 1]}{pd(d + 1)(\lambda + 1)}.$$

Then

$$\mathbb{E}(\tau_u(d)) = \int_0^1 \frac{1 - s}{G_u(s) - s} ds,$$

where

$$G_u(s) = \frac{1}{(\lambda p + 1)} \left[1 - p + \frac{(\lambda + 1)}{\lambda} \sum_{y=1}^d \left[\binom{d}{y} s^y \sum_{n=y}^{\infty} \left[\left(\frac{\lambda p}{\lambda p + 1}\right)^n \frac{\binom{n-1}{y-1}}{\binom{n+d-1}{d-1}} \right] \right] \right].$$

(2) Suppose that

$$p < \frac{d + 1}{d + \lambda(d - 1)}$$

Then

$$\mathbb{E}(\tau_i(d)) = \int_0^1 \frac{1 - s}{G_i(s) - s} ds,$$

where

$$G_i(s) = \frac{1}{\lambda p + 1} \left[1 - p + \frac{\lambda + 1}{\lambda} \sum_{y=1}^d \left[s^y \binom{d}{y} \sum_{n=y}^{\infty} T(n, y) \left(\frac{\lambda p}{d(\lambda p + 1)}\right)^n \right] \right],$$

with

$$T(n, k) = \sum_{i=0}^k \left[(-1)^i \binom{k}{i} (k - i)^n \right], n \geq k.$$

Remark 4.5. The quantity $T(n, k)$ gives the number of surjective functions $f : A \rightarrow B$, where $|A| = n$ and $|B| = k$, see Tucker [12, p. 319].

Proof. Item (1) is Proposition 3.5 with $\mathcal{N} = \mathcal{P}(\lambda)$ and $\mathcal{L} = \text{Bin}(p)$. Proof of item (2) follows in an analogous way. On this last case, the probability generating function of the number of colonies created right after a catastrophe, Y_d , is given by

$$G_i(s) = \sum_{y=0}^d \left[s^y \binom{d}{y} \sum_{n=y}^{\infty} T(n, y) \mathbb{P}[N = n] \right],$$

see Machado et al [5, proof of Proposition 4.4] for details. Finally, by using Equation (2.1) follows the result. \square

Example 4.6. Consider $U(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$ and $I(\mathbb{T}_+^d; \mathcal{P}(\lambda), \text{Bin}(p))$ with $\lambda = 1$ and $p = \frac{1}{2}$. By using Theorem 4.4 we compute the mean extinction times for both models with $2 \leq d \leq 6$, see Table 1.

TABLE 1. Mean extinction times for $U(\mathbb{T}_+^d; \mathcal{P}(1), \text{Bin}(\frac{1}{2}))$ and $I(\mathbb{T}_+^d; \mathcal{P}(1), \text{Bin}(\frac{1}{2}))$.

d	2	3	4	5	6
$\mathbb{E}(\tau_u(d))$	3.494	3.862	4.159	4.408	4.623
$\mathbb{E}(\tau_i(d))$	3.831	4.372	4.779	5.108	5.384

REFERENCES

- [1] A. Agresti. Bounds on the Extinction Time Distribution of a Branching Process. *Advances in Applied Probability*, **6** (2), 322-335 (1974).
- [2] V.V. Junior, F.P. Machado and A. Roldán-Correa. Dispersion as a Survival Strategy. *Journal of Statistical Physics* **164** (4), 937 - 951 (2016).
- [3] V.V. Junior, F.P. Machado and A. Roldán-Correa. Evaluating dispersion strategies in growth models subject to geometric catastrophes. *Journal of Statistical Physics* **183**, 30 (2021).
- [4] V.V. Junior, F.P. Machado and A. Roldán-Correa. Extinction time in growth models subject to geometric catastrophes. *Journal of Statistical Mechanics: Theory and Experiment* **2023** (4), 043501, (2023).
- [5] F.P. Machado, A. Roldán-Correa and V.V. Junior. Colonization and Collapse on Homogeneous Trees. *Journal of Statistical Physics* **173**, 1386-1407 (2018).
- [6] F.P. Machado, A. Roldán-Correa and R. Schinazi. Colonization and Collapse. *ALEA-Latin American Journal of Probability and Mathematical Statistics* **14**, 719-731 (2017).
- [7] P. Narayan. On the Extinction Time of a Continuous time Markov Branching Process, *Austral. J. Statist.* **24**, (2), 160-164, (1982).
- [8] O. Ronce. How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annu. Rev. Ecol. Evol. Syst.* **38**, 231-253, (2007).
- [9] R. Schinazi. Does random dispersion help survival? *Journal of Statistical Physics*, **159**, (1), 101-107 (2015).
- [10] D. Stirzaker. Elementary Probability, second edition, *Cambridge University Press*, (2003).
- [11] H. Thorisson. Coupling, stationarity, and regeneration. *Springer-Verlag, New York*. (2000).
- [12] A. Tucker. Applied Combinatorics 6th ed. *John Wiley & Sons, Inc.* (2012).

(Valdivino V. Junior) FEDERAL UNIVERSITY OF GOIAS, CAMPUS SAMAMBAIA, CEP 74001-970, GOIÂNIA, GO, BRAZIL.

(Fábio P. Machado) INSTITUTE OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SÃO PAULO, RUA DO MATÃO 1010, CEP 05508-090, SÃO PAULO, SP, BRAZIL.

(Alejandro Roldán-Correa) INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD DE ANTIOQUIA, CALLE 67, NO 53-108, MEDELLIN, COLOMBIA.

Email address: fmachado@ime.usp.br, alejandro.roldan@udea.edu.co, vvjunior@ufg.br