

Some three-dimensional non-nilpotent Leibniz algebras: automorphism groups

L. A. Kurdachenko, O. O. Pypka

Oles Honchar Dnipro National University, Dnipro, Ukraine

e-mail: lkurdachenko@gmail.com, sasha.pypka@gmail.com

I. Ya. Subbotin

National University, Los Angeles, USA

e-mail: isubboti@nu.edu

Abstract

The article presents the structure of the automorphism groups of two types of non-nilpotent Leibniz algebras with a dimension of 3.

Key Words: Leibniz algebra, automorphism group.

2020 MSC: 17A32, 17A36.

1 Introduction.

A *Leibniz algebra* is defined as a vector space L over a field F with a bilinear binary operation $[\cdot, \cdot]$ that satisfies the following condition:

$$[x, [y, z]] = [[x, y], z] + [y, [x, z]]$$

for all $x, y, z \in L$ [2]. The term “Leibniz algebra” first appeared in [14] and [15]. We will not provide all the definitions related to Leibniz algebras here. They can be found, for example, in the monographs [1] or [10].

Despite the many analogies between Leibniz algebras and Lie algebras, these types of algebras differ significantly from each other (see, for example, survey articles [3, 4, 12, 17]).

A bijective linear transformation f of a Leibniz algebra L is called an *automorphism* of the algebra L if it satisfies the following condition:

$$f([x, y]) = [f(x), f(y)]$$

for all $x, y \in L$. It is easy to prove that the set $Aut_{[\cdot, \cdot]}(L)$ of all automorphisms of L is a group by a multiplication (see, for example, [9]).

Describing the automorphism groups of algebraic structures is a fundamental and essential problem in algebra. Leibniz algebras are no exception. It is prudent to begin by exploring the automorphism groups of those Leibniz algebras whose structures are already well understood.

As of today, the structure of the automorphism groups of one-generated Leibniz algebras [9, 13], two-dimensional Leibniz algebras [11], and some three-dimensional Leibniz algebras [5, 6, 7, 11, 16] is known.

This article extends our research to two other types of three-dimensional non-nilpotent Leibniz algebras. It should be noted that the most comprehensive description of three-dimensional Leibniz algebras over arbitrary fields can be found in [8].

2 The automorphism group of L_1 .

Let us consider the following type of non-nilpotent Leibniz algebras:

$$L_1 = Fe_1 \oplus Fe_2 \oplus Fe_3 \text{ where } [e_1, e_1] = [e_1, e_3] = e_3, [e_1, e_2] = e_2, \\ [e_2, e_1] = [e_2, e_2] = [e_2, e_3] = [e_3, e_1] = [e_3, e_2] = [e_3, e_3] = 0.$$

In other words, L_1 is the sum of the ideal $A_2 = Fe_2$ and the non-nilpotent one-generated Leibniz algebra $A_1 = Fe_1 \oplus Fe_3$, which has dimension 2, $[A_1, A_2] = A_2$, $\text{Leib}(L_1) = Fe_2 \oplus Fe_3 = [L_1, L_1] = \zeta^{\text{left}}(L_1)$, $\zeta^{\text{right}}(L_1) = \zeta(L_1) = \langle 0 \rangle$.

The following theorem illustrates the structure of the automorphism group for these Leibniz algebras.

Denote by Ξ the canonical monomorphism of $\text{Aut}_{[\cdot, \cdot]}(L)$ in $M_3(F)$.

Theorem 2.1. *Let G be an automorphism group of the Leibniz algebra L_1 . Then G is isomorphic to a subgroup of $GL_3(F)$ that consists of matrices of the form:*

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix},$$

$\alpha_2, \alpha_3, \beta_2, \beta_3 \in F$. Furthermore, G is isomorphic to the group $GL_2(F)$. Moreover, it is a product of two subgroups $C_1 = C_G(e_1)$, $C_2 = C_G(e_2)$, and $\Xi(C_1)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta_2 & 0 \\ 0 & \beta_3 & 1 \end{pmatrix},$$

$\beta_2, \beta_3 \in F$, and $\Xi(C_2)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & 1 & \alpha_2 \\ \alpha_3 & 0 & 1 + \alpha_3 \end{pmatrix},$$

$\alpha_2, \alpha_3 \in F$. A subgroup C_1 is a semidirect product of a normal subgroup C_3 , which is isomorphic to the additive group of a field F , and a subgroup C_4 , which is isomorphic to the multiplicative group of a field F ; a subgroup C_2 is a semidirect product of a normal subgroup, which is isomorphic to the additive group of a field F , and a subgroup that is isomorphic to the multiplicative group of a field F .

Proof. Let $L = L_1$ and let $f \in \text{Aut}_{[\cdot]}(L)$. By [9, Lemma 2.1], $f([L, L]) = [L, L]$, so that

$$\begin{aligned} f(e_1) &= \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \\ f(e_2) &= \beta_2 e_2 + \beta_3 e_3, \\ f(e_3) &= \gamma_2 e_2 + \gamma_3 e_3. \end{aligned}$$

Then

$$\begin{aligned} f(e_3) &= f([e_1, e_1]) = [f(e_1), f(e_1)] \\ &= [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3] \\ &= \alpha_1^2 [e_1, e_1] + \alpha_1 \alpha_2 [e_1, e_2] + \alpha_1 \alpha_3 [e_1, e_3] \\ &= \alpha_1^2 e_3 + \alpha_1 \alpha_2 e_2 + \alpha_1 \alpha_3 e_3 \\ &= \alpha_1 \alpha_2 e_2 + (\alpha_1^2 + \alpha_1 \alpha_3) e_3, \\ f(e_3) &= f([e_1, e_3]) = [f(e_1), f(e_3)] \\ &= [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \gamma_2 e_2 + \gamma_3 e_3] \\ &= \alpha_1 \gamma_2 [e_1, e_2] + \alpha_1 \gamma_3 [e_1, e_3] \\ &= \alpha_1 \gamma_2 e_2 + \alpha_1 \gamma_3 e_3, \\ f(e_2) &= f([e_1, e_2]) = [f(e_1), f(e_2)] \\ &= [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \beta_2 e_2 + \beta_3 e_3] \\ &= \alpha_1 \beta_2 [e_1, e_2] + \alpha_1 \beta_3 [e_1, e_3] \\ &= \alpha_1 \beta_2 e_2 + \alpha_1 \beta_3 e_3. \end{aligned}$$

Thus, we can see that

$$\begin{aligned} \alpha_1 \alpha_2 e_2 + (\alpha_1^2 + \alpha_1 \alpha_3) e_3 &= \alpha_1 \gamma_2 e_2 + \alpha_1 \gamma_3 e_3 = \gamma_2 e_2 + \gamma_3 e_3, \\ \alpha_1 \beta_2 e_2 + \alpha_1 \beta_3 e_3 &= \beta_2 e_2 + \beta_3 e_3. \end{aligned}$$

It follows that

$$\alpha_1 \alpha_2 = \alpha_1 \gamma_2 = \gamma_2, \quad \alpha_1^2 + \alpha_1 \alpha_3 = \alpha_1 \gamma_3 = \gamma_3, \quad \alpha_1 \beta_2 = \beta_2, \quad \alpha_1 \beta_3 = \beta_3.$$

If we suppose that $\alpha_1 = 0$, then $f(e_1) \in Fe_2 \oplus Fe_3$. Since $f(e_2), f(e_3) \in Fe_2 \oplus Fe_3$, we obtain that $f(L) \leq Fe_2 \oplus Fe_3$ – in particular, $f(L) \neq L$ – and we obtain a contradiction with the fact that f is an automorphism of L . Hence, $\alpha_1 \neq 0$. It follows that $\alpha_2 = \gamma_2$.

If we suppose that both $\beta_2 = \gamma_2 = 0$, then $f(e_2) \in Fe_3$ and $f(e_3) \in Fe_3$, so that $f([L, L]) \leq Fe_3$. On the other hand, $\dim_F([L, L]) = 2$, and we, again, obtain a contradiction with the fact that f is an automorphism of L . Hence, $(\beta_2, \gamma_2) \neq (0, 0)$. It follows that $\alpha_1 = 1$ and then $1 + \alpha_3 = \gamma_3$.

Denote by Ξ the canonical monomorphism of $\text{Aut}_{[\cdot]}(L)$ in $M_3(F)$. Then $\Xi(f)$ is the following matrix:

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix},$$

$\alpha_2, \alpha_3, \beta_2, \beta_3 \in F$.

Let f be a linear transformation of L , having in a basis $\{e_1, e_2, e_3\}$ the matrix above. Let x, y be the arbitrary elements of L , $x = \xi_1 e_1 + \xi_2 e_2 + \xi_3 e_3$, $y = \eta_1 e_1 + \eta_2 e_2 + \eta_3 e_3$, where $\xi_1, \xi_2, \xi_3, \eta_1, \eta_2, \eta_3 \in F$. Then

$$\begin{aligned}
[x, y] &= [\xi_1 e_1 + \xi_2 e_2 + \xi_3 e_3, \eta_1 e_1 + \eta_2 e_2 + \eta_3 e_3] \\
&= \xi_1 \eta_1 [e_1, e_1] + \xi_1 \eta_2 [e_1, e_2] + \xi_1 \eta_3 [e_1, e_3] \\
&= \xi_1 \eta_1 e_3 + \xi_1 \eta_2 e_2 + \xi_1 \eta_3 e_3 \\
&= \xi_1 \eta_2 e_2 + (\xi_1 \eta_1 + \xi_1 \eta_3) e_3 = \xi_1 \eta_2 e_2 + \xi_1 (\eta_1 + \eta_3) e_3, \\
f(x) &= f(\xi_1 e_1 + \xi_2 e_2 + \xi_3 e_3) = \xi_1 f(e_1) + \xi_2 f(e_2) + \xi_3 f(e_3) \\
&= \xi_1 (e_1 + \alpha_2 e_2 + \alpha_3 e_3) + \xi_2 (\beta_2 e_2 + \beta_3 e_3) + \xi_3 (\alpha_2 e_2 + (1 + \alpha_3) e_3) \\
&= \xi_1 e_1 + (\xi_1 \alpha_2 + \xi_2 \beta_2 + \xi_3 \alpha_2) e_2 + (\xi_1 \alpha_3 + \xi_2 \beta_3 + \xi_3 + \xi_3 \alpha_3) e_3, \\
f(y) &= \eta_1 e_1 + (\eta_1 \alpha_2 + \eta_2 \beta_2 + \eta_3 \alpha_2) e_2 + (\eta_1 \alpha_3 + \eta_2 \beta_3 + \eta_3 + \eta_3 \alpha_3) e_3, \\
f([x, y]) &= f(\xi_1 \eta_2 e_2 + \xi_1 (\eta_1 + \eta_3) e_3) = \xi_1 \eta_2 f(e_2) + \xi_1 (\eta_1 + \eta_3) f(e_3) \\
&= \xi_1 \eta_2 (\beta_2 e_2 + \beta_3 e_3) + \xi_1 (\eta_1 + \eta_3) (\alpha_2 e_2 + (1 + \alpha_3) e_3) \\
&= (\xi_1 \eta_2 \beta_2 + \xi_1 (\eta_1 + \eta_3) \alpha_2) e_2 + (\xi_1 \eta_2 \beta_3 + \xi_1 (\eta_1 + \eta_3) (1 + \alpha_3)) e_3, \\
[f(x), f(y)] &= [\xi_1 e_1 + (\xi_1 \alpha_2 + \xi_2 \beta_2 + \xi_3 \alpha_2) e_2 + (\xi_1 \alpha_3 + \xi_2 \beta_3 + \xi_3 + \xi_3 \alpha_3) e_3, \\
&\quad \eta_1 e_1 + (\eta_1 \alpha_2 + \eta_2 \beta_2 + \eta_3 \alpha_2) e_2 + (\eta_1 \alpha_3 + \eta_2 \beta_3 + \eta_3 + \eta_3 \alpha_3) e_3] \\
&= \xi_1 \eta_1 [e_1, e_1] + \xi_1 (\eta_1 \alpha_2 + \eta_2 \beta_2 + \eta_3 \alpha_2) [e_1, e_2] \\
&\quad + \xi_1 (\eta_1 \alpha_3 + \eta_2 \beta_3 + \eta_3 + \eta_3 \alpha_3) [e_1, e_3] \\
&= \xi_1 \eta_1 e_3 + \xi_1 (\eta_1 \alpha_2 + \eta_2 \beta_2 + \eta_3 \alpha_2) e_2 + \xi_1 (\eta_1 \alpha_3 + \eta_2 \beta_3 + \eta_3 + \eta_3 \alpha_3) e_3 \\
&= \xi_1 (\eta_1 \alpha_2 + \eta_2 \beta_2 + \eta_3 \alpha_2) e_2 + \xi_1 (\eta_1 + \eta_1 \alpha_3 + \eta_2 \beta_3 + \eta_3 + \eta_3 \alpha_3) e_3.
\end{aligned}$$

We can see that $f([x, y]) = [f(x), f(y)]$. It follows that a linear transformation f having in a basis $\{e_1, e_2, e_3\}$ the above matrix is an automorphism of a Leibniz algebra.

Denote by Φ the mapping of $\Xi(G)$ in $GL_2(F)$ defined by the rule:

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix} \rightarrow \begin{pmatrix} \beta_2 & \alpha_2 \\ \beta_3 & 1 + \alpha_3 \end{pmatrix}.$$

We have:

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \gamma_2 & \kappa_2 & \gamma_2 \\ \gamma_3 & \kappa_3 & 1 + \gamma_3 \end{pmatrix} = \\
\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 + \beta_2 \gamma_2 + \alpha_2 \gamma_3 & \beta_2 \kappa_2 + \alpha_2 \kappa_3 & \beta_2 \gamma_2 + \alpha_2 (1 + \gamma_3) \\ \alpha_3 + \beta_3 \gamma_2 + (1 + \alpha_3) \gamma_3 & \beta_3 \kappa_2 + (1 + \alpha_3) \kappa_3 & \beta_3 \gamma_2 + (1 + \alpha_3) (1 + \gamma_3) \end{pmatrix}$$

and

$$\begin{aligned} & \begin{pmatrix} \beta_2 & \alpha_2 \\ \beta_3 & 1 + \alpha_3 \end{pmatrix} \begin{pmatrix} \kappa_2 & \gamma_2 \\ \kappa_3 & 1 + \gamma_3 \end{pmatrix} \\ &= \begin{pmatrix} \beta_2\kappa_2 + \alpha_2\kappa_3 & \beta_2\gamma_2 + \alpha_2(1 + \gamma_3) \\ \beta_3\kappa_2 + (1 + \alpha_3)\kappa_3 & \beta_3\gamma_2 + (1 + \alpha_3)(1 + \gamma_3) \end{pmatrix} \end{aligned}$$

These equalities show that a mapping Φ is a homomorphism. Clearly, $Ker(\Phi)$ consists only of the identity matrix. Finally, let

$$\begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{21} & \sigma_{22} \end{pmatrix}$$

be the arbitrary element of $GL_2(F)$. Consider the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ \sigma_{12} & \sigma_{11} & \sigma_{12} \\ \sigma_{22} - 1 & \sigma_{21} & \sigma_{22} \end{pmatrix}.$$

This matrix is a preimage of the matrix above by the mapping Φ . Moreover, both these matrices have the same determinants. It follows that the last matrix belongs to $\Xi(G)$, so that a mapping Φ is an isomorphism.

Let $C_1 = C_G(e_1)$, $C_2 = C_G(e_2)$. Then $\Xi(C_1)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta_2 & 0 \\ 0 & \beta_3 & 1 \end{pmatrix},$$

$\beta_2, \beta_3 \in F$, and $\Xi(C_2)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & 1 & \alpha_2 \\ \alpha_3 & 0 & 1 + \alpha_3 \end{pmatrix},$$

$\alpha_2, \alpha_3 \in F$. We have:

$$\begin{pmatrix} 1 & 0 & 0 \\ \sigma_2 & 1 & \sigma_2 \\ \sigma_3 & 0 & 1 + \sigma_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \kappa_2 & 0 \\ 0 & \kappa_3 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ \sigma_2 & \kappa_2 + \sigma_2\kappa_3 & \sigma_2 \\ \sigma_3 & (1 + \sigma_3)\kappa_3 & 1 + \sigma_3 \end{pmatrix}.$$

If

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix}$$

is an arbitrary matrix of $\Xi(G)$, then we obtain

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & \beta_2 & \alpha_2 \\ \alpha_3 & \beta_3 & 1 + \alpha_3 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ \sigma_2 & 1 & \sigma_2 \\ \sigma_3 & 0 & 1 + \sigma_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \kappa_2 & 0 \\ 0 & \kappa_3 & 1 \end{pmatrix}$$

where $\sigma_2 = \alpha_2$, $\sigma_3 = \alpha_3$, $\kappa_3 = \beta_3(1 + \alpha_3)^{-1}$, $\kappa_2 = \beta_2 - \alpha_2\beta_3(1 + \alpha_3)^{-1}$. Thus, we can see that a group G is a product of the subgroups C_1 and C_2 .

Furthermore, a subgroup C_1 has a normal subgroup C_3 such that $\Xi(C_3)$ is the subset of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \beta_3 & 1 \end{pmatrix},$$

which is isomorphic to the additive group of a field F , and a subgroup C_4 such that $\Xi(C_4)$ is the subset of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta_2 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

which is isomorphic to the multiplicative group of a field F . Moreover, C_1 is clearly a semidirect product of C_3 and C_4 .

Now, consider the group $\Xi(C_2)$. It is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & 1 & \alpha_2 \\ \alpha_3 & 0 & 1 + \alpha_3 \end{pmatrix},$$

$\alpha_2, \alpha_3 \in F$. Denote by Φ the mapping of $\Xi(C_2)$ in $GL_2(F)$ defined by the rule:

$$\begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & 1 & \alpha_2 \\ \alpha_3 & 0 & 1 + \alpha_3 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & \alpha_2 \\ 0 & 1 + \alpha_3 \end{pmatrix}.$$

We have:

$$\begin{aligned} & \begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 & 1 & \alpha_2 \\ \alpha_3 & 0 & 1 + \alpha_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ \gamma_2 & 1 & \gamma_2 \\ \gamma_3 & 0 & 1 + \gamma_3 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ \alpha_2 + \gamma_2 + \alpha_2\gamma_3 & 1 & \gamma_2 + \alpha_2(1 + \gamma_3) \\ \alpha_3 + (1 + \alpha_3)\gamma_3 & 0 & (1 + \alpha_3)(1 + \gamma_3) \end{pmatrix} \end{aligned}$$

and

$$\begin{pmatrix} 1 & \alpha_2 \\ 0 & 1 + \alpha_3 \end{pmatrix} \begin{pmatrix} 1 & \gamma_2 \\ 0 & 1 + \gamma_3 \end{pmatrix} = \begin{pmatrix} 1 & \gamma_2 + \alpha_2(1 + \gamma_3) \\ 0 & (1 + \alpha_3)(1 + \gamma_3) \end{pmatrix}$$

These equalities show that a mapping Φ is a homomorphism. Clearly, $Ker(\Phi)$ consists only of identity matrix. Finally, let

$$\begin{pmatrix} 1 & \sigma_{12} \\ 0 & \sigma_{22} \end{pmatrix}$$

be the arbitrary element of $GL_2(F)$. Consider the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ \sigma_{12} & 1 & \sigma_{12} \\ \sigma_{22} - 1 & 0 & \sigma_{22} \end{pmatrix}.$$

This matrix is a preimage of the matrix above by Φ . Moreover, both these matrices have the same determinants. The last matrix belongs to $\Xi(G)$, so a mapping Φ is an isomorphism.

In turn, clearly $\Xi(C_2)$ is a semidirect product of a normal subgroup that consists of matrices of the form

$$\begin{pmatrix} 1 & \sigma_{12} \\ 0 & 1 \end{pmatrix},$$

(i.e., a subgroup $UT_2(F)$) that is isomorphic to the additive group of a field F and a subgroup that consists of matrices of the form

$$\begin{pmatrix} 1 & 0 \\ 0 & \sigma_{22} \end{pmatrix},$$

which is isomorphic to the multiplicative group of a field F . □

3 The automorphism group of L_2 .

Let us consider the second type of non-nilpotent Leibniz algebras:

$$\begin{aligned} L_2 &= Fe_1 \oplus Fe_2 \oplus Fe_3 \text{ where } [e_1, e_1] = [e_1, e_3] = e_3, \\ &[e_1, e_2] = e_2 + \lambda e_3, \quad 0 \neq \lambda \in F, \\ [e_2, e_1] &= [e_2, e_2] = [e_2, e_3] = [e_3, e_1] = [e_3, e_2] = [e_3, e_3] = 0. \end{aligned}$$

In other words, L_2 is the sum of the ideal $Leib(L_2) = Fe_2 \oplus Fe_3$ and the non-nilpotent one-generated Leibniz algebra $A_1 = Fe_1 \oplus Fe_3$ of dimension 2, $[A_1, Leib(L_2)] = Leib(L_2)$, $Leib(L_2) = [L_2, L_2] = \zeta^{\text{left}}(L_2)$, $\zeta^{\text{right}}(L_2) = \zeta(L_2) = \langle 0 \rangle$.

The following theorem illustrates the structure of the automorphism group for these Leibniz algebras.

Theorem 3.1. *Let G be an automorphism group of the Leibniz algebra L_2 . Then G is isomorphic to a subgroup of $GL_3(F)$ that consists of matrices of the form*

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta_1 & 0 \\ \beta_1 - 1 & \beta_2 & \beta_1 \end{pmatrix},$$

$\beta_1, \beta_2 \in F$. Furthermore, G is abelian, $G = C \times A$ such that $\Xi(C)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \sigma & 1 \end{pmatrix},$$

$\sigma \in F$, and $\Xi(A)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & 0 & \beta \end{pmatrix},$$

$\beta \in F$. A subgroup C is isomorphic to the additive group of a field F , and a subgroup A is isomorphic to the multiplicative group of a field F .

Proof. Let $L = L_2$ and let $f \in \text{Aut}_{[\cdot]}(L)$. By [9, Lemma 2.1], $f([L, L]) = [L, L]$, so that

$$\begin{aligned} f(e_1) &= \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \\ f(e_2) &= \beta_2 e_2 + \beta_3 e_3, \\ f(e_3) &= \gamma_2 e_2 + \gamma_3 e_3. \end{aligned}$$

Then

$$\begin{aligned} f(e_3) &= f([e_1, e_1]) = [f(e_1), f(e_1)] \\ &= [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3] \\ &= \alpha_1^2 [e_1, e_1] + \alpha_1 \alpha_2 [e_1, e_2] + \alpha_1 \alpha_3 [e_1, e_3] \\ &= \alpha_1^2 e_3 + \alpha_1 \alpha_2 (e_2 + \lambda e_3) + \alpha_1 \alpha_3 e_3 \\ &= \alpha_1 \alpha_2 e_2 + (\alpha_1^2 + \lambda \alpha_1 \alpha_2 + \alpha_1 \alpha_3) e_3, \\ f(e_3) &= f([e_1, e_3]) = [f(e_1), f(e_3)] \\ &= [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \gamma_2 e_2 + \gamma_3 e_3] \\ &= \alpha_1 \gamma_2 [e_1, e_2] + \alpha_1 \gamma_3 [e_1, e_3] \\ &= \alpha_1 \gamma_2 (e_2 + \lambda e_3) + \alpha_1 \gamma_3 e_3 \\ &= \alpha_1 \gamma_2 e_2 + (\lambda \alpha_1 \gamma_2 + \alpha_1 \gamma_3) e_3, \\ f([e_1, e_2]) &= [f(e_1), f(e_2)] = [\alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3, \beta_2 e_2 + \beta_3 e_3] \\ &= \alpha_1 \beta_2 [e_1, e_2] + \alpha_1 \beta_3 [e_1, e_3] \\ &= \alpha_1 \beta_2 (e_2 + \lambda e_3) + \alpha_1 \beta_3 e_3 \\ &= \alpha_1 \beta_2 e_2 + (\lambda \alpha_1 \beta_2 + \alpha_1 \beta_3) e_3, \\ f([e_1, e_2]) &= f(e_2 + \lambda e_3) = f(e_2) + \lambda f(e_3) \\ &= \beta_2 e_2 + \beta_3 e_3 + \lambda (\gamma_2 e_2 + \gamma_3 e_3) \\ &= (\beta_2 + \lambda \gamma_2) e_2 + (\beta_3 + \lambda \gamma_3) e_3. \end{aligned}$$

Thus, we can see that

$$\begin{aligned} \alpha_1 \alpha_2 e_2 + (\alpha_1^2 + \lambda \alpha_1 \alpha_2 + \alpha_1 \alpha_3) e_3 &= \\ \alpha_1 \gamma_2 e_2 + (\lambda \alpha_1 \gamma_2 + \alpha_1 \gamma_3) e_3 &= \gamma_2 e_2 + \gamma_3 e_3, \\ \alpha_1 \beta_2 e_2 + (\lambda \alpha_1 \beta_2 + \alpha_1 \beta_3) e_3 &= (\beta_2 + \lambda \gamma_2) e_2 + (\beta_3 + \lambda \gamma_3) e_3. \end{aligned}$$

It follows that

$$\begin{aligned}\alpha_1\alpha_2 &= \alpha_1\gamma_2 = \gamma_2, \quad \alpha_1^2 + \lambda\alpha_1\alpha_2 + \alpha_1\alpha_3 = \lambda\alpha_1\gamma_2 + \alpha_1\gamma_3 = \gamma_3, \\ \alpha_1\beta_2 &= \beta_2 + \lambda\gamma_2, \quad \lambda\alpha_1\beta_2 + \alpha_1\beta_3 = \beta_3 + \lambda\gamma_3.\end{aligned}$$

If we suppose that $\alpha_1 = 0$, then $f(e_1) \in Fe_2 \oplus Fe_3$. Since $f(e_2), f(e_3) \in Fe_2 \oplus Fe_3$, we obtain that $f(L) \leq Fe_2 \oplus Fe_3$ – in particular, $f(L) \neq L$ – and we obtain a contradiction with the fact that f is an automorphism of L . Hence, $\alpha_1 \neq 0$. It follows that $\alpha_2 = \gamma_2$.

If we suppose that both $\beta_2 = \gamma_2 = 0$, then $f(e_2) \in Fe_3$, and $f(e_3) \in Fe_3$. We obtain that $f([L, L]) \leq Fe_3$. On the other hand, $\dim_F([L, L]) = 2$, and we, again, obtain a contradiction with the fact that f is an automorphism of L . Hence, $(\beta_2, \gamma_2) \neq (0, 0)$. Then, if we suppose that $\beta_2 = 0$, we will see that $\lambda\gamma_2 = 0$. Since $\lambda \neq 0$, we obtain that $\gamma_2 = 0$, which is impossible. It follows that $\beta_2 \neq 0$. Suppose that $\gamma_2 = 0$. Since $\beta_2 \neq 0$, we obtain that $\alpha_1 = 1$. If $\gamma_2 \neq 0$, then from $\alpha_1\gamma_2 = \gamma_2$, we obtain that $\alpha_1 = 1$. Thus, in every case, $\alpha_1 = 1$. Then we obtain an equation $\beta_2 = \beta_2 + \lambda\gamma_2$ or $\lambda\gamma_2 = 0$. Since $\lambda \neq 0$, we obtain that $\gamma_2 = 0$. Then $\alpha_2 = 0$, and we obtain $1 + \alpha_3 = \gamma_3$, $\lambda\beta_2 + \beta_3 = \beta_3 + \lambda\gamma_3$. The last equation implies that $\lambda\beta_2 = \lambda\gamma_3$. Since $\lambda \neq 0$, we obtain that $\beta_2 = \gamma_3$.

Denote by Ξ the canonical monomorphism of $\text{Aut}_{[\cdot]}(L)$ in $M_3(F)$. Then $\Xi(f)$ is the following matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta_2 & 0 \\ \beta_2 - 1 & \beta_3 & \beta_2 \end{pmatrix},$$

$\beta_2, \beta_3 \in F$. Since this matrix must be non-degenerate, $\beta_2 \neq 0$.

Let f be a linear transformation of L , having in a basis $\{e_1, e_2, e_3\}$ the matrix above. Let x, y be the arbitrary elements of L , $x = \xi_1e_1 + \xi_2e_2 + \xi_3e_3$, $y = \eta_1e_1 + \eta_2e_2 + \eta_3e_3$ where $\xi_1, \xi_2, \xi_3, \eta_1, \eta_2, \eta_3 \in F$. Then

$$\begin{aligned}[x, y] &= [\xi_1e_1 + \xi_2e_2 + \xi_3e_3, \eta_1e_1 + \eta_2e_2 + \eta_3e_3] \\ &= \xi_1\eta_1[e_1, e_1] + \xi_1\eta_2[e_1, e_2] + \xi_1\eta_3[e_1, e_3] \\ &= \xi_1\eta_1e_3 + \xi_1\eta_2(e_2 + \lambda e_3) + \xi_1\eta_3e_3 \\ &= \xi_1\eta_2e_2 + (\xi_1\eta_1 + \lambda\xi_1\eta_2 + \xi_1\eta_3)e_3, \\ f(x) &= f(\xi_1e_1 + \xi_2e_2 + \xi_3e_3) = \xi_1f(e_1) + \xi_2f(e_2) + \xi_3f(e_3) \\ &= \xi_1(e_1 + (\beta_2 - 1)e_3) + \xi_2(\beta_2e_2 + \beta_3e_3) + \xi_3(\beta_2e_3) \\ &= \xi_1e_1 + \xi_2\beta_2e_2 + (\xi_1(\beta_2 - 1) + \xi_2\beta_3 + \xi_3\beta_2)e_3, \\ f(y) &= \eta_1e_1 + \eta_2\beta_2e_2 + (\eta_1(\beta_2 - 1) + \eta_2\beta_3 + \eta_3\beta_2)e_3.\end{aligned}$$

Therefore,

$$\begin{aligned}
f([x, y]) &= f(\xi_1\eta_2e_2 + (\xi_1\eta_1 + \lambda\xi_1\eta_2 + \xi_1\eta_3)e_3) \\
&= \xi_1\eta_2f(e_2) + (\xi_1\eta_1 + \lambda\xi_1\eta_2 + \xi_1\eta_3)f(e_3) \\
&= \xi_1\eta_2(\beta_2e_2 + \beta_3e_3) + (\xi_1\eta_1 + \lambda\xi_1\eta_2 + \xi_1\eta_3)\beta_2e_3 \\
&= \xi_1\eta_2\beta_2e_2 + (\xi_1\eta_2\beta_3 + \beta_2\xi_1\eta_1 + \lambda\xi_1\eta_2\beta_2 + \xi_1\eta_3\beta_2)e_3, \\
[f(x), f(y)] &= [\xi_1e_1 + \xi_2\beta_2e_2 + (\xi_1(\beta_2 - 1) + \xi_2\beta_3 + \xi_3\beta_2)e_3, \\
&\quad \eta_1e_1 + \eta_2\beta_2e_2 + (\eta_1(\beta_2 - 1) + \eta_2\beta_3 + \eta_3\beta_2)e_3] \\
&= \xi_1\eta_1[e_1, e_1] + \xi_1\eta_2\beta_2[e_1, e_2] + \xi_1(\eta_1(\beta_2 - 1) + \eta_2\beta_3 + \eta_3\beta_2)[e_1, e_3] \\
&= \xi_1\eta_1e_3 + \xi_1\eta_2\beta_2(e_2 + \lambda e_3) + \xi_1(\eta_1(\beta_2 - 1) + \eta_2\beta_3 + \eta_3\beta_2)e_3 \\
&= \xi_1\eta_2\beta_2e_2 + (\xi_1\eta_1 + \xi_1\eta_2\beta_2\lambda + \xi_1\eta_1\beta_2 - \xi_1\eta_1 + \xi_1\eta_2\beta_3 + \xi_1\eta_3\beta_2)e_3 \\
&= \xi_1\eta_2\beta_2e_2 + (\xi_1\eta_2\beta_2\lambda + \xi_1\eta_1\beta_2 + \xi_1\eta_2\beta_3 + \xi_1\eta_3\beta_2)e_3.
\end{aligned}$$

We can see that $f([x, y]) = [f(x), f(y)]$. It follows that a linear transformation f having in a basis $\{e_1, e_2, e_3\}$ the matrix above, is an automorphism of a Leibniz algebra L .

The equality

$$\begin{aligned}
&\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & \gamma & \beta \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sigma & 0 \\ \sigma - 1 & \kappa & \sigma \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta\sigma & 0 \\ \beta - 1 + \beta(\sigma - 1) & \gamma\sigma + \beta\kappa & \beta\sigma \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta\sigma & 0 \\ \beta\sigma - 1 & \gamma\sigma + \beta\kappa & \beta\sigma \end{pmatrix}
\end{aligned}$$

shows that a group G is abelian.

According to the proof above, a subspace Fe_3 is G -invariant. Let $C = C_G(Fe_3)$. Then we can see that $\Xi(C)$ is the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \beta & 1 \end{pmatrix},$$

$\beta \in F$. It is not hard to see that C is isomorphic to the additive group of a field F .

Denote by A the preimage by Ξ of the set of matrices of the form

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & 0 & \beta \end{pmatrix},$$

$\beta \in F$. We have:

$$\begin{aligned}
&\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & 0 & \beta \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sigma & 0 \\ \sigma - 1 & 0 & \sigma \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta\sigma & 0 \\ \beta - 1 + \beta(\sigma - 1) & 0 & \beta\sigma \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta\sigma & 0 \\ \beta\sigma - 1 & 0 & \beta\sigma \end{pmatrix}.
\end{aligned}$$

We can now see that $\Xi(A)$ is a subgroup of $GL_3(F)$, so that A is a subgroup of G .

Moreover, the mapping, defined by the rule

$$\beta \rightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & 0 & \beta \end{pmatrix},$$

$\beta \in F$, is an isomorphism of the multiplicative group of F on $\Xi(A)$. Thus, we obtain that A is isomorphic to the multiplicative group of a field F .

Furthermore, the equality

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & \sigma & \beta \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \beta & 0 \\ \beta - 1 & 0 & \beta \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & \beta^{-1}\sigma & 1 \end{pmatrix}$$

shows that G is a product of the subgroups C and A . Moreover, this product is direct. \square

References

- [1] Ayupov, Sh., Omirov, B., Rakhimov, I.: Leibniz algebras: structure and classification. CRC Press, Taylor & Francis Group, 2020.
- [2] Blokh, A.: On a generalization of the concept of Lie algebra. Dokl. Akad. Nauk SSSR 1965; **165**(3): pp. 471–473.
- [3] Chupordia, V.A., Pypka, A.A., Semko, N.N., Yashchuk, V.S.: Leibniz algebras: a brief review of current results. Carpathian Math. Publ. 2019; **11**(2): pp. 250–257. <https://doi.org/10.15330/cmp.11.2.250-257>
- [4] Kirichenko, V.V., Kurdachenko, L.A., Pypka, A.A., Subbotin, I.Ya.: Some aspects of Leibniz algebra theory. Algebra Discrete Math. 2017; **24**(1): pp. 1–33.
- [5] Kurdachenko, L.A., Minaiev, P.Ye., Pypka, O.O.: Automorphism groups of some non-nilpotent Leibniz algebras. Res. Math., 2024; **32**(1): pp. 101–109. <https://doi.org/10.15421/242409>
- [6] Kurdachenko, L.A., Pypka, O.O., Semko, M.M.: Description of the automorphism groups of some Leibniz algebras. Res. Math. 2023; **31**(1): pp. 52–61. <https://doi.org/10.15421/242305>
- [7] Kurdachenko, L.A., Pypka, O.O., Semko, M.M.: On the structure of the automorphism groups of some low-dimensional Leibniz algebras. Ukrainian Math. J. 2024; **76**(6): pp. 864–876. <https://doi.org/10.3842/umzh.v76i6.7868>
- [8] Kurdachenko, L.A., Pypka, O.O., Subbotin, I.Ya.: On the structure of low-dimensional Leibniz algebras: some revision. Algebra Discrete Math. 2022; **34**(1): pp. 68–104. <https://doi.org/10.12958/adm2036>

- [9] Kurdachenko, L.A., Pypka, A.A., Subbotin, I.Ya.: On the automorphism groups of some Leibniz algebras. *Int. J. Group Theory* 2023; **12**(1): pp. 1–20. <https://doi.org/10.22108/IJGT.2021.130057.1735>
- [10] Kurdachenko, L.A., Pypka, O.O., Subbotin, I.Ya.: *General Theory of Leibniz Algebras*. Springer, Switzerland, 2024.
- [11] Kurdachenko, L.A., Pypka, O.O., Velychko, T.V.: On the automorphism groups for some Leibniz algebras of low dimensions. *Ukrainian Math. J.* 2023; **74**(10): pp. 1526–1546. <https://doi.org/10.1007/s11253-023-02153-2>
- [12] Kurdachenko, L.A., Semko, N.N., Subbotin, I.Ya.: Applying group theory philosophy to Leibniz algebras: some new developments. *Adv. Group Theory Appl.* 2020; **9**: pp. 71–121. <https://doi.org/10.32037/agta-2020-004>
- [13] Kurdachenko, L.A., Subbotin, I.Ya., Yashchuk, V.S.: On the endomorphisms and derivations of some Leibniz algebras. *J. Algebra Appl.* 2024; **23**(1): no. 2450002. <https://doi.org/10.1142/S0219498824500026>
- [14] Loday, J.-L.: *Cyclic homology*. Springer Verlag, 1992. <https://doi.org/10.1007/978-3-662-11389-9>
- [15] Loday, J.-L.: Une version non commutative des algèbres de Lie; les algèbres de Leibniz. *Enseign. Math.* 1993; **39**: pp. 269–293.
- [16] Minaiev, P.Ye., Pypka, O.O., Semko, L.P.: Low-dimensional nilpotent Leibniz algebras and their automorphism groups. *Algebra Discrete Math.* 2024; **37**(2): pp. 275–286. <https://doi.org/10.12958/adm2264>
- [17] Subbotin, I.Ya.: Methods of group theory in Leibniz algebras: some compelling results. *Res. Math.* 2021; **29**(2): pp. 44–54. <https://doi.org/10.15421/242108>