Is the Difference between Deep Hedging and Delta Hedging a Statistical Arbitrage?*

Pascal François^a, Geneviève Gauthier^b, Frédéric Godin^{c,d}, and Carlos Octavio Pérez Mendoza^{†c}

^aFellow, Canadian Derivatives Institute and Department of Finance, HEC Montréal, Canada

^bGERAD and Department of Decision Sciences, HEC Montréal, Canada

^cConcordia University, Department of Mathematics and Statistics, Canada

^dQuantact Laboratory, Centre de Recherches Mathématiques, Canada

August 1, 2024

Abstract

The recent work of Horikawa and Nakagawa (2024) claims that under a complete market admitting statistical arbitrage, the difference between the hedging position provided by deep hedging and that of the replicating portfolio is a statistical arbitrage. This raises concerns as it entails that deep hedging can include a speculative component aimed simply at exploiting the structure of the risk measure guiding the hedging optimisation problem. We test whether such finding remains true in a GARCH-based market model. We observe that the difference between deep hedging and delta hedging can be a statistical arbitrage if the risk measure considered does not put sufficient relative weight on adverse outcomes. Nevertheless, a suitable choice of risk measure can prevent the deep hedging agent from including a speculative overlay within its hedging strategy.

JEL classification: C45, C61, G32.

Keywords: Deep reinforcement learning, optimal hedging, arbitrage.

^{*}Gauthier is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC, RGPIN-2019-04029), a professorship funded by HEC Montréal, and the HEC Montréal Foundation. Godin is funded by NSERC (RGPIN-2024-04593).

[†]Corresponding author.

Email addresses: pascal.francois@hec.ca (Pascal François), genevieve.gauthier@hec.ca (Geneviève Gauthier), frederic.godin@concordia.ca (Frédéric Godin), carlos.octavio.perez92@gmail.com (Carlos Octavio Pérez Mendoza).

1 Introduction

The seminal paper of Buehler et al. (2019), which proposes to use deep reinforcement learning (RL) methods to obtain optimal hedging procedures for financial derivatives, initiated a whole strand of literature: Halperin (2019), Cao et al. (2020), Du et al. (2020), Carbonneau and Godin (2021), Carbonneau (2021), Horvath et al. (2021), Imaki et al. (2021), Lütkebohmert et al. (2022), Cao et al. (2023), Carbonneau and Godin (2023), Marzban et al. (2023), Mikkilä and Kanniainen (2023), Raj et al. (2023) and Wu and Jaimungal (2023). See also Hambly et al. (2023) and Pickard and Lawryshyn (2023) for related surveys. Deep RL methods are particularly well-suited to solve dynamic hedging problems as such methods can handle the curse of dimensionality, a problem that more traditional methods (e.g., finite elements dynamic programming) struggle to overcome. They can also work with very general dynamics for hedging asset prices, not being limited by mathematical tractability issues.

While the ability of deep hedging strategies to outperform standard counterparts is well-documented, the existing literature has not yet extensively analyzed the structure of optimal policies and explained how such incremental performance is attained. Neagu et al. (2024) make a step in that direction by investigating the impact of the various features on the optimal hedging decision in the presence of illiquidity market impacts.

In their recent work, Horikawa and Nakagawa (2024) investigate complete markets that allow for statistical arbitrage with respect to a specific risk measure ρ . They assert that, within this framework, hedging strategies that minimize the chosen risk metric combine the traditional delta-hedging approach with a statistical arbitrage overlay. In a more general market setting, Buehler et al. (2021) find that the optimal hedging strategy maximizing the entropy utility can also incorporate a statistical arbitrage component. Such claims raise concern about the suitability of the deep hedging approach; incorporating speculative or arbitrage-like components that do not contribute to reducing the risk exposure within hedging portfolios would be deemed undesirable in practice. Our objective is therefore to assess empirically whether deep hedging policies minimizing conventional risk metrics contain a speculative component in a GARCH-based market setting.

The paper is divided as follows. Section 2 provides the hedging problem formulation. Section 3 describes the deep hedging framework used to solve the problem, and discusses the common delta hedging benchmark. Numerical experiments assessing the behavior of the deep versus delta hedging difference strategy are provided in Section 4. Section 5 concludes.

2 Market model for hedging

This paper considers dynamic hedging strategies for European call options, which involve the construction of a self-financing hedging portfolio composed of the underlying asset S and a cash account. The portfolio is rebalanced daily to optimally offset the net risk exposure at the option maturity, denoted as T days. The time-t underlying asset price is S_t . The hedging strategy is represented by the predictable process $\delta = \{\delta_t\}_{t=1}^T$, where δ_t is the number of underlying asset shares held during the interval (t-1,t]. The time-t discounted gain made by the hedging portfolio is $G_t^{\delta} = \sum_{k=1}^t \delta_k (\beta^k S_k e^{q\Lambda} - \beta^{k-1} S_{k-1})$ with $\beta = e^{-r\Lambda}$, where r is the annualized continuously compounded risk-free rate, q is the annualized underlying asset dividend yield, and the period length is $\Lambda = \frac{1}{252}$ years. The time-t self-financing hedging portfolio value is

$$V_t^{\delta}(V_0) = \beta^{-t}(V_0 + G_t^{\delta}), \tag{1}$$

where V_0 is the initial portfolio value that we set to the option price.

The hedging problem is a sequential decision problem where the holder of a short position in a call option seeks for the best sequence of actions δ that minimizes the risk associated with the hedging error

$$\xi_T^{\delta} = \max(S_T - K, 0) - V_T^{\delta}(V_0), \tag{2}$$

where K is the call option strike price. The hedging problem is formulated as

$$\delta^* = \arg\min_{\delta} \rho\left(\xi_T^{\delta}\right),\tag{3}$$

 $^{^1}$ The Python code which allows replicating the numerical experiments from this paper can be found at $\frac{1}{\text{https://github.com/cpmendoza/DeepHedging}}$ StatisticalArbitrage.git.

where ρ is the risk measure used by the hedging agent to quantify risk. In this paper we consider the Conditional Value-at-Risk (CVaR $_{\alpha}$) defined as $\rho(\xi_T^{\delta}) = \mathbb{E}[\xi_T^{\delta} \mid \xi_T^{\delta} \geq \text{VaR}_{\alpha}(\xi_T^{\delta})]$, where $\alpha \in (0,1)$ and $\text{VaR}_{\alpha}(\xi_T^{\delta})$ is the Value-at-Risk defined as $\text{VaR}_{\alpha}(\xi_T^{\delta}) = \min_c \{c : \mathbb{P}(\xi_T^{\delta} \leq c) \geq \alpha\}$.

Each time-t action δ_{t+1} is a feedback-type decision, being a function of the information currently available on the market: $\delta_{t+1} = \tilde{\delta}(X_t)$ for some function $\tilde{\delta}$ of the state variable vector X_t .

3 Hedging strategies

3.1 Deep hedging

The deep hedging (DH) framework, introduced by Buehler et al. (2019), provides a solution to the hedging problem (3) by leveraging reinforcement learning (RL) techniques. The DH policy $\tilde{\delta}$ is approximated with a neural network δ_{θ}^{DH} with parameters θ , which returns a hedging position δ_{t+1} when provided with time-t input features X_t . The objective function to be minimized is thus

$$\mathcal{O}(\theta) = \rho \left(\xi_T^{\delta_{\theta}^{DH}} \right).$$
(4)

The neural network is optimized with the Mini-batch Stochastic Gradient Descent method (MSGD). This training procedure relies on updating iteratively all the trainable parameters of the network based on the recursive equation

$$\theta_{j+1} = \theta_j - \eta_j \nabla_{\theta} \hat{\mathcal{O}}(\theta_j), \tag{5}$$

where θ_j is the set of parameters obtained after iteration j, η_j is the learning rate (step size) which determines the magnitude of change in parameters on each time step, ∇_{θ} is the gradient operator with respect to θ and $\hat{\mathcal{O}}$ is the Monte Carlo estimate of the objective function (4) computed on a mini-batch. Automatic differentiation packages are used to compute the gradient of $\hat{\mathcal{O}}$. Additional details are provided in the appendix.

For the neural network, we employ a fully-connected Feedforward Neural Network (FFNN) architecture with four hidden layers of width 56 using a ReLU activation function. The output

FFNN layer, which maps the output of the hidden layer Z into the position in the underlying asset position δ_{t+1}^{DH} , is equipped with a dynamic upper bound on the activation function to preclude excessive leverage. Indeed, agents have finite borrowing capacity in practice. We impose that the time-t cash account value ϕ_t satisfies $\phi_t \geq -B$ for all t and for some threshold B > 0. This is achieved by setting the final output layer activation to

$$f(Z,t) = \min\left(Z, (V_t + B)/S_t\right),\tag{6}$$

which ensures that the cash amount borrowed in the hedging portfolio remains below B > 0 (see François et al. (2024)).

Agents are trained on training sets of 400,000 independent simulated paths with mini-batch size of 1,000 and a learning rate of 0.0005 that is progressively adapted with the ADAM (Kingma and Ba, 2014) optimization algorithm. The training procedure is implemented in Python, using Tensorflow and considering the Glorot and Bengio (2010) random initialization of the initial parameters of the neural network. Numerical results are obtained from test sets of 100,000 independent paths.

3.2 Delta hedging

Delta hedging aims to reduce the risk associated with price movements of an underlying asset by adjusting the hedging portfolio positions in the underlying asset based on the sensitivity (Δ) of the option price to changes in the price of the underlying asset. Specifically, the time-t position in the underlying asset is defined as the time-t sensitivity Δ_t , which is the partial derivative of the time-t option price with respect to the underlying asset value.

3.3 Statistical arbitrage

Statistical arbitrage strategies, also known as "good deals" according to the terminology of Cochrane and Saa-Requejo (2000), are profit-seeking trading strategies that capitalize on statistical anomalies in the market. Bondarenko (2003) defines statistical arbitrage as a trading strategy that makes a profit on average without requiring any initial capital investment. Assa and Karai (2013) extend this definition to offer a more nuanced and comprehensive evaluation, ensuring

that the trading strategy is not only profitable on average, but also resilient in terms of risk management. As in Assa and Karai (2013), we say that δ is a statistical arbitrage opportunity if

$$\rho\left(-V_T^{\delta}(0)\right) < 0,\tag{7}$$

which entails a trading strategy with zero initial investment that is deemed strictly less risky than a null investment according to risk measure ρ . Such definition is also in line with that of Buehler et al. (2021), who focus on the case of the entropy risk measure.

Horikawa and Nakagawa (2024) claim that in a complete market model that admits statistical arbitrage, the difference between the deep hedging and the delta hedging strategies denoted by

$$\delta^{-} = \delta^{DH} - \Delta, \tag{8}$$

is a statistical arbitrage strategy according to risk measure ρ . We wish to further extend their study and examine if the trading strategy δ^- behaves like a statistical arbitrage in a GARCH-based market model. In other words, we investigate whether the deep hedging approach typically incorporates a speculative arbitrage-like component aimed at exploiting the structure of the risk measure considered.

4 Numerical study

4.1 Stochastic market dynamics

We consider market dynamics based on a GARCH process to represent the underlying asset logreturns. The GJR-GARCH(1,1) model introduced by Glosten et al. (1993) captures time-varying volatility and accounts for the leverage effect. For t = 1, ..., T, log-returns in the model follow

$$R_t = \mu + \sigma_t \epsilon_t, \quad \sigma_{t+1}^2 = \omega + \sigma_t^2 (\alpha + \gamma \mathbb{1}_{\{\epsilon_t < 0\}}) \epsilon_t^2 + \beta \sigma_t^2, \tag{9}$$

where $\mu, \gamma \in \mathbb{R}$, ω, α, β are positive, $\mathbb{1}_A$ is the binary dummy variable indicating if event A occurs and $\{\epsilon_t\}_{t=1}^T$ are independent standard normal random variables. Parameter estimates are

obtained through maximum likelihood on a daily time series of the S&P 500 index extending from January 4, 2016, to December 31, 2020. Estimated parameters are $\mu = 0.06\%$, $\omega = 0.01\%$, $\alpha = 0.11$, $\gamma = 0.20$ and $\beta = 0.78$. Furthermore, in all experiments, the annualized continuously compounded risk-free rate and dividend yield are assumed to be constant with values set to r = 2.66% and q = 1.77%, respectively.

The initial option price is computed using Monte Carlo simulation based on the risk-neutral valuation formula

$$Call_0 = e^{-rT\Lambda} \mathbb{E}^{\mathbb{Q}}[\max(S_T - K, 0)], \tag{10}$$

where \mathbb{Q} is a risk-neutral measure.²

The call option delta is also calculated through Monte-Carlo simulation based on the relationship

$$\Delta_t = e^{-r\tau\Lambda} \mathbb{E}^{\mathbb{Q}} \left[\frac{S_{t+\tau}}{S_t} \mathbb{1}_{\{S_{t+\tau} > K\}} \mid \mathcal{F}_t \right]$$
(11)

where $\tau = T - t$ and \mathcal{F}_t represents the available information at time t, i.e., that being generated by the state X_t .

In this model the state space considered for the RL approach is represented by the vector $X_t = (V_t^{\delta}, \log(S_t), \sigma_{t+1}, \tau).$

4.2 Comparative analysis of deep hedging and delta hedging strategies

In this section, we study the relationship between delta hedging and deep hedging. More specifically, we investigate whether the difference between both strategies represents a speculative overlay reminiscent of a statistical arbitrage. The comparison is conducted by hedging at-themoney (ATM) European call options, with $S_0 = K = 100$. The option maturity is T = 63 days. The leverage constraint is B = 100.

$$R_t = (r - q)\Lambda - \frac{\sigma_t^2}{2} + \sigma_t \tilde{\epsilon}_t, \quad \sigma_{t+1}^2 = \omega + \sigma_t^2 (\alpha + \gamma \mathbb{1}_{\{\tilde{\epsilon}_t - \eta_t < 0\}}) (\tilde{\epsilon}_t - \eta_t)^2 + \beta \sigma_t^2,$$

where $\eta_t = (\mu - (r - q)\Lambda + \sigma_t^2/2)/\sigma_t$ and $\{\tilde{\epsilon_t}\}_{t=1}^T$ are independent standard normal random variables under \mathbb{Q} .

 $^{^2}$ The $\mathbb Q$ risk-neutral dynamics of the GARCH model are defined by the following equations:

Table 1 presents the hedging performance of the deep hedging agents trained with the CVaR_{α} risk measure. We consider the following confidence levels α : 1%, 5%, 10%, 20%, 50%, 85%, 90%, and 95%. High values of α only put weight on the most adverse outcomes and entail focusing purely on risk reduction. Conversely, low values for α both penalize losses and reward gains, which leads to seeking risk-reward trade-offs. As such, the CVaR_{α} with a low confidence level does not align with the objective of limiting the variability of the hedging error. Since the CVaR_{α} is an increasing function of α , there are more statistical arbitrage strategies becoming available as α decreases.

Columns labeled "Base strategies" display the risk measure applied to the hedging error for the deep hedging strategy, and the difference between the risk provided by deep hedging and that of delta hedging. Columns labeled "Difference strategy" provide statistics (hedging error risk and expectation of net cash flow) of the trading strategy δ^- representing the differential position between deep hedging and delta hedging. Since such strategy reflects a long position on the deep hedge and a short position on the delta hedge, the option payoffs from the two (long and short) positions cancel out. Hence, performance is assessed by looking at $V_T^{\delta^-}(0)$ rather than $\xi_T^{\delta^-}$.

Table 1: Performance assessment for deep hedging, delta hedging and their difference over a short position on an ATM call option with maturity T = 63 days.

	Base strategies		Difference strategy	
Metric	$\rho(\xi_T^{\delta^{DH}})$	$\rho(\xi_T^{\delta^{DH}}) - \rho(\xi_T^{\Delta})$	$\rho(-V_T^{\delta^-}(0))$	$\mathbb{E}[V_T^{\delta^-}(0)]$
$\mathrm{CVaR}_{1\%}$	-1.918	-1.368	-1.306	1.372
$\mathrm{CVaR}_{5\%}$	-1.852	-1.351	-1.167	1.372
$\mathrm{CVaR}_{10\%}$	-1.767	-1.325	-1.038	1.371
$\mathrm{CVaR}_{20\%}$	-1.575	-1.256	-0.807	1.371
$\mathrm{CVaR}_{50\%}$	-0.614	-0.791	0.071	1.370
$\mathrm{CVaR}_{85\%}$	1.505	-0.129	1.208	-0.039
$\mathrm{CVaR}_{90\%}$	2.055	-0.128	1.599	-0.225
$\mathrm{CVaR}_{95\%}$	3.102	-0.121	2.121	-0.369

Results are computed using 100,000 out-of-sample paths. The initial price of the option is 3.16. ξ_T^{δ} is the hedging error for trading strategy δ , with δ^{DH} being deep hedging and Δ being delta hedging. The strategy δ^- uses the underlying asset position defined by the difference between that of the deep hedging and the delta hedging strategies.

For all confidence levels α below 50%, the strategy δ^- exhibits both positive average profitability $\mathbb{E}[V_T^{\delta^-}(0)]$ and a CVaR value $\rho(-V_T^{\delta^-}(0))$ that is negative. This corresponds to a formal statistical arbitrage strategy. Moreover, for the case $\alpha=50\%$, even if the risk measure is positive, it is nevertheless negligible in comparison to expected profits. The strategy δ^- , though not a statistical arbitrage from the definition, exhibits a behavior that is quite similar to a statistical arbitrage. Conversely, difference strategies δ^- using $\alpha \geq 85\%$ clearly do not qualify as statistical arbitrage; the associated risk measure is high and the average profitability is negative.

The distribution of the profit and loss (P&L) for the trading strategy δ^- , which is $V_T^{\delta^-}(0)$, is depicted in Figure 1 for various confidence levels. This confirms that difference strategies associated with low confidence levels α (1%, 10% and 50%) are exactly or similar to statistical arbitrage with very high average profits and a very fat left tail (high extreme loss potential). The deep hedging agent is incorporating a strong speculative element in its trading strategy, which is unsuitable in practice. Conversely, the strategies associated with higher values for α do not

exhibit characteristics of a statistical arbitrage and do not lead to concerns about the suitability of the deep hedging strategy.




Figure 1: P&L distribution of the strategy δ^- .

Distributions are computed using 100,000 out-of-sample paths. The P&L is simply defined by the portfolio value $V_T^{\delta^-}(0)$ at maturity.

We analyze the statistical relationship between deep hedging and delta hedging strategies through
(i) sample Spearman (rank) correlations between underlying asset positions of both strategies,
and (ii) the regression model

$$\delta^{DH} = \kappa_0 + \kappa_1 \Delta + \epsilon, \tag{12}$$

with δ^{DH} and Δ being positions produced by the deep hedging and delta hedging strategies, respectively. Metrics (regressions and correlations) are computed across all rebalancing points of all paths in the test sets.

Table 2 provides the Spearman correlation coefficient ϱ , which evaluates monotonic relationships between strategies, and the coefficient of determination R^2 , which measures the strength of the linear association between the strategies. These metrics are computed for the various CVaR confidence levels.

Table 2: Statistical relationships between positions of delta hedging and deep hedging.

	Statistics		
Metric	ρ	R^2	
CVaR _{1%}	-0.270	0.003	
$\mathrm{CVaR}_{5\%}$	-0.271	0.003	
$\mathrm{CVaR}_{10\%}$	-0.272	0.003	
$\mathrm{CVaR}_{20\%}$	-0.273	0.003	
$\mathrm{CVaR}_{50\%}$	-0.273	0.003	
$\mathrm{CVaR}_{85\%}$	0.939	0.773	
${\rm CVaR}_{90\%}$	0.963	0.816	
CVaR _{95%}	0.969	0.808	

Results are for a short position on an ATM call option with a maturity of N=63 days. They are computed using 100,000 out-of-sample paths. The metric ϱ denotes the (unconditional) Spearman correlation between underlying asset positions of the delta hedging strategy and the deep hedging strategy across all rebalancing days while R^2 represents the R^2 statistic obtained after regressing deep hedging positions onto delta hedging positions.

Results presented in Table 2 show strong monotonic and linear association between deep and delta hedging positions for high confidence levels $\alpha = 85\%$, 90% or 95%. The deep hedging strategies can therefore be seen as alterations of the delta hedging procedure that improve hedging performance. Conversely, for low confidence levels (50% or below), deep hedging positions seem completely unrelated to delta hedging positions, indicating that the agent has mostly abandoned its hedging objective and is rather attempting to speculate or conduct statistical arbitrage.

5 Conclusion

Consider the trading strategy whose underlying asset positions correspond to the difference between these of deep hedging and delta hedging. What if there exist market models under which such strategy is a statistical arbitrage? This would raise concerns about the suitability of deep hedging procedures, as it raises the possibility that typical deep hedging strategies could consist of conventional hedging strategies that are enhanced with speculative overlays which are unrelated to hedging.

Our study shows that these concerns can be mitigated under GARCH-based market models; if the risk measure considered in the hedging optimization problem does not sufficiently penalize losses relative to rewards provided for gains, the deep hedging strategy attaches a statistical arbitrage strategy overlay to the delta hedging strategy. Nevertheless, when using a proper risk measure (the CVaR with sufficiently high α in our case) within the optimization problem, the difference between deep hedging and delta hedging does not exhibit statistical arbitrage-like behavior and cannot be interpreted as a speculative strategy reaping profits while exploiting blind spots of the chosen risk measure.

The two main conclusions from this study are therefore that (i) the objective function of the deep hedging problem must be carefully selected to prevent the hedging agent from abandoning its hedging objective and pursuing speculative behavior, and (ii) deep hedging can soundly achieve its hedging objectives when provided with a suitable risk measure.

References

- Assa, H. and Karai, K. M. (2013). Hedging, Pareto optimality, and good deals. *Journal of Optimization Theory and Applications*, 157:900–917.
- Bondarenko, O. (2003). Statistical arbitrage and securities prices. The Review of Financial Studies, 16(3):875–919.
- Buehler, H., Gonon, L., Teichmann, J., and Wood, B. (2019). Deep hedging. *Quantitative Finance*, 19(8):1271–1291.
- Buehler, H., Murray, P., Pakkanen, M. S., and Wood, B. (2021). Deep hedging: Learning risk-neutral implied volatility dynamics. arXiv preprint arXiv:2103.11948.
- Cao, J., Chen, J., Farghadani, S., Hull, J., Poulos, Z., Wang, Z., and Yuan, J. (2023). Gamma and vega hedging using deep distributional reinforcement learning. Frontiers in Artificial Intelligence, 6:1129370.
- Cao, J., Chen, J., Hull, J., and Poulos, Z. (2020). Deep hedging of derivatives using reinforcement learning. The Journal of Financial Data Science.

- Carbonneau, A. (2021). Deep hedging of long-term financial derivatives. *Insurance: Mathematics and Economics*, 99:327–340.
- Carbonneau, A. and Godin, F. (2021). Equal risk pricing of derivatives with deep hedging.

 Quantitative Finance, 21(4):593–608.
- Carbonneau, A. and Godin, F. (2023). Deep equal risk pricing of financial derivatives with non-translation invariant risk measures. *Risks*, 11(8):140.
- Cochrane, J. H. and Saa-Requejo, J. (2000). Beyond arbitrage: Good-deal asset price bounds in incomplete markets. *Journal of Political Economy*, 108(1):79–119.
- Du, J., Jin, M., Kolm, P. N., Ritter, G., Wang, Y., and Zhang, B. (2020). Deep reinforcement learning for option replication and hedging. *The Journal of Financial Data Science*, 2(4):44–57.
- François, P., Gauthier, G., Godin, F., and Pérez Mendoza, C. O. (2024). Enhancing deep hedging of options with implied volatility surface feedback information. *Working paper*.
- Glorot, X. and Bengio, Y. (2010). Understanding the difficulty of training deep feedforward neural networks. In *Proceedings of the thirteenth international conference on artificial intelligence and statistics*, pages 249–256. JMLR Workshop and Conference Proceedings.
- Glosten, L. R., Jagannathan, R., and Runkle, D. E. (1993). On the relation between the expected value and the volatility of the nominal excess return on stocks. *The Journal of Finance*, 48(5):1779–1801.
- Goodfellow, I., Bengio, Y., and Courville, A. (2016). Deep learning. MIT press.
- Halperin, I. (2019). The QLBS Q-learner goes NuQLear: Fitted Q iteration, inverse RL, and option portfolios. *Quantitative Finance*, 19(9):1543–1553.
- Hambly, B., Xu, R., and Yang, H. (2023). Recent advances in reinforcement learning in finance.

 Mathematical Finance, 33(3):437–503.
- Horikawa, H. and Nakagawa, K. (2024). Relationship between deep hedging and delta hedging: Leveraging a statistical arbitrage strategy. *Finance Research Letters*, page 105101.

- Horvath, B., Teichmann, J., and Žurič, Ž. (2021). Deep hedging under rough volatility. Risks, 9(7):138.
- Imaki, S., Imajo, K., Ito, K., Minami, K., and Nakagawa, K. (2021). No-transaction band network:

 A neural network architecture for efficient deep hedging. arXiv preprint arXiv:2103.01775.
- Kingma, D. P. and Ba, J. (2014). Adam: A method for stochastic optimization. arXiv preprint arXiv:1412.6980.
- Lütkebohmert, E., Schmidt, T., and Sester, J. (2022). Robust deep hedging. *Quantitative Finance*, 22(8):1465–1480.
- Marzban, S., Delage, E., and Li, J. Y.-M. (2023). Deep reinforcement learning for option pricing and hedging under dynamic expectile risk measures. *Quantitative Finance*, 23(10):1411–1430.
- Mikkilä, O. and Kanniainen, J. (2023). Empirical deep hedging. Quantitative Finance, 23(1):111–122.
- Neagu, A., Godin, F., Simard, C., Kosseim, L., et al. (2024). Deep hedging with market impact.

 To appear in proceedings of the 37th Canadian Conference on Artificial Intelligence (CAIAC 2024).
- Pickard, R. and Lawryshyn, Y. (2023). Deep reinforcement learning for dynamic stock option hedging: A review. *Mathematics*, 11(24):4943.
- Raj, S., Kerenidis, I., Shekhar, A., Wood, B., Dee, J., Chakrabarti, S., Chen, R., Herman, D., Hu, S., Minssen, P., et al. (2023). Quantum deep hedging. *Quantum*, 7:1191.
- Wu, D. and Jaimungal, S. (2023). Robust risk-aware option hedging. *Applied Mathematical Finance*, 30(3):153–174.

Appendix: Details for the MSGD training approach

The MSGD method estimates the penalty function $\mathcal{O}(\theta)$, which is typically unknown, through small samples of the hedging error called batches. Let $\mathbb{B}_j = \{\xi_{T,i}^{\delta_{\theta_j}^{DH}}\}_{i=1}^{N_{\text{batch}}}$ be the *j*-th batch where

 $\xi_{T,i}^{\delta_{j}^{DH}}$ denotes the hedging error of the *i*-th simulated path in the *j*-th batch defined as

$$\xi_{T,i}^{\delta_{\theta_{j}}^{DH}} = \max(S_{T,ij} - K, 0) - V_{T,i}^{\delta_{\theta_{j}}^{DH}}(V_{0}),$$

where $S_{T,ij}$ is the price of the underlying asset at time T in the i-th simulated path, and $V_{T,i}^{\delta_{\theta_j}^{DH}}$ is the terminal value of the hedging strategy for that path when $\theta = \theta_j$. The penalty function estimation for the batch \mathbb{B} is

$$\hat{C}^{(\text{CVaR})}(\theta_j, \mathbb{B}_j) = \widehat{\text{VaR}}_{\alpha}(\mathbb{B}_j) + \frac{1}{(1-\alpha)N_{\text{batch}}} \sum_{i=1}^{N_{\text{batch}}} \max \left(\xi_{T,i}^{\delta_{\theta_j}^{DH}} - \widehat{\text{VaR}}_{\alpha}(\mathbb{B}_j), 0 \right) ,$$

where $\widehat{\mathrm{VaR}}_{\alpha}(\mathbb{B}_{j}) = \xi_{T,\lceil\alpha \cdot N_{\mathrm{batch}}\rceil}^{\delta_{\theta_{j}}^{DH}}$ is the estimation of the VaR obtained from the ordered sample $\{\xi_{T,[i]}^{\delta_{\theta_{j}}^{DH}}\}_{i=1}^{N_{\mathrm{batch}}}$ and $\lceil \cdot \rceil$ is the ceiling function. These empirical approximations are used to estimate the gradient of the penalty function required in Equation (5).

³Details about gradient of the empirical objective function are provided in Goodfellow et al. (2016).