

# A Convex-Nonconvex Framework for Enhancing Minimization Induced Penalties

Hiroki Kuroda, *Member, IEEE*

**Abstract**—This paper presents a novel framework for nonconvex enhancement of minimization induced (MI) penalties while preserving the overall convexity of associated regularization models. MI penalties enable the adaptation to certain signal structures via minimization, but often underestimate significant components owing to convexity. To overcome this shortcoming, we design a generalized Moreau enhanced minimization induced (GME-MI) penalty by subtracting from the MI penalty its generalized Moreau envelope. While the proposed GME-MI penalty is nonconvex in general, we derive an overall convexity condition for the GME-MI regularized least-squares model. Moreover, we present a proximal splitting algorithm with guaranteed convergence to a globally optimal solution of the GME-MI model under the overall convexity condition. Numerical examples illustrate the effectiveness of the proposed framework.

**Index Terms**—Regularization, minimization induced penalty, convex-nonconvex strategy, Moreau enhancement, optimization.

## I. INTRODUCTION

Regularization is a key technique for estimation problems in signal processing and machine learning. In particular,  $\ell_1$  regularization is widely used for sparse estimation [1]–[5]. While computationally tractable,  $\ell_1$  regularization often underestimates large-magnitude components since the value of the  $\ell_1$  penalty increases with magnitude, owing to convexity. To mitigate this underestimation tendency, nonconvex penalties, e.g., [6], [7] among many others, have been proposed as better approximations of the  $\ell_0$  pseudo-norm that counts the number of nonzero components. However, nonconvex penalties usually introduce undesirable local minima, and their over-reliance on initial values of optimization algorithms is problematic.

Recently, to resolve the aforementioned trade-off, the so-called *convex-nonconvex strategy* [8]–[21], which uses a nonconvex penalty that preserves convexity of the overall regularization model, has garnered significant attention. After the pioneering works of [8], [9] and recent case studies by [10]–[15], several general frameworks [16]–[19] have been developed to remedy the underestimation tendencies of convex *prox-friendly* penalties (i.e., whose proximity operators can be computed with low complexity), including the  $\ell_1$  penalty. More precisely, these frameworks construct nonconvex penalties by subtracting the *generalized Moreau envelopes* of the convex prox-friendly penalties from them, and provide proximal splitting algorithms to compute optimal solutions of the associated regularized least-squares models under overall convexity conditions. Ex-

tensions to handle additional convex constraints have also been studied in [20], [21].

As another important direction for improving regularization, advanced convex penalties have been designed via minimization to incorporate certain signal structures flexibly. We refer to this class of penalties as *minimization induced (MI)* penalties. For example, to resolve the problem of unknown block partition in block-sparse estimation, which is encountered in various applications [22]–[28], the latent optimally partitioned (LOP)- $\ell_2/\ell_1$  penalty [29] has been designed to minimize the mixed  $\ell_2/\ell_1$  norm [30] over the set of candidate block partitions. Another prominent example is the total generalized variation (TGV) penalty [31], which is defined by minimization for computing and penalizing the magnitudes of discontinuous jumps of high-order derivatives. The TGV penalty is a sound high-order extension of the total variation penalty [32] and is widely used in image processing applications [33]–[39]. MI penalties are capable of automatically adapting to certain signal structures; however, analogous to the  $\ell_1$  penalty, they often underestimate the magnitudes of significant components. In particular, the LOP- $\ell_2/\ell_1$  penalty and the TGV penalty tend to underestimate blocks of large-magnitude components and large jumps of derivatives, respectively.

Unfortunately, existing convex-nonconvex frameworks cannot be directly used to remedy the underestimation tendencies of MI penalties. Even the most general frameworks [17], [19], [21] rely on the assumption that the convex penalty to be enhanced can be decomposed into a sum of linearly-involved prox-friendly functions. This assumption does not hold for MI penalties because the involved minimization makes their proximity operators difficult to compute. Thus, a major question arises: Can we remedy the underestimation tendencies of MI penalties while keeping computation of optimal solutions of associated regularization models tractable?

To address this question, this paper presents a novel convex-nonconvex framework to enhance MI penalties. The main contributions of this paper are summarized as follows.

- 1) After introducing a class of MI penalties defined by minimization of *seed functions*, we construct *generalized Moreau enhanced minimization induced (GME-MI)* penalties by subtracting from MI penalties their generalized Moreau envelopes. This procedure yields novel nonconvex enhancements of the LOP- $\ell_2/\ell_1$  penalty and the TGV penalty.
- 2) By applying the proposed GME-MI penalty to regularized least-squares estimation, we design the GME-MI model and present its overall convexity condition, where a mild technical assumption is imposed on the seed function. We prove that this assumption automatically

This work was supported in part by the Japan Society for the Promotion of Science under Grants-in-Aid 21K17827, 21H01592, and 24H00272.

The author is with the Department of Information and Management Systems Engineering, Nagaoka University of Technology, Niigata 940-2188, Japan (e-mail: kuroda@vos.nagaokaut.ac.jp).

holds for the seed functions used to define the LOP- $\ell_2/\ell_1$  penalty and the TGV penalty.

- 3) We present a proximal splitting algorithm which is guaranteed to converge to a globally optimal solution of the proposed GME-MI model under the overall convexity condition. We derive the proposed algorithm by carefully designing an averaged nonexpansive operator that characterizes the solution set of the GME-MI model. The proposed algorithm is composed only of simple operations: matrix-vector multiplication, the proximity operators of component functions of the seed function, and the projection onto an additional convex constraint set. All the assumed conditions are shown to be valid in the scenarios for enhancement of the LOP- $\ell_2/\ell_1$  penalty and the TGV penalty.

The remainder of this paper is organized as follows. Mathematical preliminaries are presented in Section II. In Section III, we design the GME-MI model and derive its overall convexity condition. In Section IV, we present the proposed algorithm for computing an optimal solution of the GME-MI model. Numerical examples for the enhancement scenarios of the LOP- $\ell_2/\ell_1$  and TGV penalties are presented in Section V. Finally, the paper is concluded in Section VI.

A preliminary short version of this paper is to be presented at a conference [40].

## II. PRELIMINARIES

### A. Notations

Let  $\mathbb{N}$ ,  $\mathbb{R}$ ,  $\mathbb{R}_+$ , and  $\mathbb{R}_{++}$  denote the sets of nonnegative integers, real numbers, nonnegative real numbers, and positive real numbers, respectively. For  $p \geq 1$ , the  $\ell_p$  norm of  $\mathbf{x} \in \mathbb{R}^n$  is defined by  $\|\mathbf{x}\|_p := (\sum_{i=1}^n |x_i|^p)^{1/p}$ . The cardinality of a set  $S$  is denoted by  $|S|$ . For  $\mathbf{x} \in \mathbb{R}^n$  and  $\mathcal{I} \subset \{1, \dots, n\}$ , the subvector of  $\mathbf{x}$  indexed by  $\mathcal{I}$  is denoted by  $\mathbf{x}_{\mathcal{I}} := (x_i)_{i \in \mathcal{I}} \in \mathbb{R}^{|\mathcal{I}|}$ . Let  $\mathcal{G} = (\mathcal{I}_k)_{k=1}^m$  be a partition of  $\{1, \dots, n\}$ , i.e.,  $\bigcup_{k=1}^m \mathcal{I}_k = \{1, \dots, n\}$ ,  $\mathcal{I}_k \neq \emptyset$  ( $k = 1, \dots, m$ ), and  $\mathcal{I}_{k_1} \cap \mathcal{I}_{k_2} = \emptyset$  ( $k_1 \neq k_2$ ). Then, we define the mixed  $\ell_2/\ell_1$  norm of  $\mathbf{x} \in \mathbb{R}^n$  with  $\mathcal{G}$  by  $\|\mathbf{x}\|_{2,1}^{\mathcal{G}} := \sum_{k=1}^m \sqrt{|\mathcal{I}_k|} \|\mathbf{x}_{\mathcal{I}_k}\|_2$ . For  $\mathbf{L} \in \mathbb{R}^{m \times n}$ , its transpose is denoted by  $\mathbf{L}^T$ . For  $\mathbf{L} \in \mathbb{R}^{m \times n}$  and  $S \subset \mathbb{R}^n$ ,  $\mathbf{L}(S) := \{\mathbf{L}\mathbf{x} \mid \mathbf{x} \in S\}$ . The range space of  $\mathbf{L} \in \mathbb{R}^{m \times n}$  is denoted by  $\text{ran}(\mathbf{L}) := \mathbf{L}(\mathbb{R}^n)$ . We denote the identity matrix by  $\mathbf{I}$  and the zero matrix by  $\mathbf{O}$ .

Throughout this paper, a Euclidean space  $\mathbb{R}^n$  is equipped with the standard inner product  $\langle \cdot, \cdot \rangle$  and the associated norm  $\|\cdot\|$ . For  $S \subset \mathbb{R}^n$ ,  $\text{ri}(S)$  denotes the relative interior of  $S$  (see, e.g., [41, Fact 6.14(i)]). We define the operator norm of  $\mathbf{L} \in \mathbb{R}^{m \times n}$  by  $\|\mathbf{L}\|_{\text{op}} := \max\{\|\mathbf{L}\mathbf{x}\| \mid \mathbf{x} \in \mathbb{R}^n, \|\mathbf{x}\| \leq 1\}$ . The positive definiteness and positive semidefiniteness of a symmetric matrix  $\mathbf{L} \in \mathbb{R}^{n \times n}$  are expressed by  $\mathbf{L} \succ \mathbf{O}$  and  $\mathbf{L} \succeq \mathbf{O}$ , respectively. Unless otherwise stated, we regard the product space  $\mathcal{X} := \times_{k=1}^m \mathbb{R}^{n_k}$  as a Euclidean space with the standard inner product and the associated norm. In part of this paper, with a positive definite matrix  $\mathbf{P} \in \mathbb{R}^{\dim \mathcal{X} \times \dim \mathcal{X}}$ , we use another inner product  $\langle \mathbf{x}, \mathbf{y} \rangle_{\mathbf{P}} := \langle \mathbf{x}, \mathbf{P}\mathbf{y} \rangle$  for  $\mathbf{x}, \mathbf{y} \in \mathcal{X}$ , and denote its associated norm by  $\|\cdot\|_{\mathbf{P}}$ .

### B. Tools in Convex Analysis and Optimization

A function  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$  is said to be proper if its effective domain  $\text{dom}(f) := \{\mathbf{x} \in \mathbb{R}^n \mid f(\mathbf{x}) < \infty\}$  is nonempty, lower semicontinuous if its lower level set  $\{\mathbf{x} \in \mathbb{R}^n \mid f(\mathbf{x}) \leq a\}$  is closed for every  $a \in \mathbb{R}$ , and convex if  $f(\alpha\mathbf{x} + (1-\alpha)\mathbf{y}) \leq \alpha f(\mathbf{x}) + (1-\alpha)f(\mathbf{y})$  for every  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  and  $\alpha \in (0, 1)$ . The set of all proper lower semicontinuous convex functions from  $\mathbb{R}^n$  to  $\mathbb{R} \cup \{\infty\}$  is denoted by  $\Gamma_0(\mathbb{R}^n)$ . A function  $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$  is said to be coercive if  $\lim_{\|\mathbf{x}\| \rightarrow \infty} f(\mathbf{x}) = \infty$ .

The subdifferential of  $f \in \Gamma_0(\mathbb{R}^n)$  at  $\mathbf{x} \in \mathbb{R}^n$  is defined by  $\partial f(\mathbf{x}) := \{\mathbf{u} \in \mathbb{R}^n \mid (\forall \mathbf{y} \in \mathbb{R}^n) \langle \mathbf{y} - \mathbf{x}, \mathbf{u} \rangle + f(\mathbf{x}) \leq f(\mathbf{y})\}$ .

We have the relation  $\mathbf{x}^* \in \text{argmin}_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \Leftrightarrow \mathbf{0} \in \partial f(\mathbf{x}^*)$ . If  $f$  is differentiable at  $\mathbf{x} \in \mathbb{R}^n$  with a gradient  $\nabla f(\mathbf{x}) \in \mathbb{R}^n$ , then  $\partial f(\mathbf{x}) = \{\nabla f(\mathbf{x})\}$ .

**Fact 1** (Subdifferential calculus [41, Theorem 16.47]).

- a) (Sum and chain rule). Let  $f \in \Gamma_0(\mathbb{R}^n)$ ,  $g \in \Gamma_0(\mathbb{R}^m)$ , and  $\mathbf{L} \in \mathbb{R}^{m \times n}$ . Assume  $\mathbf{0} \in \text{ri}[\text{dom}(g) - \mathbf{L}(\text{dom}(f))]$ . Then, for every  $\mathbf{x} \in \mathbb{R}^n$ ,

$$\partial(f + g \circ \mathbf{L})(\mathbf{x}) = \partial f(\mathbf{x}) + \mathbf{L}^T(\partial g(\mathbf{L}\mathbf{x})).$$

- b) (Sum rule). Let  $f \in \Gamma_0(\mathbb{R}^n)$  and  $g \in \Gamma_0(\mathbb{R}^n)$ . Assume  $\text{dom}(g) = \mathbb{R}^n$ . Then, for every  $\mathbf{x} \in \mathbb{R}^n$ ,

$$\partial(f + g)(\mathbf{x}) = \partial f(\mathbf{x}) + \partial g(\mathbf{x}).$$

- c) (Chain rule). Let  $g \in \Gamma_0(\mathbb{R}^m)$  and  $\mathbf{L} \in \mathbb{R}^{m \times n}$ . Assume  $\mathbf{0} \in \text{ri}[\text{dom}(g) - \text{ran}(\mathbf{L})]$ . Then, for every  $\mathbf{x} \in \mathbb{R}^n$ ,

$$\partial(g \circ \mathbf{L})(\mathbf{x}) = \mathbf{L}^T(\partial g(\mathbf{L}\mathbf{x})).$$

The conjugate of  $f \in \Gamma_0(\mathbb{R}^n)$  is defined by

$$f^*: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}: \mathbf{u} \mapsto \sup_{\mathbf{x} \in \mathbb{R}^n} [\langle \mathbf{x}, \mathbf{u} \rangle - f(\mathbf{x})], \quad (1)$$

and  $f^* \in \Gamma_0(\mathbb{R}^n)$  [41, Corollary 13.38]. For any  $f \in \Gamma_0(\mathbb{R}^n)$ ,  $(\partial f)^{-1} = \partial f^*$  [41, Theorem 16.29], i.e.,

$$\mathbf{u} \in \partial f(\mathbf{x}) \Leftrightarrow \mathbf{x} \in \partial f^*(\mathbf{u}). \quad (2)$$

The proximity operator of  $f \in \Gamma_0(\mathbb{R}^n)$  is defined by

$$\text{prox}_f: \mathbb{R}^n \rightarrow \mathbb{R}^n: \mathbf{x} \mapsto \text{argmin}_{\mathbf{y} \in \mathbb{R}^n} \left[ f(\mathbf{y}) + \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|^2 \right].$$

Let  $\text{Id}$  denote the identity operator. For any  $f \in \Gamma_0(\mathbb{R}^n)$ ,  $\text{prox}_f = (\text{Id} + \partial f)^{-1}$  [41, Proposition 16.44], i.e.,

$$\mathbf{p} = \text{prox}_f(\mathbf{x}) \Leftrightarrow \mathbf{x} \in (\text{Id} + \partial f)(\mathbf{p}). \quad (3)$$

We say that  $f \in \Gamma_0(\mathbb{R}^n)$  is prox-friendly if  $\text{prox}_{\gamma f}$  is available as a computational tool for any  $\gamma \in \mathbb{R}_{++}$ . Note that, if  $f \in \Gamma_0(\mathbb{R}^n)$  is prox-friendly, then  $f^*$  is also prox-friendly because the following relation holds for any  $\gamma \in \mathbb{R}_{++}$  and  $\mathbf{x} \in \mathbb{R}^n$  [41, Theorem 14.3(ii)]:

$$\text{prox}_{\gamma f^*}(\mathbf{x}) = \mathbf{x} - \gamma \text{prox}_{\frac{1}{\gamma} f}(\mathbf{x}/\gamma).$$

The indicator function of a nonempty closed convex set  $C \subset$

$\mathbb{R}^n$  is defined by

$$\iota_C: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}: \mathbf{x} \mapsto \begin{cases} 0, & \text{if } \mathbf{x} \in C; \\ \infty, & \text{otherwise,} \end{cases}$$

and  $\iota_C \in \Gamma_0(\mathbb{R}^n)$ . For any  $\gamma \in \mathbb{R}_{++}$ , the proximity operator of  $\gamma\iota_C$  reduces to the projection onto  $C$ , i.e.,

$$\text{prox}_{\gamma\iota_C}(\mathbf{x}) = P_C(\mathbf{x}) := \underset{\mathbf{y} \in C}{\text{argmin}} \|\mathbf{x} - \mathbf{y}\|.$$

### C. Fixed Point Theory of Nonexpansive Operators

Let  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathcal{H}}, \|\cdot\|_{\mathcal{H}})$  be a finite-dimensional real Hilbert space. The fixed point set of an operator  $T: \mathcal{H} \rightarrow \mathcal{H}$  is denoted by  $\text{Fix}(T) := \{\mathbf{x} \in \mathcal{H} \mid T(\mathbf{x}) = \mathbf{x}\}$ . An operator  $T: \mathcal{H} \rightarrow \mathcal{H}$  is called nonexpansive if

$$(\forall \mathbf{x}, \mathbf{y} \in \mathcal{H}) \quad \|T(\mathbf{x}) - T(\mathbf{y})\|_{\mathcal{H}} \leq \|\mathbf{x} - \mathbf{y}\|_{\mathcal{H}}.$$

A nonexpansive operator  $T: \mathcal{H} \rightarrow \mathcal{H}$  is  $\alpha$ -averaged if there exist  $\alpha \in (0, 1)$  and a nonexpansive operator  $R: \mathcal{H} \rightarrow \mathcal{H}$  such that  $T = (1 - \alpha)\text{Id} + \alpha R$ , where  $\text{Id}$  is the identity operator.

**Fact 2** (Composition of averaged operators [41, Proposition 4.44]). Let  $T_i: \mathcal{H} \rightarrow \mathcal{H}$  be  $\alpha_i$ -averaged nonexpansive for each  $i = 1, 2$ . Then,  $T_1 \circ T_2$  is  $\alpha$ -averaged nonexpansive with  $\alpha = (\alpha_1 + \alpha_2 - 2\alpha_1\alpha_2)/(1 - \alpha_1\alpha_2) \in (0, 1)$ .

**Fact 3** (Krasnosel'skiĭ-Mann Iteration [41, Theorem 5.15]). Let  $T: \mathcal{H} \rightarrow \mathcal{H}$  be a nonexpansive operator with  $\text{Fix}(T) \neq \emptyset$ . For any  $(\mu_k)_{k \in \mathbb{N}} \subset [0, 1]$  satisfying  $\sum_{k \in \mathbb{N}} \mu_k(1 - \mu_k) = \infty$  and an initial point  $\mathbf{x}_0 \in \mathcal{H}$ ,  $(\mathbf{x}_k)_{k \in \mathbb{N}}$  generated by

$$(\forall k \in \mathbb{N}) \quad \mathbf{x}_{k+1} = (1 - \mu_k)\mathbf{x}_k + \mu_k T(\mathbf{x}_k)$$

converges<sup>1</sup> to a point in  $\text{Fix}(T)$ . In particular, if  $T$  is  $\alpha$ -averaged nonexpansive,  $(\mathbf{x}_k)_{k \in \mathbb{N}}$  generated by

$$(\forall k \in \mathbb{N}) \quad \mathbf{x}_{k+1} = T(\mathbf{x}_k)$$

converges to a point in  $\text{Fix}(T)$ .

A set-valued operator  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is called monotone if

$$(\forall \mathbf{x}, \mathbf{y} \in \mathcal{H})(\forall \mathbf{u} \in A(\mathbf{x}), \forall \mathbf{v} \in A(\mathbf{y})) \quad \langle \mathbf{x} - \mathbf{y}, \mathbf{u} - \mathbf{v} \rangle_{\mathcal{H}} \geq 0.$$

A monotone operator  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is maximally monotone if there exists no monotone operator whose graph strictly contains the graph of  $A$ . A set-valued operator  $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is maximally monotone if and only if its resolvent  $(\text{Id} + A)^{-1}: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is single-valued and  $\frac{1}{2}$ -averaged nonexpansive [41, Proposition 23.10]. An important example of a maximally monotone operator is the subdifferential  $\partial f$  of  $f \in \Gamma_0(\mathcal{H})$  [41, Theorem 20.25]. The proximity operator  $\text{prox}_f$  is the resolvent of  $\partial f$  (see (3)), and thus is  $\frac{1}{2}$ -averaged nonexpansive.

## III. DESIGN OF GME-MI MODEL

Consider the estimation of an original signal  $\mathbf{x}_{\text{org}} \in \mathcal{C}$  from the observed vector  $\mathbf{y} := \mathbf{A}\mathbf{x}_{\text{org}} + \boldsymbol{\varepsilon} \in \mathbb{R}^d$ , where  $\mathcal{C} \subset \mathbb{R}^n$  is a known nonempty closed convex constraint set,  $\mathbf{A} \in \mathbb{R}^{d \times n}$  is a known measurement matrix, and  $\boldsymbol{\varepsilon} \in \mathbb{R}^d$  is a noise vector.

<sup>1</sup>Since this paper focuses on finite-dimensional real Hilbert spaces, weak convergence and strong convergence are equivalent.




Fig. 1. Overview of the proposed framework.

As outlined in Fig. 1, we design the GME-MI model as follows. In Section III-A, we introduce a class of MI penalties, which covers the important existing examples: the LOP- $\ell_2/\ell_1$  and TGV penalties. Then, in Section III-B, to remedy the underestimation tendencies of MI penalties, we design the GME-MI penalties as differences between MI penalties and their generalized Moreau envelopes. Finally, in Section III-C, we design the GME-MI model by applying the GME-MI penalty to regularized least-squares estimation, and derive its overall convexity condition.

### A. Class of MI Penalties

Before defining the MI penalty function for  $\mathbf{u} = \mathbf{L}\mathbf{x} \in \mathbb{R}^m$  with a known matrix  $\mathbf{L} \in \mathbb{R}^{m \times n}$ , we introduce a *seed function*  $\varphi \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^l)$  satisfying the following mild condition.

**Assumption 1.** For every  $\mathbf{u} \in \mathbb{R}^m$ ,  $\varphi(\mathbf{u}, \cdot)$  is proper and coercive.

**Lemma 1.** For every  $\mathbf{u} \in \mathbb{R}^m$ ,  $\text{argmin}_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma}) \neq \emptyset$ .

*Proof.* The properness of  $\varphi(\mathbf{u}, \cdot)$  and  $\varphi \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^l)$  imply  $\varphi(\mathbf{u}, \cdot) \in \Gamma_0(\mathbb{R}^l)$ , and thus its coercivity yields the claim by [41, Proposition 11.15].  $\square$

**Definition 1.** Given a seed  $\varphi \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^l)$  satisfying Assumption 1, we define the MI penalty by

$$\psi: \mathbb{R}^m \rightarrow \mathbb{R}: \mathbf{u} \mapsto \min_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma}). \quad (4)$$

**Lemma 2.**  $\psi \in \Gamma_0(\mathbb{R}^m)$  and  $\text{dom}(\psi) = \mathbb{R}^m$ .

*Proof.* Lemma 1 implies that  $\min_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma})$  is finite for every  $\mathbf{u} \in \mathbb{R}^m$ , and thus  $\text{dom}(\psi) = \mathbb{R}^m$  holds by (4). Since  $\psi$  is defined by the partial minimization of  $\varphi$ , the convexity of  $\psi$  follows from the convexity of  $\varphi$  by [41, Proposition 8.35]. The convexity of  $\psi$  and  $\text{dom}(\psi) = \mathbb{R}^m$  imply that  $\psi$  is continuous by [41, Corollary 8.40]. Altogether, we have  $\psi \in \Gamma_0(\mathbb{R}^m)$ .  $\square$

The introduced class of MI penalties covers the important existing penalties given in Example 1 below, where Assumption 1 is automatically satisfied (see Appendix A-1).

**Example 1** (Instances of MI penalties).

- (LOP- $\ell_2/\ell_1$ ) Consider the problem of estimating  $\mathbf{x}_{\text{org}}$  that is block-sparse in the range space of  $\mathbf{W} \in \mathbb{R}^{m \times n}$ . To leverage block-sparsity without the knowledge of concrete block partition, the LOP- $\ell_2/\ell_1$  penalty [29]

is designed as a tight convex approximation of the combinatorial penalty that minimizes the mixed  $\ell_2/\ell_1$  norm of  $\mathbf{W}\mathbf{x}$  over the set of possible block partitions. By setting  $\mathbf{L} = \mathbf{W}$ , the LOP- $\ell_2/\ell_1$  penalty is reproduced by (4) for  $\varphi_\alpha^{\text{LOP}} \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^m)$  defined by<sup>2</sup>

$$\varphi_\alpha^{\text{LOP}}(\mathbf{u}, \boldsymbol{\sigma}) := \sum_{i=1}^m h(u_i, \sigma_i) + \iota_{B_1^\alpha}(\mathbf{D}\boldsymbol{\sigma}), \quad (5)$$

where  $h \in \Gamma_0(\mathbb{R} \times \mathbb{R})$  is the *perspective* [42] of  $\frac{1}{2}u^2 + \frac{1}{2}$ ,  $\iota_{B_1^\alpha} \in \Gamma_0(\mathbb{R}^p)$  is the indicator function of the  $\ell_1$  norm ball,  $\mathbf{D} \in \mathbb{R}^{p \times m}$  is the first-order discrete difference operator, i.e.,

$$h(u, \sigma) := \begin{cases} \frac{u^2}{2\sigma} + \frac{\sigma}{2}, & \text{if } \sigma > 0; \\ 0, & \text{if } u = 0 \text{ and } \sigma = 0; \\ \infty, & \text{otherwise,} \end{cases} \quad (6)$$

$$\iota_{B_1^\alpha}(\boldsymbol{\xi}) := \begin{cases} 0, & \text{if } \|\boldsymbol{\xi}\|_1 \leq \alpha; \\ \infty, & \text{otherwise,} \end{cases} \quad (7)$$

$$\mathbf{D}\boldsymbol{\sigma} := (\sigma_{i_1} - \sigma_{i_2})_{(i_1, i_2) \in \mathcal{I}},$$

$\mathcal{I}$  is the set of indices of neighboring pairs,  $p = |\mathcal{I}|$ , and  $\alpha \in \mathbb{R}_+$  is a tuning parameter related to the number of blocks. See [29, Section II] for how block partition is optimized with the latent vector  $\boldsymbol{\sigma}$  using the minimization (4) for  $\varphi = \varphi_\alpha^{\text{LOP}}$ .

- b) (TGV) Consider the problem of estimating piecewise smooth  $\mathbf{x}_{\text{org}}$ . The TGV penalty [31] is introduced to leverage piecewise smoothness by penalizing the magnitudes of discontinuous jumps of high-order derivatives. Since second-order TGV is commonly used [33]–[39], we focus on it to simplify the presentation. The second-order TGV penalty is defined for  $\mathbf{x} \in \mathbb{R}^n$  by

$$\text{TGV}_\alpha(\mathbf{x}) := \min_{\boldsymbol{\sigma} \in \mathbb{R}^m} \left[ \alpha \|\mathbf{D}\mathbf{x} - \boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1} + (1 - \alpha) \|\tilde{\mathbf{D}}\boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_2} \right],$$

where  $\mathbf{D} \in \mathbb{R}^{m \times n}$  is the first-order discrete difference operator,  $\tilde{\mathbf{D}} \in \mathbb{R}^{p \times m}$  is the *symmetrized version* [31], [38] of  $\mathbf{D}$ ,  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are partitions of  $\{1, \dots, m\}$  and  $\{1, \dots, p\}$ , respectively, and  $\alpha \in (0, 1)$  is a tuning parameter. Examples of  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are provided below.

- i) (Anisotropic TGV) Set  $\mathcal{G}_1 = (\{i\})_{i=1}^m$  and  $\mathcal{G}_2 = (\{i\})_{i=1}^p$ , i.e.,  $\|\cdot\|_{2,1}^{\mathcal{G}_1} = \|\cdot\|_1$  and  $\|\cdot\|_{2,1}^{\mathcal{G}_2} = \|\cdot\|_1$ .
- ii) (Isotropic TGV) For multi-dimensional signals such as images, the isotropic version is often preferred [39], where the groups in  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are designed by collecting the directional differences.

By setting  $\mathbf{L} = \mathbf{D}$ , the TGV penalty is reproduced by (4) for  $\varphi_\alpha^{\text{TGV}} \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^m)$  defined by

$$\varphi_\alpha^{\text{TGV}}(\mathbf{u}, \boldsymbol{\sigma}) := \alpha \|\mathbf{u} - \boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1} + (1 - \alpha) \|\tilde{\mathbf{D}}\boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_2}. \quad (8)$$

See [31, Section 3.2] for how the jumps of the zeroth derivative (i.e., the signal itself) and the first derivative of  $\mathbf{x}$  are computed and penalized using the minimization (4) for  $\varphi = \varphi_\alpha^{\text{TGV}}$ , where  $\mathbf{u} = \mathbf{L}\mathbf{x}$  with  $\mathbf{L} = \mathbf{D}$ .

<sup>2</sup>While extension to complex-valued signals is straightforward, we focus on the real-valued signals to simplify the presentation.




Fig. 2. Values of TGV and GME-TGV penalties for  $\mathbf{u} = \mathbf{D}\mathbf{x}$  are plotted with respect to discontinuous jumps of a piecewise linear signal  $\mathbf{x} \in \mathbb{R}^{50}$  defined by  $x_i = 0$  ( $i = 1, 2, \dots, 25$ ) and  $x_i = s(i - 26) + r$  ( $i = 26, 27, \dots, 50$ ), where the parameters are set to be  $\alpha = 0.2$  and  $\mathbf{B} = \mathbf{I}$ .

As illustrated by Example 1, the MI penalty is able to automatically incorporate a certain signal structure (e.g., blocks and derivatives) by the minimization involved in its definition; however, it often underestimates the magnitudes of significant components because the value of the MI penalty usually increases with magnitude. In particular, the LOP- $\ell_2/\ell_1$  penalty tends to underestimate blocks of large-magnitude components because its value increases as the magnitude of components increases, owing to its coercivity [29, Theorem 1]. Similarly, the TGV penalty tends to underestimate large jumps of derivatives because it is a semi-norm [31], i.e., its value increases as the magnitude of discontinuous jumps increases (see Fig. 2(a) for an example).

### B. Design of GME-MI Penalty

To remedy the underestimation tendency of the MI penalty  $\psi$ , we design the GME-MI penalty by subtracting from  $\psi$  its generalized Moreau envelope. Our design is inspired by recent studies [14]–[21] on non-MI penalties (i.e., those defined without minimization) that have demonstrated that subtraction of the generalized Moreau envelopes from the respective penalties mitigates the increase of the penalty values, and thus remedies underestimation tendencies.

**Definition 2.** Based on the MI penalty  $\psi$  in Definition 1, we define the GME-MI penalty  $\Psi_{\mathbf{B}}: \mathbb{R}^m \rightarrow \mathbb{R} \cup \{\infty\}$  by

$$\Psi_{\mathbf{B}}(\mathbf{u}) := \psi(\mathbf{u}) - \inf_{\mathbf{v} \in \mathbb{R}^m} \left[ \psi(\mathbf{v}) + \frac{1}{2} \|\mathbf{B}(\mathbf{u} - \mathbf{v})\|^2 \right], \quad (9)$$

where  $\mathbf{B} \in \mathbb{R}^{q \times m}$  can be tuned to adjust the shape of  $\Psi_{\mathbf{B}}$ .

Applying the construction (9) to the MI penalties considered in Example 1, we derive novel nonconvex enhancements of the LOP- $\ell_2/\ell_1$  and TGV penalties.

**Example 2** (Instances of GME-MI penalties).

- a) (GME-LOP- $\ell_2/\ell_1$ ) Applying (9) to the LOP- $\ell_2/\ell_1$  penalty  $\psi_\alpha^{\text{LOP}}$  defined by (4) for the seed  $\varphi_\alpha^{\text{LOP}}$  in (5), we derive a novel GME-LOP- $\ell_2/\ell_1$  penalty  $\Psi_{\mathbf{B}, \alpha}^{\text{LOP}}$ . While it is difficult to express  $\Psi_{\mathbf{B}, \alpha}^{\text{LOP}}$  in a closed form in general, to illustrate how the underestimation is remedied, we provide an approximate relation with  $\mathbf{B} = \frac{1}{\sqrt{\gamma}} \mathbf{I}$  and  $\gamma \in \mathbb{R}_{++}$  as follows. Since  $\psi_\alpha^{\text{LOP}}$  is an approximation of the mixed  $\ell_2/\ell_1$  norm with the optimized block par-

tion [29], we expect  $\psi_\alpha^{\text{LOP}}(\mathbf{u}) \approx \sum_{k=1}^{K_\alpha^*} \sqrt{|\mathcal{B}_k^*|} \|\mathbf{u}_{\mathcal{B}_k^*}\|_2$ , where  $(\mathcal{B}_k^*)_{k=1}^{K_\alpha^*}$  denotes the optimized block partition. By further assuming a similar approximation  $\psi_\alpha^{\text{LOP}}(\mathbf{v}) \approx \sum_{k=1}^{K_\alpha^*} \sqrt{|\mathcal{B}_k^*|} \|\mathbf{v}_{\mathcal{B}_k^*}\|_2$  in (9), the *block minimax concave penalty* [7], [43] using the optimized block partition is approximately reproduced as a special instance of the GME-LOP- $\ell_2/\ell_1$  penalty, i.e.,

$$\Psi_{\frac{1}{\sqrt{\gamma}}\mathbf{I}, \alpha}^{\text{LOP}}(\mathbf{u}) \approx \sum_{k=1}^{K_\alpha^*} \rho_{\gamma, |\mathcal{B}_k^*|}(\|\mathbf{u}_{\mathcal{B}_k^*}\|_2),$$

$$\rho_{\gamma, b}: \mathbb{R}_+ \rightarrow \mathbb{R}_+: t \mapsto \begin{cases} \sqrt{bt} - \frac{t^2}{2\gamma}, & \text{if } t \leq \gamma\sqrt{b}; \\ \frac{\gamma b}{2}, & \text{otherwise.} \end{cases}$$

Based on this example, as the value of the GME-LOP- $\ell_2/\ell_1$  penalty does not increase once the magnitudes of nonzero blocks exceed certain thresholds, the underestimation of blocks of large-magnitude components is considered to have been resolved (see also numerical examples in Section V-A). Note that, since  $\psi_0^{\text{LOP}}(\mathbf{u}) = \sqrt{m}\|\mathbf{u}\|_2$  and  $\psi_\infty^{\text{LOP}}(\mathbf{u}) = \|\mathbf{u}\|_1$  by [29, Theorem 2], the equalities hold for the coarsest block partition and the finest block partition, which are optimal choices when  $\alpha = 0$  and  $\alpha = \infty$ , respectively:

$$\Psi_{\frac{1}{\sqrt{\gamma}}\mathbf{I}, 0}^{\text{LOP}}(\mathbf{u}) = \rho_{\gamma, m}(\|\mathbf{u}\|_2),$$

$$\Psi_{\frac{1}{\sqrt{\gamma}}\mathbf{I}, \infty}^{\text{LOP}}(\mathbf{u}) = \sum_{i=1}^m \rho_{\gamma, 1}(|u_i|).$$

- b) (GME-TGV) Applying (9) to the TGV penalty  $\psi_\alpha^{\text{TGV}}$  defined by (4) for the seed  $\varphi_\alpha^{\text{TGV}}$  in (8), we derive a novel GME-TGV penalty  $\Psi_{\mathbf{B}, \alpha}^{\text{TGV}}$ . To illustrate how the underestimation is remedied, we present numerically computed values of  $\Psi_{\mathbf{B}, \alpha}^{\text{TGV}}$  in Fig. 2(b). It is clear from Fig. 2(b) that the curve of the GME-TGV penalty becomes flat when the magnitudes of jumps of derivatives exceed certain thresholds, and thus the underestimation of large jumps is considered to have been resolved (see also numerical examples in Section V-B).

### C. Design of GME-MI Model and Its Convexity Condition

We apply the GME-MI penalty to regularized least-squares estimation of  $\mathbf{x}_{\text{org}} \in \mathcal{C}$  from  $\mathbf{y} = \mathbf{A}\mathbf{x}_{\text{org}} + \varepsilon$  as follows.

**Definition 3.** Using the GME-MI penalty  $\Psi_{\mathbf{B}}$  in Definition 2, we define the GME-MI model by

$$\underset{\mathbf{x} \in \mathcal{C}}{\text{minimize}} J(\mathbf{x}) := \frac{1}{2}\|\mathbf{y} - \mathbf{A}\mathbf{x}\|^2 + \lambda\Psi_{\mathbf{B}}(\mathbf{L}\mathbf{x}), \quad (10)$$

where  $\lambda \in \mathbb{R}_{++}$  is the regularization parameter.

While  $\Psi_{\mathbf{B}}$  is nonconvex in general as it is the difference of convex functions, we show that a suitable choice of  $\mathbf{B}$  can ensure convexity of the overall cost function  $J$ .

**Theorem 1.** *Under Assumption 1, we have*

$$J(\mathbf{x}) = \frac{1}{2}\langle \mathbf{x}, \mathbf{Q}\mathbf{x} \rangle - \langle \mathbf{y}, \mathbf{A}\mathbf{x} \rangle + \frac{1}{2}\|\mathbf{y}\|^2 + \lambda\psi(\mathbf{L}\mathbf{x}) + \lambda(\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^*(\mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}), \quad (11)$$

and  $J \in \Gamma_0(\mathbb{R}^n)$  if

$$\mathbf{Q} := \mathbf{A}^\top \mathbf{A} - \lambda \mathbf{L}^\top \mathbf{B}^\top \mathbf{B} \mathbf{L} \succeq \mathbf{O}. \quad (12)$$

*Proof.* By Definitions 2 and 3, since

$$\|\mathbf{B}(\mathbf{L}\mathbf{x} - \mathbf{v})\|^2 = \|\mathbf{B}\mathbf{L}\mathbf{x}\|^2 - 2\langle \mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}, \mathbf{v} \rangle + \|\mathbf{B}\mathbf{v}\|^2,$$

and  $\|\mathbf{B}\mathbf{L}\mathbf{x}\|^2$  is independent of  $\mathbf{v}$ , we obtain

$$J(\mathbf{x}) = \frac{1}{2}\langle \mathbf{x}, \mathbf{Q}\mathbf{x} \rangle - \langle \mathbf{y}, \mathbf{A}\mathbf{x} \rangle + \frac{1}{2}\|\mathbf{y}\|^2 + \lambda\psi(\mathbf{L}\mathbf{x}) - \lambda \inf_{\mathbf{v} \in \mathbb{R}^m} [\psi(\mathbf{v}) + \frac{1}{2}\|\mathbf{B}\mathbf{v}\|^2 - \langle \mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}, \mathbf{v} \rangle],$$

where we use the equality  $\langle \mathbf{x}, \mathbf{Q}\mathbf{x} \rangle = \|\mathbf{A}\mathbf{x}\|^2 - \lambda\|\mathbf{B}\mathbf{L}\mathbf{x}\|^2$  by the definition of  $\mathbf{Q}$  in (12). Moreover, since

$$\begin{aligned} & - \inf_{\mathbf{v} \in \mathbb{R}^m} [\psi(\mathbf{v}) + \frac{1}{2}\|\mathbf{B}\mathbf{v}\|^2 - \langle \mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}, \mathbf{v} \rangle] \\ &= \sup_{\mathbf{v} \in \mathbb{R}^m} [-\psi(\mathbf{v}) - \frac{1}{2}\|\mathbf{B}\mathbf{v}\|^2 + \langle \mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}, \mathbf{v} \rangle] \\ &= (\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^*(\mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}) \end{aligned}$$

due to the definition (1) of the conjugate, we have the expression (11). From  $\psi \in \Gamma_0(\mathbb{R}^m)$  by Lemma 2, we have  $\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2 \in \Gamma_0(\mathbb{R}^m)$ , and thus  $(\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^* \circ \mathbf{B}^\top \mathbf{B}\mathbf{L} \in \Gamma_0(\mathbb{R}^n)$ . Meanwhile,  $\psi \in \Gamma_0(\mathbb{R}^m)$  and  $\text{dom}(\psi) = \mathbb{R}^m$  by Lemma 2 and the condition (12) yield

$$\begin{aligned} & \frac{1}{2}\langle \mathbf{x}, \mathbf{Q}\mathbf{x} \rangle - \langle \mathbf{y}, \mathbf{A}\mathbf{x} \rangle + \frac{1}{2}\|\mathbf{y}\|^2 + \lambda\psi(\mathbf{L}\mathbf{x}) \in \Gamma_0(\mathbb{R}^n), \\ & \text{dom}[\frac{1}{2}\langle \mathbf{x}, \mathbf{Q}\mathbf{x} \rangle - \langle \mathbf{y}, \mathbf{A}\mathbf{x} \rangle + \frac{1}{2}\|\mathbf{y}\|^2 + \lambda\psi(\mathbf{L}\mathbf{x})] = \mathbb{R}^n. \end{aligned} \quad (13)$$

Altogether,  $J \in \Gamma_0(\mathbb{R}^n)$  is guaranteed.  $\square$

**Remark 1** (Design of  $\mathbf{B}$  satisfying (12)). While this paper constructs the novel GME-MI model from the MI penalty  $\psi$  defined via minimization in (4), the condition (12) is common with existing studies [17], [20], [21], [44] on enhancement of linearly involved prox-friendly penalties defined without minimization. This allows using the results of the existing studies to set  $\mathbf{B}$  satisfying (12). In particular, by [44, Theorem 1], we can set  $\mathbf{B}$  for any pair of  $\mathbf{A}$  and  $\mathbf{L}$  in the closed form via LU decomposition (see also Section V for the specific configurations in experiments).

## IV. OPTIMIZATION ALGORITHM FOR GME-MI MODEL

We present the proposed proximal splitting algorithm with guaranteed convergence to a globally optimal solution of the GME-MI model in Definition 3 under the overall convexity condition (12). Although the expression (11) represents the cost function  $J$  in (10) as the sum of convex functions under the condition (12), its minimization is still challenging because it involves non-prox-friendly functions that are difficult to handle:  $\psi$  defined by minimization in (4) and the conjugate of the sum of  $\psi$  and  $\frac{1}{2}\|\mathbf{B} \cdot \|^2$ . As a result, the standard proximal splitting algorithms (see, e.g., [45], [46]) as well as the special algorithms [16]–[21] for GME models of (linearly involved) prox-friendly penalties are not applicable to the GME-MI model. We resolve this challenge by carefully designing a

computable averaged nonexpansive operator  $T_{\text{GME-MI}}$  whose fixed point set characterizes the solution set

$$\mathcal{S} := \operatorname{argmin}_{\mathbf{x} \in \mathcal{C}} J(\mathbf{x}) = \operatorname{argmin}_{\mathbf{x} \in \mathbb{R}^n} [J(\mathbf{x}) + \iota_{\mathcal{C}}(\mathbf{x})]. \quad (14)$$

After introducing technical assumptions and lemmas in Section IV-A, in Section IV-B, we derive  $T_{\text{GME-MI}}$  via careful reformulation of  $\mathbf{x}^* \in \mathcal{S} \Leftrightarrow \mathbf{0} \in \partial(J + \iota_{\mathcal{C}})(\mathbf{x}^*)$  and establish its averaged nonexpansiveness, which yields the convergence of the proposed algorithm based on the Krasnosel'skiĭ-Mann iteration (Fact 3) of  $T_{\text{GME-MI}}$  to an optimal solution in  $\mathcal{S}$ .

#### A. Technical Assumptions and Lemmas

We adopt rather simple assumptions to reduce technical complexity, although these assumptions could be relaxed. We first impose the following condition on the seed function  $\varphi$  and the constraint set  $\mathcal{C}$ .

**Assumption 2.** The seed function  $\varphi$  can be represented as

$$\varphi(\mathbf{u}, \boldsymbol{\sigma}) = f(\mathbf{u}, \boldsymbol{\sigma}) + g(\mathbf{M}\boldsymbol{\sigma}), \quad (15)$$

where  $f \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^l)$  and  $g \in \Gamma_0(\mathbb{R}^p)$  are prox-friendly, and  $\mathbf{M} \in \mathbb{R}^{p \times l}$ . We also assume that  $\iota_{\mathcal{C}}$  is prox-friendly, i.e., the projection  $P_{\mathcal{C}}$  onto  $\mathcal{C}$  is computable.

While the form (15) may seem rather specific, it covers the seed functions used for the important instances of the GME-MI penalty presented in Example 2.

#### Example 3.

- The seed function in (5) for the GME-LOP- $\ell_2/\ell_1$  penalty can be represented in the form of (15) by setting  $f(\mathbf{u}, \boldsymbol{\sigma}) = \sum_{i=1}^m h(u_i, \sigma_i)$ ,  $g = \iota_{B_{\infty}^p}$ , and  $\mathbf{M} = \mathbf{D}$ .
- The seed function in (8) for the GME-TGV penalty can be represented in the form of (15) by setting  $f(\mathbf{u}, \boldsymbol{\sigma}) = \alpha \|\mathbf{u} - \boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1}$ ,  $g = (1 - \alpha) \|\cdot\|_{2,1}^{\mathcal{G}_2}$ , and  $\mathbf{M} = \bar{\mathbf{D}}$ .

Appendix A-2 shows that these  $f$  and  $g$  are prox-friendly. Note that the constraint set  $\mathcal{C}$  is independent of the seed function, and can be chosen based on the application.

To use the formulas of subdifferential calculus (Fact 1) in the derivation of the proposed algorithm, we further adopt the following assumption, which automatically holds for the seed functions used for Example 2 with their representations in Example 3 (see Appendix A-3).

**Assumption 3.** Define

$$\bar{\mathbf{L}} := \begin{bmatrix} \mathbf{L} & \mathbf{O} \\ \mathbf{O} & \mathbf{I} \end{bmatrix} \in \mathbb{R}^{(m+l) \times (n+l)}, \quad (16)$$

$$\bar{\mathbf{M}} := \begin{bmatrix} \mathbf{O} & \mathbf{M} \end{bmatrix} \in \mathbb{R}^{p \times (m+l)}, \quad (17)$$

and assume that the following conditions hold.

- $\varphi$  is bounded below.
- One of the following conditions holds:
  - $(\forall (\mathbf{u}, \boldsymbol{\sigma}) \in \mathbb{R}^m \times \mathbb{R}^l) \quad \varphi(-\mathbf{u}, \boldsymbol{\sigma}) = \varphi(\mathbf{u}, \boldsymbol{\sigma})$ .
  - $(\forall (\mathbf{u}, \boldsymbol{\sigma}) \in \mathbb{R}^m \times \mathbb{R}^l) \quad \varphi(-\mathbf{u}, \boldsymbol{\sigma}) = \varphi(\mathbf{u}, -\boldsymbol{\sigma})$ .
- $\mathbf{0} \in \operatorname{ri}[\operatorname{dom}(\varphi) - \operatorname{ran}(\bar{\mathbf{L}})]$ .
- $\mathbf{0} \in \operatorname{ri}[\operatorname{dom}(g) - \bar{\mathbf{M}}(\operatorname{dom}(f))]$ .

**Lemma 3.** Assumptions 3(i) and (ii) respectively imply

$$(i) \quad \operatorname{dom}\left[(\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^* \circ \mathbf{B}^T \mathbf{B} \mathbf{L}\right] = \mathbb{R}^n,$$

$$(ii) \quad \mathbf{0} \in \operatorname{ri}\left[\operatorname{dom}\left((\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^*\right) - \operatorname{ran}(\mathbf{B}^T \mathbf{B} \mathbf{L})\right].$$

*Proof.* Claim (i) holds because, for any  $\mathbf{x} \in \mathbb{R}^n$ , we have

$$\begin{aligned} & (\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^* (\mathbf{B}^T \mathbf{B} \mathbf{L} \mathbf{x}) \\ &= \sup_{\mathbf{v} \in \mathbb{R}^m} \left[-\psi(\mathbf{v}) - \frac{1}{2}\|\mathbf{B} \mathbf{v}\|^2 + \langle \mathbf{B}^T \mathbf{B} \mathbf{L} \mathbf{x}, \mathbf{v} \rangle\right] \\ &= \sup_{\mathbf{v} \in \mathbb{R}^m} \left[-\psi(\mathbf{v}) - \frac{1}{2}\|\mathbf{B} \mathbf{v} - \mathbf{B} \mathbf{L} \mathbf{x}\|^2 + \frac{1}{2}\|\mathbf{B} \mathbf{L} \mathbf{x}\|^2\right] \\ &\leq \sup_{\mathbf{v} \in \mathbb{R}^m} [-\psi(\mathbf{v})] + \frac{1}{2}\|\mathbf{B} \mathbf{L} \mathbf{x}\|^2 < \infty, \end{aligned}$$

where the last inequality holds because  $\psi$  is bounded below due to Assumption 3(i) and Definition 1.

Next, we prove claim (ii). Assumption 3(ii) and Definition 1 imply  $\psi(-\mathbf{u}) = \psi(\mathbf{u})$  for every  $\mathbf{u} \in \mathbb{R}^m$ . Thus, for any  $\mathbf{w} \in \mathbb{R}^m$ , we have

$$(\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^* (-\mathbf{w}) = (\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^* (\mathbf{w}),$$

which implies that  $\mathcal{A} := \operatorname{dom}\left[(\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^*\right]$  is symmetric, i.e.,  $\mathcal{A} = -\mathcal{A}$ . Since  $(\psi + \frac{1}{2}\|\mathbf{B} \cdot\|^2)^* \in \Gamma_0(\mathbb{R}^m)$  is proved by Theorem 1,  $\mathcal{A}$  is nonempty and convex by [41, Proposition 8.2]. Altogether,  $\mathcal{A} - \operatorname{ran}(\mathbf{B}^T \mathbf{B} \mathbf{L})$  is a nonempty symmetric convex set, and thus claim (ii) holds by [41, Example 6.10].  $\square$

We also use the following lemma, which is a refinement of [41, Proposition 16.59] for our scenario.

**Lemma 4.** For any  $(\mathbf{u}, \boldsymbol{\varsigma}) \in \mathbb{R}^m \times \mathbb{R}^l$  and  $\mathbf{r} \in \mathbb{R}^m$ ,

$$\begin{cases} \mathbf{r} \in \partial\psi(\mathbf{u}) \\ \boldsymbol{\varsigma} \in \operatorname{argmin}_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma}) \Leftrightarrow (\mathbf{r}, \mathbf{0}) \in \partial\varphi(\mathbf{u}, \boldsymbol{\varsigma}). \end{cases} \quad (18)$$

Similarly, for any  $(\mathbf{x}, \boldsymbol{\varsigma}) \in \mathbb{R}^n \times \mathbb{R}^l$  and  $\mathbf{s} \in \mathbb{R}^n$ ,

$$\begin{cases} \mathbf{s} \in \partial(\psi \circ \mathbf{L})(\mathbf{x}) \\ \boldsymbol{\varsigma} \in \operatorname{argmin}_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{L} \mathbf{x}, \boldsymbol{\sigma}) \Leftrightarrow (\mathbf{s}, \mathbf{0}) \in \partial(\varphi \circ \bar{\mathbf{L}})(\mathbf{x}, \boldsymbol{\varsigma}). \end{cases} \quad (19)$$

*Proof.* First, we prove the relation (18).

( $\Rightarrow$ ) The condition  $\boldsymbol{\varsigma} \in \operatorname{argmin}_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma})$  and Definition 1 imply  $\psi(\mathbf{u}) = \varphi(\mathbf{u}, \boldsymbol{\varsigma})$ . Thus, since  $\psi$  is proper by Lemma 2 and  $\varphi \in \Gamma_0(\mathbb{R}^m \times \mathbb{R}^l)$ , the claim follows from [41, Proposition 16.59].

( $\Leftarrow$ ) Suppose  $(\mathbf{r}, \mathbf{0}) \in \partial\varphi(\mathbf{u}, \boldsymbol{\varsigma})$ . Then, for any  $(\mathbf{v}, \boldsymbol{\sigma}) \in \mathbb{R}^m \times \mathbb{R}^l$ , we have

$$\begin{aligned} & \langle (\mathbf{v}, \boldsymbol{\sigma}) - (\mathbf{u}, \boldsymbol{\varsigma}), (\mathbf{r}, \mathbf{0}) \rangle + \varphi(\mathbf{u}, \boldsymbol{\varsigma}) \leq \varphi(\mathbf{v}, \boldsymbol{\sigma}) \\ & \Leftrightarrow \langle \mathbf{v} - \mathbf{u}, \mathbf{r} \rangle + \langle \boldsymbol{\sigma} - \boldsymbol{\varsigma}, \mathbf{0} \rangle + \varphi(\mathbf{u}, \boldsymbol{\varsigma}) \leq \varphi(\mathbf{v}, \boldsymbol{\sigma}) \\ & \Leftrightarrow \langle \mathbf{v} - \mathbf{u}, \mathbf{r} \rangle + \varphi(\mathbf{u}, \boldsymbol{\varsigma}) \leq \varphi(\mathbf{v}, \boldsymbol{\sigma}). \end{aligned}$$

In particular, by setting  $\mathbf{v} = \mathbf{u}$ , we obtain

$$(\forall \boldsymbol{\sigma} \in \mathbb{R}^l) \quad \varphi(\mathbf{u}, \boldsymbol{\varsigma}) \leq \varphi(\mathbf{u}, \boldsymbol{\sigma}),$$

i.e.,  $\boldsymbol{\varsigma} \in \operatorname{argmin}_{\boldsymbol{\sigma} \in \mathbb{R}^l} \varphi(\mathbf{u}, \boldsymbol{\sigma})$ , and thus  $\varphi(\mathbf{u}, \boldsymbol{\varsigma}) = \psi(\mathbf{u})$ . This equality and  $(\mathbf{r}, \mathbf{0}) \in \partial\varphi(\mathbf{u}, \boldsymbol{\varsigma})$  imply  $\mathbf{r} \in \partial\psi(\mathbf{u})$  by [41, Proposition 16.59].

The relation (19) can be derived similarly, since the definition of  $\bar{L}$  in (16) yields  $(\psi \circ L)(x) = \min_{\sigma \in \mathbb{R}^l} \varphi(Lx, \sigma) = \min_{\sigma \in \mathbb{R}^l} (\varphi \circ \bar{L})(x, \sigma)$ .  $\square$

### B. Design of Proposed Algorithm and Its Convergence

Now we derive the averaged nonexpansive operator  $T_{\text{GME-MI}}$  that characterizes the solution set  $\mathcal{S}$  in (14) and establish the convergence of Algorithm 1 to a globally optimal solution of the GME-MI model in Definition 3.

**Theorem 2.** Set  $\kappa > 1$  and  $\gamma_1, \gamma_2, \gamma_3, \gamma_4 \in \mathbb{R}_{++}$  satisfying<sup>3</sup>

$$\begin{cases} (1/\gamma_1)\mathbf{I} - (\kappa/2)\mathbf{A}^\top\mathbf{A} - \lambda\mathbf{L}^\top\mathbf{L} \succ \mathbf{O}, \\ (1/\gamma_2)\mathbf{I} - \mathbf{I} - \mathbf{M}^\top\mathbf{M} \succ \mathbf{O}, \\ (1/\gamma_3) \geq (\kappa/2 + 2/\kappa)\|\mathbf{B}\|_{\text{op}}^2, \\ (1/\gamma_4)\mathbf{I} - \gamma_3\mathbf{M}\mathbf{M}^\top \succ \mathbf{O}. \end{cases} \quad (20)$$

Let  $\mathcal{H} := \mathbb{R}^n \times \mathbb{R}^l \times \mathbb{R}^m \times \mathbb{R}^l \times \mathbb{R}^m \times \mathbb{R}^l \times \mathbb{R}^p \times \mathbb{R}^p$ , and define  $T_{\text{GME-MI}}: \mathcal{H} \rightarrow \mathcal{H}: (x, \sigma, v, \tau, r, \eta, \xi, \zeta) \mapsto (\hat{x}, \hat{\sigma}, \hat{v}, \hat{\tau}, \hat{r}, \hat{\eta}, \hat{\xi}, \hat{\zeta})$  by

$$\begin{aligned} \hat{x} &:= P_{\mathcal{C}}[x - \gamma_1(\mathbf{Q}x - \mathbf{A}^\top\mathbf{y} + \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}v + \lambda\mathbf{L}^\top\mathbf{r})], \\ \hat{\sigma} &:= \sigma - \gamma_2(\eta + \mathbf{M}^\top\xi), \\ (\hat{v}, \hat{\tau}) &:= \text{prox}_{\gamma_3 f}(v + \gamma_3\mathbf{B}^\top\mathbf{B}(\mathbf{L}(2\hat{x} - x) - v), \\ &\quad \tau - \gamma_3\mathbf{M}^\top\zeta), \quad (21) \\ (\hat{r}, \hat{\eta}) &:= \text{prox}_{f^*}(r + \mathbf{L}(2\hat{x} - x), \eta + 2\hat{\sigma} - \sigma), \\ \hat{\xi} &:= \text{prox}_{g^*}(\xi + \mathbf{M}(2\hat{\sigma} - \sigma)), \\ \hat{\zeta} &:= \text{prox}_{\gamma_4 g^*}(\zeta + \gamma_4\mathbf{M}(2\hat{\tau} - \tau)). \end{aligned}$$

Then, under Assumptions 1, 2, and 3 and the condition (12),

$$x^* \in \mathcal{S} \Leftrightarrow \begin{cases} \exists(\sigma^*, v^*, \tau^*, r^*, \eta^*, \xi^*, \zeta^*) \text{ such that} \\ (x^*, \sigma^*, v^*, \tau^*, r^*, \eta^*, \xi^*, \zeta^*) \in \text{Fix}(T_{\text{GME-MI}}), \end{cases} \quad (22)$$

and  $T_{\text{GME-MI}}$  is  $\frac{\kappa}{2\kappa-1}$ -averaged nonexpansive in  $\mathcal{H}$  with  $\langle \cdot, \cdot \rangle_{\mathbf{P}}$  and  $\|\cdot\|_{\mathbf{P}}$ , where  $\mathbf{P} \in \mathbb{R}^{\dim \mathcal{H} \times \dim \mathcal{H}}$  is the positive definite matrix defined by

$$\mathbf{P} \begin{bmatrix} x \\ \sigma \\ v \\ \tau \\ r \\ \eta \\ \xi \\ \zeta \end{bmatrix} := \begin{bmatrix} (1/\gamma_1)x - \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}v - \lambda\mathbf{L}^\top\mathbf{r} \\ (\lambda/\gamma_2)\sigma - \lambda\eta - \lambda\mathbf{M}^\top\xi \\ (\lambda/\gamma_3)v - \lambda\mathbf{B}^\top\mathbf{B}Lx \\ (\lambda/\gamma_3)\tau - \lambda\mathbf{M}^\top\zeta \\ \lambda r - \lambda Lx \\ \lambda\eta - \lambda\sigma \\ \lambda\xi - \lambda\mathbf{M}\sigma \\ (\lambda/\gamma_4)\zeta - \lambda\mathbf{M}\tau \end{bmatrix}. \quad (23)$$

In conjunction with Fact 3, these properties yield that  $(x_k)_{k \in \mathbb{N}}$  generated by Algorithm 1 converges to a point in  $\mathcal{S}$ , i.e., a globally optimal solution of (10), if  $\mathcal{S}$  is nonempty<sup>4</sup>.

<sup>3</sup>For any  $\kappa > 1$ , the condition (20) is satisfied by, e.g., using  $\delta \in \mathbb{R}_{++}$ ,  $\gamma_1 = 1/((\kappa/2)\mathbf{A}^\top\mathbf{A} + \lambda\mathbf{L}^\top\mathbf{L}\|\text{op} + \delta)$ ,  $\gamma_2 = 1/(\|\mathbf{M}\|_{\text{op}}^2 + 1 + \delta)$ ,  $\gamma_3 = 1/((\kappa/2 + 2/\kappa)\|\mathbf{B}\|_{\text{op}}^2 + \delta)$ , and  $\gamma_4 = 1/(\gamma_3\|\mathbf{M}\|_{\text{op}}^2 + \delta)$ .

<sup>4</sup> $\mathcal{S} \neq \emptyset$  is automatically guaranteed in many scenarios, e.g., if  $\mathcal{C}$  is bounded. The expression (11), (13), and Lemma 3(i) yield  $\text{dom}(J) = \mathbb{R}^n$ , and thus  $\text{dom}(J) \cap \mathcal{C} \neq \emptyset$ . If  $\mathcal{C}$  is bounded, then  $\mathcal{S} \neq \emptyset$  holds by [41, Proposition 11.15] under the condition (12) guaranteeing  $J \in \Gamma_0(\mathbb{R}^n)$ .

### Algorithm 1: Solver for model (10) with (4), (9), and (15)

**Input:**  $\mathbf{B}$  satisfying (12),  $\gamma_1, \gamma_2, \gamma_3, \gamma_4$  satisfying (20),

$$x_0 \in \mathbb{R}^n, \sigma_0 \in \mathbb{R}^l, v_0 \in \mathbb{R}^m, \tau_0 \in \mathbb{R}^l,$$

$$r_0 \in \mathbb{R}^m, \eta_0 \in \mathbb{R}^l, \xi_0 \in \mathbb{R}^p, \zeta_0 \in \mathbb{R}^p.$$

**for**  $k = 0, 1, 2, \dots$  **do**

$$\begin{aligned} x_{k+1} &= P_{\mathcal{C}}[x_k - \gamma_1(\mathbf{Q}x_k - \mathbf{A}^\top\mathbf{y} + \lambda\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B}v_k + r_k))]; \\ \sigma_{k+1} &= \sigma_k - \gamma_2(\eta_k + \mathbf{M}^\top\xi_k); \\ u_{k+1} &= \mathbf{L}(2x_{k+1} - x_k); \\ (v_{k+1}, \tau_{k+1}) &= \text{prox}_{\gamma_3 f}(v_k + \gamma_3\mathbf{B}^\top\mathbf{B}(u_{k+1} - v_k), \tau_k - \gamma_3\mathbf{M}^\top\zeta_k); \\ (r_{k+1}, \eta_{k+1}) &= \text{prox}_{f^*}(r_k + u_{k+1}, \eta_k + 2\sigma_{k+1} - \sigma_k); \\ \xi_{k+1} &= \text{prox}_{g^*}(\xi_k + \mathbf{M}(2\sigma_{k+1} - \sigma_k)); \\ \zeta_{k+1} &= \text{prox}_{\gamma_4 g^*}(\zeta_k + \gamma_4\mathbf{M}(2\tau_{k+1} - \tau_k)); \end{aligned}$$

*Proof.* First, we derive the relation (22). Under the condition (12), applying Fact 1(b) to  $\partial(J + \iota_{\mathcal{C}})(x)$  with the expression (11) repeatedly by the differentiability of the first three terms on the right-hand side of (11) on  $\mathbb{R}^n$ ,  $\text{dom}(\psi \circ L) = \mathbb{R}^n$  by Lemma 2, and Lemma 3(i), we obtain

$$\begin{aligned} \partial(J + \iota_{\mathcal{C}})(x) &= \mathbf{Q}x - \mathbf{A}^\top\mathbf{y} + \lambda\partial(\psi \circ L)(x) + \partial\iota_{\mathcal{C}}(x) \\ &\quad + \lambda\partial\left((\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^* \circ \mathbf{B}^\top\mathbf{B}L\right)(x). \end{aligned}$$

Furthermore, Lemma 3(ii) and Fact 1(c) yield

$$\begin{aligned} &\partial\left((\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^* \circ \mathbf{B}^\top\mathbf{B}L\right)(x) \\ &= \mathbf{L}^\top\mathbf{B}^\top\mathbf{B}\left(\partial(\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^*(\mathbf{B}^\top\mathbf{B}Lx)\right). \end{aligned}$$

In addition, by the property (2), we obtain

$$\begin{aligned} v &\in \partial(\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)^*(\mathbf{B}^\top\mathbf{B}Lx) \\ \Leftrightarrow \mathbf{B}^\top\mathbf{B}Lx &\in \partial(\psi + \frac{1}{2}\|\mathbf{B} \cdot \|^2)(v) \\ \Leftrightarrow \mathbf{B}^\top\mathbf{B}Lx &\in \partial\psi(v) + \mathbf{B}^\top\mathbf{B}v, \end{aligned}$$

where the last equivalence holds since  $\frac{1}{2}\|\mathbf{B} \cdot \|^2$  is differentiable on  $\mathbb{R}^m$ . Thus, we have

$$\begin{aligned} x^* \in \mathcal{S} &\quad (\Leftrightarrow \mathbf{0} \in \partial(J + \iota_{\mathcal{C}})(x^*)) \\ \Leftrightarrow \begin{cases} \mathbf{0} \in \mathbf{Q}x^* - \mathbf{A}^\top\mathbf{y} + \lambda\partial(\psi \circ L)(x^*) + \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}v^* + t^* \\ \mathbf{B}^\top\mathbf{B}Lx^* \in \partial\psi(v^*) + \mathbf{B}^\top\mathbf{B}v^* \\ t^* \in \partial\iota_{\mathcal{C}}(x^*). \end{cases} \end{aligned}$$

Since Lemma 1 guarantees  $\text{argmin}_{\sigma \in \mathbb{R}^l} \varphi(Lx^*, \sigma) \neq \emptyset$  and  $\text{argmin}_{\sigma \in \mathbb{R}^l} \varphi(v^*, \sigma) \neq \emptyset$ , we can introduce auxiliary variables  $\sigma^*$  and  $\tau^*$  as follows:

$$\begin{aligned} x^* \in \mathcal{S} &\Leftrightarrow \begin{cases} -\mathbf{Q}x^* + \mathbf{A}^\top\mathbf{y} - \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}v^* - t^* \in \lambda\partial(\psi \circ L)(x^*) \\ \sigma^* \in \underset{\sigma \in \mathbb{R}^l}{\text{argmin}} \varphi(Lx^*, \sigma) \\ \mathbf{B}^\top\mathbf{B}Lx^* - \mathbf{B}^\top\mathbf{B}v^* \in \partial\psi(v^*) \\ \tau^* \in \underset{\sigma \in \mathbb{R}^l}{\text{argmin}} \varphi(v^*, \sigma) \\ t^* \in \partial\iota_{\mathcal{C}}(x^*). \end{cases} \end{aligned}$$

Using Lemma 4, we further obtain

$$\begin{aligned} & \mathbf{x}^* \in \mathcal{S} \\ \Leftrightarrow & \begin{cases} (-\mathbf{Q}\mathbf{x}^* + \mathbf{A}^\top \mathbf{y} - \lambda \mathbf{L}^\top \mathbf{B}^\top \mathbf{B}\mathbf{v}^* - \mathbf{t}^*, \mathbf{0}) \\ \qquad \qquad \qquad \in \lambda \partial(\varphi \circ \bar{\mathbf{L}})(\mathbf{x}^*, \boldsymbol{\sigma}^*) \\ (\mathbf{B}^\top \mathbf{B}\mathbf{L}\mathbf{x}^* - \mathbf{B}^\top \mathbf{B}\mathbf{v}^*, \mathbf{0}) \in \partial\varphi(\mathbf{v}^*, \boldsymbol{\tau}^*) \\ \mathbf{t}^* \in \partial\iota_{\mathcal{C}}(\mathbf{x}^*). \end{cases} \end{aligned}$$

Applying Fact 1(c) under Assumption 3(iii) and Fact 1(a) under Assumption 3(iv) with the definitions of  $\varphi$ ,  $\bar{\mathbf{L}}$ , and  $\bar{\mathbf{M}}$  in (15), (16), and (17), we obtain

$$\begin{aligned} \partial(\varphi \circ \bar{\mathbf{L}})(\mathbf{x}^*, \boldsymbol{\sigma}^*) &= \bar{\mathbf{L}}^\top [\partial\varphi(\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*)] \\ &= \bar{\mathbf{L}}^\top [\partial(f + g \circ \bar{\mathbf{M}})(\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*)] \\ &= \bar{\mathbf{L}}^\top [\partial f(\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*) + \bar{\mathbf{M}}^\top (\partial g(\mathbf{M}\boldsymbol{\sigma}^*))], \end{aligned}$$

and similarly,

$$\partial\varphi(\mathbf{v}^*, \boldsymbol{\tau}^*) = \partial f(\mathbf{v}^*, \boldsymbol{\tau}^*) + \bar{\mathbf{M}}^\top (\partial g(\mathbf{M}\boldsymbol{\tau}^*)).$$

Thus, using the relations

$$\begin{aligned} (\mathbf{r}^*, \boldsymbol{\eta}^*) \in \partial f(\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*) &\Leftrightarrow (\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*) \in \partial f^*(\mathbf{r}^*, \boldsymbol{\eta}^*), \\ \boldsymbol{\xi}^* \in \partial g(\mathbf{M}\boldsymbol{\sigma}^*) &\Leftrightarrow \mathbf{M}\boldsymbol{\sigma}^* \in \partial g^*(\boldsymbol{\xi}^*), \\ \boldsymbol{\zeta}^* \in \partial g(\mathbf{M}\boldsymbol{\tau}^*) &\Leftrightarrow \mathbf{M}\boldsymbol{\tau}^* \in \partial g^*(\boldsymbol{\zeta}^*), \end{aligned}$$

since  $\bar{\mathbf{L}}^\top [(\mathbf{r}^*, \boldsymbol{\eta}^*) + \bar{\mathbf{M}}^\top \boldsymbol{\xi}^*] = (\mathbf{L}^\top \mathbf{r}^*, \boldsymbol{\eta}^* + \mathbf{M}^\top \boldsymbol{\xi}^*)$  and  $\bar{\mathbf{M}}^\top \boldsymbol{\zeta}^* = (\mathbf{0}, \mathbf{M}^\top \boldsymbol{\zeta}^*)$ , we deduce

$$\begin{aligned} & \mathbf{x}^* \in \mathcal{S} \\ \Leftrightarrow & \begin{cases} \mathbf{0} \in \mathbf{Q}\mathbf{x}^* - \mathbf{A}^\top \mathbf{y} + \lambda \mathbf{L}^\top \mathbf{B}^\top \mathbf{B}\mathbf{v}^* + \lambda \mathbf{L}^\top \mathbf{r}^* + \partial\iota_{\mathcal{C}}(\mathbf{x}^*) \\ \mathbf{0} = \lambda \boldsymbol{\eta}^* + \lambda \mathbf{M}^\top \boldsymbol{\xi}^* \\ (\mathbf{0}, \mathbf{0}) \in \lambda \partial f(\mathbf{v}^*, \boldsymbol{\tau}^*) + \lambda (\mathbf{B}^\top \mathbf{B}(\mathbf{v}^* - \mathbf{L}\mathbf{x}^*), \mathbf{M}^\top \boldsymbol{\zeta}^*) \\ (\mathbf{0}, \mathbf{0}) \in \lambda \partial f^*(\mathbf{r}^*, \boldsymbol{\eta}^*) - \lambda (\mathbf{L}\mathbf{x}^*, \boldsymbol{\sigma}^*) \\ \mathbf{0} \in \lambda \partial g^*(\boldsymbol{\xi}^*) - \lambda \mathbf{M}\boldsymbol{\sigma}^* \\ \mathbf{0} \in \lambda \partial g^*(\boldsymbol{\zeta}^*) - \lambda \mathbf{M}\boldsymbol{\tau}^*. \end{cases} \end{aligned}$$

By using the notation  $\mathbf{z} = (\mathbf{x}, \boldsymbol{\sigma}, \mathbf{v}, \boldsymbol{\tau}, \mathbf{r}, \boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\zeta})$ , the last expression can be rewritten as

$$\begin{aligned} \mathbf{x}^* \in \mathcal{S} &\Leftrightarrow \mathbf{0} \in H(\mathbf{z}^*) + \mathcal{D}(\mathbf{z}^*) + \mathbf{N}\mathbf{z}^* \\ &\Leftrightarrow \mathbf{P}\mathbf{z}^* - H(\mathbf{z}^*) \in \mathbf{P}\mathbf{z}^* + \mathcal{D}(\mathbf{z}^*) + \mathbf{N}\mathbf{z}^*, \quad (24) \end{aligned}$$

where  $H: \mathcal{H} \rightarrow \mathcal{H}$  is the affine operator,  $\mathcal{D}: \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is the set-valued operator consisting of subdifferentials, and  $\mathbf{N} \in \mathbb{R}^{\dim \mathcal{H} \times \dim \mathcal{H}}$  is the skew-symmetric matrix, respectively defined by

$$H(\mathbf{z}) := (\mathbf{Q}\mathbf{x} - \mathbf{A}^\top \mathbf{y}, \mathbf{0}, \lambda \mathbf{B}^\top \mathbf{B}\mathbf{v}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}), \quad (25)$$

$$\begin{aligned} \mathcal{D}(\mathbf{z}) &:= \partial\iota_{\mathcal{C}}(\mathbf{x}) \times \{\mathbf{0}\} \times \lambda \partial f(\mathbf{v}, \boldsymbol{\tau}) \\ &\quad \times \lambda \partial f^*(\mathbf{r}, \boldsymbol{\eta}) \times \lambda \partial g^*(\boldsymbol{\xi}) \times \lambda \partial g^*(\boldsymbol{\zeta}), \quad (26) \end{aligned}$$

$$\mathbf{N} := \boldsymbol{\Pi}^\top \begin{bmatrix} \mathbf{N}_1 & & \\ & \mathbf{N}_2 & \\ & & \mathbf{N}_3 \end{bmatrix} \boldsymbol{\Pi}, \quad (27)$$

$$\mathbf{N}_1 := \lambda \begin{bmatrix} \mathbf{O} & \mathbf{L}^\top \mathbf{B}^\top \mathbf{B} & \mathbf{L}^\top \\ -\mathbf{B}^\top \mathbf{B}\mathbf{L} & \mathbf{O} & \mathbf{O} \\ -\mathbf{L} & \mathbf{O} & \mathbf{O} \end{bmatrix},$$

$$\begin{aligned} \mathbf{N}_2 &:= \lambda \begin{bmatrix} \mathbf{O} & \mathbf{I} & \mathbf{M}^\top \\ -\mathbf{I} & \mathbf{O} & \mathbf{O} \\ -\mathbf{M} & \mathbf{O} & \mathbf{O} \end{bmatrix}, \\ \mathbf{N}_3 &:= \lambda \begin{bmatrix} \mathbf{O} & \mathbf{M}^\top \\ -\mathbf{M} & \mathbf{O} \end{bmatrix}, \end{aligned}$$

where  $\boldsymbol{\Pi} \in \mathbb{R}^{\dim \mathcal{H} \times \dim \mathcal{H}}$  is the permutation matrix defined as

$$\boldsymbol{\Pi}: (\mathbf{x}, \boldsymbol{\sigma}, \mathbf{v}, \boldsymbol{\tau}, \mathbf{r}, \boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\zeta}) \mapsto (\mathbf{x}, \mathbf{v}, \mathbf{r}, \boldsymbol{\sigma}, \boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\tau}, \boldsymbol{\zeta}). \quad (28)$$

Meanwhile, since the proximity operator is the resolvent of subdifferential (see (3)), we derive from (21) that

$$\begin{aligned} T_{\text{GME-MI}}(\mathbf{z}) &= \hat{\mathbf{z}} \\ \Leftrightarrow & \begin{cases} \mathbf{x} - \gamma_1(\mathbf{Q}\mathbf{x} - \mathbf{A}^\top \mathbf{y} + \lambda \mathbf{L}^\top \mathbf{B}^\top \mathbf{B}\mathbf{v} + \lambda \mathbf{L}^\top \mathbf{r}) \\ \qquad \qquad \qquad \in (\text{Id} + \partial\iota_{\mathcal{C}})(\hat{\mathbf{x}}) \\ \boldsymbol{\sigma} - \gamma_2(\boldsymbol{\eta} + \mathbf{M}^\top \boldsymbol{\xi}) = \hat{\boldsymbol{\sigma}} \\ (\mathbf{v} + \gamma_3 \mathbf{B}^\top \mathbf{B}(\mathbf{L}(2\hat{\mathbf{x}} - \mathbf{x}) - \mathbf{v}), \boldsymbol{\tau} - \gamma_3 \mathbf{M}^\top \boldsymbol{\zeta}) \\ \qquad \qquad \qquad \in (\text{Id} + \gamma_3 \partial f)(\hat{\mathbf{v}}, \hat{\boldsymbol{\tau}}) \\ (\mathbf{r} + \mathbf{L}(2\hat{\mathbf{x}} - \mathbf{x}), \boldsymbol{\eta} + 2\hat{\boldsymbol{\sigma}} - \boldsymbol{\sigma}) \in (\text{Id} + \partial f^*)(\hat{\mathbf{r}}, \hat{\boldsymbol{\eta}}) \\ \boldsymbol{\xi} + \mathbf{M}(2\hat{\boldsymbol{\sigma}} - \boldsymbol{\sigma}) \in (\text{Id} + \partial g^*)(\hat{\boldsymbol{\xi}}) \\ \boldsymbol{\zeta} + \gamma_4 \mathbf{M}(2\hat{\boldsymbol{\tau}} - \boldsymbol{\tau}) \in (\text{Id} + \gamma_4 \partial g^*)(\hat{\boldsymbol{\zeta}}) \end{cases} \\ &\Leftrightarrow \mathbf{P}\mathbf{z} - H(\mathbf{z}) \in \mathbf{P}\hat{\mathbf{z}} + \mathcal{D}(\hat{\mathbf{z}}) + \mathbf{N}\hat{\mathbf{z}}. \end{aligned}$$

This relation for  $\mathbf{z} = \hat{\mathbf{z}} = \mathbf{z}^*$  and (24) yield the claim (22).

Next, we prove that  $T_{\text{GME-MI}}$  is  $\frac{\kappa}{2\kappa-1}$ -averaged nonexpansive in  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathbf{P}}, \|\cdot\|_{\mathbf{P}})$ . The positive definiteness of  $\mathbf{P}$  is shown in Appendix B-1. Since  $\mathcal{D}$  in (26) is the product of subdifferentials, it is maximally monotone in  $(\mathcal{H}, \langle \cdot, \cdot \rangle, \|\cdot\|)$  by [41, Theorem 20.25 and Proposition 20.23]. Since  $\mathbf{N}$  in (27) is skew-symmetric, it is maximally monotone in  $(\mathcal{H}, \langle \cdot, \cdot \rangle, \|\cdot\|)$  by [41, Example 20.35]. Thus, since the domain of  $\mathbf{N}$  is  $\mathcal{H}$ ,  $\mathcal{D} + \mathbf{N}$  is maximally monotone in  $(\mathcal{H}, \langle \cdot, \cdot \rangle, \|\cdot\|)$  by [41, Corollary 25.5]. The positive definiteness of  $\mathbf{P}$  and the maximal monotonicity of  $\mathcal{D} + \mathbf{N}$  imply that  $\mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N})$  is maximally monotone in  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathbf{P}}, \|\cdot\|_{\mathbf{P}})$  by [41, Proposition 20.24]. Thus, its resolvent  $(\text{Id} + \mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N}))^{-1}$  is single-valued and  $\frac{1}{2}$ -averaged nonexpansive in  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathbf{P}}, \|\cdot\|_{\mathbf{P}})$ . Since  $(\text{Id} + \mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N}))^{-1}$  is single-valued, we have

$$\begin{aligned} T_{\text{GME-MI}}(\mathbf{z}) &= \hat{\mathbf{z}} \\ \Leftrightarrow \mathbf{P}\mathbf{z} - H(\mathbf{z}) &\in \mathbf{P}\hat{\mathbf{z}} + \mathcal{D}(\hat{\mathbf{z}}) + \mathbf{N}\hat{\mathbf{z}} \\ \Leftrightarrow \mathbf{z} - \mathbf{P}^{-1}H(\mathbf{z}) &\in \hat{\mathbf{z}} + \mathbf{P}^{-1}(\mathcal{D} + \mathbf{N})(\hat{\mathbf{z}}) \\ \Leftrightarrow [(\text{Id} + \mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N}))^{-1} \circ (\text{Id} - \mathbf{P}^{-1} \circ H)](\mathbf{z}) &= \hat{\mathbf{z}}, \end{aligned}$$

i.e.,

$$T_{\text{GME-MI}} = (\text{Id} + \mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N}))^{-1} \circ (\text{Id} - \mathbf{P}^{-1} \circ H).$$

In  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathbf{P}}, \|\cdot\|_{\mathbf{P}})$ , the  $\frac{1}{2}$ -averaged nonexpansiveness of  $(\text{Id} + \mathbf{P}^{-1} \circ (\mathcal{D} + \mathbf{N}))^{-1}$  and the  $\frac{1}{\kappa}$ -averaged nonexpansiveness of  $\text{Id} - \mathbf{P}^{-1} \circ H$ , proved in Appendix B-2, yield that  $T_{\text{GME-MI}}$  is  $\frac{\kappa}{2\kappa-1}$ -averaged nonexpansive by Fact 2. Finally, since Algorithm 1 is designed to have  $\mathbf{z}_{k+1} = T_{\text{GME-MI}}(\mathbf{z}_k)$ , the averaged nonexpansiveness of  $T_{\text{GME-MI}}$  and the relation (22) yield that  $(\mathbf{x}_k)_{k \in \mathbb{N}}$  converges to a point in  $\mathcal{S}$  by Fact 3.  $\square$



## V. NUMERICAL EXAMPLES

To demonstrate the effectiveness of the proposed framework, we present numerical examples of the GME-MI model corresponding to the scenarios in Examples 1–2.

### A. Estimation of Block-Sparse Signals


We conduct numerical experiments corresponding to the scenario in Examples 1–2(a) for block-sparse estimation with unknown block partition. More precisely, we consider the estimation of a block-sparse signal  $\mathbf{x}_{\text{org}} \in \mathcal{C} := \mathbb{R}^n$  from noisy compressive measurements  $\mathbf{y} = \mathbf{A}\mathbf{x}_{\text{org}} + \boldsymbol{\varepsilon} \in \mathbb{R}^d$ , where the entries of  $\mathbf{A} \in \mathbb{R}^{d \times n}$  are drawn from i.i.d. Gaussian distribution  $\mathcal{N}(0, 1)$ ,  $\boldsymbol{\varepsilon} \in \mathbb{R}^d$  is the white Gaussian noise, and  $d < n$ . The block-sparse signal  $\mathbf{x}_{\text{org}}$  with 4 nonzero blocks, 80 nonzero components, and  $n = 256$  is randomly generated by the scheme used in [29]. Note that the block partition is randomly generated for each trial to investigate the average performance for various block partitions. Amplitudes of nonzero components are drawn from i.i.d.  $\mathcal{N}(0, 1)$ .

As an instance of the GME-MI model, the GME-LOP- $\ell_2/\ell_1$  penalty  $\Psi_{\mathbf{B}, \alpha}^{\text{LOP}}$  presented in Example 2(a) is used in Definition 3. Since  $\mathbf{x}_{\text{org}}$  itself is block-sparse in the experiments, we set  $\mathbf{L} = \mathbf{I}$ . Due to  $\mathbf{L} = \mathbf{I}$ , we can set  $\mathbf{B}$  satisfying (12) by  $\mathbf{B} = \sqrt{\theta/\lambda}\mathbf{A}$  with any  $\theta \in [0, 1]$ .

We compare the proposed GME-LOP- $\ell_2/\ell_1$  penalty with the LOP- $\ell_2/\ell_1$  penalty given in Example 1(a) as well as the GME penalty  $R_{\mathbf{B}}(\mathbf{x}) := R(\mathbf{x}) - \min_{\mathbf{v} \in \mathbb{R}^n} [R(\mathbf{v}) + \frac{1}{2}\|\mathbf{B}(\mathbf{x} - \mathbf{v})\|^2]$  [17] for convex prox-friendly penalties:  $R = \|\cdot\|_{2,1}$  for block-sparsity with fixed blocks (proposed in [20], [47]) and  $R = \|\cdot\|_1$  for non-structured sparsity (proposed as GMC [14]). We also consider the  $\ell_2/\ell_1$  and  $\ell_1$  penalties for reference. All the penalties are combined with the square error  $\frac{1}{2}\|\mathbf{y} - \mathbf{A}\mathbf{x}\|^2$ .

The GME-LOP- $\ell_2/\ell_1$  model is solved by Algorithm 1 with the representation given in Example 3(a), the GME models for  $R = \|\cdot\|_{2,1}$  and  $R = \|\cdot\|_1$  are solved by the iterative algorithm given in [17], and the convexly regularized models are solved by the iterative algorithm given in [29]. We terminate the iteration when the norm of the differences between the variables of successive iterates is below the threshold  $10^{-4}$  or the number of iterations reaches 10000. For the GME-LOP- $\ell_2/\ell_1$  model, the computational complexity of Algorithm 1 per iteration is  $\mathcal{O}(dn)$  for matrix-vector multiplication and  $\mathcal{O}(n)$  for the proximity operators (see Appendix A-2a). This complexity is identical to those of the other algorithms.

We compare the models in terms of normalized mean square error (NMSE)  $\|\mathbf{x}_{\text{org}} - \mathbf{x}^*\|^2 / \|\mathbf{x}_{\text{org}}\|^2$ , where  $\mathbf{x}^*$  is an optimal solution of each regularization model. Note that the regularization parameter  $\lambda$  and other tuning parameters are adjusted to obtain the best estimation accuracy for each pair of models and experimental conditions. The tuning parameters other than  $\lambda$  are as follows:  $\alpha$  for GME-LOP- $\ell_2/\ell_1$  and LOP- $\ell_2/\ell_1$ , block-size for GME- $\ell_2/\ell_1$  and  $\ell_2/\ell_1$ , and  $\theta$  for the overall convexity conditions of GME-LOP- $\ell_2/\ell_1$ , GME- $\ell_2/\ell_1$ , and GME- $\ell_1$ . In Figs. 3(a) and (b), NMSE is plotted with respect to the number,  $d$ , of measurements and the SNR  $E[\|\mathbf{A}\mathbf{x}_{\text{org}}\|^2]/E[\|\boldsymbol{\varepsilon}\|^2]$ , respectively, where the results are averaged over 100 independent trials. As is clear from Fig. 3,



(a) NMSE versus the number of measurements, where the SNR is fixed to 40dB. (b) NMSE versus the SNR, where the number of measurements is fixed to 220.

Fig. 3. Average NMSEs for estimation of block-sparse signals.




Fig. 4. Original block-sparse signal and the estimates for a trial, where the number of measurements is 160 and the SNR is 40dB.

GME-LOP- $\ell_2/\ell_1$  model achieves the best estimation accuracy, representing a significant improvement over the LOP- $\ell_2/\ell_1$  model.

We also discuss an example involving the original block-sparse signal and the estimates for a trial in Fig. 4. The GME-LOP- $\ell_2/\ell_1$  model certainly remedies the underestimation tendency of the LOP- $\ell_2/\ell_1$  model. Meanwhile, the  $\ell_1$  and  $\ell_2/\ell_1$  models yield multiple incorrect nonzero components, possibly due to their fixed structures in contrast to the (GME-)LOP- $\ell_2/\ell_1$  model that automatically optimizes the block structure. This drawback is not resolved by the GME- $\ell_1$  and GME- $\ell_2/\ell_1$  models, while mitigating the underestimation tendency.

### B. Estimation of Piecewise Linear Signals

We conduct numerical experiments for the scenario in Examples 1–2(b) to illustrate the estimation of piecewise smooth signals. In particular, as a simple example of piecewise smooth signals, we set  $\mathbf{x}_{\text{org}} \in \mathcal{C} := [-1, 1]^n$  as the piecewise linear signal indicated by the dotted line in Fig. 5, where  $n = 128$ . Measurements are generated in the same way as in the previous section.

As an instance of the GME-MI model, the (second-order) GME-TGV penalty  $\Psi_{\mathbf{B}, \alpha}^{\text{TGV}}$  presented in Example 2(b) is used in Definition 3, where we set  $\mathbf{L} = \mathbf{D}_{1d}$  with  $\mathbf{D}_{1d} \in \mathbb{R}^{(n-1) \times n}$  defined as  $\mathbf{D}_{1d}: \mathbf{x} \mapsto (x_{i+1} - x_i)_{i=1}^{n-1}$ . In this case,  $\tilde{\mathbf{D}}$  in the seed function  $\varphi_{\alpha}^{\text{TGV}}$  in (8) becomes  $\mathbf{D}_{1d}^{\top}$ . Since only  $\mathbf{B}^{\top}\mathbf{B}$  is





Fig. 5. Original piecewise linear signal and the estimates for a trial, where the number of measurements is 100 and the SNR is 20dB.



(a) NMSE versus the number of measurements, where the SNR is fixed to 30dB. (b) NMSE versus the SNR, where the number of measurements is fixed to 100.

Fig. 6. Average NMSEs for estimation of piecewise linear signals.

used in Algorithm 1, we set  $\mathbf{B}^T \mathbf{B}$  satisfying (12) for  $\mathbf{L} = \mathbf{D}_{1d}$  as follows. We have the equality

$$\underbrace{\begin{bmatrix} -1 & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & -1 & 1 \end{bmatrix}}_{=\mathbf{D}_{1d} \in \mathbb{R}^{(n-1) \times n}} \underbrace{\begin{bmatrix} 1 & & & & \\ 1 & 1 & & & \\ \vdots & \vdots & \ddots & & \\ 1 & 1 & \dots & 1 \end{bmatrix}}_{=:\mathbf{S}_{1d} \in \mathbb{R}^{n \times n}} = \begin{bmatrix} \mathbf{0} & \mathbf{I} \end{bmatrix}.$$

Thus, based on [44, Theorem 1] (see also Remark 1), letting  $\mathbf{h} \in \mathbb{R}^d$  and  $\mathbf{H} \in \mathbb{R}^{d \times (n-1)}$  such that  $[\mathbf{h} \ \mathbf{H}] := \mathbf{A} \mathbf{S}_{1d}$ , we can show that

$$\mathbf{B}^T \mathbf{B} = (\theta/\lambda) \mathbf{H}^T (\mathbf{I} - \mathbf{h} \mathbf{h}^\dagger) \mathbf{H} \in \mathbb{R}^{(n-1) \times (n-1)}$$

with any  $\theta \in [0, 1]$  satisfies (12), where  $\mathbf{h}^\dagger = \mathbf{h}^T / \|\mathbf{h}\|^2$  if  $\mathbf{h} \neq \mathbf{0}$  and  $\mathbf{h}^\dagger = \mathbf{0}^T$  if  $\mathbf{h} = \mathbf{0}$ .

We compare the proposed GME-TGV penalty with the TGV penalty given in Example 1(b) as well as the GME-TV penalty ( $\|\cdot\|_1 \circ \mathbf{D}_{1d}$  presented in [17]). We also consider the TV penalty  $\|\cdot\|_1 \circ \mathbf{D}_{1d}$  for reference. All the penalties are combined with the square error  $\frac{1}{2} \|\mathbf{y} - \mathbf{A} \mathbf{x}\|^2$ .

The GME-TGV model is solved by Algorithm 1 with the representation in Example 3(b), the GME-TV model is solved by the iterative algorithm given in [17] with an extension [20] to handle the constraint  $[-1, 1]^n$ , and the TGV and TV models are solved as the special cases with  $\mathbf{B} = \mathbf{O}$  of the GME-TGV and GME-TV models, respectively. We terminate the iteration in the same way as in the previous section. For the GME-TGV model, the computational complexity of Algorithm 1 per iteration is  $\mathcal{O}(dn)$  for matrix-vector multiplication,  $\mathcal{O}(n)$  for the projection  $P_{[-1, 1]^n}$ , and  $\mathcal{O}(n)$  for the proximity operators

(see Appendix A-2b). This complexity is identical to that of the algorithm for GME-TV.

In Fig. 6, we compare the models in terms of NMSE, where the results are averaged over 100 independent trials, and the regularization parameter  $\lambda$  and other tuning parameters are adjusted to yield the best estimation accuracy for each pair of models and experimental conditions. The tuning parameters other than  $\lambda$  are as follows:  $\alpha$  for GME-TGV and TGV, and  $\theta$  for the overall convexity conditions of GME-TGV and GME-TV. As is apparent from Fig. 6, the GME-TGV model exhibits the best estimation accuracy, with a remarkable improvement over the TGV model.

We also discuss examples of the estimates for a trial in Fig. 5. In Fig. 5(a) for the TV model, we observe the so-called stair-casing effect. The GME-TV model does not resolve the stair-casing effect in the slanted segments in Fig. 5(b), while recovering signal discontinuity more effectively than the TV model. The TGV model depicted in Fig. 5(c) mostly does not exhibit the stair-casing effect, but underestimates the discontinuous jumps of the signal and its first derivative. In Fig. 5(d), we observe that the GME-TGV model significantly mitigates the underestimation of the jumps of the signal and its first derivative, while being free from the stair-casing effect.

## VI. CONCLUSION

This paper presented a convex-nonconvex framework to mitigate the underestimation tendencies of MI penalties. We designed the GME-MI penalty, where the generalized Moreau envelope of the MI penalty is subtracted from it. While the GME-MI penalty is nonconvex in general, we derived an overall convexity condition for the GME-MI regularized least-squares model. Moreover, we presented a proximal splitting algorithm guaranteed to converge to a globally optimal solution of the GME-MI model under the overall convexity condition. Numerical examples considered in the enhancement scenarios of the LOP- $\ell_2/\ell_1$  and TGV penalties demonstrate the effectiveness of the proposed framework.

## APPENDIX A

### VALIDITY OF ASSUMPTIONS

We prove that the assumptions adopted while designing the proposed framework automatically hold for the seed functions in Example 1 and their representations in Example 3.

#### 1) Validity of Assumption 1

a) *Example 1(a)*: Since  $\mathbf{D}$  is the discrete difference operator, for any constant vector  $\mathbf{c} = (c, c, \dots, c) \in \mathbb{R}^m$ , we have  $\mathbf{D} \mathbf{c} = \mathbf{0}$ , and thus  $\iota_{B_1^\alpha}(\mathbf{D} \mathbf{c}) = 0$  holds for any  $\alpha \in \mathbb{R}_+$  by the definition (7). Meanwhile, by the definition (6),  $\sum_{i=1}^m h(u_i, c)$  takes a finite value for any  $\mathbf{u} \in \mathbb{R}^m$  and  $c \in \mathbb{R}_{++}$ . Altogether,  $\varphi_\alpha^{\text{LOP}}(\mathbf{u}, \cdot)$  in (5) is proper for every  $\mathbf{u} \in \mathbb{R}^m$  and  $\alpha \in \mathbb{R}_+$ .

Since  $h$  is coercive by [29, Lemma 1] and  $\iota_{B_1^\alpha}$  is bounded below for any  $\alpha \in \mathbb{R}_+$ ,  $\varphi_\alpha^{\text{LOP}}(\mathbf{u}, \cdot)$  is coercive for every  $\mathbf{u} \in \mathbb{R}^m$  and  $\alpha \in \mathbb{R}_+$ .

*b) Example 1(b):* For every  $\mathbf{u} \in \mathbb{R}^m$  and  $\alpha \in (0, 1)$ ,  $\varphi_\alpha^{\text{TGV}}(\mathbf{u}, \cdot)$  in (8) is proper due to  $\text{dom}(\|\cdot\|_{2,1}^{\mathcal{G}_1}) = \mathbb{R}^m$  and  $\text{dom}(\|\cdot\|_{2,1}^{\mathcal{G}_2}) = \mathbb{R}^p$ . Since  $\|\cdot\|_{2,1}^{\mathcal{G}_1}$  is a norm, we have

$$\|\mathbf{u} - \boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1} \geq \|\boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1} - \|\mathbf{u}\|_{2,1}^{\mathcal{G}_1},$$

which implies that  $\|\mathbf{u} - \cdot\|_{2,1}^{\mathcal{G}_1}$  is coercive for any fixed  $\mathbf{u} \in \mathbb{R}^m$ . Thus, since  $\|\tilde{\mathbf{D}} \cdot\|_{2,1}^{\mathcal{G}_2}$  is bounded below,  $\varphi_\alpha^{\text{TGV}}(\mathbf{u}, \cdot)$  is coercive for every  $\mathbf{u} \in \mathbb{R}^m$  and  $\alpha \in (0, 1)$ .

## 2) Validity of Assumption 2

We show that efficient exact methods are available for computing the proximity operators of  $f$  and  $g$  in Example 3 used to represent the seed functions in the form of (15).

*a) Example 3(a):* Since  $f(\mathbf{u}, \boldsymbol{\sigma}) = \sum_{i=1}^m h(u_i, \sigma_i)$  is the separable sum of  $h$  in (6), based on [48, Example 2.4], for any  $(\mathbf{u}, \boldsymbol{\sigma}) \in \mathbb{R}^m \times \mathbb{R}^m$  and  $\gamma \in \mathbb{R}_{++}$ , we have  $\text{prox}_{\gamma f}(\mathbf{u}, \boldsymbol{\sigma}) = (\text{prox}_{\gamma h}(u_i, \sigma_i))_{i=1}^m$  with

$$\text{prox}_{\gamma h}(u_i, \sigma_i) = \begin{cases} (0, 0), & \text{if } 2\gamma\sigma_i + u_i^2 \leq \gamma^2; \\ (0, \sigma_i - \frac{\gamma}{2}), & \text{if } u_i = 0 \text{ and } \sigma_i > \frac{\gamma}{2}; \\ (u_i - \gamma t_i \frac{u_i}{|u_i|}, \sigma_i + \gamma \frac{t_i^2 - 1}{2}), & \text{otherwise,} \end{cases}$$

where  $t_i \in \mathbb{R}_{++}$  is the unique positive root of

$$t_i^3 + \left(\frac{2}{\gamma}\sigma_i + 1\right)t_i - \frac{2}{\gamma}|u_i| = 0,$$

which can be solved using Cardano's formula as follows [29], [49]. Let  $p_i = 2\sigma_i/\gamma + 1$  and  $\Delta_i = u_i^2/\gamma^2 + p_i^3/27$ . Then,

$$t_i = \begin{cases} \sqrt[3]{\frac{|u_i|}{\gamma} + \sqrt{\Delta_i}} + \sqrt[3]{\frac{|u_i|}{\gamma} - \sqrt{\Delta_i}}, & \text{if } \Delta_i \geq 0; \\ 2\sqrt{-\frac{p_i}{3}} \cos\left(\frac{\arctan(\gamma\sqrt{-\Delta_i}/|u_i|)}{3}\right), & \text{if } \Delta_i < 0, \end{cases}$$

where  $\sqrt[3]{\cdot}$  denotes the real cubic root.

Since  $\iota_{B_1^\alpha}$  in (7) is the indicator function of the  $\ell_1$  norm ball,  $\text{prox}_{\gamma \iota_{B_1^\alpha}}$  with any  $\gamma \in \mathbb{R}_{++}$  reduces to the  $\ell_1$  ball projection  $P_{B_1^\alpha}$ . There exist algorithms for computing  $P_{B_1^\alpha}$  with  $\mathcal{O}(p)$  expected complexity (see, e.g., [50]).

*b) Example 3(b):* We can represent  $f(\mathbf{u}, \boldsymbol{\sigma}) = \alpha\|\mathbf{u} - \boldsymbol{\sigma}\|_{2,1}^{\mathcal{G}_1}$  as  $f = \alpha\|\cdot\|_{2,1}^{\mathcal{G}_1} \circ U$  with  $U := [\mathbf{I} \quad -\mathbf{I}] \in \mathbb{R}^{m \times 2m}$ . Since  $UU^\top = 2\mathbf{I}$ , based on [41, Proposition 24.14], for any  $(\mathbf{u}, \boldsymbol{\sigma}) \in \mathbb{R}^m \times \mathbb{R}^m$ ,  $\gamma \in \mathbb{R}_{++}$ , and  $\alpha \in (0, 1)$ , we have

$$\text{prox}_{\gamma f}(\mathbf{u}, \boldsymbol{\sigma}) = (\mathbf{u} + \boldsymbol{\sigma} + \text{prox}_{2\gamma\alpha\|\cdot\|_{2,1}^{\mathcal{G}_1}}(\mathbf{u} - \boldsymbol{\sigma}), \\ \mathbf{u} + \boldsymbol{\sigma} - \text{prox}_{2\gamma\alpha\|\cdot\|_{2,1}^{\mathcal{G}_1}}(\mathbf{u} - \boldsymbol{\sigma}))/2.$$

Since  $\mathcal{G}_1 = (\mathcal{I}_j)_{j=1}^r$  is a partition, the proximity operator of  $\kappa\|\cdot\|_{2,1}^{\mathcal{G}_1}$  with  $\kappa = 2\gamma\alpha$  can be computed for any  $\mathbf{w} \in \mathbb{R}^m$  by

$$[\text{prox}_{\kappa\|\cdot\|_{2,1}^{\mathcal{G}_1}}(\mathbf{w})]_{\mathcal{I}_j} = \left(1 - \frac{\kappa}{\max\{\kappa, \|\mathbf{w}_{\mathcal{I}_j}\|_2\}}\right)\mathbf{w}_{\mathcal{I}_j}$$

for each  $j = 1, \dots, r$ . The proximity operator of  $g = (1 - \alpha)\|\cdot\|_{2,1}^{\mathcal{G}_2}$  can be computed similarly.

## 3) Validity of Assumption 3

*a) Example 1(a) with Example 3(a):* By the definition (5) with (6) and (7), Assumptions 3(i) and 3(ii)(a) are clear. From

the definitions (5) and (16), we obtain

$$\text{dom}(\varphi_\alpha^{\text{LOP}}) - \text{ran}(\bar{\mathbf{L}}) \supset (\mathbb{R}^m \times S_\alpha) - (\text{ran}(\mathbf{L}) \times \mathbb{R}^m),$$

where  $S_\alpha := \mathbb{R}_{++}^m \cap \{\boldsymbol{\sigma} \in \mathbb{R}^m \mid \|\mathbf{D}\boldsymbol{\sigma}\|_1 \leq \alpha\}$ . We have  $S_\alpha \neq \emptyset$  for any  $\alpha \in \mathbb{R}_+$  because  $\mathbf{D}\mathbf{c} = \mathbf{0}$  holds for any  $\mathbf{c} = (c, c, \dots, c) \in \mathbb{R}_{++}^m$ , as  $\mathbf{D}$  is the discrete difference operator. Thus, for any  $\alpha \in \mathbb{R}_+$ , we have  $\text{dom}(\varphi_\alpha^{\text{LOP}}) - \text{ran}(\bar{\mathbf{L}}) = \mathbb{R}^m \times \mathbb{R}^m$ , implying that Assumption 3(ii) holds. From the definition (17) and the setting of Example 3(a), we obtain

$$\text{dom}(g) - \bar{M}(\text{dom}(f)) = B_1^\alpha - \mathbf{D}(\mathbb{R}_+^m) = B_1^\alpha - \text{ran}(\mathbf{D}),$$

where  $B_1^\alpha := \{\boldsymbol{\xi} \in \mathbb{R}^p \mid \|\boldsymbol{\xi}\|_1 \leq \alpha\}$ , and the last equality follows from the relation  $\mathbf{D}(\mathbb{R}_+^m) = \text{ran}(\mathbf{D})$ . This relation holds because, for any  $\boldsymbol{\sigma} \in \mathbb{R}^m$ ,  $\mathbf{D}\boldsymbol{\sigma} = \mathbf{D}(\boldsymbol{\sigma} + \mathbf{c})$  and  $\boldsymbol{\sigma} + \mathbf{c} \in \mathbb{R}_+^m$  can be satisfied by setting  $\mathbf{c} = (c, c, \dots, c) \in \mathbb{R}_+^m$  with  $c \geq \max\{|\sigma_i|\}_{i=1}^m$ . For any  $\alpha \in \mathbb{R}_+$ , since  $B_1^\alpha - \text{ran}(\mathbf{D})$  is a nonempty symmetric convex set, Assumption 3(iv) holds by [41, Example 6.10].

*b) Example 1(b) with Example 3(b):* Clear from the definition (8),  $\text{dom}(\|\cdot\|_{2,1}^{\mathcal{G}_1}) = \mathbb{R}^m$ ,  $\text{dom}(\|\cdot\|_{2,1}^{\mathcal{G}_2}) = \mathbb{R}^p$ , and the setting of Example 3(b).

## APPENDIX B

### SUPPLEMENT FOR PROOF OF THEOREM 2

#### 1) Positive definiteness of $\mathbf{P}$

Using  $\mathbf{\Pi}$  in (28), we can represent  $\mathbf{P}$  in (23) as

$$\mathbf{P} = \mathbf{\Pi}^\top \begin{bmatrix} \mathbf{P}_1 & & \\ & \mathbf{P}_2 & \\ & & \mathbf{P}_3 \end{bmatrix} \mathbf{\Pi}, \quad (29)$$

$$\mathbf{P}_1 := \begin{bmatrix} (1/\gamma_1)\mathbf{I} & -\lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B} & -\lambda\mathbf{L}^\top \\ -\lambda\mathbf{B}^\top\mathbf{B}\mathbf{L} & (\lambda/\gamma_3)\mathbf{I} & \mathbf{O} \\ -\lambda\mathbf{L} & \mathbf{O} & \lambda\mathbf{I} \end{bmatrix},$$

$$\mathbf{P}_2 := \lambda \begin{bmatrix} (1/\gamma_2)\mathbf{I} & -\mathbf{I} & -\mathbf{M}^\top \\ -\mathbf{I} & \mathbf{I} & \mathbf{O} \\ -\mathbf{M} & \mathbf{O} & \mathbf{I} \end{bmatrix},$$

$$\mathbf{P}_3 := \lambda \begin{bmatrix} (1/\gamma_3)\mathbf{I} & -\mathbf{M}^\top \\ -\mathbf{M} & (1/\gamma_4)\mathbf{I} \end{bmatrix}.$$

Since  $\mathbf{\Pi}$  is the permutation matrix, we have

$$\mathbf{P} \succ \mathbf{O} \Leftrightarrow \mathbf{P}_1 \succ \mathbf{O}, \mathbf{P}_2 \succ \mathbf{O}, \mathbf{P}_3 \succ \mathbf{O}.$$

The property of the Schur complement [51, Theorem 7.7.7] yields

$$\mathbf{P}_2 \succ \mathbf{O} \Leftrightarrow (1/\gamma_2)\mathbf{I} - \mathbf{I} - \mathbf{M}^\top\mathbf{M} \succ \mathbf{O},$$

$$\mathbf{P}_3 \succ \mathbf{O} \Leftrightarrow (1/\gamma_4)\mathbf{I} - \gamma_3\mathbf{M}\mathbf{M}^\top \succ \mathbf{O},$$

and thus  $\mathbf{P}_2 \succ \mathbf{O}$  and  $\mathbf{P}_3 \succ \mathbf{O}$  are guaranteed by the second and the fourth conditions in (20), respectively. Applying the property of the Schur complement to  $\mathbf{P}_1$ , we obtain

$$\mathbf{P}_1 \succ \mathbf{O} \Leftrightarrow (1/\gamma_1)\mathbf{I} - \lambda\gamma_3\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B})^2\mathbf{L} - \lambda\mathbf{L}^\top\mathbf{L} \succ \mathbf{O} \\ \Leftrightarrow (1/\gamma_1)\mathbf{I} - (\kappa/2)\mathbf{A}^\top\mathbf{A} - \lambda\mathbf{L}^\top\mathbf{L} \\ + (\kappa/2)\mathbf{A}^\top\mathbf{A} - \lambda\gamma_3\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B})^2\mathbf{L} \succ \mathbf{O}.$$

Since  $(1/\gamma_1)\mathbf{I} - (\kappa/2)\mathbf{A}^\top\mathbf{A} - \lambda\mathbf{L}^\top\mathbf{L} \succ \mathbf{O}$  is guaranteed by the first condition in (20), it suffices to prove  $(\kappa/2)\mathbf{A}^\top\mathbf{A} -$

$\lambda\gamma_3\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B})^2\mathbf{L} \succeq \mathbf{O}$ . Since the third condition in (20) implies  $\gamma_3 \leq \kappa/(2\|\mathbf{B}\|_{\text{op}}^2)$ , for any  $\mathbf{x} \in \mathbb{R}^n$ , we obtain

$$\begin{aligned} \langle \mathbf{x}, \gamma_3\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B})^2\mathbf{L}\mathbf{x} \rangle &= \gamma_3\|\mathbf{B}^\top\mathbf{B}\mathbf{L}\mathbf{x}\|^2 \\ &\leq \gamma_3\|\mathbf{B}\|_{\text{op}}^2\|\mathbf{B}\mathbf{L}\mathbf{x}\|^2 \leq (\kappa/2)\|\mathbf{B}\mathbf{L}\mathbf{x}\|^2. \end{aligned}$$

Thus, we have

$$\begin{aligned} (\forall \mathbf{x} \in \mathbb{R}^n) \quad \langle \mathbf{x}, ((\kappa/2)\mathbf{A}^\top\mathbf{A} - \lambda\gamma_3\mathbf{L}^\top(\mathbf{B}^\top\mathbf{B})^2\mathbf{L})\mathbf{x} \rangle \\ \geq (\kappa/2)\langle \mathbf{x}, (\mathbf{A}^\top\mathbf{A} - \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}\mathbf{L})\mathbf{x} \rangle \geq 0, \end{aligned}$$

where the last inequality follows from the condition (12).

## 2) Averaged nonexpansiveness of $\text{Id} - \mathbf{P}^{-1} \circ H$

To prove the  $\frac{1}{\kappa}$ -averaged nonexpansiveness of  $\text{Id} - \mathbf{P}^{-1} \circ H$ , it suffices to prove that  $\text{Id} - \kappa\mathbf{P}^{-1} \circ H$  is nonexpansive, in  $(\mathcal{H}, \langle \cdot, \cdot \rangle_{\mathbf{P}}, \|\cdot\|_{\mathbf{P}})$ . Define  $\mathbf{R} \in \mathbb{R}^{(n+2m) \times (n+2m)}$  by

$$\mathbf{R} := \begin{bmatrix} \mathbf{A}^\top\mathbf{A} - \lambda\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}\mathbf{L} & & & \\ & \lambda\mathbf{B}^\top\mathbf{B} & & \\ & & \mathbf{O} & \\ & & & \mathbf{O} \end{bmatrix}.$$

Then,  $\mathbf{R} \succeq \mathbf{O}$  holds by the condition (12). For  $\mathbf{z} = (\mathbf{x}, \boldsymbol{\sigma}, \mathbf{v}, \boldsymbol{\tau}, \mathbf{r}, \boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\zeta}) \in \mathcal{H}$ , denote  $\check{\mathbf{z}} := (\mathbf{x}, \mathbf{v}, \mathbf{r}) \in \mathbb{R}^{n+2m}$ . From the definitions of  $H$  and  $\mathbf{\Pi}$  in (25) and (28), respectively, and the expression of  $\mathbf{P}$  in (29), for any  $\mathbf{z}_1, \mathbf{z}_2 \in \mathcal{H}$ , we have

$$H(\mathbf{z}_1) - H(\mathbf{z}_2) = \mathbf{\Pi}^\top(\mathbf{R}(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2), \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}),$$

$$\mathbf{P}^{-1}[H(\mathbf{z}_1) - H(\mathbf{z}_2)] = \mathbf{\Pi}^\top(\mathbf{P}_1^{-1}\mathbf{R}(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2), \mathbf{0}, \mathbf{0}, \mathbf{0}, \mathbf{0}).$$

Using these expressions, for any  $\mathbf{z}_1, \mathbf{z}_2 \in \mathcal{H}$ , we obtain

$$\begin{aligned} &\|(\text{Id} - \kappa\mathbf{P}^{-1} \circ H)(\mathbf{z}_1) - (\text{Id} - \kappa\mathbf{P}^{-1} \circ H)(\mathbf{z}_2)\|_{\mathbf{P}}^2 \\ &= \|(\mathbf{z}_1 - \mathbf{z}_2) - \kappa\mathbf{P}^{-1}[H(\mathbf{z}_1) - H(\mathbf{z}_2)]\|_{\mathbf{P}}^2 \\ &= \|\mathbf{z}_1 - \mathbf{z}_2\|_{\mathbf{P}}^2 - 2\kappa\langle \mathbf{z}_1 - \mathbf{z}_2, H(\mathbf{z}_1) - H(\mathbf{z}_2) \rangle \\ &\quad + \kappa^2\langle \mathbf{P}^{-1}[H(\mathbf{z}_1) - H(\mathbf{z}_2)], H(\mathbf{z}_1) - H(\mathbf{z}_2) \rangle \\ &= \|\mathbf{z}_1 - \mathbf{z}_2\|_{\mathbf{P}}^2 - 2\kappa\langle \check{\mathbf{z}}_1 - \check{\mathbf{z}}_2, \mathbf{R}(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2) \rangle \\ &\quad + \kappa^2\langle \mathbf{P}_1^{-1}\mathbf{R}(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2), \mathbf{R}(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2) \rangle \\ &= \|\mathbf{z}_1 - \mathbf{z}_2\|_{\mathbf{P}}^2 - 2\kappa\langle \check{\mathbf{z}}_1 - \check{\mathbf{z}}_2, (\mathbf{R} - \frac{\kappa}{2}\mathbf{R}\mathbf{P}_1^{-1}\mathbf{R})(\check{\mathbf{z}}_1 - \check{\mathbf{z}}_2) \rangle. \end{aligned}$$

Thus,  $\text{Id} - \kappa\mathbf{P}^{-1} \circ H$  is nonexpansive if and only if  $\mathbf{R} - \frac{\kappa}{2}\mathbf{R}\mathbf{P}_1^{-1}\mathbf{R} \succeq \mathbf{O}$ . Let  $\mathbf{R}^\dagger$  denote the Moore-Penrose pseudo-inverse of  $\mathbf{R}$ . By the property of the generalized Schur complement [52, Theorem 1.20], since  $\mathbf{R}\mathbf{R}^\dagger\mathbf{R} = \mathbf{R}$ , we have

$$\mathbf{R} - \frac{\kappa}{2}\mathbf{R}\mathbf{P}_1^{-1}\mathbf{R} \succeq \mathbf{O} \Leftrightarrow \begin{bmatrix} \mathbf{R} & \mathbf{R} \\ \mathbf{R} & \frac{2}{\kappa}\mathbf{P}_1 \end{bmatrix} \succeq \mathbf{O} \Leftrightarrow \mathbf{P}_1 - \frac{\kappa}{2}\mathbf{R} \succeq \mathbf{O}.$$

We can rewrite  $\mathbf{P}_1 - \frac{\kappa}{2}\mathbf{R}$  as

$$\begin{aligned} \mathbf{P}_1 - \frac{\kappa}{2}\mathbf{R} &= \begin{bmatrix} \frac{1}{\gamma_1}\mathbf{I} - \frac{\kappa}{2}\mathbf{A}^\top\mathbf{A} & \mathbf{O} & -\lambda\mathbf{L}^\top \\ \mathbf{O} & \mathbf{O} & \mathbf{O} \\ -\lambda\mathbf{L} & \mathbf{O} & \lambda\mathbf{I} \end{bmatrix} \\ &\quad + \lambda \begin{bmatrix} \frac{\kappa}{2}\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}\mathbf{L} & -\mathbf{L}^\top\mathbf{B}^\top\mathbf{B} & \mathbf{O} \\ -\mathbf{B}^\top\mathbf{B}\mathbf{L} & \frac{1}{\gamma_3}\mathbf{I} - \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} & \mathbf{O} \\ \mathbf{O} & \mathbf{O} & \mathbf{O} \end{bmatrix}, \end{aligned}$$

and thus it suffices to prove that each term on the right-hand side is positive semidefinite. The first term is positive semidefinite by the first condition in (20) and the property of

the Schur complement. For the second term, we have

$$\begin{aligned} &\begin{bmatrix} \frac{\kappa}{2}\mathbf{L}^\top\mathbf{B}^\top\mathbf{B}\mathbf{L} & -\mathbf{L}^\top\mathbf{B}^\top\mathbf{B} \\ -\mathbf{B}^\top\mathbf{B}\mathbf{L} & \frac{1}{\gamma_3}\mathbf{I} - \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{L} & \\ & \mathbf{I} \end{bmatrix}^\top \begin{bmatrix} \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} & -\mathbf{B}^\top\mathbf{B} \\ -\mathbf{B}^\top\mathbf{B} & \frac{1}{\gamma_3}\mathbf{I} - \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} \end{bmatrix} \begin{bmatrix} \mathbf{L} & \\ & \mathbf{I} \end{bmatrix}, \end{aligned}$$

and by the property of the generalized Schur complement,

$$\begin{aligned} &\begin{bmatrix} \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} & -\mathbf{B}^\top\mathbf{B} \\ -\mathbf{B}^\top\mathbf{B} & \frac{1}{\gamma_3}\mathbf{I} - \frac{\kappa}{2}\mathbf{B}^\top\mathbf{B} \end{bmatrix} \succeq \mathbf{O} \\ &\Leftrightarrow (1/\gamma_3)\mathbf{I} - (\kappa/2)\mathbf{B}^\top\mathbf{B} - (2/\kappa)\mathbf{B}^\top\mathbf{B} \succeq \mathbf{O}. \end{aligned}$$

Thus, the second term is positive semidefinite by the third condition in (20).

## REFERENCES

- [1] R. Tibshirani, "Regression shrinkage and selection via the lasso," *J. Roy. Statist. Soc. Ser. B*, vol. 58, no. 1, pp. 267–288, 1996.
- [2] S. S. Chen, D. L. Donoho, and M. A. Saunders, "Atomic decomposition by basis pursuit," *SIAM Rev.*, vol. 43, no. 1, pp. 129–159, 2001.
- [3] R. G. Baraniuk, "Compressive sensing [lecture notes]," *IEEE Signal Process. Mag.*, vol. 24, no. 4, pp. 118–121, Jul. 2007.
- [4] E. J. Candes and M. B. Wakin, "An introduction to compressive sampling," *IEEE Signal Process. Mag.*, vol. 25, no. 2, pp. 21–30, Mar. 2008.
- [5] A. M. Bruckstein, D. L. Donoho, and M. Elad, "From sparse solutions of systems of equations to sparse modeling of signals and images," *SIAM Rev.*, vol. 51, no. 1, pp. 34–81, 2009.
- [6] J. Fan and R. Li, "Variable selection via nonconcave penalized likelihood and its oracle properties," *J. Amer. Statist. Assoc.*, vol. 96, no. 456, pp. 1348–1360, 2001.
- [7] C.-H. Zhang, "Nearly unbiased variable selection under minimax concave penalty," *Ann. Statist.*, vol. 38, no. 2, pp. 894–942, 2010.
- [8] A. Blake and A. Zisserman, *Visual reconstruction*. MIT press, 1987.
- [9] M. Nikolova, "Markovian reconstruction using a GNC approach," *IEEE Trans. Signal Process.*, vol. 8, no. 9, pp. 1204–1220, 1999.
- [10] I. W. Selesnick and I. Bayram, "Sparse signal estimation by maximally sparse convex optimization," *IEEE Trans. Signal Process.*, vol. 62, no. 5, pp. 1078–1092, 2014.
- [11] T. Mollenhoff, E. Strekalovskiy, M. Moeller, and D. Cremers, "The primal-dual hybrid gradient method for semiconvex splittings," *SIAM J. Imaging Sci.*, vol. 8, no. 2, pp. 827–857, 2015.
- [12] I. Bayram, "On the convergence of the iterative shrinkage/thresholding algorithm with a weakly convex penalty," *IEEE Trans. Signal Process.*, vol. 64, no. 6, pp. 1597–1608, 2016.
- [13] M. Malek-Mohammadi, C. R. Rojas, and B. Wahlberg, "A class of nonconvex penalties preserving overall convexity in optimization-based mean filtering," *IEEE Trans. Signal Process.*, vol. 64, no. 24, pp. 6650–6664, 2016.
- [14] I. Selesnick, "Sparse regularization via convex analysis," *IEEE Trans. Signal Process.*, vol. 65, no. 17, pp. 4481–4494, 2017.
- [15] L. Yin, A. Parekh, and I. Selesnick, "Stable principal component pursuit via convex analysis," *IEEE Trans. Signal Process.*, vol. 67, no. 10, pp. 2595–2607, 2019.
- [16] A. Lanza, S. Morigi, I. W. Selesnick, and F. Sgallari, "Sparsity-inducing nonconvex nonseparable regularization for convex image processing," *SIAM J. Imaging Sci.*, vol. 12, no. 2, pp. 1099–1134, 2019.
- [17] J. Abe, M. Yamagishi, and I. Yamada, "Linearly involved generalized Moreau enhanced models and their proximal splitting algorithm under overall convexity condition," *Inverse Problems*, vol. 36, no. 3, Feb. 2020, Art. no. 035012.
- [18] A. H. Al-Shabli, Y. Feng, and I. Selesnick, "Sharpening sparse regularizers via smoothing," *IEEE Open J. Signal Process.*, vol. 2, pp. 396–409, 2021.
- [19] M. Yukawa, H. Kaneko, K. Suzuki, and I. Yamada, "Linearly-involved Moreau-enhanced-over-subspace model: Debiased sparse modeling and stable outlier-robust regression," *IEEE Trans. Signal Process.*, vol. 71, pp. 1232–1247, 2023.
- [20] D. Kitahara, R. Kato, H. Kuroda, and A. Hirabayashi, "Multi-contrast CSMRI using common edge structures with LiGME model," in *Proc. Eur. Signal Process. Conf.*, 2021, pp. 2119–2123.

- [21] W. Yata, M. Yamagishi, and I. Yamada, "A constrained LiGME model and its proximal splitting algorithm under overall convexity condition," *J. Appl. Numer. Optim.*, vol. 4, no. 2, pp. 245–271, 2022.
- [22] G. Yu, S. Mallat, and E. Bacry, "Audio denoising by time-frequency block thresholding," *IEEE Trans. Signal Process.*, vol. 56, no. 5, pp. 1830–1839, May 2008.
- [23] V. Cevher, A. Sankaranarayanan, M. F. Duarte, D. Reddy, R. G. Baraniuk, and R. Chellappa, "Compressive sensing for background subtraction," in *Proc. Eur. Conf. Comput. Vis.*, 2008, pp. 155–168.
- [24] Z. Gao, L. Cheong, and Y. Wang, "Block-sparse RPCA for salient motion detection," *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. 36, no. 10, pp. 1975–1987, Oct. 2014.
- [25] S. Hur, S. Baek, B. Kim, Y. Chang, A. F. Molisch, T. S. Rappaport, K. Haneda, and J. Park, "Proposal on millimeter-wave channel modeling for 5G cellular system," *IEEE J. Sel. Topics in Signal Process.*, vol. 10, no. 3, pp. 454–469, 2016.
- [26] H. Kuroda, M. Yamagishi, and I. Yamada, "Exploiting sparsity in tight-dimensional spaces for piecewise continuous signal recovery," *IEEE Trans. Signal Process.*, vol. 66, no. 24, pp. 6363–6376, Dec. 2018.
- [27] D. Kitahara, H. Kuroda, A. Hirabayashi, E. Yoshikawa, H. Kikuchi, and T. Ushio, "Nonlinear beamforming based on group-sparsities of periodograms for phased array weather radar," *IEEE Trans. Geosci. Remote Sens.*, 2022, Art. no. 4106819.
- [28] H. Kuroda, D. Kitahara, E. Yoshikawa, H. Kikuchi, and T. Ushio, "Convex estimation of sparse-smooth power spectral densities from mixtures of realizations with application to weather radar," *IEEE Access*, vol. 11, pp. 128 859–128 874, 2023.
- [29] H. Kuroda and D. Kitahara, "Block-sparse recovery with optimal block partition," *IEEE Trans. Signal Process.*, vol. 70, pp. 1506–1520, 2022.
- [30] M. Yuan and Y. Lin, "Model selection and estimation in regression with grouped variables," *J. Roy. Statist. Soc. Ser. B*, vol. 68, no. 1, pp. 49–67, 2006.
- [31] K. Bredies, K. Kunisch, and T. Pock, "Total generalized variation," *SIAM J. Imaging Sci.*, vol. 3, no. 3, pp. 492–526, 2010.
- [32] L. I. Rudin, S. Osher, and E. Fatemi, "Nonlinear total variation based noise removal algorithms," *Physica D*, vol. 60, no. 1, pp. 259–268, 1992.
- [33] F. Knoll, K. Bredies, T. Pock, and R. Stollberger, "Second order total generalized variation (TGV) for MRI," *Magn. Reson. Med.*, vol. 65, no. 2, pp. 480–491, 2011.
- [34] T. Valkonen, K. Bredies, and F. Knoll, "Total generalized variation in diffusion tensor imaging," *SIAM J. Imaging Sci.*, vol. 6, no. 1, pp. 487–525, 2013.
- [35] D. Ferstl, C. Reinbacher, R. Ranftl, M. Ruether, and H. Bischof, "Image guided depth upsampling using anisotropic total generalized variation," in *Proc. IEEE Int. Conf. Comput. Vis.*, Dec. 2013, pp. 993–1000.
- [36] S. Ono and I. Yamada, "Decorrelated vectorial total variation," in *Proc. IEEE Conf. Comput. Vis. Pattern Recognit.*, 2014, pp. 4090–4097.
- [37] C. Langkammer, K. Bredies, B. A. Poser, M. Barth, G. Reishofer, A. P. Fan, B. Bilgic, F. Fazekas, C. Mainero, and S. Ropele, "Fast quantitative susceptibility mapping using 3D EPI and total generalized variation," *NeuroImage*, vol. 111, pp. 622–630, 2015.
- [38] K. Bredies, M. Holler, M. Storath, and A. Weinmann, "Total generalized variation for manifold-valued data," *SIAM J. Imaging Sci.*, vol. 11, no. 3, pp. 1785–1848, 2018.
- [39] K. Bredies and M. Holler, "Higher-order total variation approaches and generalisations," *Inverse Problems*, vol. 36, no. 12, Dec. 2020, Art. no. 123001.
- [40] H. Kuroda, "Sharpening minimization induced penalties," in *Proc. Eur. Signal Process. Conf.*, Aug. 2024, 5 pages. (to appear)
- [41] H. H. Bauschke and P. L. Combettes, *Convex analysis and monotone operator theory in Hilbert spaces*, 2nd ed. Springer, 2011.
- [42] P. L. Combettes, "Perspective functions: Properties, constructions, and examples," *Set-Valued and Variational Analysis*, vol. 26, no. 2, pp. 247–264, 2018.
- [43] J. Huang, P. Breheny, and S. Ma, "A selective review of group selection in high-dimensional models," *Statist. Sci.*, vol. 27, no. 4, pp. 481–499, Nov. 2012.
- [44] Y. Chen, M. Yamagishi, and I. Yamada, "A unified design of generalized Moreau enhancement matrix for sparsity aware LiGME models," *IEICE Trans. Fundam.*, vol. E106.A, no. 8, pp. 1025–1036, 2023.
- [45] P. L. Combettes and J.-C. Pesquet, "Proximal splitting methods in signal processing," in *Fixed-Point Algorithms for Inverse Problems in Science and Engineering*, H. H. Bauschke, R. S. Burachik, P. L. Combettes, V. Elser, D. R. Luke, and H. Wolkowicz, Eds. Springer, 2011, pp. 185–212.
- [46] L. Condat, D. Kitahara, A. Contreras, and A. Hirabayashi, "Proximal splitting algorithms for convex optimization: A tour of recent advances, with new twists," *SIAM Rev.*, vol. 65, no. 2, pp. 375–435, 2023.
- [47] X. Liu, A. J. Molstad, and E. C. Chi, "A convex-nonconvex strategy for grouped variable selection," *Electron. J. Statist.*, vol. 17, no. 2, pp. 2912–2961, 2023.
- [48] P. L. Combettes and C. L. Müller, "Perspective maximum likelihood-type estimation via proximal decomposition," *Electron. J. Statist.*, vol. 14, no. 1, pp. 207–238, 2020.
- [49] H. H. Bauschke, M. K. Lal, and X. Wang, "Real roots of real cubics and optimization," preprint arXiv:2302.10731, 2023.
- [50] L. Condat, "Fast projection onto the simplex and the  $\ell_1$  ball," *Math. Programm.*, vol. 158, no. 1, pp. 575–585, 2016.
- [51] R. A. Horn and C. R. Johnson, *Matrix analysis*, 2nd ed. Cambridge University Press, 2012.
- [52] R. A. Horn and F. Zhang, "Basic properties of the schur complement," in *The Schur Complement and Its Applications*, F. Zhang, Ed. Boston, MA: Springer US, 2005, pp. 17–46.