

Inverse spectral problem of Sturm-Liouville equation with many frozen arguments

Chung-Tsun Shieh¹ and Tzong-Mo Tsai²

¹Department of Mathematics, Tamkang University, New Taipei City, 25137, Taiwan, ctshieh@mail.tku.edu.tw.

²General Education Center, Ming Chi University of Technology, New Taipei City, 24301, Taiwan, tsaitm@mail.mcut.edu.tw

Abstract

This research was devoted to investigate the inverse spectral problem of Sturm-Liouville operator with many frozen arguments. Under some assumptions, the authors obtained uniqueness theorems. At the end, a numerical simulation for the inverse problem was presented.

Key words: Sturm-Liouville equation, loaded differential equation, frozen argument, eigenvalues, inverse spectral problem.

2010 Mathematics Subject Classification: 34A55 34K29

1 Introduction

In this paper, the authors consider the boundary value problem $L(q(x), F_n, \alpha, \beta)$ which consists of a nonlocal Sturm-Liouville equation

$$(1) \quad -y''(x) + q(x) \sum_{i=1}^n y(a_i) = \lambda y(x), \quad 0 < x < \pi,$$

associated with the boundary conditions

$$(2) \quad y^{(\alpha)}(0) = 0 = y^{(\beta)}(\pi),$$

where λ is the spectral parameter, $F_n = \{a_i\}_{i=1}^n$, $0 < a_i < a_j < \pi$ if $i < j$, is the set of frozen arguments, $\alpha, \beta \in \{0, 1\}$ and $q(x) \in L^2(0, 1)$. Since the analyses for the problems of all cases of $\{\alpha, \beta\}$ which we concerned are similar, we shall only treat the case $\alpha = 0 = \beta$.

As applications, equation (1) occurs in the class of the so-called loaded equations [1, 2], which contain the values of the unknown function at some fixed points. Such equations have been applied for studying grounder dynamics [1, 3], heating processes [4, 5], feedback-like phenomena in vibrations of a wire affected by a magnetic field [6, 7].

Let $\{\lambda_m\}_{m \geq 1}$ be the eigenvalues of $L(q(x), F_n, \alpha, \beta)$. In this paper, the authors studied the following inverse problem:

Inverse Problem 1. Given $\{\lambda_m\}_{m \geq 1}$, F_n , α and β , determine the potential $q(x)$.

Inverse Problem 1 of the case $n = 1$ had been studied by many researchers [8, 9, 10, 11, 12, 13]. In [8, 9, 10, 16], diverse cases of the triple (a_1, α, β) with rational $a_1 = j/k$, $\gcd(k, j) = 1$, were considered. In particular, it was established that the inverse problem may be uniquely solvable or not, depending on the parameters α, β and also on (k, j) . Concerning irrational values of a_1 , it was established by Y. P. Wang et al in [11] that all pairs (α, β) , the solution of Inverse Problem 1 for $n = 1$ is always uniquely determined by the spectrum set. Recently Maria Kuznetsova [12, 13] obtained a unified asymptotic formula covering both irrational and rational cases.

The paper is organized as follows. In the next section, the uniqueness theorems were established. In section 3, numerical algorithm and simulations were presented.

2 Main Results

In this section, we shall present our approach which is different from that in [8, 9, 10] and [11]. We shall write $\lambda = \rho^2$.

Lemma 2.1. Denote $S(x, \lambda)$ the solution of (1) associated with the initial condition

$$y(0, \lambda) = 0, \quad y'(0, \lambda) = 1.$$

Then $S(x, \lambda)$ satisfies

$$(3) \quad S(x, \lambda) = \frac{\sin \rho x}{\rho} + \int_0^x \frac{\sin \rho(x-t)}{\rho} q(t) \left(\sum_{i=1}^n S(a_i, \lambda) \right) dt$$

The derivation is not complicated, we shall omit the details.

Next, we shall start with the simplest case $n = 1$ to show (3) works for our study. For $n = 1$, (3) turns to

$$(4) \quad S(x, \lambda) = \frac{\sin \rho x}{\rho} + S(a_1, \lambda) \int_0^x \frac{\sin \rho(x-t)}{\rho} q(t) dt.$$

Hence we have

$$(5) \quad \begin{cases} S(a_1, \lambda) = \frac{\sin \rho a_1}{\rho} + S(a_1, \lambda) \int_0^{a_1} \frac{\sin \rho(a_1-t)}{\rho} q(t) dt, \\ S(\pi, \lambda) = \frac{\sin \rho \pi}{\rho} + S(a_1, \lambda) \int_0^{\pi} \frac{\sin \rho(\pi-t)}{\rho} q(t) dt, \end{cases}$$

that is

$$(6) \quad \begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt - 1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^{\pi} q(t) \frac{\sin \rho(\pi-t)}{\rho} dt \end{bmatrix} \begin{bmatrix} 1 \\ S(a_1, \lambda) \end{bmatrix} = \begin{bmatrix} 0 \\ S(\pi, \lambda) \end{bmatrix}.$$

Thus λ is an eigenvalue of $L(q(x), F_1, 0, 0)$ if and only if λ is a zero of the characteristic function

$$(7) \quad \Delta_1(\lambda) = \det \left(\begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt - 1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^{\pi} q(t) \frac{\sin \rho(\pi-t)}{\rho} dt \end{bmatrix} \right).$$

One can easily see that

$$(8) \quad \Delta_1(\lambda) = \frac{\sin \rho \pi}{\rho} + \frac{\sin \rho a_1}{\rho} \int_0^{\pi} q(t) \frac{\sin \rho(\pi-t)}{\rho} dt - \frac{\sin \rho \pi}{\rho} \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt$$

After some calculation, we have

$$(9) \quad \Delta_1(\lambda) = \frac{\sin \rho \pi}{\rho} + \frac{\sin \rho a_1}{\rho^2} \int_{a_1}^{\pi} q(t) \sin \rho(\pi-t) dt + \frac{\sin \rho(\pi-a_1)}{\rho^2} \int_0^{a_1} q(t) \sin \rho t dt.$$

The right side of (9) is the form of characteristic function which appeared in [11].

To go one step further, we shall consider the case for $n = 2$. Then

$$(10) \quad S(x, \lambda) = \frac{\sin \rho x}{\rho} + (S(a_1, \lambda) + S(a_2, \lambda)) \int_0^x q(t) \frac{\sin \rho(x-t)}{\rho} dt.$$

Hence we have

$$(11) \quad \begin{cases} S(a_1, \lambda) = \frac{\sin \rho a_1}{\rho} + (S(a_1, \lambda) + S(a_2, \lambda)) \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt, \\ S(a_2, \lambda) = \frac{\sin \rho a_2}{\rho} + (S(a_1, \lambda) + S(a_2, \lambda)) \int_0^{a_2} q(t) \frac{\sin \rho(a_2-t)}{\rho} dt, \\ S(\pi, \lambda) = \frac{\sin \rho \pi}{\rho} + (S(a_1, \lambda) + S(a_2, \lambda)) \int_0^{\pi} q(t) \frac{\sin \rho(\pi-t)}{\rho} dt. \end{cases}$$

(11) leads to

$$\begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1 - t)}{\rho} dt - 1 & \int_0^{a_1} q(t) \frac{\sin \rho(a_1 - t)}{\rho} dt \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} q(t) \frac{\sin \rho(a_2 - t)}{\rho} dt & \int_0^{a_2} q(t) \frac{\sin \rho(a_2 - t)}{\rho} dt - 1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi q(t) \frac{\sin \rho(\pi - t)}{\rho} dt & \int_0^\pi q(t) \frac{\sin \rho(\pi - t)}{\rho} dt \end{bmatrix} \begin{bmatrix} 1 \\ S(a_1, \lambda) \\ S(a_2, \lambda) \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ S(\pi, \lambda) \end{bmatrix}.$$

Thus λ is an eigenvalue of $L(q(x), F_2, 0, 0)$ if and only if λ is a zero of the characteristic function

$$(12) \quad \Delta_2(\lambda) = \det \left(\begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1 - t)}{\rho} dt - 1 & \int_0^{a_1} q(t) \frac{\sin \rho(a_1 - t)}{\rho} dt \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} q(t) \frac{\sin \rho(a_2 - t)}{\rho} dt & \int_0^{a_2} q(t) \frac{\sin \rho(a_2 - t)}{\rho} dt - 1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi q(t) \frac{\sin \rho(\pi - t)}{\rho} dt & \int_0^\pi q(t) \frac{\sin \rho(\pi - t)}{\rho} dt \end{bmatrix} \right).$$

Note that one achieves

$$(13) \quad \begin{aligned} \Delta_2(\lambda) &= \begin{vmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1 - t)}{\rho} dt - 1 & 1 \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} q(t) \frac{\sin \rho(a_2 - t)}{\rho} dt & -1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi q(t) \frac{\sin \rho(\pi - t)}{\rho} dt & 0 \end{vmatrix} \\ &= \frac{\sin \rho \pi}{\rho} + \frac{\sin \rho a_1}{\rho^2} \int_0^\pi q(t) \sin \rho(\pi - t) dt - \frac{\sin \rho \pi}{\rho^2} \int_0^{a_1} q(t) \sin \rho(a_1 - t) dt \\ &\quad + \frac{\sin \rho a_2}{\rho^2} \int_0^\pi q(t) \sin \rho(\pi - t) dt - \frac{\sin \rho \pi}{\rho^2} \int_0^{a_2} q(t) \sin \rho(a_2 - t) dt \end{aligned}$$

after a simplification of (12). Denote $\Lambda(q(x), F_2) = \{\lambda_m\}_{m=1}^\infty$ the eigenvalues of $L(q(x), F_2, 0, 0)$, where $F_2 = \{a_1, a_2\}$. By Hadamard's factorization theorem, we have

$$(14) \quad \Delta_2(\lambda) = \pi \prod_{n=1}^\infty \frac{(\lambda_n - \lambda)}{n^2}.$$

From the analysis above, we have following uniqueness theorem

Theorem 2.2. *Let $F_2 = \{a_1, a_2\}$, where $0 < a_1 < a_2 < \pi$. Define $G_2(\rho) = \sin(\rho a_1) + \sin(\rho a_2)$, and denote $Z(G_2)$ the set of zeros of $G_2(\rho)$. Suppose $Z(G_2) \cap \mathbb{N} = \emptyset$ and $\Lambda(q(x), F_2) = \Lambda(\tilde{q}(x), F_2)$. Then $q(x) = \tilde{q}(x)$ almost everywhere in $(0, \pi)$.*

Note if $(a_1 \pm a_2)/\pi$ are irrational, then $Z(G_2) \cap \mathbb{N} = \emptyset$.

Proof. Denote $\hat{q}(x) = q(x) - \tilde{q}(x)$, $\Delta_2(\lambda)$ the characteristic function of $L(q(x), F_2, 0, 0)$ and $\tilde{\Delta}_2(\lambda)$ the characteristic function of $L(\tilde{q}(x), F_2, 0, 0)$. According to (13) and (14), one can obtain

$$(15) \quad \Delta_2(\lambda) - \tilde{\Delta}_2(\lambda) = 0 = \begin{vmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} \hat{q}(t) \frac{\sin \rho(a_1 - t)}{\rho} dt & 1 \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} \hat{q}(t) \frac{\sin \rho(a_2 - t)}{\rho} dt & -1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi \hat{q}(t) \frac{\sin \rho(\pi - t)}{\rho} dt & 0 \end{vmatrix}$$

The assumption $Z(G_2) \cap \mathbb{N} = \emptyset$ ensure $(\frac{\sin m a_1}{m}, \frac{\sin m a_2}{m}, \frac{\sin m \pi}{m})^t$ and $(1, -1, 0)^t$ are independent, we have

$$c_m \int_0^\pi \hat{q}(t) \sin m(\pi - t) dt = 0$$

for some $c_m \neq 0$ and for all $m \in \mathbb{N}$. We have $\hat{q}(t) = 0$ almost everywhere in $(0, \pi)$. This completes the proof. \square

Apply the same arguments, Theorem 2.2 can be generalized as following

Theorem 2.3. Let $F_n = \{a_1, a_2, \dots, a_n\}$, where $0 < a_1 < a_2 < \dots < a_n < \pi$. Define $G_n(\rho) = \sum_{i=1}^n \sin(\rho a_i)$. Suppose the following conditions

(i) $Z(G_n) \cap \mathbb{N} = \emptyset$,

(ii) $\Lambda(q(x), F_n) = \Lambda(\tilde{q}(x), F_n)$

are satisfied. Then $q(x) = \tilde{q}(x)$ almost everywhere in $(0, \pi)$. In particular for $n = 1$, condition (1) is equivalent to “ $\pi/a_1 \notin \mathbb{Q}$.”

Proof. The proof follows the same idea as that in the proof of Theorem 2.2 except the computation is slightly complicated. Since

$$S(x, \lambda) = \frac{\sin \rho x}{\rho} + \left(\sum_{i=1}^n S(a_i, \lambda) \right) \int_0^x q(t) \frac{\sin \rho(x-t)}{\rho} dt$$

we have

$$(16) \quad 0 = \frac{\sin \rho a_j}{\rho} + \sum_{i \neq j} S(a_i, \lambda) \int_0^{a_j} q(t) \frac{\sin \rho(a_j-t)}{\rho} dt + S(a_j, \lambda) \left(\int_0^{a_j} q(t) \frac{\sin \rho(a_j-t)}{\rho} dt - 1 \right)$$

and

$$(17) \quad S(\pi, \lambda) = \frac{\sin \rho \pi}{\rho} + \left(\sum_{i=1}^n S(a_i, \lambda) \right) \int_0^\pi q(t) \frac{\sin \rho(\pi-t)}{\rho} dt.$$

According (16) and (17), the characteristic function is

$$(18) \quad \Delta_n(\lambda) = \det(K_n(\lambda))$$

where

$$K_n(\lambda) = \begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt - 1 & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt & \dots & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} q(t) \frac{\sin \rho(a_2-t)}{\rho} dt & \int_0^{a_2} q(t) \frac{\sin \rho(a_2-t)}{\rho} dt - 1 & \dots & \int_0^{a_2} q(t) \frac{\sin \rho(a_2-t)}{\rho} dt \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\sin \rho a_n}{\rho} & \int_0^{a_n} q(t) \frac{\sin \rho(a_n-t)}{\rho} dt & \int_0^{a_n} q(t) \frac{\sin \rho(a_n-t)}{\rho} dt & \dots & \int_0^{a_n} q(t) \frac{\sin \rho(a_n-t)}{\rho} dt - 1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi q(t) \frac{\sin \rho(\pi-t)}{\rho} dt & \int_0^\pi q(t) \frac{\sin \rho(\pi-t)}{\rho} dt & \dots & \int_0^\pi q(t) \frac{\sin \rho(\pi-t)}{\rho} dt \end{bmatrix}$$

(18) can be simplified as

$$(19) \quad \Delta_n(\lambda) = \det(E_n(\lambda))$$

where

$$E_n(\lambda) = \begin{bmatrix} \frac{\sin \rho a_1}{\rho} & \int_0^{a_1} q(t) \frac{\sin \rho(a_1-t)}{\rho} dt - 1 & 1 & 1 & 1 & \dots & 1 & 1 \\ \frac{\sin \rho a_2}{\rho} & \int_0^{a_2} q(t) \frac{\sin \rho(a_2-t)}{\rho} dt & -1 & 0 & 0 & \dots & 0 & 0 \\ \frac{\sin \rho a_3}{\rho} & \int_0^{a_3} q(t) \frac{\sin \rho(a_3-t)}{\rho} dt & 0 & -1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & 0 & 0 \\ \frac{\sin \rho a_n}{\rho} & \int_0^{a_n} q(t) \frac{\sin \rho(a_n-t)}{\rho} dt & 0 & 0 & 0 & \dots & 0 & -1 \\ \frac{\sin \rho \pi}{\rho} & \int_0^\pi q(t) \frac{\sin \rho(\pi-t)}{\rho} dt & 0 & 0 & 0 & 0 \dots & 0 & 0 \end{bmatrix}.$$

Since $\Lambda(q(x), F_n) = \Lambda(\tilde{q}(x), F_n)$, the corresponding characteristic function $\Delta_n(\lambda) = \tilde{\Delta}_n(\lambda)$. This yields to

$$\Delta_n(\lambda) - \tilde{\Delta}_n(\lambda) = 0 = \det(H_n(\lambda)),$$

where $H_n(\lambda)$ is the matrix obtained by substituting $\hat{q}(x) = q(x) - \tilde{q}(x)$ for $q(x)$ in $E_n(\lambda)$. Thus

$$\int_0^\pi \hat{q}(t) \sin m(\pi-t) dt = 0$$

for all $m \in \mathbb{N}$. This yields to $\hat{q}(t) = 0$, which completes the proof. \square

3 numerical simulations

In this section, we shall present a numerical algorithm for reconstruction of penitential. For simplicity, we present an example for a Sturm-Liouville operator with two frozen arguments. We shall stat the algorithm first.

Algorithm. For a given spectral set $\{\lambda_n\}_{n=1}^{\infty}$ of some $L(q(x), F_2, 0, 0)$, construct the potential $q(x)$ from $\{\lambda_n\}_{n=1}^{\infty}$, where F_2 satisfies the assumption in Theorem 2.2.

One can reconstruct $q(x)$ according to the following steps.

- (1) Construct $\Delta_2(\lambda) = \pi \prod_{n=1}^{\infty} \frac{\lambda_n - \lambda}{n^2}$. In practical, we can only have a limited number of eigenvalues

$$\{\lambda_k\}_{k=1}^n. \text{ We shall substitute } \Delta_2(\lambda) \text{ with } \prod_{k=1}^n \frac{\lambda_k - \lambda}{k^2} \left(\frac{\frac{\sin \sqrt{\lambda} \pi}{\sqrt{\lambda}}}{\prod_{k=1}^n \frac{k^2 - \lambda}{k^2}} \right).$$

- (2) Apply (13), we have

$$d_n = \frac{2}{\pi} \int_0^{\pi} q(t) \sin n(\pi - t) dt = \frac{2}{\pi} \left(\frac{n^2 \Delta(n^2)}{\sin na_1 + \sin na_2} \right)$$

- (3)

$$q(t) = \sum_{n=1}^{\infty} d_n \sin n(\pi - t).$$

Example 1. Suppose we have $a_1 = 1$, $a_2 = \sqrt{2}$ and a partial spectrum $\{\sqrt{\lambda_1} = 1.98868, \sqrt{\lambda_2} = 2, \sqrt{\lambda_3} = 3.06656, \sqrt{\lambda_4} = 4, \sqrt{\lambda_5} = 5.00142, \sqrt{\lambda_6} = 6, \sqrt{\lambda_7} = 6.99975, \sqrt{\lambda_8} = 8, \sqrt{\lambda_9} = 8.99976, \sqrt{\lambda_{10}} = 10\}$. For this case, we let

$$\Delta_2(\lambda) = \pi \prod_{n=1}^{10} \left(\frac{\lambda_n - \lambda}{n^2} \right) \cdot \prod_{n=11}^{\infty} \left(\frac{n^2 - \lambda}{n^2} \right).$$

The readers can refer to figure 2. In this case, the convergence is very fast.

Example 2. Let $a_1 = 1$, $a_2 = \sqrt{2}$. A set of partial spectrum $\{\sqrt{\lambda_3} = 2.90145, \sqrt{\lambda_4} = 4.09132, \sqrt{\lambda_5} = 4.98819, \sqrt{\lambda_6} = 5.98425, \sqrt{\lambda_7} = 7.00424, \sqrt{\lambda_8} = 7.99936, \sqrt{\lambda_9} = 9.00732, \sqrt{\lambda_{10}} = 9.99528, \sqrt{\lambda_{11}} = 10.9929, \sqrt{\lambda_{12}} = 12.0105, \sqrt{\lambda_{13}} = 10.9929, \sqrt{\lambda_{14}} = 12.9998, \sqrt{\lambda_{15}} = 13.9907, \sqrt{\lambda_{16}} = 15.006, \sqrt{\lambda_{17}} = 16.0035, \sqrt{\lambda_{18}} = 16.9935, \sqrt{\lambda_{19}} = 18.0013, \sqrt{\lambda_{20}} = 19.0032, \sqrt{\lambda_{21}} = 19.9997, \sqrt{\lambda_{22}} = 20.9997\} \cup \{1.99593 \pm 0.4925i\}$ was obtained. In this case, a pair of complex eigenvalues appeared. In this case, we let

$$\Delta_2(\lambda) = \frac{((1.99593 + 0.4925i)^2 - x) \times ((1.99593 - 0.4925i)^2 - x)}{4} \cdot \prod_{n=3}^{21} \left(\frac{\lambda_n - \lambda}{n^2} \right) \cdot \prod_{n=22}^{\infty} \left(\frac{n^2 - \lambda}{n^2} \right).$$

The readers can refer to figure3 for the result of simulation, the convergence is poor and The Gibbs phenomenon appears.




Figure 2: An approximated solution for the potential $q(t) = 1 - \cos(2t)$.




Figure 3: Approximated solutions for the potential $q(t) = t$.

References

- [1] Nakhushev A.M. *An approximate method for solving boundary value problems for differential equations and its application to the dynamics of ground moisture and ground water*, Differ. Uravn. 18 (1982) no.1, 72–81.
- [2] Nakhushev A.M. *Loaded Equations and Their Applications*, Nauka, Moscow, 2012.
- [3] Nakhushev A.M. and Borisov V.N. *Boundary value problems for loaded parabolic equations and their applications to the prediction of ground water level*, Differ. Uravn. 13 (1977) no.1, 105–110.
- [4] Iskenderov A.D. *The first boundary-value problem for a loaded system of quasilinear parabolic equations*, Differ. Uravn. 7 (1971) no.10, 1911–1913.
- [5] Dikinov Kh.Zh., Kerefov A.A. and Nakhushev A.M. *A certain boundary value problem for a loaded heat equation*, Differ. Uravn. 12 (1976) no.1, 177–179.
- [6] Krall A.M. *The development of general differential and general differential-boundary systems*, Rock. Mount. J. Math. 5 (1975) no.4, 493–542.
- [7] Buterin S. and Hu Y.-T. *Inverse spectral problems for Hill-type operators with frozen argument*, Anal. Math. Phys. (2021) 11:75, 22pp.
- [8] Bondarenko N.P., Buterin S.A. and Vasiliev S.V. *An inverse spectral problem for Sturm–Liouville operators with frozen argument*, J. Math. Anal. Appl. 472 (2019) 1028–1041.
- [9] Buterin S.A. and Vasiliev S.V. *On recovering Sturm–Liouville operators with frozen argument*, J. Inv. Ill-Posed Probl. 27 (2019) no.3, 429–438.
- [10] Buterin S. and Kuznetsova M. *On the inverse problem for Sturm–Liouville-type operators with frozen argument: rational case*, Comp. Appl. Math. (2020) 39:5, 15pp.
- [11] Wang Y.P., Zhang M., Zhao W. and Wei X. *Reconstruction for Sturm–Liouville operators with frozen argument for irrational cases*, Appl. Math. Lett. 111 (2021) 106590.
- [12] Kuznetsova M.A. *Characterization of the spectrum of a Sturm–Liouville operator with frozen argument*, Proc. Int. Sci. Conf. "Ufa Autumn Mathematical School" (October 6–9, 2021, Bashkir State University, Ufa), V.1, Ufa, Aeterna, 2021, pp. 48–50.
- [13] Kuznetsova M.A. *Necessary and sufficient conditions for the spectra of the Sturm–Liouville operators with frozen argument*, Appl. Math. Lett. 131 (2022) 108035.
- [14] Albeverio S., Hryniv R.O. and Nizhnik L.P. *Inverse spectral problems for non-local Sturm–Liouville operators*, Inverse Problems 23 (2007) 523–535.
- [15] Nizhnik L.P. *Inverse eigenvalue problems for nonlocal Sturm–Liouville operators*, Meth. Func. Anal. Top. 15 (2009) no.1, 41–47.
- [16] Tsai T.-M., Liu H.-F., Buterin S., Chen L.-H. and Shieh C.-T. *Sturm–Liouville-type operators with frozen argument and Chebyshev polynomials*, Math. Meth. Appl. Soc., 45(2022),