Nonparametric Estimation of Ordinary Differential Equations: Snake and Stubble

Christof Schötz* 1,2

¹Potsdam Institute for Climate Impact Research ²Technical University of Munich

Abstract

We study nonparametric estimation in dynamical systems described by ordinary differential equations (ODEs). Specifically, we focus on estimating the unknown function $f: \mathbb{R}^d \to \mathbb{R}^d$ that governs the system dynamics through the ODE $\dot{u}(t) = f(u(t))$, where observations $Y_{j,i} = u_j(t_{j,i}) + \varepsilon_{j,i}$ of solutions u_j of the ODE are made at times $t_{j,i}$ with independent noise $\varepsilon_{j,i}$.

We introduce two novel models – the Stubble model and the Snake model – to mitigate the issue of observation location dependence on f, an inherent difficulty in nonparametric estimation of ODE systems. In the Stubble model, we observe many short solutions with initial conditions that adequately cover the domain of interest. Here, we study an estimator based on multivariate local polynomial regression and univariate polynomial interpolation. In the Snake model we observe few long trajectories that traverse the domain on interest. Here, we study an estimator that combines univariate local polynomial estimation with multivariate polynomial interpolation.

For both models, we establish error bounds of order $n^{-\frac{\beta}{2(\beta+1)+d}}$ for β -smooth functions f in an infinite dimensional function class of Hölder-type and establish minimax optimality for the Stubble model in general and for the Snake model under some conditions via comparison to lower bounds from parallel work.

Keywords: Ordinary Differential Equations, Nonparametric, Upper Bounds, Optimal Rate of Convergence, Minimax

Contents

1	Introduction						
	1.1	Contributions	3				
	1.2	Relation to Other Models	4				
	1.3	Overview	5				
2	Preliminaries						
	2.1	Ordinary Differential Equations	6				
	2.2	Formal Description of the General Model	$\overline{7}$				
	2.3	Need for Restriction of the General Model	8				
	2.4	Standard Regression Model	8				
3	The Stubble Model and Estimation of the Increment Map 10						
	3.1	Lipschitz	10				
		3.1.1 Model	10				
		3.1.2 Estimator	10				
		3.1.3 Result	10				
		3.1.4 Proof	11				
	3.2	General	12				

*math@christof-schoetz.de, 0000-0003-3528-4544

		3.2.1	Model	13
		3.2.2	Estimation	13
		3.2.3	Result	14
		3.2.4	Proof	15
4	The	e Snak	e Model and Estimation of the Observed Solutions	16
	4.1	Lipsch	nitz	16
		4.1.1	Model	16
		4.1.2	Estimator	16
		4.1.3	Result	17
		4.1.4	Proof	18
	4.2	Gener	al	19
		4.2.1	Model	20
		4.2.2	Multivariate Polynomial Interpolation	20
		4.2.3	Estimator	21
		4.2.4	Results	22
		4.2.5	Proof	24
\mathbf{A}	Der	rivative	es	25
A B	Der Loc	rivative al Pol	es ynomial Regression	25 28
A B	Der Loc B.1	rivative al Pol Mode	es ynomial Regression l	25 28 28
A B	Der Loc B.1 B.2	rivative al Pol Mode Estim	es ynomial Regression l	25 28 28 28
A B	Der Loc B.1 B.2 B.3	rivative cal Pol Mode Estim Result	es ynomial Regression l	25 28 28 28 28 29
A B	Der Loc B.1 B.2 B.3 B.4	ivative cal Pol Mode Estim Result Proof	es ynomial Regression l	25 28 28 28 29 30
A B	Der Loc B.1 B.2 B.3 B.4	ivative al Pol Mode Estim Result Proof	es ynomial Regression l	25 28 28 29 30
A B C	Der Loc B.1 B.2 B.3 B.4 Pol	ivative al Pol Mode Estim Result Proof	es ynomial Regression 1 ator ator t ator	 25 28 28 29 30 37
A B C	Der Loc B.1 B.2 B.3 B.4 Pol C.1	ivativo al Pol Mode Estim Result Proof ynomi Univa	es ynomial Regression l	25 28 28 29 30 37 37
A B C	Der Loc B.1 B.2 B.3 B.4 Pol ; C.1 C.2	ivativ al Poly Mode Estim Result Proof ynomi Univa	es ynomial Regression l	25 28 28 29 30 37 37 37
A B C	Der B.1 B.2 B.3 B.4 Pol C.1 C.2 C.3	ivativa al Pol Mode Estim Result Proof ynomi Univa Univa Multi	es ynomial Regression l ator t at Interpolation riate Polynomial Interpolation riate Polynomial Comparison variate Polynomial Interpolation	 25 28 28 29 30 37 37 37 38
A B C	Der Loc B.1 B.2 B.3 B.4 Pol C.1 C.2 C.3 Ord	ivativa al Poly Mode Estim Result Proof ynomi Univa Univa Multir linary	es ynomial Regression l	25 28 28 29 30 37 37 37 38 41
A B C D	Der Loc B.1 B.2 B.3 B.4 Pol C.1 C.2 C.3 Ord D.1	ivativo al Pol Mode Estim Result Proof Univa Univa Univa Multi linary Differ	es ynomial Regression l ator .	25 28 28 29 30 37 37 37 38 41 41
A B C D	Der Loc B.1 B.2 B.3 B.4 Pol C.1 C.2 C.3 Ord D.1 D.2	ivativa al Pol; Mode Estim Result Proof Univa Univa Univa Multir linary Differ Smoo	es ynomial Regression l ator ato	25 28 28 29 30 37 37 37 38 41 41 42

1 Introduction

In statistics, we want to extract information from observations of the world. The workings of the world are described in physics (and many other sciences) with differential equations [Str24]. Hence, statistical models of dynamical systems described by differential equations seem to be not only compelling but almost inevitable for study.

One of the simplest non-trivial instance of such a model can be described in the equations

$$Y_i = u(t_i) + \varepsilon_i, i = 1, \dots, n \qquad \dot{u}(t) = f(u(t)), \tag{1}$$

where \dot{u} denotes the time-derivative, i.e., (du)/(dt). Here, $u \colon \mathbb{R} \to \mathbb{R}^d$ is the solution to an ordinary differential equation (ODE) described by an unknown function $f \colon \mathbb{R}^d \to \mathbb{R}^d$. It is measured at times $t_1 \leq \cdots \leq t_n$, which creates our observations Y_i by adding measurement noise ε_i to the true state of the system $u(t_i)$. There are different objectives one could have for such a model: reconstructing the solution u in the observed time interval $[t_1, t_n]$ (reanalysis, borrowing a term from climate modeling), predicting a future state u(t), $t > t_n$ (forecasting, as in weather forecast), or estimation of f (learning the dynamics of the dynamical system). Here, we are interested in the estimation of f, which is fundamental also for the other tasks as an estimate \hat{f} allows create an estimate \hat{u} by solving $\dot{\hat{u}}(t) = \hat{f}(\hat{u}(t))$. We want to estimate f in a nonparametric model, i.e., we assume that f belongs to an infinite dimensional class of smooth functions, but we do not make any assumptions on its precise functional form. The parametric version of this model, in which f belongs to a finite dimensional class of functions (e.g., polynomials of fixed degree) has been studied in [Bru08; QZ10; GK12; DK15]. There, it is shown that the typical parametric \sqrt{n} -rate of convergence can be achieved. An overview of the state of the art for statistics of dynamical systems (mostly concerned with parametric settings) can be found in [MMP15; RH17; Dat21].

In the nonparametric setting, different algorithms for estimation have been proposed, e.g., [Hei+18; Che+18; GR21]. In [Mar+23], theoretical results for learning dynamics nonparametrically are shown, but in a density estimation context that is rather different from the regression-type model studied here. To the best of the author's knowledge, only [Lah+24] take a theoretical view on the ODE regression problem (additionally to proposing a new algorithm based on reproducing kernel Hilbert spaces). They show an error bound of order $n^{-1/4}$ (root mean squared error) for the reanalysis problem [Lah+24, Theorem 1]. This result cannot be optimal as standard nonparametric regression for u (ignoring the ODE constraints) yields an error of smaller order, namely $n^{-\beta/(2\beta+1)}$, where β describes the smoothness of u (in the case of [Lah+24, Theorem 1], we have $\beta = 3$).

Estimation in the nonparametric setting seem much harder than in the parametric one: If f is a smooth function with bounds on its derivative but otherwise unrestricted, an observation Y_i only yields information on f(x) for x close to $u(t_i)$, i.e., the information is local. In contrast, each observation from an ODE where f is known to be a polynomial of degree at most N carries information for all coefficients of said polynomial, i.e., the information is global. What makes estimation in a nonparametric ODE model more difficult than in a nonparametric regression model (which also suffers from locality of information) is that fact that the location of the observation $u(t_i)$ is itself depends on the unknown f.

We introduce two models, called *Snake model* and *Stubble model*, which provide a remedy of this problem. For each of the two models, we propose estimators and study their rate of convergence. We find that in both instances, one obtains an error bound (root mean squared error) of order

$$n^{-\frac{\beta}{2(\beta+1)+d}}\tag{2}$$

for β -smooth functions f under optimal circumstances. This error bound is minimax optimal as we have corresponding lower bounds in the parallel work [SS24].

1.1 Contributions

We now give more details on the contributions of this work.

A general model. In general, we assume the true model function f^* to be an element of Höldertype smoothness class denoted as $\overline{\Sigma}^{d \to d}(\beta, L_{[0,\beta]})$, which means that the k-th derivative of ffor $k \in [0,\beta] := \{0,\ldots,\beta\}$ is bounded by $L_k \in \mathbb{R}_{>0}$, i.e., $\sup_{x \in \mathbb{R}^d} \|D^k f(x)\|_{op} \leq L_k$, where $\|\cdot\|_{op}$ is the operator norm. Then, for given initial conditions x_1,\ldots,x_m , we observe the solution $t \mapsto U(f^*, x_j, t)$ of the ODE $\dot{u}(t) = f^*(u(t))$ with $U(f^*, x_j, 0) = x_j$ at time points $i\Delta t$ with time step $\Delta t \in \mathbb{R}_{>0}$, i.e.,

$$Y_{j,i} = U(f^*, x_j, i\Delta t) + \varepsilon_{j,i}, \ j \in \llbracket 1, m \rrbracket, i \in \llbracket 1, n_j \rrbracket.$$
(3)

In total, we have $n = \sum_{j=1}^{m} n_j$ observations. The noise variables $\varepsilon_{j,i}$ are assumed to be independent, centered, and to have a finite second moment. Because of the dependency of the location of the observation $u(t_{j,i})$ on f^* , in general, consistent estimation in all of $[0, 1]^d$ is impossible. Thus, we introduce the Stubble model and the Snake model, which restrict the general model.

The Stubble model. We use the notation \approx and \preccurlyeq to mean asymptotically equal and asymptotically lower than (up to a positive constant), respectively (see Notation 2.3). In the Stubble model, we observe many $(m \approx n)$ short $(n_j \preccurlyeq 1)$ solutions. We assume that their initial conditions x_j cover the domain of interest $[0, 1]^d$ suitably. For this model, we construct an estimator \hat{f} based on a multivariate local polynomial estimator and an univariate polynomial interpolation, which is

similar to the Adams–Bashforth method for numerical solutions of ODEs [But16, Chapter 24]. We obtain

$$\mathbf{E}\left[\left\|\hat{f}(x) - f^{\star}(x)\right\|_{2}^{2}\right] \preccurlyeq \left(\Delta t^{2}n\right)^{-\frac{2\beta}{2\beta+d}} + \Delta t^{2\beta}$$

$$\tag{4}$$

for all $x \in [0, 1]^d$, see Corollary 3.13. The rate is shown to be minimax optimal by comparing it to lower bounds in [SS24].

The Snake model. In the Snake model, we observe few $(m \leq 1) \log(n_j \approx n)$ solutions. We require the trajectories $U(f^*, x_j, [0, n_j \Delta t])$ to cover the domain of interest $[0, 1]^d$ suitably. For this model, we construct an estimator \hat{f} based on an univariate local polynomial estimator and a multivariate polynomial interpolation. Then

$$\sup_{x \in [0,1]^d} \left\| \hat{f}(x) - f^{\star}(x) \right\|_2^2 \in \mathbf{O}_{\mathbf{P}} \left(\delta^{2\beta} + (\Delta t \log n)^{\frac{2\beta}{2(\beta+1)+1}} \right) \,, \tag{5}$$

see Corollary 4.17, where we want to view $\delta \in \mathbb{R}_{>0}$ for now as the largest distance between a point $x \in [0, 1]^d$ and its closest state $U(f^*, x_j, t), t \in [0, T]$. This interpretation is true for the case $\beta = 1$, but if $\beta > 1$, the definition is more complex and the result more restrictive. By comparing it to lower bounds in [SS24], the rate is shown to be minimax optimal if δ and T are in a certain relation. For the rate to be optimal, we essentially require observations to be rather dense in time, and temporally distant parts of the trajectories $U(f^*, x_j, [0, T])$ to be distant enough in state space.

Connection and further results. Note the complementary nature of the two models and their estimators. In spite of this, in the optimal setting regarding Δt and δ , we obtain an error bound of order (2) in both models (up to a log-factor when considering the sup-norm), see Corollary 3.15 and Corollary 4.19. In both models, we start with the case $\beta = 1$, which allows for simple estimators and a gentle introduction of the main ideas as well as slightly stronger and more specific results. See Theorem 3.2 and Corollary 3.4 for the Stubble model, and Theorem 4.4 and Corollary 4.6 for the Snake model. For the case of general smoothness $\beta \in \mathbb{N}$, we first show results for estimation strategies that can be used with an arbitrary regression estimator, Theorem 3.11 and Theorem 4.16, respectively. If the chosen regression estimator achieves the minimax rate for a standard nonparametric regression problem, the estimation strategies achive the rates (4) and (5), respectively. In Corollary 3.13 and Corollary 4.17, we apply the general results with the (minimax optimal) local polynomial estimator as regression estimator. Furthermore, we study how the smoothness of f influences the smoothness of $t \mapsto U(f, x, t)$ and the smoothness of $x \mapsto U(f, x, t)$ in Corollary D.12, respectively.

1.2 Relation to Other Models

Regression on the solutions. The ODE model can be viewed as a regression model, with the ODE as a constraint on the regression function $u^* = U(f^*, x, \cdot)$ in [0, T]. Solutions have a smoothness parameter $\tilde{\beta} = \beta + 1$ if the model function is β -smooth. Thus, we can solve the reanalysis problem, i.e., estimation of u^* in [0, T], with a the standard nonparametric rate of convergence for the squared error

$$\mathbf{E}\Big[\|\hat{u}(t) - u^{\star}(t)\|^2\Big] \preccurlyeq \left(\frac{T}{n}\right)^{\frac{2(\beta+1)}{2(\beta+1)+1}} \tag{6}$$

by ignoring the ODE-constraint and using a suitable estimator \hat{u} , see, e.g., [Tsy09]. Furthermore, the derivative of the solution can similarly be estimated with

$$\mathbf{E}\left[\left\|\hat{u}(t) - \dot{u}^{\star}(t)\right\|^{2}\right] \preccurlyeq \left(\frac{T}{n}\right)^{\frac{2\beta}{2(\beta+1)+1}}.$$
(7)

As $\dot{u}^{\star}(t) = f^{\star}(u^{\star}(t))$, this essentially yields an estimate of f^{\star} on the set $u^{\star}([0,T])$. This view is taken for our estimator in the Snake model of section 4.

Minimax rate of nonparametric regression. For standard nonparametric estimation of the s-th derivative of a $\tilde{\beta}$ -smooth regression function $g^* \colon \mathbb{R}^d \to \mathbb{R}$ (or $\mathbb{R}^d \to \mathbb{R}^d$), we have the minimax rate

$$\mathbf{E}\left[\left\|\widehat{D^{s}g}(x) - D^{s}g^{\star}(x)\right\|^{2}\right] \preccurlyeq n^{-\frac{2(\tilde{\beta}-s)}{2\tilde{\beta}+d}},$$
(8)

see, e.g., appendix B. In the ODE model, we estimate a first derivative (s = 1) in *d*-dimensions of the solutions that have smoothness $\tilde{\beta} = \beta + 1$. So, the rate of convergence we obtain in this work (2) fits the formula of the minimax regression rate (8), even though the models are quite different.

Errors in variables. Assume that time steps are constant, i.e., $t_{i+1} - t_i = \Delta t$. Consider the so-called propagator $x \mapsto U(f^*, x, \Delta t)$. If $\Delta t \to 0$, we can infer f from it, as

$$\frac{U(f^{\star}, x, \Delta t) - x}{\Delta t} = \frac{U(f^{\star}, x, \Delta t) - U(f^{\star}, x, 0)}{\Delta t} \xrightarrow{\Delta t \to 0} f^{\star}(x).$$
(9)

Estimating the propagator from the data (Y_{i-1}, Y_i) can be viewed as an errors-in-variables regression problem, e.g., [MRS12]. In such a model one has observation of the response variable $Z_i = g(x_i) + \varepsilon_i$ and assumes the predictors x_i to be unknown but instead to be measured noisily, e.g., via $X_i = x_i + \xi_i$ for a noise variable ξ_i . With $(X_i, Z_i) = (Y_{i-1}, Y_i)$, estimation of the propagator in the ODE model gives rise to an errors-in-variables regression problem. The Stubble model of section 3 removes the problems coming with errors-in-variables, by assuming known initial conditions and observing many solutions for only a short time.

Nonparametric time series analysis. The ODE model with the sequence (Y_i) can be view from the perspective of nonparametric time series analysis, where discrete-time stochastic processes in one variable (time) are studied. See, e.g., [FY03, Chapter 6]. On one hand, the focus on discrete time and often only one state dimension, allows for a deep understanding of more complex time evolutions than in an one-dimensional ODE model. On the other hand, it seems that the structure of higher dimensional, time continuous ODEs is typically not captured in such models.

Stochastic differential equations. The noise in stochastic differential equation (SDE) models, e.g., [Bis08; CG20] can be described as *system noise* in contrast to the *measurement noise* of the ODE model: It changes the state of the system and solutions are stochastic processes. In the ODE model, the state of the system is not influenced by the noise and solutions are deterministic. In comparison to a model with observed solutions of a SDE in one variable (time), estimation in the ODE model seems more difficult: The observation Z_t at time t of the solution (Z_t) in the SDE model is the true state of the system at that time. If the SDE is Markovian, then the distribution of the future states ($Z_{t'}$)_{t'>t} only depend on the now known state Z_t . In the ODE model, we do not know the true state $u^*(t)$, but only a noisy observation of it. Even though the future evolution is deterministic, it can be hard to predict if the initial state is not know perfectly (see the topic of *chaotic systems*, e.g., in [Str24]).

Hidden markov models. As the true state of the system $u^*(t)$ is hidden from us and only noisily observed, the ODE model is connected to hidden Markov models (HMMs), e.g., [BRR98]. In HMMs, on observes Z_i which depends on the unobserved random variable X_i , which in turn depends on X_{i-1} . But Z_i is independent of X_{i-1} and Z_{i-1} given X_i . If we set $X_i = u^*(t_i)$, $Z_i = Y_i$. We can view the ODE model as a HMM. In contrast to HMMs, the transition from $u^*(t_i)$ to $u^*(t_{i+1})$ is deterministic, and the time and state domains are continuous.

1.3 Overview

The remaining article is structured as follows. In section 2, we introduce basic concepts for the study of ordinary differential equations (section 2.1), give a full formal description of the general model (section 2.2), argue why this model is not useful without further restriction (section 2.3), and introduce some notation for standard regression problems that allows us to formulate estimation strategies referring to arbitrary regression estimators (section 2.4). Sections 3 and 4 formally introduce the Stubble and Snake model, respectively, and describe the estimators \hat{f} of f^* , upper bounds on the error $\|\hat{f}(x) - f^*(x)\|_2$ and their proofs. Both sections are separated into two parts: The first one (section 3.1 and 4.1 respectively) concerns the Lipschitz-case ($\beta = 1$). The

second one (section 3.2 and 4.2 respectively) concerns the general case ($\beta \in \mathbb{N}$). The appendix makes this article largely self-contained: We first state some basics of multivariate derivatives and smoothness classes (appendix A). Then, we recall classical results on the upper error bound for multivariate local polynomial regression in appendix B. In appendix C, uni- and multivariate polynomial interpolation are discussed. And finally, appendix D provides smoothness results for the solutions of ODEs.

2 Preliminaries

In this section, we recall some basic concepts related to ODEs, we introduce a general statistical model for observing solutions of ODEs, we explain why this model requires further restriction in order to be a useful model, and we introduce some terminology for a standard regression problem, which will later allows us to construct ODE estimators in a generic fashion.

2.1 Ordinary Differential Equations

We introduce some basic terminology and fundamental properties concerning ODEs. See also [Har02; Arn06].

Notation 2.1.

- (i) Let \mathbb{Z} be the set of integers and \mathbb{R} the set of reals. For $\mathbb{K} \in \{\mathbb{Z}, \mathbb{R}\}$ and $a \in \mathbb{R}$, denote $\mathbb{K}_{>a} = \{x \in \mathbb{K} \mid x > a\}$. Define $\mathbb{K}_{>a}$, $\mathbb{K}_{<a}$, $\mathbb{K}_{<a}$ accordingly.
- (ii) Set $\mathbb{N} := \mathbb{Z}_{\geq 1}, \mathbb{N}_0 := \mathbb{Z}_{\geq 0}$. Let $a, b \in \mathbb{Z}$ with $a \leq b$. Set $\llbracket a, b \rrbracket := \mathbb{Z}_{\geq a} \cap \mathbb{Z}_{\leq b}$. Set $\llbracket a \rrbracket := \llbracket 1, a \rrbracket$.

Let $d \in \mathbb{N}$. Let $f : \mathbb{R}^d \to \mathbb{R}^d$. For a differentiable function $u : \mathbb{R} \to \mathbb{R}^d$, denote its derivative as $\dot{u} : \mathbb{R} \to \mathbb{R}^d$. Then

$$\dot{u}(t) = f(u(t)) \qquad \text{for } t \in \mathbb{R}$$
(10)

is an autonomous, first-order, ordinary differential equation. It is of first-order, as (10) only depends on the first derivative of u. It is autonomous, as the right-hand side term f(u(t)) only depends on t via u. In contrast, a non-autonomous, first-order ODE has the form $\dot{u}(t) = g(t, u(t))$ for a function $g: \mathbb{R} \times \mathbb{R}^d \to \mathbb{R}^d$. Any differentiable $u: \mathbb{R} \to \mathbb{R}^d$ that fulfills (10) is a solution to the ODE. The domain of u is called *time*. The codomain of u is called *state space*. A single element of the state space is a *state*. The image of u is called *trajectory*. We call f the model function of the ODE. Let $x \in \mathbb{R}^d$. Consider the requirement

$$u(0) = x. (11)$$

We call x the *initial conditions*. We call (10) together with (11) *initial value problem* (IVP). If u fulfills (10) and (11), it is a solution to the IVP.

Assume that f is (globally) Lipschitz continuous. Then the IVP (10), (11) has a unique solution (Picard-Lindelöf theorem, also known as the Cauchy-Lipschitz theorem). Denote this solution as $U(f, x, \cdot) \colon \mathbb{R} \to \mathbb{R}^d, t \mapsto U(f, x, t)$. The function $U(f, \cdot, \cdot) \colon \mathbb{R}^d \times \mathbb{R} \to \mathbb{R}^d$ is called *flow* of the ODE (10). Note the *semigroup property* of the flow,

$$U(f, x, 0) = x$$
, (12)

$$U(f, x, s+t) = U(f, U(f, x, s), t)$$
(13)

for all $x \in \mathbb{R}^d$, $s, t \in \mathbb{R}$. For a given time step $\Delta t \in \mathbb{R}_{>0}$, the mapping $x \mapsto U(f, x, \Delta t)$ is called *propagator*. If u is a solution to the ODE (10), then $u(t + \Delta t) = U(f, u(t), \Delta t)$ for all $t \in \mathbb{R}$. Furthermore, we call

$$\Upsilon(f, \Delta t, x) := U(f, x, \Delta t) - x = U(f, x, \Delta t) - U(f, x, 0)$$
(14)

the increment. If u is a solution to the ODE (10), then $u(t + \Delta t) = u(t) + \Upsilon(f, \Delta t, u(t))$ for all $t \in \mathbb{R}$.

The setting of autonomous, first-order ODEs is not a strong restriction: Consider the *d*-dimensional, non-autonomous, ℓ -th-order ODE

$$v^{(\ell)}(t) = g(t, v(t), v^{(1)}(t), \dots, v^{(\ell-1)}(t)),$$
(15)

where $v^{(k)}$ for $k \in [0, \ell]$ denotes the k-th derivative of the ℓ -times differentiable function $v \colon \mathbb{R} \to \mathbb{R}^d$ and $g \colon \mathbb{R} \times (\mathbb{R}^d)^\ell \to \mathbb{R}^d$ is a function. We represent the derivatives and the time variable with new variables: $u_k = v^{(k-1)}$ for $k \in [\ell]$ and $u_{\ell+1}(t) = t$. For $u(t) = (u_1(t), \ldots, u_{\ell+1}(t))^{\mathsf{T}}$, we obtain the ODE

$$\begin{pmatrix} \dot{u}_{1}(t) \\ \vdots \\ \dot{u}_{\ell-1}(t) \\ \dot{u}_{\ell}(t) \\ \dot{u}_{\ell+1}(t) \end{pmatrix} = \begin{pmatrix} u_{2}(t) \\ \vdots \\ u_{\ell}(t) \\ g(u_{\ell+1}(t), u_{1}(t), \dots, u_{\ell}(t)) \\ 1 \end{pmatrix} .$$
(16)

It is of the form $\dot{u}(t) = f(u(t))$ for a suitably chosen $f \colon \mathbb{R}^{\tilde{d}} \to \mathbb{R}^{\tilde{d}}$, where $\tilde{d} := d\ell + 1$. Hence, (16) is a \tilde{d} -dimensional, autonomous, first-order ODE. If u is a solution to the ODE (16), then u_1 is a solution of the ODE (15).

2.2 Formal Description of the General Model

We introduce a general model for observations from solutions of an ODE. It is similar to the one in [Lah+24].

Notation 2.2.

- (i) Let $a, b \in \mathbb{Z}$ with $a \leq b$. For $i \in [[a, b]]$, let x_i be some object. Set $x_{[[a, b]]} := (x_i)_{i \in [[a, b]]}$.
- (ii) Let $d \in \mathbb{N}$. For $x \in \mathbb{R}^d$, let $||x||_2$ denote the Euclidean norm. Let $k \in \mathbb{N}$. For a matrix $A \in \mathbb{R}^{k \times d}$, denote the operator norm as $||A||_{\mathsf{op}} := \sup_{v \in \mathbb{R}^d, ||v||_2 = 1} ||Av||_2$. Let $d_x, d_y \in \mathbb{N}$. Let $f : \mathbb{R}^{d_x} \to \mathbb{R}^{d_y}$. Denote the sup norm of f as $|f|_{\infty} := \sup_{x \in \mathbb{R}^{d_x}} ||f(x)||_2$. For linear operators, like the derivative Df(x), we abuse the notation slightly and set $|Df|_{\infty} := \sup_{x \in \mathbb{R}^{d_x}} ||Df(x)||_{\mathsf{op}}$. For more details on derivatives on their norms, see appendix A.
- (iii) Let $d_{\mathsf{x}}, d_{\mathsf{y}}, \beta \in \mathbb{N}$. Denote the set of β -times continuously differentiable functions $f \colon \mathbb{R}^{d_{\mathsf{x}}} \to \mathbb{R}^{d_{\mathsf{y}}}$ as $\mathcal{D}^{\beta}(\mathbb{R}^{d_{\mathsf{x}}}, \mathbb{R}^{d_{\mathsf{y}}})$. Let $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0} \cup \{\infty\}$. Denote by $\bar{\Sigma}^{d_{\mathsf{x}} \to d_{\mathsf{y}}}(\beta, L_{\llbracket 0,\beta \rrbracket}) \subseteq \mathcal{D}^{\beta}(\mathbb{R}^{d_{\mathsf{x}}}, \mathbb{R}^{d_{\mathsf{y}}})$ the set of β -times continuously differentiable functions $f \colon \mathbb{R}^{d_{\mathsf{x}}} \to \mathbb{R}^{d_{\mathsf{y}}}$ with $|D^{k}f|_{\infty} \leq L_{k}$ for $k \in \llbracket 0, \beta \rrbracket$. More details on derivatives and the smoothness class $\bar{\Sigma}^{d_{\mathsf{x}} \to d_{\mathsf{y}}}(\beta, L_{\llbracket 0, \beta \rrbracket})$ are described in appendix A.

Let $d \in \mathbb{N}$ be the dimension of the state space \mathbb{R}^d . Let $\beta \in \mathbb{N}$ be the smoothness parameter. Let $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0}$ be the Lipschitz parameters. Let $\mathcal{F}_{d,\beta} := \bar{\Sigma}^{d \to d}(\beta, L_{\llbracket 0,\beta \rrbracket})$ be the smoothness class. Let the true model function be $f^* \in \mathcal{F}_{d,\beta}$. Let $m \in \mathbb{N}$ be the number of observed solutions and let $x_1, \ldots, x_m \in \mathbb{R}^d$ be their initial conditions. For $j \in \llbracket m \rrbracket$, let $n_j \in \mathbb{N}$ be the number of observed solutions and let $x_1, \ldots, x_m \in \mathbb{R}^d$ be their initial conditions. For $j \in \llbracket m \rrbracket$, let $n_j \in \mathbb{N}$ be the number of observations for the *j*-th solution. Denote the total number of observations as $n := \sum_{j=1}^m n_j$. For $j \in \llbracket m \rrbracket$, let $T_j \in \mathbb{R}_{>0}$ be the observation duration of the *j*-th solution. For $j \in \llbracket m \rrbracket, i \in \llbracket 0, n_j \rrbracket$, let $t_{j,i} \in \mathbb{R}_{\geq 0}$ be the observation times such that $0 = t_{j,0} \leq t_{j,1} \leq \cdots \leq t_{j,n_j} = T_j$. Let $\sigma \in \mathbb{R}_{\geq 0}$ be the variance bound of the noise. Let the noise variables $(\epsilon_{j,i})_{j \in \llbracket m \rrbracket, i \in \llbracket n_j \rrbracket$ be independent \mathbb{R}^d -valued random variables such that $\mathbb{E}[\epsilon_{j,i}] = 0$ and $\mathbb{E}[\|\epsilon_{j,i}\|_2^2] \leq \sigma^2$. For $j \in \llbracket m \rrbracket$, $i \in \llbracket n_j \rrbracket$, let the observations be

$$Y_{j,i} := U(f^*, x_j, t_{j,i}) + \epsilon_{j,i}.$$

$$(17)$$

We observe $Y_{j,i}$ and know x_j and $t_{j,i}$, but f^* is unknown and to be estimated. We assume d, β , $L_{[0,\beta]}$, and σ to be fixed and we are interested in upper bounds when $n \to \infty$ for the mean squared error for estimators \hat{f} of f^* on the domain of interest $[0,1]^d$.

2.3 Need for Restriction of the General Model

We argue that the model introduced in section 2.2 is not suitable for estimation of the model function in a fixed domain of interest and give first informal descriptions of two possible remedies of the problem: the Snake model and the Stubble model.

Notation 2.3.

(i) Let $(a_n)_{n \in \mathbb{N}}, (b_n)_{n \in \mathbb{N}} \subseteq \mathbb{R}_{>0}$. Write

$$a_n \preccurlyeq b_n \text{ for } \limsup a_n/b_n < \infty \,,$$
 (18)

$$a_n \prec b_n \quad \text{for } \limsup a_n/b_n = 0 \,,$$
 (19)

$$a_n \asymp b_n \text{ for } a_n \preccurlyeq b_n \text{ and } b_n \preccurlyeq a_n.$$
 (20)

Define \succ and \succeq accordingly.

For the general model, consistent estimation is not possible when considering the domain of interest $[0,1]^d$. We can easily construct settings, where we will never make any observation in say $[-1/2, 1/2]^d$, i.e., $u(t_{j,i}) \notin [-1/2, 1/2]^d$ for all $j \in [m], i \in [0, n_j]$, no matter how large n is. As we are considering a nonparametric class of model functions $f^* \in \mathcal{F}_{d,\beta}$, we have for the maximal risk

$$\sup_{f \in \mathcal{F}_{d,\beta}} \mathbf{E}_{f^{\star}} \left[\left\| f^{\star}(0) - \hat{f}(0) \right\|_{2}^{2} \right] \succeq 1.$$
(21)

This behavior is different from parametric ODE models (finite dimensional class $\mathcal{F}_{d,\beta}$): In the nonparametric setting, observations contain only local information (information about $f^*(x)$ for x close to $u(t_{j,i})$), whereas in a parametric setting, observations typically contain global information so that consistent estimation is possible under mild restrictions [Bru08; QZ10; GK12; DK15].

In this work, we introduce two restrictions to the general nonparametric model that make consistent estimation possible:

In the Stubble model (section 3), we require the initial conditions $(x_j)_{j \in [\![m]\!]}$ to suitably cover the domain of interest $[0, 1]^d$. By doing so, we ensure that we obtain information from every part of the domain of interest. In this model, we assume to observe many short trajectories, i.e., $m \geq 1$ and $\max_{j \in [\![m]\!]} n_j \leq 1$, which looks like *stubble*.

In the Snake model (section 4), we directly require the trajectories $U(f^*, x_j, [0, T_j])$ to cover the domain of interest $[0, 1]^d$. In this model, we assume to observe few long trajectories, i.e., $\min_{j \in [m]} n_j \succ 1$. In the extreme case, we have m = 1. Consistent estimation on all of $[0, 1]^d$ with one observed solution is only possible, if its trajectory covers $[0, 1]^d$ suitably. Thus, this trajectory must not intersect itself, as otherwise it would mean that the solution is periodic. This behavior of self-avoidance in a bounded domain resembles the video game Snake [Don10, chapter 22].

2.4 Standard Regression Model

In this article, we present estimation strategies based on black-box regression estimators. To formalize what we mean by *black-box regression estimators* and to be able to talk about their properties, we introduce some terminology in this section and give some examples at the end of the section.

Notation 2.4.

(i) Let $(X_n)_{n \in \mathbb{N}}$ be a sequence of real-valued random variables. Let $(a_n)_{n \in \mathbb{N}} \subseteq \mathbb{R}_{>0}$. We use the big *O* in probability notation: By $X_n \in \mathbf{O}_{\mathbf{P}}(a_n)$, we mean that for all $\epsilon \in \mathbb{R}_{>0}$,

there are $B \in \mathbb{R}_{>0}$ and $n_0 \in \mathbb{N}$ such that

$$\forall n \in \mathbb{N}_{\geq n_0} \colon \mathbf{P}\left(\left|\frac{X_n}{a_n}\right| > B\right) < \epsilon \,. \tag{22}$$

(ii) Let $k, d \in \mathbb{N}, v = (v_1, \dots, v_d) \in \mathbb{R}^d$. Denote the projection to the k-th dimension as Π_k , i.e., $\Pi_k v = v_k$.

Let $d_{\mathbf{x}}, d_{\mathbf{y}} \in \mathbb{N}$ be the dimensions of the predictor and the target, respectively. Let \mathcal{F} be a set of functions of the form $\mathbb{R}^{d_{\mathbf{x}}} \to \mathbb{R}^{d_{\mathbf{y}}}$, the smoothness class. Let $f^* \in \mathcal{F}$ be the true regression function. Let $n \in \mathbb{N}$ be the number of observations. Let $T \in \mathbb{R}_{>0}$ be the extent of the domain of interest $[0, T]^{d_{\mathbf{x}}}$. For $i \in [n]$, let $x_i \in [0, T]^{d_{\mathbf{x}}}$. Let $\sigma \in \mathbb{R}_{>0}$ the variance bound of the noise. For $i \in [n]$, let ε_i be independent $\mathbb{R}^{d_{\mathbf{y}}}$ -valued random variables with $\mathbf{E}[\epsilon_i] = 0$ and $\mathbf{E}[\|\epsilon_i\|_2^2] \leq \sigma^2$. We call ε_i the noise. For $i \in [n]$, set the observations as

$$Y_i = f^{\star}(x_i) + \varepsilon_i \,. \tag{23}$$

The standard regression problem is to estimate f^* with an estimator \hat{f} given the data $\mathcal{D} := (x_i, Y_i)_{i \in [\![n]\!]}$. We denote this statistical model as $\mathfrak{P}^{d_x \to d_y}(\mathcal{F}, T, (x_i)_{i \in [\![n]\!]}, \sigma)$.

A regression estimator for the regression function f^* in $\mathfrak{P}^{d_x \to d_y}(\mathcal{F}, T, (x_i)_{i \in [n]}, \sigma)$ is any measurable map

$$\mathcal{E}^{d_{\mathsf{x}} \to d_{\mathsf{y}}}(\cdot, \cdot) \colon (\mathbb{R}^{d_{\mathsf{x}}} \times \mathbb{R}^{d_{\mathsf{y}}})^n \times \mathbb{R}^{d_{\mathsf{x}}} \to \mathbb{R}^{d_{\mathsf{y}}} \,. \tag{24}$$

Given the data \mathcal{D} , it estimates the regression function f^* as $x \mapsto \hat{f}(x) := \mathcal{E}^{d_x \to d_y}(\mathcal{D}, x)$. We define the term regression estimator for the derivative Df^* of the regression function accordingly.

We say $\mathcal{E}^{d_x \to d_y}$ is the *componentwise* estimator $\mathcal{E}^{d_x \to 1}$, when we apply $\mathcal{E}^{d_x \to 1}$ in each target dimension:

$$\mathcal{E}^{d_{\mathsf{x}} \to d_{\mathsf{y}}}(\mathcal{D}, \cdot) \colon \mathbb{R}^{d_{\mathsf{x}}} \to \mathbb{R}^{d_{\mathsf{y}}}, \ x \mapsto \left(\mathcal{E}^{d_{\mathsf{x}} \to 1}((x_i, \Pi_1 Y_i)_{i \in \llbracket n \rrbracket}, x), \dots, \mathcal{E}^{d_{\mathsf{x}} \to 1}((x_i, \Pi_{d_{\mathsf{y}}} Y_i)_{i \in \llbracket n \rrbracket}, x) \right) .$$
(25)

The maximal risk for the model $\mathfrak{P}_n = \mathfrak{P}^{d_x \to d_y}(\mathcal{F}, T, (x_i)_{i \in [n]}, \sigma)$ at a point $x_0 \in [0, T]^{d_x}$ of an estimator $\hat{f} = \mathcal{E}^{d_x \to d_y}(\mathcal{D}, \cdot)$ is

$$r(\hat{f}, \mathfrak{P}_n, x_0) := \sup_{\mathbf{P} \in \mathfrak{P}_n} \mathbf{E}_{\mathbf{P}} \left[\left\| \hat{f}(x_0) - f_{\mathbf{P}}^{\star}(x_0) \right\|_2^2 \right],$$
(26)

where $\mathbf{P} \in \mathfrak{P}_n$ is the probability measure that gives rise to (23) with $f^* = f_{\mathbf{P}}^*$. We say that $r_0^{\mathsf{pr},\mathsf{sup}}(\hat{f},\mathfrak{P}_n)$ is a sup-norm error bound in probability of an estimator $\hat{f} = \mathcal{E}^{d_x \to d_y}(\mathcal{D},\cdot)$ if, for all $\mathbf{P} \in \mathfrak{P}_n$, for $n \to \infty$, we have

$$\sup_{x \in [0,T]^{d_{\mathbf{x}}}} \left\| \widehat{f}(x) - f_{\mathbf{P}}^{\star}(x) \right\|_{2} \in \mathbf{O}_{\mathbf{P}} \left(r_{0}^{\mathsf{pr,sup}}(\widehat{f}, \mathfrak{P}_{n}) \right) \,. \tag{27}$$

We define the sup-norm error bound of the first derivative accordingly and denote it by $r_1^{\mathsf{pr},\mathsf{sup}}(\widehat{Df},\mathfrak{P}_n)$.

An example of a regression estimator \hat{f} suitable for the Hölder-type smoothness class $\mathcal{F} = \Sigma^{d_x \to d_y}(\beta, L)$ with $\beta, L \in \mathbb{R}_{>0}$ (see Definition A.2), is the componentwise local polynomial estimator described in appendix B. *Suitable* here means that it achieves the lowest values possible for $r(\hat{f}, \mathfrak{P}_n, x_0)$ for large enough $n \in \mathbb{N}$, up to a constant.

Further examples of (nonparametric) regression estimators that may be used in the general estimation strategies constructed in this article include orthogonal series estimators [Tsy09, chapter 1.7], wavelets [Ant07], penalized splines [Xia19], and neural networks [Sch20].

3 The Stubble Model and Estimation of the Increment Map

In the Stubble model, we observe many solutions to an ODE with different initial conditions. For each solution, we have only a few observation. The initial conditions are known and located so that they cover the domain of interest.

In this section, we first present an explicit estimation procedure for a Liptschitz-continuous class of model functions with optimal rate of convergence of the mean squared error at a point. The estimation procedure is based on a local constant estimator for the increment map and a linear interpolation for the model function. In the second part of this section, we generalize these results: For a general Hölder-smoothness class, we present an estimation strategy based on an arbitrary multivariate regression estimator for different increment maps and a univariate polynomial interpolation for the model function. If the nonparametric estimator enjoys certain optimality criteria with respect to the standard nonparametric regression problem, it induces an optimal procedure for nonparametric ODE estimation.

3.1 Lipschitz

In this specific instance of the general ODE estimation model, we consider Lipschitz-continuous functions f, initial conditions x_j on a uniform grid, and an estimation procedure for f^* that is based on the local constant estimator for the increments $\Upsilon(f^*, \Delta t, \cdot)$.

3.1.1 Model

The following is a restriction of the general model of section 2.2.

Let $d \in \mathbb{N}$. Set $\beta = 1$. Let $L_0, L_1 \in \mathbb{R}_{>0}$. Set $\mathcal{F} := \overline{\Sigma}^{d \to d}(1, L_0, L_1)$. Let $f^* \in \mathcal{F}$ and $n_0 \in \mathbb{N}$. Set $n := n_0^d$. Let $x_i \in [0, 1]^d$ for $i \in [n]$ form a uniform grid in $[0, 1]^d$, i.e.,

$$\{x_1, \dots, x_n\} = \left\{ \left(\frac{k_1}{n_0}, \dots, \frac{k_d}{n_0}\right)^\top \middle| k_1, \dots, k_d \in \llbracket n_0 \rrbracket \right\}.$$
 (28)

Let $\sigma \in \mathbb{R}_{\geq 0}$. Let $\epsilon_j := \epsilon_{j,1}, j \in [n]$ be independent \mathbb{R}^d -valued random variables such that $\mathbf{E}[\epsilon_i] = 0$ and $\mathbf{E}[\|\epsilon_i\|_2^2] \leq \sigma^2$. Let $\Delta t \in \mathbb{R}_{>0}$. Set

$$Y_j := Y_{j,1} := U(f^*, x_j, \Delta t) + \epsilon_j.$$
⁽²⁹⁾

We observe Y_j and know x_j and Δt , but f^* is unknown and to be estimated. We assume d, L_0, L_1 , and σ to be fixed. We are interested in upper bounds for the mean squared error at a point $x_0 \in [0, 1]^d$ depending on the asymptotics of n and Δt .

3.1.2 Estimator

For $f \in \mathcal{F}$, $\Delta t \in \mathbb{R}_{\geq 0}$, $x \in \mathbb{R}^d$, recall the definition of the increment map $\Upsilon(f, \Delta t, x) = U(f, x, \Delta t) - x$ and set $\iota^*(x) := \Upsilon(f^*, \Delta t, x)$.

Let $\hat{\iota}$ be the componentwise (see section 2.4) local constant estimator (Nadaraya–Watson) of ι^* with kernel K using the data $(x_j, Y_j)_{j \in [\![n]\!]}$ and a bandwidth of optimal order. See appendix B with $\beta = 1, \ell = 0, s = 0$ for details on the local constant estimator.

For $x \in \mathbb{R}^d$, define the scaled increment estimator of f^* as

$$\hat{f}(x) := \frac{\hat{\iota}(x)}{\Delta t} \,. \tag{30}$$

3.1.3 Result

Assumption 3.1.

• STRICTKERNEL: The kernel $K: \mathbb{R}_{\geq 0} \to \mathbb{R}$ is Lipschitz-continuous, nonnegative, non-

trivial $(K \neq 0)$, and there is $C_{ker} \in \mathbb{R}_{>0}$ such that

$$K(x) \le C_{\text{ker}} \mathbb{1}_{[0,1]}(x)$$
 (31)

for all $x \in \mathbb{R}_{\geq 0}$.

Theorem 3.2. Use the model of section 3.1.1 and the estimator of section 3.1.2. Assume STRICTKERNEL. Assume $\Delta t \preccurlyeq 1$. Then

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq \left(\Delta t^2 n\right)^{-\frac{2}{2+d}} + \Delta t^2.$$
(32)

Remark 3.3. By [SS24, Corollary 3.11], the error rate in Theorem 3.2 is minimax optimal.

The following corollary minimizes the error bound over Δt , i.e., it shows the asymptotic behavior of Δt that allows the best estimates of f for the same amount of data.

Corollary 3.4. Use the model of section 3.2.1 and the estimator of section 3.1.2. Assume STRICTKERNEL. Assume $\Delta t \simeq n^{-\frac{1}{4+d}}$. Then

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq n^{-\frac{2}{4+d}}.$$
(33)

3.1.4 Proof

Lemma 3.5. Assume
$$f \in \mathcal{F}$$
. Then $\Upsilon(f, \Delta t, \cdot) \in \overline{\Sigma}^{d \to d}(1, \tilde{L}_0, \tilde{L}_1)$ with
 $\tilde{L}_0 = \Delta t L_0, \quad \tilde{L}_1 = \exp(L_1 \Delta t) - 1.$
(34)

Proof. By the definition of U and the fundamental theorem of calculus,

$$U(f, x, t) = x + \int_0^t f(U(f, x, s)) ds.$$
(35)

Plugging this into the definition of ι yields

$$\iota(x) = U(f, x, \Delta t) - U(f, x, 0) = \int_0^{\Delta t} f(U(f, x, s)) \mathrm{d}s \,. \tag{36}$$

Thus, as f is uniformly bounded by L_0 ,

$$|\iota(x)| \le \int_0^{\Delta t} |f(U(f,x,s))| \,\mathrm{d}s \le \int_0^{\Delta t} L_0 \mathrm{d}s = \Delta t L_0 \,. \tag{37}$$

This is the value of \tilde{L}_0 . Lemma D.9 yields the value of \tilde{L}_1 by

$$\left|D\iota\right|_{\infty} \le \exp(L_1 t) - 1. \tag{38}$$

Lemma 3.6. Assume $\Delta t \preccurlyeq 1$. Then, for all $x_0 \in [0,1]^d$,

$$\mathbf{E}\Big[\|\hat{\iota}(x_0) - \iota^{\star}(x_0)\|_2^2\Big] \preccurlyeq \left(\Delta t^d n^{-1}\right)^{\frac{2}{2+d}} .$$
(39)

Proof. According to Lemma 3.5, ι^* is Lipschitz-continuous with constant $\tilde{L}_1 = \exp(L_1\Delta t) - 1$. As we consider L_1 to be constant and $\Delta t \leq 1$, we obtain $\tilde{L}_1 \leq \Delta t$.

By Corollary B.4 and Proposition B.9, the local constant estimator $\hat{\iota}$ with optimal bandwidth then fulfills

$$\mathbf{E}\Big[\|\hat{\iota}(x_0) - \iota^{\star}(x_0)\|_2^2\Big] \preccurlyeq \Delta t^{\frac{2d}{2+d}} n^{-\frac{2}{2+d}}.$$
(40)

Lemma 3.7. Let $\Delta t \in \mathbb{R}_{>0}$. Assume $f \in \mathcal{F}$. Let $\iota = \Upsilon(f, \Delta t, \cdot)$. Then

$$\sup_{x \in [0,1]^d} \left\| \frac{\iota(x)}{\Delta t} - f(x) \right\|_2 \le \frac{1}{2} L_0 L_1 \Delta t \,. \tag{41}$$

Proof. Let $x \in [0,1]^d$. With (36), as f is L_1 -Lipschitz,

$$\left\|\frac{\iota(x)}{\Delta t} - f(x)\right\|_{2} = \left\|\frac{1}{\Delta t} \int_{0}^{\Delta t} f(U(f, x, s)) - f(x) \mathrm{d}s\right\|_{2}$$
(42)

$$\leq \frac{L_1}{\Delta t} \int_0^{\Delta t} \|U(f, x, s) - x\|_2 \,\mathrm{d}s \,. \tag{43}$$

Using $U(f, x, s) = x + \int_0^s f(U(f, x, r)) dr$, we obtain

$$\int_{0}^{\Delta t} \|U(f,x,s) - x\|_{2} \,\mathrm{d}s \le \int_{0}^{\Delta t} \int_{0}^{s} \|f(U(f,x,r))\|_{2} \,\mathrm{d}r \mathrm{d}s \tag{44}$$

$$\leq \int_{0}^{\infty} \int_{0}^{s} L_0 \mathrm{d}r \mathrm{d}s \tag{45}$$

$$=\frac{1}{2}L_0\Delta t^2.$$
(46)

Combining the above inequalities yields

$$\left\|\frac{\iota(x)}{\Delta t} - f(x)\right\|_2 \le \frac{1}{2}L_0L_1\Delta t.$$
(47)

As this inequality holds for all $x \in [0, 1]^d$, we have finished the proof.

Proof of Theorem 3.2. Combine the bounds in Lemma 3.6 and Lemma 3.7 to obtain

$$\mathbf{E}\left[\left\|\hat{f}(x_{0}) - f^{\star}(x_{0})\right\|_{2}^{2}\right] = \mathbf{E}\left[\left\|\frac{\hat{\iota}(x_{0}) - \iota^{\star}(x_{0})}{\Delta t} + \frac{\iota^{\star}(x_{0})}{\Delta t} - f^{\star}(x_{0})\right\|_{2}^{2}\right]$$
(48)

$$\approx \frac{1}{\Delta t^2} \mathbf{E} \Big[\|\hat{\iota}(x_0) - \iota^{\star}(x_0)\|_2^2 \Big] + \left\| \frac{\iota^{\star}(x_0)}{\Delta t} - f^{\star}(x_0) \right\|_2^2$$
(49)

$$\preccurlyeq \Delta t^{-\frac{4}{2+d}} n^{-\frac{2}{2+d}} + \Delta t^2 \,. \tag{50}$$

3.2 General

We generalize the Lipschitz setting of section 3.1 in the following way: We consider an arbitrary smoothness parameter $\beta \in \mathbb{N}$ of the model function f^* . The initial conditions are not restricted to a uniform grid, but must still be *uniform enough* in some sense. We present an estimation strategy which can be used with an arbitrary regression estimator. If this estimator achieves the optimal rate of convergence for a standard nonparametric regression problem, the resulting ODE estimator is also optimal in a minimax sense.

Notation 3.8.

- (i) Let $d \in \mathbb{N}$. Let $\alpha \in \mathbb{N}_0^d$.
 - Denote $|\alpha| := \sum_{i=1}^d \alpha_i$,
 - Denote $\alpha! := \prod_{i=1}^d \alpha_i!$,
 - Let $x \in \mathbb{R}^d$. Denote $x^{\alpha} = \prod_{i=1}^d x_i^{\alpha_i}$.
- (ii) For $\ell, d \in \mathbb{N}$, denote $\mathcal{P}_{d,\ell}$ be the set of all polynomials of degree at most ℓ defined over \mathbb{R}^d ,

$$\mathcal{P}_{d,\ell} := \left\{ p \colon \mathbb{R}^d \to \mathbb{R} \middle| p(x) = \sum_{\alpha \in \mathbb{N}_0^d, |\alpha| \le \ell} \beta_\alpha x^\alpha, \ \beta_\alpha \in \mathbb{R} \right\}.$$
(51)

(iii) Let A be a set. Denote the indicator function of A as $\mathbb{1}_{A}(\cdot)$.

3.2.1 Model

The following is a restriction of the general model of section 2.2.

Let $d \in \mathbb{N}$, $\beta \in \mathbb{N}$, and $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0}$. Set the smoothness class as $\mathcal{F}_{d,\beta} := \overline{\Sigma}^{d \to d}(\beta, L_{\llbracket 0,\beta \rrbracket})$. Let $f^* \in \mathcal{F}_{d,\beta}$. Let $m \in \mathbb{N}$ and $x_1, \ldots, x_m \in \mathbb{R}^d$. Let $\Delta t \in \mathbb{R}_{>0}$. Set $n_j = \beta$ for all $j \in \llbracket m \rrbracket$. Set $t_{j,i} = t_i = i\Delta t$ for $i \in \llbracket 0,\beta \rrbracket$. Let $\sigma \in \mathbb{R}_{\geq 0}$. Let $(\epsilon_{j,i})_{j \in \llbracket m \rrbracket, i \in \llbracket \beta \rrbracket}$ be independent \mathbb{R}^d -valued random variables such that $\mathbf{E}[\epsilon_{j,i}] = 0$ and $\mathbf{E}[\|\epsilon_{j,i}\|_2^2] \leq \sigma^2$. For $j \in \llbracket m \rrbracket$, $i \in \llbracket \beta \rrbracket$, let

$$Y_{j,i} = U(f^*, x_j, i\Delta t) + \epsilon_{j,i}.$$
(52)

We observe $Y_{j,i}$, and know x_j and Δt , but f^* is unknown and to be estimated. We assume d, $L_{[0,\beta]}$, and σ to be fixed and we are interested in asymptotic upper bounds for the mean squared error at a point $x_0 \in [0,1]^d$ depending on n and Δt .

Remark 3.9.

(i) The results below, are the same (up to a constant) if we allow $n_j \ge \beta$ as long as $\max_{j \in [m]} n_j \preccurlyeq 1$.

3.2.2 Estimation

For $i \in [\![\beta]\!]$, let $\iota_i^* = \Upsilon(f^*, t_i, \cdot)$. Let $L \in \mathbb{R}_{>0}$. Let $\mathcal{E}^{d \to d}$ be an arbitrary regression estimator for the standard regression model $\mathfrak{P} := \mathfrak{P}^{d \to d}(\Sigma^{d \to d}(\beta, L), 1, (x_j)_{j \in [\![m]\!]}, \sigma)$, see section 2.4. For $i \in [\![\beta]\!]$, estimate the increment maps ι_i^* by $\hat{\iota}_i := \mathcal{E}^{d \to d}((x_j, Y_{j,i} - x_j)_{j \in [\![m]\!]}, \cdot)$. For convenience, set $\hat{\iota}_0(x) := 0$ for all $x \in \mathbb{R}^d$. Denote the maximal risk of the regression estimator, see (26), as,

$$r(d,\beta,L,\sigma,(x_j)_{j\in\llbracket m\rrbracket}) := \sup_{x\in[0,1]^d} r(\mathcal{E}^{d\to d},\mathfrak{P},x) \,.$$
(53)

For $x_0 \in \mathbb{R}^d$, let $\hat{p}(x_0, \cdot) := (\hat{p}_1(x_0, \cdot), \dots, \hat{p}_d(x_0, \cdot))$, where $\hat{p}_k(x_0, \cdot) \in \mathcal{P}_{1,\beta}$ is the (univariate) polynomial of at most degree β that interpolates $(t_i, \Pi_k \hat{t}_i(x_0))_{i \in [0,\beta]}$. To estimate f^* , we locally approximate $U(f^*, x_0, t)$ by $x_0 + \hat{p}(x_0, t)$, i.e., we consider $\hat{p}(x_0, t) \approx f^*(x_0 + \hat{p}(x_0, t))$. As $f^*(x_0) =$ $f^*(x_0 + \hat{p}(x_0, 0))$, this approach yields the estimator

$$\hat{f}(x_0) := \hat{p}(x_0, 0).$$
 (54)

Remark 3.10.

(i) The value of L is to be specified later and depends on Δt , β , and $L_{[0,\beta]}$. Typically the estimator $\mathcal{E}^{d\to d}$ does not depend on L, but the risk $r(\mathcal{E}^{d\to d},\mathfrak{P},x)$ does.

- (ii) If we set $\beta = 1$ and use the componentwise local constant estimator for $\mathcal{E}^{d \to d}$, we obtain the procedure of section 3.1.
- (iii) The polynomial interpolation in the second step can be viewed as a Adams–Bashforth method: a linear multistep method for solving ordinary differential equations numerically, see [But16, Chapter 24].

3.2.3 Result

Theorem 3.11. Use the model of section 3.2.1 and the estimator of section 3.2.2. Assume $\Delta t \leq 1$. Then

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq \Delta t^{-2} r(d, \beta, C\Delta t, \sigma, (x_j)_{j \in [m]}) + \Delta t^{2\beta}.$$
(55)

for some constant $C \in \mathbb{R}_{>0}$ depending only on β and $L_{[0,\beta]}$.

The minimax rate for the squared error in the standard regression problem with smoothness class $\Sigma^{d\to d}(\beta, L)$ is $L^{\frac{2d}{2\beta+d}}n^{-\frac{2\beta}{2\beta+d}}$ up to some constants. It is achieved, for example, by the local polynomial estimator with optimal bandwidth under some conditions on x_i , which are fulfilled by a uniform grid. See Corollary B.4. Considering $m \simeq n$, we then have

$$r(d,\beta,C\Delta t,\sigma,(x_j)_{j\in[[m]]}) \preccurlyeq \Delta t^{\frac{2d}{2\beta+d}} n^{-\frac{2\beta}{2\beta+d}}.$$
(56)

The conditions for the optimal rate for the local polynomial estimator with kernel K of appendix B are as follows.

Assumption 3.12.

• EIGENVALUE: There is $C_{egv} \in \mathbb{R}_{>0}$ such that

$$\lambda_{\min}(B(x))^{-1} \le C_{\mathsf{egv}} \tag{57}$$

for all $x \in [0, T]^d$, where B(x) is defined in (175).

• COVER: There is $C_{\mathsf{cvr}} \in \mathbb{R}_{>0}$ such that

$$\frac{1}{n} \sum_{i=1}^{n} \mathbb{1}_{\mathrm{B}^{d}(x,r)}(x_{i}) \leq \max\left(\frac{1}{n}, C_{\mathsf{cvr}}\left(\frac{r}{T}\right)^{d}\right), \tag{58}$$

for all $x \in [0, T]^d$, $r \in \mathbb{R}_{>0}$.

• KERNEL: The support of the kernel $K \colon \mathbb{R}_{\geq 0} \to \mathbb{R}$ fulfills $\operatorname{supp}(K) \subseteq [0, 1]$. Furthermore, there is $C_{\mathsf{ker}} \in \mathbb{R}_{>0}$ such that

$$K(z) \le C_{\text{ker}} \tag{59}$$

for all $z \in \mathbb{R}_{\geq 0}$.

The following corollary shows the result of Theorem 3.11 using a minimax optimal estimator $\mathcal{E}^{d \to d}$.

Corollary 3.13. Use the model of section 3.2.1. Use the estimator of section 3.2.2 with the componentwise local polynomial estimator of degree $\ell := \beta - 1$ with kernel K and optimal bandwidth as the estimator $\mathcal{E}^{d \to d}$. Assume $\Delta t \preccurlyeq 1$. Assume COVER, EIGENVALUE, and KERNEL. Then, for all $x_0 \in [0, 1]^d$,

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq \left(\Delta t^2 n\right)^{-\frac{2\beta}{2\beta+d}} + \Delta t^{2\beta}.$$
(60)

Remark 3.14.

- (i) Of course, the local polynomials can be replaced by any estimator that achieves the same (optimal) rate of convergence.
- (ii) By [SS24, Corollary 3.11], the error rate in Corollary 3.13 is minimax optimal.

The following corollary minimizes the error bound over Δt , i.e., it shows the asymptotic behavior of Δt that allows the best estimates of f for the same amount of data.

Corollary 3.15. Use the setting and assumptions of Corollary 3.13. Assume

$$\Delta t \asymp n^{-\frac{1}{2(\beta+1)+d}} \,. \tag{61}$$

Then

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq n^{-\frac{2\beta}{2(\beta+1)+d}}.$$
(62)

3.2.4 Proof

For $x_0 \in \mathbb{R}^d$, let $p^*(x_0, \cdot) \in \mathcal{P}_{1,\beta}$ be the (univariate) polynomial of at most degree β that interpolates $(t_i, \iota_i^*(x_0))_{i \in [0,\beta]}$. By the definition of \hat{f} , we can write

$$\hat{f}(x_0) - f^{\star}(x_0) = \dot{\hat{p}}(x_0, 0) - \dot{p}^{\star}(x_0, 0) + \dot{p}^{\star}(x_0, 0) - f^{\star}(x_0) \,. \tag{63}$$

Using this, we split the error into two parts,

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \le 2\mathbf{E}\left[\left\|\dot{\hat{p}}(x_0, 0) - \dot{p}^{\star}(x_0, 0)\right\|_2^2\right] + 2\left\|\dot{p}^{\star}(x_0, 0) - f^{\star}(x_0)\right\|_2^2.$$
(64)

We start with the second part: The polynomial $t \mapsto p^{\star}(x_0, t)$ interpolates the translated solution $\bar{u}^{\star}(x_0, t) := U(f^{\star}, x_0, t) - x_0$. By Corollary D.7, we have $|D^{\beta+1}\bar{u}^{\star}(x_0, \cdot)|_{\infty} \leq C_L$, where C_L depends only on β and $L_{[0,\beta]}$. By the definition of $\bar{u}^{\star}(x_0, t)$, it fulfills $\dot{u}^{\star}(x_0, 0) = f(x_0)$. Thus, Lemma C.2 provides a constant C_{β} depending only on β such that

$$\|\dot{p}^{\star}(x_0,0) - f^{\star}(x_0)\|_2 = \|\dot{p}^{\star}(x_0,0) - \dot{\bar{u}}^{\star}(x_0,0)\|_2 \tag{65}$$

$$\leq C_{\beta}C_{L}\Delta t^{\beta}.$$
 (66)

In other words,

$$\|\dot{p}^{\star}(x_0,0) - f^{\star}(x_0)\|_2^2 \preccurlyeq \Delta t^{2\beta} .$$
(67)

Let us turn our attention to the first part of the right-hand side of (64): By Corollary D.12, $|D^{\beta}\iota_{i}^{\star}|_{\infty} \leq C\Delta t$ for some constant C depending only on $d, \beta, L_{[0,\beta]}$ and the maximal Δt as $f^{\star} \in \overline{\Sigma}^{d \to d}(\beta, L_{[0,\beta]})$ and $\Delta t \preccurlyeq 1$. As $\mathcal{E}^{d \to d}$ achieves the rate of convergence $r(d, \beta, L, \sigma, (x_{j})_{j \in [m]})$ on $\Sigma^{d \to d}(\beta, L)$, we obtain

$$\mathbf{E}\Big[\|\hat{\iota}_i(x_0) - \iota_i^{\star}(x_0)\|_2^2\Big] \preccurlyeq r(d, \beta, C\Delta t, \sigma, (x_j)_{j \in \llbracket m \rrbracket}).$$
(68)

Applying Lemma C.3 with the bound (68) yields

$$\mathbf{E}\left[\left\|\dot{\hat{p}}(x_0,0) - \dot{p}^{\star}(x_0,0)\right\|_2^2\right] \preccurlyeq \Delta t^{-2} r(d,\beta,C\Delta t,\sigma,(x_j)_{j\in[[m]]}).$$
(69)

We combine (67) and (69) and get

$$\mathbf{E}\left[\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2^2\right] \preccurlyeq \Delta t^{-2} r(d, \beta, C\Delta t, \sigma, (x_j)_{j \in [m]}) + \Delta t^{2\beta}.$$

$$\tag{70}$$

4 The Snake Model and Estimation of the Observed Solutions

In the Snake model, we observe one (or a few) solution(s) to an ODE. For each solution, we have many observation. The associated estimation problem depends strongly on how the trajectories of the solutions are located in the state space.

In this section, we first present an explicit estimation procedure for a Lipschitz-continuous class of model functions and an upper bound on the maximal risk in sup norm. The estimation procedure is based on a local linear estimator for the solutions and a nearest neighbor interpolation for the model function. It is rate-optimal in some settings. In the second part of this section, we generalize these results: For a general Hölder-smoothness class, we present an estimation strategy based on a generic nonparametric estimator for the solutions and a multivariate polynomial interpolation for the model function. If the nonparametric estimator enjoys certain optimality criteria with respect to the standard nonparametric regression problem, it induces a procedure for nonparametric ODE estimation that is optimal in some settings.

4.1 Lipschitz

In this specific instance of the general ODE estimation model, we consider Lipschitz-continuous model functions f^* and an estimation procedure for \hat{f} that is based on local linear estimators \hat{u} and \hat{u} for the observed solution $U(f^*, x_1, \cdot)$ and its derivative, and a nearest neighbor interpolation of $\hat{u}(t) \mapsto \hat{u}(t)$ for the model function. We obtain an upper bound on the expected sup norm error. For this, we either require the trajectories $U(f^*, x_1, [0, T_1])$ to cover the fixed domain of interest $[0, 1]^d$ in a suitable way or we adapt our domain of interest to these trajectories, i.e., the upper bound only holds close to the trajectory.

Notation 4.1.

(i) Let $d \in \mathbb{N}$, $r \ge 0$, and $z \in \mathbb{R}^d$. Denote the closed ball with radius r around z as

$$B^{d}(z,r) := \left\{ x \in \mathbb{R}^{d} \mid ||x - z||_{2} \le r \right\}.$$
(71)

Let $\mathcal{Z} \subseteq \mathbb{R}^d$. Denote the closed ball with radius r around \mathcal{Z} as

$$B^{d}(\mathcal{Z}, r) := \bigcup_{z \in \mathcal{Z}} B^{d}(z, r) \,. \tag{72}$$

4.1.1 Model

The following is a restricting of the general model of section 2.2.

Let $d \in \mathbb{N}_{\geq 2}$ and $L_0, L_1 \in \mathbb{R}_{>0}$. Set $\mathcal{F} := \overline{\Sigma}^{d \to d}(1, L_0, L_1)$. Let $f^* \in \mathcal{F}$. Set m = 1. Let $x_1 \in \mathbb{R}^d$, $n = n_1 \in \mathbb{N}$, and $\Delta t \in \mathbb{R}_{>0}$. Set $t_{1,i} := t_i := i\Delta t$ for $i = \llbracket 0, n \rrbracket$ and $T := T_1 := n\Delta t$. Let $\sigma \in \mathbb{R}_{\geq 0}$. Let $\epsilon_i := \epsilon_{1,i}, i \in \llbracket n \rrbracket$ be independent \mathbb{R}^d -valued random variables such that $\mathbf{E}[\epsilon_i] = 0$ and $\mathbf{E}[\lVert \epsilon_i \rVert_2^2] \leq \sigma^2$. Set

$$Y_i := Y_{1,i} := U(f^*, x_1, t_i) + \epsilon_i \quad \text{for } i \in [\![n]\!].$$
 (73)

We observe Y_i and know x_1 and Δt , but f^* is unknown and to be estimated. We assume d, L_0 , L_1 , and σ to be fixed. For an estimator \hat{f} , we are interested in asymptotic upper bounds for the mean squared sup-norm of $f^* - \hat{f}$ in the domain of interest $[0, 1]^d$ depending on the asymptotics of n and Δt .

4.1.2 Estimator

Let $\hat{u}: [0,T] \to \mathbb{R}$ be the componentwise local linear estimator (as described in Appendix B, (174) with $\ell = 1, s = 0, d = 1$) of $U(f^*, x_1, \cdot)$ using the data $(t_i, Y_i)_{i \in [n]}$. Similarly, denote by \hat{u} the local linear estimator of the first derivative of $U(f^*, x_1, \cdot)$ ($\ell = 1, s = 1, d = 1$ in terms of Appendix B).

We assume that the bandwidth h is chosen of optimal order for the error bound in sup-norm, see Corollary B.7.

Let \hat{f} be the nearest neighbor interpolation of (\hat{u}, \hat{u}) : For $x \in \mathbb{R}^d$ and a continuous function $u: [0, T] \to \mathbb{R}^d$, let

$$t_{NN}(u, T, x) \in \underset{t \in [0, T]}{\arg\min} \|u(t) - x\|_2$$
(74)

with an arbitrary choice if the minimizer is not unique. Then, for $x \in \mathbb{R}^d$, we define the estimator of f^* as

$$f(x) := \dot{u}(t_{NN}(\hat{u}, T, x))$$
 (75)

4.1.3 Result

For our results in sup-norm, we require the noise to be sub-Gaussian.

Definition 4.2 (Sub-Gaussian). A real-valued random variable Z, is called *sub-Gaussian* if there is $v \in \mathbb{R}_{>0}$ such that

$$\mathbf{P}(|Z| \ge t) \le 2\exp\left(-\frac{t^2}{2v}\right) \tag{76}$$

for all $t \in \mathbb{R}_{>0}$. Then v is called *variance parameter*.

Assumption 4.3.

• SUBGAUSSIAN: The noise ε_i is sub-Gaussian with variance parameter σ^2 in each component.

For $\mathcal{X} \subseteq \mathbb{R}^d$ and a function $u: [0,T] \to \mathbb{R}^d$, define $\delta_{\mathsf{NN}}(u,T,\mathcal{X})$ as smallest radius so that the ball around u([0,T]) includes \mathcal{X} :

$$\delta_{\mathsf{NN}}(u, T, \mathcal{X}) := \sup_{x \in \mathcal{X}} \inf_{t \in [0, T]} \|u(t) - x\|_2 .$$
(77)

The results come in two versions with respect to the domain of interest: We either fix the domain of interest, e.g., $\mathcal{X} = [0, 1]^d$ and make the error bound depend on $\delta_{NN}(u^*, T, \mathcal{X})$, or we fix δ (or a sequence $\delta = \delta_n$) and adapt our domain of interest to be $\mathcal{X} = B^d(u^*([0, T]), \delta)$.

Theorem 4.4. Use the model of section 4.1.1 and the estimator of section 4.1.2. Assume STRICTKERNEL, SUBGAUSSIAN. Assume

$$\left(\frac{n}{\log(n)}\right)^{-\frac{1}{4}} \preccurlyeq T \preccurlyeq n \log(n)^{\frac{1}{4}}.$$
(78)

(i) Let $\delta \in \mathbb{R}_{>0}$, potentially changing with n. Let $\mathcal{X} := B^d(u^*([0,T]), \delta)$. Then

$$\mathbf{E}\left[\sup_{x\in\mathcal{X}}\left\|\hat{f}(x) - f^{\star}(x)\right\|_{2}^{2}\right] \preccurlyeq \delta^{2} + \left(\frac{T\log n}{n}\right)^{\frac{2}{5}}.$$
(79)

(ii) Let $\delta = \delta_{NN}(u^{\star}, T, [0, 1]^d)$. Then

$$\mathbf{E}\left[\sup_{x\in[0,1]^d}\left\|\hat{f}(x) - f^{\star}(x)\right\|_2^2\right] \preccurlyeq \delta^2 + \left(\frac{T\log n}{n}\right)^{\frac{2}{5}}.$$
(80)

Remark 4.5.

(i) Comparing the error bound in Theorem 4.4 (ii) with [SS24, Corollary 4.11], we see that

it is optimal if

$$\delta^2 \succcurlyeq \left(\frac{T\log n}{n}\right)^{\frac{1}{5}}.$$
(81)

- (ii) Note that [SS24, Corollary 4.11] uses $m \succ 1$ many solutions. But in [SS24, Appendix E], it is argued that m = 1 suffices.
- (iii) One can think of (81) as the requirement of large distances between temporally different parts of trajectory. If this is not the case, then the estimator seems suboptimal: For the estimation of \hat{u} and \hat{u} , it ignores information from observations that are distant in time but close in state space. An estimator that is optimal for regimes where different trajectory parts are not well-separated, must not ignore this information.

Corollary 4.6. Use the model of section 4.1.1 and the estimator of section 4.1.2. Assume STRICTKERNEL, SUBGAUSSIAN.

(i) Let $\delta \in \mathbb{R}_{\geq 0}$, potentially changing with n. Let $\mathcal{X} := B_{\delta}(u^{\star}([0,T]))$. Assume $T \preccurlyeq \left(\frac{n}{\log n}\right)^{\frac{d-1}{4+d}}$ and $\delta \preccurlyeq \left(\frac{n}{\log n}\right)^{-\frac{1}{4+d}}$. Then $\mathbf{E}\left[\sup_{x \in \mathcal{X}} \left\|\hat{f}(x) - f^{\star}(x)\right\|_{2}^{2}\right] \preccurlyeq \left(\frac{n}{\log n}\right)^{-\frac{2}{4+d}}.$ (82)
(ii) Let $\delta = \delta_{\mathsf{NN}}(u^{\star}, T, [0, 1]^{d})$. Assume $T \preccurlyeq \left(\frac{n}{\log n}\right)^{\frac{d-1}{4+d}}$ and $\delta \preccurlyeq \left(\frac{n}{\log n}\right)^{-\frac{1}{4+d}}$. Then

$$\mathbf{E}\left[\sup_{x\in[0,1]^d}\left\|\hat{f}(x) - f_{\star}(x)\right\|_2^2\right] \preccurlyeq \left(\frac{n}{\log n}\right)^{-\frac{2}{4+d}}.$$
(83)

Remark 4.7.

(i) The conditions on T and δ ensure an optimal trade-off between the ability to reconstruct u^* and coverage of $[0, 1]^d$. They can be fulfilled for certain f, but it is rather restrictive. Furthermore, to cover the hypercube $[0, 1]^d$ with $B_{\delta}(u([0, T]))$, where u has a speed bounded by L, we require $T \succeq \delta^{-(d-1)}$. Hence, the conditions on δ and T in Theorem 4.4 are tight in the sense that one variable determines the other up to a constant.

4.1.4 Proof

Lemma 4.8. Let
$$\delta > 0$$
. Let $x_0 \in B^d(u^*([0,T]), \delta)$. Then
 $\left\| \hat{f}(x_0) - f^*(x_0) \right\|_2 \le L_1\left(\delta + 2 \sup_{t \in [0,T]} \| \hat{u}(t) - u^*(t) \|_2 \right) + \sup_{t \in [0,T]} \left\| \hat{u}(t) - \dot{u}^*(t) \right\|_2.$ (84)

Proof. Denote $u^* := U(f^*, x_1, \cdot), \ \tau^* := t_{\mathsf{NN}}(u^*, T, x_0)$, and $\hat{\tau} := t_{\mathsf{NN}}(\hat{u}, T, x_0)$. Using the triangle inequality and the definition of \hat{f} and \dot{u}^* , we obtain

$$\left\| \hat{f}(x_0) - f^{\star}(x_0) \right\|_2 \le \left\| \hat{f}(x_0) - f^{\star}(u^{\star}(\hat{\tau})) \right\|_2 + \left\| f^{\star}(u^{\star}(\hat{\tau})) - f^{\star}(x_0) \right\|_2 \tag{85}$$

$$= \left\| \hat{u}(\hat{\tau}) - \dot{u}^{\star}(\hat{\tau}) \right\|_{2} + \left\| f^{\star}(u^{\star}(\hat{\tau})) - f^{\star}(x_{0}) \right\|_{2} \,. \tag{86}$$

Let us consider the second term of the last line. As f^* is L_1 -Lipschitz, we have

$$\|f^{\star}(u^{\star}(\hat{\tau})) - f^{\star}(x_0)\|_2 \le L_1 \|u^{\star}(\hat{\tau}) - x_0\|_2 .$$
(87)

We use the triangle inequality in $||x_0 - u^{\star}(\hat{\tau})||_2 \leq ||x_0 - \hat{u}(\hat{\tau})||_2 + ||\hat{u}(\hat{\tau}) - u^{\star}(\hat{\tau})||_2$ and bound

$$\|x_0 - \hat{u}(\hat{\tau})\|_2 \le \|x_0 - \hat{u}(\tau^*)\|_2 \tag{88}$$

$$\leq \|x_0 - u^{\star}(\tau^{\star})\|_2 + \|u^{\star}(\tau^{\star}) - \hat{u}(\tau^{\star})\|_2 , \qquad (89)$$

where we used the minimizing property of $\hat{\tau}$ and again the triangle inequality. Putting all previous bounds together, we obtain

$$\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2$$
 (90)

$$\leq L_1 \left(\|x_0 - u^*(\tau^*)\|_2 + \|u^*(\tau^*) - \hat{u}(\tau^*)\|_2 + \|\hat{u}(\hat{\tau}) - u^*(\hat{\tau})\|_2 \right) + \left\| \hat{u}(\hat{\tau}) - \dot{u}^*(\hat{\tau}) \right\|_2.$$
(91)

The claim follows by taking supremum over $\tau^*, \hat{\tau} \in [0, T]$ and noting the definition of x_0 and τ^* .

Proof of Theorem 4.4. As $f^* \in \overline{\Sigma}(1, L_0, L_1)$, we have $u^* \in \overline{\Sigma}(2, \infty, L_0, \tilde{L}_1)$, where \tilde{L}_1 depends on L_0 and L_1 , by Corollary D.7. We want to apply Corollary B.7 to obtain bounds on the sup-norm of the error for \hat{u} and \hat{u} . STRICTKERNEL together with the uniform grid $t_i = i\Delta t$ implies KERNEL, EIGENVALUE, and COVER by Proposition B.9. SUBGAUSSIAN is assumed and (78) implies (184). Thus, Corollary B.7 yields

$$\mathbf{E}\left[\sup_{t\in[0,T]}\|\hat{u}(t)-u^{\star}(t)\|_{2}^{2}\right] \preccurlyeq \left(\frac{T\log(n)}{n}\right)^{\frac{4}{5}},\tag{92}$$

$$\mathbf{E}\left[\sup_{t\in[0,T]}\left\|\hat{u}(t)-\dot{u}^{\star}(t)\right\|_{2}^{2}\right] \preccurlyeq \left(\frac{T\log(n)}{n}\right)^{\frac{2}{5}}.$$
(93)

Together with Lemma 4.8, we obtain

$$\mathbf{E}\left[\sup_{x\in[0,1]^d} \left\| \hat{f}(x) - f^{\star}(x) \right\|_2^2\right] \preccurlyeq \delta^2 + \mathbf{E}\left[\sup_{t\in[0,T]} \left\| \hat{u}(t) - u^{\star}(t) \right\|_2^2\right] + \mathbf{E}\left[\sup_{t\in[0,T]} \left\| \hat{u}(t) - \dot{u}^{\star}(t) \right\|_2^2\right]$$
(94)

$$\preccurlyeq \delta^2 + \left(\frac{T\log(n)}{n}\right)^{\frac{4}{5}} + \left(\frac{T\log(n)}{n}\right)^{\frac{2}{5}} \tag{95}$$

$$\preccurlyeq \delta^2 + \left(\frac{T\log(n)}{n}\right)^{\frac{2}{5}} . \tag{96}$$

4.2 General

We now want to consider estimation under higher order smoothness. We generalize the estimation procedure of section 4.1.2 as follows: We replace the local linear estimation of the solution u^* and its derivative \dot{u}^* by arbitrary regression estimators $\mathcal{E}^{1\to d}$ and $\tilde{\mathcal{E}}^{1\to d}$, respectively. E.g., this could be local polynomial estimators of the appropriate degree. Furthermore, the nearest neighbor interpolation is replaced by multivariate polynomial interpolation of the appropriate degree.

Notation 4.9.

(i) Let $A \subseteq \mathbb{R}^d$. Denote the diameter of A as

$$diam(A) := \sup_{a,a' \in A} ||a - a'||_2 .$$
(97)

(ii) Let $A \subseteq \mathbb{R}^d$. Denote the interior of the convex hull of A as

$$\mathring{\mathsf{ch}}(A) := \left\{ \sum_{k=1}^{K} w_i a_i \middle| K \in \mathbb{N}, \ a_1, \dots, a_K \in A, \ w_1, \dots, w_K \in (0, 1), \sum_{k=1}^{K} w_i = 1 \right\}.$$
(98)

4.2.1 Model

The following is a restricting of the general model of section 2.2.

Let $d \in \mathbb{N}_{\geq 2}$, $\beta \in \mathbb{N}_{\geq 2}$, and $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0}$. Set $\mathcal{F}_{d,\beta} := \bar{\Sigma}^{d \to d}(\beta, L_{\llbracket 0,\beta \rrbracket})$. Let $f^* \in \mathcal{F}_{d,\beta}$. Set m = 1. Let $x_1 \in \mathbb{R}^d$, $n = n_1 \in \mathbb{N}$, and $T := T_1 \in \mathbb{R}_{>0}$. Set $t_{1,i} := t_i \in [0,T]$ for $i \in \llbracket 0,n \rrbracket$ with $0 = t_0 \leq \cdots \leq t_n = T$. Let $\sigma \in \mathbb{R}_{\geq 0}$. Let $\epsilon_i := \epsilon_{1,i}$, $i \in \llbracket n \rrbracket$ be independent \mathbb{R}^d -valued random variables such that $\mathbf{E}[\epsilon_i] = 0$ and $\mathbf{E}[\|\epsilon_i\|_2^2] \leq \sigma^2$. Set

$$Y_i := Y_{1,i} := U(f^*, x_1, t_i) + \epsilon_i \quad \text{for } i \in [\![n]\!].$$
 (99)

We observe Y_i and know x_1 and t_i , but f^* is unknown and to be estimated. We assume d, $L_{[0,\beta]}$, and σ to be fixed. For an estimator \hat{f} , we are interested in asymptotic upper bounds for the mean squared sup-norm of $f^* - \hat{f}$ in the domain of interest $[0,1]^d$ depending on the asymptotics of n and T.

Remark 4.10.

- (i) It does not matter whether x_1 is known or not: Below, we derive upper bounds on the estimation error for the potentially more difficult problem of unknown x_1 . The lower bounds from [SS24, Corollary 4.11] are formulated with known x_1 , which are then also lower bounds for the case of of unknown x_1 .
- (ii) This model and following estimation procedure can easily be generalized to larger m as long as min_{j∈[m]} n_j ≻ 1 and the t_{j,i} behave in an appropriate way.

4.2.2 Multivariate Polynomial Interpolation

The estimation strategy for the model described in section 4.2.1 uses multivariate polynomial interpolation. In this section, we introduce some basics of polynomial interpolation and some further objects that are required to describe the ODE estimator. Recall Notation 3.8.

Definition 4.11 (Polynomial interpolation). Let $\ell, d_x, d_y \in \mathbb{N}$.

(i) Set

$$N := N_{d_{\mathsf{x}},\ell} := \dim(\mathcal{P}_{d_{\mathsf{x}},\ell}) = \begin{pmatrix} \ell + d_{\mathsf{x}} \\ d_{\mathsf{x}} \end{pmatrix}.$$
 (100)

(ii) For $x \in \mathbb{R}^{d_x}$, denote the vector of monomials of degree at most ℓ in d_x dimensions as

$$\psi(x) := \psi_{\ell}(x) := (x^{\alpha})_{|\alpha| \le \ell} \in \mathbb{R}^N.$$

$$(101)$$

For $M \in \mathbb{N}$, $\mathbf{x} = (x_1, \dots, x_M) \in (\mathbb{R}^{d_x})^M$, denote $\Psi(\mathbf{x}) := \Psi_\ell(\mathbf{x}) := (\psi(x_1), \dots, \psi(x_M))^\top \in \mathbb{R}^{M \times N}$.

(iii) For $x \in \mathbb{R}^{d_x}$, $\mathbf{x} \in (\mathbb{R}^{d_x})^N$ with $\Psi(\mathbf{x}) \in \mathbb{R}^{N \times N}$, $\mathbf{y} \in (\mathbb{R}^{d_y})^N = \mathbb{R}^{N \times d_y}$, denote by $I := I_\ell$ the componentwise polynomial interpolation,

$$I_{\ell}(\mathbf{x}, \mathbf{y}, \cdot) \colon \mathbb{R}^{d_{\mathbf{x}}} \to \mathbb{R}^{d_{\mathbf{y}}}, x \mapsto \begin{cases} \psi(x)^{\top} \Psi(\mathbf{x})^{-1} \mathbf{y} & \text{if } \Psi(\mathbf{x}) \text{ is invertible,} \\ 0 & \text{otherwise.} \end{cases}$$
(102)

Remark 4.12.

(i) Let $\mathbf{x} = (x_1, \ldots, x_N) \in (\mathbb{R}^{d_x})^N$ and $\mathbf{y} = (y_1, \ldots, y_N) \in (\mathbb{R}^{d_y})^N$. The function $I(\mathbf{x}, \mathbf{y}, \cdot)$ is an *interpolation* as $I(\mathbf{x}, \mathbf{y}, x_k) = y_k$. It is *polynomial* as $x \mapsto \prod_k I(\mathbf{x}, \mathbf{y}, x) \in \mathcal{P}_{d_x, \ell}$ for each $k \in [\![d_y]\!]$. It is *componentwise* as

$$I(\mathbf{x}, \mathbf{y}, x) = \left(I(\mathbf{x}, (\Pi_k y_1, \dots, \Pi_k y_N), x)\right)_{k \in \llbracket d_n \rrbracket} .$$
(103)

- (ii) The definition of $I(\mathbf{x}, \mathbf{y}, \cdot) = 0$ if $\Psi(\mathbf{x})$ is not invertible is for convenience only. The value 0 is not of importance.
- (iii) If $d_{\mathbf{x}} = 1$, the matrix $\Psi(\mathbf{x}) \in \mathbb{R}^{N \times N}$ is also called *Vandermonde matrix*. It is invertible if and only if all $x_k \in \mathbb{R}$ are distinct. In the multivariate case, conditions for invertibility of $\Psi(\mathbf{x})$ (and hence, unique existence of the polynomial interpolation) are more complex, see [Olv06, Theorem 4.1].

Definition 4.13 (Normalization). Let $d, N \in \mathbb{N}$. For $\mathbf{x} \in (\mathbb{R}^d)^N$ with diam $(\mathbf{x}) > 0$, let

$$\eta_{\mathbf{x}} \colon \mathbb{R}^d \to \mathbb{R}^d, x \mapsto \frac{x - \frac{1}{N} \sum_{k=1}^N x_k}{\mathsf{diam}(\mathbf{x})} \,. \tag{104}$$

Furthermore, denote $\eta_{\mathbf{x}}(\tilde{\mathbf{x}}) = (\eta_{\mathbf{x}}(\tilde{x}_1), \dots, \eta_{\mathbf{x}}(\tilde{x}_M))$ for $\tilde{\mathbf{x}} = (\tilde{x}_1, \dots, \tilde{x}_M) \in (\mathbb{R}^d)^M$, $M \in \mathbb{N}$.

Definition 4.14. Let $d, \ell \in \mathbb{N}$, $N = \dim(\mathcal{P}_{d,\ell})$. Let $\mathcal{X} \subseteq \mathbb{R}^d$ and $s, \delta \in \mathbb{R}_{\geq 0}$.

- (i) Define $\mathcal{G}_{\ell,\delta,s}(\mathcal{X})$ as the set of all points $x \in \mathbb{R}^d$ with following property: There is a set of points $\mathbf{x} \in \mathcal{X}^N$ with $x \in \mathring{ch}(\mathbf{x})$ such that $\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\|_{op} < s$ and $\operatorname{diam}(\mathbf{x}) \leq \delta$.
- (ii) Define $\mathcal{H}_{\ell,s}(\mathcal{X}, x)$ as the set of all sets $\mathbf{x} \in \mathcal{X}^N$ such that $x \in \mathring{ch}(\mathbf{x})$ and $\left\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\right\|_{\mathsf{op}} \leq s$.
- (iii) Let $\chi_{\ell,s}(\mathcal{X}, x)$ be an element $\mathbf{x} \in \mathcal{H}_{\ell,s}(\mathcal{X}, x)$ such that

$$\left\|\Psi(\eta_{\mathbf{x}_0}(\mathbf{x}_0))^{-1}\right\|_{\mathsf{op}}\operatorname{diam}(\mathbf{x}_0)^{\ell+1} \le (1+\epsilon)\min_{\mathbf{x}\in\mathcal{H}_{\ell,s}(\mathcal{X},x)} \left\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\right\|_{\mathsf{op}}\operatorname{diam}(\mathbf{x})^{\ell+1} \quad (105)$$

for $\mathbf{x}_0 = \chi_{\ell,s}(\mathcal{X}, x)$ and a fixed arbitrarily small $\epsilon \in \mathbb{R}_{>0}$.

4.2.3 Estimator

Let $L \in \mathbb{R}_{>0}$. Let $\mathcal{E}^{1 \to d}$ and $\tilde{\mathcal{E}}^{1 \to d}$ be arbitrary regression estimator for the regression function and its derivative in the standard regression model $\mathfrak{P} := \mathfrak{P}^{1 \to d}(\Sigma^{1 \to d}(\beta + 1, L), T, (t_i)_{i \in [\![n]\!]}, \sigma)$, see section 2.4. With these estimators, we estimate the observed ODE-solution and its derivative,

$$\hat{u}(t) := \mathcal{E}^{1 \to d} \left((t_i, Y_i)_{i \in \llbracket n \rrbracket}, t \right) \quad \text{and} \quad \hat{\hat{u}}(t) := \widetilde{\mathcal{E}}^{1 \to d} \left((t_i, Y_i)_{i \in \llbracket n \rrbracket}, t \right) .$$
(106)

Denote the maximal sup-norm risk in probability of the regression estimators, see (27), as,

$$\Gamma(n,\beta+1,L,T) := r_0^{\mathsf{pr},\mathsf{sup}}(\mathcal{E}^{1\to d},\mathfrak{P}), \qquad (107)$$

$$\Lambda(n,\beta+1,L,T) := r_1^{\mathsf{pr},\mathsf{sup}}(\widetilde{\mathcal{E}}^{1\to d},\mathfrak{P}).$$
(108)

Set $\ell := \beta - 1$. Fix $s \in \mathbb{R}_{>0}$. Let $x \in \mathbb{R}^d$. Assume $\chi_{\ell,s}(\hat{u}([0,T]), x)$ exists. Let $\hat{\mathbf{x}}(x) := (\hat{u}(\hat{\tau}_1), \ldots, \hat{u}(\hat{\tau}_N))$, where $\hat{\tau}_i$ is chosen so that $\hat{\mathbf{x}}(x) = \chi_{\ell,s}(\hat{u}([0,T]), x)$. Let $\hat{\mathbf{y}}(x) := (\hat{u}(\hat{\tau}_1), \ldots, \hat{u}(\hat{\tau}_N))$. Define the estimator

$$\hat{f}(x) := I_{\ell}(\hat{\mathbf{x}}(x), \hat{\mathbf{y}}(x), x) \,. \tag{109}$$

Remark 4.15.

- (i) The estimator is only defined for $x \in \mathcal{G}_{\ell,\infty,s}(\hat{u}([0,T]))$, which is a subset of the convex hull of $\hat{u}([0,T])$. If this is not the case, a fallback such as nearest neighbor (see section 4.1) could be applied in practice.
- (ii) The variable s is a regularity parameter for the matrix inversion executed in the multivariate polynomial interpolation. It will be assumed to fulfill $s \approx 1$ asymptotically.

4.2.4 Results

For $\mathcal{X} \subseteq \mathbb{R}^d$ and a function $u \colon [0,T] \to \mathbb{R}^d$, define

$$\delta_{\mathsf{NN}}^{\ell,s}(u,T,\mathcal{X}) := \inf\{\delta' \ge 0 \colon \mathcal{X} \subseteq \mathcal{G}_{\ell,\delta',s}(u([0,T]))\}.$$
(110)

Theorem 4.16. Use the model of section 4.2.1 and the estimator of section 4.2.3. Let $\delta \in \mathbb{R}_{\geq 0}$, potentially changing with n. Assume $T \prec n$, $\Gamma \prec 1$, $\Gamma \preccurlyeq \delta$, $\Gamma \preccurlyeq \Lambda \preccurlyeq 1$, $s \asymp 1$. Set $L := \beta! \sup_{k \in [\![\beta]\!]} L_k^{\beta+1}$.

(i) Set $\mathcal{X} := \mathcal{G}_{\ell,\delta,s}(u^{\star}([0,T]))$. Then

$$\sup_{x \in \mathcal{X}} \left\| \hat{f}(x) - f(x) \right\|_2^2 \in \mathbf{O}_{\mathbf{P}} \left(\delta^{2\beta} + \Lambda(n, \beta + 1, L, T)^2 \right) \,. \tag{111}$$

(ii) Set $\delta := \delta_{\mathsf{NN}}^{\ell,s}(u^{\star},T,[0,1]^d)$. Then

$$\sup_{x \in [0,1]^d} \left\| \hat{f}(x) - f(x) \right\|_2^2 \in \mathbf{O}_{\mathbf{P}} \left(\delta^{2\beta} + \Lambda(n, \beta + 1, L, T)^2 \right) \,. \tag{112}$$

We want to use the componentwise local polynomial estimators of appendix B as the estimators $\mathcal{E}^{1\to d}$ and $\tilde{\mathcal{E}}^{1\to d}$ (appendix B: d = 1, $\ell = \beta$, and s = 0 and s = 1, respectively). We assume KERNEL, EIGENVALUE, and COVER for the observation times t_i and SUBGAUSSIAN for the noise ε_i . Furthermore, we require (184). Then Corollary B.7 yields

$$\Gamma(n,\beta+1,L,T) \preccurlyeq \left(\frac{T\log n}{n}\right)^{\frac{\beta+1}{2(\beta+1)+1}} \tag{113}$$

$$\Lambda(n,\beta+1,L,T) \preccurlyeq \left(\frac{T\log n}{n}\right)^{\frac{\beta}{2(\beta+1)+1}},\tag{114}$$

assuming $L \simeq 1$.

Corollary 4.17. Use the model of section 4.2.1 and the estimator of section 4.2.3 with the componentwise local polynomial estimator of degree β (see appendix B) as $\mathcal{E}^{1 \to d}$ and $\widetilde{\mathcal{E}}^{1 \to d}$. Let $\delta \in \mathbb{R}_{>0}$, potentially changing with n and assume

$$\left(\frac{T\log n}{n}\right)^{\frac{\beta+1}{2(\beta+1)+1}} \preccurlyeq \delta.$$
(115)

Assume KERNEL, EIGENVALUE, COVER, where the symbols d and x_i in the assumptions are d = 1 and $x_{[1,n]} = t_{[1,n]}$. Assume SUBGAUSSIAN, $s \simeq 1$, and

$$\left(\frac{\log(n)}{n}\right)^{\frac{1}{2(\beta+1)}} \preccurlyeq T \prec n.$$
(116)

(i) Set $\mathcal{X} := \mathcal{G}_{\ell,\delta,s}(u^{\star}([0,T]))$. Then

$$\sup_{x \in \mathcal{X}} \left\| \hat{f}(x) - f(x) \right\|_{2}^{2} \in \mathbf{O}_{\mathbf{P}}\left(\delta^{2\beta} + \left(\frac{T \log n}{n}\right)^{\frac{2\beta}{2(\beta+1)+1}} \right).$$
(117)

(ii) Set $\delta := \delta_{\mathsf{NN}}^{\ell,s}(u^{\star},T,[0,1]^d)$. Then

$$\sup_{x \in [0,1]^d} \left\| \hat{f}(x) - f(x) \right\|_2^2 \in \mathbf{O}_{\mathbf{P}} \left(\delta^{2\beta} + \left(\frac{T \log n}{n} \right)^{\frac{2\beta}{2(\beta+1)+1}} \right).$$
(118)

Remark 4.18.

(i) From [SS24, Corollary 4.11], we obtain

$$\mathbf{E}\left[\sup_{x\in[0,1]^d}\left\|\hat{f}(x) - f(x)\right\|_2^2\right] \succcurlyeq \delta^{2\beta} + \left(\delta^{d-1}\frac{T\log(n)}{n}\right)^{\frac{2\beta}{2(\beta+1)+d}}.$$
 (119)

if $\log(n) \asymp \log(\delta^{-(d-1)}nT^{-1})$ and

$$\max\left(\left(\frac{T}{n}\right)^{2\beta+d+1}, T^{-1}\right) \preccurlyeq \delta^{d-1} \preccurlyeq \min\left(1, \frac{n}{T}\right) \,. \tag{120}$$

Thus, the rate given in (118) is minimax optimal if $\delta^{2\beta}$ is the dominating term the error bound, i.e., if

$$\delta \succcurlyeq \left(\frac{T\log n}{n}\right)^{\frac{1}{2(\beta+1)+1}}.$$
(121)

This condition is fulfilled in the setting of Corollary 4.19 below. See Remark 4.5 for a discussion of the condition for $\beta = 1$.

- (ii) In section 4.1 can extrapolate beyond the convex hull of $\hat{u}([0,T])$ in the case of $\beta = 1$ using nearest neighbors. Whereas our general results are only available for interpolation, i.e., for $x \in ch(\hat{u}([0,T]))$, to be precise, only for the even more restrictive assumption $x \in \mathcal{G}_{\ell,\delta,s}(u^*([0,T]))$. Extrapolation seems also possible using polynomials of degree > 0, but the technicalities seem more difficult.
- (iii) In practice, finding $\chi_{\ell,s}$ may be computationally demanding. Furthermore, using only results of interpolation may be inconveniently restrictive. Thus, one may want to replace the polynomial interpolation step by a polynomial regression step on the $k \ge N$ nearest neighbors of $\hat{u}(\tau_j)$ for some chosen time points $\tau_1, \ldots, \tau_J \in [0, T]$. Another alternative to polynomials would be Gaussian process interpolation (or regression) [RW06], potentially also restricted to k nearest neighbors for performance reasons.

The lowest upper bound on the error is obtained if the two terms in (118) are balanced. Additionally, we require $T\delta^{d-1} \geq 1$ to be able to cover the domain of interest with $B^d(U(f^*, x_1, [0, T]), \delta)$, which is necessary for $[0, 1]^d \subseteq \mathcal{G}_{\ell,\delta,s}(u^*([0, T]))$.

Corollary 4.19. Use the setting and assumptions of Corollary 4.17. Set Set
$$\delta := \delta_{\mathsf{NN}}^{\ell,s}(u^*, T, [0, 1]^d)$$

Assume $T \preccurlyeq \left(\frac{n}{\log n}\right)^{\frac{d-1}{2(\beta+1)+d}}$ and $\delta \preccurlyeq \left(\frac{n}{\log n}\right)^{-\frac{1}{2(\beta+1)+d}}$. Then
$$\sup_{x \in [0,1]^d} \left\| \hat{f}(x_0) - f(x_0) \right\|_2^2 \in \mathbf{O}_{\mathbf{P}}\left(\left(\frac{n}{\log n}\right)^{-\frac{2\beta}{2(\beta+1)+d}} \right).$$
(122)

4.2.5 Proof

As matrix inversion is continuous for the set of invertible matrices, we have following lemma.

Lemma 4.20. Let $d \in \mathbb{N}$. For $n \in \mathbb{N}$, let $A_n, B_n \in \mathbb{R}^{d \times d}$. Assume that A_n is invertible for n large enough. Assume $||A_n - B_n||_{op} \prec 1$ and $||A_n^{-1}||_{op} \preccurlyeq 1$. Then B_n is invertible for n large enough and $||A_n^{-1} - B_n^{-1}||_{op} \prec 1$.

Lemma 4.21. Let $\delta, s > 0$. Let $x_0 \in \mathcal{G}_{\beta,\delta,s}(u^{\star}([0,T]))$. Let

$$\Gamma := \sup_{t \in [0,T]} \|\hat{u}(t) - u^{\star}(t)\|_{2}, \qquad \Lambda := \sup_{t \in [0,T]} \left\|\hat{u}(t) - \dot{u}^{\star}(t)\right\|_{2}.$$
(123)

Assume $\Gamma \prec 1$, $\Gamma \preccurlyeq \delta$, $\Gamma \preccurlyeq \Lambda \preccurlyeq 1$, $s \asymp 1$. Then $\hat{f}(x_0)$ eventually (meaning for n large enough) exists and

$$\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2 \preccurlyeq \Lambda + \delta^{\beta} \,. \tag{124}$$

Proof. To show that $\hat{f}(x_0)$ eventually exists, we need to prove that $\mathcal{H}_{\ell,s}(\hat{u}([0,T]),x)$ is eventually nonempty. As $x_0 \in \mathcal{G}_{\beta,\delta,s}(u^*([0,T]))$, there is $\mathbf{x} \in u^*([0,T])^N$ with $x \in \hat{ch}(\mathbf{x})$ such that $\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\|_{op} < s$. As $\Gamma \prec 1$, the inequality in s is strict, and $\hat{ch}(\mathbf{x})$ is open, there must eventually be a $\hat{\mathbf{x}} \in \hat{u}([0,T])^N$ with $x \in \hat{ch}(\hat{\mathbf{x}})$ and $\|\Psi(\eta_{\hat{\mathbf{x}}}(\hat{\mathbf{x}}))^{-1}\|_{op} < s$. Thus, $\hat{\mathbf{x}}(x_0)$ exists and so does $\hat{f}(x_0)$.

Let us establish some notation. Let $\hat{\mathbf{x}} := \hat{\mathbf{x}}(x_0)$, Recall that $\hat{\mathbf{x}} = (\hat{u}(\hat{\tau}_1), \dots, \hat{u}(\hat{\tau}_N)) = \chi_{\ell,s}(\hat{u}([0,T]), x_0)$. Let $(u^*(\tau_1), \dots, u^*(\tau_N)) = \mathbf{x}$ be any set such that $x_0 \in \mathring{ch}(\mathbf{x})$, $\|\Psi(\bar{\mathbf{x}})^{-1}\|_{op} < s$, and $\operatorname{diam}(\mathbf{x}) \leq \delta$, which exists as $x_0 \in \mathcal{G}_{\beta,\delta,s}(u^*([0,T]))$. We summarize and add some additional notation:

$$\mathbf{x} = (u^{\star}(\tau_1), \dots, u^{\star}(\tau_N)), \qquad \mathbf{y} := (\dot{u}^{\star}(\tau_1), \dots, \dot{u}^{\star}(\tau_N)), \qquad (125)$$

$$\bar{\mathbf{x}} := \left(u^{\star}(\hat{\tau}_1), \dots, u^{\star}(\hat{\tau}_N)\right), \qquad \bar{\mathbf{y}} := \left(\dot{u}^{\star}(\hat{\tau}_1), \dots, \dot{u}^{\star}(\hat{\tau}_N)\right), \qquad (126)$$

$$\tilde{\mathbf{x}} := (\hat{u}(\tau_1), \dots, \hat{u}(\tau_N)), \qquad \tilde{\mathbf{y}} := (\hat{\dot{u}}(\tau_1), \dots, \hat{\dot{u}}(\tau_N)), \qquad (127)$$

$$\mathbf{x} = (u^{\star}(\tau_1), \dots, u^{\star}(\tau_N)), \qquad \mathbf{y} := (\dot{u}^{\star}(\tau_1), \dots, \dot{u}^{\star}(\tau_N)). \qquad (128)$$

Using the definition of \hat{f} and the triangle inequality, we can split the error into two parts,

$$\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2 \le \left\|I(\hat{\mathbf{x}}, \hat{\mathbf{y}}, x_0) - I(\hat{\mathbf{x}}, \bar{\mathbf{y}}, x_0)\right\|_2 + \left\|I(\hat{\mathbf{x}}, \bar{\mathbf{y}}, x_0) - f^{\star}(x_0)\right\|_2.$$
(129)

For first term, with Lemma C.5, we obtain

$$\|I(\hat{\mathbf{x}}, \hat{\mathbf{y}}, x_0) - I(\hat{\mathbf{x}}, \bar{\mathbf{y}}, x_0)\|_2 = \|\psi(x_0)^{\mathsf{T}} \Psi(\hat{\mathbf{x}})^{-1} (\hat{\mathbf{y}} - \bar{\mathbf{y}})\|_2$$
(130)

$$= \left\| \psi(\eta_{\hat{\mathbf{x}}}(x_0))^{\top} \Psi(\eta_{\hat{\mathbf{x}}}(\hat{\mathbf{x}}))^{-1} \left(\hat{\mathbf{y}} - \bar{\mathbf{y}} \right) \right\|_2$$
(131)

$$\leq \left\|\psi(\eta_{\hat{\mathbf{x}}}(x_0))\right\|_2 \left\|\Psi(\eta_{\hat{\mathbf{x}}}(\hat{\mathbf{x}}))^{-1}\right\|_{\mathsf{op}} \left\|\hat{\mathbf{y}} - \bar{\mathbf{y}}\right\|_{\mathsf{op}}$$
(132)

$$\leq c_N s \Lambda$$
, (133)

where we used $\|\eta_{\hat{\mathbf{x}}}(x_0)\|_2 \leq 1$, $\hat{\mathbf{x}} \in \mathcal{H}_{\ell,s}(\hat{u}([0,T]), x_0)$, and the definition of Λ in the last step. Thus, we have

$$\|I(\hat{\mathbf{x}}, \hat{\mathbf{y}}, x_0) - I(\hat{\mathbf{x}}, \bar{\mathbf{y}}, x_0)\|_2 \preccurlyeq \Lambda$$
(134)

For the second term, we apply Lemma C.6 and get

$$\|I(\hat{\mathbf{x}}, \bar{\mathbf{y}}, x_0) - f^{\star}(x_0)\|_2 \le C_{\beta}(\hat{\mathbf{x}}) \operatorname{diam}(\hat{\mathbf{x}})^{\beta}.$$
(135)

With Lemma C.7 and the minimizing property of $\mathbf{\hat{x}} = \chi_{\ell,s}(\hat{u}([0,T]), x_0)$, we obtain

$$C_{\beta}(\hat{\mathbf{x}})\operatorname{diam}(\hat{\mathbf{x}})^{\beta} \le \frac{N^{\frac{3}{2}}}{\beta!} \left\| \Psi(\eta_{\hat{\mathbf{x}}}(\hat{\mathbf{x}}))^{-1} \right\|_{\mathsf{op}} \operatorname{diam}(\hat{\mathbf{x}})^{\beta}$$
(136)

$$\leq (1+\epsilon) \frac{N^{\frac{3}{2}}}{\beta!} \left\| \Psi(\eta_{\tilde{\mathbf{x}}}(\tilde{\mathbf{x}})))^{-1} \right\|_{\mathsf{op}} \operatorname{diam}(\tilde{\mathbf{x}})^{\beta}.$$
(137)

With $\|\hat{u}(\tau_j) - u^{\star}(\tau_j)\|_2 \leq \Gamma$ and the definition of **x**, we have

$$\operatorname{diam}(\tilde{\mathbf{x}}) \le \operatorname{diam}(\mathbf{x}) + 2\Gamma \preccurlyeq \delta + \Gamma.$$
(138)

As $\|\tilde{\mathbf{x}} - \mathbf{x}\|_{op} \prec 1$, we have

$$\left\|\Psi(\eta_{\tilde{\mathbf{x}}}(\tilde{\mathbf{x}})))^{-1}\right\|_{\mathsf{op}} \preccurlyeq \left\|\Psi(\eta_{\mathbf{x}}(\tilde{\mathbf{x}})))^{-1}\right\|_{\mathsf{op}} \tag{139}$$

$$\preccurlyeq \left\|\Psi(\eta_{\mathbf{x}}(\mathbf{x})))^{-1}\right\|_{\mathsf{op}} + \left\|\Psi(\eta_{\mathbf{x}}(\tilde{\mathbf{x}})))^{-1} - \Psi(\eta_{\mathbf{x}}(\mathbf{x})))^{-1}\right\|_{\mathsf{op}}.$$
 (140)

Using $\|\tilde{\mathbf{x}} - \mathbf{x}\|_{\mathsf{op}} \prec 1$ again, we obtain

$$\|\Psi(\eta_{\mathbf{x}}(\tilde{\mathbf{x}})) - \Psi(\eta_{\mathbf{x}}(\mathbf{x}))\|_{\mathsf{op}} \prec 1.$$
(141)

Together with $\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\|_{\mathsf{op}} < s$, Lemma 4.20 then yields

$$\left\|\Psi(\eta_{\mathbf{x}}(\hat{\mathbf{x}}))^{-1} - \Psi(\eta_{\mathbf{x}}(\bar{\mathbf{x}}))^{-1}\right\|_{\mathsf{op}} \prec 1.$$
(142)

Putting the previous inequalities into (135), we obtain the following bound on the second term:

$$\|I(\hat{\mathbf{x}}, \mathbf{y}, x) - f^{\star}(x)\|_{2} \preccurlyeq (\delta + \Gamma)^{\beta} .$$
(143)

The bounds (134) and (143) on the two terms of the right-hand side of (129) together yield

$$\left\|\hat{f}(x_0) - f^{\star}(x_0)\right\|_2 \preccurlyeq \Lambda + (\delta + \Gamma)^{\beta} \preccurlyeq \Lambda + \delta^{\beta}$$
(144)

as $\Gamma \preccurlyeq \delta$.

Proof of Theorem 4.16. Set $\tilde{\beta} = \beta + 1$, $\tilde{L}_0 = \infty$ and $\tilde{L}_k = (k-1)! \sup_{\ell \in [\![0,\beta]\!]} L^k_\ell$ for $k \in [\![\tilde{\beta}]\!]$. As $f \in \bar{\Sigma}^{d \to d}(\beta, L_{[\![0,\beta]\!]})$, Corollary D.7 yields $u \in \bar{\Sigma}^{1 \to d}(\beta + 1, \tilde{L}_{[\![0,\tilde{\beta}]\!]})$. Hence, we can use

$$\Gamma = \Gamma(n, \tilde{\beta}, \tilde{L}_{\tilde{\beta}}, T) \quad \text{and} \quad \Lambda = \Lambda(n, \tilde{\beta}, \tilde{L}_{\tilde{\beta}}, T)$$
(145)

in Lemma 4.21 to obtain

$$\sup_{x \in \mathcal{X}} \left\| \hat{f}(x) - f(x) \right\|_{2}^{2} \in \mathbf{O}_{\mathbf{P}} \left(\Lambda(n, \tilde{\beta}, \tilde{L}_{\tilde{\beta}}, T)^{2} + \delta^{2\beta} \right).$$
(146)

A Derivatives

In this section, we introduce notation for derivatives in higher dimensions and a Hölder-type smoothness class. Furthermore, some elementary analytical results are shown for reference in the main proofs.

For any finite-dimensional \mathbb{R} -vector space V, we denote the Euclidean norm as $||x||_2$ for $x \in V$. For $k \in \mathbb{N}$ and finite-dimensional \mathbb{R} -vector spaces V and W, let $\mathcal{L}_k(V, W)$ be the set of k-multilinear functions from V^k to W. An element of $\mathcal{L}_k(V, W)$ is called symmetric if it is invariant under

permutations of its k arguments. A function $f: V \to W$ is differentiable at $x \in V$ if there is $A \in \mathcal{L}_1(V, W)$ such that

$$\lim_{\|v\|_{2} \to 0} \frac{\|f(x+v) - f(x) - A(v)\|_{2}}{\|v\|_{2}} = 0.$$
(147)

In this case we write Df(x) = A. The function f is differentiable if it is differentiable at all $x \in V$. Denote the set of differentiable functions from V to W as $\mathcal{D}(V, W)$. From now on, assume $V = \mathbb{R}^d$ for a $d \in \mathbb{N}$. Define $\mathbb{D}_d := \{v \in \mathbb{R}^d \mid ||v||_2 = 1\}$. The directional derivative of $f \in \mathcal{D}(V, W)$ at $x \in V$ in the direction $v \in \mathbb{D}_d$ is

$$D_v f(x) := Df(x)(v).$$
(148)

We set D^0 to the identity, i.e., $D^0 f = f$. Let $k \in \mathbb{N}$. Define the set of k-times differentiable functions $\mathcal{D}^k(V, W)$ and the k-th derivative operator D^k recursively: For k = 1, we set $\mathcal{D}^1(V, W) :=$ $\mathcal{D}(V, W)$ and $D^1 := D$. Let $k \in \mathbb{N}_{\geq 2}$. A function $f \in \mathcal{D}^k(V, W)$ is k-times differentiable at $x \in V$ if there is symmetric $A \in \mathcal{L}_k(V, W)$ such that

$$\lim_{\|v\|_{2} \to 0} \sup_{\mathbf{v} \in \mathbb{D}_{d}^{k-1}} \frac{\left\| D_{\mathbf{v}}^{k-1} f(x+v) - D_{\mathbf{v}}^{k-1} f(x) - A((\mathbf{v},v)) \right\|_{2}}{\|v\|_{2}} = 0.$$
(149)

In this case we write $D^k f(x) = A$. We may use square brackets for the arguments of the linear operator to make formulas more clear, e.g., Df(x)[v] := Df(x)(v) The function f is k-times differentiable if it is k-times differentiable at all $x \in V$. Denote the set of k-times differentiable functions from V to W as $\mathcal{D}^k(V, W)$. The k-th directional derivative of $f \in \mathcal{D}^k(V, W)$ at $x \in V$ in the directions $\mathbf{v} \in \mathbb{D}^k_d$ is

$$D_{\mathbf{v}}f(x) := D^k f(x)(\mathbf{v}) \,. \tag{150}$$

We may indicate that differentiation takes place with respect to x by $D_{x,\mathbf{v}}g(x,y) = D_{\mathbf{v}}f_y(x)$, where $f_y(x) = g(x,y)$. Let $p \in \mathbb{N}$. For $f \in \mathcal{D}^k(\mathbb{R}^d, \mathbb{R}^p)$ the derivative operator acts component-wise, i.e., for $f = (f_1, \ldots, f_p)$ with $f_\ell \in \mathcal{D}^k(\mathbb{R}^d, \mathbb{R})$, we have

$$D_{\mathbf{v}}f(x) = \begin{pmatrix} D_{\mathbf{v}}f_1(x)\\ \vdots\\ D_{\mathbf{v}}f_p(x) \end{pmatrix}, \qquad (151)$$

for $x \in \mathbb{R}^d$ and $\mathbf{v} \in \mathbb{D}_d^k$. The operator norm for $A \in \mathcal{L}_k(V, W)$ is

$$\|A\|_{\mathsf{op}} := \sup_{\mathbf{v} \in \mathbb{D}_d^k} \|A(\mathbf{v})\|_2 .$$
(152)

Define the sup-norm of the k-th derivative as

$$|D^k f|_{\infty} := \sup_{x \in \mathbb{R}^d} \|D^k f(x)\|_{\mathsf{op}} = \sup_{x \in \mathbb{R}^d} \sup_{\mathbf{v} \in \mathbb{D}^k_d} \|D_{\mathbf{v}} f(x)\|_2 .$$
(153)

Notation A.1. Let $x \in \mathbb{R}$. Denote the largest integer strictly smaller than x as $||x|| := \max \mathbb{Z}_{\leq x}$.

Definition A.2 (Hölder-smoothness classes). Let
$$d \in \mathbb{N}$$
, For $L \in \mathbb{R}_{>0}$, define

$$\Sigma^{d \to 1}(0, L) := \left\{ f \colon \mathbb{R}^d \to \mathbb{R} \mid |f|_{\infty} \leq L \right\}.$$
(154)

For $L \in \mathbb{R}_{>0}$, $\beta \in (0, 1]$, define

$$\Sigma^{d \to 1}(\beta, L) := \left\{ f \colon \mathbb{R}^d \to \mathbb{R} \middle| |f(x) - f(\tilde{x})| \le L ||x - \tilde{x}||_2^\beta \right\}.$$
(155)

Let $\beta \in \mathbb{R}_{>1}$ and $\ell := \|\beta\|$. Define

$$\Sigma^{d \to 1}(\beta, L) := \left\{ f \in \mathcal{D}^{\ell}(\mathbb{R}^d, \mathbb{R}) \, \middle| \, \| D^{\ell} f(x) - D^{\ell} f(\tilde{x}) \|_{\mathsf{op}} \le L \| x - \tilde{x} \|_{2}^{\beta - \ell} \right\} \,.$$
(156)

For $\beta \in \mathbb{R}_{\geq 0}$, use the short notation

$$\Sigma(\beta, L) := \Sigma^{1 \to 1}(\beta, L) \,. \tag{157}$$

Let $L_0, \ldots, L_\ell \in \mathbb{R}_{>0} \cup \{\infty\}, L_\beta \in \mathbb{R}_{>0}$. Define

$$\Sigma^{d \to 1}(\beta, L_{\llbracket 0, \ell \rrbracket}, L_{\beta}) := \Sigma^{d \to 1}(\beta, L_{\beta}) \cap \bigcap_{k=0}^{\ell} \Sigma^{d \to 1}(k, L_k).$$
(158)

Let $d_{in}, d_{out} \in \mathbb{N}$. Define

$$\Sigma^{d_{\rm in} \to d_{\rm out}}(\beta, L_{\llbracket 0, \ell \rrbracket}, L_{\beta}) := \left(\Sigma^{d_{\rm in} \to 1}(\beta, L_{\llbracket 0, \ell \rrbracket}, L_{\beta})\right)^{d_{\rm out}},$$
(159)

where $f = (f_1, \ldots, f_{d_{out}}) \in \Sigma^{d_{in} \to d_{out}}(\beta, L_{[0,\ell]}, L_{\beta})$ is treated as a function

$$f: \mathbb{R}^{d_{\text{in}}} \to \mathbb{R}^{d_{\text{out}}}, x \mapsto (f_1(x), \dots, f_{d_{\text{out}}}(x))^{\top}.$$
 (160)

If β is an integer, we denote

$$\bar{\Sigma}^{d_{\rm in} \to d_{\rm out}}(\beta, L_{[\![0,\beta]\!]}) := \Sigma^{d_{\rm in} \to d_{\rm out}}(\beta, L_{[\![0,\beta]\!]}) \cap \mathcal{D}^{\beta}(\mathbb{R}^{d_{\rm in}}, \mathbb{R}^{d_{\rm out}}).$$
(161)

A bound on the derivative yields the Lipschitz constant, as the next lemma shows.

Lemma A.3. Let $d \in \mathbb{N}$. Let $k \in \mathbb{N}$. Let $f \in \mathcal{D}^k(\mathbb{R}^d, \mathbb{R})$. Let $L \in \mathbb{R}_{>0}$. Assume $|D^k f|_{\infty} \leq L$. Then $f \in \Sigma^{d \to 1}(k, L)$.

Proof of Lemma A.3. Let $\mathbf{v} \in \mathbb{D}_d^{k-1}$. Then, by the mean value theorem,

$$\left| D_{\mathbf{v}}^{k-1} f(x) - D_{\mathbf{v}}^{k-1} f(\tilde{x}) \right| \le \sup_{\tilde{v} \in \mathbb{D}_d} \left| D_{(\mathbf{v},\tilde{v})}^k f \right|_{\infty} \|x - \tilde{x}\|_2$$
(162)

with $(\mathbf{v}, \tilde{v}) \in \mathbb{D}_d^k$. Thus,

$$\left\| D^{k-1} f(x) - D^{k-1} f(\tilde{x}) \right\|_{\mathsf{op}} \le \left\| D^k f \right\|_{\infty} \|x - \tilde{x}\|_2 .$$
(163)

The sub-multiplicativity of the operator norm in the following lemma is a direct consequence of its definition.

Lemma A.4. Let
$$k, d, p \in \mathbb{N}$$
. Let $A \in \mathcal{L}_k(\mathbb{R}^d, \mathbb{R}^p)$ and $v \in \mathbb{D}_d^k$. Then

$$\|A(v_1, \dots, v_k)\|_2 \le \|A\|_{\mathsf{op}} \prod_{j=1}^k \|v_j\|_2 \tag{164}$$

For reference, we state the chain and product rule as well es Taylor's approximation theorem in the following lemmas.

Lemma A.5. Let $d \in \mathbb{N}$. Let $A \colon \mathbb{R}^d \to \mathcal{L}_k(\mathbb{R}^d, \mathbb{R}^d)$ symmetric. Let $v \in \mathbb{D}_d$.

(i) Product rule: Let $f_{\ell} \in \mathcal{D}(\mathbb{R}^d, \mathbb{R}^d)$ for $\ell \in [\![k]\!]$. Then

$$D_v(A(x)(f_1(x),\ldots,f_k(x)))$$
 (165)

$$= (D_v A(x))(f_1(x), \dots, f_k(x)) +$$
(166)

$$\sum_{\ell=1}^{n} A(x)(f_1(x), \dots, f_{\ell-1}(x), D_v f_\ell(x), f_{\ell+1}(x), \dots, f_k(x)) .$$
 (167)

(ii) Chain rule: Let $f \in \mathcal{D}(\mathbb{R}^d, \mathbb{R}^d)$. Then

$$D_v(A(f(x))(a_1,\ldots,a_k)) = (DA)(f(x))(a_1,\ldots,a_k,D_vf(x)).$$
(168)

Notation A.6. Let $d, m \in \mathbb{N}$. Let $x \in \mathbb{R}^d$. Denote $\{x\}^m = (x, \dots, x) \in (\mathbb{R}^d)^m$.

Lemma A.7 (Taylor theorem with Peano remainder). Let $d \in \mathbb{N}$.

(i) Let $\ell \in \mathbb{N}_0$. Let $f \in \mathcal{D}^{\ell}(\mathbb{R}^d, \mathbb{R})$. There is $\gamma \in [0, 1]$, such that

$$f(x) = \sum_{k=0}^{\ell-1} \frac{D^k f(x_0)(\{x - x_0\}^k)}{k!} + \frac{D^\ell f(\tilde{x}) \left(\{x - x_0\}^\ell\right)}{\ell!}$$
(169)

with $\tilde{x} := x_0 + \gamma (x - x_0)$.

(ii) Let $\beta, L \in \mathbb{R}_{>0}$. Set $\ell := \|\beta\|$. Let $f \in \Sigma^{d \to 1}(\beta, L)$. Then

$$f(x) - \sum_{k=0}^{\ell} \frac{D^k f(x_0)(\{x - x_0\}^k)}{k!} \le L \frac{\|x - x_0\|_2^{\beta}}{\ell!}.$$
(170)

B Local Polynomial Regression

In order to make this article more self-contained, we provide here upper bounds on the estimation error for a multivariate regression problem using local polynomial estimators. We follow [Tsy09] and extend the proof there from the domain [0, 1] to the domain $[0, T]^d$ and also discuss estimation of derivatives. See also [RW94]. It is important for the results in the main part of the article that the dependence on T and on the Lipschitz constant L of the regression function are given explicitly.

B.1 Model

Let $d \in \mathbb{N}$. Let $\beta, L \in \mathbb{R}_{>0}$. Set $\ell := \|\beta\|$. Let $f \in \Sigma^{d \to 1}(\beta, L)$. Let $n \in \mathbb{N}$. Let $T \in \mathbb{R}_{>0}$. For $i \in [n]$, let $x_i \in [0, T]^d$. Let $\sigma \in \mathbb{R}_{>0}$. For $i \in [n]$, let ε_i be independent random variables with $\mathbf{E}[\epsilon_i] = 0$ and $\mathbf{E}[\epsilon_i^2] \leq \sigma^2$. For $i \in [n]$, set

$$Y_i = f(x_i) + \varepsilon_i \,. \tag{171}$$

We observe Y_i and know x_i . The function f is unknown. Let $s \in [0, \ell]$. Let $\mathbf{v} \in \mathbb{D}_d^s$ be s-many directions. Our target is to estimate the s-th directional derivative $D_{\mathbf{v}}f \colon \mathbb{R}^d \to \mathbb{R}$ of f. For the local polynomial estimator \hat{g} defined below, we want to show upper bounds for the mean squared error $\mathbf{E}[\|\hat{g}(x) - D_{\mathbf{v}}f(x)\|^2]$ at a point $x \in [0, T]^d$ depending on $d, \beta, L, n, T, \sigma, s$.

B.2 Estimator

Recall Notation 3.8. Set the number of scalar coefficients for polynomials $\mathbb{R}^d \to \mathbb{R}$ of degree at most ℓ as

$$N := N_{d,\ell} := \dim(\mathcal{P}_{d,\ell}) = \begin{pmatrix} \ell + d \\ d \end{pmatrix}.$$
(172)

For $x \in \mathbb{R}^d$, denote $\psi(x) := \psi_\ell(x) := ((\alpha!)^{-1} x^\alpha)_{\alpha \in \mathbb{N}^d_0, |\alpha| \le \ell} \in \mathbb{R}^N$. Let $K : \mathbb{R}_{\ge 0} \to \mathbb{R}$. Let $h \in \mathbb{R}_{>0}$. Denote the local polynomial mean squared error coefficients as

$$\hat{\theta}(x) \in \operatorname*{arg\,min}_{\theta \in \mathbb{R}^N} \sum_{i=1}^n \left(Y_i - \theta^{\mathsf{T}} \psi\left(\frac{x_i - x}{h}\right) \right)^2 K\left(\frac{\|x_i - x\|_2}{h}\right) \,. \tag{173}$$

Then the local polynomial estimator of degree ℓ for the s-th derivative in directions $\mathbf{v} \in \mathbb{D}_d^s$ is defined as

$$\hat{g}(x) := h^{-s} D_{\mathbf{v}} \psi(0)^{\mathsf{T}} \hat{\theta}(x) \,. \tag{174}$$

For $x \in \mathbb{R}^d$, define the symmetric matrix $B(x) \in \mathbb{R}^{N \times N}$ and the vector $a(x) \in \mathbb{R}^N$ as

$$B(x) := \frac{T^d}{nh^d} \sum_{i=1}^n \psi\left(\frac{x_i - x}{h}\right) \psi\left(\frac{x_i - x}{h}\right)^\top K\left(\frac{\|x_i - x\|_2}{h}\right) , \qquad (175)$$

$$a(x) := \frac{T^d}{nh^d} \sum_{i=1}^n Y_i \psi\left(\frac{x_i - x}{h}\right) K\left(\frac{\|x_i - x\|_2}{h}\right) \,. \tag{176}$$

Notation B.1. Let $k \in \mathbb{N}$ and $A \in \mathbb{R}^{k \times k}$ be a symmetric matrix. We write $\lambda_{\min}(A)$ for the smallest eigenvalue of A.

Assume $\lambda_{\min}(B(x)) > 0$. Then

$$\hat{\theta}(x) = B(x)^{-1}a(x).$$
 (177)

Furthermore, with the weights

$$w_{i}(x) := w_{i,s}(x) := \frac{T^{d}}{nh^{d+s}} D_{\mathbf{v}} \psi(0)^{\mathsf{T}} B(x)^{-1} \psi\left(\frac{x_{i}-x}{h}\right) K\left(\frac{x_{i}-x}{h}\right) , \qquad (178)$$

we can write the estimator as

$$\hat{g}(x) = h^{-s} D_{\mathbf{v}} \psi(0)^{\mathsf{T}} \hat{\theta}(x) = \sum_{i=1}^{n} w_i(x) Y_i \,. \tag{179}$$

B.3 Result

Notation B.2. All lower case c, with or without index, are elements of $\mathbb{R}_{>0}$ and universal insofar as they only depend on the variables written as index, e.g., $c_{d,\beta}$ depends only on β and d. In particular, a constant c with no index refers to a fixed positive number. Every occurrence of such a variable may refer to a different value.

Theorem B.3. Use the model of section B.1 and the estimator of section B.2. Assume KERNEL, EIGENVALUE, and COVER. Assume $n \ge C_{\mathsf{cvr}}^{-1}h^{-d}T^d$. Then, for all $x \in [0,T]^d$, we have

$$\mathbf{E}\left[\left(\hat{g}(x) - D_{\mathbf{v}}f(x)\right)^{2}\right] \leq c_{\ell} \left(C_{\mathsf{egv}}^{2}C_{\mathsf{ker}}^{2}C_{\mathsf{cvr}}^{2}L^{2}h^{2(\beta-s)} + C_{\mathsf{egv}}^{2}C_{\mathsf{ker}}^{2}C_{\mathsf{cvr}}\frac{\sigma^{2}T^{d}}{nh^{d+2s}}\right).$$
(180)

Corollary B.4. Use the model of section B.1 and the estimator of section B.2. Assume KERNEL, EIGENVALUE, and COVER. Assume $n \ge n_0$, where $n_0 \in \mathbb{N}$ depends only on $d, \beta, C_{egv}, C_{ker}, C_{cvr}, \sigma, T, L$. Then there are constants $C_1, C_2 \in \mathbb{R}_{>0}$, depending only on $d, \beta, C_{egv}, C_{ker}, C_{cvr}$, with the following property: Set the bandwidth as

$$h = C_1 \left(\frac{\sigma^2 T^d}{L^2 n}\right)^{\frac{1}{2\beta+d}} .$$
(181)

Then

$$\mathbf{E}\Big[(\hat{g}(x) - D_{\mathbf{v}}f(x))^2\Big] \le C_2 L^{\frac{2d+4s}{2\beta+d}} \left(\frac{\sigma^2 T^d}{n}\right)^{\frac{2(\beta-s)}{2\beta+d}}.$$
(182)

Remark B.5.

(i) The error rates obtained in Corollary B.4 are minimax optimal up to the constant C_2 , which can be seen by comparing them to lower bounds for nonparametric regression, e.g., [SS24, Theorem 8].

(ii) Integrating the point-wise error over $[0, T]^d$ yields the minimax optimal rate for the integrated error. For a bound on the error in sup-norm, we need to do some extra work.

Theorem B.6. Assume KERNEL, EIGENVALUE, and COVER. Assume SUBGAUSSIAN. Assume $n \ge C_{\mathsf{cvr}}^{-1}h^{-d}T^d$ Assume $T/h \ge 2$. Then, we have

$$\mathbf{E}\left[\left|\hat{g}(x) - D_{\mathbf{v}}f(x)\right|_{\infty}^{2}\right] \le c_{d,\ell,K} \left(\left(C_{\mathsf{egv}}C_{\mathsf{cvr}}Lh^{\beta-s}\right)^{2} + C_{\mathsf{cvr}}C_{\mathsf{egv}}^{2}\frac{\sigma^{2}T^{d}}{nh^{d+2s}}\log(T/h)\right).$$
(183)

Corollary B.7. Assume KERNEL, EIGENVALUE, and COVER. Assume SUBGAUSSIAN. There are $C_1, C_2, C_3, C_4 \in \mathbb{R}_{>0}$ depending only on $d, \beta, C_{egv}, C_{cvr}, K$ with the following property: Assume

$$C_1\left(\frac{\log(n)}{n}\right)^{\frac{d}{2\beta}} \le C_2\left(\frac{L^2}{\sigma^2}\right)^{\frac{d}{2\beta}} T^d \le n\log(n)^{\frac{d}{2\beta}}.$$
(184)

Set the bandwidth as

$$h = C_3 \left(\frac{\sigma^2 T^d \log(n)}{L^2 n}\right)^{\frac{1}{2\beta+d}}.$$
(185)

Then

$$\mathbf{E}\left[\left|\hat{g} - D_{\mathbf{v}}f\right|_{\infty}^{2}\right] \le C_{4}L^{\frac{2d+4s}{2\beta+d}} \left(\frac{\sigma^{2}T^{d}\log(n)}{n}\right)^{\frac{2(\beta-s)}{2\beta+d}}.$$
(186)

Remark B.8.

(i) The error rates obtained in Corollary B.7 are minimax optimal up to the constant C_4 , which can be seen by comparing them to lower bounds for nonparametric regression, e.g., [SS24, Theorem 8].

Following [Tsy09, Lemma 1.4, Lemma 1.5], we show that the uniform grid in $[0, T]^d$ fulfills COVER, and also EIGENVALUE if the kernel is not degenerated.

Proposition B.9. Let $n_0 \in \mathbb{N}$. Set $n := n_0^d$. Let $x_i \in [0, T]^d$ for $i \in [n]$ form a uniform grid in $[0, T]^d$, i.e.,

$$\{x_1, \dots, x_n\} = \left\{ \left(T\frac{k_1}{n_0}, \dots, T\frac{k_d}{n_0} \right)^\top \middle| k_1, \dots, k_d \in [\![n_0]\!] \right\}.$$
 (187)

Assume STRICTKERNEL with constants $C_{\text{ker}} \in \mathbb{R}_{>0}$. Then KERNEL, EIGENVALUE, and COVER are fulfilled with C_{ker} , $C_{\text{egv}} = c_{d,\ell,K}(1 + \frac{T}{n_0h})$, $C_{\text{cvr}} = 4^d$.

B.4 Proof

Lemma B.10. Let $q \in \mathcal{P}_{d,\ell}$. Assume $\lambda_{\min}(B(x)) > 0$. Then

$$\sum_{i=1}^{n} w_i(x)q(x_i) = D_{\mathbf{v}}q(x).$$
(188)

Proof. We set Y_i to $q(x_i)$ and consider the local polynomial fit. The coefficients $\theta_q(x)$ in the fit are

$$\theta_q(x) \in \operatorname*{arg\,min}_{\theta \in \mathbb{R}^N} \sum_{i=1}^n \left(q(x_i) - \theta^{\mathsf{T}} \psi\left(\frac{x_i - x}{h}\right) \right)^2 K\left(\frac{\|x_i - x\|_2}{h}\right) \,. \tag{189}$$

Let $z \in \mathbb{R}^d$. Since q(z) is a polynomial of degree at most ℓ and $\lambda_{\min}(B(x)) > 0$, the fit of the degree

 ℓ polynomial with base $\psi((z-x)/h)$ retrieves the original function, i.e.,

$$q(z) = \theta_q(x)^{\mathsf{T}} \psi\left(\frac{z-x}{h}\right) \,. \tag{190}$$

Denote by $D_{z,\mathbf{v}}$ the s-th directional derivative in directions \mathbf{v} with respect to z. Then,

$$D_{\mathbf{v}}q(z) = \theta_q(x)^{\mathsf{T}} D_{z,\mathbf{v}}\left(\psi\left(\frac{z-x}{h}\right)\right) = h^{-s}\theta_q(x)^{\mathsf{T}} (D_{\mathbf{v}}\psi)\left(\frac{z-x}{h}\right).$$
(191)

Thus,

$$D_{\mathbf{v}}q(x) = h^{-s}\theta_q(x)^{\top}(D_{\mathbf{v}}\psi)(0) = \sum_{i=1}^n w_i(x)q(x_i).$$
(192)

Lemma B.11. Assume KERNEL, EIGENVALUE, and COVER. Assume $n \ge C_{\mathsf{cvr}}^{-1} h^{-d} T^d$.

- (i) If $||x x_i|| \ge h$, we have $w_i(x) = 0$.
- (ii) We have

$$\sum_{i=1}^{n} |w_i(x)| \le c_\ell C_{\text{egv}} C_{\text{ker}} C_{\text{cvr}} h^{-s} \,.$$
(193)

(iii) We have

$$\sum_{i=1}^{n} w_i(x)^2 \le c_\ell C_{\mathsf{egv}}^2 C_{\mathsf{ker}}^2 C_{\mathsf{cvr}} \frac{T^d}{nh^{d+2s}} \,. \tag{194}$$

Proof. Using KERNEL and EIGENVALUE, we obtain from (178)

$$nh^{d+s}T^{-d}|w_{i}(x)| \leq \|D_{\mathbf{v}}\psi(0)\|_{2} \|B(x)^{-1}\|_{\mathsf{op}} \left\|\psi\left(\frac{x_{i}-x}{h}\right)\right\|_{2} |K|_{\infty}\mathbb{1}_{\mathrm{B}^{d}(x_{i},h)}$$
(195)
$$\leq c_{\ell}C_{\mathsf{egv}}C_{\mathsf{ker}}\mathbb{1}_{\mathrm{B}^{d}(x_{i},h)},$$
(196)

$$\int c_{\ell} C_{\mathsf{egv}} C_{\mathsf{ker}} \mathbb{1}_{\mathrm{B}^d(x_i,h)}, \qquad (196)$$

which we use in each of the following parts.

- (i) Trivial.
- (ii) By COVER,

$$(C_{\mathsf{egv}}C_{\mathsf{ker}})^{-1}\sum_{i=1}^{n}|w_{i}(x)| \le c_{\ell}T^{d}h^{-(d+s)}\frac{1}{n}\#\left\{i\in[\![n]\!]\,\middle|\,x_{i}\in\mathsf{B}^{d}(x,h)\right\}$$
(197)

$$\leq c_{\ell} T^{d} h^{-(d+s)} \max\left(\frac{1}{n}, C_{\mathsf{cvr}} \left(\frac{h}{T}\right)^{d}\right) \tag{198}$$

$$\leq c_{\ell} C_{\mathsf{cvr}} h^{-s} \,, \tag{199}$$

as we assume $n \ge C_{\mathsf{cvr}}^{-1} h^{-d} T^d$.

(iii) With

$$|w_i(x)| \le c_\ell C_{\mathsf{egv}} C_{\mathsf{ker}} \frac{T^d}{nh^{d+s}} \tag{200}$$

and (ii), we obtain

$$\sum_{i=1}^{n} |w_i(x)|^2 \le \sup_{i \in [\![n]\!]} |w_i(x)| \sum_{i=1}^{n} |w_i(x)|$$
(201)

$$\leq c_{\ell} C_{\mathsf{egv}}^2 C_{\mathsf{ker}}^2 C_{\mathsf{cvr}} \frac{T^d}{nh^{d+2s}} \,. \tag{202}$$

Proof of Theorem B.3. **Bias.** By the model equation (171) and the linear representation of the estimator (179), we have

$$\mathbf{E}[\hat{g}(x)] = \sum_{i=1}^{n} w_i(x) f(x_i) \,. \tag{203}$$

Using a Taylor expansion (Lemma A.7) of f(x) at x_i yields the existence of $\gamma_i \in [0, 1]$ such that

$$f(x) = \sum_{k=0}^{\ell-1} \frac{D^k f(x)(\{x_i - x\}^k)}{k!} + \frac{D^\ell f(x + \gamma_i(x_i - x))(\{x_i - x\}^\ell)}{\ell!}.$$
 (204)

Denote the remainder term as

$$R_i(x) := \frac{D^\ell f(x + \gamma_i(x_i - x)) \left(\{x_i - x\}^\ell \right) - D^\ell f(x) \left(\{x_i - x\}^\ell \right)}{\ell!} \,. \tag{205}$$

By applying Lemma B.10 to

$$q(x) = \sum_{k=0}^{\ell} \frac{D^k f(x)(\{x_i - x\}^k)}{k!},$$
(206)

we obtain

$$\sum_{i=1}^{n} w_i(x) f(x_i) = \sum_{i=1}^{n} w_i(x) \sum_{k=0}^{\ell} \frac{D^k f(x) (\{x_i - x\}^k)}{k!} + \sum_{i=1}^{n} w_i(x) R_i(x)$$
(207)

$$= D_{\mathbf{v}}f(x) + \sum_{i=1}^{n} w_i(x)R_i(x) .$$
(208)

Hence, we have

$$\mathbf{E}[\hat{g}(x)] - D_{\mathbf{v}}f(x) = \sum_{i=1}^{n} w_i(x)R_i(x).$$
(209)

For bounding the remainder term, we note that $f \in \Sigma^{d \to 1}(\beta, L)$, which yields

$$\left| D^{\ell} f(x + \gamma_i(x_i - x)) \left(\{x_i - x\}^{\ell} \right) - D^{\ell} f(x) \left(\{x_i - x\}^{\ell} \right) \right| \le L \left\| x_i - x \right\|_2^{\beta}.$$
 (210)

Thus, with Lemma B.11 (i), we have

$$|\mathbf{E}[\hat{g}(x)] - D_{\mathbf{v}}f(x)| \le L \sum_{i=1}^{n} |w_i(x)| \, ||x_i - x||_2^{\beta}$$
(211)

$$\leq Lh^{\beta} \sum_{i=1}^{n} |w_i(x)|$$
 (212)

Finally, by Lemma B.11 (ii), we obtain

$$|\mathbf{E}[\hat{g}(x)] - D_{\mathbf{v}}f(x)| \le c_{\ell}C_{\mathsf{egv}}C_{\mathsf{ker}}C_{\mathsf{cvr}}Lh^{\beta-s}.$$
(213)

Variance. As the noise is independent and has a second moment bounded by σ^2 , we have

$$\mathbf{E}\left[\left(\hat{g}(x) - \mathbf{E}[\hat{g}(x)]\right)^{2}\right] = \mathbf{E}\left[\left(\sum_{i=1}^{n} \varepsilon_{i} w_{i}(x)\right)^{2}\right]$$
(214)

$$=\sum_{i=1}^{n} w_i(x)^2 \mathbf{E}\left[\varepsilon_i^2\right]$$
(215)

$$\leq \sigma^2 \sum_{i=1}^n w_i(x)^2 \,. \tag{216}$$

Hence, Lemma B.11 (iii) yields

$$\mathbf{E}\left[\left(\hat{g}(x) - \mathbf{E}[\hat{g}(x)]\right)^2\right] \le c_\ell C_{\mathsf{egv}}^2 C_{\mathsf{ker}}^2 C_{\mathsf{cvr}} \frac{\sigma^2 T^d}{nh^{d+2s}} \,.$$
(217)

Together. Putting (213) and (217) together, we obtain

$$\mathbf{E}\left[\left(\hat{g}(x) - D_{\mathbf{v}}f(x)\right)^{2}\right] \leq c_{\ell} \left(C_{\mathsf{egv}}^{2}C_{\mathsf{ker}}^{2}C_{\mathsf{cvr}}^{2}L^{2}h^{2(\beta-s)} + C_{\mathsf{egv}}^{2}C_{\mathsf{ker}}^{2}C_{\mathsf{cvr}}\frac{\sigma^{2}T^{d}}{nh^{d+2s}}\right).$$
(218)

For the proof in sup-norm (Theorem B.6), we want an upper bound on the supremum of a subgaussian process using the chaining technique. In order to do this, we need some preparation:

Definition B.12 (Sub-Gaussian process). Let (\mathcal{X}, ρ) be a metric space. A real-valued stochastic process $(Z_x)_{x \in \mathcal{X}}$ is *sub-Gaussian* if

$$\mathbf{P}(|Z_x - Z_{x'}| \ge t) \le 2\exp\left(-\frac{t^2}{2\rho(x, x')^2}\right)$$
(219)

for all $t \in \mathbb{R}_{>0}, x, x' \in \mathcal{X}$.

Lemma B.13. Let ϵ_i be independent, centered, sub-Gaussian random variables with variance parameters $v_i \in \mathbb{R}_{>0}$. Let (\mathcal{X}, ρ) be a metric space. Let $b_i : \mathcal{X} \to \mathbb{R}$. Define $Z_x := \sum_{i=1}^n b_i(x)\epsilon_i$. Then Z_x is sub-Gaussian with variance parameter $v_x := \sum_{i=1}^n b_i(x)^2 v_i$. Furthermore, $(Z_x)_{x \in \mathcal{X}}$ is sub-Gaussian with $\rho(x, x')^2 := \sum_{i=1}^n (b_i(x) - b_i(x'))^2 v_i$.

Proof. The results stem from the well-known fact that a linear combination with non-negative coefficients of independent sub-Gaussian variables is again sub-Gaussian. It can be checked using the following equivalent condition for sub-Gaussian: A random variable X is sub-Gaussian with variance parameter v if and only if $\mathbf{E}[\exp(\lambda X)] \leq \exp\left(\frac{\lambda^2 v}{2}\right)$ for all $\lambda \in \mathbb{R}_{\geq 0}$.

Definition B.14 (Covering Number). Let (\mathcal{X}, ρ) be a metric space, $A \subseteq \mathcal{X}$, and $r \in \mathbb{R}_{>0}$. The covering number $N(A, \rho, r)$ with balls $B^{\rho}(x, r) := \{z \in \mathcal{X} \mid \rho(x, z) \leq r\}$ of radius r is defined as

$$N(A,\rho,r) := \min\left\{ n \in \mathbb{N} \middle| \exists (x_i)_{i \in \llbracket n \rrbracket} \subseteq \mathcal{X} \colon A \subseteq \bigcup_{i \in \llbracket n \rrbracket} B^{\rho}(x_i,r) \right\}.$$
 (220)

Lemma B.15. Let $d \in \mathbb{N}$, $a, h, T \in \mathbb{R}_{>0}$. Define the metric ρ in \mathbb{R}^d as

$$\rho(x, x') := a \min\left(1, \frac{\|x - x'\|_2}{h}\right).$$
(221)

Then

$$\int_0^\infty \sqrt{\log(N([0,T]^d,\rho,r))} \mathrm{d}r \le c_d a \max\left(1,\sqrt{\log\left(\frac{T}{h}\right)}\right) \,. \tag{222}$$

Proof. First consider a = 1. For $r \leq 1$,

$$N([0,T]^d,\rho,r) \le c_d T^d h^{-d} r^{-d} \,. \tag{223}$$

For $r \geq 1$,

$$N([0,T]^d,\rho,r) = 1.$$
(224)

Thus,

$$\int_{0}^{\infty} \sqrt{\log(N([0,T]^{d},\rho,r))} \mathrm{d}r \le \int_{0}^{1} \sqrt{\log(c_{d}T^{d}h^{-d}r^{-d})} \mathrm{d}r$$
(225)

$$\leq c_d + \sqrt{d\log(T/h)} + \sqrt{d} \int_0^1 \sqrt{\log(1/r)} \mathrm{d}r \,. \tag{226}$$

We have

$$\int_{0}^{1} \sqrt{\log(1/r)} dr = \int_{0}^{\infty} \sqrt{t} \exp(-t) dt = \Gamma\left(\frac{3}{2}\right) = \frac{1}{2}\sqrt{\pi},$$
(227)

where $\Gamma(\alpha)$ is the Gamma-function. All in all, we obtain

$$\int_0^\infty \sqrt{\log(N([0,T]^d,\rho,r))} \mathrm{d}r = c_d \max\left(1,\sqrt{\log(T/h)}\right).$$
(228)

For arbitrary $a \in \mathbb{R}_{>0}$, and any metric space $(\mathcal{X}, \tilde{\rho})$, we note that

$$\int_0^\infty \sqrt{\log(N(\mathcal{X}, a\tilde{\rho}, r))} dr = \int_0^\infty \sqrt{\log(N(\mathcal{X}, \tilde{\rho}, r/a))} dr = a \int_0^\infty \sqrt{\log(N(\mathcal{X}, \tilde{\rho}, s))} ds \,. \tag{229}$$

Lemma B.16 (Chaining). Let (\mathcal{X}, ρ) be a metric space. Let $(Z_x)_{x \in \mathcal{X}}$ be a centered, sub-Gaussian process. Let $x_0 \in \mathcal{X}$. Let $k \in \mathbb{N}$. Then

$$\mathbf{E}\left[\sup_{x\in\mathcal{X}}\left|Z_{x}-Z_{x_{0}}\right|^{k}\right] \leq c_{k}\left(\int_{0}^{\infty}\sqrt{\log(N(\mathcal{X},\rho,r))}\mathrm{d}r\right)^{k}.$$
(230)

Proof. E.g., [Tal21, chapter 2]. To be more precise, combine the results and definitions of Theorem 2.7.13, Definition 2.7.3, Exercise 2.7.6 (b), (2.40), (2.37), and Exercise 2.3.8 (b) in that reference. \Box

Proof of Theorem B.6. Bias. As

$$\mathbf{E}\Big[|\hat{g} - D_{\mathbf{v}}f|_{\infty}^{2}\Big] \le c \left(\mathbf{E}\left[\sup_{x \in [0,T]^{d}} |\hat{g}(x) - \mathbf{E}[\hat{g}(x)]|^{2}\right] + \sup_{x \in [0,T]^{d}} |\mathbf{E}[\hat{g}(x)] - D_{\mathbf{v}}f(x)|^{2}\right)$$
(231)

we can use the same bound of the bias term as in Theorem B.3, i.e., (213),

$$\sup_{x \in [0,T]^d} |\mathbf{E}[\hat{g}(x)] - D_{\mathbf{v}}f(x)| \le c_\ell C_{\mathsf{egv}} C_{\mathsf{ker}} C_{\mathsf{cvr}} Lh^{\beta-s} \,.$$
(232)

Variance. For the variance term, we have to bound the expectation of the supremum of an empirical process,

$$\mathbf{E}\left[\sup_{x\in[0,T]^d} \left|\hat{g}(x) - \mathbf{E}[\hat{g}(x)]\right|^2\right] = \mathbf{E}\left[\sup_{x\in[0,T]^d} \left|\sum_{i=1}^n w_i(x)\epsilon_i\right|^2\right].$$
(233)

Write

$$w_i(x) = \eta^{\mathsf{T}} B(x)^{-1} b_i(x)$$
(234)

with

$$b_i(x) := \psi\left(\frac{x_i - x}{h}\right) K\left(\frac{\|x_i - x\|_2}{h}\right) , \qquad (235)$$

$$\eta := \frac{T^a}{nh^{d+s}} D_{\mathbf{v}} \psi(0) \,. \tag{236}$$

Then

$$\left|\sum_{i=1}^{n} w_i(x)\epsilon_i\right| \le \left|\eta^{\mathsf{T}} B(x)^{-1} \sum_{i=1}^{n} b_i(x)\epsilon_i\right|$$
(237)

$$\leq \|\eta\| \left\| B(x)^{-1} \right\|_{\text{op}} \left\| \sum_{i=1}^{n} b_i(x) \epsilon_i \right\|_2$$
(238)

$$\leq c_{\ell} \frac{T^d}{nh^{d+s}} C_{\mathsf{egv}} \left\| \sum_{i=1}^n b_i(x) \epsilon_i \right\|_2.$$
(239)

Hence, we have to bound the expectation of

$$\sup_{x \in [0,T]^d} \left\| \sum_{i=1}^n b_i(x)\epsilon_i \right\|_2 \le \sum_{j=1}^N \sup_{x \in [0,T]^d} \left| \sum_{i=1}^n e_j^{\mathsf{T}} b_i(x)\epsilon_i \right| \,. \tag{240}$$

Fix a $j \in [\![N]\!]$ and define $Z_i(x) = e_j^{\mathsf{T}} b_i(x) \epsilon_i$. We want to find an upper bound for

$$\mathbf{E}\left[\sup_{x\in[0,T]^d}\left|\sum_{i=1}^n Z_i(x)\right|^2\right].$$
(241)

Because of SUBGAUSSIAN and Lemma B.13, $Z(x) = \sum_{i=1}^{n} Z_i(x)$ is a sub-Gaussian process on $[0,T]^d$ with metric

$$\rho(x, x')^2 := \sigma^2 \sum_{i=1}^n \left(e_j^\top (b_i(x) - b_i(x')) \right)^2 \,. \tag{242}$$

As $z \mapsto \psi(z)K(z)$ is smooth, it is Lipschitz-continuous with a constant $\tilde{L} \in \mathbb{R}_{>0}$ depending only on ℓ, d, K . If $\max(\|x_i - x\|_2, \|x_i - x'\|_2) \leq h$, then

$$\left| e_{j}^{\mathsf{T}}(b_{i}(x) - b_{i}(x')) \right| \leq c\tilde{L}h^{-1} \left\| x - x' \right\|_{2} .$$
(243)

If $\min(\|x_i - x\|_2, \|x_i - x'\|_2) \le h \le \max(\|x_i - x\|_2, \|x_i - x'\|_2)$, then

$$\left|e_j^{\mathsf{T}}(b_i(x) - b_i(x'))\right| \le c\tilde{L}.$$
(244)

If $\min(||x_i - x||_2, ||x_i - x'||_2) \ge h$, then

$$\left| e_{j}^{\top}(b_{i}(x) - b_{i}(x')) \right| = 0.$$
 (245)

Hence, counting the number of indices with x_i in a ball of radius h yields

$$\rho(x, x') \le c\sqrt{C_{\mathsf{cvr}} n h^d T^{-d}} \tilde{L}\sigma \min\left(1, \frac{\|x - \tilde{x}\|_2}{h}\right).$$
(246)

Now Lemma B.16 and Lemma B.15 yield

$$\mathbf{E}\left[\sup_{x\in[0,T]^d} \left(Z(x) - Z(x_0)\right)^2\right] \le c_{d,\ell,K} C_{\mathsf{cvr}} n h^d T^{-d} \sigma^2 \log(T/h)$$
(247)

for an arbitrary $x_0 \in [0,T]^d$. Using independence of ε_i , we have

$$\mathbf{E}\left[Z(x_0)^2\right] \le \sum_{i=1}^n (e_j^{\mathsf{T}} b_i(x))^2 \mathbf{E}\left[\epsilon_i^2\right] \le c_{d,\ell,K} C_{\mathsf{cvr}} n h^d T^{-d} \sigma^2 \,.$$
(248)

Hence, with $\log(T/h) \ge c$ as $T \ge 2h$, we get

$$\mathbf{E}\left[\sup_{x\in[0,T]^d} Z(x)^2\right] \le c_{d,\ell,K} C_{\mathsf{cvr}} n h^d T^{-d} \sigma^2 \log(T/h) \,.$$
(249)

We finally obtain a bound on the variance term (233),

$$\mathbf{E}\left[\sup_{x\in[0,T]^d}\left|\sum_{i=1}^n w_i(x)\epsilon_i\right|^2\right] \le c_{d,\ell} \left(\frac{T^d}{nh^{d+s}}C_{\mathsf{egv}}\right)^2 \mathbf{E}\left[\sup_{x\in[0,T]^d}\left|\sum_{i=1}^n Z_i(x)\right|^2\right]$$
(250)

$$\leq c_{d,\ell,K} \left(\frac{T^d}{nh^{d+s}} C_{\mathsf{egv}}\right)^2 C_{\mathsf{cvr}} nh^d T^{-d} \sigma^2 \log(T/h)$$
(251)

$$\leq c_{d,\ell,K} C_{\mathsf{cvr}} C_{\mathsf{egv}}^2 \frac{\sigma^2 T^d}{nh^{d+2s}} \log(T/h) \,. \tag{252}$$

Together. Combining the bias and variance term yields

$$\mathbf{E}\left[\left|\hat{g} - D_{\mathbf{v}}f\right|_{\infty}^{2}\right] \le c_{d,\ell,K} \left(\left(C_{\mathsf{egv}}C_{\mathsf{cvr}}Lh^{\beta-s}\right)^{2} + C_{\mathsf{cvr}}C_{\mathsf{egv}}^{2} \frac{\sigma^{2}T^{d}}{nh^{d+2s}}\log(T/h)^{2} \right).$$
(253)

Proof of Proposition B.9. STRICTKERNEL directly implies KERNEL with C_{ker} .

For COVER, we note that $B^d(z,r)$ is a subset of a hypercube with side length 2r. For each dimension, this hypercube can only contain $2n_0r/T + 1$ points separated by a distance T/n_0 . Hence

$$\frac{1}{n} \sum_{i=1}^{n} \mathbb{1}_{B^{d}(x,r)}(x_{i}) \leq \begin{cases} \frac{1}{n} & \text{if } 2n_{0}r/T < 1, \\ \frac{1}{n} \left(2n_{0}r/T + 1\right)^{d} & \text{otherwise}. \end{cases}$$
(254)

If $2n_0r/T \ge 1$,

$$\frac{1}{n} \left(2n_0 r/T + 1\right)^d \le \frac{1}{n} \left(4n_0 r/T\right)^d \le 4^d \left(r/T\right)^d \,. \tag{255}$$

Hence, we can set $C_{\mathsf{cvr}} = 4^d$.

Now consider EIGENVALUE. Let $v \in \mathbb{D}_d$ and $x_0 \in [0,T]^d$. We want to find a lower bound on $v^{\top}B(x_0)v$. Let $h \in \mathbb{R}_{>0}$ be the bandwidth. Denote $z(x) = (x - x_0)/h$ and $z_i := z(x_i)$. Denote the square with side length 2h around x_0 as $\mathcal{S}(x_0) := \mathcal{S}_h(x_0) := \mathbf{X}_{k=1}^d[\Pi_k x_0 - h, \Pi_k x_0 + h]$. Denote indices of the grid points in $\mathcal{S}(x_0)$ as $\mathcal{I}(x_0) := \{i \in [n] \mid x_i \in \mathcal{S}(x_0)\}$. From the definition of $B(x_0)$, (175), we obtain

$$v^{\mathsf{T}}B(x_0)v = \frac{T^d}{nh^d} \sum_{i=1}^n \left(v^{\mathsf{T}}\psi(z_i)\right)^2 K(\|z_i\|_2)$$
(256)

$$= \frac{T^d}{nh^d} \sum_{i \in \mathcal{I}(x_0)} \left(v^{\mathsf{T}} \psi(z_i) \right)^2 K(\|z_i\|_2) \,.$$
(257)

The x_i form a grid of side length T/n_0 in $\mathcal{S}(x_0)$, which has a volume of $(2h)^d$. Assume $2h \ge T/n_0$. Then, the Riemann sum approximation for an *L*-Lipschitz continuous function $g: \mathcal{S}(x_0) \to \mathbb{R}$ yields

$$\left|\frac{(2h)^d}{\#\mathcal{I}(x_0)}\sum_{i\in\mathcal{I}(x_0)}g(x_i) - \int_{\mathcal{S}(x_0)}g(x)\mathrm{d}x\right| \le \sum_{i\in\mathcal{I}(x_0)}\int_{\mathcal{X}_i(x_0)}|g(x_i) - g(x)|\,\mathrm{d}x\tag{258}$$

$$\leq c_d L \frac{T}{n_0} h^d \,, \tag{259}$$

where $\mathcal{X}_i(x_0)$, $i \in \mathcal{I}(x_0)$ form a suitable partition of $\mathcal{S}(x_0)$ with $x_i \in \mathcal{X}_i(x_0)$ and $\operatorname{diam}(\mathcal{X}_i(x_0)) \leq c_d T/n_0$. Applying this to $g_h(x) := \left(v^{\mathsf{T}} \psi(z(x))\right)^2 K(||z(x)||_2)$, we obtain

$$\left| \frac{nh^d(2h)^d}{T^d \# \mathcal{I}(x_0)} v^{\mathsf{T}} B(x_0) v - \int_{\mathcal{S}(x_0)} g_h(x) \mathrm{d}x \right| \le c_d \frac{T}{n_0} h^d \sup_{x \in [0,T]^d} \|Dg_h(x)\|_{\mathsf{op}} .$$
(260)

As $g_h(x) = g_1(x/h)$, with Lipschitz-continuity of K by STRICTKERNEL,

$$\sup_{x \in \mathcal{S}_h(x_0)} \|Dg_h(x)\|_{\mathsf{op}} = h^{-1} \sup_{x \in \mathcal{S}_1(x_0)} \|Dg_1(x)\|_{\mathsf{op}} \le c_{\ell,K} h^{-1}.$$
 (261)

Note that $\lfloor (2hn_0/T) \rfloor^d \leq \# \mathcal{I}(x_0) (\lfloor (2hn_0/T) \rfloor + 1)^d$. Thus,

$$\left| v^{\mathsf{T}} B(x_0) v - h^{-d} \int_{\mathcal{S}(x_0)} g_h(x) \mathrm{d}x \right| \le c_{d,\ell,K} \frac{T}{n_0 h} \,. \tag{262}$$

This means, if $T/(n_0h)$ is small enough, it only remains to show a positive lower bound on

$$h^{-d} \int_{\mathcal{S}(x_0)} g_h(x) dx = \int_{\mathbb{R}^d} \left(v^{\mathsf{T}} \psi(x) \right)^2 K(\|x\|_2) dx$$
(263)

With the same arguments as in the proof of [Tsy09, Lemma 1.4] and using STRICTKERNEL, one can show

$$\int_{\mathbb{R}^d} \left(v^{\mathsf{T}} \psi(x) \right)^2 K(\|x\|_2) \mathrm{d}x \ge c_{d,\ell,K} > 0.$$
(264)

C Polynomial Interpolation

C.1 Univariate Polynomial Interpolation

Proposition C.1. Let $\ell \in \mathbb{N}_0$. Let $u \in \mathcal{D}^{\ell+1}(\mathbb{R}, \mathbb{R})$. Let $(t_i)_{i \in [0,\ell]}$ with $t_0 < \cdots < t_\ell$. Let $p \in \mathcal{P}_{1,\ell}$ be the (Lagrange) interpolation polynomial of $(t_i, u(t_i))_{i \in [0,\ell]}$. Let $k \in [0,\ell]$. Then,

$$\sup_{t \in [t_0, t_\ell]} \left| D^k u(t) - D^k p(t) \right| \le \frac{1}{k! (\ell + 1 - k)!} \sup_{t \in [t_0, t_\ell]} \left| D^k \omega(t) \right| \sup_{t \in [t_0, t_\ell]} \left| D^{\ell + 1} u(t) \right| \,, \tag{265}$$

where $\omega(t) := \prod_{i=0}^{\ell} (t_i - t).$

Proof. [How91, Theorem 2 and 3].

In following lemma, we apply this result to the uniform grid $t_i = i\Delta t$.

Lemma C.2. Let $\ell \in \mathbb{N}_0$. Let $\Delta t \in \mathbb{R}_{>0}$. Let $u \in \mathcal{D}^{\ell+1}(\mathbb{R},\mathbb{R})$. Let $p \in \mathcal{P}_{1,\ell}$ be the interpolation polynomial of $(i\Delta t, u(i\Delta t))_{i\in[0,\ell]}$. Let $k\in[0,\ell]$. Then

$$\sup_{t \in [0,\ell\Delta t]} \left| D^{k} u(t) - D^{k} p(t) \right| \le c_{k,\ell} \Delta t^{\ell+1-k} \left| D^{\ell+1} u \right|_{\infty} .$$
(266)

C.2 Univariate Polynomial Comparison

Lemma C.3. Let $p_n, \tilde{p}_n \in \mathcal{P}_{1,\ell-1}$. Let $h_n, \delta_n \in \mathbb{R}_{>0}$. Let $c_1, \ldots, c_\ell \in \mathbb{R}$ be distinct. Let $c_{\min}, c_{\max} \in \mathbb{R}_{>0}$. Assume, for all $i = 1, \ldots, \ell$,

$$|p_n(c_ih_n) - \tilde{p}_n(c_ih_n)| \preccurlyeq \delta_n \,. \tag{267}$$

Then

$$\sup_{t \in [-c_{\min}h_n, c_{\max}h_n]} \left| D^k p_n(t) - D^k \tilde{p}_n(t) \right| \preccurlyeq h_n^{-k} \delta_n \tag{268}$$

for $k = 0, ..., \ell - 1$.

Proof. Let $p_n(t) - \tilde{p}_n(t) =: q(t) = \sum_{k=1}^{\ell} a_k t^{k-1}$. Let $y_i := q(c_i h_n)$. Then $a = V^{-1}y$, where $V \in \mathbb{R}^{\ell \times \ell}$ is the Vandermonde matrix $V_{i,j} = (c_i h_n)^{j-1}$. By the inverse formula of the Vandermonde matrix, $|(V^{-1})_{i,j}| \preccurlyeq h_n^{1-i}$. Thus,

$$|a_i| = \left| \sum_{j=1}^{\ell} (V^{-1})_{i,j} y_j \right| \preccurlyeq h_n^{1-i} \delta_n \,.$$
(269)

Furthermore, for $c \in [-c_{\min}, c_{\max}]$,

$$\left|D^{k}q(ch_{n})\right| = \left|\sum_{i=1}^{\ell-k} a_{i+k} (ch_{n})^{i-1} \prod_{s=1}^{k} (i+s)\right|$$
(270)

$$\preccurlyeq \sum_{i=1}^{c-\kappa} \left| c \right|^{i-1} h_n^{-k} \delta_n \tag{271}$$

$$\preccurlyeq h_n^{-k} \delta_n \,. \tag{272}$$

C.3 Multivariate Polynomial Interpolation

Recall Definition 4.11 for the objects $\psi(x)$, $\Psi(\mathbf{x})$, and $I(\mathbf{x}, \mathbf{y}, x)$.

Lemma C.4 (Linear Transformation). Let $\ell, d \in \mathbb{N}$. Set $N := \dim(\mathcal{P}_{d,\ell})$. Let $b \in \mathbb{R}^d$. Let $B \in \mathbb{R}^{d \times d}$. Then there is $A \in \mathbb{R}^{N \times N}$ such that $\psi(Bx + b) = A\psi(x)$ for all $x \in \mathbb{R}^d$. Furthermore, if B is invertible, then so is A.

Proof. First we need to define $\tilde{\psi}_{\ell} : \mathbb{R}^d \to \mathbb{R}^M$ as a variant of the monomial transformation $\psi_{\ell} : \mathbb{R}^d \to \mathbb{R}^N$, where commutativity is not applied, i.e., $\tilde{\psi}_{\ell}(x)$ contains a dimension for x_1x_2 and a different one for x_2x_1 . The monomial transform without commutativity $\tilde{\psi}_{\ell}$ can be expressed using the Kronecker product, see, e.g., [HJ94, Chapter 4].

Denote by \otimes the operator for the Kronecker product. For $x \in \mathbb{R}^d$, $k \in \mathbb{N}_0$, define $x^{\otimes k} \in \mathbb{R}^{d^k}$ as follows: Let $x^{\otimes 0} := 1$. For $k \in \mathbb{N}$, define $x^{\otimes k} = x^{\otimes (k-1)} \otimes x$. Denote

$$M := \sum_{k=0}^{\ell} d^k = \frac{d^{\ell+1} - 1}{d - 1} \,. \tag{273}$$

Define the monomial transform without commutativity $\tilde{\psi}_{\ell} \colon \mathbb{R}^d \to \mathbb{R}^M$ as

$$\tilde{\psi}_{\ell}(x) := \begin{pmatrix} x^{\otimes 0} \\ x^{\otimes 1} \\ \vdots \\ x^{\otimes \ell} \end{pmatrix}.$$
(274)

Our intermediate goal is to find a matrix $\tilde{A} \in \mathbb{R}^{M \times M}$ with the property

$$\tilde{\psi}(Bx+b) = \tilde{A}\tilde{\psi}(x).$$
(275)

For this, we first recall some properties of the Kronecker product: The Kronecker product is bilinear. In particular,

$$M_1 \otimes (M_2 + M_3) = M_1 \otimes M_2 + M_1 \otimes M_3 \text{ and } (M_1 + M_2) \otimes M_3 = M_1 \otimes M_3 + M_2 \otimes M_3 \quad (276)$$

for compatible matrices M_1, \ldots, M_3 . Moreover, The mixed-product property of the Kronecker product states

$$(M_1 \otimes M_2) (M_3 \otimes M_4) = (M_1 M_3) \otimes (M_2 M_4)$$
(277)

for compatible matrices M_1, \ldots, M_4 . The Kronecker product is not commutative, but $M_1 \otimes M_2$ and $M_2 \otimes M_1$ are *permutation equivalent*: There are permutation matrices P, Q such that

$$M_1 \otimes M_2 = P(M_2 \otimes M_1)Q \tag{278}$$

for compatible matrices M_1, M_2 .

Now, using the mixed-product property, we obtain by induction

$$(Bx)^{\otimes k} = B^{\otimes k} x^{\otimes k} . (279)$$

Next, the bilinearity and the permutation equivalence imply, also using induction, there are matrices $\bar{A}_j \in \mathbb{R}^{d^k \times d^j}$

$$(b+x)^{\otimes k} = \sum_{j=0}^{k} \bar{A}_j x^{\otimes k} .$$
 (280)

Thus there is are matrices $\tilde{A}_0, \tilde{A}_1 \in \mathbb{R}^{M \times M}$ such that

$$\tilde{\psi}(x+b) = \tilde{A}_0 \tilde{\psi}(x)$$
 and $\tilde{\psi}(Bx) = \tilde{A}_1 \tilde{\psi}(x)$ (281)

and, hence,

$$\tilde{\psi}(Bx+b) = \tilde{A}\tilde{\psi}(x) \tag{282}$$

with $\tilde{A} = \tilde{A}_0 \tilde{A}_1$. From (279), we can see that \tilde{A}_1 is block diagonal with blocks $B^{\otimes k}$. From (280), we can see that \tilde{A}_0 is block triangular with identity blocks $I_{d^k} \in \mathbb{R}^{d^k \times d^k}$ on the diagonal. Hence, \tilde{A} is block triangular with blocks $B^{\otimes k}$ on the diagonal. For the determinant of the Kronecker product, we have

$$\det(M_1 \otimes M_2) = \det(M_1)^{d_2} \det(M_2)^{d_1}$$
(283)

for $M_1 \in \mathbb{R}^{d_1 \times d_1}$ and $M_2 \in \mathbb{R}^{d_2 \times d_2}$. Thus,

$$\det\left(\tilde{A}\right) = \prod_{k=0}^{\ell} \det\left(B^{\otimes k}\right) = \det(B)^{\kappa}, \qquad (284)$$

where

$$\kappa := \sum_{k=0}^{\ell} k d^{k-1} = \begin{cases} \frac{\ell(1+\ell)}{2} & \text{if } d = 1\\ \frac{1+\ell d^{\ell+1} - (\ell+1)d^{\ell}}{(d-1)^2} & \text{otherwise.} \end{cases}$$
(285)

To see this, show inductively using (283) that

$$\det(B^{\otimes k}) := \det(B)^{kd^{k-1}}.$$
(286)

Now, we have the existence of $\tilde{A} \in \mathbb{R}^{M \times M}$ with

$$\tilde{\psi}(Bx+b) = \tilde{A}\tilde{\psi}(x) \tag{287}$$

and the property that if B is invertible so is \tilde{A} . In the final step, we covert from $\tilde{\psi}$ to ψ via a linear transformation of full rank: Let $Z \in \mathbb{R}^{N \times M}, \tilde{Z} \in \mathbb{R}^{M \times N}$ be linear mappings that fulfill

$$\psi_{\ell}(x) = Z \tilde{\psi}_{\ell}(x) \quad \text{and} \quad \tilde{\psi}_{\ell}(x) = \tilde{Z} \psi_{\ell}(x)$$
(288)

for all $x \in \mathbb{R}^d$. Note that the rank of Z and of \tilde{Z} is N, i.e., both matrices have full rank. Set

$$A := Z\tilde{A}\tilde{Z} \,. \tag{289}$$

Then

$$\psi(Bx+b) = Z\tilde{\psi}(Bx+b) = Z\tilde{A}\tilde{\psi}(x) = Z\tilde{A}\tilde{Z}\psi(x) = A\psi(x).$$
⁽²⁹⁰⁾

Moreover, if B is invertible, so is \tilde{A} . Then, as \tilde{A} , \tilde{Z} , and Z have full rank, A also has full rank and is invertible.

Lemma C.5 (Linear Transformation). Let $\ell, d_x, d_y \in \mathbb{N}$. Set $N := \dim(\mathcal{P}_{d_x,\ell})$. Let $b \in \mathbb{R}^{d_x}$. Let $B \in \mathbb{R}^{d_x \times d_x}$. Let $\mathbf{x} \in (\mathbb{R}^{d_x})^N$ and $\mathbf{y} \in (\mathbb{R}^{d_y})^N$. Let $x \in \mathbb{R}^{d_x}$. Assume B and $\Psi(\mathbf{x})$ are invertible. Then

(i) $\Psi(B\mathbf{x}+b)$ is invertible,

(ii)

$$I(\mathbf{x}, \mathbf{y}, x) = I(B\mathbf{x} + b, \mathbf{y}, Bx + b), \qquad (291)$$

(iii)

$$\psi(x)^{\dagger}\Psi(\mathbf{x})^{-1} = \psi(Bx+b)^{\dagger}\Psi(B\mathbf{x}+b)^{-1}.$$
(292)

Proof. Part (i) follows directly from Lemma C.4.

x

As the interpolation is done componentwise, it is enough to show the results for the case $d_y = 1$. Let us assume $d_y = 1$. The function $g: \mathbb{R}^{d_x} \to \mathbb{R}, x \mapsto I(\mathbf{x}, \mathbf{y}, x)$ is the unique interpolation polynomial of (\mathbf{x}, \mathbf{y}) with degree at most ℓ . The mapping $\tilde{g}: \mathbb{R}^d \to \mathbb{R}, x \mapsto I(B\mathbf{x}+b, \mathbf{y}, Bx+b)$ is a polynomial of degree at most ℓ and fulfills $\tilde{g}(x_k) = y_k$ for $k \in [N]$. Thus $\tilde{g} = g$. This shows (ii).

As (ii) is true for all **y** and $I(\mathbf{x}, \mathbf{y}, x) = \psi(x)^{\top} \Psi(\mathbf{x})^{-1} \mathbf{y}$, (iii) follows.

Lemma C.6 (Approximation). Let $\ell, d \in \mathbb{N}$. Set $N := \dim(\mathcal{P}_{d,\ell})$. Let $\mathbf{x} \in (\mathbb{R}^d)^N$. Let $g: \mathbb{R}^d \to \mathbb{R}$ be $(\ell + 1)$ -times continuously differentiable and set $L := \sup_{x \in \mathbb{R}^d} \|D^{\ell+1}g(x)\|_{op}$. Let $\mathbf{y} = (g(x_1), \ldots, g(x_N))$. Then

$$\sup_{\mathbf{x}\in\mathring{ch}(\mathbf{x})} \|g(x) - I_{\ell}(\mathbf{x}, \mathbf{y}, x)\|_{2} \le LC_{\ell}(\mathbf{x})\operatorname{diam}(\mathbf{x})^{\ell+1},$$
(293)

where $C_{\ell}(\mathbf{x})$ is defined as follows: For $k \in [N]$, let $e_k \in \mathbb{R}^N$ be the k-th standard base vector. Define

$$C_{\ell}(\mathbf{x}) := \begin{cases} \frac{1}{(\ell+1)!} \sum_{k=1}^{N} \sup_{x \in \mathring{ch}(\mathbf{x})} |I(\mathbf{x}, e_k, x)| & \text{if } \Psi(\mathbf{x}) \text{ is invertible,} \\ \infty & \text{otherwise.} \end{cases}$$
(294)

Proof. [CR72, Theorem 2].

Lemma C.7. Let $\ell, d \in \mathbb{N}$. Set $N := \dim(\mathcal{P}_{d,\ell})$. If $\Psi(\mathbf{x})$ is invertible,

$$C_{\ell}(\mathbf{x}) \le \frac{N^{\frac{3}{2}}}{(\ell+1)!} \left\| \Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1} \right\|_{\mathsf{op}} .$$
(295)

Proof. By Lemma C.5, we have

$$\sup_{x \in \mathring{ch}(\mathbf{x})} |I(\mathbf{x}, e_j, x)| = \sup_{x \in \mathring{ch}(\mathbf{x})} |I(\eta_{\mathbf{x}}(\mathbf{x}), e_j, \eta_{\mathbf{x}}(x))| \le \sup_{x \in B^d(0, 1)} |I(\eta_{\mathbf{x}}(\mathbf{x}), e_j, x)| .$$
(296)

For $x \in B^d(0, 1)$,

$$|I(\eta_{\mathbf{x}}(\mathbf{x}), e_j, x)| \le \|\psi(x)\|_2 \left\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\right\|_{\mathsf{op}} \|e_j\|_2 \le \sqrt{N} \left\|\Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1}\right\|_{\mathsf{op}} .$$
 (297)

Thus,

$$C_{\ell}(\mathbf{x}) \le \frac{1}{(\ell+1)!} \sum_{j=1}^{N} \sqrt{N} \left\| \Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1} \right\|_{\mathsf{op}}$$
(298)

$$\leq \frac{N^{\frac{3}{2}}}{(\ell+1)!} \left\| \Psi(\eta_{\mathbf{x}}(\mathbf{x}))^{-1} \right\|_{\mathsf{op}} \,. \tag{299}$$

D Ordinary Differential Equations and Smoothness

v

D.1 Difference Bound

Lemma D.1. Let $f : \mathbb{R} \to \mathbb{R}$ be Lipschitz continuous. Let $x \in \mathbb{R}$. Set $u := U(f, x, \cdot)$. Let $v : \mathbb{R} \to \mathbb{R}$ be differentiable with $v(0) \le x$ and $\dot{v}(t) \le f(v(t))$ for all $t \in \mathbb{R}_{\ge 0}$. Then

$$(t) \le u(t) \tag{300}$$

for all $t \in \mathbb{R}_{>0}$.

Proof. Assume there is $t_1 \in \mathbb{R}_{>0}$ with $v(t_1) > u(t_1)$. Then the first intersection time $t_0 := \inf\{t \in \mathbb{R}_{\geq 0} \mid v(t) > u(t)\}$ exists. As v and u are continuous, we have $v(t_0) = u(t_0)$. Let $t_2 = \sup\{t \in \mathbb{R}_{\geq t_0} \mid \forall s \in [t_0, t]: v(s) \geq u(s)\}$. Because of the existence of t_1 and continuity of u and v, we have $t_2 > t_0$. Let $L \in \mathbb{R}_{>0}$ be the Lipschitz constant of f. Set $\delta(t) := v(t) - u(t)$. Let $t \in [t_0, t_2]$. Then

$$\delta(t) = \int_{t_0}^t \dot{\delta}(s) \mathrm{d}s \tag{301}$$

$$\leq \int_{t_0}^t f(v(s)) - f(u(s)) \mathrm{d}s \tag{302}$$

$$\leq L \int_{t_0}^t \delta(s) \mathrm{d}s \,. \tag{303}$$

As δ is continuous, there is $t_{\max} \in \arg \max_{t \in [t_0, \min(t_0 + 1/L, t_2)]} \delta(t)$. By definition of t_0 and t_{\max} , we have $\delta(t_{\max}) > 0$. Thus, using $\delta(t_0) = 0$, the continuity of δ , and $t_{\max} - t_0 \leq 1/L$, we obtain

$$\delta(t_{\max}) \le L \int_{t_0}^{t_{\max}} \delta(s) \mathrm{d}s < L \left(t_{\max} - t_0 \right) \delta(t_{\max}) \le \delta(t_{\max}) \,, \tag{304}$$

a contradiction.

Lemma D.2. Let $d \in \mathbb{N}$, $L \in \mathbb{R}_{>0}$. Let $f, g \in \Sigma^{d \to d}(1, L)$. Let $x_0 \in \mathbb{R}^d$. Denote $u := U(f, x_0, \cdot), v := U(g, x_0, \cdot)$. Then, for $t \in \mathbb{R}_{\geq 0}$,

$$\|u(t) - v(t)\|_{2} \le \frac{\exp(Lt) - 1}{L} |f - g|_{\infty} .$$
(305)

In particular, if $t \in [0, \frac{1}{L}]$, then

$$||u(t) - v(t)||_2 \le 2t |f - g|_{\infty} .$$
(306)

Proof. Set $C := |f - g|_{\infty}$. We represent the right-hand side of (305) with the solution z of an ODE: Define $z : \mathbb{R}_{>0} \to \mathbb{R}$ as the solution of the ODE $\dot{z}(t) = Lz(t) + C$ with z(0) = 0. Then,

$$z(t) = \frac{C}{L} \left(\exp(Lt) - 1 \right) \,. \tag{307}$$

Next consider the left-hand side of (305). Define $w(t) := \|u(t) - v(t)\|_2$. Then w(0) = 0 and

$$w(t) = \left\| \int_{0}^{t} f(u(s)) - g(v(s)) \mathrm{d}s \right\|_{2}$$
(308)

$$\leq \left\| \int_{0}^{t} f(u(s)) - f(v(s)) \mathrm{d}s \right\|_{2} + \left\| \int_{0}^{t} f(v(s)) - g(v(s)) \mathrm{d}s \right\|_{2}$$
(309)

$$\leq L \int_0^t w(s) \mathrm{d}s + Ct \,. \tag{310}$$

Thus, Lemma D.1 yields, $w(t) \leq z(t)$.

D.2 Smoothness of Solutions in Time

Let $d, \beta \in \mathbb{N}$. Let $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0}$. Assuming $f \in \overline{\Sigma}^{d \to d}(\beta, L_{\llbracket 0,\beta \rrbracket})$ we want to find $\tilde{L}_{\llbracket 1,\beta+1 \rrbracket} \subseteq \mathbb{R}_{>0}$ such that $U(f, x, \cdot) \in \Sigma^{1 \to d}(\beta+1, \tilde{L}_{\llbracket 0,\beta+1 \rrbracket})$ with $\tilde{L}_0 = \infty$.

As detailed in [But16, Chapter 3], higher order derivatives of u can be expressed using tree structures. To state this result in Theorem D.6 below, we first need to introduce *increasing trees* (Definition D.3) and the derivative operator associated to such a tree (Definition D.5).

In the following the term *finite ordered set* refers to a finite set with strict total order that is denoted as <.

Definition D.3 (Increasing tree). Let S be a finite ordered set.

- (i) An increasing tree \mathcal{T} is a tuple (S, E), where $E \subseteq \{(n_1, n_2) \mid n_1, n_2 \in S, n_1 < n_2\}$ with #E = #S 1. We call S the nodes, E the edges, and min S the root. Furthermore, denote $\#\mathcal{T} := \#S$.
- (ii) Let \mathfrak{T}_S be the set of increasing trees with nodes S.
- (iii) We define $[\mathcal{T}_1, \ldots, \mathcal{T}_k]$ as the increasing tree that connects all trees \mathcal{T}_j at their roots to a new root s:

Let $k \in \mathbb{N}_0$. Let $S_0 = \{s\}$ and S_1, \ldots, S_k be disjoint finite ordered sets so that $S := \bigcup_{j \in [0,k]} S_k$ is also ordered with $s = \min S$. Let $\mathcal{T}_1 = (S_1, E_1), \ldots, \mathcal{T}_k = (S_k, E_k)$ be increasing trees. Define $[\mathcal{T}_1, \ldots, \mathcal{T}_k] := (S, E)$, where $E := \bigcup_{j \in [0,k]} E_j$ and $E_0 := \{(s, \min S_j) \mid j \in [k]\}$.

Lemma D.4. Let S be a finite ordered set. Then

$$\#\mathfrak{T}_S = (\#S - 1)!. \tag{311}$$

Proof. For $S = \{s\}$, we have $\mathfrak{T}_S = \{(\{s\}, \emptyset)\}$. Hence, $\#\mathfrak{T}_S = (\#S - 1)!$ is true. Now assume #S = k + 1, max S = s, and $S' = S \setminus \{s\}$. Then \mathfrak{T}_S can be constructed from $\mathfrak{T}_{S'}$, by adding the node s to the node n of \mathcal{T} for all trees $\mathcal{T} \in \mathfrak{T}_{S'}$ and all nodes $n \in S'$. Hence,

$$\#\mathfrak{T}_{S} = \#S'\#\mathfrak{T}_{S'} = k\,\#\mathfrak{T}_{S'}\,. \tag{312}$$

The desired result now follows by induction.

Definition D.5 (Tree derivative operator). Let S be a finite ordered set. Let $\mathcal{T} \in \mathfrak{T}_S$. Let $k \in \mathbb{N}_0$. Assume $\mathcal{T} = [\mathcal{T}_1, \ldots, \mathcal{T}_k]$. Let $d \in \mathbb{N}$. Let $f \in \mathcal{D}^k(\mathbb{R}^d, \mathbb{R}^d)$. Let $x \in \mathbb{R}^d$. Define the tree derivative of f at x, denoted as $D^{\mathcal{T}}f(x)$, as

$$D^{\mathcal{T}}f(x) := D^{k}f(x)[D^{\mathcal{T}_{1}}f(x),\dots,D^{\mathcal{T}_{k}}f(x)].$$
(313)

Theorem D.6. Let $d, k \in \mathbb{N}$. Let $f \in \mathcal{D}^{k-1}(\mathbb{R}^d, \mathbb{R}^d)$. Let $x \in \mathbb{R}^d$. Assume f is Lipschitz continuous. Set $u := U(f, x, \cdot)$. Then

$$D^{k}u(t) = \sum_{\mathcal{T}\in\mathfrak{T}_{\llbracket k \rrbracket}} D^{\mathcal{T}}f(u(t)).$$
(314)

Proof. [But16, Theorem 311B].

Corollary D.7. Let $d, \beta \in \mathbb{N}$. Let $L_{\llbracket 0,\beta \rrbracket} \subseteq \mathbb{R}_{>0}$. Let $f \in \overline{\Sigma}^{d \to d}(\beta, L_{\llbracket 0,\beta \rrbracket})$. Let $\beta \in \mathbb{N}$. Let

$$x_0 \in \mathbb{R}^d$$
. Set $u := U(f, x_0, \cdot)$. Set $L := \sup_{k=0,...,\beta} L_k$. Then, for all $k \in [\![\beta + 1]\!]$,
 $|D^k u|_{\infty} \le (k-1)! L^k$. (315)

Proof. Let S be a finite ordered set. Let $\mathcal{T} \in \mathfrak{T}_S$. We want to show by induction that

$$\left|D^{\mathcal{T}}f\right|_{\infty} \le L^{\#\mathcal{T}} \,. \tag{316}$$

If $\#\mathcal{T} = 1$, $|D^{\mathcal{T}}f|_{\infty} = |f|_{\infty} \leq L$. Now, assume $\mathcal{T} = [\mathcal{T}_1, \dots, \mathcal{T}_{\kappa}]$ and $|D^{\mathcal{T}_j}f|_{\infty} \leq L^{\#\mathcal{T}_j}$. Then,

$$\left|D^{\mathcal{T}}f\right|_{\infty} = \left|D^{\kappa}f(\cdot)[D^{\mathcal{T}_{1}}f(\cdot),\dots,D^{\mathcal{T}_{\kappa}}f(\cdot)]\right|_{\infty}$$
(317)

$$\leq L \prod_{j \in \llbracket \kappa \rrbracket} \left| D^{\mathcal{T}_j} f \right|_{\infty} \tag{318}$$

$$\leq L^{1+\sum_{j=1}^{\kappa} \#\mathcal{T}_j} \tag{319}$$

$$=L^{\#\mathcal{T}}.$$
(320)

Hence, we have shown (316). Using Theorem D.6, (316), and Lemma D.4, we obtain

$$\left|D^{k}u\right|_{\infty} \leq \sum_{\mathcal{T}\in\mathfrak{T}_{\llbracket k\rrbracket}} \left|D^{\mathcal{T}}f(u(\cdot))\right|_{\infty}$$
(321)

$$\leq \# \mathfrak{T}_{\llbracket k \rrbracket} L^k \tag{322}$$

$$\leq (k-1)!L^k. \tag{323}$$

D.3 Smoothness of Solutions in the Initial Conditions

We state a simple consequence of the basic form of Grönwall's inequality, which we will make use of below.

Lemma D.8 (Grönwall's inequality). Let $d \in \mathbb{N}$. Let $u \colon \mathbb{R} \to \mathbb{R}^d$ be differentiable. Let $L \in \mathbb{R}_{\geq 0}$. Assume $\|\dot{u}(t)\|_2 \leq L \|u(t)\|_2$ for $t \in \mathbb{R}$. (324)

Then, for all $t \in \mathbb{R}$, we have

$$\|u(t)\|_{2} \le \|u(0)\|_{2} \exp(L|t|), \qquad (325)$$

$$\|u(t) - u(0)\|_{2} \le \|u(0)\|_{2} (\exp(L|t|) - 1). \qquad (326)$$

Proof. We assume below that t > 0 and $u(t) \neq 0$ for all $s \in [0, t]$. If t < 0, consider $\tilde{u}(t) = u(-t)$ instead. If there are $s \in [0, t]$ with u(s) = 0, the proof applies to all intervals with [a, b] with $u(\tilde{s}) \neq 0$ for $\tilde{s} \in [a, b]$, which yields an upper bound that is at least as strong as claimed.

Assume t > 0 and $u(t) \neq 0$ for all $s \in [0, t]$. Set $v(t) := ||u(t)||_2$. Then

$$\dot{v}(t) = \frac{u(t)^{\top} \dot{u}(t)}{\|u(t)\|_{2}} \le \|\dot{u}(t)\|_{2} \le L \|u(t)\|_{2} = Lv(t).$$
(327)

Thus, by Grönwall's inequality in its standard form, we have

$$v(t) \le v(0) \exp(Lt), \qquad (328)$$

which is the first inequality that we want to prove. Now, we can calculate

$$\|u(t) - u(0)\|_{2} \le \int_{0}^{t} \|\dot{u}(s)\|_{2} \,\mathrm{d}s \tag{329}$$

$$\leq L \int_0^t v(s) \mathrm{d}s \tag{330}$$

$$\leq L \|u(0)\|_2 \int_0^t \exp(Ls) \mathrm{d}s$$
 (331)

$$= \|u(0)\|_{2} \left(\exp(Lt) - 1\right) \,. \tag{332}$$

Lemma D.9 (Derivative with respect to initial conditions). Let $d \in \mathbb{N}$. Let $L_1 \in \mathbb{R}_{>0}$. Let $f \in \overline{\Sigma}(1, \infty, L_1)$. Then, for all $t \ge 0$,

$$|DU(f,\cdot,t)|_{\infty} \le \exp(L_1 t) \quad \text{and} \quad |D\Upsilon(f,t,\cdot)|_{\infty} \le \exp(L_1 t) - 1.$$
(333)

Proof. Let $v \in \mathbb{D}_d$ be a direction. Set $u(t) := u_x(t) := D_{x,v}U(f, x, t)$ denote the derivative of the solution U(f, x, t) with respect to the initial conditions x. Then, using the chain rule Lemma A.5 (ii),

$$\dot{u}(t) = D_{x,v}\dot{U}(f, x, t) \tag{334}$$

$$= D_{x,v} \left(f(U(f,x,t)) \right)$$
(335)
$$= D_{x,v} \left(f(U(f,x,t)) \right)$$
(336)

$$= Df(U(f,x,t))[D_{x,v}U(f,x,t)]$$
(336)

$$= Df(U(f, x, t))[u(t)].$$
(337)

Furthermore,

$$u(0) = D_{x,v}U(f, x, 0) = v.$$
(338)

We have established the differential inequality

$$\|\dot{u}(t)\|_{2} \le \|(Df)(U(f,x,t))[u(t)]\|_{2} \le L_{1} \|u(t)\|_{2}$$
(339)

with $||u(0)||_2 = 1$. Thus, by Lemma D.8,

$$\|u(t)\|_{2} \le \exp(L_{1}|t|).$$
(340)

Hence,

$$\left|DU(f,\cdot,t)\right|_{\infty} = \sup_{v \in \mathbb{D}_d} \sup_{x \in \mathbb{R}^d} \left\|D_{x,v}U(f,x,t)\right\|_2 \tag{341}$$

$$\leq \exp(L_1 t) \,. \tag{342}$$

Set $u^{\circ}(t) := u(t) - u(0)$. Then, $\dot{u}^{\circ} = \dot{u}$ and $u^{\circ}(0) = 0$. Thus, using Lemma D.8 again,

$$\|u^{\circ}(t)\|_{2} \le \exp(L_{1}t) - 1.$$
(343)

We obtain,

$$|D\Upsilon(f,t,\cdot)|_{\infty} = |DU(f,\cdot,t) - DU(f,\cdot,0)|_{\infty} \le \exp(L_1 t) - 1.$$
(344)

For higher order derivatives of U(f, x, t) with respect to t, we used the tree derivatives in Theorem D.6. For higher order derivatives of U(f, x, t) with respect to x, we introduce partition derivatives.

Definition D.10.

- (i) Let $k \in \mathbb{N}$. Let \mathbb{A}_k be the set of partitions of $[\![k]\!]$.
- (ii) Let $k \in \mathbb{N}$. Let $\mathbf{v} = (v_1, \ldots, v_k) \in \mathbb{D}_d^k$. Define the set of partitions of \mathbf{v} , as

$$\mathbb{A}(\mathbf{v}) := \{ (v_{A_1}, \dots, v_{A_\ell}) | (A_1, \dots, A_\ell) \in \mathbb{A}_k \} .$$

$$(345)$$

(iii) Let $k \in \mathbb{N}$. Let $\mathbf{v} \in \mathbb{D}_d^k$. Let $f, g \in \mathcal{D}^k(\mathbb{R}^d, \mathbb{R}^d)$. Let $\pi = (\mathbf{v}_1, \dots, \mathbf{v}_\ell) \in \mathbb{A}(\mathbf{v})$. Denote the partition derivative of $f \circ g$ as

$$D^{\pi}(f \circ g)(x) := (D^{\ell}f)(g(x))[D_{\mathbf{v}_{1}}g(x), \dots, D_{\mathbf{v}_{\ell}}g(x)].$$
(346)

The following theorem is a version of Faà di Bruno's formula [Cra05].

Theorem D.11. Let $k \in \mathbb{N}$. Let $\mathbf{v} \in \mathbb{D}_d^k$. Let $f \in \mathcal{D}^{k-1}(\mathbb{R}^d, \mathbb{R}^d)$. Assume that f is Lipschitz continuous. Then $D_{x,\mathbf{v}}\dot{U}(f,x,t) = \sum_{\pi \in \mathbb{A}(\mathbf{v})} D^{\pi} \left(f \circ U(f,\cdot,t)\right)(x).$ (347)

Proof. For $\mathbf{v} \in \mathbb{D}_d^k$, denote $u_{\mathbf{v}}(x,t) := D_{x,\mathbf{v}}U(f,x,t)$. We prove the statement by induction over k. See the proof of Lemma D.9 for k = 1. Assume the statement is true for all $\mathbf{v} \in \mathbb{D}_d^k$. Let $v \in \mathbb{D}_d$ and $\mathbf{v}' := (\mathbf{v}, v) \in \mathbb{D}_d^{k+1}$. Then

$$\dot{u}_{\mathbf{v}'}(x,t) = D_{x,v}\dot{u}_{\mathbf{v}}(x,t) \tag{348}$$

$$= \sum_{(\mathbf{v}_1,\dots,\mathbf{v}_\ell)\in\mathbb{A}(\mathbf{v})} D_{x,v} \left(D^\ell f(U(f,x,t))[u_{\mathbf{v}_1}(x,t),\dots,u_{\mathbf{v}_\ell}(x,t)] \right)$$
(349)

By the product rule, Lemma A.5 (i),

$$D_{x,v}\left((D^{\ell}f)(U(f,x,t))[u_{\mathbf{v}_{1}}(x,t),\dots,u_{\mathbf{v}_{\ell}}(x,t)]\right)$$
(350)

$$= D_{x,v} \left(D^{\ell} f(U(f,x,t)) \right) \left[u_{\mathbf{v}_1}(x,t), \dots, u_{\mathbf{v}_{\ell}}(x,t) \right]$$
(351)

+
$$\sum_{j=1}^{\ell} \left((D^{\ell} f)(U(f,x,t))[u_{\mathbf{v}_{1}}(x,t),\ldots,D_{x,v}u_{\mathbf{v}_{j}}(x,t),\ldots,u_{\mathbf{v}_{\ell}}(x,t)] \right)$$
. (352)

By the chain rule, Lemma A.5 (ii),

$$D_{x,v}\left(D^{\ell}f(U(f,x,t))\right)\left[u_{\mathbf{v}_{1}}(x,t),\ldots,u_{\mathbf{v}_{\ell}}(x,t)\right] = D^{\ell+1}f(U(f,x,t))\left[u_{\mathbf{v}_{1}}(x,t),\ldots,u_{\mathbf{v}_{\ell}}(x,t),u_{v}(x,t)\right].$$
(353)

The set of all partitions of \mathbf{v}' can be created from the set of all partitions of \mathbf{v} as follows: Let $\mathbf{v}_1, \ldots, \mathbf{v}_\ell$ be a partition of \mathbf{v} . Then $(v, \mathbf{v}_1, \ldots, \mathbf{v}_\ell)$ and $(\mathbf{v}_1, \ldots, (\mathbf{v}_j, v), \ldots, \mathbf{v}_\ell)$, $j \in \llbracket \ell \rrbracket$ are partitions of \mathbf{v}' (the order of the partition components does not matter). Iterating this construction over all partitions of \mathbf{v} creates all partitions of \mathbf{v}' .

Now putting all previous equations in this proof together, we obtain

$$D_{x,\mathbf{v}'}\dot{U}(f,x,t) = \dot{u}_{\mathbf{v}'}(x,t) \tag{354}$$

$$= \sum_{(\mathbf{v}_1,\dots,\mathbf{v}_\ell)\in\mathbb{A}(\mathbf{v}')} D^\ell f(U(f,x,t))[u_{\mathbf{v}_1}(x,t),\dots,u_{\mathbf{v}_\ell}(x,t)]$$
(355)

$$=\sum_{\pi\in\mathbb{A}(\mathbf{v}')}D^{\pi}\left(f\circ U(f,\cdot,t)\right)(x)\,.$$
(356)

Corollary D.12. Let $\beta \in \mathbb{N}$, $L_{[1,\beta]} \in \mathbb{R}_{>0}$, $k \in [\beta]$, $d \in \mathbb{N}$. Let $f \in \overline{\Sigma}^{d \to d}(\beta, L_{[0,\beta]})$ with $L_0 = \infty$. Let $\Delta t_0 \in \mathbb{R}_{>0}$. Assume $\Delta t \in [0, \Delta t_0]$. Then

$$\left| D^{k} \Upsilon(f, \Delta t, \cdot) \right|_{\infty} \le C_{k} \left(\exp(L_{1} \Delta t) - 1 \right) , \qquad (357)$$

where $C_k \in \mathbb{R}_{>0}$ depends only on k, Δt_0 , and $L_{[1,k]}$.

Proof. Define $C_1 := 1$, $\tilde{C}_1 := \exp(L_1 \Delta t_0)$, and, for $k \in [\![2, \beta]\!]$ with an arbitrary $\mathbf{v} \in \mathbb{D}_d^k$,

$$C_k := \frac{1}{L_1} \sum_{(\mathbf{v}_1, \dots, \mathbf{v}_\ell) \in \mathbb{A}(\mathbf{v}) \setminus \{\mathbf{v}\}} L_\ell \prod_{j=1}^\ell \tilde{C}_{\#\mathbf{v}_j}, \qquad (358)$$

$$\tilde{C}_k := C_k \left(\exp(L_1 \Delta t_0) - 1 \right) \,. \tag{359}$$

Let $k \in [\![\beta]\!]$. Let $\mathbf{v} \in \mathbb{D}_d^k$. Set $u_{\mathbf{v}}(x,t) = D_{x,\mathbf{v}}U(f,t,x)$. We will show, by induction over k, that

$$\sup_{x \in \mathbb{R}^d} \|u_{\mathbf{v}}(x,t)\|_2 \le \begin{cases} \exp(L_1 t) & \text{if } k = 1, \\ C_k \left(\exp(L_1 t) - 1\right) & \text{otherwise}. \end{cases}$$
(360)

Lemma D.9 shows the case k = 1. Now let $k \ge 2$ and assume (360) for all $\mathbf{v} \in \mathbb{D}_d^{k-1}$. Let $\mathbf{v} \in \mathbb{D}_d^k$. By Theorem D.11,

$$\dot{u}_{\mathbf{v}}(x,t) = \sum_{(\mathbf{v}_1,\dots,\mathbf{v}_\ell) \in \mathbb{A}(\mathbf{v})} (D^\ell f) (U(f,x,t)) [u_{\mathbf{v}_1}(x,t),\dots,u_{\mathbf{v}_\ell}(x,t)].$$
(361)

Thus,

$$\left\|\dot{u}_{\mathbf{v}}(x,t)\right\|_{2} \leq \sum_{(\mathbf{v}_{1},\dots,\mathbf{v}_{\ell})\in\mathbb{A}(\mathbf{v})} \left|D^{\ell}f\right|_{\infty} \prod_{j=1}^{\ell} \left\|u_{\mathbf{v}_{j}}(x,t)\right\|_{2}$$
(362)

$$\leq L_1 \| u_{\mathbf{v}}(x,t) \|_2 + \sum_{(\mathbf{v}_1,\dots,\mathbf{v}_\ell) \in \mathbb{A}(\mathbf{v}) \setminus \{\mathbf{v}\}} L_\ell \prod_{j=1}^{\ell} \| u_{\mathbf{v}_j}(x,t) \|_2 .$$
(363)

Set $z(t) := ||u_{\mathbf{v}}(x,t)||_2$. Then

$$\dot{z}(t) = \frac{\dot{u}_{\mathbf{v}}(x,t)^{\top} u_{\mathbf{v}}(x,t)}{\|u_{\mathbf{v}}(x,t)\|_{2}} \le \|\dot{u}_{\mathbf{v}}(x,t)\|_{2} \le L_{1}\left(z(t) + C_{k}\right) \,. \tag{364}$$

Consider the ODE

$$\dot{v}(t) = L_1 \left(v(t) + C_k \right) \tag{365}$$

with $v(0) = ||u_{\mathbf{v}}(x, 0)||_2 = 0$ for $k \ge 2$. Then

$$v(t) = C_k \left(\exp(L_1 t) - 1 \right) \,. \tag{366}$$

By the Lemma D.1,

$$z(t) \le v(t) \,. \tag{367}$$

Hence

$$\|u_{\mathbf{v}}(x,t)\|_{2} \le C_{k} \left(\exp(L_{1}t) - 1\right) \,. \tag{368}$$

For $k \geq 2$, $D^k \Upsilon(f, t, x) = D^k U(f, x, t)$. Thus,

$$\sup_{x \in \mathbb{R}^d} \|D^k \Upsilon(f, t, x)\|_{\mathsf{op}} \preccurlyeq \sup_{x \in \mathbb{R}^d} \sup_{\mathbf{v} \in \mathbb{D}_d^k} \|u_{\mathbf{v}}(x, t)\|_2 \le C_k \left(\exp(L_1 t) - 1\right).$$
(369)

For k = 1, the result follows from Lemma D.9 with $C_1 = 1$.

References

- [Ant07] A. Antoniadis. "Wavelet methods in statistics: some recent developments and their applications". In: *Stat. Surv.* 1 (2007), pp. 16–55. DOI: 10.1214/07-SS014. MR 2520413.
- [Arn06] V. I. Arnold. Ordinary differential equations. Universitext. Translated from the Russian by Roger Cooke, Second printing of the 1992 edition. Springer-Verlag, Berlin, 2006, pp. ii+334. MR 2242407.
- [Bis08] J. P. N. Bishwal. Parameter estimation in stochastic differential equations. Vol. 1923. Lecture Notes in Mathematics. Springer, Berlin, 2008, pp. xiv+264. DOI: 10.1007/978-3-540-74448-1. MR 2360279.
- [BRR98] P. J. Bickel, Y. Ritov, and T. Rydén. "Asymptotic normality of the maximum-likelihood estimator for general hidden Markov models". In: Ann. Statist. 26.4 (1998), pp. 1614– 1635. DOI: 10.1214/aos/1024691255. MR 1647705.
- [Bru08] N. J.-B. Brunel. "Parameter estimation of ODE's via nonparametric estimators". In: Electron. J. Stat. 2 (2008), pp. 1242–1267. DOI: 10.1214/07-EJS132. MR 2471285.
- [But16] J. C. Butcher. Numerical methods for ordinary differential equations. Third. With a foreword by J. M. Sanz-Serna. John Wiley & Sons, Ltd., Chichester, 2016, pp. xxiii+513. DOI: 10.1002/9781119121534. MR 3559553.
- [CG20] F. Comte and V. Genon-Catalot. "Nonparametric drift estimation for i.i.d. paths of stochastic differential equations". In: Ann. Statist. 48.6 (2020), pp. 3336–3365. DOI: 10.1214/19-A0S1933. MR 4185811.
- [Che+18] R. T. Q. Chen, Y. Rubanova, J. Bettencourt, and D. Duvenaud. "Neural Ordinary Differential Equations". In: Proceedings of the 32nd International Conference on Neural Information Processing Systems. NIPS'18. Montréal, Canada: Curran Associates Inc., 2018, pp. 6572–6583.
- [CR72] P. G. Ciarlet and P. A. Raviart. "General lagrange and hermite interpolation in Rn with applications to finite element methods". In: Archive for Rational Mechanics and Analysis 46.3 (Jan. 1, 1972), pp. 177–199. DOI: 10.1007/BF00252458.
- [Cra05] A. D. D. Craik. "Prehistory of Faà di Bruno's formula". In: Amer. Math. Monthly 112.2 (2005), pp. 119–130. DOI: 10.2307/30037410. MR 2121322.
- [Dat21] I. Dattner. "Differential equations in data analysis". In: Wiley Interdiscip. Rev. Comput. Stat. 13.6 (2021), Paper No. e1534, 20. DOI: 10.1002/wics.1534. MR 4336079.
- [DK15] I. Dattner and C. A. J. Klaassen. "Optimal rate of direct estimators in systems of ordinary differential equations linear in functions of the parameters". In: *Electron. J. Stat.* 9.2 (2015), pp. 1939–1973. DOI: 10.1214/15-EJS1053. MR 3391125.
- [Don10] T. Donovan. Replay: The History of Video Games. Yellow Ant, 2010.
- [FY03] J. Fan and Q. Yao. Nonlinear time series. Springer Series in Statistics. Nonparametric and parametric methods. Springer-Verlag, New York, 2003, pp. xx+551. DOI: 10.1007/ b97702. MR 1964455.
- [GK12] S. Gugushvili and C. A. J. Klaassen. "√n-consistent parameter estimation for systems of ordinary differential equations: bypassing numerical integration via smoothing". In: *Bernoulli* 18.3 (2012), pp. 1061–1098. DOI: 10.3150/11-BEJ362. MR 2948913.
- [GR21] G. A. Gottwald and S. Reich. "Supervised learning from noisy observations: combining machine-learning techniques with data assimilation". In: *Phys. D* 423 (2021), Paper No. 132911, 15. DOI: 10.1016/j.physd.2021.132911. MR 4249157.
- [Har02] P. Hartman. Ordinary differential equations. Vol. 38. Classics in Applied Mathematics. Corrected reprint of the second (1982) edition [Birkhäuser, Boston, MA; MR0658490 (83e:34002)], With a foreword by Peter Bates. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2002, pp. xx+612. DOI: 10.1137/1.9780898719222. MR 1929104.

- [Hei+18] M. Heinonen, C. Yildiz, H. Mannerström, J. Intosalmi, and H. Lähdesmäki. "Learning unknown ODE models with Gaussian processes". In: International Conference on Machine Learning. PMLR. 2018, pp. 1959–1968.
- [HJ94] R. A. Horn and C. R. Johnson. *Topics in matrix analysis*. Corrected reprint of the 1991 original. Cambridge University Press, Cambridge, 1994, pp. viii+607. MR 1288752.
- [How91] G. W. Howell. "Derivative error bounds for Lagrange interpolation: An extension of Cauchy's bound for the error of Lagrange interpolation". In: *Journal of Approximation Theory* 67.2 (Nov. 1, 1991), pp. 164–173. DOI: 10.1016/0021-9045(91)90015-3.
- [Lah+24] K. Lahouel, M. Wells, V. Rielly, E. Lew, D. Lovitz, and B. M. Jedynak. "Learning nonparametric ordinary differential equations from noisy data". In: J. Comput. Phys. 507 (2024), Paper No. 112971, 22. DOI: 10.1016/j.jcp.2024.112971. MR 4729758.
- [Mar+23] Y. Marzouk, Z. Ren, S. Wang, and J. Zech. Distribution learning via neural differential equations: a nonparametric statistical perspective. 2023. DOI: 10.48550/arXiv.2309. 01043. arXiv: 2309.01043 [math.ST].
- [MMP15] K. McGoff, S. Mukherjee, and N. Pillai. "Statistical inference for dynamical systems: a review". In: Stat. Surv. 9 (2015), pp. 209–252. DOI: 10.1214/15-SS111. MR 3422438.
- [MRS12] E. Mammen, C. Rothe, and M. Schienle. "Nonparametric regression with nonparametrically generated covariates". In: *The Annals of Statistics* 40.2 (Apr. 2012). Publisher: Institute of Mathematical Statistics, pp. 1132–1170. DOI: 10.1214/12-A0S995.
- [Olv06] P. J. Olver. "On multivariate interpolation". In: *Stud. Appl. Math.* 116.2 (2006), pp. 201–240. DOI: 10.1111/j.1467-9590.2006.00335.x. MR 2200737.
- [QZ10] X. Qi and H. Zhao. "Asymptotic efficiency and finite-sample properties of the generalized profiling estimation of parameters in ordinary differential equations". In: Ann. Statist. 38.1 (2010), pp. 435–481. DOI: 10.1214/09-A0S724. MR 2589327.
- [RH17] J. Ramsay and G. Hooker. Dynamic data analysis. Springer Series in Statistics. Modeling data with differential equations. Springer, New York, 2017, pp. xvii+230. DOI: 10.1007/978-1-4939-7190-9. MR 3645102.
- [RW06] C. E. Rasmussen and C. K. I. Williams. Gaussian Processes for Machine Learning. The MIT Press, 2006.
- [RW94] D. Ruppert and M. P. Wand. "Multivariate locally weighted least squares regression".
 In: Ann. Statist. 22.3 (1994), pp. 1346–1370. DOI: 10.1214/aos/1176325632. MR 1311979.
- [Sch20] J. Schmidt-Hieber. "Nonparametric regression using deep neural networks with ReLU activation function". In: Ann. Statist. 48.4 (2020), pp. 1875–1897. DOI: 10.1214/19–A0S1875. MR 4134774.
- [SS24] C. Schötz and M. Siebel. "Lower Bounds for Nonparametric Estimation of Ordinary Differential Equations". In: (2024).
- [Str24] S. Strogatz. Nonlinear Dynamics and Chaos. With Applications to Physics, Biology, Chemistry, and Engineering. Third edition. New York: Chapman and Hall/CRC, 2024. DOI: 10.1201/9780429398490.
- [Tal21] M. Talagrand. Upper and lower bounds for stochastic processes—decomposition theorems. Vol. 60. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics. Second edition. Springer, Cham, 2021, pp. xviii+726. DOI: 10.1007/978-3-030-82595-9. MR 4381414.
- [Tsy09] A. B. Tsybakov. Introduction to nonparametric estimation. Springer Series in Statistics. Revised and extended from the 2004 French original, Translated by Vladimir Zaiats. Springer, New York, 2009, pp. xii+214. DOI: 10.1007/b13794. MR 2724359.
- [Xia19] L. Xiao. "Asymptotic theory of penalized splines". In: *Electron. J. Stat.* 13.1 (2019), pp. 747–794. DOI: 10.1214/19-ejs1541. MR 3925516.