

ALGEBRAIC RELATIONS OF SPECIAL v -ADIC ARITHMETIC GAMMA VALUES AND THEIR PERIOD INTERPRETATIONS

CHIEH-YU CHANG, FU-TSUN WEI, AND JING YU

ABSTRACT. Let v be a finite place of $\mathbb{F}_q(\theta)$. We show that algebraic relations of v -adic arithmetic gamma values over $\mathbb{F}_q(\theta)$ are explained by the standard functional equations together with Thakur's analogue of the Gross–Koblitz formula. A key step is working out a formula expressing v -adic crystalline–de Rham periods of Carlitz motives with Complex Multiplication as products of these special gamma values (a v -adic Chowla–Selberg formula). The algebraic independence of those periods in question results from determining the dimension of the motivic Galois groups, adapting and further developing existing tools.

1. INTRODUCTION

1.1. **Motivation.** This study is inspired by the Gross–Koblitz formula [GK79] and its function field analogue established by Thakur [T88]. For a prime number p , Morita [Mo75] defined a continuous function $\Gamma_p : \mathbb{Z}_p \rightarrow \mathbb{Z}_p^\times$ via a p -adic interpolation of the factorials $n!$ (up to a sign). The special p -adic gamma values we are interested in, are the values $\Gamma_p(z)$ for $z \in \mathbb{Q} \cap \mathbb{Z}_p$. Since the very unique π is missing in the p -adic domains, $\Gamma_p(1/2)$ for odd prime p , becomes algebraic. For an odd prime p , Gross–Koblitz managed to express a Gauss sum explicitly in terms of a product of specific special Morita p -adic gamma values. As a consequence, they derived the algebraicity of the special p -adic gamma value $\Gamma_p(z)$ whenever the denominator of $z \in \mathbb{Q} \cap \mathbb{Z}_p$ is a factor of $p - 1$.

Classically, one interpolates the factorials $n!$ to obtain the Euler gamma function, and a folklore conjecture asserts that all the special gamma values $\Gamma(z)$, for $z \in \mathbb{Q} \setminus \mathbb{Z}$, are transcendental. This is only known when the denominator is 2 as $\Gamma(\frac{1}{2}) = \sqrt{\pi}$, or when the denominator is 4 or 6. The latter case is due to Chudnovsky's work [Chud84, p. 6-8] (see also [BW07, p. 30]) on the algebraic independence of a nonzero period and its corresponding quasi-period of a CM elliptic curve defined over $\overline{\mathbb{Q}}$. The Lang–Rohrlich conjecture [La90] asserts that all $\overline{\mathbb{Q}}$ -algebraic relations among special gamma values and $2\pi\sqrt{-1}$ are explained by the standard functional equations satisfied by the Euler gamma function.

Back to the p -adic case, Gross–Koblitz's algebraicity result clearly reveals a deviation of the p -adic interpolations from the classical ∞ -adic interpolation. It is a natural question whether for any odd prime p , all the special Morita gamma values are transcendental except those explained in Gross–Koblitz's paper. One may ask further whether there is a ' p -adic Lang–Rohrlich phenomenon' (namely, the known existing algebraic relations arising from the standard functional equations that Morita's p -adic gamma function satisfies together with the Gross–Koblitz formula) accounting for all $\overline{\mathbb{Q}}$ -algebraic relations among the special p -adic gamma values. Our first main result (Theorem 1.2.1) is to give an affirmative answer for this question in the positive characteristic world.

In this article, we investigate the v -adic arithmetic gamma function for the rational function field k of one variable over a finite field, after L. Carlitz [Car35], D. Goss [Go80] and

Date: July 23, 2024.

2010 *Mathematics Subject Classification.* Primary 11R58, 11J93.

D. Thakur [T88], [T91]. In this case, the transcendence of ∞ -adic special arithmetic gamma values was well-known before the 21 century [T96, All96]. For any finite place v of k , we are able to show that algebraicity of special arithmetic v -adic gamma values comes only from Thakur's analogue of Gross-Koblitz. The case of special v -adic *geometric* gamma values will be treated in a forthcoming work.

Another motivation of our study on special v -adic arithmetic gamma values comes from the classical Chowla-Selberg formula, which relates a nonzero period of a CM elliptic curve over $\overline{\mathbb{Q}}$ to a specific product of special gamma values. Inspired by the p -adic Chowla-Selberg formula established by Oguz [O89], we prove in Theorem 1.3.1 that certain crystalline-de Rham periods of Carlitz motives with complex multiplication can be expressed as products of special v -adic arithmetic gamma values. It turns out that such a period interpretation plays a crucial role in the process of proving algebraic independence result of Theorem 1.2.1.

1.2. Algebraic relations among v -adic arithmetic gamma values. Let $A := \mathbb{F}_q[\theta]$ be the polynomial ring in one variable θ over a finite field \mathbb{F}_q of characteristic p with q elements, and $k := \mathbb{F}_q(\theta)$ be its fraction field. Put $D_0 := 1$, and for $i \geq 1$, $D_i := \prod_{j=0}^{i-1} (\theta^{q^i} - \theta^{q^j}) \in A$. For each $i \in \mathbb{N}$, D_i equals the product of all monic polynomials of degree i in A . Carlitz [Car35] introduced his factorial for A as follows: given a non-negative integer n , write $n = \sum_{i \geq 0} n_i q^i$ for $0 \leq n_i < q$ and put

$$\Gamma_{\text{ari},\infty}(n+1) := \prod_{i \geq 0} D_i^{n_i} \in A,$$

which becomes the positive characteristic analogue of the factorial $n!$, see (2.1.1). Taking the unit part of these factorial values, Goss [Go80] discovered that one may do interpolation in Morita style, arriving at a continuous function:

$$\Gamma_{\text{ari},\infty} : \mathbb{Z}_p \rightarrow \mathbb{F}_q((1/\theta))^\times.$$

Goss successfully applied his strategy to all places of k . Given a monic irreducible polynomial $v \in \mathbb{F}_q[\theta]$, let k_v be the completion of k at v . Let \bar{k}_v be a fixed algebraic closure of k_v , and \bar{k} be the algebraic closure of k inside \bar{k}_v . Using v -adic interpolation for the Carlitz factorials, Goss [Go88] and Thakur [T91] defined a v -adic arithmetic gamma function $\Gamma_{\text{ari},v} : \mathbb{Z}_p \rightarrow k_v^\times$, whose precise definition is given in Section 2.2.

In the ∞ -adic case, the special values $\Gamma_{\text{ari},\infty}(z)$, for $z \in \mathbb{Q} \cap \mathbb{Z}_p \setminus \mathbb{Z}$, are all transcendental over k by Thakur [T96] and Allouche [All96]. Furthermore, using transcendence theory in terms of t -motives developed by Anderson-Brownawell-Papanikolas [ABP04] and Papanikolas [P08], all algebraic relations among these special values are shown to follow from the standard monomial relations [CPTY10]. These relations come directly from the functional equations satisfied by $\Gamma_{\text{ari},\infty}$. Note that $\mathbb{Q} \cap \mathbb{Z}_p = \bigcup_{\ell \in \mathbb{N}} \frac{1}{q^\ell - 1} \cdot \mathbb{Z}$. In particular, one has:

$$\text{tr. deg}_k k(\Gamma_{\text{ari},\infty}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1)z \in \mathbb{Z}) = \ell.$$

In the v -adic case, the functional equations of $\Gamma_{\text{ari},v}$ derived in [Go88] and [T91] also give rise to standard monomial relations (2.2.3) among v -adic arithmetic gamma values at rational p -adic integers. However for the finite place v , one naturally encounters the Gross-Koblitz phenomenon, as shown by Thakur [T88]. This gives additional relations among the special values in question (see Theorem 2.2.5 and Corollary 2.2.7). The first main result of this paper is to prove the following theorem which illustrates the v -adic Lang-Rohrlich phenomenon.

Theorem 1.2.1. *Given a positive integer ℓ , we have that*

$$\text{tr. deg}_{\bar{k}}(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1)z \in \mathbb{Z}) = \ell - \gcd(\ell, \deg v).$$

Consequently, all the \bar{k} -algebraic relations among v -adic arithmetic gamma values at rational p -adic integers are explained by the functional equations in Proposition 2.2.1, the standard monomial relations (2.2.3) and Thakur’s analogue of Gross–Koblitz formula.

The decrease of transcendence degree for v -adic special values resulting from the analogue of Gross–Koblitz formula is precisely given by $\gcd(\ell, \deg v)$. Also, the above theorem complements Thakur’s theorem (cf. Remark 4.1.10):

Corollary 1.2.2. (See Corollary 4.1.9.) *Given $a, b \in \mathbb{Z}$ with $\gcd(a, b) = 1$ and $p \nmid b$, we have that*

$$\Gamma_{\text{ari},v}\left(\frac{a}{b}\right) \text{ is transcendental over } k \quad \text{if and only if} \quad q^{\deg v} \not\equiv 1 \pmod{b}.$$

Remark 1.2.3. In the case of $\deg v = 1$, Thakur [T98] initially derived the transcendence of $\Gamma_{\text{ari},v}\left(\frac{a}{b}\right)$ in the aforementioned corollary using methods from automata theory for special cases. Thakur’s approach was later generalized to encompass general $\frac{a}{b}$ in [WY02]. Also, we point out that a v -adic Anderson–Brownawell–Papanikolas criterion in [Chen23] would lead to an alternative proof of the above corollary when $\deg v = 1$.

Note that the naturally existing monomial relations indicate that the transcendence degree specified in Theorem 1.2.1 is bounded above by $\ell - \gcd(\ell, \deg v)$, see Theorem 2.2.9. To derive the opposite inequality, we shall express the v -adic arithmetic gamma values in terms of the v -adic periods of the “crystalline–de Rham comparison isomorphism” for Carlitz t -motives with complex multiplication by constant field extensions.

1.3. A v -adic Chowla–Selberg formula. Let $\mathbb{F}_q[t]$ be the polynomial ring in another variable t over \mathbb{F}_q . Given a positive integer ℓ , we let \mathbf{C}_ℓ be the Carlitz $\mathbb{F}_q[t]$ -module, then \mathbf{C}_ℓ is a CM Drinfeld $\mathbb{F}_q[t]$ -module of rank ℓ with CM field $\mathbb{F}_{q^\ell}(t)$. In [CPTY10, Theorems 3.3.1 and 3.3.2], it is shown that periods of the dual t -motive associated to \mathbf{C}_ℓ are expressed as products of special ∞ -adic gamma values (up to algebraic multiples). By [Pe08, Sec. 4.2] (see also [CP12, (3.4.5) and (3.4.8)]), periods and quasi-periods of \mathbf{C}_ℓ are explicitly related to the periods of the dual t -motive in question, and hence combining these gives a Chowla–Selberg formula for special ∞ -adic arithmetic gamma values.

Here we shall present a v -adic counterpart of the phenomenon above. Let \mathbf{C}_ℓ be the Carlitz t -motive of degree ℓ over $\mathbb{F}_q[\theta]$, and denote by $\mathbf{R}_\tau(\mathbf{C}_\ell)$ the ‘restriction of scalars’ of \mathbf{C}_ℓ (see Section 3.1). We note that the terminology ‘degree ℓ ’ indicates that \mathbf{C}_ℓ has complex multiplication by the degree ℓ constant field extension over $\mathbb{F}_q(t)$. As $\mathbf{R}_\tau(\mathbf{C}_\ell)$ is defined ‘integrally’ over $\mathbb{F}_q[\theta]$ and has ‘good reduction’ at v , one may associate its crystalline module $H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ and de Rham module $H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ over k_v (see Remark 3.5.12). Moreover, the theory of Hartl–Kim in [HaKi20] provides a crystalline–de Rham comparison isomorphism (see (3.5.13))

$$\phi_v : H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v) \xrightarrow{\sim} H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v).$$

Therefore the natural “isocrystal action” τ_v on $H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ transforms to a k_v -linear automorphism ϱ_v on $H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ through ϕ_v (see (3.7.1) for the precise definition). We show in the following theorem that ϱ_v can be expressed explicitly in terms of special v -adic arithmetic gamma values.

Theorem 1.3.1. (See Theorem 3.7.7.) *Choose the ‘global’ basis $\{\omega_0, \dots, \omega_{\ell-1}\}$ of the de Rham module $H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k)$ as in (3.7.4). For each integer s with $0 \leq s < \ell$, we write $s + d = s_0 + n'_s \ell$, where $s_0 \in \mathbb{Z}$ with $0 \leq s_0 < \ell$ and $n'_s \in \mathbb{Z}_{\geq 0}$. Then we have*

$$\varrho_v \cdot \omega_s = (-1)^{n'_s} \cdot C_s \cdot \frac{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)}{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d-1}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)^q} \cdot \omega_{s_0}, \quad \text{where } C_s := \begin{cases} -v, & \text{if } s = 0, \\ 1, & \text{if } s > 0, \end{cases}$$

and $\langle z \rangle_{\text{ari}}$ is the fractional part of z for every $z \in \mathbb{Q}$, i.e. $0 \leq \langle z \rangle_{\text{ari}} < 1$ and $z - \langle z \rangle_{\text{ari}} \in \mathbb{Z}$.

Remark 1.3.2. We may regard Theorem 1.3.1 as a function field analogue of Ogus’ p -adic Chowla–Selberg formula in [O89, Theorem 3.15].

The proof of Theorem 1.3.1 is based on an explicit description of the crystalline–de Rham comparison isomorphism ϕ_v and a product expansion formula for the special values of $\Gamma_{\text{ari},v}$ in question (see Corollary 2.3.5). By choosing a specific ‘global’ $\mathbb{F}_q[\theta][t]$ -base \mathfrak{B} of $\mathbf{R}_\tau(\mathbf{C}_\ell)$ as in (3.6.2), which induces a basis $\Lambda_v = \{\lambda_{v,0}, \dots, \lambda_{v,\ell-1}\}$ of $H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ (in (3.6.5)) and a basis $\beta_v = \{\omega_{v,0}, \dots, \omega_{v,\ell-1}\}$ of $H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v)$ (in (3.6.4)), we particularly have

$$\omega_{v,s} = \begin{cases} \left(\prod_{n=1}^{n_s-1} (\theta - \theta^{q^{\ell n+s}}) \right) \cdot \omega_s, & \text{if } s = 0, \\ \left(\prod_{n=0}^{n_s-1} (\theta - \theta^{q^{\ell n+s}}) \right) \cdot \omega_s, & \text{if } s > 0, \end{cases}$$

given in (3.7.5), and the ‘‘period matrix’’ of ϕ_v with respect to Λ_v and β_v is determined by the following result:

Theorem 1.3.3. (See Theorem 3.6.6.) *For every integer s with $0 \leq s < \ell$, we have*

$$\phi_v(\lambda_{v,s}) = \prod_{n=n_s}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{\ell n+s}}}{\theta - \varepsilon^{q^{\ell n+s}}} \right) \cdot \omega_{v,s}.$$

Here $\varepsilon = \lim_{i \rightarrow \infty} \theta^{q^{i \deg v}} \in k_v$ is the Teichmüller image of $\theta \bmod v \in A/(v)$, and n_s is the smallest non-negative integer so that $s + n_s \ell \geq \deg v$. Consequently, the period matrix of the isomorphism ϕ_v in (3.5.13) with respect to β_v and Λ_v is

$$\begin{pmatrix} \prod_{n=n_0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{\ell n}}}{\theta - \varepsilon^{q^{\ell n}}} \right) & & & 0 \\ & \ddots & & \\ & & \prod_{n=n_{\ell-1}}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{\ell n+\ell-1}}}{\theta - \varepsilon^{q^{\ell n+\ell-1}}} \right) & \\ 0 & & & \end{pmatrix}.$$

Having this period matrix at hand, we are able to realize the action of ϱ_v on the basis β_v as in (3.7.3). Combining the product formula of v -adic arithmetic gamma values in Corollary 2.3.5 enables us to derive Theorem 1.3.1.

1.4. Algebraic independence of the v -adic crystalline–de Rham periods. From the v -adic Chowla–Selberg formula in Theorem 1.3.1 and the concrete description of the period matrix in Theorem 1.3.3, our proof of Theorem 1.2.1 relies on the algebraic independence of the following v -adic periods:

Theorem 1.4.1. *Given a positive integer ℓ , the infinite products*

$$\Omega_{\ell,v,s}(v) := \prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{\ell n+s}}}{\theta - \varepsilon^{q^{\ell n+s}}}\right) \in k_v, \quad 0 < s \leq \ell,$$

are algebraically independent over \bar{k} .

Indeed, by Theorem 1.4.1, the v -adic Chowla–Selberg formula enables us to construct sufficiently many algebraically independent elements in the field

$$\bar{k}(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1)z \in \mathbb{Z}),$$

for which one ensures that its transcendence degree over \bar{k} is bounded below by $\ell - \gcd(\ell, \deg v)$ (see Theorem 4.1.4), and hence proves Theorem 1.2.1.

1.4.2. Strategy of deriving Theorem 1.4.1. In order to apply the powerful machinery of Papanikolas’ theory on Grothendieck’s period conjecture in [P08], our initial step is to regard v as the “infinite place” of the projective $\frac{1}{v}$ -line over \mathbb{F}_q . More precisely, put $\tilde{v} := \frac{1}{v}$. Then k is a finite extension of $\mathbb{F}_q(\tilde{v})$, and k_v is a finite extension of the Laurent series field $\mathbb{F}_q((\frac{1}{\tilde{v}}))$ (which is the completion of $\mathbb{F}_q(\tilde{v})$ at the place $\infty_{\tilde{v}}$ corresponding to $\frac{1}{\tilde{v}}$). So k_v is an ‘ $\infty_{\tilde{v}}$ -adic’ field. Let $\iota : \mathbb{F}_q[t] \cong \mathbb{F}_q[\theta]$ be the \mathbb{F}_q -algebra isomorphism sending t to θ , which extends to a field isomorphism $\iota : \mathbb{F}_q(t) \cong \mathbb{F}_q(\theta)$. Put $\tilde{T}_v := \iota^{-1}(\tilde{v}) \in \mathbb{F}_q(t)$. Then ι induces an \mathbb{F}_q -algebra isomorphism $\mathbb{F}_q[\tilde{T}_v] \cong \mathbb{F}_q[\tilde{v}]$. To apply Papanikolas’s theory [P08] for those special values in our $\infty_{\tilde{v}}$ -adic fields, we make the following replacements

$$\theta \leftarrow \tilde{v}, \quad t \leftarrow \tilde{T}_v$$

to switch $\bar{k}[t]$ -modules (resp. dual t -motives) to $\bar{k}[\tilde{T}_v]$ -modules (resp. dual \tilde{T}_v -motives). A detailed description of how to view v -adic fields as $\infty_{\tilde{v}}$ -adic fields is given in Sec. 5.1.

Let $d := \deg v$. Our proof of Theorem 1.4.1 is outlined in the following:

- (1) Define a “Carlitz-type” Hartl–Juschka module $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ (see Definition 6.1.3), where the infinite products $\Omega_{d\ell,v,s}(v)$ in Theorem 1.4.1 (replacing ℓ by $d\ell$), for $0 < s \leq d\ell$, show up as periods of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ (see Proposition 6.3.4).
- (2) By Papanikolas’ theorem, determining the transcendence degree of the field generated by the periods of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ is equivalent to finding the dimension of its \tilde{T}_v -motivic Galois group $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}$ (see (6.3.6)).
- (3) The “CM structure” of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ provides an upper bound of $\dim \Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}$ (cf. (5.2.6)). On the other hand, from the analysis of the Hodge–Pink weights of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$, we obtain sufficiently many “Hodge–Pink cocharacters” of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$, from which we get

$$\dim \Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})} = d^2 \ell \quad (= [\mathbb{F}_{q^{d\ell}}(t) : \mathbb{F}_q(\tilde{T}_v)]),$$

see Theorem 6.4.4.

- (4) Therefore, the period interpretation of $\Omega_{d\ell,v,s}(v)$ in (1) above for $0 < s \leq d\ell$ shows their algebraic independence over k (see Section 6.4.6). Finally, the result follows

from the identity (see (6.4.7)):

$$\Omega_{\ell,v,s}(v) = \prod_{j=0}^{d-1} \Omega_{d\ell,v,j\ell+s}(v), \quad 0 < s \leq \ell.$$

Remark 1.4.3. Put $\tilde{\theta}_v := (\theta - \varepsilon)^{-1}$, and set

$$\tilde{\Omega}_\ell(x) := \prod_{n=0}^{\infty} \left(1 - \frac{x}{\tilde{\theta}_v^{q^{\ell n}}}\right) \in k_v[[x]], \quad \forall \ell \in \mathbb{N}.$$

When $\deg v = 1$, we get $v = \theta - \varepsilon$, for which $\varepsilon \in \mathbb{F}_q$. Therefore

$$\Omega_{\ell,v,s}(v) = \tilde{\Omega}_\ell^{(s)}(x)|_{x=\tilde{\theta}_v} = \prod_{n=0}^{\infty} \left(1 - \frac{\tilde{\theta}_v}{\tilde{\theta}_v^{q^{\ell n+s}}}\right),$$

and Theorem 1.4.1 can be simply derived from the algebraic independence of periods and the quasi-periods of the Carlitz module of degree ℓ in [CPTY10] (when replacing θ to $\tilde{\theta}_v$). However, when $\deg v = d > 1$, the infinite product $\Omega_{d\ell,v,s}(v)$, for $0 < s \leq d\ell$, is actually the s th Frobenius twist of $\tilde{\Omega}_{d\ell}(x)$ evaluated at $x = \tilde{\theta}_{v,s} := (\theta - \varepsilon^{q^s})^{-1}$:

$$\Omega_{d\ell,v,s}(v) = \tilde{\Omega}_{d\ell}^{(s)}(x)|_{x=\tilde{\theta}_{v,s}} = \prod_{n=0}^{\infty} \left(1 - \frac{\tilde{\theta}_{v,s}}{\tilde{\theta}_v^{q^{d\ell n+s}}}\right).$$

Thus, to show the algebraic independence of $\Omega_{d\ell,v,s}(v)$ is equivalent to determine the algebraic relations among the evaluations of the Frobenius twists of $\tilde{\Omega}_{d\ell}(x)$ at d distinct points $x = \tilde{\theta}_{v,1}, \dots, \tilde{\theta}_{v,d}$. In order to overcome this issue, we consider the “restriction of scalars” down to $\mathbb{F}_q[\tilde{T}_v]$, so that the periods of the Hartl–Juschka $\bar{k}[\tilde{T}_v]$ -module $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ do include all of these evaluations.

Remark 1.4.4. Theorem 1.4.1 implies that

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Omega_{\ell,v,s}(v) \mid 0 < s \leq \ell \right) = \ell = \dim \Gamma_{\mathbf{R}_\tau(\mathbf{C}_\ell)},$$

where $\Gamma_{\mathbf{R}_\tau(\mathbf{C}_\ell)}$ is the “ t -motivic Galois group of the t -motive $\mathbf{R}_\tau(\mathbf{C}_\ell)$ ” (see [CPTY10, Lemma 3.2.1]). This coincidence appears to provide evidence for a suitable analogue of Papanikolas’ theorem in [P08] (Grothendieck period conjecture) associated with the v -adic periods that arise from crystalline–de Rham comparison isomorphisms.

1.5. Organization of this paper. In Section 2, we start with fixing basic notation and recalling the Carlitz factorials in Section 2.1. The definition of Goss’ v -adic arithmetic gamma function and its functional equations is given in Section 2.2, as well as Thakur’s analogue of Gross–Koblitz formula. In Section 2.3, we derive a product expansion of v -adic arithmetic gamma values, a pivotal observation that underscores their connection with the “ v -adic periods” Carlitz t -motives with complex multiplication by constant field extensions.

In Section 3, we first introduce the “integral” Carlitz t -motive \mathbf{C}_ℓ of arbitrary degree ℓ , along with its restriction of scalars $\mathbf{R}_\tau(\mathbf{C}_\ell)$ in Section 3.1. After setting up the power series rings to be used and clarifying their relations in Section 3.2, we revisit the crystalline module and de Rham module of $\mathbf{R}_\tau(\mathbf{C}_\ell)$ in Section 3.3 and 3.4, respectively. The crystalline–de Rham comparison isomorphism is explicitly constructed in Section 3.5, leading to a detailed portrayal of the crystalline–de Rham period matrix in Section 3.6. The proof of Theorem 1.3.1 is provided in Section 3.7.

In Section 4, we prove Theorem 1.2.1 on the algebraic independence of v -adic arithmetic gamma values in Section 4.1, assuming the transcendence result of crystalline periods in Theorem 1.4.1. Also, the distribution associated with v -adic arithmetic gamma values is discussed in Section 4.2.

In Section 5, we lay the groundwork for the proof of Theorem 1.4.1. Section 5.1 introduces additional notation specifically concerning the treatment of v as the infinite place of the projective $\frac{1}{v}$ -line. For completeness, Section 5.2 offers the definition of Hartl–Juschka modules, the associated motivic Galois groups, and the de Rham pairing. Also stated in Section 5.2 is Papanikolas’s theorem, which serves as the function field analogue of Grothedieck period conjecture. The Hodge–Pink cocharacters of a Hartl–Juschka module is defined in Section 5.3.

In Section 6, we introduce Carlitz-type Hartl–Juschka modules and examine their corresponding Betti modules. The definition of the module $\tilde{\mathcal{C}}_{d\ell}$, as well as its restriction of scalars $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$, is provided in Section 6.1. Section 6.2 gives a detailed account of the uniformizability of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$, after writing down explicitly the structure of the corresponding Betti modules. We articulate how the crystalline–de Rham periods of $\mathbf{R}_\tau(\mathcal{C}_{d\ell})$ can be expressed as “abelian integrals” in Section 6.3. Finally, the motivic Galois group of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ is determined in Section 6.4, culminating with the proof of Theorem 1.4.1.

2. v -ADIC ARITHMETIC GAMMA VALUES

2.1. Notation. Throughout this paper, we fix the following notation. Let p be a prime number, and let q be a power of p . Denote by \mathbb{F}_q the finite field of q elements. Let θ be a variable and let $A := \mathbb{F}_q[\theta]$ be the polynomial ring in θ over \mathbb{F}_q with quotient field $k := \mathbb{F}_q(\theta)$. For a monic irreducible polynomial $v \in A$, we still denote by v the corresponding finite place of k when it is clear from the context. The completion of k at v is denoted by k_v , and the residue field is denoted by \mathbb{F}_v . In particular, we may regard \mathbb{F}_v as a subfield of k_v via the Teichmüller embedding, which sends $\theta \bmod v$ to $\varepsilon := \lim_{i \rightarrow \infty} \theta^{q^i \deg v}$ (see [La02, Chap. XII Ex. 16]), and k_v is then identified with the Laurant series field $\mathbb{F}_v((\theta - \varepsilon))$. Note that in the case $v = \theta$ we have $\varepsilon = 0$, otherwise ε generates all the roots of unity in the local field k_v .

Choose an algebraic closure \bar{k}_v of k_v , and denote by \mathbb{C}_v the completion of \bar{k}_v . Let \bar{k} be the algebraic closure of k in \mathbb{C}_v . Let A_+ be the set of all monic polynomials in A . Put $D_0 := 1$, and for $i \geq 1$, $D_i := \prod_{j=0}^{i-1} (\theta^{q^i} - \theta^{q^j}) \in A$. For each $i \in \mathbb{N}$, we denote by $A_{+,i}$ the set of all monic polynomials of degree i , and note that $D_i = \prod_{a \in A_{+,i}} a$ (see [Car35, p. 140]). Carlitz [Car35] introduced the so-called *Carlitz factorials* as the following. For a non-negative integer n , we write $n = \sum_{i \geq 0} n_i q^i$ for $0 \leq n_i < q$ and define

$$\Gamma_{\text{ari}}(n+1) := \prod_{i \geq 0} D_i^{n_i} \in A.$$

Note that $\Gamma_{\text{ari}}(n+1)$ is analogous to the classical Euler Γ -values $\Gamma(n+1) = n!$ for $n \geq 0$ as one has the property that (due to W. Sinnott, see [Go88, Proposition 2.1.4])

$$(2.1.1) \quad \text{ord}_v(\Gamma_{\text{ari}}(n+1)) = \sum_{e \geq 1} \lfloor \frac{n}{q^e \deg v} \rfloor,$$

where $q^{\deg v}$ is the cardinality of $A/(v)$.

2.2. Definition of v -adic arithmetic gamma function. Following Goss [Go80, Go88], and Thakur [T91], we consider the v -adic arithmetic gamma function that interpolates the Carlitz factorials v -adically as follows. For each $i \geq 0$, we define

$$D_{i,v} := \prod a,$$

where a in the product above runs over all monic polynomial of degree i with $v \nmid a$. For any $z \in \mathbb{Z}_p$, the ring of p -adic integers, we express z as $z = \sum_{i \geq 0} z_i q^i$ for $0 \leq z_i < q$ and define

$$\Gamma_{\text{ari},v}(z+1) := \prod_{i \geq 0} (-D_{i,v})^{z_i}.$$

It is shown in [Go80, Appendix] that the infinite product $\prod_{i \geq 0} (-D_{i,v})^{z_i}$ converges in k_v^\times , and $\Gamma_{\text{ari},v} : \mathbb{Z}_p \rightarrow k_v^\times$ is a continuous function satisfying the following functional equations:

Proposition 2.2.1. (See [T04, Thm. 4.6.1, Lem. 4.6.2, and Thm. 4.6.3]) *Let $v \in A_+$ be irreducible. For each $z \in \mathbb{Z}_p$, we have*

(1) (Reflection)

$$\Gamma_{\text{ari},v}(z) \cdot \Gamma_{\text{ari},v}(1-z) = \Gamma_{\text{ari},v}(0).$$

(2) (Translation) *When $z \neq 0$, put*

$$r = \text{ord}_q(z) := \max \{m \in \mathbb{Z}_{\geq 0} \mid q^m | z\}.$$

Then

$$\frac{\Gamma_{\text{ari},v}(z+1)}{\Gamma_{\text{ari},v}(z)} = -\frac{D_{r,v}}{\prod_{i=0}^{r-1} D_{i,v}^{q-1}}.$$

(3) (Multiplication) *For every $n \in \mathbb{N}$ with $p \nmid n$, we have*

$$\prod_{i=0}^{n-1} \Gamma_{\text{ari},v}\left(z + \frac{i}{n}\right) = \Gamma_{\text{ari},v}(nz) \cdot \Gamma_{\text{ari},v}(0)^{\frac{n-1}{2}}.$$

As it is known that (see [T91, Theorem 4.4])

$$\Gamma_{\text{ari},v}\left(1 - \frac{1}{q-1}\right)^{q-1} = \Gamma_{\text{ari},v}(0) = (-1)^{\deg v-1},$$

the square root of $\Gamma_{\text{ari},v}(0)$ on the right hand side of the equality in Proposition 2.2.1 (3) is chosen to be

$$\Gamma_{\text{ari},v}(0)^{\frac{1}{2}} := \Gamma_{\text{ari},v}\left(1 - \frac{1}{q-1}\right)^{\frac{q-1}{2}}.$$

When taking values at a rational p -adic integer z , one has the following standard monomial relations which directly come from the q -adic expansion of z :

Proposition 2.2.2. *Let $v \in A_+$ be irreducible. Given $\ell \in \mathbb{N}$ and $c_0, \dots, c_{\ell-1} \in \mathbb{Z}$ with $0 \leq c_0, \dots, c_{\ell-1} < q$, we have*

$$(2.2.3) \quad \Gamma_{\text{ari},v}\left(1 - \sum_{i=0}^{\ell-1} \frac{c_i q^i}{q^\ell - 1}\right) = \prod_{i=0}^{\ell-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^i}{q^\ell - 1}\right)^{c_i}.$$

Remark 2.2.4. Regarding \bar{k} -algebraic relations among v -adic arithmetic gamma values at rational p -adic integers, in [T91, Sec. 2] Thakur showed that once we have the monomial relations (2.2.3) together with translation (2) of Proposition 2.2.1 (for $z \in \mathbb{Z}_p \cap \mathbb{Q}$), then the reflection (1) and multiplication (3) of Proposition 2.2.1 are valid for $z \in \mathbb{Z}_p \cap \mathbb{Q}$.

Thakur's analogue of the Gross–Koblitz formula in [T88, Theorem VI] implies:

Theorem 2.2.5. (See [T88, Corollary of Thm. V].) *Let $v \in A_+$ be irreducible. For each $z \in \mathbb{Q}$ with $(q^{\deg v} - 1) \cdot z \in \mathbb{Z}$, we have*

$$\Gamma_{\text{ari},v}(z) \in \bar{k}^\times.$$

Remark 2.2.6. In fact, Thakur's formula in [T88, Theorem VI] shows that when $z \in \mathbb{Q}$ with $(q^{\deg v} - 1) \cdot z \in \mathbb{Z}$, the special value $\Gamma_{\text{ari},v}(z)$ is related to Thakur's analogue of Gauss sums. Consequently, in this case one knows that

$$\Gamma_{\text{ari},v}(z) \in \mathbb{F}_{q^{\deg v}}(\theta) \left(\lambda_v, \sqrt[q^{\deg v} - 1]{-v} \right),$$

where λ_v is a non-zero *Carlitz v -torsion*.

As a consequence of Thakur's analogue of the Gross–Koblitz formula, we have the following monomial relations:

Corollary 2.2.7. *Let $v \in A_+$ be irreducible. Given a positive integer n with $p \nmid n$, let r be the order of $q^{\deg v} \bmod n$ in $(\mathbb{Z}/n\mathbb{Z})^\times$. For every $z \in \mathbb{Q}$ satisfying $n \cdot z \in \mathbb{Z}$, we have that*

$$\prod_{i=0}^{r-1} \Gamma_{\text{ari},v}(q^{i \deg v} \cdot z) \in \bar{k}^\times.$$

Proof. Without loss of generality, we may assume that $z = 1 - \frac{a}{n}$ with $0 \leq a < n$. The assumption of n indicates that $n \mid q^{r \deg v} - 1$. Write $q^{r \deg v} - 1 = n \cdot n'$. Then there exist $z_0, \dots, z_{r \deg v - 1} \in \mathbb{Z}$ with $0 \leq z_0, \dots, z_{r \deg v - 1} < q$ satisfying that

$$\frac{a}{n} = \frac{an'}{q^{r \deg v} - 1} = \sum_{j=0}^{r \deg v - 1} \frac{z_j q^j}{q^{r \deg v} - 1}.$$

Therefore

$$\begin{aligned} \prod_{i=0}^{r-1} \Gamma_{\text{ari},v}(q^{i \deg v} \cdot z) &= \prod_{i=0}^{r-1} \Gamma_{\text{ari},v} \left(q^{i \deg v} \cdot \left(1 - \frac{a}{n} \right) \right) \\ &\sim \prod_{i=0}^{r-1} \prod_{j=0}^{r \deg v - 1} \Gamma_{\text{ari},v} \left(1 - \frac{z_j q^{j+i \deg v}}{q^{r \deg v} - 1} \right) \\ &\sim \prod_{j=0}^{r \deg v - 1} \Gamma_{\text{ari},v} \left(1 - \frac{\sum_{i=0}^{r-1} q^{j+i \deg v}}{q^{r \deg v} - 1} \right)^{z_j} \\ &= \prod_{j=0}^{r \deg v - 1} \Gamma_{\text{ari},v} \left(1 - \frac{q^j}{q^{\deg v} - 1} \right)^{z_j} \in \bar{k}^\times \quad (\text{by Theorem 2.2.5}). \end{aligned}$$

Here, for $x, y \in \mathbb{C}_v^\times$, we denote by $x \sim y$ if $x/y \in \bar{k}^\times$. □

Remark 2.2.8. Since when $n \mid (q^{\deg v} - 1)$, the order of $q^{\deg v} \bmod n$ in $(\mathbb{Z}/n\mathbb{Z})^\times$ is 1. Therefore Corollary 2.2.7 is in fact equivalent to Theorem 2.2.5.

The above monomial relations among special values of $\Gamma_{\text{ari},v}$ at rational p -adic integers give rise to the following upper bound of the transcendence degree in question.

Theorem 2.2.9. *Let $v \in A_+$ be irreducible. Given $\ell \in \mathbb{N}$, we have*

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1) \cdot z \in \mathbb{Z} \right) \leq \ell - \gcd(\ell, \deg v).$$

Proof. The functional equation in Proposition 2.2.1 (2) and the monomial relation in Proposition 2.2.2 show that

$$\bar{k}\left(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1) \cdot z \in \mathbb{Z}\right) = \bar{k}\left(\Gamma_{\text{ari},v}\left(1 - \frac{q^i}{q^\ell - 1}\right) \mid 0 \leq i \leq \ell - 1\right).$$

Let $m = \gcd(\ell, \deg v)$ and $\ell' = \ell/m$. By Proposition 2.2.2 and Theorem 2.2.5, we get

$$(2.2.10) \quad \prod_{j=0}^{\ell'-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^{i+mj}}{q^\ell - 1}\right) = \Gamma_{\text{ari},v}\left(1 - \frac{q^i}{q^m - 1}\right) \in \bar{k}, \quad \forall i \in \mathbb{Z}, 0 \leq i < m,$$

since $(q^m - 1)$ divides $(q^{\deg v} - 1)$. The monomial relations (2.2.10) for $0 \leq i \leq m - 1$ give rise to m independent algebraic relations, from which the result follows. \square

We will prove in Section 4 that the inequality stated in Theorem 2.2.9 is, in fact, an equality. Before doing so, we first provide “period interpretations” of the v -adic arithmetic gamma values in question.

2.3. Product expansions of v -adic arithmetic gamma values. We begin with the following observation:

Lemma 2.3.1. *Let $v \in A_+$ be irreducible of degree d . Given $\ell, s \in \mathbb{N}$, let n_s be the smallest non-negative integer such that $s + n_s \ell \geq d$. We have that*

$$\frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{s-1}}{q^\ell - 1}\right)^q} = \frac{\prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}}\right)}{\prod_{n=n_s+1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell-d}}}{\theta - \varepsilon^{q^{s+n\ell-d}}}\right)} \cdot \left(\prod_{n=0}^{n_s} (\varepsilon^{q^{s+n\ell}} - \theta)\right) \cdot \begin{cases} (\theta^{q^{s+n_s\ell-d}} - \theta)^{-1}, & \text{if } s + n_s\ell > d, \\ v^{-1}, & \text{if } s + n_s\ell = d. \end{cases}$$

Proof. From the definition of $\Gamma_{\text{ari},v}$, we get

$$\frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{s-1}}{q^\ell - 1}\right)^q} = \prod_{n=0}^{\infty} \frac{D_{s+n\ell,v}}{D_{s-1+n\ell,v}^q}.$$

Note that for every $n \in \mathbb{N}$, one has that

$$\frac{D_n}{D_{n-1}^q} = (\theta^{q^n} - \theta) \quad \text{and} \quad D_{n,v} = \begin{cases} \frac{D_n}{v^{q^n-d} D_{n-d}}, & \text{if } n \geq d, \\ D_n, & \text{if } n < d. \end{cases}$$

Thus

$$(2.3.2) \quad \frac{D_{n,v}}{D_{n-1,v}^q} = \begin{cases} \frac{\theta^{q^n} - \theta}{\theta^{q^n-d} - \theta} & \text{if } n > d, \\ \frac{\theta^{q^n} - \theta}{v}, & \text{if } n = d, \\ \theta^{q^n} - \theta, & \text{if } 0 < n < d. \end{cases}$$

Observe that:

$$(2.3.3) \quad \theta^{q^n} - \theta = (\varepsilon^{q^n} - \theta) \cdot \left(1 - \frac{(\theta - \varepsilon)^{q^n}}{\theta - \varepsilon^{q^n}}\right)$$

and

$$(2.3.4) \quad \frac{\theta^{q^n} - \theta}{\theta^{q^{n-d}} - \theta} = \left(1 - \frac{(\theta - \varepsilon)^{q^n}}{\theta - \varepsilon^{q^n}}\right) \Big/ \left(1 - \frac{(\theta - \varepsilon)^{q^{n-d}}}{\theta - \varepsilon^{q^{n-d}}}\right) \quad \text{when } n > d.$$

As n_s is the smallest non-negative integer such that $s + n_s \ell \geq d$, we obtain that

$$\begin{aligned} & \frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{s-1}}{q^\ell - 1}\right)^q} = \prod_{n=0}^{\infty} \frac{D_{s+n\ell,v}}{D_{s-1+n\ell,v}^q} \\ &= \left(\prod_{n=n_s+1}^{\infty} \frac{D_{s+n\ell,v}}{D_{s-1+n\ell,v}^q} \right) \cdot \left(\prod_{n=0}^{n_s-1} \frac{D_{s+n\ell,v}}{D_{s-1+n\ell,v}^q} \right) \cdot \frac{D_{s+n_s\ell,v}}{D_{s-1+n_s\ell,v}^q} \\ &= \left(\prod_{n=n_s+1}^{\infty} \frac{\theta^{q^{s+n\ell}} - \theta}{\theta^{q^{s+n\ell-d}} - \theta} \right) \cdot \left(\prod_{n=0}^{n_s-1} (\theta^{q^{s+n\ell}} - \theta) \right) \cdot \begin{cases} \frac{\theta^{q^{s+n_s\ell}} - \theta}{\theta^{q^{s+n_s\ell-d}} - \theta}, & \text{if } s + n_s\ell > d, \\ \frac{\theta^{q^{s+n_s\ell}} - \theta}{v}, & \text{if } s + n_s\ell = d \end{cases} \\ & \hspace{25em} \text{(by (2.3.2))} \\ &= \left(\prod_{n=n_s+1}^{\infty} \frac{1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}}}{1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell-d}}}{\theta - \varepsilon^{q^{s+n\ell-d}}}} \right) \cdot \prod_{n=0}^{n_s} (\varepsilon^{q^{s+n\ell}} - \theta) \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}}\right) \\ & \cdot \begin{cases} (\theta^{q^{s+n_s\ell-d}} - \theta)^{-1}, & \text{if } s + n_s\ell > d, \\ v^{-1}, & \text{if } s + n_s\ell = d \end{cases} \hspace{5em} \text{(by (2.3.3) \& (2.3.4))} \\ &= \frac{\prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}}\right)}{\prod_{n=n_s+1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell-d}}}{\theta - \varepsilon^{q^{s+n\ell-d}}}\right)} \cdot \left(\prod_{n=0}^{n_s} (\varepsilon^{q^{s+n\ell}} - \theta) \right) \begin{cases} (\theta^{q^{s+n_s\ell-d}} - \theta)^{-1}, & \text{if } s + n_s\ell > d, \\ v^{-1}, & \text{if } s + n_s\ell = d, \end{cases} \end{aligned}$$

where $\prod_{n=0}^{n_s-1} \frac{D_{s+n\ell,v}}{D_{s-1+n\ell,v}^q}$ occurring in the second identity and $\prod_{n=0}^{n_s-1} (\theta^{q^{s+n\ell}} - \theta)$ occurring in the third identity are referred to be 1 in the case when $n_s = 0$, the fourth and fifth identities come from (2.3.4) and (2.3.3) respectively. \square

Consequently, for each $z \in \mathbb{Q}$, let $\langle z \rangle_{\text{ari}} \in \mathbb{Q}$ be the fractional part of z , i.e. $0 \leq \langle z \rangle_{\text{ari}} < 1$ and $z - \langle z \rangle_{\text{ari}} \in \mathbb{Z}$. We get:

Corollary 2.3.5. *Let $v \in A_+$ be irreducible of degree d . Given $\ell \in \mathbb{N}$ and $s \in \mathbb{Z}_{\geq 0}$ with $0 \leq s < \ell$, write $s + d = s_0 + n'_s \ell$ where $s_0 \in \mathbb{Z}$ with $0 \leq s_0 < \ell$ and $n'_s \in \mathbb{Z}_{\geq 0}$. Then*

$$\frac{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)}{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d-1}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)^q} = \frac{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s_0+n\ell}}}{\theta - \varepsilon^{q^{s_0+n\ell}}} \right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}} \right)} \cdot \left(\prod_{n=0}^{n'_s-1} (\varepsilon^{q^{s_0+n\ell}} - \theta) \right) \cdot \frac{\alpha_{s_0}}{\alpha_s} \cdot C_s^{-1},$$

where

$$(2.3.6) \quad \alpha_s := \begin{cases} 1 - \frac{(\theta - \varepsilon)^{q^s}}{\theta - \varepsilon^{q^s}}, & \text{if } s > 0, \\ \frac{1}{\theta - \varepsilon}, & \text{if } s = 0, \end{cases} \quad \text{and} \quad C_s := \begin{cases} 1, & \text{if } s > 0, \\ -v, & \text{if } s = 0. \end{cases}$$

Proof. Recall that n_{s_0} denotes the smallest non-negative integer so that $s_0 + n_{s_0} \ell \geq d$. Then $n'_s \geq n_{s_0}$ and

$$0 \leq (n'_s - n_{s_0})\ell = (s_0 + n'_s \ell) - (s_0 + n_{s_0} \ell) \leq s + d - d = s < \ell.$$

Thus $n'_s = n_{s_0}$ and $s_0 + n_{s_0} \ell - d = s$.

Suppose $s_0 > 0$. Then

$$\begin{aligned} & \frac{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)}{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d-1}}{q^\ell - 1} \right\rangle_{\text{ari}} \right)^q} = \frac{\Gamma_{\text{ari},v} \left(1 - \frac{q^{s_0}}{q^\ell - 1} \right)}{\Gamma_{\text{ari},v} \left(1 - \frac{q^{s_0-1}}{q^\ell - 1} \right)^q} \\ &= \frac{\prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s_0+n\ell}}}{\theta - \varepsilon^{q^{s_0+n\ell}}} \right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}} \right)} \cdot \left(\prod_{n=0}^{n'_s} (\varepsilon^{q^{s_0+n\ell}} - \theta) \right) \begin{cases} (\theta^{q^s} - \theta)^{-1}, & \text{if } s > 0, \\ v^{-1}, & \text{if } s = 0 \end{cases} \\ &= \frac{\alpha_{s_0} \cdot \prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s_0+n\ell}}}{\theta - \varepsilon^{q^{s_0+n\ell}}} \right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}} \right)} \cdot (\theta - \varepsilon^{q^s}) \cdot \left(\prod_{n=0}^{n'_s-1} (\varepsilon^{q^{s_0+n\ell}} - \theta) \right) \begin{cases} (\theta - \theta^{q^s})^{-1}, & \text{if } s > 0, \\ (-v)^{-1}, & \text{if } s = 0 \end{cases} \\ &= \frac{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s_0+n\ell}}}{\theta - \varepsilon^{q^{s_0+n\ell}}} \right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}} \right)} \cdot \left(\prod_{n=0}^{n'_s-1} (\varepsilon^{q^{s_0+n\ell}} - \theta) \right) \cdot \frac{\alpha_{s_0}}{\alpha_s} \cdot C_s^{-1}. \end{aligned}$$

When $s_0 = 0$, one has that $n'_s = n_{s_0} = n_0 > 0$ and $n_\ell = n_0 - 1$ (as $s + d = 0 + n_0\ell = \ell + (n_0 - 1)\ell$). Then

$$\begin{aligned}
& \frac{\Gamma_{\text{ari},v}\left(1 - \left\langle \frac{q^{s+d}}{q^\ell - 1} \right\rangle\right)}{\Gamma_{\text{ari},v}\left(1 - \left\langle \frac{q^{s+d-1}}{q^\ell - 1} \right\rangle\right)^q} = \frac{\Gamma_{\text{ari},v}\left(1 - \frac{1}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{\ell-1}}{q^\ell - 1}\right)^q} = (-1) \cdot \frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^\ell}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{\ell-1}}{q^\ell - 1}\right)^q} \\
&= \frac{\prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{\ell+n\ell}}{\theta - \varepsilon q^{\ell+n\ell}}\right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{s+n\ell}}{\theta - \varepsilon q^{s+n\ell}}\right)} \cdot \left(\prod_{n=0}^{n_\ell} (\varepsilon q^{\ell+n\ell} - \theta)\right) \cdot (-1) \cdot \begin{cases} (\theta^{q^s} - \theta)^{-1}, & \text{if } s > 0, \\ v^{-1}, & \text{if } s = 0 \end{cases} \\
&= \frac{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{n\ell}}{\theta - \varepsilon q^{n\ell}}\right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{s+n\ell}}{\theta - \varepsilon q^{s+n\ell}}\right)} \cdot \left(\prod_{n=0}^{n_0-1} (\varepsilon q^{n\ell} - \theta)\right) \cdot \frac{\theta - \varepsilon q^{s+d}}{\theta - \varepsilon} \cdot \begin{cases} (\theta - \theta^{q^s})^{-1}, & \text{if } s > 0, \\ (-v)^{-1}, & \text{if } s = 0 \end{cases} \\
&= \frac{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{n\ell}}{\theta - \varepsilon q^{n\ell}}\right)}{\prod_{n=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)q^{s+n\ell}}{\theta - \varepsilon q^{s+n\ell}}\right)} \cdot \left(\prod_{n=0}^{n'_s-1} (\varepsilon q^{n\ell} - \theta)\right) \cdot \frac{\alpha_0}{\alpha_s} \cdot C_s^{-1}.
\end{aligned}$$

Therefore the proof is complete. \square

Remark 2.3.7. Let notation be given as above. Suppose $\ell \geq d$. Since

$$\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right) = \Gamma_{\text{ari},v}\left(1 - \frac{q^{s+\ell}}{q^\ell - 1}\right) \cdot (-D_{s,v}),$$

we compute that for every integer s with $0 \leq s < \ell$,

$$\begin{aligned}
\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)^{1-q^\ell} &= \frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)^{q^\ell}} = \frac{(-D_{s,v}) \cdot \Gamma_{\text{ari},v}\left(1 - \frac{q^{s+\ell}}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^\ell - 1}\right)^{q^\ell}} \\
&= (-D_{s,v}) \cdot \prod_{j=0}^{\ell-1} \left(\frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^{s+\ell-j}}{q^\ell - 1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{s+\ell-1-j}}{q^\ell - 1}\right)^q} \right)^{q^j} \\
&= D_{s,v}^{1-q^\ell} \cdot (-D_{s+\ell,v}) \cdot \prod_{j=0}^{\ell-1} \prod_{n=2}^{\infty} \left(\frac{1 - \frac{(\theta - \varepsilon)q^{s+n\ell-j}}{\theta - \varepsilon q^{s+n\ell-j}}}{1 - \frac{(\theta - \varepsilon)q^{s+n\ell-j-d}}{\theta - \varepsilon q^{s+n\ell-j}}} \right)^{q^j}.
\end{aligned}$$

In particular, when $\ell = d$, these values are actually algebraic (the Gross–Koblitz phenomenon). We get

$$\begin{aligned}
\Gamma_{\text{ari},v} \left(1 - \frac{q^s}{q^d - 1}\right)^{1-q^d} &= D_{s,v}^{1-q^d} \cdot (-D_{s+d,v}) \cdot \prod_{j=0}^{d-1} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+d}}}{(\theta - \varepsilon^{q^{s-j}})^{q^j}}\right)^{-1} \\
&= (-1) \cdot D_s^{1-q^d} \cdot \frac{D_{s+d}}{v^{q^s} \cdot D_s} \cdot \prod_{j=0}^{d-1} \frac{\theta^{q^j} - \varepsilon^{q^s}}{\theta^{q^j} - \theta^{q^{s+d}}} \\
&= (-1)^{d-1} \cdot v^{-q^s} \cdot \prod_{j=0}^{d-1} (\theta^{q^j} - \varepsilon^{q^s}) \\
&= (-1)^{d-1} \cdot \prod_{j=0}^{d-1} (\theta - \varepsilon^{q^{s-j}})^{-q^s} \cdot \prod_{j=0}^{d-1} (\theta - \varepsilon^{q^{s-j}})^{q^j} \\
&= (-1)^{d-1} \cdot \prod_{j=0}^{d-1} (\theta - \varepsilon^{q^{s-j}})^{-(q^s - q^j)},
\end{aligned}$$

which coincides with Thakur’s formula in [T88, Theorem V], indicating the exponents of the above decomposition corresponds to the ‘Stickelberger element’ associated with the constant field extension of degree d .

In the following section, we will connect the infinite product

$$\prod_{n=0}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{s+n\ell}}}{\theta - \varepsilon^{q^{s+n\ell}}}\right), \quad s \in \mathbb{Z} \text{ with } 0 < s \leq \ell,$$

with the transcendental periods arising from the v -adic ‘crystalline–de Rham comparison isomorphism’ for the Carlitz t -motive of degree ℓ .

3. CARLITZ t -MOTIVE AND THE CRYSTALLINE–DE RHAM COMPARISON

3.1. Carlitz t -motive. For each (commutative) \mathbb{F}_q -algebra \mathcal{A} , let $\mathcal{A}[t]$ be the polynomial ring with another variable t over \mathcal{A} . Let $\mathcal{A}[t, \tau] = \mathcal{A}[t][\tau]$ be the twisted polynomial ring over $\mathcal{A}[t]$ satisfying the following multiplication law:

$$\tau \cdot f(t) = f^{(1)}(t) \cdot \tau, \quad \forall f(t) \in \mathcal{A}[t],$$

where for every $f(t) = \sum_{i=0}^r a_i t^i \in \mathcal{A}[t]$, $f^{(n)}(t)$ is the n th Frobenius twist of f :

$$f^{(n)}(t) := \sum_{i=0}^r a_i^{q^n} t^i \in \mathcal{A}[t], \quad \forall n \in \mathbb{Z}_{\geq 0}.$$

Throughout this section, we fix a positive integer ℓ .

Definition 3.1.1. *The Carlitz t -motive of degree ℓ over $\mathbb{F}_q[\theta]$ is the $\mathbb{F}_q[\theta][t, \tau^\ell]$ -module $\mathbf{C}_\ell := \mathbb{F}_q[\theta][t]$ with the following τ^ℓ -action*

$$\tau^\ell \cdot m := (t - \theta) \cdot m^{(\ell)}, \quad \forall m \in \mathbf{C}_\ell.$$

The restriction of scalars of \mathbf{C}_ℓ is the following $\mathbb{F}_q[\theta][t, \tau]$ -module:

$$\mathbf{R}_\tau(\mathbf{C}_\ell) := \mathbb{F}_q[\theta][\tau] \otimes_{\mathbb{F}_q[\theta][\tau^\ell]} (\mathbb{F}_q[\theta][t]).$$

Note that we may identify

$$(3.1.2) \quad \mathbf{R}_\tau(C_\ell) = \bigoplus_{i=0}^{\ell-1} \tau^i \otimes (\mathbb{F}_q[\theta][t]) \cong \prod_{i=0}^{\ell-1} \mathbb{F}_q[\theta][t],$$

and the equipped τ -action is given by

$$\tau \cdot (m_0, m_1, \dots, m_{\ell-1}) = ((t - \theta) \cdot m_{\ell-1}^{(1)}, m_0^{(1)}, \dots, m_{\ell-2}^{(1)}).$$

For clarity in the subsequent discussions, we set up the following notation:

Definition 3.1.3. Let d and ℓ be two positive integers. For every integer i with $0 \leq i < \ell$, we put

$$n_i = \min\{n \in \mathbb{Z}_{\geq 0} \mid i + n\ell \geq d\} \quad \text{and} \quad i_1 := i + n_i\ell - d \quad (\text{for which } 0 \leq i_1 < \ell).$$

Also, we take i_0 to be the unique integer with $0 \leq i_0 < \ell$ and $i + d \equiv i_0 \pmod{\ell}$, and set $n'_i := (i + d - i_0)/\ell \in \mathbb{Z}_{\geq 0}$.

In particular, the following explicit τ^d -action on $\mathbf{R}_\tau(C_\ell)$ will be used for period interpretation of v -adic special gamma values in the crystalline-de Rham comparison isomorphism when $d = \deg v$.

Lemma 3.1.4. Let notation be given as above. Then we have

$$\begin{aligned} \tau^d \cdot (m_0, \dots, m_{\ell-1}) &= (m'_0, \dots, m'_{\ell-1}), \\ \text{where } m'_i &= \left(\prod_{n=0}^{n_i-1} (t - \theta^{q^{i+n\ell}}) \right) \cdot m_{i_1}^{(d)}, \quad \text{for } 0 \leq i < \ell. \end{aligned}$$

Proof. For each integer i with $0 \leq i < \ell$, we write $i + d = i_0 + n'_i\ell$ for unique integers i_0 and n'_i with $0 \leq i_0 < \ell$. Then

$$m'_{i_0} = \left(\prod_{n=0}^{n'_i-1} (t - \theta^{q^{i_0+n\ell}}) \right) \cdot m_i^{(d)}.$$

Based on Definition 3.1.3, as $n'_i \geq n_{i_0}$ and

$$d \leq i_0 + n_{i_0}\ell \leq i_0 + n'_i\ell = i + d < \ell + d,$$

we get

$$0 \leq n'_i\ell - n_{i_0}\ell < (\ell + d) - (d) = \ell,$$

whence $n'_i = n_{i_0}$. Hence $i = i_0 + n_{i_0}\ell - d = (i_0)_1$ and the result holds. \square

In this section, our primary goal is to explicitly illustrate the crystalline-de Rham comparison isomorphism for $\mathbf{R}_\tau(C_\ell)$ at v .

3.2. Notation. Put $d := \deg v$ as before, and write

$$v = \theta^d + \epsilon_1\theta^{d-1} + \dots + \epsilon_{d-1}\theta + \epsilon_d, \quad \text{where } \epsilon_1, \dots, \epsilon_d \in \mathbb{F}_q.$$

Let $\iota : \mathbb{F}_q[t] \cong \mathbb{F}_q[\theta]$ be the \mathbb{F}_q -algebra isomorphism with $\iota(t) = \theta$. Set

$$T_v := \iota^{-1}(v) = t^d + \epsilon_1 t^{d-1} + \dots + \epsilon_{d-1} t + \epsilon_d,$$

which is monic irreducible of degree d in $\mathbb{F}_q[t]$. We may embed $\mathbb{F}_q[t]$ into the formal power series ring $\mathbb{F}_{q^d}[[T_v]]$ (identified with the completion of $\mathbb{F}_q[t]$ at T_v) which is compatible with ι , i.e. $t \equiv \varepsilon \pmod{T_v}$, and ι is extended to an \mathbb{F}_{q^d} -isomorphism $\mathbb{F}_{q^d}[[T_v]] \cong \mathbb{F}_{q^d}[[v]]$ sending T_v to v . These settings can be illustrated in the following diagram:

$$\begin{array}{ccc}
\mathbb{F}_{q^d}[[t - \varepsilon]] = \mathbb{F}_{q^d}[[T_v]] & \xrightarrow{\sim} & \mathbb{F}_{q^d}[[v]] = \mathbb{F}_{q^d}[[\theta - \varepsilon]] \\
& \searrow & \swarrow \\
& \mathbb{F}_{q^d}[t] \xrightarrow{\sim} \mathbb{F}_{q^d}[\theta] & \\
& \downarrow & \downarrow \\
& \mathbb{F}_q[t] \xrightarrow{\sim} \mathbb{F}_q[\theta] &
\end{array}$$

In particular, the embedding $\mathbb{F}_q[t] \hookrightarrow \mathbb{F}_{q^d}[[T_v]]$ sends t to a power series in T_v as

$$(3.2.1) \quad t(T_v) = \sum_{n=0}^{\infty} a_n T_v^n \in \mathbb{F}_{q^d}[[T_v]], \quad \text{where} \quad a_0 = \varepsilon \text{ and } a_1 = \prod_{i=1}^{d-1} (\varepsilon - \varepsilon^{q^i})^{-1},$$

$$(3.2.2) \quad \text{and} \quad t(T_v)|_{T_v=v} = \varepsilon + \left(\prod_{i=1}^{d-1} (\varepsilon - \varepsilon^{q^i})^{-1} \right) v + \cdots = \theta \in \mathbb{F}_{q^d}[[v]].$$

3.2.3. Comparison between twistings. Let $A_v := \mathbb{F}_{q^d}[[v]]$ be the valuation ring of k_v . Following Hartl–Kim we let

$$A_v[[T_v, T_v^{-1}]] := \left\{ \sum_{i \in \mathbb{Z}} b_i T_v^i \in A_v[[T_v, T_v^{-1}]] \mid \lim_{i \rightarrow -\infty} b_i v^{-ni} = 0, \quad \forall n > 0 \right\},$$

on which we define the n -th Frobenius twist as follows.

Definition 3.2.4. Given $f = \sum_{i \in \mathbb{Z}} f_i T_v^i \in A_v[[T_v, T_v^{-1}]]$, its n -th Frobenius twist is given by

$$f^{(n)} := \sum_{i \in \mathbb{Z}} f_i^{q^n} T_v^i.$$

Note that the embedding $\mathbb{F}_q[t] \hookrightarrow \mathbb{F}_{q^d}[[T_v]]$ given in (3.2.1) induces an A_v -algebra embedding $A_v[t] \hookrightarrow A_v[[T_v, T_v^{-1}]]$. However, the respective n -th Frobenius twists on $A_v[t]$ and on $A_v[[T_v, T_v^{-1}]]$ are only compatible when d divides n . Indeed, regarding $t = t(T_v)$ as a power series in $\mathbb{F}_{q^d}[[T_v]]$ given in (3.2.1), we get

$$\begin{aligned}
t^{(n)}(T_v) &= \left(\varepsilon + \left(\prod_{i=1}^{d-1} (\varepsilon - \varepsilon^{q^i}) \right) \cdot T_v + \cdots \right)^{(n)} = \varepsilon^{q^n} + \left(\prod_{i=1}^{d-1} (\varepsilon - \varepsilon^{q^i}) \right)^{q^n} \cdot T_v + \cdots \\
&= \varepsilon + \left(\prod_{i=1}^{d-1} (\varepsilon - \varepsilon^{q^i}) \right) \cdot T_v + \cdots = t(T_v) \in \mathbb{F}_{q^d}[[T_v]] \quad \text{if and only if } d \mid n.
\end{aligned}$$

We would like to remind readers to be aware of the above “non-compatibility” between these two types of Frobenius twistings.

Remark 3.2.5. The twisted polynomial ring $A_v[[T_v, T_v^{-1}]][\tau]$ over $A_v[[T_v, T_v^{-1}]]$ satisfies

$$\tau \cdot f(T_v) = f^{(1)}(T_v)\tau, \quad \forall f(T_v) \in A_v[[T_v, T_v^{-1}]].$$

Because of the non-compatibility of the Frobenius twists on $A_v[t]$ and on $A_v[[T_v, T_v^{-1}]]$, we remark that $A_v[t][\tau]$ is not a subring of $A_v[[T_v, T_v^{-1}]][\tau]$ when $d > 1$, but both do contain the subring $A_v[t][\tau^d]$.

3.3. **Crystalline module of $\mathbf{R}_\tau(\mathbf{C}_\ell)$.** Put $\theta_v := \theta - \varepsilon$. Let

$$A_v := \mathbb{F}_{q^d}[[\theta_v]] = \mathbb{F}_{q^d}[[v]] \subset k_v \quad \text{and} \quad \mathfrak{m}_v := v \cdot A_v = \theta_v \cdot A_v.$$

be the valuation ring of k_v with maximal ideal

$$\mathfrak{m}_v := v \cdot A_v = \theta_v \cdot A_v.$$

Note that as $\mathbb{F}_{q^d} \cong A_v/\mathfrak{m}_v$ is an $\mathbb{F}_q[\theta]$ -algebra, we define $\overline{\mathbf{R}_\tau(\mathbf{C}_\ell)}$ to be the reduction of $\mathbf{R}_\tau(\mathbf{C}_\ell)$ modulo \mathfrak{m}_v , i.e.

$$(3.3.1) \quad \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} = \frac{A_v}{\mathfrak{m}_v} \otimes_{\mathbb{F}_q[\theta]} (\mathbf{R}_\tau(\mathbf{C}_\ell)).$$

By (3.1.2), we have the following identification

$$(3.3.2) \quad \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} = \prod_{i=0}^{\ell-1} \mathbb{F}_{q^d}[t],$$

which becomes an $\mathbb{F}_{q^d}[t, \tau]$ -module with the τ -action given by

$$\tau \cdot (\bar{m}_0, \bar{m}_1, \dots, \bar{m}_{\ell-1}) = \left((t - \varepsilon) \cdot \bar{m}_{\ell-1}^{(1)}, \bar{m}_0^{(1)}, \dots, \bar{m}_{\ell-2}^{(1)} \right),$$

where we note that $\theta \equiv \varepsilon \pmod{v}$.

Definition 3.3.3. (See [HaKi20, Definition 3.5.14]) *The v -adic crystalline module of the t -motive $\mathbf{R}_\tau(\mathbf{C}_\ell)$ is*

$$H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), \mathbb{F}_{q^d}((T_v))) := \mathbb{F}_{q^d}((T_v)) \otimes_{\mathbb{F}_{q^d}[t]} \left(\tau^d \cdot \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} \right).$$

Remark 3.3.4. We have a natural τ^d -action on $H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), \mathbb{F}_{q^d}((T_v)))$ as follows:

$$\begin{aligned} \tau^d \cdot \left(\left(\sum_i f_i T_v^i \right) \otimes m \right) &:= \left(\sum_i f_i^{q^d} T_v^i \right) \otimes \tau^d m \\ &= \left(\sum_i f_i T_v^i \right) \otimes \tau^d m, \quad \forall \sum_i f_i T_v^i \in \mathbb{F}_{q^d}((T_v)), \quad m \in \tau^d \cdot \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)}. \end{aligned}$$

(The pair $(H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), \mathbb{F}_{q^d}((T_v))), \tau^d)$ is called the T_v -isocrystal of $\mathbf{R}_\tau(\mathbf{C}_\ell)$, see [HaKi20, Example 3.5.7].)

3.4. **de Rham module of $\mathbf{R}_\tau(\mathbf{C}_\ell)$.** Let $\mathcal{R}_\tau(\mathbf{C}_\ell)$ be the base change of $\mathbf{R}_\tau(\mathbf{C}_\ell)$ to k , i.e.

$$\mathcal{R}_\tau(\mathbf{C}_\ell) = k \otimes_A \mathbf{R}_\tau(\mathbf{C}_\ell),$$

and the τ -action is extended by

$$\tau \cdot (\alpha \otimes m) := \alpha^q \otimes (\tau \cdot m), \quad \forall \alpha \in k, \quad m \in \mathbf{R}_\tau(\mathbf{C}_\ell).$$

The *de Rham module of $\mathbf{R}_\tau(\mathbf{C}_\ell)$ over k* is defined as follows:

$$(3.4.1) \quad H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k) := \frac{\langle \tau \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k}.$$

Here $\langle \tau \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k$ is the k -subspace of $\mathcal{R}_\tau(\mathbf{C}_\ell)$ spanned by $\tau \mathbf{R}_\tau(\mathbf{C}_\ell)$.

Remark 3.4.2. The definition of the de Rham module follows from Hartl–Juschka [HaJu20, §2.3.5]. Regarding $\mathcal{R}_\tau(\mathcal{C}_\ell)$ as the t -motive associated to the “Carlitz module of degree ℓ ” over k , the de Rham module of $\mathcal{R}_\tau(\mathcal{C}_\ell)$ can be defined alternatively via the theory of biderivations developed by Anderson, Deligne, Gekeler, and Yu for Drinfeld modules (see [Ge89], and [Yu90]), and extended by Brownawell–Papanikolas for higher dimensional t -modules (see [BP02]). For the equivalence between these two definitions, we refer the reader to [NP22, Prop. 4.1.3] for a detailed comparison.

Lemma 3.4.3. *The natural inclusion $\tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \subset \tau \mathcal{R}_\tau(\mathcal{C}_\ell)$ induces the k -linear isomorphism*

$$\frac{\langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k} \xrightarrow{\sim} H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k).$$

Proof. From the description of the τ^d -action on $\mathcal{R}_\tau(\mathcal{C}_\ell)$ in Lemma 3.1.4, we observe that

$$\langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k \supset \left(\prod_{i=1}^{d-1} (t - \theta^{q^i}) \right) \cdot \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k.$$

By Chinese Remainder Theorem, we get

$$\begin{aligned} & (t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k + \langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k \\ & \supset (t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k + \left(\prod_{i=1}^{d-1} (t - \theta^{q^i}) \right) \cdot \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k \\ & = \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k, \end{aligned}$$

whence the canonical map

$$\langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k \longrightarrow \frac{\langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k} = H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k)$$

is surjective with kernel $(t - \theta) \cdot \langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k$. □

Consider the following base change:

$$H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) := k_v \otimes_k H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k),$$

called the *de Rham module of $\mathcal{R}_\tau(\mathcal{C}_\ell)$ over k_v* . Also, we put

$$(3.4.4) \quad H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v[[T_v - v]]) := k_v[[T_v - v]] \otimes_{k[t]} \langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k.$$

Then the above lemma leads to the following k_v -linear isomorphism

$$(3.4.5) \quad \frac{H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v[[T_v - v]])}{(T_v - v) \cdot H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v[[T_v - v]])} \xrightarrow{\sim} H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v).$$

This will be used in the crystalline–de Rham comparison isomorphism in the next subsection.

3.5. Crystalline–de Rham comparison isomorphism. In order to describe the crystalline–de Rham comparison isomorphism, we recall in Section 3.2.3 that following Hartl–Kim, we define

$$A_v[[T_v, T_v^{-1}]] := \left\{ \sum_{i \in \mathbb{Z}} b_i T_v^i \in A_v[[T_v, T_v^{-1}]] \mid \lim_{i \rightarrow -\infty} b_i v^{-ni} = 0, \quad \forall n > 0 \right\}.$$

We further consider the v -adic Anderson's scattering (cf. [ABP04, GL11, Har11], and [HaKi20, (3.5.3)])

$$\varpi(T_v) := \prod_{i \in \mathbb{Z}_{\geq 0}} \left(1 - \frac{v^{q^{di}}}{T_v}\right) \in A_v\{\{T_v^{-1}\}\} := A_v[[T_v^{-1}]] \cap A_v[[T_v, T_v^{-1}]].$$

Lemma 3.5.1. *There exists a unique isomorphism*

$$A_v[[T_v, T_v^{-1}]]\{\{\varpi(T_v)^{-1}\}\} \otimes_{\mathbb{F}_q[t]} \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} \xrightarrow{\sim} A_v[[T_v, T_v^{-1}]]\{\{\varpi(T_v)^{-1}\}\} \otimes_{\mathbb{F}_q[\theta][t]} \mathbf{R}_\tau(\mathbf{C}_\ell),$$

which is compatible with the τ^d -action on both sides and becomes the identity map modulo \mathfrak{m}_v .

The above lemma actually holds for other t -motives (in a more general setting), see [HaKi20, Lemma 3.5.8], [GL11, Lemma 7.4], and [Har11, Lemma 2.3.1]. We illustrate the isomorphism in Lemma 3.5.1 in concrete terms as follows. Recall $\theta_v := \theta - \varepsilon$. We first observe that:

Lemma 3.5.2. *We identify t with the power series $t(T_v)$ in (3.2.1). For every non-negative integer i and positive integer ℓ , we define*

$$\Omega_{\ell, v, i}(T_v) := \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right).$$

Then we have

$$\Omega_{\ell, v, i}(T_v) \in A_v[[T_v, T_v^{-1}]] \cap (A_v[[T_v, T_v^{-1}]]\{\{\varpi(T_v)^{-1}\}\})^\times.$$

Proof. Given a non-negative integer i , from the T_v -expansion of t in (3.2.1) and $t \equiv \varepsilon \pmod{T_v}$ we get

$$(3.5.3) \quad t - \varepsilon^i \in \begin{cases} \left(\prod_{j=1}^{d-1} (\varepsilon - \varepsilon^{q^j})^{-1}\right) \cdot T_v \cdot (1 + T_v A_v[[T_v]]), & \text{if } i \equiv 0 \pmod{d}, \\ (\varepsilon - \varepsilon^{q^i}) + T_v A_v[[T_v]], & \text{if } i \not\equiv 0 \pmod{d}. \end{cases}$$

Writing

$$\Omega_{\ell, v, i}(T_v) = \prod_{\substack{n \geq 0 \\ \ell n + i \equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right) \cdot \prod_{\substack{n \geq 0 \\ \ell n + i \not\equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right),$$

from (3.5.3) one has that

$$(3.5.4) \quad \prod_{\substack{n \geq 0 \\ \ell n + i \not\equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right) \in 1 + \theta_v A_v[[T_v]] \subset (A_v[[T_v]])^\times \subset (A_v[[T_v, T_v^{-1}]]\{\{\varpi(T_v)^{-1}\}\})^\times.$$

On the other hand, if $\gcd(d, \ell) \nmid i$, then $\ell n + i \not\equiv 0 \pmod{d}$ for every $n \in \mathbb{Z}$. Hence

$$\Omega_{\ell, v, i}(T_v) = \prod_{\substack{n \geq 0 \\ \ell n + i \not\equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right)$$

and the result holds.

Suppose $\gcd(d, \ell) \mid i$. Write $\text{lcm}(d, \ell) = d\ell_0$, and take $s_i \in \mathbb{Z}_{\geq 0}$ to be minimal so that $s_i\ell + i \equiv 0 \pmod{d}$. Put $i' = s_i\ell + i$. Then

$$\prod_{\substack{n \geq 0 \\ \ell n + i \equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n + i}}}{t - \varepsilon^{q^{\ell n + i}}}\right) = \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right).$$

Let

$$g(X) := X^d + \epsilon_1 X^{d-1} + \cdots + \epsilon_{d-1} X + \epsilon_d - v \in \mathbb{F}_q[v][X] \subset \overline{\mathbb{F}_q}[v, X] = \overline{\mathbb{F}_q}[X][v],$$

which is an irreducible element as it is of degree one in v over $\overline{\mathbb{F}_q}[X]$. It follows that $g(X)$ is the minimal polynomial of θ over $\mathbb{F}_{q^d}(v)$. Since

$$\begin{aligned} g(X) &\equiv X^d + \epsilon_1 X^{d-1} + \cdots + \epsilon_{d-1} X + \epsilon_d \pmod{v} \\ &= \prod_{i=0}^{d-1} (X - \varepsilon^{q^i}) \in \mathbb{F}_{q^d}[X], \end{aligned}$$

by Hensel's lemma we get that $g(X)$ splits completely in $\mathbb{F}_{q^d}[[v]][X]$. We denote by $\theta_j \in \mathbb{F}_{q^d}[[v]] = A_v$, $0 \leq j < d$, the roots of $g(X)$ satisfying $\theta_j \equiv \varepsilon^{q^j} \pmod{\mathfrak{m}_v}$, and take $u_j \in A_v$ so that $u_j \theta_v = \theta_j - \varepsilon^{q^j} \in \mathfrak{m}_v$. Note that $\theta_0 = \theta - \varepsilon = \theta_v$ and $u_0 = 1$. From the equalities

$$(3.5.5) \quad T_v - v = (t^d + \epsilon_1 t^{d-1} + \cdots + \epsilon_{d-1} t + \epsilon_d) - v = g(t) = (t - \theta) \cdot \prod_{j=1}^{d-1} (t - \theta_j)$$

and

$$(3.5.6) \quad T_v = \prod_{j=0}^{d-1} (t - \varepsilon^{q^j}),$$

we obtain that

$$(3.5.7) \quad h(T_v) := \frac{1 - \frac{v}{T_v}}{1 - \frac{\theta_v}{t - \varepsilon}} = \frac{t - \varepsilon}{T_v} \cdot \frac{T_v - v}{t - \theta} = \prod_{j=1}^{d-1} \frac{t - \theta_j}{t - \varepsilon^{q^j}} = \prod_{j=1}^{d-1} \left(1 - \frac{u_j \theta_v}{t - \varepsilon^{q^j}}\right) \in 1 + \theta_v A_v[[T_v]],$$

where the third identity comes from (3.5.5) and (3.5.6). Considering the Frobenius twists of the identity

$$1 - \frac{\theta_v}{t - \varepsilon} = \left(1 - \frac{v}{T_v}\right) \cdot h(T_v)^{-1},$$

we obtain that (as $i' \equiv 0 \pmod{d}$)

$$(3.5.8) \quad \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right) = \prod_{n=0}^{\infty} \left(1 - \frac{v^{q^{d\ell_0 n + i'}}}{T_v}\right) \cdot \prod_{n=0}^{\infty} h^{(d\ell_0 n + i')}(T_v)^{-1} \in A_v[[T_v, T_v^{-1}]].$$

Finally, since

$$\begin{aligned}
& \prod_{n=0}^{\infty} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) = \varpi(T_v) \in (A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}])^{\times} \\
&= \left(\prod_{\substack{n \in \mathbb{Z}_{\geq 0} \\ dn \not\equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \prod_{\substack{0 \leq n < i'/d \\ dn \equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \right) \cdot \left(\prod_{n=0}^{\infty} \left(1 - \frac{v^{q^{d\ell_0 n + i'}}}{T_v}\right) \right) \\
&= \left(\prod_{\substack{n \in \mathbb{Z}_{\geq 0} \\ dn \not\equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \prod_{\substack{0 \leq n < i'/d \\ dn \equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \right) \cdot \left(\prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right) \cdot h^{(d\ell_0 n + i')}(T_v) \right) \\
&= \left(\prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right) \right) \cdot \left(\prod_{\substack{n \in \mathbb{Z}_{\geq 0} \\ dn \not\equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \prod_{\substack{0 \leq n < i'/d \\ dn \equiv i' \pmod{d\ell_0}}} \left(1 - \frac{v^{q^{dn}}}{T_v}\right) \prod_{n=0}^{\infty} h^{(d\ell_0 n + i')}(T_v) \right),
\end{aligned}$$

one gets

$$\prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right) \in (A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}])^{\times}$$

because of (3.5.7). Together with (3.5.4) and (3.5.8), we obtain that

$$\begin{aligned}
\Omega_{\ell, v, i}(T_v) &= \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell_0 n + i'}}}{t - \varepsilon}\right) \cdot \prod_{\substack{n \geq 0 \\ \ell n + i \not\equiv 0 \pmod{d}}} \left(1 - \frac{\theta_v^{q^{\ell n + i}}}{t - \varepsilon^{q^{\ell n + i}}}\right) \\
&\in A_v[[T_v, T_v^{-1}]] \cap (A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}])^{\times}
\end{aligned}$$

as desired. \square

3.5.9. *Description of the isomorphism in Lemma 3.5.1.* From the identification of $\overline{\mathbf{R}_{\tau}(\mathbf{C}_{\ell})}$ in (3.3.2), we get

$$\begin{aligned}
& A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_{q^d}[t]} \overline{\mathbf{R}_{\tau}(\mathbf{C}_{\ell})} \\
&= \prod_{i=0}^{\ell-1} A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_{q^d}[t]} \mathbb{F}_{q^d}[t] \\
&\cong \prod_{i=0}^{\ell-1} A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}].
\end{aligned}$$

On the other hand, by (3.1.2) we have that

$$\begin{aligned}
& A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_q[\theta][t]} \mathbf{R}_{\tau}(\mathbf{C}_{\ell}) \\
&= \prod_{i=0}^{\ell-1} A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_q[\theta][t]} \mathbb{F}_q[\theta][t]. \\
&\cong \prod_{i=0}^{\ell-1} A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}].
\end{aligned}$$

Then the isomorphism in Lemma 3.5.1 is given as follows: every $(m_0, \dots, m_{\ell-1})$ in $\prod_{i=0}^{\ell-1} A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}]$ is sent to

$$(3.5.10) \quad (m_0 \Omega_{\ell,v,0}, m_1 \Omega_{\ell,v,1}, \dots, m_{\ell-1} \Omega_{\ell,v,\ell-1}).$$

To verify this, note that

$$\Omega_{\ell,v,i}(T_v) \equiv 1 \pmod{\mathfrak{m}_v} \in \frac{A_v[[T_v, T_v^{-1}]]}{\mathfrak{m}_v \cdot A_v[[T_v, T_v^{-1}]]} \cong \mathbb{F}_{q^d}((T_v)).$$

Lemma 3.5.2 ensures the bijectivity of (3.5.10), which becomes the identity map modulo \mathfrak{m}_v . Moreover, for each integer i with $0 \leq i < \ell$, write $i + d = i_0 + n'_i \ell$ where i_0, n'_i are unique integers with $0 \leq i_0 < \ell$ and $n'_i \geq 0$. Then

$$\begin{aligned} \Omega_{\ell,v,i}^{(d)}(T_v) &= \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i+d}}}{t - \varepsilon^{q^{\ell n+i+d}}}\right) = \prod_{n=n'_i}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i_0}}}{t - \varepsilon^{q^{\ell n+i_0}}}\right) \\ &= \left(\prod_{n=0}^{n'_i-1} \frac{t - \varepsilon^{q^{\ell n+i_0}}}{t - \theta^{q^{\ell n+i_0}}} \right) \cdot \Omega_{\ell,v,i_0}(T_v) \in A_v[[T_v, T_v^{-1}]], \end{aligned}$$

where the product $\left(\prod_{n=0}^{n'_i-1} \frac{t - \varepsilon^{q^{\ell n+i_0}}}{t - \theta^{q^{\ell n+i_0}}} \right)$ is defined to be 1 whenever $n'_i = 0$. We recall the notation i_1 given in Definition 3.1.3. Since $n'_i = n_{i_0}$ and $(i_0)_1 = i$, one has that for each $0 \leq i < \ell$,

$$\left(\prod_{n=0}^{n_i-1} (t - \theta^{q^{\ell n+i}}) \right) \cdot \Omega_{\ell,v,i_1}^{(d)}(T_v) = \left(\prod_{n=0}^{n_i-1} (t - \varepsilon^{q^{\ell n+i}}) \right) \cdot \Omega_{\ell,v,i}(T_v) \in A_v[[T_v, T_v^{-1}]].$$

Finally, for every $(m_0, \dots, m_{\ell-1}) \in A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_{q^d}[t]} \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)}$, by Lemma 3.1.4 we obtain that

$$\begin{aligned} &\tau^d \cdot (m_0 \Omega_{\ell,v,0}, \dots, m_{\ell-1} \Omega_{\ell,v,\ell-1}) \quad (\in A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_q[\theta][t]} \mathbf{R}_\tau(\mathbf{C}_\ell)) \\ &= \left(\left(\prod_{n=0}^{n_0-1} (t - \theta^{q^{\ell n}}) \right) \cdot m_{(0)_1}^{(d)} \Omega_{\ell,v,(0)_1}^{(d)}, \dots, \left(\prod_{n=0}^{n_{\ell-1}-1} (t - \theta^{q^{\ell n+\ell-1}}) \right) \cdot m_{(\ell-1)_1}^{(d)} \Omega_{\ell,v,(\ell-1)_1}^{(d)} \right) \\ &= (m'_0 \Omega_{\ell,v,0}, \dots, m'_{\ell-1} \Omega_{\ell,v,\ell-1}), \end{aligned}$$

where

$$\begin{aligned} (m'_0, \dots, m'_{\ell-1}) &= \left(\left(\prod_{n=0}^{n_0-1} (t - \varepsilon^{q^{\ell n}}) \right) \cdot m_{(0)_1}^{(d)}, \dots, \left(\prod_{n=0}^{n_{\ell-1}-1} (t - \theta^{q^{\ell n+\ell-1}}) \right) \cdot m_{(\ell-1)_1}^{(d)} \right) \\ &= \tau^d \cdot (m_0, \dots, m_{\ell-1}) \quad (\in A_v[[T_v, T_v^{-1}]][\varpi(T_v)^{-1}] \otimes_{\mathbb{F}_{q^d}[t]} \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)}). \end{aligned}$$

Therefore the constructed isomorphism in (3.5.10) indeed preserves the τ^d -action on both sides. \square

From the embedding $\mathbb{F}_{q^d}((T_v)) \hookrightarrow k_v[[T_v - v]]$ sending T_v to $v + (T_v - v)$, we view $\mathbb{F}_{q^d}((T_v))$ as a subfield of $k_v[[T_v - v]]$, and set

$$\begin{aligned} H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]) &:= k_v[[T_v - v]] \otimes_{\mathbb{F}_{q^d}((T_v))} H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), \mathbb{F}_{q^d}((T_v))) \\ &= k_v[[T_v - v]] \otimes_{\mathbb{F}_{q^d}[t]} \left(\tau^d \cdot \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} \right). \end{aligned}$$

On the other hand, recall that in (3.4.4) we set

$$H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]) := k_v[[T_v - v]] \otimes_{k[t]} \langle \tau^d \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k.$$

Regarding $A_v[[T_v, T_v^{-1}]] \{\varpi^{(d)}(T_v)^{-1}\}$ as a subring of $k_v[[T_v - v]]$ (see [HaKi20, (3.5.7)]), Lemma 3.5.1 induces the following crystalline–de Rham comparison isomorphism (see [HaJu20, Theorem 3.5.18]):

Theorem 3.5.11. *Let notation be given as above. Then we have the following isomorphism as $k_v[[T_v - v]]$ -modules:*

$$\Phi_v : H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]) \cong H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]).$$

Remark 3.5.12. Put

$$\begin{aligned} H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v) &:= k_v \otimes_{\theta \leftarrow t, \mathbb{F}_{q^d}[t]} \left(\tau^d \cdot \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} \right) \\ &= k_v \otimes_{v \leftarrow T_v, \mathbb{F}_{q^d}((T_v))} H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), \mathbb{F}_{q^d}((T_v))) \\ &\cong \frac{H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]])}{(T_v - v) \cdot H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]])}. \end{aligned}$$

As (3.4.5) gives us

$$\frac{H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]])}{(T_v - v) \cdot H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]])} \cong H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v),$$

the isomorphism Φ_v in Theorem 3.5.11 induces a k_v -linear isomorphism

$$(3.5.13) \quad \phi_v : H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v) \cong H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v).$$

3.6. Crystalline–de Rham period matrix. We will choose suitable “algebraic bases” and determine the crystalline–de Rham period matrix associated to the isomorphism ϕ_v in (3.5.13).

First, for each integer i with $0 \leq i < \ell$, recall that $\Omega_{\ell, v, i}(T_v) \in A_v[[T_v, T_v^{-1}]]$ is introduced in Lemma 3.5.2. Recall the notation given in Definition 3.1.3, n_i is the smallest non-negative integer so that $i + n_i \ell \geq d$, and $i_1 = i + n_i \ell - d$. In the case of $n_i > 0$, we have

$$\left(\prod_{n=0}^{n_i-1} (t - \theta^{q^{\ell n+i}}) \right) \cdot \Omega_{\ell, v, i_1}^{(d)} = \left(\prod_{n=0}^{n_i-1} (t - \varepsilon^{\ell n+i}) \right) \cdot \Omega_{\ell, v, i}.$$

Moreover, by Lemma 3.1.4 we have the following rank- ℓ module

$$\langle \tau^d \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k = \prod_{i=0}^{\ell-1} \left(\prod_{n=0}^{n_i-1} (t - \theta^{q^{i+n\ell}}) \right) \cdot k[t],$$

and the same arguments show that

$$\tau^d \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} = \prod_{i=0}^{\ell-1} \left(\prod_{n=0}^{n_i-1} (t - \varepsilon^{q^{i+n\ell}}) \right) \cdot \mathbb{F}_{q^d}[t].$$

It follows that

$$\begin{aligned} H_{\text{cris}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]) &= k_v[[T_v - v]] \otimes_{\mathbb{F}_q[t]} \left(\tau^d \overline{\mathbf{R}_\tau(\mathbf{C}_\ell)} \right) \\ &= \prod_{i=0}^{\ell-1} \left(\prod_{n=0}^{n_i-1} (t - \varepsilon^{q^{i+n\ell}}) \right) \cdot k_v[[T_v - v]] \end{aligned}$$

and

$$\begin{aligned} H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k_v[[T_v - v]]) &= k_v[[T_v - v]] \otimes_{k[t]} \langle \tau^d \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k \\ &= \prod_{i=0}^{\ell-1} \left(\prod_{n=0}^{n_i-1} (t - \theta^{q^{i+n\ell}}) \right) \cdot k_v[[T_v - v]]. \end{aligned}$$

The induced crystalline–de Rham comparison isomorphism Φ_v in Theorem 3.5.11 is given explicitly as follows:

$$\begin{aligned} \Phi_v \left(\left(\prod_{n=0}^{n_0-1} (t - \varepsilon^{q^{n\ell}}) \right) m_0, \dots, \left(\prod_{n=0}^{n_{\ell-1}-1} (t - \varepsilon^{q^{\ell-1+n\ell}}) \right) m_{\ell-1} \right) \\ &= \left(\Omega_{\ell,v,0} \cdot \left(\prod_{n=0}^{n_0-1} (t - \varepsilon^{q^{n\ell}}) \right) m_0, \dots, \Omega_{\ell,v,\ell-1} \cdot \left(\prod_{n=0}^{n_{\ell-1}-1} (t - \varepsilon^{q^{\ell-1+n\ell}}) \right) m_{\ell-1} \right) \\ (3.6.1) \quad &= \left(\Phi_{v,0} \cdot \left(\prod_{n=0}^{n_0-1} (t - \theta^{q^{n\ell}}) \right) m_0, \dots, \Phi_{v,\ell-1} \cdot \left(\prod_{n=0}^{n_{\ell-1}-1} (t - \theta^{q^{\ell-1+n\ell}}) \right) m_{\ell-1} \right), \end{aligned}$$

where $\Phi_{v,i}(T_v)$ are given by infinite products

$$\Phi_{v,i} = \left(\prod_{n=0}^{n_i-1} \frac{t - \varepsilon^{q^{n\ell+i}}}{t - \theta^{q^{n\ell+i}}} \right) \cdot \Omega_{\ell,v,i}(T_v) = \prod_{n=n_i}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}} \right), \quad 0 \leq i < \ell.$$

In particular, by (3.2.2), evaluating $\Phi_{v,i}(T_v)$ at $T_v = v$ equals

$$\Phi_{v,i}(T_v)|_{T_v=v} = \left(\prod_{n=n_i}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}} \right) \right) \Big|_{t=\theta} = \prod_{n=n_i}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{\theta - \varepsilon^{q^{\ell n+i}}} \right), \quad 0 \leq i < \ell.$$

Via the identification (3.1.2), we take $\mathfrak{B} = \{b_0, \dots, b_{\ell-1}\}$ to be the standard $\mathbb{F}_q[\theta][t]$ basis of $\mathbf{R}_\tau(\mathbf{C}_\ell)$, i.e. for $0 \leq i < \ell$,

$$(3.6.2) \quad b_i = (b_{i,0}, \dots, b_{i,\ell-1}) \in \mathbf{R}_\tau(\mathbf{C}_\ell) \quad \text{with} \quad b_{i,j} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{otherwise.} \end{cases}$$

For every integer s with $0 \leq s < \ell$, recall that n_s denotes the smallest non-negative integer with $s + n_s \cdot \ell \geq d$ and $s_1 = s + n_s \cdot \ell - d$. Then by Lemma 3.1.4 we get

$$(3.6.3) \quad b_{v,s} := \tau^d \cdot b_{s_1} = (b'_{s,0}, \dots, b'_{s,\ell-1}) \quad \text{where} \quad b'_{s,j} = \begin{cases} \prod_{n=0}^{n_s-1} (t - \theta^{q^{\ell n+s}}), & \text{if } j = s, \\ 0, & \text{otherwise.} \end{cases}$$

Recall in Lemma 3.4.3 that the inclusion $\tau^d \mathbf{R}_\tau(\mathbf{C}_\ell) \subset \tau \mathbf{R}_\tau(\mathbf{C}_\ell)$ induces a surjective map

$$\langle \tau^d \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k \twoheadrightarrow \frac{\langle \tau \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau \mathbf{R}_\tau(\mathbf{C}_\ell) \rangle_k} = H_{\text{dR}}(\mathbf{R}_\tau(\mathbf{C}_\ell), k).$$

Set

$$(3.6.4) \quad \omega_{v,s} := b_{v,s} \pmod{(t-\theta)} \in \frac{\langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k}{(t-\theta) \cdot \langle \tau^d \mathcal{R}_\tau(\mathcal{C}_\ell) \rangle_k} \cong H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k).$$

Then $\beta_v := \{\omega_{v,0}, \dots, \omega_{v,\ell-1}\}$ is a k -basis of $H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k)$. On the other hand, for each $0 \leq s < \ell$, put

$$(3.6.5) \quad \lambda_{v,s} := b_{v,s} \pmod{\mathfrak{m}_v} \in \tau^d \cdot \overline{\mathcal{R}_\tau(\mathcal{C}_\ell)}.$$

Then $\Lambda_v := \{\lambda_{v,0}, \dots, \lambda_{v,\ell-1}\}$ gives a k_v -basis of $H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v)$ through the identification in Remark 3.5.12. From the description of Φ_v in (3.6.1), we then obtain that:

Theorem 3.6.6. *For every integer s with $0 \leq s < \ell$, we have*

$$\phi_v(\lambda_{v,s}) = \prod_{n=n_s}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{\ell n+s}}}{\theta - \varepsilon^{q^{\ell n+s}}}\right) \cdot \omega_{v,s}.$$

Here n_s is the smallest non-negative integer so that $s + n_s \ell \geq d$. Consequently, the period matrix of isomorphism ϕ_v in (3.5.13) with respect to β_v and Λ_v is

$$\begin{pmatrix} \prod_{n=n_0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n}}}{\theta - \varepsilon^{q^{\ell n}}}\right) & & & 0 \\ & \ddots & & \\ & & \prod_{n=n_{\ell-1}}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+\ell-1}}}{\theta - \varepsilon^{q^{\ell n+\ell-1}}}\right) & \\ 0 & & & \end{pmatrix}.$$

3.7. Chowla–Selberg-type formula. Note that the action of $\tau_v := \tau^d$ on the T_v -isocrystal $(H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), \mathbb{F}_{q^d}((T_v))), \tau_v)$ introduced in Remark 3.3.4 is actually $\mathbb{F}_{q^d}((T_v))$ -linear. Through the isomorphism

$$k_v \otimes_{v \leftarrow T_v, \mathbb{F}_{q^d}((T_v))} H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), \mathbb{F}_{q^d}((T_v))) = H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) \xrightarrow{\phi_v} H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v),$$

there exists a k_v -linear automorphism ϱ_v on $H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v)$ satisfying

$$(3.7.1) \quad \varrho_v \cdot \omega = \phi_v \left(\tau_v \cdot \phi_v^{-1}(\omega) \right), \quad \forall \omega \in H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v).$$

That is, ϱ_v is the existence of k_v -linear automorphism so that the following diagram commutes:

$$\begin{array}{ccc} H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) & \xrightarrow{\phi_v} & H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) \\ \downarrow \tau_v & & \downarrow \varrho_v \\ H_{\text{cris}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) & \xrightarrow{\phi_v} & H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v) \end{array}$$

From Theorem 3.6.6, we are able to realize the ϱ_v -action on $H_{\text{dR}}(\mathcal{R}_\tau(\mathcal{C}_\ell), k_v)$ via v -adic arithmetic gamma values in the following.

For each integer s with $0 \leq s < \ell$, recall that n_s is the smallest non-negative integer so that $s + n_s \ell \geq d$, and we write

$$s + d = s_0 + n'_s \ell (\geq d), \quad \text{where } s_0, n'_s \in \mathbb{Z} \text{ and } 0 \leq s_0 < \ell.$$

From the inequalities

$$\ell + d > s + d = s_0 + n'_s \ell \geq s_0 + n_{s_0} \ell \geq d,$$

we get

$$(3.7.2) \quad n'_s = n_{s_0}.$$

By using argument similar to Lemma 3.1.4, we have

$$\tau^d \cdot \lambda_{v,s} = \left(\prod_{n=0}^{n_s-1} (t - \varepsilon^{q^{s+n\ell}}) \right) \cdot \lambda_{v,s_0},$$

and hence Theorem 3.6.6 implies the following identity:

$$(3.7.3) \quad \varrho_v \cdot \omega_{v,s} = \left(\frac{\prod_{n=n_{s_0}}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n + s_0}}}{\theta - \varepsilon^{q^{\ell n + s_0}}}\right)}{\prod_{n=n_s}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n + s}}}{\theta - \varepsilon^{q^{\ell n + s}}}\right)} \right) \cdot \left(\prod_{n=0}^{n_s-1} (\theta - \varepsilon^{q^{s+n\ell}}) \right) \cdot \omega_{v,s_0}$$

Finally, recall from the definition that

$$H_{\text{dR}}(\mathcal{R}_\tau(\mathbf{C}_\ell), k) = \frac{\langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k}.$$

By (3.1.2) we may identify

$$\langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k = (t - \theta) \cdot k[t] \times \prod_{i=1}^{\ell-1} k[t].$$

For $0 \leq s < \ell$, let $m_s = ((t - \theta)m_{s,0}, m_{s,1}, \dots, m_{s,\ell-1}) \in \langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k$, where $m_{s,j} = 1$ if $s = j$ or 0 otherwise, and set

$$(3.7.4) \quad \omega_s := m_s \pmod{(t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k} \in \frac{\langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k}{(t - \theta) \cdot \langle \tau \mathcal{R}_\tau(\mathbf{C}_\ell) \rangle_k} = H_{\text{dR}}(\mathcal{R}_\tau(\mathbf{C}_\ell), k).$$

Then the description of $b_{v,s}$ in (3.6.3) says in particular that

$$b_{v,s} = \begin{cases} \left(\prod_{n=1}^{n_s-1} (t - \theta^{q^{s+n\ell}}) \right) \cdot m_s, & \text{if } s = 0, \\ \left(\prod_{n=0}^{n_s-1} (t - \theta^{q^{s+n\ell}}) \right) \cdot m_s, & \text{if } s > 0. \end{cases}$$

for which we get

$$(3.7.5) \quad \omega_{v,s} = \begin{cases} \left(\prod_{n=1}^{n_s-1} (\theta - \theta^{q^{s+n\ell}}) \right) \cdot \omega_s, & \text{if } s = 0, \\ \left(\prod_{n=0}^{n_s-1} (\theta - \theta^{q^{s+n\ell}}) \right) \cdot \omega_s, & \text{if } s > 0. \end{cases}$$

By (3.7.3), we obtain that

$$\begin{aligned}
 \varrho_v \cdot \omega_s &= \left(\frac{\prod_{n=n_{s_0}}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+s_0}}}{\theta - \varepsilon^{q^{\ell n+s_0}}}\right)}{\prod_{n=1}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+s}}}{\theta - \varepsilon^{q^{\ell n+s}}}\right)} \right) \cdot \alpha_s^{-1} \cdot \omega_{v,s_0} \\
 (3.7.6) \quad &= \left(\frac{\prod_{n=1}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+s_0}}}{\theta - \varepsilon^{q^{\ell n+s_0}}}\right)}{\prod_{n=1}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+s}}}{\theta - \varepsilon^{q^{\ell n+s}}}\right)} \right) \cdot \left(\prod_{n=0}^{n_{s_0}-1} (\theta - \varepsilon^{q^{s_0+n\ell}}) \right) \cdot \frac{\alpha_{s_0}}{\alpha_s} \cdot \omega_{s_0},
 \end{aligned}$$

where α_s is given in (2.3.6). As $n_{s_0} = n'_s$ in (3.7.2), combining with Corollary 2.3.5 we arrive at the following v -adic Chowla–Selberg-type formula:

Theorem 3.7.7. *Let $\omega_0, \dots, \omega_{\ell-1}$ be the k -basis of $H_{\text{dR}}(\mathcal{R}_{\tau}(\mathbb{C}_{\ell}), k)$ chosen in (3.7.5). Given an integer s with $0 \leq s < \ell$, write $s + d = s_0 + n'_s \ell$ for unique integers s_0 and n'_s with $0 \leq s_0 < \ell$. Then*

$$\varrho_v \cdot \omega_s = (-1)^{n'_s} \cdot C_s \cdot \frac{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d}}{q^{\ell} - 1} \right\rangle_{\text{ari}}\right)}{\Gamma_{\text{ari},v} \left(1 - \left\langle \frac{q^{s+d-1}}{q^{\ell} - 1} \right\rangle_{\text{ari}}\right)^q} \cdot \omega_{s_0}, \quad \text{where} \quad C_s := \begin{cases} -v, & \text{if } s = 0, \\ 1, & \text{if } s > 0, \end{cases}$$

and $\langle z \rangle_{\text{ari}}$ is the fractional part of every $z \in \mathbb{Q}$, i.e. $0 \leq \langle z \rangle_{\text{ari}} < 1$ and $z - \langle z \rangle_{\text{ari}} \in \mathbb{Z}$.

Remark 3.7.8. Theorem 3.7.7 indicates that the matrix presentation of the automorphism ϱ_v on $H_{\text{dR}}(\mathcal{R}_{\tau}(\mathbb{C}_{\ell}), k_v)$ with respect to the basis $\beta := \{\omega_0, \dots, \omega_{\ell-1}\}$ can be concretely illustrated in terms of v -adic arithmetic gamma values.

Remark 3.7.9. The basis $\beta = \{\omega_0, \dots, \omega_{\ell-1}\}$ of the de Rham module $H_{\text{dR}}(\mathcal{R}_{\tau}(\mathbb{C}_{\ell}), k)$ is actually independent of the chosen finite place v of k . Hence Theorem 3.7.7 shows in particular that with respect to this global basis β , we are able to realize the ‘‘isocrystal’’ action ϱ_v simultaneously for every finite place v of k . In fact, this basis is also utilized implicitly in the ∞ -adic Chowla–Selberg formula in [CPTY10] (see also [Wei22]), and we would have a natural consistency between these formulas (as in [O89, Theorem 3.15] for the number field case).

4. ALGEBRAIC RELATIONS AMONG THE v -ADIC ARITHMETIC GAMMA VALUES

In the remaining sections of this paper, we will determine the algebraic relations among the v -adic arithmetic gamma values.

4.1. Transcendence result of the v -adic arithmetic gamma values. Given $\ell \in \mathbb{N}$, recall in Lemma 3.5.2 that we have defined

$$\Omega_{\ell,v,i}(T_v) = \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{t - \varepsilon^{q^{\ell n+i}}}\right) \in A_v[[T_v, T_v^{-1}]], \quad \forall i \in \mathbb{Z}_{\geq 0}.$$

In particular, one observes that

$$(4.1.1) \quad \begin{aligned} \Omega_{\ell,v,i}(v) &:= \Omega_{\ell,v,i}(T_v) \Big|_{T_v=v} = \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{\theta - \varepsilon^{q^{\ell n+i}}}\right) \in k_v^\times \\ &= \left(1 - \frac{\theta_v^{q^i}}{\theta - \varepsilon^{q^i}}\right) \cdot \Omega_{\ell,v,i+\ell}(v), \quad \forall i \in \mathbb{N}. \end{aligned}$$

We first assert that:

Theorem 4.1.2. *Given $\ell \in \mathbb{N}$, the values*

$$\Omega_{\ell,v,i}(v) = \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n+i}}}{\theta - \varepsilon^{q^{\ell n+i}}}\right), \quad 0 < i \leq \ell,$$

are algebraically independent over \bar{k} .

Remark 4.1.3. By Theorem 3.6.6, the values $\Omega_{\ell,v,i}(v)$, $0 < i \leq \ell$ are (up to k^\times -multiples) the v -adic periods of the crystalline–de Rham comparison isomorphism for the t -motive $\mathbf{R}_\tau(\mathbf{C}_\ell)$. Theorem 4.1.2 implies that

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Omega_{\ell,v,i}(v) \mid 0 < i \leq \ell \right) = \ell = \dim \Gamma_{\mathbf{R}_\tau(\mathbf{C}_\ell)},$$

where $\Gamma_{\mathbf{R}_\tau(\mathbf{C}_\ell)}$ is the “ t -motivic Galois group of the t -motive $\bar{k} \otimes_{\mathbb{F}_q[\theta]} \mathbf{R}_\tau(\mathbf{C}_\ell)$ ” (see [CPTY10, Lemma 3.2.1]). This coincidence seems to provide an evidence for a suitable v -adic analogue of Papanikolas’ theorem in [P08] (Grothendieck period conjecture).

Assuming Theorem 4.1.2, whose proof will be given in Section 6.4.6, we are able to show that:

Theorem 4.1.4. *Given $\ell \in \mathbb{N}$, we have the following equality*

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1) \cdot z \in \mathbb{Z} \right) = \ell - \gcd(\ell, \deg v).$$

Proof. First, assume that $d = \deg v \mid \ell$, and write $\ell = d\ell_0$. If $\ell_0 = 1$, then the result follows from Theorem 2.2.5. Suppose $\ell_0 \geq 2$. For every integer s with $0 \leq s < d\ell_0$, by Lemma 2.3.1 we get

$$\frac{\Gamma_{\text{ari},v}\left(1 - \frac{q^{d\ell_0-s}}{q^{d\ell_0}-1}\right)}{\Gamma_{\text{ari},v}\left(1 - \frac{q^{d\ell_0-s-1}}{q^{d\ell_0}-1}\right)^q} \sim \prod_{n=2}^{\infty} \left(\frac{1 - \frac{\theta_v^{q^{d\ell_0 n-s}}}{\theta - \varepsilon^{q^{-s}}}}{1 - \frac{\theta_v^{q^{d\ell_0 n-s-d}}}{\theta - \varepsilon^{q^{-s}}}} \right) = \frac{\Omega_{d\ell_0,v,2d\ell_0-s}(v)}{\Omega_{d\ell_0,v,2d\ell_0-s-d}(v)} =: u_s.$$

In particular, for every integer s with $0 \leq s < d(\ell_0 - 1)$, let $s_d \in \mathbb{Z}$ with $0 \leq s_d < d$ and $s_d \equiv s \pmod{d}$. Write $s = s_d + d\ell_s$ with $0 \leq \ell_s < \ell_0 - 1$. Then one has that

$$u'_s := u_s u_{s+d} \cdots u_{s+(\ell_0-2-\ell_s)d} = \frac{\Omega_{d\ell_0,v,2d\ell_0-s}(v)}{\Omega_{d\ell_0,v,d\ell_0+d-s_d}(v)} \sim \frac{\Omega_{d\ell_0,v,d\ell_0-s}(v)}{\Omega_{d\ell_0,v,d-s_d}(v)}.$$

As $d < d\ell_0 - s \leq d\ell_0$ and $0 < d - s_d \leq d$, Theorem 4.1.2 ensures the algebraic independence of u'_s over \bar{k} for $0 \leq s < d(\ell_0 - 1)$. Hence

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1) \cdot z \in \mathbb{Z} \right) \geq d(\ell_0 - 1) = \ell - \gcd(\ell, \deg v).$$

By combining with the upper bound in Theorem 2.2.9, the result follows.

Consequently, the following monomial relations

$$(4.1.5) \quad \prod_{m=0}^{\ell_0-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^{j+dm}}{q^{d\ell_0} - 1}\right) = \Gamma_{\text{ari},v}\left(1 - \frac{q^j + q^{j+d} + \dots + q^{j+d(\ell_0-1)}}{q^{d\ell_0} - 1}\right) \\ = \Gamma_{\text{ari},v}\left(1 - \frac{q^j}{q^d - 1}\right), \quad \text{for } 0 \leq j < d,$$

generate all algebraic relations among $\Gamma_{\text{ari},v}(1 - q^s/(q^{d\ell_0} - 1))$ for $0 \leq s < d\ell_0 - 1$. In particular, let $\vec{\delta}_0, \dots, \vec{\delta}_{d-1} \in \mathbb{Z}^{d\ell_0}$ be the ‘‘exponent vectors’’ corresponding to the above monomial relations, i.e.

$$\vec{\delta}_j = (\delta_{j,0}, \dots, \delta_{j,d\ell_0-1}) \quad \text{with} \quad \delta_{j,s} = \begin{cases} 1, & \text{if } j \equiv s \pmod{d}, \\ 0, & \text{otherwise.} \end{cases}$$

Then

$$\prod_{s=0}^{d\ell_0-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^{d\ell_0} - 1}\right)^{\delta_{j,s}} = \Gamma_{\text{ari},v}\left(1 - \frac{q^j}{q^d - 1}\right) \in \bar{k},$$

Moreover, as $\vec{\delta}_0, \dots, \vec{\delta}_{d-1} \in \mathbb{Z}^{d\ell_0}$ span a d -dimensional ‘‘relation subspace’’ in $\mathbb{Q}^{d\ell_0}$, for arbitrary given vectors $\vec{n}_1, \dots, \vec{n}_r \in \mathbb{Z}^{d\ell_0}$ with $\vec{n}_j = (n_{j,0}, \dots, n_{j,d\ell_0-1})$, $j = 1, \dots, r$, we have that

$$(4.1.6) \quad \text{tr. deg}_{\bar{k}} \bar{k} \left(\prod_{0 \leq s < d\ell_0} \Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^{d\ell_0} - 1}\right)^{n_{j,s}} \mid 1 \leq j \leq r \right) \\ = \dim_{\mathbb{Q}} \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{n}_1, \dots, \vec{n}_r \rangle_{\mathbb{Q}} - d.$$

Here $\langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{n}_1, \dots, \vec{n}_r \rangle_{\mathbb{Q}}$ is the subspace of $\mathbb{Q}^{d\ell_0}$ spanned by $\vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{n}_1, \dots, \vec{n}_r$ over \mathbb{Q} .

In the general case, for a given $\ell \in \mathbb{N}$, write $\text{lcm}(\ell, d) = d\ell_0 = d_0\ell$ with $\ell_0, d_0 \in \mathbb{N}$ so that $\text{gcd}(\ell_0, d_0) = 1$. Note that for an integer j with $0 \leq j < \ell$, one has

$$\Gamma_{\text{ari},v}\left(1 - \frac{q^j}{q^\ell - 1}\right) = \prod_{s=0}^{d_0-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^{j+s\ell}}{q^{d_0\ell} - 1}\right).$$

Let $\vec{\lambda}_0, \dots, \vec{\lambda}_{\ell-1} \in \mathbb{Z}^{d\ell_0}$ satisfying that

$$\vec{\lambda}_j = (\lambda_{j,0}, \dots, \lambda_{j,d\ell_0-1}) \quad \text{with} \quad \lambda_{j,s} = \begin{cases} 1, & \text{if } j \equiv s \pmod{\ell}, \\ 0, & \text{otherwise.} \end{cases}$$

As for $0 \leq j < \ell$,

$$\prod_{s=0}^{d\ell_0-1} \Gamma_{\text{ari},v}\left(1 - \frac{q^s}{q^{d\ell_0} - 1}\right)^{\lambda_{j,s}} = \Gamma_{\text{ari},v}\left(1 - \frac{q^j}{q^\ell - 1}\right),$$

by the equality (4.1.6), we obtain that

$$(4.1.7) \quad \text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari},v}(z) \mid z \in \mathbb{Q} \text{ with } (q^\ell - 1) \cdot z \in \mathbb{Z} \right) \\ = \text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari},v}\left(1 - \frac{q^j}{q^\ell - 1}\right) \mid 0 \leq j < \ell \right) \\ = \dim_{\mathbb{Q}} \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{\lambda}_0, \dots, \vec{\lambda}_{\ell-1} \rangle_{\mathbb{Q}} - d.$$

Since

$$\langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1} \rangle_{\mathbb{Q}} = \left\{ (\alpha_0, \dots, \alpha_{d\ell_0-1}) \in \mathbb{Q}^{d\ell_0} \mid \alpha_i = \alpha_j \text{ if } i \equiv j \pmod{d} \right\}$$

and

$$\langle \vec{\lambda}_0, \dots, \vec{\lambda}_{\ell-1} \rangle_{\mathbb{Q}} = \left\{ (\beta_0, \dots, \beta_{d_0\ell_0-1}) \in \mathbb{Q}^{d_0\ell} \mid \beta_i = \beta_j \text{ if } i \equiv j \pmod{\ell} \right\},$$

one has that

$$\begin{aligned} & \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1} \rangle_{\mathbb{Q}} \cap \langle \vec{\lambda}_0, \dots, \vec{\lambda}_{\ell-1} \rangle_{\mathbb{Q}} \\ &= \left\{ (\gamma_0, \dots, \gamma_{d_0\ell_0-1}) \in \mathbb{Q}^{d_0\ell} \mid \gamma_i = \gamma_j \text{ if } i \equiv j \pmod{\gcd(d, \ell)} \right\}. \end{aligned}$$

Therefore

$$\dim_{\mathbb{Q}} \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{\lambda}_0, \dots, \vec{\lambda}_{\ell-1} \rangle_{\mathbb{Q}} = \ell + d - \gcd(d, \ell),$$

for which the result follows from (4.1.7). \square

Remark 4.1.8. Theorem 4.1.4 says that all algebraic relations among v -adic arithmetic gamma values at rational p -adic integers are generated by the functional equations in Proposition 2.2.1 and Thakur's analogue of the Gross–Koblitz formula in Theorem 2.2.5.

One immediate consequence of Theorem 4.1.4 is the following.

Corollary 4.1.9. *Let a and b be nonzero integers so that $\gcd(a, b) = 1$ and $p \nmid b$. If $q^d \not\equiv 1 \pmod{b}$, then*

$$\Gamma_{\text{ari}, v} \left(\frac{a}{b} \right) \quad \text{is transcendental over } k.$$

Proof. Without loss of generality, we may assume that $\frac{a}{b} = 1 - \frac{a'}{b}$ with $0 \leq a' < b$ and $\gcd(a', b) = 1$. Let ℓ_0 be the order of $q^d \pmod{b}$ in $(\mathbb{Z}/b\mathbb{Z})^\times$. Then the assumption of b implies that $\ell_0 > 1$. Write $q^{d\ell_0} - 1 = b \cdot b'$, and take $a_0, \dots, a_{d\ell_0-1} \in \mathbb{Z}$ with $0 \leq a_0, \dots, a_{d\ell_0-1} < q$ satisfying that

$$\frac{a'}{b} = \frac{a'b'}{q^{d\ell_0} - 1} = \sum_{i=0}^{d\ell_0-1} \frac{a_i q^i}{q^{d\ell_0} - 1}.$$

Let $\vec{a} = (a_0, a_1, \dots, a_{d\ell_0-1}) \in \mathbb{Z}^{d\ell_0}$. If $\vec{a} \in \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1} \rangle_{\mathbb{Q}}$, then we get $a_i = a_j$ if $i \equiv j \pmod{d}$. Hence

$$\frac{a'}{b} = \sum_{i=0}^{d-1} \frac{a_i (q^i + q^{d+i} + \dots + q^{(\ell_0-1)d+i})}{q^{d\ell_0} - 1} = \frac{a_0 + a_1 q + \dots + a_{d-1} q^{d-1}}{q^d - 1},$$

for which $b \mid q^d - 1$, a contradiction. Therefore $\vec{a} \notin \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1} \rangle_{\mathbb{Q}}$, and by (4.1.6) we get

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\Gamma_{\text{ari}, v} \left(1 - \frac{a'}{b} \right) \right) = \dim_{\mathbb{Q}} \langle \vec{\delta}_0, \dots, \vec{\delta}_{d-1}, \vec{a} \rangle_{\mathbb{Q}} = (d+1) - d = 1$$

as desired. \square

Remark 4.1.10. Let a and b be nonzero integers so that $\gcd(a, b) = 1$ and $p \nmid b$. Combining Thakur's result in Theorem 2.2.5 and Corollary 4.1.9, we have the following equivalence:

$$b \mid q^{\deg v} - 1 \quad \text{if and only if} \quad \Gamma_{\text{ari}, v} \left(\frac{a}{b} \right) \in \bar{k}^\times.$$

4.2. v -adic arithmetic gamma distribution. Let $\mathbb{Z}_{(p)} := \mathbb{Z}_p \cap \mathbb{Q}$, and \mathcal{A}^{ari} be the free abelian group generated by all elements in $\mathbb{Z}_{(p)}/\mathbb{Z}$. We identify \mathcal{A}^{ari} with a subgroup of $\mathcal{A}_{\mathbb{Q}}^{\text{ari}} := \mathbb{Q} \otimes_{\mathbb{Z}} \mathcal{A}^{\text{ari}}$. Every element in \mathcal{A}^{ari} (resp. $\mathcal{A}_{\mathbb{Q}}^{\text{ari}}$) can be written uniquely as a formal sum

$$\mathbf{z} = \sum_{z \in \mathbb{Z}_{(p)}/\mathbb{Z}} n_z [z], \quad \text{where } n_z \in \mathbb{Z} \text{ (resp. } \mathbb{Q}) \text{ and } n_z = 0 \text{ for almost all } z.$$

Given $\ell \in \mathbb{N}$, let $\mathcal{A}_{\ell}^{\text{ari}}$ (resp. $\mathcal{A}_{\ell, \mathbb{Q}}^{\text{ari}}$) be the subgroup of \mathcal{A}^{ari} (resp. subspace of $\mathcal{A}_{\mathbb{Q}}^{\text{ari}}$) generated by $z \in \frac{1}{q^{\ell-1}}\mathbb{Z}/\mathbb{Z}$. Also, recall that the *arithmetic diamond bracket* of $z \in \mathbb{Z}_{(p)}$, introduced by Thakur [T91], comes from the fractional part of z , i.e. $\langle z \rangle_{\text{ari}} \in \mathbb{Z}_{(p)}$ is the unique number with

$$0 \leq \langle z \rangle_{\text{ari}} < 1 \quad \text{and} \quad z \equiv \langle z \rangle_{\text{ari}} \pmod{\mathbb{Z}}.$$

We may regard $\langle \cdot \rangle_{\text{ari}}$ as a function on $\mathbb{Z}_{(p)}/\mathbb{Z}$. Consider the subspace $\mathcal{R}_{\ell}^{\text{ari}}$ of $\mathcal{A}_{\ell, \mathbb{Q}}^{\text{ari}}$ spanned by

$$[z] - \sum_{i=0}^{\ell-1} z_i \left[\frac{q^i}{1-q^{\ell}} \right], \quad \forall z \in \frac{1}{q^{\ell-1}}\mathbb{Z}/\mathbb{Z} \quad (\text{see (2.2.3)}),$$

where $z_0, \dots, z_{\ell-1} \in \mathbb{Z}$ with $0 \leq z_0, \dots, z_{\ell-1} < q$ so that

$$\sum_{i=0}^{\ell-1} \frac{z_i q^i}{q^{\ell} - 1} = \langle -z \rangle_{\text{ari}}.$$

Put $\mathcal{U}^{\text{ari}} := \mathcal{A}_{\mathbb{Q}}^{\text{ari}} / \mathcal{R}^{\text{ari}}$, where $\mathcal{R}^{\text{ari}} := \cup_{\ell} \mathcal{R}_{\ell}^{\text{ari}}$. We call \mathcal{U}^{ari} the *universal distribution associated to the arithmetic diamond bracket relations* (see [Wei22, §4.3]).

Following the ∞ -adic case, we define

$$\tilde{\Gamma}_{\text{ari}, v} : \mathbb{Z}_{(p)}/\mathbb{Z} \longrightarrow \mathbb{C}_v^{\times}, \quad \tilde{\Gamma}_{\text{ari}, v}(z \pmod{\mathbb{Z}}) := \Gamma_{\text{ari}, v}(1 - \langle -z \rangle_{\text{ari}}),$$

which induces a \mathbb{Q} -linear homomorphism $\hat{\Gamma}_{\text{ari}, v} : \mathcal{U}_{\mathbb{Q}}^{\text{ari}} \rightarrow \mathbb{C}_v^{\times} / \bar{k}^{\times}$ (by the monomial relation in Proposition 2.2.2). We may call $\hat{\Gamma}_{\text{ari}, v}$ the *v -adic arithmetic gamma distribution*. From Corollary 2.2.7 and Theorem 4.1.4 together with (4.1.5), we obtain that:

Theorem 4.2.1. *The kernel of the v -adic arithmetic gamma distribution $\hat{\Gamma}_{\text{ari}, v}$ is spanned by the Gross–Koblitz–Thakur relations*

$$\sum_{i=0}^{r-1} [q^{i \deg v} z], \quad \forall z \in \frac{1}{n}\mathbb{Z}$$

where n runs through all positive integers with $p \nmid n$, and r is the order of $q^{\deg v} \pmod{n}$ in $(\mathbb{Z}/n\mathbb{Z})^{\times}$.

5. TRANSCENDENCE THEORY OF HARTL–JUSCHKA MODULES

In this section, we briefly review the necessary properties in the transcendence theory of Hartl–Juschka modules for the proof of Theorem 4.1.2. To utilize this well-developed theory in the v -adic setting, our initial step is to regard v as the infinite place of the “projective $\frac{1}{v}$ -line” over \mathbb{F}_q .

5.1. **Regarding v as the infinite place.** Fix $d = \deg v$ and a positive integer ℓ as before. Set $k_{d\ell} := \mathbb{F}_{q^{d\ell}}(\theta) = \mathbb{F}_{q^{d\ell}}(\theta_v) \subset \bar{k}$, where $\theta_v = \theta - \varepsilon$, and $k_{d\ell, v} := \mathbb{F}_{q^{d\ell}}((\theta_v))$. Let

$$\tilde{\theta}_v := (\theta - \varepsilon)^{-1} \quad \text{and} \quad \tilde{v} := v^{-1}.$$

Then we have

$$\mathbb{F}_q((v)) = \mathbb{F}_q((\tilde{v}^{-1})) \subset \mathbb{F}_{q^{d\ell}}((\tilde{\theta}_v^{-1})) = \mathbb{F}_{q^{d\ell}}((\theta_v)) = k_{d\ell, v} \subset \mathbb{C}_v.$$


The trick of viewing \mathbb{C}_v as an ‘ $\infty_{\tilde{v}}$ -adic’ field comes from the following translation.

- $\mathbb{F}_q(v) = \mathbb{F}_q(\tilde{v})$ is the fraction field of $\mathbb{F}_q[\tilde{v}]$, the polynomial ring in \tilde{v} over \mathbb{F}_q .
- The $\infty_{\tilde{v}}$ -adic valuation on $\mathbb{F}_q(v) = \mathbb{F}_q(\tilde{v})$ is given by $\text{ord}_{\infty_{\tilde{v}}} := \text{ord}_{\tilde{v}^{-1}} = \text{ord}_v$.
- $\mathbb{F}_q(v) = \mathbb{F}_q(\tilde{v})$ is the function field of the projective \tilde{v} -line over \mathbb{F}_q with $v = \tilde{v}^{-1}$ as a uniformizer of the infinite place.
- $\mathbb{F}_q((v)) = \mathbb{F}_q((\tilde{v}^{-1}))$ is the completion of $\mathbb{F}_q(v) = \mathbb{F}_q(\tilde{v})$ with respect to the $\infty_{\tilde{v}}$ -adic valuation $\text{ord}_{\infty_{\tilde{v}}} = \text{ord}_v$.
- $k_v \cong \mathbb{F}_{q^d}((v)) = \mathbb{F}_{q^d}((\tilde{v}^{-1}))$ is a finite extension of the $\infty_{\tilde{v}}$ -adic field $\mathbb{F}_q((\tilde{v}^{-1}))$.
- A fixed algebraic closure $\overline{\mathbb{F}_{q^d}((v))}$ of $\mathbb{F}_{q^d}((v))$ is the same as a particularly chosen algebraic closure $\overline{\mathbb{F}_q((\tilde{v}^{-1}))}$ of $\mathbb{F}_q((\tilde{v}^{-1}))$.
- \mathbb{C}_v is the v -adic completion of $\overline{\mathbb{F}_{q^d}((v))}$, which is the same as the $\infty_{\tilde{v}}$ -adic completion of $\overline{\mathbb{F}_q((\tilde{v}^{-1}))}$.

Since we deal with values in \mathbb{C}_v , which is viewed as an $\infty_{\tilde{v}}$ -adic field, one can apply Papanikolas’ ‘ ∞ -adic’ transcendence theory [P08]. The following comparison/replacement illustrates why we consider Frobenius modules over $\bar{k}[\tilde{T}_v]$ in the following subsection when applying Papanikolas’ theory fitting into our situation.

Over k_∞		Over k_v	
uniformizer at ∞	$\rightarrow 1/\theta$	\tilde{v}^{-1}	\leftarrow uniformizer at ∞_v
	$t \rightarrow \theta$	\tilde{v}	$\leftarrow \tilde{T}_v$
polynomial ring	$\rightarrow \bar{k}[t]$	$\bar{k}[\tilde{T}_v]$	\leftarrow polynomial ring
Tate algebra	$\rightarrow \mathbb{C}_\infty\{t\}$	$\mathbb{C}_v\{\tilde{T}_v\}$	\leftarrow Tate algebra

The relations of the field extensions to be used is illustrated in the following diagram:



Recall that in Section 3.2 we wrote

$$v = \theta^d + \epsilon_1 \theta^{d-1} + \cdots + \epsilon_{d-1} \theta + \epsilon_d, \quad \text{where } \epsilon_1, \dots, \epsilon_d \in \mathbb{F}_q.$$

As

$$(5.1.1) \quad v = \prod_{i=0}^{d-1} (\theta - \varepsilon^{q^i}) = \prod_{i=0}^{d-1} (\theta_v - (\varepsilon^{q^i} - \varepsilon)),$$

There exists $\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{d-1} \in \mathbb{F}_{q^d}$ such that

$$(5.1.2) \quad v = \theta_v^d + \tilde{\epsilon}_1 \theta_v^{d-1} + \cdots + \tilde{\epsilon}_{d-1} \theta_v \in \mathbb{F}_{q^d}[\theta_v] = \mathbb{F}_{q^d}[\theta].$$

Multiplying $\tilde{v} \tilde{\theta}_v^d$ on both sides of the equation (5.1.2), one gets

$$(5.1.3) \quad \tilde{\theta}_v^d - \left((\tilde{\epsilon}_{d-1} \tilde{v}) \cdot \tilde{\theta}_v^{d-1} + \cdots + (\tilde{\epsilon}_1 \tilde{v}) \tilde{\theta}_v + \tilde{v} \right) = 0.$$

On the other hand, recall in Section 3.2 that $\iota : \mathbb{F}_q(t) \cong \mathbb{F}_q(\theta)$ is the \mathbb{F}_q -algebra isomorphism sending t to θ , and so $\iota(T_v) = v$. Extending ι to an $\mathbb{F}_{q^{d\ell}}$ -algebra isomorphism $\iota : \mathbb{F}_{q^{d\ell}}(t) \cong \mathbb{F}_{q^{d\ell}}(\theta) = k_{d\ell}$, we set

$$\tilde{t}_v := \frac{1}{t - \varepsilon} \quad \text{and} \quad \tilde{T}_v := \frac{1}{T_v}.$$

Then $\iota(\tilde{t}_v) = \tilde{\theta}_v$ and $\iota(\tilde{T}_v) = \tilde{v}$. Moreover, the following equality holds:

$$(5.1.4) \quad \tilde{t}_v^d - (\tilde{\epsilon}_{d-1} \tilde{T}_v) \tilde{t}_v^{d-1} - \cdots - (\tilde{\epsilon}_1 \tilde{T}_v) \tilde{t}_v - \tilde{T}_v = 0.$$

In other words, the minimal polynomial of \tilde{t}_v over $\mathbb{F}_{q^{d\ell}}(\tilde{T}_v)$ is

$$\tilde{g}_{\tilde{T}_v}(X) = X^d - (\tilde{\epsilon}_{d-1} \tilde{T}_v) X^{d-1} - \cdots - (\tilde{\epsilon}_1 \tilde{T}_v) X - \tilde{T}_v \in \mathbb{F}_{q^{d\ell}}[\tilde{T}_v][X],$$

In particular, the field $\mathbb{F}_{q^{d\ell}}(\tilde{t}_v)$ is a finite separable extension of $\mathbb{F}_q(\tilde{T}_v)$ of degree $d^2 \ell$, and the integral closure of $\mathbb{F}_q[\tilde{T}_v]$ in $\mathbb{F}_{q^{d\ell}}(\tilde{t}_v)$ is $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v]$. Indeed, as

$$\tilde{T}_v = \frac{1}{T_v} = \prod_{i=0}^{d-1} \frac{1}{t - \varepsilon^{q^i}} = \tilde{t}_v^d \cdot \frac{1}{S},$$

where $S = \prod_{i=1}^{d-1} (1 - (\varepsilon^{q^i} - \varepsilon) \tilde{t}_v)$ is coprime to \tilde{t}_v^d in $\mathbb{F}_{q^{d\ell}}[\tilde{t}_v]$. Hence

$$(5.1.5) \quad \mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v] = \mathbb{F}_{q^{d\ell}}[\tilde{t}_v]\left[\frac{1}{S}\right],$$

which is a localization of the PID $\mathbb{F}_{q^{d\ell}}[\tilde{t}_v]$. It follows that $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v]$ is a PID, which implies integrally closed property, and hence it equals the integral closure of $\mathbb{F}_q[\tilde{T}_v]$ in $\mathbb{F}_{q^{d\ell}}(\tilde{t}_v)$. Moreover, since $\tilde{g}_{\tilde{T}_v}(X)$ is Eisenstein with respect to the prime ideal (\tilde{T}_v) of $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v]$, the prime ideal (\tilde{T}_v) of $\mathbb{F}_q[\tilde{T}_v]$ is totally ramified in $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v]$, and the unique prime ideal of $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v]$ lying above (\tilde{T}_v) is the principal ideal generated by \tilde{t}_v . Set

$$\tilde{t}_{v,i} := \frac{1}{t - \varepsilon^{q^{-i}}} = \frac{\tilde{t}_v}{1 - (\varepsilon^{q^{-i}} - \varepsilon) \tilde{t}_v}, \quad 0 \leq i < d\ell.$$

One checks that

- (1) $\tilde{\theta}_{v,i} := \iota(\tilde{t}_{v,i}) = (\theta - \varepsilon^{q^{-i}})^{-1}$.
- (2) $\tilde{t}_{v,i} = \tilde{t}_{v,i'}$ if and only if $i \equiv i' \pmod{d}$;
- (3) $\tilde{T}_v = \tilde{t}_{v,0} \cdots \tilde{t}_{v,d-1}$;

(4) The minimal polynomial of $\tilde{t}_{v,i}$ over $\mathbb{F}_{q^{d\ell}}[\tilde{T}_v]$ is

$$(5.1.6) \quad \tilde{g}_{\tilde{T}_v}^{(-i)}(X) = X^d - \left(\tilde{\epsilon}_{d-1}^{q^{-i}} \tilde{T}_v X^{d-1} + \cdots + \tilde{\epsilon}_1^{q^{-i}} \tilde{T}_v \cdot X + \tilde{T}_v \right) \in \mathbb{F}_{q^{d\ell}}[\tilde{T}_v][X];$$

(5) $(\tilde{t}_v) = (\tilde{t}_{v,0}) = (\tilde{t}_{v,1}) = \cdots = (\tilde{t}_{v,d\ell-1}) \subset \mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v]$.

The above setting (on the “ t -variable side”) can be illustrated by the following diagram:

$$\begin{array}{ccccccc} \mathbb{F}_{q^{d\ell}}(t) & = & \mathbb{F}_{q^{d\ell}}(\tilde{t}_v) & \supset & \mathbb{F}_{q^{d\ell}}[\tilde{T}_v][\tilde{t}_v] & \supset & (\tilde{t}_{v,0}) = \cdots = (\tilde{t}_{v,d-1}) \\ | & & | & & | & & | \\ \mathbb{F}_{q^{d\ell}}(T_v) & = & \mathbb{F}_{q^{d\ell}}(\tilde{T}_v) & \supset & \mathbb{F}_{q^{d\ell}}[\tilde{T}_v] & \supset & (\tilde{T}_v) \\ | & & | & & | & & | \\ \mathbb{F}_q(T_v) & = & \mathbb{F}_q(\tilde{T}_v) & \supset & \mathbb{F}_q[\tilde{T}_v] & \supset & (\tilde{T}_v) \end{array}$$

Now, let $\bar{k}(\tilde{T}_v)[\sigma]$ (resp. $\bar{k}(\tilde{t}_v)[\sigma]$) be the twisted polynomial ring over $\bar{k}(\tilde{T}_v)$ (resp. $\bar{k}(\tilde{t}_v)$) satisfying the following multiplication law

$$\sigma \cdot f = f^{(-1)} \cdot \sigma, \quad \forall f \in \bar{k}(\tilde{T}_v) \quad (\text{resp. } \bar{k}(\tilde{t}_v)),$$

where $f^{(n)}$ is the n -th Frobenius twist of f , i.e. raising the coefficients of powers of \tilde{T}_v of f to the q^n -power for each $n \in \mathbb{Z}$.

Lemma 5.1.7. $\bar{k}(\tilde{T}_v)[\sigma]$ is a subring of $\bar{k}(\tilde{t}_v)[\sigma]$ if and only if $\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{d-1} \in \mathbb{F}_q$.

Proof. By definition one observes that $\bar{k}(\tilde{T}_v)[\sigma]$ is a subring of $\bar{k}(\tilde{t}_v)[\sigma]$ if and only if $\tilde{T}_v \cdot \sigma = \sigma \cdot \tilde{T}_v$ in $\bar{k}(\tilde{t}_v)[\sigma]$, which is equivalent to

$$\begin{aligned} & \left(\tilde{t}_v^d - (\tilde{\epsilon}_{d-1} \tilde{T}_v) \tilde{t}_v^{d-1} - \cdots - (\tilde{\epsilon}_1 \tilde{T}_v) \tilde{t}_v \right) \cdot \sigma \\ &= \tilde{T}_v \cdot \sigma = \sigma \cdot \tilde{T}_v \\ &= \sigma \cdot \left(\tilde{t}_v^d - (\tilde{\epsilon}_{d-1} \tilde{T}_v) \tilde{t}_v^{d-1} - \cdots - (\tilde{\epsilon}_1 \tilde{T}_v) \tilde{t}_v \right) \\ &= \left(\tilde{t}_v^d - (\tilde{\epsilon}_{d-1}^{q^{-1}} \tilde{T}_v) \tilde{t}_v^{d-1} - \cdots - (\tilde{\epsilon}_1^{q^{-1}} \tilde{T}_v) \tilde{t}_v \right) \cdot \sigma. \end{aligned}$$

This equality holds if and only if $\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{d-1} \in \mathbb{F}_q$. \square

Remark 5.1.8. The above lemma indicates that $\bar{k}(\tilde{T}_v)[\sigma]$ might not be a subring of $\bar{k}(\tilde{t}_v)[\sigma]$ when $d > 1$. However, both do contain a subring $\bar{k}(\tilde{T}_v)[\sigma^d]$.

In the remaining part of this section, we will review the necessary properties of the Hartl–Juschka category \mathcal{HJ} of over \bar{k} introduced in [HaJu20, Definition 2.4.1] (with respect to the embedding $\iota : \mathbb{F}_q(\tilde{T}_v) \cong \mathbb{F}_q(\tilde{v}) \subset \bar{k} \subset \mathbb{C}_v$), as well as their corresponding “Hodge–Pink cocharacters”.

5.2. Hartl–Juschka Category. The Hartl–Juschka category over \bar{k} is defined as follows: an object of \mathcal{HJ} is a free left $\bar{k}[\tilde{T}_v]$ -module \mathcal{M} of finite rank together with an invertible σ -action on $M := \bar{k}(\tilde{T}_v) \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}$ so that M becomes a left $\bar{k}(\tilde{T}_v)[\sigma]$ -module satisfying that there exists a sufficiently large integer n so that

$$(\tilde{T}_v - \iota(\tilde{T}_v))^n \mathcal{M} \subseteq \sigma \mathcal{M} \subseteq (\tilde{T}_v - \iota(\tilde{T}_v))^{-n} \mathcal{M} \quad (\subset M).$$

We call such an object \mathcal{M} a *Hartl–Juschka module over \bar{k}* . In particular, we say that \mathcal{M} is *effective* if $\sigma \mathcal{M} \subset \mathcal{M}$. A morphism $f : \mathcal{M} \rightarrow \mathcal{M}'$ in \mathcal{HJ} is a left $\bar{k}[\tilde{T}_v]$ -module homomorphism so

that the induced $\bar{k}(\tilde{T}_v)$ -linear homomorphism $f : M = \bar{k}(\tilde{T}_v) \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M} \rightarrow M' = \bar{k}(\tilde{T}_v) \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}'$ satisfies

$$f(\sigma m) = \sigma f(m), \quad \forall m \in M.$$

Given two objects \mathcal{M} and \mathcal{M}' in $\mathcal{H}\mathcal{J}$, the set $\text{Hom}(\mathcal{M}, \mathcal{M}')$ of morphisms from \mathcal{M} to \mathcal{M}' has a natural $\mathbb{F}_q[\tilde{T}_v]$ -module structure, and elements in

$$\text{Hom}^0(\mathcal{M}, \mathcal{M}') := \mathbb{F}_q(\tilde{T}_v) \otimes_{\mathbb{F}_q[\tilde{T}_v]} \text{Hom}(\mathcal{M}, \mathcal{M}')$$

are called *quasi-morphisms from \mathcal{M} to \mathcal{M}'* . The *Hartl-Juschka Category $\mathcal{H}\mathcal{J}^I$ up to isogeny* has the same objects in $\mathcal{H}\mathcal{J}$ but the morphisms in $\mathcal{H}\mathcal{J}^I$ are given by quasi-morphisms.

5.2.1. *Uniformizability.* Let $\mathbb{C}_v\{\tilde{T}_v\}$ be the Tate algebra in the variable \tilde{T}_v over \mathbb{C}_v , i.e.

$$\mathbb{C}_v\{\tilde{T}_v\} := \left\{ \sum_{i \geq 0} a_i \tilde{T}_v^i \in \mathbb{C}_v[[\tilde{T}_v]] \mid \lim_{i \rightarrow \infty} |a_i|_v = 0 \right\}.$$

We let $\mathbb{C}_v\{\tilde{T}_v\}[\sigma]$ be the non-commutative algebra generated by σ over $\mathbb{C}_v\{\tilde{T}_v\}$ subject to the relation

$$\sigma f = f^{(-1)} \sigma, \quad \forall f \in \mathbb{C}_v\{\tilde{T}_v\},$$

where for every $f = \sum_{i \geq 0} a_i \tilde{T}_v^i \in \mathbb{C}_v\{\tilde{T}_v\}$, we define

$$f^{(n)} := \sum_{i \geq 0} a_i^{q^n} \tilde{T}_v^i, \quad \forall n \in \mathbb{Z}.$$

Given an object \mathcal{M} of $\mathcal{H}\mathcal{J}$, we set $\mathbb{M} := \mathbb{C}_v\{\tilde{T}_v\} \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}$. As $\tilde{T}_v - \iota(\tilde{T}_v) = \tilde{T}_v - \tilde{v}$ is invertible in $\mathbb{C}_v\{\tilde{T}_v\}$, one has that $M \subset \mathbb{M}$, and \mathbb{M} becomes a left $\mathbb{C}_v\{\tilde{T}_v\}[\sigma]$ -module with the σ -action given by

$$\sigma \cdot (f \otimes m) := f^{(-1)} \otimes \sigma m, \quad \forall f \in \mathbb{C}_v\{\tilde{T}_v\} \text{ and } m \in \mathcal{M}.$$

Put

$$H_{\text{Betti}}(\mathcal{M}) := \{m \in \mathbb{M} \mid \sigma m = m\}.$$

We call \mathcal{M} *uniformizable* (or *rigid analytically trivial*) if the natural homomorphism

$$\mathbb{C}_v\{\tilde{T}_v\} \otimes_{\mathbb{F}_q[\tilde{T}_v]} H_{\text{Betti}}(\mathcal{M}) \longrightarrow \mathbb{M}$$

is an isomorphism. In particular, let $\mathbb{C}_v\{\tilde{T}_v\}^\dagger$ be the algebra of rigid-analytic functions on $\mathbb{C}_v \setminus \{\tilde{v}^{q^n} \mid n \in \mathbb{N}\}$. It is known that (see [HaJu20, Proposition 2.4.27]) if \mathcal{M} is uniformizable, then $H_{\text{Betti}}(\mathcal{M})$ is contained in $\mathbb{M}^\dagger = \mathbb{C}_v\{\tilde{T}_v\}^\dagger \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}$, and the natural homomorphism

$$\mathbb{C}_v\{\tilde{T}_v\}^\dagger \otimes_{\mathbb{F}_q[\tilde{T}_v]} H_{\text{Betti}}(\mathcal{M}) \longrightarrow \mathbb{M}^\dagger$$

is an isomorphism.

5.2.2. *\tilde{T}_v -Motivic Galois group.* The full subcategory of $\mathcal{H}\mathcal{J}^I$ consisting of all uniformizable objects is denoted by $\mathcal{H}\mathcal{J}_U^I$. Let $\mathbf{Vect}_{/\mathbb{F}_q(\tilde{T}_v)}$ be the category of finite dimensional vector spaces over $\mathbb{F}_q(\tilde{T}_v)$. By [HaJu20, Theorem 2.4.23], $\mathcal{H}\mathcal{J}_U^I$ is a neutral Tannakian category over $\mathbb{F}_q(\tilde{T}_v)$ with fiber functor $\omega : \mathcal{H}\mathcal{J}_U^I \rightarrow \mathbf{Vect}_{/\mathbb{F}_q(\tilde{T}_v)}$ defined by:

$$\mathcal{M} \longmapsto H_{\mathcal{M}} := \mathbb{F}_q(\tilde{T}_v) \otimes_{\mathbb{F}_q[\tilde{T}_v]} H_{\text{Betti}}(\mathcal{M}).$$

In particular, for each object in $\mathcal{H}\mathcal{J}_U^I$, let $\mathcal{T}_{\mathcal{M}}$ be the strictly full Tannakian subcategory of $\mathcal{H}\mathcal{J}_U^I$ generated by \mathcal{M} . By Tannakian duality, we have an affine algebraic group scheme $\Gamma_{\mathcal{M}}$ defined over $\mathbb{F}_q(\tilde{T}_v)$ so that $\mathcal{T}_{\mathcal{M}}$ is equivalent to the category $\mathbf{Rep}(\Gamma_{\mathcal{M}})$ of the $\mathbb{F}_q(\tilde{T}_v)$ -finite

dimensional representations of the algebraic group $\Gamma_{\mathcal{M}}$. We call $\Gamma_{\mathcal{M}}$ the \tilde{T}_v -*motivic Galois group* of \mathcal{M} . Note that we have the following property of $\dim \Gamma_{\mathcal{M}}$ by Papanikolas' difference Galois theory [P08].

Given an object $\mathcal{M} \in \mathcal{H}\mathcal{J}_U^I$, we let r be the rank of \mathcal{M} over $\bar{k}[\tilde{T}_v]$ and fix a $\bar{k}[\tilde{T}_v]$ -basis $\{m_1, \dots, m_r\}$ of \mathcal{M} . Suppose \mathcal{M} is effective. Then the action of σ on \mathcal{M} is represented by a matrix $\Phi \in \text{Mat}_r(\bar{k}[\tilde{T}_v])$ in the sense that

$$\sigma \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix} = \Phi \begin{pmatrix} m_1 \\ \vdots \\ m_r \end{pmatrix}.$$

Since \mathcal{M} is uniformizable, by [ABP04, P08] there exists a matrix $\Psi \in \text{GL}_r(\mathbb{C}_v \{ \tilde{T}_v \})$ satisfying the system of Frobenius difference equations

$$(5.2.3) \quad \Psi^{(-1)} = \Phi \Psi.$$

In [P08], Papanikolas developed a difference Galois theory for the system of Frobenius difference equation (5.2.3) to explicitly construct an algebraic subgroup $\Gamma_{\Psi} \subset \text{GL}_r$ over $\mathbb{F}_q(t)$ for which

- $\Gamma_{\mathcal{M}}$ is isomorphic to Γ_{Ψ} as algebraic groups over $\mathbb{F}_q(t)$ (see [P08, Thm. 4.5.10]).
- $\dim \Gamma_{\mathcal{M}} = \text{tr. deg}_{\bar{k}(\tilde{T}_v)} \bar{k}(\tilde{T}_v)(\Psi)$ (see [P08, Prop. 4.3.3 and Thm. 4.3.1]),

where the notation $\bar{k}(\tilde{T}_v)(\Psi)$ is referred to be the field generated all entries of Ψ over $\bar{k}(\tilde{T}_v)$. Combining the two properties above, we have the following identity:

$$(5.2.4) \quad \dim \Gamma_{\mathcal{M}} = \text{tr. deg}_{\bar{k}(\tilde{T}_v)} \bar{k}(\tilde{T}_v)(\Psi)$$

Remark 5.2.5. Note that we have a faithful representation

$$\Gamma_{\mathcal{M}} \hookrightarrow \text{GL}(H_{\mathcal{M}}),$$

which is functorial in \mathcal{M} (see [P08, Thm. 4.5.3]). This implies that

$$(5.2.6) \quad \Gamma_{\mathcal{M}} \hookrightarrow \text{Cent}_{\text{GL}(H_{\mathcal{M}})}(\text{End}_{\mathcal{T}_{\mathcal{M}}}(\mathcal{M})) \quad (\text{cf. [CPY10, p. 1448, (6)]})$$

as we have the natural embedding $\text{End}_{\mathcal{T}_{\mathcal{M}}}(\mathcal{M}) \hookrightarrow \text{End}(H_{\mathcal{M}})$.

5.2.7. de Rham pairing and the Grothendieck period conjecture. Let \mathcal{M} be an object of $\mathcal{H}\mathcal{J}_U^I$. Recall that the *de Rham module* of \mathcal{M} over \bar{k} is

$$H_{\text{dR}}(\mathcal{M}, \bar{k}) := \text{Hom}_{\bar{k}} \left(\frac{\mathcal{M}}{(\tilde{T}_v - \tilde{v}) \cdot \mathcal{M}}, \bar{k} \right).$$

In particular, every differential $\omega \in H_{\text{dR}}(\mathcal{M}, \bar{k})$ can be extended to a \mathbb{C}_v -linear functional on $\mathbb{M}^{\dagger} = \mathbb{C}_v \{ \tilde{T}_v \}^{\dagger} \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}$ vanishing on $(\tilde{T}_v - \tilde{v}) \cdot \mathbb{M}^{\dagger}$. As the uniformizability of \mathcal{M} ensures the inclusion $H_{\text{Betti}}(\mathcal{M}) \subset \mathbb{M}^{\dagger}$, we have the following *de Rham pairing*

$$(5.2.8) \quad H_{\text{Betti}}(\mathcal{M}) \times H_{\text{dR}}(\mathcal{M}, \bar{k}) \longrightarrow \mathbb{C}_v, \quad (\gamma, \omega) \longmapsto \int_{\gamma} \omega := \omega(\gamma).$$

The primary tool in this paper of dealing with transcendence problems is the following function field version of Grothendieck's period conjecture, which was achieved by Papanikolas [P08] using the so-called ABP criterion [ABP04].

Theorem 5.2.9. (See [P08] and its refined version in [C09].) Let \mathcal{M} be a uniformizable Hartl–Juschka module \mathcal{M} over \bar{k} . Suppose that \mathcal{M} is effective. Then we have that

$$\dim \Gamma_{\mathcal{M}} = \bar{k} \left(\int_{\gamma} \omega \mid \gamma \in H_{\text{Betti}}(\mathcal{M}), \omega \in H_{\text{dR}}(\mathcal{M}, \bar{k}) \right).$$

Proof. Let $\Phi \in \text{Mat}_r(\bar{k}[\tilde{T}_v])$ be the matrix represented by the σ -action on \mathcal{M} with respect to a $\bar{k}[\tilde{T}_v]$ -basis $\{m_1, \dots, m_r\}$. Since \mathcal{M} is uniformizable, there exists $\Psi \in \text{GL}_r(\mathbb{C}_v \{ \tilde{T}_v \})$ for which $\Psi^{(-1)} = \Phi\Psi$. Put $\mathbf{m} = (m_1, \dots, m_r)^{\text{tr}}$, and note that the entries of $\Psi^{-1}\mathbf{m}$ comprises an $\mathbb{F}_q[\tilde{T}_v]$ -basis of $H_{\text{Betti}}(\mathcal{M})$ (see [ABP04, Lem. 4.4.12] for their arguments without the condition that \mathcal{M} is free of finite rank over $\bar{k}[\sigma]$.) As mentioned above that $H_{\text{Betti}}(\mathcal{M}) \subset \mathcal{M}^\dagger$, elements of Ψ^{-1} are in fact in $\mathbb{C}_v \{ \tilde{T}_v \}^\dagger$.

Our hypothesis on $\det \Phi$ ensures that the conditions of the refined version of the ABP criterion is satisfied (see [C09]), and so we have

$$\text{tr.deg}_{\bar{k}(\tilde{T}_v)} \bar{k}(\tilde{T}_v)(\Psi) = \text{tr.deg}_{\bar{k}} \bar{k}(\Psi|_{\tilde{T}_v=\tilde{v}}) = \text{tr.deg}_{\bar{k}} \bar{k}(\Psi^{-1}|_{\tilde{T}_v=\tilde{v}}),$$

where $\bar{k}(\Psi|_{\tilde{T}_v=\tilde{v}})$ (resp. $\bar{k}(\Psi^{-1}|_{\tilde{T}_v=\tilde{v}})$) denotes by the field over \bar{k} generated by all entries of Ψ (resp. of Ψ^{-1}) evaluated at $\tilde{T}_v = \tilde{v}$. It follows that by (5.2.4), we have

$$(5.2.10) \quad \dim \Gamma_{\mathcal{M}} = \text{tr.deg}_{\bar{k}} \bar{k}(\Psi^{-1}|_{\tilde{T}_v=\tilde{v}}).$$

As each $\omega \in H_{\text{dR}}(\mathcal{M}, \bar{k})$ can be viewed as a \bar{k} -linear functional on \mathcal{M} factoring through $\mathcal{M}/(t - \theta)\mathcal{M}$, a natural basis of $H_{\text{dR}}(\mathcal{M}, \bar{k})$ can be chosen as $\{\omega_j\}_{j=1}^r \subset H_{\text{dR}}(\mathcal{M}, \bar{k})$ given by $\omega_j(m_i) = \delta_{ij}$, the standard Kronecker symbol. Note that $\{\gamma_i := \sum_{j=1}^r (\Psi^{-1})_{ij} m_j\}_{i=1}^r$ is an $\mathbb{F}_q[\tilde{T}_v]$ -basis of $H_{\text{Betti}}(\mathcal{M})$, by (5.2.8) we have that

$$\int_{\gamma_i} \omega_j = ((\Psi^{-1})_{ij})|_{\tilde{T}_v=\tilde{v}} = (\Psi^{-1}|_{\tilde{T}_v=\tilde{v}})_{ij}$$

for every $1 \leq i, j \leq r$. So the desired result follows from (5.2.10). \square

5.3. Hodge–Pink cocharacters. Given an object \mathcal{M} of $\mathcal{H}\mathcal{J}_U^I$, we take a sufficiently large integer n such that

$$(5.3.1) \quad (\tilde{T}_v - \tilde{v})^n \mathcal{M} \subseteq \sigma \mathcal{M} \subseteq (\tilde{T}_v - \tilde{v})^{-n} \mathcal{M}.$$

Let r be the rank of \mathcal{M} over $\bar{k}[\tilde{T}_v]$. There exists a unique (non-ordered) r -tuple of integers (w_1, \dots, w_r) so that

$$\frac{(\tilde{T}_v - \tilde{v})^{-n} \mathcal{M}}{\sigma \mathcal{M}} \cong \bigoplus_{i=1}^r \frac{\bar{k}[\tilde{T}_v]}{(\tilde{T}_v - \tilde{v})^{n-w_i} \bar{k}[\tilde{T}_v]}.$$

We call (w_1, \dots, w_r) the *Hodge–Pink weights* of \mathcal{M} and mention that they are independent of the choices of n . Let $\{m_1, \dots, m_r\}$ be a basis of the free $\bar{k}[\tilde{T}_v]$ -module \mathcal{M} so that

$$\sigma \mathcal{M} = \bigoplus_{i=1}^r (\tilde{T}_v - \tilde{v})^{-w_i} \bar{k}[\tilde{T}_v] \cdot m_i \subset \bar{k}(\tilde{T}_v) \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{M}.$$

Put

$$H_{\mathcal{M}, \mathbb{C}_v} := \mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}_q[\tilde{T}_v]} H_{\mathcal{M}},$$

and let $\lambda_1, \dots, \lambda_r \in H_{\mathcal{M}, \mathbb{C}_v}$ be the images of m_1, \dots, m_r through the following identification

$$(5.3.2) \quad \mathbb{C}_v \otimes_{\bar{k}} \frac{\mathcal{M}}{(\tilde{T}_v - \tilde{v})\mathcal{M}} \cong \frac{\mathbb{M}^\dagger}{(\tilde{T}_v - \tilde{v})\mathbb{M}^\dagger} \cong \frac{\mathbb{C}_v \{\tilde{T}_v\}^\dagger}{(\tilde{T}_v - \tilde{v})\mathbb{C}_v \{\tilde{T}_v\}^\dagger} \otimes_{\mathbb{F}_q[\tilde{T}_v]} H_{\mathcal{M}} \\ \cong \mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}_q[\tilde{T}_v]} H_{\mathcal{M}} = H_{\mathcal{M}, \mathbb{C}_v}.$$

Then we have that:

Proposition 5.3.3. (See [Pink97, Definition 7.10] and [BCPW22, Section 7.3].) *Keep the notation as above. Let $\Gamma_{\mathcal{M}, \mathbb{C}_v} := \mathbb{C}_v \times_{\mathbb{F}_q(\tilde{T}_v)} \Gamma_{\mathcal{M}}$, the base change of $\Gamma_{\mathcal{M}}$ to \mathbb{C}_v via the evaluation map $\iota := (\tilde{T}_v \mapsto \tilde{v}) : \mathbb{F}_q(\tilde{T}_v) \hookrightarrow \mathbb{C}_v$. There exists a cocharacter $\chi_{\mathcal{M}}^\vee : \mathbb{G}_{m/\mathbb{C}_v} \rightarrow \Gamma_{\mathcal{M}, \mathbb{C}_v}$ satisfying that after composing with the embedding $\Gamma_{\mathcal{M}, \mathbb{C}_v} \hookrightarrow \mathrm{GL}(H_{\mathcal{M}, \mathbb{C}_v})$, one has that*

$$\chi_{\mathcal{M}}^\vee(x)(\lambda_i) = x^{w_i} \cdot \lambda_i, \quad \forall x \in \mathbb{G}_{m/\mathbb{C}_v}(\mathbb{C}_v) \text{ and } 1 \leq i \leq r.$$

Since $\mathrm{GL}(H_{\mathcal{M}, \mathbb{C}_v}) = \mathbb{C}_v \times_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}_q(\tilde{T}_v)} \mathrm{GL}(H_{\mathcal{M}})$, we have a natural action of $\mathrm{Aut}_{\mathbb{F}_q(v)}(\mathbb{C}_v)$ on $\mathrm{GL}(H_{\mathcal{M}, \mathbb{C}_v})$, which leaves $\Gamma_{\mathcal{M}, \mathbb{C}_v}$ invariant. Thus any $\mathrm{Aut}_{\mathbb{F}_q(v)}(\mathbb{C}_v)$ -conjugate of $\chi_{\mathcal{M}}^\vee$ is still a cocharacter of $\Gamma_{\mathcal{M}, \mathbb{C}_v}$.

Definition 5.3.4. (cf. [Pink97, Definition 7.10].) *Given an object \mathcal{M} in \mathcal{HJ}_U^J , a Hodge–Pink cocharacter of $\Gamma_{\mathcal{M}, \mathbb{C}_v}$ is a cocharacter which lies in the $\Gamma_{\mathcal{M}, \mathbb{C}_v} \rtimes \mathrm{Aut}_{\mathbb{F}_q(v)}(\mathbb{C}_v)$ -conjugate class of $\chi_{\mathcal{M}}^\vee$.*

6. CARLITZ-TYPE HARTL–JUSCHKA MODULES AND THEIR PERIODS

Throughout this section, we fix an irreducible element $v \in A_+$ of degree d as before and fix a positive integer ℓ . Having sufficient tools at hand, we are now able to interpret $\Omega_{dl, v, i}(v)$ for $0 < i \leq dl$ (occurring in Theorem 4.1.2) as the periods of the restriction of scalars of the ‘‘Carlitz-type’’ Hartl–Juschka modules, and establish their algebraic independence in this section.

6.1. Carlitz-type Hartl–Juschka modules and restriction of scalars.

Definition 6.1.1. *Let $\tilde{\mathcal{C}}_{dl}$ be the left $\bar{k}[\tilde{T}_v][\sigma^{dl}]$ -module $\bar{k}[\tilde{T}_v][\tilde{t}_v]$ whose σ^{dl} -action is given by*

$$\sigma^{dl} \cdot m = \left(1 - \frac{\tilde{t}_v}{\tilde{\theta}_v}\right) \cdot m^{(-dl)}, \quad \forall m \in \bar{k}[\tilde{T}_v][\tilde{t}_v].$$

Remark 6.1.2. By (5.1.3) and (5.1.4), one has that

$$\tilde{T}_v \equiv \tilde{v} \pmod{(\tilde{t}_v - \tilde{\theta}_v)\bar{k}[\tilde{T}_v][\tilde{t}_v]},$$

for which we get

$$(\tilde{T}_v - \tilde{v}) \cdot \tilde{\mathcal{C}}_{dl} \subset \sigma^{dl} \cdot \tilde{\mathcal{C}}_{dl} \subset \tilde{\mathcal{C}}_{dl}.$$

By (5.1.5), we have $\bar{k}[\tilde{T}_v][\tilde{t}_v] = \sum_{n=0}^{\infty} \frac{1}{S^n} \cdot \bar{k}[\tilde{t}_v]$ is not finitely generated as a $\bar{k}[\tilde{t}_v]$ -module if $d > 1$. Since $\frac{1}{S^n} \cdot \bar{k}[\tilde{t}_v] \subset \tilde{\mathcal{C}}_{dl}$ is invariant by σ^{dl} -action for every non-negative integer n , we see that $\tilde{\mathcal{C}}_{dl}$ is not finitely generated as a $\bar{k}[\sigma^{dl}]$ -module when $d > 1$ (i.e. $\tilde{\mathcal{C}}_{dl}$ is not an ‘Anderson dual \tilde{T}_v -motive’ in the terminology of [ABP04]).

Definition 6.1.3. *The restriction of scalars of $\tilde{\mathcal{C}}_{dl}$ is the following $\bar{k}[\tilde{T}_v][\sigma]$ -module:*

$$\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}) := \bar{k}[\tilde{T}_v][\sigma] \otimes_{\bar{k}[\tilde{T}_v][\sigma^{dl}]} \tilde{\mathcal{C}}_{dl}.$$

Decomposing $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ into

$$\bigoplus_{i=0}^{dl-1} \sigma^i \otimes \tilde{\mathcal{C}}_{dl},$$

we get

$$\begin{aligned} \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}) \supset \sigma \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}) &= \left(\left(1 - \frac{\tilde{t}_v}{\tilde{\theta}_v}\right) \cdot 1 \otimes \tilde{\mathcal{C}}_{dl} \right) \oplus \bigoplus_{i=1}^{dl-1} \sigma^i \otimes \tilde{\mathcal{C}}_{dl} \\ &\supset (\tilde{T}_v - \tilde{v}) \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}). \end{aligned}$$

Thus $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ is an effective Hartl–Juschka module \bar{k} .

We see from the following proposition that $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ has ‘complex multiplication’ by $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$.

Proposition 6.1.4. *Let notation be given as above. Then we have the following $\mathbb{F}_q[\tilde{T}_v]$ -algebra embedding:*

$$\phi := (\alpha \mapsto \phi_\alpha) : \mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v] \hookrightarrow \text{End}_{\bar{k}[\tilde{T}_v][\sigma]}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})),$$

where

$$\phi_\alpha := \left(\sum_{i=0}^{dl-1} \sigma^i \otimes m_i \mapsto \sum_{i=0}^{dl-1} \sigma^i \otimes \alpha m_i \right).$$

Proof. From the definition of ϕ , one sees that ϕ is a ring homomorphism. We confirm that for any $\alpha \in \mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$, ϕ_α commutes with σ as follows:

$$\begin{aligned} \phi_\alpha \circ \sigma \left(\sum_{i=0}^{dl-1} \sigma^i \otimes m_i \right) &= \phi_\alpha \left(\sum_{i=0}^{dl-2} \sigma^{i+1} \otimes m_i + 1 \otimes \sigma^{dl} m_{dl-1} \right) \\ &= \sum_{i=0}^{dl-2} \sigma^{i+1} \otimes \alpha m_i + 1 \otimes \alpha \sigma^{dl} m_{dl-1} = \sum_{i=0}^{dl-2} \sigma^{i+1} \otimes \alpha m_i + 1 \otimes \sigma^{dl} \alpha m_{dl-1} \\ &= \sigma \left(\sum_{i=0}^{dl-1} \sigma^i \otimes \alpha m_i \right) = \sigma \circ \phi_\alpha \left(\sum_{i=0}^{dl-1} \sigma^i \otimes m_i \right). \end{aligned}$$

□

Remark 6.1.5. Alternatively, we may regard $\tilde{\mathcal{C}}_{dl}$ as a $(\bar{k}[\tilde{T}_v][\sigma^{dl}], \mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v])$ -bimodule, and get the $\mathbb{F}_q[\tilde{T}_v]$ -algebra homomorphism $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v] \hookrightarrow \text{End}_{\bar{k}[\tilde{T}_v][\sigma]}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}))$ explicitly described in Proposition 6.1.4.

6.1.6. *Explicit descriptions of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$.* Let $\bar{k}[\tilde{T}_v][X]$ be the polynomial ring with variable X over $\bar{k}[\tilde{T}_v]$, and let $\tilde{g}_{\tilde{T}_v}(X)$ be the minimal polynomial of \tilde{t}_v over $\bar{k}[\tilde{T}_v]$, i.e., by (5.1.4), we have

$$\tilde{g}_{\tilde{T}_v}(X) = X^d - (\tilde{\epsilon}_{d-1} \tilde{T}_v) X^{d-1} - \dots - (\tilde{\epsilon}_1 \tilde{T}_v) X - \tilde{T}_v \in \mathbb{F}_{q^d}[\tilde{T}_v][X] \subset \bar{k}[\tilde{T}_v][X].$$

Then $\bar{k}[\tilde{T}_v][\tilde{t}_v]$ can be identified with

$$\frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}(X) \cdot \bar{k}[\tilde{T}_v][X]}.$$

We extend the Frobenius twists by fixing X to $\bar{k}(\tilde{T}_v)[X]$ as follows: for each $f(X) = \sum_m f_m X^m$ and $n \in \mathbb{Z}$,

$$f^{(n)}(X) := \sum_m f_m^{(n)} X^m.$$

We point out that when $f(X) \in \bar{k}[X]$, i.e. $f_m \in \bar{k}$ for all m , one has that $f^{(n)}(X) = \sum_m f_m^{q^n} X^m$. In the decomposition

$$\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}) = \bigoplus_{i=0}^{d\ell-1} \sigma^i \otimes \tilde{\mathcal{C}}_{d\ell},$$

for each i with $0 \leq i < d\ell$ we have the $\bar{k}[\tilde{T}_v]$ -linear isomorphism

$$\frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}^{(-i)}(X) \cdot \bar{k}[\tilde{T}_v][X]} \xrightarrow{\sim} \sigma^i \otimes \tilde{\mathcal{C}}_{d\ell} = \sigma^i \otimes \left(\frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}(X) \cdot \bar{k}[\tilde{T}_v][X]} \right)$$

defined by

$$\left(\sum_{m=0}^n f_m X^m \right) \bmod \tilde{g}_{\tilde{T}_v}^{(-i)}(X) \mapsto \sigma^i \otimes \left(\sum_{m=0}^n f_m^{(i)} X^m \bmod \tilde{g}_{\tilde{T}_v}(X) \right).$$

Therefore:

Proposition 6.1.7. *We may identify $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ with the product space*

$$\frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}^{(0)}(X) \cdot \bar{k}[\tilde{T}_v][X]} \times \frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}^{(-1)}(X) \cdot \bar{k}[\tilde{T}_v][X]} \times \cdots \times \frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}^{(1-d\ell)}(X) \cdot \bar{k}[\tilde{T}_v][X]},$$

equipped with the σ -action given by:

$$\begin{aligned} & \sigma \cdot (h_0 \bmod \tilde{g}_{\tilde{T}_v}^{(0)}, h_1 \bmod \tilde{g}_{\tilde{T}_v}^{(-1)}, \dots, h_{d\ell-1} \bmod \tilde{g}_{\tilde{T}_v}^{(1-d\ell)}) \\ & := \left(\left(1 - \frac{X}{\tilde{\theta}_v}\right) \cdot h_{d\ell-1}^{(-1)} \bmod \tilde{g}_{\tilde{T}_v}^{(0)}, h_0^{(-1)} \bmod \tilde{g}_{\tilde{T}_v}^{(-1)}, \dots, h_{d\ell-2}^{(-1)} \bmod \tilde{g}_{\tilde{T}_v}^{(1-d)} \right). \end{aligned}$$

Remark 6.1.8. As $\tilde{t}_{v,i}$ is a root of $\tilde{g}_{\tilde{T}_v}^{(-i)}(X)$ for each integer i with $0 \leq i < d\ell$, one has that

$$\bar{k}[\tilde{T}_v][\tilde{t}_{v,i}] \cong \frac{\bar{k}[\tilde{T}_v][X]}{\tilde{g}_{\tilde{T}_v}^{(-i)}(X) \cdot \bar{k}[\tilde{T}_v][X]}.$$

The above proposition allows us to identify $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ with

$$(6.1.9) \quad \bar{k}[\tilde{T}_v][\tilde{t}_{v,0}] \times \bar{k}[\tilde{T}_v][\tilde{t}_{v,1}] \times \cdots \times \bar{k}[\tilde{T}_v][\tilde{t}_{v,d\ell-1}],$$

and the σ -action is given by: for every polynomials $h_0(X), \dots, h_{d\ell-1}(X) \in \bar{k}[\tilde{T}_v][X]$,

$$\begin{aligned} & \sigma \cdot \left(h_0(\tilde{t}_{v,0}), h_1(\tilde{t}_{v,1}), \dots, h_{d\ell-1}(\tilde{t}_{v,d\ell-1}) \right) \\ & = \left(\left(1 - \frac{\tilde{t}_{v,0}}{\tilde{\theta}_v}\right) \cdot h_{d\ell-1}^{(-1)}(\tilde{t}_{v,0}), h_0^{(-1)}(\tilde{t}_{v,1}), \dots, h_{d\ell-2}^{(-1)}(\tilde{t}_{v,d\ell-1}) \right). \end{aligned}$$

6.2. Uniformizability. Recall that $\mathbb{C}_v\{\tilde{T}_v\}$ is the Tate algebra with variable \tilde{T}_v over \mathbb{C}_v . Denote by $\mathbb{C}_v\{\tilde{t}_v\}$ the Tate algebra with variable \tilde{t}_v over \mathbb{C}_v , i.e.

$$\mathbb{C}_v\{\tilde{t}_v\} := \left\{ \sum_{i \geq 0} a_i \tilde{t}_v^i \in \mathbb{C}_v[[\tilde{t}_v]] \mid \lim_{i \rightarrow \infty} |a_i|_v = 0 \right\}.$$

Put

$$\mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v] = \mathbb{C}_v\{\tilde{T}_v\} \otimes_{\bar{k}[\tilde{T}_v]} \bar{k}[\tilde{T}_v][\tilde{t}_v].$$

Lemma 6.2.1. *We have the following inclusion*

$$\mathbb{C}_v\{\tilde{t}_v\} \subset \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v].$$

Proof. For each non-negative integer i , by (5.1.4) there exist $f_{i,j} \in \mathbb{F}_{q^d}[\tilde{T}_v]$, $0 \leq j < d$, with $\deg_{\tilde{T}_v} f_{i,j} \leq i$ such that

$$\tilde{t}_v^i = f_{i,0} \tilde{t}_v^{d-1} + \cdots + f_{i,d-1}.$$

Write $f_{i,j}(\tilde{T}_v) = \sum_{s=0}^i u_{i,j,s} \tilde{T}_v^s$ where $u_{i,j,s} \in \mathbb{F}_{q^d}$. Then for each $f = \sum_{i \geq 0} a_i \tilde{t}_v^i \in \mathbb{C}_v\{\tilde{t}_v\}$, the property $\lim_{i \rightarrow \infty} |a_i|_v = 0$ implies that for $0 \leq j < d$ and $s \geq 0$,

$$b_{j,s} := \lim_{n \rightarrow \infty} \sum_{i=s}^n a_i u_{i,j,s} \quad \text{converges absolutely in } \mathbb{C}_v \text{ with } \quad \lim_{s \rightarrow \infty} b_{j,s} = 0.$$

Put

$$\begin{aligned} f_j &:= \sum_{s \geq 0} b_{j,s} \tilde{T}_v^s = \sum_{s=0}^{\infty} \left(\sum_{i=s}^{\infty} a_i u_{i,j,s} \right) \tilde{T}_v^s \\ &= \sum_{i=0}^{\infty} a_i \left(\sum_{s=0}^i u_{i,j,s} \tilde{T}_v^s \right) = \sum_{i=0}^{\infty} a_i f_{i,j}(\tilde{T}_v) \in \mathbb{C}_v\{\tilde{T}_v\}. \end{aligned}$$

We then get

$$f = \sum_{i=0}^{\infty} a_i \tilde{t}_v^i = \sum_{j=0}^{d-1} \sum_{i=0}^{\infty} a_i f_{i,j} \tilde{t}_v^j = \sum_{j=0}^{d-1} f_j \cdot \tilde{t}_v^j \in \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v]$$

as desired. \square

Remark 6.2.2. The inclusion $\bar{k}(\tilde{T}_v) \subset \bar{k}(\tilde{t}_v)$ induces a finite morphism ϕ from the projective \tilde{t}_v -line $\mathbb{P}_{\tilde{t}_v}^1$ onto the projective \tilde{T}_v -line $\mathbb{P}_{\tilde{T}_v}^1$ (over \bar{k}). In particular, $\bar{k}[\tilde{t}_v]$ and $\bar{k}[\tilde{T}_v][\tilde{t}_v]$ are the affine coordinate rings of $\mathbb{A}_{\tilde{t}_v}^1$ and $U := \mathbb{A}_{\tilde{t}_v}^1 \setminus \phi^{-1}(\tilde{v})$, respectively. Since $\mathbb{C}_v\{\tilde{t}_v\}$ and $\mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v]$ correspond to the the affinoid domains $\mathbb{A}_{\tilde{t}_v, \leq 1}^1 := \{x \in \mathbb{A}_{\tilde{t}_v}^1(\mathbb{C}_v) \mid |x|_v \leq 1\}$ and $U_{\leq 1} := \{x \in U(\mathbb{C}_v) \mid |x|_v \leq 1\}$, respectively, the natural inclusion $U_{\leq 1} \subset \mathbb{A}_{\tilde{t}_v, \leq 1}^1$ alternatively implies $\mathbb{C}_v\{\tilde{t}_v\} \subset \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v]$.

To show the uniformizability of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$, we first note that the identification of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ extends to

$$(6.2.3) \quad \mathbb{C}_v\{\tilde{T}_v\} \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}) = \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,0}] \times \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,1}] \times \cdots \times \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,d\ell-1}].$$

In particular, Lemma 6.2.1 implies that

$$(6.2.4) \quad \mathbb{C}_v\{\tilde{t}_{v,0}\} \times \mathbb{C}_v\{\tilde{t}_{v,1}\} \times \cdots \times \mathbb{C}_v\{\tilde{t}_{v,d\ell-1}\} \subset \mathbb{C}_v\{\tilde{T}_v\} \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}).$$

In particular, for every $(h_0(\tilde{t}_{v,0}), \dots, h_{dl-1}(\tilde{t}_{v,dl-1})) \in \mathbb{C}_v\{\tilde{t}_{v,0}\} \times \mathbb{C}_v\{\tilde{t}_{v,1}\} \times \dots \times \mathbb{C}_v\{\tilde{t}_{v,dl-1}\}$, one has that

$$(6.2.5) \quad \begin{aligned} & \sigma \cdot \left(h_0(\tilde{t}_{v,0}), h_1(\tilde{t}_{v,1}), \dots, h_{dl-1}(\tilde{t}_{v,dl-1}) \right) \\ &= \left(\left(1 - \frac{\tilde{t}_{v,0}}{\tilde{\theta}_v}\right) \cdot h_{dl-1}^{(-1)}(\tilde{t}_{v,0}), h_0^{(-1)}(\tilde{t}_{v,1}), \dots, h_{dl-2}^{(-1)}(\tilde{t}_{v,dl-1}) \right), \end{aligned}$$

and for $0 \leq i < dl$,

$$h_i^{(-1)}(\tilde{t}_{v,\overline{i+1}}) = \sum_m a_m^{q^{-1}} \tilde{t}_{v,\overline{i+1}}^m \quad \text{when} \quad h_i(\tilde{t}_{v,i}) = \sum_m a_m \tilde{t}_{v,i}^m \in \mathbb{C}_v\{\tilde{t}_{v,i}\}.$$

Here $\overline{i+1} = i+1$ if $i+1 < dl$ or 0 if $i+1 = dl$.

Note that since $|\tilde{\theta}_v|_v > 1$, given an integer i with $0 \leq i < dl$, we can set $\psi(\tilde{t}_v)$ to be the following infinite product

$$\psi(\tilde{t}_v) := \prod_{i=1}^{\infty} \left(1 - \frac{\tilde{t}_v}{\tilde{\theta}_v^{q^{di}}} \right)^{-1} \in \mathbb{C}_v\{\tilde{t}_v\}^\times \subset \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_v]^\times.$$

Take $\xi \in \mathbb{F}_{q^{dl}}$ so that $\mathbb{F}_{q^{dl}} = \mathbb{F}_q(\xi)$. For $0 \leq i < dl$ and $0 \leq j < d$, Put

$$\begin{aligned} \psi_{i,j} &:= \left(\xi^i \tilde{t}_{v,0}^j \psi(\tilde{t}_{v,0}), (\xi^{q^{-1}})^i \tilde{t}_{v,1}^j \psi^{(-1)}(\tilde{t}_{v,1}), \dots, (\xi^{q^{1-dl}})^i \tilde{t}_{v,dl-1}^j \psi^{(1-dl)}(\tilde{t}_{v,dl-1}) \right) \\ &\in \mathbb{C}_v\{\tilde{t}_{v,0}\} \times \mathbb{C}_v\{\tilde{t}_{v,1}\} \times \dots \times \mathbb{C}_v\{\tilde{t}_{v,dl-1}\}. \end{aligned}$$

Then

Lemma 6.2.6. *Via the decomposition (6.2.3), we have that*

$$H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})) = \bigoplus_{\substack{0 \leq i < dl \\ 0 \leq j < d}} \mathbb{F}_q[\tilde{T}_v] \psi_{i,j}.$$

Consequently, $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ is uniformizable.

Proof. We first note that

$$\psi^{(-dl)}(\tilde{t}_v) = \left(1 - \frac{\tilde{t}_{v,0}}{\tilde{\theta}_v}\right)^{-1} \cdot \psi(\tilde{t}_v).$$

From (6.2.5) we see that $\psi_{i,j} \in H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}))$ for $0 \leq i < dl$ and $0 \leq j < d$. Since $\text{rank}_{\bar{k}[\tilde{T}_v]} \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}) = d^2 \ell$, it suffices to show that the set $\{\psi_{i,j} \mid 0 \leq i < dl, 0 \leq j < d\}$ generates $\mathbb{C}_v\{\tilde{T}_v\} \otimes_{\bar{k}[\tilde{T}_v]} \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ over $\mathbb{C}_v\{\tilde{T}_v\}$. First, from the isomorphism

$$\begin{aligned} \bar{k} \otimes_{\mathbb{F}_q} \mathbb{F}_{q^{dl}} &\xrightarrow{\sim} \bar{k} \times \bar{k} \times \dots \times \bar{k} \\ \alpha \otimes 1 &\longmapsto (\alpha, \alpha, \dots, \alpha) \\ 1 \otimes \xi &\longmapsto (\xi, \xi^{q^{-1}}, \dots, \xi^{q^{dl-1}}), \end{aligned}$$

one has that for each integer i with $0 \leq i < dl$, there exist $\alpha_{i,0}, \dots, \alpha_{i,dl-1} \in \bar{k}$ such that

$$\sum_{m=0}^{dl-1} \alpha_{i,m} \xi^{mq^{-i'}} = \begin{cases} 1, & \text{if } i' = i, \\ 0, & \text{otherwise.} \end{cases}$$

Put

$$\psi'_{i,j} := \sum_{m=0}^{dl-1} \alpha_{i,m} \psi_{m,j} = \left(0, \dots, \tilde{t}_{v,i}^j \psi^{(-i)}(\tilde{t}_{v,i}), 0, \dots, 0\right).$$

We then obtain that

$$\begin{aligned} \sum_{\substack{0 \leq i < dl \\ 0 \leq j < d}} \mathbb{C}_v\{\tilde{T}_v\} \cdot \psi_{i,j} &= \sum_{\substack{0 \leq i < dl \\ 0 \leq j < d}} \mathbb{C}_v\{\tilde{T}_v\} \cdot \psi'_{i,j} \\ &= \left(\mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,0}] \cdot \psi(\tilde{t}_{v,0})\right) \times \left(\mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,1}] \cdot \psi^{(-1)}(\tilde{t}_{v,1})\right) \times \cdots \times \left(\mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,dl-1}] \cdot \psi^{(1-dl)}(\tilde{t}_{v,dl-1})\right) \\ &= \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,0}] \times \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,1}] \times \cdots \times \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,dl-1}], \end{aligned}$$

where the last equality comes from the observation:

$$\psi^{(-i)}(\tilde{t}_{v,i}) \in \mathbb{C}_v\{\tilde{t}_{v,i}\}^\times \subset \mathbb{C}_v\{\tilde{T}_v\}[\tilde{t}_{v,i}]^\times, \quad 0 \leq i < dl.$$

This completes the proof. \square

Remark 6.2.7. Since $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ has Complex Multiplication by $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$ (see Remark 6.1.5), we may regard $H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}))$ as an $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$ -module. Then the proof of the above lemma indicates that

$$H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})) = \mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v] \cdot \psi_{0,0}.$$

6.3. Periods of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$. Recall that the *de Rham module* of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ over \bar{k} is

$$H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k}) := \text{Hom}_{\bar{k}} \left(\frac{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}{(\tilde{T}_v - \tilde{v}) \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}, \bar{k} \right),$$

and the de Rham pairing is given by (see (5.2.8))

$$H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})) \times H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k}) \longrightarrow \mathbb{C}_v, \quad (\gamma, \omega) \mapsto \int_\gamma \omega := \omega(\gamma).$$

In particular, for each integer i with $0 \leq i < dl$, put

$$(6.3.1) \quad \tilde{g}_{v,i}(X) := \tilde{g}_{\tilde{T}_v}^{(-i)}(X)|_{\tilde{T}_v = \tilde{v}} = X^d - (\tilde{\epsilon}_{d-1}^{q^{(-i)}} \tilde{v})X^{d-1} - \cdots - (\tilde{\epsilon}_1^{q^{(-i)}} \tilde{v})X - \tilde{v} \in \bar{k}[X].$$

The identification in Proposition 6.1.7 leads to the following isomorphism:

$$(6.3.2) \quad \frac{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}{(\tilde{T}_v - \tilde{v}) \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})} \cong \prod_{i=0}^{dl-1} \frac{\bar{k}[X]}{\tilde{g}_{v,i}(X) \cdot \bar{k}[X]}.$$

Moreover, for each integer s with $0 \leq s < dl$, from (5.1.4) the element

$$\tilde{\theta}_{v,s} := (\theta - \varepsilon^{q^{-s}})^{-1} = \iota(\tilde{t}_{v,s}) \in \bar{k}$$

is a root of $\tilde{g}_{v,s}(X)$, which gives a differential $\tilde{\omega}_s \in H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k})$ induced from

$$(6.3.3) \quad \begin{aligned} \prod_{i=0}^{dl-1} \frac{\bar{k}[X]}{\tilde{g}_{v,i}(X) \cdot \bar{k}[X]} &\longrightarrow \frac{\bar{k}[X]}{\tilde{g}_{v,s}(X) \cdot \bar{k}[X]} \longrightarrow \bar{k} \\ \left(\bar{h}_i(X) \bmod \tilde{g}_{v,i}\right)_{0 \leq i < dl} &\longmapsto \bar{h}_s(X) \bmod \tilde{g}_{v,s} \longmapsto \bar{h}_s(\tilde{\theta}_{v,s}). \end{aligned}$$

Therefore we obtain the following period interpretation:

Proposition 6.3.4. *Recall the notation (4.1.1). Given an integer s with $0 \leq s < dl$, one has*

$$\int_{\psi_{0,0}} \tilde{\omega}_s = \frac{1}{\Omega_{dl,v,dl-s}(v)}.$$

Proof. From the definition of ω_s and the de Rham pairing, one gets that

$$\begin{aligned} \int_{\psi_{0,0}} \tilde{\omega}_s &= \psi^{(-s)}(\tilde{t}_{v,s}) \Big|_{\tilde{t}_{v,s}=\tilde{\theta}_{v,s}} \\ &= \prod_{i=1}^{\infty} \left(1 - \frac{\tilde{\theta}_{v,s}}{\tilde{\theta}_v^{q^{dli-s}}} \right)^{-1} = \prod_{i=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon^{q^{-s}})^{-1}}{((\theta - \varepsilon)^{q^{dli-s}})^{-1}} \right)^{-1} \\ &= \prod_{i=1}^{\infty} \left(1 - \frac{(\theta - \varepsilon)^{q^{dli-s}}}{\theta - \varepsilon^{q^{-s}}} \right)^{-1} = \frac{1}{\Omega_{dl,v,dl-s}(v)}. \end{aligned}$$

□

Remark 6.3.5. As $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ has CM by $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$ (see Remark 6.2.7), we have that

$$\begin{aligned} \bar{k} \left(\int_{\gamma} \tilde{\omega} \mid \gamma \in H_{\text{Betti}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})), \tilde{\omega} \in H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k}) \right) \\ = \bar{k} \left(\int_{\psi_{0,0}} \tilde{\omega} \mid \tilde{\omega} \in H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k}) \right). \end{aligned}$$

The Gronthendieck period conjecture in Theorem 5.2.9 leads to

$$(6.3.6) \quad \text{tr. deg}_{\bar{k}} \bar{k} \left(\int_{\psi_{0,0}} \omega \mid \omega \in H_{\text{dR}}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl}), \bar{k}) \right) = \dim \Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})},$$

where $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}$ is the \tilde{T}_v -motivic Galois group of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$. Next, we will determine the dimension of $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}$, and complete the proof of Theorem 4.1.2.

6.4. \tilde{T}_v -Motivic Galois group of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ and the Hodge–Pink cocharacters. We let $\mathbb{K}_{dl} := \mathbb{F}_{q^{dl}}(\tilde{t}_v)$ and $\mathbb{F} := \mathbb{F}_q(\tilde{T}_v)$. Since $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})$ has complex multiplication by $\mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v]$ by Proposition 6.1.4 and

$$\text{rank}_{\mathbb{F}_q[\tilde{T}_v]} \mathbb{F}_{q^{dl}}[\tilde{T}_v][\tilde{t}_v] = [\mathbb{K}_{dl} : \mathbb{F}] = d^2 \ell = \dim_{\mathbb{F}_q(\tilde{T})} H_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})},$$

by (5.2.6) we obtain that

$$\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})} \hookrightarrow \mathcal{R}_{\mathbb{K}_{dl}/\mathbb{F}}(\mathbb{G}_{m/\mathbb{K}_{dl}}) \subset \text{GL}(H_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}),$$

where $\mathcal{R}_{\mathbb{K}_{dl}/\mathbb{F}}(\mathbb{G}_{m/\mathbb{K}_{dl}})$ is the restriction of scalars of $\mathbb{G}_{m/\mathbb{K}_{dl}}$ to \mathbb{F} . Our goal is to show $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})} = \mathcal{R}_{\mathbb{K}_{dl}/\mathbb{F}}(\mathbb{G}_{m/\mathbb{K}_{dl}})$ by analysing the images of its Hodge–Pink cocharacters.

6.4.1. Hodge–Pink cocharacters of $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{dl})}$. For $0 \leq i < dl$, let $\tilde{\theta}_{v,i,0}(= \tilde{\theta}_{v,i}), \dots, \tilde{\theta}_{v,i,d-1}$ be the distinct roots of $\tilde{g}_{v,i}(X) \in \bar{k}[X]$ defined in (6.3.1). Comparing the minimal polynomials of $\tilde{t}_{v,i}$ in (5.1.6) and $\tilde{g}_{v,i}(X)$, we get that

$$\tilde{T}_v - \tilde{v} \in (\tilde{t}_{v,i} - \tilde{\theta}_{v,i,j}) \bar{k}[\tilde{T}_v][\tilde{t}_{v,i}],$$

whence

$$(6.4.2) \quad (\tilde{T}_v - \tilde{v}) \cdot \bar{k}[\tilde{T}_v][\tilde{t}_{v,i}] = \prod_{j=0}^{d-1} (\tilde{t}_{v,i} - \tilde{\theta}_{v,i,j}) \cdot \bar{k}[\tilde{T}_v][\tilde{t}_{v,i}].$$

On the other hand, under the identification (6.1.9), we get

$$(6.4.3) \quad \begin{aligned} \sigma \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}) &= \left((\tilde{t}_{v,0} - \tilde{\theta}_{v,0,0}) \cdot \bar{k}[\tilde{T}_v][\tilde{t}_{v,0}] \right) \times \bar{k}[\tilde{T}_v][\tilde{t}_{v,1}] \times \cdots \times \bar{k}[\tilde{T}_v][\tilde{t}_{v,d\ell-1}] \\ &\supset (\tilde{T}_v - \tilde{v}) \cdot \mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \end{aligned}$$

and the Hodge–Pink weight of $\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})$ is $(w_{i,j})_{0 \leq i < d\ell, 0 \leq j < d}$ with

$$w_{i,j} = \begin{cases} -1, & \text{if } i = j = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Moreover precisely, we may take a $\bar{k}[\tilde{T}_v]$ -base $\{m_{i,j} \mid 0 \leq i < d\ell, 0 \leq j < d\}$ satisfying that under the identification

$$\frac{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}{(\tilde{T}_v - \tilde{v})\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})} \cong \prod_{i=0}^{d\ell-1} \frac{\bar{k}[\tilde{T}_v][\tilde{t}_{v,i}]}{(\tilde{T}_v - \tilde{v})\bar{k}[\tilde{T}_v][\tilde{t}_{v,i}]} \cong \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \frac{\bar{k}[\tilde{T}_v][\tilde{t}_{v,i}]}{(\tilde{t}_{v,i} - \tilde{\theta}_{v,i,j})\bar{k}[\tilde{T}_v][\tilde{t}_{v,i}]} \cong \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \bar{k},$$

the image of $m_{i,j}$ is

$$(\delta_{(i,j),(i',j')})_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \in \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \bar{k}, \quad \text{where} \quad \delta_{(i,j),(i',j')} = \begin{cases} 1, & \text{if } i = i' \text{ and } j = j', \\ 0, & \text{otherwise.} \end{cases}$$

From the isomorphism (5.3.2), we may decompose

$$H_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \mathbb{C}_v} = \mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}} H_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})} \xrightarrow{\sim} \bigoplus_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} H_{i,j}.$$

Notice that for each $0 \leq i < d\ell$ and $0 \leq j < d$, let $\iota_{i,j} : \mathbb{K}_{d\ell} = \mathbb{F}_{q^{d\ell}}(\tilde{t}_v) \hookrightarrow \mathbb{C}_v$ be the embedding satisfying that $\iota_{i,j}(\varepsilon) = \varepsilon^{q^{-j}}$ and $\iota_{i,j}(\tilde{t}_v) = \tilde{\theta}_{v,i,j}$. Regarding $H_\sigma(\tilde{\mathcal{C}}_{d\ell})$ as a one dimensional vector space over $\mathbb{K}_{d\ell}$ (by Remark 6.2.7), we have that

$$\alpha \cdot z = \iota_{i,j}(\alpha) \cdot z, \quad \forall z \in H_{i,j} \quad \text{and} \quad \alpha \in \mathbb{F}_{q^{d\ell}}(\tilde{t}_v).$$

Therefore under the identification

$$\mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}} \mathcal{R}_{\mathbb{K}_{d\ell}/\mathbb{F}}(\mathbb{G}_m/\mathbb{K}_{d\ell}) \cong \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \mathbb{C}_v \otimes_{\iota_{i,j}, \mathbb{K}_{d\ell}} \mathbb{G}_m/\mathbb{K}_{d\ell} = \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \text{GL}(H_{i,j}),$$

the cocharacter

$$\chi_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}^\vee : \mathbb{G}_m/\mathbb{C}_v \rightarrow \Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \mathbb{C}_v} \subset \mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}} \mathcal{R}_{\mathbb{K}_{d\ell}/\mathbb{F}}(\mathbb{G}_m/\mathbb{K}_{d\ell})$$

is then realized as follows:

$$\chi_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}^\vee(x) = \begin{pmatrix} x^{-1} & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix} \in \prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \mathbb{C}_v \otimes_{\iota_{i,j}, \mathbb{K}_{d\ell}} \mathbb{G}_m/\mathbb{K}_{d\ell}.$$

Since $\text{Aut}_{\mathbb{F}_q(\tilde{v})}(\mathbb{C}_v)$ acts transitively on the coordinates of $\prod_{\substack{0 \leq i < d\ell \\ 0 \leq j < d}} \mathbb{C}_v \otimes_{\iota_{i,j}, \mathbb{K}_{d\ell}} \mathbb{G}_m/\mathbb{K}_{d\ell}$, the Hodge–Pink cocharacters of $\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})}$ generate the whole group $\mathbb{C}_v \otimes_{\tilde{v} \leftarrow \tilde{T}_v, \mathbb{F}} \mathcal{R}_{\mathbb{K}_{d\ell}/\mathbb{F}}(\mathbb{G}_m/\mathbb{K}_{d\ell})$. Consequently, we finally arrive at:

Theorem 6.4.4.

$$\Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})} = \mathcal{R}_{\mathbb{K}_{d\ell}/\mathbb{F}}(\mathbb{G}_{m/\mathbb{K}_{d\ell}}),$$

whence $\dim \Gamma_{\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell})} = [\mathbb{K}_{d\ell} : \mathbb{F}] = d^2\ell$.

Combining with (6.3.6), we finally arrive at:

Corollary 6.4.5.

$$\text{tr. deg}_{\bar{k}} \bar{k} \left(\int_{\psi_{0,0}} \tilde{\omega} \mid \tilde{\omega} \in H_{dR}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \bar{k}) \right) = d^2\ell.$$

6.4.6. *Proof of Theorem 4.1.2.* By Proposition 6.3.4 we have that

$$\Omega_{d\ell, v, i}(v) = \left(\int_{\psi_{0,0}} \tilde{\omega}_{d\ell-i} \right)^{-1}, \quad 0 < i \leq d\ell.$$

On the other hand, for $0 \leq i < d\ell$ and $0 \leq j < d$, define $\tilde{\omega}_{i,j} \in H_{dR}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \bar{k})$ as follows:

$$\begin{aligned} \prod_{i'=0}^{d\ell-1} \frac{\bar{k}[X]}{\bar{g}_{v,i'}(X) \cdot \bar{k}[X]} &\longrightarrow \frac{\bar{k}[X]}{\bar{g}_{v,i}(X) \cdot \bar{k}[X]} \longrightarrow \bar{k} \\ \left(\bar{h}_{i'}(X) \bmod \bar{g}_{v,i'} \right)_{0 \leq i' < d\ell} &\longmapsto \bar{h}_i(X) \bmod \bar{g}_{v,i} \longmapsto \bar{h}_i(\tilde{\theta}_{v,i,j}), \end{aligned}$$

where $\tilde{\theta}_{v,i,j}$ are given in the beginning of Section 6.4.1. Then

$$\{\tilde{\omega}_{i,j} \mid 0 \leq i < d\ell, 0 \leq j < d\} \quad \text{is a basis of } H_{dR}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \bar{k}),$$

and $\tilde{\omega}_i = \tilde{\omega}_{i,0}$ for $0 \leq i < d\ell$ by comparing with (6.3.3). Since

$$\dim_{\bar{k}} H_{dR}(\mathcal{R}_\sigma(\tilde{\mathcal{C}}_{d\ell}), \bar{k}) = d^2\ell = \text{tr. deg}_{\bar{k}} \bar{k} \left(\int_{\psi_{0,0}} \tilde{\omega}_{i,j} \mid 0 \leq i < d\ell, 0 \leq j < d \right),$$

we obtain that

$$\int_{\psi_{0,0}} \tilde{\omega}_{i,j}, \quad 0 \leq i < d\ell, \quad 0 \leq j < d, \quad \text{are algebraically independent over } \bar{k}.$$

Finally, as for $0 < i \leq \ell$, one has that

$$(6.4.7) \quad \prod_{j=0}^{d-1} \Omega_{d\ell, v, j\ell+i}(v) = \prod_{j=0}^{d-1} \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{d\ell n + j\ell + i}}}{\theta - \varepsilon^{q^{j\ell + i}}} \right) = \prod_{n=0}^{\infty} \left(1 - \frac{\theta_v^{q^{\ell n + i}}}{\theta - \varepsilon^{q^{\ell n + i}}} \right) = \Omega_{\ell, v, i}(v).$$

Therefore the result follows. □

REFERENCES

- [All96] J.-P. Allouche, *Transcendence of the Carlitz–Goss gamma at rational arguments*, J. Number Theory **60** (1996) 318–328.
- [A86] G. W. Anderson, *t-motives*, Duke Math. J. **53** (1986), no. 2, 457–502.
- [ABP04] G. W. Anderson, W. D. Brownawell and M. A. Papanikolas, *Determination of the algebraic relations among special Γ -values in positive characteristic*, Ann. of Math. (2) **160** (2004), no. 1, 237–313.
- [BCPW22] W. D. Brownawell, C.-Y. Chang, M. A. Papanikolas, and F.-T. Wei, *Function field analogue of Shimura’s conjecture on period symbols*, preprint (arXiv:2203.09131).
- [BP02] W. D. Brownawell and M. A. Papanikolas, *Linear independence of Gamma-values in positive characteristic*, J. Reine Angew. Math. **549** (2002), 91–148.

- [BW07] A. Baker and G. Wüstholz, *Logarithmic forms and Diophantine geometry*, New Math. Monogr., 9 Cambridge University Press, Cambridge, 2007.
- [Car35] L. Carlitz, *On certain functions connected with polynomials in a Galois field*, Duke Math. J. 1 (1935) 137–168.
- [C09] C.-Y. Chang, *A note on a refined version of Anderson-Brownawell-Papanikolas criterion*, J. Number Theory 129 (2009), 729–738.
- [CP12] C.-Y. Chang and M. A. Papanikolas, *Algebraic independence of periods and logarithms of Drinfeld modules. With an appendix by Brian Conrad*. J. Amer. Math. Soc. 25 (2012), no.1, 123–150.
- [CPTY10] C.-Y. Chang, M. A. Papanikolas, D. S. Thakur, and J. Yu, *Algebraic independence of arithmetic gamma values and Carlitz zeta values*, Adv. math. 223 (2010) 1137–1154.
- [CPY10] C.-Y. Chang, M. A. Papanikolas and J. Yu, *Geometric gamma values and zeta values in positive characteristic*, Int. Math. Res. Not. IMRN 2010, no. 8, 1432–1455.
- [Chen23] Y.-T. Chen, *A v -adic variant of Anderson–Brownawell–Papanikolas linear independence criterion and its application*, to appear in Math. Res. Letter. arXiv:2308.05825.
- [Chud84] G. V. Chudnovsky, *Contributions to the theory of transcendental numbers*, Math. Surveys Monogr., 19 American Mathematical Society, Providence, RI, 1984.
- [Ge89] E.-U. Gekeler, *On the de Rham isomorphism for Drinfeld modules*, J. Reine Angew. Math. 401 (1989), 188–208.
- [GK79] B. Gross and N. Koblitz, *Gauss sums and the p -adic Γ -function*, Annals of Math., **109** (1979), pp. 569–581.
- [GL11] A. Genestier and V. Lafforgue, *Théorie de Fontaine en égales caractéristiques*, Ann. Sci. École Norm. Supér. 44 (2) (2011) 263–360.
- [Go80] D. Goss, *Modular forms for $\mathbb{F}_r[T]$* , J. Reine Angew. Math. 317 (1980) 16–39.
- [Go88] D. Goss, *The Γ -function in the arithmetic of function fields*, Duke Math. J. **56** (1988), no. 1, 163–191.
- [Har11] U. Hartl, *Period Spaces for Hodge Structures in Equal Characteristic*, Ann. Math. 173 (3) (2011) 1241–1358.
- [HaJu20] U. Hartl and A.-K. Juschka, *Pink’s theory of Hodge structures and the Hodge conjecture over function fields in: t -Motives: Hodge Structures, Transcendence and Other Motivic Aspects*, Eur. Math. Soc., Zürich, 2020, pp. 31–182.
- [HaKi20] U. Hartl and W. Kim, *Local Shtukas, Hodge–Pink structures, and Galois representations in: t -Motives: Hodge Structures, Transcendence and Other Motivic Aspects*, Eur. Math. Soc., Zürich, 2020, pp. 183–259.
- [La90] S. Lang, *Cyclotomic Fields I and II*. Graduate Texts in Mathematics, vol. 121, Springer-Verlag, New York, 1990.
- [La02] S. Lang, *Algebra–Revised Third Ed.* Graduate Texts in Mathematics 211, Springer Science+Business Media New York 2002.
- [MFY97] M. Mendès France, J. Yao, *Transcendence of the Carlitz–Goss gamma function*, J. Number Theory **63** (1997) 396–402.
- [Mo75] Y. Morita, *A p -adic analogue of the Γ -function*, J. Fac. Sci. Univ. Tokyo, Sec. IA, Math., **22** (1975), 255–266.
- [NP22] C. Namoiyam and M. A. Papanikolas, *Hyperderivatives of periods and quasi-periods for Anderson t -modules*, to appear in Memoirs of the American Mathematical Society.
- [O89] A. Ogus, *A p -adic analogue of the Chowla–Selberg formula in: p -adic analysis*, Lecture notes in Mathematics 1454, Springer-Verlag Berlin Heidelberg 1990.
- [P08] M. A. Papanikolas, *Tannakian duality for Anderson-Drinfeld motives and algebraic independence of Carlitz logarithms*, Invent. Math. **171** (2008), 123–174.
- [Pe08] F. Pellarin, *Aspects de l’indépendance algébrique en caractéristique non nulle (d’après Anderson, Brownawell, Denis, Papanikolas, Thakur, Yu, et al.)*, Astérisque(2008), no. 317, Exp. No. 973, viii, 205–242.
- [Pink97] R. Pink, *Hodge structure over function fields*, preprint 1997, available at the following link: <https://people.math.ethz.ch/~pink/ftp/HS.pdf>.
- [Sin97] S. K. Sinha, *Periods of t -motives and transcendence*. Duke Math. J. **88** (1997), 465–535.
- [T88] D. S. Thakur, *Gauss sums for $\mathbb{F}_q[T]$* . Invent. Math. **94** (1988), no. 1, 105–112.

- [T89] D. S. Thakur, *Gross-Koblitz formula for function fields* in: *p*-adic analysis, Lecture notes in Mathematics 1454, Springer-Verlag Berlin Heidelberg 1990.
- [T91] D. S. Thakur, *Gamma functions for function fields and Drinfeld modules*, Ann. of Math. (2) **134** (1991), no. 1, 25-64.
- [T96] D. S. Thakur, *Transcendence of gamma values for $\mathbb{F}_q[T]$* , Ann. of Math. (2) **144** (1996), no. 1, 181-188.
- [T98] D. S. Thakur, *D. Thakur, Automata and Transcendence*, Number theory (Tiruchirapalli, 1996), Con-temp. Math. 210 Amer. Math. Soc. Providence, RI (1998), 387-399.
- [T04] D. S. Thakur, *Function Field Arithmetic*, World Scientific Publishing, River Edge, NJ, 2004.
- [Wei22] F.-T. Wei, *Algebraic relations among special gamma values and the Chowla-Selberg phenomenon over function fields*, preprint (arXiv:2207.01165).
- [WY02] Z.-Y. Wen and J.-Y. Yao, *Transcendence, automata and gamma functions for polynomial rings*, Acta Arith. 101 (2002), 31-51.
- [Yu90] J. Yu, *On periods and quasi-periods of Drinfeld modules*, Compositio Math. 74 no. 3 (1990) 235-245.

DEPARTMENT OF MATHEMATICS, NATIONAL TSING HUA UNIVERSITY, HSINCHU CITY 30042, TAIWAN R.O.C.

Email address: cychang@math.nthu.edu.tw

DEPARTMENT OF MATHEMATICS, NATIONAL TSING HUA UNIVERSITY, HSINCHU CITY 30042, TAIWAN R.O.C.

Email address: ftwei@math.nthu.edu.tw

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI CITY 10617, TAIWAN R.O.C.

Email address: yu@math.ntu.edu.tw