

# SQUARED DISTANCE FUNCTION ON THE CONFIGURATION SPACE OF A PLANAR SPIDER WITH APPLICATIONS TO HOOKE ENERGY AND VORONOI DISTANCE

MACIEJ DENKOWSKI, GAIANE PANINA, DIRK SIERSMA

ABSTRACT. Spider mechanisms are the simplest examples of arachnoid mechanisms, they are one step more complicated than polygonal linkages. Their configuration spaces have been studied intensively, but are yet not completely understood. In the paper we study them using the Morse theory of the squared distance function from the "body" of the spider to some fixed point in the plane. Generically, it is a Morse-Bott function. We list its critical manifolds, describe them as products of polygon spaces, and derive a formula for their Morse-Bott indices. We apply the obtained results to Hooke energy and Voronoi distance.

## 1. INTRODUCTION

It has always been fruitful to study the configuration space of a mechanical linkage using Morse or Morse-Bott theory. As an example, we would like to remind the reader that the homology groups of a polygon space in  $\mathbb{R}^3$  were computed by treating the squared length of a diagonal of a flexible polygon as a Morse function in [6]. Another example: the smoothness of polygon spaces follows from another squared distance function defined on the configuration space of a robot arm.

In this paper we study new classes of mechanical linkages playing an important and increasing role in applied problems, that is, *spider mechanisms*. At the present time the topology and control of multi-polygonal and spider mechanisms is a largely open topic. Some information about their configuration spaces can be derived from a paper of Mounoud [12] and the results of Blanc, Shvalb, Shoram [15, 16], O'Hara [14], Kamiyama and Tsukada [10]. In the present paper we study planar spider mechanisms using the Morse theory of the squared distance function from the "body" of the spider to some fixed point in the plane.

In section 2 we list the properties of spider mechanisms.

In section 3 we treat the critical point theory. Generically, the squared distance is a Morse-Bott function on the configuration space. We list its critical manifolds, show that they are equal to products of polygon spaces in Theorem 1 and derive a formula from their Morse-Bott indices in Theorem 2.

We use the Morse-Bott inequalities for the relation between critical points and the topology, and give examples where the squared distance is a perfect Morse-Bott function.

---

2020 *Mathematics Subject Classification*. 58K05, 70B15.

*Key words and phrases*. Spider linkage, work space, critical points, Morse-Bott theory, Hooke energy, Voronoi distance.

We show that (weighted) Hooke energy is equivalent to the squared distance to the (weighted) centre of gravity. That means that we also treated the critical point theory of Hooke Energy.

Another potential function is the Voronoi distance discussed in section 6. Based on the critical point theory in the plane from [17], we explore its counterpart on the configuration space.

We close in section 7 with indicating possible generalizations.

We thank George Khimshiashvili for useful discussions. This work is the result of a joint *Recherche en Residence* (RER) in CIRM (Luminy). We thank CIRM for the hospitality and excellent working atmosphere.

## 2. BASIC NOTIONS: SPIDER MECHANISM, ITS CONFIGURATION SPACE AND WORK SPACE. ROBOT ARMS AND POLYGON SPACES

**2.1. Spider mechanisms and spiders.** We will use the convention of denoting points in  $\mathbb{R}^2$  with capital letters, while the corresponding vectors will be indicated by lower-case letters; thus  $A$  is a point, while  $a$  the corresponding vector, hence the notation  $\|a - b\| = |AB|$ . We start by recalling the notion of a spider (arachnoid mechanism) in a 2-dimensional space.

**Definition 1.** An  $n$ -legged *spider mechanism* in  $\mathbb{R}^2$  consists of

- (1) a set of  $n$  fixed, pairwise distinct points (the *feet*)  $A_1, \dots, A_n \in \mathbb{R}^2$ ,
- (2) a collection of natural numbers  $p_1 > 1, \dots, p_n > 1$ , and
- (3) a collection of lengths  $\ell_{i,j+1} > 0$ ,  $j = 1, \dots, p_i$ .

An  $n$ -legged *spider* with  $(p_1, \dots, p_n)$ -articulated legs in  $\mathbb{R}^2$  is a *configuration* of a given spider mechanism. It consists of

- (1) a center point  $X \in \mathbb{R}^2$  (the *body* of the spider that will be moving around), and
- (2) the set of  $n$  *legs*  $\mathcal{L}_i$ , that are each a connected chain of  $p_i$  *edges* (rigid bars), from  $A_i$  to  $X$  i.e., for each  $i \in \{1, \dots, n\}$ , a union of segments  $[A_i^j, A_i^{j+1}] \subset \mathbb{R}^d$  of fixed lengths  $\ell_{i,j+1} > 0$ ,  $j = 1, \dots, p_i$  where  $A_i^0 = A_i$ ,  $A_i^{p_i} = X$ . The points  $A_i^j$  for  $j = 1, \dots, p_i - 1$  are called the *joints* of the  $i$ -th leg.

One imagines a spider mechanism with fixed feet and with the legs that are allowed to rotate freely around the joints. We also allow intersections and self-intersections of the legs.

**Definition 2.** The *configuration space* of a given spider mechanism is the set of all the possible positions taken by its body and joints. We call it the *spider space*  $\mathcal{S}$ :

$$\mathcal{S} := \{(X, A_1^1, \dots, A_1^{p_1-1}, A_2^1, \dots, A_n^{p_n-1}) \in (\mathbb{R}^2)^{1+N} \mid |A_i^j A_i^{j+1}| = \ell_{i,j+1}, i = 1, \dots, n, j = 0, \dots, p_i - 1\},$$

where  $N = \sum_{i=1}^n p_i - n$ . It comes together with the natural projection to  $\mathbb{R}^2$  called the *work map*

$$\pi: \mathcal{S} \rightarrow \mathbb{R}^2,$$

sending a *spider* (i.e. a configuration)  $\xi = (X, A_1, \dots, A_N) \in \mathcal{S}$  to its body  $X \in \mathcal{W}$ . The image  $\mathcal{W} = \pi(\mathcal{S})$  is called the *work space* of the spider mechanism.




FIGURE 1. Spider in the plane.

The spider space and the work map have been studied in several papers mentioned in the Introduction.

**2.2. Robot arms: spiders with one leg.** A special case is the spider with only one leg. This mechanism is also called a (*free*) *robot arm*. Its underlying graph is a path graph with start point  $A$  and end point  $X$ . It is convenient to imagine that the path graph is directed so that each edge is oriented from  $A$  to  $X$ . The spider space of the 1-leg spider does not depend on the edge lengths. It is the torus  $(S^1)^p$ , where  $p$  is the number of edges. However, people usually work with configurations of the robot arm modulo the rotation group around the point  $A$ . With this point of view the configuration space of a robot arm is the torus  $(S^1)^{p-1}$ . It is convenient to imagine  $(S^1)^{p-1}$  as the space of all configurations with a fixed first bar. And this is the point of view we adopt in the Proposition 1 below.

A configuration of a robot arm is *aligned* if it fits in a straight line. For a generic robot arm, in an aligned configuration, each edge (geometrically) is co-directed either with  $\overrightarrow{AX}$ , or with  $\overrightarrow{XA}$ . The edges co-directed with  $\overrightarrow{AX}$  are called *positively directed*, the other ones are *negatively directed*.

**Definition 3.** For an aligned configuration  $\mathcal{L}$ , denote by  $Pos(\mathcal{L})$  the number of positively directed edges. Define also

$$M(\mathcal{L}) = Pos(\mathcal{L}) - 1.$$

We are going to multiply exploit the squared distance function between the endpoints of a robot arm. In particular, we shall use the following:

**Proposition 1.** ([8]) *The squared distance between the endpoints of a robot arm  $\|a - x\|^2$  (defined on the configuration space i.e. composed with the work map, cf. Section 3) is a Morse-Bott function, whose critical points and critical manifolds are:*

- (1) *isolated critical points that correspond to aligned configurations, and*
- (2) *the set  $\{\xi \in \mathcal{S} \mid \|a - \pi(\xi)\| = 0\}$ , which exhibits the global minimum (if it is non-empty).*

*The Morse index of an aligned configuration equals  $M(\mathcal{L})$ .*

**2.3. Polygonal linkages and polygon spaces: spiders with two legs.** A spider with two legs reduces to a polygon space by adding a rigid bar connecting the two feet; now edge lengths do matter.

A *polygonal  $p$ -linkage* is a sequence of positive numbers  $L = (l_1, \dots, l_p)$ . We interpret it as a collection of  $p \geq 3$  rigid bars of lengths  $l_i$  joined consecutively in a closed chain by revolving joints. We always assume that the triangle inequality holds, so the chain can close. A *planar configuration* of  $L$  is a closed broken line whose consecutive edge lengths are  $l_1, \dots, l_p$ . As follows from the definition, a configuration may have self-intersections.

The *configuration space*, or the *polygon space* for short, is the set of all configurations of  $L$  modulo orientation preserving isometries of  $\mathbb{R}^2$ . It does not depend on the ordering of  $l_i$  but depend on the values of the  $l_i$ .

The polygon space is a closed  $(p - 3)$ -dimensional manifold iff there are no aligned configurations, that is, no configuration of  $L$  fits in a straight line. There are explicit linear conditions in  $(l_1, \dots, l_p)$  for the smoothness called *Grashof conditions*. For these facts and for more information about polygon spaces see [3] and [8].

For spiders with an arbitrary number of legs we have the following criterion:

**Proposition 2** ([15], see also [7], [12]). (1) *The singular points of the spider space  $\mathcal{S}$  correspond to:*

- *either two aligned legs lying in one and the same straight line,*
- *or three aligned legs.*

(2) *Generically, the spider space  $\mathcal{S}$  is a smooth, compact, orientable, algebraic manifold whose dimension is*

$$2 - 2n + \sum_{i=1}^n p_i.$$

**2.4. The work map.** Consider the fibers of the work map  $\pi: \mathcal{S} \rightarrow \mathcal{W}$ . We fix the point  $X$  in work space, different from the feet points  $A_i$ . We are left with the freedom of  $n$  legs with fixed starting points  $A_i$  and end point  $X$ . Each of them constitute a polygonal linkage if we add the edge  $XA_i$  to  $\mathcal{L}_i$  with length  $l_{i,0} = \|x - a_i\|$ : *the closure of the leg*. Each of these linkages defines a polygon space. If  $X = A_i$ , the  $i$ -th leg itself is closed and yields a polygonal linkage. It follows:

**Proposition 3.** (1) *The fiber of the work map over  $X \neq A_i$  is the product of the polygon spaces of the linkages, which are the closures of the legs with endpoint  $X$ .*

(2) *If  $X = A_i$ , the fiber of the work map over  $A_i$  equals the product of the polygon space of the corresponding linkage with a circle  $S^1$  and the polygon spaces of the other leg closures.*

**2.5. Discs-Annuli structure on the work space.** Fix a foot  $A_i$  and assume for the moment that the leg attached to it is moving freely in  $\mathbb{R}^2$ . Consider for the leg  $\mathcal{L}_i$  the squared distance between the foot and the body

$$\|a_i - x\|^2.$$

According to Proposition 1 its critical points correspond to aligned legs, (i.e. all the points  $A_i^j$  lie on the line  $\mathbb{R}(x - a_i)$ ). The critical points project via  $\pi$  to circles with center




FIGURE 2. Critical circles and work space.

$A_i$  and *critical radii*  $r_i^{\epsilon_i}$ :

$$C(A_i, r_i^{\epsilon_i}) := \{z \in \mathbb{R}^2 \mid \|z - a_i\| = r_i^{\epsilon_i}\}.$$

with  $r_i^{\epsilon_i} = |\sum_{j=1}^{p_i} \epsilon_{i,j} l_{i,j}|$  and  $\epsilon_i = (\epsilon_{i,j})_{j=1}^{p_i}$ ;  $\epsilon_{i,j} \in \{-1, +1\}$ . This construction includes the possibility that (for a non-generic case) a critical radius is zero and the circle degenerates to a point.

We denote by  $r_i$ , respectively  $R_i$ , the minimal and the maximal values of  $\|x - a_i\|$ . They define the zone:

$$\mathcal{Z}(A_i) := \{x \in \mathbb{R}^d \mid \rho_i \leq \|x - a_i\| \leq R_i\},$$

which is either a disc or an annulus with center  $A_i$ .

Here

$$\rho_i = \begin{cases} r_i, & \text{if the leg cannot close;} \\ 0, & \text{otherwise.} \end{cases}$$

The zone  $\mathcal{Z}(A_i)$  is the work space of the free leg.

The following three facts have obvious proofs.

**Proposition 4.** *The work space of a spider linkage with  $n$  legs is the intersection of all the zones:  $\mathcal{W} = \bigcap_{i=1}^n \mathcal{Z}(A_i)$ .*

**Proposition 5.** *Assume  $\mathcal{S}$  is smooth. The critical locus of the projection  $\pi: \mathcal{S} \rightarrow \mathcal{W}$  consists of the fibers with at least 1 aligned leg. The discriminant set coincides with those points of the work space  $\mathcal{W}$ , that lie on one of the circles  $C(A_i, r_i^{\epsilon_i})$ .*

**2.6. The Euler characteristic of the spider space.** The discriminant divides the work space in strata  $\sigma$  of dimension 0,1 and 2 in such a way that above each stratum the fibre of  $\pi$  has constant topological type.

**Proposition 6.** *In the situation considered, the Euler characteristic of the spider is*

$$\chi(\mathcal{S}) = \sum_{\sigma} (-1)^{\dim(\sigma)} \chi(\sigma) \chi(\pi^{-1}(x_{\sigma}))$$

where the sum ranges over all the strata, and  $x_{\sigma}$  is any point of the stratum  $\sigma$ .

The proof follows from the properties of the Euler characteristic. This formula extends the formula in [12] for legs with only two edges to the general case. The fibers are polygon spaces. In the smooth case their Betti numbers and Euler characteristic have been computed in [4]. See an example in section 4.

### 3. THE SQUARED DISTANCE FUNCTION AS A MORSE FUNCTION

**3.1. Squared distance function defined on the spider space.** For a given spider mechanism, let  $Z \in \mathbb{R}^2$  be a fixed point. The squared distance from the body  $X$  to  $Z$  is a function defined on the spider space. More precisely, set  $D(x) = \|x - z\|^2$ , for  $x \in \mathbb{R}^2$  and  $\mathbb{D}(\xi) = D(\pi(\xi))$ , for  $\xi \in \mathcal{S}$ . In this section we intend to study the critical point theory of  $\mathbb{D}$ . We will assume that the spider space  $\mathcal{S}$  is smooth, i.e. satisfies the conditions mentioned in Proposition 2, although the non-smooth case is also of interest.

**Definition 4.** A smooth function  $f : M \rightarrow \mathbb{R}$  on a smooth manifold  $M$  is called a *Morse-Bott function* if and only if its critical set  $\Sigma(f)$  is a disjoint union of connected submanifolds, and for each connected component  $C \subset \Sigma(f)$  the Hessian of the restriction of  $f$  to a normal slice is non-degenerate for all  $x \in C$ . The index of this Hessian is called the Morse-Bott index or transversal Morse index of the connected component  $C$ .

For properties of Morse-Bott functions we refer to Bott [1], the textbook [13] and to section 3.4 on the Morse-Bott polynomial.

We shall show that generically,  $\mathbb{D}$  is a Morse-Bott function. The *genericity* notion now becomes more restrictive. It includes the conditions of smoothness of the spider space, but also some more conditions on the relative position of  $Z$ , that will be specified later. Anyhow, the genericity condition defines an open everywhere dense subset of the parameter space of spider mechanism.

**3.2. Critical points.** Assume we have a spider mechanism with a smooth spider space  $\mathcal{S}$ . For its leg  $\mathcal{L}_i$ , define its extension  $z\mathcal{L}_i$  as the robot arm obtained by adding the edge  $ZA_i$  before the leg  $\mathcal{L}_i$ . This robot arm starts with  $Z$ , ends at  $X$ , and has  $p_i + 1$  edges.

Let us first consider mechanisms with only one leg. We assume that the following *genericity condition* holds:

( $\star$ ) no two points  $A$ ,  $X$ , and  $Z$  coincide in an aligned configuration of the leg.

Proposition 1 implies:

**Proposition 7.** *For a generic one-leg spider, the critical points of  $\mathbb{D} = \|x - z\|^2$  are:*

- (1) *A spider  $\xi$  with  $X \neq Z$  iff it is aligned, and the point  $Z$  lies on the straight line containing the spider. In this case the critical point is Morse and the Morse index is*

$$M(\xi, Z) = \text{Pos}(z\mathcal{L}) - 1$$

*In terms of the leg  $\mathcal{L}$  this can be written as*

$$M(\xi, Z) = \begin{cases} p - \text{Pos}(\mathcal{L}), & \text{if } X \text{ lies between } A \text{ and } Z; \\ \text{Pos}(\mathcal{L}) - 1, & \text{if } Z \text{ lies between } A \text{ and } X; \\ \text{Pos}(\mathcal{L}), & \text{if } A \text{ lies between } Z \text{ and } X. \end{cases}$$

- (2) The (possibly empty) submanifold of  $\mathcal{S}$  defined by  $X = Z$  is the global minimum manifold of  $\mathbb{D}$ . It is a Morse-Bott critical submanifold with Morse-Bott index 0.

The manifold is diffeomorphic to the configuration space of the polygonal linkage obtained by adding the bar  $ZX$  to the robot arm.

*Proof.* (1) The statement follows straightforwardly. One should only take into account that in the first case (when  $X$  lies between  $A$  and  $Z$ ) the positively directed edges turn to negatively directed ones. (2) The function  $\mathbb{D}$  is the composition of a submersion and the Morse function, which attains its minimum at  $X = Z$ .  $\square$

Now pass to a spider with  $n$  of legs.

**Theorem 1.** For a smooth spider space, a configuration  $\xi \in \mathcal{S}$  (satisfying  $(\star)$ ) is critical for  $\mathbb{D}$  iff it satisfies at least one of the following conditions:

- (1)  $X = Z$ ;
- (2) One leg is aligned, and the point  $Z$  lies on this line;
- (3) Two different legs are aligned.

*Proof.* For  $X = Z$  we clearly have the global minimum, so below we assume  $X \neq Z$ .

The derivative  $d_\xi \mathbb{D}$  of  $\mathbb{D}$  is a composition of

$$d_\xi \pi : T_\xi \mathcal{S} \rightarrow T_{\pi(\xi)} \mathbb{R}^2 \quad \text{and} \quad d_x D : T_{\pi(\xi)} \mathbb{R}^2 \rightarrow \mathbb{R}.$$

Here  $d_x D$  is a submersion. If  $\xi$  does not belong to the critical locus of  $\pi$  then we get the composition of two submersions, so  $\mathbb{D}$  is regular at  $\xi$ .

As soon as  $\xi$  has exactly one aligned leg, the image of  $d_\xi \pi$  is 1-dimensional (tangent space to a critical circle). The composition has maximal rank if  $ZX$  is orthogonal to this image (or: the level curves of  $\|x - z\|^2$  are tangent to the critical circle). As soon as  $\xi$  has two non-parallel aligned legs, the image of  $d_\xi \pi$  is 0-dimensional and therefore the composition is not regular. The conditions for smoothness from Proposition 2 allows us to restrict to the above cases.  $\square$

**Definition 5.** A pair (a spider mechanism, a point  $Z$ ) is called *strongly generic* if:

- (1) Its configuration space is smooth:
  - (a) No configuration has more than two aligned legs.
  - (b) Two aligned legs are never parallel.
- (2) In case of two aligned legs, the point  $Z$  never lies on any of these lines.
- (3) The point  $Z$  coincides with none of the feet  $A_i$ .
- (4) For one aligned leg,  $Z$  never coincides with  $X$ .

**3.3. Morse indices.** Assuming strong genericity, we start with some preparations. Let us have a deeper look at case 3 from Theorem 1. Assume that there are two aligned legs (say, with feet at  $A_1$  and  $A_2$ ). The three points  $A_1, A_2, X$  define a partition of the plane (cf Figure 3) into four quadrants separated by the lines  $\mathbb{R}[A_1X]$  and  $\mathbb{R}[A_2X]$ . Let  $Z$  be in one of the quadrants and define the following signs, depending on  $Z$  for the 2 aligned legs at  $X$ :

$$\sigma(Z) = (\sigma_1(Z), \sigma_2(Z)) = (\text{sign}(A_1X \cdot ZX), \text{sign}(A_2X \cdot ZX))$$




FIGURE 3. Partition of the plane.

**Theorem 2.** *For a strongly generic spider and a point  $Z$ , the function  $\mathbb{D} = \|x - z\|^2$  is a Morse-Bott function. The Morse-Bott indices  $\mu$  are as follows:*

- (1) *The condition  $X = Z$  defines a Morse-Bott critical manifold with the index*

$$\mu = 0.$$

- (2) *One leg  $\mathcal{L}$  is aligned and  $Z$  lies on this line. The Morse-Bott index of the critical manifold is*

$$\mu = M(\mathcal{L}, Z),$$

*see Proposition 7, where  $\mathcal{L}$  is the one-legged spider obtained by eliminating all the legs except for the aligned one.*

- (3) *Two legs are aligned. Then the Morse-Bott index is  $\mu = M_1 + M_2$ , where  $M_i$  is defined by*

$$M_i = \begin{cases} M(\mathcal{L}_i), & \text{if } \sigma_i(Z) > 0 \\ p_i - M(\mathcal{L}_i), & \text{if } \sigma_i(Z) < 0. \end{cases}$$

*Here  $M(\mathcal{L})$  is defined in Definition 3.*

*The critical manifolds in the three cases are diffeomorphic to:*

- (1) *The Cartesian product of  $n$  polygon spaces (obtained by adding the bar  $A_i X$  to the leg  $\mathcal{L}_i$ ),*
- (2) *The Cartesian product of  $n - 1$  polygon spaces (obtained by closing non-aligned legs in a similar way),*
- (3) *The Cartesian product of  $n - 2$  polygon spaces, also arising from closing of non-aligned legs.*

*Proof.* (1) is clear since  $X = Z$  defines the global minimum manifold.

(2) By the strong genericity assumption, only one leg of the spider is aligned. Non-aligned legs contribute to the Morse-Bott critical manifold only, which is the product of polygonal spaces, so we eliminate them and leave only the aligned leg. The statement follows from Proposition 1.

(3) By the strong genericity assumption, only two legs of the spider are aligned, and the lines that contain the legs are different. As in (2), we can eliminate the non-aligned legs and leave only the two aligned legs. Denote the configuration space of the two-legged spider by  $\mathcal{S}'$ . Its dimension is  $p_1 + p_2 - 2$ .



Let us compute the 2-jet of  $\mathbb{D}$  at the critical point. The first order terms vanish since we have a critical spider. Set  $r_i := |A_i X| = \|A_i X_0\| + \sum_j \pm \xi_{i,j}^2$  near the aligned legs, where  $X_0$  is the body of the critical spider, that is, intersection point of the two critical circles. Near  $X_0$  we use  $(r_1, r_2)$  as coordinates for  $X$ . Define the unit vectors  $e_1, e_2$  at the point  $X_0$ , having the same directions as  $A_1 X_0$  and  $A_2 X_0$  respectively.

Denote by  $\psi$  the map from local coordinates  $\xi$  to the spider space to  $X = (r_1, r_2)$ , and consider the second order term in the Taylor series expansion of  $D \circ \psi$ . Only the 2-jet of  $\psi$  and the 1-jet of  $D = \|x - z\|^2$  at  $x_0$  are important. The latter can be written as  $\|z - x_0\| + (z - x_0) \cdot x$ .

The composition of the two maps give now

$$c + (z - x_0) \cdot (x_0 + \sum_j \pm \xi_{1,j}^2 e_1 + \sum_j \pm \xi_{2,j}^2 e_2) = \|x_0 - z\| + \alpha_1 \sum_j \pm \xi_{1,j}^2 + \alpha_2 \sum_j \pm \xi_{2,j}^2$$

with  $\alpha_1 = (z - x_0) \cdot e_1$  and  $\alpha_2 = (z - x_0) \cdot e_2$ , which yields the result required.  $\square$

**3.4. Morse-Bott Polynomials.** In the case of isolated singularities the counting of critical points via a generating polynomial, *the Morse polynomial*, relates the critical points of the squared distance  $\mathbb{D}$  with the topology of  $\mathcal{S}$ .

The Morse-Bott polynomial related to the function  $\mathbb{D}$  is defined by

$$P_{\mathbb{D}}(t) = \sum_{\Sigma} P_{\Sigma}(t) \cdot t^{i(\Sigma)} := \sum \mu_k(\mathbb{D}) t^k$$

where  $P_{\Sigma}(t) = \sum b_k(\Sigma) t^k$  is the Poincaré polynomial,  $b_k(\Sigma)$  the  $k^{th}$  Betti number and  $i(\Sigma)$  the transversal Morse index of the Morse-Bott singular component  $\Sigma$ . One computes the coefficients  $\mu_k$  by performing the multiplication and summation. In the case of isolated singularities  $\mu_k$  coincides with the number of Morse points.

From [1, 13] we recall

$$P_{\mathbb{D}}(t) = P_{\mathcal{S}}(t) + (1 + t)Q(t)$$

where  $P_{\mathcal{S}}(t)$  is the Poincaré polynomial of  $\mathcal{S}$  and  $Q(t)$  is a polynomial with non-negative integer coefficients. From this follows Morse inequalities:  $\mu_k \geq b_k$ . Note that:

- $\mu_k$  depends on the position of  $Z$ , but is locally constant on the complement of a bifurcation set.
- For  $t = -1$  we get the Euler-Poincaré characteristic  $\chi(\mathcal{S}) = \sum (-1)^k \mu_k$ : independent of  $Z$ .


In the next section we will use the Morse-Bott polynomial to get information about the topology of  $\mathcal{S}$ , i.e. for computing  $\chi(\mathcal{S})$  and upper bounds for the Betti numbers of  $\mathcal{S}$ . For this we need to know  $P_{\Sigma}$  and also the Betti-numbers of spider spaces. We refer to the paper of Farber and Schütz [4], where there is a complete result in terms of short, medium and long legs. Examples can be found in section 4.

#### 4. EXAMPLES

In this section we compute critical points of the squared distance function and their indices and compare them with topological invariants of  $\mathcal{S}$ . For efficiency of the exposition we restrict to examples with a few legs only.

4.1. **A tripod.** Spider mechanisms with only two edges for each leg are easy to analyze. In strongly generic cases  $\mathbb{D}$  is a Morse function, and there are isolated critical points only.

As an example, consider a spider whose feet are positioned at the vertices of an equilateral triangle, and the edge lengths are as is depicted in Figure 4.



(Above) The work space of the tripod.  
 (Right) The critical configurations of the tripod: (a) minima in  $Z$ ,  
 (b) saddles with one aligned leg,  
 (c) maxima with one aligned leg,  
 (d) saddles with two aligned legs.

FIGURE 4. Work space and critical configurations.

The spider space  $\mathcal{S}$  is two-dimensional manifold. It is a patch of eight copies of the hexagonal work space, so its genus is 3. See [12] and Proposition 2.6.

Assume that  $Z$  is positioned at the center of the triangle. The global minimum of  $\mathbb{D}$  is attained when  $X = Z$ , which corresponds to eight different configurations.

There saddle points are of two types, with one leg is aligned (12 ones) and with two legs aligned (also 12 ones).

There are 12 points where maximum is attained: one leg is aligned. We get 44 isolated critical points and Euler-characteristic  $-4$ . Other choices of  $Z$  (e.g outside work space) will reduce the number of critical points. We will exploit this idea in the next example.

4.2. **Second Example.** We next compute Morse-Bott polynomials in the case of arbitrary number of edges for each leg. In this example we assume only two legs, with  $p_1$ , resp  $p_2$  edges (more or less of equal length). The dimension of spider space is  $p_1 + p_2 - 2$ . We assume moreover that the work space is the region bounded by the two critical circles of maximum radius and does not contain arcs of other critical circles. We will consider two different choices of the point  $Z$  (see Figures 5 and 6).

The first choice is that point  $Z$  is in the interior of the work space. According to Theorem 2 there are 5 critical components, related to the point  $Z$  and the points  $X_1, X_2, X_3, X_4$  in the Figure. The singular components are as follows:

- Type(2) For the points  $X_3$  and  $X_4$ : each just one point  $\bullet$ : isolated singularity, clearly a maximum;
- Type(1) For the points  $X_1$  and  $X_2$ : the polygon space of the two polygons, containing  $X_1$ , resp  $X_2$  and the points  $A_2$ , resp  $A_1$ . Their Poincaré polynomials can be computed by the formula in [4]. But one can see also directly that they are

homeomorphic to spheres  $S^{p_2-2}$ , resp  $S^{p_1-2}$  since the linkages contain one very long leg in comparison with the other legs (we omit details).

- Type (0) For the point  $Z$ : the product of the two polygon spaces  $S^{p_2-2} \times S^{p_1-2}$ .

The indices and the contributions to the Morse-Bott polynomial are contained in the table in Figure 5




FIGURE 5.  $Z$  lies in the interior.

It follows that the Morse-Bott polynomial can be written as:

$$P_{\mathbb{D}}(t) = 1 + (1 + t)(t^{p_1-2} + t^{p_2-2}) + (1 + t)^2 t^{p_1+p_2-4} + t^{p_1+p_2-2}.$$

As a consequence: The Euler characteristic is  $1 + (-1)^{p_1+p_2-2}$ , which is 2 if  $p_1 + p_2$  is even and 0 else. The Morse-Bott inequalities imply that several  $b_i = 0$  and the others are bounded by 1 or 2. But we can do better by another choice of  $Z$ .

The second choice is the point  $Z_1$  is outside the work space and chosen in such a way that critical points of type (1) do not occur. We have now only critical points of type (2), which are isolated: one minimum and one maximum with Morse polynomial  $P_{\mathbb{D}}(t) = 1 + t^{p_1+p_2-2}$ . Indeed since any Morse function with two critical points can only occur on a sphere, we have in our case  $S^{p_1+p_2-2}$ .





FIGURE 6.  $Z_1$  lies in the exterior.

NB. This answer can be checked since our spider with 2 legs is also a polygon space. We can also apply Proposition 2.6 to compute the Euler characteristic. This gives:  $\chi(\mathcal{S}) = 2 - \chi(S^{p_1-1}) - \chi(S^{p_2-1}) + \chi(S^{p_1-1})\chi(S^{p_2-1}) = 1 + (-1)^{p_1+p_2-2}$ .

We next consider a similar case, which includes a critical arc of non-extremal type. We get the table:



The Morse Bott polynomial is  $1 + t + t^{p_1+p_2-3} + t^{p_1+p_2-2}$ . Via Morse-Bott-‘inequalities’ it follows that in this example: the squared distance is a perfect Morse function ( $p_1 + p_2 \geq 6$ ).

## 5. HOOKE ENERGY

This was our starting point in this study of spiders. The case of 3 legs with two edges each has been studied before in [9]. It will turn out that weighted Hooke energy and squared distance give equivalent results.

The Hooke Energy potential  $H : \mathbb{R}^2 \rightarrow \mathbb{R}$  is defined by  $H(x) = \sum_{i=1}^n \|x - a_i\|^2$ . The stationary points of  $H$  are determined by:  $\nabla H = nx - 2 \sum_{i=1}^n \|x - a_i\| = 0$ , so  $x = \frac{1}{n} \sum_{i=1}^n a_i$ , the centre of gravity. There are no other stationary points on  $\mathbb{R}^2$ . Note that  $H(X) = \|x - z\|^2 + \kappa$ , where  $z = \frac{1}{n} \sum_{i=1}^n a_i$ , the centre of gravity  $Z$  and  $\kappa = \sum_{i=1}^n \|a_i\|^2 - \|z\|^2$ . All level curves are circles.

**Proposition 8.** *The Hooke energy  $H$  is an affine linear transform of the squared distance function to the center of gravity  $Z$ .*

$$H(x) = n\|x - z\|^2 + \kappa.$$

*The Hooke function is minimal in the center of gravity. Both functions have the same critical point theory.*

Given weights  $w = w_1, \dots, w_n$  to the Hooke potential, we get a weighted Hooke potential

$$H_w(x) = \sum_{i=1}^n w_i \|x - a_i\|^2.$$

NB. We will allow also negative weights. Weighted Hooke Energy is also an affine linear transform of the weighted squared distance function.

We extend the definitions to spider space:  $\mathbb{H}(\xi) = H(\pi(\xi))$  and similarly  $\mathbb{H}_w(\xi) = H_w(\pi(\xi))$ .

**Theorem 3.** *Weighted Hooke Energy and Squared Distance to the weighted center of gravity have equivalent critical point theory on spider space  $\mathcal{S}$ .*

NB. Depending on the sign of  $w_i$  the indices of critical points can become dual.

6. VORONOI DISTANCE

6.1. **Critical points.** In this section we will consider on the generic *smooth* spider space the *Voronoi distance function*  $\min_{1 \leq i \leq n} \|x - a_i\|$ . To smoothen it at the feet  $A_i$  and get a nicer gradient without modifying the level sets we will in fact work with

$$V(x) = \min_{1 \leq i \leq n} \|x - a_i\|^2 = \text{dist}(x, \{A_1, \dots, A_n\})^2.$$

This function is defined on the spider space  $\mathcal{S}$ , i.e.  $\mathbb{V}(\xi) := V(\pi(\xi))$ . A major role is naturally played here by the *Voronoi diagram* of the feet  $A_1, \dots, A_n \in \mathbb{R}^2$ , i.e. the set

$$\mathcal{V} := \{x \in \mathbb{R}^2 \mid \exists i, j: i \neq j, \text{ and } V(x) = \|x - a_i\|^2 = \|x - a_j\|^2\}$$

and its intersection with the work space:  $\mathcal{V} \cap \mathcal{W}$ . It is in fact the set of non-differentiability points of  $V(x)$ . The function itself being locally Lipschitz, it has everywhere a well-defined Clarke subdifferential  $\partial V(x)$  and thus the notion of critical point makes sense for  $V(x)$ , i.e.  $x$  is critical iff  $0 \in \partial V(x)$  which is equivalent to saying that  $x$  belongs to the convex hull of those feet  $A_i$  that realize the distance  $V(x)$ . Otherwise, the point  $x$  is called regular.

Note that  $\mathbb{V}(\xi)$  is locally Lipschitz, too and thus we can carry over the notion of the Clarke subdifferential to this situation. This is intuitively clear, but needs some extra explanation, see the Appendix. Clearly,  $\mathbb{V}$  is differentiable outside  $\pi^{-1}(\mathcal{V} \cap \mathcal{W})$ .

The critical point theory of  $V(x)$  in the plane is well understood, see [17], obviously all the feet  $A_i$  are global minima of the Voronoi potential  $V(x)$ . However, when moving to the spider space the situation presents some interesting features, the more so since there is some additional boundary information coming along.




FIGURE 7. The figure is self-explanatory: a two-legged spider with  $\mathcal{W}$  being an annulus centered at  $A$  and with several critical circles to be taken into account. The Voronoi diagram is the dashed line.

We first discuss the Delaunay and Voronoi tessellation of the affine plane. There are several ways to define this. Here we choose the *void circle property* (vc): A point  $Y$  has the  $k$ -th *vc-property* if there exist a circle with center  $Y$ , which contains  $k$  points from among  $A_1, \dots, A_n$  and no points from the feet in its interior. The centers of circles form

the  $k$ -th void circle set. We will assume as genericity condition that no points satisfy the 4-vc condition, but remark that one can adapt the theory to a more general setting.

First we define the *Delaunay tessellation*  $DT(\mathcal{V})$ : the 0-cells are the points  $A_i$ , the 1-cells are the edges which connect those vertices  $A_i, A_j$  that are related to a point satisfying the 2-vc condition, the 2-cells are the cells characterized by triples related to points having the 3-vc property. Note that these are closed sets.

Each foot  $A_i$  defines a 2-dimensional Voronoi cell or *power cell*

$$P(A_i) := \{x \in \mathbb{R}^2 \mid \|x - a_i\|^2 \leq V(x)\},$$

its open part will be denoted by  $P^\circ(A_i)$ . The 1-dimensional Voronoi cells are the nonempty intersections  $P(A_i) \cap P(A_j)$  for  $i \neq j$ , the 0-cells are the triple intersections  $P(A_i) \cap P(A_j) \cap P(A_k)$ . This tessellation is dual to the Delaunay tessellation.

**Theorem 4** (e.g. [11]). *Critical points (in the topological sense) of the Voronoi distance function are precisely the intersections between a Delaunay-cell and its dual Voronoi cell. The index of the critical point is equal to the dimension of the Delaunay cell.*

It follows that all the points  $A_i$  are minima, the intersection between Voronoi-edges and Delaunay-edges are topological saddle points and the centers of 3-void-circles are maxima; we shall use the respective notations:  $m, S, M$ . These last two types of critical points, saddles and maxima, are contained in the Voronoi diagram  $\mathcal{V}$ .

**Lemma 1.** *For a configuration  $\xi \in \mathcal{S}$  such that  $X = \pi(\xi) \notin \mathcal{V}$ , there exists exactly one power cell such that  $X \in P^\circ(A_i)$  and thus  $\mathbb{V}(\zeta) = \|\pi(\zeta) - a_i\|^2$  for  $\zeta \in \pi^{-1}(P^\circ(A_i))$ .*

Since  $\mathbb{V}$  is the composition of the work space map and the Voronoi Distance, it is good to consider its derivative  $d_\xi \mathbb{V}$ , which is well-defined at a (regular) spider  $\xi$  (i.e. with  $\pi(\xi) \notin \mathcal{V}$ ) as a composition of

$$d_\xi \pi : T_\xi \mathcal{S} \rightarrow T_{\pi(\xi)} \mathbb{R}^2 \text{ and } d_x V : T_{\pi(\xi)} \mathbb{R}^2 \rightarrow \mathbb{R}.$$

**Proposition 9** (Composition of regular maps). *If  $\xi \in \mathcal{S}$  is a regular point of the work map  $\pi$ , and  $V$  is regular at  $X = \pi(\xi)$ , then  $\mathbb{V}$  is regular at  $\xi$ .*

The proof is in the appendix. This Proposition tells us that anyhow we have regularity of  $\mathbb{V}$ , provided  $X$  is not one of the critical points of  $V$  nor lies on a critical circle. We are left with two cases for a critical point  $\xi$  of  $\mathbb{V}$ :

- (1)  $\xi$  projects on a critical point of  $V$ ,
- (2)  $\xi$  projects to a point lying on a critical circle.

**Case (1):** Since  $\pi$  is a proper mapping with continuously varying fibers, it is rather easy (see Appendix) to see that  $0 \in \partial V(\pi(\xi))$  implies  $0 \in \partial \mathbb{V}(\xi)$ . This together with the previous observations allows us to state the next result whose proof is straightforward.

**Proposition 10.** *Assume  $X = \pi(\xi)$  is a critical point of  $V$ , then so is  $\xi$  for  $\mathbb{V}$ . If moreover  $X$  does not lie on a critical circle, then  $\pi^{-1}(X)$  is a smooth component of the critical set of  $\mathbb{V}$  of codimension 2, satisfying the Morse-Bott property with transversal type, equal to the singularity type of  $V$  at  $X$  (minimum, saddle or maximum).*

**Case (2):** Inside the  $i$ -th power cell the function  $V$  reduces to the distance to the unique foot  $A_i$ , and one applies Theorem 1 with  $Z = A_i$ . But that theorem assumes that the foot is different from  $Z$  by the  $(\star)$ -condition. Therefore we have to consider two cases: On the open part of Voronoi cells of dimension 2 we consider in the work space all the non-void intersections with critical circles:  $C(A_j, r_j^{\epsilon_j}) \cap P^\circ(A_i) \cap \mathcal{W}$ . The case  $i \neq j$  is covered by Theorem 1, the case  $i = j$  gives rise to non-isolated singularities in the work space, which are discussed in the section 2.2 about robot arms.

**Proposition 11.** *For a generic smooth spider and the Voronoi distance  $\mathbb{V}$ , a configuration  $\xi \in \mathcal{S}$  with  $\pi(\xi) \neq A_j$  is critical if and only if one of the following conditions hold for  $X = \pi(\xi)$  lying in the open part of the of 2-dimensional cell  $P^\circ(A_i)$  :*

- (1) *in case  $i \neq j$ : If one leg is aligned,  $X$  is the intersection point  $C(A_j, r_j^{\epsilon_j}) \cap A_i A_j \cap \mathcal{W}$ ,*
- (2) *in case  $i \neq j$ : If two legs are aligned: the points  $X$  of intersections of 2 critical circles,*
- (3) *In case  $i = j$  we get all the intersections  $C(A_i, r_i^{\epsilon_i}) \cap P^\circ(A_i) \cap \mathcal{W}$ . These are generically arcs or circles.*

**Remark 1.** In part (1) of Proposition 11 the set  $C(A_j, r_j^{\epsilon_j}) \cap A_i A_j \cap P^\circ(A_i) \cap \mathcal{W}$  consists of isolated points. The corresponding critical set in  $\mathcal{S}$  has the Morse-Bott property. The index follows from Theorem 2(2). Similarly in (2) the Morse-Bott index follows from Theorem 2(3). In case (3) the intersections  $C(A_i, r_i^{\epsilon_i}) \cap P^\circ(A_i) \cap \mathcal{W}$ , consists of non-isolated pieces (arcs or circles). The transversal type at the intersection points is generically well defined and gives us locally the Morse-Bott property with index as in Proposition 1 about robot arms. But in general the function does not has the global Morse-Bott property. If  $X = A_j$ , then we are in Case (1): Morse-Bott with transversal minimum.

In the case of spiders, where each leg has only 2 edges, the description becomes more transparent. See the Example in Figures 8 and 9.

After studying the power cells we study next what happens with  $\mathbb{V}$  at the points of  $\pi^{-1}(\mathcal{V} \cap \mathcal{W})$ . We are left with the following two cases:

- the inverse images of the critical points of  $V$ , studied above in Proposition 10,
- the inverse images of the intersections of critical circles with the Voronoi diagram.

and of course combinations of them.

Assume a point  $X \in \mathcal{V}$ , but is not critical for  $V$ .

**Proposition 12.** *Assume a point  $X \in \mathcal{V}$  lies on a single critical circle  $C(A_k, r)$  and on the Voronoi edge of  $A_i A_j$ , but is not a critical point of  $V$ . Let  $\langle A_i X A_j \rangle$  be the affine cone around the angle  $\angle A_i X A_j$ , then*

- (1) *Assume  $i \neq k \neq j$  then*
  - *If  $X A_k$  lies in the interior of  $\langle A_i X A_j \rangle$  then  $\pi^{-1}(X)$  is a critical component of  $\mathbb{V}$ , with transversal Morse index, equal to the Morse index of  $V$  at  $X$ ,*
  - *If  $X A_k$  lies in the exterior of  $\langle A_i X A_j \rangle$  then  $\pi^{-1}(X)$  does not contain critical points, in fact  $\mathbb{V}$  is topologically regular near  $\pi^{-1}(X)$ ,*
- (2) *Assume  $i \neq k = j$  then  $\mathbb{V}$  is singular at  $\pi^{-1}(X)$  and a limit of the case (3) in Proposition 11.*

*Proof.* We follow the proof of Theorem 1; by the formula (\*) in the Appendix, any  $\ell \in \partial\mathbb{V}(\xi)$  is a composition of

$$d_\xi\pi : T_\xi\mathcal{S} \rightarrow T_{\pi(\xi)}\mathbb{R}^2 \text{ and } L \in \partial V(\pi(\xi)).$$

Here  $L$  is a submersion (see Remark 2 in the Appendix). Let  $i \neq k \neq j$ . If  $\xi$  does not belong to the critical locus of  $\pi$  then we get the composition of two submersions, so  $\mathbb{V}$  is regular at  $\xi$  (see also Lemma 4 in the Appendix). As soon as  $\xi$  has exactly one aligned leg then the image of  $d_\xi\pi$  is one-dimensional. The normal to the tangent space at  $X$  must be contained in the cone  $\langle A_i X A_j \rangle$  (related to the Clark differential) to get a critical point of  $\mathbb{V}$ .

The case  $i \neq k = j$  is left to the reader.  $\square$

Combinations of these cases can be avoided by genericity. We consider a very strong type of genericity, adapted to the Voronoi diagram: *strongly Voronoi genericity*, which means:

- we have a *strongly generic* spider,
- the critical points of  $V$  do not lie on critical circles,
- there are no 4-vc critical points,
- there are no 2-aligned legs on the Voronoi set.

Taking into account all the previous discussion, we may summarize the results of Propositions 10, 11 and 12 in the following form:

**Theorem 5.** *For a strongly Voronoi generic smooth spider and the Voronoi distance  $\mathbb{V}$ , a configuration  $\xi \in \mathcal{S}$  is critical if and only if one of the following conditions hold for  $X = \pi(\xi)$ :*

- (1) *In the case when there are no aligned legs:  $X$  is a critical point of the Voronoi distance  $V$ ,*
- (2) *In the case of one aligned leg:*
  - (a) *in an open power cell*
    - *$X$  is the intersection point  $C(A_j, r_j^{\epsilon_j}) \cap A_i A_j \cap \mathcal{W}$  if  $i \neq j$ , and*
    - *$X$  lies on the critical circle if  $i = j$  (generically arcs or the full circle),*
  - (b) *on the Voronoi diagram – if  $X$  satisfies the condition:  $X A_k$  lies in the interior of  $\langle A_i X A_j \rangle$ ,*
- (3) *in case of two aligned legs:  $X$  is the intersection point of critical circles.*

Figure 8 illustrates the general situation.

**6.2. Morse-Bott type polynomials for the Voronoi distance.** Generating polynomials for singularities of functions have been studied by describing the changes in topology when passing a critical level. This is well understood in the case of Morse functions in the isolated case and Morse-Bott functions in the non-isolated case. For other types of (transversal) singularities and other critical sets there should exist anyhow generalizations. In the preceding section we saw that for the Voronoi distance complicated situations can occur, e.g non-isolated singularities in work space (even not necessarily smooth) will get us outside the theory of Morse-Bott functions.






FIGURE 8. The Voronoi diagram is an inverse  $Y$ -fork. Its trace  $\mathcal{V} \cap \mathcal{W}$  on the work space gives three cells. We can see inside  $\mathcal{W}$  three saddle points  $s_i$ , one maximum  $M$ , non-isolated minima on the critical circles  $m_2, m_3$  while the first critical circle lying outside the work space contributes only by one point. Moreover, there are additional local maxima  $Q_1, Q_2, Q_3$  and  $R$ , while  $P_1, P_2$  are additional saddle points. Inside each cell the theory is covered by Theorem 1.

We propose to compute a Morse-Bott polynomial by using a small perturbation so that we avoid non-isolated singularities in work space. This perturbation will act as a ‘Morsification’.

Take points  $\bar{A}_i$  near to  $A_i$  and consider for the same spider the function

$$\bar{V}(x) = \min_{1 \leq i \leq n} \|x - \bar{a}_i\|^2 = \text{dist}(x, \{\bar{A}_1, \dots, \bar{A}_n\})^2,$$

and its lifting to the spider space. The new Voronoi diagram will be denoted by  $\bar{\mathcal{V}}$ .

For generic positions of the points  $\bar{A}_i$  the statements of Theorem 5 with only isolated intersections are applicable. More precisely: in the work space the non-isolated contributions are replaced by the intersections of  $A_i \bar{A}_i$  with the critical circles. As a result  $\bar{V}$  becomes a Morse-Bott function, where the index can be computed via Theorem 2 for  $\mathbb{D} = \|x - z\|^2$ . We assume in the next theorem, that the reader is familiar with the standard approach of Morse theory.

**Theorem 6.** *Consider the filtrations by lower levels of  $\mathcal{W}$  and  $\mathcal{S}$ , such that each subsequent image interval contains only one critical value in its interior. For any generic small perturbation of the  $A_i$  there is a topological equivalence of these filtrations with the induced filtration of  $\bar{V}$ , which in the work space*

- *maps critical points of isolated type to critical points of the same index.*
- *changes critical points of non-isolated type into critical points of isolated type.*

*and respects level sets outside a neighborhood of the critical sets.*



In the general  $d$ -dimensional case the Morse theoretic approach via the squared distance function could be an interesting project.

**7.2. Graph-Linkages.** A next extension could be other types of graph-linkages. A first example is a tripod spider platform, consisting of a triangular platform, where each of the vertices is connected by a robot arm with a fixed feet (and similar for a massive  $n$ -gon platform). The paper [16] contains results about the smoothness of the configuration space and Morse results for tripod spider platforms in the plane and legs with only two edges.

**7.3. Regions with stratified differential boundary.** A simpler problem, which is directly related to our spider study is to consider a region in the plane with stratified differentiable boundary together with the quadratic distance to a point in the plane. In the case of planar spiders the (stratified) critical points of quadratic distance coincide with the images of the singular sets (via work map) of quadratic distance on spider space. In [9] this was used to treat Hooke energy for tripods in the plane.

## 8. APPENDIX

In this appendix we treat the definition and certain properties of the Clark differential on a smooth manifold. They look intuitively clear, but need some extra explanation.

Note that  $\mathbb{V}(\xi)$  is also locally Lipschitz and thus we can carry over the notion of the Clarke subdifferential to this situation. Clearly,  $\mathbb{V}$  is differentiable outside  $\pi^{-1}(\mathcal{V} \cap \mathcal{W})$ .

From the general point of view, given a  $\mathcal{C}^1$ -smooth submanifold  $M \subset \mathbb{R}^N$  of dimension  $m$  and a locally Lipschitz function  $h: M \rightarrow \mathbb{R}$ , we obviously have that  $h$  is almost everywhere differentiable along  $M$  (w.r.t. the usual  $m$ -dimensional Hausdorff measure; it is enough to compose  $h$  with local parametrizations and invoke the Rademacher Theorem). Therefore, for a point  $a \in M$ , we may repeat the classic Clarke approach by taking the convex hull, denoted again by  $\partial h(a)$ , of all the possible limits of gradients  $\nabla h(x_\nu)$  when  $x_\nu$  are points at which  $h$  is differentiable and that tend to  $a$ . Here,  $\nabla h(x) \in T_x M$  is the tangential component of the gradient  $\nabla H(x)$  of any differentiable extension of  $h$  onto a neighborhood of  $a$  (it does not depend on the choice of  $H$ ). Note that the tangent spaces vary continuously, so that  $\partial h(a) \subset T_a M$ . This tangential gradient coincides with the unique  $v \in T_x M$  such that  $h(x') = h(x) + \langle v, \rho(x - x') \rangle + o(\|x - x'\|)$  where  $\rho: \mathbb{R}^N \rightarrow T_x M$  is the orthogonal projection, while the inner product is the usual one in  $\mathbb{R}^N$ .

The point  $a \in M$  is called critical, if  $0 \in \partial h(a)$ . The notion of subdifferential can be also applied to vector-valued functions once we interpret their differentials as matrices. It also can be proved that given any local parametrization  $\gamma$  of  $M$  in a neighborhood of  $a$ ,  $\partial h(a) = d_{(\gamma^{-1}(a))} \gamma(\partial(h \circ \gamma)(\gamma^{-1}(a)))$

Consider now a  $\mathcal{C}^1$  map  $g: M \rightarrow \mathbb{R}^n$  and a locally Lipschitz function  $f: U \rightarrow \mathbb{R}$  where  $U$  is a neighborhood of  $g(M)$ .

**Lemma 2.** *If  $g$  is a submersion at  $p \in M$ , then  $0 \in \partial(f \circ g)(p)$  implies  $0 \in \partial f(g(p))$ .*

*Proof.* Assume for simplicity  $g$  is everywhere a submersion. By the Rademacher Theorem,  $f$  is differentiable outside a zero measure set  $Z \subset U$ . The co-area formula (generalized Fubini Theorem) implies  $S := g^{-1}(Z)$  has measure zero, too, since  $g$  is a submersion.

Now,  $0 = \sum \lambda_i \ell_i$  is a convex combination of some  $\ell_i \in \partial(f \circ g)(p)$  and each  $\ell_i$  is the limit of some sequence  $\ell_i^\nu := d_{x_\nu^i}(f \circ g)$  where the points  $x_\nu^i \in M \setminus S$  converge to  $p$ . Putting  $y_\nu^i := g(x_\nu^i)$ , we get  $d_{x_\nu^i}(f \circ g) = d_{y_\nu^i} f \circ d_{x_\nu^i} g$  where  $d_{x_\nu^i} g: T_{x_\nu^i} M \rightarrow \mathbb{R}^n$  is surjective. The tangent planes converge to  $T_p M$  while the corresponding differentials to  $d_p g$ , and  $y_\nu^i$  to  $g(p)$ , for each  $i$ . Passing to a subsequence, if necessary, we may assume that  $d_{y_\nu^i} f$  tend to some  $\delta_i \in \partial f(g(p))$  (these sequences can be considered as elements of  $\mathbb{R}^N$ ; the sequences are bounded since  $f$  is Lipschitz). Then  $0 = \sum (\lambda_i \delta_i) \cdot d_p g$  after identification with matrices. In particular,  $d_p g$  is an  $n \times m$  matrix containing an  $m \times m$  invertible submatrix  $A$ , whereas  $\sum (\lambda_i \delta_i)$  is in fact a  $1 \times n$  matrix of ‘unknowns’. Then  $0 = \sum (\lambda_i \delta_i) \cdot A$  has zero as the unique solution.  $\square$

**Lemma 3.** *If  $M$  is compact,  $g$  is open onto its image  $W = g(M)$ , and  $g(p)$  lies in the closure of the interior of  $W$ , then  $0 \in \partial f(g(p))$  implies  $0 \in \partial(f \circ g)(p)$ .*

*Proof.* Similarly to the previous proof, we have  $0 = \sum \lambda_i \ell_i$  as a convex combination of matrices  $\ell_i = \lim \nabla f(y_\nu^i)$  for some  $y_\nu^i \in U \setminus Z$  converging to  $g(p)$ . Choosing arbitrarily  $x_\nu^i \in g^{-1}(y_\nu^i)$  and passing to convergent subsequences, we may assume that  $x_\nu^i$  converge to some  $x^i$  and thus  $g(x^i) = g(p)$ . This allows us to write  $\nabla(f \circ g)(x_\nu^i) = \nabla f(y_\nu^i) \cdot d_{x_\nu^i} g$  and these sequences converge to  $\ell_i \cdot d_{x^i} g$ . In order to avoid the problem of obtaining different points  $x^i \in g^{-1}(g(p))$  we observe that  $g$  being open onto its image, it has continuously varying fibers (see e.g. [2] Lemma 3.8):  $g^{-1}(y_\nu^i)$  converges to  $g^{-1}(p)$  in the Hausdorff metric. This implies in particular that the point  $p \in g^{-1}(g(p))$  can be obtained as the limit of some points  $z_\nu^i \in g^{-1}(y_\nu^i)$ . Then  $\nabla(f \circ g)(z_\nu^i)$  tends to  $\ell_i \cdot d_p g \in \partial(f \circ g)(p)$  and  $0 = \sum \lambda_i (\ell_i \cdot d_p g)$ .  $\square$

We observe that the work map, the spider space, and the Voronoi distance satisfy the assumptions of the Lemma above. Moreover, in our situation the spider space is an algebraic manifold and the work map is analytic. Hence, both the set of critical points of  $\pi$  (outside of which  $\pi$  is a submersion) as well as  $\pi^{-1}(\mathcal{V})$  have measure zero and so – just as in the classical case of functions defined in the Euclidean space – in order to determine  $\partial \mathbb{V}(\xi)$  it is enough to approximate  $\xi$  by sequences  $\xi_\nu$  at which  $d_{\xi_\nu} \pi = \pi|_{T_{\xi_\nu} \mathcal{S}}: T_{\xi_\nu} \mathcal{S} \rightarrow T_{\pi(\xi_\nu)} \mathbb{R}^2 = \mathbb{R}^2$  has rank 2 and  $X_\nu := \pi(\xi_\nu) \notin \mathcal{V}$ . In particular,  $d_{\xi_\nu} \mathbb{V} = d_{x_\nu} V \circ \pi|_{T_{\xi_\nu} \mathcal{S}}$  and  $d_{x_\nu} V(x) = 2\langle x_\nu - a_{i_\nu}, x \rangle$  where  $A_{i_\nu}$  is the unique point realizing the Voronoi distance  $V(\pi(\xi_\nu))$ . Passing to a subsequence, we get the same  $i_\nu = i$  for all  $\nu$ .

Let us stress once again that any non-differentiability point  $\xi_0$  of  $\mathbb{V}$  yields  $X_0 := \pi(\xi_0) \in \mathcal{V}$ . Eventually, any  $\ell \in \partial \mathbb{V}(\xi_0)$ , is a linear map  $\ell: T_{\xi_0} \mathcal{S} \rightarrow \mathbb{R}$  of the form

$$(*) \quad \ell(s) = 2\langle x_0 - a, \pi(s) \rangle, \quad s \in T_{\xi_0} \mathcal{S},$$

where  $A$  is taken from the convex hull of those feet  $A_1, \dots, A_n$  that realize the distance  $V(x_0)$ . On the other hand, any map  $\ell$  of the previous form belongs to  $\partial \mathbb{V}(\xi_0)$ , provided  $X_0$  lies in the interior of  $\mathcal{W}$ . Indeed, we may define  $L(x) = 2\langle x_0 - a, x \rangle$ ,  $x \in \mathbb{R}^2$  and we know that  $L \in \partial V(x_0)$ . Thus, it is the convex combination of the limits  $L_i$  of some  $\nabla V(x_\nu^i)$  with  $x_\nu^i \rightarrow x_0$  and all  $X_\nu^i \in \mathcal{W} \setminus \mathcal{V}$ , because  $X_0$  is an interior point of the work space. The spider space being compact and the work map open, we easily produce the corresponding required sequences  $\xi_\nu^i \rightarrow \xi_0$  to finally get the assertion.

All this implies in particular the following two remarks gathered in one obvious Lemma:

**Lemma 4.** *In the situation considered above and with its notations, if  $\xi_0$  is a critical point of  $\mathbb{V}$ , then  $x_0 - a$  is orthogonal to  $\pi(T_{\xi_0}\mathcal{S})$ , for some  $A$  in the convex hull of the feet realising the distance  $V(x_0)$ . On the other hand, any critical point  $x_0 \in \mathcal{V}$  of  $V$  and lying in the interior of the work space, is the image of a critical point of  $\mathbb{V}$  by the work map.*

**Remark 2.** According to the discussion just made, if  $0 \notin \partial V(x_0)$ , then any  $L \in \partial V(x_0)$  is a submersion.

We end with the two remaining statements in section 6:

**8.1. Proposition 9.** [Composition of regular maps] If  $\xi \in \mathcal{S}$  is a regular point of the work map  $\pi$  and  $V$  is regular at  $X = \pi(\xi)$ , then  $\mathbb{V}$  is regular at  $\xi$ .

This is a straightforward consequence of Lemma 2.

**8.2. Case 1.** It is easy to see that  $0 \in \partial V(\pi(\xi))$  implies  $0 \in \partial \mathbb{V}(\xi)$ .

Apply Lemma 3.

## REFERENCES

- [1] R.Bot, Nondegenerate critical manifolds, Ann. of Math. (2) 60 (1954), 248-261.
- [2] M. Denkowski, J.-J. Loeb, On open analytic and subanalytic mappings, Complex Var. Elliptic Eq. (2017), 62 (1), 27-46.
- [3] M. Farber, Invitation to Topological Robotics (Zurich Lectures in Advanced Mathematics), 2008.
- [4] M. Farber, D. Schütz, Homology of planar polygon spaces, Geom Dedicata (2007) 125, 75-92, DOI 10.1007/s10711-007-9139-7.
- [5] M. Goresky, R. MacPherson, Stratified Morse Theory, Springer Verlag 1988.
- [6] J.-C. Hausmann, A. Knutson, The cohomology ring of polygon spaces, Ann. Inst. Fourier (Grenoble), 48 (1998), 281–321.
- [7] J.-C. Hausmann, Sur la Topologie des Bras Articulés, In: Algebraic Topology Poznań 1989. Lecture Notes in Mathematics, vol 1474. Springer, 146-159.
- [8] M. Kapovich, J. Millson, On the moduli spaces of polygons in the Euclidean plane, Journal of Diff. Geometry, 42 (1995), 133–164.
- [9] G. Khimshiashvili, D. Siersma, Hooke and Coulomb energy of tripod spiders, Proc. I. Vekua Inst. Appl. Math. 72 (2022), 39-46.
- [10] Y. Kamiyama, S. Tsukada, The configuration space of the n-arms machine in the Euclidean space, Topology and its Applications 154 (2007) 1447-1464.
- [11] M. van Manen, D. Siersma, Power diagrams and Morse Theory, arXiv:math/0508037 2005.
- [12] P. Mounoud, Sur l'espace des configurations d'une araignée, Osaka J. Math. 48 (2011), 149-178.
- [13] L. Nicolaescu : Invitation to Morse Theory Springer Verlag 2011.
- [14] J.-I. O'Hara, The configuration space of planar spider linkages, Top. Appl. 154 (2007), 502-526.
- [15] N. Shvalb, M. Soham, D. Blanc, The configuration space of anarchnoid mechanisms, Forum Math. 17 (2005), 1033-1042.
- [16] N. Shvalb, M. Soham, D. Blanc, The configuration space of a parallel polygonal mechanism, J. Geom Topol 9 (2009) 137-167.
- [17] D. Siersma, Voronoi diagrams and Morse theory of the distance function, in: Geometry in Present Day Science, Proceedings 1999, 187-208.

M. DENKOWSKI: JAGIELLONIAN UNIVERSITY, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, INSTITUTE OF MATHEMATICS, KRAKOW, POLAND, MACIEJ.DENKOWSKI@UJ.EDU.PL; G. PANINA: ST. PETERSBURG DEPARTMENT OF STEKLOV INSTITUTE OF MATHEMATICS, GAIANE-PANINA@RAMBLER.RU; D. SIERSMA: MATHEMATISCH INSTITUUT, UNIVERSITEIT UTRECHT, D.SIERSMA@UU.NL