

NOTE ON C^* -ALGEBRAS ASSOCIATED TO BOUNDARY ACTIONS OF HYPERBOLIC 3-MANIFOLD GROUPS

SHIRLY GEFFEN AND JULIAN KRANZ

ABSTRACT. Using Kirchberg–Phillips’ classification of purely infinite C^* -algebras by K -theory, we prove that the isomorphism types of crossed product C^* -algebras associated to certain hyperbolic 3-manifold groups acting on their Gromov boundary only depend on the manifold’s homology. As a result, we obtain infinitely many pairwise non-isomorphic hyperbolic groups all of whose associated crossed products are isomorphic. These isomorphisms are not of dynamical nature in the sense that they are not induced by isomorphisms of the underlying groupoids.

1. INTRODUCTION

The Kirchberg–Phillips classification theorem [Kir94, Phi00] classifies so-called *UCT Kirchberg C^* -algebras* up to isomorphism by topological K -theory. As a consequence, this theorem produces many ‘surprising’ isomorphisms of C^* -algebras that cannot be constructed by hand, such as Kirchberg’s isomorphism $\mathcal{O}_2 \otimes \mathcal{O}_2 \cong \mathcal{O}_2$. The class of UCT Kirchberg C^* -algebras contains many examples such as the (reduced) crossed product C^* -algebra $C(\partial G) \rtimes_r G$ associated to the action of a non-elementary torsion-free hyperbolic group G on its Gromov-boundary [AD97, Tu99]. In fact, crossed products of arbitrary minimal amenable topologically free actions $G \curvearrowright X$ of acylindrically hyperbolic groups belong to this class [GGKN23]. Although this large class of C^* -algebras is *in principle* classifiable by K -theory, explicit K -theory computations remain difficult in general. For the specific example of the action of a torsion-free hyperbolic group G on its boundary, Emerson and Meyer [EM06] relate the K -theory of $C(\partial G) \rtimes_r G$ to the K -theory and K -homology of the classifying space BG . In this short note, we use their result to obtain explicit computations for fundamental groups of hyperbolic 3-manifolds. Our techniques are identical to an analogous result for non-compact manifolds [Mc20, Proposition 1.6]

Theorem A. *Let $d \geq 0$ be a fixed integer and let G be the fundamental group of a closed connected orientable hyperbolic 3-manifold M with $H_1(M) \cong \mathbb{Z}^d$. Then there are isomorphisms*

$$K_0(C(\partial G) \rtimes_r G) \cong K_1(C(\partial G) \rtimes_r G) \cong \mathbb{Z}^{2d+2}$$

that identify the class of the unit $[1] \in K_0(C(\partial G) \rtimes_r G)$ with $(1, 0) \in \mathbb{Z} \oplus \mathbb{Z}^{2d+1}$.

2020 *Mathematics Subject Classification.* Primary 46L80, 57K32, Secondary 46L35.

Key words and phrases. hyperbolic 3-manifolds, Crossed products, K -theory for C^* -algebras.

J. Kranz was supported by the Engineering and Physical Sciences Research Council [Grant Ref: EP/X026647/1]. S. Geffen was partially supported by the ERC Advanced Grant 834267 - AMAREC, by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID 427320536 - SFB 1442, as well as by Germany’s Excellence Strategy EXC 2044-390685587, Mathematics Münster: Dynamics–Geometry–Structure.

Brock–Dunfield’s work [BD15] shows that there exists a rich supply of groups that satisfy the assumptions in Theorem A. Thus, we obtain examples of isomorphisms of crossed product C^* -algebras:

Corollary B. *There are infinitely many pairwise non-isomorphic hyperbolic groups G_i , $i \in \mathbb{N}$ with pairwise isomorphic crossed products $C(\partial G_i) \rtimes_r G_i$, $i \in \mathbb{N}$.*

These isomorphisms are ‘surprising’ in the sense that they cannot be obtained from an isomorphism of the underlying groupoids.

Proposition C. *Let G_i and G_j be two non-isomorphic groups as in Theorem A. Then there is no isomorphism $C(\partial G_i) \rtimes_r G_i \cong C(\partial G_j) \rtimes_r G_j$ that restricts to an isomorphism $C(\partial G_i) \cong C(\partial G_j)$. In particular, $C(\partial G_i) \rtimes_r G_i$ contains infinitely many pairwise non-conjugate Cartan subalgebras with spectrum S^2 .*

Acknowledgments. This note arose from discussions with Johannes Ebert and is mostly based on his ideas. We are indebted to him for sharing his strategy to combine the Baum–Connes conjecture with the Atiyah–Hirzebruch spectral sequence, for educating us about hyperbolic 3-manifolds, for guiding us through the computations, and for encouraging us to use his ideas and publish this note. After this note was circulated as a preprint, Bram Mesland and Haluk Şengün brought to our attention that Theorem A can be deduced from their joint work [Mc20]. We are moreover grateful to Jakub Curda for pointing out the reference [Mat02] which drastically simplified the proof.

2. ISOMORPHISM TYPES OF CROSSED PRODUCTS

Let M be a closed connected orientable hyperbolic 3-manifold. Then the universal cover \widetilde{M} of M is isometric to the 3-dimensional hyperbolic space \mathbb{H}^3 by the Killing–Hopf theorem [Lee18, Corollary 12.5]. Let $G = \pi_1(M)$ be the fundamental group of M . Considering the natural (free) action $G \curvearrowright \widetilde{M}$ by Deck transformations, and the fact that \widetilde{M} is contractible, the orbit space $\widetilde{M}/G \cong M$ is a finite model for the classifying space BG . It follows that G is finitely generated and torsion-free [Bro82, §VIII, Exercise 4.1 and Corollary 2.5]. Moreover, since the action $G \curvearrowright \widetilde{M}$ is proper and co-compact, G is quasi-isometric to \mathbb{H}^3 by the Milnor–Švarc lemma [BH99, §I.8, Proposition 8.9]. In particular, G is a Gromov-hyperbolic group with Gromov boundary $\partial G \cong \partial \mathbb{H}^3 \cong S^2$ [BH99, §III H, Theorem 1.9 and Theorem 3.9].

We refer the reader to [CELY17] for the definition of crossed product C^* -algebras, topological K -theory for C^* -algebras, and the Baum–Connes conjecture. For a unital C^* -algebra A , we denote by $[1_A] \in K_0(A)$ the class of its unit (or equivalently the class of the rank-one A -module $[A]$).

For a connected space X and a choice of base point $x_0 \in X$, we denote by $[x_0] \in H_0(X)$ the image of the canonical generator $1 \in \mathbb{Z} = H_0(\{x_0\})$ under the natural map $H_0(\{x_0\}) \rightarrow H_0(X)$ and by $[1_X] \in K_0(X)$ the image of the canonical generator $1 \in \mathbb{Z} = K_0(\{x_0\})$ under the natural map $K_0(\{x_0\}) \rightarrow K_0(X)$.

The following computations of K -theory and K -homology of 3-manifolds using their cohomology and homology groups is an application of the Atiyah–Hirzebruch spectral sequence. The proof is given in [Mat02, Proposition 2.1], but we include a

brief introduction to spectral sequences and more detailed computations for non-experts in Appendix A. Note that a similar result was obtained in [Mc20, Section 1].

Lemma 2.1. *Let M be a closed connected orientable 3-manifold. Then there are isomorphisms*

$$\begin{aligned} (1) \quad & K_0(M) \cong H_0(M) \oplus H_2(M), \\ (2) \quad & K_1(M) \cong H_1(M) \oplus H_3(M), \\ (3) \quad & K^0(M) \cong H^0(M) \oplus H^2(M), \\ (4) \quad & K^1(M) \cong H^1(M) \oplus H^3(M), \end{aligned}$$

that identify $[1_M] \in K_0(M)$ with $[x_0] \in H_0(M)$.

Proof. The isomorphisms (1) and (2) identifying $[1_M]$ with $[x_0]$ are obtained in [Mat02, Proposition 2.1]. By Poincaré duality, we have $H^*(M) \cong H_{3-*}(M)$. Moreover, by Kasparov’s K -theoretic Poincaré duality [Kas88, Theorem 4.10]¹, we have an isomorphism $K^*(M) \cong K_{3-*}(M)$. Together with Bott periodicity, this proves (3) and (4). \square

Proof of Theorem A (cf. [Mc20, Proposition 1.6]). Using Poincaré duality together with $H^1(M) = \text{Hom}_{\mathbb{Z}}(H_1(M), \mathbb{Z})$, the assumptions in Theorem A imply the following isomorphisms:

$$\begin{aligned} H_0(M) &\cong H^3(M) \cong H^0(M) \cong H_3(M) \cong \mathbb{Z}, \\ H_1(M) &\cong H^2(M) \cong H^1(M) \cong H_2(M) \cong \mathbb{Z}^d. \end{aligned}$$

Combining with Lemma 2.1, these allow to compute

$$(5) \quad K_0(M) \cong K_1(M) \cong K^0(M) \cong K^1(M) \cong \mathbb{Z}^{d+1}$$

and identify

$$(6) \quad K_0(M) \ni [1_M] \mapsto (1, 0) \in \mathbb{Z} \oplus \mathbb{Z}^d.$$

Note that M has zero Euler characteristic by Poincaré duality. Moreover, G satisfies the Baum–Connes conjecture with coefficients by [Laf12]. Thus, [EM06, Theorem 1] applied to the action $G \curvearrowright \widetilde{M}$ produces short exact sequences

$$(7) \quad \begin{aligned} 0 \rightarrow K_0(C_r^*(G)) &\xrightarrow{u_*} K_0(C(\partial G) \rtimes_r G) \rightarrow K^1(M) \rightarrow 0, \\ 0 \rightarrow K_1(C_r^*(G)) &\xrightarrow{u_*} K_1(C(\partial G) \rtimes_r G) \rightarrow K^0(M) \rightarrow 0, \end{aligned}$$

where the map u_* is induced by the canonical inclusion $u: C_r^*(G) \rightarrow C(\partial G) \rtimes_r G$. Recall that M is a model for BG . Thus, the Baum–Connes assembly map

$$(8) \quad \mu_*^G: K_*(M) \xrightarrow{\cong} K_*(C_r^*(G))$$

is an isomorphism. By applying functoriality of the assembly map to the group homomorphism $\{e\} \rightarrow G$, we moreover conclude that

$$(9) \quad \mu_0^G([1_M]) = [1_{C_r^*(G)}] \in K_0(C_r^*(G)).$$

¹In order to apply [Kas88, Theorem 4.10], recall that as a closed oriented 3-manifold, M admits a spin^c -structure and that any spin^c -structure induces a $\mathbb{Z}/2$ -graded Morita-equivalence $C_r(M) \sim C(M) \hat{\otimes} \text{Cl}_3$, where $C_r(M)$ denotes the section algebra of the Clifford bundle of T^*M and Cl_3 the third complex Clifford algebra.

Indeed, this follows by applying commutativity of the diagram

$$\begin{array}{ccc} K_0(\text{pt}) & \xrightarrow{\mu_0^{\{e\}}} & K_0(\mathbb{C}) \\ \downarrow & & \downarrow \\ K_0(M) & \xrightarrow{\mu_0^G} & K_0(C_r^*(G)) \end{array}$$

to the element $[1_{\text{pt}}] \in K_0(\text{pt})$.

Since $K^*(M) \cong \mathbb{Z}^{d+1}$, the short exact sequences in (7) split. Using (8) at the second step and (5) at the third step, we get

$$K_0(C(\partial G) \rtimes_r G) \cong K_0(C_r^*(G)) \oplus K^1(M) \cong K_0(M) \oplus K^1(M) \cong \mathbb{Z}^{2d+2}.$$

By (6) and (9), this isomorphism identifies the class of the unit $[1_{C(\partial G) \rtimes_r G}] = u_*([1_{C_r^*(G)}]) \in K_0(C(\partial G) \rtimes_r G)$ with $(1, 0) \in \mathbb{Z} \oplus \mathbb{Z}^{2d+1}$. Analogously, we obtain an isomorphism

$$K_1(C(\partial G) \rtimes_r G) \cong K_1(C_r^*(G)) \oplus K^0(M) \cong K_1(M) \oplus K^0(M) \cong \mathbb{Z}^{2d+2}.$$

□

Proof of Corollary B. By [BD15, Theorem 2.1], there are infinitely many pairwise non-isometric closed connected orientable² hyperbolic 3-manifolds M_i , $i \in \mathbb{N}$ satisfying $H_1(M_i) \cong \mathbb{Z}^d$. In particular, the fundamental groups $G_i := \pi_1(M_i)$, $i \in \mathbb{N}$ are pairwise non-isomorphic by Mostow's rigidity theorem [Mos68].

The crossed products $A_i := C(\partial G_i) \rtimes_r G_i$ are unital simple separable nuclear purely infinite C^* -algebras by [AD97, Proposition 3.2] that satisfy the UCT by [Tu99, Proposition 10.7]. Fix $i, j \in \mathbb{N}$. Theorem A implies that there are isomorphisms $K_0(A_i) \cong K_0(A_j)$ and $K_1(A_i) \cong K_1(A_j)$ that identify $[1_{A_i}]$ with $[1_{A_j}]$. Thus, the Kirchberg–Phillips classification theorem [Kir94, Phi00] implies that A_i and A_j are isomorphic. □

Remark 2.2. We do not know, whether the reduced group C^* -algebras $C_r^*(G_i)$, $i \in \mathbb{N}$ are pairwise isomorphic, although K -theory does not distinguish them. To the best of our knowledge, it is an open problem whether the reduced group C^* -algebra of a general torsion-free discrete group G recovers the group. However, results of this type do exist for certain classes of groups (see [ER18] for instance).

As a preparation for the proof of Proposition C, recall that the *transformation groupoid* $X \rtimes G$ associated to a group action $G \curvearrowright X$ is the topological groupoid whose object space is given by $(X \rtimes G)^{(0)} := X$ and whose morphism space is given by $X \rtimes G := X \times G$, with range, source, and composition maps r, s, \cdot determined by the formulas

$$r(x, g) = x, \quad s(x, g) = g^{-1}x, \quad (x, g) \cdot (g^{-1}x, h) = (x, gh), \quad x \in X, g, h \in G.$$

The *topological full group* $[[\mathcal{G}]]$ of a topological groupoid \mathcal{G} is the group of all global bisections, that is, open subsets $U \subseteq \mathcal{G}$ of the arrow space such that the restrictions $r|_U: U \rightarrow \mathcal{G}^{(0)}$ and $s|_U: U \rightarrow \mathcal{G}^{(0)}$ are homeomorphisms³. Note that the topological full group of a topological groupoid is an isomorphism invariant.

²Note that although not stated in the theorem, the proof of [BD15, Theorem 2.1] indeed produces orientable manifolds.

³Usually, topological full groups are only defined and interesting for groupoids with totally disconnected object space, see [Mat12].

Lemma 2.3. *Let X be a connected space and $G \curvearrowright X$ an action of a discrete group. Then the natural map*

$$G \rightarrow [[X \rtimes G]], \quad g \mapsto X \times \{g\}$$

is a group isomorphism.

Proof. If $U \in [[X \rtimes G]]$ is a global bisection, then U is homeomorphic to X and therefore connected. In particular, the projection of $U \subseteq X \times G$ to G is constant. \square

Proof of Proposition C. Assume by contradiction that there is an isomorphism

$$C(\partial G_i) \rtimes_r G_i \cong C(\partial G_j) \rtimes_r G_j$$

that restricts to an isomorphism $C(\partial G_i) \cong C(\partial G_j)$. Note that the action $G_i \curvearrowright \partial G_i$ is topologically free since G_i is torsion-free. Thus, $C(\partial G_i) \subseteq C(\partial G_i) \rtimes_r G_i$ is a Cartan subalgebra (see [Ren08, §6.1]). By Renault’s reconstruction theorem [Ren08, Theorem 5.9], the isomorphism $C(\partial G_i) \rtimes_r G_i \cong C(\partial G_j) \rtimes_r G_j$ is induced by an isomorphism $\partial G_i \rtimes G_i \cong \partial G_j \rtimes G_j$ of the underlying transformation groupoids. In particular, their associated topological full groups are isomorphic. By Lemma 2.3, this implies that $G_i \cong G_j$, which is a contradiction. \square

3. K-THEORY WITH COEFFICIENTS

We end this note with a variant of Theorem A and Corollary B for 3-manifolds with torsion in their first homology. Recall that for an Abelian group A and a C^* -algebra B , the K -theory of B with coefficients in A is defined as $K_*(B; A) := K_*(B \otimes \mathcal{A})$ where \mathcal{A} is any C^* -algebra satisfying the UCT such that $K_0(\mathcal{A}) \cong A$ and $K_1(\mathcal{A}) = 0$. We refer to [RS87] for the UCT and Künneth theorem for C^* -algebras.

Theorem 3.1. *Let G be the fundamental group of a closed connected orientable hyperbolic 3-manifold. Let \mathbb{F} be a field. Then there are isomorphisms*

$$K_0(C(\partial G) \rtimes_r G; \mathbb{F}) \cong K_1(C(\partial G) \rtimes_r G; \mathbb{F}) \cong \mathbb{F}^2 \oplus H_1(M; \mathbb{F}) \oplus H^1(M; \mathbb{F})$$

which identify the class $[1] \in K_0(C(\partial G) \rtimes_r G; \mathbb{F})$ with $(1, 0) \in \mathbb{F} \oplus (\mathbb{F} \oplus H_1(M; \mathbb{F}) \oplus H^1(M; \mathbb{F}))$.

Proof. Since the proof follows the same strategy as the proof of Theorem A, we only explain the necessary modifications. Let \mathcal{A} be a C^* -algebra satisfying the UCT such that $K_0(\mathcal{A}) \cong \mathbb{F}$ and $K_1(\mathcal{A}) \cong 0$.

The K -homology of M can be computed using a version of [Mat02, Proposition 2.1] with coefficients in \mathbb{F} , which in turn can be proven in exactly the same way as the integral version (using the Atiyah–Hirzebruch spectral sequence for K -homology with coefficients in \mathbb{F}). Note that the exact sequences (7) are obtained in [EM06] as long exact sequences in K -theory associated to the short exact sequence

$$0 \rightarrow C_0(X) \rtimes_r G \rightarrow C(\overline{X}) \rtimes_r G \rightarrow C(\partial G) \rtimes_r G \rightarrow 0,$$

where $X = \widetilde{M}$ is the universal cover of M and \overline{X} is its natural compactification. Tensoring this sequence with \mathcal{A} and applying K -theory yields a new six-term exact sequence which again splits into two short exact sequences by the Künneth theorem (note that \mathcal{A} satisfies the Künneth theorem since it satisfies the UCT). In this way

we obtain the appropriate analogues of the Emerson–Meyer exact sequences with coefficients in \mathbb{F} ,

$$\begin{aligned} 0 \rightarrow K_0(C_r^*(G); \mathbb{F}) &\xrightarrow{u_*} K_0(C(\partial G) \rtimes_r G; \mathbb{F}) \rightarrow K^1(M; \mathbb{F}) \rightarrow 0, \\ 0 \rightarrow K_1(C_r^*(G); \mathbb{F}) &\xrightarrow{u_*} K_1(C(\partial G) \rtimes_r G; \mathbb{F}) \rightarrow K^0(M; \mathbb{F}) \rightarrow 0, \end{aligned}$$

which split since \mathbb{F} is a field. The left and right hand terms of these sequences are now computed in the same way as before, where Kasparov’s Poincaré duality theorem [Kas88, Theorem 4.9] and the Baum–Connes isomorphism [Laf12] have to be taken with coefficients in \mathcal{A} . \square

Corollary 3.2. *Let \mathbb{F} be a field and let \mathcal{A} be a unital simple separable nuclear C^* -algebra satisfying the UCT such that $K_0(\mathcal{A}) \cong \mathbb{F}$ and $K_1(\mathcal{A}) \cong 0$. For instance, let \mathcal{A} be the universal UHF algebra \mathcal{Q} for $\mathbb{F} = \mathbb{Q}$, or let \mathcal{A} be the Cuntz algebra \mathcal{O}_{p+1} for $\mathbb{F} = \mathbb{F}_p$. Let M be a finitely generated Abelian group. Then there are infinitely many pairwise non-isomorphic hyperbolic groups G_i , $i \in \mathbb{N}$ with $H_1(G_i) \cong M$ such that the stabilized crossed products*

$$\mathcal{A} \otimes (C(\partial G_i) \rtimes_r G_i), \quad i \in \mathbb{N}$$

are all isomorphic.

Proof. Note that the assumptions on \mathcal{A} guarantee that the stabilized crossed products in the statement are UCT Kirchberg algebras, see [KR00, Proposition 4.5] and [BO08, Proposition 10.1.7]. Thus, the corollary can be proven in the same way as Corollary B, using [BD15] and using Theorem 3.1 instead of Theorem A. \square

APPENDIX A. SPECTRAL SEQUENCES

In this appendix, we give more details on the computation K -theory and K -homology (with integer coefficients) of a closed, connected, orientable, 3-dimensional, hyperbolic manifold in terms of homology and cohomology, see [Mat02, Proposition 2.1] and Lemma 2.1. This is done using the Atiyah–Hirzebruch spectral sequence.

We recall that a *cohomological spectral sequence* (in the category of Abelian groups) is given by a collection $\{E_i, d_i\}_{i \in \mathbb{N}}$, where, for each i , the *page* E_i consists of a grid of Abelian groups, denoted $(E_i^{p,q})_{p,q \in \mathbb{Z}}$, equipped with maps $d_i^{p,q}: E_i^{p,q} \rightarrow E_i^{p+i, q-i+1}$, called *differentials*, which satisfy $d_i^{p,q} \circ d_i^{p-i, q+i-1} = 0$, or, in other words, $\text{Im}(d_i^{p-i, q+i-1}) \subseteq \text{Ker}(d_i^{p,q})$. Moreover, the “next page” E_{i+1} is given by the homology of the page E_i . That is,

$$E_{i+1}^{p,q} \cong \frac{\text{Ker}(d_i^{p,q})}{\text{Im}(d_i^{p-i, q+i-1})}.$$

The spectral sequence $\{E_i, d_i\}_{i \in \mathbb{N}}$ is called *degenerate at i_0* if the differentials $(d_i^{p,q})_{p,q \in \mathbb{Z}}$ vanish for all $i \geq i_0$. In this case, the *limiting term* can be defined as $E_\infty^{p,q} = E_{i_0}^{p,q}$, for $p, q \in \mathbb{Z}$, and some $i \geq i_0$. We will only deal with degenerate spectral sequences, although the limiting term can be defined more generally.

Assume that $(A^n)_{n \in \mathbb{Z}}$ is a collection of Abelian groups, so that each A^n admits a Hausdorff grading, which we write as

$$A^n := G^0 A^n \supseteq G^1 A^n \supseteq G^2 A^n \supseteq \dots$$

with $\bigcap_{p=0}^{\infty} G^p A^n = 0$. It is said that a (degenerate) spectral sequence $\{E_i, d_i\}_{i \in \mathbb{N}}$ converges to $(A^n)_{n \in \mathbb{Z}}$ if for every $p \geq 0$ and $q \in \mathbb{Z}$ we have $E_{\infty}^{p,q} \cong \frac{G^p A^{p+q}}{G^{p+1} A^{p+q}}$. We will use the notation $E_r^{p,q} \Rightarrow A^{p+q}$ for convergence.

Lemma. *Let M be a closed connected orientable 3-manifold. We have $K^0(M) \cong H^0(M) \oplus H^2(M)$ and $K^1(M) \cong H^1(M) \oplus H^3(M)$.*

Proof. The second page of the Atiyah-Hirzebruch spectral sequence for K -theory (which applies more generally in the setting of finite CW-complexes) is given by


$$E_2^{p,q} = H^p(M, K^q(\text{pt})).$$

We claim that the differentials $d_2^{p,q}$ vanish. Indeed, the K -theory of a point is

$$K^q(\text{pt}) = \begin{cases} \mathbb{Z}, & \text{if } q \text{ is even} \\ 0, & \text{otherwise} \end{cases}.$$

Moreover, under our assumptions on M , we have $H^n(M) = 0$ for all $n \notin \{0, 1, 2, 3\}$ (this follows, for example, from Poincaré duality and the fact that cohomology groups are defined to be zero for negative degrees).

Putting these facts together, we can visualize the second page of the Atiyah-Hirzebruch spectral sequence as follows:



It is immediately visible that the differentials d_2 either have vanishing range or vanishing domain, thus must all be the zero map. The third page in the Atiyah-Hirzebruch spectral sequence is now concluded to be identical (as a grid) to the second page (since E_3 is defined to be the homology of E_2). We will check that the differentials d_3 vanish as well. In the following figure we draw the only (recurring) differentials which could potentially be non-zero:

$$\begin{array}{cccccccc}
& \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \\
\cdots & 0 & 0 & 0 & 0 & 0 & 0 & \cdots \\
\cdots & 0 & H^0(M) & H^1(M) & H^2(M) & H^3(M) & 0 & \cdots \\
\cdots & 0 & 0 & 0 & \searrow^{d_3} & 0 & 0 & \cdots \\
\cdots & 0 & H^0(M) & H^1(M) & H^2(M) & H^3(M) & 0 & \cdots \\
& \vdots & \vdots & \vdots & \vdots & \vdots & \vdots &
\end{array}$$

Applying the Atiyah-Hirzebruch spectral sequence to a one-point space, we obtain exactly the same second and third page as obtained above for the manifold M , except $H^n(\text{pt}) = 0$ for all $n \neq 0$.

Consider the natural maps $\text{pt} \rightarrow M \rightarrow \text{pt}$. Since the Atiyah-Hirzebruch spectral sequence is functorial, we have a retract $E_3(\text{pt}) \rightarrow E_3(M) \rightarrow E_3(\text{pt})$, which is interpreted as the following commutative diagram.

$$\begin{array}{ccccc}
H^0(\text{pt}) & \longrightarrow & H^0(M) & \longrightarrow & H^0(\text{pt}) \\
\downarrow d_3 & & \downarrow d_3 & & \downarrow d_3 \\
H^3(\text{pt}) & \longrightarrow & H^3(M) & \longrightarrow & H^3(\text{pt})
\end{array}$$

As $H^3(\text{pt}) = 0$, we have that the differential maps on the left-hand side and right-hand side must vanish. The composition $H^0(\text{pt}) \longrightarrow H^0(M) \longrightarrow H^0(\text{pt})$ is the identity map. By our assumptions on M (and Poincaré duality), we have $H^0(M) \cong H^3(M) \cong H^0(\text{pt}) \cong \mathbb{Z}$, which forces the maps $H^0(\text{pt}) \longrightarrow H^0(M)$ and $H^0(M) \longrightarrow H^0(\text{pt})$ to be isomorphisms. Rewriting the diagram above, we have

$$\begin{array}{ccccc}
\mathbb{Z} & \xrightarrow{\cong} & \mathbb{Z} & \xrightarrow{\cong} & \mathbb{Z} \\
\downarrow d_3 & & \downarrow d_3 & & \downarrow d_3 \\
0 & \longrightarrow & \mathbb{Z} & \longrightarrow & 0
\end{array}$$

which now clearly forces the map $d_3: H^0(M) \rightarrow H^3(M)$ to be the zero map, as we wanted to show.

Observe that for $n \geq 3$, E_n is identical to E_2 as a grid, and it is easy to see that the differentials d_n vanish as well for all $n \geq 4$. We conclude that the Atiyah–Hirzebruch spectral sequence is degenerate at $i_0 = 2$, and so $E_\infty^{p,q} = E_2^{p,q}$, for all $p, q \in \mathbb{Z}$.

Set $A^n := K^n(M)$, for $n \in \mathbb{Z}$. By Atiyah–Hirzebruch, there exists a Hausdorff grading $A^n = G^0 A^n \supseteq G^1 A^n \supseteq G^3 A^n \supseteq \dots$, for every $n \in \mathbb{Z}$ (the concrete description of the grading will follow from the computations below), such that $E_2^{p,q} \Rightarrow K^{p+q}(M)$.

For $p, q \in \mathbb{Z}$ with $p + q = 0$ we get the following short exact sequences:

$$0 \longrightarrow G^1 A^0 \longrightarrow G^0 A^0 \longrightarrow E_\infty^{0,0} \longrightarrow 0$$

$$0 \longrightarrow G^2 A^0 \longrightarrow G^1 A^0 \longrightarrow E_\infty^{1,-1} \longrightarrow 0$$

$$0 \longrightarrow G^3 A^0 \longrightarrow G^2 A^0 \longrightarrow E_\infty^{2,-2} \longrightarrow 0$$

$$0 \longrightarrow G^4 A^0 \longrightarrow G^3 A^0 \longrightarrow E_\infty^{3,-3} \longrightarrow 0$$

⋮

The first short exact sequence reduces to

$$0 \longrightarrow G^1 A^0 \longrightarrow K^0(M) \longrightarrow H^0(M) \longrightarrow 0.$$

Since $H^0(M) \cong \mathbb{Z}$ is torsion free, we conclude that $K^0(M) \cong H^0(M) \oplus G^1 A^0$.

If n is odd, we have $E_\infty^{n,-n} = 0$, so $G^1 A^0 \cong G^2 A^0$ and $G^3 A^0 \cong G^4 A^0$. Note that $E_\infty^{n,-n} = 0$ whenever $n \geq 4$. Thus, $G^n A^0 \cong G^3 A^0$ for all $n \geq 4$. Since the grading is Hausdorff, it follows that $G^n A^0 = 0$ for all $n \geq 3$. Finally, $E_\infty^{2,-2} \cong H^2(M)$. Working inductively with the short exact sequences above, we conclude that

$$K^0(M) \cong H^0(M) \oplus G^1 A^0 \cong H^0(M) \oplus G^2 A^0 \cong H^0(M) \oplus H^2(M).$$

Next, we compute $K^1(M)$ using $E_2^{p,q} \Rightarrow K^{p+q}(M)$. For $p, q \in \mathbb{Z}$ with $p + q = 1$ we get the following short exact sequences

$$0 \longrightarrow G^1 A^1 \longrightarrow G^0 A^1 \longrightarrow E_\infty^{0,1} \longrightarrow 0$$

$$0 \longrightarrow G^2 A^1 \longrightarrow G^1 A^1 \longrightarrow E_\infty^{1,0} \longrightarrow 0$$

$$0 \longrightarrow G^3 A^1 \longrightarrow G^2 A^1 \longrightarrow E_\infty^{2,-1} \longrightarrow 0$$

$$0 \longrightarrow G^4 A^1 \longrightarrow G^3 A^1 \longrightarrow E_\infty^{3,-2} \longrightarrow 0$$

⋮

Since $E_\infty^{0,1} \cong E_\infty^{2,-1} \cong 0$, the first short exact sequence implies that $K^1(M) \cong G^1 A^1$, and the third short-exact sequence implies $G^2 A^1 \cong G^3 A^1$. Since $E_\infty^{1,0} \cong H^1(M) \cong \text{Hom}(M, \mathbb{Z})$ is torsion-free and finitely generated, we have that the second short exact sequence splits, and $G^1 A^1 \cong H^1(M) \oplus G^2 A^1 \cong H^1(M) \oplus G^3 A^1$.

Finally, $E_\infty^{3,-2} \cong H^3(M) \cong H_0(M) \cong \mathbb{Z}$ by Poincaré duality. Thus, the last short exact sequence splits as well, and we have $G^3 A^1 \cong H^3(M) \oplus G^4 A^1$. However, noticing that $E_\infty^{n,m} = 0$ for all $n \geq 4$ and all $m \in \mathbb{Z}$, we see that $G^n A^1 \cong G^4 A^1$ for all $n \geq 4$. Since the grading is Hausdorff, it follows that $G^4 A^1 = 0$, and so $G^3 A^1 \cong H^3(M)$. Combining the above,

$$K^1(M) \cong G^1 A^1 \cong H^1(M) \oplus G^2 A^1 \cong H^1(M) \oplus G^3 A^1 \cong H^1(M) \oplus H^3(M).$$

□

In a similar manner, one can apply the homological version of the Atiyah-Hirzebruch spectral sequence in order to compute the K -homology of M , $K_*(M)$. Alternatively, one can use Kasparov's K -theoretic Poincaré duality and Bott periodicity (as explained in the proof of Lemma 2.1) to conclude that $K_0(M) \cong H_0(M) \oplus H_2(M)$ and $K_1(M) \cong H_1(M) \oplus H_3(M)$.

REFERENCES

- [AD97] Claire Anantharaman-Delaroche. Purely infinite C^* -algebras arising from dynamical systems. *Bull. Soc. Math. France*, 125(2):199–225, 1997.
- [BD15] Jeffrey F. Brock and Nathan M. Dunfield. Injectivity radii of hyperbolic integer homology 3-spheres. *Geom. Topol.*, 19(1):497–523, 2015.
- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [BO08] Nathaniel P. Brown and Narutaka Ozawa. *C^* -algebras and finite-dimensional approximations*, volume 88 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [Bro82] Kenneth S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [CELY17] Joachim Cuntz, Siegfried Echterhoff, Xin Li, and Guoliang Yu. *K -theory for group C^* -algebras and semigroup C^* -algebras*, volume 47 of *Oberwolfach Seminars*. Birkhäuser/Springer, Cham, 2017.

- [EM06] Heath Emerson and Ralf Meyer. Euler characteristics and Gysin sequences for group actions on boundaries. *Math. Ann.*, 334(4):853–904, 2006.
- [ER18] Caleb Eckhardt and Sven Raum. C^* -superrigidity of 2-step nilpotent groups. *Advances in Mathematics*, 338:175–195, 2018.
- [GGKN23] Eusebio Gardella, Shirly Geffen, Julian Kranz, and Petr Naryshkin. Classifiability of crossed products by nonamenable groups. *J. Reine Angew. Math.*, 797:285–312, 2023.
- [Kas88] Gennadi G. Kasparov. Equivariant KK -theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [Kir94] Eberhard Kirchberg. The classification of purely infinite C^* -algebras using Kasparov’s theory. *Preprint*, 1994.
- [KR00] Eberhard Kirchberg and Mikael Rørdam. Non-simple purely infinite C^* -algebras. *Amer. J. Math.*, 122(3):637–666, 2000.
- [Laf12] Vincent Lafforgue. La conjecture de Baum-Connes à coefficients pour les groupes hyperboliques. *J. Noncommut. Geom.*, 6(1):1–197, 2012.
- [Lee18] John M. Lee. *Introduction to Riemannian manifolds*, volume 176 of *Graduate Texts in Mathematics*. Springer, Cham, second edition, 2018.
- [Mat02] Michel Matthey. Mapping the homology of a group to the K -theory of its C^* -algebra. *Illinois J. Math.*, 46(3):953–977, 2002.
- [Mat12] Hiroki Matui. Homology and topological full groups of étale groupoids on totally disconnected spaces. *Proc. Lond. Math. Soc. (3)*, 104(1):27–56, 2012.
- [Mc20] Bram Mesland and Mehmet Haluk Şengün. Hecke operators in KK -theory and the K -homology of Bianchi groups. *J. Noncommut. Geom.*, 14(1):125–189, 2020.
- [Mos68] G. D. Mostow. Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms. *Inst. Hautes Études Sci. Publ. Math.*, (34):53–104, 1968.
- [Phi00] N. Christopher Phillips. A classification theorem for nuclear purely infinite simple C^* -algebras. *Doc. Math.*, 5:49–114, 2000.
- [Ren08] Jean Renault. Cartan subalgebras in C^* -algebras. *Irish Math. Soc. Bull.*, (61):29–63, 2008.
- [RS87] Jonathan Rosenberg and Claude Schochet. The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K -functor. *Duke Math. J.*, 55(2):431–474, 1987.
- [Tu99] Jean-Louis Tu. La conjecture de Baum-Connes pour les feuilletages moyennables. *K-Theory*, 17(3):215–264, 1999.

SHIRLY GEFFEN, MATHEMATISCHES INSTITUT, FACHBEREICH MATHEMATIK UND INFORMATIK DER UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY.

Email address: sgeffen@uni-muenster.de

URL: <https://shirlygeffen.com/>

JULIAN KRANZ, INSTITUTE FOR ANALYSIS AND NUMERICS, APPLIED MATHEMATICS MÜNSTER, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY.

Email address: julian.kranz@uni-muenster.de

URL: <https://sites.google.com/view/juliankranz/>