

# Transport equation driven by a stochastic measure

Vadym Radchenko<sup>1 2</sup>

## Abstract

We consider the stochastic transport equation where the randomness is given by the symmetric integral with respect to stochastic measure. For stochastic measure, we assume only  $\sigma$ -additivity in probability and continuity of paths. The existence and uniqueness of the weak solution to the equation are proved.

## 1 Introduction

We consider the stochastic transport equation that formally can be written in the form

$$\begin{aligned} \frac{\partial u(t, x)}{\partial t} dt + b(t, x) \frac{\partial u(t, x)}{\partial x} dt + \frac{\partial u(t, x)}{\partial x} \circ d\mu(t) &= 0, \\ u(0, x) &= u_0(x), \quad x \in \mathbb{R}, \quad t \in [0, T]. \end{aligned} \tag{1}$$

Here  $\mu$  is a stochastic measure (SM), see Definition 1 below. We assume that  $\mu$  is defined on the Borel  $\sigma$ -algebra of  $[0, T]$ , and the process  $\mu_t = \mu((0, t])$  has a continuous paths. Assumptions on  $b$  and  $u_0$  are given in Section 3. Equation (1) we consider in the weak sense, definition of the solution is given in (6).

We will prove existence and uniqueness of the solution. Similarly to other types of stochastic transport equation, we demonstrate that the solution is given by formula  $u(t, x) = u_0(X_t^{-1}(x))$  where  $X_t(x)$  satisfies the auxiliary equation (7).

Stochastic integral with respect to  $\mu$  is defined as symmetric integral. This Stratonovich-type integral was studied in [17], we recall it's definition and basic properties in subsection 2.2. SMs include many important classes of processes, but we can prove existence of the integral only for integrands of the form  $f(\mu_t, t)$  where  $f \in \mathcal{C}^{1,1}(\mathbb{R} \times [0, T])$ . Thus, we will find our solution  $u$  having this form.

For the stochastic transport equation driven by the Wiener process, the existence and uniqueness of the solution were proved under different assumptions on  $b$  and  $u_0$ , see [3], [5], [7], [12], [24]. It was shown that stochastic term in the transport equation leads to regularization of the solution, see [1], [6], [7]. Equation in bounded domain was studied in [14]. In these papers, the stochastic term is given by Stratonovich integral and solution is considered in the weak sense. In [25] the existence and uniqueness of stochastic strong solution are obtained, the renormalized weak solution was studied in [27].

Transport equation with other stochastic integrators is less studied. The existence and uniqueness of the solution to equation driven by Lévy white noise was proved in [16], to equation driven by fractional Brownian motion – in [15]. In both papers, the Malliavin calculus approach was used.

In this paper, we consider the rather general stochastic integrator. At the same time, we need some restrictive assumptions on  $b$  and  $u_0$ , and study the case of one dimensional spatial variable.

The recent results for the equations driven by stochastic measures may be found in [2], [9], [10].

The rest of the paper is organized as follows. In Section 2 we recall the definitions and basic facts concerning stochastic measures and symmetric integral. Also we prove the analogue of the Fubini theorem for our integral that we will need below. In Section 3 we give our assumptions on the equation and formulate the main result. Section 4 is devoted to the proof of the existence of the solution, and we give the explicit formula for  $u$ . In Section 5, under some additional assumptions, we obtain the uniqueness of the solution.

---

<sup>1</sup>Department of Mathematical Analysis, Taras Shevchenko National University of Kyiv, 01601 Kyiv, Ukraine, vadymradchenko@knu.ua

<sup>2</sup>This work was supported by Alexander von Humboldt Foundation, grant 1074615. The author is grateful to Prof. M. Zähle for fruitful discussions during the preparation of this paper and thanks the Friedrich-Schiller-University of Jena for its hospitality.

## 2 Preliminaries

### 2.1 Stochastic measures

Let  $L_0 = L_0(\Omega, \mathcal{F}, \mathbb{P})$  be the set of all real-valued random variables defined on the complete probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  (more precisely, the set of equivalence classes). Convergence in  $L_0$  means the convergence in probability. Let  $X$  be an arbitrary set and  $\mathcal{B}$  a  $\sigma$ -algebra of subsets of  $X$ .

**Definition 1.** A  $\sigma$ -additive mapping  $\mu : \mathcal{B} \rightarrow L_0$  is called *stochastic measure* (SM).

We do not assume the moment existence or martingale properties for SM. In other words,  $\mu$  is  $L_0$ -valued vector measure.

Important examples of SMs are orthogonal stochastic measures,  $\alpha$ -stable random measures defined on a  $\sigma$ -algebra for  $\alpha \in (0, 1) \cup (1, 2]$  (see [20, Chapter 3]).

Many examples of the SMs on the Borel subsets of  $[0, T]$  may be given by the Wiener-type integral

$$\mu(A) = \int_{[0, T]} \mathbf{1}_A(t) dX_t. \quad (2)$$

We note the following cases of processes  $X_t$  in (2) that generate SM.

1.  $X_t$  – any square integrable continuous martingale.
2.  $X_t = W_t^H$  – the fractional Brownian motion with Hurst index  $H > 1/2$ , see Theorem 1.1 [11].
3.  $X_t = S_t^k$  – the sub-fractional Brownian motion for  $k = H - 1/2$ ,  $1/2 < H < 1$ , see Theorem 3.2 (ii) and Remark 3.3 c) in [22].
4.  $X_t = Z_H^k(t)$  – the Hermite process,  $1/2 < H < 1$ ,  $k \geq 1$ , see [8], [23, Section 3.1.3].  $Z_H^2(t)$  is known as the Rosenblatt process, see also [21, Section 3].

The detailed theory of stochastic measures is presented in [19].

The results of this paper will be obtained under the following assumption on  $\mu$ .

**Assumption A1.**  $\mu$  is an SM on Borel subsets of  $[0, T]$ , and the process  $\mu_t = \mu((0, t])$  has continuous paths on  $[0, T]$ .

Processes  $X_t$  in examples 1–4 are continuous, therefore A1 holds in these cases.

### 2.2 Symmetric integral

The symmetric integral of random functions with respect to stochastic measures was considered in [17]. We review the basic facts and definitions concerning this integral.

**Definition 2.** Let  $\xi_t$  and  $\eta_t$  be random processes on  $[0, T]$ ,  $0 = t_0^n < t_1^n < \dots < t_{j_n}^n = T$  be a sequence of partitions such that  $\max_k |t_k^n - t_{k-1}^n| \rightarrow 0$ ,  $n \rightarrow \infty$ . We define

$$\int_{(0, T]} \xi_t \circ d\eta_t := \text{p} \lim_{n \rightarrow \infty} \sum_{k=1}^{j_n} \frac{\xi_{t_{k-1}^n} + \xi_{t_k^n}}{2} (\eta_{t_k^n} - \eta_{t_{k-1}^n}) \quad (3)$$

provided that this limit in probability exists for any such sequence of partitions.

For Wiener process  $\eta_t$  and adapted  $\xi_t$  we obtain the classical Stratonovich integral. If  $\eta_t$  and  $\xi_t$  are Hölder continuous with exponents  $\gamma_\eta$  and  $\gamma_\xi$ ,  $\gamma_\eta + \gamma_\xi > 1$ , then value of (3) equals to the integral defined in [26].

The following theorem describes the class of processes for which the integral is well defined.

**Theorem 1.** (Theorem 4.6 [17]) Let A1 holds,  $f \in \mathbb{C}^{1,1}(\mathbb{R} \times [0, T])$ . Then integral (3) of  $f(\mu_t, t)$  with respect to  $\mu_t$  is well defined, and

$$\int_{(0,T]} f(\mu_t, t) \circ d\mu_t = F(\mu_T, T) - \int_{(0,T]} F'_2(\mu_t, t) dt, \quad (4)$$

where  $F(z, v) = \int_0^z f(y, v) dy$ .

Some other properties and equations with the symmetric integral are considered in [17], [18], [19].

### 2.3 Fubini theorem for symmetric integral

We will need the following auxiliary statement.

**Lemma 1.** Let  $f : \mathbb{R} \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  be measurable and has continuous derivatives  $f'_y(y, t, x)$ ,  $f'_t(y, t, x)$ . Assume that

$$|f(y, t, x)| \leq g(x), \quad |f'_y(y, t, x)| \leq g_1(x), \quad |f'_t(y, t, x)| \leq g_2(x)$$

for some  $g, g_1, g_2 \in L^1(\mathbb{R}, dx)$ . Then

$$\int_{\mathbb{R}} \int_{(0,T]} f(\mu_t, t, x) \circ d\mu_t dx = \int_{(0,T]} \int_{\mathbb{R}} f(\mu_t, t, x) dx \circ d\mu_t. \quad (5)$$

*Proof.* Denote

$$F(z, t, x) = \int_0^z f(y, t, x) dy, \quad \tilde{F}(z, t) = \int_0^z \int_{\mathbb{R}} f(y, t, x) dx dy, \quad z \in \mathbb{R}.$$

Theorem 1 and assumptions of the lemma and imply that the integrals in (5) are well-defined. Applying (4), we transform left-hand side and right-hand side of (5)

$$\begin{aligned} \int_{\mathbb{R}} \int_{(0,T]} f(\mu_t, t, x) \circ d\mu_t dx &= \int_{\mathbb{R}} \left( F(\mu_T, T, x) - \int_{(0,T]} F'_t(\mu_t, t, x) dt \right) dx, \\ \int_{(0,T]} \int_{\mathbb{R}} f(\mu_t, t, x) dx \circ d\mu_t &= \tilde{F}(\mu_T, T) - \int_{(0,T]} \tilde{F}'_t(\mu_t, t) dt. \end{aligned}$$

The equalities

$$\begin{aligned} &\int_{\mathbb{R}} F(\mu_T, T, x) dx = \tilde{F}(\mu_T, T) \\ &\Leftrightarrow \int_{\mathbb{R}} \int_0^{\mu_T} f(y, T, x) dy dx = \int_0^{\mu_T} \int_{\mathbb{R}} f(y, T, x) dx dy, \\ &\int_{\mathbb{R}} \int_{(0,T]} F'_t(\mu_t, t, x) dt dx = \int_{(0,T]} \tilde{F}'_t(\mu_t, t) dt \\ &\Leftrightarrow \int_{\mathbb{R}} \int_{(0,T]} \int_0^{\mu_t} f'_t(y, t, x) dy dt dx = \int_{(0,T]} \int_0^{\mu_t} \int_{\mathbb{R}} f'_t(y, t, x) dx dy dt. \end{aligned}$$

hold by usual Fubini's theorem. □

### 3 The problem. Formulation of the main result.

We consider equation (1) in the weak form. This means that  $u : [0, T] \times \mathbb{R} \times \Omega \rightarrow \mathbb{R}$  is a measurable random function such that for each  $\varphi \in \mathbb{C}_0^\infty(\mathbb{R})$  holds

$$\begin{aligned} \int_{\mathbb{R}} u(t, x) \varphi(x) dx &= \int_{\mathbb{R}} u_0(x) \varphi(x) dx + \int_0^t \int_{\mathbb{R}} u(s, x) \left( b(s, x) \varphi'(x) + \frac{\partial b(s, x)}{\partial x} \varphi(x) \right) dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}} u(s, x) \varphi'(x) dx \circ d\mu(s). \end{aligned} \quad (6)$$

By  $\mathbb{C}_0^\infty(\mathbb{R})$  we denote the class of infinitely differentiable functions with the compact support.

For our equation, we will refer to the following assumptions.

**Assumption A2.**  $u_0 : \mathbb{R} \times \Omega \rightarrow \mathbb{R}$  is measurable and has continuous derivative in  $x$ .

**Assumption A3.**  $|u_0(x)| \leq C(\omega)$  for some finite random constant  $C(\omega)$ .

**Assumption A4.**  $b : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous,  $\frac{\partial b(t, x)}{\partial x}$  is continuous and bounded.

**Assumption A5.**  $\sup_{t \in [0, T]} \int_{|x| \geq r} \frac{|b(t, x)|}{1 + |x|} dx \rightarrow 0$ ,  $r \rightarrow \infty$ .

Note that, by A4,  $b$  is globally Lipschitz continuous in  $x$ .

For each fixed  $\omega \in \Omega$ , we consider the following auxiliary equation

$$X_t(x) = x + \int_0^t b(r, X_r(x)) dr + \mu_t, \quad 0 \leq t \leq T. \quad (7)$$

Assumption A4 imply that (7) has a unique solution on  $[0, T]$  for each  $x$ .

By well known result of theory of ordinary differential equations, the solution has a continuous derivative

$$X'_t(x) = \frac{\partial}{\partial x} X_t(x).$$

We have

$$\begin{aligned} X'_t(x) &= 1 + \int_0^t \frac{\partial b(r, X_r(x))}{\partial x} X'_r(x) dr \\ \Rightarrow \frac{\partial}{\partial t} X'_t(x) &= \frac{\partial b(t, X_t(x))}{\partial x} X'_t(x) \quad \Rightarrow \quad X'_t(x) = \exp \left\{ \int_0^t \frac{\partial b(s, X_s(x))}{\partial x} ds \right\}. \end{aligned} \quad (8)$$

Therefore,  $X'_t(x) > 0$ , and the function  $X_t^{-1}(x)$ , where inverse is taken with respect to variable  $x$ , is well defined.

Note that  $X_t$  is the sum of a differentiable function of  $t$  and  $\mu_t$ ,  $X'_t$  is a differentiable function of  $t$ . This allows us to consider integrals

$$\int_{(0, T]} g(X_t, X'_t, \mu_t, t) \circ d\mu_t, \quad g \in \mathbb{C}^{1,1,1,1}(\mathbb{R}^3 \times [0, T]).$$

The main result of the paper is the following.

**Theorem 2.** 1) Let Assumptions A1, A2, A4 hold,  $X_t(x)$  be the solution of (7). Then the random function

$$u(t, x) = u_0(X_t^{-1}(x)) \quad (9)$$

satisfies (6).

2) In addition, let Assumptions A3 and A5 hold. Then solution (9) is unique in the class of measurable random functions  $u(t, x) = h(\mu_t, t, x)$ , such that  $h(\cdot, \cdot, x) \in \mathbb{C}^{1,1}(\mathbb{R} \times [0, T])$  for each  $x \in \mathbb{R}$ , and  $|u(t, x)| \leq C(\omega)$  for some finite random constant  $C(\omega)$ .

*Remark 1.* Note that  $u(t, x) = u_0(X_t^{-1}(x))$  has a form  $h(\mu_t, t, x)$  from the second part of the theorem. This follows from Assumption A2 and standard statements about differentiability of inverse functions. We have that  $X_t(x) = g(\mu_t, t, x)$ , where  $g \in \mathbb{C}^{1,1,1}(\mathbb{R} \times [0, T] \times \mathbb{R})$ . For the mapping

$$(\mu, t, x) \rightarrow (\mu, t, g(\mu, t, x)),$$

the matrix of the first derivatives is non-degenerated. Therefore, the inverse mapping is well-defined and smooth.

*Remark 2.* Let us compare our assumptions with those made in other papers. Usually, it is supposed that  $u_0$  is measurable and bounded (see, for example, [7], [12], [15], [24]). We additionally assume that  $u_0$  has a continuous derivative, we need this to guarantee that the symmetric integral of  $u_0(X_t^{-1}(x))$  be well-defined.

Condition of differentiability of  $b$  is standard, boundedness of  $\frac{\partial b}{\partial x}$  may be assumed in some  $L_p$  norm (see [7], [24]) or uniformly ([15]). Note that in [12] the main result was obtained for arbitrary bounded measurable  $b$ .

Our integrability condition A5 is technical and is important for our method. It is similar to respective assumptions in [1], [3], [13].

## 4 Existence of the solution

In this section, we prove the first statement of our theorem.

By the chain rule (4), for  $\varphi \in C_0^\infty(\mathbb{R})$  we have

$$\begin{aligned} d_t [X'_t(x)\varphi(X_t(x))] &= \varphi(X_t(x))d_t [X'_t(x)] + X'_t(x)d_t [\varphi(X_t(x))] \\ \stackrel{(7),(8)}{=} \varphi(X_t(x))\frac{\partial b(t, X_t(x))}{\partial x}X'_t(x) dt + X'_t(x)\varphi'(X_t(x))b(t, X_t(x)) dt + X'_t(x)\varphi'(X_t(x)) \circ d\mu(t). \end{aligned} \quad (10)$$

Applying the change of variables  $y = X_t(x)$ , we get

$$\begin{aligned} \int_{\mathbb{R}} u_0(X_t^{-1}(y))\varphi(y) dy &= \int_{\mathbb{R}} u_0(x)X'_t(x)\varphi(X_t(x)) dx \\ &= \int_{\mathbb{R}} u_0(x) [X'_t(x)\varphi(X_t(x))\Big|_{t=0} + \int_0^t d_s [X'_s(x)\varphi(X_s(x))]] dx \\ &\stackrel{(10)}{=} \int_{\mathbb{R}} u_0(x)\varphi(x) dx + \int_{\mathbb{R}} u_0(x) \int_0^t \varphi(X_s(x))\frac{\partial b(s, X_s(x))}{\partial x}X'_s(x) ds dx \\ &+ \int_{\mathbb{R}} u_0(x) \int_0^t X'_s(x)\varphi'(X_s(x))b(s, X_s(x)) ds dx + \int_{\mathbb{R}} u_0(x) \int_0^t X'_s(x)\varphi'(X_s(x)) \circ d\mu_s dx \\ &\stackrel{(5)}{=} \int_{\mathbb{R}} u_0(x)\varphi(x) dx + \int_0^t \int_{\mathbb{R}} u_0(x)\varphi(X_s(x))\frac{\partial b(s, X_s(x))}{\partial x}X'_s(x) dx ds \\ &+ \int_0^t \int_{\mathbb{R}} u_0(x)X'_s(x)\varphi'(X_s(x))b(s, X_s(x)) dx ds + \int_0^t \int_{\mathbb{R}} u_0(x)X'_s(x)\varphi'(X_s(x)) dx \circ d\mu_s. \end{aligned}$$

Lemma 1 may be applied here because  $\varphi$  has a compact support. Assumption A4 and (8) imply that  $C_1 \leq X'_s \leq C_2$  for some positive constants  $C_1$  and  $C_2$ , therefore set  $\{x : \varphi'(X_s(x)) \neq 0\}$  is bounded.

Taking the inverse change of variable  $x = X_t^{-1}(y)$ , obtain

$$\begin{aligned} \int_{\mathbb{R}} u_0(X_t^{-1}(y))\varphi(y) dy &= \int_{\mathbb{R}} u_0(x)\varphi(x) dx + \int_0^t \int_{\mathbb{R}} u_0(X_s^{-1}(y))\varphi(y)\frac{\partial b(s, y)}{\partial x} dy ds \\ &+ \int_0^t \int_{\mathbb{R}} u_0(X_s^{-1}(y))\varphi'(y)b(s, y) dy ds + \int_0^t \int_{\mathbb{R}} u_0(X_s^{-1}(y))\varphi'(y) dy \circ d\mu_s. \end{aligned}$$

Thus,  $u(t, x) = u_0(X_t^{-1}(x))$  satisfies (6).

## 5 Uniqueness of the solution

In this section, we prove the second statement of our theorem. We will follow the standard approach (see, for example, proof of the uniqueness of the solution in [3], [13]).

Let  $u(t, x)$  satisfies (6) with  $u_0(x) = 0$ . We will obtain that  $u(t, x) = 0$  what implies the uniqueness of the solution.

For this case, from (6) for  $\varphi \in \mathbb{C}_0^\infty(\mathbb{R})$  we get

$$\begin{aligned} \int_{\mathbb{R}} u(t, x) \varphi(x) dx &= \int_0^t \int_{\mathbb{R}} u(s, x) \left( b(s, x) \varphi'(x) + \frac{\partial b(s, x)}{\partial x} \varphi(x) \right) dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}} u(s, x) \varphi'(x) dx \circ d\mu_s. \end{aligned} \quad (11)$$

Denote

$$G(t, y) = \int_{\mathbb{R}} u(t, x) \varphi(x - y) dx$$

We have that  $G(0, y) = 0$  because  $u(0, x) = 0$ , and

$$\frac{\partial G(t, y)}{\partial y} = - \int_{\mathbb{R}} u(t, x) \varphi'(x - y) dx,$$

Our solution has a form  $u(t, x) = h(\mu_t, t, x)$ , and, applying (4) and (11), we obtain

$$\begin{aligned} &\int_{\mathbb{R}} u(t, x) \varphi(x - \mu_t) dx = G(t, \mu_t) \\ &= \int_0^t \int_{\mathbb{R}} b(s, x) u(s, x) \varphi'(x - \mu_s) dx ds + \int_0^t \int_{\mathbb{R}} \frac{\partial b(s, x)}{\partial x} u(s, x) \varphi(x - \mu_s) dx ds \\ &\quad + \int_0^t \int_{\mathbb{R}} u(s, x) \varphi'(x - \mu_s) dx \circ d\mu_s - \int_0^t \int_{\mathbb{R}} u(s, x) \varphi'(x - \mu_s) dx \circ d\mu_s \\ &= \int_0^t \int_{\mathbb{R}} b(s, x) u(s, x) \varphi'(x - \mu_s) dx ds + \int_0^t \int_{\mathbb{R}} \frac{\partial b(s, x)}{\partial x} u(s, x) \varphi(x - \mu_s) dx ds. \end{aligned}$$

For  $V(t, z) = u(t, z + \mu_t)$ , taking change of the variable  $x = z + \mu_t$ , get

$$\begin{aligned} \int_{\mathbb{R}} V(t, z) \varphi(z) dz &= \int_0^t \int_{\mathbb{R}} b(s, z + \mu_s) V(s, z) \frac{d\varphi(z)}{dz} dz ds \\ &\quad + \int_0^t \int_{\mathbb{R}} \frac{\partial b(s, z + \mu_s)}{\partial z} V(s, z) \varphi(z) dz ds. \end{aligned} \quad (12)$$

Let  $\phi_\varepsilon$  be a standard mollifier,

$$\phi_\varepsilon(x) = \frac{1}{\varepsilon} \phi\left(\frac{x}{\varepsilon}\right), \quad \phi \in \mathbb{C}_0^\infty(\mathbb{R}), \quad \text{supp } \phi \subset [-1, 1], \quad \phi(x) \geq 0, \quad \int_{\mathbb{R}} \phi(x) dx = 1.$$

Denote  $V_\varepsilon(t, x) := V(t, \cdot) * \phi_\varepsilon$ . Substituting  $\varphi(z) = \phi_\varepsilon(x - z)$  in (12), obtain

$$\begin{aligned} V_\varepsilon(t, x) &= \int_{\mathbb{R}} V(t, z) \phi_\varepsilon(x - z) dz \\ &= - \int_0^t \int_{\mathbb{R}} b(s, z + \mu_s) V(s, z) \phi_\varepsilon'(x - z) dz ds + \int_0^t \int_{\mathbb{R}} \frac{\partial b(s, z + \mu_s)}{\partial z} V(s, z) \phi_\varepsilon(x - z) dz ds. \end{aligned}$$

We take the derivative with respect to  $t$ , use the notation  $B(t, z) = b(t, z + \mu_t)$ , and get

$$\begin{aligned} \frac{\partial V_\varepsilon(t, x)}{\partial t} &= - \int_{\mathbb{R}} B(t, z) V(t, z) \frac{\partial \phi_\varepsilon(x - z)}{\partial x} dz + \int_{\mathbb{R}} \frac{\partial B(t, z)}{\partial z} V(t, z) \phi_\varepsilon(x - z) dz \\ &= - \frac{\partial}{\partial x} \int_{\mathbb{R}} B(t, z) V(t, z) \phi_\varepsilon(x - z) dz + \int_{\mathbb{R}} \left[ \frac{\partial}{\partial z} [B(t, z) V(t, z)] - B(t, z) \frac{\partial V(t, z)}{\partial z} \right] \phi_\varepsilon(x - z) dz \\ &\quad \stackrel{(*)}{=} - \frac{\partial}{\partial x} (BV(t, \cdot) * \phi_\varepsilon)(x) + \frac{\partial}{\partial x} (BV(t, \cdot) * \phi_\varepsilon)(x) \\ &\quad - \int_{\mathbb{R}} B(t, z) \frac{\partial V(t, z)}{\partial z} \phi_\varepsilon(x - z) dz = - \int_{\mathbb{R}} B(t, z) \frac{\partial V(t, z)}{\partial z} \phi_\varepsilon(x - z) dz. \end{aligned}$$

In (\*) we have used that  $\phi_\varepsilon$  has a compact support, and, by integration in parts,

$$\begin{aligned} & \int_{\mathbb{R}} \frac{\partial}{\partial z} [B(t, z)V(t, z)] \phi_\varepsilon(x - z) dz = \int_{\mathbb{R}} \phi_\varepsilon(x - z) d[B(t, z)V(t, z)] \\ &= - \int_{\mathbb{R}} [B(t, z)V(t, z)] d_z \phi_\varepsilon(x - z) = \int_{\mathbb{R}} [B(t, z)V(t, z)] \frac{\partial \phi_\varepsilon(x - z)}{\partial x} dz = \frac{\partial}{\partial x} (BV(t, \cdot) * \phi_\varepsilon)(x). \end{aligned}$$

Thus,

$$\frac{\partial V_\varepsilon(t, x)}{\partial t} + \left( B(t, x) \frac{\partial V(t, x)}{\partial x} \right) * \phi_\varepsilon(x) = 0. \quad (13)$$

Denote

$$\mathcal{R}_\varepsilon(B, V) = \frac{\partial V_\varepsilon(t, x)}{\partial t} + B(t, x) \frac{\partial V_\varepsilon(t, x)}{\partial x} \stackrel{(13)}{=} B \frac{\partial(\phi_\varepsilon * V)}{\partial x} - \phi_\varepsilon * \left( B \frac{\partial V}{\partial x} \right). \quad (14)$$

Lemma II.1 i) [4] gives that for each fixed  $t$

$$\mathcal{R}_\varepsilon(B, V_\varepsilon) \rightarrow 0, \quad \varepsilon \rightarrow 0 \quad \text{in} \quad \mathbf{L}_{loc}^1(\mathbb{R}, dx) \quad (15)$$

provided that  $B(t, \cdot) \in \mathbf{W}_{loc}^{1,1}(\mathbb{R})$ ,  $V(t, \cdot) \in \mathbf{L}_{loc}^\infty(\mathbb{R}, dx)$ , where  $\mathbf{W}$  denotes the Sobolev space. These conditions hold due to assumptions of our theorem.

Consider  $\pi_r(x) = \pi_1(x/r)$ , where

$$\pi_1(x) = \begin{cases} 1, & |x| < 1, \\ \in [0, 1], & 1 \leq |x| \leq 2, \\ 0, & |x| > 2. \end{cases}$$

with  $|\pi_r'| \leq \frac{C}{r}$ . We have that

$$\begin{aligned} \int_{\mathbb{R}} d_x (B(V_\varepsilon)^2 \pi_r) = 0 & \Leftrightarrow \int_{\mathbb{R}} (V_\varepsilon)^2 \pi_r d_x B + \int_{\mathbb{R}} B \pi_r d_x (V_\varepsilon)^2 + \int_{\mathbb{R}} B (V_\varepsilon)^2 d\pi_r = 0 \\ & \Leftrightarrow \int_{\mathbb{R}} B \pi_r V_\varepsilon \frac{\partial V_\varepsilon}{\partial x} dx = -\frac{1}{2} \int_{\mathbb{R}} (V_\varepsilon)^2 \pi_r \frac{\partial B}{\partial x} dx - \frac{1}{2} \int_{\mathbb{R}} B (V_\varepsilon)^2 \pi_r' dx. \end{aligned}$$

We multiply (14) by  $V_\varepsilon(t, x)\pi_r(x)$ , take the integral over  $\mathbb{R}$ , and get

$$\begin{aligned} \int_{\mathbb{R}} \mathcal{R}_\varepsilon(B, V_\varepsilon) V_\varepsilon \pi_r dx &= \frac{1}{2} \int_{\mathbb{R}} \frac{\partial V_\varepsilon^2}{\partial t} \pi_r dx + \int_{\mathbb{R}} B V_\varepsilon \pi_r \frac{\partial V_\varepsilon}{\partial x} dx \\ \Leftrightarrow \int_{\mathbb{R}} \mathcal{R}_\varepsilon(B, V_\varepsilon) V_\varepsilon \pi_r dx &= \frac{1}{2} \frac{\partial}{\partial t} \int_{\mathbb{R}} V_\varepsilon^2 \pi_r dx - \frac{1}{2} \int_{\mathbb{R}} (V_\varepsilon)^2 \pi_r \frac{\partial B}{\partial x} dx - \frac{1}{2} \int_{\mathbb{R}} B (V_\varepsilon)^2 \pi_r' dx. \end{aligned}$$

From (15) it follows that for fixed  $r$  and  $t$

$$\int_{\mathbb{R}} \mathcal{R}_\varepsilon(B, V_\varepsilon) V_\varepsilon \pi_r dx \rightarrow 0, \quad \varepsilon \rightarrow 0.$$

Therefore,

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \left( \frac{\partial}{\partial t} \int_{\mathbb{R}} V_\varepsilon^2 \pi_r dx - \int_{\mathbb{R}} (V_\varepsilon)^2 \pi_r \frac{\partial B}{\partial x} dx - \int_{\mathbb{R}} B (V_\varepsilon)^2 \pi_r' dx \right) &= 0 \\ \Leftrightarrow \frac{\partial}{\partial t} \int_{\mathbb{R}} V^2 \pi_r dx - \int_{\mathbb{R}} V^2 \pi_r \frac{\partial B}{\partial x} dx &= \int_{\mathbb{R}} B V^2 \pi_r' dx. \end{aligned} \quad (16)$$

Because  $\pi_r'(x) = 0$  for  $|x| \leq r$  or  $|x| \geq 2r$ , and  $|\pi_r'| \leq \frac{C}{r}$ , we have

$$\left| \int_{\mathbb{R}} B V^2 \pi_r' dx \right| \leq \|V\|_{\mathbf{L}^\infty}^2 \int_{r \leq |x| \leq 2r} \frac{|B(t, x)|}{1 + |x|} (1 + |x|) |\pi_r'(x)| dx \rightarrow 0, \quad r \rightarrow \infty, \quad (17)$$

where convergence holds uniformly in  $t$  for each fixed  $\omega$ .

In (17) we have used the following estimates. If  $\sup_t |\mu_t| = M(\omega)$  then

$$\begin{aligned} & \int_{r \leq |x| \leq 2r} \frac{|B(t, x)|}{1 + |x|} dx \stackrel{y=x+\mu_t}{=} \int_{r \leq |y-\mu_t| \leq 2r} \frac{|b(t, y)|}{1 + |y - \mu_t|} dy \\ & \leq \int_{|y| \geq r-M(\omega)} \frac{|b(t, y)|}{1 + |y| - M(\omega)} dy \stackrel{A5}{\xrightarrow{}} 0, \quad r \rightarrow \infty. \end{aligned}$$

Integrating (16) in  $t$  and taking into account that

$$\int_{\mathbb{R}} V^2(t, x) \pi_r(x) dx \Big|_{t=0} = \int_{\mathbb{R}} u(0, x)^2 \pi_r(x) dx = 0,$$

we get

$$\int_{\mathbb{R}} V^2 \pi_r dx = \int_0^t \int_{\mathbb{R}} V^2 \pi_r \frac{\partial B}{\partial x} dx ds + \int_0^t \int_{\mathbb{R}} BV^2 \pi_r' dx ds \quad (18)$$

Consider

$$g_r(t, x) = V^2(t, x) \pi_r(x).$$

By A4, we have  $\left| \frac{\partial B}{\partial x} \right| \leq K$  for some constant  $K$ . From (18), we get

$$\begin{aligned} \int_{\mathbb{R}} g_r(t, x) dx & \leq \int_0^t \int_{\mathbb{R}} g_r(s, x) \left| \frac{\partial B}{\partial x} \right| dx ds + R_r \leq K \int_0^t \int_{\mathbb{R}} g_r(s, x) dx ds + R_r, \\ R_r & = \sup_t \left| \int_0^t \int_{\mathbb{R}} BV^2 \pi_r' dx ds \right| \stackrel{(17)}{\xrightarrow{}} 0, \quad r \rightarrow \infty. \end{aligned}$$

From the Gronwall inequality for  $h(t) = \int_{\mathbb{R}} g_r(t, x) dx$ , we get

$$\int_{\mathbb{R}} g_r(t, x) dx \leq R_r e^{Kt}.$$

Taking  $r \rightarrow \infty$ , we get

$$\int_{\mathbb{R}} g_r(t, x) dx \rightarrow \int_{\mathbb{R}} V^2 dx, \quad R_r e^{Kt} \rightarrow 0 \Rightarrow V = 0$$

that finishes the proof of uniqueness of the solution.

## References

- [1] Beck, L., Flandoli, F., Gubinelli, M., Maurelli, M.: Stochastic ODEs and stochastic linear PDEs with critical drift: regularity, duality and uniqueness. *Electron. J. Probab.* **24**, 1–72 (2019)
- [2] Bodnarchuk, I.: Averaging principle for a stochastic cable equation. *Mod. Stoch. Theory Appl.* **7**(4), 449–467 (2020)
- [3] Catuogno, P., Olivera, C.:  $L_p$ -solutions of the stochastic transport equation. *Random Oper. Stoch. Equ.* **21**(2), 125–134 (2013)
- [4] DiPerna, R.J., Lions, P.-L.: Ordinary differential equations, transport theory and Sobolev spaces. *Invent. math.* **98**(3), 511–547 (1989)
- [5] Fang, S., Luo, D.: Flow of homeomorphisms and stochastic transport equations. *Stoch. Anal. Appl.* **25**(5), 1079–1108 (2007)



- [6] Fedrizzi, E., Flandoli, F.: Noise prevents singularities in linear transport equations. *J. Funct. Anal.* **264**(6), 1329–1354 (2013)
- [7] Flandoli, F., Gubinelli, M., Priola, E.: Well-posedness of the transport equation by stochastic perturbation. *Invent. math.* **180**(1), 1–53 (2010)
- [8] Maejima, M., Tudor, C.: Wiener integrals with respect to the Hermite process and a non-central limit theorem. *Stoch. Anal. Appl.* **25**(5), 1043–1056 (2007)
- [9] Manikin, B.: Asymptotic properties of the parabolic equation driven by stochastic measure. *Mod. Stoch. Theory Appl.*, 1–16 (2022)
- [10] Manikin, B.: Averaging principle for the one-dimensional parabolic equation driven by stochastic measure. *Mod. Stoch. Theory Appl.* **9**(2), 123–137 (2022)
- [11] Memin, T., Mishura, Y., Valkeila, E.: Inequalities for the moments of Wiener integrals with respect to a fractional Brownian motion. *Statist. Probab. Lett.* **51**, 197–206 (2001)
- [12] Mohammed, S.-E.A., Nilssen, T.K., Proske, F.N.: Sobolev differentiable stochastic flows for SDEs with singular coefficients: Applications to the transport equation. *Ann. Probab.* **43**(3), 1535–1576 (2015)
- [13] Mucha, P.B.: Transport equation: Extension of classical results for  $div b \in BMO$ . *J. Differential Equations* **249**(8), 1871–1883 (2010)
- [14] Neves, W., Olivera, C.: Initial-boundary value problem for stochastic transport equations. *Stoch. Partial Differ. Equ.: Anal. Comput.* **9**(3), 674–701 (2021)
- [15] Olivera, C., Tudor, C.: The density of the solution to the stochastic transport equation with fractional noise. *J. Math. Anal. Appl.* **431**(1), 57–72 (2015)
- [16] Proske, F.: The stochastic transport equation driven by Lévy white noise. *Commun. Math. Sci.* **2**(4), 627–641 (2004)
- [17] Radchenko, V.: Stratonovich-type integral with respect to a general stochastic measure. *Stochastics* **88**, 1060–1072 (2016)
- [18] Radchenko, V.: Averaging principle for equation driven by a stochastic measure. *Stochastics* **91**(6), 905–915 (2019)
- [19] Radchenko, V.: *General Stochastic Measures: Integration, Path Properties, and Equations*. Wiley – ISTE, London (2022)
- [20] Samorodnitsky, G., Taqqu, M.S.: *Stable Non-Gaussian Random Processes*. Chapman and Hall, London (1994)
- [21] Tudor, C.: Analysis of the Rosenblatt process. *ESAIM-Probab. Stat.* **12**, 230–257 (2008)
- [22] Tudor, C.: On the Wiener integral with respect to a sub-fractional Brownian motion on an interval. *J. Math. Anal. Appl.* **351**(1), 456–468 (2009)
- [23] Tudor, C.: *Analysis of Variations for Self-similar Processes: A Stochastic Calculus Approach*. Springer (2013)
- [24] Wei, J., Lv, G., Wang, W.: Stochastic transport equation with bounded and Dini continuous drift. *J. Differential Equations* **323**, 359–403 (2022)
- [25] Wei, J., Duan, J., Gao, H., Lv, G.: Stochastic regularization for transport equations. *Stoch. Partial Differ. Equ.: Anal. Comput.* **9**(1), 105–141 (2021)

- [26] Zähle, M.: Integration with respect to fractal functions and stochastic calculus. I. Probab. Theory Related Fields **111**(3), 333–374 (1998)
- [27] Zhang, X.: Stochastic flows of SDEs with irregular coefficients and stochastic transport equations. Bull. des Sci. Math. **134**(4), 340–378 (2010)