

First experimental test of the ratio method for nuclear-reaction analysis

S. Ota,^{1,2,*} P. Capel,^{3,†} G. Christian,^{4,5} V. Durant,³ K. Hagel,² E. Harris,^{2,5} R. C. Johnson,⁶ Z. Luo,^{2,5} F. M. Nunes,^{7,8} M. Roosa,^{2,5} A. Saastamoinen,² and D. P. Scriven^{2,5}

¹*National Nuclear Data Center, Brookhaven National Laboratory, Upton, NY 11973-5000, USA*

²*Cyclotron Institute, Texas A&M University, College Station, TX 77843, USA*

³*Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany*

⁴*Department of Astronomy & Physics, Saint Mary's University, Halifax, NS B3H 3C3, Canada*

⁵*Department of Physics & Astronomy, Texas A&M University, College Station, TX 77843, USA*

⁶*Department of Physics, University of Surrey, Guildford GU2 7XH, United Kingdom*

⁷*Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA*

⁸*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA*

(Dated: July 23, 2024)

Nuclear halos are very exotic quantal structures observed far from stability. Because of their short lifetime, they are mostly studied through reactions. The *ratio method* offers a new observable: the ratio of angular differential cross sections for breakup and scattering. It is predicted to be much more sensitive to the projectile structure than individual cross sections thanks to its independence of the reaction process. We test this new observable experimentally for the first time considering the collision of ^{11}Be on C at 22.8 MeV/nucleon. We extend this analysis to similar data recently measured on Pb at 19.1 MeV/nucleon. Both analyses confirm the theoretical predictions, which opens the door to a new era in the study of nuclear structure near the neutron dripline. This should prove invaluable in conjunction with the start of FRIB. The ratio method could also be extended to other fields of quantum physics beyond nuclear reactions.

Quantum collisions are a powerful tool in many fields of quantum physics. They provide key information about the interactions between particles (molecules, atoms, nuclei, hadrons. . .). They are also used to measure the rate of reactions that have a wide range of interests: from fundamental physics, such as astrophysics, to applied physics, such as nuclear engineering. Finally, collisions are sometimes the only way to study exotic quantal structures. This is particularly true in the realm of nuclear physics, where the development of radioactive ion beams in the 1980s has enabled the study of nuclei far from stability, leading to the discovery of *halo nuclei* [1, 2].

Halo nuclei exhibit an unusually large size compared to their isobars, breaking the empirical rule that nuclear radii scale with $A^{1/3}$, the cubic root of the mass number. They are located near the neutron dripline. In this extremely unstable region of the nuclear chart, valence neutrons can be very loosely bound to the nuclei. Therefore, they can tunnel far into the classically forbidden region to form a diffuse halo surrounding a dense and compact core [3]. This exotic structure has been the subject of many studies, both experimental and theoretical [4].

Because of their short lifetime, halo nuclei are mostly studied through reactions. Their exceptional spatial extension was first evidenced thanks to the large interaction cross sections measured in collisions with various targets [1, 2, 4]. In knockout, the halo neutrons are removed at high energies on a light target. The presence of a halo leads to a narrow momentum distribution of the outgoing core [4, 5]. A more exclusive measurement is the

diffractive breakup, where both the core and the valence neutrons are measured in coincidence [6, 7]. The corresponding cross section is large due to the fragile nature of the halo structure. In elastic scattering, the significant coupling to breakup strongly affects the differential cross section [8, 9]. Halo states can also be directly populated via (d, p) transfer reactions [10, 11].

All these reactions provide valuable information about the structure of halo nuclei. Unfortunately, inferring quantitatively accurate structure properties remains difficult because of significant ambiguities related to the reaction model. In particular, the optical potentials, which simulate the interaction between the constituents of the nucleus and the target, bear a systematic uncertainty in the analysis of data [12–15]. The *ratio method* has been proposed theoretically to circumvent this issue [16] (see Ref. [17] for a recent review). The idea is to consider the ratio of angular distributions for breakup and scattering instead of measuring cross sections for the individual channels. Within the recoil excitation and breakup model (REB) [18, 19], this observable is predicted to be independent of the reaction process, and in particular of the projectile-target interaction. Better still, according to the REB, it can be expressed entirely in terms of halo-nucleus form factors. Accordingly, the ratio should give direct access to the nuclear structure of the projectile [16, 20]. Using accurate models of the collision, various theoretical studies have confirmed these predictions and demonstrated the applicability of the ratio method beyond the range of validity of the REB. The method is expected to be valid for collisions involving neutron halos in a wide range of energies down to 20 MeV/nucleon [21] as well as for proton halos [22].

Until now, the ratio method has not been experimen-

* sota@bnl.gov

† pcapel@uni-mainz.de

tally verified because the breakup and scattering channels have not yet been measured at the same beam energy on the same target. In this letter, we report the first simultaneous measurement of both channels for ^{11}Be impinging on C at 22.8 MeV/nucleon. We analyze these new data in the framework of the ratio method. We extend this analysis to the ^{11}Be -Pb collision at 19.1 MeV/nucleon measured in Lanzhou by Duan *et al.* [23].

The experiment was performed at Texas A&M University (TAMU) using the K500 superconducting cyclotron. A ^{11}Be beam was produced by bombarding a 1 mm thick ^9Be target with a 30 MeV/nucleon ^{13}C primary beam. The fragments were selected with a dipole magnet and a velocity filter (momentum acceptance $\delta p/p = \pm 0.66\%$ (FWHM)) following the production, and then ^{11}Be (4^+) ions were separated from other isotopes using the MARS dipole magnet [24]. The ^{11}Be beam with the energy of 22.8 ± 0.3 MeV/nucleon was delivered to the target chamber. At the beginning of the experiment, the profile and purity of the ^{11}Be beam were measured at a reduced beam rate using a Si ΔE - E telescope with an active area of 5×5 cm² placed approximately 45 cm upstream from the reaction target. The ΔE detector is resistive along the MARS's dispersive axis (y) and segmented by 16 strips along the non-dispersive axis (x), leading to the position resolution of approximately $\delta y = 0.2$ mm and $\delta x = 3.1$ mm. The beam was limited with a slit (± 5 mm in y and ± 25 mm in x) placed ~ 10 cm upstream from the telescope to block other beam species by limiting the magnetic rigidity. The beam spot size was measured with the Si telescope to be $\sigma_x \sim 3$ mm and $\sigma_y \sim 2$ mm. The beam spot size remained nearly the same at the reaction target position as described below. The Si detectors were then removed from the beamline and the total intensity of the beam was determined with a phoswich detector to be typically 1.2×10^4 particles per second (pps), among which the ^{11}Be beam intensity was about 7.2×10^3 pps. Thus, the ^{11}Be beam purity was determined to be about 60% with 6% from ^8Li and the remaining from not well-focused light ions.

For the scattering/reaction target, we used natural carbon in the form of flexible graphite with 99.8% purity with a thickness of 8.7 (3) mg/cm². The thickness was determined from a weighted average of the values measured by two different weighing scales, a caliper, and the energy loss of ^{11}Be scattering events. To check the scattering yields and the thickness of the target from the energy loss, two other targets with thickness 17.4 (6) mg/cm² and 34.8 (12) mg/cm² were also used during the experiment. Consistency of the yields after normalization to the incident beam counts was confirmed within the uncertainty of the target thicknesses. The target size was limited to a 17 mm diameter and the incident beam outside the area was blocked with an aluminum target holder.

The BlueSTEAL detector system [25] was used to measure the scattering and breakup cross sections in the ^{11}Be -C collision. Because the removed neutron was not

measured, the latter corresponds to inclusive breakup. This detector comprises four layers of Micron S2 Si DSSD (one 0.5 mm thick and three 1.5 mm thick) and a phoswich plastic scintillator with 1.1 cm thickness. Two ΔE - E telescopes were built from the four Si DSSDs [0.5 mm + 1.5 mm (Si 1&2) and two 1.5 mm (Si 3&4)] (see Fig. 13(d) in Ref. [25]). In the first part of the experiment, we placed the Si 1&2 telescope at 5.08 cm and the Si 3&4 telescope at 17.78 cm downstream from the target [Config. 1]. In the second part of the experiment, we placed the Si 3&4 telescope at 10.16 cm downstream from the target [Config. 2] while the Si 1&2 telescope was placed 2.54 cm upstream from the target to monitor the beam's spatial spread and purity and to shield the Si 3&4 telescope from the defocused beam. As discussed in detail in Ref. [25], the measurements with the two configurations supplement each other, enabling us to double check the consistency of the measured cross sections in the overlapping angular range. Since the relatively broad beam-spot size causes poor angular resolution for the Si 1&2 detectors in Config. 1, it was difficult to extract precise angular cross sections. Therefore we have not used the data from that telescope.

As shown in Fig. 1(a), a clear particle identification (PID) was obtained with the Si 3&4 telescope. The numbers of ^{11}Be and ^{10}Be detected at different angles were used to obtain differential cross sections for both scattering and inclusive breakup. The beam offset from the telescope's center was deduced from the scattering measured at different azimuthal angles as explained in Ref. [25]. It was typically a few mm in x and y directions with an uncertainty $\Delta x, y < (0.5, 0.5)$ mm. During the experiment, background measurements were performed with a blank target and also with the target holder removed. Negligible ($< 1\%$) background $^{11,10}\text{Be}$ events measured with the blank target [inset of Fig. 1(a)] ensure that the observed $^{11,10}\text{Be}$ came from scattering/reaction with the target. The phoswich detector was placed about 30 cm downstream from the target to count the beam rate with clear elemental identification [Fig. 1(b)]. The PID in the phoswich's E_{short} vs E_{total} plot was made with the help of some punch-through events from the Si telescope (e.g., ^8Li , α -particles, deuterons and protons; see Ref. [25]). The beam size at the target position was checked by comparing the ^{11}Be count rates with the phoswich during the two separate runs with and without the target holder. The difference in the count rates was negligible ($< 3\%$) and was dominated by the beam-intensity fluctuation. Therefore almost all the ^{11}Be beam flux passed the target within the 17 mm diameter, without being blocked by the target holder. An additional test was made by measuring the beam halo with some inner rings of the Si 1&2 telescope without the target holder. The halo count rates measured with the Si telescope were less than 1–2% of the ^{11}Be count rates measured with the phoswich. Through GEANT4 simulations, it was confirmed that such count rates were possible only when the beam's divergence from the telescope for the beam diagnosis is negligibly small




FIG. 1. (a) Particle identification by ΔE - E plot from the downstream Si telescope (Config. 1; 14 h). Inset: same plot with a blank target (1 h), which contains $< 1\%$ of ^{11}Be observed with a 8.7 mg/cm^2 carbon target during the same time. (b) Particle identification by E_{short} vs E_{total} plot with the phoswich detector. Inset: E_{total} spectrum gated on Be in the two-dimensional plot.

($\sigma_{\theta} < 0.3^{\circ}$), leading to the beam spot size estimated at the reaction target position to be $\sigma_{x,y} < 3 \text{ mm}$.

Figure 2 displays the measured angular differential cross sections for the scattering (as a ratio to Rutherford; squares) and inclusive breakup (in mb/sr; circles) as a function of the scattering angle of ^{11}Be and ^{10}Be in the center-of-mass rest frame θ , respectively. The former cross section includes both elastic and inelastic events, since the energy resolution is not fine enough to exclude the 320 keV excitation energy of the first (and only bound) excited state of ^{11}Be . The uncertainties (1σ) of the measured cross sections are about 7–19%, depending on the angle. These uncertainties include statistical (1–17%), beam intensity ($< 0.1\%$), target thickness (3.5%), and solid angle (5.4%). The measured cross sections do not involve any data acquisition (DAQ) deadtime as both phoswich and Si arrays were run in the same DAQ system, cancelling each other's deadtime. The phoswich's efficiency for heavy ion detection was confirmed to be 100.0 (14)% by comparing Li and light ion punch-through events from the Si 3&4 telescope with those in coincidence with the phoswich. This also ensures the coincidence efficiency between the phoswich and the Si telescopes. The coincidence efficiency for $^{11,10}\text{Be}$ between Si ΔE - E detectors were confirmed to be 100 (1)% from the ratio between the coincidence-singles ratio of the E detec-

tor at the corresponding energy region ($\sim 100 \text{ MeV}$). The uncertainty on the total number of ^{11}Be beam particles was estimated by using different PID cuts in Fig. 1(b). As seen in the inset, nearly 97% of the ^{11}Be events are located in the peak position, which was used to determine the cross sections. This leads to less than 3% underestimation of the total number of beam particles. Because this uncertainty is systematic, it is not accounted for in the aforementioned estimate. However this is negligible for the total uncertainty. Note that all systematic uncertainties cancel when taking the ratio of the breakup and scattering cross sections.

In the angular range of the measurement, and with the resolution of our setup, we observe in Fig. 2 a similar diffraction pattern in scattering and breakup cross sections. This confirms the theoretical observation, upon which the ratio method is based, namely that halo nuclei are scattered similarly whether they remain bound or break up [26]. Taking their ratio therefore should remove the global angular dependence, allowing us to access much more precise structure information about the projectile [16, 20].

To analyze these data in more detail, we use the dynamical eikonal approximation (DEA) [27, 28], which provides excellent results for the elastic scattering and diffractive breakup of one-nucleon halo nuclei on both light and heavy targets. To account for the rather low beam energy of this experiment, we consider the correction analyzed in Ref. [29]. The DEA is built on the usual three-body model of reaction. The projectile P is described as a core c to which a neutron n is loosely bound. The c - n interaction is simulated by an effective potential V_{cn} . Because it reproduces the predictions of the *ab initio*




FIG. 2. Cross sections for scattering (ratio to Rutherford; squares) and inclusive breakup (in mb/sr; circles) for ^{11}Be on C at 22.8 MeV/nucleon expressed as a function of the scattering angle of ^{11}Be and ^{10}Be in the center-of-mass frame, respectively. Black and white symbols correspond to the data from Configs. 1 and 2, respectively. Calculations have been performed with different U_{cT} , see text for details.

calculation of ^{11}Be by Calci *et al.* [30] and leads to excellent agreement with various breakup [31, 32], transfer [33], and knockout [34] data, we use the Halo EFT interaction beyond next-to-leading order in partial waves $s_{1/2}$, $p_{1/2}$, and $d_{5/2}$ with the regulator $\sigma = 1.2$ fm (see Tables I, II and IV of [31]); in all other partial waves $V_{cn} = 0$, in the spirit of Halo EFT [35, 36]. The interaction of the projectile constituents with the target T are described by optical potentials U_{cT} and U_{nT} . Double-folding potentials (DFPs) are used for U_{cT} . They are built from chiral EFT nucleon-nucleon interactions at $N^2\text{LO}$ and their imaginary part is derived through dispersion relations [37]. Cutoffs $R_0 = 1.2$ and 1.6 fm are considered to estimate the sensitivity of our calculations to U_{cT} [38]. For U_{nT} , we adopt the global optical potential of Koning and Delaroche [39].

To obtain the inclusive breakup cross section, we compute the reaction without the imaginary part of U_{nT} . This way, the stripping events, in which the neutron is absorbed by the target, feed the diffractive-breakup channel. Accordingly, the DEA breakup cross section is directly comparable to our inclusive measurement. The scattering angle of the ^{10}Be core after breakup is inferred from that of the $^{10}\text{Be}-n$ center of mass by assuming the neutron to be scattered at forward angles, an approximation supported by the data of Ref. [40].

The results of these calculations are shown in Fig. 2 alongside the data. As expected for a collision of light nuclei, they strongly depend on the choice of optical potential [12]. The DFP with cutoff $R_0 = 1.2$ fm (solid red lines) provides a good agreement with the data for both the scattering and breakup angular distributions. The DFP obtained with $R_0 = 1.6$ fm predicts too high cross sections for both observables.

From these data, we can compute the ratio $\mathcal{R}_{f\text{sum}}$ (see Eq. (A3) of Ref. [20])

$$\mathcal{R}_{f\text{sum}} = \frac{\int d\sigma_{\text{bu}}/dE d\Omega dE}{d\sigma_{\text{scatt}}/d\Omega + \int d\sigma_{\text{bu}}/dE d\Omega dE}, \quad (1)$$

which is the ratio of the breakup angular cross section integrated over all $c-n$ relative energies E , divided by the sum of scattering and total breakup cross sections. In the original idea, the ratio is expressed as a function of the $^{10}\text{Be}-n$ scattering angle, which we reconstruct from the measured angle of ^{10}Be by assuming again that n is scattered at 0° . This ratio is shown in Fig. 3. It agrees very well with the calculations, independently of U_{cT} . This is one of the key advantages of the ratio: being independent of the reaction process, it is nearly insensitive to the choice of optical potential [16, 20]. This prediction is demonstrated for the first time by the present measurement.

To extend our study to Coulomb-dominated reactions, we perform a similar analysis using the data for the collision of ^{11}Be on Pb at 19.1 MeV/nucleon [23]. In Fig. 4 we show the angular cross sections for inclusive breakup (in b/sr; circles), scattering (as a ratio to Rutherford;




FIG. 3. Ratio $\mathcal{R}_{f\text{sum}}$ (1) derived from the data and calculations of Fig. 2 as a function of the scattering angle θ of the $^{10}\text{Be}-n$ center of mass.

squares) and the ratio $\mathcal{R}_{f\text{sum}}$ (divided by 5 for readability; diamonds). The DEA calculations using either DFP are in excellent agreement with the corresponding experimental cross sections (solid red and dashed green lines) [41]. Accordingly, they reproduce very well the ratio inferred from the data. Note that even the slight dependence upon the optical potential seen in each angular distribution is fully removed in the ratio.




FIG. 4. Ratio method for the collision of ^{11}Be on Pb at 19.1 MeV/nucleon. DEA calculations are compared to the data of Ref. [23] for the scattering (ratio to Rutherford) and inclusive breakup (in b/sr). The ratio inferred from the data is perfectly explained by theory, independently of U_{cT} .

In summary, the ratio has been suggested as a new reaction observable to study the structure of exotic nuclei [16, 20]. The present study is the first firm experimental evidence that the ratio method works. As predicted by the REB, we show that it removes the strong angular dependence observed in the differential cross sec-

tions for breakup and scattering. Moreover our analysis confirms that the ratio is nearly independent of the optical-potential choice. Accurate reaction calculations can explain very well these data on both C and Pb. Following these very promising results, we plan to perform an exclusive experiment, in which the valence neutron will be detected in coincidence with the core. Within the exact conditions of the REB—viz. at a higher beam energy, detecting the neutron, and considering low c - n relative energies—we will be able to compare directly the experimental ratio to the form factor of the nucleus wave function. This will provide nuclear-structure information with an accuracy unreachable in the analysis of individual cross sections. Our test of the ratio method offers the possibility to open a new era in the study of exotic quantal structures in nuclear physics as well as in other fields of quantum physics.

We express our thanks to Dr. B. T. Roeder (Texas A&M University) and the technical staff at the Texas A&M University Cyclotron Institute. We thank E. A.

Ricard from the Brookhaven National Laboratory for her constructive comments on this paper. SO acknowledges support from the Office of Nuclear Physics, Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-98CH10886 with Brookhaven Science Associates, LLC. Financial support for this work was provided by the US Department of Energy, award Nos. DE-FG02-93ER40773 and DE-SC0018980, the US National Nuclear Security Administration, award No. DE-NA0003841. This project has received funding from the Deutsche Forschungsgemeinschaft within the Collaborative Research Center SFB 1245 (Projektnummer 279384907) and the PRISMA+ (Precision Physics, Fundamental Interactions and Structure of Matter) Cluster of Excellence. GC acknowledges support from the National Sciences and Engineering Research Council of Canada, award SAPIN-2020-00052. RCJ is supported by the UKRI/STFC research grant ST/V001108/1. FMN is supported by the Department of Energy grant DE-SC0021422.

-
- [1] I. Tanihata, H. Hamagaki, O. Hashimoto, Y. Shida, N. Yoshikawa, K. Sugimoto, O. Yamakawa, T. Kobayashi, and N. Takahashi, *Phys. Rev. Lett.* **55**, 2676 (1985).
- [2] I. Tanihata, H. Hamagaki, O. Hashimoto, S. Nagamiya, Y. Shida, N. Yoshikawa, O. Yamakawa, K. Sugimoto, T. Kobayashi, D. Greiner, N. Takahashi, and Y. Nojiri, *Phys. Lett. B* **160**, 380 (1985).
- [3] P. G. Hansen and B. Jonson, *Europhys. Lett.* **4**, 409 (1987).
- [4] I. Tanihata, *J. Phys. G* **22**, 157 (1996).
- [5] T. Aumann, A. Navin, D. P. Balamuth, D. Bazin, B. Blank, B. A. Brown, J. E. Bush, J. A. Caggiano, B. Davids, T. Glasmacher, V. Guimarães, P. G. Hansen, R. W. Ibbotson, D. Karnes, J. J. Kolata, V. Maddalena, B. Pritychenko, H. Scheit, B. M. Sherrill, and J. A. Tostevin, *Phys. Rev. Lett.* **84**, 35 (2000).
- [6] R. Palit, P. Adrich, T. Aumann, K. Boretzky, B. V. Carlson, D. Cortina, U. Datta Pramanik, T. W. Elze, H. Emling, H. Geissel, M. Hellström, K. L. Jones, J. V. Kratz, R. Kulesa, Y. Leifels, A. Leistenschneider, G. Münzenberg, C. Nociforo, P. Reiter, H. Simon, K. Sümmerer, and W. Walus (LAND/FRS Collaboration), *Phys. Rev. C* **68**, 034318 (2003).
- [7] N. Fukuda, T. Nakamura, N. Aoi, N. Imai, M. Ishihara, T. Kobayashi, H. Iwasaki, T. Kubo, A. Mengoni, M. Notani, H. Otsu, H. Sakurai, S. Shimoura, T. Teranishi, Y. X. Watanabe, and K. Yoneda, *Phys. Rev. C* **70**, 054606 (2004).
- [8] A. Di Pietro, G. Randisi, V. Scuderi, L. Acosta, F. Amorini, M. J. G. Borge, P. Figuera, M. Fisichella, L. M. Fraile, J. Gomez-Camacho, H. Jeppesen, M. Lattuada, I. Martel, M. Milin, A. Musumarra, M. Papa, M. G. Pellegriti, F. Perez-Bernal, R. Raabe, F. Rizzo, D. Santonocito, G. Scalia, O. Tengblad, D. Torresi, A. M. Vidal, D. Voulot, F. Wenander, and M. Zadro, *Phys. Rev. Lett.* **105**, 022701 (2010).
- [9] A. Di Pietro, V. Scuderi, A. M. Moro, L. Acosta, F. Amorini, M. J. G. Borge, P. Figuera, M. Fisichella, L. M. Fraile, J. Gomez-Camacho, H. Jeppesen, M. Lattuada, I. Martel, M. Milin, A. Musumarra, M. Papa, M. G. Pellegriti, F. Perez-Bernal, R. Raabe, G. Randisi, F. Rizzo, G. Scalia, O. Tengblad, D. Torresi, A. M. Vidal, D. Voulot, F. Wenander, and M. Zadro, *Phys. Rev. C* **85**, 054607 (2012).
- [10] K. T. Schmitt, K. L. Jones, A. Bey, S. H. Ahn, D. W. Bardayan, J. C. Blackmon, S. M. Brown, K. Y. Chae, K. A. Chipps, J. A. Cizewski, K. I. Hahn, J. J. Kolata, R. L. Kozub, J. F. Liang, C. Matei, M. Matoš, D. Matyas, B. Moazen, C. Nesaraja, F. M. Nunes, P. D. O'Malley, S. D. Pain, W. A. Peters, S. T. Pittman, A. Roberts, D. Shapira, J. F. Shriner, M. S. Smith, I. Spassova, D. W. Stracener, A. N. Villano, and G. L. Wilson, *Phys. Rev. Lett.* **108**, 192701 (2012).
- [11] K. T. Schmitt, K. L. Jones, S. Ahn, D. W. Bardayan, A. Bey, J. C. Blackmon, S. M. Brown, K. Y. Chae, K. A. Chipps, J. A. Cizewski, K. I. Hahn, J. J. Kolata, R. L. Kozub, J. F. Liang, C. Matei, M. Matos, D. Matyas, B. Moazen, C. D. Nesaraja, F. M. Nunes, P. D. O'Malley, S. D. Pain, W. A. Peters, S. T. Pittman, A. Roberts, D. Shapira, J. F. Shriner, M. S. Smith, I. Spassova, D. W. Stracener, N. J. Upadhyay, A. N. Villano, and G. L. Wilson, *Phys. Rev. C* **88**, 064612 (2013).
- [12] P. Capel, G. Goldstein, and D. Baye, *Phys. Rev. C* **70**, 064605 (2004).
- [13] O. Sürer, F. M. Nunes, M. Plumlee, and S. M. Wild, *Phys. Rev. C* **106**, 024607 (2022).
- [14] C. Hebborn, T. R. Whitehead, A. E. Lovell, and F. M. Nunes, *Phys. Rev. C* **108**, 014601 (2023).
- [15] M. Catacora-Rios, A. E. Lovell, and F. M. Nunes, *Phys. Rev. C* **108**, 024601 (2023).
- [16] P. Capel, R. Johnson, and F. Nunes, *Phys. Lett. B* **705**, 112 (2011).
- [17] P. Capel, R. Johnson, and F. Nunes, *Eur. Phys. J. A*

- 56**, 300 (2020).
- [18] R. C. Johnson, J. S. Al-Khalili, and J. Tostevin, *Phys. Rev. Lett.* **79**, 2771 (1997).
- [19] R. C. Johnson, in *Proc. of the Euro. Conf. in Advances in Nucl. Phys. and Related Areas* (July 1997, Thessaloniki, Greece), edited by D. Brink, M. Grypeos, and S. Massen (Giahoudi-Giapouli Publishing, Thessaloniki, 1999) p. 156.
- [20] P. Capel, R. C. Johnson, and F. M. Nunes, *Phys. Rev. C* **88**, 044602 (2013).
- [21] F. Colomer, P. Capel, F. M. Nunes, and R. C. Johnson, *Phys. Rev. C* **93**, 054621 (2016).
- [22] X. Y. Yun, F. Colomer, D. Y. Pang, and P. Capel, *J. Phys. G* **46**, 105111 (2019).
- [23] F. F. Duan, Y. Y. Yang, J. Lei, K. Wang, Z. Y. Sun, D. Y. Pang, J. S. Wang, X. Liu, S. W. Xu, J. B. Ma, P. Ma, Z. Bai, Q. Hu, Z. H. Gao, X. X. Xu, C. J. Lin, H. M. Jia, N. R. Ma, L. J. Sun, D. X. Wang, G. Yang, S. Y. Jin, Z. Z. Ren, Y. H. Zhang, X. H. Zhou, Z. G. Hu, and H. S. Xu (RIBLL Collaboration), *Phys. Rev. C* **105**, 034602 (2022).
- [24] R. E. Tribble, R. H. Burch, and C. A. Gagliardi, *Nucl. Instr. Meth. A* **285**, 441 (1989).
- [25] S. Ota, G. Christian, B. J. Reed, W. N. Catford, S. Dede, D. T. Doherty, G. Lotay, M. Roosa, A. Saastamoinen, and D. P. Scriven, *Nucl. Instr. Meth. A* **1059**, 168946 (2024).
- [26] P. Capel, M. Hussein, and D. Baye, *Phys. Lett.* **B693**, 448 (2010).
- [27] D. Baye, P. Capel, and G. Goldstein, *Phys. Rev. Lett.* **95**, 082502 (2005).
- [28] G. Goldstein, D. Baye, and P. Capel, *Phys. Rev. C* **73**, 024602 (2006).
- [29] T. Fukui, K. Ogata, and P. Capel, *Phys. Rev. C* **90**, 034617 (2014).
- [30] A. Calci, P. Navrátil, R. Roth, J. Dohet-Eraly, S. Quaglioni, and G. Hupin, *Phys. Rev. Lett.* **117**, 242501 (2016).
- [31] P. Capel, D. R. Phillips, and H.-W. Hammer, *Phys. Rev. C* **98**, 034610 (2018).
- [32] L. Moschini and P. Capel, *Phys. Lett.* **B790**, 367 (2019).
- [33] J. Yang and P. Capel, *Phys. Rev. C* **98**, 054602 (2018).
- [34] C. Hebborn and P. Capel, *Phys. Rev. C* **104**, 024616 (2021).
- [35] C. Bertulani, H.-W. Hammer, and U. van Kolck, *Nucl. Phys.* **A712**, 37 (2002).
- [36] H.-W. Hammer, C. Ji, and D. R. Phillips, *J. Phys. G* **44**, 103002 (2017).
- [37] V. Durant, P. Capel, and A. Schwenk, *Phys. Rev. C* **102**, 014622 (2020).
- [38] V. Durant, P. Capel, L. Huth, A. Balantekin, and A. Schwenk, *Phys. Lett.* **B782**, 668 (2018).
- [39] A. Koning and J. Delaroche, *Nucl. Phys.* **A713**, 231 (2003).
- [40] R. Anne, R. Bimbot, S. Dogny, H. Emling, D. Guillemaud-Mueller, P. Hansen, P. Hornshøj, F. Humbert, B. Jonson, M. Keim, M. Lewitowicz, P. Møller, A. Mueller, R. Neugart, T. Nilsson, G. Nyman, F. Pougheon, K. Riisager, M.-G. Saint-Laurent, G. Schrieder, O. Sorlin, O. Tengblad, and K. Rolander, *Nucl. Phys.* **A575**, 125 (1994).
- [41] V. Durant and P. Capel, “Analysis of ^{11}Be reactions using chiral effective field theory NN interactions,” (2024), in preparation.