

# Curvature-dimension condition, rigidity theorems and entropy differential inequalities on Riemannian manifolds

Xiang-Dong Li \*

**Abstract** In this paper, we use the information-theoretic approach to study curvature-dimension condition, rigidity theorems and entropy differential inequalities on Riemannian manifolds. We prove the equivalence of the  $CD(K, m)$ -condition for  $K \in \mathbb{R}$  and  $m \in [n, \infty]$  and some entropy differential inequalities along the geodesics on the Wasserstein space over a Riemannian manifold. The rigidity models of the entropy differential inequalities are the  $K$ -Einstein manifolds and the  $(K, m)$ -Einstein manifolds with Hessian solitons. Moreover, we prove the monotonicity and rigidity theorem of the  $W$ -entropy along the geodesics on the Wasserstein space over Riemannian manifolds with  $CD(0, m)$ -condition. Finally, we compare our work with the synthetic geometry developed by Lott-Villani and Sturm.

*MSC2010 Classification:* primary 53C44, 58J35, 58J65; secondary 60J60, 60H30.

*Keywords:* curvature-dimension condition, entropy differential inequalities, rigidity theorems,  $W$ -entropy, Lott-Villani-Sturm's synthetic geometry

## 1 Introduction

In recent years, there have been extensive works in the study on curvature-dimension condition on Riemannian manifolds and metric measure spaces [1, 3, 9, 14, 33, 47, 43, 44, 45, 49, 50]. The first link between the  $K$ -geodesic convexity of the relative entropy functional on the  $L^2$ -Wasserstein space  $(P_2(M), W_2)$  of probability measures on a Riemannian manifold  $M$  and the condition on the Ricci curvature bounded below by  $K \in \mathbb{R}$  was given by Sturm and von Renesse [47]. In [33, 44, 45], Lott and Villani, Sturm independently developed a synthetic comparison geometry on metric measure spaces based on a nice definition of curvature-dimension condition, which extends the notion of the curvature-dimension condition introduced by Bakry-Emery [2] on Riemannian manifolds with weighted volume measure. By [47, 33, 44, 50], the  $CD(K, \infty)$ -condition holds on a metric measure space  $(X, d, \mu)$  if and only if the Boltzmann (relative) entropy

$$\text{Ent}(\rho) = \text{Ent}(\rho\mu|\mu) = \int_X \rho \log \rho d\mu$$

is  $K$ -convex along any geodesic  $\{\rho(t)d\mu, t \in [0, 1]\}$  on the Wasserstein space  $P_2(M, d)$  over  $(X, d, \mu)$

$$\text{Ent}(\rho(t)) \leq (1-t)\text{Ent}(\rho(0)) + t\text{Ent}(\rho(1)) - \frac{K}{2}t(1-t)W_2^2(\rho(0), \rho(1)), \quad (1.1)$$

---

\*Research supported by National Key R&D Program of China (No. 2020YFA0712700), NSFC No. 11771430 and Key Laboratory RCSDS, CAS, No. 2008DP173182.

and the  $\text{CD}(0, N)$ -condition holds on a metric measure space  $(X, d, \mu)$  if and only if the quantity<sup>1</sup>

$$S_{N'}(\rho) = S_{N'}(\rho\mu|\mu) = - \int_M \rho^{1-1/N'} d\mu$$

is convex along any geodesic  $\{\rho(t)d\mu, t \in [0, 1]\}$  on the Wasserstein space over  $(X, d, \mu)$  for all  $N' \geq N$ ,

$$S_{N'}(\rho(t)) \leq (1-t)S_{N'}(\rho(0)) + tS_{N'}(\rho(1)). \quad (1.2)$$

Recall that the Wasserstein space  $P_2(X, d)$  over  $(X, d, \mu)$  is the set of probability measures  $\gamma$  on  $(X, d)$  with finite second moment  $\int_X d^2(o, x)d\gamma(x) < +\infty$  for some (hence for all) fixed  $o \in X$ , the Wasserstein distance  $W_2(\gamma_0, \gamma_1)$  between two probability measures  $\gamma_0, \gamma_1$  on  $(X, d)$  is defined by

$$W_2(\gamma_0, \gamma_1) = \inf_{\pi} \left\{ \int_{X \times X} d^2(x, y)d\pi(x, y) \right\}$$

among all the coupling measures  $\pi(\cdot, \cdot)$  on  $X \times X$  such that  $\gamma_0, \gamma_1$  are the marginal measures of  $\pi$ , i.e.,  $\gamma_0 = \pi(\cdot, X)$  and  $\gamma_1 = \pi(X, \cdot)$ , and a geodesic on  $P_2(M, d)$  is a curve  $\gamma : [0, 1] \rightarrow P_2(M, d)$  such that

$$W_2(\gamma(s), \gamma(t)) = |s - t|W_2(\gamma_0, \gamma_1), \quad \forall s, t \in [0, 1].$$

For  $K \neq 0$  and  $N < \infty$ , the definition formula for the  $\text{CD}(K, N)$ -condition on metric measure spaces is more involved. In the literature, it was first introduced by Sturm [45] using the above mentioned quantity  $S_{N'}$  and was further extended by Lott-Villani [33] using more general functional  $U(\rho) = \int_X e(\rho)d\mu$  for continuous convex functions  $e$  in the class  $\mathcal{DC}_N$ . See Section 4.7 for the precise definition of the class  $\mathcal{DC}_N$ .

By Definition 1.3 in [45], given two numbers  $K, N \in \mathbb{R}$  with  $N \in [1, \infty)$ , we say that a metric measure space  $(M, d, \mu)$  satisfies the curvature-dimension condition  $\text{CD}(K, N)$  if and only if for each pair  $\nu_0, \nu_1 \in P_2(M, d)$  there exist an optimal coupling  $q$  of  $\nu_0 = \rho_0\mu$  and  $\nu_1 = \rho_1\mu$ , and a geodesic  $\Gamma : [0, 1] \rightarrow P_2(M, d)$  connecting  $\nu_0$  and  $\nu_1$ , with

$$\begin{aligned} S_{N'}(\Gamma(t)|\mu) \leq & - \int_{M \times M} \left[ \tau_{K, N'}^{(1-t)}(d(x_0, x_1))\rho_0^{-1/N'}(x_0) \right. \\ & \left. + \tau_{K, N'}^{(t)}(d(x_0, x_1))\rho_1^{-1/N'}(x_1) \right] dq(x_0, x_1) \end{aligned} \quad (1.3)$$

for all  $t \in [0, 1]$  and all  $N' \geq N$ . Here,  $\tau_{K, N}^{(t)}(\theta) = t^{1/N}\sigma_{K, N-1}^{(t)}(\theta)^{1-1/N}$ , with

$$\sigma_{K, N}^{(t)}(\theta) = \frac{\sin(t\theta\sqrt{K/N})}{\sin(\theta\sqrt{K/N})}, \quad t \in [0, 1]$$

when  $0 < K\theta^2 < N\pi^2$ , and  $\sigma_{0, N}^{(t)}(\theta) = t$  for  $K = 0$ , and one replaces the function  $\sin$  by the function  $\sinh$  in the definition formula of  $\sigma_{K, N}^{(t)}(\theta)$  when  $K < 0$ . See also [50].

In the  $\text{CD}(K, N)$ -condition on metric measure space,  $K$  is regarded as a lower bound of the Ricci curvature and  $N \in [1, +\infty)$  is an upper bound of the space dimension. Note that the  $\text{CD}(K, \infty)$ -condition can be viewed as the limiting case of the  $\text{CD}(K, N)$ -condition for  $N \rightarrow \infty$ . For the PDE description of the geodesic  $\{\rho(t)d\mu, t \in [0, 1]\}$  on the Wasserstein space over  $(X, d, \mu)$ , see Section 3 below.

On the other hand, the Shannon and Rényi entropies and the corresponding entropy powers have been extensively studied in the information theory. In his 1948 seminal paper

<sup>1</sup> $S_{N'}(\rho)$  is the Rényi entropy power  $N_{m, p}(\rho)$  for  $p = 1 - 1/N'$  and  $m = N'$ . See Section 2.

[41], Shannon introduced the differential entropy for an  $n$ -dimensional random vector  $X$  with probability distribution  $f(x)dx$  on  $\mathbb{R}^n$  by

$$H(X) = H(f) = - \int_{\mathbb{R}^n} f(x) \log f(x) dx,$$

and the related entropy power by

$$N(X) = N(f) = e^{\frac{2}{n}H(X)}.$$

In [41], Shannon discovered the Entropy Power Inequality (EPI): Let  $X$  and  $Y$  be two independent random vectors with probability density functions  $f$  and  $g$  respectively on  $\mathbb{R}^n$ , then

$$N(X + Y) \geq N(X) + N(Y). \quad (1.4)$$

Equivalently

$$N(f * g) \geq N(f) + N(g),$$

where  $f * g$  denote the convolution of  $f$  and  $g$ , i.e., the probability density of the law of  $X + Y$ . See Stam [42] and Blachman [4] for the complete proofs of the Shannon Entropy Power Inequality. Moreover, it is known that the equality holds in (1.4) if and only if  $X$  and  $Y$  are normally distributed with proportional covariance matrices. See [41, 42, 4, 11].

In [10], Costa proved the Entropy Power Concavity Inequality (EPCI) for the heat equation on  $\mathbb{R}^n$ . More precisely, let  $u$  be a positive solution to the heat equation on  $\mathbb{R}^n$

$$\partial_t u = \Delta u. \quad (1.5)$$

Then the Shannon entropy power  $N(u(t))$  is concave in  $t \in (0, \infty)$ , i.e.,

$$\frac{d^2}{dt^2} N(u(t)) \leq 0. \quad (1.6)$$

Using an argument based on the Blachman-Stam inequality [4], the original proof of Costa's inequality (1.6) has been simplified by Dembo et al. [11, 12, 13]. In [48], Villani gave short proof of Costa's inequality (1.6) and pointed out that one can extend it to Riemannian manifolds with non-negative Ricci curvature by the Bakry-Emery  $\Gamma_2$ -calculus.

The concavity property of the Shannon entropy power for the heat equation (1.5) has been extended to the Rényi entropy power by Savaré and Toscani [40]. More precisely, let  $u$  be a positive solution to the nonlinear diffusion equation on  $\mathbb{R}^n$

$$\partial_t u = \Delta u^p, \quad (1.7)$$

where  $p \geq 1 - \frac{2}{n}$ . Let

$$H_p(u) = \frac{1}{1-p} \log \int_M u^p dx$$

be the  $p$ -th Rényi entropy associated with (1.7), and define the Rényi entropy power by

$$N_p(u) = \exp(\sigma H_p(u)), \quad (1.8)$$

where  $\sigma = p - 1 + \frac{2}{n}$ . Then, Savaré and Toscani [40] proved that the Rényi entropy power  $N_p(u(t))$  is concave in  $t \in (0, \infty)$ , i.e.,

$$\frac{d^2}{dt^2} N_p(u(t)) \leq 0. \quad (1.9)$$

Note that when  $p \rightarrow 1$ , we have  $\sigma \rightarrow \frac{2}{n}$ ,  $H_p(u) \rightarrow H(u)$  and  $N_p(u) \rightarrow N(u)$ . In view of this, one can recover Costa's concavity result (1.6) from (1.9).

In [28], Songzi Li and the author of this paper extend the concavity of Shannon entropy power to complete Riemannian manifolds. More precisely, the Shannon entropy power is concave along the heat equation associated with the Witten Laplacian on complete Riemannian manifolds with the Bakry-Emery curvature-dimension  $\text{CD}(0, m)$ -condition. In particular, the Shannon entropy power is concave along the heat equation  $\partial_t u = \Delta u$  on complete Riemannian manifolds with non-negative Ricci curvature. Under the curvature-dimension  $\text{CD}(K, m)$ -condition, it is proved in [28] that the rigidity models of the Shannon entropy power are Einstein or quasi Einstein manifolds with Hessian solitons. Moreover, it is also proved in [28] that the Shannon entropy power is convex along the conjugate heat equation introduced by G. Perelman for Ricci flow and the rigidity models of the Shannon entropy power are the shrinking Ricci solitons. This gives a new understanding of Perelman's mysterious  $W$ -entropy functional for the Ricci flow from the information theoretic point of view.

In [29], Songzi Li and the author of this paper further proved that the Rényi entropy power is concave along the nonlinear diffusion equation  $\partial_t u = Lu^p$  associated with the Witten Laplacian on compact Riemannian manifolds with the Bakry-Emery curvature-dimension  $\text{CD}(0, m)$ -condition. In particular, for  $p \geq 1 - \frac{2}{n}$ , the Rényi entropy power  $N_p(u(t))$  is concave along the nonlinear diffusion equation  $\partial_t u = \Delta u^p$  on compact Riemannian manifolds with non-negative Ricci curvature. Under the curvature-dimension  $\text{CD}(K, m)$ -condition, it is proved that the rigidity models of the Rényi entropy power are Einstein or quasi Einstein manifolds with Hessian solitons.

In the theory of optimal transportation, the heat equation (1.5) can be realized as the gradient flow of the Boltzmann entropy on the  $L^2$ -Wasserstein space  $P_2^\infty(\mathbb{R}^n)$  of probability densities on  $\mathbb{R}^n$  equipped with Otto's infinite dimensional Riemannian metric, and the nonlinear diffusion equation (1.7) is the gradient flow of the Rényi entropy on the Wasserstein space over  $\mathbb{R}^n$ . On the other hand, by the work of McCann [35] and Benamou-Brenier [5], the geodesic flow over the  $L^2$ -Wasserstein space  $P_2^\infty(\mathbb{R}^n)$  equipped with Otto's Riemannian metric is given by the continuity equation (called also the transport equation) together with the Hamilton-Jacobi equation on  $\mathbb{R}^n$ .

The purpose of this paper is use *the information-theoretic approach* to study the curvature-dimension condition, rigidity theorems and entropy differential inequalities on complete Riemannian manifolds. We prove the equivalence of the curvature-dimension condition and the corresponding Shannon and Rényi entropy differential inequalities along the geodesics flow on the Wasserstein space over Riemannian manifolds. We prove the rigidity models of the entropy differential inequalities are the Einstein manifolds or the  $(K, m)$ -Einstein manifolds. Moreover, we prove the monotonicity and rigidity of the  $W$ -entropy along the geodesics on the Wasserstein space over Riemannian manifolds with  $\text{CD}(0, m)$ -condition. Finally, we compare our results with the synthetic geometry developed by Lott-Villani [33], Sturm [44, 45] and Villani [50].

## 2 Notation and main results

Let  $(M, g)$  be a complete Riemannian manifold,  $V \in C^2(M)$  and  $d\mu = e^{-V} dv$ , where  $dv$  is the Riemannian volume measure on  $(M, g)$ . Let  $\nabla_\mu^*$  be the  $L^2(\mu)$ -adjoint of the gradient operator  $\nabla$ , i.e., for any smooth vector field on  $M$ ,

$$\nabla_\mu^* X = -e^V \nabla \cdot (e^{-V} X) = -\nabla \cdot X + \nabla V \cdot X.$$

The Witten Laplacian acting on smooth functions is defined by

$$L = -\nabla_\mu^* \nabla = \Delta - \nabla V \cdot \nabla.$$

For any  $f, \phi \in C^\infty(M)$ , we have

$$\nabla_\mu^*(f\nabla\phi) = -fL\phi - \langle \nabla f, \nabla\phi \rangle.$$

For any  $u, v \in C_0^\infty(M)$ , the integration by parts formula holds

$$\int_M \langle \nabla u, \nabla v \rangle d\mu = - \int_M Luv d\mu = - \int_M uLv d\mu.$$

Thus,  $L$  is the infinitesimal generator of the Dirichlet form

$$\mathcal{E}(u, v) = \int_M \langle \nabla u, \nabla v \rangle d\mu, \quad u, v \in C_0^\infty(M).$$

By Itô's theory, the Stratonovich SDE on  $M$

$$dX_t = \sqrt{2}U_t \circ dW_t - \nabla V(X_t)dt, \quad \nabla_{\circ dX_t} U_t = 0,$$

where  $U_t$  is the stochastic parallel transport along the trajectory of  $X_t$ , with initial data  $X_0 = x$  and  $U_0 = \text{Id}_{T_x M}$ , defines a diffusion process  $X_t$  on  $M$  with infinitesimal generator  $L$ . Moreover, the transition probability density of the  $L$ -diffusion process  $X_t$  with respect to  $\mu$ , i.e., the heat kernel  $p_t(x, y)$  of the Witten Laplacian  $L$ , is the fundamental solution to the heat equation

$$\partial_t u = Lu. \tag{2.10}$$

In [2], Bakry and Emery proved the generalized Bochner formula

$$L|\nabla u|^2 - 2\langle \nabla u, \nabla Lu \rangle = 2|\nabla^2 u|^2 + 2\text{Ric}(L)(\nabla u, \nabla u), \tag{2.11}$$

where  $u \in C_0^\infty(M)$ ,  $\nabla^2 u$  denotes the Hessian of  $u$ ,  $|\nabla^2 u|$  is its Hilbert-Schmidt norm, and

$$\text{Ric}(L) = \text{Ric} + \nabla^2 V$$

is the so-called (infinite dimensional) Bakry-Emery Ricci curvature associated with the Witten Laplacian  $L$ . For  $m \in [n, \infty)$ , the  $m$ -dimensional Bakry-Emery Ricci curvature associated with the Witten Laplacian  $L$  is defined by

$$\text{Ric}_{m,n}(L) = \text{Ric} + \nabla^2 V - \frac{\nabla V \otimes \nabla V}{m-n}.$$

In view of this, we have

$$L|\nabla u|^2 - 2\langle \nabla u, \nabla Lu \rangle \geq \frac{2|Lu|^2}{m} + 2\text{Ric}_{m,n}(L)(\nabla u, \nabla u).$$

Here we make a convention that  $m = n$  if and only if  $V$  is a constant. By definition, we have

$$\text{Ric}(L) = \text{Ric}_{\infty,n}(L).$$

Following [2], we say that  $(M, g, \mu)$  satisfies the curvature-dimension  $\text{CD}(K, m)$ -condition for a constant  $K \in \mathbb{R}$  and  $m \in [n, \infty]$  if and only if

$$\text{Ric}_{m,n}(L) \geq Kg.$$

Note that, when  $m = n$ ,  $V = 0$ , we have  $L = \Delta$  is the usual Laplacian on  $(M, g)$ , and the  $\text{CD}(K, n)$ -condition holds if and only if the Ricci curvature on  $(M, g)$  is bounded from below by  $K$ , i.e.,

$$\text{Ric} \geq Kg.$$

Let  $P_2(M, \mu)$  (resp.  $P_2^\infty(M, \mu)$ ) be the Wasserstein space (reps. the smooth Wasserstein space) of all probability measures  $\rho(x)d\mu(x)$  with density function (resp. with smooth density function)  $\rho$  on  $M$  such that  $\int_M d^2(o, x)\rho(x)d\mu(x) < \infty$ , where  $d(o, \cdot)$  denotes the distance function from a fixed point  $o \in M$ . Similarly to Otto [34], the tangent space  $T_{\rho d\mu}P_2^\infty(M, \mu)$  is identified as follows

$$T_{\rho d\mu}P_2^\infty(M, \mu) = \{s = \nabla_\mu^*(\rho \nabla \phi) : \phi \in C^\infty(M), \int_M |\nabla \phi|^2 \rho d\mu < \infty\},$$

where  $\nabla_\mu^*$  denotes the  $L^2$ -adjoint of the Riemannian gradient  $\nabla$  with respect to the weighted volume measure  $d\mu$  on  $(M, g)$ .

For  $s_i = \nabla_\mu^*(\rho \nabla \phi_i) \in T_{\rho d\mu}P_2^\infty(M, \mu)$ ,  $i = 1, 2$ , Otto [34] introduced the following infinite dimensional Riemannian metric on  $P_2^\infty(M, \mu)$

$$\langle\langle s_1, s_2 \rangle\rangle := \int_M \langle \nabla \phi_1, \nabla \phi_2 \rangle \rho d\mu,$$

provided that

$$\|s_i\|^2 := \int_M |\nabla \phi_i|^2 \rho d\mu < \infty, \quad i = 1, 2.$$

Let  $T_{\rho d\mu}P_2(M, \mu)$  be the completion of  $T_{\rho d\mu}P_2^\infty(M, \mu)$  with Otto's Riemannian metric. Then  $P_2(M, \mu)$  is an infinite dimensional Riemannian manifold.

Similarly to Benamou and Brenier [5], we can prove that, for any given  $\mu_i = \rho_i d\mu \in P_2(M, \mu)$ ,  $i = 0, 1$ , the  $L^2$ -Wasserstein distance between  $\mu_0$  and  $\mu_1$  coincides with the geodesic distance between  $\mu_0$  and  $\mu_1$  on  $P_2(M, \mu)$  equipped with Otto's infinite dimensional Riemannian metric, i.e.,

$$W_2^2(\mu_0, \mu_1) = \inf \left\{ \int_0^1 \int_M |\nabla \phi(x, t)|^2 \rho(x, t) d\mu(x) dt : \partial_t \rho = \nabla_\mu^*(\rho \nabla \phi), \rho(0) = \rho_0, \rho(1) = \rho_1 \right\}.$$

Given  $\mu_0 = \rho(\cdot, 0)\mu$ ,  $\mu_1 = \rho(\cdot, 1)\mu \in P_2^\infty(M, \mu)$ , it is known that there is a unique minimizing Wasserstein geodesic  $\{\mu(t), t \in [0, 1]\}$  of the form  $\mu(t) = (F_t)_* \mu_0$  joining  $\mu_0$  and  $\mu_1$  in  $P_2(M, \mu)$ , where  $F_t \in \text{Diff}(M)$  is given by  $F_t(x) = \exp_x(-t \nabla \phi(\cdot, 0))$  for an appropriate Lipschitz function  $\phi(\cdot, t)$  (see [35]). If the Wasserstein geodesic in  $P_2(M, \mu)$  belongs entirely to  $P_2^\infty(M, \mu)$ , then the geodesic flow  $(\rho, \phi) \in TP_2^\infty(M, \mu)$  satisfies the continuity equation and the Hamilton-Jacobi equation

$$\partial_t \rho - \nabla_\mu^*(\rho \nabla \phi) = 0, \quad (2.12)$$

$$\partial_t \phi + \frac{1}{2} |\nabla \phi|^2 = 0, \quad (2.13)$$

with the boundary condition  $\rho(0) = \rho_0$  and  $\rho(1) = \rho_1$ . When  $\rho_0 \in C^\infty(M, \mathbb{R}^+)$  and  $\phi_0 \in C^\infty(M)$ , defining  $\phi(\cdot, t) \in C^\infty(M)$  by the Hopf-Lax solution

$$\phi(x, t) = \inf_{y \in M} \left( \phi_0(y) + \frac{d^2(x, y)}{2t} \right), \quad (2.14)$$

and solving the continuity equation (2.12) by the characteristic method, it is known that  $(\rho, \phi)$  satisfies (2.12) and (2.13) with  $\rho(0) = \rho_0$  and  $\phi(0) = \phi_0$ . See [49] Sect. 5.4.7. See also [31, 32]. In view of this, the continuity equation (2.12) and the Hamilton-Jacobi equation (2.13) describe the geodesic flow on the tangent bundle  $TP_2^\infty(M, \mu)$  over the Wasserstein space  $P_2(M, \mu)$ . Note that the Hamilton-Jacobi equation (2.13) is also called the eikonal

equation in geometric optics. In particular, when  $m = n$ ,  $V = 0$  and  $\mu = \nu$ , we have  $\nabla_v^* = -\nabla \cdot$ , and the geodesic flow on  $TP_2(M, \nu)$  is given by

$$\partial_t \rho + \nabla \cdot (\rho \nabla \phi) = 0, \quad (2.15)$$

$$\partial_t \phi + \frac{1}{2} |\nabla \phi|^2 = 0. \quad (2.16)$$

When  $M = \mathbb{R}^n$ , let  $\rho_n(x, t) = \frac{e^{-\frac{|x|^2}{4t^2}}}{(4\pi t^2)^{n/2}}$ , and  $\phi_n(x, t) = \frac{|x|^2}{2t}$ . By [27],  $(\rho_n, \phi_n)$  is a special solution to the geodesic equations (2.15) and (2.16) on  $P_2(\mathbb{R}^n, dx)$ .

Note that our definitions of  $P_2(M, \mu)$ ,  $P_2^\infty(M, \mu)$  and the geodesic equations (2.12) and (2.13) rely on the reference measure  $\mu$  on the Riemannian manifold  $(M, g)$ , while the classical definition of the Wasserstein space  $P_2(X, d)$  relied only on the distance function  $d$  on the metric space  $X$ .

Let  $(\rho, \phi)$  be a smooth geodesic flow on  $P_2(M, \mu)$ , i.e., a smooth solution to (2.12) and (2.13). The Shannon entropy associated with the geodesic flow is defined by

$$H(\rho) = - \int_M \rho \log \rho d\mu,$$

and for  $p > 1 - \frac{1}{m}$ , the  $p$ -th Rényi entropy associated with the geodesic flow is defined by

$$H_p(\rho) = \frac{1}{1-p} \log \int_M \rho^p d\mu.$$

The Shannon entropy power  $N_m$  associated with the geodesic flow is defined by

$$N_m(\rho) = \exp \left( \frac{1}{m} H(\rho) \right),$$

and the Rényi entropy power  $N_{m,p}$  associated with the geodesic flow is defined by

$$N_{p,m}(u) = \exp \left( \frac{1}{m} H_p(u) \right).$$

Let  $\theta = W_2(\rho(0)\mu, \rho(1)\mu)$ . Define

$$\sigma_{K,m,\theta}(t) := \frac{\sin(t\theta\sqrt{K/m})}{\sin(\theta\sqrt{K/m})},$$

when  $0 < K\theta^2 < m\pi^2$ ,  $\sigma_{K,m,\theta}(t) = +\infty$  when  $K\theta^2 \geq m\pi^2$ ,  $\sigma_{K,m,\theta}(t) = t$  when  $K = 0$ , and when  $K < 0$

$$\sigma_{K,m,\theta}(t) = \frac{\sinh(t\theta\sqrt{K/m})}{\sinh(\theta\sqrt{K/m})}.$$

Denote

$$N_{m,K}(t) = \sigma_{K,m,\theta}(1-t)N_m(\rho(0)) + \sigma_{K,m,\theta}(t)N_m(\rho(1)). \quad (2.17)$$

Then

$$\frac{d^2 N_{m,K}}{dt^2} = -\frac{K\theta^2}{m} N_{m,K}, \quad (2.18)$$

and

$$N_{m,K}(0) = N_m(\rho(0)), \quad N_{m,K}(1) = N_m(\rho(1)).$$

Now we state the main results of this paper.

**Theorem 2.1** *Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold, and  $K \in \mathbb{R}$ . Let  $(\rho, \phi)$  be a smooth solution to the geodesic flow on  $TP_2(M, v)$  equipped with Otto's infinite dimensional Riemannian metric, i.e.,  $(\rho, \phi)$  satisfies the continuity equation (2.12) and the Hamilton-Jacobi equation (2.13) for  $\mu = v$ . Let  $p \geq 1 - \frac{1}{n}$  and  $d\gamma = \frac{\rho^p dv}{\int_M \rho^p dv}$ . Then the following conditions are equivalent:*

- (i)  $\text{Ric} \geq K$ ,
- (ii) the Shannon entropy differential inequality holds

$$-H'' \geq \frac{1}{n} H'^2 + KW_2^2(\rho(0)v, \rho(1)v),$$

- (iii) the Shannon entropy power differential inequality holds

$$\frac{d^2 N_n}{dt^2} \leq -\frac{KN_n}{n} W_2^2(\rho(0)v, \rho(1)v),$$

- (iv) the Rényi entropy differential inequality holds

$$H_p'' + \frac{1}{n} H_p'^2 \leq -K \int_M |\nabla \phi|^2 d\gamma,$$

- (v) the Rényi entropy power differential inequality holds

$$\frac{d^2}{dt^2} N_{n,p}(\rho(t)) \leq \left( -\frac{K}{n} \int_M |\nabla \phi|^2 d\gamma \right) N_{n,p}(\rho(t))$$

- (vi) the Shannon entropy power  $N_n(\rho(t))$  satisfies

$$N_n(\rho(t)) \geq \sigma_{K,n,\theta}(1-t)N_n(\rho(0)) + \sigma_{K,n,\theta}(t)N_n(\rho(1)),$$

- (vii) For all  $N \geq n$ , it holds

$$\frac{d^2}{dt^2} S_N(\rho) + \frac{N-n}{n} [S_N(\rho)]^{-1} \left( \frac{d}{dt} S_N(\rho) \right)^2 \geq \frac{K}{N} \int_M |\nabla \phi|^2 \rho^{1-1/N} dv.$$

where  $S_N(\rho) = - \int_M \rho^{1-1/N} dv$ . In particular, it holds

$$\frac{d^2}{dt^2} S_N(\rho) \geq \frac{K}{N} \int_M |\nabla \phi|^2 \rho^{1-1/N} dv.$$

- (viii) Sturm's definition inequality (1.3) for  $CD(K, N)$  holds for all  $N' \geq N = n$ .

Moreover, the equality in one of (ii)-(vii) holds for all geodesic on  $P_2(M, v)$  if and only if  $(M, g)$  is an Einstein manifold with constant Ricci curvature, i.e.,

$$\text{Ric} = Kg,$$

and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{I}{n} g,$$

where

$$I = H'(\rho(t)) = \int_M \Delta \phi \rho dv$$



is the Fisher information associated with the geodesic flow on  $TP_2(M, v)$ , and satisfies the Riccati equation

$$-I' = \frac{I^2}{n} + KW_2^2(\rho(0)v, \rho(1)v).$$

More generally, we have the following

**Theorem 2.2** *Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold with a weighted volume measure  $d\mu = e^{-V}dv$ , where  $V \in C^2(M, \mathbb{R})$ . Let  $m \in [n, \infty)$  and  $K \in \mathbb{R}$  be two constants. Let  $(\rho, \phi)$  be a smooth solution to the geodesic flow on  $TP_2(M, \mu)$  equipped with Otto's infinite dimensional Riemannian metric, i.e.,  $(\rho, \phi)$  satisfies the continuity equation (2.12) and the Hamilton-Jacobi equation (2.13). Let  $p \geq 1 - \frac{1}{m}$  and  $d\gamma = \frac{\rho^p d\mu}{\int_M \rho^p d\mu}$ . Then the following conditions are equivalent:*

- (i)  $\text{Ric}_{m,n}(L) \geq K$ , i.e., the  $CD(K, m)$ -condition holds,
- (ii) the Shannon entropy differential inequality holds

$$-H'' \geq \frac{1}{m}H'^2 + KW_2^2(\rho(0)\mu, \rho(1)\mu), \quad (2.19)$$

- (iii) the Shannon entropy power differential inequality holds

$$\frac{d^2 N_m}{dt^2} \leq -\frac{KN_m}{m}W_2^2(\rho(0)\mu, \rho(1)\mu), \quad (2.20)$$

- (iv) the Rényi entropy differential inequality holds

$$H_p'' + \frac{1}{m}H_p'^2 \leq -K \int_M |\nabla\phi|^2 d\gamma, \quad (2.21)$$

- (v) the Rényi entropy power differential inequality holds

$$\frac{d^2}{dt^2} N_{m,p}(\rho(t)) \leq \left( -\frac{K}{m} \int_M |\nabla\phi|^2 d\gamma \right) N_{m,p}(\rho(t)) \quad (2.22)$$

- (vi) the Shannon entropy power  $N_m(\rho(t))$  satisfies

$$N_m(\rho(t)) \geq \sigma_{K,m,\theta}(1-t)N_m(\rho(0)) + \sigma_{K,m,\theta}(t)N_m(\rho(1)), \quad (2.23)$$

- (vii) For all  $N \geq m$ , it holds

$$\frac{d^2}{dt^2} S_N(\rho) + \frac{N-m}{m} [S_N(\rho)]^{-1} \left( \frac{d}{dt} S_N(\rho) \right)^2 \geq \frac{K}{N} \int_M |\nabla\phi|^2 \rho^{1-1/N} d\mu. \quad (2.24)$$

where  $S_N(\rho) = -\int_M \rho^{1-1/N} d\mu$ . In particular, it holds

$$\frac{d^2}{dt^2} S_N(\rho) \geq \frac{K}{N} \int_M |\nabla\phi|^2 \rho^{1-1/N} d\mu. \quad (2.25)$$

- (viii) Sturm's definition inequality (1.3) holds for all  $N' \geq N = m$ .

Moreover, the equality in one of (ii)-(viii) holds for all geodesic on  $P_2(M, \mu)$  if and only if  $(M, g, V)$  is a  $(K, m)$ -Einstein manifold with constant  $m$ -dimensional Bakry-Emery Ricci curvature, i.e.,

$$\text{Ric}_{m,n}(L) = Kg,$$

and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{I}{m} g,$$

where

$$I = H'(\rho(t)) = \int_M L\phi \rho d\mu$$

is the Fisher information associated with the geodesic flow on  $TP_2(M, \mu)$ , and satisfies the Riccati equation

$$-I' = \frac{I^2}{m} + KW_2^2(\rho(0)\mu, \rho(1)\mu).$$

**Theorem 2.3** *Let  $(M, g)$  be a complete Riemannian manifold and  $V \in C^2(M)$  such that  $\text{Ric}(L) = \text{Ric} + \nabla^2 V$  is uniformly bounded on  $M$ . Let  $(\rho(t), \phi(t))$  be a smooth Benamou-Brenier geodesic on  $P_2(M, \mu)$  satisfying the growth condition as required in Theorem 4.1 below. Let  $p \geq 1 - \frac{1}{m}$  and  $d\gamma = \frac{\rho^p d\mu}{\int_M \rho^p d\mu}$ . Then*

$$\begin{aligned} & \frac{d^2}{dt^2} H_p(\rho(t)) + \frac{2}{t} \frac{d}{dt} H_p(\rho(t)) - \frac{m}{t^2} \\ &= -t(p-1) \text{Var}_\gamma(L\phi) - t \int_M \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) d\gamma \\ & \quad - t \int_M \left( \left| \nabla^2 \phi - \frac{g}{t} \right|^2 + \frac{1}{m-n} \int_M \left| \nabla V \cdot \nabla \phi + \frac{m-n}{t} \right|^2 \right) d\gamma, \end{aligned}$$

where

$$\text{Var}_\gamma(L\phi) = \int_M |L\phi - \gamma(L\phi)|^2 d\gamma.$$

Let

$$H_{m,p}(\rho(t)|\rho_m(t)) = H_p(\rho(t)) - H_p(\rho_m(t)),$$

and define the  $W$ -entropy by the Boltzmann formula

$$W_{m,p}(\rho, t) = \frac{d}{dt} (tH_{m,p}(\rho(t)|\rho_m(t))).$$

Then

$$\begin{aligned} \frac{1}{t} \frac{d}{dt} W_{m,p}(\rho, t) &= -(p-1) \text{Var}_\gamma(L\phi) - \int_M \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) d\gamma \\ & \quad - \int_M \left[ \left\| \nabla^2 \phi - \frac{g}{t} \right\|^2 + \frac{1}{m-n} \int_M \left( \nabla V \cdot \nabla \phi + \frac{m-n}{t} \right)^2 \right] d\gamma. \end{aligned}$$

In particular, if  $\text{Ric}_{m,n}(L) \geq 0$ , then  $tH_{m,p}(\rho(t))$  is concave in  $t$  along the geodesic in  $P^\infty(M, \mu)$ , and  $W_{m,p}(\rho(t))$  is non-increasing in  $t$  along the geodesic in  $P^\infty(M, \mu)$ . Moreover, assume that  $\text{Ric}_{m,n}(L) \geq 0$ ,  $p \geq 1 - \frac{1}{m}$ , and  $\frac{d}{dt} W_{m,p}(\rho, t) = 0$  at some  $t = \tau > 0$ . Then  $M$  is isometric to  $\mathbb{R}^n$ ,  $m = n$ ,  $V$  is a constant and  $(\rho, \phi) = (\rho_m, \phi_m)$ .

For the significant role of the  $W$ -entropy in the proof of the non-local collapsing theorem of Hamilton's Ricci flow [18, 19] for the Poincaré conjecture and Thurston's geometrization conjecture, see Perelman [39]. See also [6, 20, 36, 7, 8]. See [37, 38] for the  $W$ -entropy formula for the heat equation  $\partial_t u = \Delta u$  on Riemannian manifolds with non-negative Ricci curvature. See [30, 23, 24, 25, 27] for the study of monotonicity and rigidity theorems of the  $W$ -entropy for the heat equation of the Witten Laplacian  $\partial_t u = Lu$  on Riemannian manifolds with  $CD(K, m)$ -condition or  $(K, m)$ -super Ricci flows and its extension to the Langevin deformation on the Wasserstein space over Riemannian manifolds with  $CD(0, m)$ -condition.

**Corollary 2.4** *The  $CD(0, m)$ -condition holds if and only if the Shannon entropy power  $N_m$  is displacement concave along any geodesic on  $P_2(M, \mu)$  in the sense that*

$$N_m(\rho(t)) \geq (1-t)N_m(\rho(0)) + tN_m(\rho(1)). \quad (2.26)$$

Moreover, if the equality in (2.26) holds for all geodesic on  $P_2(M, \mu)$ , then  $M$  is  $(0, m)$ -Einstein, i.e.,

$$Ric_{m,n}(L) = 0.$$

**Corollary 2.5** *The  $CD(K, \infty)$ -condition holds if and only if for any geodesic on  $P_2(M, \mu)$  we have*

$$\frac{d^2}{dt^2} \text{Ent}(\rho(t)) \geq K \int_M |\nabla \phi|^2 \rho d\mu = KW_2^2(\rho(0), \rho(1)). \quad (2.27)$$

Equivalently, the formal Hessian of the Boltzmann entropy functional  $\text{Ent}$  on the Wasserstein space  $P_2(M, \mu)$  is bounded below by  $K$ , i.e.,

$$\text{Hess Ent} \geq K,$$

or the Boltzmann entropy  $\text{Ent}$  is  $K$ -displacement convex along any geodesics on the Wasserstein space  $P_2(M, \mu)$  in the sense that

$$\text{Ent}(\rho(t)) \leq (1-t)\text{Ent}(\rho(0)) + t\text{Ent}(\rho(1)) - \frac{K}{2}t(1-t)W_2^2(\rho(0), \rho(1)). \quad (2.28)$$

Moreover, the equality in (2.27) holds for any geodesic on  $P_2(M, \mu)$  if and only if  $(M, g, V)$  is a  $K$ -Ricci soliton in the sense that  $Ric(L) = Kg$ , i.e.,

$$Ric + \nabla^2 V = Kg.$$

In particular, when  $\mu = v$ , then the Boltzmann entropy  $\text{Ent}$  is  $K$ -displacement convex along any geodesics on the Wasserstein space  $P_2(M, v)$  over complete Riemannian manifolds with  $Ric \geq Kg$ , and the equality in (2.27) or (2.28) holds along any geodesics on the Wasserstein space  $P_2(M, v)$  if and only if  $(M, g)$  is a  $K$ -Einstein manifold with

$$Ric = Kg.$$

Note that, in the work of Sturm and von Renesse [47], Lott-Villani [33] and Sturm [44, 45], the  $K$ -displacement convexity of the Boltzmann entropy  $\text{Ent}$  along Wasserstein geodesics (i.e., (1.1)) is used to characterize the  $CD(K, \infty)$ -condition on Riemannian manifolds  $(M, g, \mu)$  and metric measure spaces  $(X, d, \mu)$ , and the inequality (1.3) for the Rényi entropy along geodesics on the Wasserstein space over metric measure space  $(X, d, \mu)$  is used to introduce the definition of the  $CD(K, N)$ -condition on  $(X, d, \mu)$ . When restricting on Riemannian manifolds with weighted volume measures (i.e., in the setting of smooth

metric measure spaces), we will make a comparison between Lott-Villani and Sturm's definitions of their curvature-dimension  $CD(K, N)$  conditions ( $K \in \mathbb{R}$  and  $N \in [n, \infty]$ ) with our results on the concavity of the Shannon and Rényi entropy powers along the geodesic on the Wasserstein space over Riemannian manifolds with the Bakry-Emery  $CD(K, m)$ -condition ( $K \in \mathbb{R}$  and  $m \in [n, \infty]$ ) for  $N = m$ .

In summary, the novelties of our paper are:

(1) We give some new characterizations of  $CD(K, m)$ -condition which are equivalent to the ones of Sturm [45] and Lott-Villani [33] on Riemannian manifolds. Our proof uses explicit calculations and is different from [33, 45, 50]. Our work provides an *information-theoretic approach* to understand the synthetic geometry developed by Lott-Villani and Sturm [33, 50, 45].

(2) We prove the rigidity theorems for the entropy differential inequalities on Riemannian manifolds with  $CD(K, m)$ -condition, which are new in the literature.

(3) We introduce the Perelman type  $W$ -entropy associated with the Rényi entropy for the Benamou-Brenier geodesic on the Wasserstein space over a Riemannian manifold and prove its monotonicity and rigidity theorem on the Wasserstein space over complete Riemannian manifolds with  $CD(0, m)$ -condition. This is also new.

The rest of this paper is organized as follows. In Section 3, we prove the differential inequalities and rigidity theorem for the Shannon entropy and its power along geodesics on the Wasserstein space over Riemannian manifolds. In Section 4, we prove the differential inequalities and rigidity theorem for the Rényi entropy and the associated  $W$ -entropy along the Benamou-Brenier geodesics on the Wasserstein space over Riemannian manifolds. In Section 5, we compare our work with the synthetic geometry developed by Lott-Villani and Sturm and raise some problems for further study in the future.

### 3 Differential inequalities for Shannon entropy

In this section we prove the differential inequalities for the Shannon entropy and its power along the geodesic flow on the Wasserstein space over complete Riemannian manifolds with the  $CD(K, m)$  curvature-dimension condition. We prove the NIW formula which indicates the relationship between the Shannon entropy power  $N$ , the Fisher information  $I$  and the  $W$ -entropy associated with the geodesic flow on the Wasserstein space over compact Riemannian manifolds. Moreover, we prove that the rigidity models of the Shannon entropy power  $N$  are the so-called  $(K, m)$ -Einstein manifolds, i.e.,  $Ric_{m,n}(L) = Kg$ , coupled with a Hessian soliton. In particular, when  $m = n$ , the rigidity models of the Shannon entropy power  $N$  are the Einstein manifolds, i.e.,  $Ric = Kg$ , coupled with a Hessian soliton.

#### 3.1 Differential inequalities for Shannon entropy

Recall the following result which was proved in [27] on complete Riemannian manifolds. In compact case, it is well-known, see e.g. [32].

**Theorem 3.1** ([32, 27]) *Let  $M$  be a complete Riemannian manifold, and  $V \in C^2(M)$ . Suppose that  $Ric(L) = Ric + \nabla^2 V$  is uniformly bounded on  $M$ , i.e., there exists a constant  $C > 0$  such that  $|Ric(L)| \leq C$ . Let  $(\rho, \phi)$  be smooth solutions to the Benamou-Brenier geodesic which satisfies the following growth conditions*

$$\int_M [|\nabla \log \rho|^2 + |\nabla \phi|^2 + |\nabla^2 \phi|^2 + |L\phi|^2 + |\nabla L\phi|^2] \rho d\mu < \infty,$$

and for a fixed point  $o \in M$ , some functions  $C_i(t) \geq 0$  and  $\alpha_i(t) \geq 0$  on  $[0, T]$ ,

$$C_1(t)e^{-\alpha_1(t)d^2(x,o)} \leq \rho(x, t) \leq C_2(t)e^{\alpha_2(t)d^2(x,o)}, \quad \forall x \in M, t \in [0, T],$$

and

$$\int_M d^4(x, o)\rho(x, t)d\mu < \infty, \quad \forall t \in [0, T].$$

Then the entropy dissipation formulae hold

$$\frac{d}{dt}\text{Ent}(\rho(t)) = \int_M \nabla\phi \cdot \nabla\rho d\mu = - \int_M L\phi\rho d\mu, \quad (3.29)$$

$$\frac{d^2}{dt^2}\text{Ent}(\rho(t)) = \int_M \Gamma_2(\phi, \phi)\rho d\mu, \quad (3.30)$$

where

$$\Gamma_2(\phi, \phi) := \frac{1}{2}L|\nabla\phi|^2 - \langle \nabla\phi, \nabla L\phi \rangle.$$

Note that, by the generalized Bochner formula [2], we have

$$\Gamma_2(\phi, \phi) = \|\nabla^2\phi\|_{\text{HS}}^2 + \text{Ric}(L)(\nabla\phi, \nabla\phi).$$

**Theorem 3.2** *Under the assumption of Theorem 3.1, suppose that  $\text{Ric}_{m,n}(L) \geq Kg$ , where  $m \in [n, \infty)$  and  $K \in C(M, \mathbb{R})$ . Then the following Shannon entropy differential inequality holds*

$$-H'' \geq \frac{1}{m}H'^2 + \int_M K|\nabla\phi|^2\rho d\mu + \frac{1}{m} \int_M |L\phi - I|^2\rho d\mu, \quad (3.31)$$

and equivalently the Shannon entropy power differential inequality holds

$$\frac{d^2 N_m}{dt^2} \leq -\frac{N_m}{m} \int_M K|\nabla\phi|^2\rho d\mu - \frac{N_m}{m^2} \int_M |L\phi - I|^2\rho d\mu, \quad (3.32)$$

where

$$I = H'(\rho(t)) = \int_M L\phi\rho d\mu$$

is the Fisher information associated with the geodesic flow on  $TP_2(M, \mu)$ . As a consequence, if  $\text{Ric}_{m,n}(L) \geq Kg$ , where  $K \in C(M, \mathbb{R})$ , then the Shannon entropy differential inequality holds

$$H'' + \frac{1}{m}H'^2 + \int_M K|\nabla\phi|^2\rho d\mu \leq 0. \quad (3.33)$$

In particular, when  $K$  is a constant, we obtain (2.19) and (2.20).

*Proof.* Note that, for any  $m \geq n$ , we have

$$\Gamma_2(\phi, \phi) \geq \frac{|L\phi|^2}{m} + \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi).$$

Thus

$$-\partial_t^2 H(\rho(t)) \geq \int_M \left[ \frac{|L\phi|^2}{m} + \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) \right] \rho d\mu.$$

Since  $\rho d\mu$  is a probability measure, the Cauchy-Schwarz inequality yields

$$\int_M |L\phi|^2\rho d\mu \geq \left( \int_M L\phi\rho d\mu \right)^2 = (\partial_t H(\rho(t)))^2,$$

and hence

$$-\partial_t^2 H(\rho(t)) \geq \frac{(\partial_t H(\rho(t)))^2}{m} + \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi)\rho d\mu.$$

Thus, if  $Ric_{m,n}(L) \geq K$  for some function  $K \in C(M, \mathbb{R})$ , we have

$$-\partial_t^2 H(\rho(t)) \geq \frac{(\partial_t H(\rho(t)))^2}{m} + \int_M K|\nabla\phi|^2\rho d\mu.$$

This finishes the proof of (3.33). In particular, when  $K$  is a constant, we obtain (2.19).

Note that, along the geodesic flow  $(\rho(t), \phi(t))$  on  $T\mathcal{P}_2(M, \mu)$ , we have

$$\int_M |\nabla\phi(t)|^2\rho(t)d\mu = W_2^2(\rho(0), \rho(1)), \quad \forall t > 0.$$

This proves the equivalent form of the entropy differential inequality.

On the other hand, the Shannon entropy power satisfies

$$\begin{aligned} \frac{d}{dt}N_m(\rho(t)) &= N_m(\rho(t))\frac{\partial_t H(\rho(t))}{m}, \\ \frac{d^2}{dt^2}N_m(\rho(t)) &= N_m(\rho(t))\left[\frac{\partial_t^2 H(\rho(t))}{m} + \left(\frac{\partial_t H(\rho(t))}{m}\right)^2\right]. \end{aligned}$$

Thus the EPDI (2.20) follows from EDI (2.19). □

Let

$$\Theta_K(\rho(t)) = \int_M K|\nabla\phi(t)|^2\rho(t)d\mu.$$

Then we have the Riccati differential inequality along  $(\rho(t), \phi(t))$

$$H'' + \frac{1}{m}H'^2 + \Theta_K(\rho(t)) \leq 0. \quad (3.34)$$

The boundary condition for  $H'(\rho(t))$  is given by

$$H'(0+) = \lim_{t \rightarrow 0+} H'(\rho(t)).$$

Note that (3.34) is equivalent to

$$\frac{d^2 N_m}{dt^2} \leq -\frac{\Theta_K(\rho(t))}{m} N_m.$$

Let  $H'_{m,K}(t)$  be the unique solution to the Riccati equation

$$H''_{m,K} + \frac{1}{m}H'^2_{m,K} + \Theta_K(\rho(t)) = 0 \quad (3.35)$$

with the initial value  $H'_{m,K}(0) \geq H'(0+)$ , and let  $N_{m,K}$  be the unique solution to the second order differential equation

$$\frac{d^2 N_{m,K}}{dt^2} = -\frac{\Theta_K(\rho(t))}{m} N_{m,K} \quad (3.36)$$

with two-point boundary condition

$$N_{m,K}(0) = N_m(\rho(0)), \quad N_{m,K}(1) = N_m(\rho(1)).$$

By the Sturm-Liouville comparison theorem and the maximum principle for Riccati equation, we have

**Theorem 3.3** *Under the same condition as in Theorem 3.2, we have*

$$H'(\rho(t)) \leq H'_{m,K}(t), \quad (3.37)$$

and

$$N_m(\rho(t)) \geq N_{m,K}(t). \quad (3.38)$$

In the case  $K \in \mathbb{R}$  is a constant, we have  $\Theta_K(\rho(t)) = K\theta := KW_2^2(\rho(0)\mu, \rho(1)\mu)$ , and (3.34) and (3.35) read as follows

$$H''_{m,K} + \frac{1}{m}H_{m,K}^2 + K\theta^2 = 0, \quad (3.39)$$

and

$$\frac{d^2 N_{m,K}}{dt^2} = -\frac{K\theta^2}{m}N_{m,K}. \quad (3.40)$$

Hence  $N_{m,L}(t)$  is given by (2.23), i.e.,

$$N_{m,K}(t) = \sigma_{K,m,\theta}(1-t)N_m(\rho(0)) + \sigma_{K,m,\theta}(t)N_m(\rho(1)).$$

*Proof.* Note that (3.34) and (3.35) have the constant term  $\Theta_K(\rho(t)) = \int_M K|\nabla\phi(t)|^2\rho(t)d\mu$  which depends on  $(\rho(t), \phi(t))$ . Let  $u = H'(t) - H'_{m,K}(t)$ . We have

$$u' \leq -\frac{H' + H'_{m,K}}{m}u.$$

By the Gronwall inequality, we have

$$u(t) \leq u(0) \exp\left(-\frac{H(t) + H_{m,K}(t) - H(0) - H_{m,K}(0)}{m}\right).$$

In particular, as  $u(0) = H'(0) - H'_{m,K}(0) \leq 0$ , we have

$$u(t) \leq 0.$$

Integrating along geodesic from  $\rho(0)$  to  $\rho(1)$ , the Shannon entropy power inequality (2.20) implies that

$$N_m(\rho(t)) \geq N_{m,K}(t). \quad (3.41)$$

From which we derive (2.23).  $\square$

## 3.2 The $W$ -entropy and NIW formula for Shannon entropy

Recall that, in [25, 27], S. Li and the author of this paper introduced the  $W$ -entropy and proved the following  $W$ -entropy formula for the Shannon entropy along the geodesic flow on the Wasserstein space  $P_2(M, \mu)$ .

**Theorem 3.4** [25, 27] *Let  $(M, g)$  be a complete Riemannian manifold,  $V \in C^2(M)$  such that  $\text{Ric}(L) = \text{Ric} + \nabla^2 V$  is uniformly bounded. Let  $(\rho, \phi) : M \times [0, T] \rightarrow \mathbb{R}^+ \times \mathbb{R}$  be a smooth solution to the Benamou-Brenier geodesic on  $P_2(M, \mu)$  satisfying the growth condition as required in Theorem 3.1. For any  $m \geq n$ , define the  $H_m$ -entropy and  $W_m$ -entropy for the geodesic flow  $(\rho, \phi)$  on  $TP_2^\infty(M, \mu)$  as follows*

$$H_m(\rho, t) = H(\rho(t)) - \frac{m}{2}(1 + \log(4\pi t^2)),$$

and

$$W_m(\rho, t) = \frac{d}{dt}(tH_m(\rho, t)).$$

Then for all  $t > 0$ , we have

$$\begin{aligned} \frac{1}{t} \frac{d}{dt} W_m(\rho, t) &= - \int_M \left[ \left| \nabla^2 \phi - \frac{g}{t} \right|^2 + Ric_{m,n}(L)(\nabla \phi, \nabla \phi) \right] \rho d\mu \\ &\quad - \frac{1}{m-n} \int_M \left| \nabla V \cdot \nabla \phi - \frac{m-n}{t} \right|^2 \rho d\mu. \end{aligned} \quad (3.42)$$

In particular, if  $Ric_{m,n}(L) \geq 0$ , then  $W_m(\rho, t)$  is nonincreasing in time  $t$  along the geodesic flow on  $TP_2^\infty(M, \mu)$ .

Moreover, S. Li and the author [25, 27] proved the following rigidity theorem of the  $W$ -entropy for the Shannon entropy along the Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold with  $CD(0, m)$ -condition.

**Theorem 3.5** [25, 27] *Let  $(M, g, V)$  be a complete Riemannian manifold with  $Ric(L) = Ric + \nabla^2 V$  uniformly bounded, and let  $(\rho(t), \phi(t))$  be a smooth Benamou-Brenier geodesic on  $P_2(M, \mu)$  satisfying the growth condition as required in Theorem 3.1. Suppose that  $Ric_{m,n}(L) \geq 0$  and  $\frac{d}{dt} W_m(\rho, t) = 0$  at some  $t = \tau > 0$ . Then  $M$  is isometric to  $\mathbb{R}^n$ ,  $m = n$ ,  $V$  is a constant and  $(\rho, \phi) = (\rho_n, \phi_n)$ .*

As a consequence of the  $W$ -entropy formula, we can derive the following  $W$ -entropy inequality: On complete Riemannian manifold with bounded geometry condition, if  $Ric_{m,n}(L) \geq K$ , then we have

$$\frac{1}{t} \frac{d}{dt} W_m(\rho, t) \leq - \int_M K |\nabla \phi|^2 \rho d\mu - \frac{1}{m} \int_M \left| \Delta \phi - \frac{m}{t} \right|^2 \rho d\mu.$$

In particular, for  $L = \Delta$ ,  $m = n$  and  $\mu = v$ , we obtain: on any complete Riemannian manifold with bounded geometry condition, if  $Ric \geq K$ , we have

$$\frac{1}{t} \frac{d}{dt} W_n(\rho, t) \leq - \int_M K |\nabla \phi|^2 \rho dv - \frac{1}{n} \int_M \left| \Delta \phi - \frac{n}{t} \right|^2 \rho dv.$$

In this section, we prove the following NIW formula, which indicates an interesting relationship between the Shannon entropy power  $N$ , the Fisher information  $I$  and the  $W$ -entropy associated with the geodesic flow on the Wasserstein space over (weighted) complete Riemannian manifolds. More precisely, we have the following

**Theorem 3.6** *Let  $M$  be an  $n$ -dimensional complete Riemannian manifold with  $V \in C^2(M)$ . Define*

$$H(\rho(t)) = - \int_M \rho \log \rho d\mu, \quad N_m(\rho(t)) = e^{\frac{H(\rho(t))}{m}}.$$

Then

$$\begin{aligned} \frac{m}{N_m} \frac{d^2 N_m}{dt^2} &= - \frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \int_M Ric_{m,n}(L)(\nabla \phi, \nabla \phi) \rho d\mu \\ &\quad - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 \rho d\mu - \int_M \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|_{\text{HS}}^2 \rho d\mu. \end{aligned} \quad (3.43)$$



Moreover, the following NIW formula holds

$$\frac{d^2 N_m}{dt^2} = \frac{N_m}{m} \left[ \frac{1}{m} \left| I - \frac{m}{t} \right|^2 + \frac{1}{t} \frac{dW_m}{dt} \right].$$

where  $I = H'(\rho(t)) = \int_M L\phi(t)\rho(t)d\mu$  is the Fisher information. In particular, if  $\text{Ric}_{m,n}(L) \geq K$ , we have

$$\frac{1}{t} \frac{dW_m}{dt} \leq - \int_M K |\nabla\phi|^2 \rho d\mu - \frac{1}{m} \left| I - \frac{m}{t} \right|^2. \quad (3.44)$$

*Proof.* Using  $\|A\|_{\text{HS}}^2 = \frac{|\text{Tr}A|^2}{n} + \|A - \frac{\text{Tr}A}{n}g\|_{\text{HS}}^2$ , we have

$$\|\nabla^2\phi\|_{\text{HS}}^2 = \frac{|\Delta\phi|^2}{n} + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2.$$

Applying the elementary equality

$$(a+b)^2 = \frac{a^2}{1+\varepsilon} - \frac{b^2}{\varepsilon} + \frac{\varepsilon}{1+\varepsilon} \left( a + \frac{1+\varepsilon}{\varepsilon}b \right)^2$$

to  $a = L\phi$ ,  $b = \nabla V \cdot \nabla\phi$  and  $\varepsilon = \frac{m-n}{n}$ , we have

$$|\Delta\phi|^2 = \frac{n}{m} |L\phi|^2 - \frac{n}{m-n} |\nabla V \cdot \nabla\phi|^2 + \frac{m-n}{m} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2,$$

and

$$\begin{aligned} \|\nabla^2\phi\|_{\text{HS}}^2 &= \frac{|L\phi|^2}{m} - \frac{|\nabla V \cdot \nabla\phi|^2}{m-n} + \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \\ &\quad + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2. \end{aligned}$$

This yields

$$\begin{aligned} \Gamma_2(\phi, \phi) &= \frac{|L\phi|^2}{m} + \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) \\ &\quad + \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2. \end{aligned}$$

Substituting this into (3.30), we have

$$\begin{aligned} \partial_t H(\rho(t)) &= \int_M L\phi \rho d\mu, \\ \partial_t^2 H(\rho(t)) &= - \int_M \left[ \frac{|L\phi|^2}{m} + \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) \right] \rho d\mu \\ &\quad - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \rho d\mu - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \rho d\mu. \end{aligned}$$

Combining this with (3.29), we have

$$\begin{aligned}
& \partial_t^2 H(\rho(t)) + \frac{(\partial_t H(\rho(t)))^2}{m} \\
&= -\frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \rho d\mu \\
&\quad - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \rho d\mu - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \rho d\mu.
\end{aligned}$$

Combining this with (3.34) and (3.34), we have

$$\begin{aligned}
\frac{m}{N_m} \frac{d^2 N_m}{dt^2} &= -\frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \rho d\mu \\
&\quad - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \rho d\mu - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \rho d\mu.
\end{aligned} \tag{3.45}$$

By (3.45) and (3.42), we have

$$\begin{aligned}
& \frac{m}{N_m} \frac{d^2 N_m}{dt^2} - \frac{1}{t} \frac{dW_m}{dt} \\
&= -\frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \rho d\mu \\
&\quad - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \rho d\mu + \int_M \left| \nabla^2\phi - \frac{g}{t} \right|^2 \rho d\mu + \frac{1}{m-n} \int_M \left| \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right|^2 \rho d\mu.
\end{aligned}$$

Note that

$$\begin{aligned}
\left\| \nabla^2\phi - \frac{g}{t} \right\|_{\text{HS}}^2 &= \frac{1}{n} \left| \Delta\phi - \frac{n}{t} \right|^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \\
&= \frac{1}{n} \left| L\phi - \frac{m}{t} + \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right) \right|^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \\
&= \frac{1}{m} \left| L\phi - \frac{m}{t} \right|^2 - \frac{1}{m-n} \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right)^2 \\
&\quad + \frac{m-n}{mn} \left[ L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right]^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2.
\end{aligned}$$

Combining this with the fact  $I = \partial_t H(\rho(t)) = \int_M L\phi \rho d\mu$ , we have

$$\frac{m}{N_m} \frac{d^2 N_m}{dt^2} = \frac{1}{t} \frac{dW_m}{dt} + \frac{1}{m} \left| I - \frac{m}{t} \right|^2.$$

This proves the NIW formula. The  $W$ -inequality (3.44) follows from (3.35).  $\square$

In particular, when  $m = n$ ,  $V = 0$  and  $L = \Delta$ , we have the following

**Theorem 3.7** *Let  $M$  be a complete Riemannian manifold. Then*

$$\begin{aligned}
\frac{n}{N_n} \frac{d^2 N_n}{dt^2} &= -\frac{1}{n} \int_M \left| \Delta\phi - \int_M \Delta\phi \rho dv \right|^2 \rho dv - \int_M Ric(\nabla\phi, \nabla\phi) \rho dv \\
&\quad - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \rho dv.
\end{aligned} \tag{3.46}$$

Moreover, the following NIW formula holds

$$\frac{d^2 N_n}{dt^2} = \frac{N_n}{n} \left[ \frac{1}{n} \left| I - \frac{n}{t} \right|^2 + \frac{1}{t} \frac{dW_n}{dt} \right]. \quad (3.47)$$

In particular, if  $\text{Ric} \geq K$ , we have

$$\frac{1}{t} \frac{dW_n}{dt} \leq - \int_M K |\nabla \phi|^2 \rho dv - \frac{1}{n} \left| I - \frac{n}{t} \right|^2. \quad (3.48)$$

**Remark 3.8** Using again  $\|A\|_{\text{HS}}^2 = \frac{|\text{Tr}A|^2}{n} + \|A - \frac{\text{Tr}A}{n}g\|_{\text{HS}}^2$ , we have

$$\begin{aligned} \left\| \nabla^2 \phi - \frac{I}{n}g \right\|^2 &= \frac{1}{n} |\Delta \phi - I|^2 + \left\| \nabla^2 \phi - \frac{I}{n}g - \frac{1}{n} (\Delta \phi - I)g \right\|^2 \\ &= \frac{1}{n} |\Delta \phi - I|^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n}g \right\|^2. \end{aligned}$$

This yields another explicit expression of the second derivative of the Shannon entropy power along the geodesic flow on  $TP_2(M, v)$ , which is indeed equivalent to (3.46). More precisely, we have

$$\frac{d^2 N_n}{dt^2} = -\frac{N_n}{n} \int_M \left( \text{Ric}(\nabla \phi, \nabla \phi) + \left\| \nabla^2 \phi - \frac{I}{n}g \right\|_{\text{HS}}^2 \right) \rho dv.$$

Similarly, for  $m > n$ ,  $I = \int_M L\phi \rho d\mu$ , using

$$\begin{aligned} \left\| \nabla^2 \phi - \frac{I}{m}g \right\|^2 &= \frac{1}{n} \left| \Delta \phi - \frac{n}{m}I \right|^2 + \left\| \nabla^2 \phi - \frac{I}{m}g - \frac{1}{n} \left( \Delta \phi - \frac{n}{m}I \right)g \right\|^2 \\ &= \frac{1}{n} \left| \Delta \phi - \frac{n}{m}I \right|^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n}g \right\|^2, \end{aligned}$$

and

$$\begin{aligned} \frac{1}{n} \left| \Delta \phi - \frac{n}{m}I \right|^2 &= \frac{1}{m} |L\phi - I|^2 - \frac{1}{m-n} \left| \nabla V \cdot \nabla \phi + \frac{m-n}{m} \right|^2 \\ &\quad + \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2, \end{aligned}$$

we have

$$\begin{aligned} \frac{m}{N_m} \frac{d^2 N_m}{dt^2} &= - \int_M \left( \text{Ric}_{m,n}(L)(\nabla \phi, \nabla \phi) + \left\| \nabla^2 \phi - \frac{I}{m}g \right\|_{\text{HS}}^2 \right) \rho d\mu \\ &\quad - \frac{1}{(m-n)} \int_M \left( \nabla V \cdot \nabla \phi + \frac{m-n}{m} \right)^2 \rho d\mu. \end{aligned}$$

### 3.3 Rigidity of Shannon entropy power

In this subsection, we prove the rigidity part of Theorem 2.1 and Theorem 2.2.

**Theorem 3.9** Under the same condition and notation as in Theorem 2.1, we have

$$\frac{d^2 N_n}{dt^2} \leq -\frac{KN_n}{n} \int_M |\nabla \phi|^2 \rho dv - \frac{N_n}{n^2} \int_M |\Delta \phi - I|^2 \rho dv. \quad (3.49)$$

This indeed improves the entropy power concavity inequality in Theorem 2.1, i.e.,

$$\frac{d^2 N_n}{dt^2} \leq -\frac{KN_n}{n} \int_M |\nabla\phi|^2 \rho dv. \quad (3.50)$$

Moreover, the equality in (3.50) holds if and only if  $(M, g)$  is an Einstein manifold with constant Ricci curvature  $Ric = Kg$  with a Hessian soliton

$$\nabla^2\phi = \frac{I}{n}g.$$

where  $I = \int_M \Delta\phi \rho dv$  is the Fisher information associated with the geodesic flow on  $TP_2(M, v)$ , and satisfies the Riccati differential equation

$$-I' = \frac{I^2}{n} + KW_2^2(\rho(0), \rho(1)). \quad (3.51)$$

*Proof.* By (3.46) in Theorem 3.7 or (3.49), under the condition  $Ric \geq Kg$ , we have

$$\frac{d^2 N_n}{dt^2} \leq -\frac{KN_n}{n} \int_M |\nabla\phi|^2 \rho dv - \frac{N_n}{n} \int_M \left\| \nabla^2\phi - \frac{I}{n}g \right\|_{\text{HS}}^2 \rho dv. \quad (3.52)$$

Using the trace inequality  $\left\| \nabla^2\phi - \frac{I}{n}g \right\|_{\text{HS}}^2 \geq \frac{1}{n} |\Delta\phi - I|^2$ , we have

$$\frac{d^2 N_n}{dt^2} \leq -\frac{KN_n}{n} \int_M |\nabla\phi|^2 \rho dv - \frac{N_n}{n^2} \int_M |\Delta\phi - I|^2 \rho dv. \quad (3.53)$$

This improves the entropy power concavity inequality in Theorem 2.1, i.e., (2.20).

Moreover, from (3.49), we see that (3.50) is an equality if and only if

$$Ric(\nabla\phi, \nabla\phi)(x) = K|\nabla\phi(x)|^2, \quad \nabla^2\phi(x) = \frac{I}{n}g(x), \quad \forall x \in M. \quad (3.54)$$

Since the set  $\{\nabla\phi(x) : \phi \in C^\infty(M)\}$  spans the whole tangent space  $T_x M$ , we conclude that  $(M, g)$  is Einstein manifold with constant Ricci curvature

$$Ric = Kg, \quad (3.55)$$

and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2\phi = \frac{I}{n}g. \quad (3.56)$$

On the other hand, we can check that (3.50) becomes an equality if and only if the differential inequality (2.19) becomes an equality, i.e.,

$$-H''(\rho(t)) = \frac{H'^2(\rho(t))}{n} + KW_2^2(\rho(0), \rho(1)).$$

That is to say, the Fisher information  $I = H'(\rho(t))$  satisfies the differential equation (3.51). This completes the proof of Theorem 3.9.  $\square$

**Remark 3.10** *In particular, if  $Ric \geq 0$ , then  $I(t) = H'(t)$  satisfies  $I' + \frac{I^2}{n} \leq 0$ . This yields  $\frac{d}{dt} \left( \frac{1}{I} - \frac{t}{n} \right) \geq 0$ , and hence  $I(t) \leq \frac{n}{Cn+t}$ , where  $C := \lim_{t \rightarrow 0^+} \left( \frac{1}{I} - \frac{t}{n} \right) = \frac{1}{I(0)}$ .*

In particular, we have the following rigidity theorem.

**Theorem 3.11** *If  $Ric \geq 0$ , and  $p \geq 1 - \frac{1}{n}$ , we have*

$$\frac{d^2}{dt^2} N_n(\rho(t)) \leq 0, \quad (3.57)$$

*i.e.,  $N_n(\rho(t))$  is concave in  $t$  on  $[0, \infty)$ . Moreover, under the assumption  $Ric \geq 0$  and  $p \geq 1 - \frac{1}{n}$ , the equality in (3.68) holds at some  $t = \tau > 0$  if and only if  $(M, g)$  is Ricci flat, i.e.,  $Ric = 0$ , and  $\phi$  satisfies the Hessian soliton equation*

$$\nabla^2 \phi = \frac{I}{n} g.$$

*Furthermore, in the case  $(M, g)$  is a complete Riemannian manifold with bounded geometry condition as in Theorem 3.1, and  $(\rho(t), \phi(t))$  is a smooth Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold satisfying growth condition as required in Theorem 3.1. Then, under the assumption  $Ric \geq 0$  and  $p \geq 1 - \frac{1}{n}$ , the equality in (3.57) holds at some  $t = \tau > 0$  if and only if  $M$  is isometric to  $\mathbb{R}^n$ , and  $(\rho, \phi) = (\rho_n, \phi_n)$ .*

*Proof.* The last statement of rigidity theorem is indeed a reformulation of Theorem 3.5.  $\square$

In general case of weighted Riemannian manifolds, we have the following

**Theorem 3.12** *Under the same condition and notation as in Theorem 3.2, we have*

$$\frac{d^2 N_m}{dt^2} \leq -\frac{KN_m}{m} \int_M |\nabla \phi|^2 \rho d\mu - \frac{N_m}{m^2} \int_M |L\phi - I|^2 \rho d\mu. \quad (3.58)$$

*This indeed improves the entropy power concavity inequality EPCI in Theorem 3.2, i.e.,*

$$\frac{d^2 N_m}{dt^2} \leq -\frac{KN_m}{m} \int_M |\nabla \phi|^2 \rho d\mu. \quad (3.59)$$

*Moreover, the equality in (3.59) holds if and only if  $(M, g, V)$  is a  $(K, m)$ -Einstein manifold with constant  $m$ -dimensional Bakry-Emery Ricci curvature*

$$Ric_{m,n}(L) = Kg,$$

*and  $\phi$  satisfies the Hessian soliton equation*

$$\nabla^2 \phi = \frac{I}{m} g.$$

*where  $I = \int_M L\phi \rho d\mu$  is the Fisher information associated with the geodesic flow on  $TP_2(M, \mu)$ , and satisfies the differential equation*

$$-I' = \frac{I^2}{m} + KW_2^2(\rho(0), \rho(1)). \quad (3.60)$$

*Proof.* By the NIW formula (3.43) in Theorem 3.6, under the condition  $Ric_{m,n}(L) \geq Kg$ , where  $K \in C(M, \mathbb{R})$ , we have

$$\begin{aligned} \frac{m}{N_m} \frac{d^2 N_m}{dt^2} &\leq -\int_M K |\nabla \phi|^2 \rho d\mu - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 \rho d\mu \\ &\quad - \frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \int_M \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|_{\text{HS}}^2 \rho d\mu. \end{aligned} \quad (3.61)$$

This yields

$$\frac{d^2 N_m}{dt^2} \leq -\frac{N_m}{m} \int_M K |\nabla \phi|^2 \rho d\mu - \frac{N_m}{m^2} \int_M |L\phi - I|^2 \rho d\mu, \quad (3.62)$$

which implies

$$\frac{d^2 N_m}{dt^2} \leq -\frac{N_m}{m} \int_M K |\nabla \phi|^2 \rho d\mu. \quad (3.63)$$

In particular, when  $K$  is a constant, we have

$$\frac{d^2 N_m}{dt^2} \leq -\frac{KN_m}{m} \int_M |\nabla \phi|^2 \rho d\mu - \frac{N_m}{m^2} \int_M |L\phi - I|^2 \rho d\mu. \quad (3.64)$$

The inequalities (3.62), (3.63) and (3.64) are improved versions of the entropy power concavity inequality in Theorem 3.2, i.e., if  $Ric_{m,n}(L) \geq K$  for a constant  $K \in \mathbb{R}$ , then

$$\frac{d^2 N_m}{dt^2} \leq -\frac{KN_m}{m} \int_M |\nabla \phi|^2 \rho d\mu. \quad (3.65)$$

Moreover, from (3.61) we can conclude that, the equality in (3.63) or (3.65) holds if and only if

$$Ric_{m,n}(L)(\nabla \phi, \nabla \phi) = K |\nabla \phi|^2, \quad L\phi = I = \int_M L\phi \rho d\mu,$$

and

$$L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi = 0, \quad \nabla^2 \phi = \frac{\Delta \phi}{n} g.$$

Since  $\nabla \phi$  spans the whole tangent space  $TM$ , we conclude that  $(M, g)$  is a  $(K, m)$ -Einstein manifold with constant  $m$ -dimensional Bakry-Emery Ricci curvature

$$Ric_{m,n}(L) = Kg,$$

and  $\Delta \phi = \frac{n}{m} I$ . Hence  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{I}{m} g.$$

Moreover, we can check that (3.63) becomes an equality if and only if the differential inequality (3.33) becomes an equality, i.e.,

$$-H''(\rho(t)) = \frac{H'^2(\rho(t))}{m} + \int_M K |\nabla \phi(t)|^2 \rho(t) d\mu. \quad (3.66)$$

In particular, the inequality (3.65) becomes an equality if and only if the differential inequality (2.19) becomes an equality, i.e.,

$$-H''(\rho(t)) = \frac{H'^2(\rho(t))}{m} + KW_2^2(\rho(0), \rho(1)). \quad (3.67)$$

Equivalently, the Fisher information  $I = H'(\rho(t))$  satisfies the Riccati equation (3.60). This completes the proof of Theorem 3.2.  $\square$

**Remark 3.13** *In particular, if  $Ric_{m,n}(L) \geq 0$ , then  $I(t) = H'(t)$  satisfies  $I' + \frac{I^2}{m} \leq 0$ . This yields  $\frac{d}{dt} \left( \frac{1}{I} - \frac{t}{m} \right) \geq 0$ , and hence  $I(t) \leq \frac{m}{t+C_m}$ , where  $C := \lim_{t \rightarrow 0} \left( \frac{1}{I} - \frac{t}{m} \right) = \frac{1}{I(0)}$ .*

In particular, we have the following rigidity theorem.

**Theorem 3.14** *If  $\text{Ric}_{m,n}(L) \geq 0$ , and  $p \geq 1 - \frac{1}{m}$ , we have*

$$\frac{d^2}{dt^2} N_m(\rho(t)) \leq 0, \quad (3.68)$$

*i.e.,  $N_m(\rho(t))$  is concave in  $t$  on  $[0, \infty)$ . Moreover, under the assumption  $\text{Ric}_{m,n}(L) \geq 0$  and  $p \geq 1 - \frac{1}{m}$ , the equality in (3.68) holds at some  $t = \tau > 0$  if and only if  $(M, g, V)$  is quasi-Ricci flat, i.e.,  $\text{Ric}_{m,n}(L) = 0$ , and  $\phi$  satisfies the Hessian soliton equation*

$$\nabla^2 \phi = \frac{I}{m} g.$$

*Furthermore, in the case  $(M, g, V)$  is a complete Riemannian manifold with bounded geometry condition as in Theorem 3.1, and  $(\rho(t), \phi(t))$  is a smooth Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold satisfying growth condition as required in Theorem 3.1. Then, assumption  $\text{Ric}_{m,n}(L) \geq 0$  and  $p \geq 1 - \frac{1}{m}$ , the equality in (3.68) holds at some  $t = \tau > 0$  if and only if  $M$  is isometric to  $\mathbb{R}^n$ ,  $m = n$ ,  $V$  is a constant and  $(\rho, \phi) = (\rho_n, \phi_n)$ .*

*Proof.* The last statement of rigidity theorem is indeed a reformulation of Theorem 3.5.  $\square$

## 4 Differential inequalities for Rényi entropy

In this section we prove the differential inequalities for the Rényi entropy and its power along the geodesic flow on the Wasserstein space over complete Riemannian manifolds with the  $\text{CD}(K, m)$  curvature-dimension condition. We prove the NIW formula which indicates the relationship between the Rényi entropy power  $N$ , the Fisher information  $I$  and the  $W$ -entropy associated with the geodesic flow on the Wasserstein space over complete Riemannian manifolds. Moreover, we prove that the rigidity models of the Rényi entropy power  $N$  is the  $(K, m)$ -Einstein manifolds, i.e.,  $\text{Ric}_{m,n}(L) = Kg$ , coupled with a Hessian soliton. In particular, when  $m = n$ , the rigidity models of the Rényi entropy power  $N$  is the  $K$ -Einstein manifolds, i.e.,  $\text{Ric} = Kg$ , coupled with a Hessian soliton.

Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold with a weighted volume measure  $d\mu = e^{-V} dv$  satisfying the Bakry-Emery curvature-dimension  $\text{CD}(K, m)$ -condition, i.e.,  $\text{Ric}_{m,n}(L) \geq Kg$ , where  $m \in \mathbb{R}^+$  and  $K \in \mathbb{R}$ ,  $m \geq n$ .

Let  $(\rho, \phi)$  be a smooth solution to the geodesic flow on  $TP_2(M, \mu)$ , i.e.,  $(\rho, \phi)$  is a smooth solution to the continuity equation (2.12) and the Hamilton-Jacobi equation (2.13). Define the  $p$ -th Rényi entropy by

$$H_p(\rho) = \frac{1}{1-p} \log \int_M \rho^p d\mu, \quad (4.69)$$

and the  $(m, p)$ -Rényi entropy power  $N_{m,p}$  by

$$N_{m,p}(\rho) = \exp\left(\frac{1}{m} H_p(\rho)\right). \quad (4.70)$$

We need the following entropy dissipation formulas. In the case where  $M$  is compact, the result is due to Lott and Villani. See [33, 49, 50]

**Theorem 4.1** *Let  $(M, g)$  be a complete Riemannian manifold and  $V \in C^2(M)$ . Suppose that  $\text{Ric}(L) = \text{Ric} + \nabla^2 V$  is uniformly bounded on  $M$ . Let  $d\mu = e^{-V} dv$ , and*

$$U(\rho) = \int_M e(\rho) d\mu, \quad \rho \in P_2(M, \mu),$$

where  $e \in C^2(\mathbb{R}^+, \mathbb{R}^+)$ . Set<sup>2</sup>

$$p_1(r) := re'(r) - e(r), \quad p_2(r) := rp_1'(r) - p_1(r).$$

Let  $(\rho(t), \phi(t))$  be a smooth solution to the Benamou-Brenier geodesic equation on  $P_2(M, e^{-V} dv)$  satisfying the following growth condition

$$\int_M [|\nabla\phi|^2 + |L\phi|^2 + |e'(\rho)|^2 + |\nabla e'(\rho)|^2 + |L\phi|^2 |p_1'(\rho)|] \rho d\mu + \int_M \frac{|\nabla e(\rho)|^2}{\rho} d\mu < \infty,$$

and

$$\int_M [|\nabla\phi|^2 + |\nabla^2\phi|^2 + |L\phi|^2 + |\nabla L\phi|^2] p_1(\rho) d\mu < \infty,$$

Then the following entropy dissipation formulas hold

$$\frac{d}{dt} U(\rho(t)) = \int_M \nabla\phi \cdot \nabla p_1(\rho) d\mu = - \int_M L\phi p_1(\rho) d\mu, \quad (4.71)$$

and

$$\frac{d^2}{dt^2} U(\rho(t)) = \int_M \Gamma_2(\phi, \phi) p_1(\rho) d\mu + \int_M (L\phi)^2 p_2(\rho) d\mu. \quad (4.72)$$

where

$$\Gamma_2(\phi, \phi) := \frac{1}{2} L|\nabla\phi|^2 - \nabla L\phi \cdot \nabla\phi = \|\nabla^2\phi\|^2 + Ric(L)(\nabla\phi, \nabla\phi).$$

*Proof.* In the case where  $M$  is compact, see [33, 49, 50]. In the case where  $(M, g)$  is a complete Riemannian manifold and  $(\rho, \phi)$  satisfies the required growth condition at infinity, we can use the same argument as used for the proof of Theorem 3.1 in [27] to prove (4.71) and (4.72).

Let  $\eta_k$  be an increasing sequence of functions in  $C_0^\infty(M)$  such that  $0 \leq \eta_k \leq 1$ ,  $\eta_k = 1$  on  $B(o, k)$ ,  $\eta_k = 0$  on  $M \setminus B(o, 2k)$ , and  $\|\nabla\eta_k\| \leq \frac{1}{k}$ . By standard argument and integration by parts, we have

$$\begin{aligned} \partial_t \int_M e(\rho) \eta_k d\mu &= \int_M \partial_t e(\rho) \eta_k d\mu = \int_M e'(\rho) \partial_t \rho \eta_k d\mu \\ &= \int_M \nabla_\mu^* (\rho \nabla\phi) e'(\rho) \eta_k d\mu = \int_M \langle \rho \nabla\phi, \nabla(e'(\rho) \eta_k) \rangle d\mu \\ &= \int_M \rho \langle \nabla\phi, \nabla e'(\rho) \rangle \eta_k d\mu + \int_M \rho \langle \nabla\phi, e'(\rho) \nabla\eta_k \rangle d\mu \\ &:= I_1(k) + I_2(k). \end{aligned}$$

Here

$$\begin{aligned} I_1(k) &= \int_M \langle \nabla\phi, \nabla e'(\rho) \rangle \eta_k \rho d\mu = \int_M \langle \nabla\phi, \nabla p_1(\rho) \rangle \eta_k d\mu, \\ I_2(k) &= \int_M \rho \langle \nabla\phi, e'(\rho) \nabla\eta_k \rangle d\mu. \end{aligned}$$

---

<sup>2</sup>In Lott-Villani [33] and Villani [49, 50], the function  $p_1(r)$  was denoted by  $p(r)$ . Here we use the notation  $p_1(r)$  instead of  $p(r)$  to avoid the confusion of the exponent  $p$  and the function  $p(r)$ .



Under the assumption of theorem, we have  $\int_M |\nabla\phi|^2 \rho d\mu < \infty$  and  $\int_M |\nabla e'(\rho)|^2 \rho d\mu < \infty$ , and hence  $|\langle \nabla\phi, \nabla e'(\rho) \rangle| \in L^1(M, \rho\mu)$ . By the Lebesgue dominated convergence theorem, we have

$$I_1(k) \rightarrow \int_M \langle \nabla\phi, \nabla e'(\rho) \rangle \rho d\mu = \int_M \langle \nabla\phi, \nabla p_1(\rho) \rangle d\mu. \quad (4.73)$$

Note that

$$\begin{aligned} I_1(k) &= \int_M \nabla_\mu^*(\eta_k \rho \nabla\phi) e'(\rho) d\mu \\ &= - \int_M \eta_k \rho L\phi e'(\rho) d\mu - \int_M \langle \nabla\eta_k, \nabla\phi \rangle e'(\rho) \rho d\mu - \int_M \eta_k \langle \nabla\rho, \nabla\phi \rangle e'(\rho) d\mu. \\ &= - \int_M \eta_k \rho L\phi e'(\rho) d\mu - \int_M \langle \nabla\eta_k, \nabla\phi \rangle e'(\rho) \rho d\mu - \int_M \eta_k \langle \nabla e(\rho), \nabla\phi \rangle d\mu. \end{aligned}$$

Under the assumption of theorem, we have  $\int_M |L\phi|^2 \rho d\mu < \infty$ ,  $\int_M |e'(\phi)|^2 \rho d\mu < \infty$ , and  $\int_M \frac{|\nabla e(\rho)|^2}{\rho} d\mu < \infty$ ,  $\int_M |\nabla\phi|^2 \rho d\mu < \infty$ . Hence, the Cauchy-Schwarz inequality yields

$$\int_M |L\phi| |e'(\rho)| \rho d\mu + \int_M |\nabla\phi| |e'(\rho)| \rho d\mu + \int_M |\nabla e(\rho)| |\nabla\phi| d\mu < \infty.$$

Noting that  $0 \leq \eta_k \leq 1$ ,  $\eta_k \rightarrow 1$  and  $|\nabla\eta_k| \leq 1/k$ , the Lebesgue dominated convergence theorem yields

$$\begin{aligned} \lim_{k \rightarrow \infty} I_1(k) &= - \int_M \rho L\phi e'(\rho) d\mu - \int_M \langle \nabla e(\rho), \nabla\phi \rangle d\mu \\ &= - \int_M \rho L\phi e'(\rho) d\mu + \int_M e(\rho) L\phi d\mu \\ &= - \int_M L\phi p_1(\rho) d\mu. \end{aligned} \quad (4.74)$$

On the other hand, under the assumption of theorem, we have  $\int_M |\nabla\phi|^2 \rho d\mu < \infty$  and  $\int_M |e'(\rho)|^2 \rho d\mu < \infty$ , which yields  $\int_M |e'(\rho)| |\nabla\phi| \rho d\mu < \infty$ . By the Lebesgue dominated convergence theorem, as  $|\nabla\eta_k| \leq 1/k$ , we have

$$I_2(k) = \int_M \rho \langle \nabla\phi, e'(\rho) \nabla\eta_k \rangle d\mu \rightarrow 0. \quad (4.75)$$

Combining (4.73), (4.74) with (4.75), we have

$$\begin{aligned} \partial_t U(\rho) &= \lim_{k \rightarrow \infty} [I_1(k) + I_2(k)] \\ &= \int_M \langle \nabla\phi, \nabla p_1(\rho) \rangle d\mu \\ &= - \int_M L\phi p_1(\rho) d\mu. \end{aligned}$$

This completes the proof of (4.71).

Next, we prove (4.72). By standard argument, we have

$$\begin{aligned} \partial_t \int_M L\phi p_1(\rho) \eta_k d\mu &= \int_M \partial_t (L\phi p_1(\rho)) \eta_k d\mu \\ &= \int_M L\partial_t \phi p_1(\rho) \eta_k d\mu + \int_M L\phi \partial_t p_1(\rho) \eta_k d\mu \\ &= -\frac{1}{2} \int_M L|\nabla\phi|^2 p_1(\rho) \eta_k d\mu + \int_M L\phi p_1'(\rho) \nabla_\mu^*(\rho \nabla\phi) \eta_k d\mu \\ &:= I_3(k) + I_4(k). \end{aligned} \quad (4.76)$$

Here

$$\begin{aligned} I_3(k) &= -\frac{1}{2} \int_M L |\nabla \phi|^2 p_1(\rho) \eta_k d\mu, \\ I_4(k) &= \int_M L \phi p_1'(\rho) \nabla_\mu^*(\rho \nabla \phi) \eta_k d\mu. \end{aligned}$$

By the weighted Bochner formula and  $|Ric(L)| \leq C$ , under the assumption  $\int_M [|\nabla \phi|^2 + |\nabla L \phi|^2 + |\nabla^2 \phi|^2] p_1(\rho) d\mu < \infty$ , we have

$$\begin{aligned} \int_M |L |\nabla \phi|^2| p_1(\rho) d\mu &= 2 \int_M |\langle \nabla \phi, \nabla L \phi \rangle + |\nabla^2 \phi|^2 + Ric(L)(\nabla \phi, \nabla \phi)| p_1(\rho) d\mu \\ &\leq 2 \int_M [|\nabla \phi| |\nabla L \phi| + |\nabla^2 \phi|^2 + C |\nabla \phi|^2] p_1(\rho) d\mu < \infty. \end{aligned}$$

Using the fact  $0 \leq \eta_k \leq 1$  and  $\eta_k \rightarrow 1$ , the Lebesgue dominated convergence theorem yields

$$I_3(k) \rightarrow -\frac{1}{2} \int_M L |\nabla \phi|^2 p_1(\rho) d\mu. \quad (4.77)$$

On the other hand, using the integration by parts formula, we have

$$\begin{aligned} I_4(k) &= \int_M L \phi p_1'(\rho) \nabla_\mu^*(\rho \nabla \phi) \eta_k d\mu \\ &= \int_M L \phi p_1'(\rho) [-\rho L \phi - \langle \nabla \rho, \nabla \phi \rangle] \eta_k d\mu \\ &= - \int_M [|L \phi|^2 \rho p_1'(\rho) + L \phi \langle \nabla p_1(\rho), \nabla \phi \rangle] \eta_k d\mu \\ &= I_5(k) + I_6(k). \end{aligned}$$

Note that

$$\begin{aligned} I_6(k) &= - \int_M L \phi \langle \nabla p_1(\rho), \nabla \phi \rangle \eta_k d\mu \\ &= - \int_M \langle \eta_k L \phi \nabla \phi, \nabla p_1(\rho) \rangle d\mu \\ &= - \int_M \nabla_\mu^*(\eta_k L \phi \nabla \phi) p_1(\rho) d\mu \\ &= \int_M [\eta_k |L \phi|^2 + L \phi \nabla e_k \cdot \nabla \phi + \eta_k \nabla L \phi \cdot \nabla \phi] p_1(\rho) d\mu. \end{aligned}$$

Under the assumption of theorem, we have

$$\int_M |L \phi|^2 |p_1'(\rho)| \rho d\mu + \int_M [|L \phi|^2 + |\nabla \phi|^2 + |\nabla L \phi|^2] p_1(\rho) d\mu < \infty.$$

Using again the fact  $0 \leq \eta_k \leq 1$ ,  $\eta_k \rightarrow 1$  and  $|\nabla \eta_k| \leq 1/k$ , the Lebesgue dominated convergence theorem yields

$$\begin{aligned} \lim_{k \rightarrow \infty} I_4(k) &= \lim_{k \rightarrow \infty} [I_5(k) + I_6(k)] \\ &= - \int_M |L \phi|^2 [\rho p_1'(\rho) - p_1(\rho)] d\mu + \int_M \langle \nabla L \phi, \nabla \phi \rangle p_1(\rho) d\mu \\ &= - \int_M |L \phi|^2 p_2(\rho) d\mu + \int_M \langle \nabla L \phi, \nabla \phi \rangle p_1(\rho) d\mu. \end{aligned} \quad (4.78)$$

Combining (4.76), (4.77) and (4.78), we have

$$\begin{aligned}
\partial_t^2 U(\rho) &= - \lim_{k \rightarrow \infty} [I_3(k) + I_4(k)] \\
&= \frac{1}{2} \int_M L |\nabla \phi|^2 p_1(\rho) d\mu + \int_M |L\phi|^2 p_2(\rho) d\mu - \int_M \langle \nabla L\phi, \nabla \phi \rangle p_1(\rho) d\mu \\
&= \int_M |L\phi|^2 p_2(\rho) d\mu + \int_M \Gamma_2(\phi, \phi) p_1(\rho) d\mu.
\end{aligned}$$

This completes the proof of (4.72).  $\square$

**Theorem 4.2** *Under the same condition as in Theorem 4.1, we have*

$$\frac{d^2}{dt^2} U(\rho(t)) \geq \int_M \left[ \left( p_2(\rho) + \frac{1}{m} p_1(\rho) \right) (L\phi)^2 + Ric_{m,n}(L)(\nabla \phi, \nabla \phi) p_1(\rho) \right] d\mu. \quad (4.79)$$

Assume that  $p_2(r) + \frac{1}{m} p_1(r) \geq \sigma p_1(r)$  and  $Ric_{m,n}(L) \geq K$ , then

$$\frac{d^2}{dt^2} U(\rho(t)) \geq \sigma \left[ \frac{d}{dt} U(\rho(t)) \right]^2 \left[ \int_M p_1(\rho(t)) d\mu \right]^{-1} + \int_M K |\nabla \phi(t)|^2 p_1(\rho(t)) d\mu. \quad (4.80)$$

In particular, if  $p_2(r) + \frac{1}{m} p_1(r) \geq 0$  and  $Ric_{m,n}(L) \geq K$ , we have

$$\frac{d^2}{dt^2} U(\rho(t)) \geq \int_M K |\nabla \phi(t)|^2 p_1(\rho(t)) d\mu. \quad (4.81)$$

*Proof.* By (4.72) and using the Bochner inequality

$$\Gamma_2(\phi, \phi) \geq \frac{|L\phi|^2}{m} + Ric_{m,n}(L)(\nabla \phi, \nabla \phi), \quad (4.82)$$

we can derive (4.79). Under the assumption  $p_2(r) + \frac{1}{m} p_1(r) \geq \sigma p_1(r)$  and the curvature-dimension condition  $Ric_{m,n}(L) \geq Kg$ , we have

$$\frac{d^2}{dt^2} U(\rho(t)) \geq \sigma \int_M |L\phi|^2 p_1(\rho) d\mu + \int_M K |\nabla \phi|^2 p_1(\rho) d\mu.$$

By the Cauchy-Schwarz inequality

$$\left( \int_M L\phi p_1(\rho) d\mu \right)^2 \leq \left( \int_M |L\phi|^2 p_1(\rho) d\mu \right) \left( \int_M p_1(\rho) d\mu \right).$$

Since  $\sigma \geq 0$ , we have

$$\frac{d^2}{dt^2} U(\rho(t)) \geq \sigma \left( \int_M L\phi p_1(\rho) d\mu \right)^2 \left( \int_M p_1(\rho) d\mu \right)^{-1} + \int_M K |\nabla \phi|^2 p_1(\rho) d\mu.$$

Using the fact  $U'(\rho) = - \int_M p_1(\rho) L\phi d\mu$ , we then obtain (4.80)  $\square$

#### 4.1 Differential inequality for Rényi entropy

In particular, for  $e(r) = \frac{r^p}{p-1}$ , we have  $U(\rho) = \frac{1}{p-1} \int_M \rho^p d\mu$ , and

$$p_1(r) = re'(r) - e(r) = r^p, \quad p_2(r) = rp_1'(r) - p_1(r) = (p-1)r^p.$$

Moreover

$$\begin{aligned} \nabla p_1(\rho) &= \nabla \rho e'(\rho) + \rho e''(\rho) \nabla \rho - e'(\rho) \nabla \rho \\ &= \rho e''(\rho) \nabla \rho. \end{aligned}$$

In this case, we have

$$p_2(\rho) + \frac{1}{m} p_1(\rho) = \sigma p_1(\rho)$$

with

$$\sigma := p - 1 + \frac{1}{m}.$$

**Theorem 4.3** *Let  $e(r) = \frac{r^p}{p-1}$  and  $U(\rho) = \frac{1}{p-1} \int_M \rho^p d\mu$ . Assume that  $\text{Ric}_{m,n}(L) \geq Kg$ , where  $K \in C(M, \mathbb{R})$ . Then*

$$(p-1)U(\rho)U''(\rho) \geq \left(p-1 + \frac{1}{m}\right) (U'(\rho))^2 + (p-1)U(\rho) \int_M K |\nabla \phi|^2 p_1(\rho) d\mu. \quad (4.83)$$

When  $p \geq 1 - \frac{1}{m}$ , we have

$$\frac{d^2}{dt^2} N_{m,p}(\rho) \leq -\frac{N_{m,p}(\rho)}{m} \frac{\int_M K |\nabla \phi|^2 \rho^p d\mu}{\int_M \rho^p d\mu}. \quad (4.84)$$

If  $\text{Ric}_{m,n}(L) \geq K$  for  $K \in \mathbb{R}$ , we have

$$\frac{d^2}{dt^2} N_{m,p}(\rho) \leq -\frac{KN_{m,p}(\rho)}{m} \frac{\int_M |\nabla \phi|^2 \rho^p d\mu}{\int_M \rho^p d\mu}.$$

In particular, if  $\text{Ric}_{m,n}(L) \geq 0$ , i.e., if the  $CD(0, m)$ -condition holds, we have

$$\frac{d^2}{dt^2} N_{m,p}(\rho) \leq 0.$$

i.e., if  $\text{Ric}_{m,n}(L) \geq 0$ , and  $p \geq 1 - \frac{1}{m}$ , then  $N_{m,p}(\rho(t))$  is concave in  $t$  on  $[0, \infty)$ .

*Proof.* The inequality (4.83) follows from (4.80). Note that

$$H_p'(\rho) = \frac{1}{1-p} \frac{\partial_t \int_M \rho^p d\mu}{\int_M \rho^p d\mu} = \frac{1}{1-p} \frac{U'}{U},$$

and

$$H_p''(\rho) = \frac{1}{1-p} \frac{U''U - U'^2}{U^2}.$$

Thus

$$\begin{aligned} H_p'' + \frac{1}{m} H_p'^2 &= \frac{1}{p-1} \frac{U'^2 - U''U}{U^2} + \frac{1}{m(p-1)^2} \frac{U'^2}{U^2} \\ &= \frac{1}{(p-1)^2 U^2} \left[ \left(p-1 + \frac{1}{m}\right) U'^2 - (p-1)U''U \right] \\ &\leq -\frac{\int_M K |\nabla \phi|^2 p_1(\rho) d\mu}{\int_M p_1(\rho) d\mu}. \end{aligned}$$

Note that

$$\begin{aligned}\frac{d}{dt}N_{m,p}(\rho) &= \frac{1}{m}H'_p N_{m,p}(\rho), \\ \frac{d^2}{dt^2}N_{m,p}(\rho) &= \frac{1}{m}\left(H''_p + \frac{1}{m}H_p'^2\right)N_{m,p}(\rho).\end{aligned}$$

Combining the above calculations, we derive that

$$\frac{d^2}{dt^2}N_{m,p}(\rho) \leq -\frac{N_{m,p}(\rho)}{m} \frac{\int_M K|\nabla\phi|^2 p_1(\rho)d\mu}{\int_M p_1(\rho)d\mu}.$$

This proves (4.84) since  $p_1(\rho) = \rho^p$ . When  $Ric_{m,n}(L) \geq K$  for  $K \in \mathbb{R}$ , we have

$$\frac{d^2}{dt^2}N_{m,p}(\rho) \leq -\frac{KN_{m,p}(\rho)}{m} \frac{\int_M |\nabla\phi|^2 \rho^p d\mu}{\int_M \rho^p d\mu}.$$

In particular, if  $Ric_{m,n}(L) \geq 0$ , i.e., if the  $CD(0, m)$ -condition holds, we have

$$\frac{d^2}{dt^2}N_{m,p}(\rho) \leq 0.$$

□

## 4.2 The $W$ -entropy and NIW formula for Rényi entropy

In this subsection, we introduce the  $W$ -entropy and prove the  $W$ -entropy formula for the Rényi entropy along the geodesic flow on the Wasserstein space over complete Riemannian manifolds. Then we prove the NIW formula which indicates the relationship between the entropy power  $N$ , the Fisher information  $I = H'$  and the  $W$ -entropy related to the Rényi entropy for the geodesic flow on the Wasserstein space over compact Riemannian manifolds.

When  $m \in \mathbb{N}$ , let  $\rho_m(x, t) = \frac{e^{-\frac{|x|^2}{4t^2}}}{(2\pi t^2)^{m/2}}$ , and  $\phi_m(x, t) = \frac{|x|^2}{2t}$ . Then  $(\rho_m, \phi_m)$  is a special solution to the geodesic equation on  $P_2(\mathbb{R}^m, dx)$ . See [27]. The Rényi entropy of  $\rho_m(t)$  is given by

$$\begin{aligned}H_p(\rho_m(t)) &= \frac{1}{1-p} \log \int_{\mathbb{R}^m} \rho_m(x, t)^p dx \\ &= \frac{1}{1-p} \log \int_{\mathbb{R}^m} \frac{e^{-\frac{p|x|^2}{4t^2}}}{(4\pi t^2)^{mp/2}} dx \\ &= \frac{1}{1-p} \left( \log \int_{\mathbb{R}^m} e^{-\frac{p|x|^2}{4t^2}} dx - \log(4\pi t^2)^{mp/2} \right)\end{aligned}$$

Using the fact

$$\int_{\mathbb{R}^m} e^{-\frac{p|x|^2}{4t^2}} dx = (4\pi t^2 p^{-1})^{m/2},$$

we have

$$H_p(\rho_m(t)) = \frac{n}{2} \log(4\pi e t^2) + \frac{m}{2} \frac{\log p}{p-1} - \frac{m}{2}.$$

Note that

$$\lim_{p \rightarrow 1} H_p(\rho_m(t)) = H_1(\rho_m(t)) = \frac{m}{2} \log(4\pi e t^2).$$

**Proof of Theorem 2.3.** To simplify the notation, we write  $H'_p = H'_p(\rho(t)) = \frac{d}{dt}H_p(\rho(t))$ , etc. Note that

$$H'_p = \frac{1}{1-p} \frac{U'}{U} = \int_M L\phi d\gamma = I_p,$$

and

$$H''_p = \frac{1}{1-p} \frac{U''U - U'^2}{U^2} = -\frac{U''}{\int_M p_1(\rho)d\mu} + (p-1) \left( \int_M L\phi d\gamma \right)^2.$$

Now

$$\begin{aligned} U'' &= \int_M \left[ \left( p-1 + \frac{1}{m} \right) |L\phi|^2 + Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \right] p_1(\rho)d\mu \\ &\quad + \int_M \left( \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|^2 \right) p_1(\rho)d\mu. \end{aligned}$$

Therefore

$$\begin{aligned} H''_p + \frac{2}{t}H'_p - \frac{m}{t^2} &= -\int_M \left[ \left( p-1 + \frac{1}{m} \right) |L\phi|^2 + Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \right] d\gamma \\ &\quad - \int_M \left( \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|^2 \right) d\gamma \\ &\quad + (p-1) \left( \int_M L\phi d\gamma \right)^2 + \frac{2}{t} \int_M L\phi d\gamma - \frac{m}{t^2}. \end{aligned}$$

By

$$\begin{aligned} \left\| \nabla^2\phi - \frac{g}{t} \right\|_{\text{HS}}^2 &= \frac{1}{n} \left| \Delta\phi - \frac{n}{t} \right|^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \\ &= \frac{1}{n} \left| L\phi - \frac{m}{t} + \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right) \right|^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \\ &= \frac{1}{m} \left| L\phi - \frac{m}{t} \right|^2 - \frac{1}{m-n} \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right)^2 \\ &\quad + \frac{m-n}{mn} \left[ L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right]^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2, \quad (4.85) \end{aligned}$$

we have

$$\begin{aligned} H''_p + \frac{2}{t}H'_p - \frac{m}{t^2} &= -\int_M \left[ \left( p-1 + \frac{1}{m} \right) |L\phi|^2 + Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \right] d\gamma \\ &\quad - \int_M \left\| \nabla^2\phi - \frac{g}{t} \right\|^2 d\gamma - \frac{1}{m-n} \int_M \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right)^2 d\gamma \\ &\quad + \frac{1}{m} \int_M \left| L\phi - \frac{m}{t} \right|^2 d\gamma + (p-1) \left( \int_M L\phi d\gamma \right)^2 + \frac{2}{t} \int_M L\phi d\gamma - \frac{m}{t^2} \\ &= -(p-1) \left[ \int_M |L\phi|^2 d\gamma - \left( \int_M L\phi d\gamma \right)^2 \right] - \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi) d\gamma \\ &\quad - \int_M \left( \left\| \nabla^2\phi - \frac{g}{t} \right\|^2 + \frac{1}{m-n} \left( \nabla V \cdot \nabla\phi + \frac{m-n}{t} \right)^2 \right) d\gamma. \end{aligned}$$

By the definition formula of  $W_m$ , we have

$$\frac{1}{t} \frac{d}{dt} W_{m,p}(\rho, t) = H_p'' + \frac{2}{t} H_p' - \frac{m}{t^2}.$$

This completes the proof of (2.26).  $\square$

**Theorem 4.4** (NIW formula) *Under the same notations as in Theorem 4.3, we have*

$$\frac{m}{N_{m,p}(t)} \frac{d^2}{dt^2} N_{m,p}(\rho(t)) = \frac{1}{m} \left| I_p - \frac{m}{t} \right|^2 + \frac{1}{t} \frac{d}{dt} W_{m,p}(\rho, t). \quad (4.86)$$

*In particular, when  $\text{Ric}_{m,n}(L) \geq Kg$ , we have*

$$\frac{d}{dt} W_{m,p}(\rho, t) \leq -t \int_M K |\nabla \phi|^2 d\gamma - \frac{t}{m} \left| I_p - \frac{m}{t} \right|^2. \quad (4.87)$$

*Proof.* Note that

$$N_{m,p}'' = \frac{1}{m} N_{m,p} \left( H_p'' + \frac{1}{m} H_p'^2 \right),$$

and

$$\begin{aligned} H_p'' + \frac{1}{m} H_p'^2 &= H_p'' + \frac{2}{t} H_p' - \frac{m}{t^2} + \frac{1}{m} H_p'^2 - \frac{2}{t} H_p' + \frac{m}{t^2} \\ &= \frac{1}{t} \frac{d}{dt} W_{m,p}(t) + \frac{1}{m} \left| H_p' - \frac{m}{t} \right|^2. \end{aligned}$$

This completes the proof of (4.86). By Theorem 4.3 and  $H_p' = I_p$ , we obtain (4.86).  $\square$

### 4.3 Rigidity theorem of $W$ -entropy for Rényi entropy

Note that, when  $m \in \mathbb{N}$  and  $M = \mathbb{R}^m$ , we have

$$\frac{d^2}{dt^2} H_p(\rho_m(t)) + \frac{2}{t} \frac{d}{dt} H_p(\rho_m(t)) - \frac{m}{t^2} = 0. \quad (4.88)$$

and

$$W_{m,p}(\rho, t) = \frac{d}{dt} (t H_{m,p}(\rho, t)) = 0. \quad (4.89)$$

Due to this observation and inspired by Theorem 3.5, we have the following rigidity theorem of the  $W$ -entropy for the Rényi entropy along the Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold with  $\text{CD}(0, m)$ -condition.

**Theorem 4.5** *Let  $(M, g, V)$  be a complete Riemannian manifold with bounded geometry condition as in Theorem 4.1, and let  $(\rho(t), \phi(t))$  be a smooth Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold satisfying growth condition as required in Theorem 4.1. Suppose that  $\text{Ric}_{m,n}(L) \geq 0$ ,  $p \geq 1$ , and  $\frac{d}{dt} W_{m,p}(\rho, t) = 0$  at some  $t = \tau > 0$ . Then  $M$  is isometric to  $\mathbb{R}^n$ ,  $m = n$ ,  $V$  is a constant and  $(\rho, \phi) = (\rho_m, \phi_m)$ .*

*Proof.* The proof is similar to the proof of Theorem 3.5 given in [27]. For the completeness of the paper, we reproduce it as follows.

Under the condition  $\text{Ric}_{m,n}(L) \geq 0$ , and using the  $W$ -entropy formula (2.26), we see that  $\frac{d}{dt} W_{m,p}(\rho, t) = 0$  at some  $t = \tau > 0$  if and only if

$$\text{Var}_\mu(L\phi) = 0, \quad \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) = 0,$$

and

$$\left\| \nabla^2\phi - \frac{g}{\tau} \right\|^2 + \frac{1}{m-n} \left( \nabla V \cdot \nabla\phi + \frac{m-n}{\tau} \right)^2 = 0.$$

Equivalently,  $\nabla^2\phi = \frac{g}{\tau}$ ,  $\text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) = 0$  and  $\nabla\phi \cdot \nabla V + \frac{m-n}{\tau} = 0$ . By the Taylor expansion along geodesics,  $\phi(\cdot, \tau)$  has a unique minimum point  $x_0 \in M$ , and every level set of  $\phi$  is compact subset of  $M$ . Thus,  $\phi(\cdot, \tau)$  is a strongly convex exhaustion function on complete non-compact Riemannian manifold  $M$ . By Theorem 3 in [17],  $M$  is diffeomorphic to  $\mathbb{R}^n$ . Integrating  $\nabla^2\phi(\cdot, \tau) = \frac{g}{\tau}$  along the shortest geodesics between  $x_0$  and  $x$  on  $M$  shows that

$$2\tau(\phi(x, \tau) - \phi(x_0, \tau)) = d^2(x_0, x), \quad \forall x \in M.$$

This yields

$$\Delta d^2(x_0, x) = 2n, \quad \forall x \in M,$$

which implies that  $(M, g)$  is isometric to the Euclidean space  $(\mathbb{R}^n, (\delta_{ij}))$ . For this, see [30] and references therein. By the generalized Cheeger-Gromoll splitting theorem (see Theorem 1.3, p. 565 in [15]), we can derive that  $V$  must be a constant and  $m = n$ .

Thus  $\phi(\cdot, \tau) \in C^\infty(\mathbb{R}^n)$  satisfies  $\nabla^2\phi(x, \tau) = \frac{\delta_{ij}}{\tau}$ . This yields  $\nabla\phi(x, \tau) = \frac{x}{\tau}$  under the assumption  $\nabla\phi(0, \tau) = 0$ , and  $\phi(x, \tau) = \frac{\|x\|^2}{2\tau}$  up to an additional constant. By the Hopf-Lax formula for the solution to the Hamilton-Jacobi equation (2.13) with  $\phi(x, \tau) = \frac{\|x\|^2}{2\tau}$ , we have

$$\phi(x, t) = \inf_{y \in \mathbb{R}^n} \left\{ \frac{\|y\|^2}{2\tau} + \frac{\|x - y\|^2}{2(t - \tau)} \right\} = \frac{\|x\|^2}{2t}, \quad \forall t > \tau, x \in \mathbb{R}^n.$$

By the uniqueness of the smooth solution to the Hamilton-Jacobi equation (2.13), we see that  $\phi(x, t) = \frac{\|x\|^2}{2t} = \phi_m(x, t)$  for all  $t > 0$ . Solving the transport equation (2.12) with the initial data  $\lim_{t \rightarrow 0} \rho(x, t) = \delta_0(x)$ , we have  $\rho(x, t) = \frac{e^{-\frac{\|x\|^2}{4t^2}}}{(4\pi t^2)^{n/2}} = \rho_m(x, t)$  for all  $t > 0$  and  $x \in \mathbb{R}^n$ .  $\square$

#### 4.4 Rigidity of Rényi entropy power

In this subsection, using the NIW formula, we derive an explicit formula for the second derivative of the Rényi entropy power, and prove its rigidity theorem under the  $\text{CD}(K, m)$ -condition.

**Theorem 4.6** *Under the same notations as in Theorem 4.3, we have*

$$\begin{aligned} \frac{d^2}{dt^2} N_{m,p}(\rho(t)) &= - \left( p - 1 + \frac{1}{m} \right) \frac{N_{m,p}(\rho(t))}{m} \text{Var}_\gamma(L\phi) - \frac{N_{m,p}(\rho(t))}{m} \int_M \text{Ric}_{m,n}(L)(\nabla\phi, \nabla\phi) d\gamma \\ &\quad - \frac{N_{m,p}(\rho(t))}{m} \int_M \left[ \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|_{\text{HS}}^2 \right] d\gamma, \end{aligned} \tag{4.90}$$

where

$$\text{Var}_\gamma(L\phi) = \int_M |L\phi - \gamma(L\phi)|^2 d\gamma.$$



In particular, when  $Ric_{m,n}(L) \geq Kg$  with  $K \in C(M, \mathbb{R})$  and  $p \geq 1 - \frac{1}{m}$ , we have

$$\frac{d^2}{dt^2} N_{m,p}(\rho(t)) \leq -\frac{N_{m,p}(\rho(t))}{m} \int_M K |\nabla \phi|^2 d\gamma. \quad (4.91)$$

Moreover, under the assumption  $Ric_{m,n}(L) \geq Kg$  with where  $K \in C(M, \mathbb{R})$  and  $p \geq 1 - \frac{1}{m}$ , the equality in (4.91) holds at some  $t$  if and only if  $(M, g, V)$  is  $(K, m)$ -Einstein, i.e.,

$$Ric_{m,n}(L) = Kg,$$

and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{Lp}{m} g.$$

*Proof.* By (4.72), we have

$$\frac{d^2}{dt^2} U(\rho(t)) = \int_M [|\nabla^2 \phi|^2 + Ric(L)(\nabla \phi, \nabla \phi)] p_1(\rho) d\mu + \int_M (L\phi)^2 p_2(\rho) d\mu.$$

By the fact

$$\begin{aligned} \|\nabla^2 \phi\|^2 &= \frac{|L\phi|^2}{m} - \frac{|\nabla V \cdot \nabla \phi|^2}{m-n} + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|^2 \\ &\quad + \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2, \end{aligned}$$

we have

$$\begin{aligned} \frac{d^2}{dt^2} U(\rho(t)) &= \int_M \left[ \frac{|L\phi|^2}{m} + Ric_{m,n}(L)(\nabla \phi, \nabla \phi) \right] p_1(\rho) d\mu + \int_M (L\phi)^2 p_2(\rho) d\mu \\ &\quad + \int_M \left( \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|^2 \right) p_1(\rho) d\mu \\ &= \int_M \left( \frac{1}{m} p_1(\rho) + p_2(\rho) \right) |L\phi|^2 d\mu + \int_M Ric_{m,n}(L)(\nabla \phi, \nabla \phi) p_1(\rho) d\mu \\ &\quad + \int_M \left( \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|^2 \right) p_1(\rho) d\mu. \end{aligned} \quad (4.92)$$

Hence, for  $e(r) = \frac{r^p}{p-1}$ , we have

$$\begin{aligned} \frac{d^2}{dt^2} U(\rho(t)) &= \int_M \left[ \left( p-1 + \frac{1}{m} \right) |L\phi|^2 + Ric_{m,n}(L)(\nabla \phi, \nabla \phi) \right] p_1(\rho) d\mu \\ &\quad + \int_M \left( \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|^2 \right) p_1(\rho) d\mu. \end{aligned} \quad (4.93)$$

This yields

$$\begin{aligned}
H_p'' + \frac{1}{m}H_p'^2 &= \frac{1}{(p-1)^2U^2} \left[ \left( p-1 + \frac{1}{m} \right) U'^2 - (p-1)U''U \right] \\
&= \left( p-1 + \frac{1}{m} \right) \left( \frac{U'}{(p-1)U} \right)^2 - \frac{U''}{(p-1)U} \\
&= \left( p-1 + \frac{1}{m} \right) \left[ \left( \int_M L\phi d\gamma \right)^2 - \int_M |L\phi|^2 d\gamma \right] - \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi) d\gamma \\
&\quad - \int_M \left[ \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \right] d\gamma.
\end{aligned}$$

By the fact

$$\frac{d^2}{dt^2}N_{m,p}(\rho(t)) = \frac{1}{m} \left( H_p'' + \frac{1}{m}H_p'^2 \right) N_{m,p}(\rho(t)),$$

we can derive (4.90). By the same argument as used in the proof of the rigidity part of Theorem 3.2, the rigidity part of Theorem 4.6 can be easily derived from (4.90).  $\square$

In particular, we have the following rigidity theorem.

**Theorem 4.7** *If  $Ric_{m,n}(L) \geq 0$ , and  $p \geq 1 - \frac{1}{m}$ , we have*

$$\frac{d^2}{dt^2}N_{m,p}(\rho(t)) \leq 0, \tag{4.94}$$

*i.e.,  $N_{m,p}(\rho(t))$  is concave in  $t$  on  $[0, \infty)$ . Moreover, under the assumption  $Ric_{m,n}(L) \geq 0$  and  $p \geq 1 - \frac{1}{m}$ , the equality in (4.94) holds at some  $t = \tau > 0$  if and only if  $(M, g, V)$  is quasi-Ricci flat, i.e.,  $Ric_{m,n}(L) = 0$ , and  $\phi$  satisfies the Hessian soliton equation*

$$\nabla^2\phi = \frac{I_p}{m}g.$$

*Furthermore, in the case  $(M, g, V)$  is a complete Riemannian manifold with bounded geometry condition as in Theorem 4.1, and  $(\rho(t), \phi(t))$  is a smooth Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold satisfying growth condition as required in Theorem 4.1. Then, assumption  $Ric_{m,n}(L) \geq 0$  and  $p \geq 1 - \frac{1}{m}$ , the equality in (4.94) holds at some  $t = \tau > 0$  if and only if  $M$  is isometric to  $\mathbb{R}^n$ ,  $m = n$ ,  $V$  is a constant and  $(\rho, \phi) = (\rho_m, \phi_m)$ .*

*Proof.* The last statement of rigidity theorem is indeed a reformulation of Theorem 4.5.  $\square$

In particular, for  $m = n$ ,  $V = 0$ ,  $\mu = v$  and  $L = \Delta$ , we have the following

**Theorem 4.8** *Let  $N_{n,p}$  be the  $p$ -th entropy power associated with the Rényi entropy for the geodesic flow on  $TP_2(M, v)$  as in Theorem 4.3, and  $d\gamma = \frac{\rho^p dv}{\int_M \rho^p dv}$ . Then*

$$\begin{aligned}
\frac{d^2}{dt^2}N_{n,p}(\rho(t)) &= - \left( p-1 + \frac{1}{n} \right) \frac{N_{n,p}(\rho(t))}{n} \text{Var}_\gamma(\Delta\phi) - \frac{N_{n,p}(\rho(t))}{n} \int_M Ric(\nabla\phi, \nabla\phi) d\gamma \\
&\quad - \frac{N_{n,p}(\rho(t))}{n} \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 d\gamma,
\end{aligned}$$

where

$$\text{Var}_\gamma(\Delta\phi) = \int_M \left| \Delta\phi - \int_M \Delta\phi d\gamma \right|^2 d\gamma.$$

In particular, when  $\text{Ric} \geq Kg$  with  $K \in C(M, \mathbb{R})$  and  $p \geq 1 - \frac{1}{n}$ , we have

$$\frac{d^2}{dt^2} N_{n,p}(\rho(t)) \leq -\frac{N_{n,p}(\rho(t))}{n} \int_M K |\nabla \phi|^2 d\gamma, \quad (4.95)$$

and

$$\frac{d}{dt} W_{n,p}(\rho, t) \leq -t \int_M K |\nabla \phi|^2 d\gamma - \frac{t}{n} \left| I_p - \frac{n}{t} \right|^2, \quad (4.96)$$

where  $I_p = \int_M \Delta \phi d\gamma$ . Moreover, under the assumption  $\text{Ric} \geq Kg$  for  $K \in C(M, \mathbb{R})$  and  $p \geq 1 - \frac{1}{n}$ , the equality in (4.95) or (4.96) holds at some  $t$  if and only if  $(M, g)$  is  $K$ -Einstein in the sense that

$$\text{Ric} = Kg,$$

and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{I_p}{n} g.$$

In particular, if  $\text{Ric} \geq 0$ , and  $p \geq 1 - \frac{1}{n}$ , we have

$$\frac{d^2}{dt^2} N_{n,p}(\rho(t)) \leq 0, \quad (4.97)$$

and

$$\frac{d}{dt} W_{n,p}(\rho, t) \leq -\frac{t}{n} \left| I_p - \frac{n}{t} \right|^2. \quad (4.98)$$

As a consequence,  $N_{n,p}(\rho(t))$  is concave in  $t$  on  $[0, \infty)$ , and  $W_{n,p}(\rho, t)$  is nonincreasing in  $t$  on  $[0, \infty)$ . Moreover, under the assumption  $\text{Ric} \geq 0$  and  $p \geq 1 - \frac{1}{n}$ , the equality in (4.97) or (4.98) holds at some  $t$  if and only if  $(M, g)$  is Ricci flat, i.e.,  $\text{Ric} = 0$ , and  $\phi$  satisfies the Hessian soliton equation

$$\nabla^2 \phi = \frac{I_p}{n} g.$$

Furthermore, in the case  $(M, g)$  is a complete Riemannian manifold with bounded geometry condition as in Theorem 4.1, and  $(\rho(t), \phi(t))$  is a smooth Benamou-Brenier geodesic on the Wasserstein space over Riemannian manifold satisfying growth condition as required in Theorem 4.1. Then, under the assumption  $\text{Ric} \geq 0$  and  $p \geq 1 - \frac{1}{n}$ , the equality in (4.97) holds at some  $t = \tau > 0$  if and only if  $M$  is isometric to  $\mathbb{R}^n$  and  $(\rho, \phi) = (\rho_n, \phi_n)$ .

#### 4.5 The special case $p = 1 - 1/N$ as in Sturm [45]

In the special case  $e(r) = \frac{r^p}{p-1}$  with  $p = 1 - 1/N$ , we have

$$\sigma := p - 1 + \frac{1}{m} = \frac{1}{m} - \frac{1}{N} \geq 0$$

if  $N \geq m$ . Using the same notation as in Sturm [45], let

$$S_N = - \int_M \rho^{1-1/N} d\mu.$$

Then

$$U(\rho) = \frac{1}{p-1} \int_M \rho^p d\mu = N S_N(\rho).$$

Theorem 4.3 and (4.93) imply the following

**Corollary 4.9** *Under the above notation, for all  $N \geq m$ , we have*

$$\begin{aligned} N \frac{d^2}{dt^2} S_N(\rho) &= \int_M \left[ \left( \frac{1}{m} - \frac{1}{N} \right) |L\phi|^2 + Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \right] \rho^{1-1/N} d\mu \\ &\quad + \int_M \left[ \left( \frac{1}{n} - \frac{1}{m} \right) \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 + \left\| \nabla^2\phi - \frac{\Delta\phi}{n} g \right\|^2 \right] \rho^{1-1/N} d\mu. \end{aligned}$$

*Assume that  $Ric_{m,n}(L) \geq K$  for some  $K \in C(M, \mathbb{R})$ . Then, for all  $N \geq m$ , we have*

$$\frac{d^2}{dt^2} S_N(\rho(t)) + \frac{N-m}{m} [S_N(\rho(t))]^{-1} \left( \frac{d}{dt} S_N(\rho(t)) \right)^2 \geq \frac{1}{N} \int_M K |\nabla\phi|^2 \rho^{1-1/N} d\mu.$$

*In particular, for all  $N \geq m$ , we have*

$$\frac{d^2}{dt^2} S_N(\rho(t)) \geq \frac{1}{N} \int_M K |\nabla\phi|^2 \rho^{1-1/N} d\mu. \quad (4.99)$$

We will see in Section 5 that (4.99) is the differential form of (1.3). Hence (vi) in Theorem 2.2 implies (viii).

#### 4.6 Limiting case $N \rightarrow \infty$ and $p \rightarrow 1$

Letting  $N \rightarrow \infty$  in (4.99), and using

$$\lim_{N \rightarrow \infty} N[1 + S_N(\rho)] = \int_M \rho \log \rho d\mu,$$

we can derive

$$\frac{d^2}{dt^2} \int_M \rho d\rho d\mu \geq K \int_M |\nabla\phi|^2 d\mu.$$

i.e.,

$$\frac{d^2}{dt^2} \text{Ent}(\rho(t)) \geq K W_2^2(\rho(0), \rho(1)),$$

That is to say, under  $CD(K, \infty)$ -condition, we have

$$\text{Hess Ent}(\rho) \geq K.$$

## 5 Comparison with Lott-Villani-Sturm's synthetic geometry

The aim of this section is to compare the main results of this paper with the works of Lott-Villani [33], Sturm [44, 45] and Villani [50] in the setting of Riemannian manifolds with  $N = m$ .

### 5.1 Lott-Villani and Sturm

For  $N < \infty$ , Sturm [45] proved that in the Riemannian case the curvature-dimension  $CD(0, N)$  for  $N \geq n = \dim M$  holds if and only if for all  $N' \geq N$  the quantity  $S_{N'}(\cdot|m)$  is

convex on the Wasserstein space  $P_2(M, d)$ . This is basically due to the fact that the Jacobian determinant  $J_t = \det dF_t$  of any optimal transport map  $F_t = \exp(-t\nabla\phi) : M \rightarrow M$  satisfies

$$\frac{\partial^2}{\partial t^2} J_t^{1/N} \leq 0 \quad (5.100)$$

if and only if  $N \geq n = \dim M$  and  $Ric \geq 0$ . As pointed out in [45], (5.100) is essentially equivalent to the Brunn-Minkowski inequality

$$m(A_t)^{1/N'} \geq (1-t)m(A_0)^{1/N'} + tm(A_1)^{1/N'}. \quad (5.101)$$

for any  $N' \geq N$ , any  $t \in [0, 1]$  and any pair of Borel sets  $A_0, A_1 \subset M$ , where  $A_t$  denotes the set of points  $\gamma_t$  on the geodesics with endpoints  $\gamma_0 \in A_0$  and  $\gamma_1 \in A_1$ .

The curvature-dimension condition  $CD(K, N)$  for general  $K \in \mathbb{R}$  and  $N < \infty$  is more involved. In this case, the inequality (5.100) is replaced by

$$\frac{\partial^2}{\partial t^2} J_t^{1/N}(x) \leq -\frac{K}{N} J_t^{1/N}(x) d^2(x, F_1(x)). \quad (5.102)$$

When  $\mu = \nu$  is the standard volume measure, Sturm [45] proved that (5.102) is equivalent to (1.3) if and only if  $N \geq n = \dim$  and  $Ric \geq K$ . In [45], Sturm used (1.3) to introduce the definition of the  $CD(K, N)$ -condition on metric measure spaces. Independent of Sturm [45], Lott and Villani [33] introduced the curvature-dimension condition  $CD(0, N)$  on metric measure spaces in the same form as (1.2).

By Sturm [45], assume for simplicity that for  $m \otimes m$  -a.e.  $(x, y) \in M^2$  there exists a unique geodesic  $t \rightarrow \gamma_t(x, y)$  depending in a measurable way on the endpoints  $x$  and  $y$ . Then  $CD(K, N)$  states that that for any pair of absolutely continuous probability measures  $\rho_0 m$  and  $\rho_1 m$  on  $M$  there exists an optimal coupling  $q$  such that

$$\rho_t^{-1/N}(\gamma_t(x, y)) \leq \tau_{K, N}^{(1-t)}(d(x, y)) \rho_0^{-1/N}(x) + \tau_{K, N}^{(t)}(d(x, y)) \rho_1^{-1/N}(y) \quad (5.103)$$

for all  $t \in [0, 1]$  and  $q$ -a.e.  $(x, y) \in M^2$ , where  $\rho_t$  is the density of the push-forward of  $q$  under the map  $(x, y) \rightarrow \gamma_t(x, y)$ . Indeed,  $\rho_t m = (F_t)_*(\rho_0 m)$ . More precisely, we have

$$\rho_0(x) = \rho_t(F_t(x)) \det(dF_t(x)) = \rho_t(F_t(x)) J_t.$$

Equivalently

$$\rho_t(F_t(x)) = \rho_0(x) J_t^{-1}.$$

Hence

$$\begin{aligned} \frac{d^2}{dt^2} \rho_t^{-1/N}(F_t(x)) &= \rho_0^{-1/N}(x) \frac{d^2}{dt^2} J_t^{1/N} \\ &\leq \frac{K d^2(x, F_1(x))}{N} \rho_0^{-1/N}(x) J_t^{1/N}. \end{aligned}$$

That is to say

$$\frac{d^2}{dt^2} \rho_t^{-1/N}(F_t(x)) \leq \frac{K d^2(x, F_1(x))}{N} \rho_t^{-1/N}(F_t(x)). \quad (5.104)$$

Integrating in  $t$ , (5.104) implies (5.103).

Moreover, one can show that (5.104) is equivalent to (1.3). Indeed, if (5.104) hold, then

$$\begin{aligned}
\frac{d^2}{dt^2} S_N(\rho(t)) &= -\frac{d^2}{dt^2} \int_M \rho_t^{-1/N}(F_t(x)) \rho_0(x) d\mu(x) \\
&= -\int_M \frac{d^2}{dt^2} \rho_t^{-1/N}(F_t(x)) \rho_0(x) d\mu(x) \\
&\leq -\frac{1}{N} \int_M K(x) d^2(x, F_1(x)) \rho_t^{-1/N}(F_t(x)) \rho_0(x) d\mu(x).
\end{aligned}$$

Hence

$$\frac{d^2}{dt^2} S_N(\rho(t)) \leq -\frac{1}{N} \int_M K(x) d^2(x, F_1(x)) \rho_t^{1-1/N}(F_t(x)) J_t(x) d\mu(x).$$

When  $K$  is a constant, we have

$$\frac{d^2}{dt^2} S_N(\rho(t)) \leq -\frac{K}{N} \int_M d^2(x, F_1(x)) \rho_t^{1-1/N}(F_t(x)) J_t(x) d\mu(x). \quad (5.105)$$

On the other hand, if (5.105) holds for all  $\rho_0(x)$ , we can derive that (5.104) also holds.

Now we prove that

$$\int_M |\nabla \phi_t(x)|^2 \rho_t^{1-1/N}(x) d\mu(x) = \int_M d^2(x, F_1(x)) \rho_t^{1-1/N}(F_t(x)) J_t(x) d\mu(x). \quad (5.106)$$

Indeed

$$\begin{aligned}
\int_M |\nabla \phi_t(x)|^2 \rho_t^{1-1/N}(x) d\mu(x) &= \int_M |\nabla \phi_t(y)|^2 \rho_t^{1-1/N}(y) d\mu(y) \\
&= \int_M |\nabla \phi_t(F_t(x))|^2 \rho_t^{1-1/N}(F_t(x)) \det(dF_t(x)) d\mu(x) \\
&= \int_M |\nabla \phi_t(F_t(x))|^2 \rho_t^{-1/N}(F_t(x)) \rho_0(x) d\mu(x).
\end{aligned}$$

Note that  $\nabla \phi_t(F_t(x)) = \frac{d}{dt} F_t(x) = \dot{F}_t(x)$ , and  $F_t(x) = \exp_x(-t\nabla \phi(x))$  is a geodesic linking  $F_0(x) = x$  and  $F_1(x)$ . Hence

$$|\nabla \phi_t(F_t)|^2 = |\dot{F}_t(x)|^2 = |\dot{F}_1(x)|^2 = d^2(x, F_1(x)).$$

This proves (5.106). As a consequence, (2.25) for  $S_N(\rho(t))$  is the same as (5.105), which is equivalent to Sturm's definition inequality (1.3). This proves (vii) in Theorem 2.2 implies (viii) there.

Next we briefly describe the works of Lott-Villani [33] and Villani [50] on the characterization of  $CD(K, N)$ -condition by  $\mathcal{DC}_N$  class of functional. In [33, 50], Lott and Villani introduced the class  $\mathcal{DC}_N$  of general functional  $U(\rho) = \int_M e(\rho) d\mu$  on the Wasserstein space  $P_2(M)$ . By Definition 17.1 in [50], the class  $\mathcal{DC}_N$  is defined as the set of continuous convex function  $e : \mathbb{R}^+ \rightarrow \mathbb{R}$  which is  $C^2$ -differentiable on  $(0, \infty)$ ,  $e(0) = 0$  and satisfies one of the following equivalent conditions holds:

- The pressure function  $p_1(r) = re'(r) - e(r)$  satisfies

$$rp_1'(r) \geq \left(1 - \frac{1}{N}\right) p_1(r).$$

Equivalently

$$p_2(r) + \frac{1}{N}p_1(r) \geq 0.$$

- The function  $r \mapsto \frac{p_1(r)}{r^{1-\frac{1}{N}}}$  is convex.
- The function  $\delta \rightarrow u(\delta) = \delta^N e(\delta^{-N})$  is convex.

By Theorem 17.15 in [50],  $M$  satisfies the curvature-dimension condition  $\text{CD}(K, N)$  if and only if for each  $e \in \mathcal{DC}_N$ , the functional  $U(\rho) = \int_M e(\rho) d\mu$  satisfies

$$U(\mu_t) + K_{N,U} \int_0^1 \int_M \rho_s(x)^{1-\frac{1}{N}} |\nabla \phi_s(x)|^2 d\mu(x) G(s, t) ds \leq (1-t)U(\mu_0) + tU(\mu_1). \quad (5.107)$$

Here

$$K_{N,U} = \inf_{r>0} \frac{Kp_1(r)}{r^{1-1/N}}.$$

Indeed, it was proved in [50] that for any  $e \in \mathcal{DC}_N$ , it holds

$$\frac{d^2}{dt^2} U(\mu_t) \geq K_{N,U} \int_M |\nabla \phi_t(x)|^2 \rho_t(x)^{1-1/N} d\mu(x). \quad (5.108)$$

which is equivalent to (5.107) by Proposition 16.2 in Villani [50].

Now we compare Theorem 4.2 and Lott-Villani's inequalities (5.107) and (5.108). Note that, by (4.80) in Theorem 4.2, if  $p_2 + \frac{1}{N}p_1 \geq \sigma p_1$  for some constant  $\sigma \geq 0$ , it holds

$$U''(\rho_t) \geq \sigma [U'(\rho_t)]^2 \left( \int_M p_1(\rho_t(x)) d\mu(x) \right)^{-1} + \int_M K(x) |\nabla \phi_t(x)|^2 p_1(\rho_t(x)) d\mu(x).$$

In particular, for  $e \in \mathcal{DC}_N$ , we have  $\sigma = 0$ . Hence

$$U''(\rho_t) \geq \int_M K(x) |\nabla \phi_t(x)|^2 p_1(\rho_t(x)) d\mu(x).$$

Under Lott-Villani's assumption  $Kp_1(\rho) \geq K_{N,U}\rho^{1-1/N}$ , (5.108) follows.

**Remark 5.1** *Similarly to Lott-Villani [33] and Villani [50], we may introduce the class  $\mathcal{DC}_{N,\sigma}$  of functionals  $U(\rho) = \int_M e(\rho) d\mu$  such that  $p_2(r) + \frac{1}{N}p_1(r) \geq \sigma p_1(r)$  for a constant  $\sigma \geq 0$ . Note that  $\mathcal{DC}_{N,\sigma_2} \subset \mathcal{DC}_{N,\sigma_1}$  if  $\sigma_2 \geq \sigma_1 \geq 0$ , and  $e(r) = \frac{r^p}{p-1} \in \mathcal{DC}_{N,\sigma}$  with  $\sigma = p - 1 + \frac{1}{N}$ .*

## 5.2 Comparison

Now we compare our entropy power differential inequalities in Theorem 2.2 with the above mentioned results due to Sturm [45], Lott-Villani [33] and Villani [50].

Note that our entropy power inequality

$$\frac{d^2}{dt^2} N_m(\rho(t)) \leq -\frac{K}{m} N_m(\rho(t)) W_2^2(\rho(0), \rho(1)) \quad (5.109)$$

is equivalent to the entropy differential inequality

$$H'' + \frac{H'^2}{m} \leq K \int_M |\nabla\phi|^2 \rho d\mu, \quad (5.110)$$

where  $(\rho(t), t \in [0, 1])$  is a geodesic linking  $\rho(0)$  and  $\rho(1)$  on the Wasserstein space  $P_2(M, g, \mu)$ ,  $H' = \frac{d}{dt}H(\rho(t))$  and  $H'' = \frac{d^2}{dt^2}H(\rho(t))$ .

From the proof of the NIW formula, we have

$$\begin{aligned} H'' + \frac{H'^2}{m} &= -\frac{1}{m} \int_M \left| L\phi - \int_M L\phi \rho d\mu \right|^2 \rho d\mu - \int_M Ric_{m,n}(L)(\nabla\phi, \nabla\phi) \rho d\mu \\ &\quad - \frac{m-n}{mn} \int_M \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 \rho d\mu - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \rho d\mu. \end{aligned} \quad (5.111)$$

Under the assumption  $Ric_{m,n}(L) \geq K$ , we have

$$H'' + \frac{H'^2}{m} \leq -K \int_M |\nabla\phi|^2 \rho d\mu = -KW_2^2(\rho_0, \rho_1). \quad (5.112)$$

Equivalently, we have

$$N_m'' \leq -\frac{KN_m}{m} W_2^2(\rho_0, \rho_1). \quad (5.113)$$

In particular, when  $m = n$ , it holds

$$\begin{aligned} H'' + \frac{H'^2}{n} &= -\frac{1}{n} \int_M \left| \Delta\phi - \int_M \Delta\phi \rho d\mu \right|^2 \rho d\nu - \int_M Ric(\nabla\phi, \nabla\phi) \rho d\nu \\ &\quad - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 \rho d\nu \\ &= -\int_M \left[ Ric(\nabla\phi, \nabla\phi) + \left\| \nabla^2\phi - \frac{\int_M \Delta\phi \rho d\nu}{n}g \right\|_{\text{HS}}^2 \right] \rho d\nu. \end{aligned} \quad (5.114)$$

Hence, under the condition  $Ric \geq K$ , we have

$$H'' + \frac{H'^2}{n} \leq -K \int_M |\nabla\phi|^2 \rho d\nu = -KW_2^2(\rho(0), \rho(1)). \quad (5.115)$$

Equivalently, we have

$$N_n'' \leq -\frac{KN_n}{n} W_2^2(\rho_0, \rho_1). \quad (5.116)$$

Conversely, we now prove that, if (5.115) or (5.116) holds for any geodesic on the Wasserstein space  $P_2(M, v)$  over  $(M, g)$ , then we have

$$Ric \geq K.$$

Indeed, (5.114) holds at  $t = 0$  for the solution  $(\rho(t), \phi(t))$  of the continuity equation and the Hamilton-Jacobi equation with any initial data  $(\rho(0), \phi(0))$ . Taking a sequence of  $\rho_k(0)d\nu$  converges weakly to the Dirac mass at  $x_0 \in M$ , we derive that at  $t = 0$



$$H'' + \frac{H'^2}{n} = - \left[ Ric_{x_0}(v, v) + \left\| \nabla^2 \phi(x_0) - \frac{\Delta \phi(x_0)}{n} g \right\|_{\text{HS}}^2 \right].$$

For any fixed  $v \in T_{x_0}M$ , choose  $\phi \in C^2(M)$  such that

$$\begin{aligned} \nabla \phi(x_0) &= v, \\ \nabla^2 \phi(x_0) &= \frac{\Delta \phi(x_0)}{n} I_n. \end{aligned}$$

This implies that

$$H'' + \frac{H'^2}{n} = -Ric_{x_0}(v, v).$$

Therefore, if (5.115) holds, we have

$$-Ric_{x_0}(v, v) = H'' + \frac{H'^2}{n} \leq -K|v|^2.$$

That is to say  $Ric \geq K$ . Moreover, we can conclude that the equality holds in (5.115), equivalently,

$$N_n'' = -\frac{KN_n}{n} W^2(\rho_0, \rho_1)$$

holds for any  $\rho_0$  and  $\rho_1$  in the Wasserstein space  $P_2(M, g, \nu)$ , if and only if  $(M, g)$  is Einstein, i.e.,  $Ric = Kg$  and  $\phi$  is a Hessian soliton

$$\nabla^2 \phi = \frac{\Delta \phi}{n} g.$$

In the case  $m > n$ , we can use the same argument as above to prove that the inequality (5.112) or (5.113) holds if and only if the  $CD(K, m)$ -condition holds on  $(M, g, \mu)$ , i.e.,  $Ric_{m,n}(L) \geq Kg$ . Moreover, we can prove that the equality holds in (5.112) or (5.113) if and only if  $(M, g)$  is  $(K, m)$ -Einstein, i.e.,  $Ric_{m,n}(L) = Kg$  and  $\phi$  is a Hessian soliton

$$\nabla^2 \phi = \frac{\Delta \phi}{n} g,$$

and

$$\frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 = 0.$$

In general case of  $p \neq 1$ , recall that

$$\begin{aligned} H_p'' + \frac{1}{m} H_p'^2 &= \left( p-1 + \frac{1}{m} \right) \left[ \left( \int_M L\phi d\gamma \right)^2 - \int_M |L\phi|^2 d\gamma \right] - \int_M Ric_{m,n}(L)(\nabla \phi, \nabla \phi) d\gamma \\ &\quad - \int_M \left[ \frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla \phi \right)^2 + \left\| \nabla^2 \phi - \frac{\Delta \phi}{n} g \right\|_{\text{HS}}^2 \right] d\gamma. \end{aligned} \tag{5.117}$$

In particular, when  $m = n$ ,

$$\begin{aligned} H_p'' + \frac{1}{n}H_p'^2 &= \left(p - 1 + \frac{1}{n}\right) \left[ \left(\int_M \Delta\phi d\gamma\right)^2 - \int_M |\Delta\phi|^2 d\gamma \right] - \int_M Ric(\nabla\phi, \nabla\phi) d\gamma \\ &\quad - \int_M \left\| \nabla^2\phi - \frac{\Delta\phi}{n}g \right\|_{\text{HS}}^2 d\gamma. \end{aligned} \quad (5.118)$$

This yields, if  $Ric \geq K$ , we have

$$H_p'' + \frac{1}{n}H_p'^2 \leq -K \int_M |\nabla\phi|^2 d\gamma. \quad (5.119)$$

By the fact

$$\frac{d^2}{dt^2}N_{n,p}(\rho(t)) = \left(\frac{1}{n}H_p'' + \frac{1}{n^2}H_p'^2\right) N_{n,p}(\rho(t)),$$

we see that (5.119) is equivalent to

$$\frac{d^2}{dt^2}N_{n,p}(\rho(t)) \leq \left(-\frac{1}{n}K \int_M |\nabla\phi|^2 d\gamma\right) N_{n,p}(\rho(t)). \quad (5.120)$$

Conversely, we can prove that if (5.119) or (5.120) holds, then

$$Ric \geq K.$$

For this, notice that (5.118) holds at  $t = 0$  for the solution  $(\rho(t), \phi(t))$  of the continuity equation and the Hamilton-Jacobi equation with any initial data  $(\rho(0), \phi(0))$ . Taking a sequence of  $d\gamma_k(0) = \frac{\rho_k(0)^p d\nu}{\int_M \rho_k(0)^p d\nu}$  converges weakly to the Dirac mass at  $x_0 \in M$ , we derive that at  $t = 0$

$$H_p'' + \frac{1}{n}H_p'^2 = - \left[ Ric_{x_0}(v, v) + \left\| \nabla^2\phi(x_0) - \frac{\Delta\phi(x_0)}{n}g \right\|_{\text{HS}}^2 \right].$$

For any fixed  $v \in T_{x_0}M$ , choose  $\phi$  such that

$$\begin{aligned} \nabla\phi(x_0) &= v, \\ \nabla^2\phi(x_0) &= \frac{\Delta\phi(x_0)}{n}I_n. \end{aligned}$$

This implies that

$$H_p'' + \frac{1}{n}H_p'^2 = -Ric_{x_0}(v, v).$$

Therefore, if (5.119) or (5.120) holds, we have

$$Ric_{x_0}(v, v) = -H_p'' - \frac{1}{n}H_p'^2 \geq K|v|^2.$$

That is to say  $Ric \geq K$ . Moreover, we can conclude that the equality holds in (5.119), i.e.,

$$H_p'' + \frac{1}{n}H_p'^2 = -K \int_M |\nabla\phi|^2 d\gamma, \quad (5.121)$$

or equivalently, the equality holds in (5.120), i.e.,

$$\frac{d^2}{dt^2}N_{n,p}(\rho(t)) = \left(-\frac{K}{n} \int_M |\nabla\phi|^2 d\gamma\right) N_{n,p}(\rho(t)) \quad (5.122)$$

holds for any  $\rho_0$  and  $\rho_1$  in the Wasserstein space  $P_2(M, g, \nu)$ , if and only if  $(M, g)$  is Einstein, i.e.,  $Ric = Kg$  and  $\phi$  is a Hessian soliton

$$\nabla^2\phi = \frac{\Delta\phi}{n}g.$$

Similarly, we can prove that the inequality

$$H_p'' + \frac{1}{m}H_p'^2 \leq -K \int_M |\nabla\phi|^2 d\gamma, \quad (5.123)$$

equivalently,

$$\frac{d^2}{dt^2}N_{m,p}(\rho(t)) \leq \left(-\frac{K}{m} \int_M |\nabla\phi|^2 d\gamma\right) N_{m,p}(\rho(t)) \quad (5.124)$$

holds for any  $\rho_0$  and  $\rho_1$  in the Wasserstein space  $P_2(M, g, \mu)$  if and only if the  $CD(K, m)$ -condition holds on  $(M, g, \mu)$ , i.e.,  $Ric_{m,n}(L) \geq Kg$ . Moreover, we can prove that the equality holds in (5.123) (equivalently, (5.124)) if and only if  $(M, g)$  is  $(K, m)$ -Einstein, i.e.,  $Ric_{m,n}(L) = Kg$  and  $\phi$  is a Hessian soliton

$$\nabla^2\phi = \frac{\Delta\phi}{n}g,$$

and

$$\frac{m-n}{mn} \left( L\phi + \frac{m}{m-n} \nabla V \cdot \nabla\phi \right)^2 = 0.$$

We have therefore finished the proof of the rigidity part of Theorem 2.2.

### 5.3 Summary and further problems

Theorem 2.1 and Theorem 2.2 indicate the equivalence between the curvature-dimension  $CD(K, m)$ -condition, entropy differential inequalities (EDI) and entropy power differential inequalities (EPDI) along the geodesics on the Wasserstein space over Riemannian manifolds. In particular, each of the equivalent conditions (ii)-(vii) in Theorem 2.2 is equivalent to the definition inequality (1.3) (for  $N' \geq N = m$ ) which was used by Sturm [45] to define the  $CD(K, N)$ -condition on metric measure spaces. They are also equivalent to Lott-Villani's characterization of  $CD(K, N)$  condition using functionals in the class  $\mathcal{DC}_N$ . Therefore, we can use each of (ii)-(vii) or its integratral forms to introduce definition of  $CD(K, N)$ -condition on metric measure spaces. In particular, all these equivalent conditions are stable under the Gromov-Hausdorff convergence on metric measure spaces.

Theorem 2.1 and Theorem 2.2 also indicate that the rigidity models for the EDI and EPDI along all smooth geodesics on the Wasserstein space over Riemannian manifolds are  $K$ -Einstein manifolds or  $(K, m)$ -Einstein manifolds with Hessian solitons. The proofs of rigidity theorems need to use the Bochner formula (2.11) rather than the Bochner inequality (4.82).

Theorem 2.3 extends the monotonicity and rigidity theorems of the  $W$ -entropy associated with the Shannon entropy [25, 27] to the Rényi entropy for the geodesics on the Wasserstein space over complete Riemannian manifolds with  $CD(0, m)$ -condition.

In [14], Erbar-Kuwada-Sturm proved that the Shannon entropy power  $\exp(-\frac{2}{N}\text{Ent}(\rho))$  is concave along the heat equation on  $\text{RCD}(0, N)$  metric measure spaces. In [22], K. Kuwada and the author of this paper proved the monotonicity of the  $W$ -entropy for heat equation associated with the Witten Laplacian on  $\text{RCD}(0, N)$  metric measure spaces. Moreover, a rigidity theorem in a weak form was also proved for the  $W$ -entropy for heat equation on  $\text{RCD}(0, N)$  metric measure spaces.

It is interesting to raise some problems for further study in the future.

**Problem 5.2** (i) *Can we extend Theorem 2.1, Theorem 2.2 and Theorem 2.3 (in particular, the rigidity theorems) to geodesics on the Wasserstein space over  $\text{RCD}(0, N)$  or  $\text{RCD}(K, N)$  metric measure spaces?*

(ii) *Can we extend Theorem 2.1, Theorem 2.2 and Theorem 2.3 (in particular, the rigidity theorems) to geodesics on the Wasserstein space over  $\text{RCD}$  metric measure spaces with time-dependent metrics and measures?*

(iii) *In particular, can we extend EDI, EPDI and the  $W$ -entropy formula to geodesics on the Wasserstein space over metric measure spaces with  $(K, N)$ -super Ricci flows on non-smooth metric measure spaces introduced by Kopfer and Sturm [21] and Sturm [46].*

To end this paper, let us mention that in our forthcoming works in preparation, we will extend EDI, EPDI and the  $W$ -entropy formula to the Langevin deformation of flows on the Wasserstein space, which was introduced by S. Li and the author [25, 27] and can be regarded as an interpolation between the geodesics and the gradient flows of the Shannon or Rényi entropy on the Wasserstein space over a Riemannian manifold. We will also study EDI, EPDI and the  $W$ -entropy formula along geodesics on the Wasserstein space over manifolds equipped with time dependent metrics and potentials, in particular, to extend EDI, EPDI and the  $W$ -entropy formula along geodesics on the Wasserstein space over manifolds with Perelman's Ricci flow coupled with the conjugate heat equation.

**Acknowledgement.** The author would like to thank Prof. N. Mok for suggestion, Prof. B-X. Han and Dr. S. Li for helpful discussions, and Dr. R. Lei for pointing out some misprints in the earlier version of this paper.

## References

- [1] L. Ambrosio, N. Gigli, A User's Guide to Optimal Transport Theory, CIME Lecture Notes in Mathematics, Springer, 2012.
- [2] D. Bakry, M. Emery, Diffusion hypercontractives, Sémin. Prob. XIX, Lect. Notes in Maths. 1123 (1985), 177-206.
- [3] D. Bakry, Y. Gentil, M. Ledoux, Analyse and Geometry for Markov Diffusion Operators, Springer, 2014.
- [4] N.M. Blachman, The convolution inequality for entropy powers. IEEE Trans. Inform. Theory 2, 267-271, (1965).
- [5] J.-D. Benamou, Y. Brenier, A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem. Numer. Math. 84, 375-393 (2000)
- [6] H.-D. Cao, X.-P. Zhu, A complete proof of the Poincaré and geometrization conjectures: application of the Hamilton-Perelman theory of the Ricci flow, Asian J. Math. 10 (2006), no. 2, 165-492, and Asian J. Math. 10 (2006), no. 4, 663.
- [7] B. Chow, P. Lu, L. Ni, Hamilton's Ricci flow, Lectures in Contemporary Mathematics, 3, Amer. Math. Soc. Providence, Rhode Island, Beijing, 2006.

- [8] B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo and L. Ni, *The Ricci flow: Techniques and Applications, Part I, Geometric aspects*, Mathematical Surveys and Monographs, 135. American Mathematical Society, Providence, RI, 2007.
- [9] D. Cordero-Erausquin, R.J. McCann, and M. Schmuckenschläger, A Riemannian interpolation inequality à la Borell, Brascamp and Lieb. *Invent. Math.* 146, 2 (2001), 219-257.
- [10] M.H.M. Costa, A new entropy power inequality, *IEEE Trans. Inf. Theory*, IT- 31, (6) 751-760, (1985).
- [11] T. M. Cover, J.A. Thomas, *Elements of Information Theory*, Wiley-Interscines, A John Wiely and Sons, INC., Publication.
- [12] A. Dembo, A simple proof of the concavity of the entropy power with respect to the variance of additive normal noise, *IEEE Trans. Inform. Theory* 35, 887-888, (1989).
- [13] A. Dembo, T.M. Cover, and J.A. Thomas, Information theoretic inequalities, *IEEE Trans. Inf. Theory*, 37, (6), 1501-1518, (1991).
- [14] M. Erbar, K. Kuwada, K.-T. Sturm, On the equivalence of the entropic curvature-dimension condition and Bochner's inequality on metric measure spaces, *Invent. Math.* 201 (2015), no. 3, 993–1071.
- [15] F. Fang, X.-D. Li and Z. Zhang, Two generalizations of the Cheeger-Gromoll splitting theorem via the Bakry-Emery Ricci curvature. *Ann. Inst. Fourier* 59(2) (2009), 563–573.
- [16] T. Kato, The Cauchy problem for quasi-linear symmetric hyperbolic systems, *Arch Rational Mech Anal.* 58 (1975), 181–205.
- [17] R. E. Greene and H. Wu,  $C^\infty$ -convex functions and manifolds of positive curvature, *Acta Math.* 137 (1976), no. 3-4, 209–245.
- [18] R.S. Hamilton, Three manifolds with positive Ricci curvature. *J. Diff. Geom.* 17 (1982), 255-306.
- [19] R.S. Hamilton, *The formation of singularities in the Ricci flow*, *Surveys in Differential Geometry*, 2, 7-136, International Press, 1995.
- [20] B. Kleiner, J. Lott, Notes on Perelman's papers, *Geom. Topol.* 12 (2008),no. 5, 2587-2855.
- [21] E. Kopfer, K.-Th. Sturm, Heat flow on time-dependent metric measure spaces and super-Ricci flows. *Comm. Pure Appl. Math.* 71 (2018), no. 12, 2500-2608.
- [22] K. Kuwada, X.-D. Li, Monotonicity and rigidity of the  $W$ -entropy on  $RCD(0;N)$  spaces, *Manuscripta Math.* 164 (2021), 119-149.
- [23] S. Li, X.-D. Li,  $W$ -entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials, *Pacific J. Math.* Vol. 278 (2015), No. 1, 173-199.
- [24] S. Li and X.-D. Li, Hamilton differential Harnack inequality and  $W$ -entropy for Witten Laplacian on Riemannian manifolds, *J. Funct. Anal.* 274 (2018), 3263-3290.
- [25] S. Li, X.-D. Li,  $W$ -entropy formulas on super Ricci flows and Langevin deformation on Wasserstein space over Riemannian manifolds. *Sci China Math*, 2018, 61: 1385-1406.

- [26] S. Li and X.-D. Li,  $W$ -entropy, super Perelman Ricci flows and  $(K, m)$ -Ricci solitons, *J. Geom. Anal.* 30 (2020), no. 3, 3149–3180.
- [27] S. Li, X.-D. Li,  $W$ -entropy formulas and Langevin deformation of flows on Wasserstein space over Riemannian manifolds. *Probab. Theory and Related Fields*, 188 (2024), 911-955, <https://doi.org/10.1007/s00440-023-01256-y>
- [28] S. Li, X.-D. Li, On the Shannon entropy power on Riemannian manifolds and Ricci flows, arxiv2001.00410v1, 2020. To appear in *Tohoku Math. J.* 2024.
- [29] S. Li, X.-D. Li, On the Rényi entropy power and the Gagliardo-Nirenberg-Sobolev inequality on Riemannian manifolds, arXiv:2001.11184v1, 2020.
- [30] X.-D. Li, Perelman’s entropy formula for the Witten Laplacian on Riemannian manifolds via Bakry-Emery Ricci curvature, *Math. Ann.* 353 (2012), 403–437.
- [31] J. Lott, Some geometric calculations on Wasserstein space, *Comm. Math. Phys.* 277 (2008), no. 2, 423-437.
- [32] J. Lott, Optimal transport and Perelman’s reduced volume, *Calc. Var. and Partial Differential Equations* 36, 49-84 (2009).
- [33] J. Lott, C. Villani, Ricci curvature for metric-measure spaces via optimal transport, *Annals of Math.* 169, 903-991, 2009.
- [34] F. Otto, The geometry of dissipative evolution equations: the porous medium equation, *Commun. in Parial Differential Equations* 26 (1 and 2), 101-174 (2001).
- [35] R. McCann, Polar factorization of maps on Riemannian manifolds, *Geometric and Functional Analysis*, 11, 3(2001), 589-608.
- [36] J. W. Morgan, G. Tian, Ricci flow and the Poincaré conjecture, *Clay Mathematics Monographs*, 3. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007.
- [37] L. Ni, The entropy formula for linear equation, *J. Geom. Anal.* 14 (1), 87-100, (2004).
- [38] L. Ni, Addenda to “The entropy formula for linear equation”, *J. Geom. Anal.* 14 (2), 329-334, (2004).
- [39] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, <http://arXiv.org/abs/math/0211159>.
- [40] G. Savaré, G. Toscani, The concavity of Rényi entropy power, *IEEE Transactions on Information Theory* Vol 60, Issue: 5, May 2014, 2687-2693. DOI: 10.1109/TIT.2014.2309341
- [41] C. E. Shannon, A mathematical theory of communication, *Bell Syst. Tech. J.*, vol. 27, pp. 623-656, Oct. 1948.
- [42] A. J. Stam, Some inequalities satisfied by the quantities of information of Fisher and Shannon, *Information and Control*, vol. 2, pp. 101-112, June 1959.
- [43] K. T. Sturm, Convex functionals of probability measures and nonlinear diffusions on manifolds, *J. Math. Pures Appl.* 84 (2005) 149-168.
- [44] K.T. Sturm, On the geometry of metric measure spaces. I, *Acta. Math.* 196 (2006), no. 1, 65–132.

- [45] K.T. Sturm, On the geometry of metric measure spaces. II, *Acta. Math.* **196** (2006), no. 1, 133–177.
- [46] K.-Th. Sturm, Super-Ricci flows for metric measure spaces. *J. Funct. Anal.* **275** (2018), no. 12, 3504–3569.
- [47] K.-T. Sturm, M.-K. von Renesse, Transport inequalities, gradient estimates, entropy and Ricci curvature. *Commun. Pure Appl. Math.* **58**(7), 923-940 (2005)
- [48] C. Villani, A Short Proof of the Concavity of Entropy Power, *IEEE Transactions on Information Theory*, Vol. 46, No. 4, 1695-1696, July 2000.
- [49] C. Villani, *Topics in Mass Transportation*, Grad. Stud. Math., Amer. Math. Soc., Providence, RI, 2003.
- [50] C. Villani, *Optimal Transport, Old and New*, Springer, 2008.

Xiang-Dong Li, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, No. 55, Zhongguancun East Road, Beijing, 100190, China  
E-mail: xdli@amt.ac.cn

and

School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China