

LATTICE SUBSEQUENCES OF FIXED POINTS OF TOEPLITZ SUBSTITUTIONS

SHISHUANG LIU AND HUI RAO*

ABSTRACT. We define the modulo- m Toeplitz fixed point generated by Toeplitz substitution and study the lattice subsequence of such fixed point. Moreover, we provide a method to check whether one modulo- m Toeplitz fixed point is a lattice subsequence of another.

1. Introduction

Let \mathcal{A} be a finite set, and we call it an alphabet. Let $\mathcal{A}^* = \bigcup_{k \geq 0} \mathcal{A}^k$, where $\mathcal{A}^0 = \{\varepsilon\}$ and ε stands for the empty word. Let \mathcal{A}^∞ be the set of infinite sequence (or word) over \mathcal{A} . Let $\gamma\tau$ be the concatenation of words $\gamma \in \mathcal{A}^*$ and $\tau \in \mathcal{A}^* \cup \mathcal{A}^\infty$.

Let $m \geq 2$ and let $W = w_1 \dots w_{m-1}$ be a finite word over \mathcal{A} . We define the morphism σ over $\mathcal{A}^* \cup \mathcal{A}^\infty$ by

$$\sigma : a \mapsto Wa, \quad a \in \mathcal{A},$$

and call it a *Toeplitz substitution* (determined by W). Let w_1 be the initial letter of W , then

$$X = \lim_{k \rightarrow \infty} \sigma^k(w_1)$$

is called the *fixed point* of σ , and we call it a *modulo- m Toeplitz fixed point*. See [2, 3, 9].

Toeplitz fixed point is a very special type of Toeplitz words. Toeplitz words were first introduced by Jacobs and Keane in [5]. Jacobs and Keane [5] defined Toeplitz words and studied the ergodic properties related to Toeplitz words. Prodinger and Urbanek [8] and Sell [10] studied the combinatorial properties of Toeplitz words. Recently, Toeplitz words have gained increasing popularity as a model for quasicrystals, see for example [1]. Liu and Qu [6, 7] studied the spectral properties of Schrödinger operator with Toeplitz potential.

Definition 1.1. Let $X = \{X(j)\}_{j \geq 1}$ and $Y = \{Y(j)\}_{j \geq 1}$ be two sequences. Let $q \geq 1$ be an integer. We say Y is a *q -subsequence* of X if $Y(j) = X(qj)$, $\forall j \geq 1$. In this case, we also say Y is a *lattice subsequence* of X .

In Huang, Liu and Rao [4], to determine when two fractal sets are bi-Lipschitz equivalent, they arise the following question:

Question 1. *Let X and Y be two modulo- m Toeplitz fixed points. How to judge whether Y is a q -subsequence of X ?*

The main goal of the present paper is to give an answer to the above question. From now on, we shall use $X(q\mathbb{N})$ to denote the q -subsequence of X . The following lemma is obvious.

Date: July 29, 2024.

Key words and phrases: Toeplitz fixed point, lattice subsequence.

* Corresponding author.

Lemma 1.1. *Let X and Y be two modulo- m Toeplitz fixed points. Then Y is a q -subsequence of X if and only if $X(q\mathbb{N})$ is a modulo- m Toeplitz fixed point and $Y(j) = X(qj)$ for $1 \leq j \leq m-1$.*

Therefore Question 1 is converted to the following question:

Question 2. *Let X be a modulo- m Toeplitz fixed point. For which q , $X(q\mathbb{N})$ is also a modulo- m Toeplitz fixed point?*

It is easy to show that

Lemma 1.2. *Let $X = \{X(j)\}_{j \geq 1}$ be a modulo- m Toeplitz fixed point. Let $h \geq 1$ be an integer such that $\gcd\{m, h\} = 1$. Let $q \geq 1$ be an integer. Then $X(hq\mathbb{N})$ is a modulo- m Toeplitz fixed point if and only if $X(q\mathbb{N})$ is a modulo- m Toeplitz fixed point.*

Notice that $X(mq\mathbb{N}) = X(q\mathbb{N})$ (see Lemma 2.1). Therefore, to characterize all lattice subsequences of a modulo- m Toeplitz fixed point, we just need to check the q -subsequence such that $q|m^s$ for some $s \geq 1$ and $m \nmid q$.

Let $W = w_1 \dots w_n$ be a word and let $2 \leq q < n$. We say W is *almost q -periodic*, if $q \nmid j$ implies $w_j = w_{j+q}$. Similarly, if W is an infinite word, we define W is almost q -periodic in the same manner.

Let a^k be the constant word with length k and with letter a . For each word ω we use $\omega(n)$ or ω_n to denote the letter at index n .

Theorem 1.1. *Let $X = \{X(j)\}_{j \geq 1}$ be a modulo- m Toeplitz fixed point which is not a constant word. Let $q \geq 2$ be an integer such that $m \nmid q$ and $q|m^s$ for some $s \geq 1$. Then the following hold:*

- (i) *If $q \nmid m^2$, then $X(q\mathbb{N})$ is not a modulo- m Toeplitz fixed point.*
- (ii) *If $q|m$, then $X(q\mathbb{N})$ is a modulo- m Toeplitz fixed point if and only if $X(1) \dots X(m)$ is almost q -periodic.*
- (iii) *If $q|m^2$, then $X(q\mathbb{N})$ is a modulo- m Toeplitz fixed point if and only if $X(1) \dots X(m^2)$ is almost q -periodic.*

Remark 1.1. Let X and Y be two modulo- m Toeplitz fixed points. It is not hard to show that if $q|m^s$ and $Y = X(q\mathbb{N})$, then $X = Y(\frac{m^s}{q}\mathbb{N})$.

2. Lattice subsequences of Toeplitz words

We shall use \mathcal{J}_m to denote the collection of modulo- m Toeplitz fixed points (over the alphabet \mathcal{A}).

Lemma 2.1. *A word $X = \{X(j)\}_{j \geq 1}$ belongs to \mathcal{J}_m if and only if the following two conditions hold:*

- (i) *For any $j \geq 1$, $X(mj) = X(j)$;*
- (ii) *If $m \nmid j$, then $X(j) = X(j+m)$.*

Proof. Suppose $X \in \mathcal{J}_m$. Denote $a = X(1)$. Then $\sigma^n(a) = \sigma(\sigma^{n-1}(a))$, so $\sigma^n(a)$ is almost m -periodic. Therefore, X is almost m -periodic, which confirms (ii).

Since $\sigma(X(1) \dots X(j)) = \sigma(X(1))\sigma(X(2)) \dots \sigma(X(j))$, we conclude that $X(mj)$ is the last element of $\sigma(X(j)) = WX(j)$, so $X(mj) = X(j)$ and (i) is confirmed.

On the other hand, suppose that (i) and (ii) hold. Denote $W = X(1) \dots X(m-1)$. Then for any $k \geq 1$, we have

$$X(1) \dots X(m^k) = \prod_{j=1}^{m^{k-1}} WX(mj) = \prod_{j=1}^{m^{k-1}} \sigma(j) = \sigma(X(1) \dots X(m^{k-1})),$$

2

which implies that $X \in \mathcal{J}_m$. □

Proof of Lemma 1.2. Let $Y = X(hq\mathbb{N})$ and $Z = X(q\mathbb{N})$. Let $h \geq 2$.

Suppose $Z \in \mathcal{J}_m$. First, it is obvious that

$$Y(mj) = X(m(hqj)) = X(hqj) = Y(j).$$

Pick $j \geq 1$ such that $m \nmid j$. Since $\gcd\{m, h\} = 1$, we have $m \nmid hj$. Therefore,

$$Y(j) = Z(hj) = Z(hj + hm) = Z(h(j + m)) = Y(j + m).$$

By Lemma 2.1, we obtain that Y is a modulo- m Teoplitz fixed point.

Suppose $Y \in \mathcal{J}_m$. Let $p \geq 1$ be an integer such that $ph \equiv 1 \pmod{m}$. Then $\gcd\{p, m\} = 1$. By what we have just proved, $Y(p\mathbb{N}) \in \mathcal{J}_m$. Pick $j \geq 1$. Write $ph = ym + 1$ and $qj = m^t j'$ where $m \nmid j'$. Then

$$Y(pj) = X(phqj) = X((ym + 1)m^t j') = X(j') = X(m^t j') = Z(j).$$

Hence $Z = Y(p\mathbb{N}) \in \mathcal{J}_m$. □

Now we introduce words with question mark to study Teoplitz words. In other words, we study words over the alphabet $\mathcal{A} \cup \{?\}$.

For $U = u_1 \dots u_{r-1}?$ and $V = v_1 \dots v_{s-1}?$ where $u_j, v_i \in \mathcal{A}$ for $1 \leq j \leq r-1$, $1 \leq i \leq s-1$. We define *the composition of U and V* by

$$U \circ V = \prod_{j=1}^s (u_1 \dots u_{r-1} v_j),$$

where $v_s = ?$. Then all entries of $U \circ V$ belong to \mathcal{A} except the last entry. By this reason, we can define the *n -th iteration of U* by

$$(U)^{(1)} = U \text{ and } (U)^{(n)} = (U)^{(n-1)} \circ U, \quad \forall n \geq 2.$$

Clearly the operation ‘ \circ ’ is not commutative but it is associative, namely,

$$(U \circ V) \circ W = U \circ (V \circ W),$$

where $U = u_1 \dots u_{r-1}?$, $V = v_1 \dots v_{s-1}?$ and $W = w_1 \dots w_{t-1}?$. Moreover, for $\omega = (\omega_j)_{j=1}^\infty \in \mathcal{A}^\infty$, we define

$$U \circ \omega = \prod_{j=1}^\infty (u_1 \dots u_{r-1} \omega_j).$$

The following lemma is obvious.

Lemma 2.2. *If $X = \{X(j)\}_{j \geq 1}$ is a modulo- m Teoplitz fixed point. Then*

$$X = \lim_{n \rightarrow \infty} (X(1) \dots X(m-1)?)^{(n)} := [X(1) \dots X(m-1)?)^{(\infty)}.$$

Theorem 2.1. *Let $X = \{X(j)\}_{j \geq 1} \in \mathcal{J}_m$. Let $q \geq 2$ be an integer such that $m \nmid q$ and $q \mid m^s$ for some integer $s \geq 1$. Then the following three statements are equivalent:*

- (i) $X(q\mathbb{N}) \in \mathcal{J}_m$.
- (ii) X is almost q -periodic, in other words,

$$X = (X(1) \dots X(q-1)?) \circ X(q\mathbb{N}).$$

- (iii) $X(1) \dots X(m^s - 1) = (X(1) \dots X(q-1)?) \circ (X(q)X(2q) \dots X(m^s - q)?)$.

Proof. Let $Y = X(q\mathbb{N})$.

(i) \Rightarrow (ii). Since Y is a modulo- m Toeplitz word, we have

$$Y(j) = Y(j + tm^s), \quad \forall m \nmid j, t \in \mathbb{N},$$

therefore,

$$Y(jm) = Y(jm + tm^{s+1}), \quad \forall m \nmid j, t \in \mathbb{N},$$

which implies

$$X(jmq) = X(jmq + qtm^{s+1}), \quad \forall m \nmid j, t \in \mathbb{N}.$$

Denote $p = m^s/q$. Pick $1 \leq \ell \leq q - 1$ and set $j = \ell p$, then $jmq = \ell m^{s+1}$, so we obtain

$$X(\ell) = X(\ell m^{s+1}) = X(\ell m^{s+1} + qtm^{s+1}) = X(\ell + tq), \quad t \in \mathbb{N},$$

which means that X is almost q -periodic.

(ii) \Rightarrow (iii). Using (ii) to calculate the first m^s terms of X , we obtain (iii).

(iii) \Rightarrow (i). Set $U = X(1) \dots X(q-1)$? and $V = X(q)X(2q) \dots X((p-1)q)$?, we have $X(1) \dots X(m^s-1)$? $= U \circ V$. Therefore

$$X = [X(1) \dots X(m^s-1)]^{(\infty)} = (U \circ V)^{(\infty)} = U \circ (V \circ U)^{(\infty)},$$

which implies that $X(q\mathbb{N}) = (V \circ U)^{(\infty)} \in \mathcal{J}_m$. □

3. The case $q \nmid m$

In this section, we strengthen Theorem 2.1 by showing that if $X(q\mathbb{N})$ is a modulo- m Toeplitz fixed point, then necessarily $q|m^2$. Moreover, in the case that $q \nmid m$, we deduce that $X(1) \dots X(m)$ has a very special decomposition.

Let $X = \{X(j)\}_{j \geq 1}$ be a modulo- m Toeplitz fixed point. Let $q \geq 2$ be an integer such that $q|m^s$ for some integer $s \geq 2$, $m \nmid q$ and $q \nmid m$. Then $d = \gcd\{q, m\} > 1$. Denote $q_1 = q/d$, $m_1 = m/d$.

In what follows we always assume that the q -subsequence of X is a modulo- m Toeplitz fixed point.

Lemma 3.1. *The following hold.*

(i) X is almost q -periodic.

(ii) X is almost d -periodic.

Proof. (i) By Theorem 2.1, X is almost q -periodic, i.e.,

$$X(j) = X(j + q), \quad \forall q \nmid j.$$

(ii) Pick $j_0 \in \{1, \dots, d-1\}$. Write $d = xm - yq$ with $x, y \geq 1$. Then $X(j_0 + nd) = X(j_0 + nxm - nyq) = X(j_0)$, since X is almost q -periodic and almost m -periodic. The lemma is proved. □

Lemma 3.2. *If $\text{lcm}\{m, q\} \nmid jd$ and $\text{lcm}\{m, q\} \nmid (j+1)d$, then $X(jd) = X((j+1)d)$.*

Proof. $q \nmid m$ implies that $q_1 \geq 2$. Write $d = xm - yq$ with $x, y \geq 1$.

If $q \mid jd$, then $m \nmid jd$ and $q \nmid (j+1)d$, we have

$$(3.1) \quad X(jd) = X(jd + xm), \quad (\text{since } X \text{ is almost } m\text{-periodic})$$

and

$$(3.2) \quad X((j+1)d) = X((j+1)d + yq), \quad (\text{since } X \text{ is almost } q\text{-periodic})$$

which imply $X(jd) = X((j+1)d)$.

If $m|jd$, then $m \nmid jd + d$ and $q \nmid jd$, we have

$$(3.3) \quad X(jd) = X(jd + q) = X(jd + q + \frac{q-d}{d}yq)$$

and

$$(3.4) \quad X((j+1)d) = X((j+1)d + \frac{q-d}{d}xm),$$

which imply $X(jd) = X((j+1)d)$.

Now suppose $q \nmid jd$ and $m \nmid jd$. If $q \nmid (j+1)d$, we use (3.1) and (3.2), if $m \nmid (j+1)d$, we use (3.3) and (3.4). \square

Lemma 3.3. *The following hold.*

(i) *If jd is not a multiple of $\text{lcm}\{m, q\}$, then $X(jd) = X(d)$.*

(ii) *Let $j \in \{1, \dots, d-1\}$. If $q_1 \nmid j$, then $X(j) = X(d)$.*

Proof. (i) We prove the assertion by induction on j . Suppose $\text{lcm}\{m, q\} \nmid jd$.

If $\text{lcm}\{m, q\} \nmid (j-1)d$, then by Lemma 3.2, $X(jd) = X((j-1)d) = X(d)$, where the last equality is due to the induction hypothesis.

Now suppose $\text{lcm}\{m, q\} \mid (j-1)d$. Write $(j-1)d = xq = ym$.

If $q \nmid jd$, then $X(jd) = X(jd - xq) = X(d)$, since X is almost q -periodic.

If $m \nmid jd$, then $X(jd) = X(jd - ym) = X(d)$, since X is almost m -periodic. Item (i) is proved.

(ii) Clearly $X(j) = X(jm)$. That $q_1 \nmid j$ implies that $q \nmid jm$, hence, by item (i), $X(jm) = X(d)$. \square

By the following result, to check whether $X(q\mathbb{N})$ is modulo- m Teoplitz fixed point, we just need to examine the word $X(1) \dots X(m)$.

Theorem 3.1. *Let $X \in \mathcal{J}_m$ and assume that it is not a constant word. Let $q \geq 2$ be an integer such that $q \nmid m$, $m \nmid q$ and $q|m^s$ for some integer $s \geq 1$. Denote $d = \text{gcd}\{m, q\}$ and $q_1 = q/d$. Denote $a = X(1)$. Then $X(q\mathbb{N}) \in \mathcal{J}_m$ if and only if $q|m^2$, q_1 is a proper factor of d and $X(1) \dots X(m-1)?$ can be decomposed into*

$$(3.5) \quad X(1) \dots X(m-1)? = (a^{q_1-1}?) \circ (X(q_1) \dots X(tq_1)?) \circ (a^{m_1-1}?) := Q \circ T \circ D.$$

where $t = d/q_1 - 1$. In this case,

$$X(q\mathbb{N}) = (D \circ T \circ Q)^{(\infty)}.$$

Proof. First we prove the necessity. By Lemma 3.3(i), we have

$$(3.6) \quad X(jd) = X(d), \quad \text{for } 1 \leq j \leq m/d.$$

Especially, we have $X(1) = X(m) = X(d)$.

By Lemma 3.1(ii), $X(1) \dots X(m)$ is almost d -periodic, which together with (3.6) imply that $X(1) \dots X(m)$ is d -periodic, i.e., $X(1) \dots X(m) = [X(1) \dots X(d)]^{m/d}$. On the other hand, by Lemma 3.3(ii) and (3.6), $X(1) \dots X(d)$ can be decomposed into

$$X(1) \dots X(d) = [a^{q_1-1}?] \circ [X(q_1) \dots X(tq_1)a],$$

where $t = d/q_1 - 1$. Therefore,

$$X(1) \dots X(m-1)? = (a^{q_1-1}?) \circ (X(q_1) \dots X(tq_1)?) \circ (a^{m_1-1}?) := Q \circ T \circ D.$$

Next, we show that $q|m^2$ and q_1 is a proper factor of d .

If $q_1 \geq d$, then by Lemma 3.3(ii), $X(j) = a$ for all $j \in \{1, \dots, d-1\}$. Since X is almost d -periodic, we obtain that $X = a^\infty$. This contradiction proves that

$$q_1 < d.$$

Suppose on the contrary that q_1 is not a proper factor of d . Pick $\ell \geq 1$ such that $\ell q_1 < d$. Let $j = \ell q_1 + d$, then q_1 is not a factor of d implies that $q_1 \nmid j$, hence jm is not a multiple of $q_1 m = mq/d$, so by Lemma 3.3(i), we have

$$X(j) = X(jm) = X(d) = a.$$

On the other hand, since X is almost d -periodic, we have $X(\ell q_1) = X(j) = a$. Therefore, for all $j \in \{1, \dots, d-1\}$, no matter j is a multiple of q_1 or not, we always have $X(j) = a$. So X is a constant word. A contradiction. Clearly $q = q_1 d$ is a factor of d^2 , and hence it is a factor of m^2 . The necessity is proved.

Now we prove the sufficiency. Suppose (3.5) holds. Using $D \circ Q = Q \circ D$, we obtain

$$\begin{aligned} X &= \lim_{k \rightarrow \infty} (Q \circ T \circ D)^{(k)} = Q \circ T \circ \lim_{k \rightarrow \infty} (D \circ Q \circ T)^{(k)} \\ &= Q \circ T \circ \lim_{k \rightarrow \infty} (Q \circ D \circ T)^{(k)} = Q \circ T \circ Q \circ \lim_{k \rightarrow \infty} (D \circ T \circ Q)^{(k)}, \end{aligned}$$

which proves that $X(q\mathbb{N}) = \lim_{k \rightarrow \infty} (D \circ T \circ Q)^{(k)}$ is a modulo- m Teoplitz fixed point. The sufficiency is proved. \square

Example 3.1. Let $m = 12$. Let $X = [aabaaaaabaa?]^{\infty}$ be a modulo-12 Toeplitz fixed point. It is seen that

$$X = [(aa?) \circ (b?) \circ (a?)]^{\infty}.$$

Clearly $X(3\mathbb{N}) = [(b?) \circ (a?) \circ (aa?)]^{\infty}$ and $X(6\mathbb{N}) = [(a?) \circ (aa?) \circ (b?)]^{\infty}$.

Let $q = 18$, then $d = \gcd(12, 18) = 6$, $q_1 = q/d = 3$ and $m_1 = m/d = 2$. and

$$X(18\mathbb{N}) = [(a?) \circ (b?) \circ (aa?)]^{\infty}.$$

Moreover, $X(q\mathbb{N}) \in \mathcal{J}_m$ if and only if $q = m^k h p$ where $k \geq 0$, $\gcd\{h, m\} = 1$ and $p \in \{1, 3, 6, 18\}$. One can show that $X(q\mathbb{N})$ coincides with one of X , $X(3\mathbb{N})$, $X(6\mathbb{N})$ and $X(18\mathbb{N})$.

Example 3.2. Let $m = 6$. Let $X = [aaaba?]^{\infty}$. Then

$$X(2\mathbb{N}) = [ababa?]^{\infty};$$

$$X(5\mathbb{N}) = [abaaa?]^{\infty}.$$

Moreover, $X(q\mathbb{N}) \in \mathcal{J}_m$ if and only if $q = m^k h p$ where $k \geq 0$, $\gcd\{h, m\} = 1$ and $p \in \{1, 2\}$; indeed, $X(q\mathbb{N})$ coincides with one of X , $X(2\mathbb{N})$ and $X(5\mathbb{N})$.

REFERENCES

- [1] M. Baake, T. Jáger and D. Lenz, *Toeplitz flows and model sets*, Bull. Lond. Math. Soc., **48** (2016), 691-698.
- [2] J. Cassaigne and J. Karhumáki, *Toeplitz words, generalized periodicity and periodically iterated morphisms*, European J. Combin., **18** (1997), 497-510.
- [3] G. R. Goodson, *Chaotic dynamics: Fractals, tilings, and substitutions*, Cambridge University Press, (2016).
- [4] L. Y. Huang, S. S. Liu and H. Rao, *Thickness, Toeplitz words, and orientation preserving Lipschitz classification of Moran sets*, Preprint, (2024).
- [5] K. Jacobs and M. Keane, *0-1-sequences of Toeplitz type*, Z. Wahrsch. Verw. Gebiete, **13** (1969), 123-131.
- [6] Q. H. Liu and Y. H. Qu, *Uniform convergence of Schrödinger cocycles over simple Toeplitz subshift*, Ann. Henri Poincaré, **12** (2011), 153-172.

- [7] Q. H. Liu and Y. H. Qu, *Uniform convergence of Schrödinger cocycles over bounded Toeplitz subshift*, Ann. Henri Poincaré, **13** (2012), 1483-1500.
- [8] H. Prodinger and F. J. Urbanek, *Infinite 0-1 sequences without long adjacent identical blocks*, Discrete Math., **24** (1981), 277-289.
- [9] Y. H. Qu, H. Rao, Z. Y. Wen and Y. M. Xue, *Maximal pattern complexity of higher dimensional words*, J. Combin. Theory Ser. A, **117** (2010), 489-506.
- [10] D. Sell, *Combinatorics of one-dimensional simple Toeplitz subshifts*, Ergodic Theory Dynam. Systems, **40** (2020), 1673-1714.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN,
430079, CHINA
Email address: shishuangliu@mails.ccnu.edu.cn

DEPARTMENT OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN,
430079, CHINA
Email address: hrao@mail.ccnu.edu.cn